



The D. H. Hill Library



North Carolina State

QK341

B684

v.4

NORTH CAROLINA STATE UNIVERSITY LIBRARIES



S02514715 P

**This book is due on the date indicated
below and is subject to an overdue fine
as posted at the Circulation Desk.**

BOISSIER, E.

FLORA ORIENTALIS

Vol. IV

Coralliflorae, Monochlamydae (1879)

Réimpression fac-similée A. ASHER & CO. 1963

OUVRAGES DE M. ED. BOISSIER

Flora Orientalis sive enumeratio plantarum in Oriente a Græcia et Aegypto ad Indiae fines hucusque observatarum. Vol. I. *Thalami floræ.* Un gros volume in-8 de 1017 pages, 1867, Fr. 25. — Vol. II. *Calycifloræ Polypetalæ.* In-8^o de 1159 pages, 1872, Fr. 25. — Vol. III. *Calycifloræ Gamopetalæ.* In-8^o de 1033 pages, 1875, Fr. 25. — Vol. IV. *Corollifloræ et Monochlamydeæ.* In-8^o de 1276 pages, 1879, Fr. 31.

Voyage botanique dans le midi de l'Espagne pendant l'année 1837. 2 vol. grand in-4^o. — Tome I. *Narration et Géographie botanique*, x et 248 pages de texte. — Tome II. *Enumération des plantes du royaume de Grenade.* — *Additions*, 757 pages, accompagnés d'un Atlas de 206 planches dessinées et gravées par Heyland et coloriées à la main, ainsi que d'un Tableau synoptique donnant la hauteur des montagnes et les régions botaniques, 1839-1845. Prix réduits, colorié, au lieu de fr. 400, Fr. 230.
— — En noir, Fr. 150.
— — Sur papier vélin, colorié, au lieu de fr. 660, Fr. 350.

Diagnosis plantarum orientalium. 1^{re} série, 13 fascicules; 2^{me} série, 6 fascicules, in 8^o. Genève 1842-59, à 3 fr. le fascicule.
Les fascicules 1, 3, 6 et 7 sont épuisés.

Plantæ Aucherianæ; fascicule I et II in 8^o, à 3 fr. le fascicule.

Description de deux nouvelles Crucifères des Alpes et du Piémont. In 4^o, avec 2 planches, Fr. 5.

Aufzæhlung der in einer Reise durch Transkaukasien und Persien gesammelten Pflanzen, von Ed. Boissier und Buhse. In-4^o avec 10 planches et 1 carte. Moscou 1860, Fr. 10.

Diagnosis plantarum Hispaniæ, par Ed. Boissier et G.-F. Reuter. In-8^o. Genève 1842, épuisé.

Pugillus plantarum novarum Africæ borealis Hispaniæque australis. In-8^o. Genevæ 1852, Fr. 3.

Centuria Euphoriarum. Genève, 1860, Fr. 1.

Icones Euphoriarum ou figures de 122 espèces du genre *Euphorbia* dessinées et gravées par HEYLAND, avec des considérations sur la classification et la distribution des plantes de ce genre, par Ed. Boissier 1 vol. grand in-folio, 24 pages de texte et 121 planches. Genève 1866, Fr. 70.

FLORA ORIENTALIS

SIVE

ENUMERATIO PLANTARUM

IN ORIENTE

A GRÆCIA ET ÆGYPTO AD INDIÆ FINES

HUCUSQUE OBSERVATARUM

AUCTORE

EDMOND BOISSIER

SOC. PHYS. GENEV., SOC. LINN. LONDIN.

Reg. Acad. Scient., Matrit., etc., Sodali.

VOLUMEN QUARTUM

COROLLIFLORÆ ET MONOCHLAMYDEÆ

GENEVÆ ET BASILEÆ

APUD H. GEORG, BIBLIOPOLAM

LUGDUNI

APUD EUMDEM, 65, RUE DE LYON

1879

GENEVÆ, EX TYPIS CAREYANIS.

FLORA ORIENTALIS.

PLANTÆ VASCULARES.

CLASS. I. DICOTYLEDONEÆ.

Subclass. III. COROLLIFLORÆ D.C.

Calyx gamosepalus sepalis plus minus inter se coalitis. Petala plerumque coalita basi a calyce distincta. Stamina saepius corollæ adnata. Ovarium plerumque liberum.

ORD. LXIX. LENTIBULARIEÆ.

(Rich. Fl. Par. I, p. 26.)

Calyx monopetalus persistens bilabiatus quinquedentatus. Corolla monopetala hypogyna irregularis bilabiata calcarata. Stamina duo basi corollæ inserta. Ovarium liberum uniloculare multiovulatum placentâ centrali liberâ, stylus unicus vel nullus. Fructus capsularis. Semina numerosa minima exalbuminosa embryone orthotropo. — Herbæ palustres vel aquaticæ.

PINGUICULA (L. Gen. 30).

Calyx quinquepartitus subbilabiatus. Corollæ bilabiata, palato barbato, tubo hiante calcarato, labio superiore bilobo inferiore longiore

trilobo, stamna bina antheris unilocularibus transverse dehiscéntibus. Stigma sessile bilobum vel abortu unilobum. Capsula erecta unilocularis valvis duabus lateralibus dehiscens. Semina numerosa elliptica rugosa. — Herbæ perennes humiles scapiferæ.

1. *P. vulgaris* (L. Sp. 25) foliis oblongis glandulis sessilibus adspersis cæterum glabris, scapis superne minute glandulosis, calycis lobis oblongis obtusis, corollæ violaceæ lobis obovato-oblongis integris calcare linearis obtuso corollâ subduplo breviore, capsulâ ovato-conicâ acutâ ♀. *P. leptoceras* Rehb. Ic. f. 171.

Hab. in pascuis humidis regionis alpinæ, Macedonia in monte Peristeri 6500' (Griseb.), Thracia in Rhodope prope Carlova (Friv.), in monte Balkan (Jankat!), Olympus Bithynus (Auch.), Pontus Lazicus prope Djimil 6000' (Bal!), Armenia prope Erzerum (Tehih!), Taurus Cilicius ad Dasch Olugh montis Bulghardagh 8000' (Ky. 356!).

Specimina Orientalia ad formam alpinam corollâ paulo majori donatam sed ad typum intermediis transeuntem et a *P. grandiflora* Lam. corollâ angustiore et calcare brevi distinctam spectant.

Ar. Geogr. Europa omnis a Britanniâ et Scandinaviâ ad Rossiam, Sibiria, America boreali-occidentalis.

2. *P. crystallina* (Fl. Græc. Prodr. I, p. 11. Fl. Græc. I, p. 8, tab. 11) foliis oblongis superne glandulis sessilibus crystallinis obsitis, scapis minute glanduloso-hirtis, calycis glanduloso-hirtuli lacinias obovato-spathulatis, corollæ extus glandulosæ lilacinæ ad faucom flavidæ lobis labii superioris basi purpureo-striati integris inferioris latioribus et longioribus retusis, fauce luteo-barbatâ, calcare linearis subconico corollâ duplo breviore ♀.

Hab. ad fontes et in muscosis madidis regionis montanae Cypri, prope Comanderia (Bauer ex Sibth.), ad Prodromo versus fontes Francorum (Ky. 765!). Fl. Maio.

Scapi bipolligares, corolla cum calcare 7 lineas longa.

3. *P. hirtiflora* (Ten. Fl. Nap. III, p. 18, tab. 201) foliis oblongis vel linearis-lingulatis glandulis sessilibus plus minus adspersis, scapis superne præsertim glanduloso-hirtis, calycis glandulosi lobis oblongis obtusis, corollæ lilacinæ vel roseæ glabrae vel glandulis sessilibus parce adspersæ lobis labii superioris integris inferioris latioribus et longioribus retusis, fauce luteo-barbatâ, calcare subulato recto vel subincurvo corollæ subequilongo, capsulâ globosâ calycem non superante ♀. *P. Megaspila* Boiss. et Heldr. MSS. — *P. Albanica* Griseb. Spicil. II, p. 9. — *P. leta* Pant. Adn. ad Fl. Here. p. 74 non differre videtur.

Hab. ad scaturigines in saxosis et ad rupium stillicidia regionis montanae et subalpinae, Græcia supra Megaspileon Achaiae loco Vathliakia dicto 4500' (Heldr!), mons Chelmos prope Stygem 6000' (Orph. Fl. Gr. exs. 107!), Parnassi regio abietina ad Dipotaino (Heldr!), Olympus Thessalus in regione inferiore

et mediæ 500'-5000' (Heldr!, Orph. Fl. Gr. exs. 719!), insula Poros (Wied!), mons Balkan Thraciæ (Janka).

A *P. vulgari* colore corollæ, hujus lobis inferioribus retusis vel subbilobis, capsulâ globosâ obtusâ egregie distincta, sed *P. crystallinæ* valde et forsitan nimis affinis a quâ differt floribus majoribus, calycis laciniis non spathulatis, calcare longiori subulato.

Ar. Geogr. Regnum Neapolitanum, Herzegovina, Albania.

UTRICULARIA (L. Gen. 34).

Calyx bipartitus lobo superiore integro inferiore saepe retuso vel bidentato. Corolla personata infra labium inferius calcarata labio superiore erecto integro vel retuso inferiore longiore sepius trilobo. Stamina bina filamentis dilatatis, antheris unilocularibus longitudinaliter dehiscentibus. Stylus (in nostris) nullus, stigma inaequaliter bilabiatum vel unilabiatum. Capsula unilocularis demum irregulariter erupta. Semina numerosa polyedra. — Herbae (in nostris) natantes foliis saltem inferioribus multisectis capillaceis utriculiferis.

SECT. I. MEGACISTA Alph. p.c. Prodr. VIII, p. 3. — Folia superiora in verticillum unicun (involutum) disposita petiolo inflato vesicæformi natantia, inferiora multisecta utriculifera. Radices paucæ vel nullæ non adfixæ.

1. U. inflexa (Forsk. Deser. p. 9) foliis verticillatis a basi bipinnatisectis oblongo-acuminatis axi dilatata linearis segmentis pilosiusculis capillaceis prope basin utriculiferis, involuci infra medium scapi siti vesicis verticillatis oblongis utrinque attenuatis apice setiferis, scapo erecto multifloro, pedicellis bracteolâ longioribus calyce subbrevioribus, lobis calycinis ovatis obtusis, corollæ albae violaceostriatae labio superiore retuso, calcare obtuso sursum flexo labio inferiore sublongiore ♀. Ic. Del. Fl. Eg. tab. 4. — *U. stellaris* Willd. I, p. 143 ex parte.

Hab. in Ægypti inferioris aquis et oryzetis ad Damiette (Del! Ehr! Sieb!), ad Bena el Assal prope Cahiram (Schweinf!). Vulgo *Hamoud*.

Ar. Geogr. Nubia, Kordofan, Guinea, Senegalia.

2. U. stellaris (Linn. fil. Suppl. p. 86) foliis verticillatis a basi bipinnatisectis oblongis axi non dilatata segmentis pilosiusculis capillaceis prope basin utriculiferis, involuci ad medium scapi siti vesicis verticillatis ovoideo-acutis apice setiferis, scapo multifloro, pedicellis bracteolâ longioribus calyce subbrevioribus, lobis calycinis ovatis obtusis, corollæ flavæ basi albidae calyce duplo longioris labio superiore oblongo integro, calcare saccato obtuso labium inferius non æquante ♀. Ic. Wight III. p. 47, tab. 27.

Hab. in aquis Ægypti (Siebt!)

A præcedente colore corollæ et aliis notis in Diagnosi indicatis sed iterum ex vivo investigandis distincta.

Ar. Geogr. Nubia, Africa tropica, Senegalia, India.

SECT. II. LENTIBULARIA Alph. p.c. loc. cit. p. 4. — Folia omnia multisecta capillacea lateraliter utriculifera. Rædices nullæ vel paucæ non adfixæ.

3. **U. vulgaris** (L. Sp. 26) foliis undique patentibus pinnato-multipartitis laciñiis capillaribus remote subtiliter spinosis. seapo multifloro laxo, pedicellis bracteolâ et calyce longioribus. corollæ vitellinæ labio superiore palatum inflatum aurantiaco-striatum subæquante, calcare oblongo-conico descendente, antheris connatis ♀. Ic. Fl. Dan. tab. 138.

Hab. in aquis Graeciae (Auch. 2703!), Argolidis (Sibth.) et verosimiliter alibi.

Ar. Geogr. Europa omnis a Scotiâ et Sueciâ mediâ ad Rossiam septentrionalem, Sibiria, Africa borealis, America borealis.

ORD. LXX. PRIMULACEÆ.

(Vent. tab. 2. p. 285.)

Calyx monopetalus persistens quinque (rarius 4)-dentatus. Corolla monopetala hypogyna regularis quinque (rarius 4)-tida. Stamina corollæ inserta ejus lobis numero æqualia et iisdem opposita vel numero dupla, exteriora laciñiis corollæ alterna sterilia. Ovarium liberum uniculare pluriovulatum placentâ centrali liberâ, stylus I. Capsula varia dehiscens. Semina sæpius peltata, embryo rectus in centro albuminis carnosus. — Herbæ.

TR. I. SAMOLEÆ Endl. Gen. p. 734.

Capsula semisupera valvatum dehiscens. Semina ampliotropa umbilico basiliari. Embryo transversus nempe umbilico parallelus.

SAMOLUS (L. Gen. 222).

Calyx semisuperus persistens quinquefidus. Corolla perigyna tubo brevi limbo patentí quinquepartito. Stamina quinque fertilia corollæ tubo inserta ejusque lobis opposita, quinque sterilia sinibus inserta. Capsula semiquinquevalvis. — Herbæ.

1. S. Valerandi (L. Sp. 243) perennis, glaucescens, caule erecto simplici vel ramoso, foliis obovatis vel oblongo-spathulatis obtusis inferioribus in petiolum attenuatis, racemo unico vel pluribus paniculatis demum elongatis aphyllis, pedicellis setaceis erecto-patulis supra medium geniculato-incurvis et bracteolâ lanceolatâ obsitis, calycis dentibus erectis late ovatis, corollâ albâ calyce sublongiore ♀. Ic. Lam. III. tab. 101. — Rehb. t. 42.

Hab. ad fontes, stillicidia, in uliginosis subsalsis totius ditionis a Græciâ et ejus insulis ad Transcaucasiam! Ægyptum! Persiam australem (Ky!), Affghaniam! et Belutschiam (Stocks!).

Ar. Geogr. Europa omnis arcticas exceptis, Africa borealis, Sibiria, India, Caput bonæ spei, Nov. Hollandia, America.

TR. II. HOTTONIEÆ Endl. Gen. p. 734.

Capsula supera valvatim dehiscens. Semina anatropa erecta umbilico basilari.

HOTTONIA (L. Gen. 203).

Calyx quinquepartitus. Corolla hypocrateriformis tubo brevi, limbo planiusculo lobis 5 basi breviter glandulosis. Stamina 5 subsessilia inclusa. Capsula globosa demum mediis suturis dehiscens apice et basi cohærens.— Plantæ aquatice, foliis submersis verticillatis pectinatum pinnatifidis vel sectis, scapo emerso, floribus verticillato-racemosis.

1. H. palustris (L. Sp. 208), foliis approximatis in lacinias anguste lineares acutas elongatas pinnatisectis, floribus in verticillos remotos dispositis, bracteis linearibus pedicellis longis multo brevioribus, sepalis linear-lanceolatis obtusiusculis, corollâ pallide violacea calyce multo longiore laciñis oblongis subretusis ♀. Ic. Engl. bot. tab. 364.

Hab. in fossis aquâ profundâ repletis, in Bithyniâ inter Sabandja et Boli (Wied!) et prob. alibi.

Ar. Geogr. Europa sept. et media ab Angliâ et Scandinaviâ, Galliâ, Italîâ media ad regionem Danubialem, Rossiam medium et australem, Sibiriam Uralensem.

TR. III. ANAGALLIDEÆ Endl. loc. cit.

Capsula supera operculo circumscisso dehiscens. Semina amphitropa.

ANAGALLIS (L. Gen. 206).

Calyx quinquepartitus. Corolla rotata decidua quinquepartita. Stamina 5 filamentis barbatis. Capsula supera globosa circumscissa membranacea polysperma. Semina angulosa. — Herbae annuae et perennes.

1. **A. arvensis** (L. Sp. 211) annua glabra, caulis diffusis vel ascendentibus quadrangulis ramosis, foliis oppositis vel ternatis sessilibus ovatis et triangulari-ovatis, pedicellis axillaribus solitariis folia subæquantibus fructiferis reflexis, corollæ calycis lacinias acuminatas margine membranaceas parum excedente ⊖. Ic. Rehb. tab. 41.

α. phænica. — Corollæ rubræ marginibus sæpius glandulosociliatae. — *A. phanicea* Lam. Fl. Fr. II. p. 285.

β. caerulea. — Corollæ caeruleæ segmentis sæpius eglandulosis. — *A. cærulea* Lam. loc. cit.

Hab. utraque varietas in cultis regionis inferioris et montanae totius ditionis usque ad Ægyptum! Persiam australem in palmetis (Bge!), Belutschiam (Stockst.).

Ar. Geogr. Orbis totus arcticis et antarcticis exceptis.

2. **A. latifolia** (L. Sp. 212) annua vel perennans glabra, caulis diffusis compressis, foliis oppositis late ovatis cordato-amplexicanilibus obtusiusculis, pedicellis axillaribus solitariis folia subæquantibus fructiferis reflexis, corollæ caeruleæ segmentis glanduloso-ciliatis calycis lacinias setaceo-acuminatas superantibus ⊖.

Hab. in cultis præsertim regionum calidiorum, Zaconthus (Margt!), Peloponnesus (Bory!), Byzantii (Griseb!), Mesopotamia (Haussk!), Ægyptus inferior (Wiest! Samar!), Arabia petrea (Boiss!) et Mascatensis (Auch. 5228!).

Omnibus partibus *A. arvensi* major, an ab eâ sat distincta?

Ar. Geogr. Lusitania, Hispania australis, Africa borealis.

3. **A. tenella** (L. Mant. 335) perennis, glabra, caulis filiformibus basi repentinibus radicantibus, foliis oppositis parvis subpetiolatis orbiculatis, pedicellis in axillis supremis elongatis folia multo superantibus fructiferis reflexis, calycis laciñiis linear-lanceolatis subulatis margine non membranaceis, corollæ roseæ calyce duplo longioris segmentis linear-oblongis integris ♀. Ic. Fl. Dan. t. 1085. — Rehb. tab. 41. — *Jiraseckia tenella* Rehb.

Hab. in uliginosis et ad fontes regionis inferioris et collinæ Cretæ ad Enneachoria, Platania, monte Cofino prov. Messara (Sibth., Sieb., Heldr! Raul!).

Ar. Geogr. Anglia, Batavia, Germania occidentalis, Gallia, Helvetia, Lusitania et Hispania, Italia bor. et centralis, Africa borealis.

TR. IV. PRIMULEÆ Endl. loc. cit.

Capsula supera superne tantum vel longitudine totâ valvis dehiscentia.

SUBTR. I. LYSIMACHIEÆ. — Flores axillares.

GLAUX (L. Gen. 294).

Calyx campanulatus quinquefidus petaloideus. Corolla nulla. Stamina 5 hypogyna imo calyxi inserta et ejus laciniis alterna. Capsula globosa quinquevalvis oligosperma. — Herba perennis pumila fioribus axillaribus solitariis subsessilibus.

I. **G. maritima** (L. Sp. 301) glabra glauca. caulis e rhizomate crassiusculo pluribus nanis simplicibus fere a basi opposite et subdecussatim foliosis, foliis parvis carnosulis ovatis vel breviter elliptico-lanceolatis obtusis integris, floribus axillaribus racemum foliatum formantibus, calycis albo-rosei laciniis ovatis obtusis ♀. Ic. Lam. Ill. tab. 141. — Rehb. Ic. tab. 76.

Hab. in littoribus maritimis et salsuginosis interioribus etiam subalpinis, Anatolia interior in subalpinis (Noë!), paludes salsa Cappadocia ad Cæsaream (Bal. 1038!), Armenia ad Erzerum (Huett!), littora Pontica (Nordm.), et Caspia (C.A. Mey!). Persia borealis (Auch. 5229!) et australis ad fontes alpis Kuh Daena (Ky. 747!).

Ar. Geogr. Europa borealis et media ab Anglia et Scandinaviâ ad Rossiam medianam et australem et ab Hispania et Italiâ borealibus ad Dalmatiâ et regionem Danubialem, Sibiria omnis, China borealis et Japonia, America borealis maritima et interior.

LYSIMACHIA (L. Gen. 203).

Calyx quinquefidus vel partitus. Corolla quinquepartita campanulata vel rotata. Stamina (in nostris) semper 5 libera vel basi monadelpha. Capsula (in nostris) quinquevalvis. Semina levigata numerosa. — Herbae annuae vel perennes.

SECT. I. EPHEMERUM Duby. — Flores racemosi. Sinus inter corollæ laciniis edentuli. Filamenta ad basin lobi orum corollæ inserta non monadelpha. Flores albi vel rosei.

I. **L. atropurpurea** (L. Sp. 209) annua, caule erecto simplici stricte et cærebre folioso in spicam densifloram abeunti, foliis alternis minute glandulosis lanceolatis basi attenuatis imarginatis subpeti-

latis, bracteis linearis-subulatis calyce brevioribus interdum obsoletis, floribus sessilibus, calycis campanulati corollâ duplo brevioris laciniis late linearibus obtusis, corollæ rubicundæ segmentis erectis oblongis acutiusculis, staminibus exsertis, antheris atropurpureis, capsulâ a basi rumpente demum quinquevalvi ♂. Ic. Fl. Græc. tab. 187. — *Palladia atropurpurea* J. et Sp. Ill. t. 423.

Hab. in cultis derelictis et subumbrosis siccis Græcia et Anatolia, Peloponnesus (Chaub.), Taygeti radices prope Anavryti et Parnassus prope Rachova (Heldr!). Macedonia et Thracia (Friv!), Bithynia (Pest!). Lydia circa Smyrnam copiose (Boiss!), Cilicie orientalis ditio Kassan Oglu (Ky. 118!). Ex Iberia erronee indicata.

2. L. dubia (Ait. Kew I, p. 199) annua, caule erecto elato crebre foliato in racemum longum simplicem vel in paniculam longe ramosam abeunti, foliis alternis glabriusculis oblongis acutis basi attenuatis inferioribus breviter petiolatis, floribus pedicello stricto calyci æquilongo vel sublongiore suffultis laxiusculis, bracteis linearibus pedicellis subæquantibus, calycis laciniis elliptico-lanceolatis acutiusculis, corollæ rubello-albæ calyce subduplo longioris laciniis obovatis obtusis erecto-patulis. staminibus inclusis, capsulâ sero quinquevalvi ♂. Ic. Fl. Gr. tab. 188. — *L. atropurpurea* Murr. Comm. Gött. 1782, tab. 1 non L.

Hab. in dumetis humidis et ad fossas regionis inferioris, Eubœa septentrionalis (Wied!), Thracia (Friv! Janka!), Bithynia (Sibth.), Galatia et Paphlagonia (Wied!), Lydia ad Smyrnam (Heldr!), Caria (Boiss!), Phrygia (Boiss!), Cilicia (Bal!), Syria littoralis (Bl! Gaill!), Mesopotamia (Ky!), Pontus Lazicus maritimus ad Rhizé (Bal!), Transcaucasia (Hoh!), Persia bor. (Buhse!).

SECT. II. LYSIMASTRUM Endl. — Flores axillares paniculati, verticillati vel solitarii, sinus inter corollæ laciniæ edentuli. Stamina filamentis inferne monadelpha. Flores lutei.

3. L. vulgaris (L. Sp. 209) perennis, caule erecto elato, foliis breviter petiolatis et superioribus sessilibus oppositis vel 3-4-nis ovatis vel oblongo-lanceolatis subtus pubescentibus acutis, florum racemis terminalibus brevibus in paniculam dispositis, corollæ patentis flavæ laciniis ovato-oblongis obtusiusculis eglandulosis, staminibus 5 filamentis usque ad tertiam partem inferiorem connatis ovarium tegentibus ♀. Fl. Dan. tab. 689. — Rehb. t. 45.

Hab. in paludosis et ad fluviorum ripas regionis montanae et subalpinæ, Thracia circa Byzantium et in Olympo Bithynia (Sibth.). Anatolia in Lycaonia (Heldr!), monte Berytdagh Cataoniae 6000' (Haussk!), prov. Musch Armeniæ australis 5600' (Ky!), Tauria, Caucaso et Transcaucasiæ (Led.).

β. stenophylla. — Folia lanceolata. — *L. Dahirica* (Ledeb.).

Hab. ad Vodena Macedonia (Orph!), circa Byzantium (Auch. 2602!).

Ar. Geogr. Europa borealis et media a Scandinavia ad Rossiam arcticam et

ab Hispaniâ et Italiâ centralibus, Dalmatiâ, ad regionem Danubialem, Sibiria, Japonia, Africa borealis.

4. **L. punctata** (Jacq. Austr. t. 366) perennis, tota crispile papilloso-hirta et glandulosa, caule erecto simplici vel paree paniculato-ranioso, foliis oppositis 2-3-4-nis subsessilibus ovato-lanceolatis acuminatis, racemi terminalis verticillastris foliis floralibus superatis, corollæ patentis flavæ laciniis ovatis acutis glanduloso-ciliatis, filamentis 5 ad medium connatis ♀. Ic. Rehb. tab. 43.

Hab. in umbrosis humidis regionis montanæ, Macedonia in monte Korthiati et supra Bitolia 3-4000' (Heldr!, Orph. Fl. Gr. exs. 508!), Olympus Bithynus, Tmolus et alii Lydie montes in regione mediâ (Boiss!), Pisidia ad lacum Egirdir (Heldr!), Paphlagonia et Galatia (Wied!), Caucasus (herb. Petrop!), Transcaucasia (Szov!).

P. verticillata. — Folia ovato-oblonga longius petiolata obtusiora, verticillastri saepe magis multiflori floribus longius pedicellatis folia floralia saepe excedentibus. — *P. verticillata* M.B. Taur. Cauc. I, p. 641.

— Cent. Ross. tab. 32. Intermediis in typum transit.

Hab. in Anatoliâ bor. (Wied!), Armeniâ Turcicâ (Calv!), Ponto Lazico (Bal!), Tauriâ omni, Caucaso et Transcancasiam (M.B! Hoh! Rupr!). Hæc forma in parte magis orientali ditionis typo vulgatori.

Ar. Geogr. Belgium, Germania australis, Italia media, Dalmatia, regio Danubialis, Rossia media et australis.

5. **L. nummularia** (L. Sp. 211) perennis, caulis humi prostratis radicantibus, foliis glabris oppositis ovato-subrotundis brevissime petiolatis, pedunculis axillaribus solitariis folio sæpius brevioribus, calycis laciniis cordato-ovatis, corollâ luteâ calyce 2-3-plo longiore patenti segmentis ovatis punctatis et glandulosis, staminibus basi inter se breviter coalitis ab ovario discretis ♀. Ic. Fl. Dan. 539.

— Rehb. t. 43.

Hab. in pratis humidis præsertim montanis, mons Korthiati Macedoniæ 4000' (Heldr!), Byzantii ad sylvam Belgrad (Counany!), Tauria et Iberia (M.B.), Caucasus bor. orientalis (Owerin!).

Ar. Geogr. Europa borealis et media ab Angliâ et Sueciâ ad Rossiam medianam et australem, Hispaniâ ad Italiam centralem, Dalmatiam et regionem Danubialem.

SECT. III. LEROUXIA. — Flores lutei axillares solitarii vel terminales. Sinus inter corollæ laciniias edentuli. Filamenta libera.

6. **L. anagalloides** (Sibth. Fl. Græc. II, p. 74, tab. 190), perennis glabra pluriceps, caulis filiformibus basi nudis induratis procumbentibus vel ascendentibus superne laxe foliatis flexuosis, foliis parvis oppositis subsessilibus et sessilibus ovatis obtusis, pedicellis tenuibus folia 3-4-plo superantibus fructiferis incurvis, floribus minutis, calyce ad basin in laciniias angustè elliptico-lanceolatas acutas

partito, corollâ rotatâ calyce paulo longiore segmentis ovatis obtusis integris ♀.

Hab. inter saxa et ad frutices Cretæ in monte Cavoutsi prov. Sitia, in monte Idâ et in Sphatioticis 3000'-5000' (Sibth., Sieb! Heldr!), Taygeto (Bory), monte Kyllene Arcadiæ (Orph!), Parnasso et Delphi Eubeæ in regione alpinâ (Heldr!), insulâ Poros (Wied!).

Affinis *L. nemorum* sed caules parte inferiori indurati fere suffruticosi, folia quadruplo minorâ, flores submajores, calycis lacinie sublatiores non acuminatae, petala non denticulatae.

ASTEROLINUM (Link et Hoffm. Fl. Portug. 332).

Calyx quinquepartitus. Corolla calyce 3-4-plo brevior tubo brevi limbo quinquepartito campanulato. Stamina quinque filamentis corollâ longioribus. Capsula calyce è corollâ persistentibus involucrata quinquevalvis 2-3-sperma. Semina dorso rugosa ventre convexo umbilicata.— Herbula pusilla annua.

1. A. ***Linum stellatum*** (L. Sp. 211 sub *Lysimachia*) nanum simplex vel a collo ramosissimum ramis erectis filiformibus foliosis, foliis minimis sessilibus oppositis breviter lanceolatis acutis, pedicellis solitariis axillaribus folio brevioribus demum cernuis laxe racemosis, calycis laciniis stellatim patentibus lanceolato-linearibus aristatis capsulam multo superantibus ⊖. *A. Linum stellatum* L. et Hoffm. I. cit.

Hab. in rupestribus Peloponnesi (Sibth!), Atticae in Lykabeto (Heldr!), Anatoliae (Auch. 2594!), Cretæ (Sieb!), Lydiae (Fleisch!), Cypri (Sibth. i, Cilicie litt. (Bal!), Syria littoralis ad Sidonem (Bl!), Palestinae (Boiss!), Syria ad Aleppo et Mesopotamie (Haussk!). Transcaucasiæ ad Baku (C.A.M.), ad Schachbulagh (Szov!), Persiae australis ad Gere (Ky!).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, Dalmatia, Africa borealis.

SUBTR. II. ANDROSACEÆ. — Flores terminales. Folia rosulata.

CYCLAMEN (L. Gen. 204).

Calyx campanulatus quinquepartitus. Corollæ tubus ovatus brevis limbus quinquepartitus reflexus. Stamina imo tubo inserta, antheræ inclusæ sessiles cuspidatae. Capsula subcarnosa globosa apice primum demum totâ longitudine quinquevalvis.— Plantæ herbaceæ radice tuberiforni in unâquaque specie si ad sumam soli superficiem sitâ ex apice absque rhizomate folia floresque emitte. si autem humo profundius sepultâ rhizoma cylindricum tenui plus minus elongatum ex quo pedunculi et petioli prodeunt edente. — Genus inter omnia seminibus germinatione monocotyledoneis insigne.

* Corollæ faux ob petala ubi reflectuntur rectilinea edentula.

1. C. Europæum (L. Sp. 207) autumnale, tubere globoso sape depresso, foliis synanthiis ovatis basi sinu acuto profunde cordatis obtusiusculis stepe apiculatis non angulatis sed plus minus crenatis crenis muticis pedunculis denum spiraliter convolutis, calycis laciniis ovatis acutis denticulato-glandulosis, corollæ roseæ fauce edentulâ laminae laciniis oblongis acutiusculis tubo 3-4-plo longioribus ♀. Ic. Lam. III. tab. 100. — Rehb. Ic. tab. 48. — *C. astirum* Park.

Hab. prob. in Macedoniâ et Thraciâ horeali sed specimina e ditione nondum vidi. Specimina omnia e regionibus Caucasicis et Libano citata ad *C. Coum* spectant!

Ar. Geogr. Gallia, Helvetia et Germania ad radices Jurassi et Alpium, Italia borealis, regio Danubialis, Serbia.

2. C. Coum (Mill. Dict. № 6) vernale, tubere orbiculato patellari, foliis synanthiis orbiculatis vel reniformi-rotundatis sinu acuto profunde cordatis exangulatis integris vel obtuse crenato-repandis, pedunculis denum spiraliter convolutis, calycis laciniis oblongo-lanceolatis vel breviter lanceolatis acutissimis, corollæ roseæ fauce edentulâ, laminae laciniis ovatis obtusis tubo duplo longioribus ♀. Ic. Bot. Mag. tab. 4. — *C. vernum* Sweet Flow. Gard. tab. 9 non Loh.

Hab. in umbrosis rupestribus regionis montanae subalpinae et etiam alpine Thracie orientalis, Anatoliae, Syriae, Armeniae, et regionum Transcaucasicarum, in Thraciâ prope Byzantium (Auch! Duparq!), Bithyniâ (Pest!), monte Aladagh (Wied!), Lyciâ in monte Solyma (Heldr!), Armenia ad Erzerum et Karakaban (Huet!), Ponto Lazico in monte Tcharantach 7000' (Bal!), Syriae monte Amano supra Beilan (Haussk!), Libano (Lab., Post!), Imeretiâ ad Kutais (Rupr!), Iberiâ prope Tiflis 4-4500' (Hoh! Ruprecht!). Persia bor.-orientali ad Asterabad et Siaret (Bge!). Fl. in demissoribus Jan. Mart. et in excelsioribus usque ad Junium.

♂. Ibericum. — Folia ovata acutiuscula, corolla submajore. — *C. Caucasicum* W. herb. — *C. Ibericum* Stev. in herb Berol. — *C. Europæum* M.B. Taur. Cauc. — C.A. Mey. Enum. Fl. Pelop. — Sm. Prodr. non L. — *C. vernale* C. Koch Linn. XVII, p. 308. — *C. elegans* Boiss. et Buhse Aufz. p. 145.

Hab. in Transcaucasia (M.B. Hoh!), prope Tiflis (Haussk!), in ditione Talyssch (C.A. Mey!). in Persia boreali prope Asterabad (Buhse!), inter Rustamabad et Rescht (Haussk!).

A formâ typicâ parum diversum et intermediis transiens.

3. C. Cilicium (Boiss. et Heldr. Diagn. Ser. I. 11, p. 78) autumnale, tubere depresso, foliis synanthiis vel subhysteranthiis orbiculatis sinu brevi clauso cordatis obtusis integris vel obsolete repandis, petiolis pedunculisque longis minute glandulosis hisce denum spiraliter convolutis, calycis glandulosi laciniis lanceolatis acutis, corollæ parvæ roseæ fauce edentulâ laminae laciniis ellipticis acutis mucronatis basi attenuatis tubo duplo longioribus ♀.

Hab. in Ciliciâ ad Pylas Gülek Boghas (Heldr! Bal. 692!). Fl. Oct.

Facies *C. Coi* a quo differt florescentiâ autumnali, corollæ laciniis angustioribus mucronatis; folia parva 1-1 $\frac{1}{2}$ poll. diam. lata.

4. *C. latifolium* (Sibth. et Sm. Fl. Græc. II, p. 71, t. 185) hemiale et verna, tubere magno depresso, foliis synanthiis magnis ovatis acutiusculis basi sinu subaperto cordatis exangulatis circumcirca crebre et inæqualiter denticulato-crenatis, pedunculis minute glandulosis fructiferis non circinnatis, floribus majusculis, calycis laciniis ovatis acutiusculis, corollæ roseæ fauce edentulâ laciniis longe linear-ellipticis tubo quintuplo longioribus ♀. *C. hederifolium* Sm. Prodr. I, p. 270 non Ait. — *C. Aleppicum* Fisch. ubi? — *C. Persicum* Mill. (forma hortensis).

Hab. in umbrosis Græciæ (Sibth.) sed e Græciâ continentali nondum vidi, Macedonia in regione sempervirenti peninsulæ Athos (Griseb!), insulæ Chios (Orph!) et Rhodi (Heldr!), Ciliciæ (Auch. 2613!), Cypri (Ky!), totius Ciliciæ littoralis et Libani inferioris usque ad Eden (Bl! Cadet de Fontenay!). Palestina ad Hierosolymam Boiss!), Nazareth (Gaill!). Floret a Januario ad Maium.

Species foliis magnis indivisis sed multidenticulatis, corollæ laciniis elongatis, pedunculis fructiferis non spiralibus insignis. Hujus forma hortensis a me nunquam spontanea visa, foliis obtusioribus, corollæ albæ fauce roseæ laciniis brevioribus est *C. Persicum* Mill. Nec typus nec varietas in Persia crescent.

5. *C. repandum* (Sibth. Fl. Græc. II, p. 72, tab. 186) verna, tubere parvo saepè depresso, foliis synanthiis ovato-triangularibus acutiusculis basi sinu aperto cordatis angulato-repandis angulis et crenis latis obtusis mucronulatis, pedunculis minute glandulosis fructiferis circinnatis, calycis laciniis ovatis acuminatis, corollæ roseæ fauce edentulâ laciniis longe linear-ellipticis tubo quintuplo longioribus supra basin attenuatis ♀. *C. vernum* Lob. Ic. — *C. hederæfölium* Ait. Kew I, p. 196. — *C. Romanum* Gris. Spic. I, p. 5.

Hab. in sylvaticis regionis inferioris et subalpinæ Peloponnesi in Messeniâ et Laconiâ (Boiss!) usque ad 6000' in montibus Taygeto (Heldr!), Malevo et Chelmos (Orph. Fl. Gr. exs. 509!), Cretâ (Heldr!), Byzantio (Sibth.).

Ar. Geogr. Baleares, Gallia australis, Corsica, Sardinia, Italia meridionalis, Sicilia, Istria, Dalmatia.

* * Corollæ faux ob petala ubi reflectuntur lunulata corniculatum dentata.

6. *C. Neapolitanum* (Ten. Nap. III, p. 197) autumnale, tubere crasso nigro discoideo-depresso circumcircata crebre et tenuiter fibrilifero, foliis subhysteranthiis ovato-triangularibus basi sinu plus minus aperto cordatis 5-7-angulato-lobatis lobis muticis margine præterea minute crenulatis, pedunculis fructiferis circinnatis, floribus majusculis, calycis laciniis ovatis abrupte acuminatis, corollæ fauce pur-

pureo-maculatæ corniculato-10-dentatæ, laciñis albidis vel carneis oblongo-ellipticis obtusis tubo triplo longioribus ♀. Bot. Reg. tab. 49. — *C. hederaefolium* Willd. Koch. Gaud. non Ait. — Rehb. Ic. tab. 47. — *C. sicariifolium* Desmoulins. — *C. Cyprium* Kotschy Cypern p. 293.

Hab. in umbrosis rupestribus regionis montane Græciæ, Messenia circa Navarin (herb. Fauché!), Eubea (Sart!), regio media Parnethos Attice, vallis Cephissi et pineta Olympi Thessali prope Hagio Dionysi (Heldr!). Byzantii ad Buyukdere (Duparq!). Cypri (Ky!). Fl. Sept. Dec.

Ar. Geogr. Helvetia australis, Gallia australis, Corsica, Italia media et australis, Dalmatia, Serbia.

7. C. Græcum (Link in Linn. IX. p. 573) autunnale, tubere magno laterito ovato-globoso tandem irregulari basi tantum fibras radicales paucas crassiusculas edente, foliis subhysteranthiis rotundato-ovatis acutiusculis basi sinu aperto cordatis exangulatis circumcirca crebre inæqualiter et obtuse crenulato-denticulatis subtus rubris, pedunculis fructiferis circinnatis, floribus majusculis, calycis laciñis breviter lanceolatis acutis, corollæ roseæ fauce corniculato-10-dentatæ, laciñis elliptico-linearibus obtusis tubo quintuplo longioribus ♀. *C. Persicum* Sm. Prodr. I, p. 128 non Mill.

Hab. in collibus saxosis et sylvaticis partis inferioris montium Græciae 500'-1500', Peloponnesus circa Naupliam (Link), Messenia (herb. Fauché!), Attica in Hymetto, Lycaabeto et Parnes (Sprunt, Heldr!). Cretæ (Noë! Raulin!), Macedonia in peninsulâ Athoâ (Friv! Orphan!). Fl. Sept. oct.

A *C. Neapolitano* egregie distinctum radice, foliis, calycis laciñis angustioribus non abrupte cuspidatis; dies quindecim ocius floret.

ANDROSACE (L. Gen. 196).

Calyx quinquelobus. Corolla hypocateriformis tubo brevi, fauce coronulâ parvâ vel plicis coaretatæ. Capsula saepius oligosperma fere ad basin usque quinquevalvis. — Plantæ herbaceæ perennes vel monocarpicæ saepius alpineæ.

* Perennes.

1. A. villosa (L. Sp. 203) cespitosa caudiculis filiformibus fragilibus infractis, foliis parvis in rosulas apertas vel clausas congestis brevibus oblongo-vel linear-lanceolatis pilis albis obscure articulatis flexuosis villosis integris, scapo villoso sub anthesi umbellæ paucifloræ æquilongo vel longiore, pedicellis brevissimis, involueri phyllis ellipticis calyce villoso subbrevioribus, corollæ albæ limbo calyce subbreviore laciñis obovatis ♀. Ic. Jacq. Coll. tab. 12. — Rehb. tab. 71. — *A. arachnoidea* et *A. penicillata* Schott. — *A. incana* Lam. — *A. Juncquemontii* Daby in p.c. Prodr.

Hab. in rupestribus siccis regionis alpineæ, Pontus Lazicus supra Khabackar

7500' (Bal!), Armenia ad Bayazid (Auch. 5232!), Libani cacumina (Auch. 2608! Boiss! Bl!), Caucasus centralis 2500'-3500' et Orientalis 6000'-10000' (Rupr!), Cauc. occidentalis (Radde!), Tauria (M.B.), mons Demavend Persiae bor 9500' (Ky. 350!).

β. congesta. — Densius pulvinaris, folia diminuta linearia subtus carinata erectiora rosulas saepius clausas formantia, scapi abbreviati saepe subnulli, umbella ad 2 vel florem unicum reducta. — *A. dasypylla* Bunge. — *A. Olympia* Boiss. Diagn. Ser. I, 3, p. 37. — *A. odoratissima* Schreb. Dec. p. 7, tab. 4 (forma glabrescens). — *A. Orientalis foliis Coridis flore odoratissimo* Tourn. herb!

Hab. in regione alpinâ, Olympus Bithynus (Sibth., Auch. 2641!, Boiss! Pichler!), mons Davros Pisidiæ (Heldr!), Argeus Cappadociae 8500'-9000' (Bal! Ky. 204!), Taurus Cilicicus 8700' (Ky! Bal!), mons Techdagh Armeniae 10-11000' (Huet!), Caucasus in Daghestanâ (Rupr!), Affghaniae alpes ad Koh i Baba 15000' (Griff!).

Hac forma primâ facie a typo diversissima ad eum intermediis transit.

Ar. Geogr. Montes Hispaniae borealis, Pyrenei, Alpes, Jurassus, Italia centralis, Dalmatia, Transylvania, Bosnia, Songaria, Sibiria omnis, regio Himalaica.

2 A. chamæjasme (Host Syn. 95) cespitosa caudiculis filiformibus, foliis parvis ovato-oblongis in rosulas explanatas congestis viridibus glabris pilis moniliformibus ciliatis integris, scapo hirto, umbellâ pauciflorâ, involuci phyllis linearî-spathulatis pedicellos æquantibus, corollæ albæ limbo calyci viridi hirto æquilongo laciñiis obovatis ♀. *A. villosa* Jacq. Austr. tab. 352 non L. — *A. villosa* var. *latifolia* Fl. Alt. — *A. Lehmanniana* Spreng. Iris 1817, p. 1289, tab. 9 non Wall. — *A. longiscapa* C. Koch Linn. XXIII, p. 614!

Hab. in regione alpinâ Caucasi orientalis in alpe Schahdag (M.B.), centralis 6600' et orientalis in Tuschetia et Salatavia 8000'-10000' (Rupr!), in monte Kasbek 800' (Rehmann!), in Caucaso occidentali (C.A. Meyer).

Specimina Alpinis simillima.

Ar. Geogr. Alpes centrales ab Helvetia ad Austriam, Rossia arctica, Sibiriae totius alpes, America borealis.

3. A. hedreantha (Griseb. Spicil. II, p. 3) foliis expansis parvis lingulato-oblongis obtusis integris vel obsolete 1-2-dentato-repandis glabriusculis sed margine præsertim versus basin attenuatam breviter papillari-ciliatis, scapo foliis non longiori brevissime cinereo-tomentoso, umbellâ capituliformi, floribus paucis brevissime pedicellatis, involuci phyllis glabris oblongis acutiusculis dimidios calyces æquantibus, calycis glabriusculi dentibus oblongis acutiusculis tubo campanulato æquilongis, corollæ roseæ tubo calycem vix æquante lobis obovatis tubo æquilongis ♀.

Hab. in regione alpinâ montis Rilo jugi Orbelus Thraciae (Frider!). Vid. in herb. cl. Griseb. .

An cespitosa?, vidi tantum rosulas binas solitarias, folia 4 lin. longa 1 $\frac{1}{2}$ lata, scapus 4-5 lineas longus, calyx 1 $\frac{1}{2}$ lin. longus. Valde affinis *A. Wulfenii*.

nianæ Sieb. a quā differre videtur foliis majoribus præter papillas marginales glabriusculis, calycis glabri laciniis brevioribus.

4. A. cespitosa (Lehm. in Diar. Isis 1817 pag. 1288. tab. 9) dense pulvinaris, caudiculis dense et imbricatim foliosis, foliis parvis linearis-lingulatis parte inferiori membranaceis subdilatatis semiamplexicaulibus adpressis glabris superiori erecto-patulis patule albo-hirtulis obtusis, floribus terminalibus solitariis pedicellis foliis paulo longioribus glabris suffultis, calycis glabri ad $\frac{3}{4}$ partiti laciniis oblongo-linearibus obtusis, corollæ (e sicco roseæ) tubo calyce non longiore laciniis oblongis tubo æquilongis ♂.

Hab. in Alpibus provinciæ Ghilan Persiæ a Pallasio prob. lecta. Vidi specimen typicum herb. Lehmann ab am. Prof. G. Reichenb. fil. mihi benevolo dono data!

Facies *A. cylindrica* vel *Pyrenaicæ*, folia $2\frac{1}{2}$ -3 lin. longa linea dimidia vix latiora, flores magn. *A. glacialis*. Vidi in herb. Acad. Petrop. specimen sterilia in monte Antschabala ditionis Bogos Caucasi orientalis 9000'-10000' lecta quæ speciem ineditam affinem caudiculis elongatis 3-4 pollicaribus interrupte fascientifoliis, foliis mucronatis nervo crasso præter cilia marginalia glabris diversam sistunt.

* * Biennes.

5. A. multiscapa (Duby in p.c. Prodr. VIII. p. 51) pilis brevibus ramosis hirta subviscida, radice tenui verticali, foliis dense rosulatis anguste linearis-lingulatis elongatis obtusis integris vel superne minute 1-2-dentatis, scapis numerosis folia duplo superantibus eadem pubescentiis, umbellis 2-5-floris, involuci phyllis breviter oblongo-lanceolatis acutis pedicellos calyci subæquilongos æquantibus, calycis tubo campanulato glabrescenti dentibus triangulari-lanceolatis acuminatis, corollæ albæ vel carneæ limbi tubo brevioris laciniis spathulatis obtusissimis ②. *A. bidentata* C. Koch Linn. XXIII, p. 610.

Hab. in cacuminibus Libani (Auch. 320!), ad Djord Arasya 9500' (Ky. 356!), in Tauro Cilico ad Gisyl Tepe et Bulghar Maaden (Ky. 125 sub *A. Armeniaca*! Balt!), in monte Masmeneudagh Cappadociæ (Balt!), Armenia (C. Koch!).

Facies *A. carneæ* sed monocarpica, rosula unica, scapi breviores. Folia 4-8 lineas longa linea $\frac{1}{3}$ -1 lata. Species pubescentia non adpressa, foliis angustis subintegris, corone limbo brevi ab *A. Armeniaca* egregie distincta.

6. A. Armeniaca (Duby in p.c. Prodr. VIII. p. 52) pubescentia ramosa adpressa substellata sparsa ad folia præter marginem interdum evanescente obsita, radice tenui verticali, foliis rosulatis oblongo-lanceolatis basi attenuatis superne utrinque acute 2-3-laciniato-dentatis, scapis folia 2-4-plo superantibus, umbellis 5-10-floris, involuci phyllis breviter lanceolatis acuminatis pedicellis 2-3-plo brevioribus, pedicellis sub anthesi calyce vix longioribus, calycis breviter glanduloso-hirti vel glabratæ dentibus ovato-triangularibus aequaliter tubo obconico subæquilongis, corollæ albæ limbo tubo æquilongo laciniis obo-

vatis ②. *A. intermedia* Ledeb. Fl. Ross. III. p. 20. — *A. pleioscapa* C. Koch XXIII. p. 611.

Hab. in regione alpinâ Armeniæ Turcicæ, mons Techdagh prope Erzerum 8'000' (Auch. 5231!, Huet! Calv!), Armeniæ Rossicæ (Szov!), in Ponto Lazico supra Djimil 9000' (Ball!), in regione subalpinâ Guriæ (Nordm!).

Folia 5-7 lineas longa 2 lata, scapi 1-1½-pollicares fructiferi elongati 3-4-pollicares

7. *A. macrantha* (Boiss. et Huet Diagn. Ser. II, 3, p. 419) pube ramosâ viscidulâ hirtella, radice tenui verticali, foliis rosulatis oblongo-linearibus basi attenuatis utrinque parte superiori acute 3-5 dentato-laciniatis, scapis elongatis folia multo superantibus, umbellis 7-12-floris, involueri phyllis oblongis acuminatis rubro-glandulosis interdum denticulo auctis pedicellis triplo brevioribus, pedicellis sub anthesi calyce 2-3-plo longioribus, calycis stellatim hirti et rubro-glandulosi dentibus triangularibus acutissimis tubo obconico subbrevioribus, corollæ lacteæ amplæ limbo calyce duplo longiore laciniis spathulatis obtusis ②.

Hab. in Armeniâ Turcicâ prope Gumuschkhané 3500' (Huet!) et in pasenis alpinis montis Fecellem (Bourg. exs. sub *A. Armeniacâ*!).

Valde affinis *A. Armeniacæ* et ab eâ ut *A. lactiflora* ab *A. septentrionalis* discedens, forsitan ejus varietas, scapi 4-7 pollicares, corolla duplo major limbo diametro 6-7 lineas lato.

8. *A. Albana* (Stev. in Linn. Trans. II, p. 440, tab. 33) radice tenui verticali tandem in fibras divisâ, foliis rosulatis pilis papillosis basi dilatatis simplicibus præsertim versus marginem obsitis oblongo vel lanceolato-spathulatis basi attenuatis utrinque parte superiori 2-3 obtuse dentatis, scapis pube ramosâ brevi parce hirtis folia multoties superantibus, floribus brevissime pedicellatis dense umbellato-capitatis, involueri phyllis villosis oblongis acutis pedicellis superantibus, calycis albo-villosi dentibus triangularibus acutis tubo obconico-turbinato duplo brevioribus, corollæ albæ vel pallide carneæ limbo tubo æquilongo laciniis obovatis obtusis ②. *A. valerianoides* Lehm. Isis 1817 ex typo herb. Lehm. a cl. Rehb. fil. mihi benevolè largito. *A. longifolia* C. Koch loc. cit. p. 610! non Turcz.

Hab. in regione alpinâ Ponti Lazici supra Djimil 7500' (Ball!), Caucasi centralis in Ossetiâ prope Sgis 800' (Rupr!), in Caucaso Orientali frequens 8'000'-10'000' (Stev., Ruprecht!), Armeniæ Rossicæ (Szov!).

A binis præcedentibus quibus foliis accedit eorum pube simplici floribusque subsessilibus capitatis statim distinguenda, scapi 3-6-pollicares, flores magn. *A. Armeniaceæ*.

9. *A. Wiedemannii*, radice tenui verticali tandem in fibras divisâ, foliis rosulatis margine ciliatulis cæterum glabris oblongo-spathulatis basi attenuatis obtusis utrinque parte superiori obtuse et breviter 2-3-dentato-lacinulatis, scapis pube brevi ramosâ hirtis folia

multo superantibus, floribus brevissime pedicellatis capitatis, involueri phyllis oblongo-spathulatis basi attenuatis obtusis margine præsertim ciliatis calyces æquantibus, calycis dentibus viridibus pilis papillosis hirtis ovatis obtusis tubo turbinato glabriuscule æquilongis, corollæ roseæ limbo tubo sublongiore laciniis retusis ②.

Hab. in monte Ylgasdagh Anatoliæ borealis (Wied!).

Scapi 2-4-pollicares, folia et facies *A. Albanæ*.

* * * Annuæ.

10. ***A. elongata*** (L. Sp. 1668) annua, foliis rosulatis pilis simplicibus et stellatis margine præsertim parce hirtulis vel glabratis oblongo-lanceolatis acutiusculis integris vel obtuse paucidentatis, scapis brevibus erectis pube brevi ramosâ hirtulis, umbellæ 2-8-floræ pedicellis setiformibus inæqualibus longissimis, involueri phyllis pedicellis multoties brevioribus oblongo-lanceolatis linearibusque acutis, calycis breviter papilloso-hirti dentibus triangulari-lanceolatis aenimnatis tubo obconico quinquecostato æquilongis, corollâ calyce breviore limbo parvo ③. Ic. Jacq. Austr. tab. 330. — Obs. I, tab. 19. — Rehb. Ic. tab. 69. — *A. nana* Horn. En. Hafn. I. p. 184.

Hab. in Rumelia (Frvtl), Tauriâ et prov. Caucasicis (Stev.), Transeaucasiam (C. Koch, Hohen.).

Ar. Geogr. Germania, regio Danubialis, Sicilia, Rossia media et australis, Sibiria.

11. ***A. filiformis*** (Retz Obs. II, p. 10) annua glabra, foliis rosulatis ovatis vel oblongis obtusis obtuse dentatis in petiolum attenuatis, scapis elongatis erectis, umbellæ multifloræ pedicellis setiformibus inæqualibus elongatis, involueri phyllis minimis linear-i-oblongis acutis, calycis dentibus triangularibus acutis tubo hemisphaerico-obconico brevioribus, corollâ albâ calycem vix excedente lobis obtusis ④. Ic. Gmel. Sib. IV, tab. 44. — *A. radiata* Lehm. in R. et Sch. Syst. IV, p. 164!

Hab. in prov. Ghilan Persiae australis ex specimine archetypo herb. Lehmann mihi a cl. Rehb. fil. amiciss. largito!

Ar. Geogr. Songaria, Sibiria Altaica et Baikalensis, Kamtschatka, America borealis.

12. ***A. septentrionalis*** (L. Sp. 203) annua, foliis rosulatis oblongo-lanceolatis integris vel breviter paucidentatis pube brevissimâ ramosâ hirtulis vel glabratis, scapis elongatis erectis puberulis, umbellæ multifloræ pedicellis setiformibus inæqualibus elongatis, involueri phyllis pedicellis multoties brevioribus linear-i-oblongis acutiusculis, calycis minute hirti vel glabratæ dentibus triangularibus acutis tubo obconico subbrevioribus, corollâ albâ fance lutescenti calycem superante laciniis oblongo-spathulatis ⑤. Ic. Fl. Dan. tab. 7.

— Rehb. tab. 69.

Hab. in arenosis regionis alpinæ Caucasi Orientalis 9000' (C.A.M.), Caucasi centralis ad ripas fluvii Terek prope Lars 3500' (Rupr!). In Tauriā indicata sed dubia.

Ar. Geogr. Helvetia in alpibus Valesiæ, Scandinavia, Belgium, Transylvania, Rossia, omnis, Sibiria, Aimerica borealis.

13. A. maxima (L. Sp. 203) annua parce papilloso-hirtula, foliis rosulatis expansis ellipticis vel lanceolatis parte superiori serrulatis, scapis crassiusculis pilis simplicibus elongatis pubeque minimâ ramosâ hirtellis, umbellæ 3-10-floræ pedicellis calyci æquifloris vel duplo longioribus, involueri phyllis magnis obovatis subdentatis basi subcoarctatis, calycis tandem accreti hirsuti laciniis ovatis amplis tubo globoso hemisphærico longioribus, corollâ parvâ calyce multo breviore albido-rubellâ limbi concavi lobis obovatis ⊖. Ic. Jacq. Austr. tab. 337. — Rehb. Ic. t. 70.

Hab. in collibus, campis derelictis Macedoniæ (Friv!), Bithyniæ (Pest!), Pisidiæ (Heldr!), Lyciæ (Bourg!), Asie minoris borealis et centralis (Auch! Wied.), Ciliciæ orientalis ditione Kassan Oglu in subalpinis (Ky!), Syriæ ad Aleppo! Damascum!, Palestinæ ad Hierosolymam (Boiss!), Mesopotamiæ (Oliv!), Armenia (Huett! Bourg!), Tauriæ, prov. Cis et Transcaucasicæ M.B! Hohen! Rupr!), Persiæ sept. et australi (Ky!), Affghania occid. ad Herat (Bge!).

Ar. Geogr. Hispania, Gallia, Belgium, Germania occid. et australis, Italia centralis, regio Danubialis, Rossia media et australis, Songaria, Sibiria, Africa borealis interior.

DIONYSIA (Fenzl Flora XXVI, I, p. 389. — *Macrosiphonia* Duby Mém. Primul.).

Calyx profunde vel ad basin usque quinquepartitus. Corolla tubulo-hypocrateiformis tubo longo tenui vel a medio vel sub fauce subito ampliato, fauce nudâ, limbi patentis lobis obtusis vel obcordatis. Antheræ (dichogamice et prob. in speciebus omnibus) in medio tubo vel sub fauce subsessiles verticillatæ stylusque antheras non attingens vel eas superans vel exsertus; stigma globoso-capitatum; ovarium uniloculare pauciovulatum. Capsula in valvas 5 complete dehiscens. Semina 1-4 extus convexa intus carinato-angulata rugulosa. — Herbae perennes dense palvinatæ, interdum suffrutices cespitosi, rami foliorum minutorum persistentium imbricatione densâ columnares, flores in cespite sessiles vel pedunculati, lutei rarius violacei. — Genus a *Gregorid* fauce nudâ, ab *Androsace* eâdem notâ et corollæ tubo elongato distinctum.

* Folia omnia vel saltem nonnulla superiora plus minus dentata.

+ Folia margine revoluta.

I. D. revoluta (Boiss. Diagn. Ser. I, 7, p. 63) fruticulosa late cespitosa ramis elongatis dichotomis inferne denudatis superne foliis

deflexis vetustis vestitis, ramulis hornotinis brevibus densiuscule foliatis, foliis brevissime glanduloso-puberulis crassiusculis linearibus uninerviis obtusis margine revolutis et obtuse multicrenatis crenis inflatis, floribus 1-2 subsessilibus, calycis breviter hirti ad $\frac{3}{4}$ partiti laciniis anguste linearibus obtusiusculis, corollæ hirtulae luteæ tubo calyce triplo longiore, limbi laciniis obocordato-orbiculatis φ .

Hab. in fissuris rupium ad septentrionem spectantium montis Sabst Buschom prope Schiraz Persiae australis (Ky. 426!), mense Maio, in monte Sawers 11000' (Haussk!) Julio.

β. canescens. — Indumentum densius longius canescens.

Hab. in montibus Kuh Elburs, Eschker et Maregum Persiae austro-occidentalis 9-12000' (Haussk!).

Rami a caudice lignosâ 5-7-pollicares, folia 3-4 lineas longa santolinoidea, corolla 6-7 lineas longa.

+ + Folia plana.

2. D. aretioides (Lehm. Mon. Prim. p. 90, tab. 9 sub *Primula*) canescens velutina dense pulvinari-cespitsa, ramis tenuibus basi denudatis superne foliis vetustis vestitis apice rosuliferis, foliis longe et dense pilis eglandulosis crispis villosis linearispathulatis planiusculis in petiolum brevem attenuatis uninerviis acutiusculis paucidentatis, floribus solitariis pedicello calyci subaequilongo suffultis, calycis ad $\frac{3}{4}$ partiti villosuli laciniis linearibus acutissimis, corollæ luteæ superne parce puberulæ tubo limbi laciniis obocordatis sextuplo longiore φ . *D. aretioides* Boiss. loc. cit. p. 68. — *Gregoria aretioides* Duby loc. cit. p. 46.

Hab. in prov. Ghilan (Pall. in herb. Lehmt) specimina mihi a cl. Rehb. fil. data, rupibus Alpium Samamys Kuh prov. Ghilan Persiae borealis (Hablitzl in herb. Acad. Petrop.).

Folia 2½ lineas longa infra apicem fere lineam lata, calyx 2 lineas, corolla ex cl. Lehmann 12-13 lineas longa sed ex speciminiibus ejus herbarii brevior. Species indumento canescenti insignis. Icon Lehmanniana plantam speciminiibus ejus herbarii omnibus partibus majorem exhibet.

3. D. Aucheri (Duby in p.c. Prodr. VIII, p. 46 sub *Gregorid*) fruticulosa dense pulvinari-cespitsa, ramis tortuosis inferne denudatis superne ob folia vetusta imbricata columnaribus apice minime rosuliferis, foliis glanduloso-hirsutis griseis cuneato-orbiculariatis planis acutiuscule 3-7-dentatis flabellatum nervosis, floribus solitariis sessilibus, calycis hirti ad basin partiti laciniis oblongo-linearibus obtusiusculis, corollæ luteæ puberulæ tubo calyce sextuplo longiore limbi laciniis ovatis integris φ . *D. odora* Fenzl Flora 1843, p. 390. — *Aretia longiflora* Fisch. in Kotschy Kurd. exs.

Hab. in fissuris rupium montis Nalkou Assyriæ (Auch. 2610!), Kurdistanie Assyriacæ in monte Gara (Ky. 386!), in montibus Pir Omar Gudru et Jauaro Kurdistanie Persicae 6000'-7000' (Haussk!). Fl. Jul.

Rami 3-4-pollicares, folia 1-2 lineas longa apice $\frac{1}{2}$ -1 lata, viscidulo-hirta odore aromatica, corolla 10-12 lineas longa

4. **D. cespitosa** (Duby in D.C. Prodr. 8, p. 46) suffrutescens longe pulvinaris ramis foliis vetustis compacte vestitis rosulis paucifoliis; foliis viridibus minute glandulosis planis linearis vel oblongo-spathulatis obtusis integris vel utrinque obtuse 1-3-crenatis penninerviis, pedunculis longe exsertis 1-2-floris, floribus involuero tribracteato suffultis intra bracteas brevissime pedicellatis, bracteis inaequalibus ovato-oblongis dentato-incisis, calycis magni glandulosi usque ad basin partiti laciniis lanceolatis acutis integris, corollæ luteæ glanduloso-puberula tubo calyce 3-4-plo longiore limbi laciniis obovatis integris ♀. *Dionysia cespitosa* Boiss. loc. cit. — *Macrosiphonia cespitosa* Duby Mém. Prim. tab. 2, fig. 3.

Hab. in monte Elvend Persiae (Auch. 2609), in montibus tribus Bachtiaris ad occasum Ispahan (Auch. 5234! Bode!).

Pulvini 2-3 pollices alti, folia 2 $\frac{1}{2}$ lineas longa, calyx 4 lineas, corolla fere pollicaris.

5. **D. diapensiæfolia** (Boiss. Diagn. Ser. I, 7, p. 65) densissime et late cespitosa, ramis ob folia vetusta dense imbricata columnaribus, foliis minute hirto-glandulosis planis subflabellatim reticulato-venosis ovatis et oblongo-spathulatis basi attenuatis obtusissimis integris vel obtuse utrinque 1-2-crenatis, pedunculis subnullis rosulæ foliis occultatis vel paulo longioribus breviter exsertis 1-rarius 2-3-floris floribus involuero 3-5 bracteato suffultis, bracteis linearispathulatis obtusis integris calycem requantibus, calycis ad $\frac{3}{4}$ partiti laciniis linearispathulatis obtusis, corollæ luteæ glanduloso-hirtæ tubo calyce quadruplo longiore limbi ampli laciniis ovatis retusis ♀.

Hab. ad parietes rupium ad septentrionem spectantium ad Persepolin Persiae (Ky. exs. 236!).

Cespites lati 3-4 pollices elati, folia 4 $\frac{1}{2}$ -2 lineas longa, calyx 2 $\frac{1}{2}$ lineas, corolla 10-12 longa. Pedunculus exsertis et involuero affinis *P. cespitosæ* sed in hac pedunculus longus, bracteæ majores incisæ, calycis laciniæ acutæ, corollæ limbus minor.

6. **D. drabæfolia** (Bunge Bull. Acad. Petrop. VIII, p. 207) dense pulvinaris ramis vetustioribus nudatis junioribus ob folia conferta columnaribus, foliis rosulatis planis reticulato-venosis glandulosis aliis linearispathulatis integris aliis obovato-oblongis utrinque acute 1-2 dentatis acutiusculis, floribus solitariis subsessilibus bracteis tribus minutis integris inaequalibus suffultis, calycis usque ad basin partiti laciniis linearibus integris acutiusculis glanduloso-hirtis, corollæ glabra tubo calyce quadruplo longiore, limbi laciniis emarginato-bilobis ♀. *D. diapensiæfolia* var. Hohenacker in Sched.

Hab. ad ruinas Persepolis et in cacumine alpis Kuh Daëna (Ky. in Bge herb.). Non vidi.

Ex am. a Bunge a *D. diapensiifoliatâ* corollâ glabratâ imprimis distincta, folia insuper majora videntur sœpe 5 lineas longa dentibus cum adsunt acutis.

* * Folia omnia integerrima.

+ Folia plana flabellatim nervosa.

7. **D. Michauxii** (Duby in D.C. Prodr. VIII, p. 46 sub *Gregoria*) dense pulvinaris ramis brevibus columnaribus adpressissimis, foliis minimis dense imbricatis breviter velutinis elliptico-oblongis obtusissimis flabellatim nervosis integris, floribus solitariis sessilibus, calycis ad basin partiti laciniis linearis-lanceolatis acutis, corolla hirtæ calyce 4-5-plo longioris laciniis euneato-ovatis integris ♀. *D. Michauxii* Boiss. Diagn. Ser. I, 7, p. 67.

Hab. in Persiâ loco non notato (Michaux in Delessert herb! frustulum!).

Foliorum nervatione et habitu affinis *D. tapetode*, sed folia (quidquam minora) velutina, corolla brevior (4 lin. longa) valde hirta nec glabra.

8. **D. tapetodes** (Bge loc. cit. p. 210) dense et late pulvinaris, ramis brevibus adpressis tenuiter columnaribus, foliis oblongo-spathulatis integris minute glanduloso-punctatis cæterum glabris obtusis elevatim flabellato-nervosis, floribus sessilibus solitariis, calycis ad $\frac{2}{3}$ partiti laciniis minute glandulosis lanceolatis acutis, corolla glabrae flavæ tubo calyce quadruplo longiore limbi laciniis obovatis ♀.

Hab. in parietibus superpendentibus rupium supra Derrud prov. Khorassan Persie inter Nischapur et Meschhed (Bunge!), in fauibus Siri Chusma prope Kabul Affghaniae (Griff. Journ. 981!).

Rupes umbrosas quasi tapete vestiens, foliis flabellatim et elevatim nervosis fere sesquilineam longis inter affines preter *D. Michauxii* insignis, corolla 5-6 lineas longa. Ab affini *D. Michauxii* differt foliis glanduloso-punctatis nec velutinis, corollâ glabratâ.

+ + Folia plana uninervia.

9. **D. bryoides** (Boiss. Diagn. Ser. I, 7, p. 66) dense pulvinaris ramis brevibus adpressis columnaribus, foliis minimis imbricatis minute glandulosis obovatis eucullatis integris uninerviis, floribus solitariis sessilibus, calycis foliis occultati ad basin partiti laciniis linearis-spathulatis basi longe attenuatis puberulis obtusis, corolla violacea puberulae tubo calyce duplo longiore, limbi parvi lobis ovatis ♀.

Hab. in rupibus montis Kuh Ajub prope Persepolin Persie (Ky. 406!). Fl. Mai.

Facies *Androsaces Helveticae*, folia lineam vel sesquilineam, corolla 4 lineas longa.

10. **D. Kotschy** (Bunge loc. cit. p. 209) cespitosa ramis vetus-

tioribus tenuibus denudatis, hornotinis gracilibus elongatis foliis vetustis sursum imbricatis dense vestitis, foliis minute glandulosis hirtulisque oblongis obtusis integris basi attenuatis uninerviis, floribus sessilibus solitariis, calycis ad basin partiti laciniis oblongo-linearibus acutis minute glanduloso-puberulis, corollæ luteæ glabræ tubo calyce sextuplo longiore gracili limbi lobis oblongis retusis ♀.

Hab. in monte Kuh Ajub prope Persepolin in consortio *D. bryoidis* (Ky. exs. 406 b!).

Differit a *D. bryoide* ramis hornotinis elongatis pollicem et amplius longis non arete pulvinatis, foliis majoribus, corollâ semipollucari, calycis proportione multo longiore.

11. *D. curviflora* (Bunge loc. cit. p. 212) dense pulvinaris ramis tenuiter columnaribus, foliis minutissimis dense imbricatis eglandulosis albo-ciliatis obovatis obtusis obscure uninerviis integris, floribus solitariis sessilibus, calyeis foliis occultati fere ad basin partiti laciniis lanceolato-linearibus acutiusculis ciliatis, corollæ glabræ luteæ tubo calyce quadruplo longiore (an in plantâ vivâ?) incurvo, limbi laciniis cuneato-obcordatis ♀. *D. bryoidea* Boiss. et Buhse Aufz. p. 145.

Hab. in fissuris rupium montis Schirkuh prope Jezd Persiæ orientalis (Buhse!). Fl. April.

Hanc primâ facie retuleram ad *D. bryoideum* quam aspectu refert sed ab eâ optime distinguitur foliis eglandulosis ciliatis, corollâ cum limbo 7-8 lineas longâ.

+ + + Folia (saltem superiora) margine revoluta.

12. *D. rhaftodes* (Bunge loc. cit. p. 213) dense pulvinaris ramis columnaribus confertis, foliis dense et patule imbricatis hispidulis eglandulosis breviter linearibus uninerviis marginè revolutis obtusis integris. floribus solitariis sessilibus, calycis fere ad basin partiti laciniis linearibus acutis, corollæ flavæ puberulæ tubo calyce subquadruplo longiore, limbi laciniis orbiculari-obovatis ♀.

Hab. in rupibus ad septentrionem spectantibus vallis alpinæ supra Syrtsch inter Chabbise et Kerman Persiæ austro-orientalis (Bge!). Fl. Aprili.

Pulvinus 2-3 pollices altus, folia $1\frac{1}{2}$ -2 lineas longa dimidiata lineam lata, calyx sesquilineam, corolla 7 lineas longa.

PRIMULA (L. Gen. 497).

Calyx quinquefidus vel partitus. Corolla hypocraterimorpha vel infundibuliformis tubo cylindrico, fauce dilatata forniciibus brevibus saepius instructa. Ovarium multiovulatum. Capsula polysperma apice valvis 5 dehiscens. — Herbæ perennes saepius alpineæ.

SECT. I. **SPHONDYLIA** Duby in p.c. Prodr. — Folia vernatione revolutiva, flores in verticilos involucris foliiformibus suffultos dispositi. Capsula globosa.

1. **P. Boveana** (Decaisne in p.c. Prodr. VIII, p. 35) partibus junioribus minute pulverulento-floccosa, radice brevi obliquâ valde fibrillosâ, foliis tenuiter membranaceis obovato vel rhombeo-spathulatis acutis acute inciso-lobatis et minute denticulatis in petiolum planum attenuatis, scapis angulosis folia 1-3-plo superantibus florum verticilos 4-4 involucratos inferiores renitos ferentibus, involuci phyllis sessilibus basi cuneatis integris cæterum acute serratis acutis vel acuminatis verticillorum inferiorum obovato-rhombeis flores æquantibus superiorum diminutis lanceolato-linearibus, pedicellis flori æquilongis et superioribus brevioribus, calycis albo-pruinosis paulo infra medium fissi laciniis breviter lanceolatis acute 1-3-dentatis vel integris, corollæ glabrae citrinae tubo cylindrico calyce triplo longiore, limbi laciniis tubo multoties brevioribus crenulatis orbiculatis, seminibus quadrato-angulatis oculo valde armato granulosis ♀. Ic. Jaub. et Sp. tab. 439. — *P. verticillata* Decaisn. Fl. Sinaie. — Bot. Mag. tab. 2842 non Forsk.

Hab. in regione montanâ Arabiæ petreæ ad rupes stillicidiis madidas, mons Sinai ad fontem Perdicis (Boiss!), mons Horeb (Stæ Catharinæ) et Raphidim (Elirenb., Bové!, Schimp!). Fl. Maio.

Affinis *P. verticillata* Forsk Arab. p. 42. — Vahl. Symb. I, 5, ex montibus Arabiæ Felicis extra fines nostros oriundæ, foliis lanceolatis serrulatis sed non lobatis, calycis fere ad basin partiti laciniis longis lanceolatis integris diversæ. *P. Simensis* Hochst. — J. et Sp. loc. cit. tab. 440, Abyssinica, calycem habet ad medium in laciniis latae (integras) partitum *P. Boveana* sed folia lanceolata non lobata *P. verticillata*, et ab utrâque differt corollæ limbo majore tertiam tubi partem æquante laciniis latis retusis.

2. **P. Aucherii** (Jaub. et Sp. Illustr. tab. 49) partibus junioribus minute albo-pulverulenta, caudicibus elongatis tenuibus petiolorum vestigiis vestitis, foliis lanceolato-spathulatis obtusis obtuse et late crenatis in petiolum sensim attenuatis, scapis teretibus foliis 1-3-plo longioribus florum verticilos 2-6 inferiores involucratos ferentibus, involuci phyllis sessilibus lanceolatis acuminatis serratis vel integris verticillorum inferiorum pedicellos superantibus superiorum diminutis, pedicellis flore brevioribus, calycis breviter hirti ad $\frac{3}{4}$ partiti laciniis integris anguste lanceolatis acuminatis; corollæ citrinae brevissime hirtæ tubo cylindrico calyce plus duplo longiore limbi laciniis tubo multo brevioribus ovatis, seminibus (ex el. J et Sp.) orbiculatis planiusculis ♀.

Hab. ad rupes madidas montis Gebel Akadar Arabiæ Mascatensis (Auch. 5236!).

Præcedente omnibus partibus minor ad summum pedalis, folia obtusa et obtuse crenata, corolla 8-9 lineas longa.

3. **P. floribunda** (Wall. Tent. Nap. tab. 33) hirtula insuper partibus junioribus albo-pulverulenta, foliis rugulosis venosis ovato-oblongis obtusis inæqualiter dentatis in petiolum attenuatis, scapis pluribus erectis pumilis folia superantibus, florum verticillo terminali, involueri phyllis oblongis vel oblongo-lanceolatis dentatis pedicellis brevioribus, calycis hirti ultra medium partiti laciniis breviter lanceolatis integris, corollæ luteæ hirtæ tubo calyce triplo longiore limbi laciniis retusis ♀.

Hab. ad margines rivorum in Affghaniâ ad Pushut (Griff. Itin. Not. p. 338).

Specimen valde imperfectum corollis fere destructis vidi sed ut videtur plantæ Himalaicae ex viciniis Simlah in East Ind. Comp. Herb. № 3817 simile.

Ar. Geogr. Regio Himalaica.

SECT. II. PRIMULASTRUM Daby in D.C. Prodr. — Folia vernatione revolutiva rugosa non farinosa, flores umbellato-capitati, calyx argute angulatus.

4. **P. acaulis** (Jacq. Misc. I, 158) foliis obovato-oblongis rugulosis dentatis subtus ad venas vel ubique hirtis in petiolum brevem sensim attenuatis, scapis unifloris numerosis folia æquantibus vel brevioribus calyceque molliter pilosis, calycis acute prismatici dentibus lanceolatis acuminatis, corollæ flavæ limbo plano limbum æquante laciniis obcordatis ♀. Ic. Rehb. tab. 50. — *P. grandiflora* Lam. Fl. Fr. II, p. 248. — *P. sylvestris* Scop. — *P. veris* var. *acaulis* L.

Hab. in humidis ad fontes et umbrosis regionis montanæ et subalpinae, Peloponnesus in Taygeto et monte Menalo (Heldr!), Kyllene 3000'-4000' (Orph. Fl. Gr. exs. 330!), Creta in Eparchiâ Kissamos 1500' (Heldr!), Macedonia (Griseb!), Anatolia bor. in monte Aladagh (Wied!) et ad Trapezuntum (Bourg!), Tauria ad Sympheropol (Stev!), Cilicia orientalis in ditione Kassan Oghlu 4-6000' et Syriae mons Amanus 4500' (Ky!), Libanus circa Eden (Boiss! Gaill!), Persia bor.-orientalis supra Asterabad (Bge!).

β. rosea. — Corolla purpurea rosea vel alba. — Ic. Fl. Græc. tab. 184. — *P. acaulis* Iberica Hoffm. — *P. Sibthorpii* Rehb.

Hab. in Tauriâ (Stev.), circa Byzantium et Brussam copiose (Sibth., Cadet!, Barbey!, etc.), Ponto Lazico ad Rhizé (Bal!), in Transcaucasiâ totâ et Armenia Rossicâ communis!

Ar. Geogr. Europa tota a Scandinaviâ ad Rossiam australem in australioribus montana et subalpina, Africa borealis.

5. **P. officinalis** (Jacq. Misc. I, p. 159) foliis oblongo-ellipticis rugulosis in petiolum alatum sensim rarius abrupte decurrentibus subtus hirtis subconcoloribus, scapis umbelliferis floribus unilateraliter nutantibus, involueri phyllis a basi latâ subulatis, calyce tomentello cylindrico-obconico argute angulato corollæ non adpresso lobis brevibus ovatis, corollæ luteæ limbo concavo basi quinqueguttato tubo multo breviore ♀. *P. veris* var. *officinalis* L. Sp. 107, Ic. Fl. Dan. tab. 433. — Rehb. Ic. tab. 49.

Hab. in Thraciae monte Hæmo (Friv. ex Griseb. Spic.), in Tauriâ et prov. Caucasicis indicata sed prob. ex confusione cum *P. macrocalyx*. E ditione nondum vidi.

Ar. Geogr. Europa septentrionalis et media a Scandinaviâ ad Rossiam, Sibiria.

6. *P. suaveolens* (Bertol. Journ. Bot. 1813, p. 76) foliis rugulosis ovatis vel subcordatis in petiolum alatum sæpius abrupte abeuntibus subtus canescens vel niveo-tomentosis, scapis umbelliferis floribus unilateraliter nutantibus, involuci phyllis a basi latâ subulatis, calyce tomentello cylindrico obconico argute angulato corollæ non adpresso lobis breviter ovato-oblongis acuminatis, corollæ luteæ limbo concaviusculo tubo breviore ♀. Ic. Rehb. tab. 50. — *P. Columnæ* Ten.

Hab. in pratis alpinis Graeciæ in Peloponneso (Bory!), Olympo Thessalo (Heldr!), Macedonia in monte Scardo (Griseb.), Armeniâ Turciâ ad Gumusch-khanè et Zazalar chané (Bourg!), Ponto Lazico supra Djimil (Bal. exs. sub *P. macrocalyx*).

Valde affinis *P. officinali* cuius in Europâ australi vices agit et forsitan in eam transiens; differt foliis subtus plus minus canis in petiolum abruptius abeuntibus et præsertim corollæ laciniis majoribus minus concavis.

Ar. Geogr. Hispania, Gallia et Helvetia australiores, Italia, Istria, Dalmatia, Serbia.

7. *P. macrocalyx* (Bge in Fl. Ait. I, p. 209) foliis rugulosis ovatis vel cordatis in petiolum alatum abrupte abeuntibus subtus hirtis vix canescens, scapis umbelliferis floribus unilateraliter nutantibus, involuci phyllis lanceolatis, calyce tomentello a basi late obconico apertura diametro longitudini subæquilongo lobis breviter triangulari-ovatis acuminatis, corollæ luteæ limbo concavo tubo multo breviore ♀. *C. officinalis* var. *macrocalyx* C. Koch Linn. XVII. — *P. Uralensis* Fisch. ex Rehb. Pl. Crit. Cent. VII, tab. 638.

Hab. in montosis Iberiae Caucasicæ (Hoh!, Rupr!), Caucasi centralis et orientalis 3500'-6000' (Rupr!), ditione Talysch (Hoh!), Armeniâ Rossicâ ad fines Persicas (Szov!), Persiâ bor. in monte Elbrus (Ky. 132!).

Valde affinis *P. officinali* cui a nonnullis jure forsitan ut varietas adnumeratur, differt calyce breviore et ore latiore. *P. inflata* Lehm. ex Hungariâ est potius *P. officinalis* forma.

Ar. Geogr. Songaria, Sibiria Uralensis et Altaica.

8. *P. elatior* (Jacq. Misc. I, p. 158) foliis rugulosis ovato-oblongis subtus hirtis pallidioribus in petiolum alatum sæpius abrupte contractis, scapis umbelliferis floribus unilateraliter nutantibus, involuci phyllis a basi latiore subulatis, pedicellis longe et crispule hirsutis, calycis hirti anguste cylindrici corollæ adpressi dentibus lanceolatis acutis tubo duplo brevioribus, corollæ luteæ limbo piano tubo æquilongo ♀. Ic. Fl. Dan. tab. 434. — Rehb. tab. 49. — *P. veris* var. *elatior* L.

Hab. in sylvaticis Tauriæ meridionalis ad Bijuk Lambat (Stev.), et Transcaucasicæ circa Schuscha (Hohen. ex Rupr.). E ditione eam nondum vidi.

Ar. Geogr. Europa media ab Angliâ et Scandinaviâ ad Rossiam medium et ab Hispaniâ et Italî septentrionalibus ad regionem Danubialem.

9. P. Pallasii (Lehm. Prim. p. 38, tab. 3) foliis obsolete rugulosis tenuibus glabriuseulis vel subtus parce glandulosis oblongis elongatis in petiolum longe attenuatis, scapis umbelliferis floribus unilateraliter nutantibus, involucri phyllis breviter lanceolatis, pedicellis brevissime glanduloso-velutinis, calycis brevissime hirti anguste cylindrici corollæ adpressi dentibus lanceolatis acutis tubo 3-4-plo brevioribus, corollæ luteæ limbo plano tubo subbreviore ♀. *P. elatior* var. Ledeb. Fl. Ross.

Hab. in regione montanâ et alpinâ Armeniæ Turcicæ circa Erzerum (Huett!, Calv!), in Armeniae Rossicæ montibus Ararat et Adchara (Nordm.).

β. cordifolia. — Folia abrupte in petiolum contracta. — *P. cordifolia* Rupr. Prim. p. 286.

Hab. in Caucaso centrali ad Kasbek et Kobi 6000' (Rupr!), in Somchetiâ (C. Koch).

Species *P. elatiori* valde proxima et cum eâ a variis auctoribus conjuncta differt foliis glabris substantiæ tenuioris, pedunculis et scapo breviter velutinis nec longe villosulis, calycis dentibus brevioribus. Specimina Armena cum Altaicis convenient sed icon Lehmanni male calycem apertum nec angustum corollæ adpressum exhibit.

Ar. Geogr. Sibiria Uralensis.

10. P. amæna (M.B. Taur. Cauc. I, p. 138) foliis rugulosis ovatis et oblongis subtus griseo vel canô-tomentosis in petiolum sæpius sensim attenuatis, scapis umbelliferis floribus unilateraliter nutantibus, involucri phyllis breviter lanceolatis acuminatis, pedicellis crispule hirsutis, calycis breviter hirti anguste cylindrici corollæ adpressi dentibus lanceolatis acutis tubo duplo brevioribus, corollæ intense purpureæ (et interdum ex cl. Rupr. flavæ) limbo plano tubo æquilongo ♀. Ic. Lehm. Mon. tab. 3. — *P. elatior* var. *amæna* Ledeb. Fl. Ross.

Hab. in regione alpinâ Ponti Lazici supra Djimil 8000' (Bal!), in Caucasi centrali monte Beschtau (M.B.), Kasbek 6000'-40000' (Rupr!, Rehmnn!).

β. Meyeri. — Folia subcordata abrupte et longius petiolata, pedicelli glabrescentes — *P. Meyeri* Rupr. loc. cit. p. 284.

Hab. in monte Elbrus (C.A. Mey.) et ad fontes rivi Ingur (Radde!).

Species ab affini *P. elatiore* ut *P. suaveolens* ab *officinali* foliis subtus plus minus canis discedens. Varietas intermediis cum typo conjuncta floribus purpureis gaudet.

SECT. III. ALEURITIA Duby in D.C. Prodr. ex parte. — Folia vernalatione revolutiva plana vel vix rugosa subtus sæpe farniosa. Calyx teres.

11. P. megaseæfolia (Boiss. et Bal. in Bal. pl. Pont. exs. 1866)

radice obliquâ fibriferâ, foliis amplis lăvibus ovatis vel ovato-oblongis basi rotundatis vel breviter cordatis petiolis subæquilongis papilloso-hirtis suffultis circumcirca remotiuscule denticulatis denticulis latis mueronulatis supra glabris subtus ad nervos et marginem papilloso-hirtis, scapis foliis æquilongis vel brevioribus, verticillis terminalibus 1-2 approximatis, involueri phyllis a basi anguste lanceolatâ longe subulato-acuminatis pedicellos calyci æquilongos vel duplo longiores non æquantibus, calyce glabro anguste tubuloso ad tertiam partem in dentes lanceolatos acutos fisso, corollæ roseæ tubo calyce sublongiore, limbi tubo vix brevioris laciniis late obcordatis, capsulâ oblongâ calyce vix longiore, seminibus sphærico-angulatis oculo armato tuberculatis ♂.

Hab. in faucibus humidis umbrosis circa Rhizé Ponti Lazici 900' (Bal!) Fl. Mai.

Scapi semipedales, foliorum lamina 3-5 pollices longa 2-4 lata, calyx angustus et corolla magn. *P. amarae*, capsula oblonga cylindrica 7 lineas longa lacinias calycis vix superans. Species foliorum formâ insignis.

12. *P. grandis* (Trautv. Pl. Nov. p. 5 in Bull. Acad. Petr. t. VI) radice..., foliis amplis lăvibus petiolis laminâ duplo longioribus suffultis membranaceis rugulosis ovatis basi subcordatâ in petiolum breviter decurrentibus irregulariter duplicato-denticulatis utrinque viridibus subtus tenuissime pulveraceo-puberulis, umbellâ maximâ multiflorâ scapo foliis longiore suffultâ, involueri phyllis breviter lanceolato-linearibus basi non appendiculatis, pedicellis longis involucrum et flores multoties superantibus. calyce campanulato viridi sub lente pruinoso ad tertiam partem in lobos ovatos acutiusculos diviso, corollæ roseæ tubo calycem æquante limbi laciniis erectis tubo æquilongis oblongo-linearibus obtusis minute retusis sinibus latis sejunctis. stylo longe exerto, capsulis ovatis calyce vix longioribus, seminibus angulatis sub lente minute tuberculatis ♂.

Hab. in montibus Dadiash Transcaucasiae occidentalis 6-9500' (Radde!). (V. sp. incompl. in herb. Petrop!).

Folii lamina 4 poll. longa 3 lata, scapus fere pedalis, pedicelli bipollicares, calyx florifer vix tres lineas longus. Species pulchra ulterius inquirenda angustie laciniarum corollæ insignis.

13. *P. nivalis* (Pall. It. III, p. 320 — App. № 81, tab. 6) foliis planis oblongis elongatis argute vel obtusiusecule multierenatis obtusis vel acutiusculis glabris in petiolum latum brevem indistinctum sensim attenuatis, scapo crasso folia æquante vel superante, umbellâ multiflorâ (interdum binis superpositis), involueri phyllis oblongis vel lanceolatis obtusis vel acuminatis basi coadunatis, pedicellis calyce sæpius brevioribus, calycis minute puberuli corollæ tubo brevioris ad $\frac{3}{4}$, partiti laciniis linear-lanceolatis acutis, corollæ hypocrateriomorphæ purpureæ vel albæ lobis obovatis tubo vix brevioribus,

capsulâ cylindrica calyce duplo longiore, seminibus papillosois ♀. *P. crassifolia* Lehm. Prim. tab. 9.

β. farinosa Schrenk Pl. Nov. I, p. 22. — Folia secus margines, petioli, calyces, farinâ albâ aspersi. — *P. Bayerni* Rupr. loc. cit. p. 306.

Hab. ad rivos regionis alpinæ Ponti Lazici supra Djimil 8000' forma floribus roseis (C. Koch, Bal!), in Caucasi Chewsuriâ (Bayern) et Ossetiâ ad moles glaciales montis Kasbek 6000'-10000' (Rupr!, Rehm!), forma floribus albis.

A cl. Rchb. fil. accepi folium floresque binos '*P. crassifolia* ex herb. Lehm, iconi simillimos et qui omnino formam speciei hebetatam minorem floribus subdiminutis sistere videntur.

Ar. Geogr. Alpes Songariæ (var. *farinosa*), Sibiriæ (forma typica absque farinâ).

14. *P. auriculata* (Lam. Ill. № 1931) glabra viridis, collo vaginis fuscis vestito, foliis basi in petiolum brevem attenuatis planis lœvibus membranaceis lanceolato-ellipticis vel oblongo-spathulatis obtusis remotiuscule et minute denticulatis interdum subjunctis axium lateralium sterilium post anthesin elongatis, scapo florifero saepe foliis breviori sed fructifero valde elongato, umbellâ multiflorâ, pedicellis vel calyce vix brevioribus vel brevissimis, involueri phyllis pedicellis longioribus linear-lanceolatis acuminatis basi ultra insertionem in auriculam brevissimam obtusam productis, calycis glabriusculi corollæ tubo subbrevioris ad $\frac{2}{3}$ vel $\frac{3}{4}$, fidi laciniis lanceolatis obtusiusculis, corollæ roseæ vel violacea laciniis obcordatis tubo tertiatâ parte brevioribus, capsulâ ovatâ calycem parum superante, seminibus oblongis utrinque acutis compressis minute scrobiculatis ♀. Ic. Vent. hort. Cels. tab. 42. — *P. longifolia* Curt. Bot. Mag. tab. 392. — *P. pyrenorrhiza* Led. Fl. Ross. III, p. 11! — *P. macrophylla* C. Koch Linn. XXIII, p. 613. — *P. Tournefortii* Rupr. Prim. p. 293.

Hab. in pratis alpinis humidis et turfosis, Olympus Bithynus (Boiss!, Thirk!, Pichler!), mons Anemas Lycaoniac 6000' (Heldr!). Taurus Cilicie 6500'-8000' (Ky!, Bal!), Cappadocia in monte Aslandach (Bal!) et Cataoniae Berytdagh 8000' (Haussk!), montes Galatiæ et Paphlagoniae (Wied!), Pontus occid. (Tchih!), Lazicus (Bal!), Armenia Turcica omnis et Armenia Kurdica (Auch. 5247!, Calv!, Huet!, etc.), Armenia Rossica (Nordm!, Szov!), Caucasus occid. et centralis 6000'-7000' (Rupr!, Rehm!) et orientalis 8500' (Rupr!), Persia in montibus Ghilani (Gmel., Buhse!), monte Ssahend pr. Aderbibjan (Buhse!), inter Siaret et Kohrud et alibus Khorassan (Bge!), monte Elvend (Mich., Haussk!), monte Kuhdaëna Persiaæ australis (Ky. 647!), Affghaniâ ad Hajiguk 11000' (Griff!).

β. glacialis. — Folia latiora, scapus apice calycesque subfarinosi, corolla major intensius lilacina tubo elongato laciniis latioribus. — *P. glacialis* Ad. in W. herb. — Rupr. loc. cit. p. 294.

Hab. ad moles glaciales Caucasi centralis in monte Kaischaur (Ad.), ad Baidara (Rupr!), in Chewsuriâ (Bayern).

Species formâ et longitudine foliorum, pedicellis plus minus abbreviatis in australioribus præsertim brevissimis ludens. Planta Sibirica ejusdem nominis ad *P. algida* referenda est.

15. P. luteola (Rupr. loc. cit. p. 298) glabra viridis, foliis planis levibus membranaceis longis lanceolato-ellipticis obtusis minute denticulatis in petiolum sensim attenuatis, scapo florifero elato cum calycibus apice diu pulverulento-farinaceo, umbellâ multiflorâ, pedicellis calyce 2-4-plo longioribus, involueri phyllis lanceolatis basi ultra insertionem subproductis, calycis tubo corollæ duplo brevioris ad $\frac{2}{3}$ partiti laciniis lanceolatis acutiusculis, corollæ pallide sulphureæ laciniis obcordatis tubo duplo brevioribus $\frac{2}{3}$.

Hab. ad fontes et prata madida alpinâ Caucasi in distr. Alagir et Tuschetia 6000'-9000' (Rupr!). V. s. sp. et v. cult.

Folia $\frac{1}{2}$ -1-pedalia, scapus pedalis et elatior. Species characteribus affinis *P. auriculata* a quâ præter florum colorem differt pedicellis elongatis, calycis et corollæ, tubi et limbi corollini proportione diversâ.

16. P. algida (Adams in Web. et Mohr Beitr. I, p. 46) albido et saepius flavo-farinosa rarius omnino denudata, foliis oblongo-spathulatis obtusis in petiolum brevem saepe obscurum attenuatis minute et acute denticulatis saepe subserratis, scapo folia superante, pedicellis calyce saepius brevioribus rarius æquilongis vel sublongioribus, involueri phyllis lanceolatis acuminatis basi subsaccatis, calycis corollæ tubo æquilongi vel subbrevioris ad medium fissi dentibus nigris lanceolatis obtusiusculis, corollæ intense roseo-violaceaæ laciniis obcordatis tubo subbrevioribus, capsulâ oblongo-cylindricâ calycem parum superante, seminibus ovato-oblongis parvis tuberculatis $\frac{2}{3}$. Ic. Lehm. Prim. tab. 7. — *P. farinosa* M.B. Taur. Cauc. — C.A. Mey. Enum non L. — *P. Caucasicæ* C. Koch Linn. XVII, p. 308. — *P. auriculata* β *Sibirica* Led. Fl. Ross. 3, p. 12. — *P. Bungeana* C.A. Mey.

Hab. in pratis humidis alpinis Armeniæ Turcicæ (Auch. 5233!), Ponti Lazici (Bal. exs. sub. *P. farinosa*!), Caucasi centralis 6000'-40000' (C.A.M., Rupr! Rehm!), prov. Schuragel, Armeniâ Rossicâ et Karabagh (Szov!).

Valde affinis *P. farinosa* pro ejus formâ saepe habita fuit et forsitan non immrito. *P. farinosa* differt foliis obtuse crenatis, calyce non colorato dentibus ovatis obtusissimis.

Ar. Geogr. Sibiria Altaica.

17. P. farinifolia (Rupr. Prim. loc. cit. p. 303) foliis subtus albo-farinosis teneris ovato vel oblongo-spathulatis obtusis argute serratis in petiolum eis longiore attenuatis, scapis etiam fructiferis foliis brevioribus vel ea æquantibus, pedicellis longissimis inæqualibus, involueri phyllis linearibus acutis basi non saccatis pedicellis pluries brevioribus, calycis ultra medium fissi laciniis acutis, corollæ roseæ tubo calyce subduplo longiore limbi laciniis tubo duplo brevioribus, capsulâ calycem non superante $\frac{2}{3}$.

Hab. in cryptis madidis umbrosis rupium in regione sylvaticâ Caucasi Orientalis sitarum 3600'-6600' in Chewsuriâ, Daghestaniâ et ditione Tindal et Bogos (Rupr!). Vid. spec. fructifera.

Affinis *P. algidæ* a quâ differt foliis teneris longe petiolatis, pedicellis elongatis 1-2 pollicaribus, corollæ tubo calyce duplo longiore. An tamen hujus forma ex loco umbroso?

18. **P. Darialica** (Rupr. Prim. loc. cit. p. 302) glabra, foliis obovato vel lanceolato-oblongis in petiolum eis multo breviorem attenuatis argute denticulatis, scapis folia 2-3-plo superantibus, pedicellis filiformibus florem æquantibus vel eo subdupo longioribus, involucri phyllis basi subsaccatis linearibus pedicellos non æquantibus, calycis glabri ad medium fissi laciñiis oblongis acutis sub lente denticulatis, corollæ roseæ tubo calyce sesquilongiore, limbi tubo vix brevioris laciñiis obcordatis, capsulâ... ♀.

Hab. in Caucaso ad fontes fluvii Terek prope Darial 3600' (Rupr!).

Affinis *P. farinifoliae* sed glaberrima et petioli breviores. Folia 1-2-pollicaria, scapus 2-4-pollicaris, corollæ limbus sex lineas diametro latus quâ nota ad *P. frondosam* accedit.

19. **P. frondosa** (Janka Pl. Turc. Brev. II, p. 6), albo-farinosa vel denudata viridis, foliis teneris ovato vel oblongo-spathulatis obtusis argute crenatis in petiolum saepe elongatum attenuatis, scapo folia æquante vel superante, umbellæ pedicellis gracilibus calycei æquilongis vel pluries longioribus, involucri phyllis linear-lanceolatis acutis basi non sacratiss, calycis ad vel ultra medium fissi laciñiis lanceolatis acutis, corollæ cæruleo-roseæ tubo calyce triente longiore limbi laciñiis profunde emarginato-bilobis tubo æquilongis, capsulâ calyce sublongiore ♀. *P. farinosa* var. *Turcica* Friv. exs. — *P. algida* Janka pl. exs.

Hab. in saxis præruptis udis umbrosis Thraciae borealis prope Kalofer (Friv! Janka! Pichler!).

Valde affinis *P. algidæ*, *farinifoliae* et *Darialicae* et forsitan melius nota in unicam speciem cum eis conjungenda, differt a priori foliis teneris, calycis laciñiis acutis, a posteriore corollæ limbo ampio, ab ultimâ calycis laciñiis longioribus.

20. **P. capitellata** (Boiss. Diagn. Ser. I, 7, p. 64) foliis subtus parce farinosis lingulato-lanceolatis subspathulatis in petiolum attenuatis parce obsolete et obtusiusculæ denticulatis, scapo folia multo superante, floribus brevissime pedicellatis capitatis, involucri phyllis oblongo-lanceolatis obtusis basi ultra insertionem productis pedicellos superantibus, calycis ad tertiam vel dimidiam partem fissi dentibus ovatis obtusis nigricantibus, corolle roseæ tubo calyce duplo longiore limbi laciñiis obcordatis tubo duplo brevioribus, capsulâ globosâ calycei æquilongâ ♀.

Hab. in pratis humidis alpis Kuh Daëna Persiae australis (Ky. 602!), montibus prope Jezl (Buhse!), in monte Elbrus prope Derbend Persiae bor. (Ky. exs. 148 sub *P. auriculatâ*), ad Hajiguk Affghaniæ (Griff!).

Folia cum petiolo 1½-2 pollicaria superne 3-4 lin. lata. Species foliis angus-

tis, floribus ob pedicellos breves interdum subnullos capitatis, calyce breviter fisco, corollæ tubo elongato et limbo minuto, capsulâ sphæricâ a *P. auriculata* et *algidâ* discedens.

SECT. IV. ARTHRITICA Duby ex parte. — Folia vernatione con-volutiva. Umbella pauciflora vel flores solitarii.

21. P. minima (L. Sp. 203) gregaria, rhizomate obliquo subtus fibras longas emitente, foliis parvis obovato-cuneatis glabris apice truncatis obtusis acute denticulatis, pedunculis 1-rarius 2-floris foliis brevioribus vel subnullis, involueri phyllis linearibus calycem subsequantibus, calycis tubuloso-campanulati laciniis linearis-spathulatis integris vel apice denticulatis, corollæ roseæ laciniis semibifidis lobis divaricatis, seminibus parvis argute angulatis ♀. Ic. Jacq. Austr. tab. 273. — Rehb. t. 59.

Hab. in summis alpibus Thracie in monte Rhodope prope Carlova (Friv!).

Ar. Geogr. Alpes Austriae et Italiae borealis, Sudeti, Carpathi, montes altiores Transylvaniae, Croatiae, Serbiae.

ORD. LXXI. MYRSINEACE.E.

(Alph. D.C. Mém. Ann. Hist. Nat. Par. Ser. II, 15, p. 65.)

Flores hermaphroditi polygami vel dioici. Calyx 4-6-fidus vel partitus. Corolla 4-6-fida vel partita. Stamina basi corollæ adnata numero ejus partium rarius dupla eis opposita. Ovarium (in nostris) liberum uniloculare pauciovulatum, placentâ centrali liberâ, stylus cylindricus, stigma capitatum vel lobatum. Drupa globosa extus subcarnosa intus cartilaginea vel lignea indehiscens. Semina pauca (in nostris unicum), globosa. Embryo filiformis plerumque arcuatus, albumen corneum vel subcarnosum. — Frutices vel suffrutices quâ notâ et fructu drupaceo a Primulaceis distinguuntur.

MYRSINE (L. Gen. 269).

Flores polygamo-dioici. Corolle lobi 4-5 aestivatione imbricato-subquincunciales. Stamina 4-5 libera. Antheræ filamentis brevibus multo longiores. Drupa intus crustacea. Semen membranis placentaribus tenuibus indusiatum. — Frutices vel arbustive.

1. M. Africana (L. Sp. 285) fruticosa, ramulis puberulis, foliis parvis glabris serrulatis breviter petiolatis coriaceis, fasciculis axillaribus sessilibus trifloris, corollæ calycisque lobi 4 evatis acutis, antheris corollam superantibus 5. *Frutex Ethiopicus foliis Myrtilli*

Breyn. Cent. I, tab. 5. — *M. bifaria* Wall. in Roxb. Fl. Ind. II, p. 296.

Hab. ad margines lapidosos rivorum Affghaniæ in regione montanâ et subalpinâ ad Otipore, Bharowul et Chuggur (Griff! Journ., 51, 86, 103 a).

Frutex 4-6-pedalis vel in altioribus humilior.

Ar. Geogr. P. Bon. Spei, Abyssinia, insulæ Azoricæ, India borealis.

MONOTHECA (Alph. D.C. in Deless. Ic. Select. V, tab. Prodr. VIII, p. 152.)

Calyx quinquepartitus. Corolla profunde et patentim quinquefida lobis obtusis, aestivatione quincunciali. Stamina fertilia 5 exserta lobis corollæ opposita, sterilia 5 lobis corollæ alterna eisque longiora lan- ceolato-subulata ananthera. Ovarium sericeum ovulis 5. Bacca subcoriacea glabra stylo terminata. Placenta punctiformis. Semen unicum ovatum immaturum tantum notum. — Frutex sempervirens facie *Sapotacearum*. — Genus cum *Reptonia Theophrastaceis* a cl. Alph. D.C. adnumeratum.

1. **M. Mascatensis** (Alph. D.C. loc. cit. p. 152) fruticosa, ad ramos, foliorum paginam inferiorem, calyces, tomentello-cinerascens, foliis coriaceis obovatis integris breviter petiolatis pallidis, fasciculis axillaribus 8-15-floris, pedicellis brevissimis 3. Ic. Deless. V, tab. 35.

Hab. in regno Mascate Arabiæ orientalis (Auch. 4916! et 4917).

Folia 12-15 lin. longa 6-8 lata, flores 2 lineas lati, bracteolæ ad pedicellorum basin.

REPTONIA (Alph. D.C. Prodr. VIII, p. 152. — *Edgeworthia* Falcon. non Meisn.).

Calyx 5-partitus. Corolla subcampanulata acute quinqueloba aestivatione contorto-imbricata intus squamis 5 adnatis (staminibus sterilibus) cum lobis alternantibus instructa. Stamina 5 exserta basi cum squamis lateraliter coalita. Ovarium pauciovulatum. Drupa monosperma ossea. Placenta minima, semen peltatum hilo lato umbilicato excavatum. Embryo intra albumen cartilagineum ruminatum excentricus. Arbuscula sempervirens. — Genus ob corollæ tubi appendices *Theophrastaceis* a cl. Alph. D.C. adnumeratum, ex am. Decaisne *Monotheca* identicum.

1. **R. buxifolia** (Falc. Ann. Hist. Nat. 1842, p. 362 sub *Edgeworthia*) arbuscula, ramis crassiusculis divergentibus albo-tomentellis rigidis apice spinescentibus, foliis alternis coriaceis ellipticis breviter petiolatis obtusis integris margine subrevolutis, floribus parvis sub-

sessilibus in fasciculos axillares dispositis, drupâ eduli dulci p. *R. buxifolia* A. D.C. loc. cit.

Hab. ad Khyber et Pushut Affghaniæ (Griff. Journ. N° I, p. 324).

Arbuscula 10-20-pedalis atrovirens facie *Oleæ Oleastri*, fructus in Pentapotamide *Gourgoura* dicti. Praeter ramos denum spinescentes *Monothecam Mascatensem* omnino refert et ab eâ ex cl. Decaisne specifice non differt; cl. Alph. D.C. eam autem corollæ aestivatione generice distinctam autumat. Flores non vidi.

Ar. Geogr. India boreali-occidentalis.

ORD. LXXII. EBENACEÆ.

(Vent. Tabl. p. 445.)

Flores regulares dioici raro subhermaphroditi. Calyx persistens 3-7-lobus. Corolla hypogyna vel perigyna decidua lobis 3-7 aestivatione sinistrorum convolutis. Stamina flor. masc. imæ corollæ inserta rarius hypogyna 6 vel plura saepius basi geminatum comata paribus binis ante basin enjusque lobi corollæ. Ovarium liberum 3-pluriloculare loculis saepius numero duplice loborum calycis et geminatum inter lobos approximatis, ovula in loculis solitaria vel gemina collateralia ex apice anguli interioris pendula anatropa. Styli distincti vel basi coaliti. Bæcca carnosa vel coriacea cortice interdum valvatum ab apice rupta, pluri-vel abortu paucilocularis. Semen pendulum in loculis saepius solitarium, albumen cartilagineum, embryo axilis.

DIOSPYROS (L. Gen. 4451).

Flores dioici. Calyx 4-6-lobus. Corolla tubulosa vel campanulata 4-6-fida. Stamina 8-50 saepius 16, florum fæm. pectoriora antheris effletis. Ovarium 8-12-loculare in flor. masc. abortivum, styli 2-4 basi plus minus connati. Bæcca globosa calyce saepius accrescente basi tecta 4-8-locularis. Semina oblonga dorso convexa lateraliter compressa. — Arbores vel frutices.

1. D. Lotus (L. Sp. 1510) arborea, foliis oblongis acuminatis integris breviter petiolatis subtus discoloribus puberulis, floribus minutis axillaris subsolitariis subsessilibus, calyce brevi acute quadrilobo ciliato et intus villoso, corollâ masculâ calyce triplo longiore urceolato-campanulata ad tertiam partem in lobos reflexos rotundatos ciliatos fissâ, bæcâ fulvâ calyce aucto suffultâ 5. Ic. Pall. Fl. Ross. tab. 58. — Rehb. tab. 38.

Hab. in sylvaticis Anatolie borealis ad Samsun et Amasia (Wied!), Ponti ad Trapezuntein (Bonrg!) et Rhizé (Ball), Transcaucasie (Lebed!), provincie Talyssch ad Lenkoran (Hoh!), prov. Ghilan et Mazanderan Persiae bor.

(Buhse!) et Asterabad ad Siaret usque (Bge!), Affghania (Griff.). In Thraciâ (ex Griseb.), Byzantii (ex Sibth.) et Tauriâ meridionali (Stev.) sed ibi prob. introducta. Fl. Jun. Tataris Tanriæ *Churna*, Persis *Churmandu* audit.

Fructus Ceraso paulo minor, recens acerbus, siccatus sat dulcis.

Ar. Geogr. Dalmatia, Helvetia australis (an spont?), China borealis.

ORD. LXXIII. AQUIFOLIACEÆ.

(D.C. Th. Elem. Ed. I, p. 217. — *Ilicineæ* Endl.)

Flores hermaphroditi vel unisexuales. Calyx dentatus aestivatione imbricatus. Corolla partita hypogyna regularis laciñis aestivatione imbricatis. Stamina corollæ inserta et ejus laciñis alterna. Discus nullus. Ovarium 2-6-loculare loculis 1-2-ovulatis ovinis pendulis. Fructus drupaceus indehiscent, albumen magnum, embryo rectus parvus, radicula ad hilum versa. — Ordo Celastrineis affinis sed corolla monopetala.

ILEX (L. Gen. 472).

Calyx persistens 4-5-dentatus. Corolla rotata 4-5-partita. Ovarium sessile subglobosum 4-5-loculare, stigmata 4-5 subsessilia. Drupa globosa pyrenis 4-5 crustaceis. — Frutices foliis coriaceis lucidis alternis.

I. **I. Aquifolium** (L. Sp. 181) fruticosa interdum arborea, foliis ovatis acutis coriaceis nitidis undulatis spinoso-dentatis vel integerrimis, pedunculis axillaribus brevisimis multifloris, floribus subumbellatis, drupis atrosanguineis 5. Ic. Engl. bot. tab. 496. — Rehb. tab. 39.

Hab. in sylvaticis regionis montanæ, montes Thessaliæ (Hawkins), Delphi Eubœa (Fraas!), Macedonia in peninsula Athoā 1200'-3000' (Griseb., Orph!), Thracia (Friv!), Byzantii in sylvâ Belgrad (Sibth.), Bithynia (Sest.), Pontus (Tchih.), Pontus Lazicus ad Trapezuntem et supra Rhizé (Ball), Imeretia, Mingrelia, Iberia Caucasia 1800'-3000' (C.A. Mey., C. Koch), prov. Talysh (C.A. Mey., Buhse!), Persia boreali-orientalis montes supra Siaret (Bge!). Fl. Maio. Vulgo Thessalis *Leopardina*, Persis *Rahsch.*

. **Ar. Geogr.** Europa omnis media et australis a Daniâ et Norvegiâ australi ad Germaniam, Croatiam et regionem Danubialem, Africa borealis.

ORD. LXXIV. STYRACACEÆ.

(D.C. Prodr. VIII, p. 244.)

Flores regulares hermaphroditi. Calyx ovario adnatus limbo quinquefido vel subintegro. Corolla perigyna monopetala profunde 5-7-

partita lobis cum illis calycis alternantibus. Stamina basi corollæ inserta ejus laciniis alterna et numero æqualia vel dupla, libera, pentadelpha vel monadelpha. Ovarium inferum vel semiinferum 3-5-loculare loculis lobis calycis oppositis, ovulis angulo centrali affixis, stylus simplex, stigma capitatum lobatum. Fructus baccatus vel siccus interdum sero dehiscens abortu unilocularis pauci saepius monospermus, semina albuminosa, embryo axilis.

STYRAX (Tournef. Inst. p. 398).

Calyx cupuliformis repando-quinque dentatus. Corolla tubus brevis, limbi laciniæ 5-7 aestivatione sinistrorum contortæ vel subvalvari. Stamina 6-16. Fructus ovatus coriaceus calyce persistente basi adnatus valvis tribus a basi vel ab apice dehiscens. Semen magnum ovato-globosum erectum hilo infero sublaterali. — Frutices vel arbusculæ floribus *Citrum Aurantium* referentes pilis stellatis tomentosæ.

I. S. officinale (L. Sp. 535) fruticosa vel arborea humilis, foliis petiolatis ovatis integris supra glabriusculis subtus tomentosis, cymis paucifloris ramos terminantibus breviter pedunculatis pedicellis calyce longioribus eoque incanis, corollæ albae tomentosæ laciniis lanceolatis calyce truncato 3-4-plo longioribus, staminibus sub-12, fructu ab apice valvatim dehiscente 5. Ic. Fl. Gr. tab. 375. — Rehb tab. 37.

Hab. in collibus umbrosis et ad rivos regionis inferioris et montane, Messenia (Bory!), Attica ad radices montium (Boiss!, Heldr!, Orph. Fl. exs. 5!), Parnassi radices (Sprnrl!), Creta (Sieb!, Raul!), Bithynia ad radices Olympi (Auch. 2544!), ad Trojam (Schmidt!), Rhodus (Bourg!), Lydia (Sibth.), Caria (Pinard!), Pamphylia (Bourg!), Cilicia 0-4500' (Kyt., Bal!), Syria borealis prope Marasch (Hanssk!) et ad radices Amanii (Ky!), Cyprus (Sibth.), Libani regio inferior (Labill., Ehrenbl., Bl!), radices montis Gebel Cheiekh Antilibani (Gaill!), Palestina (Rauw.). Fl. Maio: Graecis *Storaki* et *Agriokydonia*. *Arabis Syriacis Abhar* et Beduinis *Libna* audit.

Resina hujus arboris Balsamum Storax dicta in pectoris morbis adhibetur.

Ar. Geogr. Dalmatia, Italiae ager Romanus, Gallia meridionalis sed ibi prob. introductum.

ORD. LXXV. OLEACEÆ.

(Lindl. Nat. Syst. Ed. II., p. 307.)

Flores hermaphroditæ regulares rarius dioici apetali. Calyx persistens quadrilobus. Corolla regularis hypogyna monopetala quadrifida vel tetrapetala petalis per paria filamento connexis. Stamina binæ corollæ tubo vel petalis adnexa. Discus nullus. Ovarium bilocularē

loculis biovulatis, ovulis pendulis collateralibus. Stylus simplex vel nullus. stigma indivisum vel bilobum. Fructus baccatus, drupaceus, capsularis vel samaroideus, bilocularis vel abortu unilocularis, interdum monospermus. Semeni albuminosum embryone recto. — Arbores vel frutices.

TR. I. OLEINEÆ. — *Fructus carnosus.*

OLEA (L. Gen. 20).

Calyx brevis quadridentatus. Corollæ tubus brevis. limbus quadri-partitus patens. Drupa abortu 1-2-sperma carne oleosâ putamine osseo. — Arbores et frutices.

1. **O. Europæa** (L. Sp. I, p. 11), fruticosa vel arborea, ramis in plantâ sylvestri plus minus indurato-spinescentibus in cultâ inermibus, foliis oblongis vel lanceolatis coriaceis integerrimis mucronulatis subtus lepidoto-canis, florum racemis axillaribus, calyce cupulari subtruncato brevissime quadridentato, corollæ albae lobis obtusis, drupâ ellipsoideâ vel globosâ 5 et 5. Fl. Græc. tab. 3. — *Olea oleaster* Hoffm. et Link. Fl. Port. I, p. 387. — Rehb. tab. 33.

Hab. ubique culta in regione inferiore interdum usque ad 4800', multis locis in rupesribus vel spontanea vel subs spontanea, v. gr. Græcia total, Macedonia, Thracia et Bithynia (Griseb., Frivt.), Cilicia (Kyt.), Syria borealis mons Giaon-dagh et mons Sindjar Mesopotamie (Haussk!), Transcaucasia (Ledeb.), Persia borealis in valle Sefirud (Buhsel); specimen fructu juniori e monte Gebel Akadar Arabiae Mascatensis Anch. 494 hinc quoque spectare videntur. Græcis *Agriodæa*, Persis *Derechti Seitan*, Turcis *Jaban Seitan agagi* audit.

Ar. Geogr. Regio mediterranea Europe et Africe borealis, in regione subalpinâ montium Nubie australis ad mare Rubrum (Schweinfurth qui hac ex ea regione forsitan oriundam putat).

2. **O. cuspidata** (Wall. Cat. N° 2847, p.c. Prodr. VIII, p. 285) arborea, foliis coriaceis lanceolatis utrinque acutis cuspido-acuminatis supra nitidis subtus lepidoto-ferrugineis, florum racemis axillaribus, calyce cupulari truncato obsolete quadrilobo, corollæ albae lobis obtusis, drupâ ellipsoideâ 5.

Hab. in Belutschia (Stocks!), in Affghania ad Pnshut (Griff!).

Valde affinis *O. Europæa* a qua differre tantum videtur foliis magis acuminatis subtus ferrugineis.

Ar. Geogr. India boreali-occid. in Pontapotamide, ditione Sindh, Napaliâ

PHILLYREA (Tourn. Inst. tab. 367).

Calyx brevis quadrilobus. Corolla subrotata quadripartita. Anthereæ subsessiles. Drupa carnosa putamine chartaceo fragili. — Arbores et frutices foliis sempervirentibus coriaceis.

1. P. Vilmoriniana (Boiss. et Bal. in Bal. exs. 1866) fruticosa glabra, foliis magnis oblongo vel elliptico-lanceolatis basi cuneatâ breviter petiolatis acuminatis integerrimis margine subrevolutis subtus minute punctatis, floribus ad axillas fasciculatis pedicellis eis sublongioribus suffultis, calyce ad medium in lobos triangulares acentos fisso, corollæ lobis oblongis, drupâ ellipsoideâ obtusâ stylo mucronulatâ 3.

Hab. in montibus Ponti Lazici supra Rhizé ad pagum Andon et in parte inferiori vallis Khabackar 3600' (Bal!). Fl. Maio, fructus maturat Septembri

Frutex 9-10-pedalis, folia omnino *Cerasi Laurocerasi* 2 $\frac{1}{2}$ -4 $\frac{1}{2}$ pollices longa 1 $\frac{1}{2}$ poll. lata. Drupa 4 $\frac{1}{2}$ lineas longa, putamen congenitibus subdurans. Species insignis foliis magnis, calycis et drupæ forma.

2. P. media (Linn. Sp. p. 40) fruticosa ramis tomentellis, foliis mediocribus vel minutis breviter petiolatis vel subsessilibus ovatis vel oblongo-ellipticis acutis vel obtusiusculis integris vel plus minus crebre et acute serrulatis, floribus ad axillas fasciculatis pedicellis brevissimis vel eis sublongioribus suffultis, calycis lobis obtusis, corollæ lobis ovatis, drupâ parvâ globosâ muticâ vel apiculatâ 5. Rehb. Ic. tab. 34. — *P. latifolia* Fl. Græc. tab. 2. Griseb. Spic. et Rehb. Ic. non L. — *P. pedunculata* Bory et Chaub. Fl. Mor. p. 2.

Hab. in regione sempervirenti frequenter, Peloponnesus (Bory!), Attica in parte inferiori montium (Heldr! Samar!). Bœotia mons Onion usque ad 3500' (Heldr!), Macedonia (Friv!), Thracia (Griseb!), Byzantium (Auch!), Archipelagus, Creta (Sieb!), Anatolia occid. (Fleisch!) et meridionalis in Cilicia (Ky!), Syria in Libano (Bl!), Palestina in monte Carmelo (Gaill!), Anatolia bor. (Wied!), Armenia prope Gumuschkhané (Bourg!).

P. latifolia L. in Oriente a variis auctoribus indicatur, sed specimina omnia quæ vidi ad *P. medianam* nec ad *P. latifoliam* in Lusitanâ, Hispaniâ, Italia, Africâ boreali obviam spectant. Hac *P. latifolia* cæterum a *P. mediâ* nisi forsan staturâ majore et foliis junioribus cordatis vix sat differre videtur. *P. angustifolia* L. erroneè in Græcia et in prov. Caucasicis indicata fuit.

Ar. Geogr. Regio mediterranea Europæ a Gallia australi ad Dalmatiam, Africa borealis.

LIGUSTRUM (L. Gen. 22).

Calyx brevis campanulatus obtuse et breviter quadridentatus deciduus. Corolla infundibuliformis tubo calyce longiore, limbo quadrifido. Bacca globosa loculis binis membranâ tenui formatis dispermis vel abortu monospermis. — Frutices foliis oppositis.

1. L. vulgare (L. Sp. 49) fruticosa, foliis sero deciduis oblongis vel ellipticis mucronatis lucidis breviter petiolatis, floribus in paniculas thyrsoides densifloras terminales dispositis, baccis nigris 5 Ic. Rehb. tab. 33.

Hab ad sepes, in dumosis, ad margines sylvarum in australioribus montana, Peloponnesus in monte Chelmos (Orph!) et Olenos supra Kalavryta

3500' (Heldr!), Macedonia (Friv!), Thracia et Byzantium (Griseb!, Sibth.), Bithynia (Thirke!), Ionia (Sibth.), Pisidia (Tchih.), Anatolia borealis (Wied!), Pontus Lazicus (Bal!, Calv!), Caucasus et Transcaucasia 1800'-4800' (Ledeb., Szov!), ditio Talyisch ad mare Caspium (Hoh!), Persia borealis in prov. Ghilan (herb. Petrop!).

Ar. Geogr. Europa sept. et media ab Angliâ, Norvegiâ et Daniâ ad Rossiam meridionalem et Lusitaniâ Hispaniâque centrali, Italia ad Dalmatiam.

TR. II. LILACEÆ Vent. — Fructus siccus.

SYRINGA (L. Gen. 22).

Calyx brevis quadridentatus persistens. Corolla hypocrateriformis tubo longo limbo in lobos 4 partito. Stamina inclusa. Ovarium biloculare ovulis in loculis geminiis ex apice septi pendulis. Stigma bifidum. Capsula coriacea subcompressa oblonga bilocularis valvis navicularibus medio septiferis. — Frutices foliis oppositis.

1. S. vulgaris (L. Sp. 41) foliis longe petiolatis ovatis cordatis integerrimis acuminatis glabris concoloribus, thyrsis terminalibus multifloris, corollæ laciniis subconcavis 5. Ic. Rehb. tab. 32. — *Lilac vulgaris* Lam.

Hab. in sylvaticis rupestribus, Bithynia circa Brussam (Thirke!). An vere spontanea?

Ar. Geogr. Hungaria, Transylvania, Serbia.

2. S. Persica (L. Sp. 41) foliis breviter petiolatis oblongo-lanceolatis basi cuneatis et interdum pinnatifidis concoloribus, thyrsis terminalibus angustatis laxifloris, corollæ laciniis planiusculis 5. *Lilac Persica* Lam.

Hab. incerta, in Daghestaniâ Caucasicâ (ex Lerche Nov. Act. Phys. Leop. Carol. V. App. 184), ex Persiâ absque loco (Derderian in herb. Hort. Petrop! An spont.?)

FONTANESIA (Labill. Syr. Dec. I, p. 9, tab. 4).

Calyx brevis quadripartitus persistens. Petala 4 hypogyna per paria basi connexa. Stamina filamentis elongatis exserta. Ovarium 2-3-lokulare ovulis in loculis geminiis pendulis, stigma bifidum. Samara compressa margine membranaceo-alata bilocularis 1-2-sperma indehiscent vel tandem septicide bipartibilis. — Frutex facie *Ligustrum* ramulis et foliis oppositis.

1. F. phillyreoides (Lab. loc. cit.) fruticosa, foliis oblongo-lanceolatis integris acutiusculis basi cuneatâ breviter petiolatis sub-persistentibus, racemis axillaribus folio brevioribus, pedicellis flore

longioribus, petalis obovatis albis calyce multo longioribus, samarâ obovatâ vel oblongâ basi retusâ apice acutiusculâ vel retusâ 3. Ic. Lam. Ill. tab. 22.

Hab. in dumosis regionis inferioris et montanæ, Anatolia australis in Lyciâ et Pamphyliâ (Heldr! Bourg!), Ciliciâ a Mersina usque ad Gulek 3800' (Ky! Bal!), Cappadociâ (Tchih!), prope Marasch (Haussk!), Syriâ boreali in monte Amano (Ky!) et Cassio (Labill! Boiss!), Anatolia bor. in Troade (Tchih!) et ad Samsun (Wied!). Flor. Apr.

Frutex 10-20-pedalis ramulis crebre foliosis, fructus 3-4 lineas longus.

Ar. Geogr. Sicilia orientalis.

FRAXINUS (L. Gen. 1160).

Flores polygami vel dioici. Calyx 4-sidus vel nullus. Petala 4 bina-
tim basi cohærentia vel nulla. Ovarium biloculare septo valvis con-
trario, loculis uniovulatis, ovulis pendulis, stylus brevis. Samara com-
pressa oblongo-vel lanceolato-spathulata compressa in alam producta
abortu unilocularis monosperma. — Arbores vel arbuseculæ.

SECT. I. ORNUS Pers. Ench. II, p. 605. — Flores calyce
corollâque prædicti.

1. **F. Ornus** (L. Sp. 1510) arborea, foliis synanthiis 3-4-jugis
foliolis petiolulatis denticulatis lanceolatis vel ovato-rotundatis acu-
miatis glabris vel subtus ad nervum barbatis, florum racemis ex
axillis supremis et terminalibus pedunculatis paniculam terminalem
foliis breviorem formantibus, petalis linearibus calyce multoties lon-
gioribus, samaris anguste linear-lanceolatis utrinque attenuatis vel
subspathulato-truncatis retusis 3. Ic. Fl. Græc. tab. 9. — Rehb. Ic.
tab. 31. — *F. rotundifolia* Lam. Dict. II, p. 546 (forma foliolis rotun-
datis). — *F. diversifolia* Rochel Bann. — *Ornus Europea* Pers.
Ench. I, p. 9. — *F. Cappadocica* Juss. herb. ex K. Koch. Dendrol. —
Fr. millelacuum C. Koch Linn. XXIII, p. 597.

Hab. in regione montanâ Græciae. Tylgetus (Boiss!), Malevo (Orph. Fl. Gr. exs. 706!), Parnassus (Sibth.), Olympus Thessalus 5000' (Orph. Fl. Gr. exs. 705!), Macedonia et Thracia 0-2000' (Griseb., Friv!), Bithynia (Thirket!), Ionia in montibus Smyrne et in Sipyo (Boiss! Bal!), Cilicia ad Tauri radices 3800'-4500' (Ky! Bal!), Libanus prope Eden (Reyg!). Vulg. Græcis *Meleos*.

Ar. Geogr. Hispania in regno Valentino, Helvetia et Germania australes,
Italia, Dalmatia, Bosnia, regio Danubialis.

SECT. II. FRAXINASTER. — Flores nudi calyce corollâque destituti.

2. **F. excelsior** (L. Sp. 1509) arborea, gemmis nigris, foliis
hysteranthiis 4-6-jugis magnis subsessilibus lanceolato-oblongis utrin-

que attenuatis serratis saepe acuminatis glabris vel subtus ad nervum puberulis, floribus polygamis in racemos breves ad ramorum apicem approximatos erectos dein pendulos dispositis, samaris ellipticis apice obtusis vel retusis rarius acutatis 5. Rehb. t. 31.

Hab. in sylvis Albaniæ et Epri (ex Griseb.) et prob. Macedoniae borealis, Ponti maritimi (Tchih.), Tauriæ (Stev.), Caucasi et Transcaucasiæ (Ledeb.), Colchidis (Rupr!).

β . *coriariæfolia*. — Folia subtus undique velutina. — *F. coriariæfolia* Scheele Linn. XVII, p. 350.

Hab. in sylvis septentrionalibus montium Talysch versus desertum Mugan (Hoh!).

γ . *petiolulata*. — Foliola magna crenulata longe attenuato-acuminata basi longiuscule petiolulata. — *F. petiolulata* Boiss. Diagn. Ser. II, 3, p. 119.

Hab. in Tauro Cilicio (Ky. 356 a!).

Varietas ob foliola magna lata 3½-4 pollices longa huc potius quam ad formas *F. oxyphyllæ* referenda, samara 15 lin longa 4 lata acutiuscula.

Ar. Geogr. Europa omnis septentrionalis et media ad Hispaniam borealem, Italiam medium et Dalmatiam usque.

3. ***F. oxyphylla*** (M.B. Taur. Cauc. II, p. 450) arbuscula, gemmis cinereo-fuscis, foliis hysteranthiis 3-4-jugis foliolis mediocribus oblongis et oblongo-lanceolatis acuminatis basi sessili attenuatis serratis supra glabris subtus ad nervum interdum hirtis, floribus polygamis in racemos breves ad ramorum apicem approximatos erectos dispositis, samaris ellipticis vel elliptico-lanceolatis ntrinque attenuatis acutis vel obtusis 5. *F. rostrata* Guss. pl. rar. 374, tab. 63. — *F. oxyacarpa* Willd. Sp. IV, p. 1100. — *F. Ornus* Pall. Ind. Taur. non L.

Hab. in sylvaticis prope Byzantium (Pest., Noé! Janka!), Bithyniæ (Wied!), Smyrnæ (Bal. 381!), Tauro Cilicio (Ky! Bal!), Cataoniæ (Haussk!), Ponto ad Trapezuntum (Bourg!) et Rhizé (Bal!), Tauriæ meridionali (Stev!), Sonchetiæ et Armeniæ (Szov!), prov. Karabagh (Hoh!).

β . *oligophylla*. — Folia glaucescentia rigidiora 5-3 interdum indivisa. Samara abbreviata oblongo-elliptica obtusa vel acuta. Forma magis australis. — *F. Syriaca* Boiss. Diagn. Ser. I, 11, p. 77.

Hab. in Syriæ ad Aleppo juxta fluvium (Ky. 22! sub *F. lentiscifolia* Boiss!, Haussk!), ad Damascum et in Antilibano (Gail!), ad Berythum (Bl!), in Kurdistanæ Persiae monte Zagros (Haussk!), in jugo Elbruseensi Persiae borealis (Auch! Buhse! Ky. 189 sub *F. rostrata*!), prov. Khorassan (Bunge!), Affghanæ (Griff!).

γ . *parvifolia*. — Foliola plurijuga minora ovata vel ovato-spathulata argute dentata saepe obtusa. Potius forma fruticosa vel junior sterilis speciei quam varietas distincta. — *F. parvifolia* Lam. Dict. II, p. 540 — *F. lentiscifolia* Desf. Arb. I, p. 102.

Hab. in montanis et alpinis, mons Balouklardan Lyciæ (Bourg!), mons

Ssoddaghi Syriae 3500' (Haussk!), Armeniae australis alpes 5500'-8500' (Koch, Ky. Suppl. 666!), mons Teng Nalli Persiae australis 700' (Haussk!).

δ. subintegra. — Fruticosa, foliola parva subintegra terna ovata vel oblonga attenuato-cuspidata. — *F. Persica* Boiss. Diagn. Ser. I, 11, p. 78.

Hab. in regione alpinâ Persiae australis, mons Kuh Daena (Ky. 646!), Kuh Nur 9000' (Haussk!), Persia (Auch. № 2548!).

Species foliolorum numero formâ et magnitudine, fructûs formâ valde variabilis et quæ notis nimis incertis a *Fraxino excelsiori* distincta hinc forsū quondam specifice conjungetur. Arbor minor tamen, foliola magis coriacea magis glauca, minora, samara minor. *F. angustifolia* Vahl Hispanica formis stenophyllis *F. oxyphylla* prob. adiunctoranda est.

Ar. Geogr. Lusitania et Hispania, Africa borealis, Gallia et Italia australes, regio Danubialis, Rossia australis.

4. F. Sogdiana (Bge Pl. Lehm. p. 390) arborea, gemmis castaneo-fuscis villosis, foliis glabris coriaceis aliis simplicibus aliis 3-5-foliolatis foliolis magnis petiolulatis ovato-oblongis basi cuneatis integrerrimis superne grosse et remote paucidentatis, racemis simplicibus et compositis in apice ramorum congestis, floribus plerisque hermaphroditis, samaris oblongo-ellipticis obtusiusculis apiculatis latitudine suâ quadruplo longioribus 5.

Hab. in rupestribus Turkestaniae (Lehm!), Buchari vulgo *Ssaur madschu*.

Ex unico et incompleto specimine mihi nota, affinis *F. oxyphyllæ* var. *oligophyllæ* a qua differre videtur foliolis grosse et paucidentatis, inflorescentiæ bracteis et bracteolis villosissimis, samaris latioribus (6 lineas latis).

5. F. xanthoxyloides (Wall. Cat. 2833. — n.c. Prodr. VIII, p. 275) fruticosa, foliis bijugis foliolis subsessilibus oblongis obtusiusculis superne crenulatis basi attenuatis, florum fasciculis lateralibus oppositis et terminalibus globosis, petalis., samaris elevatim lanceolatis elliptico-linearibus apice retusis 5. *F. Moorcroftiana* Wall. Cat. № 2834.

Hab. in montibus Belutschiae 6000'-9700' (Stocks exs. 900!).

Florum fasciculi magnitudine avellanae, foliola in specimine Stocksiiano 9-10 lineas longa, samarae 14-15 lineas longa 3 latæ. *F. Moorcroftiana* specimen ex herb. Falconer tantum fructu et foliolis majoribus differre videntur.

Ar. Geogr. Jugum Himalaicum occidentale.

ORD. LXXVI. JASMINEÆ.

(R. Br. Prodr. p. 520.)

Flores regulares hermaphroditi. Calyx persistens dentatus vel lobatus. Corolla hypogyna hypocrateriformis lobis 5-8 æstivatione spiraliter convolutis. Stamina bina inclusa tubo inserta. Ovarium disco-

destitutum bilocularē apice bilobūm, ovula in loculo erecta solitaria rarius gemina. Stylus simplex, stigma bilobūm. Fructus vel bibaccatus vel in 2 capsulas bipartibilis valvis medio septiferis. Semina albu-mine parco vel nullo, embryone recto.

JASMINUM (Tourn. Inst. 597).

Calyx campanulatus 5-8-dentatus. Corollæ tubus longus teres, limbus planus partitionibus obliquis. Bacca didyma loculis monospermis.
— Frutices dumosus vel scandentes.

1. J. fruticans (L. Sp. 9) glabrum dumosum non scandens, ramis striato-subangulatis, foliis alternis trifoliolatis nonnullis simplicibus foliolis subcoriaceis spathulato-oblongatis, floribus 2-4 ad ramulorum extremitatem, calycis dentibus subulatis tubo $1\frac{1}{2}$ -3-plo longioribus, corollæ flavæ tubo calyce duplo longiore laciniis 5 obtusis 3. Ic. Bot. Mag. tab. 461. — Rehb. tab. 36. — *J. Syriacum* Boiss. et Gaill. Diagn. Ser. II, 6, p. 120 (forma floribus submajoribus et calycis laciniis longioribus).

Hab. in collibus regionis montanæ et sempervirentis, Thracia (Friv!), Bithynia (Griseb! Thirk!), Lydia ad Smyrnam (Boiss!), Anatolia borealis usque ad 3000' (Tchih! Wiedem!), Cilicia (Kyt!), Syria borealis in montanis prope Marasch et Aintab (Hausk!), Ouadi Benka Antilibani (Gaill!), Transcaucasia (Ledeb.), Iberia prope Tiflis 2000' (Rupr!), ditio Talyssch 1800'-3000' (C.A. Mey.), Persia borealis (Bge!).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia borealis, Africa borealis.

2. J. humile (L. Sp. p. 9) glabrum dumosum non scandens, ramis teretiusculis, foliis alternis trifoliolatis foliolis ovatis terminali acutato, floribus 1-3 subterminalibus, calyce brevissime quinquedentato, corollæ flavæ tubo calyce multoties longiore laciniis 5 obtusis 5. Ic. Bot. Reg. t. 350.

Hab. in collibus siccis Messeniæ (ex Bory Fl. Pelop.), insulae Chio (ex d.c. Prodr.), Babyloniae prope Bassora (Noë 60!). An vere spontaneum?

Affine *J. fruticantis* differt ramis non angulatis, foliolis majoribus latioribus basi minus attenuatis, calycis dentibus brevissimis. In insulis Canariensis indicatum ibi ex Webb Phyt. non crescit.

3. J. revolutum (Sims Bot. Mag. tab. 1731) glabrum vel puberulum non scandens, ramis angulatis, foliis alternis pinnatis foliolis 3-5-7 ovatis aut ovato-oblongis acuminatis, paniculis terminalibus oppositifoliis corymbosis, calycis dentibus 5-6 brevissimis acutis, corollæ luteæ tubo calyce multoties longiore lobis apice subrevolutis 5.

Hab. in Affghaniâ (Griff. ex Aitch. Catal. Punjab p. 87). Editione nondum vidi.

Ar. Geogr. India borealis.

4. J. officinale (L. Sp. 9) glabrum subscandens, ramis striato-subangulatis, foliis oppositis pinnatis trijugis cum impari foliolis oblongo-lanceolatis acuminatis, paniculis terminalibus paucifloris corymbosis, calycis laciniis 5 subulatis longis, corollæ albae tubo calyce sublongiore lobis ovatis acutis 5. Ic. Bot. Mag. tab. 31. — Rehb. t. 36.

Hab. in rupestribus et dumosis Imeretiae et Iberiae (M. B., Ledeb.), Persia borealis prope Derbend (Ky. 438!), Ghilano ad sylvarum margines (Buhse!). in montanis et alpinis prov. Khorassan inter Nichapur et Mesched (Bge!). In Macedoniâ prope Vodena (Orph!) et alibi in ditione sed vix spontaneum.

Ar. Geogr. Regio Himalaica, in omni Europâ australi introductum et subspontaneum.

ORD. LXXVII. SALVADORACEÆ.

(Lindl. Nat. Syst. Ed. II, p. 269.)

Flores dioici vel subunisexuales. Calyx gamoscpalus quadrilobus. Corolla hypogyna profunde 4-fida vel 4-petala lobis cum calycinis alternantibus. Stamina 4 corollæ lobis alterna denticulis minimis interdum cum filamentis alternantibus. Discus nullus. Ovarium liberum 1-2-loculare, oviulis 1-2-4 erectis anatropis. Stigma sessile bilobum. Drupa globosa saepius monosperma, semen subglobosum, albumine nullo, cotyledonibus basi cordato-auriculatis, radiculâ inferâ. — Folia opposita petiolo articulato, stipulis minimis spinuliformibus. — Ordo transitus inter Polypetalas et Gamopetalas præbens inter priores *Celastrineis*, inter posteriores *Oleaceis* et *Jasminaceis*, hisce ovulorum situ proprietor.

SALVADORA (Garcin ex L. Gen. Ed. VI, p. 163).

Flores subunisexuales. Calyx semiquinquefidus. Corella persistens profunde quadrisida. Ovarium in floribus fertilibus uniloculare uniovulatum. Bacc paree carnosa. — Frutices foliis coriaceis pallidis.

1. S. Persica (Garcin loc. cit.) fruticosa glabra, ramis opposite ramulosis saepè dependentibus, foliis ovato-oblongis obtusis basi attenuatâ brevissime petiolatis, racemis ex axillis superioribus spiciformibus saepè compositis paniculam foliosam formantibus, floribus minimis breviter pedicellatis, calyce campanulato, corollâ calyce duplo longiore lobis revolutis 5. Ic. Lam. Ill. tab. 81. — Roxb. Corom. tab. 26. — Vahl Symb. 1, tab. 4. — Decaisne Voy. Jacquem. tab. 144, fig. I. — *Rivinia paniculata* L. Syst. 10, p. 889. — *Cissus arborea* Forsk. Descr. p. 32. — *Embelia Burmanni* ex parte et *E. Grossularia* Retz Obs. 4. p. 24.

Hab. in littore Ægyptiaco maris Rubri! (Raddi! Acerbi!), inter Kosseir et Ras Benass (Schweinf!).

Folia 1-2-pollicaria latitudine varia basi obtusa vel acuta, pedicelli 1-2 milim. longi, baccaæ parce carnosæ magnit. grani Piperis vel majores. Frutex vel arbuscula subtropica probabiliter in Belutschia calidiore quoque crescents, ut et *S. oleoides* Decaisne in Jacquem. p. 140, tab. 144 species affinis foliis angustis mucronato-acuminatis, racemis brevibus, corollæ lobis non revolutis diversa.

Ar. Geogr. Nubia, Abyssinia, Algeria interior, Arabia tropica, India boreali-occidentalis et tropica.

ORD. LXXVIII. APOCYNEÆ.

(R. Br. Prodr. 465.)

Calyx quinquepartitus persistens. Corolla monopetala hypogyna regularis quinquefida decidua laciniis æstivatione oblique contortis, fauce nudâ vel laciniis aut coronâ aucta. Stamina quinque corollæ inserta ejus laciniis alterna, filamenta libera, antheræ stigmati incumbentes, pollen granulosum. Ovaria duo unilocularia multiovulata, rarius ovarium unicum biloculare. Styli 2 in unum coaliti, stigma I. Fructus follicularis, capsularis, drapaceus vel baccatus. Semina sæpius pendula varie comosa vel alata, rarius calva, albuminosa, embryone recto.

CARISSA (L. Mant. p. 52).

Calyx quinquepartitus lobis quincuncialibus basi eglandulosis. Corolla hypocateriformpha tubo interne piloso. Antheræ filamento longiores lanceolatæ. Ovarium unicum fusiforme glabrum biloculare, stigma bilobum pilosum caducum. Ovula pauca dissepimento affixa. Bacteria dulcis bi vel unilocularis 1-3-sperma, semiua calva peltata. — Frutices vel arbusculæ lactescentes foliis oppositis integris, spinis oppositis inter folia opposita sæpe obviis, floribus albis, facie *Jasmini*.

t. C. edulis (Vahl Symb. I, p. 22) fruticosa, ramulis velutinis spinosis, foliis breviter petiolatis ovatis rigidis, pedunculis ex axillis supremis terminalibusque brevissimis dichotome 3-5-floris pedicellisque puberulis, calycis laciniis lanceolato-acuminatis ciliatis, corollæ tubo extus glabro 3. Ic. Del. pl. Caill. tab. 2. — Jaub. et Sp. Ill. tab. 495.

Hab. in Ægypto superiore (Caill.).

Flores 5 lineas longi, bacca ellipsoidea 3 lineas longa disperma.

Ar. Geogr. Nubia, Abyssinia, Arabia tropica.

VINCA (L. Gen. 295).

Calyx quinquepartitus lobis acuminatis. Corolla hypocrateriformis fauce pentagonâ nudâ, limbi laciniis apice oblique truncatis. Antheræ filamento multo longiores supra pistillum conniventes. Stylos indivisi-
sus sub apice piloso annulo stigmatifero cinctus. Folliculi bini lœves striati. Semina nuda. — Herbae foliis oppositis.

1. **V. major** (L. Sp. 304) caulis sterilibus longissimis prostratis non radicantibus, floriferis brevioribus erectis, foliis breviter petiolatis amplis ovatis apice attenuatis obtusisculis basi saepe cordatis facie glabris margine ciliatis, pedunculis ex axillis superioribus solitariis folio brevioribus, calyeis laciniis linearibus ciliatis acumi-
natis tubum corollæ subæquantibus ♀. Engl. Bot. t. 514. — Rehb. Ic. tab. 22.

Hab. in umbrosis regionis inferioris insularum Chios (Orph!), Rhodi (Auch. 1504!), Creta in saxosis umbrosis prope Candium (Heldr!), Bithynie prope Brussam (Thirkel!).

β . *pubescens*. — Folia ad nervos caulesque patule pilosi, corolla saepe minor. — *V. pubescens* Urv. Enum. p. 26.

Hab. in Ponto ad Samsun (Tehih!), Ponto Lazico ad Rhizé (Ball!). Colchide (Urv.).

Intermediis ad typum manifeste transit.

Ar. Geogr. Lusitania, Hispania, Gallia occidentalis media et australis, Italia, Helvetia australis, Istria, Africa borealis.

2. **V. minor** (L. Sp. 304) caulis sterilibus prestratis radican-
tibus, floriferis brevibus erectis, foliis brevissime petiolatis glabris ellipticis vel ovato lanceolatis utrinque attenuatis obtusis, pedunculis ex axillis superioribus solitariis folio æquilongis vel longioribus, calyeis glabri laciniis breviter lanceolatis obtusis tubo corollino multo brevioribus ♀. Engl. Bot. t. 917. — Rehb. Ic. tab. 21.

Hab. in umbrosis circa Byzantium (ex Bergg.), Bithynie prope Brussam (Thirkel!). In Tauria et Caucaso a M. B. et aliis indicata sed prob. confusione cum *V. herbacea*.

Ar. Geogr. Europa media ab Anglia et Daniâ ad Rossiam medium, Hispania boreali et Italiam ad Dalmatiam.

3. **V. herbacea** (W. K. Pl. rar. Hung. p. 8, tab. 9) caulis omni-
bus proeminentibus vel declinatis basi non radicantibus, foliis brevissime petiolatis inferioribus ovatis superioribus elliptico-lanceo-
latis vel ovato-oblongis utrinque attenuatis omnibus margine brevis-
sime sebris, pedunculis axillaribus folia æquantibus vel superan-
tibus, calycis laciniis linear-lanceolatis acutis margine breviter ciliato-
sebris corolle tubo duplo brevioribus ♀. Rehb. Ic. tab. 22.

Hab. in frutetis et umbrosis siccis regionis montanæ, Peloponnesus (Bory!), Heldr! montes Malevo et Kyllene (Orph. Fl. exs. 377!), Parnassi regio inferior (Heldr. herb. Norm. 698!), Zacynthus (Margt!), Thracia (Griseb!), montes Simyrnæ (Boiss!), Carie (Piu!), Phrygiae (Bal!), Anatolia bor. (Wied!), Taurus Cilicius ad Gülek Boghaz (Ball!), Palestina ad Ramleh (Ky. 617!) et ad meridiem Gaza (Boiss!), Armenia Turcica ad Gumuschkhané (Bourg!), Tauria (Stev!), Caucasus (M.B.), Transcaucasia (Hohenf!).

Variat foliis plus minus latis; forma latifolia in Oriente frequentior in Austria quoque occurrit!

Ar Geogr. Austria, regio Danubialis, Rossia media et australis.

. 4. **V. Libanotica** (Zucc. Act. Acad. Monac. III, p. 246. tab. 8) canibus omnibus procumbentibus vel declinatis basi non radicantibus, foliis brevissime petiolatis vel sessilibus glabris margine laevibus, pedunculis axillaribus folium æquantibus vel eo brevioribus, calycis laciniis anguste linearibus acuminatis glabris corollæ tubo duplo brevioribus, laciniis corollinis valde obliquis acutis φ . *V. sessilifolia* D.C. Prodr. VIII, p. 383.— *V. Bottae* J. et Sp. Ill. Or. tab. 186.

Hab. in monte Alemdagh Anatoliae borealis (Noë!), Cappadociâ Orientali (Auch. 1498!), Ciliciâ (Auch. 1503!), Syriâ ad Aleppo et Aintab (Ky. 51! Haussk!), Libano interiore usque ad Cedretum 6500' (Ehrenb! Boiss! Roth! Ky. 303!), Antilibano (Gail!), Mesopotamiae desertis et montibus (Botta! Ky! Haussk!), Persia boreali (Buhsel! Bge!).

Valde affinis *V. herbacea* et forsitan hujus varietas glabra, caules sapius robustiores et folia majora, corollæ laciniæ acutiores sed haec nota forsitan constans.

RHAZYA (Decaisne Ann. Sc. Nat. Ser. II, 4, p. 80).

Calyx parvus quinquepartitus. Corolla hypocraterimorpha tubo longo a medio dilatato, fance hispidâ non appendiculatâ, limbi parvi laciniis ovatis. Stamina medio tubo inserta filamentis brevissimis, antheris liberis. Stylus unicus, stigma rotundato-depressum bilobum basi membranâ cupuliformi cinctum. Folliculi bini teretes lèves erecti. Semina oblonga basi et apice nivatâ compressione membranacea. — Fructices et suffractices foliis alternis.

. 1. **R. stricta** (Decaisne loc. cit. p. 81) fruticosa glaberrima, ramis simplicibus brevibus strictis crebre foliosis, foliis coriaceo-carnosulis aveniis oblongo-lanceolatis acutissimis basi attenuatis brevissime petiolatis, cymis terminalibus densiusculis e foliis supremis vix exsertis, calycis corollâ 6-7-plo brevioris laciniis triangulari-lanceolatis acutis, corollæ albae lobis ovatis mucronatis tubo sextuplo brevioribus φ . Ic. Jacqemont Voy. tab. 411.

Hab. in Belutschia (Stocks!), in Affghaniâ ad Pushut et collum Khyber (Griff!), regno Mascate Arabie (Auch. 4922!).

Frutex tripedalis facie *Daphnes Laureole*, folia 2-3 pollices longa 5-8 lineas lata, corolla 4-5 lineas, folliculum tenue cylindricum 2 pollices longum.

Ar. Geogr. Arabia tropica (Ehrenb. sub nomine *Vince Nerii*, Schimpf.), ditiones Scinde et Pundjab Indie borealis.

2. R. Orientalis (Decaisn. Jacq. Voy. Bot. p. 105 sub *Amsonia*) fruticosa, ramis simplicibus brevibus strictis crebre foliosis, foliis membranaceis ad marginem et superne ad nervum medium pilosiusculis ovato-lanceolatis acuminatis basi rotundata subsessilibus, nervis secundariis subtus crebris, cymis terminalibus e foliis supremis non exsertis, calycis corollæ sextuplo brevioris laciniis lanceolatis aentis apice pilosulis, corollæ lobis oblongis obtusis tubo quadruplo brevioribus 5. *R. Orientalis* n.c. Prodri. VIII, p. 386.

Hab. in Bithyniâ ad Brussam et lacum Apollonias et in Phrygiâ (Auch 789 et 2488!).

Folia bipinnicaria 7-9 lineas lata, corolla pyriformis 7 lineas longa.

NERIUM (L. Gen. 484).

Calyx quinquepartitus laciniis linear-lanceolatis basi intus multiglandulosus. Corolla hypocraterimorpha limbo quinquepartito, fauce coronâ exsertâ lacero-multitudinâ aenâ. Stamina medio tubo inserta, filamenta ligulata, anthera filamento longiores medio stigmati adhaerentes basi sagittatæ apice in setas elongatas pilosas spiraliter contortas desinentes. Stylus indivisus, stigma obtusum. Folliculi bini adpressi tandem segregati. Semina numerosa comosa. — Fructus lactescens foliis oppositis vel ternatis.

1. N. Oleander (L. Sp. 305) fruticosum, ramis strictis, foliis crebris strictis oppositis vel ternatis coriaceis lanceolatis acuminatis basi breviter petiolatâ attenuatis subtus minute et obscure tomentellis, floribus terminalibus corymboso-cymosis, pedicellis calyceque tomentellis, corolla roseæ limbo patenti, appendicibus 3-4-dentatis dentibus lanceolato-acuminatis inaequalibus, antherarum setis pilosis antherâ dorso pilosâ duplo longioribus linear-spathulatis faucem vix superantibus 5. Ic. Fl. Greec. tab. 248. — Rehb. t. 23.

Hab. ad rivos regionis inferioris usque ad 4000' interdum crescentes, Graecia omnist!, Archipelagus (Urv!), Creta (Rau!), Rhodus (Bourg!), Macedonia in peninsula Athoâ (Friv!), Byzantium (G. Koch), Anatolia occidentalis a Bithyniâ ad Cariam (Tchih.), Cilicia (Ky.), Cyprus (Ky.), Syria maritima (Labili, Bl!), Palestina ad lacum Tiberiadem (Boiss!). Mesopotamia ad fl. Zab (Haussk!), Graecis *Pierodaphne* audit.

Ar. Geogr. Lusitania, Hispania australis et orientalis, Gallia austro-orientalis, Italia, Dalmatia, Africa borealis.

2. N. odoratum (Solander in h. Kew. Ed. I, p. 296) fruticosum, ramis strictis, foliis ternatis linear-lanceolatis utrinque aentis brevisime petiolatis subtus adpressissime caescenti-lepidotis, cymis terminalibus multifloris densis, pedicellis calyceque caescentibus, corollæ

roseæ limbo patenti, appendicibus loborum in laciniæ 4-7 anguste lineares ad basin usque partitis, filamentis antherisque dense pilosis eis caudâ triplo longiori e fauce exsertâ superatis ȝ. Sims Bot. Mag. tab. 2032!

β. Kotschyi. — Folia crassa firmiora margine revoluta subtus magis lepidota nervis secundariis magis prominulis, corollæ limbus diminutus subcampanulatus. — *N. Kotschyi* Boiss. Diagn. Ser. I. 7, p. 21.

Hab. ad rivos Belutschiae (Stocks!), Affghaniae (Griff.), Persie austro-orientalis inter Khabbise et Kerman (Bunge!), in lapidosis alpis Kuh Delu Persie austro-occidentalis (Ky. 558!) forma humilior foliis diminutis.

Folia eis *N. Oleandri* angustiora, flores minores.

Ar. Geogr. India borealis.

3. N. Mascatense (Alph. d.c. Prodr. VIII, p. 421) fruticosum, foliis ternatis longe lanceolatis acutis basi attenuatâ brevissime petiolatis subtus adpressissime lepidoto-tomentellis, cymâ terminali folia vix superanti, pedicellis calyceque canescensibus, corollæ roseæ vel albae limbo campanulato laciniis ante lobum unumquemque binis linearibus exsertis, filamentis antherisque dense pilosis eis caudâ duplo longiori exsertâ superatis ȝ.

Hab. in regno Mascate Arabiae (Auch 4925!), in glareosis torrentium regionis inferioris montis Kuh Nur Persie austro-occidentalis 2000'-4000' (Haussk!).

Foliorum consistentiâ floribusque eis *N. Oleandri* minoribus valde affine *N. odoro* ȝ *Kotschyi* a quo laciniis ad loborum basin binis nec 4-7 fere tantum differt An ejus varietas?

APOCYNUM (L. Gen. 187).

Calyx quinquepartitus laciniis eglandulosis. Corolla parva campanulata quinquefida. Appendices 5 triangulares e basi tubi cum lobis alternantes. Stamina basi tubi inserta, filaments ligulata, antheræ filamento longiores sagittatæ medio stigmati adhaerentes membranâ acutâ terminatae. Nectarium glandulis 5 constans. Stylus subiullus. stigma ovoideum dilatum apice subbilobum. Folliculi bini graciles elongati. Semina longe comata. — Herbæ basi suffrutescentes.

1. A. Venetum (L. Sp. 311) caulis elatis paniculatis, foliis ovato-oblongis, ellipticisque obtusis mucronulatis margine cartilagineis subdenticulatis glabris, cymis terminatibus compositis, calycis breviter canescens laciniis triangularibus acutis, corollæ purpurascens tomentelle lobis tubo æquiflongis ȝ. Ic. Lobel. 372.

Hab. in uliginosis saepius salsuginosis, in maritimis ad Byzantium (Sibth. Auch. 2818!), Smyrna (Bourg!), Syria bor. ad Marasch (Haussk!), Kurdistania (Ky. 399!), Ciscaucasia et Transcaucasia (MB., Hohen!), Persia borealis (Szov!, Buhse!, Ky!), anustralis in montibus Kuh Kilouyeh 7000' (Haussk!), Turkestania (Lehm!), Persice Murd.

Ar. Geogr. Italia boreali-orientalis, Rossia media et australis, Songaria, China borealis.

ORD. LXXIX. ASCLEPIADEÆ.

(R. Br. in Wern. Soc. I. p. 12.)

Calyx quinquepartitus persistens. Corolla monopetala hypogyna regularis decidua rotata, campanulata vel infundibuliformis, fauce nudâ vel glandulis aut appendicibus staminum tubo plus minusve adnatis auctâ. Stamina 5 basi corollæ inserta, filamentis in tubum pistillum inclientem (gynostegium) saepissime connatis, antheris erectis bilocularibus vel septis semicompletis semi quadrilocularibus, loculis in appendices coalitas stigmati applicatas productis. Styli duo apice connati, stigma unicum pentagonum ad angulos glandulis brunneis cum antheris alternantibus instructum. Pollen coalescens in massas loenorum numero æquales stigmati glandulis affixas. Ovaria duo multiovulata, placenta ad suturam ventralem. Folliculi bini. Semina pendula ad micropylem saepissime comosa. — Herbar vel frutices, plantæ saepissime lactescentes, foliis saepissime oppositis, interdum nullis.

TR. I. PERIPLOCEÆ R. Br. loc. cit. — Filamenta omnino vel superne saltem libera, antheræ dorso saepe barbatæ. Massæ pollinis granulosæ apice dilatato singulae glandulae stigmati affixaæ.

PERIPLOCA (L. Gen. 300 ex parte).

Corolla rotata laciñiis patulis vel reflexis, squamæ quinæ aristatae fauci insertæ cum laciñiis alternantes. Staminum filaments distincta, antheræ dorso barbatæ apice cohærentes ultra loculos apiculatae. Massæ pollinis solitariae e quatuor granulis cohærentibus. Folliculi cylindracei lœves. — Frutices.

1. P. **Græca** (L. Sp. 309) fruticosa elata volubilis, foliis ovatis vel ovato-oblongis obtusis membranaceis glabris breviter petiolatis, cynis ramulos terminantibus laxifloris, corolla fuscescenti-virentis laciñiis oblongis retusis margine barbatis, squamis corollinis basi dilatata biauriculatis in aristam indivisam abeuntibus, folliculis longis acuminatis subdivergentibus 5. Ic. Fl. Græc. tab. 249. — Bot. Mag. 2289.

Hab. in sylvaticis, ad sepes, juxta rivos regionis inferioris, Græcia in Etolia (Nieder!), Macedonia prope Vodena et in peninsula Athoā (Sibth., Griseb.).

prope Byzantium (Cadet!) et in Bithynia (Sibth.). Anatoliæ boreali (Tchih., Wied!), Ciliciæ (Bal!), Syriæ prope Marasch (Haussk!) et prope Tripoli (Bl!), Ponto Lazico maritimo (Bal!), omni Transcaucasia (Ledebe., Hohen!), Persie bor. prov. Ghilan (Auch. 4923!. Buhse!). Mesopotamia prope Diarbekir (Ky!).

Ar. Geogr. Italia media, Dalmatia, Montenegro.

2. *P. gracilis*. fruticosa, ramis gracilibus flexuosis elongatis scandentibus, foliis breviter petiolatis membranaceis glabris ellipticis basi acutis apice acuminatis, cymis ex axillis superioribus terminalibusque 2-4-floris pedicellis divaricatis tenuibus, corollæ luteolæ laciniis glaberrimis linearibus retusis, squamis corollinis angustis basi non auriculatis in aristam ad medium usque bifidam abeuntibus, foliulis membranaceis attenuato-acuminatis altero abortivo ♂.

Hab. in fauclibus angustis Cilicie littoralis ad Tchupurla prope Mersina (Bal. exs. 719 sub *P. levigata*!). inter Gülek et Kasauli 2000' (Ky. exs. 345 sub *P. Gracilis*!). Fl. Mai.

Folia sub antesi 12-15 lin. longa 4-5 lata, fructifera paucis aucta, flores eis *P. Graceae* dimidiis minores, folliculus 2-2½ pollicaris. Species aristis bipartitis et corollæ glaberrimæ insignis.

3. *P. levigata* (Ait. Kew I. p. 301) fruticosa humilis, ramis brevibus rigidis superioribus interdum elongatis subscendentibus, foliis glabris subcoriaceis oppositis vel fasciculatis basi attenuata subsessilibus oblongis vel elliptico-linearibus obtusis acutisve, cymis ad ramulorum apicem paucifloris foliis brevioribus, corollæ luteolæ laciniis oblongis obtusis vel retusis intus purpurascensibus praeter maculam ovatam tomentosam versus medium sitam glabris, squamis corollinis basi dilatatis biauriculatis in aristam indivisam abeuntibus, folliculis horizontalibus basi oppositis ♂. *P. angustifolia* Labill. Dec. II. p. 13. tab. 7. — *P. punicea* Cav. Ic. III. p. 9. tab. 217. — *P. rigida* Viv. Lib. p. 14. tab. 6.

Hab. ad Latakie Syriæ maritime (Labill!).

Folia vis precedantis multo minora et angustiora, flores 2-3-plo minores.

Ar. Geogr. Insulae Canarienses, Hispania austro-orientalis, Sicilia insulae, Africa borealis.

4. *P. aphylla* (Decaisne Jacquin. Bot. p. 109, tab. 116) fruticosa tandem aphylla, ramis crassiusculis rigidis striate ramulosis, ramulis elongatis interdum pendulis apice indurato-acutatis junioribus velutino-canescensibus tandem glabris, foliis minutis raris ovatis canescensibus mucronulatis subsessilibus cito deciduis, cymis secus ramulos ad axillas foliorum abortivorum subsessilibus confertilloribus fasciculiformibus, calycis hirtuli laciniis ovatis, corollæ atropurpureæ laciniis ovatis obtusis vel retusis superne a medio sursum et margine lanatis, squamis corollinis trifido-biauriculatis lobo medio in aristam indivisam abeanti, folliculis velutinis tandem glabratissimis crassis coria-

ceis apice attenuatis horizontalibus basi oppositis s. *Gomphelis riniuea* Falconer.

Hab. in desertis Sinaiticis (Bové, Auch. 1502!), in Persie australis prov. Laristan ad Dalechi (Ky. 209!), in montanis inter Kusurum et Dalechi et inter Seytum et Bebehau 300'-3000' (Haussk!), prope Bender Buschir (Auch!), in Belutschia (Stocks!), Persice *Mada*.

Florum fasciculi secus ramos pedales et longiores remotiusculi, folliculi 3-4-pollicares inferne diametro 3 lineas lati. Planta sub eodem nomine e montibus Ssoturka Nubie littoralis a cl. Schweinfurth distributa specificè differt glabritie, cymis laxilloris, folliculis duplo tenuioribus.

Ar. Geogr. India bor. occid. in regione Scinde et Pentapotamide.

TR. II. EUASCLEPLADEÆ R. Br. — Filamenta connata. Massæ pollinis decem per paria stigmatis processibus sulco longitudinali bipartitis affixa pendulae

MASTOSTIGMA (Stocks in Hook. Icon. IX. tab. 863).

Calyx quinquepartitus. Corolla expansa quinquepartita. Corona staminea petaloidea simplex profunde quinqueloba gynostegium æquans. Antheræ membranæ rotundatæ terminatae. Massæ pollinis clavatae pendulæ. Stigma obtuse conicum subbilobum. Folliculi acuminati spinis innocuis armati. — Herba ramosa incana facie *Glossonematis*.

1. **M. varians** (Stocks loc. cit.) adpresso canescens glaucum a collo ramosum, ramis crassiuseulis dichotomis abbreviatis erectis vel decumbentibus, foliis carnosulis petiolatis integris vel margine minute undulato-crispis inferioribus rotundatis superioribus ovatis acuminatis, umbellis axillaribus sessilibus 3-8-floris, pedicellis calyci subæquilongis, calycis pubescentis laciniis linearibus, corolla glabra lobis oblongis, folliculis minute canescensibus ☽.

Hab. in Persie austro-orientalis maritimis (Auch. exs. 4927! sub *Stephanidia radiante*), in rupestribus Belutschiae (Stocks!). Vulgr *Moonya*.

Rami erecti vel procumbentes? 4-5-pollicares, folia inferiora pollicem longa et lata, flores e sicco albi vel flavidi diam. 2 lineas lati, folliculi 1¹/2-2¹/2 pollices longi edules, succus dulcis innocuus.

Ar. Geogr. Regio Scinde Indie.

VINCETOXICUM (Mænch. Méth. 317).

Corolla rotata profunde quinqueloba. Corona staminea scutelliformis carnosa simplex latere interiori appendiculis destituta 5-10-loba. Antheræ membranæ terminatae. Massæ pollinis ventricosæ pendulari. Stigma apiculatum. Folliculi ventricosi pergamentacei levés patentes. — Herbae pereunes erectae interdum subvolubiles.

* Corolla intus hirsuta

1. **V. canescens** (Willd. Nov. Act. Nat. Cur. III, p. 418 sub *Asclepiade*) breviter tomentello-canescens, caulis erectis flexuosis simplicibus vel inferne ramosis, foliis late ovatis acutis basi subsessili breviter attenuatis, umbellis subsessilibus axillaribus et terminalibus multifloris, floribus brevissime pedicellatis, corollâ flavâ laciniis oblongo-triangularibus intus hirsutis, coronæ lobis 5 obtusis concaviusculis gynostegio sublongioribus, folliculis canescensibus inferne inflatis ♂. *V. canescens* Decaisne D.C. Prodr. VIII, p. 523. — *V. luteum* Sieb. exs. — *Asclepias Cretica* Clus. App. I, tom. I, CCLV. — *A. Orientalis* Buxb. Cent. III, t. 30. — *V. Anatolicum* Boiss. MSS.

Hab. in petrosis regionis inferioris montium, insula Chios 3800' (Orph!), Creta in montibus Lassiti (Sieb!), Anatolia in montibus circa Smyrnam, Tmolos, Mesogi (Boiss!), Carià (Pinard!), Lyciâ prope Elmalu (Bourg!), Pisidiâ ad Bonldour et in monte Davrosdag (Heldr!), Armeniâ Turcicâ ad Gumusch (Bourg! Tchih!) et australi (Noël), Cappadociâ (Buxb., Auch. 1496! Bal!), Tauro Cilicio ad Gulek Boghaz (Ky. 7!), monte Kechichdag supra Tokat (Wied!), Syriâ bor. circa Antiochiam (Boiss!), Mesopotamia ad Orfa (Haussk!), prov. Khorassan Persia (Bge!), Afganiâ in collibus Cabul (Griff!).

Planta Indica (*Cynanchum glaucum* Wall. Cat.) huc citata specifice differre videtur foliis angustioribus oblongo-lanceolatis, umbellis paucifloris, corollâ duplo minore laciniis ovatis.

2. **V. Arnottianum** (Wigh. Contrib. Bot. Ind. p. 58 et Icones tab. 1614) breviter canescens pubescens vel glabrescens, caulis erectis elatis stricte ramulosis, foliis inferioribus ovato-oblongis cæteris oblongo-lanceolatis apice longe basi breviter attenuatis breviter petiolatis, umbellis axillaribus paucifloris sessilibus, floribus minutis pedicellis aequilongis suffultis, calycis puberuli laciniis triangularibus, corollæ luride purpurascens laciniis ovato-oblongis obtusis, coronæ stamineæ profunde partite laciniis oblongis obtusis inter se subdistans gynostegium aequalibus, folliculis junioribus parce pubescentibus adultis glabris basi subinflatis acuminatis ♀. *V. luridum* Stocks pl. exs.

Hab. in montibus Belutschiae 3500'-5500' (Stocks!).

Sesquipedale et bipedale, folia intermedia bipinnicaria 5-7 lin. lata, flores eis *V. canescens* duplo minores.

Ar. Geogr. India in regione Himalaicâ occidentali.

3. **V. Tmoleum** (Boiss. Diagn. Ser. I, 4, p. 38) glabriuscum, caulis erectis elatis simplicibus vel stricte ramulosis, foliis brevissime petiolatis a basi rotundata vel breviter cuneata ovatis oblongisve breviter attenuato-acuminatis, umbellis axillaribus multifloris pedunculis petiolo saepius longioribus suffultis, pedicellis flore longioribus, calycis laciniis triangulari-lanceolatis, corolle virenti-flavidæ laciniis triangulari-lanceolatis acutis superne albo-hirsutis, coronæ

stamineæ quinquepartitæ laciniis ovato-triangularibus facie externâ sulcatis gynostegio subbrevioribus, folliculis glabris inferne subinflatis longe attenuato-acuminatis ♂.

Hab. gregarie in valibus alpinis Tmoli occidentalis ad Bozlagh (Boiss., Bal. 291!), in monte Gheidagh Tauri Pamphylici (Heldr!), in parte superiori montium Ciliciae littoralis prope Kechlick (Bal. exs. sub *V. contiguo.*!), et Tauri in valle Gusguta (5000' Ky. 2!, Bal!), in collibus Cappadocie circa Cesaream 4000' (Bal!), Armeniæ australi ad Gosekhar et prope Muküs 5600'-6000' (Ky. 372!) et boreali ad Kopri Chan prope Baibut (Ky. 284!).

Sesquipedale, facies *V. officinalis* sed folia crassiora non membranacea basi non cordata; a *V. canescens* glabritie, corollæ laciniis acutis, etc., discedens.

4. V. Rehmanni, secus caules pedunculos et foliorum nervos minute pubescens, caulis elongatis superne subvolubilibus, foliis subsessilibus a basi cordatâ oblongis et oblongo-lanceolatis acuminatis, umbellis axillaribus pedunculatis laxe paucifloris, pedicellis tenuibus flore parvo duplo longioribus, calycis hirti laciniis triangularibus, corollæ flavidæ laciniis oblongis obtusis calyce triplo longioribus facie internâ hirtis, coronâ stamineâ ultra medium in lobos crassos orbiculatos patentes gynostegio æquilongos fissa, folliculis glabriusculis apice valde attenuatis ♂.

Hab. in fruticetis Caucasi circa Vladikaukas (Rehm. exs. sub *V. medio.*!).

Bipedale et elatius, flores eis *V. officinalis* et *Tmolei* triplo minores, eis *V. parviflori* duplo maiores. Facies *V. Huteri* Aschers. floribus fere duplo majoribus, coronâ stamineâ deceimlobâ distincti.

5. V. speciosum (Boiss. et Sprun. Diagn. Ser. I. 4, p. 39) totum crispule velutinum, caulis elatis simplicibus, foliis amplis membranaceis brevissime petiolatis et sessilibus a basi cordatâ late ovatis superioribus oblongis acuminatis, umbellis axillaribus subsessilibus, pedicellis flori æquilongis crispo-lanatis, calyeis lanati laciniis lanceolatis, corollâ atropurpureâ majuscule laciñis lanceolatis intus hirsutis, coronæ stamineæ ad medium quinquesfidæ lobis latis obtusis apice incrassatis, folliculis tomentellis apice longe attenuatis ♀. *V. triste* Griseb. Spicil. II, p. 68. — *V. Thirkeanum* G. Koch Linn. XXIII, p. 59.

Hab. in regione montanâ et alpinâ montis Velugo (Tymphresti) Ætolia (Sprun!), Delphi Eubeæ ad Xerovoouni (Heldr!), in sylvaticis Macedoniae in Peninsulâ Athoâ (Griseb.), Bithyniâ prope Brussam (Friv!).

Caulis bipedalis et longior, folia saepè 4 pollices longa 2¹/₂ lata, corolla cæ *V. officinalis* duplo, *V. nigri* cui colore accedit quadruplo major. Species pulchra distinctissima.

6. V. nigrum (L. Sp. 315 sub *Asclepiade*) caulis erectis superne pubescentibus et saepè volubilibus, foliis subtus secus nervum et margine puberulis breviter petiolatis vel sessilibus ovalis et ovato-lanceolatis inferioribus basi interdum cordatis superioribus

acuminatis, umbellis axillaribus pedunculatis vel sessilibus paucifloris, pedicellis tenuibus flore 2-3-plo longioribus calyceque puberulis, calycis laciniis triangularibus acutis, corollæ nigræ laciniis oblongis obtusiusculis facie interiori pubescentibus, coronâ stamineâ cupuliformi decemlobâ lobis obtusis alternatim brevioribus, folliculis glabrescentibus apice longe attenuatis ♀. *V. nigrum* Mænch. Méth. 317. — *Cynanchum medium* R. Br. non Decaisne. — *V. amplifolium* C. Koch Linn. XXIII, p. 592.

Hab. in fruticetis montanis et subalpinis, Bithynia (Thirke!), Pontus Lazicus in monte Kalonoros prope Rhizé 2000' et infra Djimil 5000' (Bal!), Tauria, Caucasus et Transcaucasia (M.B., Ledeb., Hoh., Rehm!).

$\beta.$ minus. — Semipedale, folia diminuta, umbellæ sessiles, florum pedicelli abbreviati.

Hab. in alpinis Anatoliæ borealis ad Kastamouni et in monte Alidagh (Wied!), in valle Kameklu Armeniæ (Huet!), in montibus Gumuschkhané (Bourg. exs. sub *V. pumilio*).

Forma typica 2-3-pedalis est, saepe subcandens, plantæ Hispanicæ et Gallie similis. Varietas β aspectu diversa est et eam prius pro *V. pumilio* coronæ structurâ diverso habui; notas differentiales sat graves inter eam et typum non video cuius probabilissime forma alpina est.

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, Rossia media et australis.

7. **V. pumilum** (Decaisne in p.c. Prodr. VIII, p. 525) glaucum, caulis pumilis erectis simplicibus vel parce ramosis puberulis, foliis subsessilibus a basi subcordatâ ovatis attenuato-acuminatis præter nervum in paginâ inferiore puberulum glabris, florum umbellis axillaribus subsessilibus, calycis laciniis triangularibus acutis ciliatulis, corollæ luridæ laciniis oblongis supérne pubescentibus, coronâ stamineâ fere ad basin in 5 lobos oblongos obtusos partitâ, folliculis.... ♀.

Hab. ad fluvium Rezil Ouzen Persie borealis (Michaux herb. Par!), prope Schahrud (Bunge!).

Semipedale, folia non obtusa sed apice attenuata acuta vel obtusiuscula. Facies *V. nigri* var. β a quo corollæ structurâ eximie differt.

* * Corolla glabra.

8. **V. officinale** (Mænch Méth. 317) caulis elatis bifariam pubescentibus, foliis membranaceis ad nervos et margines puberulis subsessilibus a basi rotundatâ vel cordatâ ovatis superioribus oblongo-lanceolatis, umbellis axillaribus pedunculatis, pedicellis flore sublongioribus, calycis hirti laciniis lanceolatis. corollæ ochroleuce vel albae glabrae laciniis ovato-oblongis, coronâ stamineâ gynostegio equilongâ vel sublongiore patenti plus minus profunde in 5 lobos obtusissimos transverse sublatiores fissâ, folliculis glabris lanceolato-acuminatis ♀. *Asclepias Vincetoxicum* L. Sp. 314. — *Asclepias alba* Lam. — *Cynanchum laxum* Bartl. — *C. contiguum* Koch.

Hab. in Rumeлиâ (Friv. ex Griseb.), Tauriâ, Caucaso et Transcaucasiâ (M.B., C.A. Mey., Hoh.). E ditione nondum vidi.

Ar. Geogr. Europa sept. et media a Scandinaviâ ad Rossiam medium et Hispaniâ et Italiâ centralibus ad Dalmatiam et regionem Danubialem.

9. V. nivale (Boiss. et Heldr. Diagn. Ser. I, 7, p. 20) caulinis humilibus simplicibus erectis brevissime puberulo-crispis, foliis subsessilibus sessilibusque subtus ad nervos et margines puberulij cæterum glabris inferioribus a basi truncatâ vel subcordatâ ovatis superioreibus oblongis acuminatis, umbellis in axillis superioribus paucifloris pedunculatis, calycis glabri laciniis anguste lanceolatis, corollæ luteæ tandem fuscementis laciniis oblongis glabris, coronâ stamineâ gynostegium superante breviter in lobos crassiusculos obtusos fissâ, folliculis φ . *V. undulatum* Heldr. in Boiss. Diagn. Ser. II, 6, p. 421 (forma corollæ laciniis sub lente pilis nonnullis obsoletis obstitis).

Hab. in saxosis regionis alpinæ Græciæ, Taygetus ad nives cacuminis Hagios Paraskevy (Heldr!), Chelmos (Orph!), Parnassus (Sprun!, Orph!), mons Velugo (Etoliæ 5500' (Samar. et Guic!), Olympus Thessalus (Heldr!).

Semipedale vel vix pedale, folia eis *V. officinalis* minora, ab eo praeter statu ram humilem floribus tandem lividis, calycis laciniis angustioribus, coronæ lobis minus latis et magis dissitis discedens. An tamen hujus varietas?

10. V. medium (Decaisn. n.c. Prodr. VIII, p. 523 ex parte quoad plantam Orientalem) caulinis elatis subbitariam hirtulis superne interdum subvolubilibus, foliis subsessilibus ovatis basi cordatis abrupte et breviter acuminatis subtus ad nervos et margines pubescentibus, umbellis axillaribus longiusculæ pedunculatis, calycis glabriuscui laciniis triangularibus acutis, corolla glabrae atropurpurea laciniis ovato-oblongis, coronâ stamineâ gynostegio subæquiloniâ ad medium in lobos quinque triangulari-deltoides crassiusculos fissâ, folliculis glabris apice acuminatis φ . *V. latifolium* C. Koch Linn. XXIII, p. 591. — *V. funebre* Boiss. Diagn. Ser. I, 11, p. 79.

Hab. in Tauriâ (Stev.), prov. Caucasicis (C.A. Mey.), prope Tiflis (Rehm!), Persiâ boreali ad radices montis Demavend 6200' (Ky. 359!), prope Siaret (Bge!).

Planta elata facie *V. nigrum* omnino referens ab eo corolla glaberrimâ et coronâ stamineâ quinquelobâ discedens

Ar. Geogr. Rossia austro-occidentalis ad Wolgam.

11. V. fuscatum (Rehb. fil. Ic. Germ. XVII, p. 17 nec tab. 28) breviter pubescens, caulinis humilibus simplicibus, foliis breviter petiolatis inferioribus a basi rotundatâ vel subcordatâ ovatis superioribus anguste lanceolatis, umbellis paucifloris axillaribus sessilibus, pedicellis hirtis flore longioribus, calycis puberuli laciniis breviter lanceolatis acutis, corolla fuscementis glabrae laciniis oblongis, coronâ stamineâ gynostegio subæquiloniâ cupulari pentagonâ brevissime et

sæpius obsolete lobulatâ, folliculis glabris apice valde attenuatis. ♀.
V. Pisidicum Boiss. et Heldr. Diagn. Ser. I. 11, p. 79. — *V. alpinum* Schott in Ky. Sched. — *Cynanchum minus* C. Koch Linn. XIX, p. 27.

Hab. in lapidosis præsertim subalpinis et alpinis, Parnassi regio inferior ad Leucocastron supra Rachova (Heldr!), circa Byzantium et in Bithyniâ (Noë!, Coumany!, Thirkel!), Phrygia (Bal!), Lycia prope Elmalu (Bourg!), Pisidia circa Bouldour (Heldr!), Cappadocia in collibus circa Cæsaream (Bal!), Tauri Cilicici jugum Koschan 7500' (Ky. 41!).

Semipedale, folia superiora valde angusta, corolla glaberrima fusca. Species coronâ obseure lobatâ inter affines distinguenda. Synonymon *Cynanchi fuscata* Link ad plantam Lusitanicam diversam spectat, sed nostra specimina Orientalia Dalmaticis a b. Petter in subalpinis Clissa lectis simillima sunt. Icon tab. 28 Rehb. mihi dubia ob specimen nimis procerum et præsertim analysin N° 2, coronam valde lobatam exhibentem. Synonymon *A. Iberica amplissimo folio* nerb. Vail. ob corollam superne hirtam et staturam elatam ad *V. nigrum* referendum est.

Ar. Geogr. Dalmatia.

12. *V. parviflorum* (Decaisne in D.C. Prodr. VIII, p. 525) caulinibus erectis sæpius striete ramulosis puberulis. foliis carnosulis breviter petiolatis a basi rotundatâ ovato-oblongis et oblongo-lanceolatis apice attenuatis obtusis, umbellis paucifloris axillaribus sessilibus vel breviter pedunculatis, pedicellis tenuissimis pubescentibus flore minimo longioribus, calycis hirti laciñis triangularibus, corollæ lutescenti-rubellæ vel purpureæ glabrae laciñis oblongis obtusis calyce 3-4-plo longioribus, coronâ stamineâ ad 5 glandulas purpureas gynostegio multo breviores reductâ, folliculis... ♀. *V. stelliflorum* Boiss. et Ky. Diagn. Ser. II, 3, p. 120.

Hab. in Cappadociâ ad Euphratem (Auch. 4512!), in Tauri Cilicici fauceibus Gûlek Boghaz 3000' ad ripas torrentis (Ky. 88!) et in regione alpinâ vallis Kar Boghas (Bal!). Fl. Julio.

Pedale, folia inferiora 1-1 ½ poll. longa. Species floribus minimis et coronâ stamineâ brevissimâ notabilis.

SOLENOSTOMA (Hayne Arzn. IX, t. 38). — *Argelia* Decaisn. Ann. Sc. Nat. IX, p. 331, tab. 11.

Calyx quinquepartitus. Corolla quinquepartita laciñis erectis oblongo-linearibus obtusis. Corona staminea cyathiformis crassiuscula lobis 5 rotundatis brevibus gynostegii longe stipitati basin eingens eo multo brevior. Antheræ apice membranaceæ. Pollinis massæ clavatae, compressæ pendulae. Stigma pentagonum planiusculum. Folliculi ovoidei lèves cartilaginei. — Frutex.

1. *S. Argel* (Delil. Fl. Eg. p. 216, tab. 20 sub *Cynancho*) fruticosum, ramis tereftibis erectis crebre foliosis, foliis crassiusculis glaucis minute velutinis subsessilibus ovato-oblongis et ellipticis acutis, umbellis axillaribus breviter pedunculatis bracteolatis, corollâ albida 5.

Hab. in vallis deserti Arabici Ægypti superioris ad orientem Syenes (Delile, Husson!), ad Ouadi Gurra Arabiae petreæ (Schimp. 376!). Arabice *Argel*.

Folia eis *Cassia* mixta cum eis venditantur et eisdem viribus medicis gaudere videntur. Frutex 2-3 pedalis, folia magn. *Oleastri*.

Ar. Geogr. Nubia.

CALOTROPIS (R. Br. Wern. Soc. I, p. 39).

Calyx ad basin usque partitus. Corolla hemisphaerico-campanulata tubo brevi angulato angulis introrsum saccatis, limbo profunde quinquefido. Corona staminea foliolis 5 compressis carinaeformibus gynostegio verticaliter adnatis inferne gibbosus constans. Antherae membranâ latiusculâ terminatae. Massæ pollinis compressæ cuntriformes apice attenuato astixæ. Stigma pentagonum depresso apiculatum. Folliculi ventricosi herbacei. — Frutices vel arbusculæ cortice suberoso.

I. C. procera (Willd. Sp. I, p. 1263 sub *Asclepiade*) arborescens succo lacteo acri seatens. truncо ramisque tortuosus crassis, foliis amplis subsessilibus ovatis ad insertionem glandulis setaceis obsitis glaucis crassis farinoso-velutinis, umbellis ex axillis supremis et terminalibus pedunculatis, corollæ majuscule roseæ laciniis patulis ovatis margine subrevolutis, coronâ gymnostegio brevi depresso æquilongâ 5. *C. procera* R. Br. h. Kew. II, p. 78. — *Asclepias gigantea* Forsk. Eg. p. 53. — *Apocynum Syriacum* Glus. — *Ic. Bot. Reg.* 1792.

Hab. in Ægypto superiore ad deserti limites ab urbis Keneh viciniis ad Syenem (Ehr!, Sieb!, Boiss!, etc.), Arabiâ Petreâ in valle Hebræ (Schimp. 412!). Persiâ austro-occid. prope Dalachi et Schuster (Haussk!). ad Sinum Persicum (Auch. 4932!), in Persia austro-orientali prope Chabbise (Bge!). Afghanistanâ (Griff.). — Arabice *Oscher*.

Frutex 10-15-pedalis trunci et ramorum subere denso albo, folliculi maturi 5-6 poll. longi 3-4 lati.

Ar. Geogr. Senegal, Algeria subtropica interior, Nubia, Abyssinia, India borealis.

KANAHIA (R. Br. loc. cit. p. 39).

Calyx ad basin quinquepartitus. Corolla subrotata quinquepartita laciniis introrsum lanatis. Corona staminea supra medium gynostegium inserta phyllis 5 inferne caelitis superne liberis incrassatis acuminatis. Antherae membranâ terminatae. Massæ pollinis ovoideæ pendulae. Stigma planum apiculatum. Folliculi leves pergamacei. — Herbae perennes facie *Gomphocarpi fruticosi*.

1. **K. Delilii** (Decaisne D.C. Prodr. VIII, p. 537) caulibus elatis strictis simplicibus crenatis foliosis, foliis subsessilibus strictis linear-lanceolatis acuminatis, umbellis axillaribus et terminalibus pedunculo foliis dimidio breviore suffultis, floribus roseis, corollæ laciniis ovato-oblongis, corona phyllis gibbosis in acumen bifidum supra stigma arcuatum abeuntibus, folliculis ovatis breviter acuminatis ♀. *Asclepias laniflora* Delile in Caill. Cent. tab. 54 non Forsk.

Hab. in Ægypto superiore (ex D.C. Prodr.). E ditione nondum vidi.

Ar. Geogr. Nubia, Abyssinia.

PENTATROPIS (R. Br. in Salt. Abyss.).

Calyx quinquepartitus. Corolla rotata profunde quinquepartita laciniis attenuatis patulis dein reflexis. Corona staminea duplex, exterior ad denticulos 5 foliolis interioribus oppositos interdum subnullos reducta, interior phyllis 5 gynostegio verticaliter adnatis superne liberis basi calcaratis subhorizontaliter patulis. Antheræ membranæ terminatæ. Pollinis massæ ovoideæ pendulæ. Stigma planum apiculatum. Folliculi læves ovato-oblongi. — Herbae perennies volubiles.

1. **P. spiralis** (Forsk. Eg. Arab. p. 49 sub *Asclepiade*) glabra ramosissima, ramis tenuibus volubilibus intricatis, foliis minutis breviter petiolatis a basi rotundata oblongis vel ellipticis mucronatis, floribus in axillis solitariis vel paucis fasciculatis pedicellis capillaris eis sublongioribus suffultis, corollæ virentis laciniis lanceolatis elongato-acuminatis, gynostegio breviter stipitato, corona phyllis acuminatis ♀. *P. spiralis* Decaisne Ann. Sc. Nat. 1838, p. 327, tab. 11, fig. E.

Hab. in regno Mascate Arabiæ (Auch. 49261).

Folia pollicaria 3-6 lin. lata, corollæ laciniæ 4-5 lineas longæ. Species subtropica et tropica.

Ar. Geogr. Senegal. Nubia, Abyssinia, Arabia tropica.

SARCOSTEMMA (R. Br. Wern. Soc. I, p. 50).

Calyx quinquepartitus. Corolla rotata plus minus profunde quinquefida vel urceolato-rotata sinuato-quinqueloba lobis interdum denticulis interjectis. Corona staminea duplex monadelpha, exterior cyathiformis vel annularis crenata, interior pentaphylla exteriorem superans foliolis carnosis rotundatis acuminatis. Antheræ membranæ terminatæ. Pollinis massæ clavato-cylindricæ pendulæ. Folliculi graciles læves. — Suffrutices (in nostro) volubiles aphylli.

1. **S. aphyllum** (R. Br. loc. cit. p. 50) volabile, ramis tenuibus

quadrangulo-subcompressis articulatis, umbellis lateralibus 2-4-floris, pedicellis flores parvos æquantibus receptaculo brevi glabro insidentibus, calyce glabro, corollæ lacinias lanceolatis acutiusculis, cordonà stamineâ cupuliformi plicatâ quinquentatâ gynostegium aquante ♀.

Hab. in Ægypto (loco non indicato) ex Schweinfurth Beitr.

Vidi fragmenta plantæ Abyssinicae quæ habitu cum typicâ Capensi congruere videtur sed quæ ex speciminibus melioribus examini accuratiori subjicienda erit. Flores minuti.

Ar. Geogr. Abyssinia, P. B. Spei.

OXYSTELMA (R. Br. Wern. Soc. I, p. 40).

Calyx quinquepartitus. Corolla rotata patens quinqueloba lobis triangularibus ciliatis, fauce annulo expanso papilloso instructa. Gynostegium exsertum. Corona staminea pentaphylla phyllis transversis inferne inflatis apice sensim in ligulam attenuatis gynostegium superantibus. Antheræ membranæ terminatæ. Massæ pollinis clavato-compressæ apice attenuatae pendulae. Stigma muticum. Folliculi laeves. — Herbae perennes scandentes.

1. **O. Alpini** (Decaisne D.C. Prodr. VIII, p. 543) ramis tenuibus elongatis scandentibus, foliis glaucescentibus puberulis anguste linearibus acutis inferne attenuatis, racemis 2-5-floris ex axillis ortis folium æquantibus, corollæ roseæ lobis late et breviter triangularibus ciliatis, folliculis oblongis acuminatis geminis glabris ♀. I.e. Prosp. Alp. Eg. p. 62, tab. 48 (mala). — *Periploca Secamone* Del.

Hab. in Ægypto ad Nili ripas supra Edfou (Boiss!), prope Kahiram (Husson!).

Folia 2 pollices longa 2 lineas lata, corolla diametro 9-10 lineas lata

Ar. Geogr. Nubia, Abyssinia.

DÆMIA (R. Br. in Wern. Soc. I, p. 50).

Calyx quinquepartitus. Corolla hypocrateriformis tubo brevi, limbo rotato quinquepartito. Corona staminea duplex, exterior brevis annuliformis sinuato-10-lobata, interior pentaphylla phyllis inferne liberis calcaratis superne in acumen subulatum inflexum attenuatis. Pollinis massæ compressæ attenuatae processu lateraliter adnatæ. Stigma muticum. Folliculi laeves vel spinulis innocuis obsiti. — Fruticæ superne volubiles.

1. **D. cordata** (R. Br. I. cit.) fruticosa tota adpresse tomentosæ, ramis vetustis tortuosis junioribus scandentibus, foliis petiolatis a basi valde cordatâ ovatis vel reniformibus breviter acuminatis, umbellis plurifloris pedunculo folio axillari longiore tandem incrass-

sato persistente suffultis, pedicellis flore plures longioribus, corollæ margine ciliatae tubo calyce sublongiore, folliculis canis prætereaque spinis mollibus innocuis sparsim obsitis superne in rostrum curvatum attenuatis, seminibus margine inferiore inciso-crenatis ȝ. *D. incana* Decaisne. — *Pergularia tomentosa* L. Mant. p. 51.

Hab. in rupestribus desertorum, Ægyptus inter Kahiram et Suez (Boiss!) et in regione littorali ad Kosseir (Schweinf!), Arabia petrea (Schimp. 342! Boiss!), Persia australis ad Teng Tokab prope Behehan et prope Dalechi (Haussk!), ad sinum Persicum (Auch. 4928!), in Belutschia (Stocks!).

Flores albo-flavidi calyce brunneo.

Ar. Geogr. Algeria interior, Nubia, Abyssinia.

CYNANCHUM (L. ex parte — R. Br. Wern. p. 44).

Calyx quinquepartitus. Corolla rotata profunde quinquepartita. Corona staminea monophylla tubuloso-campanulata gynostegium includens laciniis 10, 5 exterioribus longioribus, 5 internis exterioribus oppositis et brevioribus. Antheræ apice membranaceæ. Pollinis massæ ovate pendulæ divergentes. Stigma pentagonum bituberculatum. — Folliculi læves abortu sæpius solitarii. — Herba perennis volubilis.

1. **C. acutum** (L. Sp. 310) volubile herbaceum inferne glabriuscum superne puberulum, foliis a basi late vel angustius cordatâ ovatis vel oblongis acutis auriculis rotundatis, floribus in umbellas vel racemos axillares pedunculatos dispositis, calycis plus minus pubescens laciniis oblongis, corollæ albæ vel roseæ laciniis oblongis obtusis, folliculis tenuibus lanceolatis acuminatis glabris ȝ. Ic. Fl. Græc. tab. 250. — *Rchb.* t. 29. — *C. Monseliacum* L. Sp. 311.

Hab. ad sepes et in humidis totius regionis inferioris præsertim maritimæ a Græciâ et ejus insulis. Macedonia (Pichler!). Thraciâ et. Byzantio (Castl.), ad Tauriam, Caucasum et Transcaucasiam (Ledeb!), Syriam maritimam (Bl!), Gaill!, Ægyptum (Husson! Schimp!), Babyloniam ad Bagdad (Noë!). Quoque in interioribus salsuginosis, Cappadocia (Bal!).

β. longifolium. — Folia angustata subtripartita laciniâ superiore lanceolata — *C. longifolium* Mart.

Hab. in Mesopotamiâ ad Charput, Diarbekir (Haussk!), Persiâ boreali (Haussk!) et in prov. Khorassan (Bge!), Turkestaniâ ad Samarkand (Lehm!).

Ar. Geogr. Regio mediterranea Europæ a Galliâ australi, Italiâ, Dalmatiâ ad Rossiam australem, Africa borealis, Sibiria Songarica et Altaica.

GLOSSONEMA (Decaisn. Ann. Sc. Nat. IX, 335).

Calyx quinquepartitus. Corolla subcampanulata profunde quinquepartita laciniis apice et superne carnoso-incrassatis. Corona staminea phyllis 5 basi corollæ adnatis planis apice in filamentum attenuatis constans. Antheræ apice membranaceæ. Massæ pollinis ovatae ex

funiculis ascendentibus pendulae. Folliculi laeves vel spinis innocuis obsiti. — Herbae perennes.

1. G. Boveanum (Decaisn. loc. cit. tab. 42. fig. D.) totum plus minus canescens a basi ramosum ramis brevibus prostratis, foliis breviter petiolatis a basi truncata vel cuneatâ ovato-oblongis lanceolatis linearibusve margine sinuatis vel undulato-crispis, floribus paucis ad axillas fasciculatis pedicellis eis subbrevioribus suffultis, calyx laciniis lanceolatis, corollæ linear-ellipticis, coronæ phyllis oblongis apice sub cuspidate retusis aut in eam attenuatis, folliculis ovatis vel oblongis acuminatis canis spinis innocuis plus minus obsitis interdum parce tuberculatis ♀.

Hab. in littore Egyptiaco maris Rubri inter Kosseir et Ras Benass (Schweinf!).

Rami semipedales, folia $\frac{1}{2}$ -2 poll. longa formâ valde variabilia, flores 2 lineas longi, folliculi 1- $\frac{1}{2}$ poll. longi.

Ar. Geogr. Arabia tropica, Nubia, Abyssinia.

COMPHOCARPUS (R. Br. Wern. Soc. I, p. 38).

Calyx quinquepartitus. Corolla quinquepartita rotata vel reflexa. Corona staminea phyllis 5 basi gynostegio adnatis cucullatis concavis urceolatis saepe verticaliter compressis navicularibus constans. Antherræ apice membranaceæ. Pollinis massæ cultriformes apice attenuatae pendulae. Stigma depresso pentagonum. Folliculi ventricosi spinis innocuis saepius echinati. — Herbae perennes vel suffrutices.

1. G. fruticosus (L. Sp. 315 sub *Asclepiade*) fruticosus glabrescens vel minute velutinus viridis, ramis virgatis simplicibus strictis crebre et stricte foliosis, foliis lanceolato-linearibus mucronatis breviter petiolatis, umbellis ex axillis supremis, pedunculo pedicellisque elongatis canis, calycis laciniis lanceolatis, corollæ albæ laciniis ovatis margine hirtis cæterum glabris, coronæ antheras æquantis phyllis urceolatis navicularibus apice truncatis et ad utrumque angulum internum bidentatis dente interno recto externo longiore extrorsum curvato, folliculis ovato-oblongis acuminatis inter setas longas glabresentibus ♂. *G. fruticosus* R. Br. loc. cit. — Ic. Jacq. Misc. tab. 2. — Rehb. Ic. tab. 30.

Hab. ad vias, rivorum margines Zacynthi (Marg!), Peloponnesi (Guérin!), insulae Naxos (Fl. Pelop.), Rhodi (Bourg!), Macedonia in Peninsulâ Athoï (Pichler!). Sæpe cultus et ideo an ubique spontaneus!

Ar. Geogr. Insulae Camarienses, Africa borealis, Hispania, Corsica, Sardinia, Dalmatia.

2. G. Sinaicus (Boiss. Diagn. Ser. I. t1. p. 80) fruticosus totus crispule canescenti s., ramis virgatis simplicibus crebre foliosis, foliis

breviter petiolatis oblongo-vel linear-lanceolatis acutis vel mucronulatis, umbellis ex axillis superioribus paucifloris pedunculis folia brevioribus suffultis, pedicellis elongatis canescensibus, calycis laciniis triangulari-oblongis obtusis, corollæ lutescentis laciniis reflexis ovatis extus pubescenti-velutinosis, coronae antheras aequantis phyllis urceolatis navicularibus apice truncatis ad utrumque angulum internum dente unico brevi erecto terminatis, folliculis oblongis in rostrum attenuatis inter setas rubras farinoso-tomentellis ♂. *G. fruticosus* Decaisne Fl. Sinaie, non R. Br.

Hab. in deserto Egyptiaco-Arabico (Husson!), in lapidosis vallium jugi Sinaitici (Ehrenb!, Bové! Anch. 4511!, Schimp. 359 sub *G. fruticosus*!), ad Ouadi Feiran (Boiss!), Arabice *Hergjel*.

Facies *G. fruticosus* a quo egregie differt indumento, corollæ extus totâ pubescente, coronae phyllorum formâ.

CIONURA (Griseb. Spic. II, p. 69).

Calyx quinquepartitus. Corolla quinquepartita laciniis patulis. Corona staminea phyllis 5 liberis oblongo-linearibus gynostegio subaequilongis constans. Antheræ membranæ terminatæ. Stigma longe rostratum bifidum gynostegium longe superans. Pollinis massæ ovatae erectæ minimæ. Folliculi leves. — Herba erecta multicaulis caulinibus basi suffrutescentibus. — Genus a *Marsdenia* coroliâ rotatâ, stigmate longe rostrato distinctum.

t. **C. erecta** (L. Hort. Cliff. p. 79 sub *Cymacho*) glabra, caulinibus interdiam superne subvolubilibus, foliis longiusculæ petiolatis cordatis ovatis breviter acuminatis, cymis pedunculatis multifloris axillaribus supraaxillaribus et terminalibus, ealycis laciniis brevissimis ovatis, corollæ albae glabrae laciniis linearibus obtusis, folliculis magnis oblongis apice attenuatis abortu sæpius solitariis ♀. *C. erecta* Griseb. loc. cit. — Ic. Jacq. Misc. tab. 2. — *Cymachum rectum* Forsk. Eg. Arab. — *Marsdenia erecta* R. Br. Wern. J. p. 29.

Hab. in siccis regionis inferioris Graecia in Cretâ (Raul!), Laconia (Heldr!), Attica (Sprun! Orph! Fl. Gr. exs. 88!), Cycladibus (Sprun!), Eubœa (Wild!), Macedonia et Thracia usque ad 4200' (Griseb!), Byzantio (Cast!), Bithynia (Pichler!), totâ Anatoliâ occidentali et australi macritinâ (Boiss! Bourg!), Galatia ad Tokat (Wied!), Tanno Cilicio usque ad 2500' (Ky! Bal!), Syriâ bor. ad Marasch et Aintab (Haussk!), Mesopotamiâ (Noë!), Persiâ austro-occid. ad Bebehau (Haussk!), ad Isghan (Anch. 4929!), Turcice *Babryk*.

LEPTADENIA (R. Br. Wern. Soc. I, p. 34).

Calyx brevis quinquepartitus. Corolla rotata quinquepartita tubo brevissimo, laciniis tomentosis marginibus revolutis basi in 5 gibbos glandulosos cum laciniis alternantes abeuntibus. Corona staminea

nulla. Antheræ contiguae sed liberae. Pollinis massa erectæ apice pelucide stipite inflato brevi affixa. — Stigma muticum. Folliculi laeves sæpius abortu solitarii. — Frutices.

2. L. pyrotechnica (Forsk. Deser. p. 53 sub *Cynancho*) fruticosa aphylla, trunco tuberoso rami ramulisque longe virgatis strictis teretibus, florum subsessilium fasciculis pedunculis brevibus lateribus post anthesin incrassatis et recurvis suffultis, corollæ parva albido-grisea, folliculis longis fusiformibus pendulis 5. *L. pyrotechnica* Dœc. loc. cit. — Ie. Del. Eg. tab. 20. — *Sarcostemma pyrotechnica* R. Br.

Hab. in desertis Arabicis Egypti (Del., Husson!) et ad littus maris Rubri prope Kosseir (Schweinf!), in Arabiæ petreâ (Bovê! Schimp. 722!), in maritimis Persie australis (Auch. 4930!), Arabice *March*

Senegal, Nubia.

BOUCEROSIA (Wight et Arn. Contrib. p. 34).

Calyx carnosus quinquepartitus. Corolla subcampanulata quinquefida sinus acutis, laciniis triangularibus introrsum verrucosis. Corona staminea gamophylla dupliæ serie gynostegium tegens phylloïdibus interioribus antheris incumbentibus indivisis, exterioribus subulatis cornitormibus divaricatis. Pollinis massa erectæ supra basin affixa. Stigma muticum. Folliculi laeves apice attenuati. — Herbe carnosa aphylla, ramulis tetragonis angulis dentatis, facie *Stapelia*.

1. B. Sinaica (Decaisne n.c. Prodr. VIII, p. 649) ramis elongatis gracilibus obsolete angulato-sulcatis, floribus ad ramulorum apicem solitariis, corollæ parvæ quinquefidæ, laciniis subdeltaoides introrsum pilosis, corolle stamineæ phyllis interioribus ligulatis acutis externa multo superantibus ♀.

Hab. in montibus Sinai Arabiae petreæ (Auch. 285!).

Ex nuce specimine nota et viva exploranda.

2. B. Aucheriana (Decaisn. n.c. Prodr. VIII, p. 649) ramis erectis tetragonis, angulis acutis inciso-dentatis dentibus subquadritatis callosis, florum fasciculis ad verticem ramorum sessilibus ♀.

Hab. in Regno Mascate Arabiae (Auch. herb. Mus. Par.).

Planta semipedalis a parte media stricte ramosa, rami breves crassitibi digiti, flores eis *B. Monbyanae* minores flavi in capitula piso subramora congesti.

3. B. Stocksiana, ramis elongatis gracilibus elongato-sulcatis, floribus in racemum terminalem laxum dispositis pedicellatis, calycis laciniis lanceolatis acutis, corolle ad medium fissa laciniis lanceolatis acutiusculis glabris ♀.

Hab. in Belutschia (Stocks. 533!).

Planta ramosa 6-10-pollicaris facie *B. Sinaicae* ramis tenuibus diametro 4-5 lineas latis; ab ea distincta racemis terminalibus bipollicaribus 4-7-floris, corollâ (diam. 2 $\frac{1}{2}$ lineas latâ) glabrâ. Coronæ formam ex speciminibus imperfectis non describere valui.

ORD. LXXX. GENTIANÆ.

(Juss. Gen. p. 141.)

Calyx liber persistens quinquefidus vel partitus. Corolla gamopetala hypogyna regularis 4-8-fida persistens. Stamina corollæ inserta tot quot corollæ laciniae et eis alterna, filamenta libera. Ovarium unicum multiovulatum uniloculare vel valvulis introflexis semibiloculare. Styli duo ex parte vel omnino connati. Ovula anatropa. Capsula polysperma bivalvis valvularum marginibus seminiferis, vel bilocularis valvulis margine introflexis dissepimentum constituentibus et placantis centralibus, rarius bacea. Semina indefinita. embryo rectus in axe albuminis carnosus.

TR. I. MENYANTHEÆ Griseb.

Corollæ æstivatio induplicativa. Ovarium disco hypogyno impositum vel glandulis stipatum. Testa lignosa. Folia semper alterna.

MENYANTHES (L. Gen. 164 ex parte).

Calyx quinquepartitus. Corolla decidua infundibuliformis limbo quinquepartito, ovarium annulo ciliato impositum. Stylus filiformis, stigma bilobum. Capsula unilocularis juxta carpellorum suturas subirregulariter rumpens valvis medio placentiferis. — Herba perennis palustris.

1. M. trifoliata (L. Sp. 208) rhizomate obliquo vestigiis petiolorum vetustorum veluti articulato radiculifero, foliis petiolo basi in vaginam amplam amplexicaulem dilatato suffultis ternatis foliolis obsovato-oblongis obtusis, floribus in racemum longe pedunculatum ex axillâ inferiori ortum dispositis, corollæ roseæ laciñiis triangularibus ciliatis intus longe barbatis ♀. Ic. Fl. Dan. tab. 541. — Rehb. t. 2

Hab. in fossis, pratis turfosis inundatis regionis alpinæ, Pontus Lazicus in monte Tcharantach 720' (Bal!), Guria Transcaucasica (Nordin!), Caucasus centralis in monte Kaischaur 600' (C.A. Mey.). Probabiliter in Macedonia et Thraciâ.

Ar. Geogr. Europa omnis borealis et media a Scandinavia ad Rossiam mediam, Hispaniæ boreali, Italiæ mediæ, Sibiria, regione Himalaicæ, America boreali.

LIMNANTHEMUM (Gmel. Act. Petrop. 1789).

Calyx quinquepartitus. Corolla decidua rotata quinquepartita. Glandulae 5 ad basin ovarii cum staminibus alternantes. Stylus filiformis, stigma bifidum. Capsula unilocularis evalvis demum mace ratione aperta, placentis suturalibus. — Herbae perennes nutantes.

1. **L. nymphoides** (L. Sp. 207 sub *Menyanthe*) foliis suboppositis suborbiculatis profunde cordatis supra nitidis subtus glanduliferis, florum umbellis axillaribus, calycis segmentis lanceolatis corollâ duplo brevioribus, corollæ membranaceæ luteæ laciniis obovatis obtusis molliter ciliatis fauce longe barbatâ, seminibus ovatis complanatis margine alato ciliolatis ♀. *L. nymphoides* Link Port. I, 344. — Ic. Rehb. tab. 1.

Hab. in aquis Græciae et Byzantii (Sibth.), Caucasi et Transcaucasiae (Led.), Persie bor. in lacubus Ghilani (Auch. 4963!) et prob. alibi.

Ar. Geogr. Europa borealis et media ab Angliâ et Belgio ad Rossiam mediam, Sibiria, China, regio Himalaica.

2. **L. Indicum** (L. Sp. I, p. 207 ex parte quoad Syn. Rumph. sub *Villarsia*) foliis solitariis orbiculatis profunde cordatis siccis membranaceis subasperis, umbellis petiolis insertis, calycis segmentis oblongis acutis corollâ duplo brevioribus, corollæ albæ segmentis margine fimbriatis intus eristatis nudis versus basin glandulâ solitaria pedicellatâ auctis, seminibus ovatis vix compressis tenuissime tuberculato-muriculatis exalatis ♀. *Villarsia Indica* Vent. Choix. p. 9. — Rumpl. Amb. tab. 72, fig. 3.

Hab. in lacubus prov. Ghilan Persiae (Auch. 4962!).

Specimen Persicum Zeylanicus habitu et seminibus simillimum; ea illis *L. nymphoides* multo minora oculo armato tuberculata.

Ar. Geogr. India orientalis.

TR. H. EUGENTIANEÆ Endl.

Corollæ aestivatio contorta. Discus hypogynus nullus. Testa membranacea. Folia sœpius opposita.

CICENDIA (Adans. Fam.).

Calyx tubulosus quadridentatus vel quadripartitus. Corolla infundibuliformis limbo quadripartito supra capsulam post anthesin con-

torto. Stylus distinctus deciduus, stigma capitulatum, capsula unilocularis vel valvulis introflexis semibilocularis. — Herbæ annuae pusillæ.

1. C. filiformis (L. Sp. 335 sub *Gentianæ*) pusilla erecta a basi stricte et dichotome ramosa, foliis parvis imis linearibus caulinis breviter subulatis, pedunculis elongatis nudis unifloris. calycis campanulati dentibus triangulari-lanceolatis, corollæ latae tubo brevi ventricoso lobis oblongis tubo æquilongis stipeius elausis, capsulâ uniloculari ♂. *C. filiformis* Delarb. Fl. Auv. I, p. 20. — Fl. Dan. tab. 324. — Rehb. tab. 4. — *Microcalyx filiformis* Link. — *Exacum filiforme* W.

Hab. in uliginosis aestate exsiccatis, Græcia in Elide (Sibth.), Bithynia prope Byzantium ad radices collis Bougourlou (Duperquet!).

Ar. Geogr. Lapponia, Dania, Anglia, Lusitania, Hispania, Gallia, Germania Italia et ejus insulæ, Istria, Transylvania.

CHLORA (L. Gen. 4258).

Calyx 8-6-partitus. Corolla hypocrateriformis marcescens tubo brevissimo inflato, limbo 8-6-partito. Stamina 6-8 fauci inserta. Stylus filiformis caducus, stigma bipartitum. Capsula unilocularis bivalvis septicida, placenta marginibus valvularum introflexis adnatæ. — Herbæ annuae flavifloræ.

1. C. perfoliata (W. Sp. II, p. 340) caule erecto folioso in cymam corymbosam plus minus multifloram abeunti rarius unifloro, foliis caulinis triangulari-ovatis totâ suâ latitudine connatis, calycis ad basin usque partiti laciniis 8 subulatis uninerviis corollâ brevioribus, corollæ laciniis oblongis obtusis ♂. Ic. Rehb. tab. 19. — *Gentiana perfoliata* L. Sp. 335.

Hab. in humidis totius ditionis a Græciâ et Macedonia ad Pontum (Tchih!), Tauriam, Abchasiam Caucasicam (Nordm.), Syriam (Ky! Haussk!), Kurdistâniam Persicam australem in monte Schahu (Haussk!).

Ar. Geogr. Europa media et australis ab Angliâ et Belgio per Germaniam australem ad regionem Danubialem et Dalmatiam.

2. C. serotina (Koch ap. Rehb. Ic. III, p. 6, tab. 351) caule erecto folioso in cymam corymbosam plus minus multifloram abeunti rarius unifloro, foliis caulinis ovatis vel ovato-lanceolatis basi subangustiore ovatâ connatis, calycis profunde partiti laciniis 8 lanceolato-subulatis sicitate subtrinerviis corollam subæquantibus, laciniis corollæ acutiusculis vel acuminatis ♂. Ic. Rehb. tab. 19. — *C. acuminata* Koch et Ziz.

Hab. in humidis, Zacynthus (Marg!), Cilicia littoralis ad Bouloukli prope Mersina (Bal!) et prob. alibi.

Ar. Geogr. Gallia, Hispania, Germania occidentalis et australis, Hispania, Italia, Dalmatia.

ERYTHR EA (Ren. Sp. 77).

Calyx tubulosus quinquefidus. Corolla infundibuliformis marcescens tubo cylindrico, limbo quinquetido. Stamina quinque corollæ tubo superne inserta, antheræ exsertæ demum spiraliter tortæ. Stylus filiformis indivisus vel bicurvis deciduus, stigma bilamellatum vel indivisum. Capsula bivalvis septicida marginibus valvarum plus minus inflexis semibilocularis, vel subunilocularis, placentis suturalibus. — Herbae monocarpicæ.

SECT. I. EUERYTHREA (Griseb. Gent. p. 136). — Corolla rosea. Stylus indivisus. Stigma bilobum vel bipartitum. Panícula fastigiata vel corymbus.

1. **E. ramosissima** (Pers. Syn. I. p. 283) caule erecto quadrangulari a basi vel superne ramosissimo laxe cymoso-fastigiato, foliis oblongis et oblongo-linearibus infimis non rosulatis superioribus acutis, floribus solitariis longiusculæ pedicellatis lateralibus a bracteis remotis, corollæ roseæ tubo sub anthesi calyce sublongiore lobis elliptico-oblongis obtusiusculis, stigmatis lobis subrotundis, capsula subuniloculari ♂. *E. pulchella* Fries Nov. p. 74. — Ic. Fl. Dan. 1637. — *Gentiana Centaurium* var. β L. Sp. I. p. 333. — *E. Caspica* Fisch. in Gris. Gent. p. 145!

Hab. in pratis sterilibus hyeme humidis totius ditionis a Graeciâ! et Macedonia! ad Arabium petream (Schimp!), Egyptum infer!, Caucasm! Persiam bor. et australem, Affghaniam (Griff!).

β. *albiflora*. — Corolla alba. — *E. Meyeri* Bge in Ledeb. Alt. I. p. 220. — Led. Ic. tab. 159.

Hab. in prov. Caucasicis (Ledeb!), Ponto Lazico ad Rhizé (Bal!), Anatoliâ boreali (Wied!), Cataonia (Haussk!), Egypto ad Kahiram (Ehr!), Arabiâ petræ (Bové!), prov. Khorassan Persiæ (Bge!).

Non minimum quidem in stigmatis structurâ inter *E. Caspicam* et *ramosissimam* discriben adest.

Ar. Geogr. Europa omnis a Scandinaviâ ad Rossiam medium, Sibiria Uralensis, Africa borealis.

2. **E. latifolia** (Sm. Angl. I. p. 321) caule erecto superne stricte ramoso cymis densis, foliis ovatis vel oblongis obtusis approximatis, floribus glomeratis basi bracteatis, corollæ tubo tenui calycem parum superante lobis lanceolatis acutiusculis, capsula subuniloculari ♂. *E. tenuiflora* Link Fl. Port. I. p. 354, tab. 67. — Engl. bot. Suppl. tab. 2719. — *E. Anatolica* C. Koch Linn. XIX. p. 27!

Hab. præsertim in maritimis, Graecia in Argolide (Orph!), Eubœa et Cephallenia (Heldr!), Attica ad Phalerum (Spr! Heldr! Orph. Fl. exs. 198!), Corinthi (Heldr!), Macedonia in Chalcide littorali (Gris!), insula Poros (Heldr!), Creta

ad Caneam (Weiss!), Lydia ad Smyrnam (Fleisch! Bal!), Syria littoralis (Bl! Gaill!), Ægyptus inferior (Husson!), Pontus Lazicus ad Batum (Calv!), Transcaucasia (Rupr! Szov!), Mesopotamia inter Bagdad et Bassora et Kurdistania Persica ad Nansud 4000' (Hausskl!).

An a *C. ramosissimâ* sat et semper distineta? Specimina quædam, saltem in herbariis, inter eas dubia.

Ar. Geogr. Auglia, Gallia occidentalis et australis, Lusitania, Hispania, Sardinia, Sicilia, Dalmatia.

3. **E. Centaurium** (Pers. Syn. I, p. 283) caule quadrangulari superne ramoso, foliis ovati-oblongis, floribus sessilibus ad apicem ramorum fasciculatis corymbum sèpius compactum formantibus omnibus basi bracteatis, corollæ roseæ tubo sub anthesi calyce sub-duplo longiore lobis ovatis obtusis, stigmatis lobis subrotundis, capsula semibiloculari ②. Ic. Fl. Dan. 617. — Engl. Bot. 417. — *Gentiana Centaurium* var. α L. Sp. 332. — *E. linearifolia* Griseb. Spicil. non Pers.

Hab. in pascuis præsertim regionis montanæ, Zacynthus (Marg!), Graecia in montibus Malevo Laconia et Parnasso (Orph!), Macedonia in Scardo et monte Athos (Griseb!), Creta (Heldr!), Bithynia (Thirke!), Caria (Pinard!), Libani regio inferior in monte Torbol et ad Eden (Boiss!), Pontus Lazicus ad Rhizé (Bal!), Tauria, Caucasus et Transcaucasia (M.B., Led., Rupr!), Persia borealis (Ky!), prov. Khorassan (Bgef!).

β *lara*. — Corymbus laxus, corollæ limbus submajus.

Hab. in insulâ Rhodo prope Salakos (Bourg!), in Libani regione inferiori supra Sidonem (Gaill!), Imeretiâ (Rupr!).

γ *acutiflora*. — Corymbus densus, corollæ laciniæ elongatæ acutæ.

Hab. in insulâ Rhodo (Cadet de Fontenay!).

Ar. Geogr. Europa omnis a Britanniâ et Sueciâ meridionali ad Rossiam medianam, Africa borealis.

SECT. II. XANTHEA (Rehb. Fl. Gerin. exs. 422). — Corolla flava.
Stylus longe bieruris. Gyna dichotoma.

4. **E. maritima** (W. Sp. I, p. 1069 sub *Chironia*) caule quadrangulari superne ramoso vel simplici unifloro, foliis elliptico-oblongis obtusis superioribus acutiusculis, cymâ pauciflorâ laxâ, floribus alaribus pedicellatis, corollæ longe tubo calycem elongatum subexcedente lobis ovatis acutiusculis, stigmatibus oblongis, capsulae valvis parum intreflexis ③. *E. maritima* Pers. Syn. I, p. 283. — Ic. Fl. Græc. tab. 237.

Hab. in pratis maritimis, Zacynthus et Peloponnesus (Sibth.), OEtolia (Nieder!), Macedonia (Pest!), Byzantium (Noë! Coumpany!), Creta (Sieb!), Rhodus (Auct!), Cilicia (Péronni!), Poros (Wied!), Syria littoralis ad Tripoli (Bl!), Pontus prope Trapezuntum (Donietti!).

Ar. Geogr. Lusitania, Hispania, Gallia occidentalis et australis, Italia, Dalmatia, Africæ borealis, Mædæra.

SECT. III. SPICARIA (Griseb. l. cit. p. 147). — Corolla rosea. Stylus indivisus stigmate infundibuliformi obsolete bilobo. Cyma spiciformes.

5. **E. spicata** (L. Sp. p. 333 sub *Gentianâ*) caule quadrangulo crebre folioso superne stricte ramoso, foliis oblongis subquinquenerviis, floribus subsessilibus strictis bracteatis in spicas longas laxiusculas simplices vel bifidas dispositis, corollæ tubo calyci æquilongo lobis lanceolatis obtusiusculis ♂. *E. spicata* Pers. Syn. I, p. 283. — Ic. Fl. Græc. tab. 238. — *D. Babylonica* Griseb. in n.c. Prodr. IX, p. 60!

Hab. in uliginosis et paludosis maritimis vel interioribus salsuginosis, Græcia in Zacyntho et Atticâ (Sibth.), Ætolia (Nied!), Cephaloniâ (Heldr!), Smyrnæ (Fleisch! Balt.), Egypto inferiore et media in prov. Fayoum (Ehrenb!), Cataoniâ ad Albistan (Haussk!), Mesopotamia (Kyt), Babyloniâ ad Bagdad (Auch!), Persiâ australi (Auch! Haussk!), Tauriâ (Lebed.), Transcaucasia ad Caspium (Hoh!), Affghanîa prope Herat (Bge!).

Ar. Geogr. Lusitania, Hispania, Gallia occidentalis et australis, Italia, Dalmatia, Rossia australis, Songaria, Africa borealis.

GENTIANA (L. Gen. 322).

Calyx 4-9-fidus rarius dimidiato-spathaceus. Corolla formâ varia limbo 4-9-fido. Stamina 5-9 corollæ tubo inserta. Stylus 1 vel subnullus persistens, stigmata duo. Capsula unilocularis bivalvis septicina, placentæ marginibus valvarum introflexis adnatae. — Herbæ.

SECT. I. ASTERIAS Ren. — Corolla rotata fauce nuda plicis destituta. Antheræ liberae. Calyx spathaceo-dimidiatus.

1. **G. lutea** (L. Sp. 329). caule crasso elato stricto simplicissimo, foliis magnis ovatis nervis 5-7 apice convergentibus radicalibus petiolatis intermediis amplexicanibus, floribus pedicellatis fasciculatis in pseudoverticillös bracteis cincinnatis involucratos dispositis, corollæ luteæ laciniis oblongo-linearibus acuminatis stellatim patentibus tubo brevissimo ♀. Rchb. Ic. tab. 18.

Hab. in pratis alpinis Macedonie in monte Peristeri supra Bitolia copiose (Orph!), Olympo Bithyno (Clem.), Lydiâ in monte Timolo supra Bozdagh (Boiss!).

Ar. Geogr. Lusitania, Hispania sept., Gallia. Italia bor. et media. Germania, Dalmatia, regio Danubialis, Bosnia.

SECT. II. CÆLANTHE Ren. — Corolla campanulata fauce nuda plicis aucta. Antheræ coalitæ. Calyx spathaceo-dimidiatus vel campanulatus.

2. **G. punctata** (L. Sp. 329) caule erecto crassiusculo simpli

eiissimo, foliis ovatis nervis 3-7 apice convergentibus radicalibus subpetiolatis superioribus amplexicaulibus, floribus sessilibus fasciculatis pseudoverticillatis bracteis involueratis, corollæ pallide flavæ purpureo-punctatae lobis ovatis obtusis tubo triplo brevioribus ♀. Ic. Jacq. Austr. App. tab. 28. — Rehb. tab. 15.

Hab. in pratis alpinis montis Peristeri Macedoniae 5000'-6500' (Griseb!).

Ar. Geogr. Gallia in Delphinatu, Helvetia, Germania in Alpibus et Sudetis, Italia bor. in Alpibus, Hungariae, Transylvaniae, Croatiae montes.

Obs. Cl. Orphanides legit in monte Pelister formam inter hanc et *G. luteam* hybridam, facie huic similem, sed corollæ plicis destituta laciniis brevioribus erecto-patulis distinctam, a *G. punctata* calyce spathaceo-dimidiato corollæque laciniis angustis elongatis discedente, *G. Charpentieri* Helveticae earumdem specierum proli hybridae fere similem.

SECT. III. AMARELLA Gaud. — Corolla hypocraterimorpha plicis destituta ad faucom barbata. Stigmata distincta ovato-oblonga. — Plantæ annue saepius ramosæ.

3. *G. obtusifolia* (Willd. Sp. I. p. 1347) humilis a basi vel superius striete et saepè pyramidatim ramosa rarius uniflora, foliis sessilibus oblongis obtusis radicalibus obovatis supremis ovato-lanceolatis acutiusculis, calycis laciniis linear-lanceolatis acutis tubo subæquilonigis sinibusque angustis, corollæ violacea quinquefidæ fauce barbatæ laciniis elliptico-lanceolatis acutis ♂. Ic. Rehb. tab. 6. — *G. Germanica* var. *minor* Griseb. Spic. II. p. 63. — *G. Germanica* var. *praecox* et var. *Persica* Griseb. in D.C. Prodr. — *G. Amarella* Sm. Prodr. Fl. Gr. et M.B. non L. — *G. Biebersteinii* Bge Gent. p. 34 cum tab.

Hab. in pratis alpinis, Olympus Bithynus (Sibth., Anch. 2423! Rayev!), Pontus Lazicus (Anch!) et in valle Djimil (Bal. exs. sub *G. Germanica*!), Caucasus occidentalis in monte Nachar (Radde!) et circa Nartzana (Bge!). Fl. Jun. Aug.

Afinis *G. Germanica* cuius ex nonnullis botanicis varietas, differt tamen foliis obtusis oblongis nec a basi latiore attenuatis, floredi tempore præcoxiore. *G. crispata* Visiani huic affinis sed laciniis calycinis eximie undulato-crispatis specificè forsitan diversa probabiliter in Macedonia alpibus observabitur.

Ar. Geogr. Germania et Helvetia in Alpibus, Sudeti, montes Hungariae, Transylvanie, Serbie.

4. *G. Caucasica* (M.B. Taur. Cane. I, p. 198) humilis a basi vel superius striete et saepè pyramidatim ramosa rarius uniflora, foliis sessilibus oblongis obtusis radicalibus obovatis superioribus cordato-amplexicaulibus, calycis ad tubi apicem subtruncati laciniis linearisubulatis subæqualibus tubo subæquilonigis sinibus latis sejunctis, corollæ violacea quinquefidæ fauce barbatæ lobis ovatis acutis ♂. Ic. Sims bot. mag. tab. 1038. — *G. Germanica* var. *Caucasia* Griseb.

Hab. in pasenis totius Caucasi 5500'-9000' (M.B., Radde!, Rupr!, Chodsko!, etc.), Georgiae Caucasicæ (Hoh!), prov. Karabagh (Szov!). Fl. Jul. Aug.

Quoad folia, habitum, affinis *G. obtusifolia*, ab eâ differt calycis laciniis

angustissimis sinibus tatis truncatis nec angustis sejunctis, corolle lobis subbrevioribus.

5. **G. minutissima**, pusilla nana ex collo ramosa ramis unifloris, foliis minimis radicalibus obovatis subpetiolatis caulinis 2-4 linearibus, pedicellis elongatis, calycis dentibus triangulari-oblongis obtusis tubo oboconico triplo brevioribus, corolla purpurea tubo calyce vix longiore lacinis lanceolatis tubo subbrevioribus, fauce barbatâ ⊖. *G. tenella* Clarke in Journ. Linn. Soc. XIV non Rottb.

Hab. in alpibus Atighanie prope Yomtt (Griff. Journ. 1024).

Mihi ex unico specimine nota vix pollicari quod primâ facie pro forma nana *G. tenella* Rottb. habui, sed præter flores triplo minores, corollamque a cl. Griffith purpuream dietam ab hac specie calyce ad tertiam partem tantum nec ad basin partito distincta.

SECT. IV. ARCTOPHILA (Griseb. Gent. p. 250). — Corolla campanulato-infundibuliformis plieis destituta non coronata sed inter basin filamentorum glandulifera. Stigmata distincta.
— Species annuae saepius ramosae

6. **G. aurea** (L. Sp. I, p. 331) a collo saepius ramosa, caule centrali argute angulato longiori stricte ramuloso internodiis distantiibus, foliis imis parvis spathulatis attenuato-subpetiolatis, caulinis sessilibus ovato-oblongis ovatis vel acutiusculis, floribus parvis breviter pedicellatis ad ramulos axillares strictos paucis fasciulis vel solitariis terminalibus interdum magis confertis, calycis corollæ tertia parte brevioris lacinis lanceolato-linearibus acutis basi attenuatis tubo suo triplo longioribus, corollæ flavæ vel pallide cærulescentis lobis ovatis acuminatis tubo subduplo brevioribus ⊖. *G. iuroluerata* Fries Act. Hafn. X, p. 434, tab. 1. — *G. umbellata* M.B. Taur. Cauc. III, p. 188. — *G. holosteoides* et *G. consobrina* Schott et Ky. in Ky. pl. Taur. 1833 (forma saepius pusilla pauciflora).

Hab. in Caucasi pratis alpinis 6000-7500', centralis (Stev., Rupr! Rehm!) et orientalis in Daghestaniâ (Rupr!), ditionis Talysh (C.A. Mey!), Tauri Cilicie 8000' (Ky. 285 et 208! Bal!).

Specimina Caucasia ne minimum quidem ab arctis et Sibiricis differunt. Cilicie interdum nana et pauciflora, interdum autem Caucasicis simillima. *G. consobrina* est forma calycinis lacinis basi non attenuatis.

Ar. Geogr. America borealis, Groenlandia, Islandia, Norvegia borealis, Lapponia, Songaria, Sibiria Altaica.

SECT. V. CHONDROPHYLLO Bge. — Corolla infundibuliformis vel hypocrateriformis plieis aueta glandulis et corona destituta. Calyx ope epidermidis in tubum connatus. Anthrae liberae. Stigmata distincta oblonga demum revoluta, capsula deumum longissime stipitata.

7. **G. Pyrenaica** (L. Mant. p. 55) cespitosa, caulis aliis flori-

feris ascendentibus unifloris aliis abbreviatis dense foliosis, folijs breviter et anguste lanceolato-linearibus mucronatis margine scabris, caulinis paucis abbreviatis distantibus, calycis dentibus lanceolatis acuminatis tubo duplo brevioribus, corollæ violaceæ lobis ovatis obtusis tubo subdupo brevioribus, plicarum appendicibus ovato-quadratis dentatis lobis subæquilongis, capsulæ stipite calycem æquante vel longiore ♀. Rchb. Ic. t. 9 — *G. Djimilensis* C. Koch Linn. XXIII, p. 583!

Hab. in pratis humidis alpinis Ponti Lazici supra Djimil 7800' (C. Koch, Bal!), ad Alischeri Khan (Ky. 283!), in Armeniæ prope Erzerum (Huet!) et in Armeniæ Rossicæ Alpe Alagös (Seidl!), Caucaso centrali 6000'-7500' (M.B., Rupr!) et orientali 8000'-9500' (Rupr!).

Ar. Geogr. Pyrenei orientales, Alpes Hungariæ.

8. ***G. prostrata*** (Hænke in Jacq. Coll. II, p. 66, tab. 17) annua nana tenuis simplex vel a collo pluricaulis caulis foliatis unifloris, folijs parvis carnosulis margine cartilagineis ovato-oblongis mucronulatis, calycis cylindrici basi attenuati dentibus ovatis vel breviter lanceolatis margine membranaceis tubo triplo brevioribus, corollæ cœruleæ tubo calyce longiore limbi laciniis ovatis acutis, plicis lobis subdupo brevioribus integris vel lobatis, capsulâ oblongo-cylindricâ basi rotundatâ stipite corollâ marcescenti demum longiore suffultâ ⊖. Ic. Rehb. tab. 8. — *G. nutans* Bge Gent. p. 232, tab. 11.

Hab. in Affghaniae alpibus circa Yonutt (Griff. Journ. 1049!), in Somchetia Caucasicâ quoque indicata sed hic locus probabiliter ad speciem sequentem referendus.

Ar. Geogr. Alpes Carinthiæ, Tyrolis australis, Sibirie Altaicæ et Baicalensis, Tibetiæ, Americae arctice, in montibus scopolosis Am. Sept.

9. ***G. humilis*** (Stev. Mém. Soc. Nat. Mosq. III, p. 258) annua pumila tenuis simplex vel a collo pluricaulis caulis laxe foliosis unifloris, folijs parvis carnosulis margine cartilagineis oblongis inferioribus praesertim apiculatis superioribus basi connatis, calycis tenuiter cylindrici basi attenuati dentibus breviter lanceolatis acutis tubo 3-4-plo brevioribus, corollæ cœruleæ tubo subexerto limbi laciniis ovato-triangularibus acutis, plicis laciniis subbrevioribus integris vel lobatis, capsulâ obovato-globosâ basi attenuatâ stipite crassiusculo corollam marcescentem demum superante suffultâ apice valvis divaricatis dehiscente ⊖. *G. aquatica* Pall. Ross. II, p. 109, tab. 97 non L.

Hab. in pratis alpinis Ponti Lazici supra Djimil 7800' (Bal. exs. sub *G. prostrata*!), Caucasi occid. et centralis 3500'-6000' (Stev! Rupr! Rehm!), orientalis in Daghestaniæ 6000'-8000' (Rupr!).

Præcedenti simillima et ab ea florifera vix distinguenda, capsulæ characteribus tamen egregie distincta.

Ar. Geogr. Sibiria Uralensis et Altaica, Dahuria, America borealis in montibus Scopolosis.

SECT. VI. *CROSSOPETALUM* Fröl. — Corolla infundibuliformis plicis et coronâ destituta inter basin filamentorum glandulifera secus margines loborum fimbriata, stigmata ovata vel orbiculata.

10. **G. ciliata** (L. Sp. 334) caule humili flexuoso unifloro vel parce et corymbose ramoso ramis foliosis unifloris, foliis inferioribus squamiformibus cæteris linear-lanceolatis acutis uninerviis, calycis campanulati laciniis lanceolatis acuminatis inferne membranaceis tubo brevioribus, corollâ cœruleâ calyce duplo longiore ad medium in lobos oblongos parte superiori denticulatos inferiori fimbriatos quadripartitâ ♀. Ic. Rehb. t. 10. — *G. fimbriata* Willd. herb.

Hab. in regione subalpinâ et alpinâ, in pratis humidis Tauri Cilicici 7000' (Ky! Bal!), Armeniâ Turcicâ prope Erzerum (Calv!), Ponto Lazico prope Khabackar 6000' (Bal!), Caucaso centrali 2500'-4000' (C.A.M., Rupr!), Daghestaniâ 6000' (Rupr!), Georgiâ Caucasicâ in monte Sarial (Hoh!), ditione Talyesch (Hoh.).

Ar. Geogr. Belgium, Gallia, Germania australis et media, Italia, Dalmatia, regio Danubialis, Rossia australis.

SECT. VII. *CYCLOSTIGMA* Griseb. — Corolla hypocaterimorpha plies aucta glandulis et coronâ destituta. Antheræ liberae. Stylus distinctus. Stigmata contigua orbiculari-infundibuliformia.

11. **G. verna** (L. Sp. 331) cespitosa, caulis pumilis unifloris, foliis sessilibus radicalibus et rosularum sterilium rosulatis breviter oblongo-lanceolatis acutis, caulinis 2-4 multo minoribus, calycis angulati lobis lanceolatis acutis tubo brevioribus, corollæ intense cœrulea lobis ovatis obtusis tubo cylindrico duplo brevioribus, plicis bitidis lobis sextuplo brevioribus. stylo indiviso ♀. Ic. Engl. bot. tab. 493. — Rehb. tab. 7.

Hab. in pascuis regionis alpinæ, Olympus Thessalus (Heldr!), Bithynus (Sibth., Boiss!), Armenia Turcica (Calv!), Caucasus mons Beschtaw (Wilh.), et Daghestania 9000'-40000' (Rupr!).

β. alata Griseb. — Major, saepe elatior, flores majores, calycis anguli in alas producti. Potius forma intermediis in typum sensim transiens quam varietas. — *G. angulosa* M.B. Taur. Cauc. I. p. 197. — Cent. tab. 47. — *G. urstiva* R. et Sch. VI, p. 156.

Hab. in pascuis siccis regionis alpine, Macedonia in Scardo (Griseb!), mons Ilgasdag Anatolie bor. (Wied!), Pontus Lazicus supra Djimil 8000' (Bal!), Caucasus omnis 5000'-8000' (M.B., Rupr! etc.).

γ. obtusifolia. — Folia obtusa, surculorum sterilium laxa, calyx parum angulatus; forma a *G. brachiphylla* Vill. foliis laxioribus obtusis diversa.

Hab. in uliginosis et spongiosis regionis alpinæ. Taurus Cilicus supra Bulghar Maaden (Bal!). Caucasus occidentalis (C.A. Mey!), mons Elbrus Persiae borealis (Ky. 696!).

Ar. Geogr. Europa media et australis, Anglia, Germania, Helvetia, Hispania, Italia, Dalmatia, regio Danubialis, Sibiria.

12. *G. nivalis* (L. Sp. 332) annua, pumila a basi stricte vel corymbose ramosa, foliis parvis sessilibus radicalibus subrosulatis obovatis obtusis, caulinis minoribus oblongis acutis, floribus ad ramorum apicem solitariis, calycis lacinias lanceolatis acutis tubo tenuiter et longe cylindrico angulato subbrevioribus, corollæ intense cœruleæ tubo e calyce vix exerto laciñis ovatis integris tubo multoties brevioribus, stigmatibus discoideo-infundibuliformibus ☺. Ic. Rehb. tab. 8.

Hab. in regione alpinâ Ponti Lazici supra Djimil 7800' (Bal!).

Specimina elongata sed ceterum plantæ Helvetice et arctice similia.

Ar. Geogr. Europæ borealis montes a Lapponiâ ad Rossiam arcticam, Scotia, Pyrenei, Jurassus, Alpes, Apennini, Carpathi et Transylvaniæ alpes, Islandia, America borealis, Groenlandia.

Obs. *G. utriculosa* L. in montibus Bosniæ et Herzegovinae crescens probabiliter quoque in alpibus Macedonie et Thracie legetur.

SECT. VIII. PNEUMONANTHE Neck. — Corolla campanulata plicis aneta coronâ atque glandulis destituta. Calyx ope epidermidis in tubum connexus. Antheræ sèpius coalitæ. Stylus nullus vel brevissimus. Stigmata distincta oblonga vel linearia demum revoluta. Capsula breviter stipitata.

13. *G. Pneumonanthe* (L. Sp. 330) caulis a collo nudo elongatis erectis strictis foliosis simplicibus, floribus axillaribus alternis et oppositis inferioribus breviter pedicellatis ceteris simplicibus rarius unico terminali, foliis oblongo-lanceolatis vel lanceolato-linearibus angusteque linearibus obtusis margine subrevolutis uninerviis infimis squamæformibus, calycis tubo obconico-tubuloso apice subtruncato laciñis linearibus aentis tubo subæquilongis, corollæ intense azureæ extus viridi-vittatæ lobis ovatis acutis plicis brevibus dentiformibus acuminatis ♀. Ic. Rehb. tab. 10.

Hab. in pratis humidis turfosis Caucasi (M.B.), in Imeretiâ et Mingrelia (Nordm.), monte Nakkerale districtu Radtscha (Rupr!).

Ar. Geogr. Europa omnis Dalmatiâ et Græciâ exceptâ, Songaria, Sibiria.

14. *G. septempida* (Pall. Ross. II, p. 101, tab. 92) caulis a collo nudo erectis vel ascendentibus simplicibus foliosis, foliis infimis squamæformibus ceteris quinquenerviis a basi ovato-subcordatâ ovato-vel oblongo-lanceolatis acutis margine sebris supremis flores subsessiles capitatos involuerantibus, calycis laciñis anguste linearis-lanceolatis

tubo æquilongis vel brevioribus sinubus truncatis sejunctis, corolla cyanæ clavatae lobis ovatis acutis plicis fimbriato-multitidis sejunctis ♀. Ic. Bot. Mag. tab. 1229 et 1410. — *G. fimbrioptera* C. Koch Linn. XXIII, p. 584!

Hab. in pratis regionis subalpinæ et alpinæ totius Caucaſi 4800'-9000' (M.B! Radde! Rupr! Rehm! etc.), Transcaucasiæ (Szov!), monte Djulfek Kuh prov. Ghilan Persiae (Auch. 4959!).

g. cordifolia. — Folia latiora et breviora obtusiora basi evidenter cordata. — *G. cordifolia* C. Koch Linn. XXIII, p. 585!

Hab. in montibus Palanteaken et Bingöldagh Armenie Turcice (Ky. 504!), in Ponto Lazico supra Djimil (7800' C. Koch, Ball!)

g. procumbens. — Caules abbreviati decumbentes, folia diminuta triangulari-ovata, flores pauci vel solitarii.

Hab. in regione alpinâ Caucaſi orientalis in Daghestaniâ et ditione Gumbet (Rupr! Chodsko!), Georgiae Caucasicæ (Hoh. sub *G. gelidâ*?), Persie borealis in monte Totschal prope Teheran (Ky. 576 et 830 sub *G. gelidâ*?).

Ar Geogr. Sibiria Altaica.

15. ***G. gelida*** (M.B. Taur. Cauc. I, p. 196) caulis a collo nudo ascendentibus simplicibus foliosis, foliis infimis squamæformibus ceteris subtrinerviis oblongis et linear-lanceolatis margine sebris, floribus subsessilibus paucis terminalibus capitatis vel insuper ex axillis supremis ortis breviter racemosis, calycis laciniis linearibus tubum æquantibus vel dimidio brevioribus, corollæ ochroleueæ clavatae lobis ovatis obtusis vel acutiusculis plicis integris vel bitidis sejunctis ♀.

Hab. in regione alpinâ Anatoliae borealis in monte Jyldisdag (Wied!), Armeniae Turcice in monte Techdag prope Erzerum (Huett!), Bingöldagh (Tchilt! Ky. 497!), Armeniâ Rossicâ (Szov! Seidl!), montibus Iberia Caucasicæ (Stev! etc.)

Valde affinis *G. septempfida*, praeter florum colorem tamen diversa videtur foliis angustis, corollæ plicis non fimbriatis.

16. ***G. calycina*** (Boiss. et Haussk.) caulis a collo nudo ascendentibus simplicibus foliosis, foliis infimis squamæformibus ceteris quinquenerviis ovatis obtusiusculis basi subcordatis, floribus terminalibus subsolitariis, calycis laciniis elliptico-linearibus obtusis basi subattenuatis subinæqualibus tubo membranaceo longioribus, corollæ cyanæ lobis ovatis obtusis plicis acute 2-3-fidis sejunctis ♀.

Hab. ad rivulos moutis Berydagh Cataoniae 7000' (Haussk!).

Quoad habitum et folia valde similis *G. septempfida* sed distincta calycis laciniis longioribus latioribus foliiformibus basi ut in *G. Boissieri* sed magis angustatis, multo majoribus et saepius inæqualibus

17. ***G. Boissieri*** (Schott et Ky. in Ky. pl. exs. 1853) caulis ex collo fibroso-squamato decumbenti-inecurvis simplicibus brevissimis crebre foliosis unifloris, foliis parvis ovatis obtusis trinerviis,

calycis laciniis anguste elliptico-linearibus acutiusculis tubo brevioribus basi subattenuatis sinibus latis sejunctis, corollæ cyaneæ clavatae lobis breviter ovatis obtusiusculis plicis breviter 3-4-dentatis sejunctis ♀.

Hab. in regione alpinâ Tauri Cilicici ad Bulghar Magara et Gisyl Tepe 8000' (Ky. 168! Bal!), in monte Aslandach Cappadociæ (Bal!)

Caules 2-3-pollicares, folia 3-4 lineas longa, corolla eā *G. septempidae* vix brevior. Ab hujus speciei formis microphyllis collo fibroso, foliis obtusis, calycis laciniis basi angustatis, plicis non fimbriato-multifidis, lobis obtusiusculis distincta. Facie *G. pneumonanthem* var. *depressam* Hispanicam omnino refert sed haec foliis linearibus uninerviis statim dignoscitur.

18. ***G. asclepiadea*** (L. Sp. 329) caulis elatis simplicibus crebre foliosis, foliis amplis sessilibus a basi ovatâ vel subcordatâ longe lanceolato-acuminatis quinquenerviis margine sebris, floribus sessilibus oppositis in axillis superioribus terminalibusque breviter racemosis foliis floralibus superatis, calycis interdum fissi dentibus anguste linearibus tubo apice truncato brevioribus, corollâ cyaneâ longe clavatâ lobis breviter lanceolatis acutis sinibus denticulatis ♀. Ic. Jacq. Austr. tab. 328. — Rchb. tab. 11. — *G. schistocalyx* C. Koch Linn. XXIII, p. 585!

Hab. in sylvaticis subalpinis, Olympus Thessalus (Heldr!), Olympus Bithynus (Noë!), Pontus Lazicus supra Trapezuntum (Bourg!) et supra Khabakhar 6000' (Bal!), in Caucaso occid. et centrali 2400'-5000' (C.A.M., Rupr!), ditione Talysh (Hoh!).

Ar. Geogr. Montes Galliae austro-orientalis, Hispaniæ borealis, Helvetiæ, Germaniæ, Corsicæ, Italiæ borealis et mediæ, Dalmatiæ, regionis Danubialis.

19. ***G. cruciata*** (L. Sp. 334) caulis extrarosularibus ascendentibus crassiusculis crebre foliosis simplicibus rarius ramosis, foliis oblongo-lanceolatis obtusis trinerviis radicalibus basi attenuatis caulinis in vaginam inferiorum longiusculam connatis, floribus sessilibus ad axillas superiores fasciculatis vel tantum terminalibus capitatis foliis floralibus longe superatis, calycis breviter campanulati interdum lateraliter fissi membranacei dentibus brevibus linear-lanceolatis acutis inæqualibus, corollæ tubulosæ cyaneæ extus griseæ lobis quaternis ovatis erectis plicis breviter lanceolatis dentatis ♀. Ic. Jacq. Austr. t. 372. — Rchb. tab. 11.

Hab. in pascuis siccis regionis subalpinæ et alpinæ in Tauriâ (Stev.), Anatoliâ boreali prope Samsun et Tokat (Wied!), Armeniâ Turcicâ prope Erzerum (Huet!), Baibut (Bourg!), ad radices australes Bingöldagh (Ky. 361!), Caucasi centralis et orientalis 3000'-8000' (M.B., C.A.M., Rupr!), Transcaucasiæ et prov. Talysh (C.A.M.).

Ar. Geogr. Europa media et australis a Belgio, Germaniâ, Hispaniâ boreali et Italia bor. et mediæ ad regionem Danubialem, Rossiam medium, Sibiria omnis.

20. ***G. Olivieri*** (Griseb. Gent. p. 278) collo fibroso, caulis

extrarosularibus ascendentibus simplicibus, foliis radicalibus oblongo vel lanceolato-spathulatis obtusis in petiolum sensim attenuatis quinque nerviis, caulinis paucis anguste linearibus remotis connatis, floribus breviter pedicellatis vel sessilibus 3-7 in cymam corymbosam simplicem vel compositam approximatis, calycis lacinias lanceolatis acutis tubo æquilonigis vel brevioribus, corolla cyanea obconicae calyce 2 $\frac{1}{2}$ -plo longioris lacinias oblongis obtusiusculis plicas triangulares denticulatas duplo superantibus, antheris liberis ♂.

Hab. in graminosis et collibus regionis inferioris et subalpinis usque ad 5000', Cappadocia ad radices Tauri (Auch. 2429!), Mesopotamie deserta ab Aintab et Nisib usque ad Mossul et montem Sindjar (Oliv! Haussk!), Kurdistania (Barre!), Persia austro-occid. ad Kotel Mallu (Haussk!) et prope Gere (Ky. 49!), Ispahan (Auch. 4957!), Persia bor. in prov. Aderbidjan (Szov!), ad radices montis Demavend (Ky. 327!), in prov. Khorassan (Bge!), Affghania (Griff!), Belutschia (Stocks!), Armenia Turcica circa Erzerum (Huet! Tchih!), Pontus prope Niksar (Tchih!). Fl. Apr. Mai.

Inter congeneres statione in regione calida insignis, variat foliis radicalibus 4-8 lineas latis 2-3-pollicaribus, caules semipedales raro pedales, corolla 10-13 lineas longa.

Ar. Geogr. Turkestania, Songaria.

PLEUROGYNE (Escâh. in Linn. I, p. 487. — *Lomatogonium* A Br.
Fl. 1830, I, p. 221)

Calyx quinquepartitus. Corolla marcescens rotata profunde quinquepartita glandulis, plicis, foveisque destituta, segmentis expansis basi fimbriarum serie coronatis. Stylus nullus. stigmata bina lateralia secus valvularum margines decurrentia. Capsula unilocularis septicida bivalvis, placentis ad vel juxta suturam decurrentibus. — Herbae annuae.

I. P. Carinthiaca (Wulf. in Jacq. Misc. II, 53, tab. 6 sub *Swertia*) pumila pluricaulis et basi ramosa, foliis minutis radicalibus obovatis obtusis cæteris ovatis, pedicellis elongatis nudis, calycis lacinias oblongis acutis corollæ pallide cerulea extus olivaceo-cyanea parte dimidiâ brevioribus, corollæ lacinias ellipticis acutis ovario obtuso, seminibus ad suturam insertis ②. **P. Carinthiaca** Griseb. Gent. p. 310. — Rehb. tab. 4. — *Lomatogonium Carinthiacum* A. Br. loc. cit. — *Swertia rotata* Pall. Ross. tab. 89 non L. — *Gentiana Stelleriana* Ch. et Schl. — *G. rotata* M.B.

Hab. in pratis alpinis Caucasi (M.B.), prope Kobi (Rehm!), in monte Schorudagh Daghestaniæ (Chodsko!), in Affghania alpibus prope Yomut (Griff!). Fl. aut. Specimina Caucasicæ Carinthiacis similissima.

Ar. Geogr. Alpes Valesiae, Carinthia, Tyrolis australis, Sibiriae Altaiæ et Baicalensis, Dahuria, regio Himalaiae. Planta Americana citata ad *P. rotata* potius spectare videtur.

SWERTIA (L. Gen. N° 322 ex parte).

Calyx 5-rarius 4-partitus. Corolla rotata limbo plano 5-rarius 4-partito laciniis basi foveis binis vel unicâ ambitu fimbriatis instructis. Stylus nullus. stigma terminale continuum vel bilobum. Capsula unilocularis bivalvis septicida placentis juxta suturas expansis. — Herbæ (in nostris) perennes.

1. S. punctata (Baumg. in Rehb. Crit. tab. 463) caule elato in racemum longum strictum compositum abeunte, foliis radicalibus longe petiolatis amplis ovato-oblongis obtusis et saepius mucronatis, caulinis omnibus alternis diminutis superioribus sessilibus semiamplexicaulibus, floribus pentameris, calycis laciniis lanceolatis, corollæ flavidæ nigro-punctatæ laciniis elliptico-lanceolatis acutis calyce duplo longioribus, foveis binis oblongis longe fimbriatis, seminibus late alato-marginatis ♀. *S. perennis* M.B. non L. — *S. Iberica* Fisch. MSS. — *S. obtusa* β *albiflora* Ledeb. ex parte. — *S. stigmantha* C. Koch Linn. XXIII, p. 586.

Hab. in pratis humidis subalpinis et alpinis, Caucasi occid. (Radde!), centralis et orientalis 480'-7500' (Rupr!, Ower!), Iberiæ Caucasicæ (Hohen!).

Sæpe bipedalis et elatior, differt a *S. perenni* foliis caulinis alternis, corollæ foveis magis elongatis, a *S. obtusa* Ledeb. Altaicâ præter florum colorem et folia radicalia ampla longe petiolata laciniis corollinis acutis.

Ar. Geogr. Alpes Hungariæ et Transylvaniæ.

2 S. Balansæ, caule elato in racemum elongatum strictum compositum foliosum abeunte, foliis omnibus oppositis radicalibus longe petiolatis amplis elliptico-oblongis acutiusculis, caulinis diminutis superioribus sessilibus florum racemulos superantibus et æquantibus, floribus tetrameris, calycis laciniis lanceolatis longe acuminatis corollâ quartâ parte brevioribus, corollæ ochroleucæ concoloris laciniis oblongis acutiusculis, foveis binis rotundatis longe fimbriatis ♀.

Hab. in pratis alpinis Ponti Lazici prope Djimil 6000' (Bal! exs. sub *S. Aucheri*):

Species non absque dubio proposita *S. perenni* multo robustior et ab ea corollæ ochroleucæ laciniis abbreviatis latis discedens; facies *S. punctata* foliis caulinis alternis corollâque punctatâ et *S. Aucheri* corollæ laciniis unifoveatis distinctæ; affinis quoque *S. connata* Schrenck, sed folia caulinæ non connata.

3. S. Aucheri (Boiss. Diagn. Ser. I, V, p. 90 (1844) caule crasso elato erecto, foliis omnibus oppositis inferioribus oblongis obtusiusculis in petiolum latum attenuatis, caulinis diminutis oblongo vel linearí-lanceolatis plus minus connatis, floribus tetrameris simpli-

citer racemosis vel ample et stricte paniculatis, sepalis lanceolatis acutis corollâ parte dimidiâ brevioribus, corolle ochroleucæ concoloris laciinis oblongis acutiusculis, foveis in unicam orbiculatam coalitis ♀. *S. Persica* Griseb. in p.c. Prodr. IX, p. 132 (1843). — Jaub. et Sp. Illustr. Or. t. 235. — *S. obtusa* C. Koch Linn. XXIII, p. 586 non Ledeb.

Hab. in pratis humidis alpinis Armeniae (Auch!), ad Zylzylkhan inter Baibut et Erzerum 5000' (Ky. 665 sub *S. longifoliâ*!), inter Sepigor et Erzinghan (Tchih!), in Persia (Auch. 2433!).

Planta 1 ½-2-pedalis, folia inferiora ampla.

4. ***S. longifolia*** (Boiss. Diagn. Ser. I, 5, p. 90) caule elato simplici in racemum valde interruptum superne abeunti, foliis omnibus oppositis radicalibus longis lanceolatis vel lanceolato-spathulatis basi longe attenuatis obtusis vel acutiusculis, caulinis paucis valde diminutis linearibus basi connatis, floralibus racemos breves fasciculiformes vix æquantibus, floribus tetrameris, sepalis breviter lanceolatis corollâ duplo brevioribus, corolle flavidæ concoloris laciinis ovatis acutiusculis, foveis in unicam transverse ovatam longe fimbriatam coalitis ♀.

Hab. in Persia borealis monte Elbrus (Auch. 4960!), in uliginosis alpinis Sehir Pela montis Totschal prope Teheran (Ky. 575!).

Bi-tripedalis, folia radicalia 4-10 pollices longa 6-8 lineas lata; species foliis radicalibus elongatis angustis non distincte petiolatis, inflorescentia remote pseudoverticillata, petalis latis a *S. Aucheri* distincta.

5. ***S. lactea*** (Bge Pl. Lehni. p. 393), caule erecto in racemum valde interruptum superne abeunti, foliis omnibus oppositis basi liberis infinitim anguste spathulato-oblongis superioribus diminutis linearibus obtusiusculis, florum fasciculis oppositis inferioribus longe superioribus breviter pedunculatis, floribus tetrameris, sepalis lanceolato-linearibus inaequalibus acutiusculis dimidiâ corollâ longioribus, corolle (lacteo-lilacinæ) laciinis ellipticis obtusis vel retusis, foveis in unicam orbiculatam fimbriatam coalitis ♀.

Hab. in monte Karatau Turkestaniæ (Lehm!), in regione alpinâ prov. Khorassan (Bge!).

Sesquipedalis, differt a *S. Aucheri* et *longifoliâ* foliis angustioribus omnibus liberis, corolle laciinis angustioribus.

ORD. LXXXI. BIGNONIACEÆ.

(R. Br. Prodr. 470.)

Calyx gamophyllum quinquelobus vel integer truncatus vel spatha-
ceus aut bilabiatus. Corolla hypogyna decidua quinqueloba sœpissime

irregularis bilabiata. Stamina 5 corollæ tubo adnata sæpius 4 fertilia didynama quinto sterili. Discus glandulosus circa ovarii basin. Ovarium biloculare pluriovulatum, stylus filiformis, stigma bifidum. Fructus (in nostris) capsularis bilocularis bivalvis, septo valvis nunc parallelo nunc contrario demum libero ad commissuram cum valvulis seminifero. Semina transversa vel pendula compressa exalbuminosa alata. Embryo rectus.

TECOMA (Juss. Gen. 439).

Calyx campanulatus quinquelobus. Corolla campanulata fauce valde ampliata. Antheræ biloculares loculis divergentibus. Capsula (in nostrâ) siliquæformis elongata bilocularis bivalvis septo valvis contrario. Semina transversa alata. — Arbores vel frutices interdum scandentes..

1. **T. undulata** (Roxb. Fl. Ind. III, p. 101 sub *Bignonia*) arbuseula ramosa non scandens ramis divaricatis cinereis, ramulis novellis abbreviatis fasciculatim foliosis, foliis oppositis glaucis subcoriaceis breviter petiolatis oblongis vel cuneatis integerrimis interdum subundulatis, racemis paucifloris terminalibus, pedicellis brevisimis, calyce campanulato breviter et late quinquelobo, corollâ amplâ puniceâ lobis rotundatis subæqualibus, capsulâ longâ linearî lœvi siliquæformi subcompressâ ♂. *T. undulata* Don Gen. Syst. IV, p. 223. — Smith Exot. bot. I, p. 19. — *Bignonia glauca* Decaisne Jacq. Voy. p. 137, tab. 142.

Hab. in Belutschia (Stocks!), in regno Mascate Arabiæ (Auch.).

Ar. Geogr. India boreali-occidentalis.

AMPHICOME (Lindl. Bot. Reg. tab. 49).

Calyx oblongo-campanulatus 5-fidus lobis acutis. Corolla longe infundibuliformis subbilabiata. Antheræ per paria stylo arcte adpressæ lobis pilosis calcarato-aristatis. Capsula siliquæformis bilocularis suturâ alterâ dehiscens. Semina pendula utrinque comosa. — Herbae erectæ.

1. **A. Emodi** (Royl. et Lindl. loc. cit.) glabra, foliis alternis petiolatis pinnatisectis segmentis cum impari 2-3-jugis ovato-lanceolatis petiolatis non confluentibus serratis, racemo terminali laxo paucifloro, calycis lobis abbreviatis acutiusculis, corollâ roseâ ♀.

Hab. in Affghania (Griff.).

Ar. Geogr. Regio Himalaica occidentalis.

ORD. LXXXII. SESAMEÆ.

(D.C. Théor. Ed. II, p. 247. — *Pedalineæ* Lindl.)

Calyx 5-partitus vel fidus subæqualis. Corolla hypogyna irregularis fauce ventricosa limbo quinquelobo saepius bilabiato. Discus hypogynus carnosus-glandulosus. Stamina quina corollæ inserta superiori sterili cæteris didynamis fertilibus, antheræ biloculares connectivo ultra loculos producto. Ovarium 2-4 rarius 8-loculare, stylus unicus, stigma lobatum. Fructus capsularis dehiscens vel drupaceo-nucamentaceus indehiscens carpellis normaliter 5 sed abortu 2-4. Semina secus valvularum margines vel columnam centralem valvis adnatam uniseriata exaluminosa. Embryo rectus.

SESAMUM (L. Gen. 782).

Calyx persistens lobo superiore minore. Corolla tubo sensim ampliato, limbo bilabiato lobis superioribus binis brevioribus inferioribus tribus. stigma bilamellatum. Capsula prismaticæ tetragona quadrisulcata bilocularis sed valvis intus recurvis parte superiori exceptâ quadrilocularis apice septicide dehiscens. Semina numerosa aptera columnæ centrali affixa. — Herba annua.

1. **S. Indicum** (L. Sp. 884) caule erecto pubescente simplici vel stricte ramoso, foliis inferioribus oppositis ovatis trisectis vel trilobis lobis integris vel dentatis, superioribus alternis oblongis vel linear-lanceolatis breviter petiolatis integris, floribus in axillis solitariis breviter pedicellatis, capsulâ styli basi persistente mucronatâ velutino-pubescente ⊙. Ic. Endlich. Ic. tab. 70. — *S. Veterum* Bauh. — *S. Orientale* Cham.

Hab. in calidioribus ditionis cultum et hinc inde in agris spontaneum, v. gr. in Argolide (Sprun!), Lydiâ prope Sardes (Boiss!), Ægypto! Susianâ (Haussk!).

Ar. Geogr. Ex Indiâ orientali prob. oriundum in regionibus subtropicis et tropicis totius orbis cultum.

ORD. LXXXIII. CYRTANDRACEÆ.

(Jack Trans. Linn. Soc. XIV, p. 23.)

Calyx 5-fidus vel 5-partitus saepius æqualis. Corolla hypogyna basi tubulosa superne plus minus ampliata limbo 4-5-lobo saepius inæquali bilabiato. Stamina corollæ adnata 2, 4, rarius 5 fertilia, didynama

postico saepius sterili vel libera. Ovarium liberum annulo glanduloso cinctum, stylus simplex. Fructus rarius baccatus saepius capsularis bilocularis vel unilocularis, dehiscentia loculicida vel septicida. Semina numerosa exalbuminosa, embryo minutus.

HABERLEA (Friv. in Act. Hung. 1835, II, p. 249. tab. 1).

Calyx profunde quinquefidus, subbilabiatus labio superiore quinquentato inferiore paulo infra medium bifido. Corolla infundibuliformis bilabiata labio superiore bilobo inferiore trilobo. Stamina inclusa quatuor didynama filamentis apice inflexis, antheris per paria connatis, stamen quintum rudimentarium (interdum fertile) filamentis duobus brevioribus corollæ parte superiori adnatis interpositum. Stigma capitatum. Ovarium marginibus carpellorum introflexis et adnatis biloculare. Capsula obconica stylo mucronata bilocularis septicida. Semina numerosa oblonga. — Herbæ scapiferae foliis et facie *Ramondiae* vel *Rehmanniæ*.

1. **H. Rhodopensis** (Friv. loc. cit.) saepius dense et late cespitosa rosulis confertis, foliis crassiusculis oblongo-spathulatis basi attenuatis obtusis profunde crenato-serratis superne viridibus utrinque sed subtus copiosius setis basi subincrassatis hirsutis et ciliatis, scapis hirtis foliis sublongioribus subumbellatim 2-5-floris rarius unifloris, bracteis linear-setaceis, pedicellis calyci æquilongis vel longioribus, calycis hirti nigricantis laciniis triangulari-lanceolatis acutiusculis, corollæ violaceæ lobis rotundatis labii superioris abbreviatis, capsulæ calyci æquilonga 2. Ie. Endlich. Icon. tab. 69 (corollæ formâ male exhibitâ).

Hab. in saxis schistosis umbrosis faucium Akdere montis Balkan Thraciæ prope Kalofer in regione sylvaticâ (Friv! Janka! Pichler!). Fl. Junio.

Folia 2-3 pollices longa saepè in speciminibus pusillis multo minora, corolla pollicem aut amplius longa. Floræ Thracice eximium decus. Corollæ formam ex litteris cl. a Janka qui plantam vivam observavit restitui.

JANKÆA (*Jancæa* Boiss. Dec. Prima, p. 4).

Calyx quinquepartitus. Corolla breviter campanulato-infundibuliformis subæqualiter quadriloba. Stamina quatuor libera filamentis rectis. Capsula ob valvulas breviter introflexas unilocularis. — Genus facie *Haberlean* et *Ramondiam* omnino referens; prior differt corollâ bilabiata quinquelobâ, staminibus didynamis et filamentis apice inflexis, capsulâ ob valvulas longe introflexas biloculari; posterior corollâ rotatâ, staminibus quinis, filamentis brevissimis distinguitur. — Genus cl. Victor de Janka Flora regionis Danubialis scrutatori strenuo, qui primus hanc plantam ab *Haberlea* specificie differre cognovit dicatum.

t. J. **Heldreichii** (Boiss. Dec. Prim. p. 5) rosulis solitariis, collo dense rufo-lanato, foliis crassiuseulis ovatis integris obtusis basi breviter attenuatis adpresso et dense superne albo-sericeis subtus ferrugineo-lanatis, scapis rosulâ duplo longioribus glanduloso-hirtis apice 1-2-floris ebracteatis, pedicellis calyce sublongioribus, calyce glanduloso fere ad basin in lacinias ovatas obtusas partito: corollâ violacea campanulatâ ad medium in lacinias 4 obovatas subæquales partitâ, staminum filamentis rectis antherâ non longioribus, capsulâ calyce duplo longiore ♀. *Haberlea Heldreichii* Boiss. Diagn. Ser. II, 3, p. 141. — *Ramondia Heldreichii* Janka in litt.

Hab. in rupibus regionis mediae et superioris montis Olympi Thessaliae supra Hagios Dionysios 4000'-8000' (Heldr! Orph. Fl. Gr. exs. 625!).

Folia pollicem longa, corolla jam a basi ampliata 8-9 lineas tantum longa.

ORD. LXXXIV. POLEMONIACEÆ.

(Vent. Tabl. II, p. 398.)

Calyx liber quinquefidus vel partitus persistens. Corolla monopetala hypogyna regularis vel subirregularis quinquefida lobis aestivatione contortis. Stamina 5 tubo corollæ inserta ejus lacinis alterna. Ovarium disco hypogyno cinctum biloculare, stylus unicus, stigma trifidum. Ovula in loculis solitaria vel plura angulo centrali lateraliter affixa. Capsula trilocularis trivalvis valvis medio septiferis ab axi centrali trigonâ plus minusve complete solutis. Semina testâ spongiosa donata, albuminosa, embryone axilli recto.

POLEMONIUM (L. Gen. 217).

Calyx campanulatus quinquefidus. Corolla rotato-campanulata tubo brevissimo, fauce filamentis basi dilatatis declinatis clausâ. Discus cupuliformis crenatus. — Herbae foliis pinnatis.

1. **P. cæruleum** (L. Sp. 230) glabrum vel viscoso-pubescent, caule elato folioso, foliis pinnatis segmentis ovato-lanceolatis acuminatis acutis integris, paniculâ corymbosâ glanduloso-pilosâ, calycis ad medium quinquefidi lacinis ovatis acutis, corollâ cæruleâ calyce triplo longiore, antheris ovatis, ovarii loculis 6-10-ovulatis ♀. Lam. III. tab. 106.

Hab. in pratis paludosis, faucibus umbrosis, Caucasi regio subalpina et alpina (C.A. Mey!), circa Kasbek (Rehm!).

Ar. Geogr. Europa a Scandinaviâ ad Rossiam arcticam et Hispaniâ et Italia borealis ad regionem Danubialem, Sibiria omnis, America borealis.

ORD. LXXXV. CONVOLVULACEÆ.

(Juss. Gen. 132.)

Calyx liber quinquespalus vel 4-5-sidus persistens. Corolla gamopetala hypogyna regularis decidua tubulosa campanulata vel infundibuliformis limbo quinquelobo longitudinaliter plicato. Stamina 5 fundo corollæ inserta ejus lobis alterna. Nectarium annulare plerumque circa ovarium. Ovarium (in nostris) unicum simplex liberum 2-4-loculare, rarius subuniloculare, loculis 1-2-ovulatis, ovula erecta. Stylus 1 apice bifidus vel styli 2. Fructus capsularis varie dehiscens vel sicco-baccatus indehiscens. Semina basi angulorum centralium fructus inserta dissepimentis incompletis fundum capsule occupantibus sejuncta. albumen parcum.

TR. I. EUCONVOLVULEÆ.

Plantæ foliis et cotyledonibus præditæ. Sepala 5.

CONVOLVULUS (L. Gen. 215).

Calyx quinquespalus. Corolla infundibuliformi-campanulata angulo-quinquloba vel repanda quinqueplicata. Stylus unicus stigmatibus binis linearis-cylindricis sœpe revolutis. Ovarium biloculare loculis biovulatis. Capsula bilocularis loculis dispermis bivalvis. — Herbae vel frutices et suffrutices. — Genus in sectiones naturales ægre dividendum: greges hic propositæ sœpe nimis artificiales, nam limites inter species ramis tandem spinescentibus vel inermibus, caulinis scandentibus vel subscendentibus, rhizomate plus minusve suffrutescenti donatas interdum incerti sunt.

Specierum Orientalium distributio.

SERIES A. — Perennes.

§ 1. **Acanthocladi.** — Fruticosi vel suffruticosi, rami vetusti tandem spinosi vel spinescentes. Ovarium hirsutum.

C. leiocalycinus, acanthocladus, spinosus, fruticosus, erinaceus, oxyphyllus.

§ 2. **Spinescentes.** — Fruticosi vel suffruticosi, ramis vetustis tandem spinosis vel spinescentibus vel saltem persistentibus. Ovarium glabrum.

C. hystrix, virgatus, lanatus, ulicinus, Scindicus, oxysepalus, eremophilus, leptocladus, evolvuloides, Dorycnium.

§ 3. **Inermes.** — Fruticosi vel basi tantum suffrutescentes inermes. Ovarium hirsutum.

* Rami scroparii elongati persistentes.

C. sarothrocladus, chondrilloides.

* * Rami fruticosi

C. Cneorum, oleifolius, Cyprinus, Schirazianus.

* * * Rami e rhizomate suffrutescenti herbacei.

C. Calverti, commutatus, holosericeus, Cantabrica, cardiosepals, Ancherei, Ruprechtii, lineatus, Libanoticus, Cataonicus

§ 4. **Compacti.** — Pulvinares acaules. Ovarium hirsutum

C. compactus, cochlearis, Assyriens, radicosus

§ 5. **Pannosi.** — Rami inermes crassiusculi e rhizomate indurato herbacei stipes ascendentes vel decumbentes. Ovarium glabrum. — Plantæ villosissimæ vel tomentosæ

C. Persicus, secundus, reticulatus, waltherioides, sericeus, Schimperi, Kotschyanus, Hansknechtii, cephalophorus

§ 6. **Diffusi.** — Rami tenues herbacei prostrati vel ascendentes non scandentes. Ovarium glabrum.

C. glomeratus, pilosellifolius, microphyllus, Ammocharis

§ 7. **Volubiles.** -- Caules herbacei scandentes vel subscandentes. Ovarium hirsutum.

C. Galaticus, Germanicus, hirsutus, Armenus, stenophyllus

§ 8. **Scandentes** — Rami herbacei plus minus scandentes Ovarium glabrum.

C. ulthroides, tenuissimus, stachydifolius, Palestinus, Scammonia, arvensis

SERIES B. — Annui.

§ 9. **Siculi** — Capsula glabra.

C. Fatmensis, Cirelesyriacus, Siculus, pentapetaloides, Stocksie

§ 10. **Undulati.** — Capsula hirsuta

C. undulatus.

SERIES A. — Perennes.

§ 1. **Acanthocladi.**

1. **C. leiocalycinus** (Boiss. Diagn. Ser. I, 7, p. 28) fruticosus, ramis elongatis patule ramulosissimis ramulis brevibus rigidis junioribus adpresso sericeis velutinis denudatis apice pedunculisque in spinas breves acutatis, foliis parvis breviter petiolatis paree puberulis vel glabris a basi obtuse cordato-sub hastata ovato vel oblongo-triangularibus obtusis interdom obtuse repandis superioribus angustis, floribus ad ramulorum axillas inferiores solitariis pedunculis crassinervulis folio brevioribus versus medium articulatis et bracteolatis suffultis, sepalis glabris seariosis ovatis obtusis mucronulatis, corollâ albâ glabrinervulâ calyce 5-6-plo longiore, ovario villoso ♂.

Hab. in Persia australi (Auch. 4954!), in rupes tribus apriis faneum inter Abchir et Schiraz (Ky. 39!), in apriis montium Sawers, Nur, Eschker 6-8000 (Hanssk!), in Persia austro-orientali interiori inter Khabbise et Kerman (Bgeb. Fl. Mart. Jun).

♂. *Stocksii* Boiss. Diagn. Ser. II, 3, p. 423. — Folia angustiora basi interdom cuneata sericea. — *C. lasiophorus* Janb. et Sp. III, tab. 368

Hab. in Persia australi prope Schiraz (Auch. 4935!), in Belutschia inferiore (Stocks!)

♀. *lycoides*. — Glabrescens viridis, folia superiora anguste linearia acutiuscula, pedunculi folio sublongiores, sepala oblonga acutiuscula.

C. lycoides Boiss. Diagn. Ser. I, 7, p. 29

Hab. in collibus calcareis prope Schiraz (Ky!).

Frustr 3-4-pedalis, folia majora 4-6 linea longa, corolla pollicem aut tere amplius longa. Species foliis cordato-hastatis a sequentibus distincta.

2. **C. acanthocladus** (Boiss. Diagn. Ser. I, 7, p. 27) foliis adpresso-erectis enim suffruticosus ramosissimus erinaceus, ramis intricatissimis brevibus rigidis floriferis apice dichotome cymosis velutinis in spinas tenues breves alienantibus, foliis parvis inferioribus oblongo-spatulatis obtusis basi longe attenuatis, floralibus anguste linearibus, floribus parvis ad cyme ramulos brevissimos terminalibus brevissimis peticello articulato curvato brevissimo suffultis, sepalis patule et longe hirsutissimis oblongo-lanceolatis breviter acuminatis corollâ albâ calyce triplo longiore ad angulos longe hispida, ovario villoso ♂. Janb. et Sp. III, tab. 367.

Hab. in collibus aridis Persiae australis (Auch. 4934!), circa Dalechi et Schiraz 1000-3000' (Ky. 208! et 352! Hanssk!) Fl. Apr.

Suffrutex sespipedalis demum spinis ramosis erinaceus, foliis inferioribus 2-4 linea longa, calyx 4 linea longus. Species floribus patule hirsutissimis notabilis.

3. **C. spinosus** (Lurm. Ind. 47, tab. 19, fig. 4 et herb! non Desr. nec Eichw.) totus breviter et adpressissime cinereo-sericeus suffruticosus ramosissimus erinaceus ramis creberrimis intricatis rigidis apice tenuiter spinosis, foliis minutis ellipticis acutiusculis basi attenuatis subpetiolatis superne concavo-subuplicatis superioribus linearibus squamiformibus, floribus ad ramorum juniorum partem inferiorem in apice ramuli patentis rigidi calyce sublongioris articulatis brevissime pedicellatis erectis solitariis rarius 2-3-nis, bracteolis binis minimis ad pedicelli basin, ramulis floriferis vetustis superioribusque sterilibus abbreviatis spinescentibus, calycis oblongi adpresso hirsuti sepalis ovatis cuneatibus arce imbricatis obtusis interdum submucronatis, corollæ soriceo-hirsutæ lobatæ calyce 3-4-plo longiore, ovario hirsuto 3. *C. genistoides* J. et Sp. Ill. t. 370.

Hab. in desertis Persiae australis (Burm!, Auch 4933!), in Belutschia inferiore (Stocks exs. 695!). Affghania (Griff!).

Frutex ut videtur humilis sed intricatissimus, folia 4-3 lineas longa, flores in ramulo 3-4 lineas longo subhorizontaliter patenti ob pedicellum sursum infracto-curvatum erecti 8-9 lineas longi Rami, ramulis florales et etiam pedicellarum sterilius sub flore rudimenta spinescentia.

4. **C. fruticosus** (Pall. It. III, p. 144, table V) totus adpresso argyreus, fruticosus, ramis erectis divaricatim et breviter ramulosus, foliis inferioribus longe linearispathulatis obtusis inferne longe attenuatis, floralibus subsessilibus anguste elliptico-linearibus acutis, floribus ad ramulorum floralium brevium horizontalium apicem breviter pedicellatis articulatis erectis solitariis rarius 2-3-nis, ramulis floralibus pedicellisque tandem in spinas breves tenues mutatis, bracteis linearibus elongatis, calycis adpresso hirsuti sepalis laxe imbricatis ellipticis acutis, corollæ extus sericea obscure lobata calyce triplo longiore, ovario hirsuto 3. *C. spinosus* Desr. Eneyel. III, p. 548 et Eichw. Casp. tab. 22. — *C. Eichwaldii* Boiss. Diagn. Ser. I, 7, p. 27 in notâ. — *C. frutescens* Pall. — *C. brerispinus* Jaub. et Sp. Ill. IV, tab. 369.

Hab. in aridis Persiae australis prope Schiraz (Auch. 4937 ex J. et Sp.), prope Ispahan inter Gias et Murtsehehar (Bge!), inter Ispahan et Kaschan (Buhset!), Persia boreali prope Schahrod (Bge!), Turcomania littoralis ad Caspium et desertis Kisilkum atque ad fluvium Syr Darja Turkestanie (Lehm!).

Inflorescentia *C. spinosi* sed rami elongati, folia multo majora, inferiora sesquipollicem longa, bracteæ elongate calyce sèpè longiores, sepalæ acutiora laxe nec arce imbricata, corolla vix lobata.

Ar. Geogr. Songaria, Sibiria Uralensis et Altaica.

5. **C. erinaceus** (Ledeb. in Eichw. Casp. p. 44, tab. 7) adpressissime canescens-soriceus suffruticosus ramosissimus erinaceus ramis longissimis ramulisque rectangule patentibus tenuibus apice aphyllis et abortivis nec spinosis, foliis anguste et longe linearibus acutiusculis floralibus brevibus squamiformibus, cymis 3-5-pedunculis axillaribus rigidis folio longioribus suffultis superioribus brevius

pedunculatis abortu unifloris, pedicellis calyce brevioribus supra medium articulatis et bracteolatis, sepalis sericeis ovato-spathulatis obtusissimis, corollâ extus sericeâ calyce duplo longiore, ovario hirsuto ♂.

Hab. in littore orientali maris Caspii (Eichw., Karel!), in desertis Kisylkum et inter Buchara et Kermine Turkestaniæ (Lehml!).

Basi lignosus, rami bipedales subflexuosi divaricatum ramosissimi, folia lineam lata inferiora sesquipollucaria, calyx minutus sesquilineam longus.

Ar. Geogr. Sibiria Uralensis.

6. *C. oxyphyllus* (Boiss. Diagn. Ser. I, 7, p. 26) patule tomentosus suffruticosus a basi ramosissimus erinaceus, ramis virgatis rigidis dichotome et divaricatum ramulosus apice demum subspinescentibus, foliis minutis sessilibus oblongis et ellipticis acutis mucrone subspinoso, superioribus ad squamas reductis, fasciculis 1-3-floris sessilibus dissitis spicas interruptas longas formantibus, bracteis ovatis, sepalis hispidissimis lanceolatis acuminatis, corollâ hispidâ calyce subdupo longiore, ovario villoso ♂.

Hab. in arenosis Assyriæ (Auch. 3840! a cl. Choisy ad *C. sericeum* erroneè adductus), prope Kerkuk (Haussk!), in Persia australi (Auch. 4950!), inter Seytun et Behehan (Haussk!). Fl. Jun.

Rami virgati paniculato-spicati sœpe pedales et sesquipedales interdum breviores, folia inferiora 4-6 lineas longa superiora valde diminuta, calyces 4 lineas longi, habitus erinaceus. Species foliis apice mucronato-subspinosis notabilis.

§ 2. Spinescentes.

7. *C. hystrix* (Vahl Symb. I, p. 16) fruticosus ramosissimus erinaceus pilis adpressiusculis vel patulis plus minus obsitus, ramulis numerosis brevibus rigidis apice spinosis, foliis parvis subsessilibus basi truncato-subauriculatis oblongis acutis ramulorum squamiformibus, floribus ad ramulorum basin et axillas inferiores sessilibus solitariis vel 2-3-nis bracteis oblongis apice subrecurvis involucratis, sepalis scariosis hirsutis obovatis cucullatis mucronatis internis angustioribus, corollâ hirsutâ angustâ calyce vix duplo longiore, ovario glabro ♂. *C. spinosus* Forsk. Cat. p. 106 non Burm. — *C. armatus* Del. Fl. Eg. p. 201, tab. 18 quoad analyses pessima.

Hab. in desertis Ægyptiaco-Arabicis (Del.), ad littus Ægyptiacum maris Rubri (Schweinf!), in valle Abraham deserti Sinaitici (Auch. 1391!). Fl. Mart. Apr.

Frutex erinaceus sœpe hemisphæricus 4-5-pedalis ramis primariis sœpe 2-3-pedalibus, folia inferiora 4-5 lineas tantum longa.

Ar. Geogr. Nubia ad mare Rubrum, Arabia subtropica.

8. *C. virgatus* (Boiss. Diagn. Ser. I, 7, p. 24) suffruticosus glaucus ramosissimus, ramis tenuibus rigidulis elongatis præter pilos

sparsissimos glabris, foliis paucis subsessilibus præter pilos nonnulos glabris brevissime petiolatis anguste linearibus basi minute angulato-subhastatis, capitulis 3-5-floris pedunculis axillaribus folio longioribus suffultis racemos longos laxos formantibus, bracteis hirsutis ellipticis acuminatis calyces subæquantibus, sepalis hispidis a basi lanceolatâ subulato-acuminatis, corollâ hispidâ calyce subduplo longiore, ovario glabro 5.

Hab. in desertis Persiae australis (Auch. 4955!).

B. subaphyllus. — Rami abbreviati intricati ob folia ad squamulas axillares reducta subaphylli. — *C. Mascatensis* Boiss. Diagn. Ser. I, 7, p. 25.

Hab. in aridis regni Mascate Arabiae (Auch. 4938!).

Rami in typo pedem et amplius longi fere a basi racemosi foliis inferioribus 8-12 lineas longis basi lineam latis, in varietate divaricatum et dichotome ramo-siores, pedunculi polliceni vel sesquipollicem longi, corolla e sicco 5-6 lineas longa. Species cum præcedenti foliis minute auriculato-hastatis a sequentibus distincta.

9. **C. lanatus** (Vahl Symb. I, p. 16) fruticosus humilis ramosus, ramis rigidis erecto-patulis junioribus tomentosis vetustis denudatis spinescentibus, foliis tomento brevissimo cinerascentibus non rugosis inferioribus oblongo-spathulatis et linearis-spathulatis in petiolum attenuatis, superioribus sessilibus breviter lanceolatis acutiusculis, fasciculis dense capituliformibus 5-10-floris sessilibus ad axillas foliorum superiorum spicas interruptas formantibus, bracteis ovatis, sepalis hirsutissimis lanceolatis inæqualibus, corollâ sericeâ albo-roseâ calyce 2 $\frac{1}{2}$ -plo longiore, ovario glabro 5. *C. Cneorum* Forsk. Eg. p. 63 non L. — *C. Forskahlei* Del. Fl. Eg. p. 203, tab. 18. — Fl. Græc. tab. 202. — *C. sericeus* Choisy in D.C. Prodr. IX, p. 400 non Burm.

Hab. in desertis Arabicis Ægypti inferioris (Forsk., Del., Wiest 545! Boiss! etc.), in desertis Arabiae Petreæ (Auch. 1400! Bové! Schimp. 127!), in Creta (Sibth.), Arabice *Brehema*.

Frutex humilis pecoribus depastus, folia inferiora 1-2-pollicaria 3-4 lineas lata, floralia glomerulis florum breviora, corolla 7-8 lineas longa. Nullum plane discriminem inter *C. lanatum* Vahlii et *C. Forskahlei* Del. qui cæterum pro plantâ suâ Vahlii synonymon citat observavi; specimina ex Arabiâ petrea Ægyptiacis interdum paulo densus tomentosa sunt.

10. **C. ulicinus** (Boiss. Diagn. Ser. I, 7, p. 26) suffruticosus humili erinaceus totus brevissime tomentosus cinereus, ramis intricatis rigidis brevibus apice spinosis, foliis ad ramulorum partem inferiorem paucis parvis crassiunculis oblongo-vel linearis-spathulatis obtusis vel retusis basi longe attenuatis, fasciculis 3-4-floris ad ramulorum partem inferiorem paucis axillaribus sessilibus bracteis ovato-rotundis cœcum lati involucratis, sepalis rufescenti-hispidissimis triangulari-lanceolatis acutis, corollâ sericeâ quinquelobâ calyce duplo longiore, ovario glabro 5.

Hab. in regno Mascate Arabiæ (Auch. 4936!).

Suffrutex ut videtur humillimus ramis crassis, ramulis floriferis semipollinariibus crassis, folia majora 6-8 lineas cum petiolo longa apice 1-2 lata, superiore squamiformia minima, flores parvi e sicco 5-6 lineas longi. E grege *C. lanati*, multo humilior et spinosior, folia minuta, capitula paucissima, flores minores.

11. ***C. Scindicus*** (Stocks in Hook. Journ. Bot. 1852, p. 173 non Boiss. Diagn.) fruticosus humilis dichotome ramosissimus, ramis brevissime tomentellis intricatis subflexuosis remote foliosis tandem denudatis subspinescentibus, foliis parvis tomentosis ovato-spathulatis brevissime petiolatis secus costam et nervos plicatis obtusissimis, florum capitulis terminalibus, bracteis ovatis calyces aequalibus, sepalis hirsutissimis lanceolatis inaequalibus, corollâ sericeâ calycem parum excedente, ovario glabro 5. *C. brachyphyllus* Boiss. Diagn. Ser. II, 3, p. 122.

Hab. in Belutschia (Stocks exs. 433 sub *C. Forskahlei*?).

C. lanato ramosior ramis tenuioribus et magis flexuosis, folia majora cum petiolo 5-6 lineas tantum longa valde plicata nec plana, capitula avellanâ majora terminalia nec axillaria; calyx 4 lineas longus.

Ar. Geogr. Ditio Scinde.

12. ***C. oxysepalus*** (Boiss. Diagn. Ser. I, 7, p. 23) totus brevissime tomentosus cinereus suffruticosus a basi ramosissimus humili, ramis basi ramulosis ramulisque tenuibus rigidulis paucifoliis aliis sterilibus apice attenuatis aliis capitulo 5-6-floro terminatis, foliis minimis ovato- et oblongo-spathulatis obtusis inferioribus subpetiolatis, bracteis ovatis abrupte acuminatis calyce subæquantibus, sepalis longe hispidis a basi lanceolatâ longe setaceo-acuminatis, corollâ... 5.

Hab. in Persiâ australi inter Taroun et Darap (Auch. 4948!).

Suffrutex 4-5-pollicaris ramis subdivaricatis ramulosis ramulisque tenuissimis rigidulis 2-3 pollices tantum longis; plerique ramuli steriles persistunt sed apice attenuati subnudi potius quam spinescentes, pauci florum capitulo terminati, folia inferiora 2-3 lineas tantum longa superiora minima, calyx 5 lineas longus. Species ex specimine unio ex descripta bracteis latis acuminatis insignis.

13. ***C. eremophilus*** (Boiss. et Buhse Aufz. p. 148) breviter pubescens cinerascens suffruticosus ramosissimus erinaceus, ramis flexuosis ramulis rectangule patentibus rigidis apice dentatis subspinescentibus, foliis inferioribus parvis sessilibus breviter oblongo-lanceolatis acutis superioribus squamiformibus, cymis axillaribus longe pedunculatis trifloris vel abortu saepius 2-4-floris, pedicellis brevissimis incurvis, sepalis tomentellis elliptico-lanceolatis acuminatis, corollâ villosâ calyce 2 1/2-plo longiore, ovario glabro 5. *C. erinaceus* C A. Mey. Enum. p. 102 ex calycis descript. non ædebat.

Hab. in desertis lapidibsis salsis Persiae prope Kaschan (Buhse!), inter Kaschan et Nasrabad, inter Tabriz et Teheran et in ditione Khorassan, inter Massinon

et Mehr (Bge!), in arenâ mobili Transcaucasiae ad Caspium ad Eszek Karawanserai inter Baku et Sallian (C.A. Mey.).

Frutex subaphyllus facie et characteribus *C. erinacei* a quo praeter ovarium glabrum ramulis apice subspinescentibus, foliorum forma et præsertim sepalis elongatis acuminatis specifice differt.

14. *C. leptocladus* (Boiss. Diagn. Ser. I, 7, p. 25) glabriuscultus pilis adpressis brevibus sparsissime obsitus suffruticosus ramosissimus, ramis flexuosis ramulisque tenuissimis denum persistentibus, foliis ad squamulas breviter et anguste lineares reductis, cymis trifloris pedunculo filiformi rigido longo axillari rigidulo suffultis, pedicello floris centralis brevissimo lateralibus flore longioribus, sepalis glabriusculis scariosis ellipticis acuminatis, corollâ hirtâ calyce subtriplo longiore, ovario glabro ♂.

Hab. in Persiâ australi (Auch. 4942!).

Afinis *C. eremophilo* sed fere glaber, rami ramuli et pedunculi multo tenuiores, folia ad squamulas reducta, calyx sesquilineam longus.

15. *C. evolvuloides* (Boiss. Diagn. Ser. I, 7, p. 25) totus patule et brevissime tomentellus basi suffrutescens, ramis herbaceis erectis tenuiter virgatis inferne striete ramosis, foliis inferioribus ad rhizomatis collum anguste linearispathulatis obtusis longe basi attenuatis, rameis minimis ellipticis sessilibus, pedunculis strictis filiformibus squamâ axillari multo longioribus racemos laxos formantibus tri vel siue abortu 2-4-floris pedicellis brevissimis basi bracteolatis, sepalis adpresso hirsutis ellipticis acutis, corollâ hirsutâ calyce duplo longiore, ovario glabro ♂.

Hab. in prov. Mekran Persiae australis (Auch. 4952!), circa Jezd (ex Buhse).

Ex specimine nimis imperfecto descriptus. rami pedales et longiores parte inferiori divisi, in racemum laxum semipedalem abeuntes, basi foliis pollicariis attenuato-subpetiolatis apice lineam latius stipati, pedunculi 6-8 lineas longi, calyx sesquilineam longus. Facies *Evolvuli* sed stylus unicus. Afinis *C. chondrilloidi* a quo totius plante tomento, sepalis ellipticis acutis differt.

16. *C. Dorycnium* (L. Sp. 224) erinaceo-dunosus, ramis herbaceis induratis erectis elongatis inferne crassiusculis foliosis patule villosis superne dichotome et divaricatim ramentosissimis amplissime paniculatis adpresso hirsutis, foliis sessilibus inferioribus villosis obtusis oblongo-spathulatis et ellipticis, superioribus et floralibus ad dichotomias adpresso hirtis valde diminutis anguste linearibus acutis, cymis terminalibus trifloris pedunculo longo rigido suffultis abortu siue unifloris, flore ad basin pedicelli brevissimi bracteolato, sepalis adpresso sericeis obovato-rotundis obtusis vel retusis breviter mucronatis, corollâ roseâ extus ad angulos hirsutâ calyce multo longiore, ovario glabro ♀. Ic. Fl. Græc. tab. 201.

Hab. in aridis et ad vias regiomis inferioris, Peloponnesus ad Corinthum (Sibth.), Attica inter Athenas et Phalerum (Sprn! Heldr!). Creta (Sieb!). Fl. Jun

β. oxysepalus. — Sepala subangustiora in mucronem longiorem attenuata.

Hab. in Cypro (Ky!), Ciliciâ littorali (Bal!), Syriâ littorali ad Tripolin et Tyrum (Bl! Gaill!), Palestinâ (Auch. 1347!) et ad Tiberiadem (Boiss!), Antilibani vallibus 4000' (Gaill!) et ad Balbeck (Ehrenb!), Syriâ boreali ad Marasch et Aintab (Haussk!), Persiâ australi prope Schiraz (Ky. 436!) et boreali in ditione Khorassan (Bge!).

Caules e radice plures cum panicula multiflorâ pedali et ampliori 2-3-pedales, folia inferiora $1\frac{1}{2}$ -2-pollicaria 6-7 lineas lata, calyx 2- $2\frac{1}{2}$ lineas tantum longus, corolla magnit. *C. arvensis*. Varietas β formis intermediis ex Atticâ et Syriâ cum typo conjungitur. *C. dianthoides* Kar. et Kir. = *pseudocantabrica* Schrenck ex Songariâ valde affinis foliis inferioribus multo angustioribus, calyce glabro, ovario hirto differt.

§ 3. Inermes.

17. *C. sarothrocladus* (Boiss. et Haussk.), adpressissime hirtus pallide virens basi suffrutescens scoparius, ramis numerosis rigidis erectis elongatis a basi dichotome ramosissimis, foliis rigidulis inferioribus obovatis subspathulato-oblongis et elliptico-lanceolatis basi attenuatis subsessilibus obtusiusculis, superioribus anguste linearibus acutis, cymis terminalibus trifloris vel abortu 2-1-floris longiuscule pedunculatis, pedicellis calyce subbrevioribus basi bracteolatis, sepalis glaberrimis omnino scariosis ellipticis abruptiuscule mucronatis, corollâ albâ extus ad angulos sericeâ calyce subtriplo longiore, ovario superne hirto 5.

Hab. in arenosis Derbent i Basian et inter Kerkuk et Sulimanieh Assyriæ (Haussk!).

Rami sesquipedales fere a basi alterne et dichotome ramulosi, ramuli elongati erecto-patentes, folia inferiora $1\frac{1}{2}$ pollicaria $1\frac{1}{2}$ -pollicem lata, cymæ ramulos terminantes sœpius unifloræ, calyx basi attenuatus $2\frac{1}{4}$ lineas longus, corolla eâ *C. Dorycnii* minor. Species affinis *C. dianthoides* Kar. et Kir. foliis omnibus anguste linearibus corollâ roseâ distincto et *C. Dorycnio* basi vix suffrutescenti et ovario glabro donato.

18. *C. chondrilloides* (Boiss. Diagn. Ser. I, 11, p. 83) basi suffrutescens, ramis herbaceis numerosis glabris tenuiter virgatis erectis subaphyllis stricte ramosissimis ramulis filiformibus rigidulis elongatis, foliis inferioribus ad ramorum basin et partem inferiorem tomentellis breviter lanceolato-linearibus, rameis glabriusculis linearisubulatis, floribus ad ramulos terminalibus solitariis et lateralibus pedicellis brevissimis sub anthesi incurvis suffultis parvis, sepalis tomentellis ovatis vel ellipticis obtusis interdum obscure mucronulatis, corollâ albâ patente ad angulos sericeâ calyce triplo longiore, ovario superne hirto 5. *Evolvulus virgatus* Choisy v.c. Prodr. IX, p. 446.

Hab. in montibus Kurdistaniæ inter Bagdad et Kermanchah (Auch. 1410!), montibus Avroman et Schahu 5000' (Haussk!), Persiâ australi (Auch. 4941!), montibus prov. Susiana et Kuh Kilouyeh 9000' (Haussk!), monte Elbrus Persiæ bor. prope Derbend (Ky. 421!).

Caules et rami tenues stricti intricati, planta bipedalis facie *Chondrilloë* vel *Lini*, flores cymarum abortione saepè secus ramulos superiores laxe subspicati, calyx lineâ unâ vix longior, corolla eâ *Lini tenuifolii* minor. Stylus unicûs. A *C. Dorycnio* parte plantae inferiore suffruticosâ, ramis tenuibus strictis, floribus triplo minoribus distinctus.

19. C. Cneorum (L. Sp. 224 ex parte) adpressissime argenteo-sericeus fruticosus, ramis erectis inferne denudatis superne dense foliosis, foliis elongatis oblongo-lanceolatis et lanceolato-linearibus basi attenuatis apice latioribus acutiusculis, floribus fasciculato-capitatis pedunculo terminali folia suprema parum superante suffultis, bracteis linearibus calyces æquantibus vel superantibus, sepalis patule hirsutis oblongis breviter attenuato-acuminatis, corollâ albâ ad angulos extus sericeâ calyce triplo longiore, ovario apice hirto 5. Ic. Fl. Græc. tab. 200. — Bot. Mag. tab. 459. — *C. argenteus* Desr. Ency. III, p. 552.

Hab. in scopolis Samo vicinis (ex Sibth.).

Frutex saepè tripedalis ramis vetustis elongatis lignosis. Hanc e ditione non vidi sed icon Flora Grecæ speciem omnibus notam cum speciebus affinibus saepè confusam exacte exhibet Planta Cretica huc relata est *C. oleæfolius* Desr.

Ar. Geogr. Sicilia et Dalmatia littorales.

20. C. oleæfolius (Desr. Encycl. III, p. 552) adpressissime argyreus basi fruticosus, ramis herbaceis erectis elongatis totâ longitudine foliosis simplicibus vel stricte ramosis, foliis imis linear-i-spathulatis cæteris anguste linearibus obtusiusculis, floribus terminalibus laxe cymoso-subcapitatis, pedicellis calyce brevioribus, bracteis subulatis, sepalis hirsutissimis lanceolatis, corollâ roseâ ad angulos sericeâ calyce quadruplo longiore, ovario hirsuto 5. *C. Tournefortii* Sieb. exs. — *C. linearis* Bot. Mag. tab. 289 non D.C. — *C. lineatus* Fl. Græc. tab. 199 non L.

Hab. in collibus siccis regionis inferioris Græciæ (Auch. 1381), Atticæ (Sprunt! Heldr! Orph!), Eubeæ (Wild!), Cretæ (Sieb! Raul!), Archipelagi (Urv!), Rhodi (Bourg!), Cypri (Ky. 480!) Fl. Mai. Jun.

Affinis *C. Cneoro*, minus fruticosus, rami novelli herbacei $\frac{1}{2}$ -1 pedales vel etiam longiores, folia angustiora, inflorescentia laxior, sepala lanceolata nec oblonga abrupte et breviter mucronata. *C. saxatilis* Vahl varietas *sericea* = *C. lanuginosus* var. *argenteus* Choisy = *C. linearis* D.C. ex Hispaniâ et Gallia australi cum *C. oleæfolio* saepè confusus differt ramis tenuioribus, floribus dense capitatis bracteis triangulari-lanceolatis involucratis, sepalis angustius et longius acuminatis.

21. C. Cyprius, fruticosus humilis adpressissime sericeus, ramis annotinis tenuibus brevibus, foliis anguste lanceolato-linearibus acutiusculis basi attenuatis etiam superioribus elongatis, cymis subtri-floris capituliformibus terminalibus et ad ramulos breves axillares bracteis linearibus subrecurvis calyci subæquilongis involucratis, floribus breviter pedicellatis, calycis angusti adpressissime sericei

sepalis oblongis in caudam brevem linearem obtusam subcontractis, corollâ roseâ extus sericeâ calyce quadruplo longiore, ovario hirsuto 5. *C. lineatus* var. *angustifolius* Kotschy Cypr. p. 285.

Hab. in fissuris rupium regionis inferioris Cypri prope Lamuias ad promontorium Capo Gatto (Ky. 627!). Fl. Mai.

Rami hornotini 4-5-pollicares tenues crebre et stricte foliosi, vetusti exsiccati persistentes nec spinescentes, folia etiam superiora sesquipollucaria $1\frac{1}{2}$ -2 lineas lata, calyces $2\frac{1}{4}$ lineas longi. A formis stenophyllis *C. oleifolii* calycis induimento adpressissimo sepalisque minoribus caudato-contractis nec lanceolatis specifice distinctius videtur, ex paucis speciminibus tamen notus et ulterius observandus. Nulla cum *C. lineato* specie non frutescenti affinitas.

22. *C. Schirazianus* (Boiss. Diagn. Ser. I, 11, p. 82) adpressissime argyreо-sericeus suffruticosus, ramis novellis herbaceis tenuibus virgatis stricte ramulosis exsiccatis subpersistens, foliis longe et anguste linearibus acutis, capitulis terminalibus longiusculae pedunculatis, bracteis anguste linearи-lanceolatis, sepalis hispidis lanceolatis acuminatis, corollâ adpresso hirtâ calyce subtriplo longiore, ovario hirsuto 5.

Hab. in Persiaustrali (Auch. 4945!), in monte Sabst Buschom prope Schiraz (Ky. exs. 379! sub *C. saxatili* var.).

Rami pedales, folia inferiora $1\frac{1}{2}$ -2-pollicaria 1 $\frac{1}{2}$ -2 lin. lata, calyx $3\frac{1}{2}$ -4 lineas tantum longus. Affinis *C. Calverti* et *commutato* ab utroque ramis tenuibus elongatis virgatis subpersistens et transitum ad species subspinosas præbentibus diversus. Inter species subspinescentes *C. oxysepalus* foliis minutis brevibus, bracteis ovato-orbiculatis differt.

23. *C. Calverti* (Boiss. Diagn. Ser. II, 3, p. 124) adpresso sericeus et præterea patule hirsutus, caulis ex rhizomate suffrutescenti humilibus simplicibus erectis vel ascendentibus foliosis basi vestigiis foliorum vetustorum squamosis, foliis inferioribus anguste linearи-oblongis cæteris anguste linearibus acutiusculis, floribus præter 1-2 interdum ex axillis supremis ortis capitatis terminalibus subsessilibus bracteis linearibus calycem aequantibus suffultis, sepalis hispidissimis oblongis in cuspidem linearи-setaceam eis longiorem abeuntibus, corollâ roseâ ad angulos sericeâ calyce duplo longiore, ovario piloso 5. *C. saxatilis* M.B. I, p. 146 non Vahl. — *C. Cneorum* Habl. Taur. non L. — *C. pusillus* Pall. Index Taur. MSS. — *C. lanuginosus* Ledeb. Fl. Ross. III, p. 88 non Desr. — *C. triquetus* Rehm. exs.

Hab. in Taurie calcareis saxosis circa Karassabazar et Sympheropol (M.B., Stev!), in rupestribus ad Bairdar (Rehm!), Anatoliâ boreali ad Kastamouni (Wied. exs. 201!), Armeniâ Turcicâ ad Tortum (Calv!), ad Baibut (Bourg!).

Caules 3-6-pollicares, folia 1-2 lineas lata, flores parvi, calyx 3-4 lineas tantum longus. Species a *C. saxatili* Vahl = *lanuginoso* Desr. caulis humilibus, floribus minoribus, sepalis abrupte et longe caudatis discedens.

24. *C. commutatus* (Boiss. Diagn. Ser. I, 11, p. 81) adpressissime cinereo-canus, caulis ex rhizomate suffrutescenti pluribus

ascendentibus pumilis crassiuseulis stricte et breviter ramosis, foliis radicalibus spathulatis oblongo-lanceolatis in petiolum attenuatis, caulinis sessilibus elliptico-linearibus acutis utrinque attenuatis, floribus capitatis terminalibus sessilibus, bracteis oblongis acuminatis, sepalis hispidissimis a basi lanceolata longe subsetaceo-acuminatis, corollâ extus sericeâ calyce duplo longiore, ovario piloso 3. *C. lanuginosus* var. β Desr. Enc. p. 552. — *C. modestus* Boiss. loc. cit. p. 82.

Hab. in siccis Mesopotamiæ (Oliv. et Bourg!), Auch. 1411!), Kurdistaniæ Armenie inter Van et Ischri (Noë!), Persiae bor. prope Sisian (Buhse!), prov. Aderbidjan (Auch. 4947!), ad Seidabad prope Tabris et inter Sultanieh et Sengan (Bgel).

Caules semipedales, a *C. saxatili* foliis lato-ribus et sepalis longius setaceo-acuminatis discedens, magis affinis *C. Calverti* a quo tamen indumenti patuli defectu, caulinis crassioribus, foliis lato-ribus, floribus majoribus, sepalis 5-6 lineas longis sat differre videtur.

25. ***C. holosericeus*** (M.B. Taur. Cauc. I, p. 147) totus adpresso-simile sericeo-nitens basi suffrutescens, caulinis ascendentibus vel procumbentibus tenuiter virgatis, foliis inferioribus lanceolatis in petiolum attenuatis cæteris linear-lanceolatis et anguste linearibus acutis sessilibus, floribus in axillis superioribus subsolitariis breviter vel longiuscula pedunculatis laxe racemosis terminalibus 1-3 approximatis, pedicellis calyce subbrevioribus, bracteis subulatis, sepalis adpresso hirtis ovatis breviter acuminatis externis parte inferiori inflato-gibbis, corollâ lacteâ extus holosericeâ calyce quadruplo longiore, capsulâ hirtâ 5. Ic. M.B. Cent. Ross. tab. 24.

Hab. in collibus siccis Anatoliæ bor. ad Kadikeui (Wied!). Phrygiæ ad Ouchak (Bal!), Lydiæ ad radices Mesogis prope Derbent (Boiss!), Lyciæ ad Elmalu (Bourg. 2411!), Pisidiæ (Heldr!), Isauriæ (Péron!), Ponti (Tchih!), Capadociæ (Auch. 1409!), Syriæ bor. ad Aintab et Mesopotamiæ conterminæ (Haussk!), Armenia ad Baibut (Bourg!), Tauriæ (Stev!).

Caules semipedales rarius pedales, flores magn. *C. arvensis*. Folia variant valde angusta. Species calyce (ut in *Helianthemi* speciebus nonnullis) inflato insignis.

26. ***C. Cantabrica*** (L. Sp. 225) caulinis herbaceis e rhizomate indurato decumbentibus vel erectis elongatis fere a basi ramosis patule vel adpresso hirsutis, foliis viridibus utrinque hirtis inferioribus oblongo-lanceolatis in petiolum attenuatis superioribus sessilibus lanceolatis et linearibus, cymis trifloris vel abortu unifloris longe pedunculatis terminalibus paniculam laxam patulam plus minusve multifloram formantibus, pedicellis calyce brevioribus, bracteis linear-subulatis, sepalis hirsutis lanceolatis acutis, corollâ roseâ extus lineatim hirsutâ calyce duplo longiore, capsulâ hirtâ 2. Ic. Jacq. Austr. tab. 296. — Rchb. tab. 135.

Hab. in collibus siccis Peloponnesi (Bory), Atticae (Boiss! Orph! Heldr!), Cretæ (Noë!), Macedoniae et Thraciæ (Griseb., Thirke!). Anatoliæ borealis

(Wied!), occidental is et australis (Pinard! Ky!), Syriæ littoralis (Boiss!), Libani (Auch. 1390! Bl!), Tauriæ (M.B., Rehml!), Transcaucasiæ totius (Hoh! C.A. Mey!, Szov!), Persiæ borealis (Buhse! Bge!). Fl. vere.

Variat indumento patulo vel adpresso, sepalis plus minus elongatis sed semper acutis vel acuminatis.

Ar. Geogr. Europa australis a Galliâ mediâ, Austriâ australi ad regionem Danubialem, Africa borealis.

27. *C. cardiosepalus*, totus pilis adpressis et patulis demum rufescentibus hispidus, caulis herbaceis ascendentibus a parte mediâ ramulosis elongatis, foliis inferioribus oblongo-spathulatis in petiolum attenuatis superioribus breviter linearibus acutis, floribus 3-5 ad ramos elongatos nudos erecto-patentes terminalibus cymosocapitatis subsessilibus parvis paniculam laxam formantibus, sepalis binis externis breviter triangularibus acutis, internis obcordatis mucronatis præter mucronem herbaceum scariosis, corollâ extus lineatim hirsutâ patenti calyce quadruplo longiore, capsulâ hirtâ ♀.

Hab. in calcareis Cilicie littoralis prope Boulokli (Bal. exs. sub *C. Cantabrica!*). Fl. Mai.

Pedalis, folia et inflorescentia *C. Cantabrici* pro cuius formâ hunc habui, sed præter cymas ob flores subsessiles capitatas distinctus calyce multo breviore vix 2 lineas longo, sepalis internis late obcordatis; corollæ 6-7 lineas longæ latæque color e sicco incertus.

28. *C. Aucheri* (Choisy in D.C. Prodr. IX, p. 402) totus pilis brevibus adpressiusculis aliisque longioribus patulis tomentoso-villosus, caulis e rhizomate indurato elatis fere a basi dichotome et patule ramosissimis, foliis inferioribus ovatis superioribus sessilibus lanceolatis acutis, floralibus brevibus oblongo-lanceolatis, cymis 3 vel abortu saepius unifloris ad ramulos terminalibus paniculam amplam laxam formantibus, pedicellis calyce vix brevioribus, sepalis villosis oblongis abrupte in cuspidem eis subbreviorem abeuntibus, corollâ extus hirsutâ angustâ calyce quadruplo longiore, ovario hirsuto ♀.

Hab. in Syriâ boreali prope Aintab (Auch. 1405!).

Caules saepe bipedales, folia intermedia sesquipollucaria 4 lineas lata, calyx 2 lineas, corolla (e sicco alba?) 9 lin. longa. Ex paucis et imperfectis speciminiibus notus, valde affinis *C. Cantabrico* a quo tamen distinctus videtur caule proceriore multo ramosiore, foliis etiam supremis latioribus, sepalis abruptius mucronatis, præsertim corollâ longiore angustiore tubuloso-attenuata.

29. *C. Ruprechti*, adpressissime argyreus, rhizomatis suffrutescentis ramis brevibus crassis, caulis floriferis erectis vel ascendentibus pumilis, foliis lineatis subsessilibus basi attenuatis inferioribus et surculorum sterili um ovatis vel obovato-oblongis, superioribus diminutis oblongis vel oblongo-linearibus acutis, floribus (præter 1-2 interdum ex axillâ superiori) 3-5 terminalibus subsessilibus capitatis bracteis elliptico-oblongis involucratis, sepalis longe hirsutissimis

oblongis attenuato-acuminatis, corollâ roseâ extus hirsutâ calyce $2\frac{1}{2}$ -plo longiore, ovario hirto ♂.

Hab. in rupestribus Caucasi orientalis, ditio Gumbet inter Danuch et Arguani 4800'-5000' et Daghestania prope Kutuschi 4500' (Rupr!). Fl. Jun.

Caules 3-4-pollicares, folia inferiora 5-6 pollices longa tres lata, sepala $3\frac{1}{2}$ lineas longa. Aflinis *C. Calverti* differt foliis linearatis ovatis et oblongo-spatulatis nec linearibus, indumento adpressissime argyreo. *C. convolutatus* differt ramis floriferis longioribus crassioribus, foliis acutis; *C. lineatus* rhizomate non suffruticoso longius distat.

30. ***C. lineatus*** (L. Sp. 224) adpresso sericeus, caulis herbaceis et rhizomate pluricipiti ascendenteribus vel procumbentibus pumilis simplicibus vel ramulosis interdum brevissimis, foliis oblongis inferioribus in petiolum longe attenuatis superioribus saepe angustatis linearibus, cymis ad ramulorum apicem 3 vel 2-1-floris subsecundis terminalibusque, bracteis linearibus basi attenuatis calyces superantibus, floribus breviter pedicellatis, sepalis oblongo-lanceolatis basi membranaceis apice patulo herbaceis, corollâ roseâ calyre quadruplo longiore, ovario villoso ♀. Ie. Rehb. tab. 134. — *C. spicafolius* Desr. Enc. III p. 549 — *C. intermedius* Lois. — *C. Besseri* Spreng. Syst. I, p. 610. — *C. nitens* C. Koch Linn. XXIII, p. 595.

Hab ad vias et in collibus siccis regionis inferioris Anatoliae occid. a Bithynia ad Ciliciam et interioris in Cappadociâ (Bal!), Syrie ad Libani radices (Auch. 441!), Antilibano ad Rascheya (Boiss!), deserto ad Palmyram (Bil!), Mesopotamiâ (Haussk!), Armeniâ Turcicâ (Huet! Bourg!), Tauriâ (M. B., Stev!), prov. Cis et Transcaucasicis (M. B., Hoh! etc.) et Caucaso orientali in Daghestaniâ ad 4800' usque (Rupr!), Persia boreali (Auch. 4946!), mediâ (Bge!) et australi ad Schiraz (Ky. 371!), Afghaniâ (Griff!), Belutschia (Stocks!), Ägypto ad Alexandriam (Ehrenb!). Fl. Äst.

Ar Geogr. Europa australis a Gallia mediterraneâ et Hispania ad Italiam, Rossia media et australis, Songaria et Sibiria Altaica, Africa borealis.

31. ***C. Libanoticus*** (Boiss. Diagn. Ser. I, 11, p. 82) cespitosus adpresso canescens-griseus rhizomatis lignosi ramis crassis denudatis prostratis, caulis decumbentibus herbaceis brevibus simplicibus vel ramulo auctis, foliis radicalibus oblongis vel oblongo-linearibus obtusis basi attenuatis caulinis linearibus acutis, floribus 3-5 laxiuscula cymosis vel solitariis pedicellis calyce aequilongis suffultis, sepalis hirtis oblongis apice herbaceis acutinseculis, corollâ carneâ extus hirtâ calyre quadruplo longiore, capsulâ apice pilosâ ♀. *C. cespitosus* Ehrenb. MSS.

Hab. in siccis regionis alpinæ Libani supra Cedros (Boiss!, Auch. 1399!), inter Zachle et montem Sannin (Ehrenb!), inter Dimam et Yamouny (Bil!), Antilibano in cacumine Hermonis (Boiss! Gaill!), in montibus Zebdani (Ky. 541).

Folia radicalia 4-6 lineas longa apice $1\frac{1}{2}$ -2 lata, caules longiores 2-3-pollicares, sepala vix 2 lineas longa. A *C. compacto* et *cochleari* indumento non sericeo-argyreo, sepalis non caudatis discedens; hic sepalorum forma accedens ad *C. lineatum* rhizomate non lignoso foliis-que ceterum diversissimum.

32. C. Cataonicus (Boiss. et Haussk.) adpresso hirtus pallide virens cespitosus. rhizomate valde fibrilloso breviter ramoso, caulinis floriferis brevibus ascendentibus vel procumbentibus simplicibus vel ramulosis, foliis facie superiori glabris anguste linearis-spathulatis basi longe attenuatis inferioribus obtusis superioribus linearibus acutiusculis saepe recurvis, floribus 1-4 sessilibus ad ramulos terminalibus capitatis, bracteis anguste linearibus calycem æquantibus, sepalis hirsutis inferne pallidis oblongis in caudam herbaceam subaequilongam contractis, corollâ albâ extus hirsutâ calyce triplo longiore, ovario hirto ♀.

Hab. in agris regionis inferioris montis Berytdagh Cataonie 5000' (Haussk!).

Caules floriferi 4-6-pollicares, folia surculorum cespitis et caulina inferiora conferta 4-1½-pollicaria apice 2-2½ lata, superiora angustiora, sepalâ tres lineas longa. Affinis *C. Libanotico* a quo caulinis elongatis sepalisque caudatis differt. a *C. holosericei* formis pumilis indumento non sericeo calyce que non inflato facile distinguendus.

§ 4. Compacti.

33. C. compactus (Boiss. Diagn. Ser. I, 4, p. 40) dense pulvinatus adpressissime argyreo-canus, rhizomatis ramis denudatis, caulinis nullis vel brevissimis. floribus e cespite non exsertis apice cymose 3-5-floris, foliis confertis anguste linearibus vel linearis-spathulatis acutis plicatis. floralibus saepe spathulato-dilatatis. sepalis hirsutissimis oblongis longe caudato-acuminatis, corollâ roseâ ad angulos sericeâ calyce subtriplo longiore, capsulâ hirsutissimâ ♀.

Hab. in montanis et subalpinis siccis Anatoliae, mons Cadmus Cariæ supra Gheyla (Boiss!), colles tophacei prope Bouldour Pisidiae et prope Koniah (Heldr!), ad Kadikiöi (Wied!), Cappadocia in montibus Maßmeneudagh et Aslandach (Ball!). Fl. Ang.

Pulvini densissimi, flores in cespite sessiles vel rarius caules ½-1-pollicares cynam capituliformem ferentes, folia 5-9 lineas longa apice sesquilineam rarius lineam lata.

34. C. cochlearis (Grisebach Spicil. II, p. 76 in notâ) acaulis dense pulvinatus adpresso argyreo-sericeus, rhizomate erasso lignoso breviter ramoso, foliis minimis ovatis cuneatis acutiusculis longiuscule petiolatis paginâ superiori lineatis et secus nervum plicatis, floribus solitariis in cespite subsessilibus, sepalis hirsutissimis ovatis plus minus longe caudato-acuminatis, corollâ purpureâ extus lineatum sericeâ calyce triplo longiore, ovario hirsuto ♀. *C. Parvassicus* Boiss. et Orph. Diagn. II, 3, p. 125.

Hab. in aridis montanis et subalpinis, Parnassus regio media ad Hagios Nicolaos et supra Carcaria 4500'-5000' (Guice! Oph. Fl. Gr. exs. 480!), Phrygia circa Onchak 3000' (Bal!), Cataonia inter Aintab et Marasch et in monte Berytdagh 7000' (Haussk!), mons Amanus Syrie bor. in subalpinis Tolos 4000' (Ky. 4391 sub *C. compacto*).

Foliorum majorum limbus vix tres lineas longus et latus, flores eis *C. lineati*

et *C. nitidi* submnores Huic et *C. compacto* affinis, foliis distinctus, an tamen hujus varietas?

35. **C. Assyricus** (Griseb. Spicil. II, p. 75 in notâ) dense pulvinatus compactus acaulis totus pilis nitidis rigidis erecto-patulis strigulosus, rhizomatis lignosi ramis tortuosis denudatis, foliis dense cespitosis anguste linearibus elongatis plicatis inferne attenuatis apice interdum spathulato-subdilatatis, floralibus calycent longe superantibus longe pectinato-ciliatis, sepalis breviter ovatis muticis, corollâ roseâ extus valde hirsutâ tubuloso-attenuatâ calyce sextuplo longiore, ovario hirto ♀. *C. strigulosus* Boiss. Diagn. Ser. I, 11, p. 83.

Hab. in montanis Anatoliae inter Angora et Tokat (Auch 4939!), in Cappadocie monte Aslandagh (Bal. 973!), ad radices Argæi 5000' Ky. 179!, in Mesopotamia inter Bir et Surug (Hanskt!).

Folia 5-9 lineas longa, facies *C. compacti* a quo indumento striguloso et præsertim sepalis linea vix longioribus muticis nec caudatis corollaque elongata distinguitur.

36. **C. radicosus** (Heldr. et Sart. in Boiss. Diagn. Ser. II, 3, p. 124) caulis e rhizomate lignoso ramoso pumilis cespitosis dense foliosis, foliis parvis subsessilibus oblongo-spathulatis supra glabris subtus albo-strigosis, cymis trifloris brevissime pedunculatis, pedicellis strigoso-hirtis calyce multoties brevioribus, bracteis folio conformibus, sepalis sericeo-strigosis ovato-oblongis apice patentibus foliaceis acutiusculis, corollâ rubicundâ intus maculis 5 atropurpureis pictâ extus lineatim sericeo-strigosâ calyce duplo longiore ♀.

Hab. in pascuis siccis Livadi montis Kyllenes Peloponnesi supra Trikala 5000' (Heldr. spec. unicum). Non vidi.

Rhizoma pennam anserinam crassum, folia majora 4-5 lineas longa 1-1 1/2 lata, sepalâ 3 lineas longa denum aucta, corolla semipollicaris. Ulterius investigandus. ex descriptione affinis *C. cochlearia*, ab eo et affinibus foliis pagina superiori glabris discedens.

§ 5. Pannosi.

37. **C. Persicus** (Linn. Am. Acad. II, p. 341) dense albo-pannosus, caulis e rhizomate duro ramoso decumbenti-ascendentibus subsimplicibus inferne nudis dein crebre foliosis tomentosis crassis, foliis majuseulis crassis non rugosis brevissime petiolatis obtusis inferioribus ovatis superioribus ellipticis, pedunculis axillaribus unifloris infra calycent bibracteatis folio subaequilongis, sepalis æqualibus ovatis sericeo-tomentosis obtusis obsolete mucronulatis, corollâ amplâ albâ extus sublanatâ breviter late et obtuse quinquelobâ calyce triplo longiore, ovario glabro ♀. Ic. Gmel. It. III, t. 7.

Hab. in littoribus arenosis maritimis, Byzantii (Sibth.), prope Kila ad Euxinum (Commany!), Abchasia et Mingrelia (Nordm.), ad Poti (Rehm!), in littoribus Caspiis prope Derbent, Baku, Lenkoran (M. B., C. A. M., Hoh!). in littoribus

prov. Ghilan Persiae (Auch. 4940! Buhse!), Turcomaniâ littorali (herb. Petrop!). Fl Jun. Aug.

Caules parte foliosâ 6-10-pollicares, folia inferiora 4-4 $\frac{1}{2}$ poll. longa, corolla diametro sesquipollicaris et amplior. In Sibiriâ erroneâ indicatus.

38. **C. secundus** (Desr. Encycl. III, p. 553 non R. P.) adpresse sericeo-tomentosus demum (in herbarii specim. vetustis) ferrugineus, caulis e rhizomate duro ramoso prostratis elongatis simplicibus vel inferne parce ramosis crassis tomentosis crebre foliosis, foliis secundis crassis nervoso-rugosis inferioribus lanceolatis in petiolum brevem attenuatis cæteris anguste oblongis obtusis sessilibus, floribus 2-5 in fasciculos brevissime pedunculatos vel subsessiles axillares unilaterales dispositis, bracteis et sepalis breviter lanceolatis subæqualibus hirsutissimis, corollâ albâ extus villosâ calyce 2 $\frac{1}{2}$ -polo longiore breviter quinquelobâ, ovario glabro ♀. *C. salviefolius* Sieb. exs!

Hab. in littoribus arenosis maritimis Ægypti ad Suez (Ky. 890!), Palestinæ ad Gaza (Boiss!), Kaifa (Schlumberger!), Joppæ (Sieb.), Ky. 652!. Syriâ littorali (Auch. 4398!), ad Sidonem (Pestal!, Bl!) Fl. Apr. Jul.

Caules virgati crassi pedales et longiores fere totâ longitudine laxiuseule floriferi, folia inferiora cum petiolo 12-16 lin. longa 3-4 lata, corolla fere pollicem longa

39. **C. reticulatus** (Choisy in D.C. Prodr. IX, p. 399) adpresse tomentosus demum rufescens, caulis e radice durâ longâ prostratis vel ascendentibus longis crassis tomentosis simplicibus vel inferne parce ramosis, foliis crassis reticulato-rugosis infimis et turionum steriliū oblongo-lanceolatis in petiolum brevem attenuatis cæteris sessilibus ovatis cordatis acutiusculis, superioribus valde diminutis, floribus 2-5 in fasciculos axillares breviter pedunculatos vel sessiles dispositis, bracteis sepalisque subæqualibus hirsutissimis lanceolatis acutis, corollâ albâ extus hirsutâ breviter lobatâ calyce duplo longiore, ovario glabro ♀. *C. ascendens* Steud. in Kotschy exs.

Hab. in cretaceis aridis Syriae borealis ad Aintab (Auch. 1407!, Haussk!), Mesopotamia ad Euphratem inter Malatia et Karput et ad Derbent i Basian (Haussk!), ad Bauerd et inter Mardin et Assuaner (Ky. 338! et 410!), ad Amadiam Kurdistanæ Assyriace (Ky. 348!). Fl Jun. Jul.

Caules 1-2-pedales longe et laxe floriferi, folia caulinâ inferiora 1-2 pollices longa 12-15 lineas lata, turionum multo angustiora, corolla semipollicem longa.

40. **C. waltherioides** (Boiss. et Haussk.) totus adpresse tomentosus demum rufescens, caulis e radice durâ ascendentibus crassis parte inferiori sœpius ramulosus, foliis crassis superne lœviuseculis subtus reticulato-rugosis inferioribus ovatis in petiolum longiusculum attenuatis, caulinis subsessilibus et sessilibus ovato-suborbiculatis, floribus 5-8 dense fasciculato-capitatis pedunculo axillari crasso folio æquilongo vel longiore suffultis, bracteis oblongis, sepalis breviter lanceolatis acutiusculis subinæqualibus hirsutissimis, corollâ albâ

extus hirsutâ breviter lobatâ calyce subduplo longiore, ovario glabro ♀.

Hab. in graminosis Teng Tokab et Teng Biresa Persie austro-occidentalis 3-4000' et in rupibus gypsaceis ad Bebehān (Haussk!). Fl. Jun. Jul.

Caulis pedales, folia caulinata intermedia pollicem longa lataque. Affinis *C. reticulato* sed racemi multo breviores, pedunculi longiores, sepala obtusiora, corolla paulo minor angustior, folia pagina superiore non rugosa basi non vel obscure cor-lata.

41. **C. sericeus** Burm. Ind. 49, tab. 19, f. 3 et herb! non L. et excl. synon. Prodri. Cand.) totus breviter tomentosus, caulis decumbentibus vel erectis simplicibus, foliis minutis oblongo-spathulatis securis eostam et nervos eximie plicato-rugosis margine suberispis, inferioribus brevissime petiolatis cæteris sessilibus, floralibus planisculis, capitulis globosis pedunculis axillaribus folio longioribus et æquilongis suffultis racemos iaxos terminales formantibus, bracteis oblongis acutis calyces aequalibus, sepalis hirsutissimis lanceolatis acutis, corollâ extus tomentosâ calyce duplo longiore, ovario glabro ♀. *C. Persicus sericeus Sideritidis folio sinuato* (herb. Garcin!). — *C. cephalopodus* Boiss. Diagn. Ser. I, 7, p. 24.

Hab. in desertis provincie Mekran Persie australis (Auch. 4949!, Garcin!).

Caulis ex rhizomate duro 6-8-pollicares ut videtur decumbentes, folia inferiore 7-9 lineas longa, racemus semipedalis 6-12-cephalas, capitula 3-5-flora avellanae magnitudine, calyx vix 3 lineas longus. Haec species a cl. Choisy in **C. lanato** ramis rigidis crassis denum spinoscentibus diversissimo confusa fuit Planta herb. Burmanniani *C. cephalopodo* meo simillima est, sed ejus icon flores solitarios erroneè exhibet.

42. **C. Schimperi** (Boiss. Diagn. Ser. I, 11, p. 81) caulis prostratis? inferne ramosis ramisque longe et patule tomentoso-hispidis, foliis dense tomentoso-hispidis denum rufis oblongo-spathulatis basi attenuatis obtusis valde plicato-rugosis anargine undulato-crispis, capitulis 3-5-floris axillaribus pedunculis folio axillari sèpius brevioribus suffultis breviter racemosis, bracteis sepaliisque a basi lanceolatâ longe linear-i-caudatis hispidissimis, corollâ angustâ pilosâ calycem breviter excedente, ovario glabro ♀.

Hab. in Arabiâ subtropicâ (Schimp. pl. Arab. sub *C. salviifolio*!).

Foliis rugoso-plicatis et inflorescentiâ affinis *C. sericeo*, tomentum densius longius, folia majora, inferiora 1-1 1/2-pollicaria 4 lineas lata, pedunculi breviores, in primis sepala longe caudata 5 lineas longa. Species vix nostræ ditionis ad comparationem cum *C. sericeo* admissa.

43. **C. Kotschyanius** (Boiss. Diagn. Ser. I, 7, p. 23) caulis ex rhizomate duro pluricipiti erectis vel ascendentibus simplicibus præter indumentum breve glandulosum patule et longe hispidis, foliis hirsutis tenuibus radicalibus plus minusve rugoso-plicatis anguste oblongo-linearibus linearibusque obtusis basi longe attenuatis, caulinis sessilibus saepe latioribus sèpius planis, fasciculis 3-4-floris inferio-

ribus brevissime pedunculatis cæteris sessilibus spicas longas interruptas formantibus, sepalis subinaequalibus lanceolato-acuminatis, corollâ roseâ ad angulos hirtâ calyce triplo longiore, ovario glabro ♀.

Hab. in desertis et arenosis Persie australis ad Belutschiae fines (Auch 4944!), in collibus prope Schiraz (Ky. 357!), inter Seytun et Bebehān (Haussk!) Specimen Noé exs. 4177 ex Karput Armeniæ australis non differre videtur. Fl. Mai. Jun.

Caules $\frac{1}{2}$ -4-pedales, folia radicalia 1-2-pollicaria 2-5 lin. lata, racemi pedem dimidium et amplius longi, corolla magn. *C. arvensis*. A *C. Schimperi* caulibus simplicibus, foliis angustis substantiae tenuis, corollâ calycem plures superante distinctus.

44. ***C. Haussknechtii***, indumento dupli altero brevissimo viscidulo altero pilis longis patulis constante hispidum, caulis, e rhizomate pluribus ascendentibus simplicissimis, foliis omnibus membranaceis non plicatis inferioribus oblongis obtusis in petiolam longe attenuatis margine interdum obscure undulato-repandis, floralibus sessilibus triangulari-lanceolatis acuminatis, fasciculis 2-3-floris sessilibus axillaribus spicas longas interruptas formantibus, sepalis plumoso-hispidae subinaequalibus anguste lanceolatis acuminatis vel acutis, corollâ roseâ extus hirsutâ calyce subtriplo longiore, ovario glabro ♀. *C. gonoclados* Boiss. Diagn. Ser. I, 7, p. 22. — *C. pyrrhotrichus* Boiss. Diagn. Ser. II, 3, p. 122.

Hab. in graminosis gypsaceis inter Dalechi et Kasrun Persie australis (Ky. 27!, Haussk!), in regno Cabulico ad Khuggur (Griff 5879!) Fl. Apr.

Caules fere pedales, folia inferiora cum petiolo æquilongo tripollicaria 4-5 lineas lata, spica valde interrupte 6-9 pollicares. A *C. Kotschyano* valde affini differt foliis planissimis nec rugoso-plicatis, caulinis latioribus; an tamen hujus varietas? Nomen specificum mutavi nam ex speciminum copiâ majore caules exceptione tantum angulatos esse compcri.

45. ***C. cephalophorus*** (Boiss. Diagn. Ser. I, 7, p. 22) adpressissime et brevissime tomentellus, caulis humilibus erectis simplicissimis, foliis non rugosis dissimilibus radicalibus et inferioribus in petiolam longe attenuatis anguste lanceolatis, superioribus a basi sessili late ovatis vel oblongis breviter acuminatis, fasciculis 3-4-floris ad folia superiora breviter pedunculatis et sessilibus racemum oblongum inferne interruptum superne densum formantibus, bracteis sepalisque longe linear-setaceis plumosis, corollâ albâ extus rosellâ et hispidâ calyce duplo longiore breviter lobata, ovario glabro ♀.

Hab. in Persie austro-occid. monte Kotele Malo prope Dalechi (Ky. 138!), inter Dalechi et Kaserun 1-3'000' (Haussk!). Fl. Apr.

Semipedalis, facies *Phlomidis*, indumento brevissimo cinereo obductus, folia inferiora 2-3-pollicaria 4-6 lineas lata, caulinâ 1-1 $\frac{1}{2}$ pollicaria 9-12 lin. lata, sepalâ tenuissima 6-7 lineas longa.

§ 6. Diffusi.

46. ***C. glomeratus*** (Choisy n.c. Prodr. IX, p. 401) caulis c.

radice verticali pluribus prostratis tenuissime virgatis longissimis simplicibus, foliis viridibus subtus adpresse hirtis parvis brevissime petiolatis minute et obtuse cordato-hastatis inferioribus ovalis cæteris breviter oblongis acutis, florum capitulis globosis pedunculis axillaribus folio longioribus suffultis longe et laxissime racemosis, bracteis foliaceis acuminatis calyce longioribus, sepalis villosissimis oblongis longe acuminatis, corollâ albâ glabrâ calycem vix accedente, ovario glabro ♀. Wight Icon. tab. 1366. — *C. capitatus* Schimp. exs. non Vahl.

Hab. in Belutschia calidore (Frere ex Aitch. Cat.).

Ar. Geogr. Abyssinia, Arabia subtropica, ditio Scinde et India bor.-occidentalis.

47. *C. pilosellæfolius* (Desr. Encycl. III, p. 551) caulis tenuibus virgatis ascendentibus vel prostratis valde elongatis adpresse hirtis a medio vel a parte inferiore ramulos floriferos clementibus, foliis pallide virentibus utrinque sparsim et adpressiuseule vel patule pilosis margine saepe undulato-repandis inferioribus oblongo-lanceolatis in petiolatum attenuatis, caulinis sessilibus abbreviatis acutis breviter lanceolatis et triangularibus basi interdum subcordatis, floribus ad ramulorum apicem 5-1 laxe cymulosis pedicellatis parvis racemum longum laxum formantibus. sepalis adpresse et sparsim hirtis ellipticis acutiusculis, corollâ roseâ calyce triplo longiore ad angulos hirtâ, capsulâ oblongâ glabrâ ♀. *C. Sogdianus* Bge. pl. Lehm. p. 395.

Hab. in incultis, ad vias, desertis Mesopotamiae et Assyria (Oliv. et Brugt Auch. 1412! Ky. 405!), prope Kerkuk, Mossul et in monte Sindjar (Hausk!), Persia austro-orientali in palmetis Khabbise, orientali ad Schurab et in ditione Khorassan (Bge!), Armeniâ Rossicâ et Persiâ boreali (Szov. 409! et 607!). Turkestania inter Buchara et Samarkand (Lehm!), Affghania (Griff!).

Planta perennis nec annua, caules tenues saepe bipedales et longiores, flores dimidio minores eis *C. Cantabrica* cuius formis parvifloris facie affinis sed sepalâ omniuino diversa, capsula glabra, etc.

48. *C. microphyllus* (Sieb. exs. — Visian. Ill. Eg. p. 12) caulis ex radice verticali prostratis tenuiter virgatis elongatis simplicibus villosis, foliis hirsutis linearis-lingulatis basi attenuatis obtusis infimis subpetiolatis cæteris sessilibus minutis, pedunculis axillaribus brevissimis vel subnullis 1-2-floris, floribus parvis, sepalis villosis oblongo-lanceolatis breviter attenuato-acuminatis, corollâ carneâ extus ad angulos hirsutâ calyce duplo longiore, ovario glabro ♀.

Hab. in Ægypto ad deserti limites prope Achmin (Sieb!), ad Assouan (Kralik!).

Caules tenues $\frac{1}{2}$ -1-pedales, folia inferiora 8-10 lin. longa $2\frac{1}{2}$ lata, flores 4-5 lineas tantum longi. *C. deserti* Hochst. et St. ex Arabiâ tropicâ huc a cl. Choisy citatus specifice dilert indumento parco, floribus 5-7 cymosis pedunculis folio axillari multo longioribus suffultis, sepalis longe acuminatis. *C. Scindicus* Boiss. Diagn. Ser. II, 3, p. 123 non Stocks e ditione Scinde Indiæ est forsitan

tantum *C. microphylli* varietas foliis radicalibus amplioribus oblongis, floribus majoribus.

Ar. Geogr. Nubia.

49. C. Ammocharis (Boiss. et Haussk.) viridis pilis adpressis sparsis hirtus, caulinibus ex rhizomate tenuiter ramoso procumbentibus vel ascendentibus tenuibus, foliis inferioribus linear-i-oblongis basi attenuatis superioribus linear-i-lanceolatis et linearibus abbreviatis, pedunculis axillaribus filiformibus folio longioribus rigidulis 2-4-floris, floribus parvis brevissime pedicellatis bracteolatis, sepalis subscariosis hirtis ciliatulis ellipticis acutis, corollâ roseâ ad angulos hirtâ calyce duplo longiore, ovario glabro ♀.

Hab. in arenosis Persiae australis prope Abuchir (Haussk !). Fl. Mai.

Caules 6-10-pollicares fere a basi ramulos floriferos 4-1½-pollicares ex axillis edentes, folia infraiora 9-10 lineas longa 2 lata, calyx sesquilineam corolla 3½ lineas longa. Planta pusilla elegans.

§ 7. Volubiles.

50. C. Galaticus (Rostan in D.C. Prodr. IX, p. 408) totus dense tomentosus subsericeus. caulinibus procumbentibus flexuosis, foliis plus minusve ruguloso-plicatis et margine sinuatis insimis ovatis basi cuneatis petiolo subæquilongo suffultis, cæteris breviter petiolatis cordatis obtuse auriculatis ovatis saepe obscure et obtusissime 3-5-lobis, summis sensim diminutis acutis, pedunculis axillaribus crassiulus folio æquilongis vel longioribus 3-4-floris, pedicellis calyci æquilongis vel longioribus, bracteis brevibus setaceis, sepalis tomentosis ovato-ellipticis breviter acuminatis, corollâ purpureâ extus ad plicas hirsutâ calyce quadruplo longiore, ovario hirsuto, seminibus elevatim et minute subreticulato-rugulosis ♀. *C. agrophilus* C. Koch Linn. XXIII, p. 745!

Hab. in graminosis collium, agris argillosose in cultis Bithyniæ ad Bolu (Pest!), Lyctiæ ad Elmali (Bourg!), Galatiæ ad Angora (Rostan! Wiedem. 1981), Ponti australis (Tchih!), Armeniæ ad Baibut (Bourg!), Cappadociæ ad Bereketly et in regione inferiori montis Masmeneudagh (Ball!), Syriae borealis ad Aintab (Auch. 1406! Haussk!), Mesopotamiæ ad Oïfa et Suerek (Ky. 73!).

Caules ½-1-pedales interdum sublongiores non volubiles nec scandentes, folia majora circ. pollicem longa, corolla eâ *C. arvensis* submajor; species elegans habitu malvaceo.

51. C. Germaniciæ (Boiss. et Haussk.) patule tomentellus cinereus, caulinibus diffusis flexuosis tenuibus elongatis, foliis breviter petiolatis ovatis cordatis obtusis subrepandis non lineato-plicatis, pedunculis tenuibus 1-2-floris folio longioribus, pedicellis calyci æquilongis, bracteis minimis setaceis, sepalis oblongis breviter acuminatis, corollâ albâ extus lineatim hirtâ calyce triplo longiore, capsula hirsutâ, semine tuberculis conicis aspero ♀.

Hab. in agris ad Giaur Göl prope Marasch (Germanicam) Cataonie (Haussk!). Fl. Jul.

Caules sesquipedales, folia majora pollicaria. An varietas *C. Galatici?* sed indumentum multo tenuius, folia non rugoso-plicata, flores minores albi et seminis sculptura tuberculis nec rugulis irregularibus constans.

52. *C. hirsutus* (Stev. in M.B. Taur. Cauc. I, p. 422) pilis fulvescentibus hirtus vel plus minus hirsutus, caulis crassiusculis elongatis decumbentibus vel scandentibus simplicibus, foliis petiolo limbo 3-4-plo breviore suffultis cordato-hastatis acutis rarius obtusiusculis margine obscure sinuatis auriculis basilaribus integris vel lobatis, foliis superioribus sensim diminutis, pedunculis folio multo longioribus 3-t-floris, pedicellis calyci aequilongis vel longioribus, bracteis anguste linearibus, sepalis hirsutis ovato-oblongis acuminatis, corollâ calyce quadruplo longiore extus ad plicas pilosâ albo-roseâ vel flavidâ, capsula hirsutissimâ, seminibus concoloribus elevatim ruguloso-tuberculatis ♀. Ic. M.B. Cent. rar. tab. 6. — *C. betonicifolius* Mill. Diet. № 20. — *C. pubescens* Russ. Alep. p. 152. — *C. sagittifolius* Sibth. Fl. Gr. tab. 193. — *C. atriplicifolius* Poir. Diet. Suppl. III, p. 467. — *C. lanuginosus* Vahl Symb. III, p. 23 ex descript. non Desr. — *C. tomentosus* Choisy in D.C. Prodr. IX, p. 413 non L. — *C. amoenus* C. Koch Linn. XIX, p. 19. — *C. Sibthorpii* R. et Sch.

Hab. in agris, inter segetes, ad dumeta, Graecia in Peloponneso (herb. Fauché! Sprunt!), Bœtiâ (Sprunt!), Phtiothide (Heldr!), Archipelago (Sil.th.), insula Chio (Orph!), Rhodo (Bourg!), Thracia (Friv. exs. sub *C. Scammonia!*), Bithyniâ (Thirk!), Galatiâ (Wied. 149!), Lyciâ (Bourg!), Ciliciâ (Ky!), Syriâ boreali, Mesopotamiâ (Ky. 76! et 232!), Anch. 1392!, Haussk!), Syriâ littorali (Boiss! Blt Gaill!), Cælesyriâ (Ehrenb!), Palestinâ (Boiss!), Tauriâ merid. (Steven!).

β. *virescens*. — Folia minora obtusiora, corolla minor luteo-virescens.

Hab. in campis Pisidiæ prope Egirdir (Heldr!).

γ. *tomentosus*. — Folia dense tomentoso-pannosa subsericea. Præter indumentum a typo non diversus et ad eum formis intermediis transiens. — *C. peduncularis* Boiss. Diagn. Ser. I, 11, p. 84.

Hab. in Mesopotamiâ inter Orfa et Suerek (Ky. 58!), inter Bir et Teharmelik (Haussk!), in Libano ad Bezulmar et prope Cedros (Bl!).

Ar. Geogr. Gallia australis orientalis sed-prob. introductus.

53. *C. Armenus* (Boiss. et Ky. pl. exs. 1859) parce et breviter hirtus pallide virens, caulis elongatis volubilibus, foliis a basi sagittato-hastata trianguli-lanceolatis inferne subdentatis auriculis profunde et obtuse 2-3-lobis inferioribus longiuseule petiolatis, superioribus sensim diminutis et angustatis anguste hastato-lanceolatis acutissimis, pedunculis folio multo longioribus saepe longissimis 3-7-floris, pedicellis calyci aequilongis vel longioribus, bracteis setaceis, sepalis patule hirtis oblongo-lanceolatis tenuiter acuminatis, corollâ

pallide roseâ extus lineatim hirtulâ calyce quadruplo longiore. capsulâ hirsutâ, seminibus tuberculis et rugulis albidis elevatis obsitis ♀. *C. batataefolius* Boiss. et Haußk. MSS.

Hab. in Armeniae australis lapidosis prope Goschkar prov. Musch 5600' et ad Gumgum prope Warto 4500' (Ky. 373!), ad ripas et sepes humidas Syriæ borealis et Cataoniæ ad Aintah et inter Göksu et Behesne (Haussk!).

Caules longissimi, folia inferiora saepè magna tripollicaria, corolla pollicem et amplius longa sed ea *C. hirsuti* manifeste angustior. A formis hujus speciei foliis magis elongatis et lanceolatis, bracteis sepalsique apice setaceis, indumento brevissimo sparsa, corollæ formâ et seminis sculpturâ differt.

54. *C. stenophyllus*, adpresso et brevissime tomentellus, caulis tenuibus decumbentibus volubilibus elongatis, foliis auriculato-cordatis crenato-dentatis margine undulatis inferioribus oblongis acutiusculis longiuscule petiolatis superioribus vel omnibus anguste lanceolato-linearibus linearibusque acutissimis, pedunculis 3-4-floris folio æquilongis vel longioribus, pedicellis calyce longioribus basi minutissime bracteolatis, sepalis adpresso hirtis ovatis obtusis interdum mucronulatis, corollâ sulphureâ extus lineatim hirtulâ calyce quadruplo longiore, ovario hirsuto ♀. *C. Palæstinus* var. *diversifolius* et var. *stenophyllum* Boiss. Diagn.

Hab. in Syriâ (nec in Bithyniâ) (Pestalozza!), inter Tripoli et Deir Keftim (Bl. 1351!) et in Libano inter Khan Medeiridj et Khan Djedid et inter El Kleâ et Djedaïde (Gaill. exs. 2054! et 2057!).

Caules bipedales et longiores, folia superiora et saepè omnia angustissima 4-1½ poll. longa 2-4 lineas lata, flores eis *C. Scammoniae* et *hirsuti* paulo minores. Prius pro varietate *C. Palæstini* habueram sed certe differt capsulâ hirsutâ et foliorum formâ.

§ 8. Scandentes.

55. *C. althæoides* (L. Sp. 422) patule rarius adpressiuscule villoso-hirsutus, caulis diffusis vel scandentibus, foliis inferioribus longius petiolatis cordato-ovatis obtuse crenatis vel lobatis superioribus subpedatim palmatifidis partitivis laciniis linearibus vel lanceolatis integris vel dentatis terminali longiore, pedunculis axillaribus folio multo longioribus uni rarius bifloris, pedicello calyce longiore, bracteis anguste setaceis, sepalis ovato-oblongis obtusis margine scariosis interdum mucronulatis, corollâ roseâ calyce 4-5-plo longiore extus zonatim hirsutâ, capsulâ glabrâ. ♀. Ic. Fl. Græc. tab. 194.— *C. hirsutus* Ten. Nap. I, p. 60, tab. 15. — *C. Italicus* R. Sch. Syst. IV, p. 266. — *C. bryoniifolius* Rehb. exc. p. 348. — *C. alceaefolius* Fl. Pélop. p. 68 non Lam.

Hab. in cultis et dumosis regionis inferioris, Peloponneso (Bory, Sprun!), Archipelago (Urv! Orph!), Cretâ (Heldr!), Rhodo (Auch!), Bithyniâ (Pestal.), Cypro (Ky!), Syriâ littorali (Bl! Gaill!), deserto Palestinae ad meridiem Gaza (Boiss!). Ægypto inferiore ad Alexandriam (Ehrenb!).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, Africa borealis, Madera, insulæ Canarienses.

56. **C. tenuissimus** (Fl. Graec. II, p. 79, tab. 195) adpressissime sericeo-argyreus, caulis diffusis vel scandentibus, foliis inferioribus longius petiolatis lineato-plicatis cordato-ovatis obtuse repando-crenatis, caulinis plerisque in lacinias lineares elongatas palmatim vel pedatim partitis, pedunculis unifloris axillaribus folio multo longioribus, pedicello ealyce aequilongo vel longiore, bracteis brevissimis, sepalis ovato-oblongis obtusis margine scariosis, corollâ roseâ calyce 4-5-plo longiore extus zonatum hirsutâ, capsulâ glabra ♀. *C. argyreus* D.C. Fl. Fr. Suppl. 423.

Hab. in rupestribus et collinis regionis inferioris, Zaconthus (Marg!), Peloponnesus (Bory!), Attica (Spr! Boiss!), Macedonia (Friv!), Creta (Heldr!), Anatolia in Pisidiâ (Tchih!), ditio Kesraouan Libani (Bl!).

Indumento et foliis caulinis angustius et profundius divisis a precedente specifice distinctus videtur.

Ar. Geogr. Italia australis, Sicilia, Dalmatia, Africa borealis.

57. **C. stachydifolius** (Choisy in D.C. Prodr. IX, p. 408) caulis retrorsum pubescentibus prostratis flexuosis tenuibus, foliis brevissime et adpresso-crenatis pubescentibus paginâ superiori interdum glabratis petiolatis cordatis ovatis circumflexa obtuse crenato-lobatis supremis sensim diminutis acutis acute dentieulatis, pedunculis 3-4-floris folio axillari multo longioribus, pedicellis calyce 2-3-plo longioribus tandem refractis, bracteis longis setaceis, sepalis ellipticis margine scariosis obtusissimis vel retusis muticis glabris vel dorso parce hirtis, corollâ intense purpureâ patenti ealyce quadruplo longiore, capsulâ mucronatâ glabrâ, seminibus levibus ♀. *C. Damascenus* Boiss. et Gaill. Diagn. Ser. II, 6, p. 122. — *C. venulosus* Ehr. MSS.

Hab. in graminosis collinis et agris Syriae borealis ad Aleppo (Ky. 431! Haussk!), inter Aintab et Nisib (Haussk!), circa Damascum copiose (Gaill!) in Cœlesyriâ (Ehreub!), in Mesopotamia (Oliv! Rousseau!), circa Mossul (Noë! Haussk!).

Nomen specificum omnino improprium, facies *Erodii*, caules 1-2-pedales, folia majora pollicaria, corolla diametro 12-15 lineas lata. Habitus *C. Galatici* a quo egregie differt indumento brevi parco caulum deflexo, pedunculis et pedicellis longissimis, sepalis obtusis muticis margine scariosis, capsula glabrâ, seminibus levibus.

58. **C. Palæstinus** (Boiss. Diagn. Ser. I, II, p. 84) adpresso et breviter pubescenti-tomentellus, caulis tenuibus prostratis vel decumbentibus subvolvilibus elongatis, foliis ovato-oblongis obtuse auriculato-cordatis obtuse undulato-crenatis inferioribus longe petiolatis obtusis, superioribus sensim diminutis acutis, pedunculis axillaribus 3-4-floris folio brevioribus vel vix longioribus, pedicellis ealycei

æquilongis vel brevioribus, bracteis setaceis minimis, sepalis hirtis oblongo-spathulatis obtusissimis muticis, corollâ ex sicco luteâ calyce quadruplo longiore ad angulos extus hirtâ, ovario glaberrimo ♀.

Hab. in Palestinâ loco non notato (Boiss!).

Caules 2-4-pedales, folia majora pollicem longa basi 9 lineas lata; formis minoribus *C. hirsuti* affinis ab eo differt foliis crenato-lobatis infimis petiolo sœpe bipinnicili suffultis et praesertim ovario glaberrimo et sepalis obtusis muticis. A *C. Scammoniâ* ovario glabro et sepalis truncatis quoque gaudenti distinguitur tomento totius plantæ, foliis parvis obtusis et sepalis non mucronatis.

59. C. Scammonia (L. Sp. 218) glaberimus, radice longâ fusiformi, caulis longissimis scandentibus simplicibus, foliis petiolo limbo breviore suffultis sagittatis acuminatis basi cordatis auriculis divergentibus acutis integris vel remote angulato-dentatis, superioribus angustatis interdum linearibus pedunculis longissimis erecto-patentibus 3-1-floris, pedicellis calyce subbrevioribus basi bracteis minutis subscariosis ovatis obsitis, sepalis laxiusculis ovatis margine scariosis obtusissimis vel truncaitis muerone brevi interdum extus flexo, corollâ glabrâ patenti calyce 4-5-plo longiore ochroleucâ extus ad plicas rubellâ, capsulâ glabrâ, seminibus obscure tuberculatis ♀. Ic. Fl. Græc. tab. 192. — *C. pseudoscammonia* C. Koch Linn. XXIII, p. 598.

Hab. inter segetes et praesertim ad sepes et dumeta scandens in regione calidiori, Lydia circa Smyrnam (Boiss! Ball!), insula Cos (Urv.), Rhodus (Sibth., Bourg!), Caria (Pinard!), Cilicia littoralis (Bal!) et usque ad Gülek 3800' (Ky!), Syria borealis (Haussk!) et in regione inferiore Amaní (Ky. 16!), Syria littoralis in regno infer Cassii (Boiss!), prope Tortose (Gail!), et in Libano prope Eden (Reyg!), Galilea ad Banias (Boiss!), Armenia prope Ispir (Huet!), Tauria meridionalis in valle Laspi (Stev.). Fl. Mai. Jun.

Radix longissima succum concretem officinalem *Scammoniâ* nomine notum præbens, caules sœpe 10-15 pedes longi, flores magni, capsula pisum æquans.

60. C. arvensis (L. Sp. 218) glaber vel pubescens, caulis elongatis prostratis vel volubilibus, foliis petiolatis obtuse vel acute sagittato-auriculatis acutis triangulari-ovatis vel oblongis vel plus minus anguste linearibus, pedunculis axillaribus solitariis petiolo longioribus unifloris rarius 2-3-floris, bracteis ad pedicelli elongati basin linearibus vel linear-i-oblongis angustis, sepalis ovato-rotundis obtusis margine scariosis, corollâ albâ vel roseâ glabrâ calyce pluries longiore, capsulâ glabrâ ♀.

Hab. ad vias, in cultis totius regionis ad Egyptum (Gail!), Babyloniam (Noë!). Arabiam petream (Schimp. 278!), Persiam australē (Haussk!), Affghaniam et Belutschiam (Stocks!) usque. Crescit a regione inferiori ad alpinam, mons Makmel Libani (Ecknbl.), Kuh Delu Persiæ australis (Ky!) et Kuh Nur alt. 12000' (Haussk!). Arabice *Alek*.

Ar. Geogr. Orbis fere totus.

SER. B. ANNUL.

§ 9. **Siculi.**

61. C. Fatmensis (Kze Flora 1840. I. p. 172) viridis, sub lente minute puberulus vel glabratus, radice tenui, caulis diffusis elongatis tenuibus flexuosis, foliis petiolis in inferioribus aequilongis suffultis cordato-oblongis obtusis late obtusissime et inaequaliter dentato-lobatis. pedunculis axillaribus 2-1-floris petiolum aequantibus vel brevioribus, pedicello calyce subbrevoire tandem recurvo, bracteis minimis setaceis, sepalis ovatis obtusissimis margine scariosis, corollâ parvâ glabrâ calyce subdupo longiore, seminibus levibus ☽. *C. hastatus* Sieb. exs. — *C. ambiocalyx* Steud.

Hab. ad margines agrotum Ægypti ad Saovieh (Sieb!), ad Choubrah (Husson!), ad Esneh (Kralik!).

Caulis filiformis saepe bipedales et longiores, folia majora pollicaria, corolla parva 4 lineas longa.

Ar Geogr. Arabia tropica, Algeria in ditione Saharensi.

62. C. Cælesyriacus (Boiss. Diagn. Ser. I. 41, p. 85) viridis glabrescens vel sub lente parce hirtulus, caulis decumbentibus et ascendentibus tenuibus subvolubilibus, foliis inferioribus longe petiolatis cordato-reniformibus obtusissimis indivis subrepandis. intermediiis quinquelobis vel partitis laciniis oblongo-linearibus terminali elongata. summis trilobis vel anguste linear-i-hastatis, pedunculis axillaribus unifloris folio brevioribus, pedicello calyce et saepius pedunculo ipso longiore fructifero incrassato et recurvo. sepalis patule hirtis obovatis obtusis abrupte et longe mucronatis, corollâ roseâ ad angulos extus puberâ calyce triplo longiore, capsulâ glabrâ, seminibus levibus ☽. *C. heterophyllus* Ehrenb. MSS.

Hab. in Cælesyriâ inter Hasbeya et Rascheya (Boiss!), in cultis et rupestribus prope Sidonem (Gall!), Tripoli (Bl!), ad Arissam (Ehrenb!).

Affinis *C. Fatmensi*, corolla major 7 lineas longa, sepala longe mucronata nec mutica, insignis insuper foliis inferioribus et superioribus valde dissimiliis.

63. C. Siculus (L. Sp. 223) pallide virens plus minusve pubescenti-hirtus. caulis procumbentibus vel erectis simplicibus saepe elongatis flexuosis, foliis petiolatis cordato-ovatis acutis supremis oblongis basi cuneatis. pedunculis axillaribus solitariis unifloris flore sublongioribus folio brevioribus sub flore bracteis binis lanceolatis calycem aequantibus vel superantibus obsitis fructiferis recurvis, sepalis ciliatis ellipticis attenuato-acutis, corollâ calyce vix duplo longiore limbo cyaneo extus lineatim hirto profunde quinquelobo, capsulâ glabrâ ☽. Ic. Fl. Græc. tab. 196.

Hab. in saxosis apriis rarius in cultis, Argolis et Attica (Sprun!), Messenia ad Pylos (herb. Fauché!), Archipelagus (Wied!), Creta (Sibth.), Cyprus (Ky!), Ægyptus ad Alexandriam (Ehrenb!).

Ar. Geogr. Hispania australis et orientalis, Gallia austro-orientalis, Corsica, Sardinia et Sicilia, Italia australis, Africa borealis, Nudia littoralis, Madera, insulae Canarienses.

64. **C. pentapetaloides** (L. Syst. Nat. III, p. 229) pilosus viridis, caulis ascendentibus et prostratis simplicibus saepè elongatis, foliis obtusis radicalibus oblongo-spathulatis in petiolum longe attenuatis, superioribus latiuscule linear-lanceolatis sessilibus, pedunculis unifloris axillaribus solitariis flore longioribus folio brevioribus ad medium minutissime bracteolatis demum patentibus, floribus parvis, sepalis subscariosis ovato-oblongis acutis, corollâ calyce subtriplo longiore tubo flavido limbo cyaneo profunde quinquelobo, capsulâ glabra ☽. Ic. Flor. Græc. tab. 197. — Cavan. Ic. rar. tab. 123.

Hab. in rupestribus siccis et graminis, Peloponnesus in Messeniâ (Bory!) et Argolide (Sprun!), Archipelago (Sibth.), Rhodi (Bourg!), Cypro (Ky!), Ciliciâ littorali (Bal!), Syriâ interiori ad Aleppo (Haussk!) et littorali (Bl!), Palestina (Boiss! Ky. 515!). Fl. vere.

Ar. Geogr. Hispania, Italia australis et insulæ.

65. **C. Stocksii.** adpressissime hirtus pallide virens, caulis ascendentibus et erectis tenuibus simplicibus stricte foliosis, foliis anguste et longissime linear-lanceolatis acutis basi attenuatis, intimis brevibus oblongo-lanceolatis subpetiolatis, cymis 1-3-floris pedunculis tenuissimis strictis folio axillari longioribus suffultis, pedicellis calyci æquilongis basi bracteis linear-setaceis suffultis, floribus minimis, sepalis subscariosis glabris ovatis acutis, corollâ roseâ calyce duplo longiore extus ad angulos sericeâ, ovario glabro ☽. *C. tenellus* J. E. Stocks in Hook Journ. Bot. 1852, p. 172 non Lam.

Hab. ad Schah Bilawul Belutschiae (Stocks!).

Planta semipedalis, folia 4-4½-pollicaria sesquilineam lata, pedunculi fere bipollicares stricti rigiduli tenuiter filiformes, sepala vix 2 lineas longa. Species valde distincta facie *Evolvuli* sed stylus unicus apice bifidus.

§ 10. Undulati.

66. **C. undulatus** (Cav. Ic. III, p. 39, tab. 277, 1) patule hirtus pallide virens, caulis ex collo pluribus crassinerviis saepè longinerviis simplicibus decumbentibus centrali erecto, foliis spathulato-oblongis obtusis subundulatis inferioribus basi attenuatis subpetiolatis superioribus semiamplexicanibus, floribus in axillis superioribus solitariis subsessilibus terminalibus subcapitato-congestis foliis floribus subsuperatis, sepalis hirtis breviter oblongo-lanceolatis acutis, corollâ calyce subduplicite longiore tubo cylindrico glabro limbo violaceo expanso extus ad angulos adpresso hirtâ, capsulâ magnâ sphæricâ

hirsutæ ♂. *C. humilis* Jacq. Coll. IV, tab. 32. — *C. crotaloides* Desf. Atl. I, p. 176, tab. 49. — Fl. Graec. tab. 198.

Hab. in Cypro (Sibth.), in Ægypto (ex auct.). E ditione nondum vidi.

Ar. Geogr. Hispania, Liguria, Sicilia, Africa borealis.

Species Convolvuli in ditione ut videtur erronee indicate.

C. suffruticosus Desf.

Hab. in Græciâ (ex herb. Sibth.).

Species ex Africâ boreali tantum nota.

C. farinosus L.

Hab. in arvis et sepibus Peloponnesi et Asiae minoris in Mysiâ (Sibth.).

Species Africana tropica et Capensis.

C. tricolor L.

Hab. in Bithyniâ (C. Koch Linn. 19, p. 19).

Ex hortis aufuga; species regionis mediterraneæ magis occidentalis.

CALYSTEGIA (R. Br. Prodr. p. 483).

Corolla infundibuli-campanulata limbo quinqueplicato repando. Stylus unicus stigmatibus binis ovatis planiusculis. Ovarium semibiloculare loculis biovulatis. Capsula subunilocularis bivalvis. — Herbae repentes vel volubiles flores basi 2 bracteis dilatatis cincti.

1. **C. sepium** (L. Sp. 218 sub *Convolvulus*) caulis proceris scandentibus, foliis cordato-oblongis vel sagittato-hastatis auriculis basilaribus integris angulatis vel truncatis, pedunculis longis unifloris, bracteis ovatis subcordatis planiusculis apice attenuatis, sepalis oblongis acutis, corollâ albâ magnâ ♀. *Cal. sepium* R. Br. Prodr. p. 483. — Ic. Fl. Dan. tab. 458. Ic. Rehb. tab. 139.

Hab. ad sepes et fossas humidas, Attica ad Phalerum (Orph! Heldr!), Byzantii (Bergg.), Creta ad Candia (Heldr!) et ad Platania (Raui!), Rhodus ad Salakos (Bourg!), Syria littoralis ad Tripoli (Bl!), Anatolia sept. (Wied!), Tauria et prov. Caucasicæ (Ledeb.), littora Caspia ad Lenkoran (Hoh. exs. sub *C. sylvatica*!).

Ar. Geogr. Europa omnis a Britannia et Scandinaviâ ad Rossiam, Sibiria, America borealis et ex d.c. Prodr. Australia.

2. **C. sylvatica** (W. K. pl. rar. Hung. III, p. 390, tab. 264 sub *Convolvulus*) caulis proceris scandentibus, foliis cordato-ovatis vel oblongis auriculis basilaribus integris vel truncatis, pedunculis longis unifloris, bracteis amplis cuneatatis basi ventricosis ovatis obtusis, sepalis ovatis obtusiusculis mucronatis, corollâ albâ amplissimâ ♀. Ic.

Rchb. tab. 140. — *Cul. sylvatica* Chois. in p.c. Prodr. — *Conv. sylvestris* Willd. Enüm. I, p. 202.

Hab ad sepes humidas et in sylvaticis, Laconia prope Kastanitza (Heldr!), Macedonia, Thracia et Bithynia (Griseb!), Pontus ad Samsun (Tchih!), ad Trapezuntum (Bourg!) et Rhizé (Bal!), in prov. Caucasicis (ex C.A. Mey.).

Vix foliis sed bracteis amplioribus obtusioribus magis patentibus a præcedente (an satis?) distincta.

Ar Geogr. Regio Danubialis, Italia australis et Sicilia. Dalmatia, Africa borealis.

3. C. Soldanella (L. Sp. 226 sub *Convolvulo*) caulis repen-tibus inferne denudatis, foliis carnosulis reniformi vel rotundato-cordatis obtusissimis petiolatis integris vel subrepandis. pedunculis axillaribus unifloris folio longioribus, bracteis ovatis obtusis concavis, sepalis ovatis obtusis vel retusis mucronulatis, corollâ purpureâ calyce quadruplo longiore ♀. Ic. Engl. Bot tab. 314. Rchb. Ic. 140. — *Cal. Soldanella* R. Br.

Hab in arenis maritimis, Zaconthus (Marg!), Peloponnésus (Bory!), Attica (Sprn!), Macedonia et Bithynia (Sibth.), littora Ponti (Tchih! Bal!), Guriæ (Rupr!), Taurie (M.B.), littora Caspia Persiae borealis (Szov!), Cilicia (Tchih.).

Ar. Geogr. Littora totius Europæ septent. a Galliâ occidentali ad Bataviam et Germaniam, littora mediterranea, California, America australis, Nova Hollandia.

IPOMÆA (L. Sp. 227).

Calyx quinquesepalus. Corolla infundibuliformi-campanulata quinqueloba vel repanda quinqueplicata. Stylus unicus stigmate capitato saepius bilobo. Ovarium 2 rarius 3-4-loculare loculis dispermis. Capsula bilocularis rarius 3-4-locularis. — Herbæ vel frutices facie *Convoluti*.

1. I. littoralis (L. Sp. 227 sub *Convolvulo*) glabra, caulis repentibus subtus fibrillosis stolones elongatos edentibus, foliis carnosulis aliis ovato-panduratis basi cordatis apice retusis saepe sinuatis, aliis trilobis vel tripartitis lobis obtusis lateralibus interdum bifidis terminali majore elliptico-oblongo, pedunculis ex axillis 1-2 superioribus unifloris folio brevioribus, pedicello calyce longiore superne incrassato basi minute bracteolato. sepalis ellipticis obtusis mucronulatis margine membranaceis. corollâ magnâ ochroleucâ glabrâ calyce 4-5-plo longiore. stigmatis lobis rotundatis, capsulâ magnâ biloculari, seminibus binis magnis villosissimis ♀. *Convolvulus Imperati* Vahl Symb. I, p. 47. — *C. arenarius* Vahl Symb. I, p. 48. — *Conv. stoloniferus* Cyr. Pl. rar. I, tab. 5. — *Conv. sinuatus* Petagna. — *Batatas littoralis* Chois. p.c. Prodr. IX, p. 337. — *Bat. sinuata* Guss.

Hab. in arenosis maritimis Palestinae (Auch. 1401!), ad Gaza (Bové!), Syriae ad Sidonem (Pest! Gaill!), ad Berythum (Ehrenb! Bl!), ad Tripoli (Boiss!), ad Alexandrette (Haussk!). Fl. aest.

Hab. in regno Neapolitano, Africâ boreali, insulis Azoricis, Americâ boreali, Antillis.

2. I. sagittata (Desf. Atl. I, p. 177) glabra, caulis tenuibus scandentibus elongatis. foliis inferioribus cordato-suborbiculatis ceteris sagittatis sensim angustioribus lanceolatis et linear-lanceolatis acutis auriculis linear-lanceolatis sepe longissimis. pedunculis axillaribus folio brevioribus 1-2-floris versus medium minutissime bibracteolatis. sepalis ellipticis obtusis mucronulatis, corollâ amplissimâ roseâ glabrâ, stigmate capitato retuso, seminibus tomentosis ♀. Cav. Ic. rar. II, tab. 207. — Lam. Ill. tab. 104. — Conv. Wheleri Vahl Symb. II, p. 36. — Conv. sagittifolius Mich. Am. I, p. 138.

Hab. ad fossas et in dumetis humidis maritimis, Cilicia Trachea inter Selefke et Perchende (Tchih!), Syria ad Alexandrette (Ky. 154!).

Ar. Geogr. Hispania orientalis, Sicilia, Africa borealis, America borealis.

3. I. palmata (Forsk. Descr. p. 43) glabriuscula, caulis proceris valde scandentibus, foliis petiolatis quinatis partitionibus ellipticis interdum subpetiolulatis glabris vel saepius acutis et acuminatis margine minute serrulato-asperis lateralibus interdum bifidis. pedunculis 1-3-floris foliò axillari longioribus, pedicellis calyce longioribus incrassatis, sepalis oblongis obtusis margine membranaceis, corollâ campanulatâ purpureâ glabrâ calyce 5-6-plo longiore, stigmatis lobis subrotundis, seminibus tomentosis ♀. Conv. Caiicus L. Sp. 222. Bot. Mag. 699. — Ipom. Senegalensis Lam. Ill. I, p. 464.

Hab. ad sepes Ægypti et inter arundines ad Nili ripas (Forsk., Del., Auch. 1396!), ad Rosette (Ehrenb!), ad Benisouef (Sieb!), ad Philæ (Boiss!), in Syria littorali ad Ras el Ain prope Tyrum (Gaill!).

Ar. Geogr. Nubia, Abyssinia, Senegalia.

EVOLVULUS (L. Sp. 394).

Calyx quinquespalus. Corolla campanulata vel infundibuliformis ore repanda vel quinqueloba. Styli duo bifidi. Ovarium biloculare quadriovulatum. Capsula bilocularis. — Herbae vel suffrutices.

1. E alsinoides (L. Sp. 392) patule villosus a collo ramosus ramis tenuiter virgatis decumbentibus simplicibus subdistiche et crebre foliosis, foliis subpetiolatis obovato-oblongis obtusis, pedunculis filiformibus 2-4-floris folio brevioribus, pedicellis nutantibus calyce subaequilongis basi bracteatis, sepalis breviter lanceolatis acutis hispidis, corollâ parvâ hirsutâ calyce duplo longiore, capsulâ seminibusque glabris levibus ♀. Ic. Burm. Zeyl. tab. I, fig. 61. — E. linifolius L. — E. fugacissimus Hochst. in Schimp. exs.

Hab. in Belutschia (Stocks!).

Ar. Geogr. Abyssinia et Africa tropica, India Orientalis, America tropica.

SEDDERA (Hochst. et St. — Choisy in D.C. Prodr. IX, p. 440).

Calyx ad basin quinquepartitus. Corolla tubuloso-infundibuliformis. Stamina inclusa. Styli duo. stigmatibus capitatis. Ovarium biloculare loculis dispermis. — Suffrutices ramosissimi.

1. **S. latifolia** (Hochst. et St. in Schimp. exs. № 883!) ramosissima ramulis intricatis deum subspinescentibus tenuibus brevibus, foliis minutis adpresso cinereo-sericeis ovatis brevissime petiolatis, floribus in axillis superioribus solitariis sessilibus breviter spicatis, sepalis oblongis acuminatis, corollâ calycem paulo superante, capsulâ glabrâ calyce sublongiore 5.

Hab. in collibus lapidosis Belutschiae (Stocks!).

Ar. Geogr. Nubia, Abyssinia, Arabia tropica, ditio Scinde Indiæ.

CRESSA (L. Sp. 325).

Calyx ad basin quinquepartitus. Corolla infundibuliformis ultra medium quinquepartita. Stamina 5 imae corollæ affixa exserta. Styli 2 stigmatibus capitatis Ovarium biloculare quadriovulatum. Capsula irregulariter dehisceens abortu saepius monosperma. — Herba humilis.

1. **C. Cretica** (L. Sp. 325) tota pilis adpressissimis aliisque patulis obsita cinerea, caulinis ex collo pluribus brevibus erectiusculis vel diffusis a basi ramosissimis ramisque cerebellime foliosis, foliis minutis sessilibus acutiusculis apice callosis caulinis a basi subcordatâ ovato-oblongis ramulorum elliptico-lanceolatis, axillis gemmiferis, floribus brevissime pedunculatis spicas breves densas oblongas bracteatas terminales formantibus, calycis lacinias eucullatis ellipticis obtusis, corollæ calyce vix duplo longioris lacinias recurvis subtus et versus apicem pilosis vel glabratis, capsulâ ovatâ acutiusculâ, semine levâ 2. Ic. Fl. Græc. tab. 256. — Lam. Ill. tab. 183.

Hab. in argilloso salsis maritimis et desertoram, Peloponnesus (herb. Fauché!), Attica ad Phalerum raro florens! (Heldr!), Macedonia (Friv!), Archipelagus (Sibth!), Creta (Sieb!), Palestina prope Joppen (Roth!), Egyptus ad Alexandriam et desertis Suez (Sieb! Ehr! Gaill! etc.), Arabia petrea (Schimp. 720! Boiss!), Babylonia (Noë! Haussk!), Persia australis ad Abuchir et interior inter Teheran et Isphahan (Haussk!) et inter Herat et Tebes (Bge!), Transcaucasia ad Caspium (Ledeb.), Belutschia (Stocks.).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, Africa borealis, Nubia, Abyssinia, India, Australia et prob. America tropica.

TR. II. CUSCUTEÆ. — Plantæ aphyllæ parasitæ embryone spirali acotyledoneo.

CUSCUTA (L. Gen 170).

Calyx 4-5 fidus rarius partitus. Corolla tubulosa vel campanulata 4-5-fida post anthesin saepe urceolata sub staminibus squamulis fimbriatis vel dentatis sapissime instructa. Stamina corollæ tubo adnata ejus lobis alterna. Ovarium biloculare quadriovulatum, styli 2 liberi vel in unicum concreti. Fructus saepius capsularis membranaceus circumcisus vel transverse et irregulariter lacerus, rarius baccatus. — Herbae filiformes ramosissimæ parasitæ. Genus ob specierum polymorphiam (ex loco, plantâ nutrici, oriundam) difficile, infoste ex speciminibus siccis elaborandum. Specierum imprimis Gerontogearum sectionis *Eucuscute* circumscriptio adhuc imperfecta, sic v. gr. diagnoses comparatativas ex vivo *C. Epithymi* et *planiflora* nullibi reperies. Monographiam accuratissimam amic. Engelmann paucis mutatis secutus sum.

SECT. I. CUSCUTA Engelm. — Styli duo, stigmata elongata.

§ 1. **Eucuscuteæ** Engelm. — Styli stigmatibus filiformibus æquilungi vel longiores. Capsula regulariter circumscissa.

* Styli ovario longiores.

1. **C. Epithymum** (Linn. Syst. Ed. Murr. p. 140) caule ramoso, floribus sessilibus vel subsessilibus in capitula globosa dispositis, calycis lobis ovatis acutis patulis, corollæ tubo cylindrico calycem superante, limbi tubo brevioris lobis ovatis acutis, squamulis conniventibus faucentibus clandentibus fimbriatis ♂. Ic. Engl. bot. tab. 55. — Rehb. Ic. V, tab. 499. — *C. minor* D.C.

Hab. *Labiatarum*, *Genistcarum*, *Poterii*, multarumque aliarum parasitica in totâ fere ditione a Græciâ et Macedoniâ ad regiones Caucasicas, Asiam minorem.

Species valde variabilis quâcum praeter *C. Trifolii* Bab. in ditione nostrâ nondum observatam formas sequentes cl. Engelmann conjungit.

$\beta.$ *macranthera*. — *C. macranthera* Heldr. et Sartori in Boiss. Diagn. Ser. I, 11, p. 126. Calyx basi longius obconicus laciniis obtusioribus, antheræ submajores magis exsertæ.

Hab. in regione superiore Taygeti et Parnassi ad *Daphne buxifoliam*, *Sideritides* etc., parasitica (Heldr.).

$\gamma.$ *Calliopes*. — *C. Calliopes* (sphalmate *Cassiopes*) Heldr. et Sart.

loc. cit. p. 128. — Praecedenti affinis sed calyx corollæ proportione brevior et flores distinctius pedicellati.

Hab. in regione alpinâ Parnassi ad *Daphnem buxifoliam* parasitica (Orph! Guicci).

δ. micrantha. — Flores minores semper sessiles, calyx corollæ tubum subæquans, ejus et corollæ lobi acutiores. — *C. alba* Presl Del. Prag. 87. — *C. planiflora* Koch Syn. ex Engelmann non Tenore. — *C. Kotschy* Desmoul. non Boiss. Planta sœpe rubella.

Hab. in montibus Græciæ et Macedoniae (ex Engelmann).

Ar. Geogr. Europa omnis ab Angliâ et Daniâ ad Rossiam medium, Sibiria Altaica, Africa borealis.

2. C. planiflora (Ten Syll. p. 128. — Fl. Neap. tab. 220, fig. 3 — Engelmann. Cusc. p. 14) caule ramoso, floribus sessilibus in capitula globosa multiflora dispositis, calycis lobis obtusiusculis, corollæ scutellatæ limbo longiore valde expanso rotato lobis ovatis breviter acuminatis obtusis, squamulis conniventibus fimbriatis ⊖.

Species ex cl. Engelmann omnium maxime variabilis et cuius formæ nonnullæ ex ipso a *C. Epithymo* ægre discernuntur; differt corollæ subtortatæ limbo tubi proportione longiore valde expanso lobis obtusioribus, staminibus et stylis ob corollam expansam magis exsertis. Probabilissime hujus varietas, sed eas vivas non comparavi et auctores sequor. Ea in Oriente multo vulgatior et formæ sequentes occurruunt plus minusve altera in alteram transeuntes.

α. Tenorii Engelm. loc. cit. — Calycis et corollæ lobi crassi turgidi apice sœpe cucullati, flores minores, styli ovario longiores capsulæ æquilongi vel breviores. — *C. alba* Auct. et collectorum plurimorum non Presl.

β. approximata Eng. loc. cit. — Calycis lobi membranacei vix carnosæ, corollæ lobi apice tantum turgidi, flores et capitula majora, styli ovario multo longiores. — *C. approximata* Babingt. — *C. urceolata* Kunze Flor. 1846 p. 651. — *C. Epispertos* Boiss. et Orph. MSS. — *C. leucosphæra* Boiss. et Heldr. pl. exs. — *C. Schiraziana* Boiss Diagn. Ser. I, 11, p. 86 (forma calycis et corollæ lobis angustioribus).

Hab. in totâ ditione, in regione montanâ frequentior a Græciâ! Thraciâ! Asiâ minore! ad Syriam et Mesopotamiam (Ky! Haussk!). Ægyptum (Fischer), Persiam borealem et australem (Ky! Bge!), regiones Caucasicast, Belutschiam (Stocks!), Affghaniam (Griff!).

Ar. Geogr. Insulæ Canarienses, Africa borealis, Hispania, Gallia australis, Italia et Sicilia, Rossia australis, Songaria, Sibiria Altaica, regio Himalaica occidentalis. In Europam medium interdum introducta.

C. Palæstina (Boiss. Diagn. Ser. I, 11, 86) caulinibus ramosis capillaceis valde intricatis, floribus sessilibus in capitula minima glomeratis, calycis carnosuli lobis latis brevibus carinato-costatis obtusis, corollæ calycem vix excedentis lobis breviter ovatis obtusis, squamulis majusculis late spathulatis denticulatis, stylis ovario sublongioribus ⊖. *C. Cretica* Tourn. Cor. 45. — *C. micrantha* Tineo. — *C.*

capillaris Rehb. Crit. V, 64. — *C. globularis* Bertol. Fl. Ital. VIII, p. 625.

Hab. in collibus ad *Poterium spinosum*, *Genisteas* et *Labiatas* suffruticosas, etc. parasitica, Peloponnesus (Bory), ad Trikala (Heldr!), et in monte Malevo (Orph!), Attica in Lycabetto (Heldr!), Creta (Sieb! Raul!), Archipelagus (Engelm.), Cyprus (Ky!), Cilicia (Bal!), Syria littoralis (Bl!) et ad Damascum (Gaill!), Palestina (Boiss!).

Hanc in Diagn. Ser. II, 3, p. 127 ad *C. planifloræ* varietates reduxi sed cl. Engelmann eam speciem distinctam esse autem; flores eis *C. planiflora* dimidio minores capitula diametro $1\frac{1}{2}$ -2 lineas tantum lata formantes, calyx corollæ vix brevior, corollæ lobi apice eucullati, styli ovario sublongiores.

Ar. Geogr. Sicilia.

* * Styli ovario æquilongi vel breviores.

+ Flores sessiles.

4. **C. brevistyla** (A. Braun in Schimp. exs. et Rich. Tent. Abyss. II, p. 79) caule ramoso, floribus sessilibus, calyce corollæ parum breviore fere ad basin in laciniis ovatas obtusas partito, corollæ laciniis ovatis expansis, squamulis brevibus tenuibus truncatis interdum bilobis, stylis capsulæ multo brevioribus stigmata subclavata subæquantibus ☐.

Hab. in monte Sinai Arabiae Petree (Botta ex Engelm.), Persiæ australi et Kurdistaniæ conterminâ (Haussk!), boreali in monte Totschal prope Teheran (Ky. 580!), Affghaniæ (Griff. 686 ex Engelm.), Assyriæ ad Tigridem (Noë 518 sub *C. elegans* ex Engelm.).

Ut cl. Engelmann animadvertisit a quibusdam *C. planifloræ* formis stylis brevibus tantum distinguenda.

Ar. Geogr. Abyssinia! Tibetia.

5. **C. globulosa** (Boiss. et Reut. Diagn. Ser. II, 3, p. 126), sæpius rubella, caulis tenuissime capillaris, floribus minutis sessilibus capitula minima pauciflora globosa basi bracteata formantibus undique papillosis, calycis purpurei breviter obconici corollæ tertiaræ parte brevioris ad tertiam partem fissi lobis triangularibus crassiuseculis, corollæ ad medium in lobos ovatos obtuse mucronatos fissâ fructiferâ ob lobos suprà capsulam connientes globosâ, stylis stigmate et capsulæ multo brevioribus ☐. *C. Balansæ* Boiss. et Reut. MSS.

Hab. ad Bozdaghi Tmoli occid. in Lydiâ *Genista Lydia* parasitica (Bal. 413!), in Tauro Cilicio supra Bulgharmaaden ad *Hyperica* (Bal!), in monte Alidagh Cappadociae (Bal!), in monte Ssahend Persiae borealis ad *Tragacanthas* (Buhse!), montibus Kellal et Ssebekuh Persiae australis (Haussk!).

Ex cl. Engelmann dubitanter forma *C. brevistylæ* sed omnino distincta videatur capitulis minimis paucifloris, calycis et corollæ lobi formâ, præsertim corollæ fructiferæ lobis non patentibus sed supra capsulam clausis. Facies et flores minutæ *C. Palestine* a quâ stylis brevissimis, etc., differt.

6. **C. Europæa** (L. Sp. 180) caulis ramosis filiformibus, flori-

bus sessilibus globoso-capitatis, calyce obconico basi carnoso lobis membranaceis ovatis obtusis, corollæ membranaceæ tubo calyce sublongiore lobis tubo brevioribus ovatis obtusis, squamulis parvis tenuissimis tubo adpressis paucierenatis stylis ovario magno stigmatibusque multo brevioribus ⊖. Ic. Rehb. Icon. V, tab. 497. — *C. major* D.C. — *C. brachystyla* G. Koch Linn. XXII. p. 747. — *C. hyalina* Boiss. in Bal. exs. non Roth. — *C. Schkuhriana* Pfeiff.

Hab. in *Urticâ*, *Cannabi*, *Salicibus*, *Umbelliferis*, *Galiis*, etc., probab. in totius regionis parte magis boreali. habeo ex montibus Peloponnesi (Heldr!), Mace-donie (Frvt!), Tauro Cilicico (Bal!), Ponto Lazico (Bal!), Armeniâ australi (Ky. 306!), montibus Persiae borealis (Ky!), ditionis Khorassau (Bge!), Tur-kestaniâ (Lehm!). Præterea in Anatoliâ, Caucaso, Tauriâ, Affghania ab Engel-mann, Steven, etc., indicata.

Ar. Geogr. Europa sept et media, India borealis, Africa borealis.

7. *C. Kurdica* (Engelm. Cusc. p. 21) caulis capillaceis, floribus sessilibus saepius tetrameris, glomerulis parvis paucifloris bracteâ ovatâ acuminatâ suffultis, calycis fere ad basin partiti lobis oblongis acutis tubum corollæ superantibus. corollæ tenuiter membranaceæ pellucido-punctatæ lobis ovato-lanceolatis acutis tubo sublongioribus, squamulis basi tubi adfixis parvis hyalinis tenuissimis truncatis tubo adpressis, stylis capsulâ et stigmatibus multo brevioribus ⊖. *C. alpina* Hohen. in Ky. exs.

Hab. in monte Gara Kurdistaniæ in *Daphnes* sp. parasitica (Ky. 588 b!), in montibus Schahu et Juanro ad nives 12000' (Haussk!).

Texturâ et facie affinis *C. Europæ* sed flores panciores, calycis et corollæ lobi acuti, corolla in fructu globosa totam capsulam arcte cingens nec tantum ejus verticem obtegens.

8. *C. Epilinum* (Weihe Prodr. Monast. 75) caulis filiformi-bus, floribus sessilibus in capitula densissima ebraeteata congestis, calyce corollâ adpresso et ejus tubo æquilongo in lobos ovatos latos profunde partito. corollæ tubo subgloboso. limbi tubo duplo brevioris lobis triangularibus acutis patulis, squamulis parvis tubo adpressis. stylis ovario multo brevioribus, stigmatibus stylo duplo longioribus tandem subelavatis ⊖. *C. densiflora* Soy. Willem. Rehb. Ic. V. tab. D. — *Epilinella cuscutooides* Pleiff.

Hab. in *Lino usitatissimo* parasitica, vidi ex Ægypto (Kralik!) ex regione Bachtiaris Persiæ (Haussk!) et probab. alibi.

Ar. Geogr. Insulae Canarienses, Europa fere omnis.

+ + Flores pedicellati.

9. *C. Babylonica* (Auch. MSS. — Choisy Mém. Soc. Phys. Gen. IX. p. 270, tab. 1) flava, caulis capillaceis ramosis elongatis laxe intricatis, floribus pedicellis eis saepius longioribus suffultis umbellatis, calyce hemisphærico-cupulato obtusissime et obsolete repando-

lobato corollâ subduplo breviore, corolle campanulatae ad medium fissæ lobis breviter ovatis obtusissimis, squamulis oblongis vix dentatis, stylis ovario stigmateque multo brevioribus ⊖.

Hab. in Babyloniâ ad Bagdad (Auch. 4120! et 3183!), ad Kutt juxta Tigridem in *Acacia* sp. (Noë 7 et 905!), ad Derbent i Basian Assyria et ad radices montis Avroman in *Vitice Agno casto* (Haussk!). in monte Gara Kurdistaniae (Ky. 588 a!).

Ex florum plurium observatione styli ovario manifeste et stigmata plus duplo breviores.

10. *C. elegans* (Boiss. et Bal. Diagn. Ser. II, 3, p. 129) rubella, caulis capillaceis brevibus intricatis, floribus pedicellis calyci aequilongis vel duplo longioribus suffultis solitariis vel 3-5 umbellulatis papilloso, calyce carnosò brevissime obconico subpentagono corollæ tubum aequante truncato obsoletissime quinquedentato, corollâ ultra medium in laciniis lanceolatas acutas valde patentas fissâ, squamulis parvis ovatis denticulato-laceris, antheris subexsertis, stylis capsula et stigmatibus longissimis multo brevioribus ⊖.

Hab. in regione alpinâ, mons Masmeneudagh Cappadocia in *Gennista* sp. et *Convolvulus compactus* parasitica (Balt.) in montibus Sawers et Nur Persie australis 12000' ad varias *Labiatas* suffrutescentes (Haussk!).

A *C. Babylonica* cui a el. Engelmann conjungitur calyce pentagone obconico, corolla laciniis triangulari-lanceolatis acutis præter colorem et ramos confer-tos certe distincta.

§ 2. *Epistigmæ* Engelm. — Styli tuberculo brevi constantes vel nulli, stigmata cylindrica vel subulata ovario saepius aequilonga. Capsula transverse et irregulariter rumpens.

* Flores sessiles vel subsessiles.

11. *C. Kotschyana* (Boiss. Diagn. Ser. I, 7, p. 29) caulis capillaceis, floribus sessilibus vel subsessilibus arete glomeratis bracteâ ovatâ suffultis, calycis scutellæformis lobis ovatis acuminatis corollæ tubum aequantibus vel superantibus acumine crassifuso, corollæ demum fauce constrictæ laciniis abrupte acuminatis acutis demum patulis, squamulis spathulatis profunde fimbriato-laceris majusculis faucem aequantibus capsulae apicem demum obtegentibus, stigmatibus cylindricis sessilibus, capsula tenuissimâ totâ corollâ persistente obtectâ, seminibus sub lente reticulatis ⊖. *C. Persica* Decaisne in Engelm. Cusec. p. 22!

Hab. in regione alpinâ montis Kuh Daëna Persie australis (Ky. 749!) et montium Avroman et Schahu (Haussk!) *Phlomidis* et *Marrubii* parasitica, ad Ispahan Auch. in Mus. Par. et ditione Khorassan (Bge!) in *Pharnopo*, prope Murtehehar in *Pegan*o (Bge!).

Species floribus mediocribus, insignis calyce corollæ proportione magno et ejus corollæque laciniis valde acuminatis, squamulis majusculis fimbriis capsule verticem obtegentibus. Stigma sessilia etiam in floribus speciminis typici

Aucheriani *C. Persicæ* semper vidi. Corolla firma persistens totam capsulam tenuissimam depresso obtegit.

* * Flores distincte pedicellati.

12. **C. pulchella** (Engelm. loc. cit. p. 24) caulis capillaribus, umbellis laxis paucifloris bracteâ ovatâ acuminatâ suffultis, pedicellis calyci aut flore aequilongis, calycis carnosus breviter obconici vix ad medium fissi lobis ovatis acutiusculis, corollæ tubo calyce vix longiori laciis tubo aequilongis ovatis acutis papillosum erectis, squamulis angustis fimbriatis incurvis faucem attingentibus, stigmatibus sessilibus ovarium æquantibus, capsulâ corollâ connivente totâ involutâ ⊖.

α. Affghana Engelm. loc. cit. — Tota splendide rubra.

Hab. in Affghanâ ad Tajie in *Artemisiis*, *Alhagi*, etc. (Griff. Journ. 770!).

Umbellæ 4-6-floræ, flores $1\frac{1}{2}$ lineas longi. Cl. Engelmann habet var. *β Altai-cam* in Rossiaustrali et Sibiriâ Altaicâ crescentem calyce profundius et in lacinias acutiores fisso.

13. **C. Arabica** (Fresen. pl. Eg. p. 95) caulis capillaceis ramiessimis intricatis, capitulis paucifloris, pedicellis calyci aequilongis vel longioribus, calyce subcarnoso cupuliformi ad medium in lacinias ovalis fisso, corollâ calyce vix duplo longiore semiglobosâ laciis ovatis acutis rectis, squamulis minimis dentatis, staminibus subexsertis, stigmatibus sessilibus ovario brevioribus ⊖. Ic. Mém. Soc. Phys. Gen. IX, tab. 1.

Hab. in Leguminosis, *Pegano*, etc., parasitica. Arabia Petrea (Rüppell, Bové, Schimp. 140!), in Egypto ad Alexandriam, Kahiram, Siout, Keneh, etc. (Ehrenb! Kralik!).

Planta Sinaitica paululum differt floribus brevius pedicellatis in capitulo pluribus.

SECT. II. GRAMMICA Engelm. — Styli duo. Stigmata capitata.

§ 3. **Eugrammicæ** Engelm. p. 28. — Capsula plus minus irregulariter circumcissa.

14. **C. Chinensis** (Lam. Encycl. II, p. 229) caulis tenuiter filiformibus, floribus in umbellulas 3-7-floras basi squamâ triangulari suffultas dispositis, pedicellis calycem æquantibus vel eo vix longioribus, calyce capsulari ad medium in lobos triangulari-ovatos fisso, corollâ calyce duplo longiore campanulatâ in lacinias erectas oblongas obtusiusculas infra tertiam partem partitâ, squamulis majusculis profunde fimbriatis, stylis ovario subbrevioribus, stigmatibus capitatis ⊖. Ic. Choisy loc. cit. tab. 3. — *C. hyalina* Wight Icon. 1372. — *C. ciliaris* Hohen. in Boiss. Diagn. Ser. II, 3, p. 129.

Hab. in Assyriâ prope Mossul in *Alhagi* parasitica (Ky. 631!), in Kurdistaniâ (Grant), Affghanîâ ad Candahar (Griff. 685! ex Engelm.).

Ar. Geogr. China, Asia tropica, Nova Hollandia.

15. C. hyalina (Roth Nov. Sp. p. 100 non Wight) albido-flavescens. caulis capillaceis valde intricatis, floribus in umbellas aut racemulos saepe compositos dispositis bracteolatis, pedicellis inæqualibus calyce 2-4-plo-longioribus. calyce obconico corollâ tertiatâ parte breviore ad medium in lobos triangulares aentos fisso. corolle ultra medium partitæ laciñiis lanceolatis acutissimis, squamulis nullis, stylis inæqualibus ovario longioribus demum divaricatis, stigmatibus globosis ⊖. *C. Arabica* Wight Icon. tab. 1371 non Fresen. — *C. oxy-petala* Boiss. Diagn. Ser. II, 3, p. 130. — *C. acutissima* Buchinger. — *C. Boissieri* Stocks Hook. Journ. Bot. 1852. p. 173.

Hab. in Belutschia inferiore *Trianthematis*, *Tribuli*. *Amaranthi* parasitica (Stocks 478!).

Ar. Geogr. India Orientalis, Abyssinia.

§ 4. **Clistogrammicæ** Engelm. p. 47. — Capsula baccata.

16. C. obtusiflora (H.B.K. Nov. Gen. III, p. 122) caulis capillaceis aurantiacis, capitulis paucifloris. pedicellis calyci æquilongis vel longioribus, calycis ad medium fissi tubo breviter obconico in pedicellum producto limbi lobis ovatis obtusis, corollæ calyce sublongioris tubo brevi scutellato lobis æquilongis ovatis obtusis mox reflexis, squamulis ovatis vel spathulatis, stylis brevibus crassis, ovario magno depresso, capsulâ depressâ baccatâ nudâ ad basin corollâ emarcidâ obsitâ ⊖.

Var. *breviflora* Engelm. — Flores eis typi submajores, saepe tetrameri, squamulæ minimæ saepe bifidæ. — *C. breviflora* Vis. Dalm. II, p. 231. — *C. Tinei* Inseng. — *C. aurantiaca* Req. — *C. chrysocoma* Welw. — *C. Rogovitschiana* Trautv. — *C. Cesatiana* Bertol.

Hab. in variis locis *Ocymo Basilico* parasitica, Græcia (Zucc. ex Eng.), Athenis (Orph!), Byzantii (Boissl).

Ar. Geogr. Lusitania, Gallia australis, Italia, Dalmatia, Serbia, Hungaria, Rossia australis; typus in Americâ et aliæ hujus speciei polymorphæ varietates in Indiâ, Novâ Hollandiâ.

SECT. III. MONOGYNA Engelm. — Stylus unicus. Capsula regulariter circumcissa.

§ 5. **Monogynellæ** Engelm. loc. cit. p. 74. — Stigmata capitata vel ovata. Corolla emarcida in capsulæ summo vertice remanens.

17. **C. monogyna** (Vahl Symb. II, p. 32. — Engelm. p. 76)

caule funiformi crassiusculo ramoso, floribus racemoso-spicatis, calycis laciniis ovatis imbricatis obtusis, corollæ sub anthesi cylindricæ tubo calyce vix longiore laciniis tubo dimidio brevioribus ovatis obtusis crenulatis erectis, antheris cordato-ovatis paulo sub fauce subsessilibus. squamulis medio tubo adnatis hippocrepidoformibus denticulatis. stylo stigmati subgloboso-bilobo æquilongo ovario globoso-ovato multo breviore, seminibus ovatis rostratis ⊖. Ic. Fl. Græc. tab. 257. — *C. Orientalis* Tourn. Cor. 45. — *C. astyla* Engelm. *C. Lehmanniana* var. *exquamata* Engelm. mihi eadem videtur squamulis magis adnatis.

Hab. ad arbores et frutices parasitica in regione totâ a Græciâ et Thraciâ! per Asiam minorem ad Caucasum et Transcaucasiam! Syriam! Persiam borealem et australem in alpibus (Ky. 743!), Affghaniam (Griff. ex Engelm.).

Flores $1\frac{1}{4}$ - $\frac{1}{2}$ lineas longi, capsula 2-3 lineas longa sœpius ovalis obtusa.

Ar. Geogr. Lusitania, Gallia australis, Serbia.

18. ***C. Lehmanniana*** (Bge Pl. Lehm. pag. 396) caule funiformi. floribus racemoso-spicatis, calycis laciniis ovatis obtusis imbricatis, corollæ tubo calyce duplo longiore limbi tubo dimidio brevioris laciniis oblongis obtusis crenulatis patentibus, squamulis in medio tubo adnatis rotundatis fimbriatis, antheris cordatis sub fauce insertis. stylo ovario multo breviore stigmate ovato bipartito æquilongo ⊖.

Hab. ad *Tamarices* et alios frutices in Turkestaniâ prope Buchara (Lehm!). Vulgo *Asgyschtsch*.

Flores $2\frac{1}{4}$ lineas longi. An a *C. monogyna* sat differt corollæ tubo longiore laciniisque patentibus nec erectis, squamulis margine excepto adnatis? Ejus varietas *esquamata* Engelm. p. 78 ad *C. monogynam* mihi reducenda videtur.

19. ***C. lupuliformis*** (Krocker Siles. p. 261. tab. 36) caule funiformi. floribus racemoso-spicatis, calycis laciniis ovatis obtusis interdum mucronatis. corollæ cylindricæ tubo calyce duplo longiore limbi laciniis tubo duplo brevioribus oblongis obtusis erectis. antheris oblongo-linearibus sub fauce sessilibus, squamulis brevibus bifidis ad partem inferiorem tubi adnatis. ovario ovato-conico in stylum gracilem stigmate globoso-ovato 3-4-plo longiore attenuato, seminibus triangulari-ovatis rostratis ⊖. *C. monogyna* Auct. Fl. Germ. non Vahl.

β . *Asiatica* Engelm. loc. cit. — Flores longiores et graciliores. laciniæ subpatulæ, squamulæ integræ late ovatæ.

Hab. in Caucaso (herb. Hooker ex Engelm.).

Facies omnino *C. monogyna* a quâ (an satis?) notis indicatis distinguitur.

Ar. Geogr. Germania, Austria, Hungaria, Rossia centralis. Varietas in Rossia austro-orientali. Songariâ. Sibiria Altaica.

20. ***C. gigantea*** (Griffith Notulæ I. p. 243) caulis filiformibus, floribus in spicas racemosas breves dispositis brevissime pedi-

cellatis, calycis partitionibus ovatis imbricatis amplis dimidium tubum corollinum æquantibus, corollæ laciniis patulis vel reflexis oblongis obtusis crenulatis tubo parum brevioribus, squamulis parti inferiori tubi insertis ejus medianam partem attingentibus, antheris magnis cordato-ovatis subsessilibus subexsertis, ovario conico, stylo stigmatibus oblongo-subligatis æquilongo ♂.

Hab. in faucibus Siahsung Affglaniae 40,500', ex cl. Engelm. *Tamarici*, ex meo specimine *Cichoraceæ* Sp. parasitica (Griff!).

Diffr. a precedentibus floribus majoribus $2\frac{1}{2}$ - $3\frac{1}{2}$ lineas longis, corollæ laciniis longis patentí-reflexis, stigmate.

Species dubia.

C. subuniiflora C. Koch Linn. XXII, p. 748.

Hab. in Ponto Lazico *Verbasci* parasitica (C. Koch!).

Forma junior non sat evoluta floribus solitariis vel 2-3 nis sessilibus *C. planiflora* vel *brevistyle*?

ORD. LXXXVI. BORRAGINEÆ.

(Juss. Gen. p. 128.)

Calyx liber persistens quinquetidus vel partitus. Corolla gamopetalæ hypogyna quinquefida regularis rarius inæqualis fauce nudâ vel pilis aut fornicibus obsitâ. Stamina quinque (rarius unicum fertile) corollæ inserta et ejus laciniis alterna. Ovarium rarius iudivisum, saepius bina bilocularia vel quatuor unilocularia uniovulata, ovaria pendula. Stylus unicus terminalis vel basilaris in medio ovariorum. Fructus rarius baccatus, saepius pyrenis binis bilocularibus vel ueuallis quatuor unilocularibus constans. Semina exalbuminosa, rarius tenuiter albuminosa, embryone recto.

Ta. I. CORDIEÆ p.c. — Ovarium indivisum stylo terminali.

Fructus baccato-drupaceus. Semina 4 exalbuminosa.

CORDIA (L. Gen. 256).

Calyx tubulosus vel campanulatus 5 vel irregulariter 3-8-dentatus. Corolla infundibuliformis limbo 5 rarius 6-8-partito. Stamina exserta vel inclusa. Ovarium quadriloculare ovulis solitariis pendulis. Stylus terminalis bis furcatus. Drupa pulposa endocarpio duro. Cotyledones longitudinaliter plicatæ. — Arbores vel suffrutices glabri.

1. **C. Myxa** (L. Sp. 273) arborea glabra vel seabriuscula, foliis longe petiolatis ovato-orbiculatis obtusis interdum mucronatis, floribus laxe cymosis polygamis, calycis breviter campanulati intus sericei dentibus brevibus ovatis, corollæ tubo calycem æquante lobis oblongo-linearibus recurvis, antheris ovatis exsertis, stigmatibus 4 flabellatim dilatatis erosionis, drupâ ovatâ mucronatâ ȝ. Ic. Delile Fl. Eg. tab. 19.

Hab. in Babyloniam circa Bassra et in insulâ Karrak Persie australis; an vere spontanea? culta in Cypro (Ky!), Ægypto (Delile), Arbor in Babyloniam Bombar dictus, fructus in Ægypto Mokhayet nominati.

C. crenata Del. Eg. tab. 20 in Ægypto quoque colitur et *C. subopposita* D.C. probab. in regno Mascate crescit.

Ar. Geogr. India.

EHRETIA (L. Gen. 257).

Calyx quinquepartitus. Corolla hypocrateriformis vel subrotata tubo brevi lobis ovatis imbricatis. Stamina saepius exserta. Stylus filiformis bifidus vel bipartitus. Ovarium bi vel quadriloculare. Drupa pulposa vel exsueca 2-4-pyrena. Cotyledones non plicatae. Frutices.

1. **E. obtusifolia** (Hochst. in Schimp. exs. — D.C. Prodr. IX, p. 507) fruticosa, ramis inermibus teretibus glabris glaucis, gemmis lateralibus et terminalibus folia 3-5 fasciculata cymamque ferentibus ebracteatis tandem in ramulum elongatis, foliis pubescenti-velutinis obovato-oblongis obtusis in petiolum brevem attenuatis, cymis breviter pedunculatis divaricatis filiformibus, calycis hispiduli laciniis breviter triangulari-oblongis obtusis, corollæ breviter campanulatae tubo calyce duplo longiore, lobis oblongis tubo æquilongis ȝ. *E. obovata* R. Br. in Salt (nomen solum).

Hab. in Belutschia (Stocks exs. 539!).

Ex specimine unico descripta cum plantâ Abyssinica congruere videtur.

Ar. Geogr. Abyssinia, Scinde.

2. **E. floribunda** (Royle Ill. p. 306) arborea, ramis teretibus canescensibus glabris, gemmis lateralibus et terminalibus folia solitaria vel bina cymasque sessiles a basi divaricatum dichotomas edentibus, foliis ovatis vel oblongis basi cuneatâ breviter petiolatis glaucis puberulis tandem glabratis, calycis velutini laciniis ovatis obtusis, corollæ tubo calycem parum excedente lobis obovato-oblongis tubo æquilongis ȝ.

Hab. in Belutschia (Stocks exs. 607!).

Plantam Royleanam non vidi, Brahuica a cl. Stocks determinata *E. obtusifolia* valde affinis est.

Ar. Geogr. India borealis.

TR. II. HELIOTROPEÆ. — Ovarium stylo terminali et suturis 4 præditum. Fructus drupaceus vel siccus, maturitate in 2 pyrenas vel nuces 4 aut abortu pauciores secedens. Semina tenuiter albuminosa.

TOURNEFORTIA (L. Gen. 192).

Calyx quinquepartitus. Corolla hypocraterimorpha fauce nuda. Stamina inclusa. Stylus simplex. Stigma a basi annulo carnoso cincta indivisum vel 2-4-lobum. Fructus drupaceus bicarpellaris in 2 pyrenas biloculares tandem solutus.

1. **T. Arguzia** (Linn. fil. Suppl. 132 sub *Messerschmidia*) pilis adpressis basi subtuberculatis hirsuta pallide virens e rhizomate duro pluricaulis, caulinibus ascendentibus foliosis simplicibus vel parce ramosis, foliis sessilibus oblongis vel lanceolatis basi attenuatis obtusis, spicis brevibus scorpioideis in corymbum terminalem dispositis, calycis corollæ albae tubo sericeo 3-4-plo brevioris laciniis lanceolatis obtusis, limbi ampli lobis ovatis, antheris oblongis acutis, stylo brevi, stigmate breviter pileiformi, drupâ ovatâ apice subquadrilobâ adpresso hirtâ ♀. *T. Arguzia* R. et Sch. Syst. IV, p. 540.

Hab. præsertim in littoribus arenosis maritimis, Asia minor (Auch. 23521), Tauria ad Sulak (Stev., Rehm!), in Caucasiam ad Caspium prope Petrowsk (Rupr!), prope Kisilagatsch et Lenkoran (Hohen!), ad littora Ghilani (Buhse!).

Ar. Geogr. Rossia media, Turcomania, Sibiria, Mongolia Chinensis, China borealis, Japonia.

HELIOTROPIUM (L. Gen. 139). — *Heliophytum* D.C.

Calyx quinquepartitus persistens vel cum fructu deciduus. Corolla infundibuliformis vel hypocraterimorpha limbo quinquelobo vel partito saepe 5 laciniis intercalaribus e sinubus ortis aucto Antheræ subsessiles connectivo interdum in apiculum terminalem abeunte. Stylus simplex. Stigma a basi latiori annulo carnoso cincta breviter pileiforme, vel conicum, vel subulato-elongatum, integrum vel bifidum. Nucule quatuor (interdum abortu pauciores) distinctæ vel diutius cohærentes, vel per paria connatae, vel rarius in caryopsem monospermam concretæ. — Herbæ vel suffrutices. — Multa ex egregiâ amic. a de Bunge monographiâ Bull. Mosc. 1869 exscerpsi, nonnullis tamen in Sectionum et specierum delimitatione mutatis.

Specierum Orientalium dispositio.

SERIES A. — ANNUA.

* Corollæ lobi aestivatione imbricati.

+ Calyx cum fructu deciduus.

SECT. I. PIPTOCHLÆNA Don. — Nuculæ in caryopsem unicam concretæ.

*H. spinum.*SECT. II. HEDYOSMA Bge. — Nuculæ quatuor (abortu interdum 2-3).
Stigma conico-elongatum.*H. crassifolium, Noëanum, calcareum, Chorassanicum.*

+ + Calyx nuculis deciduis persistens.

SECT. III. AGORÆA Bge. — (*Orthostachys* Bge ex parte.) — Stigma ex basi conicâ attenuatum.

× Racemi basi bracteati.

H. Aleppicum, Kunzei.

× × Racemi ebracteati.

H. Europeum, Bocconi, Eichæaldi, lasiocarpum, pallens, Szovitsii, dissitiflorum, gracillimum.

SECT. IV. PILOSTYLIS Stev. — Stigma pileiforme.

H. suaveolens, villosum, confertiflorum, Agdense.

** Corollæ lobi aestivatione margine plicati vel convoluti irregulariter et subvalvatim conniventes, recti, incurvi vel circinnati.

SECT. V. BUCANION Sczegl. — Calyces fructiferi immutati persistentes.

× Corollæ limbus quinquelobus.

H. grandiflorum, Griffithii, biannulatum, Bovei, Mamamense, Teheranicum.

× × Corollæ limbus ob lacinias intercalares decemlobus.

H. circinatum, Mesinanum, samoliflorum, Sultanense.

SECT. VI. PSEUDONOSMA Bge. — Calyx fructifer auctus.

H. murranthos, acutiflorum.

SERIES B. — PERENNIA.

SECT. VII. CATOXY'S Bge. — (*Catimas* Alph. D.C. ex parte.) —
Corollæ lobi angusti acuti aestivatione margine involuti
subvalvatum conniventes.

× Corolla intus hirta.

H. Transoxanum, Kumense, trichotomum.

× × Corolla intus glabrae.

H. dasycarpum, minutiflorum, eremobium, Halane, luteum, Aucheri, Carmanicum.

SECT. VIII. RADULA Bge. — (*Heliophyti* Sect. *Pterotropium* D.C. — *Orthostachys* ex parte.) — Corollæ lobi ovati vel oblongi aestivatione imbricati.

× Corolla intus hirta.

H. Cabulicum, Affghanum, strigosum, rariflorum.

× × Corolla intus glabra.

H. rotundifolium, Radula, brevilinæ, ophioglossum, denticulatum, Sogdianum, Arbainense, undulatum, Persicum, tuberosum.

SERIES A. — ANNUA.

SECT. I. PIPTOCHLENA.

* Corollæ lobi aestivatione imbricati.

1. **H. supinum** (L. Sp. 187) annum, pube ramorum patenti foliorum adpressâ velutinum cinereum, collo pluricaule, caulinis prostratis centrali saepè erecto omnibus dichotome ramosis, foliis omnibus petiolatis ovatis obtusis facie superiori plicato-lineatis, racemis densifloris lateralibus et terminalibus, floribus minutis, corolla extus hirtæ tubo calyce subbreviore lobis ovatis, antheris linearibus, calyce hirsuto fructifero lobis conniventibus fructum amplectente et cum eo caduco, stigmate elongato-conico stylo æquilongo, nuculâ magnâ unicâ glabrâ hinc planâ illine convexâ ⊖.

Hab. in arenosis et lutosis hyeme inundatis Graeciae (Sprun! Heldr!), Archipelagi! Macedonie (Friv!), Crete (Heldr!), Anatoliae occidentalis et australis! Syriae littoralis (Bl!), Ægypti mediae et superioris! Mesopotamiae (Ky!). Babylonie ad Bagdad (Noë!), Transcaucasia ad Caspium (Lebed.).

Ar. Geogr. Regio mediterranea Europæ et Africæ borealis, Senegalia, Abyssinia.

SECT. II. HEDYOSMA.

2. **H. crassifolium** (Boiss. Diagn. Ser. II, 3, p. 131) annum totum pilis albis adpressis dense tomentosum canum, caule elato dichotome ramoso, foliis breviter petiolatis ovatis obtusis basi rotundatis vel truncatis superne eximie lineato-depressis subtus elevatim nervosis, racemis demum longissimis laxis, calycis velutini laciniis oblongo-linearibus obtusis. corollæ tubo calyce subdupo longiore extus retrorsum velutino intus minute hirtulo, limbi tubo triplo brevioris lobis ovatis sinubus edentulis, antheris prope corollæ basin insertis longe linearibus connectivo apice in apiculum introflexum abeunte, stylo subnullo, stigmate a basi latiori longe linearis hirto, nuculis oblongis laeviusculis glabris ⊖.

Hab. in subalpinis Persiæ occidentalis prope Haneky (Chanekin) et versus Kermanchah (Noë 1001! et 1039!).

Sesquipedale, folia inferiora saepè sesquipollicaria, corolla tres lineas longa, facies *H. rotundifolii* radice perenni, etc., distinctissimi. Species antheris longe linearibus insignis.

3. **H. Noëanum** (Boiss. Diagn. Ser. II, 3, p. 132) annum totum pilis longis adpressiusculis tomentosum canum, caulibus elatis dichotome ramosissimis, foliis ovato-rotundatis obtusis breviter petiolatis superne depresso-lineatis subtus elevatim nervosis, spicis simplicibus vel conjugatis demum elongatis laxifloris, calycis sessilis tomentosi laciniis oblongis obtusis, corollæ extus hirtæ intus glabrae tubo calycem non excedente limbo infundibuliformi concavo lobis semiorbiculatis, sinubus laciniis intercalaribus acutissimis inter se apice contiguis faucom claudentibus obsitis, antheris paulo infra medium tubum insertis faucom attingentibus breviter linearibus connectivo apice in apiculum intus flexum abeunte; stigmate a basi latiori linearis hirto stylum hirtum longitudine superante, nuculis glabris rugulosis ⊖.

Hab. in Persiæ occidentali prope Haneky (Noë 981! Auch. 5003!).

β. edentulum. — Corollæ sinus intus plicati laciniis intercalaribus obsoletis

Hab. in cultis vallis Chyrsan Susianæ, prope Bebehan, Masibin, circa Sulimanieh, ad radices montium Avroman, Schahu, Pir Omar Gudrun 3-6000' (Haussk!).

Pedale vel sesquipedale, folia valde pannosa, facies *H. crassifolii* antherarum formâ quoque affinis et a quo differt corollâ dimidio minore vix $1\frac{1}{4}$ lineam longâ ejus tubus calycem non excedit, stylo. Corollæ limbus variat tubo non multum brevior vel brevissimus, laciniis intercalaribus manifestis vel plicæformibus fere obsoletis.

4. **H. calcareum** (Stocks Hook. Journ. Bot. IV (1852) p. 174) annum totum pubescenti brevi densâ adpressâ basi minute tuberculata

canescens, caule erecto, foliis longiusculæ petiolatis ovatis acutiusculis basi subcuneatis subitus prominule nervosis superne non lineatis, spicis solitariis et conjugatis demum valde elongatis remotifloris, calycis tomentosi laciniis linear-i-oblongis tubum corollæ aequantibus, corollæ tubo piloso intus glabro limbi tubo triplo brevioris lobis late et breviter ovatis, sinuum dentibus accessoriis ovatis introflexis, antheris paulo infra medianam corollam insertis linearibus connectivi apiculo introflexo, stylo a basi conice attenuato-elongato hirto stylo hirto duplo longiore, nuculis hispido-pilosis apice acuto ancipitibus ⊖.

Hab. in collibus Belutschiae inferioris (Stocks exs. 630!).

Affine *H. Noëano* differt foliis acutioribus non depresso-lineatis, limbi corollæ dentibus accessoriis pliciformibus obtusissimis, nuculis hirsutis acutis. Corolla extus ad faucem foveolis 5 ex plicis introflexis ortis insculpta.

5. H. Chorassanicum (Bge loc. cit. p. 23) annum molliter et patule villosum, caulis elongatis diffusis, foliis longe petiolatis cordato vel ovato-subrotundis obtusis superne lineato-plicatis, spicis 2-6-nis confertifloris, calycis pedicellati patule hirsuti laciniis linearibus corollæ tubo subbrevioribus, corollæ tubo extus retrorsum piloso intus hirto, limbo patentissimo tubo subdupo breviore, antheris oblongis supra basin insertis, stigmate conico longe subulato-attenuato glanduloso-hirto stylo glabro 2-2 $\frac{1}{2}$ -plo longiore, nuculis minutis oblongis velutinis ⊖.

Hab. in prov. Khorassan Persiæ prope Nichapur, Achmedabad et Affghania prope Herat (Bge!).

β. leiocarpum Bge loc. cit. — Caules breves erecti. corolla major limbo latiore, nuculæ glabræ.

Hab. in montibus argillosis Tabat Kuh ad orientem urbis Mesched (Bge!).

Caulis in typo pedalis et longior, in varietate humilis. In typo corollæ tubus 3 lin. longus limbus diam. 2 $\frac{1}{2}$ lineas latus, in varietate limbus 3 $\frac{1}{2}$ -4 lineas latus. Estivatione quincunciale! Facies *H. Ajdensis* et *confertiflori* a quibus stigmate subulato-attenuato statim distinguitur.

SECT. III. AGORÆA.

× Racemi bracteati.

6 H. Aleppicum (Boiss. Diagn. Ser. I, 11, p. 88) dense et patule setis albis basi tuberculatis tomentoso-strigulosum a basi dichotome ramosissimum, foliis oblongis et oblongo-linearibus acutis basi attenuatis inferioribus breviter petiolatis, spicis unilateralibus laxiusculis parte inferiori bracteatis, floribus parvis, calycis extus albostrigosissimi intus albi laciniis anguste linear-i-lanceolatis, corollæ extus hirsutæ intus glabré tubo calyce breviore, limbi tubo subbrevioris lobis ovatis erosulis sinibus denticuliferis, antheris linea-

ribus paulo supra tubi basin insertis, stylo subnullo, stigmate a basi conicâ subulato apice albo-penicillato, nuculis tuberculatis glabris ⊖.

Hab. in planicie elata inter Antiochiam et Aleppum (Boiss!), in rupestribus calcareis ad Magdouché prope Sidonem (Gaill!), in Cataonia circa Malatia et Adamian (Haussk!).

Planta 3-5-pollicaris sed interdum elongata pedalis, folia 8-12 lineas longa 3-4 lata. calyx persistens, corolla vix lineam longa.

H. Kunzei (Lehm. Ic. St. p. 19, tab. 29) annum pilis tuberculatis aliisque minutis omnibus adpressis sericeum. caulinibus decumbentibus, foliis ovato-ellipticis inferioribus breviter petiolatis, spicis 2-3-nis densifloris secundis ebracteatis, floribus minimis. calycis sericeo-hispidi laciniis oblongo-lanceolatis, corollæ extus sericeo-hispidae tubo calyxi æquilongo limbi parvi lobis ovatis, antheris ovato-oblongis infra medium tubum insertis apice puberulis, stigmate sessili breviter conico hirtulo, nuculis subglobosis hirsutis ⊖. *H. syenites* Spr. Syst. I, p. 388. — *H. Niloticum* Alph. D.C. Prodr. IX, p. 341.

Hab. in Egypto ad Syenem (Sieb!), ad Esneh (Boiss!).

Valde affinis *H. ovalifolio* Forsk. Descript. p. 38. — *H. Coromandeliana* Retz quod facie simillimum differt tantum antheris stigmateque sublongioribus, ambo probabilissime in unicam speciem conjungenda sunt.

Ar. Geogr. Nubia, Abyssinia, Senegalia.

× × Racemi ebracteati.

H. Europeum (L. Sp. 187) annum viride pilis brevibus adpresso-basi tuberculatis aspernatum, caule erecto ramoso, foliis petiolatis ellipticis obtusis, spicis solitariis vel geminatis densifloris, calycis hirti laciniis linearibus fructiferis patulis, corollæ tubo hirto calycem tenuiante limbo tubo subbreviore, antheris in parte inferiori tubi insertis, stigmate elongato-conico basi tenuissime puberulo cæterum glabro stylum glabrum brevem multo superante, nuculis glabris vel parce hispidulis serobiculato-rugosis ⊖. Ic. Jacq. Austr. 207. — Rehb. tab. 93. — *H. subcanescens* Stev. loc. cit. p. 364.

Hab. in ruderatis et cultis Graecie in Atticâ et Eubœâ (Heldr!), Tauriæ (Stev!), Caucasi (M.B., Ledeb.).

H. tenuiflorum. — Magis canescens. corollæ limbus minor erectiusculus nec patulus simum dentibus majoribus, stigma magis pubescens, nuculae magis hispidæ. — *H. tenuiflorum* Guss. Syn. I, p. 216. — Fl. Inar. tab. 7. — *H. commutatum* C. Koch Linn. XXIII, p. 628! — *H. Carduchorum* C. Koch loc. cit!

Hab. in ruderatis et cultis, in ditione tipo frequentius et magis australe, Graecia in insula Cephaloniâ (Heldr!), Thessaliâ (Orph!), Rhodus (Bourg!), Armenia Turcica (Ky!), Mesopotamia ad Mossul (Ky. 659!), ad Diarbekir et Mardin (Ky. 373!), Syria ad Damascum (Gaill!). Egyptus ad Alexandriam (Gaill!), Assyria et Babylonia. Persia australis (Haussk!), Persia borealis (Buhse!).

Hanc formam que in Europae regione mediterranea australiori fere omni occurrit ingenti specimini numero comparato ab *H. Europeo* typico specifice distinguere nequeo. In *H. Carduchorum* specimine typico staminum insertionem eamdem quam in typo observo.

Ar. Geogr. Europa media et australis a Belgio et Germania media ad regionem Danubialem et Rossiam australem, Africa borealis.

9. H. Bocconi (Guss. Ind. Bocc. 1825 — Prodr. p. 205) annum viride, pilis brevibus adpressis basi minute tuberculosis obsitum, caule erecto ramoso, foliis petiolatis ovatis obtusis, spicis solitariis et geminatis, calycis hirsuti laciniis linearibus corollae tubo subbrevioribus apice sub anthesi squarrosis fructiferis patulis, corollae tubo hirsuto limbi ampli tubo longioris laciniis obtusissimis, antheris ad partem inferiorem tubi insertis, stigmate a basi conica longe subulato-attenuato stylo triplo longiore utroque glaberrimis; nuculis majusculis glabris valde rugosis ☽.

Hab. in insula Cephalonia et in Etolia circa Missolonghi (Heldr!).

Ab *H. Europeo* floribus odoris, corollae limbo duplo maiore, fauce sensim nec abrupte ampliata, nuculis majoribus, ab *H. suaveolenti* cui facie et corollis magnis affinis stigmate longe subulato-attenuato nec breviter conico diversum.

Ar. Geogr. Sicilia.

10. H. Eichwaldi (Steud. Nom. — n.c. Prodr. IX, p. 535) annum breviter et adpresso canescens, caule erecto dichotome ramoso, foliis ovato-ellipticis obtusis basi rotundatis petiolatis minutissime et crenate tuberculatis. spicis solitariis et conjugatis, calycis laciniis oblongis tubum corollae extus hirta aequantibus fructiferis fructu applicatis, corollae sinibus dentiferis fauce glabra, antheris sub medio tubo insertis, stigmate elongato-conico hispidulo, stylo subnullo, nuculis glabris minutissime verruculosis ☽. *H. ellipticum* Ledeb. in Eichw. Casp. tab. 4, p. 6 non Salt. — *H. strictum* Ledeb. Fl. Ross. 3, p. 100. — *H. littorale* Stev. Obs. p. 565. — *H. dolosum* Notar. Repert. Lig. p. 319. — *H. macrocarpum* Guss. Inarinn. p. 214, tab. 7. — *H. Stevenianum* Andr. — *H. parviflorum* Stev. in litt.

Hab. in littoribus maritimis et quoque in interioribus, Cappadocia ad Caesaram (Bal. exs. sub *H. tenuifloro*!). Cilicia Trachea (Pérón!), Tauria in littore ad Sudak et prope Sympheropol (Stev!), insula Syra, Attica ad Phalerum et Phocis ad Amphissam (Heldr!), littora peninsulae Athoae (Janka!), littora Caspia (Eichw!), Transcaucasia (Hoh.), Persia borealis et Afghanistan (Szov! Bge!), Kurdistania Persica ad Awiheng (Haussk!).

Ar. Geogr. Italia, Serbia, Asia media, Songaria.

11. H. lasiocarpum (F. et M. Ind. Petrop. IV, p. 38) annum piloso-villosum, caule erecto, foliis petiolatis ovato-ellipticis vel subrotundis sublineatis minutissime et crenate tuberculatis, spicis solitariis et conjugatis densifloris, calycis laciniis oblongo-linearibus corollae tubo extus hirto sublongioribus, limbo corollae tubo subbreviore den-

tibus sinuum minimis, antheris sub medio tubo insertis, stigmate subsessili elongato-conico hispidulo, nuculis hispido-velutinis parce rugulosis ♂.

Hab. in littore orientali Caspii (Hansen), Turkestaniā (Lehm.), Persiā boreali et ditione Khorassan (Bge!), Affghaniā prope Herat (Bge!), prope Kabul (Griff. exs. 5931!), regnum Masecate Arabiæ (Auch. 5005!).

Ab affini *H. Eichwaldi* hirsutie majori patula, nuculis pubescentibus, distinctum.

Ar. Geogr. Arabia felix, India boreali-occidentalis.

12. *H. pallens* (Del. pl. Afriq. Caill. p. 69, tab. 64) annum molliter tomentosum, caule elato, foliis petiolatis ovatis obtusis tenuiter nervosis, spicis 2-4-nis elongatis densifloris, calycis hirsuti lobis linearibus corollæ tubum æquantibus, corollæ extus hirtæ fauce nudâ lobis suborbiculatis, antheris in parte inferiori tubi insertis oblongo-linearibus muticis, stigmate elongato-conico bifido stylo hirto breviore, nuculis glabris minute tuberculatis ♂.

Hab. in Egypto superiori ad Esneh (Acerbi ex D.C.).

Sesquipedale et elatius, spicæ 3-5-pollicares, corolla 2 $\frac{1}{2}$ lineas longa.

Ar. Geogr. Nubia, Africa tropica.

13. *H. Szovitsii* (Stev. Bull. Mosq. 1854, I, p. 568) annum pallide virens, caule subdivaricatum ramoso adpressissime hirto, foliis superne glabriusculis subtus pilis minimis adpressis e tuberculis ortis aspersis ovatis et oblongis obtusis petiolatis, spicis 2-3-nis elongatis laxifloris, calyeis pedicellati parce et adpresso strigulosi corollæ triplo brevioris lacinias linearis-lanceolatis, corollæ intus extusque glabrâ infundibuliformi tubo in limbum eo longiore breviter late et obtusissime quinquelobum sensim ampliato, antheris prope basin corollæ sitis breviter linearibus apiculo terminali introrsum flexo, stylo brevissimo, stigmate longe conico hirto, nuculis levissimis glabris ♂. *H. Kowalewskyi* Stegl. — *Bucanion Szovitsii* Stev. loc. cit. p. 567.

Hab. in Armeniā Rossicā ad Nackitcewan et in prov. Aderbidjan (Szov. 428!), prope Ordubad (Kowal!).

Pedale, corolla demum tres lineas longa. Facies *H. grandiflori* et *circinnati* sed prater inomentum sparsum adpressum corollæ lobis ovatis brevibus et aestivatione imbricata ab eis alienum, corollæ ampliæ infundibuliformi glaberrimâ ab omnibus distinctum.

14. *H. dissitiflorum* (Boiss. Diagn. Ser. I, 11, p. 88) annum subcinerascens totum pilis brevissimis adpressiusculis glandulisque sessilibus in tuberculos albos minimos demum mutatis obsitum asperulum, caulibus virgatis ramosis, foliis parvis petiolatis ovatis vel oblongis obtusis, spicis gracilibus demum valde elongatis dissitifloris uniseriatis, calycis sessilis lacinias linearis-oblongis obtusis corollæ

extus hirtæ intus glabré tubo æquilongis, limbo tubo plus duplo breviore minimo decemlobo lobis exterioribus obovato-orbiculatis internis intercalaribus ovatis dimidio minoribus conniventí - incurvis, stigmate elongato-conico stylum retrorsum hispidum longitudine vix superante, antheris vix infra medium tubum insertis oblongo-linearibus apiculo uncinatim introrsum flexo, nuculis lèviusculis glabris ♂.

Hab. in lapidosis prope Sonak ad radices montis Elbrus (Ky. 583 a!).

Pedale vel sesquipedale, racemi tenues 6-8-pollicares, flores minimi, corolla vix lineam longa. Èstivatio manifeste quincuncialis.

15. **H. gracillimum** (Bge Heliotr. p. 16) annum minutum stri-gilloso-canescens, caule erecto parce ramoso, foliis ellipticis utrinque rotundatis breviter petiolatis, spicis simplicibus gracillinis. calycis cano-villosi laciniis oblongis corollæ tubo pubescente brevioribus fructiferis patentibus, limbi sinuum denticulis triangularibus, antheris infra medium tubi insertis ad faucem prominulis divaricato-hiapiculatis, stylo brevissimo hispido, stigmate glaberrimo e basi disciformi elongato subulato brevissime bilobo, nuculis minutis glabris levibus ♂.

Hab. ad Salinas Mamam Persiae in consortio *H. Mamamensis* (Seidlitz). Non vidi.

Ex cl. Bunge nulli affine, spicarum formâ *H. longiflorum* æmulat. Planta 2 ½-3 ½-pollicaris, spica 1 ½-2-pollicaris, flores 1 ½ lin. longi.

SECT. IV. PILOSTYLIS.

16. **H. suaveolens** (M.B. Taur. Caue. III, p. 116) annum pilis brevissimis basi minute tuberculatis obsitam pallide virens, caule erecto dichotome ramoso, foliis ovatis obtusis petiolatis, spicis 2-4-nis densifloris. calycis hirsuti laciniis linearibus corollæ tubo hirsuto subbrevioribus fructiferis erecto-patulis, corollæ limbo tubo æquilongo sinibus obsolete erosio-dentalis, antheris sub medio tubo insertis, stigmate e stylo brevissimo breviter et acute conico utroque glabris, nuculis verrucosis glabris ♂. *H. intermedium* Andr. — *H. Turcicum* Friv. Fl. 1836, p. 436. — *H. odoratum* Stev. Bull. Mesq. 1851, p. 567.

Hab. in collibus et lapidosis, Macedonia (Friv!, Orph!), Thracia ad Philippopolin (Janka!), Anatolia in valle Meandri (Boiss!), Isauria (Heldr!), Lycia (Bourg!), Cappadocia ad Cæsaream (Bal!), Cilicia (Ky. 263!), Armenia Rossica (Szov!), Tauria (Rehmt!), Daghestaniæ (Rupr!), Transcaucasia (C. A. Mey., Hohl), Persia bor. (Buhse!), Assyria (Haussk!).

Corollæ magnæ et facies *H. Bocconi*, stigma diversissimum.

17. **H. villosum** (Willd. Sp. I, p. 741) annum pilis brevibus

adpressis et longioribus patulis molliter tomentosum, caulinis elongatis erectis et decumbentibus ramosis foliis petiolatis ovatis obtusis, spicis solitariis et conjugatis densifloris tandem longis, calycis villosi corollæ tubo hirsuto brevioris laciniis linearibus fructiferis erectiusculis, corollæ tubo intus glabro vel hirto costato costis infra faveum in dentem abeuntibus, limbo ampio patentí, antheris infra medium tubum adnatis, stigmate subsessili glabro conico-pileiformi, nuculis glabris tuberculatis ⊖. Ic. Desf. Cor. tab. 16.

Hab. in cultis Cephaloniae (Heldr!), Peloponnesi (Heldr!, Sprun!), Atticæ (Spr!. Heldr!), Archipelagi (Tourn., Auch!, Bourg!), Cretæ (Sieb!), Byzantii (Grisel!), Lydie (Boiss!), Cataoniae (Haussk!), Syrie littoralis (Ehrenb!, Bl! Gaill!), Palestine (Gaill!).

β. brevilimbe. — Corollæ limbus diminutus cæterum typo similimum.

Hab. in Syriâ ab Tripoli (Boiss!).

Species floribus magnis, tubo intus costis in denticulos abeuntibus insignis. Corollæ tubus variat intus hirtus vel glaber.

18. *H. confertiflorum* (Boiss. et Noë Diagn. Ser. II. 3, p. 132) annum patule et dense tomentoso-canescens, caule elongato ramoso, foliis breviter petiolatis ovatis, spicis 2-3-nis longis densissimis, calycis hispidi laciniis linearibus corollæ tubo hirsuto subbrevioribus fructiferis erectis, corollæ limbo minuto tubo duplo breviore lobis ovatis, tubo intus non costato, antheris prope basin corollæ insertis, stylo subnullo glabro, stigmate conico-pileiformi hispidulo, nuculis minimis glabris ruguloso-tuberculatis ⊖.

Hab. in subalpinis ad Haneky Persia austro-occidentalis (Noë exs. 4040!).

Caules empi spicis pedales demum indurati, folia eis *H. villosi* minora, nuculae duplo minores, flores minores adpressissimi, corolla intus costis destituta.

19. *H. Agdense* (Bunge loc. cit. p. 19) annum patule villosum pallide virens, caule pumilo divaricatim ramoso, foliis petiolatis ovatis basi et apice rotundatis, spicis solitariis vel conjugatis sub anthesi abbreviatissimis tandem parum elongatis laxis, calyce fructifero pedicellato corollæ tubo glanduloso-hirto breviore laciniis linearibus, corollæ tubo intus non costato limbo patentissimo tubo subbreviore, antheris prope corollæ basin insertis, stigmate pileiformi glabro, stylo brevissimo hispidulo, nuculis oblongis glabris levissimis ⊖.

Hab. in margine australi et occidentali deserti salsi magni Persiae prope Agda et Schenrab (Bge!).

Planta florifera 4½-3 pollices, fructifera 6 pollices longa. Folia 7-8 lineas longa, racemi fructiferi cum pedunculo bipollicares, flores 3 lineas longi. Affine *H. confertifloro* a quo notis in diagnosi indicatis differt

* * Corolla lobi aestivatione plicati subvalvatim conniventes

SECT. V. BUCANION.

* Corolla limbis quinquelobus.

20. **H. grandiflorum** (Auch. exs. D.C. Prodr. IX. p. 532) annum molliter et adpresso villosum canescens, caule erecto ramoso, foliis petiolatis ovatis obtusis, specie solitariis vel conjugatis demum elongatis laxiseulis, calycis hirsuti corollæ quintuplo brevioris laciniis lanceolatis obtusis, corolla extus hirtæ intus glabra tubo tenui sensim in fauce elongatam infundibuliformem ampliato, limbi lobis elongatis oblongis aestivatione margine convolutis apice subinflexis, sinibus edentolis, antheris linearibus paulo supra corollæ basin sitis apieno brevi intus flexo, stylo subnullo, stigmate a basi latiore longe subulato hirto, nuculis vix rugosis glabris ⊖.

Hab. in Armenia (Auch. 2376!).

Caulis 6-7-pollicaris, prob. quoque longior, calyx linea longus, corolla 5 lineas longa

21. **H. Griffithii** (Boiss. Diagn. Ser. II, 3. p. 131) annum molliter hirsutum, caule dichotome ramoso, foliis petiolatis ovatis obtusis nervis subtus prominentibus, racemis crassiuseulis densis, calycis dense tomentosi laciniis linearibus, corollæ calyce duplo longiore anguste tubulosâ extus et intus tomentellâ limbi minutissimi lobis 3-oblongis erecto-patulis, antheris versus corollæ basin insertis connectivo brevissime apiculatis, stylo stigmate glabro breviter conico sublongiore, nuculis oblongis glabris levibus ⊖.

Hab. in Affghania: ruderatis arcis Gulgula et in collibus Barrea (Griff!).

Inter affinia corollæ angustæ et longe tubulosâ limboque parvo insignis, ejus lobi aestivatione distincte longitudinaliter plicati sunt et juxtapositi subcontorti.

22. **H. biannulatum** (Bunge loc. cit. p. 22) annum molliter tomentoso-vilosum, foliis petiolatis ovatis utrinque obtusissimis, specie denique elongatis dense et biseriatim floriferis, calycis laciniis linearibus villosissimis dimidium corolle tubum aequantibus, corolla tubo extus villoso intus supra antheras pilis subreversis hispidulo abhinc glabro et rursus ad faucom annulo denso pilorum erectorum obsito, limbi brevis lobis oblongis obtusis, antheris oblongis versus corollæ basin insertis, stylo breviter conico glabro stylum glabrum longitudine superante, nuculis ovatis minutis tenuissime et seriatim punctulatis ⊖.

Hab. in regno Cabulico (Griff. Cat. 5933). Non vidi.

Ex descriptione *H. Griffithii* valde affine videtur.

23. **H. Bovei** (Boiss. Diagn. Ser. I, 11, p. 87) annum pilis albis adpressissimis canescens vel canum a basi dichotome ramosum, foliis petiolatis ovatis obtusis superne lineato-depressis subtus elevatim nervosis, spicis solitariis vel conjugatis demum elongatis uniserialibus laxis, calycis albo-tomentosi laciniis lanceolatis obtusis corollæ tubo hirto subæquilongis, corollæ intus glabræ limbo tubo æquilongo lobis ovatis obtusis sinibus vel denticulis minimis instructis vel omnino nudis, antheris infra medium tubum affixis faucem attingentibus breviter linearibus acumine intus flexo, stylo subnullo, stigmate longe conico-attenuato superne hispido, nuculis tuberculato-rugosis minute hispidulis vel glabris ⊖. *H. Prusianum!* *H. Smyrnæum!* et *H. Kotschyanum!* Bge loc. cit. p. 33 et 34.

Hab. in cultis Syriæ prope Sidonem (Bl!, Gaill!), ad Balbeck (Ky. 940!), in Ciliciâ ad Tarsus (Ky. 277! Cadet!), in vinetis ad Aintab (Haussk!). Specimen a cl. Bové lectum probabilissime Syriacum est et ex schedulæ errore ex Arabiâ felici indicatum; loci Brussa et Smyrna in schedulis indicati probabiliter quoque erronei sunt.

Planta demum pedalis et sesquipedalis, spicis fructiferis 4-6-pollicaribus, calycibus deciduis vel diutius persistentibus, corolla $1\frac{1}{2}$ -2 lineas longa lobis aestivatione convolutis erecto-sulcavatis. Species sœpe missa sub nomine *H. rotundifoliū* speciei perennis facie affinis sed diversissimæ.

24. **H. Mamamense** (Bunge loc. cit. p. 30) annum villosocanum, caule a basi ramoso, foliis breviter petiolatis ovatis basi et apice rotundatis, spicis solitariis vel geminatis demum elongatis laxis, calycis sessilis villosi laciniis lanceolatis obtusis corollæ extus adpresso hirtæ tubo subbrevioribus, corollæ longe infundibuliformis tubo limboque subæquilongis hocce in lobos oblongos obtusos aestivatione convolutos apice subflexos profunde partito, antheris versus basin corollæ insertis linearibus faucem subattingentibus apiculo introrsum flexo subhamato terminatis, stylo brevissimo, stigmate longe conico hirto, nuculis tenuissime rugoso-verrucosis ⊖.

Hab. ad salinas Mamam Persiæ boreali-occidentalis (Seidl!).

Planta 6-9-pollicaris, spicæ fructiferæ 3-6-pollicares. Corolla tres lineas longa.

25. **H. Teheranicum** (Bunge loc. cit. p. 31) annum adpresso canescens asperum, foliis oblongis obtusis basi cuneatis petiolatis, spicis solitariis vel geminatis tandem elongatis laxis, calycis sessilis canescentis laciniis lanceolatis obtusiusculis corollæ extus adpresso hirtæ tubo brevioribus, corollæ infundibuliformis limbo tubo breviore in lobos breves ovatos aestivatione convolutos fisso, antheris versus corollæ basin insertis linearibus breviter apiculatis, stylo brevissimo, stigmate longe conico hirto, nuculis (junioribus) glabris ⊖.

Hab. in apricis siccis ad Sergendeh prope Teheran (Bunge!).

Valde affine *H. Mamamensi* et forsitan ejus varietas, ab eo differt tantum

indumento adpresso, foliis basi cuneatis, corollæ lobis brevioribus, antheris brevius apiculatis.

× × Corollæ limbus decemlobus.

26. *H. circinatum* (Griseb. Spicil II, p. 78) annum molliter et adpresso villosum canescens, caule elato ramosissimo, foliis ovatis et oblongis obtusis breviter petiolatis subtus elevatim lineatis, spicis solitariis et conjugatis demum valde elongatis laxis, calycis corollæ subtriplo brevioris laciniis oblongo-linearibus, corollæ extus hirtæ intus glabré infundibuliformis tubo in limbum eo longiore sensim ampliato, limbi laciniis 10 a basi triangulari subulato-attenuatis in aestivatione circinnatim curvatis intercalaribus paulo brevioribus, antheris linearibus prope basin corollæ insertis apiculo introrsum flexo, stylo brevissimo, stigmate conico valde elongato hirto, nuculis laevibus glabris ⊖. *H. neglectum* et *H. Haussknechtii* Bge loc. cit. p. 32 et 34.

Hab. in glareosis Assyriæ ad Tigridem prope Mossul (Ky. exs. 436! sub *H. grandiflora*), in schistosis Karduchiaæ Armenie ad Mükus '4000' (Ky. Suppl. 700!), in Cataonia prope Marasch (Haussk!).

Pedale et sesquipedale, racemi demum pedales, calyces decidui; facies *H. grandiflora* a quo differt corollæ tres lineas tantum longe tubo multo breviore, limbo acute decemlobo, que nota in spec. typicis *H. circinatum* æque ac in *H. neglecto* et *H. Haussknechtii* Bge observatur.

27. *H. Mesinanum* (Bge loc. cit. p. 35) annum, humile a basi spicigerum divaricato-ramosum retrorsum villosulum, foliis petiolatis subrotundis obtusissimis supra atrovirentibus strigilosis subtus canescensibus, spicis elongatis laxissimis uniseriatis, calycis laciniis linear-oblängis erectis corollæ tubum retrorsum hispidum intus glabrum superantibus, limbi infundibularis extus glabri lobis 10 acuminatis exterioribus incurvo-involutis, intercalaribus multo brevioribus erectis, antheris in medio tubo sitis apice non hamato-mucronatis, stigmate subsessili a basi dilatata undulata breviter conico apice hispidulo, nuculis laevissimis glabris ⊖.

Hab. in Persiæ boreali magis orientali inter Mesinon et Mehr (Bunge). Non vidi.

Ex amic. auctore affine *H. neglecto* quod ad *H. circinatum* adduxi, flores in spicis inter se distantes, eorum longitudine non indicatur.

28. *H. samoliflorum* (Bunge loc. cit. p. 36) annum a basi divaricatum ramosum pilis patulis ad ramos et petiolos retrorsis villosum, foliis longe petiolatis ovato-orbiculatis obtusis subtus præsertim crenate et minute tuberculatis, spicis solitariis uniseriatis elongatis tandem laxis, calycis laciniis oblongo-linearibus obtusis corollino tubo hirto intus glabro subbrevioribus, corollæ limbo tubo subæquilongo fere ad basin in 5 lobos spathulato-oblängos et 5 lacintas intercalares subulatas partito, antheris oblongo-linearibus infra medium tubum

insertis apiculo obsoleto, stigmate conico-subulato puberulo stylum pluries superante, nuculis glabris tenuissime et seriatim granulosis ⊖.

Hab. in apricis limoso-salsis prope Schurab, inter Kaschan et Kom Persiae mediae occidentalis (Bge!), in collibus ad Kohnud (Haussk. exs. sub *H. dissitiflora*!).

Caulis primarius ramis lateralibus divaricatis sa-pe brevior, folia inferiora pollicaria, spicæ 3-4-pollicares, corolla 2½ lin. longa. Species limbi corollini structura distinctissima

29. *H. Sultanense* (Bunge loc. cit. p. 57) annum a basi divaricatum ramosum pilis patulis ad petiolas et ramos retrorsis villosum, foliis longe petiolatis ovato-orbiculatis obtusis subtus præsertim crebre et minute tuberculatis, spicis solitariis uniseriatis elongatis tandem laxis, calycis laciniis oblongo-linearibus obtusis corollinum tubum extus hirtum intus glabrum æquantibus, limbo tubo duplo breviore decemlobo lobis ovatis exterioribus acutiusculis extus hirtis interioribus intercalariibus glabris subbrevioribus obtusis, antheris paulo infra medium insertis faucem attingentibus apiculo subincurvo, stigmate conico hispidulo stylum longitudine superante, nuculis glabris tenuissime et seriatim granulosis ⊖.

Hab. in limoso-salsis ad Hous-i-Sultan inter' Kum et Teheran (Bge!).

Præcedenti facie, foliorum indumento simillimum, flores subminores sessilineam longi. Ab eo limbo corollino in 10 dentes omnes ovatos partito distinctum.

SECT. VI. PSEUDONOSMA.

30. *H. micranthos* (Pall. It. II, p. 485. App. p. 734, tab. L sub *Onosmate*) annum, caule dichotome ramoso glabriusculo, foliis longe petiolatis a basi rotundata vel subcordata ovato-oblongis obtusiuseulis supra strigillosis lineatis subtus dense canescensibus, racemis oppositifoliis laxis brevibus 3-5-floris, pedicellis patentissimis flore 3-4-plo longioribus, calyce canescenti-piloso laciniis oblongis acutiusculis fructifero acreto laciniis stellatim patentibus, corolla calycem vix æquante extus retrorsum sericea, limbi minutissimi tubo quadruplo brevioris lobis lanceolatis acutis aestivatione inflexis, sinus edentulis, antheris medio tubo insertis apiculatis, stigmate conico apice hispido stylo glabro longiore, nuculis linearis-oblongis retrorsum sericeo-lauatis ⊖. *H. micranthum* Bge Pl. Lehm. p. 399. — *Tournefortia micranthos* Alph. D.C. Prodr. X. p. 67.

Hab. in deserto arenoso Kisilkum Turkestaniae (Lehm!).

Planta 6-9-pollicaris, folia majora pollicaria, corolla linea brevior, calyx fructifer 2 lineas longas, nucullæ 2 lineas longæ.

Ar. Geogr. Rossia australis, Songaria, Sibiria Altaica.

31. **H. acutiflorum** (Kar. et Kir. Enum. Song. N° 1841) annum, caule dichotome ramoso hispidulo-pubescente, foliis longe petiolatis ovalo-ellipticis basi subattenuatis supra parce et breviter strigilosis subtus pubescentibus, racemis oppositifoliis 4-6-floris, pedieillis florem vix superantibus, ealyce patule hispido laciinis ovatis acutis denum subaucto laciinis stellatim patentibus, corollæ ealyce sublongioris tubo extus retrorsum sericeo limbi tubo quadruplo brevioris laciinis lanceolatis acutis aestivatione inflexis sinubus dentiferis, antheris breviter linearibus medio tubo insertis apiculatis, stigmate conico apice hispidulo stylum subæquante, nuculis oblongis retrorsum sericeo-lanatis ⊖.

Hab. in arenosis Turkestaniae inter flumina Kuwan et Jan darja (Lehm!).

Affine praecedenti, caules hispiduli, racemi minus laxi, pedicelli breviores, corolla longior ad sinus appendiculata, nuculae dimidio breviores.

Ar. Geogr. Songaria.

SERIES B. — PERENNIA.

SECT. VII. CATOXYS.

✗ Corolla intus hirta.

32. **H. Transoxanum** (Bge Pl. Lehm. p. 402) basi suffruticosum multicaule, caulis virentibus dichotome et subdivaricatim ramosissimis glabratis vel superne hispidulis, foliis ovatis vel oblongis petiolatis aentis superne impresso-lineatis seabridulis viridibus subtus canescensibus, spicis conjugatis divaricatis laxissime 6-8-floris, calycis canescens laciniis oblongis, ealyceis tubo sericeo ealyce 2-3-plo longiore intus supra antherarum insertionem villosulo, limbi laciinis hispidulis anguste lanceolato-linearibus intus flexis tubo quadruplo brevioribus, sinubus plicis dentiformibus interstinetis, antheris paulo infra medium tubum insertis oblongo-linearibus retusis, stigmate elongato-conico styloque æquilongis, nuculis sericeo-villosis 5. *H. desertorum* Boiss. et Buhse Enum. p. 130.

Hab. in deserto salso Persiae mediae prope Enarek (Buhse!), et in toto ambitu deserti magni (Bge!), in arena mobili deserti Kisilkum Turkestaniae (Lehm!).

Facie et characteribus *H. dasycarpa* valde affine sed corolla intus hirta nec glabra, lacinia breviores.

33. **H. Kumense** (Bunge loc. cit. p. 42) suffruticosum, caulis elatis albis fere a basi dichotome et strictiuscule ramosis hirtulis, foliis subsessilibus minimis ovato-oblongis margine subrevolutis utrinque hispidulo-canescensibus, spicis geminatis denique elongatis laxissimis, calycis laciniis ovatis aentis subpatulis corollæ triplo brevioribus, corollæ tubo extus sericeo intus totâ longitudine hispido, limbi lobis

breviter triangulari-lanceolatis intus flexis tubo quintuplo brevioribus, antheris sub medio tubo insertis apice retusis connectivo brevissime incurvo, stigmate conico-elongato stylum triplo superante utroque hirsuto, nuculis sericeis ♂.

Hab. prope Pal i Dallak in deserto ad septentrionem urbis Koum Persiæ (Bge!).

Pedale vel sesquipedale, rami graciles albi, folia majora semipollucaria, corolla lineam unam vix excedens.

34. **H. trichostomum** (Bge loc. cit. p. 41) perenne multicaule, caulis albidis herbaceis diffusis ramosissimis villosulis, foliis parvis breviter petiolatis ovato-suborbiculatis obtusissimis superne pallide virentibus hispidulis subtus canescenti-villosis, spicis numerosis abbreviatis confertifloris, calycis laciiniis oblongis acutiusculis erecto-patulis fructiferis patentissimis corollæ tubo retrorsum sericeo sub-brevioribus, limbi corollini brevissimi laciiniis triangularibus subulato-acuminatis intus inflexis, fauce minute puberula, antheris oblongo-linearibus sub medio tubo insertis faucem attingentibus apiculo infexo, stigmate conico seriatim puberulo stylum retrorsum hispidum æquante, nuculis pube retrorsâ tandem secedente sericeis ♀.

Hab. ad margines australiores magni deserti salsi Persiæ ad Kermusch inter Kerman et Jezl (Bge). Non vidi.

Vix 6-pollicare, folia semipollucaria, affine dicitur *H. Halame* sed pubescen-tia fauciæ corollæ distinctum; a ceteris sectionis caulis non suffruticosis quoque differre videtur.

× × Corolla intus glabra.

35. **H. dasycarpum** (Ledeb. Fl. Ross. III, p. 101) suffruticosum mul'icaule, caulis viridibus dichotome ramosissimis patule plus minus hispidis vel glabratis, foliis parvis supra impresso-nervosis et parce strigillossis subtus canescenti-pilosulis vel hispidis acutiusculis basi cuneatis inferioribus longius cæteris brevius pedunculatis, racemis geminatis paucifloris laxis, calycis hispiduli vel glabrati laciiniis ovalis vel oblongis, corollæ tubo calyce duplo longiore extus retrorsum plus minus sericeo intus glabro laciiniis anguste lanceolato-subulatis intus reflexis dimidium tabum æquantibus vel brevioribus, plieis retusis ad sinus, antheris medio tubo insertis, stigmate elongato-conico styloque æquilongis hispidulis, nuculis ovatis sericeis vel glabratis ♂. Ic. Eichw. Casp. tab. 5. — *H. Schrenkianum* Led. Fl. Ross. III. p. 102. — *H. Brahuicum* Stocks Hook. Journ. Bot. 4, p. 173.

Hab. in Persiæ orientali (Bge), Turkestaniæ in littore orientali Ca-piæ (Eichw.) et interiore (Lehm!), Belutschia superiore 4000' (Stocks!), Affghania (Griff.).

Semipedale vel pedale, multum variat indumento hispidulo vel fere evanido. Corolla 2 lineas longa.

Ar. Geogr. Songaria.

36. **H. minutiflorum** (Bunge loc. cit. p. 46) fruticosum dense canescens ramosum, foliis parvis breviter petiolatis ovatis acutiusculis supra virentibus adpresso strigulosis subtus incanis, spicis solitariis vel conjugatis dense paucifloris, calycis laciniis ovato-oblongis corollæ tubum extus retrorsum sericeum intus glaberrimum subæquantibus, limbi laciniis triangulari-lanceolatis acuminatis tubo dimidio brevioribus sub anthesi erecto-patulis, antheris medio tubo insertis apice retusis apiculo hamato, stigmate abbreviatum conico stylum retrorsum hispidum æquante, nuculis sericeo-villosis ♂.

Hab. ad Safar Khodje non procul ab Kasbin Persiae (Bienert specimina pauca primâ anthesi). Non vidi.

Folia parva semipollicaria ad summum 5 lineas lata margine rigide ciliata. Ex amic. auctore prope *H. dasycarpum* collocandum.

37. **H. eremobium** (Bunge loc. cit. p. 43) suffruticosum caulinibus dichotome et strictiuseule ramosissimis ut et rami albis hispidulis, foliis brevissime petiolatis parvis ovato-oblongis acutiusculis supra impresso-lineatis sebris subtus canescensibus, spicis paucifloris gracilibus laxis, calycis sessilis hirsuti laciniis lanceolatis obtusiusculis, corollæ extus hirtæ intus glabræ tubo calyce sesquilociore, limbi tubo duplo brevioris laciniis patentibus a basi triangulari abrupte et anguste lanceolatis, antheris medio tubo insertis linearibus apice retusis et apiculo intus flexo obsitis, stigmate pyramidato-conico stylo retrorsum hispido sublongiore, nuculis sericeo-villosis ♂.

Hab. ad orientem et septentrionem magni deserti salsi Persiae inter Bedescht et Meiomei et inter Kum et Pull i Dallak.

Pedale. folia 4-6 lineas longa, corollæ aurantiacæ sesquilineam longæ.

38. **H. Halame** (Boiss. et Buhse Enum. p. 150) fruticosum caulinibus albis pumilis divaricatum ramulosis hispidis, foliis parvis breviter petiolatis ovatis obtusis undulato-repandis patule hispidis griseis supra depresso-lineatis, spicis brevissimis paucifloris capitato-fasciculatis floribus 1-2 inferioribus interdum remotiusculis, calycis patule hirsuti laciniis lanceolatis acutis, corollæ extus hirsutæ intus glabræ tubo calyce duplo longiore limbi brevissimi lobis a basi oblongâ in euspidem subulatam intus flexam attenuatis, antheris oblongis versus medium tubum insertis, stigmate conico-pyramidalito hirtulo stylo glabriusculo æquilongo, nuculis sericeo-villosis ♂.

Hab. in aridis deserti salsi Persiae prope Djendack et prope Jezd (Buhsel), ad Meibut inter Jezd et Ispahan (Bge!), in monte Sefa (Bode!).

Semipedale. folia majora sex lineas longa, corolla vix 2 lineas longa. Ab affinibus floribus capitato-congestis discedens.

39. **H. luteum** (Poir. Suppl. 3, p. 22) fruticosum dumulosum ramosissimum ramis rigidis brevibus albidis retrorsum asperulis, foliis parvis sessilibus oblongis margine sœpe undulatis supra lineato-

depressis et pilis basi tuberculatis asperis subtus canescensibus, spicis terminalibus conjugatis brevibus rigidis densifloris, calycis hirsuti laciniis lanceolatis, corollæ tubo extus sericeo rarius glabrato calyce sesquilociore limbi brevis laciniis anguste linearibus introrsum curvatis sinuum plicis retusis, antheris oblongis muticis retusis prope faucem sitis, stylo retrorsum hispido, stigmate longe conico hirsuto stylo duplo breviore, nuculis magnis dense et longe albo-sericeis rarius glabratis ♂. *H. lineatum* Del. Fl. Eg. 37, tab. 17 non Vahl. — *H. eriocarpum* Lehm. Asp. p. 55. — *Lithospermum digynum* Forsk. — *H. callosum* Spreng. Nov. Prov. 21.

Hab. in desertis arenosis Egypti (Forsk., Del!, Sieb!, Boiss!, etc.).

Fruticulus 1½-1-pedalis, folia majora 8-9 lineas longa, corolla tres lineas longa.

Ar. Geogr. Africa borealis.

40. *H. Aucheri* (p.c. Prodr. IX, p. 533) perenne imà basi suffruticosa, caulis subherbaeis ramosis erectis flexuosis albidis tomentellis sebris setis nonnullis longioribus patulis, foliis petiolatis majusculis ovatis vel orbiculatis saepe subcordatis supra pallide virentibus depresso-lineatis tuberculatis breviter setiferis et pube brevi adpresso scaberrimis subtus molliter canescenti-tomentosis, spicis solitariis et conjugatis brevibus densifloris fructiferis laxiusculis, calycis hirsutissimi laciniis elliptico-linearibus obtusis. corollæ extus retrorsum sericeæ intus glabrae tubo calyce duplo longiore limbi laciniis tubo 3-6-plo brevioribus sub anthesi patentibus angustissime linearis-lanceolatis acutis, antheris medio tubo insertis linearibus apice retusis cum apiculo subobsoleto, stigmate elongato-conico stylo retrorsum hispido subæquilongo apice hispidulo, nuculis sericeo-hirsutis ♂. *H. seriocarpum* Bunge loc. cit. p. 44!

Hab. in Persia (Auch!), in sterilibus aridis Persie mediae ad occidentem deserti salsi magni ad Schurab inter Kaschan et Koum (Bge!).

Pedale, folia inferiora 9-10 lineas longa interdum transverse latiora, spicæ sub anthesi 6-7 lineas longe demum 1-2-pollicares, corolla sesquilineam longa. Species foliis superne scaberrimis insignis. Planta Aucheriana schedulae commutatione erronee ex Mossul Assyriæ indicata Bungeanæ omnibus notis similis est.

41. *H. Carmanicum* (Bunge loc. cit. p. 46) perenne, caulis herbaceis diffusis albis flexuosis dichotoine ramosisque villosulis, foliis subtus patule supra adpresso pubescenti-griseis depresso-lineatis petiolatis late ovatis vel subcordatis obtusis supremis mucronulatis, spicis brevibus paucifloris tandem laxe uniseriatis. calycis albo-villosi laciniis acute lanceolatis corollæ tubo extus retrorsum sericeo intus glabro vix brevioribus, limbi brevis laciniis a basi triangulari subulatis intus flexis, antheris paulo infra medium tubum insertis oblongis retusis apiculo flexo, stigmate conico stylum glabrum longitudine superante, nuculis glaberrimis ♀.

Hab. in Persia austro-orientali provincia Carmaniæ ad orientem et occidentem urbis Kerman ad Chabbise, Feisabad, etc. (Bunge!).

Caulis jam sub anthesi 1-1 $\frac{1}{2}$ -pedales, folia inferiora pollicaria tere *H. Aucheri* sed mollia nec scaberrima, spicæ sub anthesi pollicares fructiferae 2-3-pollicares, corolla 2 lineas longa.

SECT. VIII. RADULA.

× Corolla intus hirta

42. *H. Cabulicum* (Bge loc. cit. p. 39) virens multicaule basi suffruticosum tuberculis brevissime microniferis asperum, caulinis herbaceis duris erectis ramosis, foliis obovato-oblongis in petiolam brevem attenuatis, racemis paucifloris uniseriatis brevibus tandem laxiusculis, calycis intus extusque hirsuti laciniis oblongis, corollæ tubo extus hirsuto calyce duplo longiore intus totâ parte dimidiâ superiore hirsuto sed inferne glaberrimo, lobis obovato-oblongis utrinque hirtis tubo quadruplo brevioribus, antheris infra medium tubum insertis oblongis, stigmate glabro ovato-conico subsessili, nunculis (jnnioribus) glabris 5.

Hab. in arenosis ad Putullah Affghaniae (Griff. Journ. 528! exs. 5030!).

Pedale et elatus, folia majora 10 lineas longa, corolla inter affinia majuscula fere 3 lineas longa. Tubo intus parte superiori hirsutissimo insigne.

43. *H. Affghanum*, suffruticosum setalis albis basi tuberculatis incurvis et adpressis griseum asperulum, ramis erectis flexuosis tenuibus, foliis parvis sessilibus breviter linear-lanceolatis utrinque attenuatis margine undulatis et subrevolutis, spicis conjugatis brevibus laxiusculis saepe cymæformibus floribus terminalibus subfasciellatis, calycis hirsuti intus glabri laciniis oblongis obtusis encultatis margine anguste membranaceis, corollæ parvæ calyce sesquilociore extus sericeâ intus dimidiâ parte superiore parce papillosâ lobis ovato-oblongis brevissimis, antheris oblongis versus medium tubum sitis, stigmate conico sessili acuto, nuculis sericeis 5.

Hab. in Affghania (Griff. N° 5936!).

Semipedale vel longius, folia inferiora 4-6 lineas longa 1-1 $\frac{1}{2}$ lim. lata, flores vix lineas longi. Affine *H. brevilimbii* et *Cabulico* sed characteribus distinctum.

44. *H. strigosum* (Willd. Sp. I. p. 743) basi suffrutescens ramosissimum totum adpresso strigulosum, foliis anguste linear-lanceolatis vel linearibus acutis margine subrevolutis, ramis tenauer virgatis in spicas laxas tenues inferne bracteatas abenitibus, calycis adpresso strigosi laciniis oblongo-lanceolatis acutis, corollæ extus stri-gulosae tubo calyce subbreviore lobis ovato-oblongis, fauce hirtulâ, antheris versus medium tubum insertis apice attenuato puberulo primum cohaerentibus, stigmate conico-attenuato apice hirto stylo

brevi longiore, nuculis globosis strigulosis ♂. *H. bicolor* Hochst. et Steud. n.c. Prodr. IX, p. 546. — *H. Cordofanum* H. et Steud.

Hab. in Belutschia (Stocks exs. 311!).

Variat ramis erectis rarius decumbentibus, indumento rariori vel sericeo. *H. thymoi* 'es Jaub. et Spach Illustr. tab. 362 videtur hujus forma minor.

45. ***H. rariflorum*** (Stocks Hook. Journ. Bot. IV (1852) p. 174) suffruticosum setis adpressis strigulosum canescens, ramis erectis crebre foliatis in spicas laxas solitarias vel conjugatas bracteatas abeuntibus, foliis anguste oblongo-linearibus linearibusque utrinque attenuatis sessilibus margine subrevolutis, calycis strigulosi laciniis oblongis acutiusculis, corollæ extus strigulosæ intus undique sparsim papillari-hirtulae tubo calycem æquanti lobis semiorbiculatis, antheris supra medium tubum insertis apiculo hirto primum coherentibus, stylo glabro ovario sublongiore, stigmate hirto brevissime globoso-conico, nuculis..... ♂.

Hab. in Belutschia inferiore (Stocks exs. 492!).

Semipedale, hoc prius pro *H. strigosii* formâ habui sed præter folia sublitoria differt corollæ intus undique hirtæ lobis brevissimis styloque stigmate triplo longiore.

Ar. Geogr. Ditio Scinde Indiæ.

× × Corolla intus glabra.

46. ***H. rotundifolium*** (Sieb. exs.) fruticosum totum tomento brevi mollissimo canum, ramis ascendentibus vel erectis elongatis ramosis, foliis majusculis orbiculatis vel ovatis obtusissimis supra valde depresso-lineatis subtus elevatim nervosis inferioribus longe superioribus breviter petiolatis, racemis longis uniserialibus tandem laxiuseculis, calycis tomentosi corollæ tubo subduplo brevioris laciniis elliptico-linearibus obtusis, corollæ tubo extus sericeo intus glabro limbi tubo æquilangi patentis lobis breviter ovatis obtusis, antheris oblongis apice acentatis infra medium tubum sitis, stigmate subsessili conico-elongato apice hirto, nuculis minute tuberculatis glabris ♂.

Hab. in collibus Palestine circa Hierosolymam et Bethleem (Sieb! Boiss! Gaill!), in Antilibano ad faucom Barrada et circa Damascum (Gaill!).

Rami cum spicis 4-½-pedales, spicæ 2-3-pollicares, corolla lineas tres vel paulo amplius longa limbo amplio, folia inferiora diametro 7-9 lineas lata. Corollæ aestivatio ob laciniias subconvolutas intermedia.

47. ***H. radula*** (Fisch. et Mey. ex Ledeb. Fl. Ross. III, p. 101) basi suffrutescens caulinibus herbaceis sed induratis ramosissimis ramisque albis hirtulis, foliis elliptico-ovatis obtusis basi rotundatis breviter petiolatis supra strigulosis depresso-lineatis subtus elevatim nervosis canescensibus, spicis brevissimis ovato-oblongis densifloris, calycis albo-hirsuti laciniis ovatis, corollæ calyce vix sesquialongioris

tubo extus hirsuto intus glabro. limbi brevissimi lobis breviter ovatis, antheris oblongis supra medium tubum insertis. stigmate conico stylo sublongiore, nucibus albo-villosis ȝ.

Hab. in arenâ mobili deserti Kisylkum Turkestaniae (Lehm!).

Pedale et longius. folia majora 6-9 lineas longa, flores parvi, corolla lineam longitudine vix excedens.

48. *H. brevilimbe* (Boiss. Diagn. Ser. I, 11, p. 89) basi suffruticosum erectum pluricaule ramosissimum totum pilis basi subtuberculatis ad ramos retrorsis dense canescenti-velutinum asperum. foliis oblongo-lanceolatis in petiolum brevem attenuatis obtusiusculis superne depresso-lineatis subtus elevatim nervosis margine subundulatis, racemis brevibus unilateralibus tandem laxiusculis, calycis extus valde hirsuti laciniis oblongis acutiusculis intus apice puberulis, corollæ tubo calyci æquilongo extus sericeo intus glabro lobis obovatis obtusis tubo quadruplo brevioribus, antheris parvis ovato-oblongis infra medium tubum sitis, stigmate elongato-conico glabro, stylo brevissimo, nuculis sericeis ȝ.

Hab. in Persiâ loco non notato (Auch. 505 A!), in montibus Sofi prope Ispahan et Kohrud prope Kaschan (Bode! Kapherr!).

Strictiuscule ramosum 9-10-pollicare, folia inferiora pollicaria 2-3 lineas lata cetera multo breviora et angustiora, flores parvi vix sesquilineam longi.

49. *H. ophioglossum* (Stocks MSS. ex Aitch. Cat. Punj. p. 94) suffruticosum ramosum dupli indumento altero pube tomentellâ altero setis rigidis longis pellucidis basi tuberculatis constante griseum asperulum, foliis oblongis et oblongo-lanceolatis inferioribus in petiolum brevem attenuatis superioribus sessilibus acutis, spicis longis uniseriatis densiusculis. calyce setoso-hispidissimo laciniis anguste linearibus acutis corollæ tubum subæquantibus, corollæ tubo hirsuto intus glabro, limbi tubo subtriplo brevioris erecto-patentis lobis breviter ovatis, sinibus lobos accessorios totidem minores edentibus, antheris paulo sub medio tubo insertis longe linearibus apice subexsertis, stigmate glabro stylo brevi suffulto a basi breviter conicâ longissime subulato apice longe bilobo lobis exsertis, nuculis ovatis laeviusculis glabris ȝ. *H. ditrichum* Stocks exs.

Hab. in Belutschia (Frere ex Aitch. Cat. p. 94).

Ar. Geogr. Ditio Scinde.

Fere pedale, folia fere *H. undulati* et *Arbainensis* sed latiora vix undulata, spicæ demum 4-5-pollicares, flores vix 2 lineas longi. Species inter omnes stigmatis subulati longissimi (fere 2 lineas longi) lobis longis subulatis apice exsertis insignis.

50. *H. denticulatum* (Boiss. et Haussk.) basi suffruticosum multicaule totum adpressissime tomentosum canum, caulis erectis parce ramosis, foliis supra impresso-lineatis subtus elevatim nervosis subsessilibus et sessilibus oblongis obtusiusculis sursum diminutis,

spicis subsolitariis uniserialibus demum laxiusculis, calyce breviter extus et intus tomentoso corollæ tubo plus duplo breviore in lobos ovatos obtusos paulo infra medium fisso, corollæ tubo extus adpresso tomentoso intus glabro, limbo tubo quadruplo brevioris lobis 10, externis 5 ovatis eximie denticulatis, internis 5 accessoriis vix brevioribus a basi lanceolatâ canaliculatâ in cuspidem intus recurvam attenuatis, antheris oblongo-linearibus apiculatis in parte inferiore tubi sitis, stigmate conico-attenuato apice hirto sessili, ovario glabro 3.

Hab. in rupestribus calcareis montanis ditionis Elymaitidis Persiae austro-occid. prope Malamir (Haussk!). Fl. Jul. Aug.

Dumulosum 8-10-pollicare, folia inferiora 8-10 lin. longa 4-5 lata, spicæ 2-3-pollicares, calyx demum deciduus lineâ vix longior, corollæ 3½ lineas longæ tubus angustus. Species inter perennes limbo decemlobo et ab omnibus calyce minus profunde partito distincta.

51. H. Sogdianum (Bge Pl. Lehm. p. 403) perenne, caulis herbaceis albis demum ramosis parce hispidis, foliis pallide viridibus supra tuberculis sparsis aculeolatis strigulosis subtus hispidulis, spicis geminatis paucifloris laxiusculis, calycis hispida tubo corollino glaberrimo triplo brevioris lacinii lanceolato-linearibus acutis, corollæ limbo extus piloso tubo quadruplo breviore lobis rotundatis undulato-crispis, antheris ovatis prope faucem sitis apiculo intus flexo, stylo elongato stigma breviter conicum apice truncatum et hispidum longitudine quintuplo superante, nuculis... ♀

Hab. in desertis argillosis Turkestaniæ ad Batkakkum et Kisilkum (Lehm!).

Caules semipedales vel longiores, erecti?, folia inferiora 9-10 lin. longa, spicæ demum pollicares 10-15-floræ, corolla 3½-4 lineas longa. Species ab omnibus congeneribus longitudine stylî (fere 2 lineas longi) distinctissima.

52. H. Arbaïnense (Fresen. Beitr. p. 168) suffruticosum totum pube brevi viscidâ pilisque longioribus patulis glanduloso-tomentellum flavo-vires, caulis pumilis crassiusculis simplicibus vel ramosis, foliis oblongis in petiolum attenuatis acutinseculis margine undulatis et superne saepius depresso-lineatis, spicis solitariis et conjugatis densifloris uniserialiatis sub anthesi valde incurvis, calycis intus et extus hirsuti lacinii linear-lanceolatis, corollæ tubo calyce subduplo longiore extus hirto intus glabro et costis 5 prominulis infra faucem in denticulum abeuntibus obsito, limbi patentis tubo sesquibrevioris lobis ovatis repandis, antheris oblongo-linearibus infra medium tubum sitis, stigmate pileiformi glabro subsessili, nuculis reticulato-rugosis glabris 5.

Hab. in desertis Aegypti (Radde!), ad Kosseir (Acerbi, Schweinf.), Arabia petraea ad Ouadi Arbaïn (Ruppell.), ad Ouadi Hebran (Schimp. 225!), ad Ouadi Scheick (Boiss!).

Semipedale vel parum longius, folia majora 8-9 lineas longa rarius pollicaria, flores majusculi, corolla 4 lineas longa.

Ar. Geogr. Nubia littoralis. Arabia tropica

53. H. undulatum (Vahl Symb. I, p. 13) adpressiusculæ tuberculato-setosum griseum asperulum basi suffruticosum, caulinis herbaceis elongatis procumbentibus, foliis in petiolum brevem attenuatis oblongis et oblongo-lanceolatis obtusiusculis margine undulato-crispis, spicis densis biseriatis, calycis hirsuti laciniis linearis-oblongis obtusis, corollæ tubo calyce sublongiore extus hirto, lobis ovato-orbiculatis patentibus tubo subbrevioribus sinus intreflexis, antheris linearibus medio tubo adnatis, stigmate hirto elongato-conico quadrilobo stylo sublongiore, nuculis 4 hirtulis separabilibus 5. *H. crispum* Desf. Atl. I, p. 151, tab. 41. — *Lithospermum hispidum* Forsk.

Hab. in cultis et arenosis desertorum Ægypti ad Kahiram et Suez (Forsk., Del., Ehrenb!, Boiss!, etc.), ad mare Rubrum (Schweinf!), Palestinam ad Gaza (Boev!).

Species affines cum *H. undulato* a me in Diagn. II, p. 50 conjuncte sunt *H. Nubicum* Bge loc. cit. p. 52 nuculis ad dorsum gibbo inflato auctis et *H. erosum* Lehm. e Senegaliam et Canariis nuculis per paria concretis rectius distingueda.

Ar. Geogr. Africa borealis.

54. H. Persicum (Lam. Diet. III, p. 94) setulis basi tuberculatis asperum griseum suffruticosum caulinis erectis, foliis subsessilibus basi attenuatis lanceolatis acutiusculis margine undulatis, spicis solitariis vel conjugatis siccipius laxiusculis subuniserialibus, calycis hirsuti laciniis oblongis obtusis, corollæ tubo extus hirto calyce sublongiore, lobis ovatis erosulis tubo sublongioribus, antheris ovatis paucis sub medio tubo adnatis, stigmate glabriusculo breviter conico stylo brevi longiore, nuculis 4 hirtulis separabilibus 5. *H. ramosissimum* Sieb. exs. — *H. eriocarpum* Del. in Lehm. Asp. p. 55. — *H. Persicum* Burm. Ind. p. 41, tab. 29.

Hab. in desertis Ægypti (Del.), Arabiae Petreæ (Schimp. 726!), Assyriae (Auch 2078!), Persiae mediae ad limites deserti magni (Oliv!, Bge!), australis ad Abuchir et Dalechi (Ky. 134!, Haussk!) et in insulis sinus Persici (Ky. 17!), Afghanistanæ ad Dair Haj (Griff!), Belutschia (Stocks!).

Ab affini *H. undulato* characteribus datis differre videtur.

Ar. Geogr. Arabia tropica, Pentapotamis.

55. H. tuberculatum (Boiss. in Ky. exs. Pers. austr. 1843) fruticosum dumulosum setulis albis basi tuberculatis asperum siccitate nigrescens, ramis erectis cibbre foliosis, foliis parvis angustis et breviter lineari-lanceolatis semiamplexicaulinibus acutis margine revolutis, spicis solitariis vel conjugatis brevissimis ovatis distiche densifloris, calycis laciniis obtusis ciliatis corollæ tubo subbrevioribus, corollæ extus adpresso sericeæ lobis tubo quadruplo brevioribus ovatis sinus intreflexis, antheris oblongis sub medio tubo insertis, stigmate hirto conico stylo aquilongo, nuculis glabris per paria connatis 5. *H. undulatum* var. *tuberculatum* Boiss. Diagn. II, p. 89. — *Heliophytum Kotschyi* Bge Pl. Lehm. p. 404.

Hab. ad oras Persiae australis (Auch. 5002!), in arenosis insularum Karak et Korgo (Ky. 22!).

Fruticuli ramosissimi $\frac{1}{2}$ -1-pedales, folia 3-4 lineas longa 1 lata, racemi brevissimi usque ad flores foliati.

TR. III. BORRAGEÆ D.C. Prodr. — Ovarium constans carpellis binis bipartitis rarius bilocularibus. Stylus a basi in centro gynobaseos ortus. Nuculæ quatuor uniloculares rarius duo biloculares. Semina exalbuminosa.

SUBTR. I. CERINTHEÆ D.C. — Nuculæ duo biloculares areolâ planâ toro incidentes a gynobasi et stylo liberæ.

CERINTHE (Tourn. Inst. 79, tab. 56).

Calyx profunde quinquepartitus laciniis foliaceis inæqualibus. Corolla tubulosa fauce nuda. Antheræ hastatæ lobis basilaribus divergentibus apice appendice acutâ auctæ. Stylus filiformis capitatus. Nuces 2 ovatæ subosseæ biloculares e nuculis duabus geminatum connatis constantes basi areolâ planâ toro affixæ. — Herbæ.

1. **C. minor** (L. Sp. 772) glaucescens glabra, caulis erectis ramosis, foliis levibus vel verruculis nonnullis obsitis fasciculorum steriliū et inferioribus oblongo-spathulatis in petiolam attenuatis, superioribus cordato-oblongis auriculis rotundatis, racemis fructiferis valde elongatis bracteis cordatis calyces superantibus obsitis, calycis laciniis lanceolatis inæqualibus plus minus manifeste ciliatis fructiferis auctis basi rotundatis vel subcordato-auriculatis, corollâ calyce sesquilongiore ultra tertiam partem in laciniis lanceolatas acuminatas erecto-conniventes fissâ, antheris filamento brevissimo suffultis appendice lanceâ corollam æquante vel breviore terminatis, carpellis levibus ② vel ȝ. Ic. Jacq. Austr. t. 224. — Rehb. t. 94. — *C. maculata* M.B. Taur. Cauc. № 336. — *C. auriculata* Ten. — *C. cleistostoma* Boiss. et Spr. Diagn. Ser. I. 4, p. 44. — *C. macrophylla* Boiss. et Heldr. pl. exs. — *C. Cilicica* Ky. exs.

Hab. in fruticetis, rupestribus umbrosis interdum in cultis regionis inferioris et sæpius montanae, in subalpinam et alpinam ascendens, Græcia in regione alpinâ Taygeti (Heldr! forma *macrophylla*). Arcadiâ (Heldr!), in fauibus Hymetti (Spr! Boiss!), Olympo Thessalo (Heldr!), Macedoniâ (Orph!), Thraciâ (Griseb!), Bithyniâ (Thirke!), Paphlagoniâ et Galatiâ (Wied!), montibus Lydiæ (Bal!), Pamphyliæ (Heldr!), Lyciæ (Bourg!), Ciliciæ in Tauro usque ad 7500' (Ky!), Cappadociæ (Bal!), Libano ad Cedros (Boiss!), Antilibano (Ky. 415!), Syria boreali et Kurdistania Persicâ (Haussk!), Armeniâ (Huet! Bourg!), Tauriâ et Caucaso onni usque ad 7200' (M.B.. Rupr!) Persiâ boreali (Szov! Buhse!).

Multum variat foliis majoribus et minoribus interdum albo-maculatis, supe-

rioribus nomumquam ciliatis, calycis laciniis basi plus minus dilatatis, corolla omnino flavâ vel ad faucem purpureo-maculatâ vel laciniis purpureis, antherarum appendicibus plus minus elongatis.

Ar. Geogr. Gallia in Alpibus, Germania, Corsica, Italia, Dalmatia, regio Danubialis, Rossia media et meridionalis.

2. C. Alpina (Kit. ap. Schult. Oestr. I, p. 353) glabra, caulis erectis, foliis lœvibus glabris fasciculorum sterilium et inferioribus oblongo-spathulatis, caulinis cordato-oblongis auriculis rotundatis, racemis fructiferis valde elongatis bracteis cordatis calyces demum deflexos superantibus obsitis, calycis laciniis inaequilatis lanceolatis obtusis, corollâ calyce quartâ parte longiore rubellâ vel flavo-virenti intra faucem purpureo-maculatâ dentibus brevissimis ovatis obtusis apice recurvatis, antheris subsessilibus inclusis appendice brevi, carpellis lœvibus ♀. *C. glabra* D.C. Fl. Fr. III, p. 619. — Ic. Rehb. t. 94.

Hab. in regione alpinâ Ponti Lazici supra Djimil 7800' (Bal!).

A speciminiis Europæis non differre videtur nisi calyce majori et corollâ rubellâ.

Ar. Geogr. Gallia in Pyrenæis et Sabaudia, Helvetia, Germania australis, Dalmatia, regio Danubialis.

3. C. major (Lam. Diet. IV, p. 67. — L. ex parte) plus minus tuberculata et ciliata vel glabrata, caule erecto simplici vel ramoso, foliis inferioribus obovato-spathulatis in petiolum attenuatis superioribus ovato-oblongis obtusis basi cordatis rotundato-auriculatis, racemis scorpioideis brevibus densis, bracteis cordatis calyces æquantibus vel superantibus, calycis laciniis oblongis subinæqualibus acutiusculis ciliatis, corollâ flavâ vel purpureâ rectâ clavatâ latâ calyce plus duplo longiore dentibus brevibus latissimis acuminatis recurvis, antheris inclusis filamento brevioribus appendice brevi acutâ, carpellis magnis lœvibus ♂. *C. aspera* Roth Cat. I, p. 33. Fl. Graec. tab. 170. — *C. strigosa* Rehb.

Hab. in cultis et inter segetes Zacynthi (Sibth., Marg!), Peloponnesi (Sibth.), Atticæ (Sprunt!), Bœotiae (Orph!), Macedoniæ (Sibth.), Byzantii (Coumany!), Cycladum (Sart! Orph!), Cretæ (Sieb!).

Ar. Geogr. Regio Mediterranea totius Europæ et Africa borealis.

4. C. retorta (Sibth. et Sm. Fl. Graec. II, p. 60, tab. 171) glabriuscula, caule erecto ramoso, foliis membranaceis lœvibus obtusisimis insimis oblongo-spathulatis attenuato-subpetiolatis cæteris ovatis vel fere orbiculatis late et obtusissime cordato-auriculatis, racemis terminalibus capitato-scorpioideis dein laxis bracteis foliis similibus sed minoribus obsitis, calycis laciniis inaequalibus linearibus obtusis post anthesin auctis basi auriculato-cordatis, corollâ calyce duplo longiore tubuloso-clavatâ albido-ochrolencâ ad basi attenuatam et rursus sub apice violaceo constricto bis curvatâ ad faucem constrictâ

dentibus parvis triangularibus reflexis, antheris corollam æquantibus filamento infracto-ascendente sublongioribus, carpellis lævibus nitidis fusco-maculatis ⊖. Ic. Rehb. tab. 95. — *C. purpurea* Visian. Fl. Dalm. tab. 21.

Hab. in lapidosis umbrosis faucium Messeniae in monte Ithome (Heldr!), Laconiae in monte Malevo (Sprun!), ad Acrocorinthum (Boiss!), in faucibus montis Hymetti (Sprun! Boiss! Heldr!), Acarnaniæ ad Astacos (Nieder!), in Cariâ (ex Sibth.).

Ar. Geogr. Dalmatia.

SUBTR. II. ANCHUSEÆ D.C. — Nuculæ quatuor uniloculares toro insidentes basi excavatae et ibidem sæpius margine tumido cinctæ a gynobasi vel stylo liberae.

BORRAGO (L. Gen. 182).

Calyx quinquepartitus. Corolla rotata tubo brevissimo vel subnullo limbo patenti quinquelobo, fornicibus glabris retusis. Filamenta brevia crassa sub apice appendice obtusâ extus aucta, antheræ exsertæ in conum conniventæ. Nuculæ 4 liberæ basi excavatæ et annulo plicato cinctæ. — Herbæ annuæ vel perennæ.

1. **B. officinalis** (L. Sp. 197) longe et patule setosa substrigosa, caulinis erectis vel ascendentibus crassiusculis brevibus superne paniculato-corymbosis, foliis inferioribus ovatis petiolatis floralibus sessilibus parvis oblongis, pedunculis elongatis fructiferis apice nutantibus, racemis simplicibus vel conjugatis, calycis lacinias linearibus, corollæ cœruleæ basi intus replicatae et villosæ tubo subnullo lobis ovatis acutis, fornicibus triangulari-ovatis retusis, filamentis basi dilatatis, antheris mucronatis, nucibus rectis grosse rugulosis et tuberculatis apice anicipitibus ⊖. Ic. Engl. Bot. tab. 36. — Rehb. tab. 101.

Hab. in ruderatis et cultis Græciæ!, Anatoliæ occidentalis et borealis (Thirke! Wied!), Cypri (Sibth.) et forsitan alibi.

Ar. Geogr. Europæ regio mediterranea et quoque in Europâ media ubi forsitan introducta, Africa borealis.

ANCHUSA (L. Gen. 182). — *Lycopsis* L.

Calyx quinquesfidus vel partitus post anthesin sæpe accretus. Corollæ infundibuliformis tubo recto vel curvato et gibbo fauce fornicibus 5 obtusis papillosum clausâ. Nuculæ 4 liberæ basi excavatæ ibidemque margine tumido sæpius plicato cinctæ. — Herbæ perennæ vel monocarpicæ.

*Specierum Orientalium distributio.***§ 1. Perennes.**

* Calyx quinquefidus. Nuculæ in rostrum obliquum curvatae.

A. officinalis, hybrida, undulata, obliqua, ochroleuca.

* * Calyx quinquepartitus. Nuculæ rectæ

A. Italica, macrocarpa, strigosa, Barrelieri.

* * * Calyx quinquepartitus. Nuculæ in rostrum obliquum abeuntes
vel horizontaliter curvatae.

A. cespitosa, limbata, neglecta, myosotidiflora.

§ 2. Annuæ. Calyx quinquepartitus.

* Tubus corollæ rectus.

+ Nuculæ horizontaliter curvatae.

A. aggregata, hispida, Aucheri.

+ + Nuculæ in rostrum obliquum abeuntes.

A. Egyptiaca, Milleri, stylosa.

+ + + Nuculæ rectæ.

A. Thessala.

* * Corollæ tubus curvatus.

A. arvensis, Orientalis, variegata.

§ 1. Perennes.

1. A. officinalis (L. Sp. 191) perennis, caulis erectis patule hispidis superne paniculatis, foliis oblongo-lanceolatis et lanceolatis breviter hispidis integris vel subrepandis inferioribus in petiolam attenuatis, racemis bifidis tandem laxis, bracteis triangulari-lanceolatis, pedicellis calyce brevioribus fructiferis nutantibus, calyeis quinquefidi fructiferi accreti campanulati laeiniis lanceolatis, corollæ azureæ tubo calycem æquante vel superante, fornicibus ovatis velutinis, nuculis oblique ovatis et rostratis latere altero tuberculato-rugosis supra annulum basilarem constrictis ♀. Ic. Fl. Dan. tab. 572. —

Engl. bot. tab. 662. — *A. angustifolia* D.C. Fl. Fr. — *A. arvalis* Rehb. Pl. Crit. III, fig. 470. — *A. Parnassica* Boiss. et Orph. Diagn. Ser. II, 3, p. 134.

Hab. in regione montanâ et subalpinâ Græciæ, Taygetus ad sylvarum margines prope Hagios Dimitri 600' (Heldr!), mons Kyllene 2500' et Parnassus 5000' (Orph!), Veluchi (Etoliae regio summa (Samar. et Guicci!), mons Delphi Euheæ (Orph!), Olympi Thessaliæ radices (Heldr!), Macedonia prope Vodéna (Griseb! Orph!), mons Korthiatî (Heldr!).

β. ochroleuca. — Flores ochroleuci, cæterum foliis mollibus subrepandis typo simillima.

Hab. prope Bitolia Macedoniae (Orph!).

Hæc varietas eadem est ac planta Banatûs sub *A. ochroleucâ* M.B. distributa. Species Biebersteinii differt foliis strictis rigidioribus integrerrimis angustioribus linear-lanceolatis acutissimis calloso-punctatis setulis adpressioribus asperis.

Ar Geogr. Europa borealis et media ab Anglia et Scandinaviâ ad Hispaniam et Italiam boreales atque Rossiam septentrionalem et medium.

2. A. hybrida (Ten. Fl. Nap. I, p. 45, tab. 11) biennis villosa-hispida, caulis erectis superne paniculatis, foliis oblongo-lanceolatis sinuato-dentatis subundulatis, floralibus ovato-cordatis vel breviter lanceolatis, racemis erectis tandem laxis, pedicellis calyce brevioribus fructiferis nutantibus, calyce fructifero inflato campanulato ad medium in lacinias lanceolatas fisso, corollæ cœruleæ tubo calyce longiori, forniciis ovatis velutinis, nuculis oblique ovatis latere altero crebrius tuberculato-rugosis supra annulum subconstrictis ②. *A. undulata* Fl. Græc. tab. 165, Griseb. Spic. II, p. 96 non L. — *A. amplexicaulis* Sm. Prodr. Fl. Græc. I, p. 116!

Hab. in vineis et cultis totius Græciæ in regione inferiori (Sprun!, Heldr!) Thraciæ, Byzantii et Bithyniæ (Griseb., Wied!), insulæ Poros (Wied!), Rhodi (Bourg!), Cretæ (Weiss!), Cyprî (Ky!), Anatoliæ occid. et australis ad Ciliciam usque (Ky!, Ball!), Syriæ in Libano (Bl!, Gaill!), Palestinæ ad Hierosolymam (Röth!).

β. Pinardi. — Corollæ ochroleucæ. — *A. Pinardi* Boiss. in Pin. exs.

Hab. ad vias et ad culta prope Adalia Pamphyliæ (Pinard!, Heldr! Bourg. 191!).

Species inter *A. officinale* et *undulatum* quasi intermedia, a priori radice bienni et foliis undulato-dentatis distincta, posteriori magis affinis et foliorum formâ fere tantum discedens, nam nota e stylo plus minusve elongato omnino inanis est. Cl. Bertoloni eas in unicum conjungit sed cl. Gussone et Heldreich qui omnes formas in locis natalibus observaverunt eas specifice distinctas dicunt; item alii componant.

Ar. Geogr. Italia, Sicilia.

3. A. undulata (L. Sp. 181) biennis cinerascens adpressiuscule tomentello-aspera, caulis paniculatis, foliis linear-oblongis et anguste linearibus margine undulato-crispis obtusis, superioribus

abbreviatis basi breviter decurrentibus, racemis erectis tandem laxiusculis, pedicellis calyce brevioribus tandem nutantibus, calyce fructifero inflato campanulato ad tertiam partem dentato dentibus triangularibus, corollæ cœruleæ tubo calyce longiore, fornicibus ovatis velutinis, nuculis oblique ovatis latere altero crebrius tuberculato-rugosis supra annulum non constrictis ②. Ic. Guss. pl. rar. tab. 16. — Hoffm. et Link Fl. Port. tab. 22. — *A. nonneoides* Fisch. et Mey. Ann. Sc. Nat. 4 Ser. vol. I, p. 35. — *A. clarata* Viv. Eg. Dec. № 23 et herb! (forma pumila). — *Lycopsis elongata* Lehm. Asp. II. p. 264!

Hab. in cultis et collibus Attice (Sprunt), Heldri!, Bœotiae (Sprunt), Byzantii (Wied!), Archipelagi (Urv., Orph!), Rhodi (Heldri!), Smyrnæ (Bal!), Isauriæ (Heldri!), Syrie littoralis in Libanum usque (Gail!), Ægypti ad Alexandriam (Ehr!).

Specimina Orientalia Hispanicis a Linneo descriptis similia sunt.

Ar. Geogr Lusitania, Hispania, Gallia australis, Corsica, Sardinia, regnum Neapolitanum

4. **A. obliqua** (Visian. Ind. Sem. Patav. 1836) biennis ? hirsuta et setuloso-aspera, caule erecto vel ascendentem elato paniculato-ramosissimo, foliis oblongo-lanceolatis repandis acutis inferioribus in petiolum attenuatis, intermediis oblique semidecurrentibus, bracteis calyce duplo brevioribus subcordato-triangularibus, racemis demum longissimis curvatis densiusculis, pedicellis brevissimis demum nutantibus, calycis fructiferi inflati campanulati ad medium quinquefidi lobis obtusis corollæ cœruleæ tubo calyce sublongiore, fornicibus ovatis velutinis, nuculis oblique ovatis et rostratis latere altero crebrius rugosis supra annulum basilarem vix constrictis ②? Ill. Ale. P. Græc. tab. 5.

Hab. ad littus Europæum Bosphori prope Phanaraki (Parolini!), ad littora et in vineis insulae Tenos (Sartorii)

Hujus speciei specimina incompleta et absque fructu maturo tantum vidi; valde affinis *A. officinali* et *hybride*, ab eis praesertim differre videtur racemis multo longioribus flexuosis etiam post anthesin magis densifloris, calycibus minoribus, foliis caulinis inæqualiter decurrentibus sed an hæc nota sat firma? *A. microcalyx* Vis. *A. officinali* forsan nimis affinis huic quoque proxima est sed folia non decurrentia et calycis dentes acuti.

5. **A. ochroleuca** (M.B. Taur. Cauc. I, p. 125) perennis, tota setis adpressis aspera, caulis erectis ramosis paniculatis, foliis setularum tuberculis plus minus calloso-punctatis integerrimis insimis linearis-oblongis basi attenuatis cæteris anguste lanceolatis et linearibus acutis, racemis demum elongatis strictis laxis, bracteis minutis triangularibus striguloso-ciliatis, pedicellis brevissimis vix cernuis, calyce fructifero inflato campanulato ad medium in dentes obtusos fiso, corollæ ochroleucæ, albae vel cœruleæ tubo calyce longiore, fornicibus ovatis tomentosis, nuculis valde oblique rostratis latere altero imprimis areolato-rugosis crebre et minute tuberculatis ȝ. Ic. Bot. Mag.

tab. 1608. — *A. angustifolia* Fl. Græc. tab. 164. — *A. leptophylla* Ledeb. Fl. Ross. III. p. 118! an R. et Sch? — *A. linearifolia* Urv. Enum. p. 20. — *A. angustissima* C. Koch Linn. XXIII, p. 483! — *A. Gmelini* Ledeb. — *A. Lorentii* Alph. D.C. Prodr. X, p. 52.

Hab. in neglectis, collibus argillosis, ad ripas, Græciæ insulæ (Sibth.), Anatolia borealis tota a Bithyniâ ad Armeniam (Wied!), Phrygia (Tchih!), Isauria (Heldr!), Cappadocia (Ky. 187!) et in cacamine montis Alidagh 5000' (Bal. 905!), Tauria (Stev!).

β. canescens. — Copiosius setosa vel tomentosa canescens vel cana. — *A. incana* Ledeb. Fl. Ross. III. p. 117! — *A. tomentosa* Boiss. Diagn. Ser. II. 3. p. 133. — *A. aspera* Boiss. Ann. Sc. Nat. Ser. IV, 2, p. 243.

Hab. in incultis Anatoliæ (Noë!), Armeniæ (Auch. 2329!) et ad Baibut (Bourg!), Cataonie in monte Scherdagh supra Albistan 3-4000' (Haussk!), Cilicie Tracheæ (Tchih!).

Species indumento magis aspero setulis brevioribus adpressioribus basi magis tuberculatis constante, foliis angustioribus acutioribus strictis integrerimis, calycibus vix cernuis ab *A. officinali* distinguenda. Corollæ in Oriente sæpius ochroleucæ quam allæ vel cæruleæ. *A. leptophylla* R. et Sch. et *A. angustifolia* L. ex locis natalibus potius formas angustifolias *A. officinalis* sistere videntur.

Ar. Geogr. Rossia australis.

6. A. Italica (Retz Obs. I, p. 12) perennis setis albis patulis e tuberculo ortis hispida aspera, caule elato ramoso paniculato, foliis oblongis et lanceolatis inferioribus in petiolum attenuatis superioribus sessilibus angustis. ramis floriferis erectis plerumque bifidis, bracteis anguste linearibus, pedicellis erectis calycei æquilongis, calycis profunde partiti laciñiis erectis anguste linearí-acuminatis fructiferis auctis latioribus. corollæ cæruleæ tubo calyce non longiore limbum subæquante, forniciis dense penicillatis exsertis, nuculis erectis angulato-trigonis elevatim et grosse rugosis areolis inter rugas tenuissime pappulosis ♀. *A. paniculata* Ait. Kew I, p. 289. — Fl. Græc. tab. 163. — *A. azurea* Rehb.

Hab. ad vias et in cultis totius ditionis a Zacinthro (Marg!), Græciâ omni! ad Thraciam (Griseb.), Syriam et Libanum ad 5800' usque (Ky!, Bl., Gaill!), Tauriam Caucasum et prov. Transcaucasicas (Ledeb!). Persiam hor. (Szov!), in monte Elbrus (Ky!) et prope Siaret (Bge!), centralem prope Ispahan (Bge!).

Ar. Geogr. Europa australis omnis ab Hispaniâ, Galliâ, Helvetiâ et Germaniâ australibus ad regionem Danubialem et Rossiam australem, Sibiria Uralensis, Africa borealis, Madera

7. A. macrocarpa (Boiss. et Hoh. Diagn. Ser. I, 4, p. 42) perennis tota setis rigidis flavidis tuberculo insidentibus strigulosa, caule elato ramoso paniculato, foliis oblongis inferioribus in petiolum attenuatis superioribus diminutis semiamplexicauli-subcordatis acutis, ramis floriferis erectis bifidis demum elongatis, bracteis lanceolatis

brevibus, pedicellis erectis calyce brevioribus, calycis profunde partiti laciniis strigosissimis lanceolatis demum auctis, corollæ albae tubo calyce subæquanti limbo subæquali longo, forniciis penicillato-pilosissimis exsertis, nuculis erectis oblongo-subtriquetris grosse et obtuse rugoso-tuberculatis ♀.

Hab. in monte Gara Kurdistanæ ad scaturigines (Ky. Alepp - Kurd. exs. 364!).

Valde affinis *A. Italicæ* a qua differt indumento flavido, pedicellis abbreviatis, foliis superioribus oblongis subcordatis nec lanceolato-linearibus acuminitatis, floribus albis, nuculis non areolato-rugosis sed irregulariter et obtuse tuberculatis.

8. *A. strigosa* (Labill. Syr. Dec. III. p. 7. tab. 4) perennis tota setis albis rigidis e tubereulo crasso ortis aculeato-strigosissima, caule elato paniculato, foliis eroso-subdentatis radicalibus obovatis vel oblongis petiolatis superioribus lanceolatis et linearibus acutis, racemis bifidis demum longis, bracteis pedicelli longitudine, pedicellis calyce brevioribus, calycis profunde partiti laciniis obtusis demum auctis, corollæ cœruleo-violaceæ tubo calyce et limbo longiore, forniciis papillis elongatis obsitis, nuculis oblongis erectis triquetris longitudinaliter et elevatim nervoso-costatis ♀. *A. echinata* Lam. Ill. 1821.

Hab. ad vias et culta Syriae (Auch. 2325!) ad Aleppo (Ky!), Damasenm (Gail!), Tripolin, Berythum, Sidonem (Labill!, Bl!, Gail!), Palestinæ ad Gaza (Boiss!). Hierosolymam (Roth!), Naplone (Boiss!), Mesopotamia (Ky. 381!), inter Orfa et Surug (Haussk!), Persiæ australi (Auch. 4981!).

Bipedalis et elatior, setæ eis præcedentium multo strigosiores, nuculæ majoræ longitudinaliter costatae.

9. *A. Barrelieri* (All. Fl. Pedem. N° 184 sub *Buglosso*) perennis, ad caules retrorsum ad folia adpresso et molliter hirsuta pilis minute tuberculatis, caulis stricte foliosis in paniculam foliosam abeuntibus, foliis subdentatis inferioribus oblongis obtusis in petiolum attenuatis, superioribus sessilibus semiamplexicaulibus lanceolatis acutis, racemis conjugatis demum laxis, bracteis squamaformibus, pedicellis calyce brevioribus strictis, calycis quinquepartiti demum accreti laciniis erecto-patulis strigulosis linearibus obtusissimis, corollæ cernue parvæ tubo brevissimo, forniciis semiobtusatis glabris ciliatis, nuculis rectis oblongo-cylindricis longitudinaliter et grosse costato-rugosis ad areolas minute tuberculatis, areolæ basilari laevi ♀. *A. Barrelieri* n.c. Fl. Fr. N° 2731, Bot. Mag. tab. 2349. — *Myosotis obtusa* W. K. Pl. rar. I. tab. 100

Hab. inter frutices et ad ripas herbidas Byzantii (Auch. 2321!), Bithyniae (Wied!), Phrygiae (2321!), Pisidie (Heldr!), Cariæ in monte Cadmo supra Denisleh (Boiss!), Cilicie littoralis (Bal!) et in montibus 4500' (Ky!).

Ar. Geogr. Italia bor. et media, Dalmatia, regio Danubialis, Bosnia, Rossia austalis.

10. **A. cespitosa** (Lam. Encycl. I, p. 498) perennis, e rhizomate sublignoso pluricipi dense cespitosa nana tota setulis e tuberculo ortis obsita et inter eas glanduloso-seabrida. foliis radicalibus crebris linearibus obtusis patentibus caules et flores superantibus, caulinis brevissimis racemo 3-5-floro interdum in rosula subsessili terminatis, calycis quinquepartiti laciniis linearibus obtusis. styllo calyce longiore, corollæ cyaneæ tubo calyce subduplo longiore limbo majusculo, forniciis exsertis oblongis penicillatis, nuculis grosse et areolatim serobiculato-rugosis oblique ovatis rostro elongato ancipi fere horizontali ♀. Ic. Fl. Græc. tab. 169.

Hab. in saxosis montium summorum Sphacioticorum Cretæ, Stravopodia, Volakia, Hagios Theodoros, Hagion Pneuma 6000'-7000' (Tournef., Sibth., Heldr!, Raul!). Fl. Julio.

Folia radicalia 1 1/2-2-pollicaria sesquilineam lata, corollæ eis *A. officinalis* non minores, nuculæ parvæ. Species insignis.

11. **A. limbata** (Boiss. Diagn. Ser. I, 11, p. 99) perennis? tota setis albis basi tuberculatis patulis strigoso-hispida, radice verticali, caulinis erectis brevibus foliosis breviter dichotomis corymbosis, foliis radicalibus numerosis inferioribusque longe linearibus basi subattenuatis, superioribus a basi rotundatâ lanceolatis, racemis per anthesin brevibus densifloris, bracteis triangularibus, pedicellis calyce multo brevioribus, calyce ad medium quinquelobo fructifero inflato campanulato lobis obtusiusculis, corollæ tubo calyce sesquilongiore limbo atropurpureo anguste crateriformi integro subcrenato, forniciis oblongis dense penicillatis antherisque breviter exsertis, nuculis (junioribus) laevibus transverse ovato-subglobosis obtuse et vix rostratis ♀.

Hab. rarissima in colle calcareo Pino Alepensi adumbrato inter Adalia et Jenidje Khan Pamphyliæ (Heldr!). Fl. Marte.

Caules floriferi 4-5-pollicares, folia radicalia 2-3-pollicaria 2 lineas lata, racemi sub anthesi densi scorpioidei serius laxi, calyx florifer 2 1/2 lin. longus, corolla 3 1/2 lin. longa limbo vix lineam dimidiâ lato crenulato nec lobato erecto-patulo insignis. Facies *Alkannæ*, sed nuculæ (juniiores) perforatæ omnino videntur.

12. **A. neglecta** (Alph. D.C. Prodr. X, p. 49) perennis vel biennis viridis tenera setulis brevissimis e tuberculo ortis aliisque minutissimis comparsa asperula, caulinis elatis flexuosis ramosis, racemis ebracteatis demum laxis secundis ex axillis superioribus terminalibusque paniculatis. foliis amplis obsolete repandis radicalibus ovatis vel ovato-oblongis basi rotundatâ vel subattenuatâ petiolatis intermediis basi attenuatâ sessilibus, superioribus acutis sœpe subamplexicaulinibus, pedicellis demum patentibus vel reflexis calyci fructifero subæquilongis glandulosis, calycis quinquepartiti laciniis breviter lanceolatis fructiferis valde accretis longe linearis-lanceolatis nuculas multo superantibus, corollæ azureæ tubo brevi calycem non excedente,

fornicibus transverse ovatis velutinis, limbi explanati lobis latis, nuculis parvis semiovatis rugoso-areolatis et parce tuberculatis rostro brevi obliquo ancipiū ♀. *Myosotis phanicea* Ehrenb. MSS. — *Myosotis Orientalis* Schenk Pl. Sp. p. 26.

Hab. in rupestribus umbrosis, ad rupium verticalium pedem regionis subalpinæ et alpinæ Libani ad Eden, Kanobin, circa Cedros (Auch. 2420! Boiss! Blt!), Antilibano circa Rascheya (Boiss!), Dimar (Gaill!), ad Bludan 3'000'-8000' (Ky. 441), in monte Cassio (Boiss!), Amano (Auch. 1594! Ky. 241!), Mesopotamia prope Diarbekir (Ky!), in monte Sindjar (Haussk!), Kurdistanæ Persie monte Pir Omar Gudrun (Haussk!), Persie occidentalis montibus Elvend et Zerdkou (Auch!).

β. stenocarpa. — Nuculæ angustiores minores semioblongæ

Hab. ad margines umbrosos fluvii Guzeleré supra Sedichig in Ciliciâ littorali (Ball).

Planta bi vel tripedalis, folia tenera inferiora 3-4 pollices longa, corolla brevis et late limbata *Omphalodis* vel *Cynoglossi*, nuculæ in plantâ elatâ parvæ, majores 1½ lineam vix excedentes.

13. *A. myosotidiflora* (Lehm. Asp. № 175) perennis adpresso hirta pallide virens, caulis erectis simplicibus foliosis, racemis ebracteatis tandem laxis ex axillis supremis terminalibusque breviter paniculatis, foliis radicalibus magnis longe petiolatis ovatis vel reniformibus acutis basi profunde et obtuse cordatis, caulinis valde diminutis sessilibus oblongis, pedicellis fructiferis calyce pluries longioribus erecto-patulis, calycis quinquepartiti laciniis lanceolatis obtusis fructiferis parum auctis nuculas non excedentibus, corollæ cæruleæ tubo calycis longitudine, fornicibus transverse ovatis, limbi expansi lobis latis, nuculis semiovatis areolato-rugulosis rostro brevi obliquo ancipiū ♀. *Myosotis macrophylla* Adam in Web. et Mohr. — M B. Cent. Pl. Ross. tab. 14.— *Brunnera myosotidiflora* Stev. Bull. Mosq. 1851, p. 582.

Hab. in umbrosis rupestribus Caucasi centralis in Ossetiâ, distr. Alagir et Radscha alt. 2000'-5400' (M B., Stev! Rupr!)

Caulis 4-6-pollicaris rarius pedalis, foliorum radicalium limbus sæpe 3-4 pollices longus, nuculæ magnit grani miliacei.

Ar. Geogr. Sibiria Altaica.

§ 2. Annuæ.

14. *A. aggregata* (Lehm. Asp. p. 219, tab. 47) annua, setis crebris a bis verrucis insidentibus strigosissima, caulis erectis et diffusis dichotome ramosis, foliis linear-oblängis linearibusque repandis inferiорibus in petiolum attenuatis superioribus sessilibus angustis, racemis terminalibus densissimis corymboso-capituliformibus, floribus subsessilibus, calycis bracteis linearibus brevioris albo-strigosisissimi post anthesiu accreti laciniis deinde triangulari-lanceolatis, corollæ cyaneæ parvæ tubo e calyce breviter exerto limbo parvo,

fornicibus exsertis oblongis penicillatis, nuculis horizontalibus transverse latioribus breviter rostratis rugulosis annulo basili membranaceo inflato laevi ⊖. *A. parviflora* Fl. Græc. p. 57, tab. 167 non Willd. — *A. micrantha* Ræm. et Sch. 4, p. 98. — *Lycopsis glomerata* Urv. Enum. p. 22. — *Echium humile* Desf. Atl. I, p. 65 et herb! ex cl. Guss.

Hab. in arenosis maritimis prope Athenas (Sibth.), insulam Cos (Urv!), Rhodum (Heldr! Bourg!), Cyprum (Ky. 580!), Anatolian occid. (Auch. 2408!), Ciliciam (Bali), Syriam littoralem ad Tortosam et Sidonem (Bl! Gaill!), Pales-tinam ad meridiem Gaza (Boiss!), Ægyptum ad Alexandriam (Ehrenb! Samar!).

Variat subacaulis et caulinus plus minus elongatis interdum pedalibus.

Ar. Geogr. Sicilia, Africa borealis orientalis.

15. **A. hispida** (Forsk. Eg. p. 40) annua tota setulis inæqualibus basi tuberculatis ad folia adpressiusculis aspera, caulinus e collo pluribus prostratis simplicibus vel dichotome ramosis, foliis remote denticulato-repandis margine saepe undulato-crispis lanceolatis obtusis inferioribus basi longe attenuatis, floribus ad axillas solitariis minimis pedicellis brevissimis tandem cernuis suffultis, calycis setoso-hispidissimi demum accreti quinq̄repartiti lacinias lanceolatis denum patulis, corollâ tubulosâ violacea calycem non superante limbo minimo fornicibus papillosis obtusis, nuculis transverse ovatis pendulo-subhorizontalibus angulo acuto rostratis areolato-rugosis ad areolas minute tuberculatis, annulo basili rugoso ⊖. *A. deflexa* Lehm. Ind. Hamb. 1823 et herb!

Hab. in arenâ desertorum Ægypti ad Alexandriam (Ehr! Samar!), inter Kahiram et Suez (Boiss!), Arabiâ Petreâ (Auch. 2332! Schimp. 224! Boiss!), Persiâ in toto ambitu deserti salsi (Bge!). Afghanistan ad Herat (Bge!) et orientali (Griff!), Belutschia (Stocks!), Turkestaniâ in alveo fluv. Jandarja (Lehm!). Arabie *Gahalle*.

Nuculae eis *A. Anchera* et *aggregata* similes paululum magis inclinatae et subpendulæ. Sæpe et in eodem specimine adsunt nuculae solitariae vel binæ abortivæ persistentes et ut in *Paracaryo* calathiformes concavæ margine albido inflato.

Ar. Geogr. Africa borealis in interioribus, Nubia, Songaria, India borealis in Pentapotamide.

16. **A. Aucheri** (Alph. n.c. Prodr. X, p. 49) annua pumila viridis setis albis patulis basi tuberculatis strigosa, collo pluricaulis caulinus erectis vel diffusis, foliis oblongis vel linearis-oblongis obtusis subrepandis basi attenuatis inferioribus petiolatis floralibus ellipticis flores superantibus, floribus axillaribus racemos breves foliatos densiusculos formantibus, pedicellis calyce saepius brevioribus tandem cernuis, calycis albo-hispidissimi quinq̄repartiti denum valde accreti lacinias denum patentibus lanceolatis, corollæ roseo-cærulescentis tubo calycem vix superante limbo parvo, fornicibus oblongis exsertis velutinis, nuculis transverse oblongis subcurvatis horizontaliter rostratis areolato-rugosis ad areolas minute tuberculatis, annulo basili rugoso ⊖. *A. hyalotricha* Boiss. MSS.

Hab. in rupestribus Syriae bor. in monte Cassio (Boiss!), ad Beilan (Hanssk!), in ditione Kassan Oglu (Ky. Suppl. 164!), Ciliciâ (Auch. 230!) et ad Guzeldere (Bal!), Mesopotamia ad Terek (Haussk!), ad radices montis Pir Omar Gudrun Assyria (Haussk!), Armenia Turcica prope Erzerum ad radices Techdagh (Huet!), ad Gumuschkhané (Bourgt!).

Planta 4-6-pollicaris, flores parvi, corolla tubulosa tres lineas longa, nucula horizontalis 2 lineas transverse longa et facies omnino *Nomiae* e sect. *Cryptanthera*, sed ob fornices *Anchuse* adnumerata.

17. A. Ægyptiaca (L. Sp. Ed. I, p. 138 sub *Lycopsis*) annua pallide virens tota setis brevibus e verrucis albis validis ortis strigoso-asperrima, caulinibus diffusis vel ascendentibus dichotome et divaricatim ramosis flaccidis, foliis denticulatis oblongo-lanceolatis utrinque attenuatis infimis obtusis superioribus acutis, racemis laxissimis foliosis, pedicellis supraaxillaribus calyci æquifloris vel longioribus fructiferis sepe recurvis, calycis quinquepartiti lacinias lanceolatis fructiferis valde auctis papilloso-strigosis nuculas duplo superantibus, corollæ sulphureæ tubo calyce breviore, forniciis exsertis apice capitatis velutinis, nucibus crassis semiovatis erectis facie internâ planis rectis externâ convexis grosse et elevatim reticulatis apice acutis annulo basilari rugoso ⊖. *A. Ægyptiaca* n.c. Prodr. X, p. 48 Jaub. et Sp. Ill. tab. 418. — *A. flava* Forsk. Eg. p. 40. — *Asperugo Ægyptiaca* Linn. Jacq. Vind. III, tab. 21. — *A. verrucosa* Lam. Ill. N° 1819.

Hab. ad muros, in neglectis, in cultis Ægypti ad Alexandriam et Cahiram (Ehr! Kral! Sam!), Palestinae (Boiss!), ad Hebron (Ky. 638!), Syria littoralis ad Berythum et Sidonem (Bl!), Damasci (Gaill!), Mesopotamia in monte Sindjar (Haussk!), Cypri (Ky!), Cretæ ad Girapetra (Heldr!), insulae Astypaleæ (Urv!), Persie australis ad Gere (Ky. 36 sub *A. Milleri*!).

Caules interdum pedem et amplius longi fere a basi laxissime floriferi, nuculae 2½ lin. longæ 2 latæ.

18. A. Milleri (Willd. Enum. I, p. 179) annua patule hispida, caule erecto vel subdiffuso, foliis repandis oblongis utrinque attenuatis inferioribus petiolatis obtusis cæteris acutis, floribus ad vel supra axillas racemos laxos foliatos firmantibus, pedicellis calyci æquifloris vel longioribus rectis, calycis hispidissimi qui quinquepartiti lacinias demum auctis linearibus nuculas duplo superantibus, corollæ albo-roseæ tubo calyce subduplo longiore, forniciis exsertis oblongis apice glabris, nuculis rectis semioblongis facie internâ planis externâ subconvexis irreguläreriter rugoso-areolatis apice acutis annulo basilari rugoso ⊖. *Lycopsis Milleri* Horn. H. Hafn.

Hab. in cultis Ægypti ad Alexandriam (Martins! Cadet! Duparc!), in arenosis vallium jugi Sinaïtici (Bové! Schimp. 168! Auch. 233! Boiss!), deserto Syriaco inter Hauna et Palmiyram (Bl!). Arabie *Torr.*

Affinis *A. Ægyptiaca* differt foliis non verrucosis indumento vix strigoso, corollæ tubo exerto, nuculis dimidio minoribus multo angustioribus.

19. A. stylosa (M.B. Taur. Cauc. I, p. 123) annua viridis setis e tuberculo ortis parce obsita, caule gracili erecto angulato superne

ramoso-paniculato, foliis subundulato-repandis inferioribus oblongis in petiolum attenuatis, caulinis sessilibus semiamplexicaulibus, racemis erectis tandem laxis saepius conjugatis. bracteis pedicello brevioribus, pedicellis etiam fructiferis rectis calyce longioribus, calycis quinquepartiti laciniis linearibus obtusis fructiferis auctis patulis, corollæ cæruleæ tubo recto calyce triplo longiore, forniciis albidis obtusis breviter papillosis subexsertis. stylo valde elongato. nuculis oblique ovatis in rostrum breve anceps subcurvatis parce areolato-rugosis ad rugas nitidis ad areolas minute tuberculatis ⊖. Ic. Cent. Pl. Ross. tab. 23. — Rchb. Pl. Crit. VIII, fig. 966.

Hab. in collibus lapidosis et inter segetes Tauriae (M.B., Stev!).

β. major D.C. Prodr. — Indumentum magis strigosum, folia sublatiora, corollæ tubus calycis proportione brevior et limbus major, sed formas intermedias observavi. — *A. Spruneri* Boiss. Diagn. Ser. I, 11, p. 98.

Hab. in Atticâ (Sprun! Heldr!), in vineis et montanis Thraciæ (Friv!), ad Philipoppolin (Janka!), Asiâ minore in Bithyniâ ad Boli (Wied!), ad Sarular (Auch. 499!), in campis ad Buldur Pisidiæ (Heldr!), in Cappadociâ ad Cæsarream (Bal. exs. sub *Lyc. erectâ*!).

20. **A. Thessala** (Boiss. et Sprun. Diagn. Ser. I, 11, p. 99) annua viridis setis e tuberculo ottis aspera, caule gracili erecto parce ramoso, foliis integris vix repandis inferioribus oblongis in petiolum attenuatis superioribus linear-lanceolatis sessilibus acutis, racemis saepius conjugatis demum laxis. calycis strigulosi quinquepartiti laciniis demum accretis linearibus nuculas superantibus, corollæ pallide cæruleæ tubo recto calyce sesquiongiore, forniciis ovatis papillosis subexsertis, nuculis rectis semioblongis parce rugosis et tuberculatis annulo basiliari laeviusculo ⊖.

Hab. inter segetes in Thessaliâ prope Zeitoun (Sprun!).

Species ex unico specimine descripta facie et characteribus valde affinis *A. styllose* a qua sat differre videtur stylo calyce fructifero breviore nec longiore, et præsertim nuculis angustioribus rectis apice sursum directo, nec oblique ovatis rostro horizontali.

21. **A. arvensis** (L. Sp. 199 sub *Lycopside*) annua patule setoso-aspera, caule erecto ramoso corymboso-paniculato, foliis lanceolatis repando-dentatis inferioribus in petiolum attenuatis obtusis cæteris basi sessili semiamplexicaulibus acutis. racemis confertifloris tandem laxis bracteis calyces superantibus vel æquantibus foliatis, calycis quinquepartiti demum subaucti laciniis lanceolatis, corollæ cæruleæ tubo calyce sublongiore medium versus curvato, forniciis papillosis, nuculis transverse ovatis valde curvato-rostratis elevatim reticulato-areolatis ad areolas tuberculatis annulo basiliari rugoso ⊖. Ic. Fl. Dan. t. 435. — *A. arvensis* M.B. Taur. Cauc. I, 123.

Hab. in arvis et cultis, Peloponnesus inter Nisi et Calamata (herb. Fauché!).

Macedonia ad Thessalonicam (Friedr. ex Griseb.), Tauria, Caucasus et Transcaucasia (Ledeb.).

Cl. Steven Enum. Taur. p. 257, *L. Tauricam* prope Sympheropol et Sudak Tauriae crescentem corollâ majore ab *A. arvensi* distinguit sed ipsius Stevenii specimen ex Sudak in Hohen. Pl. rar. exs. *L. Orientalem* sistit.

Ar. Geogr. Europa borealis et media usque ad Hispaniam et Italiam centrales, regionem Danubiale, Rossia omnis, Sibiria Uralensis.

22. A. Orientalis (L. Sp. 199 sub *Lycopside*) annua breviter tuberculato-setulosa, caulis erectis striete et subdichotome ramosis, foliis obsolete repando-dentatis ovato-oblongis inferioribus obtusis in petiolum attenuatis superioribus sessilibus abbrevialis acutis, racemis elongatis laxis foliatis, calycis quinquepartiti demum aucti lacinias linearis-lanceolatis, corollæ cœruleæ tubo calycei æquilongo infra medium curvato et infra curvaturam angustato, forniciis papillosis, nuculis transverse ovatis valde curvato-rostratis opacis elevatim reticulato-areolatis ad areolas valde tuberculatis annulo basilaris rugoso ⊖. *A. Orientalis* Rehb. fil. Ic. Fl. Germ. tab. 109. — *A. ovata* Lehm. Asp. № 266. — *L. micrantha* Ledeb. in Pand. Beitr. I, p. 65. — *L. erecta* Urv. Enum. p. 22! — *L. Taurica* Stev. Taur. p. 257.

Hab. in cultis et inter segetes Anatoliae borealis (Wied!), Cariae (Sibth.), Lyciae ad Elmaliu (Bourg!), Tauriae ad Theodosiam (Urv!) et Sudak (Stev. exs. sub *L. Tauricā*!), Armeniae in parte inferiori montis Bingoldagh 5000' (Ky. 295!), prope Baibut (Bourg!) et Erzerum (Huet!), Ponto Lazico ad Djimil (Balti), Caucaso (Rupr!), Iberia Caucasicæ (Wilh.), ditione Karadagh (Buhsel), Kurdistanæ Persicæ prope Awiheng (Haussk!), Persia boreali in prov. Aderbidjan (Buhsel), Khorassan (Bgel), Persia australi prope Schiraz (Haussk!), Affghania (Griff.).

Folia eis *A. arvensis* latiora et subintegra, racemi laxiores, corolla minor tubo infra medium tubum curvato et superne ampliato, nuculae crebrius granulato-tuberculatae.

Ar. Geogr. Hispania, Rossia australis, Belgium sed ibi prob. cum frumento introducta, Tibetia.

23. A. variegata (L. Sp. 198 sub *Lycopside*) annua hispidula setisque sparsis e tuberculo ortis strigulosa, caulis erectis vel ascendentibus abbreviatis, foliis repando-dentatis dentibus setoso-spinulosis oblongis inferioribus in petiolum attenuatis caulinis semianplexicaulibus suminis acutis, racemis solitariis brevibus laxiusculis inferne bracteatis, calycis strigulosi quinquepartiti tandem subacereti lacinias lanceolato-linearibus, corollæ cyaneæ albo-variegatae tubo calycei æquilongo antrorsum curvato limbi lobis posticis intensius coloratis, nuculis globoso-depressis præter reticulum tenuem laevibus rostro brevi laterali annulo basilaris corrugato ⊖. Ic. Fl. Græc. t. 178. — *Anch. variegata* Lehm. Asp. 223. — *A. perlata* Lam. Ill. 1820. — *Lyc. bullata* Cyr. Pl. rar. I, tab. 11.

Hab. in apricis, ad vias totius regionis inferioris Peloponnesi et Atticae (Sprn!, Orph!, Boiss!), Etoliae (Nieder!), Macedoniae (Friv!), Anatoliae

orientalis (Auch.) et ad Smyrnam (Fleisch!), Iberiæ et Armeniæ Rossicæ (ex C. Koch).

Ar. Geogr Istria, Dalmatia.

Species obscuræ a me non visæ probab. synonyma negligenda.

A. Cesatiana (Fenzl in Frieder. Reise 282) densissime strigosa, foliis radicalibus longe petiolatis ellipticis acutis, caulinis sessilibus oblongis seniamplexicaulibus, floralibus ovatis cordatis, floribus paniculatis, calycis pedicello æquilongi campanulati lobis ovatis acutis, calyce fructifero inflato reflexo pube aureo-ferruginea subglandulosâ strigoso, corollæ atrœcæruleæ infundibuliformis tubo parum exerto limbi lobis rotundatis, fornicibus villosis obtusissimis bifidis ♀.

Hab. in sylvis Arcadie prope Bassæ (Fried.).

Nuculae ignotæ; an forsitan *Pulmonaria mollis* quam ex eisdem locis habeo?

A. parviflora (Willd. Sp. I. p. 759 et herb!) ex unico ramo floribus orbato ramosissimo in quo calyces in foliola mutata sunt et *Echii* ejusdam formam monstrosam exhibente descripta.

NONNEA (Medic. Phil. Bot. I, p. 31). — *Echioides* Desf.
Fl. Atl.

Calyx quinquefidus. Corolla infundibuliformis vel tubulosa tubo recto medium versus squamulis pilosis vel laciniatis obsito. Nuculae 4 liberae basi excavatae ibidemque margine tumido saepius plicato cinctæ. — Herbae perennes et annuae. — Genus ab *Anchusa* squamis fancialibus minoribus non ad faucem sitis sed e medio tubo oriundis tantum et forsitan non sat distinctum. Habitus tamen proprius.

Specierum Orientalium distributio.

SECT. I. ORTHOCARYUM Alph. D.C. Prodr. X. p. 28. — Nuculae reclæ vel subgibbæ rostro sursum directo.

* Annuae.

N. lutea, *obtusifolia*, *rosea*.

* * Perennes.

N. alpestris.

SECT. II. CRYPTANTHERA Alph. D.C. loc. cit. — Nuculae transverse

ovatæ vel oblongæ rostro laterali ad angulum internum directo.

§ 1. **Squamulæ fauciales breves velutinæ.**

* Annuæ.

N. Vivianii, melanocarpa, Philistaea, picta.

* * Perennes.

N. pulla, Persica.

§ 2. **Squamulæ fauciales magnæ longe laciniatæ. —** Grex ad *Anchusam* accedens.

N. intermedia, anchusoides, pulmonarioides, decurrentis, macrosperma, stenosolen.

SECT. III. CYRTOCARYUM Alph. D.C. loc. cit. p. 32. — Nuculae reniformi-curvatæ horizontales obtuse ad medium sinūs renalis affixæ.

N. ventricosa.

SECT. I. ORTHOCARYUM.

1. **N. lutea** (Lam. Ill. N° 1828 sub *Lycopsiside*) annua, pube brevi papillari setisque longioribus hirsuta, caulinis erectis et ascendentibus simplicibus vel ramosis, foliis radicalibus oblongo-lanceolatis in petiolum attenuatis obtusis, caulinis sessilibus basi amplexicaulibus linearilanceolatis acutis integris vel repando-subdentatis, racemis floriferis confertis fructiferis laxiusculis unilateralibus, bracteis oblongo-lanceolatis acuminatis, calyce hispido ad medium quinquefido fructifero valde aucto campanulato nutanti lobis acuminatis, corollæ luteæ rarius cœrulecentis tubo calycem æquante vel excedente, nuculis rectis semioblongis obtusis parce et longitudinaliter nervosis hirtis ♂. *N. lutea* Rehb. Fl. exc. I, p. 338. — *Anchusa lutea* M.B. Taur. Cauc. I, p. 321. — *Lycopsis ciliata* Willd. Sp. I. p. 780 et herb! — *Lycopsis setosa* Lehm. Asp. p. 269. — *Nonnea setosa* R. et Sch.

Hab. in sterilibus saxosis Georgiae et Iberiae Caucasicæ (M.B., Hoh! Szov!), in Persiâ hor. (Buhsel!), prov. Asterabad (Bge!).

β. parviflora. — Corollæ calycem vix superantis limbus diminutus. — *Lycopsis flavescens* C.A. Mey. Enum. p. 98. — *Nonnea flavescens* F. et Mey.

Hab. in proclivibus lutosis ad Sallian et Lenkoran juxta mare Caspium nec non in insula Sara (C.A. Mey! Hohen!).

Corolla variat tubo calycem æquante vel eum parte dimidiâ superante, limbus vel tubo æquilongus vel diminutus, luteus sed interdum (ex cl. Ruprecht. Sched.) margine rubellus vel totus cæruleus.

2. N. obtusifolia (Willd. Sp. I, p. 780 et herb! sub *Lycopside*) annua, viridis flaccida setulis brevibus basi tuberculatis aspera, foliis obovato-oblongis subintegris radicalibus basi attenuatis obtusis, caulinis sessilibus breviter decurrentibus, floribus axillaribus foliis superatis racemos laxos breves formantibus. calycis fere ad medium quinquefidi fructiferi cernui inflati campanulati dentibus acutis, corollæ calycem vix superante albo-cærulescente, nuculis puberulis atris nitidis rectis acutiusculis lœviusculis ♂. *N. obtusifolia* Ræm. et Sch. Syst. IV, p. 71. — *N. lamprocarpa* Griseb. Spic. II, p. 93. — *Anchusa leiosperma* Fl. Pélop. tab. 7, fig. 3.

Hab. in rupestribus umbrosis, ad vias, Peloponnesus (Bory!), Attica ad Kephissum et ad radices Hymetti et Parnes (Sprun!, Heldr!, Orph. Fl. Gra. 166!), Chios et Lesbos (herb. Willd!), Smyrnæ (Heldr!, Bal!), Cilicia littoralis (Bal!), Syria borealis ad Cassii radices (Boiss!), Syria littoralis et Libanus inferior (Bl! Gaill!), Palestina ad Hierosolymam (Roth!).

3. N. rosea (M.B. Taur. Cauc. I, p. 125 sub *Anchusæ*) annua pube brevi viscidâ setisque longioribus hirsuta, caulinis erectis et decumbentibus ramosis, foliis integris vel repandis inferioribus oblongis in petiolum attenuatis. intermediis sessilibus basi latioribus acutis, floralibus triangulari-acuminatis, racemis tandem laxis, calyce ad tertiam partem quinquefido fructifero pendulo valde aucto et inflato campanulato dentibus acuminatis, corollæ roseæ violaceæ vel leucophæ tubo calycem non excedente in limbum majusculum ampliato, nuculis hirtulis gibbis semi-ovatis elevatim areolato-rugosis et subtuberculatis ♂ rarius ♀. Ic. M.B. Cent. tab. 43. — *N. rosea* F. et M. Cat 1835. — *Anchusa versicolor* Stev. Act. Mosq. p. 21, Bot. Mag. tab. 3477. — *Nonnea versicolor* Sweet. — *Echioides grandiflora* Desf. Cat.

Hab. in ruderatis et cultis regionis subalpinæ Caucasi centralis ad fluv. Terek (M.B.), ad Kasbeck (Rehm!), ad Kaischausr (Owerin!), Daghestaniæ 5000'-6000' (Rupr!), Caucasi Iberici (Hohen!), ditionis Talyssch (C.A. Mey.), Ponti Lazici ad Djimil inter messes (Bal!), Armeniæ ad Erzerum (Huet!).

Nuculae eis *N. luteæ* duplo maiores et gibbosiores tuberculatae, corollæ limbus multo amplior. Specimina ex Erzerum Armeniæ radicem validiorem habent et biennia videntur.

4. N. alpestris (Steven in Traus. Linn. Soc. XI, p. 409. tab. 32 sub *Anchusæ*) perennis, pube brevissimâ viscidulâ setisque e tuberculo ortis hispida, caulinis ascendentibus et procumbentibus, foliis eroso-dentatis inferioribus oblongis obtusiusculis in petiolum attenuatis superioribus sessilibus semiamplexicaulibus oblongis acu-

tis, racemis floriferis confertis dein elongatis laxiusculis, calyce albo-hispido ad tertiam partem quinquedentato fructifero pendulo aucto campanulato, corollæ luteæ fauce fulvæ tubo calyce sublongiore limbo amplio, nuculis semiovatis gibbis elevatim areolato-rugosis minute albo-maculatis ♀. *N. alpestris* Don.—*Lycopsis alpestris* Lehm. Asp. p. 260.

Hab. in regione alpinâ Caucasi orientalis (C.A. Meyt), alpes Schahdag ad nives (Stev.), Daghestania 6000'-9000' (Rupr!).

Affinis *N. lutea* sed facile distincta radice perenni, calycibus hirsutioribus minus profunde divisis dentibus abbreviatis et fructiferis minus inflatis, nuculis magis gibbis areolato-scrobioulatis.

SECT. II. CRYPTANTHERA.

§ 1. **Squamulæ fauciales breves velutinæ.**

5. ***N. Vivianii*** (Alph. n.c. Prodr. X. p. 31) annua, brevissime pubescenti-glandulosa prætereaque setis tuberculatis strigulosa, caulinibus ramosis patentí-diffusis, foliis radicalibus obtusiusculis oblongis basi angustatis, caulinis oblongo-ellipticis floralibus triangulari-oblongis acutis calyces superantibus. racemis foliatis fructiferis laxiusculis, calycis quinquefidi fructiferi nutantis inflati campanulati dentibus triangularibus acutis, corollæ calyce sublongioris tubo in limbum atrovioletaceum abeunti, nuculis majusculis glabris transverse oblongis leviusculis albo-maculatis supra annulum basilarem constrictum dentatum concentrice plicatis ♂. *Anchusa venericosa* Vivian. Lib. p. 10, tab. 6, fig. I.

Hab. ad Alexandriam Ægypti versus Campum Cæsarî (Ehrenb!, Samart!).

Facie et characteribus valde affinis *N. nigricanti* quæ tamen differt induimento minus aspero, præsertim nuculis oblique subglobosis rugosioribus.

6. ***N. melanocarpa*** (Boiss. Diagn. Ser. I, 11, p. 96) annua setulis tuberculatis inæqualibus brevibus asperula, caulinibus erectis vel decumbentibus, foliis integris inferioribus oblongo-lanceolatis obtusis basi longe attenuatis cæteris ellipticis basi subcordatis. floralibus breviter triangulari-lanceolatis acuminatis flores superantibus, racemis foliosis tandem laxiusculis, calycibus ultra tertiam partem quinquefidi fructiferis deflexis campanulatis inflatis dentibus triangularibus acutis, corollæ rubellæ vel pallide violaceæ tubo calyce sublongiore limbo minuto, nuculis nigris puberulis transverse oblongis tenuiter areolato-rugosis margine basilari plicatulo ♂. *N. Armena* Stev. Bull. Mosq. 1851. p. 574.

Hab. in Palestinâ circa Hierosolymam (Boiss!), deserto Syriaco inter Hama et Palmyram (Bl!), (ad Aleppo Haussk!), Mesopotamia (Haussk!), Persia australi (Ky. 107 sub *N. pullâ*!, Haussk!), Persia occid. ad Urmiah (Szov!).

Affinis *N. Vivianii*, minus strigosa, corolla pallidior, nuculæ pubescentes elevatim reticulatae.

7. **N. Philistæa** (Boiss. Diagn. Ser. I, 44, p. 96) annua, breviter pubescenti-viscida prætereaque setis basi tuberculatis aspera, caulinibus decumbentibus ramosis, foliis inferioribus lanceolato-oblongis basi attenuatis floralibus triangularibus acuminatis calyces superantibus, racemis foliatis fructiferis densiusculis, calycis ad tertiam partem quinquefidi fructiferi erecto-patentis campanulati dentibus triangularibus acentis, corollæ pallide aurantiacæ tubo calycem æquante limbo mediocri erecto-patulo, nuculis obsolete puberulis transverse ovatis subcurvatis acute areolatis annulo basilari plicato dentato ⊖.

Hab. in Palestinâ australi circa Gaza (Boiss!).

Affinis *N. Vivianii* et *melanocarpe* a quibus corollæ colore, racemis etiam fructiferis densis, nuculis subcurvatis acutius areolato-nervosis differt.

8. **N. picta** (M.B. Taur. Cauc. I, p. 127 sub *Anchusâ*) annua, setis basi tuberculatis strigulosa prætereaque pilis brevissimis viscidulis hirta, caulinibus decumbentibus simplicibus vel divaricatim ramosis, foliis erosâ subdentatis inferioribus oblongo-lanceolatis basi attenuatis, caulinis subdecurrentibus, floralibus anguste lanceolato-linearibus valde acuminatis flores multo superantibus, racemis foliosis tandem laxis abbreviatis, calycis ad medium quinquefidi fructiferi unilateraliter penduli valde aucti campanulati dentibus subulatis, corollâ cœruleâ saturatius pictâ tubo e calyce subexerto limbo parvo, nuculis glabris vel basi puberulis transverse oblongis tenuiter areolato-rugosis annulo basilari rugoso vel laeviusculo ⊖. *N. picta* F. et M. Ind. 1835, p. 43. — *Lycopsis Caspica* Lehm. Asp. 256 et *N. Caspica* Don. — *N. sordida* F. et M. Ind. 1835, p. 43. — *N. diffusa* Boiss. et Buhse Aufz. p. 152 — *N. Szovitsiana* Stev. Bull. Mosq. 1851, p. 574.

Hab. in agris, ad vias, Cappadocia ad Everek (Ky. 234!), ad Caesaream (Bal!), Armenia Rossica (C Koch), Caucasus et prov. Transcaucasicae (M.B., Hohen!, Szov!), Persia borealis (Buhset, Bge!), Persia media (Bge!), Affghania (Griff!), Belutschia (Stocks!), Turkestania (Lehm!).

Omnibus partibus *N. Vivianii* et *melanocarpâ* minor, caules breviores, folia angustiora, floralia imprimis angustissima et valde acuminata, calycis dentes setaceo-acuminati nec triangulares acuti.

Ar. Geogr. Rossia australis orientalis, Songaria, Sibiria Uralensis, India borealis in Pentapotamide.

9. **N. pulla** (L. Sp. 196 sub *Lycopside*) perennis, pube brevi glandulosâ setis basi tuberculatis plus minus copiosis intermixtis pilosa, caulinibus erectis foliosis superne breviter paniculatis, foliis inferioribus oblongo-lanceolatis in petiolum attenuatis, caulinis superioribus lanceolatis basi semiamplexicaulibus, racemis brevibus, floribus secundis, calyce ad medium quinquefido fructifero acreto campanulato dentibus triangulari-lanceolatis acutis, corollæ brunneæ calyce sesquilongioris tubo in limbum campanulatum breviter lobatum erectum ampliato, nuculis depressis transverse breviter ovatis

rostro lateralí rugoso-areolatis interdum puberulis φ . *N. pulla* D.C. Rehb. t. 101. — *N. Taurica* Ledeb. Fl. Ross. IV, p. 112, Stev. Verz. p. 255. — *N. atra* Griseb. Spic. II, p. 94. — *N. Armena* Boiss. et Huet MSS. — *Lycopsis ciliata* C.A. Mey. Enum. p. 98! non Willd.

Hab. in cultis derelictis Thraciae (Griseb.), Bithyniae (Pest!), Byzantii (Barbey!), Pouti supra Niksar (Tchiht!), Armenia Turcica (Calv!), Huet!, Bourg!, prope Van (Noë!), Tauria (Stev!), prov. Caucasicis (M.B.), Persiae prov. Aderbidjan (Szov!), ditionis Talysch (C.A. Mey!).

Variat caulinus abbreviatis vel elatioribus, inflorescentia plus minus condensata vel laxa; specimina Orientalia Germanie ob tuberculos setarum saepe crassiores saepius magis aspera sunt sed non semper.

Ar Geogr. Germania orientalis, regio Danubialis, Rossia media et australis.

10. N. Persica (Boiss. Diagn. Ser. I, 7, p. 32) pereunis, humilis, a collo pluricaulis vel ramosa setis brevibus e tuberculo crassiusculo ortis aspera pube breviori interdum interjecta, foliis radicalibus lanceolatis acutis in petiolum attenuatis, ceteris anguste linearibus sessilibus, floralibus calyce longioribus, panicula brevis ramis divergentibus saepe flexuosis, racemis abbreviatis tandem laxiusculis, calycis ad medium quinquefidi fructiferi nutantis aucti breviter campanulati lobis triangularibus acutis, corollæ tubulosæ calyce sublongioris limbo rubro-violaceo vix distincto, nuculis parvis transverse breviter ovatis lateraliter rostratis elevatim rugoso-cristatis et sub lente tuberculatis φ .

Hab. in montibus Persiae austro-occidentalis, monte Kuh Barfi prope Schiraz (Ky. 328!), montibus Sawers et Nur 41000'-42000' (Hausk!), monte Ssahend prov. Aderbidjan (Szov! Buhsel), ad Ischredabad inter Yezd et Ispahan (Bge!), in monte Dehballo prope Jezd (Buhse!), in monte Totschal prope Teheran (Ky. 588!).

Planta 3-5-pollicaris caulinus erectis vel divergenti-decumbentibus; differre videtur a *N. pullæ* formis foliis angustis, floribus minoribus, corollâ tubulosâ, nuculis minoribus elevatius cristato-rugosis et ad areolas tuberculatis. Ejus statio imprimis alpina videtur.

§ 2. **Squamulæ fauiales magnæ longe laciniatæ.**

11. N. intermedia (Ledeb. Fl. Ross. III, p. 111) perennis, pilis brevibus basi vix tuberculatis tomentella, caulinus erectis simplicibus foliosis supernæ breviter corymbosis, foliis radicalibus oblongis utrinque attenuatis, caulinis mediis et floralibus a basi subcordatâ semi-amplexicauli oblongo-lanceolatis acuminatis, racemis abbreviatis, calycis ad medium quinquefidi fructiferi accreti campanulati nutantis dentibus lanceolatis acuminatis, corollæ violaceæ tubo calyce breviore limbo tubo æquilongo patenti planiusculo, nuculis globoso-depressis elevatim reticulato-rugosis puberulis rostro lateralí φ .

Hab. in prov. Transcaucasia Guriâ (Nordm! Radde!), in Transcaucasia loco non notato (Szov!), ad fluvium Ingour fontes in Suanetiâ Caucasicâ (Radde!).

Pedalis, folia intermedia 2 pollicies longa 5-6 lin. lata. A *N. pullâ* facile distinguitur indumento molliuscule, corollæ limbo multo ampliando nec campanulato erectiusculo, squamarnum fauialium structurâ.

12. *N. anchusoides* (Boiss. et Buhse Aufz. p. 152) perennis, setulis basi tuberculatis hispida, caulis erectis simplicibus stricte foliosis apice breviter paniculatis, foliis linear-lanceolatis acutis sessilis superioribus abbreviatis semianplexicaulibus, racemis etiam fructiferis abbreviatis. calycis ad tertiam partem quinquefidi fructiferi modice aucti campanulati nutantis dentibus lanceolatis, corollæ violaceous tubo calyce sublongiore in limbum eo breviorem erecto-patulum ampliato, nuculis glabris transverse ovatis elevatim rugosis rostro elongato laterali ♂.

Hab. in monte Ssahend prov. Aderbidjan Persiae prope Schoch Jordi 8400' (Buhse!).

Pedalis, folia intermedia bipinnicaria 3-4 lineas lata. A *N. pullæ* formis foliis angustis, squamarum facialium structuræ; a *N. intermediæ* corollæ tubo longiori limboque minori erecto nec patulo, calyce fructifero modice aucto distincta.

13. *N. pulmonarioides* (Boiss. et Bal. MSS. — Dec. 2, p. 4) perennis, pilis brevibus basi subtuberculatis molliusculæ hirta, caule erecto simplici folioso superne breviter corymboso-paniculato, foliis radicalibus lanceolatis sessilis utrinque attenuatis. caulinis superioribus et floralibus a basi subcordato-semianplexicauli subdecurrenti breviter lanceolatis, racemis brevibus, calyce ad quartam partem in dentes etiam sub anthesi breviter triangulares fiso fructifero acreto campanulato nutanti. corollæ rubro-violaceæ tubo calyce breviore limbo tubo æquilongo patenti planiuseculo, nuculis glabris transverse ovatis inter rugas elevatas tuberculatis rostro laterali ♂.

Hab. in regione alpinâ Ponti Lazici supra Djimil 7000' (Bal!).

Pedalis, facies *Pulmonariæ*; affinis *N. intermediæ* a qua specifice differre videtur foliis subdecurrentibus, calycis minus profunde fissi dentibus etiam sub anthesi late et breviter triangularibus nec lanceolatis.

14. *N. decurrens* (C.A. Mey. Enum. p. 98 sub *Lycopside*) perennis, pilis brevibus nonnullis basi tuberculatis asperulo-velutina, caule erecto simplici crebre folioso angulato superne corymbose paucifloro, foliis lanceolatis integris radicalibus basi attenuatis, caulinis anguste decurrentibus acutis floralibus abbreviatis, racemis brevissimis. calyce ultra partem tertiam in dentes lanceolatos acutos fiso fructifero acreto breviter campanulato nutanti, corollæ purpureæ tubulosæ calyce vix longioris tubo in limbum brevem erectum subampliato. nuculis maximis dense velutinis semiglobosis elevatim et acute areolato-rugosis nostro laterali ♂.

Hab. in herbidis alpinis ditionis Talysch 5400'-6000' (C.A. Mey!).

Pedalis, folia intermedia 2-3 pollices longæ, calyx florifer corollaque 3 lineis vix longiores, nucula saepius unica piso vix minor. Facies *N. pulmonarioides* et *intermediæ* sed foliis longiusculæ decurrentibus, corollæ tubulosæ limbo fere indistincto, nuculis magnis velutinis insignis.

15. *N. macrosperma* (Boiss. et Heldr. Diagn. I, 11, p. 97)

perennis, setis albis brevissimis basi crasse tuberculatis pube viscidula sœpe intermixtis scabrida, caulis e radice crassâ erectis pumilis crassis sœpius a basi ramosissimis corymbosis ramorum decurrentiâ angulato-compressis, foliis radicalibus magnis rosulatis oblongis basi longe attenuatis, caulinis valde diminutis lanceolatis basi semiamplexicauli breviter decurrentibus, racemis brevibus tandem laxiusculis, calyce ultra tertiam partem in dentes lanceolatos fiso fructifero acreto campanulato nutanti, corollæ luteæ tubo calyce sublongiore limbo tubo breviore erecto-patulo, nuculis crassis semigloboso-depressis glabris grosse scrobiculato-rugosis albo-maculatis rostro laterali ✘. *N. Euphratica* Boiss. loc. cit. (spec. juniora).

Hab. in collibus incultis Phrygiae ad Ouchak 2800' (Balt!), Anatoliæ bor. ad Beybasar et in monte Elmadagh (Wied!), Pisidiae ad Egirdir (Heldr!), Lyciae ad Elmali (Bourg!), Cappadociae (Ky. 1881), Armeniæ ad Euphratem (Auch. 4978 B!).

Planta semipedalis vel humilior, folia radicalia corymbum sœpe subæquantia. Inter affines caule a basi subdivaricatum ramoso, corollis flavis, etc., insignis.

16. ***N. stenosolen*** (Boiss. et Bal. Diagn. Ser. II. 6, p. 123) perennis, tota setis albis basi tuberculatis pube minimâ intermixtis strigosa pluricaulis, caulis pumilis erectis et diffusis ramosis, foliis brevibus lanceolatis acutis, floralibus angustis acuminatis longe stri-goso-ciliatis flores superantibus, racemis etiam fructiferis densiusculis, calyce dense et adipesse hirsuto prætereaque longe strigoso ad quartam partem in dentes ovato-triangulares fiso fructifero modice aucto, corollâ rubrâ anguste tubulosâ calyce vix sesquiongiore limbo brevi erecto fere indistincto, nuculis parvis semigloboso-depressis glabris tuberculato-rugosis albo-maculatis rostro laterali ✘.

Hab. in Cappadociâ ad radices montis Karamasdag (Balt!).

Caules cum inflorescentiâ semipedales, racemi etiam fructiferi tres pollices non excedunt, folia brevia inferiora pollicaria 3-4 lineas lata, flores 3 lineas longi. Ex unico specimine mihi nota, an forsitan varietas parviflora et microcarpa *N. macrocarpa*?, sed calyces multo magis strigosi et corollæ color diversus.

SECT. III. CYRTOCARYUM.

17. ***N. ventricosa*** (Sibth. et Sm. Fl. Græc. II, p. 58, tab. 168 sub *Anchusæ*) annua, pube brevi adipressâ setisque sparsis tuberculatis asperula, caulis a collo pluribus prostratis vel ascendentibus, foliis oblongis et oblongo-linearibus obtusis radicalibus basi attenuatis, racemo terminali simplici vel bifido tandem laxiusculo secundo, calyce ad quartam partem quinquefido dentibus lanceolatis acutis tandem valde acreto vesicario nutante, corollæ albidae minute tubo calyce sesquiongiore limbo brevi patenti, nuculis reniformibus curvatis utrinque obtusis tenuiter areolato-rugosis ⊖. *N. ventricosa* Griseb.

Spic. II, p. 93. — *N. alba* D.C. Fl. Fr. V, p. 420. — *Lycopsis Sibthorpiana* R. et Sch. Syst. IV, p. 770. — *Nonnea Sibthorpiana* Don — *Nou. amplexicaulis* Hochst. in Ky. Sched. — *Campylocaryon album* D.C. in Ky. Sched. — *N. macrosperma* Gris. Spic. II, p. 93 non Boiss.

Hab. in cultis et ad vias Peloponnesi et Atticæ (Sprun! Heldr!), Thraciæ (Griseb!), totius Anatoliae occid. et australis a Mysiâ ad Ciliciam (Boiss! Heldr! Bal!), Archipelagi (Auch!), Rhodi (Bourg!), Cypri (Sibth.), Syriæ littoralis (Gaill!) et interioris ad Damascum (Gaill!) et Aleppo (Ky. 35!), Mesopotamie (Haussk!).

Ex. am. Rehb. fil. Ic. Germ. 18 p. 60 planta Europæ occidentalis differt ab orientali calyce fructifero ampliori, corollæ tubo basi ampliato, squamis fauclibus brevius fimbriatis, sed eas omnes notas etiam in speciminiis ex Oriente variare comperti.

Ar. Geogr. Hispania, Gallia australis, Dalmatia.

PULMONARIA (L. Gen. 184).

Calyx quinquefidus post anthesin campanulatus. Corolla infundibuliformis fornicibus destituta sed quinque fasciculis pilorum cum staminibus alternantibus instructa. Stylus filiformis stigmate bicapitato. Nuculae 4 leves ovatae basi excavatae supra annulum basilarem levem constrictæ. — Herbæ perennes, collo fasciculos steriles foliorum et caules floriferos edente. Flores in fere omnibus speciebus pseudopolygamî, sexu altero antheris in medio tubo insertis et stylo longo, altero staminibus ad faucem sitis et stylo brevi instructo. Mihi prorsus inexplicabile auctores multos hoc genus *Lithospermeis* nec *Anchuseis* adnumeravisse.

1. **P. officinalis** (L. Sp. 294) foliis radicalibus exterioribus cordatis petiolatis acutis, caulinorum inferiorum petiolo brevi alato superioribus oblongis sessilibus, caule pilis patulis sparsis obsito, corollæ tubo sub aunculo pilorum glabro, nuculis acutis ♀. Ic. Fl. Dan. tab. 482.

Hab. in sylvaticis agri Byzantini (Andr. ex Griseb.), in Caucaso demissiore (M.B.).

Ar. Geogr. Europa sept. et media ab Angliâ et Scandinavia ad Hispaniam et Italiam centrales, regionem Danubialem, Bosniam! Rossiam.

2. **P. mollis** (Wulf. ex Hell. Fl. Wire. Suppl. 13) foliis radicalibus elliptico-lanceolatis acutis in petiolum attenuatis, caulinis sessilibus superioribus semiamplexicaulis, caulis et plantæ totius præsertim in parte superiori pilis molibus flexuosis viscidis, corollæ tubo sub annulo pilorum hirto, nuculis obtusis ♀. Ic. Rehb VI, tab. 696. — *Nonnea auriculata* Boiss. Diagn. Ser. I. 11, p. 95.

Hab. in umbrosis humidis vallis fluvii Neda Messeniæ in via ad Phigaleam (Bory!), in sylvis subalpinis Gurie (Nordm., Szov!), Caucasi septentrionalis (Ledeb.).

Ar. Geogr. Gallia in Pyreneis, Germania australis, Italia borealis, regio Danubialis, Rossia media et australis, Sibiria tota.

3. **P. azurea** (Bess. Prim. Gall. I, 150) setis rigidulis ad caulem patulis ad folia adpressis obsita, foliis anguste lanceolatis acutis radicalibus in petiolum attenuatis caulini semiamplexicaulinibus, corolle tubo sub pilorum fasciculis glabro, nuculis obtusis ♀. Ic. Rehb. VI, tab. 694.— *P. angustifolia* Wahlenb., Schrank, non Mert. et Koch (quæ *P. tuberosa* Schrank), an L. Sp?

Hab. in alpinis, Olympus Bithynus (Noë!), Caucasus septentrionalis (M.B.), ad Kaischaur 6000' (Rupr!).

Ar. Geogr. Galliae et Hispaniae borealis montes, Germania, Dania, Helvetia in Alpibus, Regio Danubialis, Istria et Dalmatia, Rossia media et australis.

SYMPHYTUM (Tourn. Inst. 438, tab. 56).

Calyx quinquesfidus vel partitus. Corolla cylindrico-infundibuliformis limbo subventricoso late et breviter quinquelobo, forniciis longis cum antheris alternantibus. Stamina inclusa. Nuculae 4 liberae recte oblique vel horizontaliter curvatae basi excavatae et ibi annulo cinctæ. — Herbae perennes.

* Fornices inclusæ.

+ Radix fusiformis vel ramosa.

1. **S. officinale** (L Sp. 495) radice fusiformi ramosâ, totum pubeminitatâ aspersum præterea ad caules, foliorum nervos, calyces setis albis basi dilatatis asperum, caulis erectis, foliis inferioribus ovato-lanceolatis in petiolum contractis, superioribus lanceolatis decurrentibus, calycis laciniis lanceolatis acuminatis, corollæ albide vel violaceæ calyce duplo longioris oboconico-tubuloso dentibus breviter triangularibus obtusis recurvis, forniciis inclusis antheras filamentis 2-3-plo longiores superantibus, nuculis ovato-trigonis levibus lucidis supra basin non contractis apice obliquo ♀. Ic. Engl. bot. tab. 817.— Rehb. Ic. tab. 102.

Hab. in fossis, ad aquas, Græcia (ex Sibth.), Bithynia (Thirke!).

Ar. Geogr. Europa fere omnis a Scandinaviâ et Angliâ ad Rossiam, Sibiria Uralensis.

2. **S. Orientale** (L Sp. 495 totum pilis flexuosis molliter hirsutum, radice ramosâ, caulis erectis ex axillis breviter ramosis, foliis subtus densius pilosis oblongis cordatis præter summa sessilia basi rotundata vel breviter cuneata petiolatis, calycis præter indumentum brevius viscidum setulosi laciniis tubo brevioribus lanceolatis acutiusculis, corollæ albæ calyce duplo longioris infundibuliformi-

tubulosæ lobis latis brevibus ovatis erectis, fornicibus inclusis antheras filamentis subæquilongas paulo superantibus, nuculis minutis levibus obliquis apice laterali subcarinato supra annulum basilarum constrictis ♀. Ic. Tourn. Voy. I, p. 524. — Jacq. fil. Ecl. tab. 82.

Hab. in umbrosis Byzantii (Tourn., Cast! Noë!), ad radices Olympi Bithyni prope Broussam (Auch. 2349! Boiss! Barbey!), in Bithynia (Thirkel!), prope Angora (Wied!).

3. **S. Caucasicum** (M.B. Taur. Cauc. I, p. 128) radice fusiformi ramosâ, molliter piloso-hirtum cinerascens setis præter calyces nullis vel paucis, caule erecto paniculato-corymbose, foliis inferioribus oblongis in petiolum attenuatis superioribus anguste et breviter decurrentibus, calyce viscidulo et setuloso ad tertiam partem in dentes lanceolatos fiso. corollæ cœruleæ tubuloso-infundibuliformis calyce triplo longioris lobis breviter et late ovatis erecto-patulis, fornicibus inclusis antheras filamento æquilongas vix excedentibus, nuculis obliquis longitudinaliter areolato-rugosis minuteque tuberculatis ♀. *S. racemosum* Steph. in Willd. herb!

Hab. in dumosis Caucasi sept. ad Mosdok et Kislar (M.B.), Iberia Caucasicæ (Hohen! Rupr!), Sonchetia et prov. Karabagh (Szov!), Caucasi orientalis in Daghestanijā 6500' (Rupr!).

Indumentum molle *S. Orientalis* sed folia inferiora non cordata et superiora decurrentia, corolla calycis proportione longior cœrulea, etc.

4. **S. Tauricum** (Willd. Nov. Act. Berol. III, p. 120, tab. 6) viride piloso-hirtum et secus caules et foliorum nervos setis sparsis basi tuberculatis obsitum, caulibus gracilibus ex axillis ramosis, foliis ovatis acutis inferioribus petiolatis basi cordatis vel rotundatis summis sessilibus. racemis tandem laxis, calycis setosi ultra medium quinquefidi laciniis anguste linearibus obtusiusculis, corollæ albidae infundibuliformi-tubulosæ calyce triplo longioris lobis ovatis erecto-patulis, fornicibus antheras filamento æquilongas æquantibus inclusis, nuculis grosse areolatis sub lente minute tuberculato-punctatis apice subobliquo ♀. Ic. Bot. Mag. tab. 1787. — *S. Orientale* M.B. Taur. Cauc. I, p. 129 non L.

Hab. in umbrosis Tauriae, circa Sympheropol (Stev!), in monte Tchatyr Dagh (Rehm!), Ponto (Tchih!), Ponto Lazico ad Rhizé (Ball), Armenia Turcica (Calv!), Abchasiâ et Imeretiâ (Rupr!).

Affine *S. Orientali* a quo differt indumento non denso molli sed sparso et setis intermixto. foliis brevioribus, racemis tandem laxis, calyce corollæ proportione breviori profundius fiso laciniis angustioribus, nuculis tuberculato-punctatis.

5. **S. sylvaticum** (Boiss. in Bourg. exs. Armen. 1862 — Dec. 2, p. 4) viride, caulibus gracilibus ramosis setis retrorsis basi tuberculatis strigosis, racemis terminalibus brevibus paucifloris basi foliolosis, foliis teneris oblongo-lanceolatis acuminatis denticulatis in

petiolum brevem attenuatis paginâ superiore sparsim et subtus ad nervos breviter tuberculato-setosis, summis diminutis sessilibus, calyce ad medium in dentes lineares obtusos fiso fructifero parum aucto campanulato, corollæ albæ tubuloso-infundibuliformis calyce subtriplo longioris lobis ovatis brevibus, fornicibus inclusis antheras filamentis æquilongas superantibus, nuculis parvis subglobosis obliquis areolato-rugulosis lævibus nitidis ♀.

Hab. in sylvis Ponti ad Calia supra Trapezuntem (Bourg!).

Folia ea *Impatiens noli tangere* referentia, inferiora $2\frac{1}{2}$ - $3\frac{1}{2}$ pollicaria 12-45 lin. lata, racemi breves 5-7-flori foliolosi, nuculæ lineam late apice obliqua. Forma foliorum, nuculis lævibus, etc., ab affini *S. Taurico* distinctum.

6. **S. Anatolicum** (Boiss. Diagn. Ser. I, 4, p. 43) viride præter pubem sparsam minutissimam setis e tuberculo ortis rigidis apice saepe incurvis scabrum, radice ramosâ, caulibus flaccidulis fere a basi subdichotome ramosis, foliis teneris inferioribus oblongis subrepandis in petiolum brevem attenuatis, summis sessilibus internum breviter et anguste decurrentibus, racemis tandem laxis, calyce viscido et setoso-strigoso ad $\frac{1}{2}$, in lacinias lanceolato-lineares obtusiusculas partito basi attenuato fructifero deflexo acreto laciñis patentibus, corollâ albâ vel violacea calyce duplo longiore tubuloso-infundibuliformi lobis brevibus ovatis, fornicibus inclusis antheras filamento æquilongas vix superantibus, nuculis elongatis rectiusculis obsolete areolato-nervosis sub lente tuberculatis ♀. *S. asperrimum* Urv. Cat. p. 20 non M.B.

Hab. in umbrosis montium circa Smyrnam, Mesogi, Tmolo (Boiss! Bal. 308!), in summis insulae Cos (Urv!).

Planta flaccidula, folia majora 2 pollices vix excedentia, a *S. asperrimo* indumento aculeis non constante, corollæ quoad calycem proportione, etc., longe distans, magis affinis *S. Taurico* setis non aduncis, foliis ovato-cordatis, nuculis obliquis diverso.

7. **S. brachycalyx** (Boiss. Diagn. Ser. I, 4, p. 43) viride, pubescentia brevi setis patulis longis secus caules nervos et calyces intermixta obsitum, caulibus erectis subdichotome paniculato-corymbosis, foliis teneris subrepandis inferioribus oblongis in petiolum latum attenuatis ceteris sessilibus, racemis tandem laxis. calyce fructifero parum aucto breviter campanulato deflexo ad tertiam partem in dentes triangulares fiso. corollâ albâ calyce subtriplo longiore late et breviter lobatâ, fornicibus inclusis antheras filamento sesquibreviores vix æquibus, nuculis rectiusculis elevatim rugosis ♀.

Hab. in Cariæ montibus (Pinard exs. sub *S. Orientali*!).

Foliorum inferiorum lamina 2-2 $\frac{1}{2}$ poll. tantum longa A. congeneribus calyce breviter quinquefido etiam fructifero parum acreto et breviter campanulato distinctum

8. **S. Palæstinum** (Boiss. Diagn. Ser. I, 11, p. 94) pallide

virens brevissime pubescenti-viscidum setis paucis patulis intermixtis, radice ramosâ, caulinibus petiolorum decurrentia angulatis, foliis repandis inferioribus obtusis ovatis vel oblongis basi rotundatis vel cordatis, superioribus breviter petiolatis sessilibusque, racemis fructiferis vix laxis, calyce viscido et parce setoso florifero tubuloso ad quartam partem quinquefido fructifero valde aucto patentí-deflexo late oblongo-cylindrico dentibus late campanulatis, corollæ albæ calyce duplo longioris lobis ovatis, fornicibus inclusis antheras filamento sublongiores æquantibus, nuculis rectiusculis elevatim rugosis minute tuberculatis ✘.

Hab. in umbrosis, circa Hierosolymam (Boiss! Ky. 446! Roth), in Antilibano ad Rascheva (Boiss! Gaill!), monte Amano (Haussk!), Ciliciâ ad Tauri radices (Auch. 2351!, Ky. 125! et Bal! sub *S. brachycalyce* et Ky. Suppl. 44! sub *S. Orientali*).

β ? *dentatum*. — Folia acute denticulata.

Hab. in Lyciâ prope Kourmalu ad occidentem urbis Adalia et in monte Solyma (Heldr! Specimina juniora qua tamen calyce convenire videntur).

Indumentum *S. Orientalis* sed brevius parcius et subsetosum; species insignis calycibus fructiferis elongatis cylindricis 6-7 lineas longis; folia superiore non revera decurrentia sed eorum petiolus in caulis angulos productus.

9. **S. Kurdicum** (Boiss. et Haussk. Dec. II, p. 5) pallide virens setis aculeiformibus recurvis ad caulem ramos et nervos foliorum parce obsitum insuper sub lente parce hirtulum, caule elato ramoso, foliis membranaceis repando-denticulatis inferioribus..., intermediis et superioribus petiolatis ovato-oblongis obtusiusculis basi rotundatis vel breviter cuneatis, racemis fructiferis furcatis brevibus densiusculis, calyce strigoso florifero ad medium quinquefido fructifero erecto ampliato acreto oblongo-campanulato ad tertiam partem in dentes late triangulares obtusos fisso, corollæ albæ tubuloso-infundibuliformi calyce duplo longiori lobis breviter ovatis, fornicibus inclusis antheras filamentis breviores vix æquantibus, nucibus supra annulum basilem valde obliquis subhorizontalibus elongatis tenuiter areolato-rugosis minute tuberculatis ✘.

Hab. in Kurdistanâ Per-iæ conterminâ inter Sulimaniën et Sikna (Haussk!).

Ex unico specimine fructifero foliis radicalibus orbato descriptum sed distinctissimum, caulis elatus videtur, folia etiam superiora 3 poll. longa, calyx fructifer ut in *S. Palestino* elongatus amplius 6 lineas longus, nuculae 2 lineas longæ supra basin horizontaliter curvatae.

10. **S. sepulcrale** (Boiss. et Bal. in Bal. pl. Pont. 1866) viride, caulinibus elatis sulcatis ramulosis breviter asperulis, racemis terminalibus brevibus densiusculis bifidis, foliis membranaceis subtus hirtellis superne tuberculatis sparsis muticis vel breviter setigeris obsitis amplis ovato-oblongis acutis vel acuminatis basi rotundatis vel subcordatis inferioribus breviter petiolatis superioribus sessilibus, summis saepius oppositis racemos subæquantibus, calyce ultra medium in

dentes lanceolato-lineares obtusos partito fructifero acreto campanulato, corollæ violaceæ intundibuliformis calyce quadruplo longioris lobis breviter ovatis, forniciis inclusis antheras filamentis æquilongas æquantibus, nucibus obliquis subcompressis lœvibus nitidis 1-2 rugis tantum obsitis annulo basilaris rugoso denticulato ♀.

Hab. in umbrosis subalpinis Ponti Lazici, Djimil ad sepulchra (Bal!).

Planta elata, folia inferiora semipedalia 3-4 poll. lata, racemi fructiferi tripliocares. Facie, foliis amplis, corollis elongatis et calyce profunde partito *S. asperrino* affine sed indumentum non asperrimum, calyx fructifer campanulatus, nuculæ lœves.

11. *S. asperrimum* (Sims Bot. Mag. tab. 929) radice ramosa, caulis elatis ramosis retrorsum setis aculeiformibus muricatis, foliis ad petiolos et nervos tuberculato-setosis subtus iusuper pubescentibus ovato-oblongis acuminatis inferioribus petiolatis basi rotundatis vel subcordatis summis sessilibus cuneatis, calyce setoso-aculeolato ad $\frac{3}{4}$ longitudinis in lacinias lanceolatas obtusas fructiferas elongatas fiso, corollæ celestinæ calyce 4-5-plo longiore infundibuliformi lobis ovatis, forniciis longis inclusis antheras filamentis subaequilongas superantibus, nucibus gibbosis valde rugosis et ad areolas tuberculato-punctatis rostro obliquo anicipiti ♀. *S. aspernum* Lepech. — *S. peregrinum* et *S. echinatum* Ledeb. Sem. Dorp. (formæ hortenses forsitan hybridae).

Hab. in sylvaticis, dumosis, ad rivos regionis subalpinæ Caucasi septentrionalis et centralis 2400' (M.B., Rupr! Rehm!), Daghestaniae 6000' (Rupr!), Guriae (Szov!), Armeniæ, ditionis Talysh et Persiae borealis (Buhse!).

Planta sœpe humanae altitudinis.

+ + Radix tuberosa.

12. *S. tuberosum* (L. Sp. 195) pube minutâ et setis sparsis brevibus basi tuberculatis hirsutum, rhizomate carnoso tuberculiformi obliquo præmorsæ antice fibras edenti, caulis simplicibus, foliis insimis minoribus sub anthesi emarginatis inferioribus ovato-oblongis in petiolum contractis superioribus oblongo-lanceolatis semi-decurrentibus, racemis terminalibus brevibus, calyce fere ad basin in lacinias lanceolatas ciliatas partito, corollæ ochroleucæ infundibuliformis calyce triplo longioris lobis breviter triangularibus extus curvatis, forniciis inclusis, antheris filamento duplo longioribus, nuculis globoso-trigonis obtusis tuberculatis supra annulum basi denticulatum contractis ♀. Jacq. Austr. tab. 225. — Rehb. tab. 103.

Hab. in sylvaticis ad Byzantium (Coumany!), in regione alpinâ montis Scardi Macedoniae (Griseb!).

Ar. Geogr. Hispania borealis et media, Anglia, Gallia media et australis, Helvetia et Germania australes, Italia, Dalmatia, regio Danubialis, Rossia media et australis.

13. *S. grandiflorum* (D.C. Prodr. X, p. 40) pube minutâ et ad

caulem, petiolos, foliorum margines et nervos setis paucis basi tuberculatis hirtum, rhizomate.... caule simplici, foliis infimis minoribus sub anthesi emarcidis inferioribus subcordato-ovalis acutis longe petiolatis summis diminutis basi breviter attenuatâ in petiolum brevem decurrentibus, racemo folio supremo approximatio confertim paucifloro, calyce hirsuto ciliato ad $\frac{1}{2}$, in lobos breviter lanceolatos patentes fiso, corollæ ochroleucae infundibuliformis calyce 5-6-plo longioris lobis brevissime et late semiorbiculatis, forniciis inclusis, filamentis antherâ vix longioribus, stylo incluso, nuculis.... ♀. *S. cordatum* M.B. I, p. 130 non Willd. — *S. Ibericum* Stev. in litt. M.B. Taur. Cauc. III, p. 647.

Hab. in Georgiâ Caucasica (Wilhm. in D.C. herb!).

Pedale, foliorum inferiorum limbus triplicaris, corolla 9-10 lineas longa. Radice ignotâ incomplete descriptum, facie et indumento affine videtur *S. tuberoso* quod differt foliis breviter petiolatis basi attenuatis superioribus magis elongatis, corollâ 6-7 lineas tantum longâ, filamentis longioribus.

* * Fornices longe exsertæ.

14. ***S. bulbosum*** (Schimp. Bot. Zeit. VIII, I, p. 17) pubescens et setis longioribus hirtum, rhizomate cum stolonibus subterraneis hinc inde in tubera subglobosa inflato, caulis simplicibus vel parce ramosis, foliis ovato-oblongis infimis in petiolum attenuatis superioribus sessilibus saepe semidecurrentibus, racemis brevibus paucifloris, calyebus in lacinias lanceolatas semiquinquefidis, corollæ tubulosoinfundibuliformis calyce duplo longioris lobis ovatis rectis, forniciis linearibus denticulatis longe exsertis dimidiâ corollâ longioribus, filamentis antheras aequantibus, nuculis obliquis tuberculatis annulo basilari basi dentato ♀. Ic. Rehb. tab. 104. — *S. filipendulum* Bisch. — *S. Clusii* Gmel. — *S. punctatum* Gaud. — *S. macrolepis* Gay in Rehb. — *S. Zeyheri* Schimp. — *S. brochum* Bor. et Chaub. Pl. Pel. tab. 7.

Hab. in umbrosis regionis montanæ et subalpinæ totius Peloponnesi (Bory! Boiss! Orph!), Atticæ et Bœotiae (Heldr!, Orph!), ad radices Parnassi (Guicci!).

Ar Geogr. Corsica, Italia australis, Germania occidentalis, Helvetia australis, Dalmatia.

15. ***S. Ottomanum*** (Friv. Fl. 1836, p. 439) pube brevi viscidulâ setisque longis patulis hirsutum, rhizomate..., caulis elongatis flexuosis ramosis, foliis ovato-oblongis inferioribus petiolatis cæteris sessilibus, racemis fructiferis brevibus laxis, calyce setosostrigoso ad medium quinquefido, corollæ ochroleucae calyce sesquilongoris lobis ovatis erectis, forniciis linearibus longe exsertis corollæ aequilongis, antheris filamento longioribus, nuculis parvis obliquis parce et elevatim areolato-rugosis asperulis annulo basi non dentato ♀. Ic. Rehb. tab. 104.

Hab. in castanetis umbrosis montis Athos (Griseb!), montis Corthiati (Orph!), in vineis Thraciae prope Carlova (Frivl.).

Ar. Geogr. Banatus.

PSILOSTEMON (D.C. Prodr. X, p. 35. — *Trachystemon* Don — *Nordmannia* Ledeb.).

Calyx quinquefidus. Corolla tubo brevi, limbo in lacinias lineares tubo plures longiores partito. Fornices breves vel elongatæ. Filamenta longe exserta basi ad insertionem papillosa vel hirta. Antheræ distantes. Nuculae 4 libere, obliquæ vel rectæ, basi excavatæ et annulo tævi cinctæ. — Herbæ perennes vel annuæ.

1. **P. Orientale** (L. Sp. 197 sub *Borrago*) rhizomate carnosò cylindrico stolonifero, foliis superne adpresse strigosis subtus ad nervos et venas hirtis radicalibus longe petiolatis amplis cordatis breviter acuminatis, caulis setoso-hispidis folia radicalia vix superantibus erectis nudis in paniculam multifloram foliis parvis sessilibus ovato-oblongis bracteisque parvis ovatis obsitam abeuntibus, calyce ultra medium in lacinias oblongas obtusas partito, corollæ cœruleo-purpurascens tubo calyce sublongiore intus medium versus 5 fasciculis pilorum instructo, limbo tubo triplo longiore in lacinias lineares extus pilosas horizontales revoluto-contortas partito, forniciis parvis ovatis retusis, filamentis longe exsertis pilosulis basi hispidis, nucibus in calyce parum ampliato parvis nigris nitidis transverse ovatis rostro horizontali acuto rugâ fere unicâ annulo tenui integro ♀. *P. Orientale* D.C. Prodr. X, p. 36. — *Borago cordifolia* Mænch. — *Nordmannia cordifolia* Ledeb. Bull. Petersb. 2, p. 312. — Ic. Tourn. Voy. p. 523. — Fl. Græc. t. 175. — Bot. Reg. tab. 288. — *Trachystemon Orientale* Don. — Jaub. et Sp. Ill. tab. 419.

Hab. in sylvis umbrosissimis humidis Byzantii (Tourn., Sibth! Cast! Auch!), radices Olympi Bithyni (Boiss! Barbey!), Bithynia et Paphlagonia (Wied!), Pontus ad Trapezuntum (Bourg!) et Rhizé (Bal!), in Colchide, Abcasia et Imeretiâ (Nordm! Rupr! usque ad 1900!), Guria (Stev!).

2. **P. Creticum** (Willd. Sp. I, p. 778 sub *Borrago*) molliusculæ setuloso-hispida, rhizomate fusiformi, caulis erectis vel ascendentibus flexuosis foliosis fere a basi ramosis, foliis a basi ovatâ vel subcordatâ ovatis et oblongis obtusis repando-petiolatis inferioribus et intermediis petiolatis summis diminutis sessilibus, racemis junioribus scorpioideo-capitatis demum elongatis laxis, calyce ad medium in lacinias lanceolatas fisso, fructifero valde acreto nutanti campanulato, corollæ albidae tubo calyce breviore imâ basi glandulosò limbo tubo $2\frac{1}{2}$ -plo longiore in lacinias lineares erecto-patulas glabriusculas partito, forniciis linearibus papillosis longe exsertis sed laciniis corollæ brevioribus, filamentis glabris longe exsertis fornices superan-

tibus rubris breviter ad insertionem papillosis, nuculis oblongis rectis areolato-rugosis sub lente punctatis annulo lævi ♀. Ic. Desf. Coroll. tab. 17. — Fl. Græc. tab. 176. — *P. Creticum* D.C. Prodr. X, p. 36.

Hab. in rupestribus umbrosis, speluncis Cretæ (Sibth., Heldr!), Rhodiæ (Bourg!), Zacynthi (Sibth.), Peloponnesi in Messeniâ (Bory! Heldr!), Eubeæ (Auch. 2341!). Specimina ex Athone a Sibthorpio citata ad *Sympyrum Ottomanum* facie simile prob. spectant.

SUBTR. III. LITHOSPERMEÆ Koch. Syn. — *Echieæ* et *Lithospermæ* D.C. Prodr. — Nuculæ quatuor rarius binæ uniloculares areolâ planâ toro insidentes a gynobasi vel toro liberæ.

PODONOSMA (Boiss. Diagn. Ser. I, 11, p. 113).

Calyx quinquepartitus. Corolla tubulosa, fauce nuda basi nectario-circulare instructa. Filamenta ad medium partem usque corollæ adnata, antheræ elongatae basi sagittatae et liberæ apice membranâ linearî retusâ appendiculatæ exsertæ. Stylus glaber stigmate indistincto. Nuculæ valde curvatæ areolâ basiliari planâ substipitata affixæ rostro horizontaliter flexo — Herba perennis. Genus nuculis et facie *Alkannæ* donatum sed ab eâ fauce nudâ et præsertim antheris longis apice appendiculatis *Onosmatis* distinctum.

1. **P. Syriacum** (Labill. Syr. Dec. III. p. 8, tab. 5 sub *Onosmatis*) totum pube minutâ glandulosâ setis basi tuberculatis tenuibus plus minus intermixtâ obsitum viride fragile, ramis elongatis flexuosis ramulosis crebre foliosis, foliis sessilibus oblongo-lanceolatis acutis inferioribus basi attenuatis superioribus semiamplexicaulibus, racemis foliosis demum laxis, bracteis lanceolatis calyce longioribus, calycis demum accreti laciniis a basi latiore anguste linearibus, corollâ glabra caerulecente laciniis lanceolatis acutis demum reflexis, nectario hirsuto, antheris ultra medium exsertis, nuculis crebre tuberculato-asperis ♀.

Hab. e fissuris rupium verticalium et murorum dependens, Cilicia (Ky!), Syria omnis calidior et montana ad Tripolin (Lab!), Sidonem (Gaill!), Nahr el Kelb prope Berythum (Bl!), Darkusch ad Orontem (Boiss!), montem Gebel Abiad deserti ad Palmyram (Bl! forma magis setosa foliis diminutis), Hierosolymæ ubique *Hyoscyami auræ* et *Parietariae Judaicæ* socium (Boiss! Gaill!); Assyria prope Mossul (Auch.), Persia australis prope Gere (Ky. 68!), circa Kaserun, Schahpour, Kumaredj, etc. 2000'-6000' (Ky!). Fl. vere et aestate.

ONOSMA (L. Gen. 159).

Calyx quinquepartitus in unicâ specie quinquetidus. Corolla tubulosa vel clavata regularis fauce nuda apice breviter quinquedentata

intus imâ basi squamâ nectariformi lobulatâ semper instructa. Filamenta corollæ medio tubo adnata superne plus minus longe libera. Antheræ longæ apice membranâ pellucidâ appendiculatæ basi sagittato-bicalcaratâ sæpius inter se cohærentes, rarius liberæ vel totâ longitudine concretæ. Stylus longus, stigma subbilobum. Nuculæ rectæ acutæ triquetro-ovatae vel oblongæ nitidæ læves rarius tuberculato-scorbiculatae, areolâ basilari planâ. — Herbae vel suffrutices floribus sæpe anthesi peractâ versicoloribus. — Species in Florâ Orientali numerosissimæ, interdum quoad characteres difficile limitandæ: induimenti structura divisionem facilem præbet, sed species nonnullæ quoad tuberculos setarum glabros vel stellatim pilosos intermediæ sunt.

Specierum Orientalium distributio.

§ 1. **Haplotricha.** — Foliorum setæ tuberculis glabris insidentes.

* Biennia.

+ Nectarium glabrum.

O. polyanthum, echoïdes, setosum, tinctorium, Græcum, Taygeteum, stenosiphon, longilobum.

+ + Nectarium hispidum.

O. Aleppicum, bulbotrichum.

* * Perennia.

+ Antheræ fere ad apicem connatæ (*Macrostemon* Bge).

O. stamineum, rostellatum.

+ + Antheræ basi cohærentes vel omnino liberæ.

A. Nectarium glabrum.

✗ Foliorum setæ adpressissimæ.

O. fruticosum, sericeum, flavum, pachypodium, Bodeanum, liparioides, polyphyllum, simplicissimum, halophilum.

✗ × Foliorum setæ adpressiusculæ vel patentes, caulis et inflorescentiae patentissimæ.

1. Humilia, caules floriferi semipedales vel breviores.

O. nanum, velutinum, Troodi, cespitosum, frutescens, strigosissimum, rupestre.

2. Elatiora.

O. microspermum, Kotschyï, Cassium, orcodocum.

B. Nectarium hispidum.

O. procerum, gracile.

§ 2. **Heterotricha.** — Foliorum setæ tuberculis ad folia inferiora saepius glabris, ad superiora præsertim in paginâ inferiori stellatim pilosis incidentes.

+ Nectarium pilosum.

O. lanceolatum.

+ + Nectarium glabrum.

O. Kilouyense, chlorotrichum, Olivieri, hebebulbum

§ 3. **Asterotricha.** — Foliorum setæ tuberculis stellatim pilosis incidentes. — Nectarium semper glabrum. Antheræ basi exceptâ semper liberæ.

* Perennia.

+ Elatiora, caules floriferi pedales vel longiores.

O. Isauricum, Bourgæi, ovalifolium, molle.

+ + Humiliora. Caules vel rami floriferi semipedales vel breviores.

O. dasytrichum, latifolium, albo-roseum, cœrulescens, mutabile, Rascheyanum, mite, Armenum, Roussæi, stellulatum.

* * Biennia.

O. Spruneri, giganteum, auriculatum.

§ 1. **Haplotricha.**

* Biennia.

1. **O. polyanthum** (D.C. Prodr. X. p. 63) pallide virens totum setis brevissimis tuberculo glabro incidentibus adpressis asperulum inter setas minutissime papillosum. caule unico crebre et stricte folioso, foliis linear-lanceolatis acutis radicalibus basi attenuatis sub anthesi exsiccatis, caulinis sensim abbreviatis basi rotundatis, racemis terminalibus, capitulis fructiferis non elongatis corymbosis, calycis fructiferi vix aucti lacinias linear-lanceolatis obtusis viscido-hirtis sparsim setulosis, corollæ albida viscido-puberula calycem quartâ parte excedente lobis breviter ovato-triangularibus recurvis, nectario glabro, antheris filamento vix longioribus inclusis, nuculis ovato-triquetris nitidis acutis subtuberculatis ②.

Hab. in Cappadocia ad Euphratem (Auch. 2361).

Sesquipedate, facie *Cynoglossum* referens, folia radicalia tripollicaria 4 lineas lata sursum sensim abbreviata, corymbus fere capituliformis, corolla cylindrica superne parum ampliata 7 lineas longa. Radix cylindrica verticalis et collum vaginis foliorum radicalium squamosum hanc speciem setis brevissimis insignem biennem esse suadent.

2. **O. echooides** (L. Sp. 196) totum setis e tuberculo glabro ortis ad folia adpressiusculis obsitum inter eas puberulum vel glabrum, caule pyramidatum ramoso, foliis linear-lanceolatis obtusis inferioribus basi longe attenuatis, superioribus et bracteis oblongo-linearibus, racemis simplicibus vel bifidis fructiferis laxis, calycis fructiferi vaide accreti lacinis linearibus, corollæ ochroleucæ pubescenti superne dilatata calyce quartâ parte longiore dentibus brevibus ovatis, nectario glabro, antheris inclusis filamento sesqui vel duplo longioribus, nuculis ovato-trigonis acutiusculis levibus lucidis ②. Ic. Jaq. Austr. tab. 195. — *O. arenarium* W.K. Hung. tab. 279. — *O. Gmelini* Ledeb. Alt. I, p. 184. — Icon. tab. 280.

Hab. in Tauriæ et Caucaso (M.B., Stev.). Saepè cum *O. setoso* et *microcarpo* ab auctoribus confusum in ditione rarius videtur. Vidi specimen authentica ex Caucaso septentrionali ad fluvium Terek (Ower. Herb. Acad. Petrop!). *O. arenarium* W.K. ab *O. echoide* Gallico et Germanico distinguere non valeo, rhizoma nec plus nec minus biennæ vel perennans, character antheræ margine glabrae vel scabrae omnino incertus, sed *O. echooides* ex Hispaniæ et Africæ boreali specific differt nuculis majoribus malleato-scorbiculatis interdum lateraliter subbicornutis, an *O. triceropspnum* Lag!. *O. Transylvanicum* Schur differre quoque videtur.

Ar. Geogr. Gallia australis, Germania media occidentalis et australis, Helvetia australis, Dalmatia, regio Danubialis, Rossia media et australis, Sibiria Uralensis et Altaica.

3. **O. setosum** (Ledeb. Fl. Alt. I, p. 181. — Icon. tab. 196) totum setis e tuberculo glabro ortis ad folia adpressiusculis strigosum cinerascens inter eas glabrum vel asperulum, caulis elatis stricte foliosis superne stricte et breviter ramuloso-pyramidalis, foliis linearibus obtusis margine revolutis inferioribus et intermediis elongatis, floralibus abbreviatis lanceolatis acutis, racemis etiam fructiferis brevibus parum laxis, calycis demum accreti lacinis linear-lanceolatis, corollæ ochroleucæ velutinae calyce tertiatâ parte longioris apice sub-dilatata lobis brevibus ovatis, nectario glabro, antheris apice sub-exsertis filamento suo sublongioribus, nuculis magnis trigonis acutis malleatis et sepius obtuse rugoso-tuberculatis ②. *O. Visianii* Clem. Bull. Mosq. excl. Syn. Visian. Fl. Dalm. et *O. calycinum* Stev. Ind. Petrop. IX.

Hab. in agris aridis Tauriæ (Stev., Beaupré!), Georgiæ Caucasicæ (Hohen!), Armeniae ad Nackitschewan (Szovt!).

β. dichroanthum. — Inter setas pubescens, corolla versicolor demum cærulescenti-violaceens. — *O. dichroanthum* Boiss. Diagn. Ser. I, 11, p. 107. — *O. strigosum* Stev. MSS. — *O. Caspium* Gruner Bull. Mosq. 1867, p. 441.

Hab. in Transcaucasiâ ad Baku (C.A. Mey!), Persiâ bor. ad radices montis Demavend (Ky. 388!), inter Siaret et Schahrud (Bge!), Affghaniâ (Grif!), Belutschia ad Zahree et Doubund (Stocks!).

O. echoide elatius, folia magis stricta, corolla longior magis exserta, nuculæ duplo maiores tuberculatæ vel saltem malleatæ.

Ar. Geogr. Rossia australis!

4. **O. tinctorium** (M.B. Taur. Cauc. I. p. 131) totum setis tenuibus patentibus basi tuberculo glabro minuto insidentibus pubeque brevi intermixtâ cinerascens, caulis crassiusculis modice elatis pyramidatim ramosis, foliis radicalibus et inferioribus late linearibus elongatis obtusis, superioribus et floralibus abbreviatis oblongis, racemis simplicibus vel bifidis paucifloris fructiferis elongatis laxis, calycis hispidissimi fructiferi accreti laciniis sublinearibus, corollâ albida dein roseâ miniatâ tandem violacea velutinâ calyce tertiatâ parte longiore tubulosâ superne parum dilatata dentibus triangularibus minutis, nectario glabro. antheris inclusis filamento brevissimo, nuculis triangulari-ovatis acutis lœvibus nitidis marmoratis ②. Ic. M.B. Cent. Pl. Ross. tab. 79. — Lehm. Asp. tab. 11. — *O. echoides* var. *parviflorum* Ledeb. Fl. Ross. III. p. 125.

Hab. in Tauria (M.B., Stev.), Caucaso et Iberia Caucasicâ (M.B., herb. Petrop!).

Ab *O. echoide* valde distinctum corollâ cylindrica duplo minore, filamenti parte liberâ brevissimâ. Varietas β *flavicomâ* D.C. Prodr. p. 64 est planta perennis ab *O. tinctorio* distincta sed ex specimine juniori et incompleto non rite determinanda.

Ar. Geogr. Rossia australis et orientali-australis, Songaria.

5. **O. Græcum** (Boiss. Diagn. Ser. I. 11, p. 106) totum setis tenuibus adpressis tuberculo minuto glabro insidentibus pubeque brevi velutinâ cinerascens vix strigosum. caule crassiusculo erecto mediocri parce et striete ramoso subpaniculato, foliis radicalibus crebris lanceolato-linearibus obtusis basi attenuatis superioribus et floralibus a basi rotundata oblongis, racemis brevibus paucifloris fructiferis laxiusculis, calycis fructiferi valde accreti laciniis linearibus obtusiusculis, corollâ pubescenti anguste tubulosâ calyce subdupo longiore luteâ demum violascenti dentibus brevissimis triangulari-ovatis. nectario glabro. antheris summo apice vix exsertis filamento brevissimo, nuculis trigonis acutis profunde serobiculatis et tuberculatis ②. *O. tinctorium* Fl. Pelop. non M.B.

Hab. in siccis Peloponnesi ad Navarin (herb. Fauché!), Hymetti (Boiss!), Cretæ (Sieb. sub *O. echoide*) ad Caneam (Weiss!) et Malaxa (Raul!), insulæ Syra (Boiss! Weiss!).

Rarius pede dimidio elatius, corollâ anguste tubulosâ insigne hâcque notâ ad *O. tinctorium* accedens nuculis lœvissimis alienum.

6. **O. Taygeteum** (Boiss. et Heldr. Diagn. Ser. I. 11, p. 101)

flavido-virens totum setis longis tenuibus patentibus e tuberculo glabro ortis strigosissimum inter setas scabridum, canle a basi subdivaricatum ramosissimo, foliis linearibus, floralibus abbreviatis pedicello longioribus, racemis paucifloris saepius cymose bifidis fructiferis laxis, calyce strigosissimo fructifero valde acreto laciniis lanceolatis binis vel ternis in unam latiorem saepissime coalitis 3-4-partito, corollâ albida clavata velutinâ calyce vix longiore dentibus ovatis brevissimis, nectario glabro, antheris subsessilibus apice breviter exsertis, nucibus in rostrum anceps acutum abrupte angustatis opacis sub lente tenuiter granulatis ②.

Hab. inter alvei exsiccati lapides regionis alpinæ Taygeti inter Varvara et Hagios Paraskevy (Heldr!).

Ex unico specimine hucusque notum et ulterius investigandum, affine videtur *O. Graeco* sed corollâ clavata et seminum sculpturâ discedens. In eodem specimine occurunt calyces (pauciores) quinquepartiti, sectio *Aponosma* igitur nullo pacto servanda.

7. *O. stenosiphon* (Boiss. Diagn. Ser. I, 7, p. 35) setis e tuberculo glabro ortis longis patulis aspernum inter eas patule hirtum, caulinibus e rhizomate violaceo pluribus pumilis superne vel fere a basi subcorymbose ramulosis, foliis radicalibus anguste oblongo-lanceolatis obtusis basi longe attenuatis, caulinis et bracteis breviter linearibus, racemis scorpioideis fructiferis subelongatis densis, calyce albo-hirsutissimo laciniis anguste linearibus fructifero elongato tenui longe pedicellato patulo, corollæ e sicco rubræ tomentellæ anguste tubulosæ calyce vix quartâ partè longioris lobis minute triangulatis obtusis, nectario hispido nuculis parvis subtriquetris acutis levibus nitidis ②.

Hab. in Persiæ australis alpe Kuh Daëna (Ky. 721!), prope Ispahan (Auch. 4994! Derderian!), ad Gerdekuh inter Kerman et Ispahan (Bge!), in montibus prope Jezd (Buhse!), in Affghaniæ jugo Kuschlak (Griff!), Belutschia ad Doubound (Stocks!).

Semipedale pluricaule, fructiferum interdum 9-10-pollicare, folia radicalia 1-1½ poll. longa 3-5 lin. lata, racemi fructiferi 2-3 pollicares densiusculi, calyx fructiferi tenuiter pedicellatus 5 lineas longus. Species corollâ angustissime tubulosâ apice nou inflatâ 4-5 lineas longâ valde insignis.

8. *O. longilobum* (Bge Helioe. p. 12) setis e tuberculo glabro ortis patulis hirsutissimum inter eas patule et dense pubescens, caulinibus e rhizomate violaceo pluribus pumilis, foliis linearispathulatis inferioribus basi attenuatis, racemis terminalibus 2-3 capitatis scorpioideis in corymbum confertis fructiferis vix elongatis, ealycis albo-hirsutissimi fructiferi subelongati et longiusculi pedicellati laciniis anguste linearibus, corollâ pallide violascenâ glabra cylindrica calycem vix excedente ultra medium in laciniis lanceolatas erectas partitâ, nectario hirsuto, antheris subliberis apice subexsertis, filamentis brevissimis, nuculis ovatis subcompressis abrupte acutatis nitidis levibus ②.

Hab. in rupestribus apricis Persiae borealis prope Schahrud (Bge!). Fl. Jun. Semipedale, facies et indumentum albo-pilosissimum *O. stenosiphon*. Calyx fructifer 7 lineas longus, corolla 5 lineas longa. Species inter omnes corollam profunde partita insignis.

9. *O. Aleppicum* (Boiss. Diagn. Ser. I, 41, p. 107) totum setis albis longis patulis e tuberculo glabro ortis strigosum inter setas patule puberulum cinerascens, caulis erectis crassis a basi vel a medio stricte ramosis paniculatis, foliis oblongo-linearibus inferioribus obtusis elongatis basi attenuatis, floralibus et bracteis anguste linear-lanceolatis acutis, racemis paucifloris cernuis fructiferis laxiusculis. calycis valde setosi fructiferi elongati laciniis laxiusculae lanceolato-linearibus, corollæ albidae glabrae calyce quartâ parte longioris cylindrico-subclavatae lobis latis brevissimis obtusis. nectario ciliato, antheris apice subexsertis filamento subæquilongis, nuculis magnis triquetris carinatis acutis grosse tuberculatis ②. *O. echinatum* D.C. Prodr. (quoad pl. Syriacam).

Hab. in collibus aridis Syriae ad Aleppo (Ky. 465!), sub *O. echoide*, Auch. 1732!, in Palestinâ inter Hierosolynam et Jericho (Boiss!), in desertis inter Damascum et Antilibanon (Boiss!), in Mesopotamiâ ad Aintab, Bir, Orfa, desertum Chabur (Haussk!).

β. xanthotrichum. — Setæ flavidæ tuberculis minoribus. — *O. xanthotrichum* Boiss. l. cit. p. 109.

Hab. in Antilibano ad Zebdani (Boiss! Ky. 455!), circa Damascum (Gaill!).

Pedale, ample paniculatum, facies *O. echoidea* a quo nectario ciliato, nuculis tuberculatis, etc., longe distat, ab *O. bulbotricho* corollâ glabra discedens.

10. *O. bulbotrichum* (D.C. Prodr. X, p. 64) totum setis longis patulis albis vel flavidis e tuberculo glabro ortis strigosum pubere breviore intermixta, caulis crassis erectis vel ascendentibus fere a basi ramosis saepè divaricatis, foliis oblongo-lanceolatis vel linearibus radicalibus obtusis basi longe attenuatis, superioribus brevibus acutis, racemis demum valde elongatis semper densiusculis, calycis fructiferi aucti laciniis lanceolatis, corollæ lutescentis velutinae clavatae calyce tertia parte longioris lobis breviter triangularibus subreflexis, nectario hirsuto, antheris apice subexsertis filamento subæquilongis, nuculis magnis trigonis parce et grosse tuberculatis ②. *O. echinatum* D.C. Prodr. X, p. 64 ex parte quoad pl. Persicam. — *O. Chorassanicum* Bge Helioc. p. 12 (nomen solum).

Hab. in Persia (Auch. 2297! et 4993!), prope Schiras (Ky. 350!), inter Seytun et Bebehan (Haussk!), inter Teheran et Tabris (Buhse!), in ditione Khorassan (Bunge!).

Strigiosus *O. Aleppico*, corolla major magis exserta pubescens nec glabra, nectarium longius hirsutum.

* * Perennia.

11. *O. stamineum* (Ledeb. in Eichw. Casp. p. 10, tab. 3) basi

suffruticosum multiceps setis patulis tuberculo glabro insidentibus strigulosum inter eas pubescens, caulinis rigidis erectis simplicibus pumilis, foliis parvis sessilibus inferioribus anguste linearibus superioribus brevioribus oblongis acutis, racemo simplici brevi paucifloro, calycis demum vix aucti longe setosi laciniis lanceolatis, corollæ cylindricæ cyaneâ glabrâ calyce subdupo longiore lobis triangularibus reflexis, nectario glabro, antheris connatis omnino exsertis, nuculis acutis levibus nitidis ♀.

Hab. in Turcomaniâ ad mare Caspium (Eichw! Karel! Lehm!).

Caules semipedales vel parum longiores, folia inferiora 6-7 lin. longa sessilis lineam lata superiora breviora et sublatiora

12. O. rostellatum (Lehm. Asper. II. p. 374) basi suffruticosens totum setis adpressiusculis tuberculo glabro insidentibus et prope brevi canescens, ramis numerosis fragilibus pumilis ramosum, foliis parvis sessilibus oblongis et oblongo-lanceolatis utrinque attenuatis, racemo simplici brevi tandem laxiuseculo, calycis hirsutissimi demum vix acereti laciniis lanceolatis, corollæ cylindricæ pubescentis albide vel cœrulecentis calyce dimidiâ parte longioris dentibus breviter triangularibus erectis, nectario hirsuto, antheris connatis ultrâ medium exsertis appendicibus longis cœruleis, nuculis acutis leviusculis opacis ♀. *O. setigerum* Spreng Neu Entd. III, p. 159. — *O. incanescens* D.C. Prodr. X. p. 58.

Hab. ex rupium abscissarum fissuris dependens. Mesopotamia in monte Sindjar (Haussk!). Assyria prope Mossul (Auch. 2315!), inter Bagdad et Kermandchah (Oliv!), prope Schiraz Persiae (Auch. 4983! Ky. 298!). Fl. Mai.

Vix semipedale, folia majora 8-9 lin. longa, corolla vix 4 lineas longa.

13 O. fruticosum (Labill. Syr. Dec. III. p. 10, tab. 6) totum setis adpressissimis tuberculo glabro insidentibus obsitum inter eas pubescens canum fruticosum ramosissimum humile dumosum, ramis crassis rigidis intricatis demum denudatis subspinescentibus, foliis parvis sessilibus breviter oblongo-lanceolatis margine subrevolutis, floribus 1-3 approximatis, calycis adpresso setosi laciniis linear-lanceolatis obtusis, corollæ (e sicco albâ vel ochroleucâ) cylindricæ glabrâ calyce duplo longiore lobis ovatis erectis, nectario glabro, antheris basi coherentibus cæterum liberis ultra medium exsertis filamento antherâ subbreviore, nuculis (immaturis) subtuberculatis ♂. Ic. Fl. Græc. tab. 174.

Hab. in Cypri monte S^{te} Crucis (Labill! Sibth.), ad Chrysostomo prope Buffavento 2000' (Ky. 445!). Fl. Apr.

Dumus pedalis et ultra, folia majora 5-6 lin. longa, corolla 5½ lin. longa. Unica prater *O. dasytrichum* species generis ob ramos lignosos persistentes fruticosa dicenda. *O. stamineum* ob antheras exsertas refert sed illæ basi tantum coherent.

14. O. sericeum (Willd. Sp I, p. 774) totum setulosis adpres-

sissimis e tuberculo minimo glabro ortis densissimis sericeo-cinereum, rhizomatis suffruticosi ramis procumbentibus ramos breves steriles rosuliferos caulesque floriferos in racemum bifidum vel in paniculam brevem abeentes edentibus, foliis rosularum et inferioribus ovato-et oblongo-spathulatis obtusis in petiolum attenuatis, superioribus sessilibus lanceolatis, racemis demum laxiusculis, bracteis linearis-setaceis, calyce pedicellato laciniis latiuscule linearibus acutis vel ad basin usque distinctis quinquepartito vel inter se plus minusve coalitis quadri-vel tripartito, fructifero valde aucto basi indurata carinatum quinquegibbo laciniis accretis late lanceolatis basi subangustatis, corollâ ochroleucâ adpresso velutinâ clavatâ calyce tertia vel quartâ parte longiore lobis brevissimis latis obtusis, nectario glabro, antheris inclusis filamento longioribus, nuculis ovatis acutis subbicarinatis laevissimis nitidis albis ♀. Ic. Lehm. tab. 16. — *O. elegans* C. Koch Linn. XVII, p. 306. — *O. tripartitum* Hochst. in Lorent Reise, p. 334.

Hab. in siccis collinis, montanis et subalpinis Anatoliæ borealis ad Amasia, Tokat, Mersiwan (Wied!), Armeniæ Turcicæ ad Baibut (Bourg!), Erzerum (Calv! Huet!), Musch (Ky!), Armeniæ Rossicæ (Buhse!), Transcaucasiæ occid. ad Borschom (Radde!), Iberiæ Caucasicæ (M.B., Hohen!), Persiæ prov. Aderbidjan (Szov!, Auch. 4988!).

Calyces 5 vel 4-3-partiti in eodem specimine variant. Variant quoque corollæ longitudo, filamenta parte liberâ plus minus elongata.

15. ***O. flavum*** (Lehm. Mag. Ges. Nat. Berl. 8, p. 92. tab. 4 sub *Colsmanniæ*) totum setuloso adpressissimum e tuberculo minimo glabro ortis densissime sericeum saepius flavidum, rhizomatis suffruticosi ramis procumbentibus ramos breves steriles rosuliferos caulesque floriferos in racemum bifidum vel in paniculam brevem abeentes edentibus, foliis rosularum et inferioribus ovato-et oblongo-spathulatis obtusis in petiolum attenuatis, superioribus sessilibus oblongo-lanceolatis, racemis demum subelongatis, bracteis linearibus, calyce 5 vel laciniarum coalitione 4-3 partito fructifero valde aucto basi indurata acute 5-carinato-gibbo laciniis late oblongo-lanceolatis basi attenuatis, corollâ ochroleucâ apice velutinâ clavatâ calyce tertia vel quartâ parte longiore lobis brevibus reflexis, nectario glabro, filamento anthera multo breviore, nuculis ovatis acutis subbicarinatis scrobiculato-subtubercolatis nitidis ♀. *Onosma flavum* Vatke Dissert. Ined. Jan. 1855. — *O. flavidum* Boiss. Diagn. Ser. I. 11, p. 100. — *O. brachysolen* Boiss. loc. cit. p. 104. — *O. trachytrichum* Boiss. loc. cit. p. 103.

Hab. in siccis collinis et montanis ditionis Kassan Oglu Ciliciæ orientalis (Ky. 136! sub *O. sericeo*), Libani (Labill! Bl!), Antilibani circa Rascheya, Zebdani etc. (Boiss! Gaill!, Ky. 139!), Syriæ bor. ad Aleppo, Aintab etc. (Auch. 2301!), Haussk!), Mesopotamia (Ky!), monte Sindjar (Haussk!), monte Avroman Kurdistaniae Persicæ (Hausek!). Persiæ in montibus Teheran (Ky. 399!) et prov. Aderbidjan (Auch. 4987!).

Forsitan *O. sericei* varietas, ab eo nuculis obsolete scrobiculatis nec laevibus

tantum distinctum, nam cæteræ notæ, filamenti longitudo, indumentum flavidum vel cinereum ex speciminibus numerosis collatis valde variant.

16. O. pachypodium (Boiss. Diagn. Ser. I, 11, p. 104) pallide virens totum setis sparsis e tuberculo glabro ortis adpressissimis pubeque minimâ adpressâ obsitum basi suffrutescens, caulis erectis flexuosis ramosis, racemis demum elongatis laxis, foliis radicalibus..., caulinis oblongis acutis utrinque attenuatis, bracteis lanceolato-setaceis minimis, calyce in lacinias lineares quinquepartito fructifero acreto pedicello sensim incrassato suffulto tubo obconico lœvi, corollâ extus velutinâ cylindrico-clavatâ calyce duplo longiore lobis brevibus ovatis, nectario glabro, antheris inclusis subsessilibus, nuculis oblongis acutis obsolete scrobiculatis ♂.

Hab. in schistosis faucis Murch Mahal montis Elbrus Persiæ bor. (Ky. 458!).

Caules pedales, corollæ pollicares, calyx fructifer 9-10 lineas longus. Etiam si *O. sericeo* et *flavo* affine specifice differre videtur setis sparsis, corolla longiori, calycis fructiferi tubo non quinquangulari sed lœvi et pedicello incrassato suffulto, lacinias etiam fructiferis anguste linearibus nec dilatatis.

17. O. Bodeanum, basi suffrutescens multicaule pallide virens setis e tuberculo glabro ortis pilisque intermixtis omnibus adpressissimis obsitum, caulis gracilibus firmis simplicibus foliosis racemo simplici sub anthesi capitato paucifloro terminatis, foliis parvis acutis vel acuminatis inferioribus lanceolatis basi attenuatis superioribus diminutis semiamplexicaulibus, calyce adpresso setoso etiam sub anthesi elongato lacinias lanceolatis, corollâ clavatâ brevissime tomentellâ calycem tertiatâ parte excedente (e sicco) violacea lobis breviter triangularibus demum reflexis, nectario glabro, antheris apice tantum subexsertis, nuculis... ♂.

Hab. in montibus Bachtiaricis Persiæ (Bode!).

Caules 6-9-pollicares numerosi rigidi, folia inferiora 1-1 ½-pollicaria 4-5 lineas lata, calyx florifer 8 lineas longus lacinias lineam latis, corolla 11 lineas longa; affine *O. sericeo* et *flavo* a quibus corollâ violacea vel versicolori differre videtur.

18. O. liparioides (D.C. Prodr. X, p. 65) totum setulis basi minute tuberculatis pube brevi intermixtis adpressissimis canescens sericeum, caulis ascendentibus pumilis foliosis in racemum pauciflorum sub anthesi capitatum interdum bifidum abeuntibus, foliis parvis oblongo-spathulatis obtusis prater floralia oblongo-linearia acutiuscula sessilia basi attenuatis subpetiolatis, calycis patule ciliato-plumosi lacinias linearibus, corollæ hirtæ albidae cylindricaæ calyce vix longioris lobis ovatis revolutis, nectario glabro, antheris inclusis filamento brevissimo, nuculis... ♂.

Hab. in monte Keschischdag Cappadociæ (Auch. 2307! et 2433!).

Caules e rhizomate verticali 4-5-pollicares, folia rosularum sterilium et inferiora 6 lineas longa, caulina 8 lineas longa, calyx 6 lineas longus lacinias vix

lineam dimidiā latis, corolla vix 7 lineas longa. Indumentum ut in *O. sericeo* adpressissimum nitidum sed ab ejus formis caulinis nanis, calycis plumosi laciniis angustiis, corollā brevissimā alienum.

19. ***O. polyphyllum*** (Ledeb. in Pand. Beitr. I, p. 72) basi suffrutescens cespitosum, caulinis pumilis patule villosis in racemum capitatum saepius bifidum abeuntibus, foliis rosularum confertissimis adpressissime setulosis canis longe et angustissime linearibus acutis margine revolutis, caulum floriferorum et bracteis brevioribus patule villosis non revolutis, floribus magnis. calycis plumosi laciniis longis lanceolato-linearibus, corollā albidā brevissime velutinā clavatā calyce quartā parte longiore dentibus triangularibus reflexis, nectario glabro, antheris apice subexsertis filamento æquifolius, nuculis triquetris abrupte acutatis lèvibus ♀. Ic. Ledeb. tab. 24.

Hab. in rupestribus Tauriae meridionalis ad Bachtchisarai, ad fluvium Alma, in valle Sudak et circa Simeis (Ledeb., Steven!).

Caulis floriferi 3-5-pollicares, folia rosularum $1\frac{1}{2}$ -2-pollicaria linea longa, corolla 9-10 lineas longa; species indumento duplice foliorum inferiorum adpressissimo cano, partis superioris plantae patulo flavidio, foliis longis angustissimis insignis.

20. ***O. simplicissimum*** (L. Sp. 196 excl. synon.) setis sparsis basi tuberculatis pilisque brevioribus omnibus adpressissimis obsitum flavidō-virens a basi suffrutescenti multicaule, caulinis aliis brevibus sterilibus aliis floriferis foliosis rigidis simplicibus in racemuni bifidum sub anthesi capitatum fructiferum laxum brevem abeuntibus, foliis anguste linearibus margine revolutis acutiusculis caulum sterilium longioribus confertis fertilium strictis superne sensim abbreviatis, bracteis calyce brevioribus, calycis adpresso hirsuti fructiferi vix aucti laciniis lanceolatis brevibus. corollā albidā brevissime velutinā valde clavatā calyce triplo longiore dentibus late et breviter ovatis. nectario glabro, antheris inclusis filamento lato brevioribus, nuculis ovato-triquetris carinatis lèvibus acutis ♀. Ic. M.B. Cent. tab. 57 (folia nimis acuta).

Hab. in Iberia (M.B.), Georgiā Caucasicā (Hohen.). E ditione nondum vidi.

Caulis floriferi $\frac{1}{2}$ -1-pedales, folia inferiora sesquipollīcaria 1-1 $\frac{1}{2}$ lin. lata, corolla valde clavata 10 lin. longa.

Ar. Geogr. Rossia media et australis, Turcomania borealis, Songaria, Sibiria Uralensis.

21. ***O. halophilum*** (Boiss. et Heldr. Diagn Ser. I, 11, p. 105) pallide virens totum setis adpressissimis subdistantibus aliis majoribus e tuberculo glabro ortis sub lente obsitum, caulinis e rhizomate verticali infrarosularibus ascendentibus simplicibus in racemum bifidum fructiferum elongatum abeuntibus, foliis radicalibus et inferioribus lanceolatis obtusiusculis basi longe in petiolum attenuatis, caulinis et bracteis brevibus lanceolato-linearibus acutis, calycis

adpresso setulosi lacinis linear-lanceolatis acutis, corollæ albidae breviter velutinæ clavatae calyce 2 $\frac{1}{2}$ -pollicare longioris dentibus ovatis subreflexis, nectario glabro, antheris apice subexsertis filamento æquilongis, nuculis parvis ovato-trigonis acutis lævibus nitidis ♀.

Hab. in salsis hyeme inundatis planitiei Lycaonicæ inter Koniah et montem Karadagh sitæ (Heldr!). Fl. Jun.

Caules cum racemo-6-10-pollicares, folia radicalia cum petiolo 2-2 $\frac{1}{2}$ -pollicaria tres lineas lata, racemi fructiferi 3-4-pollicares, corolla 9 lineas longa. Affine *O. simplicissima* sed basi non suffrutescens. caules infrarosulares, folia latiora.

22. *O. nanum* (B.C. Prodr. X, p. 63) totum setis flavidis longis creberrimis e tuberculo glabro ortis patulis ad folia adpressiuseculis hispidum cespitosum, rhizomate pluricipiti dense squamoso, caulis erectis nanis in racemum capitatum pauciflorum abeuntibus, foliis linearibus longiuseculis obtusis margine subrevolutis nervo medio subtus prominenti. floralibus angustatis acutis, floribus majusculis, calycis villosissimi lacinis anguste linearibus, corollâ brevissime velutinâ albida tandem cærulecente cylindrico-subclavata calyce subdupo longiore dentibus ovato-triangularibus. nectario glabro, antheris apice vix exsertis, antheris filamento subæquilongis, nuculis.... ♀. *O. decipiens* Schott et Ky. Sched.

Hab. in regione alpinâ, Cappadocia ad Euphratem (Auch. 2304!), Taurus Cilicicus supra Bulgharmaaden (Balt!), ad Gusguta et Gisyl Tepe 6000' (Ky. 61).

Folia inferiora 12-16 lin. longa 2 lata. caules 2-4-pollicares, flores pro plantæ magnitudine ampli 10-11 lineas longi.

23. *O. velutinum* (Boiss. Diagn. Ser. I, 11. p. 105) totum pilis longis basi vix tuberculatis patulis ad folia adpressiuseculis obsitum vix asperum cespitosum. rhizomate crasso dense squamato albo-hirsuto, caulis erectis nanis in racemum dense capitatum saepius bifidum abeuntibus, foliis planis non elevatim nervosis radicalibus linear-spatherulatis obtusiusculis basi attenuatis. floralibus anguste linearibus acutis. calycis villosi fructiferi vix aucti lacinis anguste linearibus, corollâ albida demum cærulea glabriuseculâ calyce duplo longiore cylindrica subclavata dentibus brevibus ovatis obtusis, nectario glabro. antheris inclusis filamento æquilongis, nuculis parvis ovato-trigonis lævibus nitidis carinatis apice acuto anicipiti-compres- sis ♀.

Hab. in regione alpinâ Olympi Bithyni (Boiss!).

Caules 3-4-pollicares, folia radicalia bipollicaria 2-3 lineas lata, corolla 9 lineas longa. Valde affinis *O. nano* a quo differre videtur indumento molli non strigoso, foliis nec margine revolutis nec elevatim nervosis An tamen hujus varietas?

24. *O. Troodi* (Ky. Cypr. p. 283) totum setis albis basi subtu-

berculatis adpressis dense vestitum asperulum canum a basi sustru-
tescente cespitosum multicaule nanum, caulinis crebre foliosis in
racemum dense capitatum abeuntibus, foliis minimis infimis rosulatis
ovato-oblongis spathulatis, caulinis strictis lanceolatis acutis basi atte-
nuatis. calycis laciniis linearis-lanceolatis, corollæ parvæ flavidæ caly-
cem quartâ parte excedentis hirtæ cylindricæ dentibus triangularibus
acutis. nectario glabro, antheris apice subexsertis filamento quadruplo
longioribus. nuculis... ♀.

Hab. in fissuris rupium montis Troodos insulae Cypri (Ky. 754!). Fl. Junio.
Caules 2-3-pollicares, folia radicalia 4-5 lineas caulina 6-8 longa $1\frac{1}{2}$ lata.
Flores 10-15 conferti, calyx 4-5 lineas, corolla 6-7 longa. Differt ab affini *O. nano* indumento multo densiori adpressissime cano non strigoso, foliis minutis,
corollâ cylindricâ minutâ, filamentis brevibus.

25. *O. cespitosum* (Ky. Cypr. p. 282) basi suffrutescens ces-
pitoso-pluricaule setis albis basi subtuberculatis erecto-patulis den-
sissimis pube brevi intermixtis canum. caulinis nanis ascendentibus
densissime foliosis in racemum pauciflorum capitatum abeuntibus,
foliis confertis crassis margine subrevolutis inferioribus et rosularum
oblongo-spathulatis cæteris lanceolatis. calycis laciniis plumosis lan-
ceolato-linearibus, corollæ hirtæ citrinæ parvæ calycom quartâ parte
superantis cylindricæ dentibus triangularibus, nectario glabro, anthe-
ris inclusis filamento brevissimo, nuculis... ♀.

Hab. in fissuris rupium verticalium Cypri montis Buffavento sub ruinis
Castello della Regina locis imperviis 20' (Ky. 445! in herb. Vindob.). Fl. fine
Aprilis.

Planta 2-3-pollicaris dense foliosa, folia rosularum et inferiora 7-8 lineas
longa 3 lata, superiora angustiora. calyx florifer 6 lineas longus. Species ex
speciminibus imperfectis tantum nota ab *O. Troodi* foliis duplo majoribus et
imprimis crassioribus distincta.

26. *O. frutescens* (Lam. Ill. N° 1837) basi suffruticosum et
ramosissimum totum setis e tuberculo glabro ortis ad folia adpressis
asperum inter setas saepius puberulum, ramis ascendentibus flexuoso
foliosis, foliis radicalibus oblongo-spathulatis, caulinis oblongo-et
linearis-lanceolatis obtusiusculis, racemis 1-2 terminalibus capitatis
fructiferis vix elongatis, calyce setoso fructifero valde ampliato laci-
niis lanceolatis conniventibus. corollâ glabrâ rarius puberulâ superne
ampliata luteâ calyce tertia parte longiore lobis breviter triangularibus
demum reflexis. nectario glabro, antheris breviter exsertis fila-
mento longioribus. nuculis magnis ovatis acutiusculis laevissimis
nitidis marmoratis ♀. *O. echooides* Fl. Greec. tab. 172 non L. — *O. Tournefortii* Griseb. Spic. II. p. 80 ex parte quoad citat. Sibthorpii.
— *O. Orientale* Lehm. Asp. II. p. 376 et herb!

Hab. ad rupes Peloponnesi, in Messeniâ, ad Spartam, Leondari, Naupliam,
Acrocorinthum (Sprn! Boiss!), in montibus Malevo et Kyllene (Orph!), Atticâ
in Hymetto et Lykabetto (Boiss!, Heldr!), Acarnania (Nieder!), Anatolia ad
Euxini littora (Sibth.), Caria (Pin!), Lyciâ ad Elmalu (Bourg!), Syriâ ad Antio-

chiam (Bl!), Sidonem et Libanum (Gaill!), Antilibanum ad Ainhata (Gaill!), Palestinæ ad Banias et in Carmelo (Boiss!). Fl. vere.

Species pulchra ramis semipedalibus, floribus capitatis, corollæ magnæ exsiccatæ sæpe rubellæ vel violascentes.

27. O. strigosissimum (Boiss. Diagn. Ser. I, 11, p. 102) setis e tuberculo glabro ortis longis patentibus strigosum inter eas glabrum, a collo ramosum ramis erectis et ascendentibus pumilis foliosis, foliis radicalibus oblongo-spathulatis in petiolum brevem attenuatis, caulinis breviter lanceolatis, racemo sæpius bifido capitato fructifero vix elongato, calyce setoso fructifero valde accreto laciniis lanceolatis, corollâ glabrâ flavâ superne ampliatâ calyce florifero sesquiongiore lobis parvis triangularibus demum reflexis, nectario glabro, antheris apice exsertis, nuculis magnis globosis acutis nitidis malleatis obsolete granuloso-tuberculatis ♂.

Hab. in Pamphyliâ (Pestal!), in rupestribus maritimis ad Adalia (Bourg. 1881).

Semipedale, valde affine *O. frutescens* sed differre videtur collo non suffrutescenti, foliis longe et patule strigosis inter setas glabris, nuculis malleatis subtuberculatis.

28. O. rupestre (M.B. Taur. Cauc. I, p. 132) totum setis albis e tuberculo ortis pube brevi intermixtâ hispidum cinerascens pluricaule, caulinis pumilis cæsiis ascendentibus simplicibus vel parce ramulosis, foliis parvis breviter linearis-spathulatis obtusis floralibus acutis, racemo simplici vel bifido sub anthesi scorpioideo-capitato demum subelongato, calycis fructiferi vix accreti laciniis anguste linearibus, corollâ cylindricâ flavidâ brevissime velutinâ calyce tertia parte longiore dentibus triangularibus rectis, nectario glabro, antheris inclusis filamento lato duplo longioribus, nuculis minimis triangulari-oblongis abrupte et longe acutatis lævibus nitidis ♀. Ic. Lehm. tab. 12. — *O. tenuiflorum* Willd. Sp. I, p. 775. — *O. diffusum* Boiss. in Huet pl. exs. (forma subramosa).

Hab. in rupestribus Iberiae Caucasicæ circa Tiflis (M.B., Hohen! Rupr!), Imeretiae ad Borschom (Radde!), Armeniæ Turcicæ ad Bayazid (Auch. 4990!), inter Ispir et Tortum (Huet!), Cappadocie ad rivum Kamechly Tchai (Bal!).

Caules 4-6-pollicares fructiferi interdum sublongiores et magis ramosi, folia $\frac{1}{4}$ -1 pollicem longa $1\frac{1}{2}$ -2 lin. lata, corolla angusta apice vix dilatata 5 lin. longa. Species elegans cum nullâ aliâ nisi formis minutis *O. microspermum* commutanda a quibus tamen corollis cylindricis angustis, calyce fructiferi vix aucto et nuculis longe acutatis distinguitur.

29. O. microspermum (Stev. in Hoh. Talysh p. 75) totum setis albis patulis e tuberculo glabro ortis plus minus copiosis et validis strigosum imo strigosissimum, caulinis sæpius numerosis ascendentibus simplicibus vel ramosis, foliis sæpius brevibus inferioribus linearis-spathulatis in petiolum attenuatis superioribus et floralibus breviter lanceolatis, racemis demum elongatis laxiusculis, calycis fructiferi modice accreti laciniis lanceolatis, corollâ breviter velutinâ

ochroleucâ clavatâ calyce duplo longiore dentibus triangularibus reflexis, nectario glabro, antheris inclusis aut apice breviter exsertis filamento sublongioribus, nuculis parvis breviter triquetris acutis levissimis nitidis ♀? *O. microcarpa* D.C. Prodr. X, p. 62. — *O. concinnum* D.C. Prodr. loc. cit. p. 64. — *O. Gmelini* β *microcarpum* Ledeb. Fl. Ross.

Hab. in siecis Armeniæ Turcicæ ad Ispir (Huet!), Erzerum (Calvi), Rossicæ (Seidl! Szov!), Imeretiæ (Radde!), Iberiæ Caucasicæ, Karabagh et ditionis Talyisch (C.A. Mey! Hoh!), Persiæ bor. in prov. Aderbidjan (Szov!), ad radices montis Elbrus (Ky. 437! et 476! sub *O. bulbotricho*), ad Kohrud et Teheran (Bge!), ad Ispahan (Auch. 4995!), ad Persepolim (Ky. 395! sub *O. echinato*), in monte Avroman Kurdistanie Persiæ (Haussk!).

Ab *O. echooidis* formis et *O. setoso* distinctum foliis abbreviatis, calyce fructifero dimidio minore, corollâ calycis proportione longiore et evidentius clavatâ, nuculis tandem duplo vel triplo minoribus abbreviatis acutissimis; videatur insuper perennis nec biennis, caulinis e rhizomate sœpe plurimis et ascendentibus. Variat plus minus strigosum et tunc *O. bulbotrichum* referens, sed hoc semper præter calyces fructiferos majores et nuculas magnas tuberculatas, nectario hirsuto distinguitur.

30. ***O. Kotschy*** (Boiss. Diagn. Ser. I, 7, p. 34) basi induratum pluricanle, caulinis tenuibus rigidis albidis elongatis simplicibus vel parce ramulosis, foliis pallide virentibus setis brevibus e tuberculo glabro ortis asperrimis inter eas glabris breviter oblongo-lanceolatis inferioribus basi attenuatis superioribus semiamplexicaulibus acutiusculis, racemis paucifloris demum laxiusculis, calycis laciniis tenuissime linearibus, corollæ albidae tomentellæ calyce parum longioris lobis breviter triangularibus acutis, nectario hirsuto, antheris inclusis vel apice subexsertis filamento triplo longioribus, nuculis.... ♀. *O. trachyanthum* Boiss. in Ky. exs.

Hab. in Persiæ australi ad Perezend (Auch. 4985! et 4992!), in fauibus alpis Kuh Delu et ad radices montis Sabst Buschom prope Schiraz (Ky. 411! et 503!).

Caules pedales et sesquipedales, folia majora pollicem longa 5 lineas lata, corolla 5-6 lineas longa. Insigne caulinis tenuibus sed duris, foliis breviter setosis sed asperrimis.

31. ***O. Cassium*** (Boiss. Diagn. Ser. I, 11, p. 102) caulinis ascendenteribus simplicibus elatis foliosis setis patulis sparsis obsitis, foliis viridibus setis minutis tuberculis parvis glabris insidentibus parce obsitis inter eas glabris oblongis obtusis superioribus semiamplexicaulibus, racemis paucis terminalibus paniculato-corymbosis fructiferis elongatis densiusculis, calyce fructifero pedicellato erecto et ampliato laciniis lanceolatis, corollâ glabrâ flavidâ superne ampliatâ calyce tertia parte longiore lobis late et breviter ovatis obtusis demum revolutis, nectario glabro, antheris apice subexsertis filamentis brevissimis, nuculis parvis ovatis carinatis acutissimis nitidis marmoratis ♀.

Hab. in regione montanâ Cassii Syriae borealis (Boiss!), Amani prope Beilan (Ky. 42!).

Planta 1-1 $\frac{1}{2}$ -pedalis, folia infer. 1-1 $\frac{1}{2}$ -pollicaria 6-8 lin. lata, corolla 7-8 lineas longa, calyx fructifer 7 lineas longus.

32. O. oreodoxum (Boiss. Diagn. Ser. I, 41, p. 103) caulis erectis vel ascendentibus simplicibus elatis setis longis basi tuberculatis hispidis inter eas puberulis, foliis eodem indumento obsitis radicalibus oblongo-lanceolatis obtusis subpetiolatis, superioribus abbreviatis semiamplexicaulibus ovato-oblongis acuminatis, racemis paniculato-corymbosis fructiferis elongatis, calycis albo-hirsutissimi laciniis anguste linearis lanceolatis, corollæ albidae glabrae calyce quartâ parte longioris lobis triangularibus acutissimis denum reflexis, nectario glabro, antheris ultra medium exsertis filamentis brevibus, nuculis ovatis carinatis acutissimis laevibus nitidis marmoratis ♂.

Hab. in fanebus montis Climax et in fauce Tsimboukkhan Pisidiae inter Isbarta et Adalia (Heldr.). Fl. Mai.

Sesquipedale vel bipedale, facies et characteres plures *O. Cassii*, sed praeter calyces longius et densius albo-pilosos differt corollæ lobis longioribus acutis, antheris exsertis.

33. O. procerum, viride setis sparsis a tuberculo glabro ortis ad folia adpressiusculis obsitum inter eas glabriusculum, caulis elatis, foliis lanceolatis inferioribus acutiusculis basi attenuatis floralibus a basi semiamplexicauli attenuato-acuminatis, racemis paniculatis post anthesin elongatis, calycis fructiferi modice accreti laciniis lanceolatis, corollæ albidae pubescenti calyce duplo longiore lobis ovatis obtusis apice subreflexis, nectario hirto, antheris inclusis filamento duplo longioribus, nuculis junioribus ovalis laevibus ♀.

Hab. in Armeniâ loco non indicato. (Cult. Hort. Genev. e seminibus Kotschyani).

Bipedale, folia inferiora 3-4-pollicaria 7-9 lineas lata, corolla 8-9 lineas longa. Facies et characteres *O. Cassii* et *oreodoxi* a quibus corollæ nectario hirto nec glabro differt.

34. O. gracile (Trautv. Stirp. Nov. Descr. p. 10) basi e radice indurata verticali multicaule pallide virens totum setis rigidis patentibus e tuberculo magno glabro ortis seberrimum inter setas glabrum, caulis tenuibus rigidis albidis aliis sterilibus brevioribus aliis floriferis parce ramosis, racemis axillaribus et terminalibus paucifloris laxiusculis brevibus, foliis radicalibus..... caulinis lanceolatis utrinque attenuatis acutiusculis, floralibus anguste linearibus, calycis patule setoso-hirsuti laciniis linearibus, corollæ crebre pubescente ochroleucâ calycem subæquante dentibus ovatis obtusis revolutis, nectario hirsutissimo, antheris inclusis filamento triplo longioribus, nuculis parvis late ovatis breviter acuminatis laevibus nitidis ♀.

Hab. in Aranenâ Rossicâ prope pagum Belaw (Raddel). Vid. in herb. cl. a Trautvetter.

Caules cum racemis 6-10-pollicares, folia caulina inferiora 15-16 lineas longa 4 lata, caulum steriliū minora angustiora, calyx et corolla $5\frac{1}{4}$ lin. longi. Species setis brevibus sed strigosissimis, floribus parvis notabilis.

§ 2. **Heterotricha.**

35. **O. lanceolatum** (Boiss. et Haussk. pl. exs. 1868. — Dec. II, p. 1) basi suffrutescens totum setis adpressissimis tuberculatis cinereum, tuberculis præter nonnullos in paginâ inferiori foliorum superiorum pilosulos glabris, caulis erectis superne parce racemosis, racemis fructiferis parum elongatis densiusculis secundis, foliis radicalibus.... cæteris sessilibus lanceolatis acutis trinerviis, bracteis anguste lanceolato-subulatis, calye hirsuto laciniis lanceolatis acutis fructifero subacereto pedicellato subpatulo tubo hispidissimo laciniis dilatatis, corollâ e sicco ochroleucâ cylindricâ superne subdilatata hirtâ calyce quartâ parte longiore dentibus late triangularibus, nectario hispido, nuculis ovato-triquetris laevissimis acutis nitidis ♀.

Hab. in herbidis montis Pir Omar Gudrun Kardistaniae Persicæ 3-6000' (Haussk!).

Caules fere pedales, folia caulina inferiora 2-2 $\frac{1}{2}$ poll. longa 5-6 lineas lata, calyx florifer 7 lineas corolla 9-10 longa. Facie et calyce fructifero affine *O. Olivieri* sed folia subtus trimervia, nectarium hirsutum, etc.

36. **O. Kilouyense** (Boiss. et Haussk. MSS. — Dec. II, p. 1) basi suffrutescens pluricaule, caulis rigidis albidis sparsim et patule setulosis inferne subaphyllis, foliis setis brevibus patulis tuberculatis asperis tuberculis paginâ superioris glabris inferioris stellatim pilosis lanceolatis acutis basi subattenuatis inferioribus cito exsiccatis, racemis 2-3 terminalibus fructiferis secundis vix elongatis, calycis breviter albo-hirsuti laciniis longis linearibus acutis fructifero pedicellato laciniis valde dilatatis oblongis, corollâ e sicco purpureâ velutinâ cylindricâ superne subattenuatâ calyce quartâ parte longiore dentibus ovato-triangularibus, nectario glabro, antheris inclusis filamento brevissimo, nuculis ovatis acutis obsolete scrobiculato-rugulosis ♀.

Hab. in graminosis montium Sawers, Nur, districtus Kuh Kilouyeh Persicæ australis 41000' (Haussk. pl. exs. sub *O. Kotschyi*!).

Caules tenues rigidi 1-1 $\frac{1}{2}$ -pedales, folia caulina intermedia 1-1 $\frac{1}{2}$ -pollicaria 5-6 lineas lata, calycis 6 lineas longi laciniæ lineam fructiferae 2-3 lineas latæ, corolla 10 lineas longa. Facie et caulis rigidis refert *O. Kotschyi* sed hoc honotrichum est, asperrimum. hujus flores multo minores sunt, calycis fructiferi laciniæ vix dilatatae, etc.

37. **O. chlorotrichum** (Boiss. et Noë Diagn. Ser. II, 3, p. 137) totum setis flavidis patentibus e tuberculo ortis densissime hirsutum inter setas velutinum intense flavidò-virens, tuberculis præter non-

nullos paginae inferioris foliorum superiorum glabris, caule erecto parere ramoso, foliis radicalibus late lanceolatis acutiusculis in petiolum longe attenuatis caulinis diminutis sessilibus, racemis floriferis densis scorpioideo-capitatis, calycis flavo-hirsutissimi laciniis breviter lanceolatis obtusiusculis tubo basi attenuato parum longioribus, corollæ ochroleucæ breviter velutinæ calyce duplo longiore elevata lobis triangularibus reflexis, nectario glabro, antheris subexsertis filamento longioribus, nuculis... ♀.

Hab. in Kurdistaniâ inter Van et Ischri (Noë 162!).

Specimen unicum vidi, planta verosimiliter basi suffruticosa, caulis e rhizomatis duri ramo ad latus rosulae nascens 10-pollicaris, folia rosulae 5 pollices longa pollicem lata, calyx florifer 5 lineas, corolla 8-9 longa. Species indumento flavo et calyce minus profunde partito insignis.

38. **O. Olivieri** (Boiss. MSS. in Oliv. pl. exs. Dec. II, p. 2) basi suffruticosum totum setis flavidis adpressiuseculis basi tuberculatis dense hirsutum asperum, tuberculis praeter nonnulla paginae inferioris foliorum superiorum substellatum pilosula glabris, caulis elatis, foliis radicalibus oblongis vel oblongo-spathulatis in petiolum longiusculum attenuatis, caulinis sessilibus lanceolatis acutiusculis, racemis 1-2 axillaribus terminalique bifido fructiferis elongatis densiusculis secundis, calycis flavo-hirsutissimi laciniis lanceolatis acutis, fructifero pedicellato erecto acreto laciniis dilatatis, corollæ e sicco ochroleucæ glabrescente calyce tertia parte longiore subelavata lobis triangularibus reflexis, nectario glabro, antheris inclusis filamento brevissimo, nuculis immaturis majusculis ovatis acutis ♀.

Hab. in Babyloniâ inter Bagdad et Kermanchah (Oliv!), in montibus Avroman et Schahu (Haussk!).

Pedale et procerius, foliorum radicalium lamina 2-2 1/2-pollicaris, calyx florifer 6-7 lin. longus, corolla 9-10 lineas longa. Indumento flavo insigne, affine *O. chlorotricho* sed praeter indumentum adpressius et folia radicalia breviter nec longe attenuata flores maiores, calyx profunde partitus.

39. **O. hebebulbum** (D.C. Prodr. X, p. 60) setis crebris patentibus e tuberculis praeter eos foliorum inferiorum interdum glabratos stellatum pilosis ortis asperum canescens, caulis simplicibus, racemis 2-3 brevissimis in capitulum terminale densum congestis, foliis radicalibus... caulinis oblongo-lanceolatis acutis sursum sensim diminutis inferioribus basi attenuatis, calycis albo-villosissimi laciniis linearibus, corollæ albida pubescente calyce tertia parte longiore dentibus triangularibus reflexis, nectario glabro, antheris apice summo exsertis filamento brevissimo, nuculis... ♀.

Hab. in Mesopotamiâ (Auch. 2314! et 2719 ex Persiâ sed ut videtur per errorem indicatum).

Caules pedales, folia inferiora bipollicaria 5 lineas lata, corolla 8 lineas longa. Species capitulo albo-hispidissimo insignis transitum ab *Heterotricha* ad *Asterotricha* ut jam Candolleus observaverat præbens.

§ 3. *Asterotricha.*

* Perennia.

40. **O. Isauricum** (Boiss. et Heldr. Diagn. Ser. I, 11, p. 101) totum setis adpressissimis e tuberculo stellatim piloso ortis cinereum subsericatum basi suffruticosum, caulis ascendentibus basi saepe ramos breves steriles edentibus superne paniculato-subcorymbosis, foliis inferioribus ovato- et oblongo-spathulatis obtusis in petiolam attenuatis superioribus semiamplexicaulibus sessilibus, bracteis anguste linearibus, racemis fructiferis demum subelongatis. calyce 5 vel laciniarum coalitione 4-3-partito laciniis linearibus acutis fructifero acreto basi angulato-quinquegibbo laciniis subdilatatis. corollæ oclu-
lencæ adpresso velutinæ clavata calyce sublongiore dentibus late ovatis subreflexis. nectario glabro, antheris apice vix exsertis subsessilibus, nuculis levissimis oblongis subrostrato-acutatis ♀.

Hab. in rupestribus apriis Isauriæ inter Karaman et Ermenek (Heldr!), Paphlagoniæ et Galatiæ ad Mersiwan et Tokat (Wied!), Cappadociaæ prope Berekely et circa Cæsaream (Bal. exs. sub *O. sericeo*!).

Valde affine *O. sericeo* a quo artificialiter tuberculis stellatis removetur et cum eo facile confusum; filamenta subnulla, nuculæ magis acutatæ.

41. **O. Bourgæi** (Boiss. Dec. II, p. 2) totum setis mollibus adpressinsculis e tuberculo stellatim piloso ortis obsitum inter. ^{te} tuber-
culos adpresso pubescens tomentosum basi e rhizomate brevi indurato cespitoso-pluricaule, caulis crebre foliosis in racemum bifidum circinnatum densum abeuntibus, foliis lanceolatis radicalibus in petiolam brevem attenuatis obtusiusculis cæteris strictis diuinutis sessilibus acutis, bracteis triangulari-lanceolatis, floribus pedicellatis, calycis albo-hirsutissimi laciniis lanceolatis acutiusculis, corollæ pubescentis calyce quartâ parte longioris dentibus triangularibus. nectario glabro, antheris inclusis, nuculis... ♀. *O. circinnatum* Bourg. Sched. 1863.

Hab. in regione alpinâ montis Fecelem prope Gumuschkhané Armeniae Turcicae.

Pedale, folia fere *Hieracii cymosi* inferiora 2-2½-pollicaria 4-5 lineas lata, racemus fructifer subelongatus, pedicelli hispidissimi calycem tandem subæquautes, calyx 5 lin. longus ut videtur fructifer parum auctus, corolla 6 lin. longa.

42. **O. ovalifolium** (Ky. et Boiss. pl. exs. 1859) totum setis tenuibus adpressissimis e tuberculis minimis non contiguis stellatim pilosis asperulum griseum basi breviter suffrutescens ramis floriferis numerosis crebre foliosis, foliis basi attenuatis radicalibus subpetiolatis ovato-spathulatis obtusissimis superioribus sensim diminutis oblongis et ellipticis summis acutiusculis, racemis 2-3 sub anthesi

brevisimis scorpioideis in capitulum congestis, bracteis lanceolatis acutis, calycis longe et adpressiuscule albo-plumosi laciniis anguste lanceolatis acutissimis, corollæ puberulæ albidae cylindrica superne parum dilatatae calyce quartâ parte longioris dentibus triangularibus, nectario glabro, antheris inclusis filamentis brevibus latis, nuculis... ♀.

Hab. in valle versus fodinas vallis Gorumse in ditione Kassan Oglu Ciliciæ orientalis 4500' (Ky. 1351).

Pedale, folia inferiora 1-1 1/2 pollicem longa 7-11 lata, calyx florifer 5 1/2, corolla 7 1/2 lin. longa. Insigne tuberculis minimis dissitis nec contiguis, indumento brevi asperulo, foliis radicalibus latis brevibus.

43. O. molle (D.C. Prodr. X, p. 61) pluricaule cespitosum totum pilis stellatis contignis dense pannosum canum setâ terminali vix longiore et validiore adpressâ, foliis late linearibus obtusis radicalibus creberrimis in petiolum longe attenuatis, caulinis sessilibus, racemis 1-3 terminalibus fructiferis elongatis rigidis subaxialis, calyce tomentoso-cano laciniis longis lanceolatis acutis fructifero subdilatato pedicello brevi incrassato stricto suffulto, corollâ ochroleucâ velutinâ clavata calyce tertiatâ parte longiore lobis ovatis reflexis, nectario glabro, antheris inclusis filamento duplo longioribus, nuculis ovato-triquetris acutis laevibus maculatis ♀. *O. ciliatum* D.C. loc. cit. (eadem prorsus planta fructifera!). — *Alkanna tomentosa* Griseb. Spicil. II, p. 88 in notâl (ex specimine submonstroso corollis orbato descripta).

Hab. in cretaceis Cappadociae ad Euphratem (Auch. 3202!), prope Orfaï Biredjick! (Haussk!), ad Gebel Abiad deserti Syriaci (Bl.).

Tandem pedale racemis fructiferis crassis strictis 5-6-pollices longis, folia radicalia 4 pollices longa superne 2-2 1/2 lineas lata, calyx florifer 5 fructifer 7 lineas longus laciniis basi latis, corolla 10-11 lineas longa. Species indumento pannoso denso distinctissima nullibi ciliata.

44. O. dasytrichum (Boiss. Diagn. Ser. I, 7, p. 33) fruticosum totum setis brevibus adpressissimis e tuberculis contiguis stellatum lepidotis ortis sericeo-canum. ramis brevibus aliis sterilibus aliis floriferis, foliis ramorum sterilium majoribus ovatis brevissime petiolatis, ramorum fertilium elliptico-oblongis, floribus 3-5 subsessilibus capitato-congestis magnis, calycis adpressiuscule et longe hirsuti laciniis lanceolatis acutis, corollæ tomentellæ e sicco violaceæ calyce subduplo longioris dentibus longiusculæ triangularibus, nectario glabro, antheris apice subhexsertis filamento longioribus, nuculis (junioribus) glabris 5.

Hab. in Persiae australis fauibus prope Kaserun (Ky. 146!), ad Kotel Kumaredj et prope Bebehau 2900' (Haussk!). Fl. Apr.

Rami erecti lignosi parce ramulosi 6-10-pollicares, folia sterilium 1-1 1/2 pollicem longa 9-11 lineas lata, fertilium 4-6 lineas longa 1 1/2-2 lineas lata. Corolla pollicem longa. Species elegans indumento et capitulis paucifloris insignis.

45. **O. latifolium** (Boiss. et Haussk. in Pl. Pers. Sched. 1868. — Dec. 2, p. 3) suffruticosum totum setis longis adpressissimis e tuberculis stellatim pilosis ortis tenuiter tomentosum griseum, foliis ramorum steriliū majoribus obovatis in petiolum brevem attenuatis, ramorum fertiliū parvis oblongo-lanceolatis acutiusculis, racemo terminali bifido dense capitato fructifero non elongato, calycis inferne albo-pilosī fructiferi acreti laciniis virentibus late lanceolatis acutis, corollā lutescenti patule puberulā subelavatā calyce tertiatā parte longiore dentibus triangularibus, nectario glabro, antheris apice exsertis, nuculis magnis oblongo-triquetrīs acutis levibus nitidis ♀.

Hab. in herbidis montium Avroman et Schahu Kurdistaniae Persicæ (Haussk !)

Ramorum steriliū folia 1 1/2-2-pollices longa 1-1 1/2, lata, fertiliū minorū et angustiora, calyces fructiferi pollicem longi laciniis basi demum 2-2 1/2, lineas latis, corolla 10 lin. longa, nuculae fere tres lineas longae. Veluti *O. dasytrichum* heterophyllum et *cephalanthum*, sed indumentum opacum. rami (semipedales) fere herbacei nec fruticosi et calycis laciniæ multo latiores.

46. **O. albo-roseum** (Fisch. et Mey. Ind. V, Petrop. p. 36) basi suffrutescens totum tomento densissimo stellato et setulis tenuibus adpressissimis dense pannosum canum minime strigosum, rhizomatis cespitosi ramis procumbentibus foliorum vestigiis vestitis, foliis oblongis obtusis in petiolum brevem attenuatis, caulinis sessilibus abbreviatis late linearibus, ramis floriferis pumilis in racemum simplicem rarius bifidum etiam fructiferum congesto - capitatum abeuntibus, floribus magnis subsessilibus, calycis tomentosi vel hispidae laciniis late linearī lanceolatis obtusiusculis fructiferis dilatatis. corollā pruinoso-velutinā albā dein sanguineā tandem cœruleā calyce tertiatā parte longiore dentibus triangularibus reflexis, nectario glabro, nuculis magnis oblongis acutiusculis nitidis levibus vel obsolete rugosis ♀. *O. congestum* D.C. Prodr. X. p. 64.

Hab. in fissuris rupium calcarearum, Galatia ad Amasia (Wied!), Cilicia Trachea ad Anamur (Peron!). Taurus Cilicicus (Auch. 2399!, Bal!), montes Nur et Tschoch Dagh Ciliciæ orientalis (Ky. 96!), montes Ssodagh et Akdaghi Syriæ borealis, Terek, Mardin et mons Sindjar Mesopotamiae (Haussk!).

Species pulcherrima habitu *O. frutescens* referens sed stellatim pannosa, calycis fructiferi laciniis latissimis obtusis insignis. Corolla 11 lineas longa. Calycis pili grisei vel flavi.

47. **O. cœrulescens** (Boiss. Diagn. Ser. I, 11, p. 110) basi valde suffruticosum multicaule totum tomento stellato cano setisque creberrimis brevibus patulis asperum griseum, caulis floriferis pumilis in 2-3 racemos in capitulum congestos fructiferos non elongatos abeuntibus, foliis obtusis inferioribus oblongo-spathulatis in petiolum breve attenuatis, caulinis diminutis basi attenuatā sessilibus, calycis longe albo-hispidae laciniis longe et anguste linearibus obtusis, corollā glabrā albida tandem cœrulecenti anguste cylindricā vix ampliatā calyce quartā parte longiore dentibus triangularibus

revolutis, nectario glabro, antheris apice subexsertis, nuculis medio-cribus oblongo-triquetris acutis laevissimis nitidis ♀.

Hab. in Antilibano inter Rascheya et Damascum (Boiss!), in latere orientali montis Gebel es Scheick (Gailli!), in monte Schergi supra Bludan 5000' (Ky. 7 et 1221).

Caules floriferi 5-10-pollicares, folia majora pollicem longa 3-4 lineas superne lata, corolla 7 lineas tantum longa. Ab *O. albo-roseo* et *Rascheyano* corollâ multo breviori tubulosâ nec clavatâ indumentique naturâ abunde distinctum.

48. *O. mutabile* (Boiss. et Haussk. Sched. 1866. — Dec. 2, p. 3) basi suffrutescens ramosum totum tomento tenui stellato et setis adpressiusculis tomentellum griseum, ramis floriferis numerosis erectis pumilis, foliis inferioribus oblongo spathulatis obtusis breviter attenuato-spathulatis, caulinis sessilibus late linearibus acutisusculis, racemis 2-3 terminalibus congestim capitatis fructiferis vix elongatis, calycis hirsuti laciniis longissime lineari-lanceolatis acutissimis fructiferis vix auctis, corollâ adpresso hirtâ subclavatâ flavidâ dein roseâ siccâ cœruleâ calyce tertiatâ parte longiore dentibus obtuse triangularibus subreflexis, nectario glabro, nuculis majusculis triquetris abrupte acutatis nitidis obscure ruguloso-tuberculatis ♀.

Hab. in regione superiore montis Berydagh Cataoniæ 6000'-9000' (Haussk!). Fl. Aug.

Caules floriferi semipedales, facie et florum magnitudine *O. albo-roseo* affine et inter hoc et *O. Rascheyanum* intermedium; prius differt indumento crasso panuoso, calycis laciniis latis obtusis; posterius indumento patenti strigoso, corollis pruinoso-glabrescentibus, nuculis laevissimis distinguitur.

49. *O. Rascheyanum* (Boiss. Diagn. Ser. I, 11, p. 410) basi suffrutescens ramosum totum setis patulis e tuberculo stellatum piloso ortis strigulosum pallide virens, ramis aliis decumbentibus sterilibus foliosis aliis paulo longioribus in racemum simplicem vel bifidum capitatum fructiferum non elongatum abeuntibus, foliis ramorum sterilium anguste oblongo-spathulatis in petiolum longe attenuatis, caulinis elliptico-linearibus sessilibus, bracteis anguste lanceolatis acutis, calycis adpressiuscule et longe hirsuti laciniis longis lanceolatis acutissimis, corollâ clavatâ breviter pruinoso-velutinâ calyce tertiatâ parte longiore dentibus triangularibus revolutis, nectario glabro, antheris apice subexsertis filamento longioribus, nuculis oblongis attenuato-acutis laevissimis nitidis ♀.

Hab. in Antilibano inter Rascheya et Damascum (Boiss!), in monte Pir Omar Gudrun Kurdistaniæ Persicæ (Haussk!). Fl. Mai.

Caules floriferi vix semipedales, folia inferiora et ramulorum cum petiolo 4-4 1/2 poll. longa apice 3 lineas lata. Calyx florifer 9 lineas, corolla 12-13 longa. Species pulchra ab *O. stellulati* formis rhizomate magis suffruticoso, floribus capitatis, corollâ versicolore, calycis laciniis elongatis, etc.; ab *O. Armeno* et *Roussæi* corollâ majore clavato-inflata versicolori, nuculis majoribus elongatis, etc., distincta. Specimina ex Antilibano tantum florifera mihi nota cum Kurdicis fructiferis et floriferis optime congruere videntur.

50. **O. mite** (Boiss. et Heldr. Diagn. Ser. I, 11, p. 111) basi suffrutescens totum setis tenuibus brevibus adpressissimis e tuberculis contiguis stellatim pilosis ortis pannoso-canum, ramis aliis sterilibus brevibus dense foliosis aliis in racemum simplicem brevem tandem laxiusculum abeuntibus, foliis margine revolutis ramorum sterilius anguste oblongo-linearibus basi attenuatis, caulum linearibus, bracteis lanceolatis calycem æquantibus, calyce patule flavidohirsuto laciniis linearibus acutis fructifero parum acreto, corollâ pubescenti vel glabriuseculâ ochroleucâ calyce tertiatâ parte longiore sursum subdilatata lobis ovatis acutis erectis, nectario glabro, antheris filamento triplo longioribus appendice longe exsertâ, nuculis ovatis acutiusculis lăvibus nitidis ♀.

Hab. in pinetis apricis ad radices montis Taktali Lyciae supra portum Tcherali (Heldr!), in Cypri declivibus inter Prodromo et Fini (Ky. 705!).

Rami steriles bipollicares, floriferi cum racemo fructifero 7-9-pollicares, folia majora 9-10 lineas longa superne lata, calyx 5 lineas, corolla 6 longa. Indumento niti, corolla lobis erectis et antherarum appendice exsertâ ab *O. stellulato* et affinibus longe alienum.

51. **O. Armenum** (D.C. Prodr. X, p. 59) multicaule totum setis tuberculo stellatim piloso insidentibus patulis ad folia adpressiusculis canescens, caulinis pumilis ascendentibus simplicibus, foliis parvis breviter linearibus margine revolutis inferioribus et rosularum sterilius obtusis floralibus angustis acutis, racemis 2-3 terminalibus brevissimis in capitulum conglobatis fructiferis vix elongatis, calycis plumosi fructiferi parum aucti laciniis longis anguste linearibus acuminatis, corollâ subvelutinâ pallide ochroleucâ cylindricâ calycem vix vel tertiatâ parte excedente dentibus ovatis reflexis, nectario glabro, nuculis minimis triquetris apice abrupte acutatis nitidis lăvibus ♀. *O. versicolor* Schott in Ky. Sched

Hab. in montibus Armeniae (Auch. 2358!). Cappadociae ad Euphratem (Auch. 2305!) et in monte Alidagh (Bal!). Cataoniae monte Berytdagh 7000' (Haussk!). Anatoliae prope Hamanli, Safranbol, Tokat (Wied!), Pisidiæ in montibus tophaceis ad Buldur (Heldr!), Lyciae supra Elmalu (Pest!).

Semipedale, folia sex lineas longa 1 1/2 lata, calyx 6-7 lineas longus. Affine formis pumilis *O. stellulati* var. *pallidi* sed differre videtur racemis brevibus in capitulum confertis, corollâ cylindricâ superne vix dilatata.

52. **O. Roussæi** (D.C. Prodr. X, p. 49) cespitoso-multicaule totum setis tuberculo stellatim piloso insidentibus patulis vel adpressiusculis canum, caulinis pumilis ascendentibus, foliis linearibus margine revolutis obtusis, racemis 4-3 terminalibus brevibus in capitulum confertis fructiferis parum elongatis densis, bracteis calyce brevioribus, calycis cano-hirsuti fructiferi subaccreti laciniis linearibus brevibus, corollâ velutinâ ochroleucâ calyce duplo longiore dentibus brevibus reflexis, nectario glabro, nuculis parvis triquetris apice abrupte acutatis nitidis lăvibus ♀.

Hab. in Syriâ circa Aleppo (nec ad Bagdad) (Rousseau!), inter Aleppo et Orfa (Ky. 3!), in cretaceis inter Nisib et Aintab et ad Mardin (Haussk!), in Libano supra Cedros, forma minus cana setis patulis, caulinibus cæsiis (Boiss!), in monte Sannin et virtus Ainete (Ehrenb. 162!).

Semipedale vel humilius, folia 1-4 $\frac{1}{2}$ -pollicaria. Facie et characteribus affine *O. Armeno* a quo tamen differt foliis longioribus et præsertim calyce 4-5 lineas tantum longo laciniis multo minus acuminatis, corollâ magis exsertâ.

53. *O. stellulatum* (W. K. Pl. rar. Hung, II, p. 189, tab. 173) basi suffrutescens multicaule totum setis tuberculo stellatum piloso insidentibus adpressis vel patulis hispidâ - sebrum vel asperum, caulinibus erectis vel ascendentibus foliosis racemo bifido rarius simplici terminali vel parce corymboso-paniculatis ramis fructiferis elongatis, foliis strictis linearí-oblongis planis vel margine revolutis radicalibus et inferioribus subpathulatis obtusis, superioribus lanceolatis aculis basi interdum semiamplexicaulibus, calycis fructiferi parum acreti laciniis anguste lanceolatis acutis, corollâ albida flavâ vel citrinâ clavatâ calyce 2-3-plo longiore brevissime velutinâ vel glabrescente dentibus late et breviter ovato-triangularibus reflexis, necario glabro, antheris inclusis, nuculis triquetris abrupte acutatis nitidis levibus ♀.

Species statuâ indumento patenti vel adpresso, corollæ colore et longitudine, antheris filamento parum vel multum longioribus, etc., mire varians. Formæ altera in alteram transeunt; evidentiores sunt sequentes.

z. genuimum. — Planta viridis, setae foliorum adpresso-patulae, calycis sæpius patentes. Corolla flava. — *O. Helveticum* Boiss. Diagn. Ser. I, 11, p. 111. — *O. heterophyllum* Griseb. Spie. II, p. 80.

Hab. in siccis regionis inferioris et montanae in parte magis septentrionali ditionis, Macedonia et Thracia (Griseb! Orph!), Byzantium (Wied!). Anatolia borealis tota (Wied!), Pontus Lazicus 6000' (Bal!), Armenia prope Baibut (Bourg. exs. sub *O. Armeno*), Caucasus omnis 2000'-6000' (Hoh! Rupr! etc.).

β. pallidum. — Pallide virens totum patule plus minus strigosum, folia sæpius abbreviata. Corolla albida. Forma a præcedenti sæpe ægre distinguenda. — *O. pallidum* Boiss. Diagn. Ser. I, 11, p. 112. — *O. Aucherianum* D.C. Prodr. X, p. 60. — *O. rigidum* Ledeb. Ic. tab. 238 (forma autumnalis elongata macra). — *O. scaberrimum* Boiss. et Heldr. MSS. — *O. Trapezuntinum* Huet pl. exs.

Hab. in siccis Peloponnesi (Orph!), Phocidis et Beotiae (Sprun! Orph! Guicci!), Macedonia (Orph!), Thracie (Jank!), insulæ Chio (Orph!), Byzantii (Cast!), totius Anatoliae occidentalis a Bithyniâ et Lydiâ (Boiss!), etc., ad Ciliciam (Bal. 101! sub *O. Armeno*) et interioris in Lycœnia (Heldr!), Cappadociâ (Bal!), Tauriâ (Stev!), Caucaso centrali (R.-hm!).

γ. angustifolium. — Planta cinerascens setis ad folia adpressissimis, inflorescentiæ adpressis vel patulis. Folia angusta margine revoluta sæpe elongata. Corolla citrina. — *O. angustifolium* Lehm. Asp. p. 363, tab. 14. — *O. Tauricum* Pall. Nov. Act. p. 306. — *O. paradoxum* Janka Brev. p. 7. — *O. aurantiacum* Janka exs. — *O. Orphaideum* et *O. Laconicum* Boiss. MSS. — *O. scorpioides* Huet MSS.

Hab. in montibus Græciae, Taygeto et ad Megaspileon (Heldr!), Kyllene Malevo, Citheron (Orph!), Macedoniae in monte Athone (Griseb!), Bithynia (Griseb!), Anatoliæ boreali (Wied!), Cariæ in Cadmo (Boiss!), Ciliciæ (Bal!), Armeniæ Tureicæ (Huet!), Tauriæ (Turcz! Rehm!), Transcaucasiæ (Szov!).

O. brevifolium. — Indumentum adpressum præcedentis sed folia brevia Forma montana sæpius humilior. — *O. brevifolium* D c. MSS.

Hab. in montibus Cariæ (Boiss! Pinard!), Lyciae (Bourg!), Pisidiæ (Heldr!), Tauri Cilicio (Bal!), Syria in Cassio (Boiss!), Libano (Bl!).

O. erectum. — Humile, folia angusta margine revoluta adpresso cana. racemus brevis simplex vel bifidus pilis adpressis et aliis sub-patulis canus et plus minus plumosus. Forma in præcedentem sensim transiens. — *O. erectum* Fl. Græc. II, p. 62, tab. 173 et *O. montanum* Prodr. I, p. 121.

Hab. in Cretæ saxosis maritimis ad Hagio Rumeli et in monte Idæ 5000' (Heldr! Sibth.), Attica in Hymetto (Sprun! Boiss! Heldr! Orph. Fl. Gr. exs. 601).

Ar. Geogr. Sabaudia, Helvetia australis, Germania australis, Italia, Dalmatia, regio Danubialis.

* * Biennia.

54. *O. Sprunerri* (Boiss. Diagn. Ser. I, 11, p. 109) totum setis brevibus tenuibus e tuberculo minuto stellatim piloso ortis adpressissime et tenuiter cinereum, caulis e radice verticali simplici 1-3 erectis elatis apice parce et stricte racemosis, foliis radicalibus longe et late linear-lanceolatis obtusiusculis in petiolam longe attenuatis, superioribus semiamplexicaulibus oblongo-lanceolatis acutis, racemis fructiferis elongatis laxis rigidis, bracteis calyce brevioribus, calyce adpressissime setoso laciinis breviter lanceolatis fructifero acreto pedicello incrassato strictissimo insidente obconico, corollâ albidâ adpresso pilosâ superne ampliatâ calyce triplo longiore dentibus triangulari-ovatis, nectario glabro, antheris inclusis, nuculis magnis oblongis obtusiusculis obscure granulato-tuberculatis nitidis ②.

Hab. in Bœotia ad radices montis Citheron Bœotiae (Sprun! Orph. Fl. Gr. exs. 744!), in regione inferiori montis Kyllenes Peloponnesi 2500' (Orph!). • Onosmatum rex, ut reges optimi rarissimum. ▶ Cl. Orph. in litt.

Tandem sesquipedale, folia radicalia 6-7-pollicaria 5-6 lineas lata, corolla pollicem longa. Ab *O. stellulati* formis indumento multo tenuiore et adpresso et præsertim radicis indole longe remotum.

55. *O. giganteum* (Lam. Ill. 1840) totum setis adpressis e tuberculo stellatim piloso ortis griseum, caule e radice longâ fusiformi unico crasso elato ample ramoso-paniculato. foliis lanceolatis radicalibus longissimis obtusiusculis longe petiolato-attenuatis, caulinis crebris strictis valde abbreviatis et angustatis acutis, paniculâ albo-vel flavidâ hispidissimâ, racemis fructiferis elongatis densis secundis, bracteis minutis triangularibus acuminatis, calycis pedicello tenui erecto-patulo fructifero non incrassato suffulti fructiferi parum accreti hispidissimi laciinis anguste linearibus, corollâ minute velutinâ pal-

lidle ochroleucâ calyce tertiatâ parte longiore superne subampliatâ lobis obtusissimis, nectario glabro, antheris inclusis filamento brevi, nuculis parvis ovato-triquetris acutis lœvibus nitidis ②. Ic. Tausch Hort. Canal. fasc. I. — *O. astroites* Stev. Bull. Mosq. 1851, p. 585 (in notâ).

Hab. in collibus et agris derelictis Palestinæ in planitie Esdraelon (Boiss!), ad Hierosolymam et Ramleh (Post!), Ouadi el Melek (Gaill!), in Syria (Auch. 2298!), monte Amano (Kyt!), in monte Ssoffdagh 5000' (Haussk!), Cilicia (Ky. 2471).

β. hispidum. — Setæ patentes.

Hab. ad Bouloukli Ciliciæ (Bal!), Cypro (Ky. 680!), valle Ermenek Ciliciæ Tracheæ (Heldr!).

Planta sœpe bipedalis multiflora, folia inferiora $\frac{1}{2}$ -1 pedem longa 6-10 lin. lata, corollæ plantæ proceræ minutaæ 7 lineas tantum longæ.

56. ***O. auriculatum*** (Auch. pl. exs. p.c. Prodr. X, p. 61) totum setis ad folia adpressis tenuibus e tuberculo stellatim piloso ortis pallide virens vel griseum, caule e radice fusiformi crasso elato cærebre et stricte folioso superne ample paniculato, foliis inferioribus oblongo-lanceolatis obtusiusculis in petiolum attenuatis intermediis basi dilatatâ auriculatis, bracteis subcordato-triangularibus parvis, racemis fructiferis elongatis laxiusculis secundis, pedicellis hispidissimis tenuibus tandem patulis calyce vix brevioribus, calyce hispido laciniis linearispathulatis obtusis fructiferis subdilatatis, corollâ paleide ochroleucâ minute velutinâ calyce tertiatâ parte longiore lobis obtusis, nectario glabro, antheris inclusis filamento æquilongis, nuculis ovato-triquetris oblique acutis lœvissimis nitidis ②.

Hab. in Mesopotamiæ humidis inter Surug et Orfa (Haussk!), ad Ninivam (Auch. 2313!), ad Diarbekir (Ky. 249!).

Planta 1- $1\frac{1}{2}$ -pedalis, corolla 7-8 lineas longa. Facies omnino *O. gigantei* a q. 10 differt foliis caulinis auriculato-dilatatis, lacinarum calycis formâ, nucula subobliquâ.

Species incerta.

O. Orientale (L. Amæn. 4, p. 267 non Pall. nee Lehm.) caulinis hispidis, foliis lanceolatis hispidis, racemo bifido, bracteis anguste lanceolatis, calycis laciniis a basi cordatâ linearibus, corollæ laciniis subulatis acutis patulis, staminibus corollæ æquilongis, fructibus pendulis.

Hab. in Ægypto (Hasselquist).

MEGACARYON (Boiss. Dec. I, p. 7).

Calyx quinquepartitus laciniis fructiferis longissimis. Corolla subregularis hirsuta tubuloso-infundibuliformis sursum ampliata fauce

nudâ, lobis brevibus ovatis erectis. Stylus longus hispidus apice bifidus. Antheræ breviter oblongæ exappendiculatæ nutantes medio dorso filamentis longis inæqualibus sub corollâ dimidiâ insertis apice incurvis affixæ. Nucula abortu sœpius solitaria magna ovata subbicarinata et apice anceps lœvissima nitida areolâ basilarî planâ ovatâ. — Herba elata pilosa. — Genus *Echio* affine, ab eo corollâ fere regulari et præsertim nuculâ magnâ ovatâ lœvissimâ ancipiti nec ut in omnibus *Echiis* turbinato-triquetra tuberculatâ vel rugoso-scrobiculatâ diversum. Calyx fructifer valde auctus et ut fructus habitum præ se ferens *Onosmatis* antheris longis appendiculatis diversi.

t. M. **Orientale** (L. Sp. Ed. 11, I, p. 200 sub *Echio*) caule crasso elato pyramidatum ramoso ut et rami setis patulis basi tuberculatis rigidalis strigoso, foliis radicalibus et inferioribus amplis late oblongo-lanceolatis utrinque attenuatis acuminatis adpresso et breviter canescenti-velutinis, superioribus sensim abbreviatis setosis, floralibus anguste linearis-lanceolatis calycem subæquantibus, spicis primum compactis subcapitatis dein fructiferis elongatis, calycis laciniis longis anguste linearibus acuminatis corollâ parum brevioribus fructiferis dilatatis strigosissimis dorso uninerviis, corollâ griseo-cæruleâ ♀ vel ♂. M. **Armenum** Boiss. Decad. Or. p. 7. — *Echium Orientale* *verbascifolio flore maximo campanulato* Tourn. Cor. 6. — Voy. 14, p. 248 Icon. — *Onosma megalospermum* Boiss. in Bourg. exs.

Hab. in Armenia Turcica (Tourn!), in arvis prope Macka (Bourg!), in Anatoliâ boreali inter Aradsch et Kastamouni (Wiedem!).

Radix ex Tournef. pedem longa, caulis tripedalis et fructifer procerior, folia inferiora mollia *Cynoglossi*, 1-1½ pedes longa 2-3 pollices lata, calyx florifer 12-13 lineas longas, fructiferi laciniæ sesquilineam latæ, corolla 15-16 lineas longa, nucula piso parum minor.

ECHIUM (L. Gen. 191).

Calyx (in nostris) quinquepartitus. Corolla infundibuliformis vel tubuloso-obconica, fauce ampliatâ nudâ, limbo (in nostris) plus minusve irregulari quinquelobo subbilabiato. Stamina 5 inæqualia filamentis longis, antheræ ovatæ dorso affixæ. Stylus filiformis longus. Nuculae 4 liberae fundo calycis arcâ basilari triangulari planâ affixæ ovatæ vel triquetrae tuberculato-rugosæ. — Herbæ (nostræ) perennes vel monocarpicæ.

* Stylus apicæ indivisus vel capitato-bilobus.

t. E. **rubrum** (Jacq. Fl. Austr. 5. App. p. 27, tab. 3) bienne, setis albis erecto-patulis basi tuberculatis hispidum cinereo-virens, caule erecto simplicissimo modice elato in thyrsus spiciformem spiculis brevibus simplicibus constantem abeunte, foliis crebris strictis

lineari-lanceolatis acutis superne sensim angustatis et abbreviatis, calycis corollâ 3-4-plo brevioris laciniis linearis lanceolatis albo-setosis, corollæ adpresso hirtæ purpureæ exsiccatione violaceæ tubo angusto, staminibus exsertis, stylo indiviso ②. *E. acutifolium* Lehm. № 321, tab. 2. — *E. clavatum* Willd. herb. Lehm. Asp. p. 440 (forma macra).

Hab. in pratis siccis et in lapidosis regionis montanæ et subalpinæ, Tauria (Stev!), Caucasus omnis 3000'-6000' (C.A. Mey., Rupr!, etc.) et in montanis Georgiae Caucasicæ (Hohen!), Armeniae Rossicæ (Buhse!, Szov!).

Planta 1½-1 pedalis rarius bipedalis, spica semi-pedalis vel brevior, corolla angusta sex lineas longa.

Ar Geogr. Austria, Moravia, regio Danubialis, Rossia media et australis.

2. *E. linearifolium* (C. Koch Linn. XXIII, p. 629) bienne, setis homomorphis e tuberculo ortis tenuibus caulis patulis foliorum adpressiusculis cinerascens non strigosum, caule erecto superne ramoso, foliis uninerviis radicalibus longe lanceolatis in petiolam longe attenuatis, caulinis sessilibus elliptico-triangularibus acutis, floralibus linearibus calyce longioribus, spiculis axillaribus simplicibus densifloris racemum thyrsoidem brevem formantibus, calycis laciniis linearibus albo-setosis, corollâ calyce triplo longiore tubulosâ adpresso hirtâ, staminibus exsertis, stylo longo exerto apice capitato-bilobo ②. *E. papillosum* C. Koch Linn. 17, p. 307 an Lehm? — *E. glomeratum* Ledeb. Fl. Ross. III, p. 106 non Poir.

Hab. in districtu Daratchitchak Armeniae Rossicæ (C. Koch!). Specimen unicum herb. Acad. Petr. a cl. Ruprecht ad Re'dan inter Wladikaukas et Dolta Caucasicæ ntralis lectum ejusdem speciei forma spiculis laxifloris videtur.

Specimen imperfectum in herb. cl. Koch vidi, folia radicalia cum petiolo 7-10 poll. longa 10 lineas lata, spiculae 1-1½ poll. longæ densifloræ, corolla angusta 6-7 lineas longa. Affine *E. Italico* et *E. glomerato* Poiret sed non strigosum et ab eis præterea stylo capitato-bilobo nec longe bifido distinctum, nuculæ ignota.

* * Stylus apice longe bifidus.

+ Stamina exserta.

3. *E. Italicum* (L. Sp. 139) bienne, caule erecto elato setosohispidissimo in paniculam amplam pyramidatam vel ob ramos florales abbreviatos in racemum spiciformem abeunti, foliis setis basi tuberculatis adpressiusculis asperis acutis radicalibus lanceolatis in petiolam attenuatis floralibus anguste linearibus, spiculis bifidis, calycis albo-setosissimi laciniis lanceolatis, corollâ albido-carneâ extus hirtâ calyce duplo longiore angustâ, staminibus exsertis, stylo bifido ②. Ic. Engl. Bot. tab. 2081. — *E. altissimum* Jacq. Austr. V, tab. 16. — *E. Pyrenaicum* Desf. Atl. I, p. 164. — *E. pyramidatum* D.C. Prodr. X, p. 23. — *E. aspernum* Lam. III, № 1854 — *E. aspernum* M.B. Taur. Cauc. I, p. 135.

Hab. in aridis et cultis derelictis, Zacynthus, (Marg!), Græcia (herb. Fauché!).

Macedonia et Thracia 0-1400' (Griseb!), Cyclades (Urv., Sart!), Anatolia borealis! et australis!, Syria hor. et Cataonia (Haussk!), Syria littoralis (Gaill.), Mesopotamia (Ky. 437!), Assyria (Hausk!), Tauria (M.B.), Caucasus et Transcaucasia (Buhse!), Turkestania ad Samarkand (Lehm!), Persia prov. Aderbidjan (Szov!) et Khorassan (Bge!).

Corolla pro plantæ magnitudine parva 6-7 lineas longa.

Ar. Geogr. Europa australis a Gallia occidentali et australi, Helvetia et Germania australibus ad regionem Danubialem et Rossiam medium, Africa borealis.

4. **E. glomeratum** (Poir. Dict. VIII, p. 670) bienne, cinereum, caule elato teretiusculo simplici setis longis palulis basi tuberculatis pubē brevissimā intermixtis hispido in spicam longissimum tenuem basi interruptam abeunti, foliis adpresso setosis acutis radicalibus lanceolatis floralibus anguste linearis-lanceolatis basi non dilatatis spiculas sessiles breves fasciculiformes superantibus, calycis æqualiter albo-strigosissimi laciniis lanceolatis, corollæ hirtæ albido-carneæ tubo calyce vix longiore limbo intus glabro, staminibus exsertis, stylo bifido ②. *E. elongatum* Poir. Dict. 669! — *E. columnæ* Ehrebr. MSS!

Hab. in pascuis aridis regionis inferioris, Cilicia ad Mersina (Ky. 351!), Cataonia ad Marasch (Hausk!), Syria (Labill!), ad Aleppo (P. Jean!), ad Berythum (Ehrebr!), Sidonem (Bl!), Galilea in valle Jordani (Boiss!).

Planta 3-4-pedalis, spica 1-3-pedalis, corolla minuta 4-5 lineas longa. Species ab *E. Italico* inflorescentia distinctissima, affinis *E. pomponio* Boiss. = *E. Boissieri* Stend. Hispanico et boreali-Africano foliis floralibus basi subcordatis, spiculis elongatis, bracteis et calycis laciniis pectinato-ciliatis, corolla limbo intus hirto specifice distincto.

5. **E. vulgare** (L. Sp. 200) bienne, viride setis homomorphis adpresso e tuberculo ortis obsitum, caule unico vel a collo pluribus strictis in racemum thyrsoidem abeuntibus, foliis uninerviis lanceolatis radicalibus basi attenuatis superioribus acutissimis, florum spiculis brevibus indivisis, calycis adpresso setoso-hirsuti laciniis erectis linearis-lanceolatis obtusiusculis, corollæ cæruleæ extus hirtæ calyce subtriplo longiore. staminibus exsertis. stylo bifido, nucib⁹ dorso subdepressis aentis tenuiter et inæqualiter tuberculatis ②. Ic. Engl. bot. 181. — Rehb. tab. 97.

Hab. in siccis Peloponnesi (ex Sibth.), Thraciae (ex Sibth.), Ponti Lazici prope Of (Bal!). Armeniæ ad Gumuschkhane et Erzerum (Bourg! Calv!) Tauria et Caucasi (M.B., Stev., Rupr!).

Ar. Geogr. Europa borealis et media ab Anglia et Scandinavia ad Rossiam septentrionalem et Hispaniam boreali et centrali per Italiam medium et Dalmatiam ad regionem Danubialem, Sibiria Uralensis.

6. **E. pustulatum** (Fl. Græc. II, p. 68, tab. 180) bienne, cinerascens setis dimorphis aliis longioribus e tuberculo valido ortis strigoso-hispidum. caule unico vel pluribus in paniculam pyramidalam laxiusculam abeuntibus, foliis lanceolatis inferioribus obtusius-

culis floralibus anguste linearibus acutissimis, spiculis pedunculatis laxis indivisis, calycis albo-strigosi laciniis anguste linearibus patentibus, corolleæ caeruleæ hirtæ tubo calyce duplo longiore, staminibus exsertis, stylo bifido, nuculis acutis grosse serobiculato-tuberculatis ♂. *E. tuberculatum* Link et Hoffm. Fl. Port. — *E. pictum* Jan.

Hab. in campestribus insulae Zacynthi (Margot).

Affine *E. vulgari* differt setis dimorphis, indumento strigoso, inflorescentia laxiore paniculato-pyramidalata, calycis laciniis patentibus, corolla tubo elongato, etc.

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia et ejus insulae, Dalmatia, Algeria occidentalis!

7. ***E. sericeum*** (Vahl Symb. II, p. 35) perenne, pluricaule indumento minuto adpresso canum prætereaque setis e tuberculo ortis saepius adpressis crebre obsitum, caulis basi interdum suffrutescentibus decumbentibus vel prostratis, foliis linearibus latiusculis vel angustis margine saepius subrevolutis et interdum subundulatis inferioribus in petiolam attenuatis, floralibus triangulari-lanceolatis calycem subæquantibus, spicis paniculatis patule albo-hispidi demum elongatis laxis, calycis laciniis linearibus obtusiusculis, corollæ hirtæ prætereaque setis longioribus obsita punicea sed post excitationem violacea calyce $2\frac{1}{2}$ -plo longiore, staminibus exsertis, stylo bifido, nuculis griseis acutis acute cristulato-tuberculatis ♀. *E. prostratum* Del. Eg. p. 193, tab. 17, fig. I. — *E. rubrum* Forsk. Deser. p. 41. — *E. distachyum* Viv. Lib. ex el. Cosson.

Hab. in arenosis maritimis et desertorum Ægypti ad Alexandriam et Rosettam (Forsk., Del., Auch. 2394!, Kraft, Gaill! etc.), in deserto inter Egyptum et Palestinam (Boiss!), Rhodi in maritimis (Bourg!), Cretæ ad Cydoniam (Heldr!),

β. diffusum. — Caules ut in formâ typicâ diffusi sed setæ totius plantæ erecto-patentes nec adpresso. — *E. diffusum* Fl. Græc. II, p. 69, tab. 182.

Hab. in arenosis maritimis Cretæ (Sibth., Heldr! Raul!), Cypri (Haussk!), insulae Naxos (Sart!), Cos (Urv!).

γ. hispidum. — Caules erecti proceriores saepe 1-2-pedales præter indumentum canum ut et folia patule hispidi. — *E. hispidum* Fl. Græc. p. 68, tab. 181. — *E. Sibthorpii* R. et Sch. IV, p. 26. — *E. elegans* Lehm. Asp. № 339. — Jaub. et Sp. Ill. tab. 417. — *E. setosum* Fl. Pelop. non Vahl. — *E. expansum* Boiss. et Haussk. MSS.

Hab. in arenosis præsertim maritimis et alveis Atticae (Sprun! Boiss! etc.), ad Corinthum et Sicyonem (Boiss!), Argolide (Sprun!), Byzantii (Noe), Chios (Orph!), Rhodus (Bourg!), Cyprus (Ky!), Bithynia et Paphlagonia (Wied!), Pamphylia (Heldr!), Cilicia Trachea (Péron.), Syria littoralis omnis (Ky! Bl!), Palestina ad Gaza (Boiss!), Hierosolymam (Roth!), Hebron (Ky!), ad Giaur Göl prope Marasch Syriae bor. (Haussk! forma ramis magis expansis).

Species pulchra indumento cano, corollis versicoloribus, nuculis acute tuberculatis insignis, caulis prostratis vel erectis brevibus vel proceris, foliis plus minus angustis, setarum directione valde variabilis sed formâ

omnes evidentissime in typum unicum confluunt. Corolla 7-9 lineas longa. *E. maritimum* Willd. in arenosis maritimis Hispaniæ, Galliæ australis, Italiae et insularum, Africæ borealis copiose crescens formis pumilis *E. sericei* facie accedit sed specifice differre videtur staminibus subinclusis nec longe exsertis, corollâ ex auctoribus violacea nec versicolori.

Ar. Geogr. Italia australis ad mare Adriaticum, Africa borealis in Cyrenaicâ.

8. ***E. longifolium*** (Del. Fl. Eg. p. 192, tab. 16, fig. 3) annum, caulis herbaceis erectis pube brevi et setis longioribus patentibus hispidis ramosis, foliis pallide virentibus setis e tuberculo ortis sparsis pube minutâ in paginâ inferiore intermixiâ obsitis. radicalibus lingulatis obtusis basi longe attenuatis caulinis et floralibus valde diminutis. spicis ad ramos terminalibus demum valde elongatis et laxis, calycis crebre albo-setosi laciniis anguste linear-lanceolatis, corollâ pubescenti violacea infundibuliformi calyce subtriplo longiore, staminibus breviter exsertis, stylo hispido longe bifido stigmatibus capitulatis, nuculis griseis tetragono-conicis acute et parce tuberculato-rugosis echinatis ♂.

Hab. ad culta in Ægypto prope Kahiram (Del., Kral! Schweinf!) ad littus Ægyptiacum maris Rubri ad Ouadi Gadireh (Schweinf!), in cultis Sinaiticis Arabiae Petreeæ (Auch. 2401!).

Planta 1-2-pedalis, folia radicalia saepius elongata 4-6-pollicaria, corolla 8-12 lineas longa.

Ar. Geogr. Nubia.

9. ***E. Rauwolfii*** (Del. Fl. Eg. p. 212, tab. 19, fig. 3) annum totum setis rigidis albis e tuberculis ortis obsitum, caulis erectis ramosis ramisque strigoso-subaculeatis. foliis inferioribus in petiolum attenuatis obtusis oblongo-vel linear-lanceolatis floralibus diminutis sessilibus calyces æquantibus vel superantibus, spicis paniculatis fructiferis longissimis rectis laxis, calycis laciniis tuberculato-strigosissimis lanceolatis inæqualibus, corollâ albidâ vel pallide roseâ patule hirtâ campanulatâ calyce duplo longiore, staminibus subexsertis, stylo hirsuto bifido lobis capitatis, nuculis albis acutis nitidis laevibus obsolete tuberculatis ♂. *Lycopsis Dioscoridis* Rauwolf It. tab. 22. — *E. tinctorium* Vivian Eg. Déc. № 24 et herb!

Hab. ad vias, cultura et in arenosis ad Nilum circa Alexandriam, Kahiram et in Ægypto media et superiori (Delt! Ehrenb! Boiss! Kral! Schweinf! etc.), ad Ouadi Hebrae Petreeæ (Schimp!).

Caules 1-2-pedales, corolla 6-7 lineas longa. Species setis strigosissimis, corollâ albidâ, nuculis nitidis laevibus insignis.

Ar. Geogr. Nubia (Ky. exs. 318 sub *E. setosa*!).

10. ***E. plantagineum*** (L. Mant. 202) biennne setis homomorphis adpressis e tuberculo minuto ortis tomentosum, caulis erectis vel ascendentibus laxe paniculatis, foliis radicalibus ovatis vel oblongis in petiolum attenuatis subtus parallele lineato-nervulosis, florali-

bus valde diminutis cordato-lanceolatis, spiculis simplicibus pedunculatis, calycis laciniis adpresso albo-setulosis pube brevissimâ immixtâ, corollæ violaceæ calyce 3-4-plo longioris tubo in limbum amplum superne paucisetosum abrupte ampliato, staminibus tandem breviter exsertis, stylo bifido, nuculis grosse tuberculato-asperis obtusiusculis ②. Ic. Fl. Græc. tab. 179. — *E. macranthum* Viv. Cors. — *E. violaceum* Koch. Syn. et p.c. Prodr. (ex parte) non Linn.

Hab. in pratis ruderatis regionis inferioris copiose, Græcia (Sprunt! Heldr!), Macedonia et Thracia 0-600' (Griseb!), Byzantium (Cast!), Bithynia (Clem!), Paphlagonia (Tchiht), Lydia, Lycia et Caria (Pin! Heldr!), Syria littoralis (Bl! Gaill!), Palestina (Boiss!), Pontus (C. Koch), Transcaucasia ad fines Turcicas (Ledeb.), Tauria (Stev.).

Ar. Geogr. Europæ regio mediterranea a Galliâ australi. Italiâ, ad Dalmatiam, Africa borealis, Madera, insulæ Canarienses.

+ + Stamina inclusa.

11. ***E. amœnum*** (Fisch. Mey. Ind. IV, Petrop. p. 35) biennæ vel perenne, pube molli setisque longioribus e tuberculo ortis ad caules patulis ad folia adpressiusculis obsitum, caulis erectis vel ascendentibus-decumbentibus superne ramosis, foliis uninerviis inferioribus oblongo-lanceolatis obtusis petiolatis, superioribus a basi ovatâ sessili acuminatis, spicis simplicibus demum valde elongatis, calycis strigosi laciniis lanceolatis acutis demum valde accretis, corollæ magnâ cœruleo-violacea adpresso et parce hirtulâ a basi sensim ampliatâ calyce triplo longiore, staminibus inclusis, stylo hirsutissimo bifido, nuculis magnis acutis crenatibus tuberculatis ♀. *E. Orientale* C A. Mey. Enum. p. 97 non L.

Hab. in arenâ fluvii Lenkoran ad mare Caspium et in montibus Talyshch 5400' (C.A. Mey! Hoh!, in montibus Ghilani (Auch. 5008! Buhse!), in monte Ssahend prov. Aderbidjan (Buhsel), in prov. Asterabad prope Siaret (Bge!).

Folia radicalia *E. plantaginei* sed non lyrata, rami terminales pauci in spicas demum fructu maturante pedales abeuntes, corolla maxima 42-45 lineas longa. Facies *E. grandiflori* De-f = *E. australis* Lam. = *E. Cretici* Moris Sard. tab. 97 an Linn? canthæ valde strigoso, calyce vix acercenti, corollæ dimidio minore et angustiore, nucularum minorum tuberculis paucis crassis diversi.

12. ***E. setosum*** (Vahl Symb. II, p. 35) annuum, setis e tuberculo ortis tenuibus foliorum adpressiusculis caulis et spicarum patentibus obsitum pallide virens, caulis procumbentibus hispidis parte inferiori sepe ramosis, foliis radicalibus oblongo-spathulatis in petiolum brevem attenuatis, caulinis sessilibus acutiusculis, spicis terminalibus scorpioideis demum valde elongatis albo-hirsutissimis densifloris, foliis floralibus lanceolatis albo-ciliatis calycis nou acercentis laciniis lineares aequantibus, corollæ cœruleæ adpresso hirtâ obconico-tubulosâ calyce duplo longiore limbo parum ampliato, staminibus inclusis, stylo bifido lobis apice capitulatis, nuculis minimis ovato-

triquetris acutis tenuiter tuberculatis ♂. Ic. Del. Eg. tab. 17, fig. 2. — *E. verecundum* Vivian. Eg. Dec. N° 25 et herb!

Hab. in aridis et arenosis Ægypti inferioris ad Alexandriam (Del., Ehrenb! Samar! Gaill! etc.), ad Rosette (Ky. 593!).

Caules 4-6-pollicares tandem elongati, spicæ demum 4-6-pollicares, corolla 7 lineas longa, nuculæ e minimis lineam dimidiata latæ.

13. ***E. arenarium*** (Guss. Ind. Hort. Boccad. 1825) annum et biennie, setis e tuberculo ortis foliorum adpressis caulis et ramorum patulis obsitum, caulibus brevibus ascendentibus vel prostratis simplicibus vel ramosis, foliis radicalibus oblongo-spathulatis obtusis in petiolum attenuatis, caulinis sessilibus a basi subdilatatae sæpe semiamplexicauli lanceolatis, spicis demum elongatis laxis, bracteis subcordato-oblongis, floribus inferioribus sæpe extraaxillaribus, calycis strigosi fere immutati laciniis lanceolatis obtusis, corollæ minutæ tubuloso-obconicæ hirtæ calyce subduplo longiore, staminibus non exsertis, stylo longe bifido, nuculis acuminatis acute tuberculatis ♂. Ic. Guss. Pl. rar. tab. 17.

Hab. in littoribus Atticæ (Sprun! Boiss! Sart!), Argolidis (Sprun!), Cretæ (Heldr!), insulæ Syra (Weiss!).

Ab *E. calycinum* calycibus fructiferis non ampliatis, ab *E. setoso* spicis laxis minus longe et dense setosis et floribus inferioribus extraaxillaribus, calycis laciniis latioribus, nuculis acuminatis majoribus egregie distinctum.

Ar. Geogr. Corsica, Italia, Sicilia, Africa borealis, insulæ Canarienses.

14. ***E. calycinum*** (Viv. Ann. bot. I, p. 164) annum, setulis basi tuberculatis caulis et ramorum patentibus foliorum adpressis obsitum, caulibus ascendentibus vel prostratis in spicam simplicem rarius parce ramosam foliosam demum laxam abeuntibus, foliis radicalibus rosulatis oblongis vel oblongo-spathulatis obtusis in petiolum brevem attenuatis, superioribus sessilibus interdum semiamplexicaulis, floribus inferioribus sæpe extraaxillaribus, calycis adpresso setosi laciniis lanceolatis fructiferi valde auctis foliaceis oblongis, corollæ anguste tubuloso-obconicæ cæruleæ calyce subduplo longiore extus hirtæ limbo vix ampliato, staminibus inclusis, stylo longe bifido ♂. Ic. Viv. Fl. Ital. Fragm. II, tab. 4. — *E. Creticum* Fl. Græc. tab. 183 non L. — *E. prostratum* Ten. Fl. Nap. tab. 12 non Desf. — *E. parviflorum* MENCH.

Hab. ad vias, in collibus siccis rupestribus et arenosis maritimis, regio inferior Graeciae totius et ejus insularum! Thraciæ et Bithyniæ (Griseb.), Syriæ ad Aleppo (Auch. 2412!).

Ar. Geogr. Hispania littoralis, Gallia australis, Corsica, Sardinia, Sicilia, Italia, Dalmatia, Africa borealis.

ECHIOCHILON (Desf. Atl. I, p. 167).

Calyx 5-partitus laciniâ alterâ breviore sæpe evanidâ. Corolla tubo gracili subarcuato, fauce nudâ, limbo patente bilabiato, labio supe-

riore obsolete bilobo inferiore trilobo. Antheræ 5 intra tubum subsessiles. Stylus inclusus, stigma bilobum. Nuculae 4 ovatæ tuberculatæ areæ basilari planâ fundo calycis affixæ. — Suffrutex.

1. **E. fruticosum** (Desf. loc. cit. tab. 47) fruticosum a basi ramosissimum, totum setulis albis adpressis obsitum, ramis tortuosis divaricatis intricatis cortice albâ vestitis, foliis minutis oblongis vel linearibus obtusiusculis, axillis sâpe gemmiferis, floribus paucis ad axillas supremas solitariis subsessilibus breviter et laxiusculè spicatis, calycis laciniis lanceolatis, corollâ cœruleâ 5. *Lithospermum divaricatum* Sieb. in Spr. Syst. I. p. 543.

Hab. in arenosis maritimis et desertorum, Ægyptus ad Alexandriam (Cadet! Sam! Gaill!), deserta inter Cahiram et Suez, Arabia Petrea (Boiss!), Syria littoralis ad Joppen (Bové!), ad Berythum (Pestalz). Calessyria prope Zachle (Ehrenb!).

Ar. Geogr. Africa borealis, Nubia littoralis.

MACROTOMIA (D.C. Prodr. X, p. 26. — *Munbya* Boiss. Diagn. Ser. I, 11, p. 114 — *Arnebia* species perennes D.C. Prodr.

Calyx quinquepartitus laciniis fructiferis valde elongatis. Corolla infundibuliformis tubo longissimo, fauce nudâ, limbo patent. Stylus bifidus vel stigma dicephalum. Antheræ oblongæ inclusæ subsessiles. Nuculae ovatae vel triquetrae rectæ acutæ tuberculato-rugosæ areolâ basilari subconcavâ. — Herbæ perennes floribus magnis capitato-congestis. — Genus ab *Arnebia* facie, statione alpinâ, inflorescentiâ, corollâ basi annulo nectariformi destitutâ distinctum.

1. **M. echiooides** (Linn. Sp. p. 199 sub *Lycopsis*) e rhizomate squamoso pluricaulis, pilis brevibus patentibus molliter hirta pallide virens, caulis elatis extraosularibus a basi ascendentè erectis foliosis, foliis radicalibus oblongo-lanceolatis obtusis utrinque attenuatis, caulinis et bracteis calyces superantibus a basi latiore semiamplexicauli oblongis acutis, racemis 2-3 floriferis in capitulum congestis fructiferis valde elongatis laxiusculis, calycis fructiferi valde dilatati laciniis tandem lanceolatis elongatis obtusis, corollæ flavæ pubescens tubo calyce 2-2 1/2-plo longiore limbo amplio lobis rotundatis, stylo bifido vel stigmatibus 2 sessilibus globosis, nuculis ovatis acutis obsolete ruguloso-serobiculatis ♀. Ic. Buxb. Cent. I, tab. 4. — *Anchusa echiooides* M.B. I, p. 123. — *Lycopsis setosa* Lehni. Asp. p. 269. — *Lithospermum setosum* Fisch. et Mey. Ind. I. — *Arnebia echiooides* Alph. D.C. Prodr. X, p. 96. — *Arnebia longiflora* C. Koch Linn. XXIII, p. 640. — *Aipyanthus echiooides* Stev. Bull. Mosq. 1854, p. 600.

Hab. in regione alpinâ Ponti Laziei supra Djimil 7500' (Ball!). Armenia ad

Ispir (Huet!), Guriæ (Nordm.), Suaniae (Radde!), Caucasi centralis 6-7000' (Stev! Rehm!) et orientalis in Daghestaniâ 6500'-8500' (Rupr!), Persiae bor. (Auch. 5000!, Buhse!).

2. M. cephalotes (Alph. D.C. Prodr. X. p. 96 sub *Arnebiā*) collo crasso valde squamoso fasciculos foliorum et caulem foliosum edente, foliis adpressissime setosis canescensibus vel canis radicalibus longe linear-lanceolatis vel lanceolatis, caulinis brevioribus acuminatis, inflorescentiâ patule hispidâ flavidâ, racemis in capitulum amplum conglobatis fructiferis parum elongatis, calycis longissimi lacinii linear-lanceolatis acutis, corollæ aureæ pubescentis tubo calyce parum superante vel sesquiolongiere limbo ampio, stigmate bipartito, nuculis magnis serobiculato-punctatis obsolete tuberculatis triquetro-ovatis apice attenuatis acutis ancipitibus ♀. *Munbya cephalotes* Boiss. Diagn. I. cit. p. 116 et *Munbya conglobata* Boiss. — *Arnebia densiflora* et *Lith. densiflorum* Ledeb. Fl. Ross. et herb!

Hab. in rupestribus regionis alpinæ, mons Chelmos Peloponnesi prope Stygem (Orph. Fl. Gr. exs. 112), Olympus Bithynus (Clem!), montes Anatoliæ borealis (Wied! Noë!), Lyciæ (Pestal!), Tauri Cilicici 7000'-8000' et ditionis Kassan Oglu (Ky! Bal!), montis Aslandach Cappadocie (Bal!), Akdag (Auch!) et Berytdagh Cataoniæ (Hau-sk!), Transcaucasiae ad fines Turcicas (Nordm.).

Alpium Orientis decus, caulis 5-6-pollicaris sed etiam pedalis, folia radicalia 4-5 poll. longa 3-6 lin. lata, corymbus capituliformis ovo major, calyx 9-15 lineas sæpe longus proportione cum tubo corolla varians. Radix secca chartam colore violaceo tingens.

3. M. cyanochroa (Boiss. Diagn. Ser. I, 7, p. 33 sub *Lithospermo*) setis patulis basi tuberculatis asperillum, collo crasso valde squamoso, caulinis pumilis folia radicalia non multum superantibus, foliis radicalibus numerosis anguste lanceolato-linearibus, caulinis abbreviatis a basi latiore oblongo-lanceolatis, racemis 2-3 terminalibus floriferis brevibus in capitulum confertis fructiferis parum elongatis, calycis strigosi lacinii linearibus post anthesin valde elongatis, corollæ cyanæ puberulae tubo calyce dimidiâ parte longiore limbi brevis lobis ovatis, nuculis ovato-trigonis apice ancipitibus grosse cristato-tuberculatis ♀. *Arnebia tingens* Alph. D.C. Prodr. X. p. 96. *Munbya cyanochroa* Boiss. Diagn. Ser. I, 11, p. 115.

Hab. in rupestribus regionis alpine Persiae australis, mons Kuhdaëna (Ky. 5000!), mons Zerdou (Auch. 2302!). montes Kellal et Ssebsekuh (Haussk!).

Planta 6-8-pollicaris, folia radicalia 2-3 pollices longa tres lineas lata, corolla 7-8 lineas longa, calyx fructifer fere pollicaris. Radix chartam violaceo colore tingens. Species affinis *M. perenni* (D.C. sub *Arnebiā*) a quâ *M. endochroma* Indica non sat differre videtur; ab ea distinguitur indumento asperimo, foliis caulinis dilatatis, etc.

ARNEBIA (Forsk. Eg. p. 62).

Calyx quinquepartitus. Corolla infundibuliformis tubo elongato inâ basi annulo nectariformi obsito, sauce nudâ, limbi lobis paten-

tibus. Antheræ breviter tubo affixæ inclusæ. Stylus apice bifidus ramis sæpius bilobis. Nuculae triquetrae acutæ tuberculatae vel sericeulatae-rugosæ rarius levæ. — Herbae annuae flavigloræ desertorum incoleæ. — Genus stylo bifido a *Lithospermo* distinctum.

* Calyx fructifer vix accretus.

1. **A. hispidissima** (Spreng. Syst. I. p. 556 sub *Diocled*) annua, tota setis albis patulis basi grosse tuberculatis seberrima a basi ramosissima pumila, foliis lanceolatis obtusiusculis, floralibus anguste linearis-lanceolatis acutis, racemis secundis densis, calycis fructiferi vix aucti lacinias angustæ lanceolatis, corollæ flavæ patule pubescentis tubo calyce duplo longiore, stylo bipartito stigmatibus complanatis, nuculis minimis trigonis acutis tuberculatis ⊖. *A. hispidissima* D.C. Prodr. X. p. 94. — Ic. Jaub. et Sp. III. tab. 363. — *Lith. hispidissimum* Lehm. Ic. tab. 39. — *Lith. vestitum* Wall. — *Anchusa asperrima* Del. Ill. № 210.

Hab. in desertis Ægypti ad Abukir (Del!), prope Cabiram (Sieb!), inter Suez et Gaza (Bové), in Arabiâ Petréa ad Ouadi Hamine et Ouadi Feiran (Schimp. 213!), inter Keneh et Kosseir (Schweinf!), in desertis Kischma Persiae australis (Auch. 4981!), in Belutschâ (Stocks!), Arabice *Attahn*.

Setis albis strigosissimis, calyce fructifero non aucto 3 lineas longo distinctissima, corolla sex lineas fere longa.

Ar. Geogr. Nubia tropica, regio Scinde, India boreali-occidentalis.

2. **A. Griffithii** (Boiss. Diagn. Ser. II, 3. p. 135) annua, setis albis patulis basi tuberculatis hirsuta, caulinis pumilis, foliis oblongis et oblongo-lanceolatis obtusis basi longe attenuatis, racemis 2-3 brevibus densis, bracteis angustis calyce vix brevioribus, calycis albo-hirsuti fructiferi fere immunitati lacinias angustissime linearis-subulatis, corollæ hirsutæ flavæ tubo calyce duplo longiori a parte mediâ sensim in limbum hypocrateriformem ampliato, stylo bipartito, nuculis parvis triquetris acutis elevatim rugoso-cristatis subtuberculatis ⊖.

Hab. in Affghanâ (Griff. exs. 5953!).

Planta 3-4-policaris, calyx fructifer 5 lineas tantum longus, corolla 9 lineas longa. Ab *A. hispidissimâ* strigositate minore, corollâ longâ sensim dilatata limbo ampio, nuculis rugoso-suberistatis nec tuberculis distinctis obsitis differt.

* * Calyx fructifer valde accretus tubo cornuto-cristato vel tuberculato.

3. **A. cornuta** (Ledeb. Pl. Alt. I. p. 175, Icon. I, tab. 25 sub *Lithospermo*) annua, pumila erecta setulis aliis adpressis aliis patulis flavicantibus hispida, caule a basi sæpe ramoso, foliis inferioribus linearis-oblongis cæteris lanceolato-linearibus acutis, racemis fructiferis elongatis laxiusculis, bracteis calyces æquantibus vel superantibus, calyce hispido in lacinias lineares fisso fructifero valde acreto basi indurata incassato pentagono angulis gibbosso-prominentibus

plus minus cristatis laciinis elongatis anguste linearibus uninerviis conniventibus, corollæ luteæ hirsutæ tubo calycem vix vel duplo superante, stigmate bi vel quadrisido, nuculis parvis triquetris acutis inaequaliter tuberculatis ♂. *A. cornuta* F. et M. Ind. Petr. 1835. — *Onosma Orientalis* Pall. It. tab. 50. — *Lithospermum decumbens* Vent. Cels. tab. 37 (nomen antiquius sed inaptum). — *Lith. micranthum* Viv. Lib. IV, tab. 1. — *Lith. tubatum* Bertol. Mise. I, p. 14. — *Onosma asperifolium* Lehm. Asp. II, p. 372. — *Arnebia Vivianii* Coss. et Dur. (forma calyce majore). — *A. calycina* Stev. Bull. Mosq. 1831, p. 398.

Hab. in arenosis et desertis Arabiae petreæ ad Ouadi Uscit et in deserto Tih (Boiss!), deserto Syriaco ad Palmyram (Bl!), ad Aleppo et in Mesopotamia (Haussk!), Babyloniâ (Auch. 2359!), Armeniâ Turcicâ prope Van (Noë!) et Rossicâ (C. Koch), Georgiâ Caucasicâ (C.A. Mey! Hoh!), Persiâ boreali (Szov! Buhset!), media (Bge!), australi (Ky. 84!), Turkestaniâ ad Caspium et prope Buchara (Lehm!).

Multum variat antherarum insertione, styli divisione, corollæ magnitudine et formâ (nunc 4 linea, nunc 8-9 longâ); formæ subbisexuales prob. adsunt.

Ar. Geogr. Africa borealis interiore ab Algeriâ ad regnum Tripolitanum, Rossia austro-orientalis, Songaria, Sibiria Altaica.

4. *A. linearifolia* (D.C. Prodr. X, p. 95) annua, tota setis flavescentibus ad caules foliorum margines et inflorescentiam patulis strigulosa, caulis pumilis rectis a collo saepius pluribus ramosis ramis a basi in racemos densissimos breves distichos abeuntibus, foliis radicalibus parvis oblongo-spathulatis subpetiolatis, caulinis latiuscula oblongo-linearibus obtusis, bracteis linearibus acutis calyce subbrevioribus, calyce fructifero valde accreto basi inflato vix indurato ad nervos prominentes longe setoso et albo-tuberculato laciinis longissimis late linear-lanceolatis obtusiusculis trinerviis. corollæ luteæ villosæ tubo ealyce (florifero) sesquiloniore limbo parvo, nuculis triquetris acutis grosse serobiculatis et tuberculatis ♂. *A. flavescens* Boiss. Diagn. Ser. I, 41, p. 117.

Hab. in arenosis desertorum, Ægyptus prope Kahiram (Husson!), Arabia petrea in regione Hauara (Schimp. 398 sub *Lith. Arnebiâ!* Boiss!, Auch. 2368!), in deserto Tih (Boiss!), deserto Syriaco ad Palmyram (Bl!), Armeniâ Rossicâ ad Nachitchewan (Szov!), Persiâ ad Firouz Kuh (Bode!) et inter Ispahan et Teheran (Bge!), Belutschia (Stocks!).

Differit ab *A. cornuta* racemis fructiferis brevibus densissimis, calycis fructiferi multo magis aucti tubo non indurato pentagono cornuto et cristato sed membranaceo tuberculato et hispido, laciinis multo majoribus (sæpe pollicariibus 2-3 linea lati) trinerviis. Nuculae duplo majores.

* * * Calyx fructifer accretus tubo lævi laciinis elongatis.

5. *A. tinctoria* (Forsk. Eg. p. 63) annua, dense et adpresso setis tuberculatis cinerascens nana a basi breviter ramosa, foliis oblongo-lanceolatis obtusis basi attenuatis, racemis etiam fructiferis brevibus densis secundis, bracteis linearibus obtusis calyces subsuperantibus,

calycis hirsuti fructiferi aucti tubo kevi laciniis elongatis anguste linearibus obtusis subtrinerviis, corollæ glaberrimæ obscure violaceæ tubo calyce nou longiore limbo minuto, stigmate quadrifido, nuculis triquetris acutis laeviuseulis nitidis ☺. *Lithospermum tetragynum* Lam. Encycl. III, p. 30. — *Lith. tectorium* Vahl Symb. II, 33, t. 28. — *Lith. Arnebia* Del. III, № 203.

Hab. in desertis Ægypti prope Kahiram (Forsk.), Arabiæ petreæ (Boiss!), in regione Hauara (Schimp. 203!) mixta cum *A. linearifoliâ*. Arabice *Sagaret el arneb*.

Planta 2-3-pollicaris ab *A. linearifoliâ* calyce fructifero multo minus acreto 5 lineas tantum longo basi laevi, corollæ brevi glabrâ violaceâ, nuculis levibus distincta.

6. A. Bungei, annua, setis brevissimis basi tuberculatis ad foliorum margines longioribus obsita, caule pumilo hirtulo fere a basi ramoso, foliis oblongo-lanceolatis et linearibus obtusis inferioribus basi longe attenuatis, racemis etiam fructiferis brevibus densissimis distichis, bracteis linearibus obtusis calyce brevioribus, calycis fructiferi laciniis elongatis anguste linearibus obtusis uninerviis, corollæ luteæ hirtæ tubo calyce florifero subdupo longiore, limbi ampli lobis repando-crenatis, nuculis (junioribus) scrobiculato-rugosis ☺.

Hab. ad Ssertschah Persiæ mediae orientalis (Bge!). Fl. Mart.

Planta 3-4-pollicaris, facies et corollæ limbus amplius *A. fimbriopetalæ* a quâ differre videtur pilis praeter cilia foliorum brevissimis, laciniis calycis fructiferi elongatis quidem (9 lineas longis) sed angustis lineam dimidiata latis, limbi corollini lobis obsolete crenulatis nec longe fimbriatis.

7. A. fimbriopetala (Stocks in Hook Journ. et Kew Misc. III, p. 180, tab. 6) annua, setis adpressis basi tuberculatis aspera ad foliorum margines longius ciliata, caule humili erecto, foliis longe et anguste linearibus obtusis, racemis 2-5 etiam fructiferis prævious densissimis distichis, bracteis linearibus obtusiuseulis calyce fructifero brevioribus, calycis laciniis linearibus obtusis corollinum tubum dimidium æquantis fructiferis valde elongatis subtrinerviis reticulatum nervulosum, corollæ villosulæ flavae limbo ample lobis inciso-fimbriatis sinibus intruso-plicatis, stylo vix bifido nuculis verruculosis triquetris acutis basi acute bigibbis ☺.

Hab. in Belutschia superiore in rupestribus ad Doubund. Chehel Tun (Stocks!), in Persiæ australis graminosis gypsaceis prope Kumaredj prov. Schiraz (Haussk!).

Planta 4-6-pollicaris, folia longa 4 $\frac{1}{2}$ lin. lata, calycis laciniæ sex lineas fructiferæ pollicem et amplius longæ, corollæ tubus saepè pollicaris, limbus diametro 8-10 lineas latus lobis fimbriatis quâ notâ hec species elegantissima ab affinis differt.

* * * * Species anomala.

8. A. obovata (Bge pl. Lehm. p. 407) perennis, molliter setuloso-canescens, caulinis dichotome et divaricatum ramosis, foliis

adpresso strigulosis obovatis basi cuneato-attenuatis, racemis terminalibus solitariis confertifloris fructiferis elongatis laxiusculis secundis, bracteis linear-lanceolatis acutiusculis calyce brevioribus, calycis laciinis linearibus demum elongatis, corollæ violaceæ pubescentis tubo calyce duplo longiore limbi patentis lobis rotundatis, filamentis paulo infra faucem insertis antheræ exsertæ brevioribus, stylo breviter bitido, nuculis acutatis lateribus bigibbis tenuissime granulatis 2.

Hab. ad rupes Turkestaniæ ad fortalitium Fon prope fluvium Sarafschian (Lehm!). Non vidi.

Planta 6-9-pollicaris. folia intermedia 1 1/2-pollicaria superne semipollicem lata, racemi fructiferi semipedales, calyx florifer 4 lineas laciinis fructiferis 7 lineas longis. Corollæ tubus circ. 7 lineas longus gracilis. Species mihi descriptione tantum nota ab *Arnebiâ* cui eam am. auctor adnumerat radice perenni et imprimis antheris exsertis discedens. Corollæ tubus ut in *Arnebiâ* basi annulo nectarifero instructus. An genus proprium?

LITHOSPERMUM (L. Gen. 181).

Calyx quinquepartitus. Corolla infundibuliformis fauce perviâ vel nudâ, vel plicis velutinis longitudinalibus inter stamina obsita. Antheræ incluse filamentis brevissimis. Stylus indivisus stigmate capitulo interdum subbilobo. Nuculæ trigonæ rugoso-tuberculatae vel ovatae laevissimæ areolâ basilari planâ. — Herbæ vel suffrutesc.

SECT. I. RHYTISPERMUM Link. — Nuculæ ovato-triquetrae scrobiculatae vel echinatae. Corollæ faux plicis longitudinalibus velutinis obsita.

1. **L. arvense** (L. Sp. 190) annum, adpresso strigulosum, caulis erectis, foliis inferioribus obovato-oblongis in petiolum attenuatis superioribus lanceolatis, racemis fructiferis laxissimis, pedicellis fructiferis brevissimis vix incrassatis, bracteis flores superantibus, corollæ albæ rarius cærulescentis tubo hirto calyce non longiore, nuculis ovoideo-trigonis acutis scrobiculatis et acute tuberculatis ♂. *L. bicolor* Bert. Misc. I, p. 14. — *L. Rochetii* Friv. Flor. 1835, p. 336 (forma foliis latioribus et corollæ limbo submajore, specimina cæterum non fructifera). — *L. Leithneri* Heldr. (forma montana minor).

Hab. in arvis et etiam sterilibus montanis totius ditionis a Græciâ! Tauriâ, regionibus Caucasicis! ad Persiam borealem! Syriam! Affghaniam et Belutschiam!

Ar. Geogr. Europa tota a Scandinavia ad Rossiam septentrionalem, Sibiria tota, Japonia, Africa borealis. In Americam migravit.

2. **L. Sibthorpiatum** (Griseb. Spicil. II, p. 86) annum, adpresso strigulosum canescens, a collo ramosum ramis brevibus prostratis, foliis anguste lanceolatis, racemis fructiferis laxis, bracteis

calyces parum superantibus, pedicellis fructiferis brevissimis vix incrassatis, corollæ albæ rarius cærulecentis tubo hirto calyce non longiore, nuculis lœvibus ovoideo-triquetris acutis scrobiculatis subtuberulatis ⊖. *L. tenuiflorum* Fl. Græc. tab. 159 et Ledeb. Fl. Ross. non L.

Hab. in collibus, cultis regionis montanæ et etiam in arenosis maritimis, Attica ad Phalerum (Heldr! Clem!) et in montibus Parnes, Pentelico, Hymetto (Sprn! Boiss! Heldr!), in Bithynia maritimis (Griseb!). Smyrnae (Bal!), Cypro (Sibth.), Lycia montibus in regione alpinâ (Bourg!), Ciliciae ditione Kassan Oglu (Ky. 63!). Mesopotamiâ (Haussk!), Georgiâ Caucasicâ (Hoh. exs. sub *L. tenuifloro!*), Persiâ boreali (Auch. 5013!).

Corollæ tubus a cl. Grisebach inflatus dictus mihi nec magis nec minus cylindricus ac in *L. arvensi* videtur a quo haec species caulinis prostratis abbreviatis, nuculis minoribus (an satis?) differt.

3. **L. incrassatum** (Guss. Prodr. Fl. Sic. I, p. 211) annum, virens adpresso strigulosum, caulinis flexuosis erectis vel decumbentibus, foliis radicalibus latioribus obovatis superioribus linearibus obtusis, racemis fructiferis elongatis laxissimis, pedunculis fructiferis valde incrassatis obconicis apice calyci aequilatis, corollæ cærulecentis rarins albidae tubo hirto calycem non superante, nuculis parvis ovoides acutis scrobiculatis subtuberulatis ⊖. *L. Gasparrinii* Heldr. in Guss. Syn. I, p. 217 forma albiflora.

Hab. in arenosis et cultis montanis et subalpinis, Taygetus 6500', mons Panachaicon Achaiae, Parnes (Heldr!), mons Ida Cretæ (Heldr!), mons Peristeri Macedoniae (Griseb.), Libanus ad Cedros et alibi (Ehrenb! Bl! Gaill!), Antilibanus (Ky!), Anatolia orientalis in montibus Mesogi et Cadmo (Boiss!), Phrygiâ supra Ouchak (Bal!). Crescit quoque in demissioribus, Mesopotamia ad Biredjik (Haussk!), Cyprus (Ky!), Egyptus ad Alexandriam (Ehrenb!).

Ar. Geogr. Galliæ austro-orientalis et Hispaniæ australis montes, Italiæ meridionalis et Siciliæ, Dalmatiæ, Africæ borealis.

4. **L. tenuiflorum** (L. fil. Suppl. 130) striguloso-hispidum sæpe fulvescens, caule erecto saepius pumilo superne ramoso, foliis radicalibus oblongo-spathulatis breviter petiolatis, caulinis oblongo-linearibus, bracteis calyces subsuperantibus, racemis fructiferis tandem laxiusculis, pedicellis brevissimis immutatis, corollæ cæruleæ rarius albæ tubo angusto hirsuto, nuculis minimis tuberculatis trigonis laterolifer sæpe bigibbis in rostrum breve contractis ⊖. *L. fulvidulum* Bert. Misc. I, p. 13. — Jacq. Ic. rar. II, tab. 313. — *Myosotis tenuiflora* Viv. Lib. p. 9, tab. I.

Hab. in collibus et cultis Græciae in Argolide et ad Chalcidem Eubœa (Sprn!). Atticae circa urbem et in monte Hymetto (Boiss! Heldr! Orph. Fl. exs. 151), Phrygiâ ad Ouchak (Bal!), Syriâ littorali (Bl!) et interiori ad Damascum (Gaill!). Aleppo et Aintab (Ky. 4! Haussk!), Hierosolymæ (Boiss!), Egypto (ex Delite), Mesopotamiâ (Chesn! Haussk!), Persiâ occid. ad Mendeli (Noël), et mediâ (Derderian!).

A præcedentibus nuculis minute tuberculatis nec scrobiculatis et formæ diversæ egregie distinctum. *L. minimum* Moris planta Sardoa nuculis scrobiculatis omnino alienum est.

Ar. Geogr. Dalmatia, Africa borealis.

5. **L. Apulum** (L. Sp. 189 sub *Myosotide*) annum. setis tuberculatis hirsutum, caule pumilo rigidulo simpliciusculo, foliis linearibus angustis uninerviis radicalibus subspathulatis caulinis strictis, racemis subcorymbosis fructiferis densis parum elongatis foliaceo-bracteatis, corollæ flavæ tubo calycem superante, fauce puberulo-annulata, nuculis ovato-triquaetris lucidis tuberculatis gibbosis ⊖. *L. Apulum* Vahl Symb. II, p. 32. — Ie. Fl. Græc. tab. 158. — *L. strigosum* M.B. Taur. Cauc. III, p. 121. — *L. Mesopotamicum* D.C. Prodr. X, p. 75.

Hab. in collibus siccis regionis inferioris, Peloponnesus (Sprun!), Attica (Heldr!), Creta (Sieb!), Archipelagus (Sibth. Auch!). Anatolia orientalis (Fleisch! Auch. 2329!). Bithynia (Wied!), Cyprus (Ky!), Syria bor. ad Aintab (Haussk!), Mesopotamia (Auch. 1484!), Tauria (Stev!).

Ar. Geogr. Europa et Africa borealis regio Mediterranea. Planta sub hoc nomine in insulis Canariensisibus indicata est species diversa, *L. microspermum* Boiss. Diagn. Ser. II, 3, p. 435 = *L. arvense* var. *carrulescens* Bourg. exs. 318 = *L. Webbii* Cossion et Dur. in Bal. Marocc. 1867.

SECT. II. EULITHOSPERMUM D.C. — Nuculae ovatæ levissimæ. Corolla faux sub lobis gibbis vel plicis velutinis instructa.

6. **L. officinale** (L. Sp. 189) perenne. indumento adpresso aspernum, caule firmo elato folioso superne striete ramosissimo, foliis oblongo-lanceolatis oblongisque acutis venosis, racemis fructiferis laxis bracteis calyces longe superantibus obsitis, pedicellis non incrassatis, calycis fructiferi vix aucti setosi laciniis lanceolatis obtusis, corollæ albido-flavidæ extus hirsutæ tubo calycem non superante limbo parvo, nuculis ovatis obtusiusculis albis nitidis levibus ♀. Ic. Fl. Dan. tab. 1084.

Hab. in pratis humidis dumosis, sylvaticis, pomariis totius ditionis praesertim septentrionalis a Peloponneso (Bory!), ad Thraciam et Bithyniam (Thirke!), prov. Caucasicas et Iberiam!, Persiam borealem in prov. Aderbidjan (Szov!) et Khorassan (Bge!), Kurdistaniæ Persiae monte Avroman (Haussk!). Ex Anatoliâ australi et Syriâ nondum notum.

Ar. Geogr. Europa omnis a Scandinavia et Rossiâ ad Hispaniam centralem, Italiam, Dalmatiam, Sibiria tota, Japonia, America borealis.

7. **L. purpureo-cæruleum** (L. Sp. 190) perenne, adpresso hirtum, surculis repentibus, caulis floriferis erectis apice 2-3-fidis, foliis lanceolatis acutis basi attenuatis, racemis fructiferis demum laxissimis, pedicellis brevibus fructiferis incrassatis, calycis laciniis anguste linearibus fructiferis valde elongatis, corollæ purpureæ tandem azureæ hirtæ calyce longiore limbo ampio, nuculis ovatis obtusis albis nitidis levibus ♀. Ic. Jacq. Austr. tab. 14. — Rehb. t. 112.

Hab. in sylvaticis Peloponnesi (Bory!), Bithyniae (Calvi!) et in Olympo Bithyno (Barbey!), Phrygiæ (Auch!), totius Anatoliæ bor. (Wied!), Ponti et

Armeniæ Turcicæ (Tohih! Bourg!), in monte Djebel Nur Ciliciae (Ky. 48!), Tauriæ, Caucaso, Iberiæ Caucasica (M.B., Ower!, Szov!), Persia bor. prov. Asterabad (Bge!). Fl. vere.

Ar. Geogr. Europa media et australis ab Anglia, Gallia et Germania media ad Rossiam medium.

Sect. III. *LITHODORA* Griseb. Spicil. II, p. 85. — Nuclei ovato-triquetrae laeves vel minutissime tuberculati. Corollæ fauæ plicis destituta glabra vel hirtula.

8. L. callosum (Vahl Symb. I, p. 14) suffruticosum, a rhizomate lignoso ramosissimum dumulosum ramis crassis pumilis erectis vel ascendentibus ramulosis, totum setis albis adpressis inaequalibus strigosissimum canescens vel canum, foliis breviter lanceolatis acutiusculis calloso-marginatis, racemis brevibus spiciformibus densifloris tandem elongatis laxiusculis, bracteis anguste lanceolatis calyce longioribus, calycis lacinias linearibus strigosissimis, corollæ cœruleæ calyce $2\frac{1}{2}$ -3-plo longioris tubo patule hirta a basi angustiori sensim ampliato limbi brevis lobis oblongis, fauce hirtulæ, nuculis ovoideo-triquetris acutis nitidis præter 1-2 tuberculos saepe deficientes laevibus ȝ. Ic. Del. Eg. tab. 16, fig. 2. — *L. angustifolium* et *L. latifolium* Forsk. Deser. 39. — *L. ciliatum* Lehm. Asp. p. 314 et herb!

Frutex semipedalis, corollæ sex lineas longæ.

Hab. in Ægypti desertis arenosis ad Kahiram Forsk. Del., Auch! Boiss! etc., ad Rosette (Ehr!), in Arabiæ Petreæ (Boiss!), ad Gaza Palestine (Bové!), in Persiæ australi ad Abuchir et Dalechi (Haussk!). Fl. Apr. Mai.

Ar. Geogr. Africa borealis in regione Saharensi orientali.

9. L. Kotschyï (Boiss. et Hoh. Diagn. Ser. I, 4, p. 49) suffruticosum pumilum, totum setis adpressissimis æqualibus canum a basi ramosissimum ramis ascendentibus ramulisque crebre foliosis, foliis minimis squamæformibus a basi semiamplexieauli linearis-oblongis patulo-incurvis canaliculato-plicatis acutiusculis, floribus ad axillas foliorum superiorum conformium paucis breviter spicatis, calycis lacinias oblongis obtusis, corollæ cyaneæ hirtulae tubo calyce vix longiore limbi parvi lobis ovatis, fauce hirtulæ, nuculis ovato-triquetris acutis laevibus paucis punctis impressis notatis ȝ.

Hab. in fissuris rupium arenacearum insulae Karrak Persiæ australis (Ky!), in prov. Mekran Persiæ austr. (Auch. 5001!). Fl. Jan.

Suffrutex semipedalis ramis foliis squamiformibus $1\frac{1}{2}$ -2 lineas longis crebre obsitis eis *L. callosi* tenuioribus, corolla 2 lineas tantum longa.

10. L. hispidulum (Sibth. et Sm. Fl. Græc. II, p. 53, tab. 162) suffruticosum, intricatim ramosissimum ramis brevibus siccis subpersistibus, totum setulis adpressissimis et ad margines et nervum foliorum validioribus basi tuberculatis obsitum, foliis pallide virenti-

bus elliptico-oblongis obtusis basi attenuatis, floribus 1-4 inter folia superiora sessilibus congestis, calycis fructiferi vix accreti laciniis lanceolato linearibus adpresso setosis, corollæ glabrae tubo calyce subdupo longiore limbi lobis breviter ovatis, fauce glabra, nuculis parvis ovato-triquetris albis nitidis sub lente minutissime tuberculatis 5.

Hab. in saxosis apricis maritimis Rhodi (Auch! Heldr!), Cretæ in littore meridionali inter Hagios Paulos et Hagio Rumeli (Heldr!), Cypri versus Nicomia (Ky. Suppl. 489!), Cilicie ad occid. Mersina (Bal!. Fl. Mart. Apr.

Dumi semipedales vel paulo elatiore, folia brevia, corolla fere sex lineas longa.

11. **L. rosmarinifolium** (Ten. Fl. Nap. III, p. 178, tab. 114) fruticosum ramosissimum ramis erectis vel dependentibus vetustis denudatis novellis crebre foliosis, foliis longis oblongo- et linearilanceolatis margine subrevolutis supra parcissime subtus dense et adpresso canescensibus. floribus 1-3 terminalibus congestis foliis supremis calyce longioribus bracteatis, calycis laciniis anguste linearibus, corollæ cœruleæ adpresso pubescentis calyce triplo longioris limbo ampio lobis rotundatis, fauce glabra, nuculis albis levibus nitidis 5. *L. fruticosum* Fl. Græc. tab. 161 non L.

Hab. in montosis Græciæ et in Archipelagi insulis (Sibth.), in insula Samo (Urv.). E ditione non vidi.

Ar. Geogr. Regnum Neapolitanum, Sicilia.

Species mihi ignotæ, probabititer synonyma.

L. crassifolium (Lehm. Asp. p. 324) caulinis herbaceis ascendentibus incanis, foliis lanceolato-spathulatis obtusis adpresso canescensibus. floralibus cordatis acutis subglabris, spicis terminalibus conjugatis elongatis. corollis glabris calyce sublongioribus, nuculis rugosis.

Hab. in Oriente.

Deest in herb. Lehmann nunc cl. Sonder.

L. Pythicum (Clarke ex Spr. Neu. Entd. III, p. 165) foliis undulatis inferioribus lanceolato-oblongis superioribus bracteisque cordato-oblongis hirsutis.

Hab. in Oriente.

Prob. *Alkanna Orientalis*.

MOLTKIA (Lehm. Act. Scrut. Hal. 3, f. 2).

Calyx quinquepartitus. Corolla inferne cylindrica superne inflata infundibuliformis lobis rectis, fauce nudâ. Staminum filamenta longa exserta, antheræ oblongæ curvatæ prope basin filamento insertæ. Sty-

Ius filiformis acutus. Nuculae rectæ vel subinecurvæ ovato-triquetrae vel angulatae areolâ basilarî planâ. — Suffrutices. — Genus a *Lithospermî* sectione *Lithodorâ* antheris exsertis et corollæ lobis erectis, antheris curvatis fere basilixis distinctum.

§ 1. **Lithospermoideæ**. — Nucleæ lèvissimæ.

1. **M. petræa** (Trattin. Thes. Bot. p 8 sub *Echioj*) suffruticosum ramosissimum adpressissime sericeo-canescens, ramis tenuibus erectis brevibus crebre et stricte foliosis, foliis margine subrevolutis obtusis inferioribus anguste oblongo-linearibus superioribus linearibus sensim diminutis, racemis 2-4-floriferis in capitulum densiflorum congestis fructiferis subelongatis laxiusculis, calyceis laciniis linearibus, corolla cœruleâ glabrâ cylindrico-obconicâ calyce subtriplo longiore tubo in limbum sensim ampliato, filamentis exsertis, nuculis parvis albis lèvissimis nitidis ovato triquetris apice attenuatis anicipitibus 5. *Lithospermum petraeum* Alph. n.c. Prodr. X, p. 82. — *Lith. rosmarinifolium* Rehb. Fl. exc. I, 337 non Ten.

Hab. in fissuris rupium montis (Eta Thessalicae prope Thermopylas (Sprun!)).

Suffrutex elegantissimus semipedalis vel pedalîs, species inter *Lithospermî* sectionem *Lithodoram* et *Moltkiam* intermedia illi nuculis lèvibus nitidis, huic corollæ lobis erectis nec patentibus, filamentis exsertis et halitu magis affinis.

Ar. Geogr. Dalmatia, Montenegro, Bosnia.

§ 2. **Eumoltkiæ**. — Nucleæ scrobiculato-rugosæ.

2. **M. cœrulea** (Willd. Sp. I, p. 775 sub *Onosmate*) suffruticosa multicaulis setis basi tuberculatis adpressissimis aspera canescens, caulinis erectis punnulis rigidis crassiuseculis crebre foliosis, foliis radicalibus oblongo-spathulatis obtusis basi longe attenuatis, caulinis sessilibus strictis linear-lanceolatis acutiuseculis, racemis 2-5 primum scorpioideis fructiferis elongatis laxiusculis rigidis, bracteis lanceolatis calyces vix superantibus, calycis strigulosi demum accreti laciniis linear-lanceolatis, corollæ cœruleæ parce setulosæ calyce quadruplo longioris tubo tenui in limbum ampullaceum eo breviorem abrupte ampliato, filamentis exsertis, nuculis crassis albis obsolete et grosse rugosis minute depresso-punctatis incurvis rostro brevi horizontali 5. *M. cœrulea* Lehm. I. cit. tab. 44. — *Cynoglossum rugosum* Willd. herb! et *Lith. rugosum* n.c. Prodr. X, p. 83. — *Moltkia punctata* Lehm. loc. cit. et tab. 44 (forma longistyla antheris inclusis). — *M. Anatolica* Boiss. Diagn. Ser. I, 41, p. 114.

Hab. in collibus et incultis Cariae (Boiss! Pinard!), Lyciae (Bourg!), Pisidiæ (Tchih!), Cappadociæ (Bal!), Bithynia (Wied!). Galatiae (Sestini!), Ponti (Tchih!). Armeniae Turcicae ad Baibut (Bourg!), Karpot (Noe!), Erzerum (Calv!), Syria ad Antiochiam (Auch. 2403!).

β. *suberistata*. — Nucleæ grosse tuberculato-rugosæ suberistatae. Videtur intermediis ad typum transire.

Hab. in Syriâ prope Ain'ab et Nisib (Haussk!), ad Djebel Abiad deserti Syriaci ad Palmyram (Bl!), Armeniâ Rossicâ (Buhse!), Iberiâ (Hohen!), Persiæ prov. Aderbidjan (Szov!), inter Teheran et Tabris (Bge!).

In hæc specie ut prob. in congeneribus omnibus forma vulgatior habet filamenta exserta stylo vix superata, sed rarius occurrit forma brachyantha corollâ fere duplo breviore, antherarum apice corollam vix superante et stylo elongato; hæc *M. punctatam* Lehm. sistit quæ nequidem varietas.

3. *M. angustifolia* (D.C. Prodr. X, p. 72) suffruticosa multi-caulis setis adpressissimis asperula, caulis erectis rigidis tenuiter virgatis, foliis anguste linearibus obtusis elongatis, racemis paucis terminalibus fructiferis subelongatis densis, bracteis linearibus calyce vix longioribus, calycis laciniis linearibus obtusis, corollæ cœruleæ glabriuscule calyce quadruplo longioris tubo tenui in limbum breviorum sensim ampliato, filamentis exsertis. nuculis (junioribus) scrobiculato-rugosis 5. *Echium longiflorum* Bertol. Misc. I, p. 45!

Hab. in Assyriâ (Oliv., Auch. 2300!), ad Euphratem (Chesn!), in cretaceis deserti ad radices montis Sindjar (Haussk!).

Difert a præcedente caulis tenuioribus fere pedalibus, foliis angustioribus longioribus (1-1 1/2 poll. longis 4-1 1/2 lin. lati), corollæ tubo sensim nec abrupte ampliato.

4. *M. aurea* (Boiss. Diagn. Ser. I, 4, p. 49) suffruticosa multi-caulis tota adpressissime setulosa asperula, caulis erectis pumilis striete et crebre foliosis, foliis radicalibus oblongis in petiolum attenuatis, caulinis sessilibus linear-lanceolatis acutiusculis, racemis terminalibus 2-3 floriferis capitato - scorpioideis fructiferis elongatis laxiusculis, bracteis calyci æquilonigis, calycis laciniis breviter linearibus obtusis. corollæ aureæ glabré tubo calyce vix duplo longiore in limbum ovatum eo subbreviore abrupte ampliato, filamentis exsertis. nuculis crassis albidis serobiculato-punctatis vix rugosis angulato-ovatis vix rostratis basi angustatis 5.

Hab. in collibus Cariae (Pinard!), Lycie prope Elmalu (Bourg. 492!), Pisidiæ prope Bouldour et ad radices Davrosdagh (Heldr!), Phrygiæ prope Ouchak (Bal!). Fl. Mai.

Species pulchra a præcedentibus colore florum, corollæ tubo abbreviato et nuculis non rostrato-incurvis distincta.

ZWACKHIA (Sendtn. in Rehb. Ic. Germ. XVIII, p. 65, t. 445).

Calyx quinquepartitus. Corolla infundibuliformis limbo profunde et subinequaliter quinquefido, tubo inter filamentorum bases plieis minutis transversis obsito. Filamenta in ostio brevissima ad antheram paulo supra basin affixa, antheræ ovatae basi obcordatae apice connectivo apiculare obsitæ ad loculorum margines crenulato-fimbriatae. Stigma minute capitato-truncatum. Nuculae ovoideo-trigonæ apice in rostrum anceps attenuatae minute tuberculato-reticulatae areolâ basi-

lari ovatâ planâ. — Herba perennis facie *Moltkiae aureæ*. — Genus corollæ laciniis erectis *Moltkiae* afinitis, ab eâ ut a *Lithospermo* praeter corollam subirregularē profunde quinquefidam antheris connectivo apiculatis et ad dehiscentiæ margines fimbriatis distinctum.

I. **Z. aurea** (Sendtn. loc. cit.) parce et adpresse setulosa viridis, rhizomate verticali squamoso, foliis radicalibus fasciculatis anguste et longe linear-i-lanceolatis obtusis basi longe attenuatis, caulinis flori-feris extrarosularibus ascendentibus in racemum simplicem scorpioideum fructiferum elongatum laxum abeuntibus, foliis caulinis diminutis sessilibus, calycis fructiferi accreti laciniis late lanceolatis acutis, corolle flavæ glabré tubo calyce sublongiore limbo tubo æquilongo ad medium in lobos oblongos quinquefido ♀. *Moltkia Sendtneri* Boiss. Diagn. Ser. II, 3, p. 138.

Hab. prob. in Albaniâ et Macedonia sed e ditione nondum vidi.

Ar. Geogr. Bosnia, Serbia.

ALKANNA (Tausch Flor. 1824, p. 234. — *Campylocaryum* D.C.
Mss. — *Camptocarpus* C. Koch Linn. 1843 p. 304.

Calyx quinquepartitus. Corolla regularis infundibuliformis plicis destituta rugis transversis glabris minimis interdum subobsoletis ad faucem perviam instructa basi nectario annuliformi glabro minimo donata. Stamina versus medium tubum spiraliter vel verticillatim inserta inclusa filamentis brevissimis. Stigma capitatum vel subbilibum. Nuculæ 4 vel abortu 2-1 verruculosæ vel scrobiculato-rugosæ a basi anticâ vel modice curvatae sinu basilari tunc lato et rostro horizontali vix descendantí, vel curvatissimæ sinu basilari angusto et rostro fere verticaliter flexo; areolâ planâ sœpe stipitatâ basilari, subbasilari vel in medio sinu sitâ afflxæ. Pericarpium fragile. Semen curvatum. — Herbæ facie *Nonneæ*.

Specierum Orientalium distributio.

* Corolla extus hirta.

A. *Pulmonaria, Scardica, tubulosa.*

* * Corolla extus glabra.

+ Nuculæ reticulatæ vel tuberculis basi confluentibus scrobiculato-rugosæ.

✗ Corollæ cyaneæ.

A. *macrosiphon, strigosa, Syriaca, megacarpa, tinctoria.*

× × Corollæ xanthinæ (Conf. A. Græcam).

1 Nuculæ parum curvatae sinu basili lato rostro horizontali vel deorsum parum inclinato.

A. hirsutissima, Orientalis, cordifolia, Sieberi, Sartoriana, primuliflora

2 Nuculæ valde curvatae sinu basili angusta rostro fere verticaliter deorsum flexo.

A. Pinardi, Kotschyana, Græca, Galilæa, frigida, bracteosa, viscidula

+ + Nuculæ verruculis basi non confluentibus obsitæ.

A. incana, areolata, rostellata, macrophylla.

* Corolla extus hirta.

1. **A. Pulmonaria** (Griseb. Spicil. II, p. 90) præter pubem breviorem setis basi tuberculatis ad foliorum paginam superiorem adpressis aspera. caulinibus rectis brevibus stricte ramosis, foliis rosularum lanceolatis acutiusculis basi attenuatis, caulinis bracteisque anguste lanceolatis calyces superantibus, racemis fructiferis brevibus laxinseculis, calycis fructiferi acretri deflexi laciniis a basi lanceolatâ subulatis, corollæ flavæ vel cœrulecentis extus pilosæ tubo calyce subduplo longiore limbo parvo. acheniis serobiculato-rugosis subincurvis rostro horizontali areolâ basili ȝ. *A. nonneiformis* Griseb. loc. cit.

Hab. in herbidis regionis subalpinæ montis Nidja Macedonie supra Vodena 2700'-3000' (Griseb!) et inter Perlepe et Trojazchan (Friedr.). Fl. Jun.

Species binæ a cl. Grisebach propositæ corollæ colore in *A. pulmonariâ* flavo. in *A. nonneiformi* cœrulecente tantum differre videntur. Planta vix semipedalis. species affinis *A. Græca* (colore corollæ quoque varianti) sed distincta corollâ pilosâ, corollæ tubo minuto, nuculis parum curvatis. *A. tubulosa* corollâ pilosâ quoque donata distat radice bienni, indumento cinereo, nuculâ curvatissimâ.

2. **A. Scardica** (Griseb Spic. II, p. 91) pilis tenuibus e tuberculo ortis ad foliorum paginam superiorem adpressis hirsuta, caulinibus pumilis erectis crebre foliatis apice 2-3-sidis, foliis lanceolatis inferioribus acutiusculis basi attenuatis caulinis sensim decrescentibus sessilibus. bracteis anguste linearí-lanceolatis calyces superantibus, racemis etiam fructiferis brevibus densis. calycis fructiferi parum ancti erecto-patuli laciniis lanceolatis. corollæ cyaneæ hirtæ tubo calyci aequilongo limbo duplo longiore. nuculis ruguloso-subtuberulatis incurvis rostro deorsum flexo areolâ versus sinum basilarem sitâ ȝ.

Hab. in Macedoniae regione alpinâ Scardi et Bertisci 5-7000 (Griseb!).

Palmaris, folia inferiora sesquipollicaria 3-4 lin. lata, caulina multo breviora. Affinis *A. pulmonaria* a qua floribus cyaneis, præsertim racemis brevissimis fructiferis densis et nuculis magis incurvis differt.

Ar. Geogr. Montenegro.

3. A. tubulosa (Boiss. Diagn. Ser. I, 4, p. 48) pube brevi setisque basi valde tuberculatis obsita aspera cinereo-virens, caule humili crassiusculo a basi stricte ramoso paniculato. foliis oblongo-vel linear-lanceolatis inferioribus basi attenuatis superioribus et bracteis linearibus calyces superantibus, racemis fructiferis densiusculis, calycis fructiferi valde accreti deflexi laciniis lanceolatis obtusis, corollæ luteæ hirtæ tubo calyce duplo longiore in limbum parvum subdilatato, nuculis magnis tuberculatis confluentibus reticulatis curvatissimis sinu basilari angusto, rostro verticaliter descendantem, areola basilari sessili ②.

Hab. in sylvaticis regionis montanae Cariæ inter Jenidje et Gheyra et in monte Cadmo inter Gheyra et Denisleh (Boiss!), ad Sabounjou Kaivé inter Smyrnam et Magnesiam (Bal!). Fl. Jun.

Planta 6-10-pollicaris, caulis unicus vel plures e collo saepius stricte fere a basi ramosi. Species radice bienni verticali simplici, calycibus fructiferis valde accretis laciniis elongatis, corollâ tubulosâ, nuculis magnis insignis.

* * Corolla extus glabra.

+ Nuculae reticulatae vel scrobiculato-rugosæ.

4. A. macrosiphon (Boiss. et Heldr. Diagn. Ser. I, 11, p. 120) tota præter pubem brevem viscidulam setis patulis basi tuberculatis strigulosa, caulis ascendentibus vel prostratis, foliis obtusis radicalibus linear-lanceolatis basi attenuatis caulinis sessilibus abbreviatis, bracteis calyces non superantibus, racemis fructiferis elongatis laxis, calycis fructiferi accreti elongati aucti laciniis lanceolatis obtusis, corollæ purpureæ glabræ tubo calyce plus duplo longiore in limbum parvum subdilatato, nuculis majusculis profunde scrobiculato-reticulatis valde incurvis sinu basilari angusto, rostro verticaliter descendantem, areola basilari valde stipitata ȝ.

Hab. ad Alaya Pamphyliæ in rupestribus (Heldr!). Fl. medio April.

Planta cum racemis demum semipedalibus pedalis et longior. Species corollis tubulosis elongatis, nuculis valde curvatis *A. tubulosa* affinis sed distincta indumento strigoso, corollâ glabrâ purpureâ, nuculae areola stipitata

5. A. strigosa (Boiss. Diagn. Ser. I, 4, p. 46) tota setis basi tuberculatis patulis rigidis strigosa canescens a basi suffrutescenti multicaulis, caulis erectis crassiusculis saepè ramosis, foliis linearibus abbreviatis sessilibus nervo medio subtus prominente inferioribus basi subattenuatis superioribus semiamplexicaulis. racemis fructiferis tandem laxis, bracteis calyce sublongioribus, calycis strigo-

sissimi fructiferi parum accreti laciniis lanceolatis acutis, corollæ azureæ glabrae tubo calyce subduplo longiore limbo ample, nuculis crebre et elevatim reticulatis non tuberculatis incurvis rostro obtuso elongato deorsum subcurvato, areolâ sessili triangulari prope cavitatem curvaturæ sitâ ♀.

Hab. in collibus siccis Palestine ad Hierosolymam (Auch. 2364! Boiss!), in cultis derelictis ad Merré prope Damascus et Antilibano prope Dimar (Gailli!), inter Damascum et Zebdani (Boiss!).

Caules crassi rigidi 6-9-pollicares paniculato-corymbosi, folia inferiora 1-4 $\frac{1}{2}$ -pollicaria 2-3 lineas lata, nuculae transverse $1\frac{3}{4}$ lineas late. Species rhizomate suffrutescenti et indumento valde strigoso insignis.

6. A. Syriaca (Boiss. et Hohen. in Ky. Pl. Alep. Kurd. sub *Campylotropis*) pube minutâ viscidulâ setisque patulis basi tuberculatis obsita, a basi suffrutescenti pluricaulis caulis erectis vel ascendentibus saepè ramosis paniculato-corymbosis, foliis inferioribus linearibus acutiuseulis, superioribus et floralibus calyces superantibus a basi latiore semiamplexicanli lanceolatis acutis, racemis fructiferis laxis, calycis demum accreti et deflexi strigosi laciniis anguste linearilanceolatis, corollæ caeruleæ glabrae tubo calyce vix longiore limbo ample, nuculis crebre et elevatim reticulatis non tuberculatis subincurvis rostro obtuso elongato deorsum subcurvato, areola stipitata prope curvaturam medium sitâ ♀. *A. Aucheriana* D.C. Prodr. X, p. 102 (ex parte quoad plantam montis Akdaghi Cataoniae!).

Hab. ad Aleppo Syriae in collibus aridis gregatim (Ky. 40! Haussk!), in monte Akdaghi Cataoniae (Auch. 2358!). Fl. Marte.

An ab *A. strigosa* quâcum eam olim combinavi sat distincta sit ulterius observandum; non strigosa, folia superiora magis dilatata, calycis laciniæ angustiores, corollæ tubis brevier, nuculae sculpturâ similes sed areola (an constanter?) stipitata.

7. A. megacarpa (Alph. D.C. Prodr. X, p. 101) præter tomentum breve pilis longis tenuibus hispida canescens vel cana, caule a collo crasso vaginato sæpius unico erecto vel ascendentí crasso fere a basi ramoso, foliis inferioribus et rosulae sterilis lanceolatis elongatis basi attenuatis, floralibus abbreviatis calyces superantibus, racemis demum laxiuseulis, calycis hispidissimi demum accreti deflexi laciniis late lanceolatis, corollæ caeruleæ tubo calyce sesqui-longiore limbo ample, nuculis magnis valde serobiculatis et obtuse tuberculatis valde curvatis sinu angustissimo rostro deorsum curvato, areolâ sessili basilari ♀. *A. Cappadocica* Boiss. et Bal. Diagn. Ser. II, 6, p. 123.

Hab. in Cappadociâ ad Euphratem (Auch. 2275!), in monte Alidagh Cappadociae (Bal.), ad Gormise (Ky. Suppl. 203! et 204!), ad Bakker Maaden Armeniae australis (Noë. 858!).

Semipedalis vel paulo elatior, indumentum fere *A. Syriaca* sed longius magis canum, collum non suffrutescens sed caulem fere unicum et rosulam sterili ebulens, nuculae fere $2\frac{1}{2}$ lin. longæ valde nec parum curvatae serobiculato-tuberculatae nec reticulatae.

8. **A. tinctoria** (L. Sp. Ed. I, p. 432 sub *Lithospermo*) breviter velutina prætereaque setis ad ramos patulis ad folia adpressiusculis basi saepe tuberculatis plus minus strigosa cinerea vel canescens, caulis numerosis ascendentibus vel prostratis, foliis lanceolatis obtusis superioribus basi subdilatata semiamplexicaulibus, racemis demum elongatis, bracteis calyces fructiferis deflexos aceretos superantibus, calycis lacinias lanceolato-linearibus acutis, corolla cœrulea extus glabrae tubo ealycem æquanti vel paulo longiore muculis tuberculatis basi confluentibus reticulatum asperis valde incurvis, areola parva substipitata basilari, rostro deorsum incurvo ♀. *A. tinctoria* Tausch Flor. 1824, p. 234. — *A. Matthiolii* Tausch loc. cit. — *Anchusa tinctoria* Desf. Atl. I, p. 156. Fl. Græc. tab. 166. — *Anch. tuberculata* Forsk. Eg. p. 51. — *Anch. rhizochroa* Viv. Dec. N° 20. — *Buglossum tinctorium* Lam. — *Lith. Lehmanni* Tineo in Guss. Syn. II, p. 791.

Hab. in arenosis maritimis, collibus calcareis Peloponnesi (Bory! Sprunt!), Attice (Spr! Heldr! Boiss!), Macedonia (Griseb.), Cycladum (Urv., Weiss!), Cretæ (Heldr!), Rhodi (Bourg!), Bithynia (Griseb.), Lydiae (Fleisch!), Lyciae et Pamphyliae (Bourg!), Cilicie (Bal!), Palestine ad Gaza (Boiss!), Egypti ad Alexandriam (Forsk., Ehrenb! Samar! Gaill! etc.).

Specimina Egyptiaca nonnulla corollis ex secco flavidis occurunt.

Ar. Geogr. Hispania, Gallia australis, Italia, Sicilia, Dalmatia, Hungaria, Africa borealis.

9. **A. hirsutissima** (Alph. n.c. Prodr. X, p. 101) præter pubem brevem viscidulam setis albis patulis tuberculatis hispida canescens a basi suffrutescenti pluricaulis, caulis brevibus ascendentibus rigidulis parce ramosis, foliis anguste linear-lanceolatis acutis floralibus acutissimis calyces superantibus, racemis fructiferis elongatis rectis densiusculis, calycis fructiferi acereti deflexi lacinias lanceolatis longe attenuatis, corolla glabrae luteæ vel dilute griseæ tubo ealyce sublongiore limbo ampliuscule, muculis magnis cebre et elevatim reticulatis subincurvis rostro superne subcarinato horizontali-subdeflexo, areola sessili prope medianam curvaturam sita ♀. *Lithosp. hirsutissimum* Bertol. Miscell. I, p. 14.

Hab. in Syriâ boreali et Mesopotamiâ inter Aintab, Arkûk, Bir, Orfa (Haussk! Ky exs. 20!), in viâ ad Bagdad (Rousseau!), in deserto ad Iherium Chabur (Haussk!), ad Sedjim Kala prope Euphratem (Chesney 75!), ad montem Sindjar in cretaceis (Haussk!).

Affinis A. *Syriace* differt foliis præsertim floralibus angustioribus acutioribus, colore florum, calycis fructiferi amplioris lacinias longioribus. Mucularum reticulus magis irregularis. Flores odorem gratissimum spirant (Haussknecht).

10. **A. Orientalis** (L. Syst. 156 sub *Lithospermo*) glandulosopubesca viscida insuper setis brevibus tuberculatis ad paginam superiorum sparsis obsita, caulis elatis erectis vel ascendentibus, foliis inferioribus oblongo-lanceolatis inferne longe attenuatis plerumque dentatis et undulatis, intermediis semiamplexicaulibus, bracteis

a basi rotundatâ vel subcordatâ ovatis obtusis calyces superantibus, racemis fructiferis densiusculis distichis, calycis laciniis lanceolatis acutis fructiferi modice accreti deflexi conniventibus, corollæ glabrae luteæ tubo calyce duplo longiore limbo majuscule, nuculis parvis scrobiculato-reticulatis interdum subtuberculatis parum incurvis rostro subdescendente, areolâ subbasilari ♀. Ic. Fl. Græc. tab. 160, Bot. Mag. tab. 515. — *Alkanna Orientalis* Boiss. Diagn. Ser. I, 4, p. 46. — *Anchusa Orientalis* L. Sp. 191. — *Campnocarpus Orientalis* C. Koch Linn. XVII, p. 304. — *Lycopsis? mollis* Visian. Ill. p. 14, tab. 6 (ex Icone et loco). — *Lith. Pythicum* Clarke in Spr. Neu Entd. III. p. 165? (ex descriptione).

Hab. in Archipelago (Sibth.), Cariâ (Tchihi!), Ciliciâ Tracheâ (Pér!), Lycaoniæ monte Karadagh (Heldr!), Galatiâ et Paphlagoniâ (Wied!), Cappadociâ ad Argæum 6000' (Bal!), Syriâ ad Damascum (Gaill!), in Cælesyriâ et Antilibano 5000' (Post! Ky!), Libano ad Cedros (Boiss!), ad Rascheya (Boiss!), in monte Sinai (Ehrenb! Schimp! Boiss!), Armeniâ Turcicâ ad Van (Noë!), Erzelen (Huet!). Gumuschkhane (Bourg!), Armeniâ Rossicâ (Buhsel!), Persiæ prov. Aderbidjan (Szov! Bgel!), Arabice *Lebett*.

B. Hellenica. — Bractæ breviter lanceolatæ acutæ, racemi fructiferi laxiores. Nuculæ magis incurvæ rostro deorsum subverticaliter flexo.

Hab. in Græciæ monte Malevos Laconiæ (Sprun!), Acrocorinthus (Boiss!), Messenia (Bory), Ætolia prope Missolonghi (Nieder!).

Pedalis et sesquipedalis, indumento glanduloso-viscido et foliis sœpe undulatis insignis.

11. **A. cordifolia** (C. Koch Linn. XXIII, p. 640) eglandulosa setis e tuberculo ortis ad foliorum paginam superiori adpressis aspera, caulis erectis, foliis integris acutis rosularum lanceolatis elongatis utrinque attenuatis, caulinis brevibus cordatis oblongis, bracteis cordato-triangularibus acutissimis imbricatis, racemis fructiferis dense et distiche bracteatis, calycis fructiferi deflexi accreti laciniis lanceolatis acutis, corollæ luteæ tubo calycem parum superante limbo ampio tubo æquilongo, nuculis mediocribus tenuiter areolato-reticulatis curvatis rostro brevi deorsum subflexo, areolâ basilari ♀.

Hab. in Ponti Alpibus 5500' (C. Koch!), in incultis circa Djimil Ponti Lazici 6000' (Bal. pl. exs. sub *A. Orientali*!).

Facie, caulis pedalibus, etc., affinis *A. Orientali* sed certe distincta indumento aspero nullo modo viscido, foliis rosularum integerim angustioribus (4-6 pollic. 5-6 lin. latis), bracteis cordato-triangularibus acutissimis imbricatim distichis, corollæ tubo multo ampliore, nuculis majoribus reticulo tenui areolatis.

12. **A. Sieberi** (Alph. D.C. Prodr. X, p. 99) præter pubem brevissimam patule et crebre albo-setulosa, caulis e rhizomate verticali pluribus prostratis abbreviatis in racemos simplices vel binos fructiferos elongatos laxiusculos abeuntibus, foliis parvis oblongis

radicalibus basi attenuatis, caulinis oblongo-linearibus obtusis basi semiamplexicaulibus, bracteis calyces non superantibus, calycis fructiferi deflexi parum acereti laciniis breviter lanceolatis obtusiusculis, corollæ glabrae tubo fusco calyce duplo longiore in limbum luteum eo breviorem sensim ampliato, nuculis minutis tuberculatis sparsis inaequalibus basi confluentibus obsitis modice curvatis rostro horizontali, areolâ basilari ♀.

Hab. in prom. Maleca Cretæ (Sieb!), in arenosis maritimis peninsulae Akrotiri et prope Frankocastro (Heldr!). Fl. Mart. Apr.

Caules cum racemis 4-6-pollicares tenues; ab *A. tinctoria* indumento magis patent, foliis abbreviatis, bracteis calyces non superantibus, corollæ tubo elongato, nuculae minus incurvæ rostro horizontali nec deorsum flexo, etc., descendens.

13. A. Sartoriana (Boiss. et Heldr. Diagn. Ser. II, 3, p. 134) præter indumentum breve tota setis sparsis basi tuberculatis patentibus strigulosa, caulis procumbentibus ramosis ramis fructiferis tandem valde elongatis laxiusculis, foliis inferioribus subrepandis oblongis obtusis in petiolum attenuatis superioribus bracteisque calyces superantibus a basi rotundato-cuneatâ breviter oblongo-lanceolatis obtusis, calycis fructiferi subaucti non deflexi laciniis lanceolatis, corollæ glabrae luteæ tubo calycem non superante, nuculis parvis tuberculatis in rugas subquinas concentricas confluentibus obsitis serobiculatis rostro carinato deorsum subflexo, areolâ basilari substipitatâ ♀.

Hab. in arenosis ad viam inter Naupliam et Porto Tolon Argolidis (Heldr!).

Affinis *A. Graecæ* a quâ differt caulis prostratis, pilis strigosis, racemis tandem longissimis, foliis repandis, corollæ tubo brevi, nuculis non uniformiter reticulatis et minus curvatis.

14. A. primuliflora (Griseb. Spic. II, p. 89) glanduloso-pubescentis pilisque longioribus e tuberculo sœpe ortis sparsis hispida, caulis erectis pumilis superne bifidis, foliis parvis linear-lanceolatis inferioribus obtusis superioribus et bracteis calyces superantibus a basi rotundata lanceolatis acutis, racemis fructiferis laxiusculis, calycis fructiferi aucti patent-deflexi laciniis lanceolato-attenuatis, corollæ glabrae luteæ tubo calyci æquilongo limbo ampio patulo, nuculis parvis tuberculatis obtusis subconfluentibus obsitis curvatis rostro horizontali deorsum subflexo, areolâ basilari substipitatâ ♀ vel ♂. *Lithosp. apulum* Friv. pl. exs. non L.

Hab. in Thraciâ boreali prope Philipopolin (Friv!).

Planta 4-6-pollicaris foliis parvis inferioribus 6-9 lineas tantum longis 2 lin. latit. hæc notâ indumento viscido, nuculis minimis parum curvatis non serobiculatis ab *A. Graecâ* distincta.

15. A. Pinardi (Boiss. Diagn. Ser. I, 4, p. 48) tota setis patulis basi sœpius tuberculatis hispida flavo-virens, caulis pumilis ascen-

dentibus superne 2-3-ramosis, foliis inferioribus oblongo-lanceolatis obtusis basi attenuatis, superioribus basi subdilatata semiamplexicaulibus, bracteis calyces fructiferos vix superantibus, calycis hispido fructiferi aucti deflexi laciniis lanceolatis obtusis, corollæ glabré luteæ tubo calyci æquilongo limbo ampio patulo, nuculis serobiculato-tuberculatis valde incurvis rostro deorsum incurvo, areolâ substipitatâ basilari ♀. *A. filuginoides* Alph. D.C. Prodr. X, p. 98.

Hab. in Cariâ (Pinard!), in agris incultis ad Adalia Pamphyliæ (Heldr!), in monte Tauro (Auch!). Fl. Mart. Mai.

Valle affinis *A. primuliflora* sed non glandulosa et indumentum magis asperum, bracteæ breviores, nuculæ majores serobiculato-rugulosæ valde incurvæ.

16. A. Kotschyana (D.C. Prodr. X, p. 98) canescens breviter asperulo-hirta setis brevibus basi tuberculatis intermixtis, caulis infra rosulam ascendentibus elongatis apice bifidis vel parce ramosis, foliis obtusis inferioribus oblongo-spathulatis basi longe attenuatis intermediis et floralibus basi subcordatâ semiamplexicaulibus, bracteis linearibus obtusis calyces multo superantibus, racemis fructiferis densiusculis distichis, calyce tomentello fructifero deflexo subacereto laciniis lanceolatis apice longe attenuatis, corollâ glabrâ luteâ tubo fuscescens-rubello calyce subduplo longiore limbo parvo, nuculis majusculis serobiculato-tuberculatis valde curvatis rostro deorsum directo, sinu angusto, areolâ basilari substipitatâ ♀ vel ♂. *A. amplexicaulis* D.C. l. cit. p. 101! — *A. bicolor* Boiss. et Ky. in Ky. exs. 1859. — *A. brachysolen* Boiss. MSS.

Hab. in Tauro (Ky. 361!), ad Gülek Maaden 6000' (Ky!), fauce Guzelderé Ciliciæ (Bal!), prope Messis et in ditione Kassan Oglu (Ky. 40! et Suppl. 47!), Syriâ et Cappadociâ (Auch!), Mesopotamiâ ad Terek (Haussk!).

Fere pedalis. Species bracteis obtusis, corollæ tubo elongato et limbo minuto ab *A. Græca* distineta.

17. A. Græca (Boiss. et Sprun. Diagn. Ser. I, 4, p. 47) pilis adpressis tuberculo insidentibus aspera, caulis rectis inter pilos pube viscidulâ obsitis bifidis, foliis rosularum oblongo-lanceolatis acutiusculis basi attenuatis, caulinis et bracteis a basi rotundatâ vel semiamplexicauli-subcordatâ breviter lanceolatis acutis hisce calyces superantibus, racemis fructiferis laxiusculis, calyceis fructiferi accreti et deflexi laciniis lanceolatis, corollæ glabré tubo calyce longiore fuso in laminam aurantiacam amplam dilatato, nuculis serobiculato-reticulatis curvatissimis rostro carinato-ancipiti deorsum curvato sinu angustissimo, areolâ basilari stipitatâ ♀. *Nonnea lutea* Fl. Pelop.! non D.C.

Hab. in saxosis montanis Argolidis et Laconiae (Boiss! Heldr!), Messeniæ (Pl. Pélop!), Laconiae in monte Malevo (Orph!). Fl. Apr. Mai.

β. hispidior. — Inflorescentia magis albo-setosa. — *A. Bæotica* D.C. Prodr. X, p. 98!

Hab. in monte Parnes Atticae (Orph!), Delphi Eubeæ (Sprun! Orph! Heldr!), Parnasso infer. (Guicci!), Bœtiâ (Auch. 2275!), monte Citheron (Boiss! Heldr. Herb. Norm. I.).

γ. versicolor. — Corolla cœruleo-lutescens.

Hab. in monte Malevo cum typo (Orph!).

Species ab *A. Orientali* egregie distincta indumento aspero, caulis humilioribus, foliis angustioribus non undulato-dentatis, corollæ tubo fusco, nuculis.

18. **A. Galilæa** (Boiss. Diagn. Ser. I, 11, p. 118) breviter velutina prætereaque setis tuberculatis patulis secus ramos margines foliorum et in parte superiori plantæ copiosioribus aspera, caulis ascendentè-procumbentibus elongatis, foliis acutiusculis inferioribus oblongo-lanceolatis basi breviter attenuatis cæteris bracteisque calyces multo superantibus oblongis et ovatis basi cordato-subauriculatis, racemis fructiferis longis laxissimis foliosis, calycis elongati fructiferi deflexi et aucti laciniis lanceolatis non conniventibus, corollæ flavourantiacæ glabrae tubo calyce vix longiore limbo ampio, nuculis maximis valde sericeo-fuso-reticulatis valde incurvis rostro deorsum flexo, areolâ subbasilari ♀.

Hab. in agris derelictis Galileæ in descensu ad urbem Tiberiadem (Boiss!). Fl. Mai.

Caules pedales et longiores fere a basi tandem laxissime racemiteri. Insignis bracteis magnis latis dissitis fere pollicem longis, calyce eo congenerum longiori (florifero 6-7 lineas longo), nuculis pisum parvum aequalibus.

19. **A. frigida** (Boiss. Diagn. Ser. I, 7, p. 32) pallide virens tota pube brevissimâ setisque brevibus sparsis basi tuberculatis aspera, caulis e rhizomate verticali pluribus erectis pumilis folia radicalia non vel vix superantibus in racemum simplicem vel bifidum fructiferum laxum brevem abeuntibus, foliis radicalibus oblongis acutiusculis basi longe attenuatis, caulinis bracteisque calyces longe superantibus linear-lanceolatis obtusiusculis semiamplexicaulibus, calycis fructiferi modice accreti pedicellati deflexi laciniis anguste lanceolatis, corollæ glabrae brunneo-luteæ tubo calyce $2\frac{1}{2}$ -plo longiore limbo majusculo, nuculis tuberculatis acutis in cristulas basi connatis crebre echinatis valde incurvis rostro deorsum directo, areolâ basilari ♀.

Hab. ad nives regionis alpinae Persie austro-occidentalis, Kuhdaëna (Ky. 573!), montes Sawers, Nur, Eschker, Kellal et alii districtus Kuh Kilouyeh et montes Avroman et Schahu alt. 9000'-13000' (Haussk!), mentibus Bachtiaricis (Loftus!), monte Elbrus Persie bor. (Ky!).

Caules cum racemis fructiferis 4-6-pollicares, folia radicalia saepe caulis aequilonga. Variant 6-45 lineas lata, corolla 5 lineas longa. Species indumento brevi aspero et nuculis acute tuberculato-cristatis insignis.

20. **A. bracteosa** (Boiss. Diagn. Ser. I, 11, p. 118) viridis setis brevibus basi regulariter tuberculatis asperrima, caulis e rhizomate crasso squamoso brevibus fere a basi bifidis, foliis radicalibus lanceo-

latis acutis basi longe attenuatis caules æquantibus, bracteis elliptico-lanceolatis amplis calyces multo superantibus, racemis fructiferis laxis, calycis strigoso-hirsuti fructiferi accreti deflexi laciniis lanceolatis, corollæ aurantiacæ glabræ tubo calyce sesquialiore limbo ampio, nuculis magnis tuberculis minutis in rugulas plus minusve confluentibus scabridis rostro longo deorsum curvato, areolâ magnâ basilari ♀.

Hab. in declivibus orientalibus faucis Dudera montis Elbrus Persiae borealis (Ky. 269!), in valle Lar (Buhse!).

Folia radicalia cum petiolo 3-5 poll. longa, caules cum racemis 5-6-pollicares, bracteæ 4-4 ½-pollicares 4-5 lineas latæ, flores magn. *A. Orientalis*. Affinis *A. frigida* a quâ nuculis minute scabridis nec elevatim tuberculato-cristatis eximie differt.

21. A. viscidula (Boiss. Diagn. Ser. I, 11, p. 119) tota molliter tomentella viscidula grisea, caulis ascendentibus fragillimus, foliis inferioribus oblongo-lanceolatis basi attenuatis, superioribus et bracteis ovato-oblongis obtusis calyces fructiferos superantibus, racemis fructiferis laxis, calycis pedicellati tandem deflexi laciniis a basi latiore linear-setaceis, corollæ glabræ flavæ tubo calyce parum longiore limbo parvo, nuculis tuberculis basi confluentibus asperis et scrobiculatis valde curvatis rostro deorsum incurvo, areolâ basilari stipitata ♀. *Nonnea pulla* γ *mollis* Alph. d.c. Prodr. X, p. 32.

Hab. in monte Akdaghi Cataoniae (Auch. 2414!).

Caulis fragiles et indumentum molle *A. incana* et *A. macrophylla* sed viscidum et ab eis insuper colore florum et nuculâ non sparsim verruculosâ sed crebre scrobiculato-tuberculatâ discedens.

+ + Nuculæ verruculosæ.

22. A. incana (Boiss. Diagn. Ser. I, 4, p. 45) tomento brevi denso pilisque longioribus omnibus adpressis molliter cana, caulis pumilis gracilibus, foliis inferioribus lanceolatis basi attenuatis, caulinis et bracteis a basi rotundatâ anguste oblongis hisce calyces fructiferos patulos parum accretos superantibus, racemis fructiferis densis, calycis dense sericei laciniis lanceolatis acutis, corollæ glabræ cæruleæ tubo calyce duplo longiore limbo magno expanso, nuculis parvis tuberculis basi distinctis conicis exasperatis valde incurvis areolâ oblongâ sinum basilarem fere totum occupante, rostro deorsum incurvo ♀.

Hab. in fissuris rupium Cariæ (Pinard!), Pisidiæ ad Tcheltikchi (Heldr!). Fl. Mai.

Planta semipedalis vel vix longior, corollæ magn. *Lith. purpureo-cerulei*. Ab *A. tectoria* indumento cano molli, caulis fragilibus, calyce fructifero et nuculis duplo minoribus hisce conico-verrucosis et areolâ basilari versus rostrum dilatata distincta.

23. A. areolata (Boiss. Diagn. Ser. I, 4, p. 45) virens tota

præter pubem brevem pilis rigidis basi tuberculatis aspera, caulinibus humilibus ascendentibus vel decumbentibus superne parce ramosis, foliis lanceolatis subrepandis inferioribus obtusis basi attenuatis superioribus bracteisque longioribus sessilibus sensim diminutis acutis, racemis fructiferis tandem laxiusculis, calycis asperi laciniis a basi latiori linearisubulatis fructiferis patulo-deflexis, corollæ glabrae cœruleæ tubo calyce sesquilongiore, nuculis parvis verruculis minimis sphæricis nitidis obsitis valde incurvis rostro deorsum spectante, areolâ triangulari-dilatata fere totum sinum occupante carinâ elevatâ versus medium percursâ ♀. *A. azurea* Boiss. in Bourg. exs.

Hab. in rupestribus umbrosis Cariæ in monte Cadmo supra Colossam (Boiss!), Lyciae in montibus Elmali (Bourg!), Sipyli supra Magnesiam (Bal!). Fl. Jun.

Affinis *A. incanæ* a quâ differt indumento aspero, calycis laciniis linearisetaceis, nucularum verrucis minimis globosis, etc.

24. A. rostellata (Boiss. Diagn. Ser. I, 11, p. 119) setis adpressis nonnullis basi tuberculatis dense cana basi suffrutescens ramis pumiliis, foliis omnibus acutis breviter lanceolatis inferioribus et ramorum sterilium basi attenuatis floralibus et bracteis sessilibus linearis-lanceolatis, racemis demum elongatis densiusculis, calycis fructiferi patent-deflexi laciniis lanceolatis acutis, corollæ flavæ glabrae* tubo calyce subdupo longiore limbo parvo, nuculis verrucis parvis globosis sparsis obsito angulo recto curvato rostro attenuato horizontali acutiusculo, areolâ basilari marginatâ ovatâ ♀. *A. Aucheriana* d.c. Prodr. ex parte.

Hab. in Tauro (Auch. 2278 bis!), in fauce Guzeldere supra Sedichig Ciliciae (Bal!).

Planta 4-6-pollicaris, folia inferiora $1\frac{1}{2}$ -2 pollices longa 4-5 lineas lata indumentum canum *A. incanæ* sed non molle, corollæ luteæ, nuculae rostrum elongatum horizontale nec deorsum curvatum, etc.

25. A. macrophylla (Boiss. et Heldr. Diagn. Ser. I, 11, p. 120), caulinibus ascendentibus superne ramosis fragillimis elongatis ciliato-villosis, foliis amplis breviter velutinis et insuper setis tuberculatis sparsis obsitis teneris magnis inferioribus oblongo-lanceolatis basi attenuatis, superioribus bracteisque calyce longioribus cordato-oblongis obtusiusculis, racemis fructiferis elongatis densiusculis, calycis fructiferi nutantis parum aucti laciniis linearis-lanceolatis acutis, corollæ azureæ tubo calyce duplo longiore intus 5 fasciis purpureis percurso. nuculis parvis verruculis minimis conicis parce obsitis incurvis dorso subcarinatis rostro subhorizontali. areolâ ovatâ dilatata fere sinum totum occupante ♀.

Hab. in rupestribus maritimis ad Adalia Pamphyliæ (Pestal! Bourg. 197!), ad muros amphitheatri Perge (Heldr!), in Libano superiore (Gaill. 421 fide specim. cæteram nimis incompletæ).

Caules sœpe pedales, folia inferiora $2\frac{1}{2}$ -4 pollices longa, corolla speciosa

sex lineas longa. Nuculis *A. incanæ* et *areolatae* affinis differt indumento cinea-
reo-argyreo molli setulis intermixto, foliis magnis floralibus cordatis.

MYOSOTIS (L. Gen. 480).

Calyx campanulatus quinquesfidus vel partitus. Corolla hypocrate-
rimorpha vel infundibuliformis, fauce (in nostris) fornicibus obtusis
instructa, limbi plani vel concavi lobis aestivatione contortis. Stamina
inclusa filamentis brevibus, antheræ oblongæ medio dorso affixæ
apiculo membranaceo vel calloso. Stigma obtusum subbilobum.
Nuculæ levissimæ ovatae vel oblongæ subcompressæ, areolæ basilari-
sublateralis nigræ subelevatus. — Herbæ perennes vel monocarpicæ.

Specierum Orientalium distributio.

SECT. I. EUMYOSOTIS Alph. D.C. — Flores racemoso-scorpioidei.
Nucularum areola ecarunculata.

* Calycis pili adpressissimi.

M. palustris, cæspitosa, Sicula, littoralis, Idæa, pusilla.

* * Calycis pili patuli baseos uncinati.

+ Perentes vel biennes.

M. sylvatica, platyphylla, alpestris, Olympica, intermedia.

+ + Annuæ.

M. hispida, stricta, uncata, refracta.

SECT. II. STROPHIOSTOMA Endl. — Flores racemosi. Nucularum areola
ad centrum carunculâ albâ prominenti obsita.

M. sparsiflora, propinqua, amœna.

SECT. III. PHYLLOCEPHALUM. — Flores in capitula cymis repetitive
dichotomis constantia congesti. Nucularum areola centro
carunculâ albâ prominenti obsita.

M. involucrata.

SECT. I. EUMYOSOTIS.

1. ***M. palustris*** (With. Arr. Brit. Pl. II, p. 225) perennis, pilis
totius plantæ adpressis in parte inferiori caulis patulis, rhizomate

interdum stolonifero, caule angulato foliis oblongo-lanceolatis inferioribus in petiolum attenuatis superioribus acutiusculis, racemis demum elongatis laxis, calyce adpresso piloso ad tertiam partem fisso fructifer campanulato patent, corollæ cœruleæ vel albae limbo plano tubo longiore, stylo calyci æquilongo, pedicellis fructiferis horizontalibus, carpellis ovatis lucidis ♀. Ic. Rehb. tab. 119. — Engl. Bot. tab. 1973. — *M. scorpioides* var. *palustris* L. — *M. perennis* var. L. D.C.

Hab. ad fossas, ripas, paludes, Thracia (Friv! Griseb.), Armenia Turcica ad Erzerum (Calvt!), Caucasus, Iberia et Imeretia (Ledeb.), Affghaniae alpes (Griff!) et prob. alibi.

Ar. Geogr. Europa tota a Scandinaviâ ad Rossiam et Lusitaniam ad Dalmatiam, Sibiria et Dahuria, Madera.

2. ***M. cæspitosa*** (Schultz Fl. Starg. Suppl. II) biennis, tota adpresso hirta, caule tereti, foliis glabriusculis oblongis vel linguiformibus basi attenuatis, racemis demum elongatis laxis basi subfoliatis, pedicellis fructiferis patulo-reflexis inferioribus calyce 2-3-plo longioribus, calyce adpresso hirto semiquinquefido fructifero campanulato aperto, corollæ parvæ cœruleæ limbo plano tubo calyce breviori æquilongo, stylo brevissimo, nuculis late ovatis basi truncatis ⊖. *M. lingulata* Lehm. Asp. p. 110. — Rehb. tab. 120. — *M. uliginosa* Schrad.

Hab. in paludibus et fossis Ponti Lazici ad ostium fluvii Kalopotamos (Balt!), Caucasus (Ledeb.), Abchasiæ et Mingrelia (Nordm.), ad fluv. Hessawi inter Nisibin et montem Sindjar Assyriæ et ad rivulos alpinos montis Sawers Persie australis (Haussk!). Prob. quoque alibi.

E. macrocalyx F. et M. Index III, Petrop. p. 40. — Calyx fructifer major interdum 4 lineas longus. Cum typo formis intermediis conjugitur.

Hab. in Transcaucasiâ prope Lenkoran ad Caspium (C.A. Meyer!).

Ar. Geogr. Europa borealis et media a Scandinaviâ ad Rossiam, Hispaniam et Italiam centralibus ad regionem Danubialem, Sibiria, Madera, Africa boreali, America borealis.

3. ***M. Sicula*** (Guss. Syn. I, p. 214) annua, parce et adpressissime hirta, caule tereti basi sœpe radicante, foliis oblongo-linearibus obtusis, racemis demum laxis basi non foliatis, pedicellis fructiferis patulo-reflexis inferioribus calyce parum longioribus, calyce adpressissime et parce hirto fructifero campanulato-cylindrico dentibus obtusis subconniventibus, corollæ parvæ limbo concavo tubum longitudine non aquante, stylo carpellis æquilongo ⊖. Ic. Rehb. tab. 120. — *M. micrantha* Guss. Prodr. I, p. 207 non Pall.

Hab. in pratis humidis turfosis, Peloponnesus ad Tyrinthum (Orph!), Byzantium (Wied!), Pamphylia ad fluvium Catarrhaicum (Heldr!), Pontus Lazicus ad Djimil (Balt!).

Valde affinis *M. cæspitosa*, ar. notis indicatis sat distincta.

Ar. Geogr. Gallia media, Sardinia, Sicilia, Serbia, Algeria.

4. **M. littoralis** (Stev. Cat. Gorenk. 1812) annua, nana subcaulis ad folia patule secus inflorescentiam adpresso hispida, foliis radicalibus parvis aggregatis ovato-oblongis obtusis, racemis fere radicalibus numerosis etiam fructiferis brevibus imbricatim densifloris secundis incurvis ebracteatis, pedunculis fructiferis incrassatis strictis calyce brevioribus, calyce adpresso hispido fructifero cylindrico laciniis conniventibus, corollæ exiguæ cœruleæ vel albae limbo concavo tubum non æquante ⊖.

Hab. in arenis maritimis Tauriæ meridionalis (Stev!), Bithyniæ ad sinum Nicodemicum prope Hersek (Griseb!), Attice ad Phalerum (Heldr. Herb. Norm. 554! Orph. Fl. Gr. exs. 220! sub *M. pusillæ*).

Plantula 1-2-pollicaris a collo sœpe ramosissima, racemi fructiferi densiflori semipollicem vel ad summum pollicem longi.

5. **M. Idæa** (Boiss. et Heldr. Diagn. Ser. I, 11, p. 121) annua, inferne subpatentim superne adpressiuscule hirta pumila caule brevi ramoso. foliis radicalibus obovato-spathulatis, caulinis paucis elliptico-linearibus, racemis fructiferis elongatis laxis secundis ebracteatis, pedunculis fructiferis plus minus incrassatis erecto-patulis vel patenteflexis calyce aequilongis vel sesquilongioribus, calyce adpresso piloso fructifero anguste campanulato laciniis erectis non conniventibus, corollæ exiguæ cœruleæ limbo concavo limbum non æquante ⊖.

Hab. in littore maris Tauriæ prope Sudak (Rehm!), in monte Apos insulae Chios (Orph!), in saxosis montis Ida et montium Lassiti Cretæ 5-6000' (Heldr!), Hymetto orientali in Attica (Heldr!), Malevo Laconiæ (Orph!), Lyciæ montibus supra Elmalu (Bourg!).

β . *grandiflora* Boiss. herb. — Corollæ cœruleæ vel lacteæ limbus major tubo tamen vix aequilongus. — *M. Kiesenwetteri* Heldr. in litt.

Hab. in Attica in monte Lycabetto, ad lapidicinas Pentelici, ad cacumen Hymetti et montis Parnes Heldr! (Orph. Fl. Gr. 219!), in arenosis et alveis prope Athenas (Heldr. et Orph. Fl. exs. 218! sub *M. Creticæ* non Boiss. et Heldr.), in monte Malevo Laconiæ (Orph!), regione inferiori Parnassi (Guicci!), ad radices montis Delphi Eubeæ et in insula Chios (Orph!).

γ . *Cadmea*. — Corolla cœrulea magna limbo tubum superante, pedicelli fructiferi vix incrassati, valde patentes. — *M. Cadmea* Boiss. loc. cit. p. 122. — *M. Bœotica* Reut. MSS.

Hab. in monte Helicon Bœotiae (Orph!), in monte Cadmo Cariæ supra Colossas (Boiss!), prope Byzantium (Wied!).

Planta 2-6-pollicaris a *M. littorali* pilis brevioribus, racemis fructiferis laxis, pedicellis longioribus magis patentibus, calyce fructifero non clauso; a *M. pusillæ* racemis ebracteatis discedens. An tamen omnes ad speciem polymorpham reducendæ? Var. β et γ sistunt formas corollæ majori, variationes parallelæ in aliis speciebus generis observantur.

6. **M. pusilla** (Lois. in Desv. Journ. II, p. 260, tab. 8) annua, pilis erecto-patulis ad inflorescentiam adpresso hispida pumila a collo brevissime ramosa, foliis radicalibus obovatis rameis paucis ellipticis obtusis, racemis fructiferis laxis parte inferiore bracteatis, pedicellis

fructiferis vix incrassatis calyci æquilongis vel sublongioribus erecto-patulis, calyce adpresso hirti fructifero campanulato laciniis erectis, corollæ exiguae cœruleæ vel albae limbo concavo tubum non æquante ♂. Ic. Mor. Fl. Sard. tab. 96 Opt. — *M. Cretica* Boiss. et Heldr. loc. cit. p. 121. — *M. Enesia* Heldr. MSS. — *M. incrassata* Guss. Syn. I, p. 214.

Hab. in regione abietinâ montis Ænos Cephaloniæ (Heldr!), in monte Idæ Cretæ 5000'-6000' (Heldr!), monte Troodos Cyprî (Ky. exs. sub *M. Idæa*!).

A præcedentibus racemis bracteatis vix secundis et calyce fructifero breviori distincta. Cl. Gussone erronee racemos ebracteatos dicit.

Ar. Geogr. Gallia australis, Corsicæ, Sardiniae, Siciliæ montes et regio littoralis, Dalmatia, Algeria interior.

7. ***M. sylvatica*** (Hoffm. Deutsch. Fl. I, p. 85) biennis, pluri-caulis, caulinis sæpius elatis inferne hispidis parce ramosis, foliis molliter hirsutis radicalibus oblongo-ovatis in petiolum longum sensim vel abruptiusculæ attenuatis, caulinis oblongo-lanceolatis acutiusculis, racemis sæpius bifidis fructiferis laxis ebracteatis, pedicellis fructiferis patentibus calycem æquantibus vel eo duplo longioribus adpresso hirtis, calycis tubo pilis patulis vel reflexis plerisque uncinatis obsito laciniis fructiferis erectis non conniventibus, corollæ intense cœruleæ rarius albae tubo calycem æquante limbo magno plano, nuculis ovatis submarginatis alterâ facie obscure marginatis ♂. Ic. Rchb. tab. 121. — *M. perennis* β *sylvatica* D.C. — *M. montana* Besser Galic. № 491. — *M. cyanea* Reut. MSS. — *M. lithospermifolia* Guss. Syn. I, p. 213. — *M. densiflora* et *M. Pontica* C. Koch Linn. XXIII, p. 641. — *M. amena* Schott Österr. Bot. Woch. 1857, p. 206.

Hab. in sylvaticis et umbrosis humidis regionis montanæ et subalpinæ fere ubique, montes totius Græciae a Taygeto, Parnes Atticæ, etc., ad Olympum Thessalum (Heldr! Orph!), Macedoniæ et Thraciæ (Griseb.), totius Anatoliæ ab Olympo Bithyno (Boiss!), ad montes Lycie (Bourg. exs. sub *M. alpestri*!), Ciliciae (Ky! Balt!), ad montes Ponti Lazici (Bal. sub *M. alpestri*!), Armeniae Turcicæ (Ky!), totius Caucasi usque ad 7200' (M B, Ruprl), Iberie Cauca-icæ (Hoh!). Persiæ bor. in monte Elbrus (Ky. 130!) et montibus supra Asterabad (Bge!).

Formæ minores e locis elatioribus facile cum *M. alpestri* confunduntur a quâ pedicellis minus strictis, calyce breviori, radice tenuiori bienni tamen distinctæ sunt.

Ar. Geogr. Europa tota ubi in australioribus montana, Sibiria, regio Hima-laica, Insulæ Canarienses, Abyssinia.

8. ***M. platiphylla***, biennis, caulinis erectis elatis striatis gla-bris simplicibus, foliis tenerrimis præter ciliola pauca marginalia glabris radicalibus oblongis petiolatis superioribus latis a basi sessili rotundata oblongis acutis, racemis fructiferis brevibus laxis, pedicellis fructiferis adpresso hirtis tenuibus strictissimis inferioribus calyce triplo longioribus, calycis fructiferi laciniis lanceolatis acutis erecti-

patulis margine præsertim pilis rectiusculis erecto-patulis obsitis, corollæ limbo ampio tubo longiore plano, nuculis ovatis acutiusculis alterà facie carinatis ②. *M. latifolia* Boiss. et Ky. Sched. 1859 non Poiret.

Hab. in valle Teng Armeniæ Turcicæ prov. Musch 600° (Ky. Suppl. 699!).

Pedalis, insignis glabritie, foliis tenerimis, intermediis bipinnaribus, supremis 9-10 linearibus longis 6-8 latis, corolla eā *M. sylvaticæ* duplo major, calyx fructifer major et magis apertus. Ulterius observanda.

9. ***M. alpestris*** (Schm. Bohem. III, p. 26) perennis, adpresso-hispida, caulis pumilis rigidulis, foliis oblongo-lanceolatis inferioribus in petiolum attenuatis, racemis fructiferis brevibus strictis laxiusculis ebracteatis, pedicellis fructiferis erectis inferioribus calyce sublongioribus, calycis fructiferi pilis adpresso-hispida lacinia elongatis erecto-patulis, corollæ intense azureæ limbo plano tubo longiore, nuculis rotundato-ovatis ♀. *M. suaveolens* W. K. — *M. lithospermifolia* Hornem. — *M. sylvatica* var. *alpestris* Koch Syn. — *M. Pyrenaica* Pourr.

Hab. in pascuis alpinis, Macedonia 5000'-7000' (Griseb., Orph!), regio alpina Ponti Lazici supra Djimil 6000'-8000' (Bal!), Tauriæ mons Tchatyrdagh (Rehm!), Caucasus totus (M.B., C.A. Mey!), Caucasus orientalis 7000'-11000' frequentissime (Rupr!).

Nonnulli eam pro merâ *M. sylvaticæ* formâ e locis editioribus habent, radice duratio culturâ probanda.

Ar. Geogr. Montes Scandinaviæ, Scotiæ, Germaniæ, Galliæ, Helvetiæ, Corsicæ, Italiæ, regionis Danubialis, Songariæ, Sibiriæ, Americæ borealis.

10. ***M. Olympica*** (Boiss. Diagn. Ser. I, 4, p. 50) perennis, cæspitosa pilis adpresso-patulis foliorum basi tuberculatis hirsuta, caulis nanis, foliis radicalibus oblongo-spathulatis in petiolum longe attenuatis, caulinis parvis sessilibus oblongo-linearibus, racemis 2-4 terminalibus brevibus densiusculis, pedicellis fructiferis axi strictissimis calyce subbrevioribus, calyce pilis patulis uncinulatis albo-strigoso lacinia fructiferis erectis subconniventibus, corollæ azureæ limbo plano tubo longiore, nuculis oblongis compressis utrâque facie subcarinatis ♀. *Myosotis nana* Sm. Prodr. I, p. 112 quoad pl. Olympicam non Vill.

Hab. in cacuminibus Olympi Bithyni (Auch. 2269! Boiss! Clem!).

β. laxa. — Racemi fructiferi longiores laxiores. — *M. modesta* et *M. microsepula* Schott et Ky. in Ky. exs. 1853.

Hab. in regione alpinâ Tauri Cilicici (Ky! Bal!), ditionis Kassan Oglu (Ky. 103!).

Valde affinis *M. alpestris* a quâ præsertim differt pilis paginæ superioris crassius tuberculatis, calycis patulis valde uncinatis nec adpressis paucis uncinatis, nuculis oblongis nec rotundato-ovatis. Racemi fructiferi in typo breves densiori sed in varietate laxi quamobrem species mihi nunc subdubia.

11. M. intermedia (Link Enum. Berl. I, p. 164) biennis, villosa, caule parce ramoso plus minus elato, foliis radicalibus spathulato-oblongis in petiolum attenuatis caulinis oblongo-lanceolatis, racemis fructiferis aphyllis laxis longis, pedicellis fructiferis patentibus calyce duplo longioribus, calycibus profunde quinquefidis fructiferis sub-clausis pilis patulis vel reflexis uncinatis obsitis, corollæ rosetæ tandem cœruleæ tubo calyce breviori limbo parvo concavo ②. Ic. Rehb. tab. 122. — *M. scorpioides* var. *arvensis* L. — *M. arvensis* Roth. — *M. heteropoda* Trautv. Stirp. Nov. p. 11.

Hab. in cultis et arenosis montanis, Laconia in monte Malevo (Orph!), Attica mons Parnes (Heldr!), Thracia (Griseb.), Anatolia borealis omnis (Wiedem!), Caucasus et Transcaucasia (Lebed.), Persia borealis ad radices Elbrus (Ky. 741).

Ar. Geogr. Europa tota, Sibiria Altaica, P. B. Spei.

12. M. hispida (Schlecht. Mag. Nat. Ber. VIII, p. 229) annua, tota pilis patulis non uncinatis hispida, a collo saepius ramosa, racemis fructiferis aphyllis laxis caule vel ramo longioribus, foliis oblongis radicalibus in petiolum attenuatis, pedicellis patentibus adpresso hirtis calyci aequilongis vel brevioribus, calyce patentib; brevi quinque-fido fructifero aperto pilis patentibus uncinatis obsito, corollæ cœruleæ tubo calyce breviore limbo tubo breviore ③. Rehb. tab. 122. — *M. collina* Rehb. — *M. pygmaea* Bert. Amæn. I, p. 13.

Hab. in collibus et graminosis præsertim montanis, Græcia in montibus Hymetto, Parnes, insula Salami (Heldr!), Eubea (Orph!). Thraciæ (Friv!). Anatolia boreali (Wied.) et orientali ad Smyrnam (Heldr!), in monte Cadmo (Boiss!), Lyciæ (Bourg!), ins. Chios (Orph!), Cypro (Ky!), Syriæ ad Aleppo (Haussk!), Libano et Antilibano (Gailly), Mesopotamia (Chesn.), Transcaucasiæ (Lebed., Hohen!), Persia bor. ad Teheran (Ky!).

M. grandiflora Diagn. Ser. I, 11, p. 123. — Corollæ limbus major tubo subæquilongus.

Hab. in agris et olivetis Cretæ ad Cydoniam (Heldr!), Ponti Lazici ad Rhizé (Ball!).

Ar. Geogr. Europa fere omnis a Scandinaviâ ad Rossiam, Africa borealis.

13. M. stricta (Link Enum. Berol. I, p. 164) annua, tota breviter et patule pilis ad foliorum insertionem saepius uncinatis hispida collo multicaulis pumila, foliis oblongis obtusis radicalibus rosulatis brevissime petiolatis, racemis fructiferis rigidis elongatis laxis inferne foliatis, pedicellis strictis calyce brevioribus, calyce quinquefido fructifero clauso piloso pilis baseos divaricatis uncinatis, corollæ minimæ cœruleæ tubo calyce breviore limbo minuto concavo ④. Ic. Rehb. tab. 123. — *M. areuaria* Schrad. — *M. minutiflora* Boiss. et Reut. Pag. p. 80 (forma minor calyce abbreviato magis hispido et minus clauso). — *M. micrantha* Pall. ex Lebed.

Hab. in arenosis regionis inferioris et saepius montane et alpinæ, Byzantii (Bergg.), Lydiæ (Ball), Phrygiæ prope Ouchak (Bal!) forma nuculis majoribus

et calyce fructifero longiore 2 lineas longo), Cappadociæ ad Argæum (Ky. Suppl. 282!), Cariae regione alpinæ in monte Cadmo (Boiss!), Armeniæ ad Bai-but (Bourg!), Ponto Lazico ad Djimil 6900' (Bal!), Libano ad Cedros (Bl!), Antilibano supra Zebdani (Ky!), Armeniæ Rossicæ et Transcaucasiæ (C.A M., Koch), Tauriæ ad Sudak (Rehm!), Affghania (Griff!), Belutschia (Stocks!).

β. speluncicola. — Tenera, nana, rami et racemi flexuosi laxi, folia tenuiter membranacea ovata obtusa brevius et parcius hirta, pedicelli inferiores calyce sublongiores patuli interdum subnutantes, calyx minus clausus. — *M. speluncicola* Schott in Ky. Sched. 1853.

Hab. in rupestribus opacis et speluncis vallis Karli Boghas Tauri Cilicici 6000' (Ky. 42!), ad Pylas Cilicias (Bal. 554!).

Species calycis longitudine, racemis plus minus laxis valde variabilis; specimina ex alpibus Granatensibus parviflora olim sub *M. minutifloræ* nomine distinxii sed formas intermedias nuper vidi. Varietas β foliis dilatatis teneris, racemis flexuosis, pedicellis inferioribus interdum elongatis et subnutantibus printâ facie distinctissima videtur, tamen omnino forma e loco umbroso, similem vidi in Galloprovinciâ a cl. Marcilly lectam.

Ar Geogr. Europa tota a Scandinaviâ ad Rossiam, Sibiria Uralensis et Altaica.

14. **M. uncata** (Boiss. et Bal. Diagn. Ser. II, 3, p. 139) annua, patule et longe piis non uncinatis hispida, foliis infimis oblongis in petiolum brevem attenuatis caulinis oblongo-linearibus, caule brevi dichotome ramoso ramis in racemos fructiferos longissimos ebracteatos laxos abeuntibus, pedicellis calyce brevioribus patulis vel cernuis, calyce fructifero pilis longis patulis uncinatis hispido campanulato subclauso, corollæ minimæ azureæ tubo calyce sublongiore, limbo minimo concavo \odot .

Hab. in umbrosis faucis Guzeldere supra Sedichig Cilicie littoralis (Bal!). Fl. Apr.

Caulis 2-3-pollicaris, racemi tandem semipedales, calyces fructiferi parvi et corollæ *M. strictæ* var. *minutifloræ*. Ab eâ sat differre videtur pilis caulis et foliorum non uncinatis, racemis longissimis ebracteatis, pedicellis brevissimis nutanti-recurvis.

15. **M. refracta** (Boiss. Voy. Esp. p. 433, tab. 125) annua, pilis albis patulis ad caulem sœpe retrorsis hispida collo vel caule brevissimo ramoso, ramis in racemos aphyllos demum valde elongatos flexuosos sœpe tortuosos unilaterales laxos abeuntibus, foliis minutis oblongo-lanceolatis radicalibus subpetiolatis spathulatis, pedicellis fructiferis calyce multo brevioribus incrassatis recurvo-refractis, calyce florifero semiquinquefido breviter campanulato patule et dense hispido fructifero elongato basi attenuato pyriformi-clavato toto vel superne tantum unciato-piloso, corollæ minimæ cœruleæ tubo calyce non longiore, limbo concavo, nuculis acutis \odot .

Hab. in lapidosis regionis subalpinæ et alpinæ, mons Parnes Atticæ in abietinæ (Heldr!). Creta in monte Idæ 5-6.000' (Heldr!). Cyprus (Ky. 904!), Lydiæ et Cariae montes Tmolus et Cadmus (Boiss! Auch. 2247!), Phrygia ad Kaiagheulderé prope Ouchak (Bal!), Lyciæ montes Elmalu (Bourg. 496!), Syria in cacu-

mine Cassii (Boiss!), Libano ad Hasrun (Bl!), Antilibano in montibus Zebdani (Boiss!), Mesopotamiā prope Mardin (Haussk!), Turkestaniā (Lehm.).

Racemi floriferi brevissimi fructiferi sœpe semipedales. Species calycibus fructiferis basi longe attenuatis et refractim axi plus minus arce incubentibus insignis.

Ar. Geogr. Montes Hispaniae australis.

SECT. II. STROPHIOSTOMA.

16. **M. sparsiflora** (Mik. in Hoppe Taschenb. 1807, p. 74) annua, viridis pilis adpressiusculis sparsis obsita, caulis flaccidis laxe ramosis subdiffusis, foliis teneris inferioribus ovato-spathulatis obtusis petiolatis cæteris oblongo-lanceolatis acutis, racemis brevibus paucifloris basi foliatis, pedicellis fructiferis inferioribus calyce multo longioribus refractis, calyce adpressiuscule setuloso pilis baseos uncinatis fructifero aucto campanulato, corollæ cœruleæ limbo tubum calyce subbreviore non æquante ♂. *Strophiostoma sparsiflorum* Turcz.

Hab. in dumosis humidis Ponti Lazici ad Castellum Rhizé (Bal!), Taurie (Stev!), Caucasi occid. et centralis 2-30° (M.B., C.A.M., Rupr!) et orientalis 60° (Rupr!), Iberiae et prov. Talysh (C.A. Mey.), Persiae prov. Aderbidjan (Szovl.).

Ar. Geogr. Scandinavia, Germania borealis, orientalis et australis, regio Danubialis, Bosnia, Rossia, Songaria, Sibiria.

17. **M. propinqua** (F. et M. in Karel. Enum. Turcom. Bull. Mosq. 1839, p. 163) annua, viridis setis patulis e tuberculo sœpe ortis hispidula, caulis erectis dichotome ramosis, foliis teneris inferioribus obovato-oblongis basi breviter attenuatis cæteris sessilibus oblongis, racemis demum longis ebracteatis unilateraliter multifloris laxis, pedicellis erecto-patulis rarius subrecurvis inferioribus calyce vix duplo longioribus, calyce patule setoso pilis baseos uncinatis fructifero aucto campanulato, corollæ cœruleæ tubo calyce subbreviore limbo minimo ♂.

Hab. in Ponto Lazico prope Djimil 6000' (Bal!), in montibus Persiae borealis supra Asterabad (Karel!), prope Jezd (Buhse!).

Valde affinis præcedenti, flores et corollæ erroneè majores dictæ, ab ea differt tantum racemis ebracteatis 10-20-floris tandem 3-4 pollices longis et pedicellis brevioribus, pubescens paulo copiosior et *Lagis* patula.

18. **M. amæna** (Ruprecht MSS. sub *Strophiostoma*) perennis, patule et tenuiter pilosa, surculis repentibus basi fibrillosis, caulis erectis elatis parce ramosis, foliis oblongis obtusis basi attenuatis, inferioribus et surculorum in petiolum attenuatis, racemis demum elongatis laxissimis aphyllis pedicellis tenuibus erecto-patulis calyce plures longioribus, calyce pilis erecto-patulis vix uncinatis hispido

fructifero subaccreto campanulato, corollæ cœruleæ tubo calyci æquilongo, limbo amplio plano tubo æquilongo lobis rotundatis ȝ.

Hab. in fruticetis Imeretiæ Caucasicæ inter Sekari et Saleschkro 2000', Caucaſi centralis prope Kaischaur 4800' et districtu Alagir 2000' (Rupr!).

Planta $\frac{1}{2}$ - $\frac{1}{2}$ -pedalis, a precedentibus eximie distincta rhizomate surculosorepenti, corolle limbo diam. 2 lineas longo; racemi fructiferi 4-5-pollicares pedicellis tandem 6-7 lineas longis, nuculae ut in *M. sparsiflorū* acutæ oblongæ nitidæ strophiolatae.

SECT. III. PHYLLOCEPHALUM.

19. **M. involucrata** (Stev. Mem. Soc. Mosq. III, p. 253) annua, tota setis rigidis basi tuberculatis adpressiusculis hispida aspera, radice longâ tenui subsimplici, foliis oblongis obtusis basi longe attenuatis, caule brevissimo paulo supra collum sub cymâ centrali brachiatum ramoso ramis lateralibus decumbentibus elongatis subaphyllis apice iterum sub cymis terminalibus brachiatis, cymâ centrali et terminalibus sessilibus brevissime et pluries dichotomis cum flore in dichotomiâ capituliformibus, foliis floralibus flores longe superantibus, pedicellis crassiusculis calyci æquilongis tandem recurvis, calyce obconico-campanulato hirsutissimo semiquinquefido laciniis acutis, corollâ albidâ vel cœruleâ minimâ calycem parum superante lobis ovato-orbiculatis tubo æquilongis, nuculis maximis abortu subsolitariis ovatis subcompressis facie interiori carinatis, areolâ concavâ cum umbone ⊖.

Hab. inter schisti fragmenta in ditionibus Tindal, Bogos, Daghestan Caucaſi orientalis 4000'-9000' (Stev! C.A.M! Rupr!).

Rarius simplex pollicaris, saepius ramose brachiata ramis lateralibus 3-4 poll. longis, folia majora pollicaria apice 3 lineas lata, nuculae $1\frac{3}{4}$ lin. longæ lineam late, flores minimi. Planta inter *Myosotides* inflorescentiâ cymosa et nuculis maximis anomala. An *Eritrichii* species? sed corollæ aestivationem ut el. Alph. D.C. contortam quoque vidi et nuculae areola basilaris.

ERITRICHIUM (Schrad. Diss. Asperif. 1820).

Calyx quinquepartitus. Corolla hypocrateriforma fauce fornicibus obtusis saepè biserialibus superpositis clausâ, limbi lobis aestivatione quincuncialibus. Stamina inclusa filamentis brevibus. Antheræ oblongæ medio dorso affixæ. Nuculae triquetrae antice planæ et margine prominenti cinctæ postice obtusiores et areâ minutâ ad dorsi basin receptaculo affixe. — Herbæ facie *Myosotidis*.

1. **E. nanum** (Vill. Delph. II, 459 sub *Myosotide*) dense cespitosa pilis longis albis hirsuta, foliis minutis oblongo-linearibus, ramis erectis minute foliatis in racemos paucifloros fasciculiformes demum subelongatos abeuntibus, calycis hispido-setosi laciniis tubum corollæ

æquantibus, corollæ azureæ limbo ampio tubo longiore fornicibus duplice ordine superpositis superioribus transverse latioribus inferioribus punctiformibus. nuculis ovato-trigonis ad angulos alâ dentatâ vel integrâ marginatis ♀. *E. nanum* Schrad. — Ic. Rehb. tab. 124.

Hab. in regione alpinâ Caucasi occidentalis 7000'-9.000' (G.A. Meyer) Non vidi.

Ar. Geogr. Alpium totus tractus, Transylvanie montes.

2. **E. Persicum** (Boiss. Diagn. Ser. I, 14, p. 123) totum brevissime et crispule velutinum cinereum cespitosum, collo petiolis vetustis dense stipato, foliis radicalibus numerosis obovatis obtusis in petiolum longe attenuatis, caulinis anguste linear-lanceolatis, ramis procumbentibus vel ascendentibus cespite subduplo longioribus, racemis 1-3 sub anthesi scorpioideo-capitatis, calycis laciniis elliptico-linearibus obtusis, corollæ tubo calyce breviore limbo parvo, fornicibus uniseriatis, nuculis.... ♀.

Hab. ad nives deliquescentes in alpibus Hasartschal montis Elbrus occidentalis Persiae borealis 1100' (Ky. 500').

Fructu ignoto inter *Eritrichium* et *Echinospermum* subdubium sed priori ob habitum et stationem valde alpinam potius adnumerandum. Folia radicata pollicem cum petiolo longa apice 3-4 lineas lata, rami 2-3-pollicares, corolla ea *E. nani* multo minor, ab eo indumento diversissimum.

SUBTR. IV. CYNOGLOSSÆ D.C. — Nuculæ quatuor vel abortu pauciores gynobasi vel stylo areolâ planâ elongatâ plus minus laterali insertæ et cum eo sèpius coalitæ.

ROCHELIA (Rehb. Flora 1824, p. 243).

Calyx quinquepartitus laciniis post anthesin sèpius auctis. Corolla infundibuliformis minima calyci subæqualis sub medio tubo fornicibus obseuris instructa. Stamina 5 inclusa filamentis brevissimis, antheris oblongis basi retusis apice connectivo brevi exerto. Stigma capitatum. Nuculæ binæ (rarius abortu unica) styli basi adnatae obliquæ basi latiores apice acutatæ sèpius glochidiato-tuberculatae. — Herbæ facie *Lithospermi* sect. *Rhytispermi*.

* Calycis laciniæ 9-10.

1. **R. cancellata** (Boiss. et Bal. Diagn. Ser. II, 3, p. 136) setis longis adpressis basi tuberculatis canescens, caule fere a basi divaricatum ramoso, foliis inferioribus oblongis basi attenuatis caulinis linearibus, racemis demum elongatis laxis, bracteis linear-ellipticis pedicellos superantibus, pedicellis fructiferis recurvis calyce duplo brevioribus, calycis fructiferi pilis uncatis hispidissimi laciniis 9-10

lineari-subulatis basi patulo-reflexis dein curvatis et cancellatim apicibus subconiventibus fructu duplo longioribus, nuculis binis glochidibus stellatis sessilibus crebre tuberculatis ♂.

Hab. in valle Yachamichlar Keui prope Ouchak in Phrygiâ 4000' (Bal!), in regionis montanæ superioris Tauri Cilicici locis hyeme inundatis supra Bulgharmaaden (Bal!).

• • Calycis laciniæ 5.

+ Calycis fructiferi laciniæ arcuato-incurvæ.

1. Pedicellus fructifer calyci æquilongus vel brevior.

2. **R. stellulata** (Rehb. loc. cit.) a collo pluricaulis vel superne breviter paniculato-ramosa setulis tuberculatis adpressis canescens, foliis inferioribus oblongo-spathulatis petiolatis cæteris linearibus sessilibus, racemis fructiferis elongatis, bracteis linearibus calyces ad vel supra axillas sitos superantibus, pedicellis calycem æquantibus demum deflexis, calycis fructiferi laciniis subquinis linearibus adpresiuseule hamato-pilosus basi patulis laxis sursum hamato-incurvis fructu duplo longioribus, nuculis binis (rarius solitariis) glochidibus stellatis sessilibus dense tuberculatis ♂. Ic. Rehb. Ic. Crit. 2, tab. 123. — Ic. Germ. tab. 132. — *Lithospermum dispermum* L. Sp. 191. — *L. retortum* Pall.

Hab. in arenosis et cultis fere totius ditionis, in australioribus montana et alpina, mons Ossa Thessaliæ (Sibth.), Anatolia in Phrygiâ ad Ouchak (Bal!), Lyciâ ad Elmalu (Bourg!), Ciliciâ (Ky! Bal!), Caria in monte Cadmo (Boiss!), Pisidiâ (Heldr!), Cappadociâ (Ky! Bal!), Syriâ interiore ad Aleppo (Auch!), Damascum (Gaill!), in Antilibano (Boiss!), Mesopotamiâ (Auch!), Armeniâ Turcicâ (Huet!), Tauriâ (Stev!), Transcaucasiâ et Armeniâ Rossia (Szov! Hoh!), Persiâ bor., (Auch!), Belutschia (Stocks!), Affghania (Griff.).

Nuculas vidi in specimine Persico (Ky. 89) binas et solitarias.

Ar. Geogr. Hispania centralis, regio Danubialis, Rossia australis, Turcomania borealis, Songaria, Regio Himalaica occid., Algeria interior.

3. **R. incana** (Kar. et Kir. Enum. 603) a collo pluricaulis vel simplex setulis minute tuberculatis canescens, caulis parce ramosis, foliis linearibus, racemis fructiferis laxis, bracteis linearibus calyces superantibus, pedicellis calyce longioribus tandem refractis, calycis fructiferi laciniis 5 anguste linearibus hamatim et patule setulosis basi nuculis adpressis sursum hamato-incurvis fructu vix longioribus, nuculis binis glochidibus stellatis sessilibus parce obsitis ♂.

Hab. in rupibus graniticis Turkestaniæ prope Bakali (Lehn. ex Bge).

Præcedente minus cana, gracilior; an ab ea characteribus sat firmis distincta?

Ar. Geogr. Songaria.

4. **R. Persica** (Bunge in litt.) setulis basi tuberculatis erecto-

patulis hirta, a collo ramosa vel simplex, caulinis gracilibus parce ramosis, foliis inferioribus linearispathulatis basi attenuatis cæteris linearibus, racemis fructiferis elongatis laxis, pedicellis recurvis calyce brevioribus, calycis fructiferi laciniis 5 linearibus patentibus setis rectis hirsutis fructu sublongioribus, nuculis glochidibus stellatis sessilibus parce tuberculatis ⊖.

Hab. in Persiâ prope Teheran et ad Ssertschah Persiæ mediae orientalis (Bunge!).

Plantula tenuis 4-6-pollicaris, racemi fructiferi 2-3-pollicares. A *R. stellata* et *R. incana* calycis subsessilis pilis rectis nec apice hamatis distincta; nuculae multo faciliter secedentes.

5. ***R. mollis*** (Bunge in litt.) tota pilis basi minute tuberculatis patulis villosa collo ramosa, foliis linearibus inferioribus basi attenuatis, racenis fructiferis elongatis densiusculis, bracteis linearibus calyces æquantibus, pedicellis calyce fructifero brevioribus erecto-patulis, calycis fructiferi laciniis 5 elongatis linearibus pilis patulis longis apice uncatis hispidissimis basi patulis sursum incurvis fructu longioribus, nuculâ unicâ glochidibus stellatis crebris tuberculatâ ⊖.

Hab. in Persiâ austro-orientali prope Kerman (Bge!).

Calycis fructiferi laciniis longis hamato-curvatis *R. stellatae* affinis, ab eâ differt non tantum nuculis solitariis (quæ in eâ interdum quoque occurunt) sed pilis totius plantæ et præsertim calycis patentissimis pedicellisque brevibus.

6. ***R. leiocarpa*** (Ledeb. Fl. Alt. p. 172) setis tenuibus adpressis basi tuberculatis hirta, caule gracili superne parce ramoso, foliis linearibus infimis subspathulatis, racemis fructiferis laxis, bracteis fructus vix superantibus, pedicellis refractis calyce subæquilongis, calycis fructiferi laciniis 5 parum elongatis anguste linearibus patentim hamato-pilosis incurvis laxiuseculis fructu vix longioribus, nuculis lævissimis nitidis ⊖. Ic. Ledeb. tab. 244.

Hab. in Turkestaniâ prope Bakali (Lehm.).

Facies *R. incane* sed fructus lævissimi.

Ar. Geogr. Songaria.

2 Pedicellus fructifer calyce brevior.

7. ***R. stylaris***, pilis basi tuberculatis strigulosa canescens, caule tenui pumilo superne paniculato-corymboso, foliis inferioribus linearispathulatis basi attenuatis cæteris linearibus, racemis fructiferis laxis unilateralibus, bracteis brevissimis squamiformibus, pedicellis irnec-tiferis tenuissimis reactis calyce fructifero vix acreto sesquilongioribus, calycis laciniis 5 linearibus pilis rectis hirsutis patentibus apice curvatis fructu subduplo longioribus, nuculis binis glochidibus stellatis sessilibus crebre tuberculatis, stylo nuculas valde superante ⊖.

Hab. in Affghaniâ circa Pushut (Griff.).

Fructifera semipedalis, a *R. stellulata* et affinibus pedicellis fructiferis multo longioribus, calycis pilis non hamatis, stylo elongato distincta.

8. ***R. peduncularis*** (Boiss. Diagn. Ser. I, 7, p. 35) setis basi tuberculatis striguloso-canescens, caule gracili tenui simplici vel 2-3 racemoso, foliis parvis linearibus, racemis demum elongatis laxis, bracteis brevibus, pedicellis fructiferis refractis clavato-incrassatis calyce duplo longioribus, calyce fructifero valde aucto laciniis lanceolatis carinatis erecto-patentibus fructu duplo longioribus, nuculis binis verrucis stellatim lepidotis obsitis ⊖.

Hab. in humidis alpis Kuh Delu Persiæ australis (Ky. 531!).

Florifera 2 pollices longa fructifera 6-8-pollicaris, pedicelli fructiferi 4-5 lineas longi apice in calycis tubum coriaceum sensim incrassati.

+ + Calycis fructiferi laciniæ rectæ.

9. ***R. macrocalyx*** (Bge Pl. Lehm. p. 419) collo ramosa sub-patentim hirsuta canescens, foliis inferioribus obovato-oblongis basi attenuatis, cæteris bracteisque pedicellos æquantibus vel superantibus linearibus, racemis demum elongatis laxiusculis, pedicellis rectis non incrassatis calyci æquilongis et longioribus, calycis fructiferi valde aucti laciniis 5 rectis lanceolatis basi subattenuatâ bigibbis pilis hamatis hispidis fructu triplo longioribus, nuculis glochidiat-saccharatis ⊖. *R. rectipes* Stocks in Hook. Journ. Bot. IV, 1852, p. 176.

Hab. prope Ssertschah Persiæ mediae orientalis (Bge!), in rupestribus graniticis Turkestanie ad Bakali (Lehm.), in Belutschiæ superiore prope Gurghina (Stocks!), Affghania ad Achulzey (Griff.).

Planta florifera bipollicularis fructifera 5-6-pollicaris, fere a basi racemifera. Species laciniis calycinis rectis latis, fructiferis 2 $\frac{1}{2}$ lin. longis distinctissima.

10. ***R. cardiosepala*** (Bunge Pl. Lehm. p. 420) setis basi tuberculatis strigillosa canescens, caule humili superne parce ramoso, foliis radicalibus ellipticis subsessilibus cæteris lineari-spathulatis, bracteis flori æquilongis, racemis fructiferis elongatis laxis, pedicellis fructiferis vix incrassatis calyci subæquilongis patulo-subdeflexis, calycis sub anthesi basi truncati pyramidati laciniis pilis hamatis hispidis lanceolatis fructiferis auctis a basi cordato-auriculatâ lanceolatis erectis fructu duplo longioribus, nuculis binis glochidibus sessilibus stellato-tuberculatis ⊖.

Hab. in rupibus graniticis ad Bakali Turkestanie (Lehm.). Vid. spec. junius ab am. a Bge commun.

Species a *R. macrocalyx* pedicellis refracto-patentibus et calycis laciniis basi cordato-auriculatis distincta.

ECHINOSPERMUM (Swartz in Lehm. Asp. p. 413. — *Heterocaryum* Alph. D.C.).

Calyx quinquepartitus. Corolla hypocraterimorpha fauce fornicibus

minutis clausâ. Stamina inclusa antheris filamento brevi longioribus. Nuculae quatuor lateraliter columnæ centrali adnatae et interdum cum ea et inter se arete concretae erectæ trigone vel compressæ, disco tuberculato vel laevi aculeis glochidiatis uni vel pluriseriatis basi liberis vel connatis marginato. — Herbae minutifloræ saepius alpresso strigulosæ. — Genus affine *Paracaryo* ad quod species posteriores aculeis marginalibus nucularum basi connatis accedunt.

Specierum Orientalium distributio.

SECT. I. HETEROCARYUM Alph. D.C. Prodr. — Nuculae elongatae linearis-oblongæ toto ventre et marginibus inter se et cum stylo in fructum non partibilem concretae facie dorsali marginatae et aculeatae.

E. Szovitsianum, oligacanthum, loriyatum.

SECT. II. SCLEROCLARYUM Alph. D.C. — Nuculae acie internâ stylo adnatae marginibus liberæ in fructum vix partibilem concretae, facie dorsali carinatae scrobiculato-nuricatae non distincte marginatae.

E. spinocarpos.

SECT. III. LAPPULA. — Nuculae breves basi stylo adnatae partibiles areolâ dorsali et margine distincto præditæ.

* Nuculae margine 2-3-serialiter glochidiatae.

E. Lappula, heteracanthum, barbatum.

* * Nuculae margine uniserialiter glochidiatae aculeis basi distinctis.

E. patulum, microcarpum, Sinaicum. semiglabrum, macranthum.

* * * Nuculae margine uniserialiter glochidiatae aculeis basi plus minus concretis.

E. marginatum, Bungei, sessiliflorum.

SECT. I. HETEROCARYUM.

1. **E. Szovitsianum** (F. et M. Ind. II, Petrop. 1833) setis brevibus patulis e tuberculo ortis hispidum asperum a basi ramosum, ramis demum incrassatis tortuosis in racemos fructiferos crassos laxos foliosos abeuntibus, foliis linearis-lanceolatis, floralibus acutis pedicel-

los et fructus superantibus, pedicellis fructiferis clavato-incrassatis patent-i-incurvis vel recurvis fructu 2-3-plo longioribus, calycis fructiferi laciniis patentibus linearibus fructu longioribus, nuculis linearis-oblengis in fructum prismaticum concretis facie linearis tuberculatis utrinque aculeis 5-6 basi triangulari inter se concretis elongatis glochidiatis marginatis, nuculæ alteræ (rarius duabus oppositis) alâ membranacea latiori dentata obsoletis ⊖. *Heterocaryum Szovitzianum*, *H. pachypodium* et *H. rigidum* Alph. d.c. Prodri. X, p. 144.

Hab. in aridis, ad vias, inter segetes, Lycia circa Elmalu (Bourg!), Pisidia ad lacum Buldur (Heldr!), Cappadocia (Bal!), Transcaucasia ad Persiae fines (Szov!), Persia borealis et orientalis (Szov! Buhse! Bunge!), Turkestania (ex Bge.).

Affine *E. minimo* Lehm. ex Rossia australi et Sibiria Uralensi oriundo et a quo differt (an satis?) nuculæ alteræ saepius alata et aculeis omnibus basi connatis nec fere liberis.

Ar. Geogr. Songaria.

2. E. oligacanthum, setis brevibus patulis e tubculo ortis asperum, ramis demum divaricatim ramulosis diffusis in racemos fructiferos incrassatos elongatos laxos abeuntibus, foliis linearibus acutiusculis floralibus calyce longioribus, pedicellis fructiferis incrassatis fructu brevioribus vel subnullis, calycis fructiferi laciniis linearibus patentibus fructum æquantibus, nuculis in fructum breviter prismaticum concretis linearis-oblengis facie tuberculatis vel lœvibus utrinque aculeis 2-3 subliberis remotis a basi late triangulari elongatis glochidiatis marginatis, nuculæ alteræ margine saepè in alam membranaceam remote dentatam dilatato ⊖.

Hab. in Persiâ orientali prope Schahrud et Ssertschah (Bge!), Affghanâ ad Puschut et Seriab (Griff!), Belutschia (Stocks!).

Differt ab *E. Szovitsiano* ramis minus incrassatis, fructu sessili dimidio minore 2 nec 4 lineas longo, aculeis marginalibus nucularum multo paucioribus.

3. E. lœvigatum (Kar. et Kir Enum. № 583) præter setas basi tuberculatas breves adpressas ad foliorum margines et ad calyces et pedicellos glabrinsculum, caule tenui gracili pumilo simplici vel parce et bifide ramoso in racemos paucifloros fructiferos breves laxissimos abeunti, foliis oblongis floralibus linearibus acutis pedicello brevioribus, pedicellis fructiferis patentibus subcurvatis clavato-incrassatis fructu plures longioribus, calycis demum patentis laciniis fructu sublævioribus, nuculis in fructum pyramidatum concretis disco adpresso pubescenti-asperis rarius glabris aculeis apice glochidiatis basi dilatatis liberis vel inter se plus minus concretis marginatis ⊖. *Heterocaryum lœvigatum* Alph. d.c.

Hab. in Persiâ orientali inter Chabbise et Kerman (Bge!), Turkestaniâ ad Bakali (Lehm!), Belutschia ad Doubund (Stocks!).

Ar. Geogr. Songaria, Sibiria Altaica.

SECT. II. SCLEROCARYUM.

4. E. spinocarpos (Forsk. Descr. 41 sub *Anchusá*) totum setis basi tuberculatis adpressis rarius subpatulis canescens rarius virescens pumilum ad vel paulo supra collum divaricatum et dichotome ramosissimum tandem diffusum, racemis crassiuseculis demum induratis laxis flexuosis, foliis linearis-spathulatis, pedicellis fructiferis incrassatis brevissimis erectis, bracteis linearibus fructu longioribus, corollæ minutæ albidae vel cærulecentis limbo calycem vix excedenti, calyce fructifero valde acreto laciniis linearibus crassis fructum pyramidatum longe stylosum superantibus erecto-patulis, nuculis erectis triangularibus carinatis aculeis conicis crassis versus basin nuculæ elongatis muricatis et scrobiculatis ♂. *Myosotis spinocarpos* Vahl Symb. II, p. 32. — *Echinospermum Vahlianum* Lehm. Asp. N° 103.

Hab. in desertis et incultis, Cappadocia in planitie Karahissar (Bal!), Egyptus inter Kahiram et Suez (Boiss!), ad Alexandriam (Forsk., Samarl), deserta Sinaica (Schimp. 178! Boiss!), Gebel Abiad in deserto ad Palmyram (Bl!), Iberia Caucasicæ et ditio Talyesch (Hohl, C. A. Mey!), Turcomania ad Caspium (C. A. M!), Persia media Orientalis (Bge!), Affghania (Griff!), Belutschia (Stocks!), Arabice *Kleiterai*.

SECT. III. LAPPULA.

* Nuculæ 2-3-serialiter glochidiatae.

5. E. Lappula (L. Sp. 189 sub *Myosotide*) pilis basi tuberculatis adpressis hirsutum, caule virgato rigido elato paniculato, ramis in racemos demum elongatos unilaterales abeuntibus, foliis lanceolatis caulinis strictis, bracteis inferioribus calyce longioribus, pedicellis fructiferis erectis calyce brevioribus, calycis fructiferi laciniis patulis linearis-oblongis, corollæ cæruleæ tubo calyce subbreviore, limbo tubo breviore, nuculis facie tuberculatis margine 2-3-serialim glochidiatis ♂ vel ♀. *Ech. Lappula* Lehm. Asp. N° 94. — Ic. Rchb. tab. 128. — *Cynoglossum Lappula* Scop. — *Myosotis squarrosa* Retz. — *Echin. squarrosum* Rchb. Fl. exc. — *E. Ispahanicum* Boiss. Diagn. Ser. I, 11, p. 123.

Hab. in siccis, ad vias, Peloponnesus (Bory!), Bithynia (Pestal.), Anatolia bor. (Wied!). Armenia Turcica (Huet!), Pontus Lazicus (Bal!), Tauria Stev!, Caucasus et Transcaucasia (Ledeb., Hohen.). Persia borealis (Buhsel) australis ad Ispahan (Auch. 5018!).

Ar. Geogr. Europa omnis a Scandinaviâ ad Rossiam medianam et ab Hispaniâ et Italiâ mediis ad Dalmatiâ et regionem Danubialem, Sibiria omnis, Africa borealis.

6. E. heteracanthum (Ledeb. Suppl. Sem. Dorpat 1823) pilis basi tuberculatis patulis et adpressis obsitum, caule elato virgato

superne in racemos paucos demum rigidos elongatos parum laxos abeunti, foliis linear-lanceolatis superioribus strictis, pedicellis erectis calyce brevioribus, bracteis calyci æquifloris, calycis fructiferi laciniis expansis linearibus, corollæ minute limbo calyce parum longiore, nuculis margine biseriatim glochidiatis seriei intericris aculeis longioribus basi triangulari inter se connatis, faciebus tuberculatis ⊖. Ic. Eichw. Casp. tab. 21.

Hab. in Caucasi lapidosis ad fluvium Kuban et circa Thermas Nartzana (C.A. Mey.), in Sionchetiâ (Eichw.). Editione nondum vidi.

Ar Geogr. Rossia australi-orientalis, Turcomania borealis ad Caspium.

7. **E. barbatum** (M.B. Taur. Cauc. I, p. 421 sub *Myosotide*) adpressissime canescens et setis basi tuberculatis subpatulis plus minus ciliatum a collo saepe pluricaule, caulinis elatis virgatis paniculatis, racemis demum valde elongatis laxissimis foliis radicalibus linearispathulatis caulinis linearibus apice saepe recurvis, bracteis minutis, pedicellis erectis calyce non longioribus, calycis fructiferi laciniis linearibus patulis, corollæ albide vel caeruleæ tubo calycem æquanti, limbo expanso magno tubo longiore, nuculis facie tuberculatis margine subbiseriatim et breviter aculeato-glochidiatis ⊖ vel ⊙. Ic. Cent. Ross. tab. 36. — *E. barbatum* Lehm. — *Myosotis saxatilis* Pall. — *Anchusa leucocephala* Lehmann Asper. p. 186. — *Lithospermum orientale leucocephalum* flore caruleo parvo Tournef. Cor. p. 6 et herb!

3. Nucularum discus secus mediam faciem quoque glochidiatus. — *E. Cariense* Boiss. Diagn. Ser. I. 4. p. 40.

Hab. forma typica ut et varietas β in campis sterilibus, lapidosis totius Anatoliae et Armeniae Turcicæ!, Tauriae (Stev!), Transcaucasie (M.B., Hoh! etc.), ad Bakn (C.A.M!) forma procumbens). Ciliciae (Ky!), Syriae in Antilibano (Boiss! Gaill!), Persia borealis (Buhse! Bunge!).

Species ab *E. Lappula* indumento adpressissimo canescente, corollis duplo vel triplo majoribus facile distinguenda. Inter typum et varietatem intermedia frequentia sunt. Planta Tournefortii a cl. W. Barbe et Jules Poisson benevolè comparata hinc certe spectat

Ar Geogr. Bulgaria, Rossia australis, Turcomania, Sibiria Uralensis.

* * Nuculae uniserialiter glochidiatae aculeis distinctis.

8. **E. patulum** (Lehm. Asper. N° 95) pilis basi tuberculatis patulis pilosum, caule erecto corymbose paniculato, foliis inferioribus oblongo-lanceolatis caulinis linearibus strictis, racemis demum strictis laxis unilateralibus, bracteis calyces superantibus, calycis laciniis fructiferis linearibus patulis, pedicellis brevissimis erectis, corollæ caeruleæ limbo minuto calycem parum excedente, nuculis marginè longe et simplici serie glochidiatis dorso tuberculatis ⊖. *Myosotis squarrosa* M.B. Taur. Cauc. I. p. 120 non Retz.

Hab. in ruderatis, agris incultis, etc., Bithynia (Pestal. ex Griseb.), Anatolia borealis (Wiedem!), Cilicia in regione superiori montana Tauri supra Bulgharmaaden (Bal!), Armenia Turcica ad Baibut (Bourg. exs. sub *E. squarrosa*!).

Armeniâ Rossicâ (Kowal!), Caucaso et Transcaucasiâ (Ledeb., Hohen!), Tauriâ (Stev.).

Ar. Geogr. Hispania australis, Rossia australis, Songaria, Sibiria Altaica, Africa borealis interior.

9. **E. microcarpum** (Led. Fl. Alt. I, p. 202) pilis basi sèpius tuberculatis adpressis canescens a collo sèpe pluricaule, caulis ramosis ramisque tenuiter virgatis, racemis fructiferis laxissimis elongatis, foliis inferioribus oblongo-linearibus basi attenuatis cæteris linearibus, corollæ mediocris cœruleæ tubo limboque æquilongis, pedicellis fructiferis strictissimis fructu æquilongis vel vix longioribus, calycis fructiferi laciniis fructu adpressis brevibus, fructù pyramidati stylo longiusculo superati nuculis margine breviter et simplici serie aculeato-glochidiatis undique cœbre tuberculatis, carinâ sèpius longius tuberculatâ ♂. Ledeb. Ic tab. 183. — *E. rigidum* D.C. Prodr. X, p. 141. — *E. filiforme* Godet in D.C. Prodr. p. 140!

Hab. in Persiâ (Auch. in D.C. herb!), prov. Aderbidjan, circa Jezd et Ispahan (Buhse!), sept. in prov. Khorassan septentrionali ad Schahrud et mediâ orientali (Bge!), australi in regione alpinâ montium Kuh Nur et Sawers (Haussk!) et Kurdistaniæ Persicæ in montibus Avroman et Schahu ad 10000' usque (Haussk!), in Caucaso orientali (ex Godet).

Indumentum adpressum canescens *E. barbati* a quo differt caulis ramisque tenuioribus, racemis tandem laxissimis, calycis fructiferi laciniis erectis, nucularum aculeis uniserialibus brevioribus sèpe brevissimis, corolla minore. Variat multum fructù magnitudine, ejus formâ plus minus elongatâ, aculeis brevissimis (in iconè citata nimis tenuibus). *E. brachycentrum* Led ab *E. microcarpo* non distingnere valeo.

Ar. Geogr. Songaria, Sibiria Altaica.

10. **E. Sinaicum** (D.C. Prodr. X, p. 141) patule hispidulum, caulis tenuibus flexuosis simplicibus vel parce ramosis, foliis inferioribus oblongis in petiolum attenuatis cæteris ellipticis, racemis paucifloris demum laxissimis basi sèpe foliatis, pedicellis calyce fructifero demum longioribus incrassatis strictis vel sèpius recurvis, calycis fructiferi laciniis reflexis breviter linearibus, corollæ cœruleæ minutæ limbo parvo calycem parum excedente, nuculis cœbre et minute tuberculatis facie angustâ, marginis tubercululis interdum apice glochidiatis ♂. *E. Kotschyi* Boiss. Diagn. Ser. I, 7, p. 29.

Hab. in arenosis deserti Arabiæ petree et horto Conventi Sinaitici (Auch. 2289!), Boiss!), in rupestribus arenosis Persiæ australis ad Persepolin (Ky. 258!), ad Ispahan (Auch 4970!), prope Jezd (Buhse!), in monte Elbrus prope Derbend Persie borealis (Ky. 87!), circa Schahrud (Bge!), montibus Afghauiaæ (Griff!), Belutschiaæ (Stocks!).

Planta pusilla 6-9-pollicaris, racemi demum 3-5-pollicares fructibus remotissimis, pedicelli fructiferi fructu 1½-4-plo longiores, fructus minutus margine non vel vix glochidiatus.

11. **E. semiglabrum** (Ledeb. Fl. Alt. I, p. 204) pallide virens setis sparsis brevibus patentibus basi grosse tuberculatis asperum,

caule superne divaricatum ramoso ramis demum elongatis patentissimis elongatis laxis, foliis oblongis inferioribus basi attenuatis, bracteis oblongo-linearibus calyce longioribus, pedicellis brevissimis demum incrassatis patentibus, calycis fructiferi laciniis breviter et latifuscule linearibus patentibus, corollæ minutæ albidae vel cœruleæ tubo calycem æquanti limbo parvo, nuculis aculeis distinctis longissimis glochidiatis uniserialiter marginatis disco secus lineam medium tuberculato vel aculeolato ⊖. Ic. Ledeb. tab. 28. — *E. Caspium* F. et M. Ind. V, Petrop. non sat differre videtur.

Hab. in desertis Karakum Turkestaniae (Lehm!).

Ar. Geogr. Turcomania, Sibiria Altaica.

12. *E. macranthum* (Ledeb. Fl Alt. I. p. 205) pilis longis tenuibus cinerascens, caule erecto folioso superne paniculato-ramoso, racemis fructiferis strictis elongatis laxis, foliis oblongo-linearibus obtusis caulinis apice saepe recurvis, bracteis oblongis calyce sublongioribus, pedicellis fructiferis erectis calyci subæquilongis calycis laciniis linearibus patentibus, corollæ cœruleæ limbo ampio tubo multo longiore, nuculis serie simplici aculeorum lanceolatorum elongatorum apice glochidiatorum marginatis carinâ faciali aculeolatis vel laeviusculis ⊖. Ic. Ledeb. tab. 29.

Hab. in arenosis Kuwandarja et Syr Darja Turkestaniae (Lehm!).

Ab *E. semiglabro* aculeis longis quoque donato corollis magnis facile dignoscendum.

Ar. Geogr. Sibiria Altaica.

* * * Nuculæ uniserialiter glochidiatæ aculeis basi concretis.

13. *E. marginatum* (M.B. Taur. Cauc. I. p. 120 sub *Myosotide*) pilis saepius adpressiusculis canescens, caule humili superne vel a basi ramoso parce ramoso subpaniculato, racemis fructiferis brevibus laxis, foliis anguste oblongo-linearibus obtusis, bracteis parvis, pedicello fructifero erecto calyce subbreviore, calycis fructiferi laciniis linearibus patentibus, corollæ cœrulecentis limbo minuto tubo breviore, nuculis serie simplici aculeis a basi triangulari attenuatis glochidiatis basi inter se ad tertiam partem longitudinis connatis marginatis facie laeviusculis ⊖. *E. marginatum* Lehm. Asper. p. 130.

Hab. in Caucaso sept. ad fluvium Terek (M.B.), in Kachetiâ et Somchetiâ (Eichw.), Turcomaniâ (Karel.). E ditione nondum vidi.

Ar. Geogr. Rossia australis, Sibiria Uralensis.

14. *E. Bungei*, totum adpressissime canescens, caule elato paniculato ramisque elongatis tenuissimis fructiferis laxissimis, foliis anguste linearibus, bracteis squamiformibus pedicello brevioribus, floribus minutissimis, pedicellis fructiferis erectis fructu triplo brevioribus, corollæ cœruleæ calycem parum excedente, nuculis oblongo-

triangularibus præter discum parce tuberculatum glabris dentibus triangularibus basi inter se coalitis apice glochidiatis versus faciem nuculæ incurvis uniserialiter marginatis ⊙ vel ⊖.

Hab. in Persiâ boreali prope Schahrud (Bge!).

Sesquipedale, tenuissime virgatum, rigidulum, flores e minimis generis, nuculæ in pyramidem vix lineam longam approximatae. Ob nuculas membranâ profunde dentatâ marginatas, mediante *E. marginato* transitum ab *Echinospermo* ad *Paracaryum* præbet, illi tamen omnibus notis adnumerandum.

15. E. sessiliflorum (Boiss. Diagn. Ser. I, 11, p. 124) pumilum adpressissime sericeo-canum, caule tenui simplici vel parce ramoso fere a basi racemos foliosos tenues fructiferos valde elongatos laxos unilaterales edenti, foliis breviter linearibus obtusis, bracteis flores superantibus, floribus sessilibus minimis, corollâ cœruleâ calyce vix longiore, calycis fructiferi subacereti laciniis patentibus, fructu sessili patenti pyramidato. nuculis acute triangularibus undique tuberculatis margine crassiusculo unilateraliter glochidiato-aculeato. nuculæ superioris margine in membranam latiorem calathiformem dentalam interdum dilatato ⊙. *Heterocaryum divaricatum* Stocks.

Hab. in Persiâ ad Isfahan (Auch. 5017!), ad Ssertschah et in toto ambitu deserti magni (Bunge!), in Belutschia ad Doubund (Stocks!).

Floriferum biplicare, fructiferi racemus 4-6-pollicaris, flores et fructus e minimis, nuculæ ut in *E. Vahliano* in pyramidem concretae sed partibiles lineæ $\frac{1}{4}$, vix longæ; altera (superior) interdum membranâ calathiformi obsita nixum ad *Paracaryum* demonstrat. Nuculis heteromorphis ad sectionem *Heterocaryum* accedit sed habitu omnino alienus et nuculæ separabiles

PARACARYUM (Boiss. Diagn. Ser. I, 11, p. 128). — *Omphalodes* Sect. I. — *Pseudanchusa* et Sect. II. *Paracaryi* § 1, atque *Mattia* Sect. 3 d.c. Prodr.

Calyx quinquepartitus. Corolla infundibuliformis tubo plus minus elongato, limbo quinquelobo erecto-patulo. Stamina inclusa antheris ellipticis filamento longioribus, stigma capitatum. Nuculæ stylo pyramidato lateraliter affixa ascendentes disco tuberculatae rarius lèves marginatae membranâ vel inflatâ introflexâ et tunc urceolatae vel calathiformes, vel planâ aut concaviusculâ non introflexâ. — Herbae facie *Cynoglossi*. — Genus *Cynoglosso* corollâ simile, ab eo fructu ut in *Omphalode* et *Mattiâ* membranaceo-marginato distinctum. *P. mucranthum* et *calathicarpum* transitum præbent inter sectiones binas et quoque ad *Echinospermum* cuius species nonnullæ nuculas habent dentibus basi subconnatis marginatas sed non membranâ auctas

Specierum Orientalium distributio.

SECT. I. EUPARACARYUM Boiss. Diagn. Ser. I, 11, p. 129. — Nucu-

lae membranâ marginali inflatâ introflexâ urceolatæ vel cyathiformes.

* Annuæ.

P. calathicarpem, microcarpum, P. micranthum, salsum.

* * Biennia vel perennia.

P. rugulosum, undulatum, strictum.

SECT. II. MATTIASTRUM Boiss. l. cit. p. 130. — Nuculæ membranâ concaviusculâ vel planâ marginatæ.

* Stylus fructu 3-5-plo brevior.

P. myosotoides, Reuteri, lamprocarpum, cristatum, leptophyllum, Ponticum, erysimifolium, Cappadocicum, stenolophum, modestum.

* * Stylus fructu æquilongus vel longior.

× Fornices apice capitato-inflatæ subincurvæ saepe retusæ vel bilobæ.

P. Ancyritanum, Aucheri, calycinum, longipes, asperum.

× × Fornices tridentatæ dentibus lateralibus gibbiformibus intermedio longo lineari.

P. incanum, glastifolium, corymbiforme, angustifolium.

SECT. I. EUPARACARYUM.

* Annuæ.

1. **P. calathicarpum** (Stocks in Hook. Journ. IV (1852) p. 175 sub *Echinospermo*) annum, pilis brevibus basi tuberculatis cinerascens, caule pumilo fere a basi racemos paucos ebracteatos fructiferos elongatos laxos edente, foliis radicalibus oblongo-spathulatis, caulinis breviter oblongo-lanceolatis, pedicellis fructiferis inferioribus calycem æquantibus subpatulis, corollæ cæruleæ minutissimæ limbo tubo subbreviore, nuculis heteromorphis, axili parvâ oblongâ ad discum et marginem angustissimum non membranaceum echinulatâ, cæteris cyathiformibus membranâ introflexâ valde inflatâ radiatim bullatrorugosissimâ circumcirca in tuberculos conicos abeunte marginatis aperturâ centrali minimâ disco parce aculeolato ⊖.

Hab. in Belutschia superiore (Stocks!).

Fructiferum 6-7-pollicare racemis 3-6 pollices longis, flores eis *P. micranthi* non majores, nuculæ anticæ duplo majores cyatho magis clauso membranâque multo magis rugosâ, rugis ad periphæriam in tuberculos conicos elongatis.

Echinospermī speciebus aculeis basi connatis sane affinis, sed facie et membranā marginanti rugoso-bullatā *Paracaryo* adnumerandum.

2. **P. microcarpum** (Boiss. Diagn. Ser. II, 3, p. 139) annum vel biennium, molliuscule et breviter hirsutum cinerascens, caulinibus paniculatis, foliis radicalibus oblongis petiolatis caulinis lanceolatis, racemis bifidis fructiferis elongatis laxis, pedicellis fructiferis calycei aequilongis tenuibus patentि-recurvis, corollæ cœruleæ limbo tubo longiore, nuculis minimis ovatis planis disco parœ echinulato margine introflexo levè tumido cylindrico ⊖.

Hab. in Affghaniâ circa Choky, ad Otipore et in sylvaticis Bharowul (Griff Journ. 50 et 972).

Pedale vel humilius, flores magn. *P. myosotoides*, racemi fructiferi 3-4-pollicares. Insigne nuculis minimis planis lineam vix longis margine convexo sed levissimo non rugoso nec plicato. Facies omnino *Cynoglossi micranthi* Desf. sed pedicelli longiores et fructus diversissimus

3. **P. micranthum** (Boiss. Diagn. Ser. I, 11, p. 129) annum, pilis brevibus basi tuberculatis velutino-asperulum cinereum, caulinibus e collo saepius pluribus pumilis tenuibus erectis et ascendentibus parce ramosis fere a basi racemiferis, foliis radicalibus oblongis obtusis, cæteris oblongo-linearibus, racemis fructiferis elongatis tenuibus laxis unilateralibus, pedicellis fructiferis recurvis calycei aequilongis vel brevioribus, corollæ cœruleæ minime limbo tubum subæquante, nuculis calathiformibus disco echinulatis rarius levibus margine membranâ vel latâ valde introflexâ bullato-rugosâ vel breviore vix introflexâ denticulatâ obsitis ⊖. *O. micrautha* et *O. intermedia* B.C. Prodr. X, p. 159. — *Cynoglossum intermedium* Fresen. Beitr. 169. — *O. glochidiata* et *O. physodes* Bge Lehm. p. 413.

Hab. in arenosis et desertis, Ouadi Sanour et Ouadi Arabah Ægypti superioris (Husson!), valles Sinaiticae (Rüpp., Schimp. 123!, Boiss!), Persia ad Ispahan (Derderian!) et Persia orientalis tota (Buhse! Bge!), Belutschia (Stocks!), Turkestania (Lehm!).

Floriferum 2-3-pollicare, fructiferum semipedale, nuculae diametro sesquilineam latæ marginis formâ et aculeolis mire variantes, saepius membranâ marginali valde inflatâ sphæricæ rarius eâ angustâ concavæ.

4. **P. salsum**, annum, viride pumilum a basi dichotome ramosum tuberculis crassis sparsis ad foliorum margines copiosioribus breviter setiferis asperum, foliis carnosulis oblongis et oblongo-lanceolatis basi attenuatis inferioribus subsessilibus, racemis fructiferis elongatis laxis unilateralibus, corollis minimis cœruleis calycem hispidum vix excedentibus, pedicellis fructiferis brevissimis vel nullis, nuculis globoso-calathiformibus disco echinulatis membranâ inflatâ v. id introflexâ marginé rugulosâ integrâ ore angusto, stylo brevi ⊖.

Hab. ad margines magni deserti salsi Persiae in districtu Veronine ad meridiem Teheran (Bode!).

Tandem semipedale, folia inferiora pollicaria tres lineas lata, racemi fructi-

feri 4-5-pollicares, flores eis *P. micranthi* non maiores sed nuculæ triplo maiores $2\frac{1}{2}$ lineas diam. latae.

* * Biennia vel perennia.

5. **P. rugulosum** (D.C. Prodr. X, p. 161) perenne, e radice verticali simplici cespitoso-pluricaule totum indumento brevi denso albo saepius retrorso velutinum canescens vel canum, foliis inferioribus oblongo-lanceolatis basi longe attenuatis, caulinis lanceolatis et linearibus strictis, caulibus saepius pumilis parce et stricte paniculatis, racemis fructiferis brevibus laxis, pedicellis fructiferis patulis non incrassatis calyces æquantiibus vel inferioribus sublongioribus, corollæ parvæ violaceæ vel purpureæ limbo tubum calyci subæquilonium non æquante, forniciis inclusis ovatis obtusissimis, nuculis sphærico-calathiformibus disco echinulatis vel lævibus, membranâ bullatrorugosâ margine breviter denticulato-introflexâ ♀. *P. rugulosum* Boiss. loc. cit. p. 131. — *Omphalodes Persica* Boiss. Diagn. Ser. I. 7, p. 30 et *Paracaryum Persicum* Boiss. Diagn. I. — *P. rubriflorum* Stocks in Hook. Journ. Bot. IV, 1852, p. 175. — *Omphalodes myosotoides* Fresen. Beitr. 170 non Labill.

Hab. in desertis arenosis lapidosis deserti Sinaitici et in jugo Sinai (Schimp. 401! Auch. 2286! Boiss!), Persiâ australi ad Persepolin (Ky. 240!), Ispahan (Auch. 4969!), orientali ad Rischm, Djendack, Jezd (Buhse!), ad Chabbise et Kerman et prope Ssertschah (Bge!), boreali orientali ad Schahrud (Bge!), Afganîa (Griff!), Belutschiâ (Stocks!).

5. *cyclhymenium* — Margo nucularum magis rugosus extus 1-2 membranis circularibus crispulis eo brevioribus auctus.

Hab. in Persiâ prope Schahrud (Bge!), in monte Sofi prope Ispahan (Bode!), in monte Kohrud prope Kaschan (Derd!), Persiâ australi (Auch. 4971!).

Semipedale, rarius longius, flores et nuculæ eis *P. micranthi* saltem duplo maiores. Notæ quibus plantam Persicam ab Arabicâ distinxeram speciminibus plurimis confertis variables sunt.

6. **P. undulatum** (Boiss. Diagn. Ser. I, 11, p. 129) perenne, virens pilis e tuberculo ortis crebris patulis asperulum, caulibus elatis gracilibus paniculatis, foliis oblongo-lanceolatis acutis undulato-crispis radicalibus in petiolum attenuatis, racemis fructiferis elongatis densiusculis secundis, pedicellis fructiferis calyce subbrevioribus, corollæ cæruleæ limbo tubum calyci subæquilonium non æquante, nuculis sphærico calathiformibus disco tuberculato, membranâ valde bullatrorugosâ ante marginem introflexum denticulatum tuberculatâ ♀.

Hab. in Persiæ borealis monte Elbrus prope Derbend (Ky. 436!), prope Rudbar (Buhse!), in prov. Aderbidjan (Auch. 4972!), ad Seidabad prope Tabris (Bge!).

Pedale, affine *P. ruguloso* a quo distinctum videtur foliis margine undulato-crispis, indumento patentí aspero Stylus inter nuculas brevior.

7. **P. strictum** (C. Koch Linn. 1843, p. 302 sub *Omphalode*)

perenne pilis longis patulis basi tuberculatis asperum, caulinis stricte ramosis, foliis lanceolatis radicalibus basi attenuatis caulinis linearibus acutis strictis, racemis fructiferis elongatis unilateralibus laxis, pedicellis fructiferis erectis calyce subbrevioribus, corollae minuta tubo calyce subbreviori limbo tubum non æquante, nuculis urceolatis disco longe echinatis membranâ inflatâ parum intreflexâ margine valde echinulatâ, stylo nuculis quadruplo breviore ♀. *Omphalodes hirsutus* D.C. Prodr. X, p. 159. — *Paracaryum hirsutum* Boiss. Diagn. II, p. 130.

Hab. in Armeniâ australi ad Karput (Noë! Auch. 2283! et 1936!), in graminis Ssoffdaghs Cataoniæ (Haussk!), Armeniâ Rossicâ ad Araxem (G. Koch!).

Pedale, racemi demum semipedales, a *P. ruguloso* et *undulato* nucularum membranâ marginali breviter nec valde intreflexâ disco longe echinato distinctum; *P. cristatum* nucularum membranâ planitculâ, etc. longius differt.

SECT. II. MATTIASTRUM.

+ Stylus fructu 3-4-plo brevior.

8. **P. myosotoides** (Labill. Decad. II, p. 6, tab. 2 sub *Cynoglosso*) pilis albis longis adpressis basi tuberculatis cinerascens, rhizomate pluricipiti, caulinis ascendentibus pumilis simplicibus vel fere a basi racemiferis, foliis radicalibus oblongis petiolatis, caulinis paucis angustis sessilibus, racemis fructiferis elongatis laxis, corollae cœruleo-violaceæ parvæ limbo tubo subbreviore, pedicellis fructiferis patentirecurvis calyce brevioribus, nuculis disco echinatis rarius levibus membranâ subconcavâ nitidâ margine denticulis horizontaliter curvalis obsitâ rarius integrâ instruetis, stylo nuculâ quintuplo breviore ♂. *Omphalodes myosotoides* Schrank. — *Paracaryum myosotoides* Boiss. loc. cit. — *Omphalodes Cariensis* Boiss. Diagn. Ser. I, 4, p. 41 et *Paracaryum Cariense* Boiss. loc. cit.

Hab. in regione alpinâ montium Sphacioticorum Cretæ 70° (Sibth., Heldr!). Lycie in montibus Akdag et Bereket-dagh (Pestal! Bourg!), Cariæ monte Cadmo (Boiss!), monte Anemas Lycaoniæ (Heldr!), Cyprî monte Troodos (Ky. 741!), Tauro Cilicico 8000' (Ky! Bal!), Libano supra Cedros (Labill! Boiss! Bl!), Hermone 9-9500' (Boiss! Ky. 246!).

Planta 4-6-pollicaris rarius elatior. *P. Cariense* formam typicâ subproceriorum sistit sed characterem nucularum marginis integrâ vel denticulati variabilem esse certior factus fui.

9. **P. Reuteri** (Boiss. et Haussk. Decad. N° 2, p. 6) perenne, adpressissime et dense velutinum canum, rhizomate pluricipiti cespitoso, foliis radicalibus oblongis obtusis in petiolum attenuatis, caulinis parvis strictis oblongo-lanceolatis, caulinis pumilis erectis in racemum simplicem vel fureatum fructiferum laxum brevem abeuntibus, pedicellis fructiferis erecto-patulis fructu æquilongis vel brevioribus, corollâ violacea calyce albo-velutino sesquiflongiore, forniciibus

tubum vix excedentibus ovatis apice inflatis, nuculis disco glochidiatis membranâ planiusculâ nitidâ margine brevissime denticulis horizontaliter curvatis minimis obsitâ instructis, stylo nuculâ triplo breviore ♀.

Hab. in rupestribus regionis alpinæ montis Berytlagh Cataoniae 6000'-40000' (Haussk!).

Planta floritera semipedalis, fructifera subprocerior. flores eis *P. myosotoides* submajores; hanc speciem nuculis refert sed ab eâ et omnibus affinibus foliis ut in *Solenantho lanato* molliter canis differt.

10. *P. lamprocarpum* (Boiss. Diagn. Ser. I, 11, p. 431) perenne, pilis longis albis patulis basi tuberculatis asperum griseum, caulinis elatis, foliis lanceolato-linearibus inferioribus in petiolum attenuatis cæteris acutis, racemo terminali sæpius bifido vel pluribus fructiferis elongatis laxis, pedicellis calyce subbrevioribus, corollâ cœruleâ calyce dense albo-hispido sesquiloniore, fornicibus limbo brevioribus ovatis apice inflatis recurvis, nuculis nitidis levibus membranâ planâ ovatâ latâ margine integrâ vel obsolete repandâ obsitis, stylo fructu quadruplo breviore ♀.

Hab. ad rupes Antilibani ad Sonk Ouadi Barrada inter Zebdani et Damascum (Boiss!), in monte Garbi supra Zebdani 5000' (Ky. 129!).

Planta 1-2-pedalis, folia radicalia 3-5 pollices longa 3-5 lineas lata, flores magn. *P. myosotoides*, nucule diametro 5 lineas latae levissimæ et integrae.

11. *P. cristatum* (Lam. Ill. N° 1800 sub *Cynoglosso*) perenne, pilis tenuibus brevibus e tuberculo ortis adpressiusculis asperulum cineraceens, foliis radicalibus anguste linear-i-oblongis basi attenuatis caulinis strictis linearibus acutiusculis, caulinis apice furcatis interdum ramis axillaribus acutis, racemis fructiferis longis secundis rigidis laxis, pedicellis erectis calyce brevioribus, corollæ mediocris intenue cœruleæ limbo tubo subbreviore, fornicibus ovatis obtusis, nuculis disco valde echinatis membranâ planiusculâ ovatâ echinulatâ et margine in dentes triangulares abeunti obsitis, stylo nuculis quadruplo breviore ♀. *P. cristatum* Boiss. loc. cit. — *Mattia cristata* Don. — *Paracaryum denticulatum* Boiss. in Huet exs.

Hab. in Armenia (Tourn., Auch. 2284!), prope Erzerum (Huet!), Baibut (Bourg!).

Pedale et sesquipedale, membrana mucula 5 lineas longa 4 lata.

12. *P. leptophyllum* (Alph. D.C. Prodr. X, p. 170 sub *Mattia*) bienne, pilis adpressis brevibus e tuberculis majusculis ad margines foliorum lineatim copiosioribus ortis strigillosem, caulinis humilibus parce ramosis, foliis linear-i-lanceolatis inferioribus in petiolum longiusculum attenuatis superioribus basi attenuatâ sessilibus, racemis fructiferis brevibus laxis, calycis adpresso hirsuti faciniis oblongis, corollâ calyce sesquiloniore cœruleâ, pedicellis fructiferis fructu æquilongis

vel brevioribus, nuculis disco echinulatis membranà planiusculà latà margine triangulari-denticulatà, stylo fructu quadruplo breviore ②. *P. leptophyllum* Boiss. loc. cit. — Ic. Jaub. et Spach. III. Orient. bib. 364.

Hab. in Armeniâ (Auch. 2281!).

Planta $\frac{1}{2}$ -4-pedalis, folia cum petiolo 1 $\frac{1}{2}$ -2-pollicaria 2 lineas lata, racemi fructiferi 2-3-pollicares, corolla vix 2 lineas longa, calyx sesquilineum tere longus, nuculae cum membranâ 4 lineas circiter longae et late

13. P. Ponticum (C. Koch Linn. XXIII sub *Omphalodes*) biennie, tuberculis erebris brevissime pilosis asperum, caulinis paniculatis, foliis oblongis radicalibus petiolatis, racemis fructiferis elongatis laxis, calyeis laciniis oblongis, pedicellis erecto-patulis fructu brevioribus, corollâ ..., nuculis disco tuberculatis membranâ planâ margine horizontaliter flexo breviter glochidiato-denticulatâ, stylo fructu quadruplo breviore ②.

Hab. in districtu Sber Ponti Lazizi (C. Koch!).

Ex speciminiibus valde mancis semipedalibus descriptum. Valde affine *P. leptophyllum* pro eius formâ in Pl. Or. Decad. II hoc habui, tamen ob folia breviora latioraque et nuculae dentes marginalis non erectos sed ut in *P. moyosotoide* et *P. calycino* horizontaliter flexos donec melius notum sit seorsim descripti.

14. P. erysimifolium, biennie, pilis sparsis brevissimis adpressis e tuberculo minuto ortis obsitum pallide virens, foliis inferioribus oblongo-lanceolatis obtusis longe petiolatis, superioribus linearibus acutis basi attenuatis, caule puncto paree ramoso, racemis tandem laxis, calyeis adpresso hirsuti laciniis linearibus obtusis, corollâ e sicco carneâ calycem vix excedente, forniciis lobis subaequilongis linearibus apice subinflatis incurvis, pedicellis anthesi peractâ calyce sublongioribus stylo calycem vix excedente, nuculis (junioribus) disco et ad membranâ marginem echinulatis ②.

Hab. in monte Coureoulizos prope Gumushkhané Armenie Furecio Bourg. exst sub *P. corymbiformi* var.).

Semipedalis, facies *P. leptophylli* pro quo habui sed foliis magis oblong. 3-4 lineas lata tuberculis minoribus obsita, calyx duplo longior (tres lineas longus), corolla albida vel carnea calycem non excedens.

15. P. Cappadocicum (Boiss. et Bal. Diagn. Ser. II. 6. p. 124) biennie, pilis brevibus patulis basi tuberculatis asperum, caule elato ample et striete paniculato, foliis oblongis radicalibus in petiolum attenuatis obtusis, caulinis strictis acutiusculis, racemis fructiferis longis laxis, pedicellis fructiferis erecto-patulis calyce sublongioribus, corollis parvis azureis limbo a calyce subexerto, forniciis ovatis obtusissimis, nuculis ovatis disco glochidiato-aculeatis membranâ concaviuseculâ margine minute denticulatâ instructis, stylo fructu quintuplo breviore ②.

Hab. in monte Alidagh Cappadociae 5500' (Bal!).

Planta 2-2 1/2-pedalis, racemi fructiferi semipedatales, flores eis *P. myosotoides* submajores, nuculae paulo minores structuræ similes sed denticuli marginales conici recti nec horizontales; ab eo cæterum radice bienni et caule procero paniculato alienum.

16. ***P. stenolophum*** (Boiss. Diagn. Ser. II, 6, p. 124) bienne, pilis tenuibus patulis hirsutum, caule elato paniculato, foliis inferioribus oblongo-lanceolatis petiolatis superioribus anguste linearibus acutis, racemis fructiferis filiformibus erecto-patulis calyce sublongioribus, nuculis lœvibus membranâ concaviusculâ subintegrâ angustâ marginatis, stylo fructu quadruplo breviore ②.

Hab. in Cappadociae collibus ad occasum Cæsareæ 3600' (Bal!).

Bipedale, facie *P. Cappadocico* simile sed pedicelli tenuissimi sublongiores, nuculae minores margine duplo angustiori integro, corolla minor. An nihilominus hujus varietas?

17. ***P. modestum*** (Boiss. et Haussk. Pl. Or. Dec. 2, p. 5) bienne, pilis patulis basi tuberculatis hirsutum, caulis tenuibus ascendentibus brevibus 2-3-racemosis, racemis brevibus 3-7-floris fructiferis laxis, foliis angustis acutis inferioribus oblongo-lanceolatis longe petiolatis caulinis lanceolato-linearibus, corollâ intense cæruleo-violaceâ tubo calycem subæquante limbo tubo æquilongo, fornicibus ovatis inclusis bilobis, calycis laciniis oblongo-lanceolatis, pedicellis fructiferis patulis tenuibus fructu æquilongis et inferioribus longioribus, nuculis disco glochidiatis membranâ latâ planâ suborbiculari intra marginem repando-subdenticulatanam lineâ concentricâ tuberculatâ obsitâ, stylo fructu quintuplo breviore ②.

Hab. in graminosis prope Nackschi Timur prope Kaserun Persiæ australis 3000' (Haussk.).

Caules cum racemis 6-8-pollicares, folia 2-3-lineas lata, corolla 2 1/2 lineas longa limbo latiusculo.

* * Stylus fructu æquilongus vel longior.

18. ***P. Ancyritanum***, bienne, pilis brevibus adpressis e tuberculo ortis obsitum pallide virens, foliis inferioribus oblongo-lanceolatis obtusis longe petiolatis superioribus acutis, caule parce ramoso, racemis fructiferis laxis, calycis adpresso hirsuti laciniis linearibus obtusis, corollâ e secco carneâ vel rubellâ calyce sesquilociore, fornicibus corollam æquantibus linearibus apice inflatis obtusis, pedicellis calyce sublongioribus, nuculis (junioribus) disco et ad membranæ planiusculâ marginem echinulatis, stylo fructu longiore ③.

Hab. in Anatoliâ boreali prope Angora et Kastamouni (Wied!, Auch. 4978 A).

Planta 1-2-pedalis, facies et folia *P. erysimifolii* sed corolla calyce longior, stylus e corollâ exsertus fructu longior. Calyx 3 lineas, corolla 4-4 1/2 longa.

19. ***P. Aucheri*** (Alph. D.C. Prodr. X, p. 169 sub *Mattiæ*) bienne?

totum pilis longis tenuibus adpressis vix tuberculatis molliter canescens, foliis radicalibus linear-i-oblongis longe petiolatis caulinis strictis longe linearibus acutiusculis, racemis 2-4 subcorymbosis demum elongatis laxis secundis, calycis adpresso hirsuti laciniis linearibus obtusis fructiferi elongatis patulo-deflexis, corollæ cœruleæ calyce parum longioris tubo calyce breviore, fornicibus lobos corollæ æquantibus linear-i-lanceolatis apice capitato-inflatis, pedicellis fructiferis calycis longitudinis, nuculis (junioribus) disco securus lineam medium tuberculatus membranâ planâ margine integrâ, stylo fructu æquilongo ②. Ic. Jaub. et Spach III. Or. tab. 365.

Hab. in montibus Cariæ supra Moglah (Auch. 2295).

Sesquipedale, folia 2½-3 lineas lata, caulina vix abbreviata 1½-2-pollicaria, corolla tres lineas longa. Species indumento tenui molli *Cynoglossi clandestini* et nuculæ membranâ margine integrâ et sparsim glochidiata ab affinibus distincta.

20. P. calycinum (Boiss. et Bal. Diagn. Ser. II, 6, p. 125) bienne, tuberculis crebris brevissime et adpresso pilosis asperulum, caule elato paniculato-ramoso, foliis inferioribus oblongis in petiolum attenuatis rameis linearibus, racemis demum elongatis laxis secundis, calycis fructiferi laciniis valde elongatis linearibus deflexis, pedicellis erecto-patulis inferioribus fructu sublongioribus. corollæ cœruleæ tubo calyci æquilongo limbo dimidio breviore, fornicibus corollæ lobos æquantibus linearibus apice incurvis capitato-inflatis, nuculis disco parce echinatis membranâ latâ planâ margine horizontaliter flexo minute denticulatâ, stylo fructu æquilongo vel longiore ②.

Hab. in Cappadociâ ad radices Argæi et in collibus prope Cæsaream 3600' (Bal!).

Pedale et sesquipedale, corolla 4 lineas longa, nuculae 4-5 lineas longæ. A *P. leptophyllo* et *P. Pontico* affinibus præter stylum longum calycis fructiferi laciniis longis reflexis tres lineas longis distinctum.

21. P. longipes (Boiss. Diagn. Ser. II, 6, p. 126) bienne. pilis adpresso usculis longis basi tuberculatis hirsutum, caule elato paniculato, foliis lanceolato-linearibus inferioribus in petiolum attenuatis superioribus anguste linearibus acutis, racemis fructiferis laxissimis tenuibus flexuosis, calycis adpresso hirsuti laciniis breviter linearibus, corollâ intense cœruleo-violaceâ calyce vix sesquilongiore, fornicibus loborum basin non attingentibus linearibus truncato-retusis, pedicellis fructiferi arcuatis fructu duplo longioribus, nuculis disco parce echinatis membranâ latâ planâ nitidâ margine crebre crispulodenticulatâ, stylo fructu æquilongo ③.

Hab. in monte Alidagh Cappadociae 4200' (Bal!).

Bipedale, folia 4-5 lineas lata, racemi fructiferi 6-8-pollicares, calyx 2 lineas, corolla 3 lin. longa, pedicelli fructiferi 6-8 lineas longi.

22. P. asperum (Stocks Hook. Journ. Bot. IV (1852), p. 175)

bienne, pilis rigidis patulis e tuberculo ortis asperum cinereum, caulinibus erectis paniculatis, foliis lanceolatis aculis inferioribus margine subundulatis in petiolum longum attenuatis superioribus linearibus acutis, racemis fructiferis elongatis laxissimis, calycis laciñiis anguste lanceolatis, corollæ purpureæ-violaceæ tubo calyce subduplo longiore limbo tubo duplo breviore, fornicibus ad medianam corollam bilobis, pedicellis fructiferis erectis fructu brevioribus, nuculis disco aculeatis membranâ planiusculâ latâ margine grosse dentatâ dentibus glochidiatis, stylo fructu vix breviore ②.

Hab. in Belutschia superiore (Stocks 906!), Affghaniâ ad Choky (Griff. 491!).

Planta sesquipedalis, folia radicalia 2-3-pollicaria 4-5 lin. lata, corolla 5 lineas longa. Species tubo corollæ calycem superante insignis.

23. *P. incanum* (Ledeb. Fl. Ross. III. p. 173 sub *Mattia*) adpresso tomentosum incanum, caule elato stricte folioso paniculato-subcorymboso, calyeis valde albo-tomentosi laciñiis linearibus, corollâ calyce sesquilociore e sicco purpureo-violacea, fornicibus infra medianam partem utrinque gibbis inde abrupte attenuatis, pedicellis inferioribus jam anthesi vix peractâ calyce pluries longioribus nuculis (junioribus) disco echinatis margine denticulatis, stylo e corollâ longe exerto ④?

Hab. in Transcaucasiâ ad fines Turcicas (Nordm!), in Anatoliâ boreali ad Angora (Wied!).

Planta etiam florifera sesquipedalis, folia inferiora 2-2½-pollicaria 3-4 lineas lata, corolla magna sex lineas longa, pedicelli infimi jam corollâ vix delapsa 1-1½-pollicares. Species ab affinibus tomento cano distinguenda

24. *P. glastifolium* (Willd. Sp. 764 sub *Cynoglosso*) glabrum totum tuberculis depressis obsitum, caule elato superne paniculato, foliis lanceolatis margine denticulato-scabris inferioribus in petiolum attenuatis, caulinis sessilibus acutis, calycis laciñiis linearibus obtusis margine breviter ciliatis, corollâ cœruleâ calyce subduplo longiore, pedicellis fructiferis fructu longioribus, nuculis latis ovato-subcordatis disco levibus vel 2-3 aculeis obsitis membranâ planâ latâ integrâ, stylo e corollâ exerto fructu longiore ②. *P. glastifolium* Boiss loc. cit. — *Mattia glastifolia* Don — *Rindera glastifolia* R. et Sch.

Hab. in Armeniâ (Tourneft). Vid. frustul. in Willd. herb.

Inter affinia tuberculis totius plantæ glabris nec setiferis notabile, ex specimine valde manco herb. Willd. mihi tautum notum.

25. *P. corymbiforme* (D.C. sub *Mattia*) bienne, pluricaule pilis brevibus adpressis basi tuberculatis canescens asperulum, foliis inferioribus linearí-oblongis in petiolum attenuatis caulinis linearí-lanceolatis strictis, racemis fructiferis laxis corymbosis erecto-patulis, calycis hirsuti laciñiis linearibus fructiferis deflexis non auctis, pedicellis gracilibus patentibus fructu longioribus, corollâ violaceo-rubellâ

calyce quartâ parte longiore, forniciis infra medium utrinque gibbosis dente intermedio subulato longissimo, nuculis disco echinatis membranâ latâ planâ margine obsolete tuberculatâ, stylo fructu longiore ②. *P. corymbiforme* Boiss. loc. cit.

Hab. in Armeniâ (Anch. 2292! et 2329!), ad Erzerum (Calv!), in Ponto australi inter Agdja et Chabbane Karahissar (Tchih. 748!).

Pedale, folia 4-5 lineas lata, corymbus sub anthesi contractus, flores sex lineas longi, racemi fructiferi 3-4-polligares pedicellis inferioribus 8-9 lineas longis, nuculae 5 lineas longae. Affine *P. glastifolium* sed tuberulii non mutici, inflorescentia corymbosa, corolla calyce non duplo longior.

26. **P. angustifolium** (Willd. Sp. I. p. 763 sub *Cynoglosso*) perenne cespitosum, pilis longis adpressis non tuberculatis canescens saepe sericeum, foliis radicalibus confertis linearibus et linear-oblongis longe petiolatis, caulinis paucis anguste linearibus, caulinibus pumilis, racemis 2-3 corymbosis fructiferis brevibus laxis, calycis laciniis latiuscula oblongo-linearibus, corollâ azureâ calyce plus duplo longiore, forniciis apice tridentatis dentibus lateralibus gibbiformibus intermedio linearis duplo longiore, pedicellis fructiferis tenuibus erecto-patulis fructu 2-3-plo longioribus, nuculis disco tuberculatis vel laeviusculis membranâ late ovata planâ nitidâ integrâ, stylo fructu longiore ♀. *P. angustifolium* Boiss. loc. cit. — *Cynoglossum emarginatum* Lam. Ill. 1799. — *Cynoglossum racemosum* Schreb. — *Paracaryum azureum* Boiss. et Heldr. Diagn. Ser. I. II. p. 131. — *P. erigerifolium* Schott in Ky. Sehed. — *P. corymbiforme* Tchih. As. Min. tab. 21 non Boiss.

Hab. in collibus subalpinis et alpinis Lycaonie (Heldr!), Galatiæ (Wied!), ad Sivas (Noë!), Tauri Cilicici 8000' (Ky!). Cappadocie (Ky. 180!), Armeniae (Tourn.), ad Balbut et Gumuschkhané (Bourg!).

Affine *P. corymbiforme* differt indumento, caulinibus pedem dimidium raro superantibus, corymbi ramis paucis, corollâ calyce plus duplo longiore, fornicium formâ. Species elegans.

CYNOGLOSSUM (L. Gen. 483).

Calyx quinquepartitus. Corolla infundibuliformis tubo calyci subæquilongo limbo quinquelobo patenti. Fornices faucem claudentes. Stamina inclusa antheris oblongis. Stigma integrum vel retusum. Nuculae 4 ovatae depresso-undique echinato-glochidiatae margine erassimulo echinato cinctæ vel immarginatae. — Herbæ.

1. **C. Sphacioticum** (Boiss. et Heldr. Diagn. Ser. I. II. p. 125) perenne molliter hirsutum cinerascens, rhizomate squamoso ramoso, caulinibus nanis in racemum simplicem vel bifidum brevem pauciflorum laxum abeuntibus, foliis parvis inferioribus oblongis in petiolum breve attenuatis cæteris a basi semiamplexicauli oblongo-linearibus, corollâ atroviolacea calyce albo-hirsuto sesquilociore, pedicellis erec-

tis fructu subæquilongis, nuculis plano-convexis immarginatis undique dense glochidiatis ②. Ic. Raulin Descr. Cret. tab. 16.

Hab. rarum in summo cacumine Stravopodia montium Sphacioticorum Cretæ 7000' inter saxa (Heldr!).

Planta 3-4-pollicaris habitu *Paracaryi myosotoidis*, folia majora cum petiolo 10 lineas longa, racemi fructiferi sesquipolligares, nuculæ eis *C. officinalis* subminores.

2. ***C. officinale*** (L. Sp. 192) biennne, caule elato paniculato, foliis tomento tenui molli canescentibus acutis, inferioribus ellipticis in petiolum attenuatis caulinis strictis a basi semiamplexicauli lanceolatis, racenis fructiferis erectis brevibus laxiusculis unilateralis ebracteatis, corollâ sordide rubrâ, pedicellis fructu subbrevioribus arcuatis, nuculis facie plano-depressis margine crassiusculo prominulo cinctis glochidiatis ②. Ic. Engl. Bot. tab. 921. — Rchb. tab. 129.

Hab. in incultis, ad vias, Græcia (ex Sibth.), Byzantium (ex Bergg.), Anatolia borealis ad Samsun (Tchih!), Pontus Lazicus ad Rhizé (Bal!), Armenia Turcica ad Baibut (Bourgl.) et Erzerum (Calv!), Tauria (Stev.), Caucasus totus 800'-4500' (C.A. Mey.), Imeretia, Iberia, ditio Talyisch (Ledeb.), Persia bor. (Szov! Buhse!).

Ar. Geogr. Europa fere omissis australioribus exceptis à Scandinaviâ ad Rossiam, Sibiria tota, America borealis.

3. ***C. holosericeum*** (Stev. Mem. Mosq. III, p. 255) biennne cespitosum, caulis elatis, foliis acutis adpresso-sanguineâ et tenuiter canis radicalibus crebris longe lanceolatis, caulinis strictis abbreviatis a basi semiamplexicauli oblongo-lanceolatis acuminatis, floribus dense capitatis, racemis paucifloris fructiferis brevissimis subcorymbosis, corollâ calycem parum superante purpureo-sanguineâ post anthesin sordide violaceâ, pedicellis fructu æquilongis et brierioribus erecto-patulis, nuculis planis canescentibus aculeis glabris glochidiatis undique adspersis margine acuto angusto ②.

Hab. in Caucaso orientali inter Chinalag et Wanda (Stev!), ad moles glaciales Djultidagh 8500'-9600' (Rupr!).

Cespites latos formans, sesquipedale. folia radicalia 6-8-pollicaria 5-7 lineas lata, corolla eâ *C. officinalis* subbrevior, racemi fructiferi 1-2-pedales fructibus 3-5, terminalibus saltem confertis.

4. ***C. montanum*** (Lam. Dict. II, 237) biennne viride, caule elato paniculato-corymbose, foliis membranaceo-pellucidis supra glabris subtus pilis brevibus sparsis basi tuberculatis asperulis inferioribus ellipticis breviter petiolatis, cæteris sessilibus basi semiamplexicaulis intermediis supra basin dilatatam angustatis, corollâ violaceo-cærulescenti. racemis fructiferis longis laxiusculis ebracteatis, nuculis undique glochidiatis plano-convexis margine obscurâ aculeis basi latiore inter se subconnatis constante ②. Engl. bot. tab. 1642. Rchb. tab. 132. — *C. pellucidum* Lapeyr. — *C. sylvaticum* Hænke.

Hab. in sylvis montanis, mons Ilghasdagh Anatolice borealis (Wied!), Taurice mons Babugan (Stev!), Caucasus orientalis 2400'-3600' (C.A. Mey!), ditio Karabagh et Persiae prov. Ghilau (Buhse!).

Ar. Geogr. Britannia, Germania, Gallia, Helvetia, Italia borealis et media, regio Danubialis.

5. C. Nebrodense (Guss. Prodr. I, p. 216) biennie, pallide virens pilis interdum basi tuberculatis pubescens vel hirsutum, caule paniculato, foliis inferioribus oblongis petiolatis, caulinis a basi semiamplexicauli lanceolatis acutis, corollæ rubræ, racemis fructiferis elongatis laxiusculis, pedicellis deflexis fructu subbrevioribus, nuculis parvis immarginatis aculeis glochidiatis basi dilatata confluentibus crebre obsitis ①. Ic. Rehb. tab. 131. — *C. pustulatum* Boiss. Elench. — *C. Diocoridis* Ledeb. Fl. Ross. III, p. 166! — *C. parrifolium* C. Koch Linn. XXII, p. 645.

Hab. in rupestribus subalpinis et alpinis, mons Chelmos Peloponnesi (Orph!), Anatolia borealis ad Mersivan et Tokat (Wied!). Lydia et Caria in montibus Tmolo, Mesogi, Cadmo (Boiss!), Pisidia (Heldr!), Lycia (Bourg!), Ciliciae ditione Kassan Oglu (Ky!), Syrie monte Cassio (Boiss!) et Libano ad Cedros (Boiss! Bl!), Antilibano (Gail!), Cataoniae Berytdagh (Haussk!), Cappadociæ monte Aslandach (Bal!), Armeniæ ad Gumuschkhané (Bourg!), Ponto Lazico (Bal!), Caucaso ad Pâtigorsk (Ledeb.), Persiæ boreali ad Siaret (Bge!).

β. areolatum. — Nuculæ areola laevis vel glochidiis paucis obsita, margo aculeatus.

Hab. in Daghestaniæ prope Schoralo 6000' (Rupr!).

Species affinis *C. montano* a quo hirsuti e majori, foliis angustioribus supra basin non angustatis, nuculis minoribus differt, sed nimis affinis *C. Diocoridis* quod foliis etiam radicalibus angustioribus linear-lanceolatis tantum differt et forsitan hujus varietas.

Ar. Geogr. Hispania australis, Sicilia, Africa borealis.

6. C. pictum (Ait. Kew I, p. 179) biennie, tomento tenui molli canescens, caule elato paniculato, foliis inferioribus oblongis obtusis longe petiolatis, caulinis approximatis strictis a basi cordato-semiamplexicauli lanceolatis, corollæ pallide cœruleæ reticulatim pictæ, racemis fructiferis elongatis densiusculis, pedicellis reflexis fructu brevioribus, nuculis plano-convexis immarginatis undique crebre glochidiato-aculeolatis ②. Ic. Rehb. tab. 130. — *C. cheirifolium* Scop. — *C. Creticum* Vill.

Hab. in derelictis, ad vias totius regionis inferioris a Græciâ et ejus insulis! Macedonia et Thraciâ (Griseb.), ad Anatoliam maritimam borealem! occidentalem et australem! Cretam (Raul!), Rhodum (Bourg!), Syriam littoralem (Bl! Gail!), Palestinam (Boiss!), Tauriam (Stev.), Transcancasiam (M B!, Hohen!).

Ar. Geogr. Regio mediterranea totius Europæ, Africæ borealis, insulae Canarienses.

7. C. Columnæ (Bivon. Manip. II, p. 3) biennie, molliter tomentosum canescens vel canum, caule superne parce ramoso, foliis infe-

rioribus petiolatis elliptico-oblongis, caulinis a basi dilatata cordato-semiamplexicauli breviter lanceolatis acutis, corollæ rubræ tubo calyce tomentoso breviore, racemis ebracteatis fructiferis elongatis densiusculis, pedicellis erecto-patulis fructu brevioribus, nuculis magnis areola planâ aculeolata marginè elevato denticulato-glochidiato cinctâ ②. Ic. Ten. Fl. Nap. tab. 116.

Hab. ad vias, in regione inferiori montium Peloponnesi, in Laconia, Arcadiâ, Argolide (Boiss!), ad radices Kyllenes (Orph!), Pentelici et montis Onion (Heldr!), Delphi Eubeæ (Orph!). Ætolia (Nieder!), Zacyntho (Marg!), Syra (Orph!), Cretâ (Sieb. exs. sub *C. cheirifolio*), ad Cydoniam (Heldr!), Asia minore (Auch. 2290!).

Racemis ebracteatis a *C. cheirifolio* statim distinguendum.

Ar. Geogr. Sicilia, regnum Neapolitanum, Dalmatia.

8. C. macrostylum (Bge Pl. Lehm. p. 412) bienne, caule elato patentim hispido paniculato, foliis viridibus pilis sparsis adpressis basi tuberculatis strigillosis radicalibus oblongo-lanceolatis longe petiolatis, caulinis lanceolatis breviter petiolato-attenuatis, racemis brevibus laxis paucifloris bracteolatis, pedicellis calyce longioribus demum recurvis, corollæ pallide violaceæ tubo calycem adpresso canescentem æquante limbo tubo subdupo longiore, nuculis subdepressis undique dense glochidiatis marginatis, aculeis marginis pluriserialibus longioribus basi confluentibus ②.

Hab. ad fossas ad Katte Kurgan inter Buchara et Samarkand Turkestaniae et in vallis montis Karatau (Lehm!). Vid. frustulum absque flore et fructu.

Tantum ex speciminibus autumnalibus mancis notum, ut videtur elatum; folia radicalia ampla cum petiolo sæpe pedialia, flores laxè cymoso-racemosi, corolla 5 lineas longa, fornices obtuse trilobæ, nuculae 2 lineas late. Species a congeneribus ditionis nostræ racemis bracteolatis atque limbo tubi proportione longiore distincta, ex cl. auctore *C. anchusoidi* Indico affinis.

OMPHALODES (Tourn. Inst. 140. — *Omphalodes* Sect. *Paracaryi* § 2. — Sect. *Maschalanthus* et *Euomphalodes* D.C. Prodr.).

Calyx quinquepartitus. Corolla rotata tubo brevissimo fauce clausâ, forniciis obtusis. Stamina inclusa antheris ellipticis filamento longioribus. Stylus apice capitatus. Nuculae depressæ marginatæ vel calathiformes horizontales puncto subbasili styli basi affixæ. — Herbæ.

1. O. Cappadocica (Willd. Sp. I, p. 767) perennis, rhizomate repenti fasciculos foliorum et caules floriferos hinc inde edente, foliis minute sparsim et adpressissime hirtis viridibus radicalibus amplis longe petiolatis ovato-cordatis acuminatis arcuatim 5-7 nerviis ad nervos sericeis, caulinis floriferis ascendentibus tenuibus inferne foliis paucis parvis subsessilibus ovato-oblongis subuninerviis instrutis in racemum longum laxum nudum abeuntibus, pedicellis calyce

longioribus filiformibus demum refractis, calycis fructiferi valde accreti patentis lacinias ovato-oblongis acutis, corollæ cœruleæ lobis orbiculatis calyce triplo longioribus, nuculis calyce multo minoribus puneto basali affixis cyathiformibus sphærico-depressis margine introflexo valde dentato ♀. *O. Cappadocica* D.C. Prodr. X, p. 161. — *Cynoglossum Omphalodes* var. β Lam. — *Omph. cornifolia* Lehm. — *Omph. Orientalis cornifolio* Tourn. Cor. p. 7. — *O. Wittmanniana* Stev. Bull. Mosq. 1851, p. 607.

Hab. in umbrosis, Pontus Lazicus ad Rhizé (Bal!), Cappadocia (Tournef.), Guria (Nordm!), Abchasia (Wittm.).

Foliorum radicalium lamina 3-3 $\frac{1}{2}$ -pollicaris, corolla eam *O. vernæ* aquans a quâ foliorum nervis arcuatî sericeis, racemis elongatis, carpellorum membranâ introflexâ dentatâ, etc., eximie differt.

2. *O. rupestris* (Rupr. MSS.) cespitosa adpressissime et breviter sericeo-pubescentis, foliis parvis omnibus petiolatis radicalibus a basi truncatâ oblongis obtusiusculis subuninerviis petiolo longo. caulinis 1-2 minoribus ovatis obtusis petiolo laminæ aequilongo. caulinibus ascendentibus nanis folia radicalia vix superantibus paucifloris, pedicellis filiformibus longissimis, calycis sericei lacinias ovatis, corollæ cœruleæ laminâ calyce triplo longiore, nuculis.... ♀.

Hab. in fissuris rupium ad Bolta Caucasi inter Wladikaukas et Tiflis 2500' (Rupr!).

Planta bipollicaris, folia majora 9-10 lineas longa, flores magn. *Omph. Cappadociae* a quâ differt foliis subuninerviis obtusis non sericeo-multinerviis, caulinis ad racemi basin petiolatis.

3. *O. Luciliæ* (Boiss. Diagn. Ser. I, 4, p. 44) dense cespitosa glabra viridis glauco-virens, rhizomate crasso folia radicalia racemosque simplices ascendentis foliatos demum laxos edente, foliis calcareo-punctatis subuninerviis radicalibus longe petiolatis a basi rotundata vel cuneatâ ovatis vel oblongis obtusiusculis, foliis caulinis diminutis oblongo-ellipticis sessilibus utrinque attenuatis acutis, pedicellis folio axillari longioribus fructiferis valde elongatis arcuato-recurvis, calycis fructiferi aucti lacinias ovato-oblongis acutiusculis, corollæ cœruleæ limbo calyce 4-5-plo longiore, nuculis ovatis depressis ventre puncto affixis scrobiculato-rugosis dorso concavis levibus linea longitudinali bipartitis margine brevissimo erecto non introflexo integro ♀. Ic. Jaub. et Sp. Ill. tab. 366.

Hab. ex fissuris parietum verticalium erumpens et cespites sa-pe latissimus compactos formans in regione alpinâ, Græcia in Parnasso supra Mana Arachova (Orphl), supra Gourna (Guicci!), Anatolia in monte Sipylo Lydie (Auch! Bal!), Cadmo supra Geyra et supra Khonas (Boiss!), Tauro Cilicie (Ky! Bal!).

Foliorum radicalium lamina 9-15 lineas longa, caules floriferi cum racemo 2-3-pollicares fructiferi semipedales, corolla eâ *O. vernæ* duplo major, nuculæ fere 2 lineas longæ.

SOLENANTHUS (Ledeb. Fl. Alt. I, p. 193).

Calyx quinquepartitus. Corolla tubulosa lobis obtusis erectis. Fornices obtusæ sæpius villosæ. Stamina 5 filamentis infra medium corollam insertis longis sæpius longe exsertis. Antheræ oblongæ integræ. Stigma punctiforme. Nuculæ 4 ovatae depresso-echinato-glochidiatae, vel marginatae areola concavæ, vel convexæ immarginatae. — Herbae facie *Cynoglossi*. — Genus fructu *Cynoglosso* simile, corollæ ab eo distinctum. Species binæ priores ex speciminibus incompletis notaæ staminibus inclusis a cæteris discrepant et forsitan ad *Cynoglossum* revocandæ erunt.

* Stamina corollam non excedentia. Corollæ lobi oblongo-spathulati.—Species a *Cynoglosso* corollæ tubulosa tantum distinctæ.

1. S. brachystemon (F. et Mey. in Hohen. Talysh Enum. p. 76) virens, villis elongatis sparsis lanuginosus, rhizomate elongato, caule parce ramoso, foliis inferioribus oblongo-lanceolatis semiamplexicaulibus, racemis longe pedunculatis capitatis etiam post anthesin brevibus confertifloris, pedicellis calyci subæquilongis, calycis lanuginosi laciniis oblongo-linearibus obtusis, corollæ calyce vix longiore tubulosâ lobis oblongo-spathulatis obtusissimis, forniciis ad faucem oblongis obtusis glabris, filamentis corollæ æquilongis, antheris oblongis, nuculis marginatis echinato-glochidiatis ② vel ♀.

Hab. in sylvis prope Lenkoran ad Caspium sub *Cynoglosso stamineo* (Hohen!), ad Rescht Persiae borealis maritinae (Buhse!).

Planta bipedalis, folia radicalia 6-8-pollicaria sesquipollicem lata longe petiolata, racemi 2-3 axillares capitatum 8-10-flori, calyx sub anthesi 4 lineas longus.

2. S. eriocalycinus (Boiss. et Buhse Aufz. p. 157) viridis, praeter flocculos sparsos deciduos glaber, caule crebre et stricte folioso, foliis obtusiusculis oblongo-lanceolatis inferioribus in petiolum longe attenuatis superioribus late linear-lanceolatis semiamplexicaulibus, floribus in capitulum terminale foliis supremis involucratum conglobatis, pedicellis calyce subbrevioribus, calycis dense lanati laciniis linearibus obtusis, corollæ tubulosâ calyce sesquilongiore laciniis oblongo-linearibus obtusis, forniciis ad plicas versus faucem sitas reductis, antheris linearibus corollam æquantibus, fructu..... ♀.

Hab. in montibus ditionis Transcaucasicae Talysh ad Massula 6000' (Buhse!).

Pedalis, folia inferiora 3-4-pollicaria 15-20 lineas lata longe petiolata, calyx 4 lineas, corolla 5½ longa. Affinis *S. brachystemone* a quo differt glabritie, forniciis ad plicas reductis, antheris duplo longioribus.

* * Stamina longe exserta. Corollæ lobi oblongo-lineares.

+ Folia basi attenuata.

3. S. Biebersteinii (D.C. Prodr. X, p. 165) biennis, molliter pubescens vel canescens, caule crasso elato sulcato folioso, foliis radicalibus ovatis longe petiolatis, caulinis elongatis anguste oblongo-lanceolatis in petiolum breviorem attenuatis, racemis numerosis sub anthesi capitato-scorpioideis paniculam foliosam strictam thyroideam formantibus, calycis lanuginosi laciniis oblongis obtusis, corollæ calyce vix longioris (albido-rubellæ?) lobis oblongis obtusis, forniciis oblongo-lanceolatis villosis supra medium corollam insertis, staminibus longe exsertis, nuculis facie anticâ areolatis planis paree glochidiatis margine elevato rotundato pluriserialiter aculeolis confluentibus glochidiato ②. *Cynoglossum stamineum* M.B. III, p. 127 non Desf. — *Solenanthus dubius* Fisch. et Mey. in Talyseh Eium, p. 76.

Hab. in umbrosis montium Taurie meridionalis in valle sup. fluv. Salgir inter Kisilkoba et Kurnosen (Stev. in D.C. herbt.).

Sesquipedalis, fol. radicalium limbus 6-pollices longus, calyx 3 lineas, corolla glabra nec ut cl. Bieb. pubescens $3\frac{1}{2}$ longa. Valde affine *S. Apennino* quod tamen ut videtur foliis caulinis angustioribus acuminatis, nuculis sub-majoribus areola depressiore, margine magis elevato a nostro differt.

4. S. mollissimus (Lehm. Asper. p. 165 sub *Cynoglosso*) biennis, caule elato crasso piloso folioso, foliis supra glabriusculis subtus molliter canescens radicalibus ovato-oblongis in petiolum attenuatis, caulinis lanceolatis utrinque attenuatis, racemis numerosis sub anthesi capitato-scorpioideis paniculam ovatam foliosam formantibus, calycis hirsuti laciniis lanceolatis, corollæ (albidæ?) calyce vix longioris lobis oblongis obtusis, forniciis oblongo-lanceolatis villosis infra medium corollam insertis, staminibus breviter exsertis, nuculis.... ②. *S. mollissimus* D.C. Prodr. X, p. 166. — *Cynoglossum Gileadense* Willd. herb! (spahmate pro *Ghilanico*).

Hab. in montibus prov. Ghilan Persiae (Pall!). (Vid. in herb. Berol.)

Simillimus *S. Biebersteinii* a quo staminibus brevius exsertis et forniciis inferioris corollæ insertis tantum differt. An eadem species? Nucule ignotæ et donec immotuerint sub judice lis est.

5. S. Tournefortii (D.C. Prodr. X, p. 164) molliter et adpresso canescens-velutinus, caule folioso crassiuseculo breviter paniculato, foliis parallele trinerviis inferioribus oblongis obtusis basi longe attenuatis petiolatis, caulinis strictis linearis-lanceolatis basi semiamplexi-cauli subdecurrentibus, racemis floriferis brevissimis densifloris fructiferis elongatis, calycis albo-tomentosi laciniis oblongo-linearibus, corollæ calyce sublongiore rubro-cærulescenti lobis triangularibus, forniciis ovatis sub medio tubo sitis, filamentis corollæ plus duplo longioribus, nuculis aculeis glochidiatis basi dilatatis uniformiter obs-

tis margine indistincto ♀. *S. conglobatus* Alph. d.c. Prodr. X, p. 166 (specimen junius). — *Cynoglossum stamineum* Desf. Cor. p. 28, tab. 19. — *Mattia staminea* R. et Sch. — *Moltzia Libanotica* Zucc. Abhandl. Monach. III, p. 246, tab. 3.

Hab. in rupestribus regionis subalpinæ et alpinæ, Armenia Turcica (Huet!), Cappadocia ad Euphratem (Auch. 2293!), Berytdagh Cataoniae 8000' (Haussk!), Taurus Cilicus (Bal!) et ditio Kassan Oghlu 6000' (Ky. 93!), Libanus ad Hasrun (Bl!), Antilibani vallis Mandschura (Ky!), Persia borealis in monte Ssahend (Buhse!) et monte Totschal prope Teheran (Ky. 200!).

+ + Folia basi cordata.

6. *S. circinnatus* (Ledeb. Fl. Alt. I, p. 194) viridis pubescens, caule elato folioso, foliis membranaceis radicalibus oblongis subcordatis longe petiolatis, caulinis oblongo-lanceolatis acutis sensim diminutis basi rotundatâ semiamplexicaulibus, racemis brevibus striete et longe paniculatis fructiferis elongatis laxiusculis, calycis hirsuti laciniis oblongis obtusis, corollâ e sicco albida vel rubella calyeem parum excedente lobis triangularibus, forniciibus triangularibus hirsutis sub medio tubo sitis, staminibus longe exsertis, pedicellis calyce et fructu brevioribus, nuculis areolâ planâ parce echinulatâ margine prominenti pluriseriatim et breviter echinulato ♀. Ic. Ledeb. tab. 26.

Hab. in montosis Persie boreali-orientalis supra Asterabad, Radkann (Bunge!).

Ar. Geogr. Songaria, Sibiria Altaica.

7. *S. petiolaris* (d.c. Prodr. X, p. 164) viridis adpresso et parce pubescens, caule elato folioso, foliis radicalibus et caulinis inferioribus longe petiolatis oblongis obtusis cordatis vel subcordatis, superioribus sessilibus sensim diminutis summis semiamplexicaulibus, racemis paniculam longam angustam formantibus fructiferis laxiusculis, calycis adpresso hirsuti laciniis oblongis obtusis, corollâ cœruleâ calyce sessili longiore dentibus triangularibus, forniciibus hirsutis sub medio tubo sitis, staminibus longe exsertis, nucularum areâ echinatâ depressâ margine aculeis basi dilatata subconnatis pluriserialibus cristatâ ♀. *S. ruminifolius* Boiss. in Ky. Pl. Pers.

Hab. in Kurdistaniâ prope Van (Noë!), Mesopotamiâ (Auch. 2295!), montibus Pir Omar Gudrun, Avroman et Schahu 6000'-7000' (Haussk!), monte Sawers Persie australis usque ad 43000' (Haussk!), montibus ad Schiraz (Ky. 325!), ad Isphahan et in prov. Aderbidjan (Auch 4973! et 4974!), monte Ssahend (Buhse!), monte Elbrus ad Derbend (Ky. 129!).

Affinis *S. circinnata* differt floribus cœruleis, nucularum aculeis elongatis marginalibus in cristam coalitis.

8. *S. amplifolius* (Boiss. Diagn. Ser. I, 11, p. 126) parce et minute pubescens viridis, caule elato sultato superne ample paniculato, foliis tenuiter membranaceis radicalibus longe petiolatis amplis ovato-oblongis profunde cordatis, superioribus ovatis et suborbiculari-

tis ample et obtusissime auriculatis, racemis demum elongatis laxis, calycis hirsuti laciniis oblongo-spathulatis obtusissimis, corollæ cernæ calycem vix excedente lobis rotundatis, forniciis ovatis ciliatis sub medio tubo sitis, filamentis longe exsertis, nuculis disco tuberculato-echinulatis marginæ aculeis pluriserialibus basi dilatatis constante ♀.

Hab. in umbrosis rupestribus Antilibani inter Zebdani et Zachle (Boiss'!). Fl. Maio.

Foliorum radicalium laminae 9-10 pollices longæ; planta bipedalis affinis *S. petiolaria* a quo foliis caulinis late auriculatis, calycis laciniis spathulatis, forniciis formâ, nuculis majoribus differre videtur.

TRACHELANTHUS (Kunze Bot. Zeit VIII, p. 663. — *Cerinthopsis* Kotschy, MSS.).

Calyx quinquepartitus. Corolla calycem parum excedens cylindrica dimidiâ parte superiori dilatata profunde quinqueloba lobis erecto-couniventibus linearis-oblongis obtusis et extus ad loborum basin gibbis 5 retusis prominentibus instructa. Fornices elongatæ oblongæ. Stamina 5 filamentis latis brevissimis sub fauce insertis, antheris longe linearibus basi sagittatis. Stigma capitatum. Nucule 4 ovatae convexo-depressæ columnæ centrali adfixæ undique glochidiato-echinatae immarginatae. — Herbæ glabrae elatae grandifoliae, floribus *Cerinthes*. — Genus affine *Solenanthe* a quo corellis sub mediâ parte gibbis quinque prominentibus donatis et antheris longis linearibus differt.

1. **T. cerinthoides** (Boiss. Diagn. Ser. I, II, p. 127 sub *Solenanthe*) glaber glaucescens, caule crasso elato in paniculam amplam foliosam abeunti, foliis subtus tuberculis parvis calcareis sparsis obsitis oblongis radicalibus amplis basi angustatis vel truncatis longe petiolatis, caulinis et rameis diminutis a basi obtuse subcordatâ oblongis, racemis brevibus subcorymbose paucifloris, pedicellis gracilibus calyce pluries longioribus, calycis laciniis ellipticis obtusis, nuculis convexis immarginatis tuberculatis brevissimis glochidiatis echinulatis ♀. *T. cerinthoides* Kunze loc. cit. p. 663.

Hab. ad radices montis Demavend Persie borealis prope Ask (Ky. 181!), prope Kohrud (Bge!), in monte Dehbatlo prope Jezd (Buhse!), ad Gulfur districtus Kuh Kilouy Persie occidentalis 7000' (Haussk!).

Planta bipedalis et procerior, folia radicalia magnitudine variantia inter clara $\frac{1}{2}$ -1-pedalia, calyx florifer tres lineas longus tandem accretus, corolle lata lobis linea rubra notatis.

2. **T. Kurdicus** (Ky. Sched. 1859 sub *Cerinthopsis*) glaber, glaucescens, caule crasso elato in paniculam corymbosem foliosam abeunte, foliis membranaceis radicalibus amplis ovatis basi breviter attenuatis, caulinis et floralibus a basi obtuse cordato-auriculata late

ovatis obtusissimis, racemis brevibus floriferis umbellatis dein parum elongatis, pedicellis calyce duplo longioribus, calycis laciniis oblongo-linearibus obtusis, nuculis convexis undique longe et crebre echinato-glochidiatis ♀.

Hab. in angustiis vallis Teng provinciae MuschArmeniae Kurdieæ 6500' (Ky. 493!).

Affinis præcedenti a qua differre videtur foliis caulinis et rameis multo latioribus et obtusioribus late-auriculatis, nuculis longius echinatis.

CYPHOMATTIA.

Calyx quinquepartitus. Corolla calycem parum excedens cylindrica in lacinias linearis-subulatas rectas conniventes ultra medium partita supra basin gibbis quinque inflatis obsita et infra gibbos abrupte et breviter constricta. Fornices suprabasiliare. Filamenta inclusa, antheræ lineares. Stigma punctiforme. Nuculae compressæ planæ ventre columnâ centrali pyramidatâ adfixæ alâ planâ latâ marginatâ laevissimæ. — Herba Orientalis facie *Mattiae*. — Genus a *Mattiæ* ut *Trachelanthus* a *Solenanthe* corollæ tubo supra basin abrupte angustatum gibbis quinis instructo, lobis conniventibus, antheris elongatis diversum; a *Rinderæ* non fornicibus (quæ in hâc rudimentariæ sæpe adsunt) sed corollæ tubo gibboso distinctum.

1. **C. lanata** (Lam. Ill. № 1802 sub *Cynoglosso*) rhizomate crasso squamato, caulibus crassis a parte inferiori sæpe ramosis corymbose paniculatis, foliis firmis pilis adpressis basi tuberculatis canescensibus interdum glabratis viridibus tuberculato-punctatis radicalibus petiolatis oblongis basi cuneatis, caulinis oblongis vel lanceolatis inferioribus basi attenuatis superioribus strictis semiamplexicaulibus. inflorescentia tomentosa, racemis floriferis densis capitatis fructiferis subelongatis. calycis dense lanati laciniis linearis-lanceolatis, corollæ roseâ demum cærulescente ultrâ medium in lacinias angustas linearis-lanceolatas partitâ. fornicibus oblongis prope corollæ basin sitis, staminibus corollâ multo brevioribus, pedicellis fructu longioribus erecto-patulis, nuculis planis laevissimis alâ eis æquilitatâ vel latiore orbiculatâ radiatim subplicatâ margine integrâ vel repandâ cinctis, stylo fructu æquilongo vel longiore ♀. *Mattia lanata* Schult. Obs. p. 31. — *M. punctata* Alph. d.c. (forma foliis glabrescentibus) et *M. canescens* Alph. d.c. Prodr. X. p. 167 et 168. — *M. umbellata* C. Koch Linn. XVII, p. 302 non Schult. — *M. eriantha* Ledeb. Fl. Ross. III, p. 173. — *Rinderæ pubescens* C. Koch Linn. XXIII. p. 648. — *Cyn. lanatum* Desf. Choix tab. 20 (flores nimis cernui).

Hab. in collibus, rupestribus, regionis subalpinæ et alpinæ, Lycaonia in sylvaticis Karadagh (Heldr!), Cappadocia ad Caesaream (Ball!), Cilicieæ ditio Kassan Oghlu (Ky. 56!), Syria borealis ad Aintab (Auch. 2296!), Assyria ad Diarbekir (Ky. 236!), Armenia ad Erzerum (Huet!, Calv!), Gumuschkhané

(Tehi!), Baibut et Ispir (Bourg!), Armenia Rossica (Szov!), Persia borealis (Buhs! Bge!), ad Schahrund (Bge!), Aderbidjan (Auch. 4975!), montes Schahu, Pir Ounar Gudrun 6-10000' (Haussk!), Sawers 10-11000' (Haussk!).

β. brachyantha. — Calycis laciniæ abbreviatae oblongæ. Corolla laciniæ breviores obtusiores. — *M. brachyantha* Boiss. Diagn. Ser. I, 11, p. 127.

Hab. in Galatiâ prope Angora (Wied!), in Persiâ (Auch. 4976!), in monte Kuh Barti prope Schiraz (Ky. 332!).

Planta pulchra statura ($\frac{1}{2}$ - $\frac{1}{2}$ -pedalis), foliis oblongis vel anguste lanceolatis, calycis laciniis plus minus abbreviatis et obtusis, corollâ $5\frac{1}{2}$, vel 3 lineas longâ valde varians. Nuculæ cum alis diametro 6-9 lineas late.

MATTIA (Schult. Obs. p. 30. — *Mattiæ* Sect. I et II, d.c. Prodr.).

Calyx quinquepartitus. Corolla cylindrica tubo aequali laciniis ovatis vel linearibus rectis. Fornices versus basin tubi vel sub fauce sitæ. Stamina inclusa vel exserta antheris ovato-oblongis. Stigma punctiforme. Nucule compressæ planæ ventre columnâ centrali pyramidatâ adfixæ alâ planâ latâ marginatae levissimæ. — Herbae perennes calyce lanato. — Genus corollâ *Solenautho* proximum, fructu *Paracaryi* sectione *Mattiastro*; a *Rinderâ* non tantum fornicibus quæ in eâ quoque sœpe rudimentaria adsunt sed antheris diversum.

1. **M. cespitosa** (Alph. n.c. Prodr. X, p. 168) pilis adpressis caulis retroflexis canescens cespitosa, rhizomate valde squamato, caulis simplicibus crebre foliosis corymboso-umbellatis, foliis radicalibus lineari-lanceolatis acutis basi longe attenuatis caulinis linearibus strictis superne abbreviatis, corymbi floriferi compacti racemis fructiferis subelongatis, pedicellis floriferis calyci æquifloris fructiferis fructu multo longioribus, calycis lanati laciniis anguste linearibus, corollâ calyce sesquilociore ad tertiam partem in lacinas lineares partitâ, fornicibus ad faucem brevibus transverse pliciformibus, antheris subexsertis, nuculis levibus alâ æquilaterâ membranaceâ ovalo-orbiculari radiatim subplicatâ erosulâ marginatis, stylo fructu longiore ♀.

Hab. in regione alpinâ Armeniæ Turcicæ, mons Techdagh prope Erzerum 9-10000' (Huet!), Bingöldagh (Ky. Suppl. 695!), Berytdagh Cataonia (Haussk!), Cappadocia ad Euphratem (Auch. 2282!), Aslandach (Balz.).

Caulis florifer 3-5-pollicaris, fructifer 6-10-pollicaris, folia rosularum sterilium tandem 6-8-pollicaria 3-4 lin. lata, corolla 4 lineas longa, nuculæ cum alâ tandem 6 lineas longæ, fornices interdum deficiunt!

2. **M. Græca** (Boiss. et Heldr. Diagn. Ser. I, 7, p. 30 sub *Rinderâ*) adpresso argenteo-sericea cespitosa, rhizomate squamato, foliis anguste linearibus radicalibus basi longe attenuatis, caulinis preter infima paucis squamiformibus, caulis simplicibus pumilis corymboso-umbellatis, racemis fructiferis brevibus laxis, pedicellis flori-

feris brevibus fructiferis fructu longioribus, calycis lanati laciniis linearibus, corollâ purpureâ calyce sesquiloniori ultra tertiam partem in laciniis oblongo-spathulatas fissâ, forniciis sub fuce triangularibus obtusis, antheris inclusis, nuculis levibus alâ eis sublatiore ovatâ margine repandâ cinctis, stylo fructu longiore ♀. *M. Græca* Alph. n.c. Prodr. X, p. 168. — *M. Schmidtii* Heldr. Index Sem. Athen. 1860. — *M. bracteata* Orph. MSS.

Hab. in rupestribus regionis alpinae Graeciae, mons Malevo Laconiae supra Kastanitzza 6000' (Heldr!), Chelmos prope Stygem 5000' (Orph. Fl. Gr. exs. 168!), Kyllene 7000' (Heldr!), insula Poros (Wied!), Citheronis et Parnassi regio alpina (Guice. in Heldr. Herb. Norm. 575!), cacumen montis Delphi Eubeæ (Schmidt, Illeades!).

Caulis florifer 2-4-pollicaris fructifer subelongatus, folia 4 1/2-2 lineas lata, corolla 4 1/2-5 lin. longa, nuculae sex lineas latae. Affinis *M. cespitosæ* differt fornicum et corolla loborum formâ, staminibus inclusis.

3. ***M. Bungei***, cespitosa adpressissime cana, rhizomate verticali ad collum vix squamoso, foliis radicalibus numerosis linear-lanceolatis aentis basi longe attenuatis, caulinis paucis brevissimis, caulibus simplicibus pumilis corymboso-umbellatis, racemis fructiferis brevibus paucifloris laxis, pedicellis fructu 2-3-plo longioribus, calycis lanati laciniis linearibus, corollâ calyce quartâ parte longiore ad quintam partem in lobos ovatos fissâ, forniciis prope basin corollæ sitis ovatis, filamentis corollâ duplo longioribus, nuculis levibus margine membranaceo æquilato radiatim subplicato integro vel subrepando cinctis, stylo fructu subbrèviore ♀.

Hab. in montibus Persiae bor.-orientalis prope Schahrud (Bge!).

M. cespitosæ et *Græca* affinis, folia radicalia 2-3-pollicaria 2-3 lineas lata, caules 2-4-pollicares, corolla 2 1/2 lineas longa; ab eis staminibus longe exsertis, corollâ brevis lobatâ et forniciis subbasilaribus distincta.

4. ***M. Schlumbergeri***, cespitosa adpressissime cana, foliis radicalibus ovato-oblongis spathulatis obtusis in petiolum æquilonatum attenuatis, caulinis linear-lanceolatis basi sessili attenuatis, caulibus fere a basi laxe racemosis pumilis folia radicalia vix duplo excedentibus, pedicellis inferioribus axillaribus cæteris basi nudis omnibus strictis fructu 2-3-plo longioribus, calycis cani corollâ duplo brevioris laciniis linearibus, corollâ purpureâ longe tubulosâ ad sextam partem usque in lobos ovatos fissâ, forniciis transverse pliciformibus filamentisque corollâ subbrevioribus paululum sub fuce insertis, nuculis (junioribus) membranâ denticulatâ marginatis, stylo fructu longiore ♀.

Hab. in Libano ubi cl. Schlumberger legit vere 1872.

Planta 3-4-pollicaris, foliorum radicalium lamina 9-10 pollices longa 5-6 lata, pedicelli post anthesin 8-9 lineas longi, corolla fere sex lineas longa. Species elegantissima ex unico specimine mihi nota.

RINDERÆ (Pall. Itin. I, p. 486).

Calyx quinquepartitus. Corolla cylindrica in lacinias anguste lineares erectas subcoinvientes profunde partita tubo aequali, fauce undâ vel obscure et transverse plicatâ. Stamina filamentis brevibus latis sub fauce insertis, antheris linearibus basi sagittatis. Stylus elongatus, stigma punctiforme. Nuculae compressæ planæ ventre columnâ centrali pyramidatâ adfixæ alâ planâ latâ marginatae levissimæ. — Herbae facie *Cyphomattiae* et *Mattiae*, a priori tubo corollæ non gibbosæ, a posteriore antheris elongatis sagittatis, ab utrâque fornicibus multis vel ad plieas transversas obsoletas reductis diversæ.

I. R. cyclodonta (Bunge Pl. Lehm. p. 415) caule humili corymboso, foliis radicalibus glabris parce tuberculato-punctatis linearioribus oblongis in petiolum aequilongum attenuatis, caulinis semiamplexicaulis oblongis summis sublanatis, inflorescentia cymoso-racemosa ramis brevibus paucifloris laxis, pedicellis calyce longioribus fructu brevioribus, calycis lamati lacinias oblongo-linearibus, corollâ calyce vix sesquiloniore ad medium partitâ, filamentis fauce longioribus corollâ brevioribus, fornicibus ad plieas prope faucem sitas vix perspicuas reductis, nuculis planis membranâ orbiculari aequilaterâ radiatim plicatulâ margine minute denticulatâ obsitis, stylo fructu subaequilongo ♂.

Hab. in rupestribus graniticis ad Bakali et Tiumenbaitan Turkestaniae (Lehm!).

Semipedalis, foliorum radicalium lamina bipinnicaris 3-4 lineas lata, corolla 5 lineas longa, nucula diam. 9-10 lineas late. *R. tetraspis* Pall = *R. larigata* R. et Sch. = *R. corymbulosa* C. Koch Linn. XXIII, p. 648, in Rossia australi, Songariâ, Sibiria nec in nostrâ ditione crescentis est planta clatior, foliis latioribus, corollâ calyce duplo longiore et tantum ad tertiam partem partitâ, fructu alis integris.

ASPERUGO (L. Gen. 489).

Calyx florifer quinquetidus lobis basi sinuato-dentatis, fructifer valde accretus compressus bivalvis valvis latis oppositis alterâ 6 — alterâ 7 — dentato-lobatâ. Corolla parva calyci subaequilonga subinfundibuliformis fauce fornicibus obtusis elansâ, limbo quinquelobo. Antheræ inclusæ subsessiles ovatae utrinque retusæ. Stigma capitatum. Nuculae lateraliter compressæ pyriformes verruculosa supra medium hilo laterali columnæ centrali affixæ. — erba annua flaccida.

I. A. procumbens (L. Sp. 498) annua, setis deorsum directis aculeolata a basi ramosa decumbens, foliis elliptico-oblongis radicalibus petiolatis, superioribus basi attenuatis 2-4-nis, floribus axillaribus

brevissime pedicellatis 3-4 fasciculatis foliis floralibus oppositis, pedicellis fructiferis cernuis, calycis fructiferi reticulato-venosi lobis acutis ciliatis ♂. Ic. Fl. Dan. tab. 552. — Rchb. tab. 126.

Hab. in ruderatis, cultis, totius ditionis ab Attica (Sibth., Heldr!), Macedoniâ et Thraciâ! ad Arabiam petream (Auch!), Persiam borealem (Bge!) et australem (Ky!), Affghaniam (Griff!), Belutschiam (Stocks!), Turkestaniâ (Lehm!). In australioribus sâpe montana.

β. peduncularis. — Flores solitarii, pedicelli calyci fructifero (juniori) æquilongi.

Hab. ad Ssertschah Persiæ mediæ orientalis (Bge!).

Specimen non sat ætate provectum.

Ar. Geogr. Europa omnis, Sibiria Uralensis et Altaica, Africa borealis.

SUCHTELENIA (Karelin Bull. Soc. Mosq. 1841, p. 16, tab. 2).

Calyx quinquelobus fructifer valde accretus expansus. Corolla parva hypocaterimorpha tubo limbū obtuse quinquelobum non æquante, fauce fornicibus obtusis clausâ. Antheræ 5 ovatæ subsessiles inclusæ. Stylus brevis apice capitellatus. Nuculae 1-4 oblongæ apice attenuatæ erectæ dorso convexo echinatæ vel læves ventre concavo toto stylopodio magno globoso inflato cavo adfixæ et interdum cum eo concretæ. — Herba annua glauca pumila facie *Omphalodes linifoliae*. — Genus stylopodio insigne, calyx fructifer *Caccinia* et *Heliocaryæ*.

1. S. calycina (C.A. Mey. Enum. p. 100 sub *Cynoglosso*) annum pumilum glaucum carnosulum a collo sepius pluricaule, caulis inferne foliosis in racemos 2-3 laxos abeuntibus, foliis integris oblongis glabris vel margine præsertim tuberculato-ciliatulis inferioribus basi attenuatis, bracteis oblongis pendulis, pedicellis filiformibus fructiferis cernuis fructu brevioribus, calycis fructiferi lobis ovatis obtusis, fructifero valde acreto stellatim expanso, corollâ cœruleâ calyce sublongiore lobis ovatis obtusis, nuculis dorso vel lævibus, vel scabridis vel aculeolato-glochidiatis ♂. *S. calycina* Alph. D.C. Prodr. X, p. 163. — *S. acanthocarpa* Kar. — *S. uniserialis* Led. Fl. Ross. III, p. 169.

Hab. in argillosis thermalibus in deserto Transcaucasico inter Baku et Sallian ad diversorium Kæte Koili (C.A. Mey!), in littore orientali Caspii (Kar.), Turkestaniâ in desertis Karakum et Kisilkum (Lehm!).

Fructus ex cl. a Bunge in eisdem locis læves vel echinati mire variant. Planta semipedalis, flores vix lineaæ $\frac{3}{4}$ longi, calyx fructifer diam. 5 lineaes latus, nuculae vix 2 lineaes longæ.

CACCINIA (Savi Cose bot. p. I, tab. I. — *Anisanthera* Rafin.).

Calyx ultra medium quinquefidus, fructifer valde accretus stellatim patens. Corolla hypocaterimorpha tubo angusto calycem superante,

fauce abrupte ampliatâ intus fornicibus obtusis clausâ laciniis patentibus subinæqualibus linearibus vel oblongis. Stamina apicí tubi inserta, antheræ exsertæ anticae et laterales minores inæquales subsessiles, posticum majus oscillatorium longius exsertum. Stylus filiformis longe exsertus, fructiferi basis in cupulas crustaceo-membranaceas nuculas foventes expansus. Nuculae fertiles 2 vel 4 compressie ovatae vel orbicularis horizontali-subpendulae. — Herbae valde strigosæ.

SECT. I. EUCACCINIA (*Phyllobotrys* et *Gymnobotrys* Bge Helioc. p. 6.
— Antherarum filamenta recta.

1. **C. glauca** (Savi Cos. bot. pag. 4, tab. 1) perennis glaucescens carnosa, caule florifero humili a basi ramoso paniculato, fructiferi angulati ramis elongatis laxis inferne tantum fructiferis superne bracteas steriles ferentibus, foliis oblongis et oblongo-lanceolatis margine et paginâ superiore aculeis albis conicis basi tuberculatis muricatis, bracteis late lanceolatis, calyce florifero in lacinias angustæ-linearæ strigosas fisso, fructifero valde ampliato stellato-patenti lobis late et breviter triangularibus, corollæ cœruleo-scapulæ tubo calyce sesqui vel duplo longiore laciniis elliptico-linearibus acutis, antheris a posticâ maximâ ad anticas sensim diminutis, nuculis ovatis disco planiusculis grosse et siépe obsolete rugosis margine brevissime triangulari-denticulatis ♀. *C. Celsii* Boiss. Diagn. Ser. 1, 41, p. 133. — *Borago crassifolia* Vent. Hort. Gels. tab. 100.

Hab. in ditione Talusch 5000' (Hohen!), prov. Aderbidjan Persiae (Szov! Auch. 5006!), Persia boreali ad radices montis Demavend prope Ask (Ky!), prope Schahrud, Khabbise, Kerman Persiae orientalis et inter Teheran et Ispahan (Bgel!), inter Hamadan et Teheran (Oliv.), in Belutschia ad Doubund (Stocks!) forma nucularum margine obtusius dentato

Tubi corollini quoad calycem ex speciminibus nuper observatis et dentium nuculae longitudo variat, ideo *C. Celsii* non specifice distinguenda videtur.

2. **C. Rauwolfii** (C. Koch Linn. XVII, p. 303 non p.c. Prodr.) perennis viridis carnosa, caule florifero elato et superne corymbose paniculato, foliis lanceolatis basi attenuatis superioribus saltem utrinque strigoso-tuberculatis, bracteis late et breviter lanceolatis, calyce florifero in lacinias breviter triangulari-lanceolatas fisso fructifero valde aucto stellatimi patenti lobis late ovatis, corollæ cœruleæ tubo calyce duplo longiore laciniis angustæ linearibus acutis, antheris lateralibus minutis, anticas duplo majoribus, posticâ eis vix duplo longiore, nuculis orbiculatis planis disco et margine cristæformi dentato tuberculatis stellatim glochidiatis creberriniis obsitis ♀.

Hab. in Armenia Rossicâ ad Nackitchevan (Szov!), ad Erivan (Boiss! Seidl!), in tractu Lori (C. Koch!), in Persia australi orientali prope Ssertschah (Bgel).

Ex cl. Bge loc. cit. planta sub anthesi *C. glauca* procerior 4-4¹/2-pedalis superne tantum ramosa, viridis, calycis dentes breviores, anthera postica eā *C. glauca* brevior; ab eā nuculis undique crebre stellatum glochidiatis distinctissima.

3. C. strigosa (Boiss. loc. cit. p. 134), perennis glaucescens, carnosa, caulinibus decumbentibus elongatis superne breviter paniculatis, foliis aculeis conicis basi albo-tuberculatis strigosis amplis ovato-oblongis obtusissimis basi attenuatis aculeato-ciliatis, racemis brevibus fructiferis laxis pedicellisque tandem recurvis aculeatissimis, bracteis minutis linearis-filiformibus, calycis floriferi dentibus triangularibus valde strigosis, fructifero ampliatostellato dentibus brevissime et late triangularibus introflexis, corollæ cœruleæ tubo calyce plus duplo longiore laciniis elliptico-linearibus, nuculis utrâque facie convexis exteriori levibus elevatim lineatis margine angustissimo crassiusculo aculeis minutis obsito ♀.

Hab. in Persiâ boreali ad radices montis Demavend prope Ask 6000' (Ky. 368!). Vid. sicc. et viv. cult.

Folia inferiora 4-5 pollices longa 2 1/2-3 lata. A precedentibus bracteis non late foliaceis sed angustissime linearibus, nucularum margine crassiusculo distincta.

4. C. Kotschy (Boiss. Diagn. Ser. I, 7, p. 31) perennis? caulinibus decumbentibus foliosis simplicibus in racemum simplicem breveni panicolorum abeuntibus, foliis pilis brevibus e tuberculo ortis strigulosis ovato-oblongis inferioribus in petiolum longiusculum attenuatis, superioribus ad racemi basin sessilibus, bracteolis paucis angustis linearibus vel nullis, calycis floriferi laciniis angustis linearibus strigosissimis, fructifero valde ampliato patenti lobis late ovatis acuminatis, corollæ tubo calyce sesquilociore laciniis anguste linearibus, antheris . . ., nuculis facie exteriori planis eximie rugulosis immarginatis ♀.

Hab. in glareosis ad parietes rupium montis Kuh Ajub prope Persepolin Persiae (Ky. 4-21).

C. strigosa racemo bracteis filiformibus obsito vel nudo affinis differre videtur nuculis (sex lineas longis) apice subincurvis facie non levibus et grosse lineatis sed rugulosis marginaque dentato destitutis.

5. C. Russelii (Boiss. loc. cit. p. 134) biennis, caule crasso simplici supra basin foliosissimum: in thyrsum pyramidatum ramosissimum abeunte, foliis facie superiore et margine tuberculatis albis breviter aculeatis obsitis inferioribus lanceolatis elongatis acutis basi attenuatis, raneis et floralibus linearis-lanceolatis brevibus, ramis erecto-patulis demum laxis rigidis, calycis floriferi dentibus ciliatis lanceolatis, fructifero ampliatostellato-patenti lobis triangularibus, corollæ tubo calyce duplo longiore laciniis longis linearibus obtusis, anthera posticâ aliis subaequalibus triplo majore, nuculis ovatis disco planis et lineâ elevatâ carinali percursis margine erecto angusto deu-

ticulato ②. *C. Rauwolfii* Alph. n.c. Prodr. X, p. 167 non C. Koch. — *Lycopsis Dioscoridis* Rautw. Besch. p. 120, tab. 22, — *Borrigo marranthera* Russ. Alepp. II, p. 246. — *Cynoglossum longiflorum* Lehm. Asp. p. 162.

Hab. in cultis et aridis ad Aleppo (Russell, Ky. 257!), inter Aleppo et Biredjick (Auch. 2348!), circa Damascum inter Triticæ (Gailb.).

Florifera semipedalis, fructifera 1-1½-pedalis. Inflorescentia thyrsoides. *Saxifraga pyramidalis* ut recte est n.c. notat et radice bienni insignis.

SECT. II. ACTINOBOLE Bge Helioc. p. 8. — Antheræa postica majoris filamentum apice ad insertionem versus medium antheram sitam angulo recto infractum.

6. **C. Actinobole** (Bunge loc. cit.) perennis, tota setis crassis basi tuberculatis albo-strigosa, caulinis erectis in paniculam thyrsoidem sub anthesi strictam oblongam abeuntibus, foliis oblongo-spatulatis in petiolum attenuatis, racemis numerosis brevibus tandem laxis, bracteis minutissimis filiformibus, pedicellis sub anthesi filiformibus, calycis lobis ovatis acutis fructifero aneto expanso, corolla purpurascens tubo calyce sesquilociore laciniis oblongis subimmaculatis, tubo 2½-plo brevioribus, antheræa postica e ceteris subimmaculatis subdupo longiore, nucula (juniore) solitaria disco hispidâ aculeis elongatis glochidiatis basi in alam confluentibus cinetâ ♀.

Hab. in præruptis montium prope Schahrud Persie borealis (Bgef).

Caules ½-1-pedales, folia inferiora 2-3-pollicaria, thyrsus sub anthesi 4-6 pollices longus, flores eis præcedentium duplo breviores. Facies *Heliocarya* a quâ characteribus genericis *Cuccinia* differt.

HELIOCARYA (Bge Mem. 1874).

Calyx ad medium quinquelidus fructifer aceretus stellatum patens. Corolla infundibuliformis tubo angusto calycem superante in faucom intus nudam sensim ampliato lobis erecto-patulis brevibus inaequalitatis subbilabiatis. Anthera fertilis unica postica filamento brevi piano inâ fauce inserta. Stylus filiformis inclusus, stigma punctiforme. Nucula abortu solitaria horizontalis obovato-orbicularis plana radiatim glochidiato-aculeata. — Herba perennis albo-strigosa. — Genus a *Cuccinia* corolla tubo in faucom sensim nec abrupte ampliato, limbo subbilabiato, fornicibus nullis, antheræa unicæ fertili subsessili inclusâ eximie distinctum.

1. **H. monandra** (Bge Helioc. p. 4) perennis, tota aculeolis basi tuberculatis albo-strigosa, caule crasso angulato brevi pumilio mox in thyrsum ovato-oblongum denum multiflorum tandem racemis fructiferis elongatis laxis corymbosum abeunti, foliis late obovatis in

petiolum attenuatis, bracteolis minutis brevibus filiformibus, calycis albo-setosissimi laciniis breviter oblongis, fructifero acreto stellatum patente obtuse quinquelobo, corollæ cœruleæ tubo calyce longiore apice sensim in limbum breviter et obtuse quinquelobum ampliato, nuculâ planâ disco hirtâ margine disco subangustiore in dentes lanceolatos glochidiatōs ad basin partito ♀.

Hab. in Persia mediae montosis prope Meibut et Aghda inter Jezd et Ispahan (Bge!), in montibus Sofi et Kohrud (Bode! Derderian!).

Seinipedalis vel parum elatior, folia et facies *Borraginis officinalis* sed flores multo minores, calyx florifer $2\frac{1}{2}$ lineas longus fructifer rotatus diam. 5-6 lineas latus, corolla 5 lineas longa, nucula diam. cum cristâ $3\frac{1}{2}$ lin. lata.

TRICHODESMA (R.Br. Prodr. 496).

Calyx profunde 5-sidus demum accrescens. Corolla calyce vix longior campanulata lobis latis subulato-acuminatis, fauce nudâ. Stamina tubo inserta filamentis brevissimis antheris longis dorso velutinis vel lanatis in appendices sabulatas in conum dextrorum contortum conniventes abeuntibus. Stylus filiformis. Nuculæ ascendentí-rectæ columnæ centrali quadrangulari ventre toto adnatæ marginatæ vel immarginatæ. — Herbae scabrae.

SECT. I. LEIOCARYUM Alph. D.C. — Nuculæ lèves non marginalæ.

4. T. **Indicum** (L. Sp. 197 sub *Borragine*) annum, a basi dichotome ramosum erectum vel diffusum setis adpressis alteris tuberculatis asperum, foliis plerisque oppositis lanceolatis inferioribus basi attenuatis caeteris basi semiamplexicaulibus, pedicellis supra-axillaribus calyce sublongioribus demum cernuis, calycis laciniis triangularibus acuminatis basi acute auriculatis, corollæ lobis acuminatis patentí-reflexis, antheris exsertis dorso velutinis apice elongato contortis, nuculis albis lapideis lèvissimis oblongis dorso convexis immarginatis ♂. T. *Indicum* R.Br. Prodr. 496. — Ic. Act. Par. 1718, tab. 10.

Hab. in Affghania (Griff.), Belutschia (Frere).

Ar. Geogr. India orientalis.

SECT. II. FRIEDERICHSTHALIA Alph. D.C. — Nuculæ granulatæ vel echinatæ margine denticulato cinctæ.

2. T. **Africanum** (L. Sp. 197 sub *Borragine*) annum vel perennans, caulis ramosis strigoso-aculeatis demum induratis, foliis seberrimis inter tuberculatos aculeiferos glabris plerisque oppositis oblongis acutis basi cuneatis inferioribus petiolatis, racemis termi-

nalibus laxis nudiusculis, pedicellis filiformibus strigosissimis calyci æquilongis demum cernuis, calycis basi truncati vix auriculati laciniis triangularibus acutis, corollæ lobis patentibus subbulato-attenuatis, antheris exsertis dorso hirsutis, nuculis oblongis acutiusculis dorso granulatis margine recto denticulato ♂. *T. Africanum* R. Br. loc. cit.

Hab. in saxis desertorum, Egyptus in Thebaide (Boiss! etc.) ad littus maris Rubri (Schw!), Ouadi Hebrae petreæ (Schimp 226!), valles Sinaiticae (Boiss!), regnum Mascate (Auch. 5023!) Belutschia inferior (Stocks!).

Ar. Geogr. India borealis, insulæ Capitis Viridis, Senegal, Caput B. Spei.

3. T. Ehrenbergii (Schweinfurth in Sched. pl. Eg. 1857) annuum setulis brevibus basi tuberculatis asperum inter setas ad foliorum paginam inferiorem pubescens, caulis ramosissimus, foliis oppositis ovatis obtusis basi truncatis inferioribus petiolatis, racemis brevibus laxiusculis paniculatis basi tantum minute foliolosis, pedicellis filiformibus hispidis calyce sublongioribus patentibus, floribus parvis, calycis basi subtruncati fructiferi acreto laciniis triangularibus acutis, corollæ lobis patentibus attenuatis, antheris dorso villosis exsertis apice contortis, nuculis oblongis disco echinulatis margine recto denticulato ♂. *Borrago Arabica* Ehrenb. MSS.

Hab. in littore Ægyptiaco inter Kosseir et Ras Benass (Schweinf!)

Ab affini *T. africano* distinctum pilis brevioribus tenuioribus minus strigosis, foliis latioribus basi truncatis (infer. 2 pollices longis 12-15 lineas latis) fere *Boerrhavia*; flores vix 2 lineas longi.

Ar. Geogr. Arabia.

SECT. III. OMMATOCARYUM. Alph. D.C. — Nuculæ urceolatæ membranâ introflexâ marginatae.

4. T. molle (D.C. Prodr. X, p. 174) perenne? totum molliter et adpresso incanum, caulis elongatis ramosissimus, foliis oppositis et alternis ovatis acutiusculis sessilibus inferioribus basi rotundatis cæteris subcordato-auriculatis, racemis paucifloris subpaniculatis terminalibus, bracteolis squamiformibus linearibus, pedicellis calyce longioribus demum cernuis, floribus magnis, calycis laciniis a basi rotundata oblongis longe setacco-acuminatis, fructifero acreto patenti, corollæ cæruleæ lobis triangularibus acuminatis reflexis, antheris extus parte inferiori dense et longe lanatis apice contortis, nuculis magnis cum columnâ subconcretis ovatis acutis dorso grosse serbiculato-rugosis non marginatis ♀. *Spirocoron glaucus* Stev. loc. cit. p. 576.

Hab. in apricis schistosis, glareosis torrentium Persiæ borealis in prov. Aderbijan (Szov!), montibus Ssamam (Buhse!), valle Loura montis Elbrus (Ky!), prov. Chorassan ad Nichapur (Bge!), inter Teheran et Tabris (Bge!),

ad Ispahan (Auch. 5024!), in monte Kuh Daëna Persiae australis (Ky. 618), ad Kermanchah, Sihna, in montibus Aviheng et Pir Omar Gudrun Kurdistaniae Persiae 7000' (Haussk!), Affghaniæ (Griff!).

Caules crassi 2-3-pedales an procumbentes? folia majores 1-1 $\frac{1}{2}$ -polli. longa, calyx fructifer diam. 15 lineas latu, nuculæ fere 3 lineas longæ.

5. **T. incanum** (Bge pl. Lehm. p. 417) perenne? totum dense pannosum incanum, caulis elongatis ramosissimis demum induratis, foliis inferioribus oppositis superioribus sparsis oblongis sessilibus utrinque attenuatis acutis sensim versus ramorum apicem diminutis, racemis terminalibus brevibus paucifloris, bracteis squamiformibus, pedicellis calyci fructiferi subæquilongis demum cernuis, calycis laciniis a basi rotundata ovato-lanceolatis acuminatis, fructifero valde aucto patent, corollæ cæruleæ lobis subcordato triangularibus longe acuminatis demum reflexis, antheris extus et inferne longe lanatis apice contortis, nuculis cum columnâ subconcretis oblongis acutis dorso-scabridis linea elevata longitudinali percursis ♀.

Hab. in Persiae prov. Khorassan prope Meschhed (Bge!), in rupestribus Turkestaniæ prope Samarkand et in valle fluvii Sarafshan (Lehm!).

Facie et characteribus *T. molli* affine differt foliis omnibus acutis basique attenuatis, nuculis non serobiculato-rugosis sed tenuiter tuberculato-scabridis.

6. **T. Aucherii** (d.c. Prodr. X, p. 174) perenne pilis brevibus retrorsis basi tuberculatis asperulum viride, rhizomate indurato pluricauli, caulis erectis simplicibus pumilis foliosis in racemum corymbose 3-5-florum laxum abeuntibus, foliis anguste lanceolatis acuminatis sessilibus basi subattenuatis infimis sensim abbreviatis squamiformibus, bracteis brevissimis linearibus, pedicellis calyce sublongioribus demum cernuis, calycis laciniis oblongis lanceolatis, fructifero parum acreto non expanso, corollæ campanilate pumiceæ lobis brevibus ovatis abrupte acuminatis non reflexis, antheris latere exteriori lanatis apice contortis, nuculis ureolatis disco brev carinato margine crasso radiatum ruguloso in membranam introllexo ♀.

Hab. in lapidosis calcareis alpinis Persiae australis (Auch. 2340!), alpes Kuh Daëna (Ky. 706!), montes calcarei ditionis Kuh Kilouyeh 40° 00' (Haussk!).

Caules rigiduli inferne denudati cum racemo 6-8-pollicares, folia majora 1-1 $\frac{1}{2}$ -pollices longa 3 lineas lata, corolla 7 lineas longa

7. **T. Stocksii** (Boiss. Diagn. Ser. II, 3, p. 140) perenne? totum setis brevissimis basi grosse tuberculatis aliis tenuissimis intermixtis omnibus adpressis cauescens asperum, caulis elongatis deennibentibus ex axillis breviter ramosis, foliis oblongo-ellipticis obtusis basi attenuatis inferioribus breviter petiolatis catenis sessilibus, racemis subcorymbose 3-5-floris, bracteis minutis lineari-lanceolatis, pedicellis fructu longioribus cernuis, calycis laciniis a basi rotundata lanceolatis acutis fructiferis acretis a basi cordata triangularibus

non expansis, corollæ campanulatae calyce vix longioris lobis breviter ovatis abrupte acuminatis non reflexis, antheris latere exteriori lanatis apice subcontortis, nuculis brevibus irregolatis globosis margine membranaceo valde introflexo ore poriformi ȝ.

Hab. in Belutschia inferiore (Stocks!), regno Mascate Arabie (Auch. 50251), huc quoque spectare videntur specimenia florifera ex Ssertschah Persie mediae orientalis (Bge!).

Caules pedales et longiores, folia majora 4-4 ½-pollicaria 8-9 linea ̄s lata Differt insuper a *T. Aucheri* sepalis fructiferis cordatis, nuculis globosis membranâ introflexâ præter porum apicalem omnino clausis, piso subuinoribus

ORD. LXXXVII. SOLANACEÆ.

(Juss. Gen. p. 124.)

Flores saepissime hermaphroditi regulares vel subirregulares. Calyx quinquefidus vel partitus persistens vel basi circumscissâ persistente deciduus. Corolla gamopetala formâ varia limbo recto vel obliquo et subirregulari saepius quinquelobo. Stamina quina laciniis corollæ alterna. Antheræ biloculares. Ovarium liberum bilocularē loculis multiovulatis, ovulis anatropis vel subamphitropis. Stylus 1, stigma integrum vel bilamellatum. Fructus nunc baccatus indehiscentis nunc capsularis dehiscens. Semimum testa reticulata vel foveolata. Embryo in centro albuminis carnosí cyclicus, spiralis, vel rectus.

TR. I. SOLANEÆ

Corollæ limbus subæqualiter plicatus vel in lobos valvatos vel induplicatos divisus. Fructus baccatus. Embryo periphericus cycliens vel subspiralis, cotyledonibus semiteretibus radiculâ non latioribus.

SOLANUM (L. Gen. 234).

Calyx campanulatus vel patens (in nostris) quinquedentatus vel partitus, fructifer parum angustus. Corolla rotata tubo brevissimo, quinqueloba. Stamina 5 fauci affixa brevissima, antheræ conniventes vel cohærentes apice poro interdum in rimam dilatato deliscentes. Ovarium bilocularē, stylus simplex. Bæcca polysperma, semina compressa granulata vel scrobicalata.

SECT. I. PACHYSTEMONUM Dun. — Inermia. Antheræ breviores crassiores poris introrsum spectantibus sæpe in rimas continuatis dehiscentes.

1. S. nigrum (L. Sp. 266) annum vel perennans glabrum vel brevissime pubescens, ramorum angulis prominulis sæpe tuberculatis, foliis ovatis vel deltoideo-rhombeis integris vel angulatis vel sinuato-dentatis, corymbis paucifloris umbelliformibus breviter pedunculatis extraaxillaribus, pedicellis fructiferis apice incrassatis reflexis, calycis laciniis brevissimis rotundatis, corollâ albâ pubescente calyce duplo longiore, baccis nigris ♂. *S. Dillenii* Schult. — *S. parviflorum* Bad. — *S. hirsutum* Dunal. — *S. Judaicum* Besser — *S. moschatum* Presl, cum innumeris aliis synonymis.

Hab. in ditionis totius ruderatis.

β. induratum. — Rami perennantes basi indurati sæpe elongati. — *S. nigrum* var. *suffruticosum* Moris Fl. Sard. III, p. 148. — *S. suffruticosum* Schousb. in Willd. Enum. Berl. I, 1236. — *S. Memphiticum* Mart? Dunal! Forma in Lusitaniam! Hispaniam australi! regno Maroccano! quoque occurrens.

Hab. in Judeâ ad Tiberiadem (Boiss!), Persia australi (Auch. 5030!)

γ. chlorocarpon Spenn. — Baccæ viridi-lutescentes vel luteæ. — *S. ochroleucum* Bast. — *S. flavum* Kit.

Hab. cum typo, habeo specimina ex Atticâ et Cephaloniâ (Heldr!), Transcaucasiâ (Ledeb.)

ε. humile. — Rami minus angulati. Baccæ ceraceo-flavæ. Cum præcedente confluit. — *S. humile* Bernh. in Willd. Emm. Berol. I, 236.

Hab. cum typo, Græcia (Heldr!) Creta (Raul!), Persia (Haussk!), Transcaucasia (C. Koch) et alibi.

Species valde polymorpha et a quâ binæ sequentes forsitan specifice non sat distinctæ.

Ar. Geogr. Fere ubique borealibus et antarcticis exceptis.

2. S. miniatum (Berh. ex Willd. Emm. Ber. I, 236) annum pubescens vel villosum, ramorum angulis prominulis tuberculatis, foliis ovato-deltoideis sinuato-dentatis, corymbis paucifloris breviter pedunculatis extraaxillaribus. pedicellis fructiferis apice incrassatis deflexis, corollâ albâ pubescente calyce duplo longiore, baccis miniatis ♂. \approx *villosum* Mill. non Lam. — *S. puniceum* Gmel.

Hab. in ruderatis, Græcia (Heldr!) Zæcynthus (Marg!), Syria (Gail!), Persia (Haussk!) et alibi.

Ar. Geogr. Europa media et australis, Nubia.

3. S. villosum (Lam. Diet. 4, p. 289) annum undique patule tomentoso-villosum, ramorum angulis fere obsoletis, foliis ovato-deltoides sinuato-dentatis, corymbis paucifloris breviter pedunculatis extraaxillaribus, pedicellis fructiferis apice incrassatis deflexis, corollâ albâ pubescente calyce duplo interdum 2 1/2-plo longiore, baccis luteis ♂. *S. nigrum* var. *villosum* L. Sp. 266.

Hab. in ruderatis totius ditionis a Cretâ! et Græciâ! ad Egyptum! Arabiam petream! Mesopotamiam! Persiam occidentalem!

Ar. Geogr. Europa media et australis, Africa borealis.

4. S. Sinaicum (Boiss. Diagn. Ser. I. 11. p. 135) basi suffrutescens dumulosum ramosissimum partibus junioribus herbaceum viride adpressiusculæ pubescens, ramis tenuibus ascendentibus, foliis oblongis et rhombicis irregulariter sinuatâ dentato-paucilobatis rariis indivisis inferioribus longe superioribus breviter petiolatis, cymis longis pedunculatis extraaxillaribus 2-5-floris, pedicellis deflexis vix incrassatis elongatis, calycis setulosi laciniis oblongis obtusis, corollâ albidâ calyce triplo longiore, baccis nigris ȝ.

Hab. in fissuris rupium vallis Ouadi es Scheick jugi Sinaitici (Boiss!) Fl. Mart.

Rami novelli 6-10-pollicares, folia majora pollicaria. A *S. nigro* var. *indurato* ramis vetustis non induratis sed lignosis pennae anserinae crassitie, indumento setuloso sparso adpresso, calycis laciniis elongatis distinctum.

5. S. Dulcamara (L. Sp. I, p. 264) caule fruticoso sarmientoso, ramis novellis elongatis herbaceis, foliis ovatis cordatis petiolatis minute pubescentibus vel tomentellis superioribus hastatis vel utrinque lobulo auctis, cymis extraaxillaribus longe pedunculatis plures et late dichotomis, pedicellis articulatis, calycis lobis brevisimis late triangularibus, corollæ violaceæ laciniis lanceolatis saepe reflexis, baccis ovatis rubris ȝ. Ic. Rehb. Ic. tab. 12.

β. indirizum. — Folia etiam superiora ovato-cordata indivisa. — *S. Persicum* Willd. ex Ræm. et Sch. Syst. IV, p. 662. — *S. assimile* Friv. Flora XIX, 439 (forma tomentella).

Hab. in umbrosis humidis, ad rivos, varietas β multo vulgatior, in totâ ditione a Græciâ! et Macedoniâ! ad Caucasum et Transcaucasiam! Persiam borealem! Anatoliam! Libanum!

Ar. Geogr. Europa tota arcticis exceptis, Sibiria, Japonia, China, Africa borealis.

6. S. Kieseritzkyi (C. A. Mey. Enum. p. 113) caulinis a rhizone indurato tenui repente fibrillifero ascendentibus subsimplicibus apice foliosis, foliis paucis sparsis sed approximatis magnis membranaceis longiusculæ petiolatis a basi cuneatâ ovato-oblongis acuminatis glabriusculis, cymâ subaxillari breviter pedunculatâ 2-3-florâ, pedicellis articulatis calyce brevioribus, calycis truncati

lobis latissimis fere obsoletis mucronulatis, coroliæ lacinij a basi lanceolatâ linearibus elongatis reflexis ȝ.

Hab. in sylvaticis ad radices montium regionis inferioris ditionis Transcaucasicae Talysh prope Lenkoran (C. A. Mey! Hohèn!).

Caules 6-8-pollicares, folia 4-5 laminâ 3-4 pollices longâ sesquipollicem latâ, corollæ laciniae eâ *S. Dulcamaræ* augustiores et magis reflexæ. Ab eâ vegetatione distinctissimum.

SECT. II. LEPTOSTEMONUM Dun. — Aculeata. Antheræ elongatae poris minutis sursum vel extrorsum spectantibus dehiscentes.

7. *S. gracilipes* (Decaisne in Jacqueæn. Voy. p. 113 tab. 119) suffruticosum, ramis teretiusculis elongatis macilentis cicatricosis inermibus, novellis herbaceis aculeis brevibus tenuibus patentibus obsitis lepidotis, foliis squamis stellatis lepidotis petiolatis cordato-ovatis vel orbiculatis obtusis subrepandis, floribus ex axillis superioribus solitariis pedicello folium aequante suffultis, calyce subcampanulato in lacinias lanceolatas lepidotas profunde partito, corollâ purpureâ calyce duplo longiore lacinij oblongis obtusis ȝ.

Hab. in Belutschia (Stocks!).

Ar. Geogr. India boreali-occidentalis.

8. *S. coagulans* (Forsk. Fl. Eg. Arab. p. 47) fruticosum totum adpresso tomentosum canum, ramis aculeis brevibus recurvis basi dilatatis obsitis, foliis petiolatis subtus elevatim nervosis et ad costam medianam plus minus aculeatis ovatis margine subundulatis repandis vel obtuse sinuato-lobatis, cymis extraaxillaribus breviter pedunculatis bifidis pedicello infimo florem fertilem cæteris steriles gerente, calycis campanulati lobis anguste triangularibus obtusis, corollâ purpureâ tomentosa calyce triplo longiore lacinij oblongis acutis, baccâ globosâ cerasiformi ȝ. Ic. Del. Eg. tab. 23. fig. I. — *S. sanctum* L. Sp. I, 269. — *S. Hierochuntinum* Duval loc. cit. p. 369.

Hab. in calidis Palestinæ prope Jericho et ad mare Mortuum (Boiss!), Egypto superiore (Del! Schimp! Boiss!), Persia australi (Auch. 5029!) Belutschia (Stocks!).

Ar. Geogr. Nubia, Abyssinia, Arabia tropica.

PHYSALIS (L. Gen. 250).

Calyx campanulatus quinquefidus fructifer valde inflatus membranaceus 5-10-angulato-costatus fructum includens dentibus conniventibus. Corolla subrotata limbo plicato 5-angulato vel fido. Stamina 5 prope corollæ basin inserta filamentis filiformibus, antheræ loculis paral-

lulis longitudinaliter dehiscentes. Ovarium bilocularis, stylus filiformis, stigma bilamellatum. Baccæ globosa calyce inclusa et eo multo minor. — Herba.

1. Ph Alkekengi (L. Sp. I, p. 262) herbacea perennis viridis parce hirtula, radice repente, caulinis erectis simplicibus vel parce ramosis flexuosis, foliis petiolatis ovatis acuminatis repandis superioribus geminatis, floribus in axillis solitariis pedunculo apice nutanti tandem reflexo suffultis, calycis villosuli lobis breviter triangularibus acuminatis patulis, fructifero ample vesiculoso ovato basi umbilicato igneo, corollæ squalide rubrâ fance virenti ♀. Ic. Fl. Græc. tab. 234. — Rchb. Ic. tab. 9

Hab. in umbrosis pinguis regionis montanae, Græcia in montibus Taygeto et Malevo Laconiae (Heldr! Orph!), Olympo Thessalo et Parnasso (Sibth!), Byzantii (Sibth!), Anatoliæ boreali (Wied!), Ponto littorali ad Rhizé (Bal!), Armeniæ Turcicæ ad Ispir Huet!, Ciliciæ prope Gülek (Bal!), Tauriæ, Caucaso et Transcaucasiæ! Persiæ boreali (Auch! Buhse!).

Ar Geogr Europa media a Belgio et Germaniæ ad Rossiam medianam et ex Hispaniæ boreali et Italiæ mediâ ad Dalmatiam et regionem Danubialem, Siberia Uralensis.

Obs. *Nicandra physalodes* L. I, p. 230 (sub *Atropâ*) Rchb. Ic. tab. 5 est planta Peruiana nunc variis locis v. gr. Caucaso (Rehm!), Transcaucasiæ (Hohen!) introducta.

WITHANIA (Pauq. ex Endl. Gen. 666. — *Puneeria* Stocks).

Calyx campanulatus quinquefidus fructifer valde auctus baccam includens. Corolla anguste campanulata quinquefida lobis valvatis. Stamina prope corollæ basin inserta quina, filamenta plana, antheræ erectæ loculis parallelis. Ovarium bilocularis, stylus filiformis, stigma breviter bilamellatum. Baccæ globose polysperma. Semina compressa reniformia. — Frutices.

1. W. somnifera (L. Sp. I, p. 261 sub *Physalide*) fruticosa, tomentosa ramis rectis teretibus flexuosis, foliis ovatis integris petiolatis superioribus saepius oppositis, floribus hermaphroditis brevissime pedicellatis axillaribus aggregatis rarius solitariis, calycis albo-lanati laciniis linearibus brevibus muticis, corollæ luteo-virentis calyce subduplo longioris laciniis ovato-triangularibus patulis, baccæ globosæ calyce fructifero vesieoso apice clauso inclusâ ♂. Cav. Ic. tab. 103. — Fl. Græc. t. 233. — *Physalis flexuosa* L. Sp. p. 261. — *P. arborescens* Thunb. Fl. Cap. — *Hypnoticum somniferum* Rodrig.

Hab. in collibus ad vias, in neglectis regionis inferioris, Peloponnesus in Messeniâ (Fl. Pelop!), Eubœa (Silth! Sprunt! Orph! Fl. exs. 723!), Creta (Heldr!), Phrygia (Auch. 2405!), Rhodus (Heldr! Bourg!), Cyprus (Sibth!), Cilicia (Bal!), Syria littoralis (Bl!), Egyptus (Ehr! Boiss!), Arabia petræa

(Schimp! Boiss!), Arabia Mascatensis (Auch!), Babylonia ad Bagdad (Noë!), Persia australis (Haussk!), Belutschia (Stocks!).

Ar. Geogr. Hispania australis, Sardinia, Sicilia, India orientalis, Arabia, Abyssinia, Africa borealis, Insulae Canarienses et Capitis Viridis, P. B. Spei.

2. W. coagulans (Stocks in Wight Ill. 4, tab. 1616 sub *Puneerid*) tota adpressissime lepidota cana fruticosa ramis strictis, foliis crebris supremis oppositis carnosulis strictis petiolo brevi in laminam elliptico-oblongam obtusiusculam integrum abeunte, floribus abortu dioicis breviter pedunculatis in axillis fasciculatis nutantibus, calycis lepidoti lobis triangularibus obtusiusculis, corollæ flavidæ calyce subdupo longioris lobis ovatis patentibus, baccâ calyce acreto sed apice aperto breviore ♂. Ic. Hook. Ic. tab. 801. — *W. coagulans* Dunal in D.C. Prodr. XIII, p. 685.

Hab. ad Shukarpore, Julalabad, Puschut Affghaniae (Griff!), in Belutschia (Stocks!).

Frutex bipedalis stricte ramosissimus; baccis ad lac coagulandum utuntur.

Ar. Geogr. Ditio Scinde Indiæ.

TR. II. ATROPEÆ.

Corollæ lobi imbricati. Fruetus baccatus. Embryo periphericus cyclicus vel spiralis cotyledonibus semiteretibus radiculâ non latioribus.

LYCIUM (L. Gen. 262).

Calyx poculiformis campanulatus vel tubulosus apice subtruncatus irregulariter dentatus saepè subbilabiatus fructifer non auctus. Corolla infundibuliformis tubo superne ampliato limbi lobis 5 patentibus imbricatis. Stamina 5 tubo medio inserta, antherarum loculi paralleli. Ovarium disco impositum biloculare, stylus filiformis, stigma capitatum vel bilamellatum. Bacca globosa vel ovata polysperma. — Frutices ramis gemmiferis, saepius spinosi spinis e gemmis vel terminalibus.

* Filamenta basi glabra.

1. L. Europæum (L. Syst. I, 228) fruticosum spinosum glabrum vel pruinosum glaucescens ramis erectis, foliis fasciculatis petiolatis obovato-lanceolatis obliquis obtusis vel acutiusculis, floribus ex fasciculis solitariis vel geminis longiusculi pedunculatis, calyce breviter poculiformi margine membranaceo primum æqualiter quinquedentato demum subbilabiato dentibus obtusiusculis, corollæ violaceæ tubo infundibuliformi, limbi lobis breviter ovatis tubo

triplo brevioribus, staminibus inclusis filamentis glabris, baccâ globosâ ȝ. Ic. Fl. Græc. tab. 236. — Jaub. et Sp. Ill. tab. 403. — Rehb. Ic. tab. 15. — *L. Mediterraneum* Dunal in D.C. Prodr. XIII, p. 523. — *L. Orientale* Miers Ill. II, p. 99 ex parte quod plantam Smyrenensem.

Hab. ad sepes Atticæ, Isthmi Corinthiaci (Heldr!), Anatoliæ in Lydia (Boissi!), Aegypti ad Alexandriam (Samar! Gall!) forma saepe foliis diminutis spinis crebroribus brevioribus).

Ar. Geogr. Europa mediterranea, Africa borealis, Insulae Canarienses.

2. *L. Arabicum* (Schweinfurth in Sched.) fruticosum ramosissimum spinosissimum ramis elongatis tenuibus erectis, foliis saepius cinereo-tomentellis oblongo-spathulatis in petiolum brevem attenuatis obtusis vel acutiusculis fasciculatis tandem sparsis, floribus solitariis vel geminis longiusculæ pedunculatis, calycis hirti breviter tubulosi dentibus ciliatulis brevissime triangularibus obtusis inæqualibus, corollæ violaceæ tubo infundibuliformi tenui, limbi lobis ovatis limbo quadruplo brevioribus, filamentis glabris, antheris subinclusis, baccâ globosâ nigrâ ȝ. *L. Mediterraneum* var. ȳ, ε et ζ Dunal in D.C. Prodr. loc. cit. p. 524. — *L. Orientale* Miers loc. cit. ex parte quoad spec. Arabica. — *L. Persicum* Miers loc. cit. pag. 100. tab. 65 (nomen erroneum, nam planta ex ipso auctore non ex Persiâ sed Arabiâ tropicâ oriunda).

Hab. in desertis Egypti inter Keneh et Kosseir (Schw!), Arabiæ petreæ ad Ouadi Mokatteb et in planitiæ Ramla (Boissi!).

A *L. Europeo* non tantum indumento plerumque tomentello et foliis minoribus (quæ tandem plantâ fructiferâ majora evadunt) sed presertim calycis formâ, corollæ tubo tenuiore et lobi proportione longiore distinctum; antheræ variant subapiculatae et muticæ. Huic ut forma foliis latioribus omnino adnamerandum videtur. — *L. saevum* Miers loc. cit. p. 95, tab. 64. — *L. abelieflorum* Rehb. Ic. XX, p. 10, tab. 15 ex Arabiâ tropicâ (Schimp. 855!) quod non differre videtur, et *L. Indicum* Wight Ic. tab. 1403 ex India boreali.

Ar. Geogr. Nubia! Arabia tropica, India borealis

3. *L. Barbarum* (L. Sp. 277 ex parte) fruticosum spinosum glabrum glaucescens, ramis flexuosis erectis albidis, foliis fasciculatis oblongo-spathulatis vel linearibus obtusis in petiolum attenuatis, floribus ad gemmas fasciculatis longiusculæ pedunculatis, calyce poculiformi margine membranaceo subbilabiato dentibus brevibus latis inæqualibus triangularibus acutis, corollæ rubrae lobis ovato-oblongis tubo vix brevioribus, filamentis basi glabris, antheris exsertis ȝ. — *L. foliosum* Stocks Kew. Gard. Journ. IV, p. 179 et *L. deppressum* Stocks eodem loco forma foliis angustioribus.

Hab. in Galatiæ montibus calcareis Dykmen prope Angora (Wied!), ad sepes et in agris prope Aleppo Syriæ (Ky. 55!), ad Orfa Mesopotamiae (Haussk!),

Babyloniae ad Mohammera (Noë. 136! et 998!), Persiae australis ad Dalechi (Ky. 166!), et borealis (Buhse! Bge!), in Belutschia superiore et circa Kelat (Stocks exs. 1117! et 995!).

Ne confundas cum *L. vulgari* Dunal in Europâ et Africa boreali frequenter culto, ramis pendulis, foliis longioribus, filamentis basi pilosis alieno et quod prob forma stenophylla *L. Chinensis* Mill. tantum est. Formæ foliis anguste linearibus *L. Barbari* cum *L. Ruthenico* facile confunduntur sed stamina semper basi glabra.

• • Filamenta basi hirsuta.

4. ***L. Ruthenicum*** (Murr. Comm. Gött. 1779. pag. 2, tab. 2) fruticosum glabrum glaucum, ramis erectis flexuosis spinosis albidis, foliis fasciculatis carnosis anguste linearis-spathulatis obtusis basi attenuatis floribus fasciculatis longiuscule pedunculatis, calyce membranaceo campanulato bilabiato labiis obtusis irregulariter et breviter dentatis, corollæ lobis ovatis tubo sesquibrevioribus, filamentis basi cum corollæ tubo hirtis, antheris exsertis, baccis nigris 5. — *L. Tataricum* Pall. Fl. Ross. I, p. 78, tab. 49. — *L. glaucum* Miers Ill. II, p. 104, tab. 66.

Hab. in desertis salsis Iberiae Transcaucasiæ ad Sallian et in tractu Swant (C. A. Mey! Hoh!), in Persia boreali (Auch. 503!), Turkestania (Lehm!).

Ar. Geogr. Rossia meridionalis, Songaria, Sibiria Uralensis.

5. ***L. Turcomanicum*** (Turcz. MSS. et Miers Ill. loc. cit. p. 118' tab. 69 D ubi filaments erronee glabra!) fruticosum glaucum glabrum, ramis erectis divaricatum ramulosis et spinosis, foliis carnosis soli tariis et fasciculatis oblongo-lanceolatis obtusis vel acutiusculis in petiolum attenuatis, floribus fasciculatis pedunculatis, calyce breviter campanulato superne membranaceo bilabiato labiis integris vel breviter bifidis, corollæ roseæ lobis oblongis tubo sesquibrevioribus, filamentis basi villosis, antheris exsertis, baccis coccineis 5.

Hab. in desertis argillosis salsis Turkestaniae ad fluvios Syr et Jandarja et circa Buchara (Lehm!).

Affine *L. Ruthenico* sed rami crassiores, folia multo latiora, baccæ coccinea nec nigra.

Ar. Geogr. Turcomania, Songaria.

ATROPA (L. Gen. 249 ex parte).

Calyx foliaceus profunde quinquesfidus fructifer parum auctus. Corolla late tubuloso-campanulata lobis quinque latis imbricatis. Stamina quinque basi corollæ affixa, filaments imâ basi incrassata inclusa, antheræ loculi distincti. Ovarium biloculare disco impositum, stylus subexsertus, stigma dilatatum breviter bilamellatum. Baccæ globosa succosa, semina numerosa compressa scrobiculata. — Herbæ perennæ.

1. A. Belladonna (L. Sp. 260) viscidulo-pubescent, caule herbaceo crasso erecto superne 2-3-tomo, foliis ovatis et oblongis integris acuminatis in petiolum brevem angustatis superioribus geminatis inaequalibus, floribus axillaribus solitariis rarius geminatis pedunculatis nutantibus, calycis lobis triangulari-ovatis acuminatis, corollâ squalide brunneâ lobis rotundatis, baccâ globosâ rubra φ . Ic. Rehb. tab. 8.

Hab. in sylvaticis montanis, Thessalia in monte Olympo (Heldt!), Macedonia in Athone (Sibth! Auch! Griseb!), Cilicia ad Tauri radices 3000' (Ky. 35 d.), Tauria, Caucasus et Transcaucasia (Lebed! Ower!), Persia borealis in monte Djulfek (Auch!) et circa Siaret (Bunge!).

Ar. Geogr. Europa media et australis ab Anglia et Daniâ ad Rossium, Africa borealis.

MANDRAGORA (Juss. Gen. 125).

Calyx subfoliacens quinquefidus vel partitus fructifer auctus. **Corolla** campanulato-semiquinquefida sinubus inter lobos induplicatis. **Stamina** 5 infra medium tubum affixa inclusa, filamenta filiformia, antherae oblongæ loculis subparallelis, ovarium disco impositum biloculare, stylus filiformis, stigma capitatum subbilobum. **Bacca** succosa poricarpio tenui, polysperma, semina punetato-scrobiculata. **Herbe** acaules radice crassâ.

1 M. officinarum (L. Sp. I, 481) radice crassâ carnosâ verticali simplici vel bi-trifidâ fusiformi, foliis rosulatis late ovato-oblongis magnis corrugato-crispis sepe sinuato-dentatis pilosulis vel glabris inferioribus obtusis cæteris acutis vel acuminatis, pedunculis e collo fascieculatis unifloris calyce longioribus hirtulis, calycis laciniis lanceolatis baccâ magnâ globosâ subbrevioribus, corolle albo-virentis extus hirtulæ laciniis oblongis vel lanceolatis calyce 2¹/₂-plo longioribus φ . Ic. Rehb. tab. 7. — **M. vernalis** Bertol. Comm. Maud. p. 9, tab. I. — **Atropa Mandragora** L. Sp. p. 259 et herb!

Hab. in cultis derelictis, aī pedem rupium regionis inferioris, Grœcia in Thessaliâ prope Zeitun (Sprunt!) et prob. alibi, insulâ Syra (Heldt!), Cretâ (Sieb! Raulin!), Pamphyliâ ad Adalia (Bourg!), Cypro (Ky. 446!), Syria littorali ad radices Libani circa Sidonem (Gauß!), Palestina prope Ramla (Boiss!). Floret a Februario ad Martem — Arabice *Rabouhe*.

Ar. Geogr. Hispania, Italia borealis, Dalmatia, Africa borealis ?

2. M. autumnalis (Spreng. Syst. I, p. 693) radice crassâ carnosâ verticali simplici fusiformi vel ramosâ, foliis rosulatis magnis ovato-oblongis subrepandis supra glabriusculis subtilis puberulis intimis obtusis cæteris acutis vel acuminatis, pedunculis e collo fascieculatis unifloris calyce longioribus glabriusculis, calycis hirtuli laciniis lanceolatis baccam ovato-oblongam æquantibus, corollæ violacea

calyce 2 $\frac{1}{2}$ -plo longioris laciinis oblongo-lanceolatis ♀. *Atropa Mandragora* Fl. Græc. III, p. 26, tab 232. — Rehb. Ic. tab. 6. — *M. officinarum* Bertol. Comm. pag. 10, tab. 2 et *M. microcarpa* Bert. loc. cit. p. 12. tab. 3.

Hab. in cultis, ad vias, Attica (Sprun! Heldr! Orph! Fl. exs. 75!), Peloponnesus (Bory!), Archipelagus (ex Sibth!). Fl. Oct. Nov.

Ab auctoribus saepe cum precedente confusa a quā differt foliis minoribus angustioribus non bullatis sub anthesi 3-4 polices tantum longis, corollā violaceā, baccā fere dimidio minore, florendi tempore.

Ar. Geogr. Hispania australis, Italia australis, Sardinia, Sicilia, Africa borealis.

TR. III. HYOSCYAMEÆ.

Corollæ lobi plicati vel imbricati. Fructus capsularis. Embryo periphericus cyclicus vel subspiralis cotyledonibus semiteretibus radiculâ non latioribus.

DATURA (L. Gen. 246)

Calyx longe tubulosus angulatus apice quinquedentatus vel spathaceus, fructifer supra basin persistentem interdum patentem et auctam circumscissus. Corolla infundibuliformis apice quinquedentata cœstivatione contortuplicata. Stamina quina prope basin tubi affixa inclusa, antheræ longe lineares loculis parallelis introrsum rimosis. Ovarium biloculare vel septis spuriis quadrilocellatum, ovula numerosa ad septum vel ad septa supria. Stylus filiformis apice dilatatus bilamellatus. Capsula exsucca oblonga semiquadrilocularis loculicide et septicide quadrivalvis. — Herbæ.

I. **D. Stramonium** (L. Sp. I, p. 179 annua, sparsim et minute papillosa viridis, caule laevi crasso erecto tereti superne dichotomo, foliis petiolatis ovatis glabris inæqualiter sinuato-dentatis acutiusculis, corollâ albâ calyce pentagono triangulari-quinquedentato subdupo longiore lobis latis brevibus abrupte et tenuiter mucronatis, capsulis erectis æqualiter aculeatis, seminibus nigris reniformibus alveolatis ♂. Ic. Engl. bot. tab. 1288. — Rehb. Ic. XX, tab. 3.

Hab. in ruderatis et ad vias totius ditionis a Græciâ! et Macedoniâ! ad Transcaucasiam, Syriam! Egyptum! Patria vera incognita.

Ar. Geogr. Europa media et australis, Asia et Africa, America borealis.

Obs. *D. Metel* L. Americana ad vias Syriae maritimæ circa Berythum introducta frequens (Boiss! Bl!), et *D. fastuosa* L. Indica in cultis Arabiæ Masantensis (Auch. 5038!) et Egypti occurunt.

PHYSOCALINA (G. Don Gen. Syst. IV, 470).

Calyx tubuloso-campanulatus apice breviter quinquefidus, fructifer auctus costato-reticulatus capsulam includens. Corolla campanulata-infundibuliformis lobis brevibus imbricatis. Staminæ 5 in medio tubi affixa filiformia, antheræ ovatae loculis parallelis. Ovarium biloculare multiovulatum, stylus elongatus stigmate dilatato obscure bitubo. Capsula bilocularis supra medium circumscissa. Semina compressa granulosa vel serobiculata. — Herbae facie *Hydropitheli*. — Genus affine *Scopolia*, ab eâ floribus parvis corymbosis, corollæ non plicatae lobis imbricatis distinctum.

1. **P. Orientalis** (M. B. Taur. Cœc. I. 164 sub *Hyoscyamo*) caulinis crassis simplicibus a basi foliosis ex rhizomate tuberoso-incrassato ascendentibus, foliis longe petiolatis majusculis teneris viridibus subpuberulis a basi subcordatâ vel cuneatâ ovatis vel oblongis acutiusculis subrepandis integris, floribus in cymam terminalem densiusculæ corymbosam pedunculatam congestis, calycis glanduloso-hirti dentibus triangularibus obtusiusculis tubo triplo brevioribus, corollâ violacea calyce $2\frac{1}{2}$ -plo longiore violacea, stylo exerto, antheris subexsertis ♀. *P. Orientalis* Don loc. cit. p. 470. — Ic. Bot. Mag. tab. 2414. — *Hyoscyamus Orientalis* Benth. *folio tuberosâ radice* Tourn. Cor. p. 5.

Hab. ad rupium verticalium radices et in cryptis, Armenia Turcica ad Teutera prope Gumusch Khane (Bourg!), in Caucaso ad Nartzana (M. B.), inter Kasbeck et Darial 3600'-4600' (Rupr!), in Alchasia et Somchetum (Lederb.).

Pedalis vel sesquipedalis, folia post anthesin aneta, calyx florifer tres lineas, corolla 6 longi

HYOSCYAMUS (L. Gen. 247).

Calyx tubuloso-campanulatus breviter quinquefidus fructifer auctus rigidus costatus capsulam includens. Corolla infundibuliformis interdum lateraliter tissa limbo obliquo quinquefido lobis latis imbricatis. Stamina 5 tubo affixa saepius exserta, antheræ oblongæ loculis parallelis. Ovarium biloculare multiovulatum, stylus filiformis, stigma capitatum. Capsula bilocularis apice vel prope apicem circumscissa operculo integro vel valvatum fisso. Semina compressa tuberculosa vel serobiculata. — Herbae.

1. **H. muticus** (L. Mant. 45) perennis, viridis, pubescens vel pilis crispis viscidulis flavidis tomentellus, caulinis crassis ascenden-

tibus et erectis angustis simplicibus vel stricte ramosis, foliis carnosulis a basi cuneatâ ovatis oblongo-rhombeis acutis superne acutè angulatis vel paucilobatis rarius integris, inferioribus late petiolatis superioribus subsessilibus, floribus breviter vel longiuscule pedunculatis in racemum scorpioideum brevem tandem elongatum rectum secundum dispositis, foliis floralibus sessilibus oblongis vel ellipticis integris acutis calyce æquifloris vel brevioribus, calyce fructifero aucto longe obconico dentibus late et breviter triangularibus acutis muticis, corollæ albidæ violaceo-maculatæ tubo breviter exerto limbo infundibuliformi subirregulari fisso, staminibus declinatis subexsertis ♀. Ic. Jaub. et Sp. Ill. tab. 415. — *H. Datura* Forsk. Eg. Arab. 43. — *H. betæfolius* Lam. Dict. III, 329. — *Scopolia mulica*, *S. Datura* et *S. Boveana* Dun. in D.C. Prodr. XV, 551. — *Hyoscyamus insanus* Stocks Kew Journ. Bot. IV, p. 178.

Hab. in desertis Egypti inferioris et superioris ad El Chargeh in Oasi magna (Schw! Del! Auch! Wiest! Boiss! etc.), Arabiae petreæ (Bové! Boiss!), Cataoniae prope Malatia (Auch. 2472!), Persiae australis ad Kaserun, Gere (Ky. 38! Haussk!), orientalis inter Khabbise et Kerman (Bge!), in rupibus Belutschiae (Stocks!).

Omnino contra affinitatem a Dunalio *Scopoliae* adnumeratus, calyce corollâ *Hyoscyamus* legitimus. Species a Dunalio propositæ nequidem varietates sunt.

2. *H. pusillus* (L. Sp. 258) annuus crispule papilloso-hirtus viscidulus, caule nano eretto demum crassiusculo simplici vel a collo ramoso, foliis oblongis vel oblongo-lanceolatis cuneatis acutis vel integris vel saepius sinuato-repandis vel acutè pinnatifidis partitisve radicalibus longe petiolatis, floribus subsessilibus calyce longioribus integris vel paucilobis, floribus minutis subsessilibus demum spicas secundas rigidas laxiusculas formantibus, calyce fructifero stricto infundibuliformi dentibus triangularibus pungentibus patulis tubo 4-5-plo brevioribus, corollæ flavidæ tubo brevi inclusa fance violascenti limbi subobliqui lobis subæqualibus ♂. Ic. Jaub. et Sp. Ill. tab. 414. — *H. micranthus* Ledeb. ex Don Syst. 4, p. 472. — *H. puugens* Griseb. Spic. II, p. 52.

Hab. in cultis, ruderatis, ad messes, Phrygia ad Bulghas Kieui (Ball!), Lycaonia (Heldr!), Cappadocia (Ball!), Armenia Turcica (Huett!), Galatia (Noël), Arabia petreæ (Boiss! Schimpt!), Egyptus inferior (Wiest! Kralik!), ditio Talysh (C.A. M!), Persia septentrionalis et media (Auch. 5041! Ky! Bge!), Turkestania (Bge!), Belutschia (Stocks!).

Ar. Geogr. Songaria, Sibiria, Altaica, Tibetia.

3. *H. niger* (L. Sp. 257) annuus vel biennis villosus viscidulus, caule eretto ramoso, foliis teneris villosis inferioribus petiolatis ovato-oblongis sinuato-pinnatifidis, caulinis a basi sessili semiamplexicauli oblongo-lanceolatis lobis triangulari-lanceolatis acutis, floribus sessilibus in spicam foliosam unilateralem scorpioideam

tandem elongatam rectam dispositis, foliis floralibus oblongo-lanceolatis calyces superantibus superioribus integris, calyce fructifero stricto supra medium constricto dentibus erecto-patulis triangulatis mucronatis, corollâ subaequali pallide flavidâ violaceo-reticulatâ fundo violaceo, staminibus fauce parum excedentibus ⊖ vel ⊙. Ic. Engl. Bot. tab. 591. — Rehb. Ic. tab. 2, fig. 2. — *H. agrestis* Kit. — *H. pallidus* W. K. (forma reticulatis violaceis corolla obsoletis). — *H. pictus* Bernh. — *H. Sysspirensis* C. Koch Linn. XXII, p. 736. — *H. Persicus* Boiss. et Buhse Enum. p. 158.

Hab. in ruderatis praesertim partis septentrionalis ditionis, *H. albo* rarior, Graecia (Sibth!), Macedonia et Thracia (Griseb.), Anatolia borealis (Thirkel! Wied.), Lycia (Heldr.), Armenia (Bourg!), Tauria, Caucasus et Transcaucasia (Ledeb!), Persia borealis (Buhse!).

Ar. Geogr. Europa fere omnis, Sibiria, Africa borealis in Atlante, India borealis.

4. *H. reticulatus* (L. Spec. 257 et Auct.) annuus vel biennis pube adpressâ arachnoidens, caule crasso elato simplici vel superne striate ramoso folioso, foliis oblongis vel lanceolatis plus minus profunde pinnatifidis radicalibus in petiolum brevem attenuatis, caulinis sessilibus oblongis, floribus sessilibus in spicas foliosas unilaterales scorpioideas demum elongatas rectas dispositis, foliis floralibus oblongis vel lanceolatis integris aentis flores superantibus, calyce fructifero stricto supra medium non constricto dentibus patentibus recurvis lanceolatis spinosis, corollâ aequali purpureo-violacea reticulata, staminibus fauce sublongioribus, stylo exerto ⊖ et ⊙. Ic. Janb. et Sp. tab. 416. — *H. Camerarii* F. et M. Ind. 10. Petrop. p. 38. — *H. pimatifidus* Schlecht. Linn. 1843, p. 127. — *H. purpureus* Griseb. MSS.

Hab. in ruderatis, cultis, Pisidia (Heldr.), Lycia (Bourg. 140!), Anatolia borealis (Wied!), Cappadocia (Bal!), Armenia ad Baibut Bourg!) et Erzerum (Huet!), Syria ad Aleppo (Ky. 467! Haussk!), in Cœlesyriâ, Damasco et Antilibano (Boiss! Gaill!), Mesopotamia (Hau-sk!), ditione Transcaucasicâ Talysh (C. A. M! Hoh!), Persia boreali (Buhse! Ky. 762! Bunge!), australi ad Persepolin (Ky. 808!).

β. *integrifolius*. — Folia etiam radicalia sinuata vel subintegra.

Hab. in Armenia australi prope Karput (Noë 1190!).

Valde affinis *H. nigra* a quo differt foliis caulinis sessilibus sed vix amplexicaulibus, calyce fructifero ad medium non constricto et ejus dentibus multo longioribus subrecurvis, corollâ purpureo-violacea. Specimina hujus speciei in herb Linnaeano non adsunt et ideo cl. auct. descriptio inter hunc et *H. album* subdubia.

5. *H. albus* (L. Sp. 257) annuus et perennans villoso-viscosus, caule erecto ramoso, foliis mollibus omnibus petiolatis a basi subcordata vel breviter cuneata ovato-orbiculatis grosse triangulirobatatis, floribus præter infima saepe breviter pedunculata sessilibus

in speciem unilateralem foliosam primum scorpioideam dein elongatam erectam dispositis, bracteis petiolatis flores superantibus cordatis ovatis dentatis vel sepius oblongis integris basi attenuatis, calycis fructiferi stricti villosi dentibus brevissime triangularibus acutis corollæ pallide luteæ non reticulatae fundo virentis vel atropurpureæ limbo obliquo lobis subimæqualibus, staminibus fane sublongioribus ♂ vel ♀. Ic. Fl. Graec tab. 230. — *H. major* Mill. Dict. II. — *H. aurens* All., Ten., non L. — *H. varians* Vis. Fl. Dalm. tab. 24. — *H. minor* Mill. Dict. 4. — *H. Canariensis* Ker Bot. Reg. 180

Hab ad muros, vias, in ruderatis totius ditionis vulgatissimus a Græciâ et Macedoniâ! ad Anatoliam borealem! Pontum circâ Trapezuntum (Huet!), Anatoliam australem! Syriam littoralem! Egyptum!

β. desertorum. Ascherson in litt. — Flores minores, inferiores pedunculo calyce tandem 2-3-plo longiore suffulti, calycis dentes abbreviati. Corollæ fane cum staminibus violascens.

Hab. ad Ouadi Askar deserti Ægyptiaco-Arabici Egypti mediæ (Schweinf. exs. 108!).

Formæ a Millero et Duvalio ut species propositæ nequidem ut varietates distinguendæ sunt.

Ar. Geogr. Europa meridionalis a Gallia australi ad Dalmatiam, Africa borealis, insulae Canarienses.

6. ***H. aureus*** (L. Sp. 257) perennis patule villoso-viscosus, caulis decumbentibus vel pendulis elongatis fragilibus ramosis, foliis petiolatis cordatis ovatis vel orbiculatis triangulari-lobatis lobis acute dentatis floribus omnibus pedunculis calyce brevioribus suffulti racemos demum longissimos laxos foliosos unilaterales formantibus, foliis floralibus petiolatis triangulari-ovatis vel oblongis acutissime dentatis flores superantibus, calyce fructifero erecto-patulo vel nutante obconico-campanulato membranaceo dentibus erectis triangulari-lanceolatis vix pungentibus, corollæ tubo limbo longiore subexerto declinato limbo aereo fundo violaceo obliquo fisco segmentis binis superioribus multo minoribus, genitalibus exsertis ♀. Ic. Fl. Graec. tab. 231 — *H. Creticus* Tourn. Inst.

Hab. ad mœnia et muros, insula Tenedos (Forsk.), Anatolia occid. ad Smyrnam (Ky!), in Cariâ (Pinard!), ad Ephesum (Boiss!), Pamphyliâ (Heldr! Bourg!), Cretâ (Siebt!), Rhodô (Auch. 2171!), Cypro (Ky!), Cilicia littoralis (Bul!), Syria ad Aleppo (Haussk!) et Damascum (Wettst! Gaill!), Berythum (Bl!). Palestinâ ad Hierosolymam (Auch! Boiss!), Mesopotamiâ in monte Sindjar (Haussk!).

A *H. albo* radice perenni, ramis decumbentibus fragilibus, foliis acute dentatis, floribus pedunculatis, calyce membranaceo vix pungente, corollæ tubo tenui elongato et limbi lobis valde inæqualibus distinctus. Semina duplo minora.

7. H. Senecionis (Willd. Enum. 4, p. 228) perennis crispule hirsutus et papillaris viscidulus, caulinis e rhizomate elongato indu- rato tenui ramoso decumbentibus elongatis ramosis, foliis longe petiolatis a basi truncatâ vel subcordatâ ovatis profunde 3-5-partitis lobis ovatis plus minus acute dentatis, floribus parvis ad axillas pedunculis calyce longioribus suffultis remotis paucis laxissime spicatis, foliis floralibus conformibus diminutis breviter petiolatis, calyce fructiferò pedunculo reflexo contorto suffulto obconico campanulato dentibus ovato-triangularibus acutis, corollæ aureæ fundo violas- centis tubo recto limbo tubo sublongiore fiso subobliquo segmentis subæqualibus, antheris e fauce subexsertis ♀.

Hab. in Persie montibus (nec in Egypto!) (Oliv! Auch. 5042!), Persia boreali (Buhsel!) in monte Totschal prope Teheran (Ky. 195!), in montibus Kuh Sawers et Kuh Nur Persie austro-occidentalis ad nives 12-13000' (Haussk!).

3. bipinnatisectus. — Folia in lacinias oblongas dentatas saepe in lacinulas minimas bipinnatisecta. — *H. bipinnatisectus* Boiss. Diagn. Ser. I, 7, p. 36.

Hab. in Persie montibus prope Jezl et inter Ispahan et Kaschan (Buhsel!), prope Ischredabad et inter Ssof et Kohrud (Bge!), in glareosis regionum altiorum alpis Kuh Delu Persie australis (Ky. 466!).

Species a *H. aureo* affini caulinibus tenuibus, foliis dissectis, pedunculis elongatis fructiferis contortis, corollæ duplo minoris limbo vix inaequidi, antheris vix exsertis distincta. Varietas ad typum manifeste transit.

ORD. LXXXVIII. SCROPHULARIACEÆ.

(Lindl. Veg. Kingd. 681.)

Flores hermaphroditi sœpissime irregulares. Calyx inferus liber persistens lobis vel segmentis 3 rarius 4. Corolla gamopetala campanulata vel rotata, limbo 3-4-lobo æqualiter patenti vel bilabiata et tunc labio postico integro retuso vel bilobo, antico trilobo fauce interdum bigibbo, lobis aestivatione varie imbricatis. Stamen sœpius 4 didynama, vel bina tubo corollæ affixa et ejus lobis alterna, quintum posticum deficiens vel ad staminodium reductum rarius perfectum, filamenta libera, antheræ vel distinete biloculares loculis longitudinaliter dehiscentibus vel loculis apice confluentibus uniloculares. Ovarium liberum biloculare loculis multiovulatis rarissime bioculatis, ovulis anatropis vel amphitropis. Stylus simplex vel bilobus. Fructus capsularis varie dehiscentis vel rarius baccatus, placentæ quatuor tandem liberae vel inter se aut cum valvularum marginibus vel cum columnâ centrali coalitæ. Semina sœpius numerosa albuminosæ, embryo rectus rarius curvatus, rarissime (*Campylanthus*) cyclicus.

TR. I. VERBASCEÆ Bth. in D.C. Prodr. — Corolla rotata subirregularis. Stamina declinata, antheræ loculorum confluentia uniloculares, stamen quintum interdum perfectum adest. Inflorescentia simpliciter centripeta. Folia omnia alterna.

VERBASCUM (L. Gen. p. 97).

Calyx 5-fidus vel partitus. Corolla explanato-rotata rarius concava lobis 5 plus minus inæqualibus. Stamina 5 filamentis inæqualibus posticis tribus vel omnibus lanato-barbatis. Antheræ uniloculares omnes filamento transverse insertæ reniformes vel illæ filamentorum binorum inferiorum filamento longitudinaliter adnatæ decurrentes. Stylus unicus, stigma clavatum. Capsula septicida apice bivalvis bilocularis. Semina lacunoso-rugosa. — Herbæ monocarpicæ vel perennes rarius suffrutices. — Genus vastum ob formarum hybridarum (hic neglectarum) copiam difficile. Calycis longitudinem semper ex flore nec ex fructu indicavi.

Specierum Orientalium distributio.

SECT. I. THAPSUS Bth. in D.C. Prodr. X, p. 225. — Staminum longiorum antheræ adnato-decurrentes.

§ 1. **Euthapsi.** — Flores fasciculati vel glomerulati in racemos dispositi.

* Filamentorum lana albida vel lutescens.

+ Flores subsessiles.

V. Thapsus, thapsiforme, phlomoides, campestre, Lagurus.

+ + Florum pedicelli calyci æquilongi vel longiores.

V. Prusianum, Guicciardii, Armenum, longifolium.

* * Filamentorum lana purpurea vel violacea

V. epixanthinum, fœtidum, malacotrichum.

§ 2. **Blattariæ.** — Flores solitarii racemosi vel spicati.

* Filamentorum lana albida vel lutescens.

V. subnivale.

* * Filamentorum lana purpurea vel violacea.

+ Flores subsessiles.

V. ovalifolium, Georgicum, alceoides.

+ + Flores pedicello calyci æquilongo vel longiore suffulti.

V. spectabile, Punalense, Wiedemannianum, macrocarpum, Blattaria, microcarpum.

SECT. II. LYCHNITIS Bth. in p.c. Prodr. X, p. 230. — Antheræ omnes reniformes subæquales.

§ 3. Thapsoidæ Bth. loc. cit. char. mutato. — Flores fasciculati vel glomerati sessiles vel subsessiles in racemos dispositi. Plantæ plus minusve tomentosæ.

* Calyces majusculi 3-5 lineas rarius $2\frac{1}{2}$ lin. longi.

+ Filamentorum lana albida vel lutescens.

V. glomeratum, *Antiochium*, *Amanum*, *salviafolium*, *simplex*, *Syriacum*, *leptocladum*, *pinnatifidum*, *plicatum*.

+ + Filamentorum lana purpurea vel violacea.

V. ptychophyllum, *Hohenackeri*, *Boerrhavii*, *voleanicum*, *Lydium*, *Casarum*, *Galileum*.

* * Calyces 2 lineas longi vel breviores.

V. mucronatum, *Antinori*, *detersile*.

§ 4. Glomerulosa. — Flores pedicello calycis longitudinem non excedente siffulti fasciculati vel glomerulati in racemos dispositi. Calyces saepius parvi raro 3 lineas longi. Plantæ tomentosæ.

+ Filamentorum lana albida vel flavescentia.

V. gnaphalodes, *eriorhabdon*, *pycnostachyum*, *Sinaiticum*, *Daenense*, *Khorassanicum*, *stachydiforme*, *lasianthum*, *erianthum*, *leucophyllum*.

+ + Filamentorum lana purpurea vel violacea.

V. pumilum, *elegans*, *Tauri*, *Adeliae*, *Tripolitanum*, *sinuatum*.

§ 5. Lychnitidea Bth. l. cit. p. 255 char. submutato. — Flores fasciculati pedicellis calyce longioribus siffulti, racemosi. Plantæ plus minusve tomentosæ vel tomentellæ. C. rotula extus plus minus pilis stellatis tomentella.

* Calyx minutus ($1\frac{1}{2}$ - $1\frac{1}{2}$ lin. longus).

+ Filamentorum lana albida vel flavescentia.

× Capsula globosa, ovata vel oblonga.

V. pulverulentum, *mallophorum*, *Olympicum*, *orgyale*, *Lychnitis*, *Delticium*, *speciosum*, *cheiranthifolium*, *humile*, *Heldreichii*, *petiolare*.

× × Capsula oblongo-cylindrica.

V. stenocarpum, *myriocarpum*, *Cedreti*.

+ + Filamentorum lana purpurea vel violacea.

V. Damascenum, *nigrum*, *glabratum*, *Bithynicum*, *Orientale*.

* * Calyx major $2\frac{1}{2}$ -4 lin. longus. Filamentorum barba albida vel flava

V. myrianthum, *sphenandroides*, *cestroides*, *phyllostachyum*

§ 6. **Racemifera.** — Flores fasciculati, panicula ad racemos axillares sessiles reducta. Filamenta glabra.

V. racemiferum.

§ 7. **Triflora.** — Pedunculi racemosi sæpius bini elongati, altero unifloro, altero trifloro et opposite bibracteolato.

V. Mesopotamicum, globiflorum, Szovitsianum.

§ 8. **Leiantha** Bth. loc. cit. p. 239. — Flores fasciculati, bini, vel solitarii in racemos dispositi. Calyx minimus glaber vel glabrescens lineam vel sesquilineam rarius excedens. Plantæ parte inferiori et ad folia sæpius valde tomentosæ paniculæ glabré sæpius viscidâ subglandulosâ. Corolla glaberrima sæpius glandulis pellucidis punctata. Grex species floribus fasciculatis et solitariis sed corollæ charactere et habitu conjunctas amplectens.

* Capsula globosa vel ovata.

V. parviflorum, nupifolium, Germanicum, denudatum, Isauricum, kochiae-forme, rubricaulis, adenocarpon.

* * Capsula oblongo-cylindrica vel cylindrica.

V. Græcum, cylindrocarpum, Smyrneum, Cilicum, leianthum, calvum, lœtum.

* * * Capsula non nota.

V. Pmardi, cerinum.

§ 9 **Blattarioidea** Bth. loc. cit. p. 241 (char. metato). — Flores solitarii, rarissime 1-2 inferiores racemi subbini. Plantæ herbaceæ. Axis determinata caulis vel ramis in racemos abeuntibus.

* Flores sessiles.

V. Steveni.

* * Flores pedicellis calyce brevioribus vel eo æquilongis suffulti.

V. pyramidatum, chrysorrhacos, Schimperianum, Basianicum, Tiberiadis, Aleppense, infideatum, digitalifolium, alyssitolum.

* * * Flores pedicellis calyce pluries longioribus suffulti.

V. Euphraticum, leuconerum, lyratifolium, Berytem, Blancheanum, adenophorum, biscutellafolium, phœnicum, scaposum, xanthophœnicum

§ 10 **Spinosa.** — Pedicelli solitarii racemosi demum cum axi racemis persistentes indurati spinosi. — Suffrutex erinaceus

V. spinosum.

§ 11. **Phlomopsidea.** — Pedicelli solitarii breves racemosi. Axis indeterminata ramis lateraliter ex axillis superioribus racemos paucos edentibus.

V. Pestalozza.

SECT. I. THAPSUS.

§ 1. Euthapsi.

Filamentorum lana albida vel lutescens.

1. **V. Thapsus** (L. Sp. p. 252 ex Fries) biennne dense lutescenti-tomentosum simplex vel parce ramosum, foliis radicalibus oblongis crenulatis petiolatis, caulinis a folio ad folium decurrentibus, floribus fasciculatis brevissime pedicellatis spicam densam vel basi interruptam formantibus, calyce tomentoso (circ. 4 lin. longo) ad $\frac{2}{3}$ in lacinias lanceolatas fisco, corollâ parvâ concavâ, antheris staminum inferiorum longiorum glabrecentium breviter adnato-decurrentibus, breviorum albo-lanatorum reniformibus, capsulâ ovatâ calyce sublongiore ②. Ic. Engl. bot. tab. 540. — *V. Schraderi* Mey. Chl. Hann. p. 326 Rehb. Ic. tab. 16.

Hab. in siccis specie sequenti multo rarius, insula Zacynthus (Marg!), Caucasicus sept. (C.A. Mey!), Ossetia (G. Koch!), Transcaucasia (Ledeb!).

Ar. Geogr. Europa præsertim borealis et media, jugum Himalaicum (ex cl. Bth).

2. **V. thapsiforme** (Schrad. Mon. I, p. 24) biennne, dense lutescenti-tomentosum simplex vel rarius ramosum, foliis radicalibus oblongo-ellipticis vel elliptico-lanceolatis crenulatis in petiolum attenuatis, caulinis a folio ad folium decurrentibus, floribus fasciculatis brevissime pedicellatis spicam longam densam formantibus, calyce tomentoso (4-5 lin. longo) ad $\frac{2}{3}$ in lacinias lanceolatas fisco, corollâ magnâ rotatâ planiusculâ, antheris staminum inferiorum longiorum glabrecentium longe adnato-decurrentibus breviorum albolanatorum reniformibus, capsulâ ovatâ calyce sublongiore ②. Ic. Rehb. tab. 17. — *V. cuspidatum* Schrad. Mon. I p. 23 tab. I. — *V. macrurum* Ten. Nap. III, p. 216. Fl. Nap. tab. 214.

Hab. in siccis montanis, mons Malevo Laconiae (Heldr! Orph! Fl. Gr. exs. 735!), Creta (Heldr! sub *V. Schraderi*), Tauria (M. B. Stev!).

Ar. Geogr. Europa tota media et australis a Daniâ et Gothiâ ad Rossiam mediam.

3. **V. phlomoides** (L. Sp. 253) biennne dense albido vel lutescenti-tomentosum simplex vel superne ramosum, foliis radicalibus crenulatis ellipticis vel oblongo-lanceolatis in petiolum attenuatis, caulinis amplexicaulibus sape cordatis vel brevissime decurrentibus, floribus fasciculatis brevissime pedicellatis racenum longum parte

inferiori interruptum et saepe laxum formantibus, calyce tomentoso (circ. 4 lin. longo) ultrâ medium in lacinias triangulare-lanceolatas fiso, corollâ explanatâ, filamentis brevioribus albido-lanatis, longiorum antheris decurrentibus, capsulâ ovatâ calyce sesquiloniore ②. Ic. Rehb. tab. 18. — Fl. græca tab. 224 ob lanam filamentorum purpuream subdubia. — *V. australe* Schrad Mon. tab. 2. — *V. montanum* Griseb. Spic. II, p. 49! non Schrad.

Hab. in siccis montanis, Thessalia (Heldr! Orph!), mons Dirphys Eubeæ (Heldr!), Macedonia prope Vodena (Orph! Griseb!) in monte Korthiati (Heldr!), insulâ Tenos Archipelagi (Sart!), Tauriâ (M.B. Stev!), prov. Caucasica (M.B., C.A. Mey!).

β. Sartorii. — Florum glomeruli omnes dissiti saepe densius tomentosi. — *V. Sartorii* Boiss. et Heldr. Diagn. Ser. I, 7 p. 38.

Hab. in olivetis regionis inferioris montis Ithomes Messeniae (Heldr!), monte Pentelico Attice (Heldr!), insulâ Zacinthro (Marg!), Macedonia (Friv. exs. sub *V. Bocrrhavii*!).

Varietas 3 habitu plane refert *V. glomeratum* Boiss. a quo antherarum et calycis characteribus plane differt. — *Verbascum crenatifolium* Boiss. Diagn. Ser. I, p. 51 extra fines ditionis nostræ in Bulgariâ ad Danubium observatum foliis profunde erenatis, caulinis numquam decurrentibus specifice differre videtur. *V. phlomoides* Fl. Gr. tab. 224 ob filamentorum lanam purpuream et antheras omnes reniformes forsitan non spectat.

Ar. Geogr. Europa media et australis a Galliâ et Germaniâ ad Rossiam medianam.

4. *V. campestre* (Boiss. et Heldr. Diagn. Ser. I, 12 p. 5) biennne tenuiter tomentoso-canescens simplex, foliis minute crenulatis inferioribus oblongo-lanceolatis in petiolum attenuatis, caulinis parvis strictis oblongo-linearibus basi subattenuatis brevissime decurrentibus vel basi rotundatis, floribus subsessilibus fasciculatis spicam simplicem basi interruptam formantibus, bracteis linearibus calyce aequantibus, calyce tomentoso (circ. 4 lin. longo) ad $\frac{3}{4}$ in lacinias lanceolatas acutas partito, filamentis flavido-lanatis, antheris staminum longiorum breviter decurrentibus, capsulâ globosâ calyci aequi-longâ detersile tomentosâ ②.

Hab. in campus prope pagum Karkin ad orientem arbis Kouiah Lycaoniæ in Anatoliâ (Heldr!).

Affine *V. phlomoides* et præsertim *V. viminali* Guss., planta cum racemo pedalis vel vix sesquipedalis, folia radicula cum petiolo semipedalia 1- $\frac{1}{2}$ poll. lata. Foliis vix decurrentibus et calyce dimidio minori a *V. viminali* distinctum.

5. *V. Lagurus* (F. et M. Ind. v. Petrop. p. 42) biennne tomento niveo denso in parte inferiori plantæ tandem subdetersili obsitum, foliis radicalibus ovato-oblongis in petiolum attenuatis acutiusculis obsolete crenatis, caulinis sensim diminutis ovatis acutis basi rotundatis vel subcordatis, spicâ simplici vel ramosâ interrumpâ lanâ longissimâ niveâ calyces et bracteas omnino occultante vestitâ, floribus subsessilibus fasciculatis, bracteis ovatis acutis calyces aequantibus,

calyce (3 lin. longo) fere ad basin in lacinias late lanceolatas elevatim plurinervias acutas partito, corollâ planâ calyce quadruplo majore, filamentis sordide albo-lanatis longiorum antheris adnatis, capsulâ ovatâ lanatâ calyci æquilongâ ②. *V. bombyciforme* Boiss. Diagn. Ser. I, 4 p. 52.

Hab. circa Byzantium (Wied!), supra Broussam ad Olympi radices (Auch. 2441! Boiss! Pichlert!).

Planta 3-4-pedalis, folia radicalia cum petiolo saepe pedalia, corolla diam. 4-4 $\frac{1}{2}$ -pollicaris. Species lanâ niveâ demissimâ totam inflorescentiam corollis exceptis occultante insignis.

6. V. Prusianum (Boiss. Diagn. Ser. I, 7 p. 37) biennne elatum superne ramosum indumento floccoso detersili obsitum tandem virescens, foliis amplis subrepandis ovatis vel oblongis petiolatis preter infima acuminatis, superioribus valde diminutis acuminatis-simis breviter auriculato-decurrentibus, racemis longissimis virgatis glanduloso-viscidis laxis, floribus fasciculatis pe. icellis calycei subæquilongis suffultis, bracteis auguste linear-i-lanceolatis pedicello longioribus, calyce (3 lineas longo) ad basin usque in lacinias anguste linear-i-lanceolatas acutas partito, corollâ amplâ explanatâ, filamentis omnibus flavescenti-lanatis antheris longiorum adnatis, capsulâ majusculâ oblongâ mucronatâ glabrescente calyce tertiatâ parte longiore ②. *V. Garganicum* var. *heterophyllum* Griseb. Spic. II, p. 49 (forma alpina caule simplici, foliis minoribus).

Hab. in castanetis ad radices Olympi Bithyni (Boiss!), in monte Scardo Macedoniae (Griseb!).

Caulis 3-4-pedalis, inflorescentiae rami bipedales, folia inferiore saepe pedalia. Species visciditate, dentibus calycinis longis, capsulâ elongatâ 4-4 $\frac{1}{2}$ lineas longâ inter affines notabilis.

Ar. Geogr. Montenegro.

7. V. Guicciardii (Boiss. et Heldr. Diagn. Ser. II, 6 p. 127) biennne simplicissimum indumento tenui subdetersili obsitum viridi-cinerascens, foliis subrepandis infimis oblongis cuneatis longe petiolatis, caulinis sessilibus basi rotundatis, floribus breviter pedicello calycei æquilongo suffultis dissite glomerulatis in spicam longam laxam dispositis, bracteis calyces æquantibus lanceolatis, calyce tomentoso (3 lineas circ. longo) ad medium in lacinias triangulares acutas fiso, corollâ amplâ explanatâ, filamentis fere ad apicem albo-barbatis majorum binorum antheris decurrentibus, stylo clavato, stigmate decurrente, capsulâ glabratâ sphærica calyce sublongiore ②.

Hab. in dumosis ad radices Parnassi loco Pagera dicto 2000' (Guicc. in Heldr., Herb., Norm. 696!).

Foliorum inferiorum limbus 4-5 pollicaris, spica saepe bipedalis, corolla diam. 15-16 lineas lata, capsula magu. Pisi. Racemo longo laxo affine *V. Prusiano*, sed calycis minoris laciniae latæ breves, capsula sphærica dimidio minor.

8. V. Armenum (Boiss. et Ky. in Ky. exs. 1859) totum adpresso sissime et molliter canescens, caule elato crasso angulato superne thyrsoides ramoso. foliis subduplicatim crenatis radicalibus amplis oblongo-lanceolatis basi breviter attenuatis, caulinis diminutis acuminatis, floralibus parvis ovatis in acumen contractis, floribus fasciculatis in racemos longos inferne interruptos dispositis, pedicellis calyci æquilongis, calyce dense tomentoso (3 lin. circ. longo) fere ad basin in lacinias lanceolatas partito, corollâ flavâ mediocri, filamentis (e sicco) griseo-lanatis antheris longiorum decurrentibus, capsulâ ... ②.

Hab. in alveis torrentium inter Kirikan et montem Bingöldagh in districtu Warto Armeniae australis 7000' copiosum (Ky. 518!).

Valde elatum, folia inferiora interdum sesquipedalia 4-5 pollices lata, corolla non magna. Facies *V. glomerati* et affinum sed antheræ binæ decurrentes.

9. V. longifolium (Ten. Fl. Nap. I, p. 89 tab. 21) bienne lutescenti vel albido-tomentosum, spicâ simplici vel breviter et stricte ramulosâ, foliis inferioribus longe oblongo-lanceolatis, caulinis oblongis acuminatis subcordato-amplexicaulibus, floralibus fasciculatis dense spicatis, pedicellis calyce 2-3-plo longioribus, calyce tomentoso (tres lineas longo) ad ¾ in lacinias lanceolatas acutas partito, corollâ explanato-concavâ, filamentis superioribus albido-lanatis inferiorum antheris longe decurrentibus. capsulâ ovoideâ calyce sublongiore tomento detersili obtectâ ②. *V. pannosam* Vis. et Panç. Plantæ Serbiceæ exs.

Hab. in regione inferalpinâ montis Scardi (Peristeri) prope Bitolia in Macedonia 3500'-4600' (Griseb!).

A *V. phlomoide* foliorum radicalium longitudine et floribus longiuscule pedicellatis nec subsessilibus distinctum.

Ar. Geogr. Italia australis, Serbia.

** Filamentorum lana purpurea vel violacea.

10. V. epixanthinum (Boiss. et Heldr. Diagn. Ser. I, 7 p. 89) bienne, caule subsimplici, foliis tomento denso adpresso superne æruginoso-aureo subtus cano-grisco obsitis radicalibus oblongis in petiolum brevem attenuatis, caulinis paucis sessilibus aentis, bracteis linearibus pedicellatos æquantibus, floralibus fasciculatis subsessilibus laxo et longe spicatis, calyce tomentoso (2 lineas tantum longo) fere ad basin in lacinias lanceolatas acutas partito, corollâ basi violaceo maculatâ, filamentis violaceo-barbatis. longiorum antheris modice decurrentibus, capsulâ ovatâ tomentosâ mucronatâ calyce tertiatâ parte longiore ②.

Hab. in regione superiori Taygeti et montis Olenos Peloponnesi (Heldr!), Parnassi in regione abietinâ et alpina (Heldr! Orph!).

β. *Samaritanii*. — Indumentum tenuius subdetersile, flores paulo longius pedicellati. — *V. Samaritanii* Heldr. in Boiss. Diagn. Ser. II, 6 p. 127.

Hab in pascuis montis Tymphresti (Velugo) .Etolim 5000' (Samar. et Guice!).
Pedale et bipedale, fola radicalia cum petiolo 4-5-pollicaria, corolla diam.
pollicaris.

11. V. foetidum (Boiss. et Heldr. Diagn. Ser. II, 3, p. 141) bienne, tenuiter tomentosum flavo-virens, caule elato ramoso, foliis radicalibus amplis repando-crenatis ovatis vel oblongis breviter petiolatis, superioribus basi rotundata vel cordata sessilibus, floribus fasciculatis racemos longes laxos formantibus, pedicellis calyci aequi-longis, bracteis linearibus, calyce lanâ detersili obsito (3 lineas longo) fere ad basin in lacinias lanceolatas partito, filamentis violaceo-barbatis longiorum antheris totâ longitudine adnatis, capsula ovatâ mucronatâ calycinis laeiniis duplo longiore ②.

Hab. in regione alpinâ superiore Parnassi ad Dipotamo (Heldr!).

Pluripedale, odor totius plantæ fere *Daturæ Stramonii*, racemi 1-2-pedales Valde affine *V. epixanthino* a quo differt ramositate, foliis majoribus, floribus pedicellatis, calyce majori et præsertim antheris duabus totâ longitudine nec breviter adnatis. An tamen hujus varietas sit ex loco pinguiori et numeroso orta in locis natalibus investigandum.

12. V. malacotrichum (Boiss. et Heldr. Diagn. Sc. II, 3, p. 142) bienne, totum lanâ albâ floccosâ tandem detersit absitum elatum ramosum, foliis suberuatis inferioribus oblongis in petiolum brevem attenuatis cæteris basi rotundata vel cordata sessilibus, floribus fasciculatis breviter pedicellatis racemos longos laxos virgatos formantibus, bracteis lanceolatis calyces aequantibus, calyce (2 1/2 lin. longo) lanâ detersili obsito ad 3/4 in lacinias elliptico-lanceolatas acutas partito, filamentis violaceo-barbatis, antheris longiorum longe adnato-decurrentibus, capsula globosâ calycem non suprante ②. *V. heteromallum* Pančić MSS.

Hab. in regione sylvaticâ Olympi Thessali prope Hagio Dionysi (Heldr!), in fauces montis Korthiati Macedoniae (Heldr! Orph. Herb. Norm. № 734!).

Affine *V. foetido* quod capsulis duplo majoribus ovatis calycem excedentibus differt

Ar. Geogr. Serbia.

§ 2. Blattariæ.

13. V. subnivale (Boiss. et Haussk.) bienne, totum tenuiter tomentosum lutescens, caule procero simplici in racemum longum laxum abeunte, foliis obtuse crenatis subtus intensius canescens radicalibus amplis oblongis basi breviter cuneatis longe petiolatis, canlinis 2-3 diminutis præter sumimum sessile breviter petiolatis, floribus præter infima rarius bina solitariis pedicello calyci aequilongo suffultis, bracteis linearibus pedicello subaequilongis, calyce tomentoso (3 lineas longo) ad 2/3 in lacinias oblongo-lanceolatas acutas fissis, corollâ luteâ tomentellâ, filamentis sordide albo-barbatis anticis duobus elongatis præter basin barbatam nudis antheris longe decurrentibus, capsula... ②.

Hab. in graminosis regionis alpine montis Berydaghi Cataoniæ 8-9000. (Haussk!) Fl. Ang.

Bi vel tripedale, racemus tandem $\frac{1}{2}$ -1-pedalis, folia inferiora $\frac{1}{2}$ -1 pedem longa 3-4 pollices lata. Affine *V. ovalifolio* et *Georgico* a quibus præter alias notas floribus pedicellatis differt.

14. ***V. ovalifolium*** (Don in Sims Bot. Mag. tab. 1037) biennæ tomentoso-griseum subsimplex, foliis subtus incanis grosse et sub-duplicatim crenatis ovatis magnis cordatis petiolatis, superioribus cordato-amplexicaulibus, floribus subsessilibus solitariis magnis spicam basi interruptam formantibus, bracteis ovato-triangularibus acuminatis calycem subæquantibus, calyce dense albo-tomentoso (5 lin. longo) ad $\frac{3}{4}$ in lacinias late lanceolatas partito, corollâ amplâ, filamentis purpureo-lanatis longiorum glabrescentium antheris adnatis, capsulâ... ②. *V. compactum* M.B. Taur. Cane. I, p. 159 — *V. formosum* Fisch. Cat. Gor. — Bot. Reg. tab. 558. — *V. Orientale betonicæfolio flore maximo* Tourn. Cor. sed *V. betonicæfolium* Desf. Gor. tab. 23 ob flores parvos videtur potius *V. nigrum*. — *V. saccatum* C. Koch Linn. XVII, p. 283.

Hab. in Armeniâ Turcicâ (Tourn.), Rossicâ prope Erivan (C. Koch!), Iberiâ Caucasicâ (M.B. Hohen!). in Tauriâ (ex M.B.).

Spica et calyx crassi tomentosi fere *Alceæ*, corolla diam. sesquipollicaris, species distinctissima.

15. ***V. Georgicum*** (Bth. in D.C. Prodr. X, p. 228) biennæ tomento adpresso canescens subsimplex, caule mediocri in spicam brevem densam abeunte, foliis integris inferioribus petiolatis obovatis vel oblongis, superioribus brevibus oblongo-lanceolatis semiamplexicaulibus, floribus subsessilibus solitariis approximatis, calyce valde tomentoso ($3\frac{1}{2}$ lin. longo) in lacinias lanceolatas acutas partito, filamentis purpureo-lanatis longiorum glabrescentium antheris decurrentibus ②. *V. molle* C. Koch Linn. XVII, p. 284!

Hab. in Georgia Caucasicâ (Wilhelm. in D.C. herb!), Armeniâ Rossicâ in latere orientali montis Alagos (Koch!). Habeo specimen ex Armeniâ circa Erzerum (Huet!) et Persicum ex viciniis Persepolis (Ky. 806!) quæ eamdem speciem sistere videntur.

Sesquipedale, differt a *V. ovalifolio* foliis integris multo minoribus (radicalibus cum petiolo 3 pollices longis), flores præterea dimidio minores esse videntur. Planta ut videtur rara, an species legitima, an *V. ovalifolio* proles hybrida sit investigandum.

16. ***V. alceoides*** (Boiss. et Haussk.) perenne pluricaule, totum tomento albo vel flavido ad folia radicalia densiore pannoso ad caulinam tenuiore obsitum, caulis inferne foliosis cito in racemum unicum vel 2-3 longos crassos spicæformes rigidos abeuntibus, foliis oblongo-ellipticis subcrenatis obtusis subtus elevatim nervosis radicalibus petiolatis caulinis sessilibus, floribus subsessilibus solitariis bracteâ linearî in inferioribus calycem æquante suffultis, calyce crasse

tomentoso ($3\frac{1}{2}$ lineas longo) ultrâ medium in lacinias oblongas acutiusculas partito fructifero aucto, corollâ tomentosâ, filamentis violaceo? barbatis, antheris binis decurrentibus, capsulâ ovato-globosâ demum glabrâ reticulatâ calyce subbreviore ♂.

Hab. in graminosis montis Pîr Omar Gudrun Kurdistaniae Persiae (Haussk.)

Folia inferiora 3-4 pollicaria 12-15 lineas lata, racemi crassi rigidi pedales, capsula piso sub major. Species insignis calycibus fructiferis magnis tomentosis *Alcea* alabastra referentibus.

17. V. spectabile (M.B. Taur. Can. III, p. 158) biennie laxe et tenuiter tomentellum cinereo-virens elatum simplex vel parce ramosum, foliis radicalibus et inferioribus subduplicato-crenatis longe petiolatis oblongis cordatis obtusis, superioribus sessilibus sensim diminutis cordatis longe acuminatis, racemo longo laxinseulo spiciformi viscoso-villosulo, pedicellis filiformibus præter intimos subgeminos solitariis calyce sublongioribus, bracteis inferioribus triangularibus cuspidatis supremis subulatis pedicello brevioribus, calyce demum glabrescente (3 lin. circ. longo) ad $\frac{3}{4}$ in lacinias lanceolatas acutas partito, corollâ magnâ luteâ basi purpureo-maculatâ, filamentis purpureo-lanatis longiorum antheris decurrentibus, capsulâ ovato-oblongâ tomentosâ mucronatâ calyce sublongiore ②.

Hab. in montanis asperis Tauriae meridionalis circa podium Sobla, ad fontes fluv. Salgir, supra Ulu Usen (M.B. Stev. Rehm!).

Planta 2-4-pedalis, foliorum inferiorum limbus 2-4-pollicaris, capsula 4 lin. longa.

18. V. Punalense (Boiss. et Buhse Aufz. p. 161) biennie elatum pyramidatim ramosum, foliis crenatis supra virentibus subtus cinereis, radicalibus..., caulinis a basi sessili cordatâ oblongis acutiusculis, paniculâ glanduloso-viscidâ pubeque patulâ crispâ hispidulâ, floribus solitariis pedicello calyce æquilongo suffultis laxiuscule racemosis, calyce ($2\frac{1}{2}$ lineas longo) ad $\frac{3}{4}$ in lacinias oblongas partito, filamentis purpureo-lanatis, daobus longioribus superne nudis antheris adnatis, capsulâ... ②.

Hab. in prov. Ghilan Persiae borealis prope Punal ad rivum Tchafrud (Buhse).

Folia radicalia probabiliter cordata, facies *V. pyramidati* vel *V. nigri*, sed ob antheras binas decurrentes affine *V. spectabile* a quo differt paniculâ pyramidatâ, calycis laciiniis latioribus brevioribus, etc.

19. V. Wiedemannianum (Ind. IV Petrop. p. 51) biennie lanâ arachnoideâ adpressâ tandem subdetersili obsitum, caule crasso elato inferne folioso in racemum simplicem vel in paniculam parce ramosam abeunti, foliis lanuginosis radicalibus oblongis vel ellipticis crenatis vel subintegris in petiolum attenuatis, caulinis diminutis oblongis sessilibus, floribus solitariis in racemos laxos tandem longissimos dispositis, pedicellis calyce et bracteâ lanceolato-lineari brevioribus, calyce dense glanduloso-hirto ($4\frac{1}{2}$ lin. longo) in laci-

nias lanceolato-lineares fere ad basin partito, corollâ magnâ intense purpureâ glabrâ, filamentis longioribus præter basin nudis antheris longe adnatis, brevioribus purpureo-barbatis, capsulâ pedicello incrassato stricto sublongiore ovato-oblongâ acutiusculâ hirtâ calycem quartâ parte superante ②. *V. insigne* Boiss. et Bal. Diagn. Ser. II, 6, p. 129.

Hab. in Anatoliâ boreali prope Toptscha, Tokat et in monte Aladagh (Wied!), Ponto ad meridiem Samsun (Tchih!), Armeniâ prope Baibut (Bourg!), Cappadociâ prope Karahissar et ad radices montis Dededagh (Bal!).

Planta 4-3-pedalis, folia inferiora 3-5-pollicaria, racemos fructifer sœpe bipedalis, corolla diam. 15 lin. lata, capsula piso duplo major. Species floribus intense purpureis elegantissima.

20. ***V. macrocarpum*** (Boiss. Diagn. Ser. I, 12, p. 6) biennie inferne pube papillosâ crispulâ obsitum inflorescentiâ valde glanduloso-viscidâ, caule tereti valde elato folioso in racemum tandem longissimum abeunte, foliis tenuiter membranaceis viridibus radicalibus oblongis in petiolum attenuatis grosse et subduplicatim dentatis versus basin sepe lobatis, caulinis crebris sursum diminutis, superioribus ovato-oblongis cordato-auriculatis serratis longe attenuato-acuminatis, floribus solitariis laxiuscule racemosis pedicellis calyce subbrevioribus suffultis, bracteis triangulari-lanceolatis acuminatis inferioribus calyce ræquantibus vel superantibus; calyce valde glanduloso-hirto (tres lineas longo) in lacinias ellipticas acutiusculas profunde partito, corollâ luteâ extus glandulosâ, filamentis purpureo-barbatis longiorum antheris decurrentibus, capsulâ magnâ ovatâ glandulosâ calyce quartâ parte longiore ②.

Hab. in regione inferiori montis Elbrus Persiæ hor. prope Derbend (Ky. 715! Buhsé!), in Persiâ australi (Ky. 926!), in pomariis montis Avroman 4-5000' et in Elymaitide (Haussk!).

Planta 4-8-pedalis, folia inferiora 4-6-pollicaria, racemos fructifer 1 $\frac{1}{2}$ -3-pedalis, capsula piso duplo major. Aflne *V. virginato* = *V. Celsiae* Boiss. foliis parvis rigidis, bracteis minutis, calyce multo minore, etc. distincto.

21. ***V. Blattaria*** (L. Sp. 254) biennie glabrum viride, caule elato tereti folioso simplici vel superne parce ramoso, foliis oblongis et oblongo-lanceolatis dentatis, inferioribus in petiolum attenuatis indivisis vel sinuato-pinnatifidis, superioribus sessilibus cordato-amplexicaulis acutis, racemo longissimo laxo minute glanduloso, floribus solitaris pedicello calyce 2-3-plo longiore suffultis, bracteis lanceolatis acuminatis pedicello brevioribus, calyce glanduloso ($2\frac{1}{2}$ lin. longo) ad $\frac{3}{4}$ in lacinias linear-lanceolatas partito, corollâ majusculâ flavâ fauce violacea rarius albâ, filamentis violaceo-barbatis, antheris duabus decurrentibus, capsulâ globosâ pedicello eâ longiore suffulta calyce longiore ②. Ic. Engl. Bot. tab. 393. — Rchb. Ic. 32. — *V. repandum* Willd. Enum. p. 226 (forma ramosior corollâ et capsulis subminoribus).

Hab. in pinguibus et agris humidis regionis inferioris et montane totius ditionis a Grecia et Macedonia ad Tauriam, Transcaucasiam! Persiam borealem (Buhsel!), Affghaniam (Griff!). E Syria et Egypto non vidi.

Ar. Geogr. Europa omnis borealis et media ab Anglia et Belgio ad Rossiam medium, Hispaniam et Italiam boreales, Dalmatiam, Sibiria Altaica, Africa borealis.

22. V. microcarpum (Bth. in D.C. Prodr. X, p. 230) biennie, caule inferne pubescenti superne glabro a basi valde ramoso ramis in racemos virgatos laxissimos longos rigidos abeuntibus, foliis pilis stellatis sparsis adpressis obsitis pallide virentibus obovato-oblongis obtusis inferioribus in petiolum brevem longe attenuatis, superioribus paucis sessilibus, floribus solitariis parvis remotis pedicello calyci aequilongo demum sursum arcuato et rachidi adpresso suffultis, bracteis minimis triangulari-ovatis, calyce paree glanduloso ($1\frac{1}{4}$ linea longo) in lacinias lineares acutas partito, filamentis violaceo-lanatis brevissimis, antheris inferioribus (ex cl. Bth.) breviter decurrentibus, capsula parvâ anguste oblongo-cylindrica obtusa calyce quartâ parte longiore ②.

Hab. in Mesopotamiâ (Auch. 2458).

Ex speciminibus imperfectis descriptum, sesquipedale racemis tenuibus fere pedalibus, facies *Hirschfeldiae* vel *Rapistri*, folia inferiora bipinnata pollicem lata, capsula 2 lineas longa. Antheras ex cl. Bth. descripti.

SECT. II. LYCHNITIS.

§ 3. Thapsoidea.

* Calyces majusculi (3-5 lineas, rarius $2\frac{1}{2}$ longi).

+ Filamentorum lana albida vel lutescens.

23. V. glomeratum (Boiss. Diagn. Ser. I, 4, p. 52) biennie totum dense pannosum griseo-flavidum elatum crassum thyrsoideo-paniculatum, foliis subintegris subsessilibus oblongo-lanceolatis basi attenuatis crassis superioribus a basi cordato-subauriculata ovatis acuminatis, floribus subsessilibus in glomerulos 3-5-floros bractea triangulari calyce breviore stipatos in spicas interruptas laxas dispositis, calyce (3-6 lin. longo) pannoso fere ad basin in lacinias crassas lanceolatas partito, corollâ luteâ extus valde tumentosâ, filamentis villo flavescente barbatis, antheris omnibus reniformibus, capsula parvâ ovalâ calycem non excedente ②. *V. Chium* Boiss. et Orph. MSS.

Hab. in collinis sicceis, ad vias Lydiae ad Smyrnam et Magnesiam (Boiss!), Pisidiâ in monte Boudroum (Heldr!), Lyciâ ad Elmalu (Bourg!), insula Chio (Orph!), Phrygiâ ad Ouchak (Bal!) prope Iconium (Heldr!), in Syria boreali prope Aintab et monte Ssoddagh (Haussk!).

Planta 4-6-pedalis, folia inferiora pedem et amplius longa 3-4 pollices lata, capsula piso dimidio minor.

Obs. *V. splendidum* Boiss. Diagn. Ser. I, 4, p. 53 et *V. ringens* Boiss. et Heldr. loc. cit. p. 13 videntur omnino formæ hybridæ (steriles) inter hanc speciem et *V. pycnostachyum* vel *V. mucronatum*, forsitan *V. lasianthum*. *V. protractum* Fenzl in Tchih. vol. 2, p. 6 = *V. lychnitis* var. *grandiflora* Bth. in D.C. Prodr. est forma hybrida *V. ringenti* similis.

24. **V. Antiochium** (Boiss. Diagn. Ser. I, 12, p. 9) biennie, caule rubello glabrescenti simplicissimo in racemum longissimum valde interruptum abeunte, foliis obsolete crenatis supra tomentellis virentibus subtus griseo-tomentosis, radicalibus oblongis basi cuneatis petiolatis, caulinis parvis subcordato-amplexicaulibus, glomerulis 4-6-floris dissitis, floribus sessilibus, calyce (3-4 lineas longo) pube stellata flavidâ tomentello et glanduloso ad $\frac{2}{3}$ in lacinias rigidulas erector-patentes triangulari-lanceolatas acutas partito, corollâ mediocri luteâ, filamentis flavidâ-lanatis, capsulâ majusculâ ovatâ mucronatâ calycem parum excedente ②.

Hab. circâ Antiochiam ad rupes et mænia (Auch. 2449 sub *V. simplici!*, Boiss!) in Libani ditione Daunié (Bl!), in Antilibano inter Rascheya et Damascum (Boiss!), in Syriâ boreali (Haussk!).

Planta cum racemo saepe bipedali quadripedalis, foliorum radicalium lamina 4-5-policaris, capsula piso major. Specimina Aucheriana e Syriâ in D.C. Prodr. ad *V. gnaphalodem* citata huc vel ad *V. Syriacum* spectant. *V. gnaphalodes* indumento detersili niveo aliisque notis longe differt.

25. **V. Amanum**, biennie inferne tomento tenui ad paginam inferiorem copiosiore virenti-griseum secus paniculam et flores pube ramosâ longâ laxâ hispidissimum, caule elato ample paniculato racemis ramulosis, foliis ovato-oblongis obsolete crenatis inferioribus petiolatis cordatis, caulinis diminutis cordato-amplexicaulibus acuminate, florum subsessilium fasciculis præsertim inferioribus dissitis, calyce hispido (2½ lineas longo) fragili ad basin usque in lacinias linear-lanceolatas acutissimas basi subangustatas fisso, corollâ parvâ, filamentis omnibus sordide et longe albo-hirsutis, capsulâ minutâ oblongâ calyce paulo breviore ②. *V. Antiochium* var. β. in Ky. exs. 1862.

Hab. in calcareis humidis ad templum destructum prope pagum Attyk montis Amani Syriæ bor. prope Beilan (Ky. 191).

Planta elata, folia radicalia pedem dimidium longa poll. 3-4 lata, capsula angusta 3 lineas vix longa. Species a *V. Antiochio* cui cam prius retuleram foliis cordatis, caule paniculato, capsulâ, etc. diversa.

26. **V. salviæfolium** (Boiss. Diagn. Ser. I, 4, p. 55) perenne pluricaule totum tomento niveo non detersili pannosum, caulis erectis parte inferiori foliosis in racemum simplicem vel parce ramosum abeuntibns, foliis sub tomento nervoso-reticulatis crenulatis radicalibus oblongis et oblongo-lanceolatis acutiusculis vel obtusis in petiolum longiusculum attenuatis, caulinis diminutis lanceolatis decurrentibus, fasciculis 2-3-floris remotiusculis bracteâ triangulari suffultis in racemum laxum dispositis, floribus subsessilibus, calyce

dense albo-tomentoso (4 lin. longo) ad medium in lacinias lanceolatas fisco, corollâ intense luteâ calyce $2\frac{1}{2}$ -plo latiore, filamentis usque ad apicem aurantiaco barbatis, capsulâ... ♂.

Hab. in collibus siccis Cariae circa Yenidje et Gheyra (Boiss!) inter Karagatsch et Koniah Lycaoniae (Heldr!).

Caudex induratus ramosus vestigiis petiolorum cicatricosus, caules cum racemo 6-10-pollicari pedales et sesquipedales, folia inferiora cum petiolo 3-5-pollicaria 8-12 lineas lata. Facies *V. simplicis* a quo eximie differt foliis caulinis decurrentibus, calyce majori, etc. Species corollis intense aureis elegantissima.

27. *V. simplex* (Labill. Dec. IV, p. 10, tab. 5) basi suffrutescens multicaule dense niveo vel flavescenti-pannosum, caulis inferiorne foliatis dein in racemum simplicem longum valde interruptum rarius ramulosum abeuntibus, foliis radicalibus crenulatis parvis ovato vel oblongo-spathulatis obtusissimis longe petiolatis, caulinis subsessilibus et sessilibus 2-5 in glomerulos dissitos dispositis, calyce (2 $\frac{1}{2}$ lin. longo) albo-tomentoso ad $\frac{2}{3}$ in lacinias oblongas mucronatas partito, corollâ flavâ, filamentis albo-barbatis, capsulâ tomentellâ globosâ mucronatâ calycem non excedente ♀.

Hab. ad radices orientales Libani ad Ainhata (Bl!), circa Balbeck in Cœlesti-syriâ inter Antilbanum et Libanum (Ehrl Boiss! Gaill!), prope Dimar (Gaill!), in planicie circa Aleppo (Boiss!), in deserto Syriaco ad Palmyram (Bl!).

Planta subbipedalis racemo $\frac{1}{2}$ -4-pedali, foliorum inferiorum lamina 1-2 pollicaris. Capsula Cannabeos semine vix duplo major.

28. *V. Syriacum* (Schrad. Mon. II, p. 6, tab. I, fig. I) perenne pluricaule tomento tenui adpresso cinereo obsitum, caulis superne paniculato-ramosis, foliis subcrenatis vel integris obovatis et oblongis inferioribus petiolatis superioribus sessilibus basi rotundatis acutis vel acuminatis, floribus brevissime pedicellatis 3-6 fasciculatis fasciculis remotiusculis racemos laxos formantibus, calyce tomentello (3 lin. longo) ad $\frac{2}{3}$ in lacinias lanceolatas acutas fisco, corollâ luteâ, filamentis albo-barbatis, capsulâ globosâ tuberculato-punctatâ calycem non excedente ♀. *V. Kotschyi* Boiss. Diagn. Ser. I, 4, p. 56.

Hab. in sterilibus, inter culta, circa Aleppum (Ky. 171! Boiss!), circa Marasch Aintab, Nisib, Surug et Orfa. Planta planicie et desertorum (Haussk!).

Valde affine *V. simplici* sed basi non suffruticosum videtur, caules plus minus paniculati, indumentum multo tenuius, folia majora, inferiora limbo interdum 4-pollicari, flores majores, calycis dentes non mucronati, capsula paulo major. An tamen cum *V. simplici* formis intermediis confluat ulterius investigandum.

29. *V. leptocladium* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 10) perenne pluricaule totum tenuiter et adpresso arancoso-canum, caulis inferiorne foliosis superne paniculato-corymbosis ramis longe et tenuiter virgatis, foliis radicalibus numerosis et inferioribus tenuiter crenulatis oblongo-lanceolatis acutiusculis in petiolum longe

attenuatis, superioribus paucis sessilibus linearibus, floribus 3-4 sub-sessilibus in fasciculos minutos laxe spicatos dispositis. bracteis linearibus calyces subæquantibus, calyce tomentoso (flor. $2\frac{1}{2}$ lineas fruct. 3 lineas longo) in lacinias linear-lanceolatas apice tandem glabratas partito, corollâ parvâ tomentosâ flavâ, filamentis sordide flavescenti-lanatis antheris reniformibus, capsulâ ovatâ canâ pedicello clavato triplo longiore calyci æquilongâ ȝ.

Hab. in pinetis apricis arenosis maritimis Pamphyliæ ad orientem urbis Adalia (Heldr!) Fl. Jun.

Sesquipedale specis fere pedalibus, folia inferiora cum petiolo 3-4 pollices longa 7-10 lineas lata, capsula fere 3 lineas longa. Facies *V. Aleppensis* et *V. infidellum* sed flores subsessiles fasciculati.

30. ***V. pinnatifidum*** (Vahl Symb. 2, p. 39) biennie junius floccoso-tomentosum tandem plus minusve glabratum, radice verticalli, caule humili erecto rigido a basi pyramidatim paniculato, foliis radicalibus rosulatis petiolatis oblongo-lanceolatis pinnatifidis partitique segmentis oblongis profunde et obtuse lobatis rachide lobulata, foliis caulinis et rameis parvis sessilibus oblongis et linear-lanceolatis pinnatifidis vel dentatis, floribus 2-3 sessilibus in fasciculos remotiusculos bracteâ suffultis dispositis, calyce tomentoso ($3\frac{1}{2}$ lin. longo) ultra medium in lacinias lanceolatas fisso, corollâ luteâ punctatâ, filamentis longe flavid-barbatis, capsulâ majuscûla glabratâ ovatâ acutâ calyce sublongiore ȝ. Ic. Fl. Græc. tab. 228 — *V. bipinnatifidum* Bot. Mag. tab. 1777. — *V. ceratophyllum* Schrad. Mon. 2, p. 7, tab. I, fig. 2.

Hab. in arenosis maritimis Atticæ ad Phalerum et alibi (Sprun! Heldr!), Thessalæ (Auch. 245!), insularum Eubea, Ægina et Hydra (Sprun!), Scyros (Urv!), Macedoniâ (Friv!), Byzantii (Cast!), Bithyniâ (Sibth.), Lydiâ prope Smyruam (Boiss!), Tauriâ (Stev!).

Erronea a nonnullis auctoribus basi suffruticosum ab aliis stoloniferum dictum.

31. ***V. plicatum*** (Sibth. et Sm. Fl. Gr. III, p. 21, tab. 226) biennie totum adpresso nivo vel lutescenti-tomentosum saepius paniculatum, foliis radicalibus oblongo-spathulatis undulato-plicatis sinuato-pinnatifidis lobis rotundatis subimbricatis, foliis caulinis remotiusculis cordato-amplexicaulibus ovatis acutis, floribus sessilibus 2-3 fasciculatis racemos longos interruptos formantibus, calyce tomentoso (4 lin. circ. longo) ad $\frac{3}{4}$ in lacinias lanceolatas partito, filamentis flavid-lanatis, capsulâ ovatâ acutâ laciñis calycinis æquilongâ ȝ. *V. Græcum fruticosum folio sinuato candidissimo* Toara. Voy. I, p. 335. Ic. — *V. sinuatum* var. β . L. Sp. — *V. undulatum* Lam. Diet IV, p. 221 ex parte quoad Syn. Tournef. — Rehb. Ic. tab. 38. — Jaub. et Spach. III, tab. 408.

Hab. in lapidosis collium regionis calidæ, Græcia circa Athenas (Sprun! Heldr!), insula Hydra (Sibth!), Argolis (Boiss!), Cyciales (Tourn.).

β . *rigidum*. — Indumentum tenuius griseum, folia inferiora

remote sinuata siepe vix undulata corolla subminor. — *V. rigidum* Boiss. et Heldr. Diagn. Ser. II, 3, p. 143. — *V. Gallicum* Heldr. exs.

Hab. ad radices montis Korthiati Mace-touiae (Heldr! Orph! Fl. exs. 732!), in insulâ Poros (Calauria) (Heldr!).

Præter alias notas calyce triplo majore a *V. sinuato* distinctum. Nomen Sibthorpiatum prætuli antiquiori cl. Lunarek qui in descriptione hanc speciem cum *V. ptychophyllum* Syriacō commixxuit.

Ar. Geogr. Dalmatia.

++ Filamentorum lanâ purpurea vel violacea.

32. *V. ptychophyllum* (Boiss. Diagn. Ser. I, 12, p. 8) biennne totum adpresso lutescenti-pannosum a collo pluricaule caulinis simplicibus vel 1-2 ramis brevibus auctis in racemos simplices longissimos laxos abeuntibus, foliis radicalibus ovato-oblongis obtusis in petiolum brevem attenuatis valde undulatis et sinuato-lobatis, caulinis oblongis crenato-dentatis et integris basi longe decurrentibus, floribus sessilibus 3-4 fasciculatis fasciculis remotiusculis bracteâ subcordatâ triangulari suffultis, calyce (4 1/2 lin. longo) pannoso ad 3/4 in laciniis lanceolatas partito, filamentis violaceo-lanatis, capsulâ ovalâ inuernatâ calyce vix longiore ②. *V. undulatum* Lam. Dict. IV, p. 221 et p.c. Prodr. ex parte quoad plantam Syriacam.

Hab. in Syria interiore circa Rascheyâ et Zebâlani Antilibani (Boiss! Ky. exs. 133!) prope Dommar (Gaill!), inter Antilibanum et Damascum (Boiss!), in deserto ad Palmyram (Bl!).

Folia radicalia 3-4-pollicaria, caules cum racemo tandem bipedali 8-pedales. A *V. sinuato* foliis radicalibus brevibus, calyce triplo magiore, a *V. plicato* præter alias notas staminum lanâ purpureâ distinctum

33. *V. Hohenackeri* (F. et M. Index V Petrop. p. 42) biennne, tomentosum canum simplex, foliis crenulatis inferioribus oblongis in petiolum attenuatis superioribus semidecurrentibus ovatis vel subcordatis cuspidatis, floribus 2-3 fasciculatis laxe racemosis, pedicellis brevissimis, corollâ magâ flavâ, filamentis superioribus violaceenti-harbatis, inferioribus binis sublongioribus violaceo-barbatis, antheris omnibus reniformibus, capsulâ oblongâ calycis laciniis linear-i-lanceolatas superante (②).

Hab. in montibus ditionis Tadysch (Hohen.) Specimen in Hort. Petrop. ex semiinibus Hohenack. cultum in Herb. Hort. Petrop. vidi.

Caulis elatus, facies *V. montana* et *phlomoides*, sed ab eis staminibus abhorrens. Corolla diam. 10 lineas lata, calyx 2 1/2 lineas tantum longus quâ notâ ab affinibus differt. Species ex specimine culto tautum et imperfecte notâ

34. *V. Boeerhavii* (L. Mant. p. 45) biennne, tomento floccoso tandem detersili obsitum, caule rubello elato folioso in racemum spiciformem longum rigidum inferne interruptum abeunti, foliis crenato-dentatis, inferioribus ovatis obtusis petiolatis, superioribus

cordato-amplexicaulibus acutis vel acuminatis, floribus fasciculatis subsessilibus flocco tandem evanido immersis, calyce (4 lin. longo) preter tomentum glanduloso-viscoso fere ad basin in lacinias lanceolatas fisso, corollâ magnâ luteâ ad faucem violaceo-maculatâ, filamentis violaceo-barbatis, capsulâ magnâ tandem glabratâ elliptico-ovatâ mucronatâ calyce tertiatâ parte longiore ②. Ic. Rehb. tab. 33. — *V. maiale* D.C. Fl. Fr. 6, p. 415.

Hab. in Græciâ circa Calamata (Gittard in herb. Fauché!) sed schedulis hujus collectoris pauca fides habenda.

Icon. *V. phlomoides* Fl. Græc. tab. 224 ex Byzantio ob stamna purpurea et antheras omnes reniformes huc potius spectat.

Ar. Geogr. Hispania, Gallia australis, Corsica, Sicilia, Italia, Africa borealis.

35. *V. vulcanicum* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 6) bienne, caule mediocri in racemum tomento crasso flavido obsitum simplicem vel parce ramoso-paniculatum abente, foliis parvis pilis stellatis parce et adpresse canescentibus minute crenulatis inferioribus oblongo-spathulatis in petiolum attenuatis, supremis amplexicauli-subauriculatis, floribus 2-4 sessilibus fasciculatis breviter et laxiuscule racemosis, bracteis lanceolatis acuminatis fasciculos subæquantibus, calyce dense tomentoso (3 lin. longo) in lacinias lanceolatas acutas ad $\frac{2}{3}$ partito, filamentis violaceo-lanatis, longioribus apice nudis, capsulâ..... ②.

Hab. in arenosis vulcanicis regionis inferioris montis Karadagh Lycaoniæ prope Masehri (Heldr!).

Planta pedalis, folia inferiora cum petiolo 3-4 pollices longa $1\frac{1}{2}$ lata, racemus 8-9 pollicaris. Ob capsulam ignotam imperfecte notum sed facie et filamentis purpureo-lanatis affine videtur *V. Boerhavii* a quo indumento denso non defersili totius intlorescentiæ, calyce multo breviori, etc. differt.

36. *V. Lydium* (Boiss. Diagn. Ser. I, 4, p. 62) bienne, totum tomento denso tandem in parte inferiori plantæ detersili floccosum, caule elato in racemum subsimplicem vel in paniculam abeunte, foliis tenuiter membranaceis duplicatim crenatis radicalibus amplis ovato-oblongis sessilibus basi longe angustatis, superioribus oblongis acutis late cordato-auriculatis, glomerulis 3-5 floris laxe racemosis, pedicellis calyce brevioribus, bracteis linear-lanceolatis glomeralos saepe æquantibus, calyce (3 lin. longo) floccoso prætereaque glandulosobrito in lacinias lanceolatas ad $\frac{3}{4}$ partito, corollâ flavâ basi violaceo-maculatâ, filamentis violaceo-lanatis, longioribus apice nudis, capsulâ ovatâ calyce breviore ②.

Hab. in Lydiâ a regione inferiori prope Smyrnam ad valles subalpinas Tmoli circa Bozdagh (Boiss!).

Planta 3-4-pedalis, folia inferiora pedalia 4-5 poll. lata, corolla diam. pollicaris, capsula piso tertiatâ parte minor, Indumentum floccosum.

37. *V. Cæsareum* (Boiss. Diagn. Ser. I, 12, p. 7) bienne,

caule crasso elato tereti in racemum longum crassum simplicem vel basi 1-2 ramulis auctum abeundi, foliis viridibus subtus tenuiter flavido-tomentellis elevatim nervosis radicalibus amplis oblongo-ellipticis basi rotundatis petiolatis obtusis repando-crenatis, caulinis subcordato-amplexicaulibus acutis, fasciculis 3-5-floris in racemum basi exceptâ densum dense flavido-tomentosum dispositis, floribus sessilibus, bracteis lanceolatis calycem æquantibus, calyce (4 lin. longo) ad $\frac{2}{3}$ in lacinias lanceolatas acutas partito, corollâ luteâ extus valde tomentosâ, filamentis violaceo-lanatis, capsulâ..... ②.

Hab. in vallibus circa Banias (Caesaream Philippi) in Galilæa (Boiss!) Fl. Mai.

Planta 3-4-pedalis racemo sesquipedali, foliorum radicalium lamina 10-12 pol. longa 4-5 lata. E grege *V. Boeerhavii* et *Lydii*, formâ foliorum, indumento non floccoso nec detersili ab eis distinctum.

38. V. Galilæum (Boiss. Diagn. Ser. I, 12, p. 8) biennie, tomento tenui secus inflorescentiam crassiori obsitum, caule elato in paniculam longam abeunte, foliis membranaceis viridibus amplis oblongo-lanceolatis irregulariter repando-lobatis et denticulatis insimis basi breviter attenuatis, caulinis late cordato-aureculatis et supra basin sæpe angustatis, floribus 3-5 sessilibus fasciculatis fasciculis remotis bracteis ovatis acuminatis involucratis longe et laxissime racemosis, calyce tomentoso (4 $\frac{1}{2}$ lin. longo) in lacinias lanceolatas ad $\frac{3}{4}$ partito, filamentis violaceo-lanatis, capsulâ..... ②. *V. macrophyllum* Ky. exs.

Hab. in umbrosis et pinguisibus Galilee in montibus Carmel, Thabor et ad lacum Tiberiadis (Boiss!), in Syriâ littorali circâ Tripoli (Bl!) et Tortosam (Gail!) circa Damascum (Gail!), in monte Amano prope Beilan (Ky. 17!), Ciliciâ ad cataractas Cydni prope Tarsum (Ky. 24!).

Pluripedale, folia inferiora interdum sesquipedalia, paniculæ racemi pedem et amplius longi, corolla magna. Differt a *V. Caesareo* foliis membranaceis basi attenuatis, fasciculis remotis latè bracteatis.

• Calyces 2 lineas longi vel breviores.

39. V. mucronatum (Lam. Dict. IV, p. 218) tomento denso niveo pannosum, caule elato crebre folioso ample paniculato, foliis crassis radicalibus oblongis crenatis interdum sublobatis subpetiolatis, caulinis oblongis longiuscule decurrentibus, superioribus suborbiculatis abrupte mucronatis, florum glomerulis globosis valde dissitis racemos longos virgatos rigidos interruptos formantibus, floribus subsessilibus, calyce (2 lin. longo) lanâ bombycinâ occultato fere ad basin in lacinias late lanceolatas partito, filamentis albido-lanatis, capsulâ parvâ globosâ calyce non longiore ②. *V. auriculatum* Fl. Gr. III, p. 20, tab. 225. — *V. candidissimum* D.C. Fl. Fr. Suppl. pag. 413.

Hab. in collibus, ad vias, insula Samos (Sibth!), Creta (Labill!), prope Smyrnam et Magnesiam copiosum (Boiss!), Pisidia (Heldr!), Armenia prope Karput (Haussk!).

Pluripedale, folia radicalia magnitudine valde variant, glomeruli avellanâ minores sœpe pollicem distantes, capsula semine Cannabeos non major. Cum *V. glomerato* formas gignit hybridas, v. gr. *V. astrophyllosum* Boiss. et Heldr. Diagn. Ser. I, 12, p. 43 in Isauriâ observatum.

40. V. Antinori (Boiss. et Heldr. Diagn. Ser. II, 3, p. 144) bienne, preter inflorescentiam viscidulo-hirtum adpresso pannosotomentosum a collo pluricaule, caulis rigidis creberrime et stricte foliosis lateralibus simplicibus centrali pyramidato foliis radicalibus oblongo-spathulatis obtusis profunde et obtuse sinuatim undulatis, caulinis confertissimis oblongis acutiusculis breviter decurrentibus, floribus 2-4 glomerulatis subsessilibus racemos laxos formantibus, calyce (2 lineas longo) ultra medium tisso virenti glanduloso-hirto laciinis ovatis acutiusculis, corollâ parvâ, filamentis paree albo-barbatis, capsulis ovatis acutiusculis mueronatis calyce sesquilongoribus ②.

Hab. in saxosis vulcanicis ad Cordileon juxta Smyrnam (Heldr!) Fl. Jun, Jul.

Cum paniculâ pedali bipedale, folia inferiora fere omnino *V. villicati* a quo haec species paniculâ non tomentosa sed viscidâ et calycis multo minoris laciinis latis eximie differt. Tantum defloratum et ex paucis speciminibus notum et ulterius explorandum. Capsula fere 3 lineas longa. An forma hybrida v. gr. *V. sinuati* cum specie quâdam ex *Thapoideis?*, sed capsulae et semina omnino normalia videntur.

41. V. detersile (Boiss. et Heldr. Ser. I, 12, p. 15) bienne, foliis radicalibus tomento lutescenti-griseo tenui obsitis oblongis grosse et obtuse sinuato-sublobatis, caule elato folioso laxe et longe paniculato cum ramis jam sub anthesi denudato nigro nitido, foliis caulinis cito detersilibus oblongis et ovato-orbiculatis basi late auriculato-cordatis abrupte acuminatis, glomerulis 2-4-floris distantibus laxissime racemosis, pedicellis brevissimis, calyce ut et bracteæ lanceolatae flocco lombacino cito deciduo prætereaque stipitato glanduloso (2 lineas longo) in laciinas linearis-lanceolatas patentes partito, corollâ luteâ amplâ tomento detersili obsitâ, filamentis sordide albo-barbatis, capsulâ.... ②

Hab. in rupestribus apricis faucis montis Climax Lyciae supra Kourmalu cum *V. orgyalis* rarius (Heldr!) Fl. Maio.

Tripedale, folia radicalia 7-8 pollices longa 2-3 lata, flores eis *V. orgyalis* et *pulverulenti* duplo maiores, ab eis insuper floribus subsessilibus remotum et ob indumenti naturam et inflorescentiam demum omnino glabratam et nitidam *Leianthis* accedens, sed corolla major et tomentella. Capsula inquinenda.

§ 4. Glomerulosa.

+ Filamentorum lana albida vel flavescens.

42. V. gnaphalodes (M. B. Taur. Cauc. III, p. 152) bienne,

lanâ niveâ floccosâ tandem ad foliorum paginam superiorem et caulem deciduâ obsitum, foliis crenulatis vel integris ovato-ellipticis oblongisve inferioribus in petiolum breveni attenuatis, superioribus diminutis sessilibus cordato-amplexicaulibus sepe acuminatis, caule simplici vel parce ramoso, florum fasciculis densis 5-7-floris demum remotiusculis spicas longas laxas formantibus, pedicellis aequalibus longioribus calycei æquilongis, calyce (2 $\frac{1}{2}$ lin. circ. longo) praeter dentes apice glabratos albo-lanato fere ad basin in lacinias lanceolatas acutissimas partito, corollâ parvâ, filamentis sordide albo-lanatis, antheris reniformibus, capsulâ glabratâ ovatâ calyce vix longiore ②.

Hab. in Tauriâ meridionali (Stev!), Ponto Lazico in abietinis supra Khabachkar (Bal!) et in regione inferiori (C. Koch!), Mingrelia et Guriâ (Ledeb!), Imeretiâ (Rupr!). Specimina Aucheriana Syriaca et Graeca in d.c. Prodr. citata ad alias species spectant.

Facies *V. lychnitidis*, indumentam *V. laguri* et *V. malacotrichi* inter *Thapsus*.

43. ***V. eriorrhabdon*** (Boiss. in Tchih. As. Min. 2, p. 4) biennne, lanâ niveâ floccosâ tandem ad folia et caulem deciduâ obsitum, caule rarius in racemum simplicem saepius in paniculam abeunt, foliis crenulatis vel integris inferioribus ovatis vel oblongis in petiolum breviter attenuatis, superioribus sessilibus rotundatis vel semi-amplexicaulibus, florum fasciculis bracteâ linearî suffultis in racemos longos laxiusculos dispositis, pedicellis calycei æquilongis, calyce bombycino tandem glabratâ (2 lineas longo) ad basin in lacinias anguste lineares partito, corollâ parvâ flavâ, filamentis flavidolanatis, capsulâ oblongâ mucronulatâ calyce sesquilongiore ②. *V. Schneideri* et *V. alpestre* Boiss. et Haussk. MSS. — *V. macrosepalum* Boiss. et Ky. (forma autumnalis elongata). — *V. oxycarpum* C. Koch Linn. XXIII, p. 581.

Hab. in Bithyniâ (C. Koch!), in Paphlagoniâ ad Cerasontem (Tchih!), Cataoniâ in monte Berytdagh (Haussk!), circa Bitlis (Ky. 720!), ad Arablar prope Marasch (Haussk!).

E. insulare. — Omnibus partibus majus, tripedale, fasciculi multiflori — *V. insulare* Boiss. et Heldr. Diagn. Ser. I, 12, p. 14.

Hab. in insulâ lacûs Beychehr Pisidae (Heldr!).

Species indumento et characteribus affinis *V. gnaphalode* a quo differt calycis lacinias anguste linearibus, capsulâque oblongo-cylindrica 3 lin. longâ.

44. ***V. pycnostachyum*** (Boiss. et Heldr. Diagn. Ser. I, 12, p. 11) biennne, totum tomento albo nou detersili dense pannosum, paniculae rami congestis lateralibus brevibus terminali elongato, foliis sub-integris oblongis inferioribus in petiolum brevem attenuatis, caulinis crebris imbricatis versus basin paniculae sensim diminutis acuminatis basi auriculatis interdum decurrentibus, fasciculis 2-3-floris dense racemosis, pedicellis calyce brevioribus, calyce deuse tomentoso (3 lineas longo) paulo infra medium in lacinias triangulares fisso

filamentis dense albo-barbatis, capsulâ lanâ detersili vestitâ globosâ calyci æquilonigâ ② *V. pilematophorum* C. Koch Linn XXIII, p. 571.

V. Daënense Boiss. quoad spec. Anatolicum non Boiss. Diagn. — *V. millelacuum* Boiss. in Tchih. As. Min. 2, p. 3.

Hab. in agris, ad vias Pisidiæ prope Egirdir (Heldr!), prope Ermenek Ciliciæ Tracheæ (Peronin!), in saxosis Isaurie (Heldr!), Cappadociæ ad Cæsaream (Bal!), monte Jildisdagh Galatiæ (Wiedem!), Armeniâ Turcicâ (C. Koch! Tchih!).

Planta 1 $\frac{1}{2}$ -3-pedalis, folia inferiora 3-5 pollices longa, caulinâ valde congesta et imbricata conum formantia, corolla 8-10 lineas diam. lata, capsula piso fere dimidio minor. Inter omnia *V. glomerato* præsertim affine egregie differt staturâ minore, florum fasciculis non vel vix dissitis, calyce et corollâ dimidio minoribus. *V. Daënense* cui hoc prius retuleram distinguuntur foliis radicalibus lanceolato-linearibus elongatis, floribus longius pedicellatis, calyce profundius partito.

45. V. Sinaiticum (Bth. in D.C. Prodr. X, p. 236) bienne, totum tomento aureo vel ferrugineo non detersili dense pannosum, caule elato in paniculam longissime ramosam abeunti. foliis crassis crenatis inferioribus oblongis elongatis basi longe attenuatâ subpetiolatis, superioribus crebris oblongis subcordato-amplexicaulibus interdum breviter decurrentibus. florum fasciculis remotis racemos longissimos laxos formantibus. pedicellis tomentosis inæqualibus longioribus calycem vix superantibus, calyce tomentoso (2 $\frac{1}{2}$ longo) ad $\frac{2}{3}$ in laciniis oblongo-lanceolatas tisso, filamentis albo-lanatis, capsulâ glabratâ ovatâ subretusâ calyce sublongiore ② *V. fasciculatum* Ehr. exs! — *V. Barradense* Boiss. olim.

Hab. in deserto Sinaitico et in montis Sinai humidis (Ehrenb! Bové! Schimp! Boiss!), in Antilibano variis locis (Gaill! Kotschy exs. 231 sub *V. Daënensi*). Arabice *Cherma*.

Folia radicalia pedem et amplius longa, racemi fructiferi saepè bipedales, capsula tres fere lineas longa piso minor. Corolla ex speciminibus cultis alba!

46. V. Daënense (Boiss. Diagn. Ser. I, 7. p. 38) bienne, elatum tomento griseo ad folia persistenti ad partem superiorem plantæ subfloccoso obsitum, caule pyramidatim paniculato. foliis subintegris radicalibus longe et late linear-lanceolatis acutiusculis, caulinis diminutis basi attenuatis vel subauriculato-amplexicaulibus subtus elevatim nervosis, florum fasciculis subdissitis racemos laxos formantibus, pedicellis calyce vix brevioribus, calyce tomentoso (2 $\frac{1}{2}$ lineas longo) fere ad basin in laciniis lanceolatas acutissimas fisso, filamentis sordide albo-barbatis, capsulâ glabrescente ovatâ calyce vix longiore

Hab. in dumosis montium Persiae austro-occidentalis, Kuh daëna (Ky. 692!), Sawers, Eschker. Nur 3000' (Haussk!).

Folia radicalia pedalia 2 pollices tantum lata. Species affinis *V. Sinaitico* a quo sàt differre videtur indumento minus denso, foliis angustioribus, impressis calycis laciniis valde acutis et capsulâ majore pisum æquante.

47. **V. Khorassanicum**, elatum tenuiter stellato-tomentellum pallide virens, canle paniculato ramis longis virgatis, foliis oblongis obsolete undulato-sublobatis repandis breviter in petiolum attenuatis rameis parvis sessilibus basi truncato-subcordatis, florum fasciculis subdissitis laxe racemosis, pedicellis calyce vix brevioribus, calyce tomentoso (2 lineas longo) fere ad basin in lacinias lanceolatas acutas fisso. filamentis sordide albo-barbatis, capsulâ.... (2).

Hab. in pratis montanis prov. Khorassan Persiae prope Dchegar et Chanlug in viâ inter Nichapur et Meschhed (Bge!).

Affine videtur *V. Daenensi* a quo differt indumento tenui, foliis brevibus, floribus fere dimidio minoribus; facies *V. erianthi* calyce breviter dentato diversi.

48. **V. stachydiforme** (Boiss. et Buhse Aufz. p. 159) perenne? pluricaule, totum tomento denso griseo-flavido pannosum, caulis foliatis in racemum simplicem vel breviter et parce ramulosum abeuntibus, foliis oblongis crenulatis sub indumento reticulato-nervosis inferioribus in petiolum brevem attenuatis cæteris sessilibus amplexicaulibus, floralibus lanceolatis longe acuminatis flores superantibus, spicis densiusculis, floribus 2-3-fasciculatis, pedicellis calyce brevioribus, calyce pannoso (2 1/2 lin. longo) fere ad basin in lacinias linearis-lanceolatas obtusiusculas fisso, corollâ parvâ flavâ, filamentis omnibus flavo-barbatis, capsulâ.... (2)?

Hab. in agris incultis ditionis Talysh prope Aschewlar (Buhse!).

Caules cum racemo semipedali sesquipedales, folia inferiora 4 poll. longa 12-15 lineas lata. Facies *Stachydis lanata*. Duratio incerta, ex specimine juniori descriptum ulterius observandum.

49. **V. lasianthum** (Boiss. in Pinard. exs. 1843. — Bth. in D.C. Prodr. X, p. 234) tomento denso flavicante asperulo pannosum, caule valde elato superne ample et stricte paniculato, foliis integris radicalibus lanceolatis saepè elongatis, caulinis diminutis a basi amplexicauli oblongis vel ovatis, floribus 4-5 pedicellis calyce subbrevioribus suffultis fasciculatis, racemis virgatis laxis, calyce tomentoso pentagono (2 1/2 lineas longo) ad tertiam partem in dentes triangulares fisso, filamentis omnibus sordide albo-lanatis, capsulâ.... (1). *V. phlomoides* Urv. Enum., p. 25 non L. — *V. brevidens* Bth. loc. cit. p. 235.

Hab. in Cariâ (Pinard!), insulâ Samos (Urv!), monte Balloucklardan Lyciae (Bourg. exs. sub *V. Pisidico*), in agris prope Sam (Haussk!).

Planta saepè 7-8-pedalis foliis radicalibus pedem et amplius longis. Species pulchra facie *V. glomeratum* referens sed ab eo et affinibus calyce multo minore campanulato et breviter dentato discedens. Planta ex insulâ Samo a Cariensi floribus paululum minoribus tantum differt.

50. **V. erianthum** (Bth. in D.C. Prodr. X, p. 236) biennne, elatum tomento tenui cinereum, caule nigricanti superne paniculato,

foliis crenatis radicalibus oblongis in petiolum attenuatis, caulinis brevibus cordato-auriculatis, floribus remote fasciculatis laxissime racemosis tomentosis, pedicellis inæqualibus calyce subbrevioribus, calyce (vix 2 lineas longo) ad medium in dentes triangulares fisco, filamentis sordide albo-lanatis, capsula ovatâ calyce duplo longiore ②.

Hab. in regno Cabulico (Griff. 630), in Belutchiâ (Stocks 1043!).

A præcedentibus floribus minoribus, calycis dentibus brevibus distinctum, a *V. brevidente* floribus longius pedicellatis aliisque notis. Capsula 2½ lineas longa. Specimen Armenum Aucherianum valde incompletum huc a cl. Bth. adnumeratum ad *V. erianthum* non spectat.

51. *V. leucophyllum* (Griseb. Spic. II, p. 46!) biennie, caule crassiusculo jam ex parte inferiori in paniculam ramosissimam rigidam abeunte, foliis adpressissime canis radicalibus consertis petiolatis oblongis profunde sinuato-lobatis lobis rotundatis sinibusque obtusissimis, foliis caulinis paucissimis minimis sessilibus obtuse dentatis, floribus 2-5 fasciculatis vel superioribus solitariis in racemos laxos dispositis, pedicellis calyce et capsula vix brevioribus, bracteis squamiformibus triangulari-ovatis, calycis tomentelli tandem glabratî (sesquilineam longi) laciniis linear-i-lanceolatis acutiusculis, corollâ tomentellâ, filamentis albo-barbatis, capsula ovoideo oblongâ mucronatâ glabratâ calyce subduplo longiore ②.

Hab. in collibus et agris sterilibus Macedoniae inter Thessalonicam et montem Korthiati (Griseb! Heldr!).

Pedale vel sesquipedale, folia inferiora cum petiolo 3-4 pollicaria, paniculae rami numerosi rigidi erecto-patentes 5-6 pollicares, capsula fere 3 lineas longa pedicello incrassato suffulta. Folia fere *V. plicati* a quo calyce et floribus minutis longe differt, a *V. Heldreichii* indumento cano, racemis brevibus rigidis, pedicellis brevioribus, capsula multo majore distinctum.

+ + Filamentorum lana purpurea vel violacea.

52. *V. pumilum* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 20) biennie, caule pumilo fere a basi in paniculam abeunti, foliis cano-tomentosis minute crenulatis parvis radicalibus rosulatis ovatis et oblongis petiolatis, caulinis paucis acutis, paniculae ramis brevibus glabrescentibus, floribus 2-3 fasciculatis laxiusculis. pedicellis calyci æquilongis, calyce tomentello (sesquilineam longo) ad basin in laciniis linear-i-subspathulatas obtusas partito, corollâ parvâ flavâ violaceo-maculatâ, filamentis violaceo-barbatis, antheris aurantiacis, capsula ovato-oblongâ calyce tertiatâ parte longiore ②.

Hab. in montibus aridis Isauriae inter Beychehr et Koniah versus planitiem Iconiensem descendendo (Heldr!), in Cappadociâ ad Talasse prope Caesaream (Bal!).

Pianta semipedalis rarius pedalis, folia inferiora 1-2 polices longa, corolla diametro sex lineas lata, capsula tres lineas longa. Teste amico de Heldreich florum colore elegantissimum.

53. V. elegans (Boiss. et Heldr. Diagn. Ser. I, 12, p. 12) bienne, tomento denso non detersili tomentosum æruginoso-canum, caule pumilo inferne crebre folioso in paniculam angustam parce et breviter ramosam abeunte, foliis parvis oblongis acutiusculis sub-integris radicalibus in petiolum attenuatis subtus reticulato-rugosis, caulinis oblongis strictis basi sessili attenuatis, floribus solitariis vel 2-3 fasciculatis secus paniculæ ramos laxis, pedicello calyce sub-breviore, calyce ($2\frac{1}{2}$ lineas longo) tomentoso ultra medium in lacinias linear-lanceolatas obtusas fisso, corollâ luteâ, filamentis purpureo-lanatis, capsulâ.....

Hab. in montibus Pisidiæ vel Isauriæ (Heldr! spec. unicum absque loco natali proprio).

Planta vix pedalis paniculâ semipedali, folia radicalia cum petiolo sesquipollucaria 6-7 lin. lata. Affine *V. chrysorrhacos* differt indumento denso pannoso, foliis multo minoribus formæ diversæ, floribus subfasciculatis, filamentis purpureo-lanatis. Facie simile *V. pumilo* quocum probab. crescit et a quo differt indumento densius pannoso, pedicellis brevibus, calyce fere duplo majori. An forsan hujus probes hybrida?

54. V. Tauri (Boiss. et Ky. Diagn. Ser. II, 3, p. 145) bienne, totum indumento flavido-griseo ad folia pannoso persistenti ad caules et ramos subdetersili obsitum, caule brevi pyramidatum paniculato, foliis radicalibus ovatis et oblongis in petiolum attenuatis subtus reticulatis, caulinis ad ramorum ortum parvis subcordato-amplexicaulibus ovatis, floribus subbinis fasciculatis laxe racemosis, pedicellis sub anthesi calyci æquilongis capsulâ 2-3-plo longioribus, calyce (circ. 2 lin. longo) ad basin in lacinias lineares partito, filamentis pallide violaceo-barbatis, capsulâ oblongâ tomentellâ calyce tertiatâ parte longiore pedicello breviore ②.

Hab. in regione alpinâ Tauri Cilicici in monte Gisyl Tepe 8000' (Ky. 235! Bal!), ditione Kassan Oglu Cilicie orientalis (Ky. 175!).

Planta 1-1 $\frac{1}{2}$ -pedalis, folia radicalia laminâ 2-3 pollices longâ, capsula 2 $\frac{1}{2}$ lin. longa. Affine *V. eleganti* a quo differt caule elatiore, foliis majoribus, floribus longiuscule pedicellatis et imprimis laciñis calycis angustioribus.

55. V. Adeliæ (Heldr. in Boiss. Diagn. Ser. II, 3, p. 145) bienne, tenuiter tomentosum griseo-flavidum, caule parum elato crebre folioso in paniculam brevem confertam densifloram abeunti, foliis subintegris inferioribus ovato-oblongis obtusis in petiolum attenuatis, superioribus ovatis breviter acuminatis cordato-auriculatis, florum glomerulis paucifloris approximatis, pedicellis calyci æquilongis, calyce tomentoso (lineam circ. longo) fere ad basin in lacinias triangulari-lanceolatas acutas partito, filamentis violaceo-barbatis, capsulâ oblongâ obtusâ calyce duplo longiore ②.

Hab. in valle Aperantos ad radices montis Za insulæ Naxi Cycladum (Heldr!).

Caules 1-2-pedales, foliorum inferiorum lamina tripollucaris, capsula 2 lineas

longa, facies *V. Lychnitis* sed panicula etiam fructifera 4-5 pollices tantum longa ramis brevibus, bracteæ breves et filamenta violaceo-barbata. A *V. Boegerhavii* et affinibus calyce minimo alienum.

56. V. Tripolitanum (Boiss. Diagn. Ser. I, 12, p. 9) bienne, ad rosulas steriles dense æruginoso-pannosum in plantâ floridâ tenuius flavidio-virescens, caule elato nigricanti in paniculam amplam abeunti, foliis inferioribus oblongis vel oblongo-lanceolatis petiolatis crenatis et saepius præsertim basin versus grosse lobatis, caulinis diminutis sessilibus non decurrentibus, floribus 3-4 in glomerulos dissitos bracteâ obcordatâ acuminatâ suffultos dispositis, pedicellis calyce brevioribus, calyce patule hirsuto (3 lin. longo) ultra medium in lacinias oblongas acutiusculas partito, filamentis violaceo-lanatis longioribus basi nudis, capsulâ ovatâ tomentellâ calyce vix longiore ②.

Hab. ad radices montis Torbol ad Tripoli Syrie (Boiss! Bl. 434!), prope Broummana Libani (Bl!), supra Saida (Gaill. 1023!), ad Nanlu montis Amani 3000! (Ky. exs. 246!).

Facies *V. sinuati*, planta 3-4-pedalis, folia inferiora 4-7-pollicaria, paniculæ racemi virgati longissimi glomerulis distantibus. Præter folia non undulata et caulinæ non decurrentia calycis duplo majoris formâ ab eo statim distinguendum.

57. V. sinuatum (L. Sp. p. 254) bienne, plus minus dense tomentosum lutescens vel griseum, paniculæ pyramidatae ramis tenuiter virgatis elongatis erecto-patulis, foliis radicalibus oblongo-spathulatis subsessilibus sinuato-pinnatifidis saepè undulatis, caulinis superioribus oblongis acutis integris breviter decurrentibus, florum glomerulis dissitis, pedicellis inæqualibus majoribus calycem aequantibus, calyce tomentello (sesquilinea vix longiore) ultra medium in lacinias breviter lanceolatas fisso, corollâ parvâ, filamentis purpureo-lanatis, capsulâ minimâ globosâ calycem subæquante ②. Ic. Fl. Græc. tab. 227. — Relib. Ic. tab. 24. — *V. Ceccarinianum* Boiss. et Heldr. MSS. — *V. Gaillardotii* Boiss. Diagn. Ser. II, 6, p. 128.

Hab. in sterilibus, ad vias, totius fere ditionis a regione inferiore ad montanam a Græciâ! Macedonia! et Thraciâ! ad Syriam littoralem et interiore! Palestinam, Assiriâ et Kurdistaniâ Persicam in regione inferiori montis Avroman 5000' (Haussk!), montibus Persiæ austro-occid. (Ky. 587! Haussk!), Tauriâ (Stev!), Transcaucasiâ (Ledeb!), Armeniâ Rossicâ (Szov!).

Indumento foliisque interdum tantum sublobatis caulinisque vix decurrentibus valde variabile. Capsula *Cannaleos* semen aequans.

Ar. Geogr. Regio mediterranea Europæ et Africæ borealis, insulæ Canarienses.

§ 5. *Lychnitidea.*

* Calyx minutus (1-1 1/2 lin. longus).

+ Filamentorum lana albida vel flavescens.

58. V. pulverulentum (Vill. Delph. II, p. 490) bienne, to-

mento denso niveo floccoso tandem deciduo obsitum, caule elato in paniculam pyramidatam amplam abeunte, foliis integris vel obsolete crenatis radicalibus oblongis vel ellipticis in petiolum brevem attenuatis, superioribus ovatis vel rotundatis amplexicaulibus plus minus abrupte acuminatis, florum fasciculis parum dissitis bracteâ linearî suffultis longe racemosis, pedicellis calyce sublongioribus, calyce dense floccoso (sesquilineam longo) in lacinias linearî-lanceolatas partito, corollâ parvâ, filamentis albido-lanatis, capsulâ parvâ ovato-globosâ retusâ calyce longiore (2). Engl. bot. tab. 487. — *V. floccosum* W. K. I, p. 81, tab. 79. — Ic. Rehb. tab. 26.

Hab. in Macedonie peninsula Athoâ et monte Korthiati prope Salonicam (Griseb!).

Ar. Geogr. Europa media et australis ab Anglia et Belgio ad Dalmatiam et regionem Danubialem

59. ***V. mallophorum*** (Boiss. et Heldr. Diagn. Ser. I, 7, p. 39) biennium, tomento denso niveo floccoso plus minus persistente vestitum, caule elato in paniculam pyramidatam amplam abeunte, foliis radicalibus in petiolum brevem attenuatis ovato-oblongis integris vel obsolete crenatis, canlinis basi rotundatis ovatis acutis, florum fasciculis fere contiguis bracteâ linearî suffultis, pedicellis calyce sublongioribus, calyce dense floccoso (sesquilineam longo) in lacinias anguste lineares partito, corollâ parvâ, filamentis rufescenti-lanatis, capsulâ parvâ oblongâ obtusâ mucronatâ calyce triplo longiore (2).

Hab. in regione moutanâ et subalpinâ Græciae 4000'-5500', Taygetus, Olenos et Kyllene (Heldr!), Malevo Laconia (Orph!), Arapocephala (Etolia) (Sam. et Guic!), Delphi Eubea (Heldr!), Parnassus (Orph!).

Valde affine *V. pulcherrima* a quo differre videtur tomento vix et superne tantum detersili, foliis caulinis non abrupte acuminatis, filamentorum lanâ rufescente et praesertim capsulâ oblongâ (2½ lin. longâ) nec subglobosâ retusa.

60. ***V. Olympicum*** (Boiss. Diagn. Ser. I, 4, p. 54) biennium, elatum, totum tomento denso niveo vix detersili canum, paniculae pyramidatae racemis longissimis laxis, foliis inferioribus oblongo-lanceolatis lanceolatisque elongatis integris basi attenuatis, superioribus breviter lanceolatis acutis, glomerulis multifloris bracteâ linearî suffultis, pedunculis tomentosis calyce duplo longioribus, calyce tomentosissimo (2 lin. longo) in lacinias longe et anguste linearî-lanceolatas partito, corollâ majusculâ, filamentis albo-barbatis, longioribus duobus totâ parte superiore nudis, capsulâ oblongâ tandem glabrescente calyce duplo longiore (2).

Hab. in regione mediâ Olympi Bithyni in pratis (Auch. 2455! Boiss! Pichler!).

Caules 4-5-pedales, folia interiora sœpe sesquipedalia, corolla diam pollicaris, capsula tres lineas longa. Affine *V. mallophoro* a quo differt foliis et pedicellis elongatis, corolla capsulaque (3 lin. longâ) majoribus. Facies *V. orgyalis* quod pedicellis brevibus et calyce minimo longius differt.

61. **V. orgyale** (Boiss. et Heldr. Diagn. Ser. I, 12, p. 15) bieinne, valde elatum inferne tomento denso non detersili cinereo obsitum, caule ample pyramidato-paniculato, foliis radicalibus longe lanceolatis acutis basi sensim angustatis, caulinis superioribus oblongis et ovatis basi amplexicauli subauriculatis acutis vel acuminatis, paniculae ramis longis tenuiter virgatis nigricantibus tandem glabratibus, fasciculis 2-4-floris bracteâ setaceaâ suffultis remotiusculis, pedicellis calyce demum sublongioribus, calyce tenuiter floccoso minimo (lineam vix longo) in lacinias subulatas acutas partito, corollâ parvâ luteâ calyce sextuplo majore, filamentis rufescenti-barbatis, antheris fuscis, capsulâ... (2).

Hab. in rupestribus apricis montis Climax Lyciae supra Kourmalu 1000' copiosum (Heldr!).

Planta 6-7 pedes alta, folia inferiora pedalia 3 pollices lata, corolla diametro 9-10 lineas lata, paniculae rami saepe sesquipedales. Ob capsulam non notam affinitas dubia, facie accedit ad *V. speciosum* calyce duplo majori, pedicellis longioribus, etc. diversum.

62. **V. Lychnitis** (L. Sp. 253) bieinne, caule paniculato ramis erecto-patulis angulatis, foliis siccitate nigricantibus supra viridibus subtus tenuiter canis vel canescentibus crenatis radicalibus oblongo-ellipticis in petiolum angustatis, superioribus lanceolatis sessilibus, florum fasciculis laxis bracteis linear-lanceolatis eis saepe longioribus suffultis, pedicellis calyce longioribus, calyce tomentoso (sesquilineam longo) in lacinias lanceolatas acutas apice glabras ad $\frac{3}{4}$ partito, corollâ parvâ luteâ vel albâ, filamentis albo-lanatis, capsulâ parvâ ovatâ obtusâ calyce duplo longiore (2). Ic. Engl. bot. tab. 58. Rchb. tab. 27. — *V. pulverulentum* M.B. I, p. 160 non Vill. — *V. Biebersteinii* Besser.

Hab. in Ponto ad Trapezuntem (ex Urv!), Tauriâ (Stev!), prov. Caucasicis (M.B!), Iberiâ Wilhelm. (ex Ledeb.).

Ar. Geogr. Europa omnis a Sueciâ et Hispaniâ Italiâque centralibus ad Ros-siam medium.

63. **V. Delphicum** (Boiss. et Heldr. Diagn. Ser. II, 3, p. 146) bieinne, caule glabrescenti in racemum simplicem vel inferne parce ramosum abeunte, foliis supra viridibus subtus tenuiter tomentosocanescentibus radicalibus petiolatis amplis ovalis basi rotundatis vel subcordatis obtuse crenatis, superioribus sessilibus diminutis cordatis ovatis, florum glomerulis racemos densiusculos formantibus, pedicellis calyce longioribus calyceque tomento detersili niveo obsitis, calyce (sesquilineam longo) fere ad basin in lacinias late lanceolatas partito, corollâ flavâ, filamentis albo-barbatis, capsulâ demum glabrâ oblongâ obtusâ calyce duplo longiore (2).

Hab. in regione sylvaticâ et superiore montis Delphi Eubeæ 3000'-5500' (Heldr. Herb. Norm. N° 799!).

Bipedale, foliorum inferiorum limbus semipedalis, racemi saepe pedales.

Facie et characteribus affine *V. nigro* L. et *lanato* Schrad. a quibus differt filamentorum lanâ albida nec violacea, laciniis calycinis latis, capsula paulo majore.

64. V. speciosum (Schrad. Hort. Gött. fasc. 2, p. 22, tab. 16) bienne, elatum indumento tenui persistente tomentosum vel griseum, foliis subintegris radicalibus amplissimis oblongo-lanceolatis utrinque attenuatis saepe acuminatis, caulinis brevibus cordato-amplexicaulibus, paniculae amplissime pyramidatae ramis elongatis, florum fasciculis taxis, pedicellis calyce longioribus, calyce (2 lin. longo) in laciniis anguste lanceolatas ad basin partito, corollâ parvâ, filamentis albo-lanatis, capsula ovato-oblongâ obtusâ calyce subdupo longiore ②. Ic. Rehb. tab. 25. — *V. longifolium* D.C. Fl. Fr. non Ten.

Hab. in Armenia Turcicâ prope Ispir (Bourg! C. Koch!), Cataoniâ prope Arablar (Haussk!), prov. Talysh (C.A. Mey!), Armenia Rossicâ et Persiâ boreali (Buhse! Szov! Bge!).

β. megaphlomos Boiss. et Heldr. Diagn. Ser. II, 2, p. 144. — Folia angusta crasse lutescenti-pannosa.

Hab. in regione abietinâ Parnassi (Heldr!).

γ. microcarpum. — Capsula minor ovato-oblonga.

Hab. in lapidosis Lyciae ad Ballouklardan supra Elmalu (Bourg!), in Tauro Isaurico inter valles Tourtchalar et Ermenek 3000' (Heldr!).

Planta pluripedalis, folia radicalia saepe bipedalia 3-5-pollices lata, paniculae rami saepe sesquipedales, capsula in typo fere 3 lineas longa, in varietate microcarpâ vix 2 lineas. *H. hyoseridifolium* Schrad. Mon. II, p. 31, tab. 3 ex descriptione et icone foliis repando-subsinuatibus tantum differt et hujus varietas vel forma hybrida videtur.

Ar. Geogr. Austria, regio Danubialis, Rossia australis.

65. V. cheiranthifolium (Boiss. Diagn. Ser. I, 4, p. 56) tomento stellato tenui saepe adpressissimo cinerascens vel flavidum, caule in paniculam pyramidatam ramosissimam multifloram abeunti, foliis radicalibus lanceolatis vel oblongo-lanceolatis in petiolum attenuatis subintegris, caulinis oblongis a basi suburiculato-cordatâ acuminatis, floribus 2-5 in fasciculos remotiusculos bracteolis linearibus stipatos dispositis, paniculae ramis tenuibus strictis, pedicellis calyci æquilonigis vel longioribus, calyce tomentello (vix sesquilineam longo) fere ad basin in laciniis linear-lanceolatas partito, corollâ flavâ, filamentis sordide flavidio-barbatis, capsula oblongâ obtusâ mucronulatâ calyce tertiatâ parte longiore ②. *V. asperulum* Boiss. loc. cit. (forma foliis radicalibus latioribus ovato-oblongis). — *V. Pisidicum* Boiss. et Heldr. Diagn. Ser. I, 42, p. 45. — *V. monticolum* et *V. Seytunense* B. et Haussk. exs. (formæ forsitan hybridae).

Hab. in pratis regionis subalpine Anatolie hinc inde in valles descendens, Tmolus ad Bozdag (Boiss!), Cadmus supra Denisleh (Boiss!), Sipylus supra Magnesiam (Boiss!), mons Mouraddagh Phrygiae et Alidagh Cappadociae (Bal!), Taurus Isauricus 3500'-500' (Heldr!), Mesogis colles circa Derbend

(Boiss!), mons Boudroun Pisidiae in regione inferiore (Heldr!), prope Seytun et in monte Berytdagh (Haussk!).

β. Heldreichii. — Folia angustissima $\frac{1}{2}$ -1-pollicem lata, indumentum paulo densius, racemi densiores, capsula paulo major. — *V. stenophyllum* Boiss. et Heldr. Diagn. Ser. I, 12, p. 17.

Hab. in regione superiori montis Boudroun Pisidiae 3000'-6000', montibus Isauriae supra Ermenek, monte Anemas Lycaonia ad nives 8000' (Heldr!).

Species omnium montium Anatoliæ australis incola, bi-tripedalis, folia inferiora 5-8 pollices longa pollicem vel quoque 2 pollices lata, capsula $2\frac{1}{2}$ lineas longa. Affine *V. specioso*, folia multo minora et angustiora, calyces minores, capsula minor magis angustata. Formas verosimiliter hybridas gignit.

66. *V. humile* (Janka Breviar. 2, p. 6) biennie, pallide virens, foliis firmis radicalibus rosulatis oblongo-lanceolatis profunde et obtuse crenatis supra sub lente puberulis subtus adpresso canescens, caule humili striete paniculato inferne foliis paucis angustatis lanceolatis obsito, paniculâ sub lente adpresso puberulâ, floribus 2-3 fasciculatis in racemos breves dispositis pedicellis calyce sesquilon-gioribus suffultis, bracteis linearibus pedicellos æquantibus, calyce adpresso canescenti (sesquilineam longo) ad $\frac{3}{4}$ in lacinias anguste lineari-spathulatas partito, corollâ parvâ luteâ tomentellâ, filamentis e sicco albido-barbatis, capsulâ tomentellâ ellipsoideo-ovatâ obtusâ mucronatâ calyce sublongiore ②.

Hab. in lapidosis calcareis regionis mediæ montis Tchatalkaja prope Sliwno Thraciæ borealis (Janka!).

Specimen unicum, caulis cum paniculâ semipedali pedalis, folia radicalia 5-7 pollices longa pollicem vel paulo amplius lata, racemi 3-4 pollicares, flores e minimis, capsula Cannabeos semine duplo major. Ex affinitate *V. cheiranthifolia* et *Heldreichii*.

67. *V. Heldreichii* (Boiss. Diagn. Ser. II, 3, p. 147) biennie, indumento tenui obsitum flavo-virens, caule rubello elato a medio paniculato ramis tenuiter virgatis in racemos longissimos laxos abe-untibus, foliis inferioribus petiolatis lyratis segmentis lateralibus utrinque 1-3 parvis oblongis irregularibus terminali magno oblongo lobato, nervis paginæ inferioris albis prominentibus, foliis superioribus et rameis parvis ovatis acuminatis, florum glomerulis subdissitis, pedicellis inæqualibus, calyce ($1\frac{1}{3}$ lin. longo) tomentoso fere ad basin in lacinias anguste lanceolatas apice glabrescentes partito, corollâ luteâ tomentellâ, filamentis sordide albo-barbatis, capsulâ ovato-oblongâ obtusâ mucronatâ calyce subduplo longiore ②.

Hab. in faucibus montis Korthiati Macedoniae 2000' (Heldr!), ad Keretzkiöi supra Thessalonicam (Orph. Fl. Gr. exs. 733!).

Bipedale et procerius, folia inferiora cum petiolo semipedalia segmentis lateralibus 5-10 lineas terminali 3-4 pollices longo, racemi 4-4 $\frac{1}{2}$ -pedales, capsula Cannabeos semine minor. Facies *V. sinuati* a quo insuper foliis et characteribus longe differt; accedit ad *Verbasca leiantha*, sed panicula non ut in illis glabra et viscosa et corolla tomentosa nec glabra.

68. V. petiolare (Boiss. et Ky. in Kotschy Sched. 1859) bienne, foliis supra adpresso hirtis virentibus subtus tenuiter canis radicalibus subcrenatis ovatis late cordatis petiolatis obtusis, caulinis paucis oblongis basi attenuatis breviter petiolatis et sessilibus, caule rubello tandem glabrecente ramosissime et longe paniculato ramis virgatis elongatis, fasciculis plurifloris remotis bracteâ linearî suffultis in racemos laxissimos dispositis, pedicellis calyci æquilongis fructiferis incrassatis, calyce hispido-lanato subdetersili (lineam longo) ad basin in lacinias lineares obtusinusculas partito, corollâ parvâ luteâ tomentella, filamentis sordide lutescenti-lanatis, capsulâ oblongâ obtusâ mucronatâ calyce duplo longiore ②.

Hab. in glareosis faucium ad pagum Gorumse districtus Kassan Oglu Ciliciæ orientalis (Ky. 1711) Fl. Mai.

Caulis pluripedalis racemis pedalibus et longioribus, folia radicalia petiolo 3-4-pollicari suffulta 4-6 pollices longa interdum suborbiculata. Fasciculus plurifloris et habitu affine *V. leiantho* cuius folia radicalia nondum nota conferre non licet, sed illis neglectis ab eo differt fasciculus florum remotis, calycis lanuti nec glabri laciniis linearibus nec ovato-oblongis. Facies *Leianthorum* sed corolla tomentella.

69. V. stenocarpum (B. et Heldr. Diagn. Ser. I, 42, p. 18) bienne, tomento ad folia radicalia æruginoso persistente ad partem superiorem plantæ griseo tandem detersili obsitum, caule superne stricte et pyramidati ramosissimo, foliis inferioribus oblongis basi attenuatis valde et obtuse sinuato-lobatis undulatisque, caulinis oblongis integris acuminatis, fasciculis 3-5-floris remotiusculis racemos longos laxiusculos strictos formantibus, pecciculis calyci subæquilongis, bracteis linearî-setaceis, calyce (sesquilineam longo) fere ad basin in lacinias lineari-subulatas partito, corollâ parvâ luteâ, filamentis omnibus griseo-lanatis, capsulâ anguste oblongo-cylindricâ obtusâ mucronatâ calyce subtriplo longiore ②.

Hab. in rupestribus Pamphyliæ ad Alaya inter urbem et castellum (Heldr!) Fl. Apr.

Folia inferiora *V. sinuati*, caulis cum paniculâ 2-3-pedalis, corolla diam. 6 lineas lata. Capsula *Verbasci Cedreti* et *Damasceni* sed tenuior 2½ lin. longa vix lineam lata. Fructu affine *V. cheiranthifolio* sed foliis diversissimum.

70. V. myriocarpum (Boiss. et Heldr. Diagn. Ser. I, 42, p. 21) bienne, ad caulis partem inferiorem et folia etiam ramealia dense et adpresso argyreo-tomentosum, foliis radicalibus amplis elliptico-lanceolatis acutiusculis basi cuneatis et breviter petiolatis crenatis et grosse undulato-repandis, caulinis et rameis valde diminutis sessilibus ovati-oblongis integris, caule fere a basi ramosissime paniculato paniculâ glabrâ, fasciculis plurifloris bracteâ subulatâ suffultis laxiuscule ramosis, pedicellis calyci æquilongis, calyce (vix lineam longo) puberulo in lacinias lineari-setaceas partito, corollâ minimâ luteâ, filamentis albo-barbatis, capsulâ glabratâ minutâ oblongo-cylindricâ obtusâ mucronatâ calyce 3-4-plo longiore ②.

Hab. in pinetis regionis mediae montium Pamphyliæ inter Marla et Adalia 4000' (Heldr!).

Sesquipedale, folia radicalia fere pedalia 4 poll. lata, flores in grege minutissimi, capsulae numerosissimæ sesquilineam longæ lineâ dimidiâ vix latiores. Indumento argyreo quod etiam folia ramealia vestit, corollâ extus tomentellâ a *Leianthorum* grege distinctum, ejus minutie et cum calyce proportione a *V. Cilicico* diversum.

71. V. Cedreti (Boiss. Diagn. Ser. I, 12, p. 19) biennie, tomento griseo ad folia adpresso ad partem superiorem plantæ tandem detersili obsitum, caule stricte et pyramidatim paniculato, foliis radicalibus oblongis obtusis cuneatis petiolatis crenulatis sæpius grosse et obtusissime sinuato-lobatis, caulinis parvis oblongis acuminatis, floribus 2-3-nis fasciculatis vel solitariis pedicellis calyce æquilongis suffultis. bracteis setaceis, calyce (1 1/2-2 lin. longo) demum glabrato ad basin in lacinias subulatas partito, corollâ parvâ luteâ, filamentis albo-barbatis, capsulâ oblongo-cylindricâ obtusâ mucronatâ calyce plus duplo longiore ②. *V. Rascheyanum* Boiss. loc. cit. p. 18.

Hab. in regione subalpinâ et alpinâ Libani ad Cedros, circa Eden et Bescherré (Boiss! Bl!), circa Dimam (Bl!), in Antilibano circa Rascheya (Boiss!), ad el Keneise (Gaill!), in monte Garbi supra Zebdani (Ky. 6721).

Planta 1 1/2-2-pedalis, capsula 3-3 1/2 lin. longa 1 1/4 lata. Facie et fructu affine *V. Damasceno* differt indumento tenuiori, foliis sæpè sinuato-lobatis et præsertim lacinias calycis subulatis et staminibus albo-barbatis. Ex cl. et am. Blanche qui plantam in locis natalibus observavit *V. Cedreti* et *Rascheyanum* intermedii confluent.

+ + Filamentorum lana purpurea vel violacea.

72. V. Damascenum (Boiss. Diagn. Ser. I, 12, p. 19) biennie, ad folia inferiora præsertim rosularum densissime pannosum lutescens, caule pyramidato-ramoso cum paniculâ tomento, tandem detersili obsito, foliis radicalibus ovatis obtusis petiolatis, caulinis diminutis basi sessili attenuatis acutis, floribus pedicellis calyce sublongioribus suffultis in fasciculos dissitos 2-4-floros dispositis vel ad racemi partem superiorem solitariis, bracteis linear-setaceis, calyce minimo (sesquilineam vix longo) floccoso demum denudato ad 3/4 in lacinias lanceolatas acutas partito, corollâ parvâ luteâ, filamentis violaceo-lanatis, capsulâ oblongo-cylindricâ mucronatâ calyce aucto duplo longiore ②.

Hab. in collibus aridis ad Salehieh prope Damascum (Boiss!). Specimina florifera e deserto Chabur Mesopotamiæ (Haussk!) hue quoque spectare videntur.

Planta pedalis vel sesquipedalis, foliorum radicalium lamina 1 1/2-3-pollicaris, capsula 3-3 1/2 lin. longa 1 1/2 lata.

73. V. nigrum (L. Sp., p. 353) biennie, caule elato superne angulato in racemum elongatum simplicem vel inferne parce ramo-

sunt abeunti, foliis supra virentibus subtus plus minusve tomentosis crenatis inferioribus longe petiolatis ovalis et oblongis basi cordatis, supremis sessilibus basi rotundatis acuminatis, florum fasciculis laxis multifloris, pedicellis calyce longioribus, calyce hirsuto (sesquilineam longo) in lacinias lanceolato-subulatas fere ad basin partito, corollâ parvâ luteâ fauce violaceâ, filamentis omnibus violaceo-barbatis, capsulâ parvâ ovato-globosâ calyce sublongiore ②. Ie. Engl. bot. tab. 59. — Rehb. tab. 28. — Fl. Dan. tab. 1088. — *V. macrophyllum* C. Koch Linn. XXIII, p. 580.

Hab. in monte Kyllene Peloponnesi (Orph!), Macedonia in monte Korthiati 4000' (Heldr!), Caucaso (M.B!).

V. hypoleucum Boiss. et Heldr. Diagn. Ser. II, 3, p. 147 e faucibus montis Kyllenes differt caule paniculato, foliis radicalibus grosse lobatia, supremis cordatis et videtur omnino probes hybrida *V. nigri* cum aliâ specie, forsas *V. sinuato*? *V. Hinkei* Friv. Flora 1836, p. 440 = *V. Wierzbickii* Heuffel in Rumelia indicatum est forma hybrida ex icone mihi tantum nota facie *V. nigrum* referens.

Ar. Geogr. Europa media et australis a Scandinavia ad Rossiam medium, Sibiria

74. *V. glabratum* (Friv. Flora 1836, p. 440) biennne, caule elato stricto simplici nudiusculo inferne floccoso superne glaberrimo in racemum simplicem vel parce ramosum abeunte, foliis superne virentibus subtus discoloribus villosis duplicato-crenatis inferioribus breviter petiolatis subcordatis ovato-oblongis, superioribus remotis ovato-lanceolatis, florum fasciculis distantibus, pedicellis glaberrimis inæqualibus, calycis glaberrimi laciinis (lineâ vix longioribus) lanceolatis, corollâ parvâ calyce triplo longiore luteâ fauce purpureâ, filamentis purpureo-lanatis, capsulâ ovato-globosâ calyce duplo longiore ②. *V. leiostachyon* Griseb. Spic. II, p. 43 et *V. leiocaulon* Henffel ex cl. Borbas.

Hab. in Rumelia (Friv.) Vidi iconem MSS. a cl. Arch. Haynald mihi benigne communicatam

Affine *V. nigro* a quo characteribus indicatis differre videtur.

Ar. Geogr. Banatus, Albania.

75. *V. Bithynicum* (Boiss. Diagn. Ser. I, 4, p. 63) biennne, caule elato folioso superne ramosissime paniculato obtuse angulato, foliis supra glabrescentibus viridibus subtus canescentibus radicalibus amplis oblongis duplicato-crenatis inferne longe attenuatis subsessilibus, superioribus parvis ovatis cordato-auriculatis abrupte acuminatis, fasciculis multifloris laxe racemosis, pedicellis calyce longioribus, calyce tomentello (sesquilineam longo) in laciinas linear-lanceolatas partito, corollâ parvâ luteâ, filamentis violaceo-barbatis, capsulâ parvâ ovatâ hirtâ calyce tertiatâ parte longiore ②.

Hab. circa Byzantium (Noë! Castagne), in sylvaticis regionis inferioris Olympi Bithyni (Auct. 2454! Boiss!).

Planta 2-4-pedalis, folia radicalia pedalia pedem dimidium lata, panicula ramosissima multiflora stricta, flores *V. nigri* a quo differt foliorum formâ, caule ample paniculato, lacinis calycinis majoribus. *V. Orientale* foliis parvis, lacinis calycinis angustis, præsertim capsulâ oblongâ distinguitur.

76. *V. Orientale* (M. B. Taur. Cae. I, p. 160) bienne, caule elato folioso superne paniculato, foliis hirtis subtus sepe magis crenescens inferioribus petiolatis oblongis crenatis acutis basi sepe contractis, fasciculis paucifloris in racemos tenuiter virgatos laxos dispositis, pedicellis calyce longioribus, calycis tomentelli (vix sessilineam longi) lacinis subulatis, corollâ parvâ luteâ, filamentis violaceo-barbatis, capsulâ parvâ oblongâ hirtâ calyce duplo longiore ②. *V. Wilhelmsianum* C. Koch Linn. XXIII, p. 574!

Hab ad vias et in locis aridis Georgiae Caucasicæ (Hohen!), Caucasi orientalis (Rupr!), in Tauriâ (ex Stev.).

Differt a *V. nigro* et *V. Bithynico* foliis minoribus angustioribus, capsulæ formâ. *V. Chaixi* Vill. planta Gallica est species diversa. Auctores plures *V. Austriacum* Schrad. et *V. Bannaticum* Schrad. *V. Orientali* adnumerant quod ulterius et præsertim ex speciminiis fructiferis examinandum. *V. Bannaticum* Friv. pl. exs. et Griseb. Spic. II, p. 44 est proles hybrida *V. nigri* prob. cum *V. sinuato*.

* * Calyx major $2\frac{1}{2}$ - $4\frac{1}{2}$ lin. longus. Filamentorum lana albida vel flava.

77. *V. myrianthum*, bienne, multicaule canibus strictis rubellis inferne foliosis superne striete thyrsoideè vel corymbose ramosis, foliis adpresso arachnoideo-canescens inferioribus subtus elevatim nervosis oblongis inferioribus pinnatifolatis in petiolum longe attenuatis lobis irregularibus triangulari-ovatis crenatis, superioribus sessilibus summis linear-lanceolatis integris, fasciculis plurifloris remotis racemos longos spiciformes laxos formantibus, pedicellis calyci aequilongis et longioribus bracteisque linearibus tomentosis, calyce tomentoso ($2\frac{1}{2}$ lineas longo) angusto fere ad basin in lacinias lineares obtusas partito, corollâ minutâ flavâ extus hirtâ, filamentis sordide albo-lanatis longioribus apice nudis, capsulâ parvâ oblongâ tomentellâ acutinerculâ mucronatâ calyce sublongiore ②.

Hab. in Anatoliâ boreali in Paphlagoniâ et Galatiâ circâ Tokat et Kastamouni (Wiedem!).

Sesquipedale, caules rigidi numerosi intricati stricti $\frac{1}{2}$ -1-pedales, folia inferiores 3-5-pollices longa 2 lata, superiora angusta sed sepe elongata, corolla e minimis in secco 4 lin. diam. lata, capsula vix 3 lineas longa. Floribus innumeris insigne, a *V. sphenandroide* foliis lobatis, pedicellis multo brevioribus, etc., discedens.

78. *V. sphenandroides* (C. Koch Linn. XXII, p. 583) bienne, pluricaule caulis strictis inferne foliosis in racemum simplicem vel thyrsoideo-paniculatum abenitibus, foliis radicalibus adpresso, canis vel canescens inferioribus crenatis et sepe tenuiter rugulosis lanceolatis acutis in petiolum breviter attenuatis, caulinis diminutis angustis.

tatis sessilibus, floribus 2-5 fasciculatis superioribus interdum solitariis pedicellis tenuibus calyce duplo longioribus suffultis laxe racemosis, bracteis linear-setaceis, calyce tomentello ($2\frac{1}{2}$ lin. longo) in lacinias oblongo-lanceolatas acutiusculas ad $\frac{3}{4}$ partito, corollà luteâ extus tomentosâ, filamentis aurantiacis sordide albo-lanatis, capsulâ ovato-oblongâ acutiusculâ tomentellâ tandem glabrata calyce tertia parte longiore ②. *V. stachyldifolium* Boiss. et Heldr. Diagn. Ser. I, 12, p. 23.

Hab. in Armeniâ Turcicâ, vallis Tchabantz 6-7000' (C. Koch!), in collibus aridis circa Ispir (Bourg!), Ponto meridionali (Tchih!), in saxosis planicie Iconiensis Lycaoniae inter Karkin et Suliman hatus (Heldr! forma magis cana).

Pedale vel sesquipedale, folia radicalia 2-4 pollicaria 6-12 lin. lata, capsula tres lineas longa pedicello eâ 2-3-p'ō longiore suffulta. Foliæ et habitus *V. biscutellafolii* a quo differt floribus fasciculatis, pedicellis multo brevioribus, calyce non glanduloso, etc.

79. *V. cestroides* (Boiss. et Haussk.) perenne, pluricaule tenuiter et stellatim tomentellum cinereo-virens, caulis inferne foliatis in racemum laxinseulum simplicem vel basi 1-2 ramiis auctum abeuntibus, foliis obsolete repandis oblongis inferioribus in petiolum attenuatis, superioribus sessilibus diminutis ovatis acutis, fasciculis 3-4-floris bracteâ anguste lanceolatâ suffultis, pedicellis inaequalibus longioribus calyceem duplo excedentibus, calyce stellatim et dense tomentoso ($2\frac{1}{2}$ lineas longo) ad $\frac{2}{3}$ in lacinias oblongas obtusiusculas fisso, corollâ luteâ extus tomentellâ, filamentis lateo-lanatis, antheris reniformibus, capsulâ... ②.

Hab. ad Tell Bagdad Mesopotomiae inter Orfa et Kharran (Haussk!) Fl. Maio.

Pedale, racemi semipedales, affine *V. Mesopotamico* a quo differt inflorescentia, tomento crassiore, laciinis calycis longioribus.

80. *V. phyllostachyum* (Boiss. et Haussk.) biennie? totum tomento intescenti aspero densissime pannosum, caulis foliosis crassis fere a basi florum fasciculos axillares racemum longum foliatum formantes edentibus, foliis crebre crenatis et subtus elevatim nervulosis radicalibus oblongis basi cuneatis petiolatis, caulinis cuneatis basi valde cordato-semiamplexicaulibus ovatis abrupte attenuato-acuminatis, pedicellis crassis calyce longioribus, calyce panno (4 lineas fere longo) fere ad basin in lacinias lanceolatas partito, corollâ parvâ valde tomentosâ luteâ, filamentis omnibus flavo-barbatis, antheris reniformibus, capsulâ.... ②?

Hab. in montibus Avroman et Schahu Kurdistanie Persicae 60 0' (Haussk!).

Tomentum crassum *Crozophora*, caules crassi sesquipedales, folia inferiore semipedalia, 2 poll. lata. Species ex speciminibus junioribus imperfecte no a insignis floribus ad axillas longe et inaequaliter pedicellatis folio axillari multo brevioribus, corolla calycis proportione parva videtur.

§ 6. **Racemifera.**

81. **V. racemiferum** (Boiss. et Haussk.) biennne ? adpresso tomentosum cinereum, caule folioso ex omnibus axillis racemos sessiles paniculam foliosam formantes edente, foliis integris ovato-oblongis inferioribus breviter petiolatis obtusis, superioribus semi-amplexicaulibus acutis, floribus in racemis laxiusculis 2-3 fasciculatis bracteis tomentosâ linearis-lanceolatâ acuminatâ fasciculum superante suffultis, pedicellis calyce dimidio brevioribus, calyce tomentoso-hispido ($3\frac{1}{2}$ lineas longo) ad basin in lacinias linearis-lanceolatas aculas partito, corollâ flavâ glabrâ, filamentis glabris, antheris reniformibus, capsulâ... ②?

Hab. in Assyriâ prope Mardin (Haussk.).

Hanc speciem insignem ex unico specimine descripsi; pedalis, folia tres pollices longa, floralia inferiora parum diminuta, racemi axillares circ. 2 pollices longi florum fasciculis subdistantibus, corolla parva videtur. Species filamentis glabris ab omnibus mihi notis discedens.

§ 7. **Triflora.**

82. **V. Mesopotamicum** (Boiss. Diagn. Ser. I, 12, p. 24) perenne ? totum pilis stellatis adpressissimis non contigiis cinerascenti-vires, caulis erectis inferne foliosis in paniculam parce ramosam abeuntibus, foliis parvis oblongis obtusis integris inferioribus in petiolum brevem attenuatis superioribus cordato-ovatis acuminatis, pedunculis racemi centralis subbinis altero unifloro altero trifloro opposite bibracteolato, racemorum lateralium solitariis calyce subduplo longioribus, calyce cinerascenti ($2\frac{1}{2}$ lineas longo) patenti ad medium in lacinias triangulares acutiusculas fisco, corolla dense tomentosâ, filamentis pallide violaceo-lanatis antheris reniformibus, capsula ovata hirtâ calycem vix excedente ④?

Hab. in Mesopotamiâ inter Orfa et Suerek (Ky. exs. 67!), ad Kharput Armeniae australis in collibus arenosis (Haussk. spec. fructiferum foliis orbatum huc spectare videtur).

Pedale, folie inferiora pollicem longa 5-6 lineas lata. Insigne racemi centralis pedunculo altero (ramo abbreviato) opposite trifloro et bibracteolato, pedicellis calyce brevioribus. Capsula piso tertia parte minor.

83. **V. globiflorum** (Boiss. et Noë Diagn. Ser. II, 3, p. 148) perenne, pluricaule totum albo-tomentosum, caulis simplicibus inferne foliosis superne in racemum abeuntibus, foliis inferioribus suberentatis oblongis obtusis in petiolum attenuatis, superioribus et floralibus valde diminutis ovato-oblongis acuminatis, racemi laxiusculi pedunculis ex eâdem bracteis subbinis altero unifloro longitudine calycis altero trifloro bibracteolato pedicello intermedio calyce longiore, calyce dense albo-lanato (4 lin. longo) ad medium in lacinias

ovatas acutæ fissa, corollæ flavæ extus tomentosâ, filamentis sordide albo-lanatis antheris reniformibus, capsulâ... ♀.

Hab. in Galatiâ prope Sivas (Noë 1086!) Fl. Junio.

Pedale, folia inferiora 3-4-pollicaria 12-15 lineas lata, racemus 6-8 pollicaris, alabstra globosa magna, pedunculi saepe pollicares. Inflorescentiâ et characteribus affine *V. Mesopotamico* differt racemo simplici, calyce dense lanato duplo majore, etc.

84. V. Szovitsianum, perenne? totum adpresso et molliter arachnoideo-canum vel canescens, caule elato simplici vel ramoso in paniculam simplicem vel compositam thyrsoidem abeunte, foliis inferioribus oblongis in petiolum attenuatis grosse et obtusiuscale dentatis vel paucilobatis, caulinis paucis ovatis vel oblongis sessilibus, pedunculis subbinis longis basi bracteâ lanceolatâ squamiformi suffultis altero unifloro interdum deficiente altero trifloro bibracteolato pedicello medio calyci subæquilongo lateralibus multo longioribus, calyce tomentoso (tres lineas longo) ultra medium in lacinias late oblongo-lanceolatas acutas fissa, corollæ luteâ tomentellâ, filamentis albo-lanatis, capsulâ... ♀?

Hab. in ditione Transcaucasica Karabagh orientali et in Persiae prov. Aderbidjan (Szovt!).

Planta ut videtur elata pluripedalis inflorescentiâ *V. Mesopotamico* et globo-floro affinis a quibus foliis inferioribus majoribus (fere semipedalibus) grosse dentato-lobatis, pedunculis et pedicellis multo longioribus (pollicaribus) differt.

§ 8. Leiantha.

85. V. parviflorum (Lam. Diet. 4, p. 222) biennne, foliis omnibus radicalibus tenuiter tomentoso-canescens ambitu lanceolatis runcinato-pinnatisectis segmentis numerosis ovatis vel oblongis acutiusculis dentatis saepe sinuato-paucilobis, caule elato tereti purpureo nudo glabro in paniculam ramosissimam corymbosam glabram abeundi, ramis strictis tenuibus virgatis, floribus 2-3 fasciculatis vel superioribus solitariis pedicellis calyce sublongioribus suffultis laxe spicatis, bracteolis minimis ovalis, calyce glabrescenti (vix lineam longo) ad basin in lacinias linearis-stellulatas partito, corollæ parvæ, filamentis usque ad apicem villo lato dense barbatis, capsula ovato-globosâ tomentosâ tandem glabrâ obtusâ calyce sesquialiore ②. *V. taraxacifolium* Boiss. Diagn. Ser. I, 4, p. 59.

Hab. in collibus Lyliae circa Smyrnam (Auch. 1206! Boiss! Bat. 307!), ad Brussa Bithynia (Thirke!) Fl. Mai.

Folia radicalia 6-8-pollicaria 1-2 poll. lata, caulis cum paniculâ 2-3-pedalis, corolla diam. 8-9 lineas lata, magn. *Coriandri*.

86. V. napifolium (Boiss. Diagn. Ser. I, 4, p. 60) biennne, foliis radicalibus rosulatis tomentosis canis breviter petiolatis lyratis

segmentis lateralibus utrinque 1-3 minimis ovatis vel oblongis terminali maximo ovato obtuso basi rotundato omnibus integris vel obsolete crenatis, caulinis paucis parvis sessilibus ellipticis, caule rubello in paniculam ramosissimam corymbosam glabrescentem abeunti, ramis tenuibus virgatis strictis, floribus 2-3 fasciculatis vel superioribus solitariis pedicellis calyce sesquilonioribus suffultis in spicas laxas dispositis, bracteis minimis ovatis, calyce puberulo (lineam longo) in laciniis obtriangulari-spathulatas obtusas partito, corollâ parvâ luteâ, filamentis usque ad apicem flavidо-barbatis, capsulâ tomentellâ tandem glabrescente ovatâ obtusâ calyce $2\frac{1}{2}$ -plo longiore ②.

Hab. in collibus dumosis Cariæ interioris circa Gheyla (Boiss!) Fl. Jun.

Foliorum 2-3-pollicarium segmenta sæpe confluentia terminale 1-2-pollicare, caulis cum paniculâ bipedalis Differt a *V. parrifloro* foliis brevioribus lyratis, caule basi folioso, calycinis laciniis obtriangularibus, capsula paulo major $2\frac{1}{4}$ lin. longa.

87. V. Germanicæ (Haussk. pl. exs.) bienne, pluricaule, caulinis rubellis paulo supra basin in paniculam glabram tenuiter ramosam abeuntibus, foliis inferioribus... caulinis et ramis pubescentibus sessilibus oblongis crenatis, floribus solitariis bracteâ minimâ ovatâ acutâ suffultis pedicellis filiformibus calyce duplo longioribus longe et laxe racemosis, calyce glabro (lineam circ. longo) in laciniis oblongo-spathulatas obtusas acutatas partito, corollâ parvâ flavâ, filamentis sordide purpureo-lanatis, capsulâ pedicello incrassato et elongato arenato eâ triplo longiore suffultâ oblongâ obtusâ mucronulatâ calyce subdupo longiore ②.

Hab. prope Arablar inter Aintab et Marasch Cataoniæ florif. (Haussk!), circa Marasch (Germaniciam) fructif. (Haussk!).

Sesquipedale, paniculæ rami tennes semipedales, folia inferiora jam emarginata, caulina 4-4½-pollicaria, capsula sesquilineam longa. Incomplete notum sed calycis laciniarum formâ ab omnibus affinibus et præsertim a *V. Euphratico* cui facie accedit distinctum.

88. V. denudatum (Boiss. et Heldr. Diagn. Ser. I. 42, p. 23) bienne, foliis fere omnibus radicalibus pannosis griseo-canis ambitu oblongo-spathulatis subruncinatim lyratis petiolatis, segmentis lateralibus parvis utrinque 4-5 oblongis et triangularibus a superioribus confluentibus ad inferiora diminutis terminali ovato vel oblongo multo majore 3-4-lobato vel repando, caule rubello nudo glabro in paniculam ramosam glabram longam abenante, floribus binis vel solitariis pedicello calyci æquilongo suffultis laxe racemosis, bracteis minimis oblongis acutis, calyce subglanduloso (sesquilineam longo) in laciniis late lineares obtusas partito, corollâ flavâ, filamentis lutescenti barbatis longioribus apice nudis, capsulâ glabrâ ovato-globosâ obtusâ mucronulatâ calyce sesquiloniore ②.

Hab. in rupestribus fœncis Tsimbukchan Pamphyliæ et in collibus tophaceis prope Buldur Pisidia (Heldr!), in Lyciâ prope Elmalu (Bourg!).

Planta 3-4-pedalis, corolla diam 12-15 lin. lata, capsula piso minor. Affine *V. Smyrneo* differt foliorum segmentis lateralibus pluribus, capsula subglobosa nec oblongo-cylindrica.

89. V. Isaureum (Boiss. et Heldr. Diagn. Ser. I, 12, p. 22) bienne, pluricaule præter paniculam glabratam viscidulo-hirtam inferne indumento denso albo vel flavido pannosum, foliis radicalibus ovato-oblongis basi cuneatis petiolatis, caule inferne crebre folioso et tomentoso superne ramosissime paniculato, foliis caulinis pannosis diminutis auriculato-amplexicaulibus, floribus binis et solitariis secus paniculae ramos tenuiter virgates laxe spicatis pedicellis calyci subæquilongis suffultis, bracteis minimis ovatis acutis, calyce glanduloso (sesquilineam longo) ad $\frac{3}{4}$ in lacinias lanceolatas acutas partito, corollâ luteâ glabrâ, filamentis sordide flavo-lanatis, longioribus superne nudis, capsula glabrescente ovatâ acutiusculâ longe mucronatâ ②.

Hab. in saxosis apries prope urbem Ermenek Isauriae (Heldr! Peron!) Fl. Jun.

Folia radicalia triplicaria, caules bipedales paniculae ramis erecto-patulis, corolla diam. pollicaris, capsula piso subminor. Affine *V. Pinardi* differt foliis brevibus dense pannosis, canle inferne dense foliato, calyce.

90. V. kochiaeforme (Boiss. et Haussk.) perenne, pluricaule dense pannosam canum, caulis interne foliosis superne in paniculam corymbosam abeuntibus, foliis integerrimis obtusiusculis parvis oblongis et ovatis inferioribus basi cuneatis petiolatis, superioribus crebris sessilibus subcordato-amplexicaulibus, floribus solitariis secus paniculae ramos gracillimos glabratos laxissime racemoso-spicatis, pedicellis calyce subbrevioribus bracteas squamiformes superantibus, ealyce (lineam longo) glabro glandulis subsessilibus sparsis obsito ad basin in lacinias ellipticas obtusas partito, corollâ parvâ luteâ parce glandulosâ, filamentis purpureo-lanatis antheris reniformibus, capsula globosâ tomentellâ tandem glabratâ longe mucronatâ calyce sublongiore ȝ.

Hab. in montibus culcareis Persiae australis inter Tchinar et Maregun et Inter Seytun et Bebehan (Haussk!).

Sesquipedale, foliorum inferiorum lamina 1½-2-pollicaris, paniculae ramí tenuissimi 4-6-pollicares. Facies *V. Syriaci* et *V. simplicis*, sed flores solitarii, calyx triplo minor etc.

91. V. rubricaulé (Boiss. et Heldr. Diagn. Ser. I, 12, p. 23) bienne, foliis parvis albo-tomentosis integris vel obsolete crenatis radicalibus et inferioribus oblongis obtusis in petiolum brevem attenuatis, caulinis crebris strictis oblongis obtusiusculis basi subauriculatis superioribus glabrecentibus, caule rubro elato in paniculam glabrescentem subglandulosam longe striete et tenuiter ramosam abeundi, floribus solitariis pedicello ealyce multo breviore suffultis

laxe spicatis, bracteis minimis ovatis acutis, calyce viscidulo-glanduloso (sesquilineam longo) in lacinias elliptico-ovatas acutas ad $\frac{2}{3}$ partito, corollâ luteâ, filamentis flavidо-lanatis, capsulâ ovatâ obtusâ glabrâ calyce sesquilongiore ②.

Hab. in fruticetis apricis Tauri Isaurici inter Karaman et Ermenek in valle Djoksu prope Pichacktschi (Heldr!) Fl. Jun.

Gracile quadripedale, folia inferiora $2\frac{1}{2}$ -3-pollices longa pollicem vix lata, caps. magn. Pisi. Affine *V. Isaurico* specifice differre videtur foliis minoribus, ramis longius virgatis, floribus brevius pedicellatis, capsulâ majore obtusâ.

92. V. adenocaulon (Boiss. et Bal. Diagn. Ser. II, 6, p. 128) bienne, caule inferne folioso et arachnoideo-lanato dein nudo paniculato ut tota inflorescentia præter glandulas minimas sparsas glabro, foliis adpresso arachnoideo-canis brevibus radicalibus petiolatis ovatis obtusissimis obsolete repandis, caulinis paucis confertis diminutis oblongis sessilibus, floribus solitariis pedicello calyce breviore suffultis laxe racemosis, bracteis linearibus minimis, calyce (sesquilineam longo) in lacinias lineares acutiusculas partito glandulis stipitatis obsito, filamentis sordide albo-barbatis, longioribus apice nudis, corollâ parvâ, capsulâ glabrâ ovatâ obtusâ mucronatâ calyce duplo longiore ②.

Hab. in collibus ad rivum Kamechly Tchaï prope Berekely Cappadociæ (Bal!).

Ex unico specimine sesquipedali notum, caulis inferne brevi spatio foliosus dein paniculatus nudus, folia radicalia cum petiolo 2-3 pollices longa, capsula piso subminor. Affine *V. rubricauli* ab eo et cæteris gregis indumento arachnoideo-lanato foliorum discrepans, bractearum et calycis laciniarum forma aliena.

93. V. Græcum (Heldr. et Sart. in Boiss. Diagn. Ser. II, 3, p. 148) bienne, caule crasso præter basin foliosam albo-tomentosam glabro cito in paniculam amplam ramosissimam glabram abeunte, foliis integris vel subrepandis inferioribus dense pannosis niveis obovatis vel oblongis in petiolum attenuatis, caulinis paucis diminutis sessilibus tomento detersili tandem glabratris, fasciculis 4-5-floris subdissitis bracteis linearibus suffultis racemos longos laxos formantibus, pedicellis calyci æquilongis vel longioribus, calyce glabriuscule (lineam lengo) ad basin in lacinias lineares obtusas partito, corollâ parvâ luteâ, filamentis sordide albo-barbatis longioribus apice nudis, capsulâ demum glabratâ cylindricâ calyce triplo longiore ②.

Hab. in regione inferiori et mediâ montis Parnes Atticæ 2000'-2500' (Heldr!).

β. Ætolicum. — Folia majora tenuius tomentosa subvirentia, capsulae paulo majoras.

Hab. in monte Arapocephala Ætoliae (Samar. et Guicci!).

γ. Zuccarinii. — Folia radicalia obtusiora spathulato-suborbiculata, flores secus paniculæ ramos bini vel solitarii, calycis laciniae sub-

abbreviatæ subobtusiores. — *Celsia tomentosa* Zuccar. Pl. Nov. fasc. 2, p. 22 !

Hab. in Græciâ sept. inter Negrepont (sphalmate Megara) et Zeitun Thesalîæ (Berger!) Vidi in Herb. Reg. Monac.

Bipedale, folia radicalia semipedalia et longiora, in var. γ 2-3-pollicaria, paniculæ rami pedales, corolla diam. 8 lineas latâ, capsula 2½-3-lineas longa. Affine *V. myriocarpo*, folia non undulato-subblobata, corolla glabra, capsulae majores. Amiciss. Heldreich legit in insulâ Cephalloniâ specimen nimis incompletum et floribus orbatum huic speciei affine sed capsulâ ovatâ diversum

94. V. cylindrocarpum (Griseb. Spic. II, p. 46) bienne, caule atro glaberrimo viscidulo paniculato, foliis radicalibus subpetiolatis oblongis undulato-sinuatis, rameis parvis integris ovato-oblongis sessilibus omnibus utrinque sed subtus densius et adpressissime stellato-pubescentibus glaucis, paniculæ amplæ ramis erecto-patentibus ramulosis, pedicellis calyce triplo longioribus, calyce glabro viscido (1 lineam longo) ad basin in lacinias breviter lanceolatas capsulâ triplo breviores partito, corollâ parvâ flavâ glabriusculâ, capsulâ cylindricâ glabrâ viscidâ obtusâ subretusâ ②.

Hab. in arenâ littorali insulæ Thasos Macedoniae prope Panagia versus Kastro (Griseb!) Vid. fruct. in herb. cl. Griseb.

Planta elata, rami sesquipedales, folium (ex inferioribus) 7 pollicare 3 pollices latum obtusum, panicula pedalis, capsula 2 lineas longa. Affine *V. Graecum* a quo foliorum formâ, indumento, etc. differt.

95. V. Smyrneum (Boiss. Diagn. Ser. I, 4, p. 61) bienne, foliis fere omnibus radicalibus rosulatis argyreo-panniosis lyratis segmentis lateralibus utrinque 1-2 paucis auriculiformibus ovalis interdum evanidis terminali maximo orbiculari vel ovato sinuato-dentato basi truncato vel rotundato, foliis caulinis paucis ovatis, caule rubro glabrescenti paniculato, paniculæ ramis virgatis, floribus 2-3 fasciculatis vel solitariis secus ramos laxè spicatis pedicellis calyce sublongioribus suffultis, bracteis minimis triangularibus, calyce (lineam longo) glabrescenti in lacinias lanceolato-lineares partito, corollâ parvâ luteâ, filamentis flavidò-barbatis junioribus superne glabrescentibus, capsulâ tomentellâ dein glabrescente oblongo-cylindricâ obtusâ calyce 2 ½-plo longiore ②.

Hab. in declivibus dumosis montis Corax Lydiæ ad occasum Smyrnæ (Boiss!) Fl. Mai.

Facies *V. napifolii*, folia magis pannosa, ab eo egragie differt calyeis formâ, capsulâ elongatâ 3 lin. longâ lineam latâ.

96. V. Cilicicum (Boiss. Diagn. Ser. II, 3, p. 143) bienne, foliis indumento stellato tenui canescens radicalibus oblongis obtusis integris in petiolum longiusculum breviter attenuatis, caulinis paucis sessilibus breviter petiolatis, caule rubello fere a basi in paniculam ramosissimam glabratam abeunte, fasciculis 5-7-floris bracteis parvis linearibus suffultis, pedicellis calyce sublongioribus,

calyce puberulo (lineam longo) ad basin in lacinias linearis-oblongas partito, corollâ luteâ minutâ, filamentis albo-lanatis, capsulis pedicellis arcuatis 2-3-plo longioribus suffultis oblongis obtusis mucronulatis calyce tertiatâ parte longioribus ②.

Hab. ad ruinas castelli Gülek Boghas Tauri Cilicici (Bal!) Fl. Jun.

Foliorum radicalium lamina 2-3-pollicaris sesquilineam lata, panicula pedalis, capsula 2 lineas longa. Affine *V. Pinardi* cuius fructus ignotus et a quo distinctum videtur indumento tenuissimo nec pannoso niveo, fasciculis plurifloris, bracteolis linearibus nec ovatis, calyce puberulo.

97. *V. leianthum* (Bth. in D.C. Prodr. X, p. 239) biennne, foliis superne adpresso hirtis virentibus subtus tenuiter canescentibus oblongis obtusis crenatis, caulinis inferioribus basi attenuatis superioribus basi semiamplexicauli subdecurrentibus, paniculâ amplâ glabrâ ramis longissime virgatis, fasciculis plurifloris bracteâ triangulari suffultis laxiuscule et longissime racemosis, pedicellis inaequalibus calyce longioribus, calyce glabro (lineam longo) ad $\frac{3}{4}$ in lacinias ovato-oblongas acutas partito corollâ parvâ, filamentis sordide violaceo-lanatis, capsulâ oblongâ retusâ mucronatâ calyce duplo longiore ②. *V. microcalycinum* Fenzl MSS.

Hab. ad Tauri Cilicici radices (Ky. 388!).

Specimina incompleta in quibus plantæ pars inferior deficit, folia inferiora semipedalia $2\frac{1}{2}$ pollices lata, racemi tenues sesquipedales, corolla e minimis gregis; inter affinitatem fasciculis plurifloris approximatis racemum multilorum formantibus insigne.

98. *V. calvum* (Boiss. et Ky. in Ky.-Sched. 1859) paniculâ amplâ glabrâ viscidâ ramis longissime virgatis, fasciculis plurifloris bracteâ brevi triangulari suffultis laxe et longe racemosis, pedicellis inaequalibus calyce 3-4-plo longioribus, calyce glabro ad basin usque in lacinias lanceolatas partito, corollâ majusculâ, filamentis albo-lanatis, capsulâ ovato-oblongâ retusâ calyce triplo longiore ②.

Hab. in aridis a.1 Bitlis Kurdistanæ (Ky. Suppl. 721!).

Ex frustulis incompletis in quibus plantæ tota pars inferior et omnia folia desunt descriptum. Paniculæ rami stipe bipedales, facies *V. leianthi*, flores longius pedicellati, corolla major, filamenta albo nec violaceo-lanata. Capsula $2\frac{1}{2}$ lineas longa.

99. *V. lætum* (Boiss. et Haussk.) perenne? pluricaule, caulinibus glabrescentibus inferne foliatis cito in paniculam ramosam abeuntibus, foliis integris indumentostellato tenui obtusis cinereis inferioribus oblongis subspathulatis obtusis in petiolum attenuatis, superioribus sessilibus strictis acutis, floribus solitariis pedicello calyce subbrevisse suffultis laxiuscule racemosis, bracteis oblongis acutis calyce longioribus, calyce ut et bracteæ glabro glandulis sparsis stipitatis asperulo (2 lineas vix longo) ad basin in lacinias longas lineares acentiusculas partito, corollâ luteâ rarius albidâ, filamentis purpureo-

Ianatis antheris reniformibus, ovario glabro oblongo-cylindrico 2 $\frac{1}{2}$.
V. Pinardi D.C. Prodr. quoad pl. ex Mesopotamiâ.

Hab. in graminosis montis Gebel Taktak Mesopotamiae (Haussk! Aucher. 31281 et 2459!) erronee e Persia bor. indic. Fl. Mai.

Sesquipedale vel bipedale, folia radicalia semipedalia, caulinis valde diminuta, panicula floribunda ramis semipedalibus, corolla diametro 9 lineas lata

100. ***V. Pinardi*** (Boiss. Diagn. Ser. I, 4, p. 62) biennie, foliis pannoso-canis indivisis inferioribus oblongo-lanceolatis in petiolum longe angustatis integris vel obsolete repandis, caulinis sessilibus ellipticis et linearibus, caule inferne tomentello in paniculam corymbosam glabram abeunte ramis tenuibus virgatis, floribus 2-3 fasciculatis vel solitariis laxe spicatis pedicellis calyce sublongioribus suffultis, bracteis minimis ovatis acutis, calyce (lineam longo) glabro in lacinias lineares acutas partito, corollâ luteâ, filamentis dense flavido-barbatis, longioribus apice nudinseculis, capsulâ.... (2).

Hab. in Cariâ (Pinard!).

Folia inferiora argyrea cum petiolo 5-6-pollicaria 1-2 poll. lata nervis subtus prominentibus; a precedentibus foliis indivisis discrepans. Corolla paulo major 10-12 lin. diametro lata.

101. ***V. cerinum*** (Boiss. et Heldr. Diagn. Ser. I, 12, p. 21) biennie, foliis fere omnibus radicalibus densissime pannosis flavescens-tibus asperis integris oblongis et oblongo-lanceolatis acutiusculis in petiolum brevem sensim attenuatis, caule fere a basi paniculato praeter folia pauca hracteiformia nudo cum paniculâ glabro ceruo-nitido, floribus 3-5 fasciculatis subsessilibus, fasciculis secus paniculæ ramos remotis bracteâ parvâ ovatâ suffultis, calyce glabro (sesquilineam longo) ad $\frac{3}{4}$ in lacinias ovatas acutas partito, corollâ luteâ mediocri, filamentis flavo-barbatis, capsulâ... (2).

Hab. in collibus aridis Lycaoniæ ad occasum urbis Koniah leucâ una remotis (Heldr!) Fl. Jun.

Folia radicalia crebra 6-8 poll. longa $1\frac{1}{2}$ -2 lata, caulis cum paniculâ, $1\frac{1}{2}$ -2-pedalis. A *V. Pinardi* foliorum indumento densiore et aspero, floribus sessilibus, calyce, etc. distinctum.

§ 9. Blattarioidea.

* Flores sessiles.

102. ***V. Steveni*** (Boiss. et Buhse Aufz., p. 160) biennie, totum tomento stellato ad folia tenui cinerascenti ad racenum et calyces denso pannoso obsitum, caule erecto crassiuseulo in racemum simplicem vel basi parce et breviter ramulosum abeunti, foliis crenato-dentatis acutis oblongis inferioribus in petiolum attenuatis superioribus sessilibus, floribus solitariis subsessilibus laxe spicatis bracteâ

anguste lanceolatâ longe setaceo-acuminatâ calyce longiore suffultis, calyce dense cano-tomentoso (fere 5 lineas longo) in lacinias late triangulari-ovatas acutas ad medium fisso, corollâ flavâ extus tomentosâ, filamentis flavescenti-barbatis, antheris reniformibus, capsulâ juniore tomentosâ globosâ ②.

Hab. in valle Alyndsha prope Nachitschewan Armeniæ Rossicæ (Buhse!) Fl. Maio.

Planta 1 $\frac{1}{2}$ -2-pedalis, foliorum radicalium lamina 3-4-pollicaris pollicem fere lata. Affine *V. alceoidi* differt foliis radicalibus tenuiter tomentosis acutis, bracteis setaceo-acuminatis, antheris omnibus reniformibus, etc. Inter *Blattarioida* ob flores solitarios militans sed ob calyces magnos artificialiter; habitus *V. salviifolia*.

* * Flores pedicellis calyce brevioribus vel eo æquilongis suffulti.

103. ***V. pyramidatum*** (MB. Taur. Cauc. I. p. 161) bienne, elatum, tomento tenui obsitum, caule crasso longe pyramidato, foliis supra viridibus subtus canescensibus radicalibus magnis oblongis basi longe angustatis duplicatim et obtuse lobatis vel crenatis, superioribus sessilibus ovatis auriculato-cordatis longe acuminatis, paniculâ canescente ramis tandem laxiusculis, floribus solitariis vel binis pedicello calyci æquilongo suffultis, calyce (fere tres lineas longo) tomentoso in lacinias oblongas vel ovatas ultra medium partito, corollâ amplâ, filamentis purpureo-lanatis, capsulâ ovatâ magnâ tomentellâ acutiusculâ pedicello æquilongâ calyce tertia parte longiore ②. Ic. Sw. Brit. Fl. Gard. tab. 31. — *V. alpigenum* C. Koch Linn. XXIII, p. 574.

Hab. in regione alpinâ Ponti Lazici 5500'-7000' (C. Koch! Bal!), subalpinis Guria (Nordm.), Georgiæ Caucasicæ et prov. Talysh (M B., Hoh!), Daghستانie 6000' (Rupr!).

Planta 2-4-pedalis paniculâ interdum bipedali, folia inferiora pedalia et longiora, capsula 3 $\frac{1}{2}$ lin. longa.

104. ***V. chrysorrhacos*** (Boiss. Diagn. Ser. I, 4, p. 58) bienne, totum indumento stellato aureo adpressissimo tomentosum, caule pumilo a basi pyramidatum paniculato, foliis inferne confertis integris oblongo-lanceolatis acutiusculis basi longe attenuatis, floralibus parvis oblongo-triangularibus acuminatis, floribus secus paniculæ rames erecto-patentes breves solitariis remotiusculis pedicello calyce sub-breviore bracteâque oblongâ acuminatâ pedicello æquilongâ suffultis, calyce chryseo (3 $\frac{1}{4}$ lin. longo) ad medium in lacinias ovato-triangulares fisso, corollâ intense flavâ calyce triplo majore, filamentis aureo-barbatis, longioribus apice nudis, capsulâ.. ②.

Hab. in regione alpinâ Cadmi orientalis supra Colossam in Cariâ (Boiss!) Fl. Jun.

Pedale, paniculæ rami 3-5-pollicares, folia inferiora cum petiolo 3-5 pollicaria 10-15 lineas lata, corolla diam. fere pollicaris. Species indumento inten-^se aureo pannoso elegantissima.

Hoc inter et *V. cheiranthifolium* plantæ hybridae occurruunt = *V. Cadmeum* Boiss. olim.

105. V. Schimperianum (Boiss. Diagn. Ser. I, 42, p. 11) pannosum flavidо-аeruginosum, foliis fere omnibus prope collum confertis oblongis obtusis in petiolum brevem attenuatis obtuse sinnatolobatis marginе valde undulatis et crispis, caule fere a basi paniculatum ramosissimo præter 1-2 folia squamuformia ovato-cordata aphylla ramis tenuiter virgatis patentibus in racemos longos gracillimos abeuntibus, floribus solitariis breviter pedicellatis, calyce tomentoso (sesquilineam vix longo) in lacinias ovatas ad medium fissō, filamentis sordide albo-lanatis, capsulā ovatā mucronatā calyce sublongiore ②. *V. crispum* Ehrenb. MSS.

Hab. in Arabiā petreā loco non notato (Schimp!), in viā inter Tor et montem Sinai (Ehr!).

Folia radicalia 2-4-pollicaria, panicula patula ramosissima ramis gracilibus cum racemo saepē pedalibus. Affine *V. plicato* et *V. sinuato* sed insigne paniculā gracili aphylla, floribus minimis solitariis; a *V. sinuato* insuper lana staminum albida et capsulā majore ovatā distinctum.

106. V. Basianicum (Boiss. et Hassk.) biennne, totum tomento adpresso tenui canescens, caule inferne foliato eito in paniculam ramosissimam corymbosam abeunte, paniculae rauis virgatis tenuibus erecto-patulis iterum saepē ramulosis, foliis inferioribus breviter oblongo-lanceolatis obtusiusculis grosse dentato-sublobatis basi breviter cuneatis et petiolatis, caulinis paucis subintegris ovatis auri culato-amplexicaulibus, floribus secus paniculae ramos solitariis laxe racemosis pedicellis calyci subæquilongis suffultis, bracteis tomentosis pedicello brevioribus, calyce tomentoso (sesquilineam longo) ad basin in lacinias oblongas acutiusculas partito, corollā parvā luteā extus tomentosā, filamentis sordide luteo-lanatis, capsulā... ②.

Hab. ad Derbent i Basian Mesopotamiae (Haussk!).

Sesquipedale, folia inferiora 2-3-pollices longa pollicem lata, paniculae racemi 6-8-pollicares, facies *V. sinuati* sed folia non sinuato-lobata et flores solitarii. A *V. Schimperiano* floribus solitariis quoque donato foliis non rugosis nec valde undulato-crispis, etc. alienum. *V. Tiberiadis* magis affine differt inumenti naturā, foliis multo minoribus formae diversæ, etc.

107. V. Tiberiadis (Boiss. Diagn. Ser. I, 42, p. 26) biennne, totum inferne papillis longis adpressiusculis albis obsitum partibus floralibus tomentello-glandulosum, caule crasso parte inferiori crebre foliato superne in paniculam ampliam ramosissimam abeunte, foliis radicalibus oblongis vel oblongo-lanceolatis acutis in lobos magnos irregulares triangulari-lanceolatos dentatos ad medium profunde incisis in petiolum brevem sensim attenuatis, caulinis inferioribus sessilibus, supremis sensim diminutis a basi cordatā acuminitatis dentulatis, floribus solitariis pedicellis calyce brevioribus suffallis racemos virgatos laxissimos formantibus, bracteis parvis triangulari-

cordatis acutis. calyce dense glanduloso-tomentello (sesquilineam longo) in lacinias ovatas fere ad basin partito, corollâ flavâ extus hirtâ, filamentis purpureo-lanatis antheris reniformibus, capsulâ hirtâ ovato-globosâ obtusâ longe mucronatâ calyce sublongiore pedicello incrassato erecto-patulo subaequilongo suffultâ ②.

Hab. in Galilæa circa lacum Tiberiadis et infra Banias (Boiss!) Fl. Mai.

Folia inferiora cum petiolo $\frac{1}{2}$ -1 pedem longa, caulis cum paniculâ 2-3-pedalis, corollæ eis *V. lyratifolii* minores, capsula pisum fere æquans.

108. ***V. Aleppense*** (Benth. in D.C. Prodr. X, p. 241) biennie? caule crassiuscule simplici inferne folioso tomentello superne in racemum simplicem vel in paniculam parce ramosam strictam thyrsoidream glabram abeunte, foliis rosulæ sterilis canis imbricatis oblongis valde undulato-sinuatis, fertilis tenuiter subtus densius canescentibus, inferioribus oblongo-lanceotatis dentatis et interdum inciso-lobatis in petiolum attenuatis, superioribus sessilibus subcordato-amplexicaulibus plicato-subcanaliculatis oblongo-lanceolatis acuminatis, floribus solitariis laxe racemosis pedicellis calyci æquilongis, suffultis, bracteis triangulari-oblongis acutis pedicello æquilongis, calyce ut et pedicelli et bracteæ crebre stipitato-glanduloso ($2\frac{1}{2}$ lin. longo) in lacinias lanceolatas acutas ad $\frac{3}{4}$ partito, corollâ luteâ extus hirtulâ tandem glabrata, filamentis sordide purpurascenli-lanatis, antheris reniformibus, capsulâ ovatâ crasse mucronatâ calycem vix excedente ②.

Hab. in Syria prope Aleppo (Auch. 1839!), in Mesopotamia prope Tcharmelik, Biredjick, Erbil et ad fluvium Chabur (Haussk!).

Pedale vel sesquipedale, folia inferiora 3-4 pollicaria 9-12 lin. lata, paniculæ rami crassi breves rigidi, corolla diam. 9 lin. lata, capsula piso subminor.

109. ***V. infidelium*** (Boiss. et Haussk.) biennie? pluricaule caulinis inferne foliosis in racemum simplicem abeuntibus ant parte inferiori parce ramosis ut et inflorescentia tota glandulis sessilibus pilisque longis patulis rigidis obsitis, foliis crenatis radicalibus albo-pannosis oblongis acutis in petiolum brevem attenuatis, caulinis cinerascentibus a basi sessili semiamplexicauli lanceolatis attenuato-acuminatis, floribus solitariis pedicello calyce breviore et bracteæ lanceolatae acutæ calycem subaequante suffultis, calyce hispido in lacinias lanceolatas acutas vel acuminatas partito (circ. $2\frac{1}{2}$ lin. longo), corollâ luteâ sparsim hirsutâ, filamentis longis flavidio-lanatis superne nudis, capsulâ ovatâ hirsutâ mucronatâ calyce subbreviore ②.

Hab. in pascuis ad lacum Giaurgöl ad radices montis Giaurdagh Cataoniae non procul a Marasch (Haussk!).

Affine characteribus et habitu *V. Aleppensi*. differt foliis non inciso-dentatis, inflorescentiæ hispidâ nec glandulosâ glabrâ, calycis laciñis acutioribus, filamentorum lanâ flavâ.

110. ***V. digitalifolium*** (Boiss. et Haussk.) perenne? totum

praeter paniculæ ramos glabratos tomentosum griseo-canum, caule parte inferiori folioso in paniculam thyrsoidream abeunti, foliis sub-integris subtus prominenter costatis et nervosis oblongis acutis radicibus in petiolum angustatis, caulinis sensim diminutis sessilibus semiamplexicaulibus, floribus praeter inferiores rami centralis interdum binos solitariis pedicello calyce subbreviore suffultis laxiusculæ racemosis, bracteis tomentosis oblongis acuminatis pedicello longioribus, calyce stellato-tomentoso (fere tres lineas longo) in lacinias elliptico-lanceolatas acutas ad basin partito, corollâ amplâ pilis stellatis obsitâ, filamentis violaceo-barbatis, antheris reniformibus, ovario juniore ovato tomentello, capsulâ... ♀?

Hab. in monte Gebel Taktak Mesopotamiae in consortio *V. laeti* (Haussk!).

Multis notis *V. laeto* affine etsi inter *Leiantha* non militans, folia radicalia breviora, caulina numerosiora omnia densius tomentosa, calyces majores tomentosi lacinii latioribus, corolla diam. pollicaris extus parce tomentella.

111. ***V. alyssifolium*** (Boiss. Diagn. Ser. I, 4, p. 58) perenne pluricaule totum adpressissime canum, rhizomate crasso duro pellicula reliquiis vestito, caulis humilibus simplicibus vel paree ramosis in racemos laxos abenntibus, foliis radicalibus consertis erassiusculis linearis-spathulatis et linearibus obtusis subcanaliculatis basi subattenuatis, caulinis paucis sensim brevioribus et acutatis, floribus subsolitariis pedicellis tenuissimis tandem incrassatis calyce duplo longioribus suffultis, bracteis squamiformibus minimis, calyce cano (lineam vix excedente) in lacinias lanceolato-lineares fere ad basin partito, corollâ tomentellâ parvâ, filamentis sordide barbatis, antheris reniformibus, capsulâ ovalâ acutiusculâ demum glabratâ calyce tertiatâ parte longiore ♀.

Hab. in Armeniâ (Auch. N° 2366! spec. florifera). Hunc quoque spectare videntur spec. fructifera ex monte Berydaghi Cataoniae 6'00' (Haussk! *V. Armenum* MSS.).

Caules semipedales demum pedales tenues, folia 2-3 pollices longa 2-2 $\frac{1}{2}$ lineas lata, corolla 5-6 lineas diam. lata, capsula piso dimidio minor. Species foliis insignis

* * * Flores pedicellis calyce pluries longioribus suffulti.

112. ***V. Euphraticum*** (Blh. in D.C. Prodr. X, p. 241) biennne? pluricaule, pilis stellatis sparsis adpressis canescens-virens, foliis inferioribus obtuse repando-crenatis spathulato-ellipticis in petiolam longe attenuatis, caulinis et rameis sessilibus linearibus integris, paniculâ ramosissimâ ramis elongatis tenuibus flexuosis, floribus solitariis inferioribus rarius geminis pedicellis calyce subduplo capsulâ 3-4-plo longioribus suffulti, calyce tomentoso (lineam longo) ad $\frac{3}{4}$ in lacinias ovato-triangulares acutiusculas partito, corollâ tomentosâ luteâ parvâ, filamentis sordide purpureo-lanatis, capsulâ parvâ tomentellâ ovato-globosâ calyce subduplo longiore ②.

Hab. in Cappadociâ ad Euphratem (Auch. 2445!).

Ex speciminibus imperfectis tantum notum, sesqui vel bipedale, folia inferiora 2-3-pollicaria 6-8 lineas lata longiuscule petiolata, caulinâ angustissima, bracteæ squamiformes acutæ brevissimæ, pedicelli fructiferi 6 lineas longi arcuati, capsula piso dimidio minor.

113. V. leuconeurm (Boiss. et Heldr. Diagn. Ser. I, 42, p. 27) bienne, caule tereti elato superne rubello glabro in paniculam compositam amplissimam glabram abeunti inferne folioso cum foliis lanâ albâ adpressâ arachnoideâ ad nervos copiosiore obsitum, foliis integris oblongis obtusis inferioribus in petiolum attenuatis, superioribus sensim diminutis strictis truncato-subcordatis, floribus solitariis pedicello tenui erecto-patulo incurvo calyce 4-5-plo longiore suffultis laxe et longe racemosis, bracteis miniumis ovato-triangularibus, calyce glanduloso (vix 2 lineas longo) in lacinias ovato-oblongas partito, corollâ, capsulâ parvâ ovato-globosâ calyce sublongiore pedicello 4-5-plo breviore ②.

Hab. in pinetis partis superioris vallis Ermenek Tauri Isaurici (Heldr.).

Ex specimine deflorato et fructifero descriptum, 4-5-pedale, folia et indumentum *V. Wiedemannianii*, inflorescentia et pedicelli elongati *V. lyratifolii*. Stamina non vidi et *Verbasco* tantum ex habita ad numero.

114. V. lyratifolium (Köchel in Ky. exs. d.c. Prodr. X, p. 242) bienne, totum papillis longis hirtum preterea ad inflorescentiam glandulosum, caule elato gracili basi folioso superne simplici vel ramoso in racemos multifloros longissimos graciles abeunti, foliis viridibus radicalibus et inferioribus petiolatis ambitu ovatis vel oblongis lyrato-pinnatifidis segmentis acute dentatis lateralibus utrinque 2-4 parvis lanceolatis terminali oblongo dentato et triangulari-löbato. foliis superioribus sessilibus ovatis denticulatis acutis, floribus solitariis pedicellis tenuibus patentibus calyce multoties longioribus suffultis bracteolâ minimâ linearî obsitis laxe racemosis, calyce parce glanduloso (2 lin. longo) fere ad basin in lacinias oblongo-lineares acutas partito, corollâ flavâ glabrâ, filamentis purpureo-lanatis, stamine quinto brevissimo, capsulâ globosâ glabrâ calyce duplo longiore ②. *V. Köchelii* Fenzl MSS.

Hab. in Ciliciâ ad radices Tauri (Ky. 335!), ad Kechlick prope Mersina (Bal.) Fl. Mai.

β. linearilobum Boiss. Diagn. Ser. II, 3, p. 150. Folia inferiora in lacinias anguste lineares elongatas pinnatum vel bipinnatum secta.

Hab. ad Boulokli prope Mersina Ciliciæ (Bal. 678!), in valle Su Nedere propè Gülek (Ky. Suppl. 348!).

Planta 3-4-pedalis, folia inferiora 4-6-pollicaria, pedicelli fructiferi pollicem longi, corolla diametro pollicaris, capsula piso tertiatâ parte minor. Specimen Aucheri 846 e Phrygiâ huc in d.c. Prodr. citatum est *V. xanthophœnicum*

115. V. Berytheum (Boiss. Diagn. Ser. I, 42, p. 28) bienne, totum virescens breviter hirtellum glandulosum radice brevi oblongâ

tuberiformi, caule elato tereti laxe paniculato oligophyllo, foliis radi-calibus rosulatis petiolatis ambitu oblongo-spathulatis pinnatisecto-lyratis segmentis inferioribus utrinque 2-3 ovato-oblongis parvis interdum nullis superiore maximo ovato vel oblongo irregulariter sinuato-lobato omnibus denticulatis obtusis. foliis caulinis paucis sessilibus oblongis et lanceolatis basi dentatis, floribus solitariis pedicello erecto-patenti arcuato calyce 4-6-plo longiore suffultis racemos longos laxos formantibus, bracteis parvis lanceolatis acuminatis. calyce glanduloso sesquilineam longo in lacinias elliptico-lineares acutas partito, corollâ glabrâ luteâ, filamentis violaceo-tanatis superioribus apice nudis. capsulâ ovato-globosâ glabrâ calyce sesquilongiore ②. *V. bulbosum* Ehrenb. MSS.

Hab. ad sepes et margines viarum cavarum circa Bezzylthum Syriæ (Ehr! Boiss! Gaill!) Fl. vere.

Caules 2-3-pedales racemis tenuibus longissimis, folia radicalia 3-6-pollicaria, pedicelli tandem pollicares et longiores, corolla diam. 9-10 lineas lat., capsula piso minor. Affine *V. lyratifolio* quod differt indumento longe papillari, foliorum magis divisorum segmentis et lobis acutis, corollâ majore etc.

116. *V. Blancheanum*, bienne, viride parce papilloso-hispidum superne subglandulosum, radice brevi subcylindricâ, caule in racemum longum basi breviter et stricte ramulosum abeunte. foliis radi-calibus rosulatis petiolatis oblongis triangulari-lobatis lobis denticulatis, caulinis paucis valde diminutis sessilibus triangularibus auriculatis acuminatis inciso-dentatis, floribus solitariis pedicello arcuato calyce triplo longiore suffultis laxe racemosis, bracteis parvis triangularibus acuminatis, calyce parce glanduloso $2 \frac{1}{2}$ lineas longo in lacinias ellipticas basi attenuatas acutissimas mucronatas sœpe denticulatas partito, corollâ luteâ, filamentis purpureo-barbatis longioribus apice nudis, capsulâ globosâ tomentellâ laciinis calycinis accretis superatâ ②.

Hab. ad vias, in cultis derelictis prope Tripolin Syriæ (Bl!), prope Amrith et Tortosam (Gaill!).

Valde affinē *V. Berytheo*, sed differt foliis non lyratis, indumento parce papilloso et in primis calycis duplo majoris laciinis elliptico-rhombeis mucronatis basi vaide attenuatis fructiferis calycem superantibus.

117. *V. adenophorum* (Boiss. Diagn. Ser. I, 4, p. 64) bienne, canle crassiusculo suhangulato rubello simplici vel ad basin racemi terminalis stricte et breviter ramoso, foliis subintegris tomento brevi stellato lutescente obsitis nervis subtus prominulis inferioribus oblongo-lanceolatis et lanceolatis acutiusculis in petiolum brevem longe attenuatis, superioribus sessilibus strictis anguste linearibus. floribus solitariis longe et laxe racemosis pedicello patenti calyce 3-4-plo longiore tandem apice sursum arcuato suffultis, bracteis longis linearibus setaceo-acuminatis, calyce glanduloso-hirto ($2 \frac{1}{2}$ lin. longo in lacinias anguste linearis-lancolatas partito. corollâ flavâ basi viola-

ceo-maculatâ glabrâ. filamentis violaceo-barbatis, ovario tomentoso capsulâ.... (2).

Hab. in Cariâ (Pinard!).

Bipedale, folia radicalia cum petiolo 4-5-pollicaria vix pollicem lata, racemus florifer 8-9-pollicaris multilorus, pedicelli 7-9 lineas longi; corolla magn. *V. Blattarie.*

118. *V. biscutellæfolium* (Bth. in D. C. Prodr. X, p. 241) bienne, caule erecto tenui inferne tomentello cano superne in racemum simplicem vel ramo anctum elongatum laxum stipitato-glandulosum abeunte, foliis omnibus ad collum et caulis partem inferiorem confertis adpresso et stellatim niveis subrepandis anguste oblongo-spathulatis obtusis basi longe in petiolum attenuatis subpetiolatis, superioribus paucis ad squamas lanceolatas breves reducitis, pedicellis solitariis filiformibus bracteâ linearî calyceque multo longioribus erecto-patulis fructiferis elongatis, calyce ($1 \frac{1}{4}$ lin. longo) papillis glandulosis obsito in lacinias oblongas fere ad basin partito, filamentis (ex. cl. Bth.) purpureo-lanatis, capsulâ ovatâ longe mucronatâ glabrâ calyce duplo longiore (2).

Hab. in Armeniâ (Auch. 2371! et 2435!).

Bipedale, racemus 6-9-pollicaris, folia cum petiolo 2-3-pollicaria 6-8 lineas lata, pedicelli fructiferi 8-10 lineas longi, capsula piso submuor. Ex affinitate *V. phœnicei* et præsertim *V. adenophori*, indumento adpressissimo niveo foliorum insigne.

119. *V. phœnicium* (L. Sp. 254) bienne, radice brevi incrassatâ, caule in racemum simplicem gracilem laxum glandulosum abeunte rarius parce ramoso, foliis superne glabratris subtus brevissime puberulis radicalibus rosulatis ovato vel oblongo-rhombeis in petiolum brevem attenuatis obtusis obtuse crenatis vel repandis, caulinis paucis sessilibus sensim diminutis oblongis et lanceolatis acutis, pedicellis filiformibus solitariis erecto-patentibus calyce et bracteâ minute linearî pluries longioribus, calyce glanduloso (sesquilineam longo) in lacinias ellipticas obtusas partito, corollâ violacea vel rubrâ glabrâ. filamentis purpureo-lanatis, capsulâ ovatâ acutiusculâ calyce sublongiore (2). Ic. Jacq. Austr. tab. 125 — Rchb. Ic. tab. 31. — *V. triste* Sm. Prodr. I, p. 151. — *V. ferrugineum* And. Rep. tab. 162.

Hab. in Macedonia monte Athone (Sibth!), Armeniâ Turcica (C. Koch), ad Gümusch Khané (Bourg!), Tauriâ, Caucaso et Transcaucasiâ (Led., Rupr!), Persiâ boreali (Szovl Auch. 5050! Buhse!).

β. flavidum. — Corolla lutea ant lutescenti-virens. — Forma hybrida inter hanc speciem et *V. nigrum* est *V. rubiginosum* W. K. tab. 197 in Caucaso observatum.

Hab. in Macedonia (Friv!), Armeniâ ad Ortus (Huet!), ad Erzerum (Calv!).

Ar. Geogr. Germania, regio Danubialis, Italia, Dalmatia, Rossia meridionalis, Sibiria Uralensis et Altaica.

120. V. scaposum (Boiss. Diagn. Ser. I. 12. p. 26) glabrum, radice ovato-oblongâ subtuberosa, caule præter 2-3 folia minima linearia auriculato-subhastata nudo scapiformi in racemum simplicem laxum abeunte, foliis radicalibus rosulatis parvis ovatis obtusissimis rotundato-crenatis subpetiolatis, pedunculis solitariis basi bracteâ brevissimâ ovato-triangulari acutâ suffultis calyce 4-6-plo longioribus erecto-patulis crenatis, calyce (sesquilineam longo) ad basin in lacinias oblongas acutas margine parce glandulosas partito, corollâ.... capsulâ ovato-sphæricâ glabrâ mucronatâ calyce duplo longiore ②.

Hab. in pinetis regionis inferioris montis Cassii Syrie borealis ad meridiem pagi Cassab (Boiss!) Fruct. Jun.

Folia radicalia pollicem longa, caulis cum racemo pedalis, pedunculi 7-8 lineas longi, capsula piso minor. Affine *V. phœniceo* a quo distinctum videtur foliis parvis glaberrimis, bracteis triangularibus subauriculatis nec linearibus. Capsula piso minor.

121. V. xanthophæniceum (Griseb. Sp. II, p. 42) bienne, caule præter 1-2 folia minima oblonga denticulata sessilia nudo scapiformi simplici vel 2 3-ramoso ramis in racemos longos graciles laxos abeuntibus, foliis radicalibus 4-5 rosulatis ovatis subsessilibus rugosis grosse obtuse vel acute crenato-dentatis pilis papillaribus longis crispulis obsitis, pedicellis filiformibus erecto-patulis calyce bracteâque setacea 4-6-plo longioribus, calyce glanduloso (lineam longo) in lacinias ellipticas obtusas partito, corollâ luteâ glabrâ, filamentis violaceo-lanatis, capsulâ.... ②. Ic. Buxb. Cent. I, tab. 21.

Hab. ad Zekerakieui juxta torrem Ovidis non procul a Byzantio (Duparq!), in Thraciâ (Griseb!), in Anatoliâ boreali ad Kastamouni (Wied!), in Phrygiâ (Auch. 846!).

Folia 2 1/2-3-polices longa 2 lata, caulis cum racemis sesquipedalis, pedicelli 6-8 lineas longi. A *V. lyratifolio* caule napiformi, foliis subsessilibus indivisis omnibus radicalibus, a *V. phœniceo* flavilloro cui magis affine foliis papillari-hispidis dentatis rugosis sessilibus, caule subnudo scapiformi.

§ 10. **Spinosa.**

122. V. spinosum (L. Sp. 254) suffruticosum ramosissimum humile erinaceum tomento tenuissimo adpresso canescens, ramis intricatis parte inferiori foliosis in paniculas breves abeuntibus, paniculæ ramis brevibus paucifloris rigidis crassiusculis apice demum pedicellisque in spinas tenues induratis, foliis parvis oblongis obtusis in petiolum attenuatis aliis obtuse sinuato-paucilobatis aliis integris, floribus solitariis paucis pedicello calyci subæquilongo demum elongato suffultis, bracteis minimis, calyce tomentello (sesquilineam longo) ad 2/3 in lacinias oblongo-lanceolatas obtusinseculas partito, corolla parvâ luteâ extus tomentellâ, filamentis sordide lutescenti-lanatis,

capsulâ parvâ tomentellâ oblongâ acutiusculâ calyce duplo longiore 3. Ic. Sieb. Reis. tab. 7.

Hab. ad rupes regionis inferioris et montanæ prov. Sphakia Cretæ 0-5000' (Belli, Tourn., Sibth! Raul! Heldr!).

Semipedale, folia pollicem vel paulo amplius longa, capsula piso minor et angustior. Species inclyta facie *Poterii spinosi*.

§ 11. Phlomopsidea.

123. V. Pestalozzae (Boiss. Diagn. Ser. I. 12, p. 25) perenne, nanum totum tomento crasso flavido pannosum collo ramosum, ramis ascendentibus brevissimis crebre foliosis, foliis parvis elevatim reticulato-nervosis oblongis et ellipticis integris longiuscule petiolatis, axi indeterminata. racemis 1-3 ex axillis superioribus pedunculo folia vix superante suffultis patule et longe lanato-hirsutis brevibus densiusculis, floribus solitariis pedicellis calyci æquilongis suffultis, bracteis lanceolato-linearibus pedicello sublongioribus, calyce (4 lineas longo) in lacinias lineares villo connexas ad basin partito, corollâ luteâ tomentosâ, filamentis flavidio-lanatis, capsulâ globosâ tandem glabra calyce breviore ȝ.

Hab. in montibus elatis Lyciae (Pestalozza).

Planta 4-6-pollicaris, foliorum inferiorum lamina pollicaris 4-6 lineas lata, folium in seapo unum alterutrumve sessile oblongo-lineare, racemus fructifer 3-4-pollicaris. Species staturâ nanâ, foliis rugosis fere *Phlomidis*, racemis axillaribus insignis.

Species dubiæ incomplete descriptæ.

V. gossypinum (M. B. Fl. Taur. Cauc. III, p. 153) lanâ niveâ floccosum, foliis oblongis crenulatis supra viridibus sublanatis, caulinis decurrentibus, racemo elongato simplici, fasciculis densis multifloris demuni remotis, filamentis binis superioribus lanatis tribus inferioribus nudis ②.

Hab. in Iberiâ ex M.B. in cuius herbario deesse videtur.

Planta ex cl. auctore suo *V. gnaphalodi* affinis sed ex filamentorum descriptione (de antheris nihil) *Thapsis* adnumeranda, forsitan proles hybrida. *V. gossypinum* C.A. Mey. Enum. p. 111 ex montibus Talysch prope Perimbali ex ejus herbario speciminibus binis constat fructiferis descriptioni Marschalianæ nullo modo quadrantibus, tomentoso-canescensibus, pedalibus, facie *V. campestre* referentibus, sed folia longe decurrentia. Cæterum corolla staminibus deficiens nihil certi de eis dicendum.

V. chryserium (Schrad. Mon. 2, p. 3) biennie, dense flavidio-tomentosum. caule simplici vel superne paniculato, foliis integris inferioribus obovato-lanceolatis obtusis in petiolum attenuatis, superioribus sessilibus cordato-amplexicaulibus, paniculæ ramis erecto-patulis, floribus 4-5-fasciculatis. pedicellis calycem æquantibus, calycis

laciniis oblongo-ovatis acutiusculis, corollâ flavâ, filamentis glabris antheris reniformibus, capsulâ sphærico-ovatâ calyceem superante ②.

Hab. in Palestina prope Arimathæam (Sieber). Non vidi.

Ex cl. auctore facies *V. longifolii*. In loco natali inquirendum et filamentis glabris agnoscendum.

V. oreophilum (C. Koch Linn. XXII, p. 578) pubescenti-asperulum, foliis ovato-oblongis acutis subsimplicibus crenatis, caule paniculato, floribus solitariis pedicello bracteam ovato-acuminatam æquante suffultis. calyce in lacinias ellipticas partito, filamentis violaceo-barbatis quinto sterili, capsulâ ovato-retusâ calyce tertia parte longiore.

Hab. in districtu Artanutsch Armeniæ C. (Koch).

Affine dicitur *V. pyramidato* a quo calyce profundius partito et stamine quinto sterili (an satis?) differt.

V. strictum (Clarke Trav. Ed. II. 2, p. 262) caule fruticoso paniculato elato, foliis inferioribus spathulato-ovatis petiolatis superioribus ovato-oblongis stellatum canescensibus pedicellis calyce longioribus divaricatis, corollâ magnâ flavâ.

Hab. in insula Rhodo (Clarke).

V. diversifolium
V. geminiflorum
V. obovatum } Hochst. in Lorent Reise p. 336.

Hab. prope Nisib Mesopotamiæ (Lorent).

Species sicut fere omnes ejusdem operis ob diagnoses incompletissimas nullò modo recognoscendæ et merito oblivioni tradendæ.

CELSIA (L. Gen. p. 312. — *Janthe* Griseb. Spicil. II, p. 40).

Calyx quinquefidus vel partitus. Corolla explanato-rotata lobis 5 subinæqualibus. Stamina 4, superiore quinto omnino deficiente. Filamenta barbata antheris omnibus reniformibus vel filamentorum binorum anticum longiorum adnato-decurrentibus. Stylus unicus, stigma clavatum. Capsula apice septicide bivalvis bilocularis. Semina lacunoso-rugosa. — Herbae vel suffrutescens. — Genus charactere mere artificiali staminis quinti defientis a *Verbascu* distinctum.

Specierum Orientalium dispositio.

SECT. I. ARCTURUS Bth. in D.C. Prodr. X, p. 244. — *Thapsandra* Griseb. Spic. II, p. 40. — *Staminum* longiorum antheræ adnato-decurrentes.

* Folia opposita.

C. glandulosa, alpina.

* * Folia alterna.

× Flores subsessiles.

C. bugulifolia, Pontica, Cretica.

× × Flores longe pedicellati

C. Arcturus, Dænzeri, speciosa, Boissieri.

SECT. II. NEFFLEA Bth. in D.C. Prodr. X. p. 246. — Antheræ omnes reniformes mediofixæ.

* Folia opposita.

C. parviflora.

* * Folia alterna.

1. Pedicelli calyce 2-pluries longiores.

a. Perennes.

C. acaulis, Cyllenea, pyroliformis, Persica, coronopifolia, Luciliae, Lycia, anceolata.

b. Biennes.

C. Sucarowiana, pinetorum, Cilicia, Coromandeliana, heterophylla, intricata, Aucheri.

c. Annuæ

C. Orientalis, brachysepala

2. Pedicelli calyci subæquilongi.

C. aurea.

SECT. I. ARCTURUS.

1. **C. glandulosa** (Bouché Linn. V. Litter. p. 12) biennis, tota patula et breviter glanduloso-pubescent, caule elato longe folioso in racemum simplicem longum laxum abeunte, foliis omnibus oppositis præter supraem petiolatis lyratis acutiusculæ duplicato-dentatis segmentis utrinque 3-4 ovato-oblongis saepe remotis minutis terminali ovato obtuso, foliis supremis diminutis sessilibus ovatis acuminatis, pedicellis filiformibus longissimis horizontalibus basi bracteâ triangulari-ovatâ acute denticulatâ acuminatâ eis multo breviore suffultis, calyce glanduloso (2 lin. longo) ad $\frac{3}{4}$ in lacinias oblongas acutas partito, corollâ glabrâ flavâ, filamentis violaceo-barbatis posticis apice breviter nudis antheris adnatis, capsulâ ovatâ glabrâ calyce duplo

longiore ②. *C. Arcturus* Jacq. Vindob. II, tab. 107. — *C. Arcturus*
β. oppositifolia F. et M. Ind. Petr. IX, p. 65.

Hab. ad rupes et muros, Lycia ad Elmalu (Bourg!), Pamphylia ad mœnia urbis Adalia (Heldr!), Cyprus ad mœnia Chrysostomo (Ky. 389!) Fl. Mart., Apr.

A *C. Arcturo* egregie distincta indumento, foliis oppositis, segmentis numerosioribus et radice quæ in *C. Arcturo* perennis videtur, in hac manifeste biennis. Planta procerior, caule longius folioso, capsula piso subminor.

2. **C. alpina** (Boiss. et Ky. Diagn. Ser. II. 3, p. 150) perennis, patula hirta viscidula pallide virens, rhizomate sublignoso, caulis fragilibus pumilis simplicibus foliosis in racemum laxum abeuntibus, foliis oppositis inferioribus longiuscule petiolatis ovatis obtusis subduplicatim crenato-dentatis interdum ob segmenta minima ovata sub laminâ obviâ lyratis, superioribus sessilibus ovatis, pedicellis filiformibus horizontalibus vel deflexis calyce 4-5 plo longioribus, bracteis cordatis ovatis acuminatis integris pedicello 2-4 plo brevioribus, calyce glanduloso (sesquilineam longo) ultra medium in lacinias oblongas fisco, corollâ glabrâ, filamentis violaceo-barbatis duobus longioribus superne denudatis antheris adnato-decurrentibus, capsula ... ♀.

Hab. in præruptis vallis Mantchura Antilibani supra Zebdani 5500' (Ky. 114!) Fl. Jun.

Caules cum racemo 6-10-pollicares, foliorum lamina 1-1½-pollicaris, pedicelli 6-8 lineas longi. Foliis oppositis affinis *C. glandulosæ*, differt rhizomate perenni lignoso, foliis parvis, bracteis integris, etc.

3. **C. bugulifolia** (Lam. Dict. IV, p. 226 sub *Verbascœ*) perennis, radice brevi oblongo-cylindricâ, foliis crassiuseculis parvis minute et sparsim puberulis vel glabratris, radicalibus longiuscule petiolatis rosulatis ovato-orbiculatis vel oblongis dentato-crenatis vel repando-subintegris, caulinis diminutis sessilibus oblongo-lanceolatis acutis, caule simplicissimo in racemum demum longum laxissimum glandulosum abeunte, floribus subsessilibus remotis, bracteis linearilanceolatis calycem superantibus, calyce valde glanduloso (3 lineas fere longo) in lacinias ellipticas acutas partito, corollâ majusculâ glabrâ metallicæ cyaneo-viridi intus purpureo-maculatâ, filamentis purpureo-barbatis longiorum antheris longe adnatis posticis brevibus minute antheriferis, capsulâ majusculâ adpresse hirtâ strictâ ovatâ acutiusculâ mucronatâ calycem parum superante ♀. — *C. bugulifolia* Jaub. et Sp. Ill. tab. 407. — *Janthe bugulifolia* Griseb. Spic. II, p. 40. — *Verbascum Osbeckii* L. Sp. p. 255 (ex parte quoad plantam Orientalem.) Ic. Tourn. II, p. 181. — Buxb. cent. 5, tab. 33. — *Trigadera baccata* Dunal in D.C. Prodr. XIII, p. 22.

Hab. in pascuis umbrosis Thraciæ et Bithyniæ circa Byzantium (Tourn! Cast! Auch! Barbeyl) Fl. Maio.

Pedalis et fructifera sesquipedalis, foliorum lamina 1-1½-pollicaris, corolla diam. pollicaris, capsula piso submajor. Planta corollis metallico-nitentibus

margine concentrica luteo-fasciatis elegans. Genus *Janthe* a *Celsia* staminibus posticis minute antheriferis non sat differt.

4. C. Pontica (Boiss. in Tchih. As. Min. II, p. 11) biennis, pa-pillis sœpe crispis hirsuta superne insuper glandulosa, caule tereti elato folioso in racemum tandem longissimum abeunte, foliis tenuiter membranaceis viridibus inferioribus amplis oblongis obtusis cordatis petiolatis late et obtuse crenatis, superioribus valde diminutis ovato-oblongis cordato-amplexicaulibus acutis vel acuminatis, floribus subsessilibus solitariis longe et densiuscule racemosis bracteis a basi triangulari acuminatis calyce longioribus suffultis, calyce valde glanduloso-hirto ($3\frac{1}{2}$ lin. longo) in lacinias ellipticas obtusas profunde partito, corollâ luteâ extus puberulâ tandem glabratâ, filamentis duobus longioribus basi exceptâ glabris antheris longe adnatis, cæteris purpureo-barbatis antheris reniformibus, capsulâ magnâ ovatâ acutiusculâ mucronatâ hirtâ calyce sublongiore ②.

Hab. in Ponto ad meridiem urbis Samsun (Tchih!), Armeniâ Turcicâ (Ky. exs. 993!).

Planta pluripedalis, foliorum inferiorum lamina 6-9-pollicaris, racemus fructifer 1-2 pedalis, capsula piso major. Species præter folia inferiora cordata obtuse crenata nec basi dentato-lobatâ et attenuata, stamna 4 nec 5 et filamenta longiora nuda *Verbasco macrocarpo* simillima et generum nimiam affinitatem demonstrans.

5. C. Cretica (L. fil. Suppl. p. 281) biennis, pubescenti-hirta, caule crasso simplici crebre foliato in racemum densiflorum longe bracteatum glandulosum abeunte, foliis inferioribus petiolatis lyrato-pinnatisectis vel indivisis ovatis vel oblongis sublobatis et dentatis, superioribus sensim diminutis cordato-auriculatis oblongis dentatis acuminatis, floribus subsessilibus, bracteis lanceolatis caudato-acuminatis calyces superantibus, calyce lato (fere 6 lineas longo) in lacinias foliaceas serratas ovato-oblongas inæquilatas partito, corollâ flavâ purpureo-maculatâ amplâ, filamentis violaceo-barbatis antheris binis longe adnatis, capsulâ magnâ globosâ calyce breviore ②. Ic. Jacq. Schænb. tab. 480. — Bot. Mag. tab. 964. *Verb. lyratum* Lam. Dict. IV, p. 223 non huc sed ad *C. Cavanillesii* spectat

Hab. in Cretâ (ex frustulo in d.c. herb. sed a recentioribus ibi non lecta).

Ar. Geogr. Hispania australis, Sardinia, Sicilia, Italia australis.

6. C. Arcturus (L. Sp. pag. 254 sub *Verbasco*) perennis, inferne laxe lanata, caulis e rhizomate indurato basi crebre foliatis cito in racemum longum laxum parce stipitato-glandulosum abeuntibus, foliis omnibus alternis subtus canescentibus vel canis breviter petiolatis lyratis segmentis lateralibus utrinque 1-2 minimis ovatis approximatis terminali maximo oblongo obtuso obtuse late et subduplicatim crenato, petiolis basi subdilatatis, foliis caulinis paucis oblongis minutis semiamplexicaulibus, racemi pedicellis filiformibus longissimis horizontalibus basi bracteâ triangulare acuminatâ basi denticulatâ

reflexâ eis multo breviore suffultis, calyce glabriusculo ($2\frac{1}{2}$ lin. longo) ad $\frac{3}{4}$ in lacinias ellipticas acutas partito, corollâ glabrâ flavâ, filamentis violaceo-barbatis, posticis longis superne nudis antheris longe adnatis, capsulâ glabrâ subglobosâ calyce subtriplo longiore ♀. *C. Arcturus* Murr. Syst. p. 469. — Bot. Mag. tab. 1962. — Jaub. et Sp. Ill. tab. 406. — *C. sublanata* Jacq. Fragm. 79, tab. 126.

Hab. ad rupes et muros regionis inferioris et collinæ Cretæ (Tourn! Sibth! Sieb! Raul! Heldr!). Fl. Mart., Apr.

7. ***C. Dænzeri*** (Bor. et Chaub. Fl. Pelop. N° 980) biennis, inferne parce pilosula superne parce glandulosa, caule gracili in racemum laxum simplicem vel parce ramulosum elongatum abeunte, foliis radicalibus rosulatis petiolatis ambitu oblongo-lanceolatis in segmenta oblongo-lanceolata dentata ab inferioribus ad supremum aucta pinnatisectis vel saepius lyratis segmento terminali ovato-oblongo majore lobato, foliis caulinis alterniis paucis sessilibus triangulari-lanceolatis inciso-dentatis, pedicellis solitariis filiformibus subpatulis calyce pluries longioribus fructiferis vix incrassatis, bracteis brevissimis lanceolatis et linearibus inferioribus basi dentatis, calyce (sesquilineam longo) in lacinias oblongas margine subglandulosas partito, corollâ flavâ glabrâ, filamentis violaceo-barbatis, longioribus apice nudis antheris decurrentibus, capsulâ glabrâ globosâ obtusâ mucronatâ calyce subdupo longiore ②.

Hab. in regione montanâ Peloponnesi, Messenia ad Phygaleam (Dænz!), radices Taygeti supra Mistra (Boiss!) et Anavryti (Heldr!), montes Malevo et Kyttene (Orph!).

Planta 1 $\frac{1}{2}$ -3-pedalis, folia 3-5 pollices longa, pedicelli 12-15 lin. longi, corolla diametro pollicaris, capsula piso vix minor.

8. ***C. speciosa*** (Fenzl in Friedriesth. Reise p. 280) biennis, caule glandulosso-puberulo basi parce ramoso cereum folioso in racemum longum simplicem foliosum abeunte, foliis glabriusculis triangulari-ovatis acutis anguste lobato-dentatis et serratis inferioribus breviter petiolatis, floralibus sensim diminutis sessilibus amplexicaulibus cuspidatis, pedicellis filiformibus erecto-patulis folio axillari duplo longioribus, calyce breviter glanduloso in lacinias anguste lanceolatas patentí-recurvatas inferne 1-2 dentatas partito, corollâ flavâ glabrâ, filamentis violaceo-barbatis longioribus apice nudis antheris decurrentibus, capsulâ.... ②.

Hab. in Arcadiâ circâ Carithenam (Frider. in herb. Vindob!) In eâdem plaga specimen alterum ex Insula Tassos est *Verb. Blattaria*.

Bipedalis, racemos tandem sesquipedalis axi tenui, pedunculi sesquipollitares, corolla e seco magnitudine *V. Blattaria*. Affinis *C. Dænzeri* differt foliis indivisis, bracteis triangularibus nec lanceolato-linearibus, calycis laciniarum formâ.

9. ***C. Boissieri*** (Heldr. et Sart. MSS.) perennis, glabra superne parce glandulosa, rhizomate crasso vestigiis petiolorum vestito saepe

pluricauli, caulis rigidis crassiusculis in racemum longum laxum simplicem abeuntibus, foliis radicalibus petiolatis rosulatis oblongo-lanceolatis brevibus lobato-dentatis vel pinnatifidis lobis brevibus acutè inciso-dentatis, caulinis alternis sessilibus paucis anguste lanceolatis inferioribus pinnatifidis superioribus acute dentatis, pedicellis solitariis calyce 3-4-plo longioribus erecto-patulis fructiferis sub-incrassatis, bracteis minimis integris linear-lanceolatis acuminatis, calyce parce glanduloso (2 lin. circ. longo) in lacinias lanceolatas acutas partito, corollâ flavâ glabré, filamentis violaceo-barbatis longioribus apice denudatis antheris adnatis, capsulâ glabré ovatâ acentius-enâ mucronatâ calyce duplo longiore ȝ. *C. Dænzeri* var. *lyrata* Boiss. et Heldr. Diagn. Ser. II, 3, p. 151.

Hab. in regione mediâ montis Parnes Atticæ (Boiss! Heldr! Sart! Orph! Fl. Mai.

Affinis *C. Dænzeri* pro cuius varietate olim habui sed differre videtur radice perenni, caulis simplicibus rigidioribus, foliis brevioribus minus divisis, præsertim bractearum calycisque characteribus et capsulâ acutiusculâ.

SECT. II. NEFFLEA.

10. *C. parviflora* (Decaisn. Fl. Sinaic. p. 16) perennis, suffruticosa pumila basi opposite ramosa dumulosa, foliis parvis oppositis infimis canescenti-tomentosis oblongo-lanceolatis subrepando-dentatis in petiolum brevem attenuatis cæteris sessilibus linearibus strictis ut et rami glabris fusco-purpurascensibus, racemis oppositis vel alternis brevibus laxe paucifloris, pedicellis solitariis patentibus calyce 2-3-plo longioribus filiformibus, bracteis linearibus brevissimis, calyce (lineam longo) in lacinias ellipticas acutas margine glandulosas parvito, corollâ minimâ fusco-luteâ hirsutâ, filamentis lutescenti-barbatis longioribus apice nudis, antheris reniformibus, capsulâ globosâ ȝ.

Hab. in deserto Sinaitico (Bové!), ad latus septentrionale montis Stæ. Catharinae (Schimp. 232!). — Arabice *Nefflē*.

Suffrutex semipedalis aut paulo elatior ramulis strictis rigidis persistentibus, folia inferiora 6-7 lineas longa 2 lata, racemi triplicares, corolla diam. 2½ lin. lata.

11. *C. acaulis* (Bory et Chaub. Fl. Pelop., p. 40, tab. 18, fig. 2) perennis, minuta, radice crassiusculâ fibrosâ sæpe 2-3-cipiti, foliis parvis omnibus radicalibus et rosulatis oblongis et ellipticis obtusis obtuse dentatis utrinque sed subtus copiosius glandulis sessilibus aspersis in petiolum limbo æquilongum vel longiore attenuatis, racemi radicalis sessilis rachide brevissimâ, pedicellis longe filiformibus basi bracteâ linear-lanceolatâ suffultis, floribus minimis, calyce glanduloso (2 lineas longo) in lacinias lineares acutas basi subangustatas partito, corollâ glandulosâ luteâ, filamentis sordide

luteo et rubello-barbatis, antheris reniformibus, capsula glabra acutâ mucronatâ calyce duplo longiore ♀.

Hab. in regione alpinâ superiori Taygeti ad nives persistentes (Bory! Heldr! Orph!), monte Chelmos supra fontem Stygis 7000' (Heldr!). Fl. Jul.

Facies *Erodii cicutarii*, foliorum lamina 9-15 lineas longa 4-6 lata, pedicelli 1-2-pollicares, corolla diam. 4 lineas lata. Racemi axis aedeo constricta et inter folia occultata ut pedicelli solitarie a collo oriri videantur.

12. C. Cyllenea (Boiss. et Heldr.) perennis, pumila minute glanduloso-puberula, foliis omnibus radicalibus parvis oblongis obtusis obtusiuscule crenato-dentatis petiolo aequilongo vel longiori suffultis, caule brevi paulo supra basin in racemum laxum pauciflorum abeunte, pedicellis glandulosis calyce 2-4-plo longioribus basi bracteâ brevi linearis suffultis, calyce glanduloso (sesquilineam longo) in laciniis lineares acutiusculas partito, corollâ parvâ luteâ glandulosâ, filamentis sordide violaceo-barbatis, antheris reniformibus, capsula.... ♀.

Hab. in regione alpinâ montis Kyllene Peloponnesi 6500' (Heldr!). Fl. Jul.

Foliorum lamina 10-12 lineas longa basi truncata vel vix cuneata, caulis cum racemo 3-5-pollicaris. An varietas magis evoluta *C. acaulis* cui facie et foliis similis est investigandum, praeter racemum elongatum differt foliis in petiolum non attenuatis, pedicellis multo brevioribus.

13. C. pyroliformis (Boiss. et Heldr. Diagn. Ser. I, 42, p. 28) perennis glaberrima, radice verticali, collo incrassato cespitoso, caulis ascendentibus simplicissimis imâ basi parce foliatis superne in racemum laxum abeuntibus, foliis radicalibus confertis carnosulis oblongo-lanceolatis integris subcanaliculatis obtusiusculis subuninerviis in petiolum longe attenuatis. caulinis valde diminutis sessilibus linearis lanceolatis summis minimis squamiformibus, pedicellis solitariis filiformibus calyce 4-5-plo longioribus, bracteis minimis lanceolatis, calyce glabro (lineam longo) in laciniis triangulari-lanceolatas acutas partito, corollâ parvâ luteâ glabra, filamentis sordide albo-lanatis, antheris reniformibus, capsula ovata laciniis calycinis triplo longiore ♀.

Hab. in salsis hyeme inundatis planicie Koniah Lycaoniæ ad radices montis Karadagh (Heldr!). Fl. Jun.

Folia radicalia cum petiolo 2-3-pollicaria 3-6 lineas lata, caules cum racemo 1/2-1-pedales, pedunculi 6-8 lineas longi, corolla 4 lin. diam. lata. Species insignis.

14. C. Persica (C. A. Mey. Enum. p. III) perennis, inferne tomentella partibus floralibus parce glandulosa, foliis inferioribus confertis brevibus breviter petiolatis in laciniis breves anguste lanceolatas pinnatifidas pinnatisectis, caulinis ad caulum partem inferiorem paucis sessilibus lanceolato-linearibus acute incisis, caulis subsimplicibus rigidis in racemum longum laxum abeuntibus, pedicellis rigidis erecto-patulis calyce pluries longioribus, bracteis minimis

squamiformibus linearibus, calyce glanduloso (lineam longo) in lacinias lanceolato-lineares partito. corollâ parvâ glabrâ luteâ, filamentis sordide albo-lanatis longioribus apice denudatis antheris reniformibus, capsulâ glabrâ ovato-pyriformi obtusâ calyce tertiatâ parte longiore ♀.
Scrophularia nudicaulis Wydl. Mon.

Hab. in aridis saxosis regionis montanæ et subalpinæ, ditio Talyesch (C.A. Mey! Hoh!) mons Ssahend prov. Aderbidjan Persicæ (Buhse!), prope Tabris (Bgel), in monte Elvend et prope Ispahan (Oliv., Auch. 2461!).

Planta 1-4½-pedalis racemo fructifero sœpe pedali, folia 1-2-pollices longa 5-6 lineas lata. Corolla diam 6-7 lin. lata.

15. C. coronopifolia (Boiss. et Bal. Diagn. Ser. II, 6, p. 126) perennis glabra. rhizomate indurato vaginis vestito pluricauli, foliis radicalibus congestis angustissime linearispathulatis in petiolum attenuatis in dentes triangulares breves ascendentis acute incisis, caulibus præter 1-2 folia basin versus radicalibus similia nudis simplicibus vel parce ramosis in racemos longos laxos parce glandulosos abeuntibus, pedicellis solitariis tenuiter filiformibus rigidulis erecto-patulis calyce plures longioribus, bracteis minimis linearibus, calyce (lineam longo) ad basin in lacinias linearis-lanceolatas acutiusculas partito, corollâ parvâ luteâ glabrâ, filamentis violaceo-lanatis antheris omnibus reniformibus, capsulâ oblongâ apice subattenuatâ breviter mucronatâ calyce 2 ½-plo longiore ♀.

Hab. in regione montana jugi Mouraddagh Phrygiæ (Bal!).

Pedalis vel sesquipedalis racemo 6-9-pollicari, folia cum petiolo sesqui-pollicaria 2 lineas lata ea *Menonvilleæ* referentia, corolla diam. 6-7 lin. tantum lata, capsula 2-2½ lin. longa. Affinis *C. Persicæ* eximie differt foliis glabris anguste linearibus non bipinnatifidis, filamentis violaceo-lanatis, etc.

16. C. Luciliæ (Boiss. Diagn. Ser. I, 4, p. 65) perennis, pluricaulis pube tomentella pilis glandulosis intermixta cinerascens, caulinis erectis vel ascendenteribus foliosis rigidulis in racemuni simplicem abeuntibus vel paniculatis, foliis alternis inferioribus petiolatis lyratis segmentis obtuse dentatis lateralibus utrinque 2-4 oblongis terminali majore, superioribus sessilibus indivisis oblongo-lanceolatis acutis acute dentatis, racemo brevi laxiusculo glanduloso. pedicellis solitariis patentibus vel deflexis calyce 4-6-plo longioribus, bracteis brevissimis inferioribus lanceolatis denticulatis superioribus linearis-lanceolatis setaceo-acuminatis integris, calyce glanduloso (linea vix longiore) in lacinias ovatas acutiusculas partito, corollâ flavâ glabrâ, filamentis flavidò-barbatis antheris omnibus decurrentibus, capsulâ glabrâ sphærica calyce sublongiore ♀.

Hab. in fissuris rupium circa Gheyla (Aphrodisiam) Cariæ (Boiss!). Fl. Jun.

Planta 1-2-pedalis, folia *Scrophularie* 1½-2½-pollicaria, racemi 3-5-pollicares, pedicelli 7-9 lineas demum longi, corolla diam 6-8 lin. lata, capsula piso dimidio minor; facies *C. heterophyllæ* sed perennis, tomentella, bracteæ dissimiles, corolla duplo major, etc.

17. **C. Lycia.** perennis pumila, rhizomate breviter ramoso vestigiis petiolorum fibroso cespitoso pluricauli, caulis rigidis tenuibus simplicibus vel inferne ramo auctis in racemos laxissimos paucifloros glandulosos cito abeuntibus, foliis ad caulinum basin congestis breviter tomentello-viscidulis parvis oblongo-linearibus obtusis obtusissime crenatis longiuscule petiolatis, caulinis paucis alternis sessilibus linearibus obtusis integris, pedicellis tenuiter filiformibus calyce 3-4-plo longioribus basi bracteâ minimâ linearî suffultis, calyce (sesquilineam longo) glanduloso in lacinias linearî-spathulatas obtusas partito, corollâ parvâ luteâ hirtâ, filamentis lutescenti-lanatis, antheris omnibus reniformibus, capsulâ. . . . 2.

Hab. in regione alpinâ montis Akdaghi Lyciae (Bourgl.).

Semipedalis, rhizoma durum fere suffrutescens, foliorum inferiorum lamina 9-12 lineas longa 2-3 lata, racemi 8-12-flori sub anthesi 3-4-pollicares, corolla diam. 7-8 lin. lata. Species distinctissima ex unico specimine nota.

18. **C. lanceolata** (Vent. Hort. Cels tab. 27) perennis, adpresse et stellatum arachnoideo-cana, rhizomate suffrutescenti pluricipiti, caulis pumilis crassiusculis rigidis paulo supra partem inferiorem foliatam ramosis ramis in racemos elongatos laxos flexuosos duros post anthesim cum pedicellis induratos et persistentes abeuntibus. foliis alternis inferioribus oblongis et oblongo-lanceolatis obtuse incisodentatis vel subsinuato-lobatis, superioribus sessilibus linearî-lanceolatis crenatis et integris, pedicellis solitariis calyce 2-3 plo longioribus horizontalibus vel deflexis bracteâ linearî minimâ suffultis, calyce tomentoso (2 lin. fere longo) in lacinias ellipticas obtusas partito, corollâ luteâ parvâ extus tomentellâ, filamentis sordide rubello-barbatis, antheris reniformibus, capsulâ tomentosâ glabrâ calycem non superante 2. *C. incana* Bth. in p.c. Prodr. X, p. 247.

Hab. in Mesopotamiâ (Auch!), in deserto flavii Chabur (Haussk!), inter Mossul et Bagdad (Oliv!), Kurdistanâ Persicâ ad Mendeli (Noë 1165!), inter Marand et Djulf (Bge!).

Semipedalis, fructifera fere pedalis, folia 1 $\frac{1}{2}$ -2-poll. longa 5-8 lineas lata, pedicelli 3-4 lineas longi demum spinescentes. Icon Horti Cels ex specimine culto elongato habitum speciei pessime exhibit.

19. **C. Suwarowiana** (C. Koch Linn. XVII, p. 264) biennis, tomentella superne glandulosa, caule crassiusculo rigido inferne plurifoliato in racemum longum abeunte, foliis inferioribus oblongis in petiolum latum attenuatis grosse et argute serratis nonnullis lyratopinnatisectis, caulinis sessilibus anguste lanceolatis acuminatis, pedicellis solitariis calyce pluries longioribus demum patentissimis, bracteis triangulari-lanceolatis valde acuminatis pedicello 2-3 plo brevioribus, calyce glandulis sessilibus obsito (2 lineas longo) in lacinias ovato-ellipticas partito, corollâ glabrâ flavâ, filamentis violaceo-lanatis longioribus apice denudatis, antheris reniformibus, capsulâ subglobosâ ②. *C. collina* Bth. in p.c. Prodr. X, p. 247.

Hab. in calcareis salsuginosis prov. Eriwan Armeniæ Rossicæ (C. Koch!), prov. Aderbidjan Persiae (Szov! Auch. 5046!), inter Teheran et Ispahan (Bge!).

Pedalis, folia radicalia cum petiolo 3-4-pollicaria 7-12 lineas lata, differt a *C. Persicæ* præter formam foliorum et radicem quæ biennis videtur bracteis multo majoribus.

20. *C. pinetorum* (Boiss. Diagn. Ser. I, 12, p. 30) biennis, inferne cum foliis papillari-hispida, caule gracili tereti in paniculam patule glandulosam abeunte, foliis alternis inferioribus petiolatis pinnatisecto-lyratis segmentis lateralibus paucis minimis triangularibus terminali ovato oblongo acute inciso-lobato et dentato, foliis superioribus indivisis subsessilibus summis amplexicauli-cordatis triangulari-ovatis acutissime dentatis acuminatis, panienlæ racemis longis laxis tenuibus, pedicellis solitariis capillaribus patentibus calyce pluries longioribus, bracteis pedicello multo brevioribus integris a basi ovatâ cuspidatis, calyce subglanduloso (sesquilineam longo) in laciniis elliptico-lanceolatas acutas partito, corollâ luteâ glabrâ, filamentis violaceo-barbatis longioribus apice nudis, antheris reniformibus, capsulâ globosâ glabrâ calyce sesquiloniore ②.

Hab. in pinetis vallium regionis inferioris Cassii Syriæ borealis inter Cassab et Suadieh (Boiss!). Fl. Jun.

Bipedalis, folia radicalia 2 $\frac{1}{2}$ -pollicaria, pedicelli 8-11 lin. longi, corolla diam. fere pollicaris, capsula piso minor. Species elegans facie *C. Arcturi* et *C. Denzieri* a quibus præter alias notas antheris reniformibus differt.

21. *C. Cilicia* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 30) biennis, inferne patule papilloso-hispida, caule paucifolio divaricatim ramosissimo, foliis radicalibus rosulatis breviter petiolatis ovatis obtusis acutiusculæ dentatis, caulinis paucis parvis sessilibus suborbiculatis, racemis longis tenuibus laxissimis, pedicellis filiformibus patulis calyce pluries longioribus, bracteis vix perspicuis ovatis squamiformibus, calyce glanduloso (lineâ vix longiore) in laciniis ovatas acutiusculas partito, corollâ luteâ glabrâ, filamentis violaceo-lanatis apice nudis, antheris reniformibus, capsulâ ovato-globosâ calyce sesquiloniore ②.

Hab. in Ciliciâ ad radices montis Bulghardagh prope diversorium Tchiftikhan Sept. jam deflorata (Heldr!).

Ex speciminibus imperfectis descripta, affinis *C. pinetorum* a quâ differt foliis indivisis, bracteis ad squamulas vix perspicuas reductis, calycis laciniis latioribus. An tamen hujus varietas?

22. *C. Coromandeliana* (Vahl Symb. 3, p. 79) biennis, pubescenti viscidutâ tomentella, caule paniculato-ramoso, foliis alternis inferioribus petiolatis lyratis segmentis denticulatis lateralibus parvis lanceolatis terminali multo majore oblongo, foliis caulinis sessilibus oblongis vel ovatis dentatis acuminatis superioribus subauriculatis, racemis longis laxiusculis, pedicellis solitariis calyce subdupo longioribus, bracteis pedicello multo brevioribus inferioribus triangulari-

ovatis acuminatis superioribus lanceolatis, calyce glanduloso lineam longo in lacinias ovatas partito, corollâ parvâ luteâ glabrâ, filamentis sordide albo-barbatis antheris reniformibus, capsulâ globosâ calyce sublongiore ②. *C. viscosa* Roth Cat. 2, p. 69. — Hook. Journ. Bot. I. tab. 129.

Hab. in regno Cabulico infra Dair Hadj (Griff. 564!).

Species aliæ distinctæ sub eodem nomine ex Indiâ latere videntur.

Ar. Geogr. India Orientalis Cis- et Transgangetica.

23. *C. heterophylla* (Desf. in Pers. Syn. II. p. 161) biennis, præter caules interdum basi parce hispidulos glaberrinia, caule elato ramoso in racemos elongatos vel paniculas ramosas abeunte, foliis alternis inferioribus pinnatisectis petiolatis segmentis oblongis crenatis remotis lobulis saepe intermixtis terminali submajore oblongo lanceolato, foliis intermediis subsessilibus segmentis et lobis angustioribus acutis, superioribus semiamplexicaulibus ovatis acuminatis integris, pedicellis solitariis tenuiter filiformibus calyce 4-5-plo longioribus tandem horizontalibus, bracteis deitoideo-cordatis indivisis valde acuminatis integris inferioribus pedicellos aequantibus superioribus eis multo brevioribus, calyce glaberrimo (lineam longo) in lacinias oblongas aetas partito, corollâ minimâ luteâ glabrâ, filamentis flavo-barbatis antheris omnibus reniformibus, capsulâ parvâ globosâ glabrâ calyce tertiatâ parte longiore ②. Ic. Jaub. et Sp. III. tab. 404. — *C. chrysurus* Fenzl MSS. — *C. agrimonialfolia* G. Koch Linn. 22, p. 732!

Hab. ad aquas a regione inferiori ad montanam et subalpinam, Galilea ad lacum Phiatla prope Banias (Boiss!), Antilibanus et Libani radices orientales (Ky! Gaill!), Syria bor. ad Aintab et Marasch (Auch. 2433! Ky!), Mesopotamia et Assyria (Ky! Haussk!), Armenia australis (Noë!), Kurdistania Persica in montibus Avroman et Schahu (Haussk!), Persia bor. in monte Elbrus (Ky. 714!), in prov. Ghilan (Auch. 5045!), ditio Talysh (Hohen!).

Planta 2-4-pedalis, racemi tenuissimi demum saepe pedales et longiores, corolla e minimis 3-4 lineas diam. lata, capsula piso duplo minor.

24. *C. intricata* (Bth. in p.c. Prodr. X, p. 247) biennis, caule fere a basi glandulis sessilibus asperso divaricatim et dichotome ramosissimo aphylo, ramis tenuiter virgatis glabris longis intricatis in racemos laxissimos paucifloros subflexuosos abenentibus, foliis omnibus radicalibus parvis rosulatis oblongis vel ovato-orbiculatis crenato-dentatis breviter petiolatis, bracteis ad dichotomias et ad pedicellorum basin minimis squamiformibus, pedicellis filiformibus patulis calyce 3-5 plo longioribus capsulâ delapsâ cum ramis persistentibus, calyce glandulis sessilibus obsito ($1\frac{1}{4}$ lin. longo) in lacinias ellipticas aetas basi subangustatas partito, corollâ minimâ luteâ, filamentis albo-barbatis longioribus superne nudis, antheris reniformibus, capsulâ minimâ globosâ glabrâ calyce duplo longiore ②.

Hab. in Persiae monte Sofi, prope Ispahan (Auch. 2463! et 5050!, Derderian!), prope Schurab inter Ispahan et Teheran (Haussk!), inter Yezd et Ispahan (Bge!).

Pedalis, a basi intricatim ramosissima ramis et ramulis tenuibus, folia radi-calia 6-9 lineas longa petiolo brevi, corolla 3 lineas diam. lata, capsula magn. seminis *Cannabeos*.

25. C. Aucheri (Boiss. Diagn Ser. I 12, p. 29) biennis, caulinibus fere a parte inferiore tomentellâ ramosis ramis in racemos laxissimos longos abeuntibus, foliis cano-tomentosis radicalibus petiolatis ovatis vel oblongis crenato-dentatis vel lyratis laciniis laterilibus utriuque 1-2 minimis ovatis, foliis superioribus et rameis ovatis acutis denticulatis sessilibus saepe amplexicauli-cordatis, pedicellis tenuiter filiformibus calyce pluries longioribus patentibus, bracteis minimis triangularibus acutis, calyce crebre glanduloso-papillari ($1\frac{1}{4}$ lin. longo) in laciniis oblongas acutas partito, corollâ parvâ luteâ extus hirsutâ, filamentis sordide albo-lanatis longioribus apice nudis, antheris reniformibus, capsulâ ovatâ mucronatâ calyce tertiatâ parte longiore ②. *C. intricata* var? *macrocarpa* Bth. in D.C. Prodr.

Hab. in Persia prope Teheran (Auch. 5043!), prope Schurab (Haussk! Bge!), ad radices montis Elbrus prope Derbend (Ky. 581!), ad Hous i Sultan (Bge!).

Sesquipedalis, folia sesquipollicaria, pedicelli pollicem et amplius longi. A *C. intricata* distinctissima caulinibus et ramis inferne foliatis, capsulâ triplo majore pisum æquante, etc

26. C. Orientalis (L. Sp. p. 866) annua, minute tomentella viridis, caule crebre folioso gracili simplici vel superne parce paniculato-ramoso, foliis ambitu ovato-oblongis in laciniis divergentes bipinnatisectis inferioribus petiolatis segmentis oblongis superioribus sessilibus laciniis anguste linearibus, floribus pedicello calyce breviori suffultis longissime et laxe spicatis, bracteis inferioribus pinnatisectis cæteris oblongo-lanceolatis acutis calyce longioribus, calyce tomentello ($2\frac{1}{2}$ lin. longo) in laciniis lanceolatas acutas fisso, corollâ luteâ glabra, filamentis flavidio-barbatis antheris reniformibus, capsulâ subsessili strictâ ovatâ calycis laciniis subbreviore ③. Ic. Hort. Ups. tab. 2. — Fl. Græc. tab. 605. — Rehb. Ic. tab. 50.

Hab. in collibus regionis inferioris, Peloponnesus in Argolide (Sibth! Sprun!), Macedonia et Bithynia (Griseb! Friv!), Paphlagonia et Galatia (Wied!), Lycia (Bourg!), Pamphyliâ (Heldr!), Syria maritima usque ad regionem montanam Libani (Bl! Gaill!), Palestinâ (Boiss!), Mesopotamiâ (Haussk!), Kurdistaniâ Persicâ (Noë 1199!), Tauriâ (M.B!), Transcaucasiâ (Hoh! Rupr!).

Ar. Geogr. Dalmatia.

27. C. brachysepala (F. et Trautv. Ind. Petrop. 10, p. 33) annua, minutissime tomentella, caule gracili crebre folioso simplici vel superne parce paniculato-ramoso, foliis inferioribus petiolatis ovatis obtuse inciso-lobatis vel sublyratis, caulinis subsessilibus et sessilibus in laciniis lineares subdentatas pinnatisectis, floribus pedi-

cellis solitariis calyce triplo longioribus filiformibus suffultis longe et laxe racemosis, bracteis lanceolatis pedicello 2-3-plo brevioribus, calyce glanduloso (2 lin. longo) in lacinias lanceolatas partito, corollâ luteâ glabrâ, filamentis luteo-barbatis, antheris reniformibus, capsulâ pedicello erecto-patulo triplo breviore suffultâ ovatâ calyce tertiatâ parte longiore ♂. *C. lepturus* Schott et Ky. Est. Woch. IV.

Hab. in saxosis regionis inferioris, Anatolia loco non notato (Fisch!), Lycia ad Elmalu (Bourgt!), Pisidia prope Tcheltickchi (Heldr!), Cilicia ad Gülek Boghaz (Ky. 79! Balt.).

Affinis *C. Orientali* a quâ foliis minus dissectis et pedicellis elongatis egregie differt, capsula minor pisum dimidium æquans.

28. *C. aurea* (C. Koch Linn. XXII, p. 583) biennis, caule elato angulato superne in paniculam amplam elongatam sœpe compositam pilis minimis aureis stellatis sparsis pulverulentam abente, foliis viridibus subtus pube stellatâ minimâ lutescente adspersis inferioribus amplis acutiuscule crenato-dentatis elongato-oblongis acutiusculis vel acuminatis in petiolum brevem latum subalatum attenuatis subtus elevatis reticulatis, superioribus parvis sessilibus lanceolatis acutis subintegris, floribus solitariis numerosissimis parvis pedicello calyce sublongiore et bracteâ lanceolatâ pedicello breviore suffultis secus ramulos tennes laxiuscule racemosis, calyce aureo (vix 2 lineas longo) angulato ad medium in lacinias ovalas fisso, corollâ citrinâ extus aureo-tomentellâ, filamentis flavo-barbatis antheris omnium reniformibus, capsulâ pedicello tenui stricto suffultâ oblongâ tomentosâ calyce tertiatâ parte longiore ♀. — *V. Armeniacum* Boiss. in Huet exs.

Hab. in subhumidis regionis montanæ et alpinæ Armeniae Turcicæ 4-600! (C. Koch!), ad Erzerum (Huet!), Baibut (Bourg!), circa montem Bingöldagh (Tchih! Ky. 324!). Fl. Aug.

β. *buddleiaefolia*. — Folia multo minora 3-4-pollicaria subtus valde reticulata, panicula simplex anguste thyroidea.

Hab. in monte Jyldisdagh Anatoliæ borealis (Wiedem!).

Typus pluripedalis foliis radicalibus 1-1½-pedalibus fere *Cochleariae Armoracice*, paniculæ pedalis et longioris multifloræ rami virgati longi, corolla diam. pollicaris, capsula 2½ lin. longa. Facies omnino *V. pyramidati* et *V. nigri*.

STAUROPHRAGMA (Fisch. et Mey. Ind. IX, Petrop. p. 90).

Calyx quinquepartitus. Corolla explanato-rotata lobis subinæqualibus. Stamina 4 filamentis barbatis inferioribus longioribus, antheris omnibus reniformibus. Stylus simplex apice in stigma incrassatus. Capsula cylindrica subindehiscens valvis margine involutis subquadriangularis, placentis 4 distinctis marginalibus. Semina lacunosa-rugosa. — Herba facie *Verbasci* et *Celsiae*, ab eis non formâ externâ

capsulæ in eorum generum speciebus pluribus sœpe obviâ, sed loculis 4 nec 2 distincta.

1. S. Natolicum (F. et M. loc. cit.) bienne, caule crassiusculo tereti rubello inferne tomentello mox in paniculam longam parce glandulosam strictam abeunte, foliis radicalibus et inferioribus congestis denge tomentosis flavidis oblongo-lanceolatis acutiusculis integris basi longe in petiolum attenuatis subtns elevatim nervosis, caulinis paucis parvis lanceolatis strictis, pedicellis calyci æquiforis demum incrassatis secus paniculæ ramos virgatos strictis longe et laxe racemosis, calyce glanduloso (sesquilineam longo) in laciniæ lanceolato-lineares partito, corollâ majusculâ luteâ glabrâ. filamentis sordide rubello-barbatis. capsulâ tenuiter cylindricâ obtusâ mucronatâ glabrâ calyce quintuplo longiore ♂.

Hab. in Anatoliâ loco non notato (Fisch!), in collibus aridis supra pagum Talasse Cappadociae 4200' (Bal!). Fl. Junio.

Bi vel tripedale, folia radicalia numerosa 4-6-pollicaria 6-12 lineas lata, corolla diametro 15-16 lineas lata, racemi fructiferi 1-1½-pedales, capsula 6-7 lineas longa diam. sesquilineam lata.

TR. II. ANTIRRINEÆ. (Chav. Mon.) — Corolla tubulosa vel campanulata tubo basi sœpe saccato vel calcarato. Capsula poris dehiscens. Inflorescentia centripetâ uniformis. — Folia saltem inferiora opposita vel verticillata.

ANARRHINUM (Desf. Atl. II, p. 54).

Calyx profunde quinquesfidus. Corolla tubulosa bilabiata tubo basi breviter calcarato vel (in nostris) ecalcarato, fance æquali perviâ, labii superioris laciniis erectis dein reflexis, inferioris patentibus. Antheræ reniformes loculis confluentibus. Capsula globosa loculis æqualibus ad dorsum poro oblongo univalvulato dehiscentibus. Semina ovoidea aptera tuberculata vel muricata. — Herbæ biennes vel perennes erectæ.

1. A. Orientale (Bth. in D.C. Prodr. X, p. 289) glabrum, ex rhizomate demum indurato multicaule. caulis erectis elatis virgatis inferne crebre foliosis, spicis terminalibus longis multifloris laxiusculis simplicibus vel paniculatis, foliis infimis petiolatis obovato-oblongis integris vel dentatis, caulinis sessilibus lanceolatis vel linearibus strictis indivisis et triseptis acutis, bracteis linearibus inferioribus flores æquantibus, calycis pedicello longioris laciniis lanceolatis glabris, corollæ ecalcaratae labio superiore breviore fere bipartito, capsulâ retusa glabrâ ♀. Ic. Jaub. et Sp. III. tab. 450. — *Cardiotheca virgata* Ehrenb. MSS. — *Anarrhinum ecalcaratum* Fenzl MSS.

Hab. in rupes tribus siccis Arabiae petreæ (Bové ex Bth.), Palestine ad Hierosolymam (Roth), Syriæ in regione inferiori et mediâ, Libani, Antilibani et ad Danas cum (Ehr! Boiss! Gaill! Bl!), Mesopotamia ad Orfa, Mardin (Ky! Haussk!), Tauri Cilicieci (Ky. 394! Bal!), Cappadociae (Bal! Haussk!), Armeniae Turcicæ australi, (Tchih! C. Koch!), Anatoliæ ad Tokat, Amasia, Toptscha (Wied!).

Planta bipedalis a Billardiero et aliis cum *A. bellidifolio* corollâ calcaratâ alieno confusa. *A. bellidifolium* in Græciâ ex schedulâ mendaci in Bory Fl Pelop. erroneâ indicatur. Planta Abyssinica in d c. Prodri. *A. Orientali* adnumerata est *A. Abyssinicum* J. et Sp. Ill. tab. 447-447 ex claris auctoribus labio superiore bilobo nec bipartito distinctum, *Anarrh. fruticosum* Desf. Atl. tandem ab hoc et specie sequenti preter folia indivisa labio superiore æquilongo et staminodio elongato capillari differt.

2. *A. pubescens* (Fresen. Mus. Senckenb. p. 89) patule præsertim inferne glandulosohirtum, e rhizomate denum indurato pluricaule caulibus erectis elatis virgatis parce foliatis, spicis terminalibus longis multifloris laxinsealnis simplicibus vel paniculatis, foliis infimis cuneato-spathulatis in petiolum attenuatis apice dentatis vel lobatis, caulinis linearibus indivisis et trifidis, bracteis calycem vix excedentibus, pedicellis brevissimis, calycis laciniis lanceolatis glandulosociliatis, corollæ exalcaratæ labio superiore breviore fere bipartito, capsulâ retusâ glabrâ ♀. Ic. J. et Sp. Ill. tab. 451. — *Cardiotheca pubescens* Ehr. MSS.

Hab. in rupibus ad radices montis Sinai (Ehr! Bové! Schimp. 354!) Arabice *Arphaije*.

Affine præcedenti differt indumento, foliis remotis, floribus minoribus, calyce ciliato.

LINARIA (Juss. Gen. p. 420).

Calyx quinquepartitus laciniis subæqualibus. Corolla personata tubo basi exalcarato, palato prominenti nunc faucem claudente, nunc depresso lancee perviâ, labio superiore erecto bifido inferiore reflexo trilobo. Stamina 4 didynama, antherarum loculi oblongi Capsula globosa vel ovata bilocularis loculis æqualibus vel parum inaequalibus utroque vel altero tantum apice varie dehiscentibus vel ruptilibus. Semina ovata aut prismatica testâ varie exculptâ, vel discoidea membranaceo-marginata. — Herbæ vel suffrutices.

Specierum Orientalium distributio.

SECT. I. CYMBALARIA Chav. Mon. p. 98. — Corollæ faux palato clausa, capsulæ loculi valvulis tribus dehiscentes, semina oblonga rugosa. Folia palminervia.

L. microcalyx, longipes.

SECT. II. ELATINOIDES Chav. l. cit. p. 103. — Corollæ faux palato clausa, capsulæ loculi operculis binis circumscisis vel valvæformibus dehiscentes. Semina ovata vel oblonga. Folia penninervia.

* Annuæ.

L. floribunda, Acerbiana, spuria, lanigera, Elatine, Græca, cirrhosa.

** Perennæ.

L. ramosissima, Cabulica, ovata, Aegyptiaca, macilenta.

SECT. III. LINARIASTRUM Chav. loc. cit. p. 114. — Corollæ faux palato clausa, capsulæ loculi valvulis dentiformibus 4-6 ab apice dehiscentes. Folia penninervia.

§ 1. **Discoideæ.** — Semina discoidea plana membranaceo-marginata.

* Perennæ.

L. pyramidata, Griffithii, confertiflora, fastigiata, Kurdica, Aucheri, pycnophylla, vulgaris, macroura, odora, Damascena, Nurensis, Michauxii.

** Annuæ.

L. Pelisseriana, arvensis, micrantha.

§ 2. **Oblongæ.** — Semina oblonga reniformia vel angulato — triquetra non marginata.

* Perennæ.

L. Dalmatica, Macedonica, præalta, genistifolia, Sibthorpiana, lineolata, coridifolia, Iconia, purpurea.

** Annuæ.

L. triphylla, reflexa, Chalepensis, Armeniaca, Hælava, Ascalonica, albitrons.

SECT. IV. CHÆNORRHINUM D.C. — Corollæ faux palato depresso pervia, capsula dentibus dehiscens vel irregulariter rupta. Folia penninervia.

* Capsulæ loculi inæquales, inferior indehiscens, superior valvula dehiscens.

L. rubrifolia.

** Capsulæ loculi inæquales, uterque apice valvulis tribus dehiscens.

L. cryptarum, minor, pterospora.

*** Capsula indehiscens irregulariter ruptilis loculis æqualibus.

L. Persica.

SECT. I. CMBALARIA.

1. L. microcalyx (Boiss. Diagn. Ser. I. 4, p. 72) tenerrima lanâ longâ albâ crispâ villosa, ramis diffusis filiformibus valde intricatis, foliis longe petiolatis parvis orbiculatis infimis integris cæteris grosse et obfuscissime trilobis raro quinquelobis, pedunculis folia superantibus, calycis glabriusculi laciniis minutissimis ovatis obtusis, corollâ glaberrimâ pallide violaceâ palato aurantiaco. calcare recto corollâ dimidio breviore ♀ *Ant. pilosum* Fl. Pelop. p. 39 non L.

Hab. in fissuris rupium regionis montane inferioris et alpinæ, Taygetus supra Mistra (Boiss!), mons Malevo Laconiae prope Platanos 350' (Orph. Fl. exs. 712!), mons Olenos 5500'-6000' (Heldr!). Fl. Apr., Jul.

β. acutiloba Boiss. Diagn. Ser. I, 12, p. 40. — Foliorum præsentim superiorum lobi acutiores.

Hab. ad rupes promontorii Alaya Pamphyliæ (Heldr!).

Affinis *L. pilosæ* D.C. indumento saepius breviori, foliorum lobis mucronatis numerosioribus, calycis pilosi laciniis lanceolatis ($\frac{1}{2}$ lin. longis) duplo longioribus, corollâ minore puberulâ distinctæ.

2. L. longipes (Beiss. et Heldr. Diagn. Ser. I, 12, p. 40) glaberrima, ramis feneris elongatis filiformibus diffusis intricatis, foliis longissime petiolatis orbiculatis infimis integris cæteris late et obtusissime 3-5-lobatis, pedunculis longissimis, calycis minutissimi laciniis oblongis obtusis, corollâ pallide violaceâ striatâ palalo flavo, calcare recto corollâ subdimidio breviore ♀.

Hab. ad muros et rupes regionis inferioris, insulæ Chios (Auch. 1905!), in rupibus maritimis insulæ Salamis (Heldr!) et Rhodi (Bourg!), Cretæ (Sieb!), ad Adalia Pamphyliæ (Heldr!).

Affinis *L. cymbalarie* differt foliorum lobis minus profundis obtusissimis et calycis laciniis oblongis obtusis lineaæ tertiam partem longis nec lanceolatis acutis fere lineaem longis. Corolla eâ *L. cymbalarie* paulo major. Calyce parvo brevi affinis *L. microcalyci*, sed glabra et folia majora.

SECT. II. ELATINOIDES.

* Annuæ.

3. L. floribunda (Boiss. Diagn. Ser. I. 12, p. 40) patule hirsuta cinerascens, rhizomate indurato ramoso, caulinibus prostratis basi denudatis dein crebre foliatis in spicas foliosas densas cylindricas elongatas abeuntibus, foliis parvis in petiolum brevem attenuatis inferioribus elliptico-rhombeis integris vel interdum medium versus denticulo auctis, floralibus linearibus flores superantibus, pedicellis capillaribus calyce brevioribus, calycis villosi laciniis anguste lanceolatis acuminatissimis herbaceis, corollâ hispidulâ flavâ calyce duplo longiore, calcare tenui valde uncato-recurvo corollæ subæquilongo,

capsulâ parvâ sphæricâ glabrâ calyce subbreviore, seminibus paucis ovatis tuberculatis ♀.

Hab. in agris derelictis regionis Amalecitarum ad meridiem urbis Gaza (Boiss!), in deserto Judæa (Roth 578!).

Spicæ sœpe semipedales, folia 5-6 lineas longa 1-2 lata, corolla cum calcare 7 lineas longa, capsula (an normaliter?) Cannabeos semine non major. Affinis *L. scariosæ* Desf. foliis majoribus et latioribus, spicâ laxiore, calycis laciniis scariosis ovato-oblongis, calcare subarcuato corollâ breviore, seminibus foveolatis, etc. distinctæ.

4. L. Acerbiana, patule pubescenti-viscidula, annua vel indurata perennans, a collo multicaulis caulinis ramosis ramis tenuiter virgatis demum valde elongatis, foliis integrerrimis inferioribus longe petiolatis majusculis ovatis obtusis subcordatis, floralibus valde diminutis ovatis mucronulatis petiolo limbo breviore suffultis, floribus axillaribus laxissime et longe spicatis pedicellis petiolo brevioribus suffultis, calycis hirsuti laciniis anguste lanceolatis acutissimis, corollæ flavæ velutinae calcare gracili acuto subincurvo, capsulâ parvâ globosâ glabrâ calyce breviore, seminibus minutissimis ovatis glanduloso-tuberculatis ♂. *L. alsinefolia* Bth. in p.c. Prodr. 10, p. 269 (ex parte quoad pl. Egyptiacam non Vivian. Fl. Corsica nec Spreng.)

Hab. in deserto Egypti ad mare Rulrum (Acerbi!), inter Kosseir et Ras Benass (Schweinf!).

Rami demum valde elongati decumbentes 2-3-pedales, folia inferiora pollicem floralia 2-1 lineas longa, capsula magn. sem. Cannabeos. Planta Corsica ab auctoribus pluribus cum hac nostrâ confusa longe differt indumento villoso, pedicellis folio longioribus, etc., et ex cl. Moris Fl. Sard. sectioni *Chænorhino* adnumeranda.

5. L. spuria (L. Sp. 851 sub *Antirrhino*) villosula, a basi ramosa ramis prostratis elongatis subsimplicibus, foliis alternis breviter petiolatis orbiculari-ovatis obtusis subintegris vel infinitis subdentatis, floralibus diminutis mucronulatis, pedicellis axillaribus villosis filiformibus patentibus folio axillari 2-3-plo longioribus, calycis laciniis ovato-oblongis acutis basi subcordatis, corollæ flavæ labio superiore violascenti, calcare incurvo acuto corolle æquilongo, capsulâ glabrâ, seminibus crebre gyroso-alveolatis ♂. *L. spuria* Mill. Dict. 15. — Ic. Fl. Dan. tab. 913. — Rehb. tab. 59.

Hab. in cultis Graeciæ et insularum (Marg! Heldr. herb. Norm. 656! etc.), Anatolia borealis (Wied!). Bithyniæ (Forsk!), Tauriæ et prov. Caucasicarum (M.B., Ledeb.). Persie borealis in prov. Ghilan (Haussk!).

Ar. Geogr. Europa media et australis ab Angliâ et Germaniâ ad Rossiam mediam, Africa borealis, Insulae Canarienses.

6 L. lanigera (Desf. Atl. II, p. 38, tab. 130) lanato-villosa, multicaulis caulinis prostratis longissimis superne breviter et patule ramulosis, foliis alternis præter inferiora breviter petiolata subsessilibus ovatis superioribus sensim diminutis orbiculari-cordatis acutis,

floribus in ramulo axillaribus paucis paniculam longam angustam foliosam formantibus, pedicellis pilosis calyci æquilongis vel vix longioribus, calycis laciniis lanceolatis acutis, corollæ lutescenti-violetaceæ villosæ tubo tenui incurvo corollâ vix breviore, capsulâ globosâ glabrâ, seminibus minimis reticulato-loveolatis ②.

Hab. in cultis Ciliciae littoralis ad Mersina (Bal!), Syriae littoralis prope Berythum et Tripolim (Bl!).

Specimina Hispanicis et boreali-Africanis similia. Species indumento et facie *L. Elatines villosæ* a quâ foliis floralibus non triangularibus nec hastatis differt; a *L. spuria* distinguuntur foliis inferioribus minoribus, pedicellis brevibus, floribus et seminibus dimidio minoribus; ab utrâque inflorescentiâ paniculatâ. An tamen *L. spuriae* varietas vel forma autumnalis?

Ar. Geogr. Lusitania, Hispania, Africa borealis, Madera.

7. *L. Elatine* (L. Sp. 821 sub *Antirrhino*) pilosa, a basi ramosa caulis et rami prostratis tenuibus elongatis, foliis præter infima opposita alterris breviter petiolatis ovatis, superioribus sensim diminutis hastatis acutis, pedicellis axillaribus setaceis patentibus glabris folio axillari multo longioribus, calycis laciniis lanceolatis acutissimis, corollâ albida labio superiore violascente inferiore lutescente, calcare subulato subincurvo corollæ subæquilongo, capsulâ globosâ, seminibus alveolato-reticulatis ③. *L. Elatine* Mill. Dict. № 16. — Ic. Fl. Dan. tab. 426. — *L. Caucasica* Muss. in Spreng. Syst. II, p. 790.

Hab. in cultis regionum septentrionalium ditionis, Thracia (Friv! Pichler!), Anatolia borealis Tauria et prov. Caucasicæ (Ledeb!) Persia borealis (Szov!).

$\beta.$ *villosa*. — Dense villosa sepe bombycina, folia floralia ut in typo sed saepè minus manifeste hastata, pedicelli glabri vel hirsuti saepè minus elongati. — *L. bombycina* Boiss. et Bl. Diagn. Ser. II, 3, p. 101. — *L. Prestandreae* Tineo in Guss. Syn. II, p. 842. — *L. Sieberi* Rchb. Fl. exc. p. 374. — Ic. Guss. Fl. Ischia, tab. 9, fig. 1.

Hab. in cultis ditionis calidioris frequentissima a Græciâ et ejus insulis (Heldr! Orph!), Cretâ (Sieb! Heldr!), Anatoliâ australi, ad Syriam (Bl! Gaill!), Mesopotamiam (Ky!), Babyloniam et Persiam australem (Haussk!), Persiam borealem (Szov!), Arabiam Mascatensem (Auch. 5066!), Egyptum inferiorem (Del.) et in Oasi magnâ (Schweinf!).

Varietas β ad typum intermediis manifeste transit.

Ar. Geogr. Europa media et australis ab Angliâ et Gothiâ ad regionem Danubialem, Africa borealis, Abyssinia, Madera, Insulæ Canarienses.

8. *L. Græca* (Bory et Ch. Fl. Pelop. № 796, tab. 21 sub *Antirrhino*) annua vel perennans pilosa, multicaulis caulis prostratis elongatis prope basin saepè radicantibus, foliis breviter petiolatis inferioribus ovato-oblongis oppositis, superioribus sensim et valde diminutis triangulari vel lanceolato-hastatis, pedicellis axillaribus capillaribus patentibus folio multo longioribus cirrhiformiter curvatis glabris vel hirtis, calycis hirti laciniis lanceolatis acutissimis, corollæ flavidæ-cærulescentis vel flavæ breviter hirtulae calcare tenui rectius-

culo subæquilongo, capsulâ glabrâ coriaceâ, seminibus tuberculatis ⊖. *L. Graeca* Chav. Mon. p. 108. — *L. commutata* Bernh. in Rehb. Ic. IX, p. 6, tab. 815.

Hab. in collibus herbidis et montosis aridis, Græcia in Peloponneso (Bory!), Atticâ et Ætolia (Heldr!), Macedonia (Janka!), Byzantii (Commany!), Cyclades (Urv!), Creta (Heldr!), Rhodus (Bourg!), Lydia ad Smyrnam (Boiss!), Cilicia (Per!), Syria littoralis et Libani regio inferior (Boiss! Bl! Gaill!), Palestina in monte Carmel (Boiss!).

Planta Orientalis ab Italica et Dalmatica a quâ botanici nonnulli eam distinctam antumaverunt differt tantum caulis interdum magis elongatis, foliis floralibus et floribus paulo minoribus, probabiliter e locis siccioribus, sed omnino eadem species. Specimina ex Canariis, Gallia occidentali et australi, *L. caulinrhiza* Del., formam sistunt foliis obsoletis et saepe vix hastatis.

Ar. Geogr. Insulæ Canarienses, Gallia occidentalis et australis, Dalmatia, Africa borealis.

9. **L. cirrhosa** (L. Mant. p. 249 sub *Antirrhino*) annua vel perennans, hirsuta multicaulis caulis gracillimis elongatis ascendentibus vel procumbentibus, foliis parvis petiolatis radicalibus ovato-oblongis cæteris triangulari-lanceolatis et linear-lanceolatis acute hastatis alternis, pedicellis ex omnibus axillis capillaribus folio multo longioribus subinde cirrhiformibus, floribus minimis, calycis laciniis linear-lanceolatis acutis, corollâ cœrulecente calyce triplo longiore, calcare recto corollæ æquilongo, capsulâ globosâ calyce longiore, seminibus ovatis subcompressis crebre tuberculatis ⊖ *L. cirrhosa* Willd. Enum. 639. — Ic. Jacq. Hort. Vindob. tab. 82. — Rehb. Ic. tab. 60.

Hab. in vineis Cretæ (Sibth.), in Egypto (L.). E ditione non vidi.

Ar. Geogr. Hispania et Lusitania australes, Gallia australis, Corsica, Sardinia, Sicilia, Italia australis, Dalmatia, Africa borealis.

* * Perennes.

10. **L. ramosissima** (Wall. Pl. Asiat. rar. II, p. 43, tab. 153) glabra siccitate nigricans, basi suffrutescens, ramis herbaceis decumbentibus longissimis tenuibus, foliis petiolatis triangulati-hastatis infimis saepe 5-7-lobatis superioribus lanceolato-sagittatis, racemis longissimis foliosis laxis, pedicellis filiformibus rigidis petiolo longioribus folio brevioribus, calycis laciniis a basi lanceolatâ membranaceo-marginatâ setaceis acutissimis, corollæ luteæ calcare tubo breviore arcuato, capsulâ globosâ glabrâ, seminibus minimis echinatis ♂.

Hab. in Affghania (Griff. Herb. Ind. Comp. № 3858!).

Pedicelli eis *L. Cabulicae* subbreviores, planta glabra.

Ar. Geogr. Regio montana et alpina Indie Orientalis.

11. **L. Cabulica** (Bth. in D.C. Prodr. X, p. 270) patule hirta, basi suffrutescens ramosissima ramis herbaceis valde elongatis flexuosis,

foliis inferioribus longe petiolatis ovato-triangularibus grosse 2-3-lobato-hastatis superioribus sensim diminutis breviter lanceolato-hastatis acutis, racemis longis laxis, pedicellis axillaribus filiformibus rigidulis folio multo longioribus, calycis laciniis a basi lanceolata marginata setaceis acutissimis, corollæ breviter puberulae luteæ rubro-punctatæ calcare tubo breviore, capsula parvâ globosâ glabra calyce subbreviore, seminibus echinatis 5.

Hab. in Affghanist. (Griff!), in Belutschia ad Chehel Tun (Griff!), in Persia australi ad Kotel Mallu (Haussk!).

Rami tenues longe racemosi, folia inferiora pollicaria, floralia parvula, flores cum calcare 4 lineas longi. Probabiliter varietas hirsuta *L. ramosissima* Wallich; rhizomate suffruticoso et laciniis calycinis longis setaceo-attenuatis a formis *L. Elatines* et *Græcæ* distincta

12. *L. ovata*, patule hispida basi suffrutescens ramosissima ramis herbaceis tenuibus elongatis laxe et longe racemosis, foliis omnibus petiolatis ovatis integris, floralibus minimis acutis, pedicellis filiformibus rigidulis folio axillari multo longioribus, calycis hispidulii laciniis a basi lanceolata marginata subulatis, corollæ luteæ puberulae calcare tenui recurvo tubo sublongiore, capsula sphaericâ glabra calyce subbreviore, seminibus minimis echinulatis 5. — *L. ramosissima* var ? *ovata* Bth. in D.C. Prodr. X. p. 270.

Hab. in regno Cabulico ad Kandahar (Griff.).

Pedicellis setiformibus, indumento, valde affinis *L. Cabulicae*. sed folia ovata integra parva *L. Acerbiana*.

13. *L. Ægyptiaca* (L. Sp. 851 sub *Antirrhino*) pubescens, primum herbacea inermis demum suffruticosa ramosissima ramis rigidis patule ramulosis tandem spinulosis, foliis parvis petiolatis inferioribus oblongis obtusis ceteris diminutis acute triangulari hastatis, pedicellis axillaribus patentibus rigidis folio longioribus apice flexis demum spinescentibus, calycis laciniis lanceolatis acutis, corollæ flavâ paleo purpureo-maculato, calcare incurvo corollæ subæquilongo tenui, capsula parvâ globosâ velutinâ calyci subæquilongâ, seminibus ovatis granuloso-tuberculatis 5. — *L. Ægyptiaca* Dum. Courset. — Ic. Del. Eg. tab. 32. — *Antirrh. spinescens* Viv. Fl. Lib. p. 22, tab. 27.

Hab. in desertis Egypti medie (Del! Auch! Boiss! etc.), Arabiae petreæ (Schimp! Boiss!), Palestine inter Hierosolymam et Jericho atque circa Nazareth (Boiss!). Fl. Mart., Jun.

Ar. Geogr. Africa borealis orientalis.

14. *L. macilenta* (Decaisn. Flor. Sin. p. 14) glabra, suffruticosa ramosissima ramis rigidis subaphyllis elongatis breviter ramulosis demum spinescentibus, foliis inferioribus petiolatis ovato-vel lineari-lanceolatis integris vel basi sagittatis supremis subulatis integris, racemis terminalibus laxis breviter bracteolatis, pedicellis ca-

pillaribus calyce 2-4-plo longioribus, calycis glabri laciniis subulatis crassiunculis, corollâ hirtulâ calcare tenui subincurvo eæ æquilongo, capsulâ globosâ glabrâ parvâ calyce subbreviore, seminibus ovatis tuberculato-echinatis ♂.

Hab. in deserto Sinaico (Bové! Auch! Schimp. 147!), in monte Gheno Persiæ australis (Auch. 5069! Specimina manca).

Suffrutex ramis numerosis tenuiter virgatis, folia inferiora pollicaria 2-2 $\frac{1}{2}$ lin. lata, flores 7-8 lineas longi, (in spec. Boveanis multo minores sed subabortivi videntur), capsula magn. seminis Cannabeos. Species a *L. Ägyptiacâ* glabritie, formâ foliorum, ramis (vix pedicellis) spinescentibus diversa. *L. hastata* R. Br. affinis quoque pedicellis setaceis longissimis, floribus triplo minoribus differt.

Ar. Geogr. Nubia.

SECT. III. LINARIASTRUM.

§ 1. Discoideæ.

* Perennes.

15. ***L. pyramidata*** (Lam. Dict. IV, p. 360 sub *Antirrhino*) caulinibus erectis elatis crassis totâ longitudine crebre foliosis, foliis alternis glaberrimis semiamplexicanibus ovato-vél oblongo-lanceolatis acutis subquinquenerviis superne sensim diminutis, spicâ etiam fructiferâ densâ sèpius solitariâ, floribus magnis subsessilibus, bracteis lanceolatis, calycis dense papitloso-lanati laciniis brevibus late ovato-oblongis, corollâ luteâ palato flavo-lanato, calcare recto acuto corollâ subbreviore, capsulâ globosâ calyci subæquilongâ, seminibus marginatis disco tuberculatis nigris ♀. *L. pyramidata* Spr. Syst. II, p. 796. — *L. Orientalis* *Lilii Persici* foliis florum spicâ densissimi et p. *ramidata* Tourn. Cor., p. 9.

Hab. in Armeniâ (Tourn!), provinciâ Transcaucasicâ Talyssch (Hoh! C. A. Mey!), Karadagh (Buhse!).

β. *Kotschyana*. — Flores subminores, calyx parcius papillosus.

Hab. in valle Teng prov. Musch Armeniæ (Ky. 513!).

Planta 2-3-pedalis, folia inferiora 2-3-pollicaria 6-9 lineas lata, spica fructifera 3-5-pollicaris, corolla cum calcare in typo sesquipollicaris. in varietate pollicaris. Nomen specificum improprium.

16. ***L. Griffithii*** (Bth. in D.C. Prodr. X, p. 272) glabra, caulinibus erectis elatis crebre foliatis, foliis alternis glaucis semiamplexicaulinibus oblongis acutis 5-7-nerviis, spicâ elongatâ laxiusculâ, floribus magnis sessilibus, bracteis calycisque laciniis linear-lanceolatis acutis villosis, corollæ luteæ calcare tenui subincurvo corollâ subbreviore, capsulâ globosâ, seminibus discoideis marginatis ♀.

Hab. in regno Cabulico (Griff. 608!). Vidi in herb. Kew.

Affinis *L. pyramidata* differt spicâ tenuiore et laxiore, calycis laciniarum formâ.

17. L. confertiflora (Bth. in D.C. Prodr. X, p. 272) caulis pluribus a basi ascendentib; crebre et stricte foliosis, foliis glabris anguste oblongo-lanceolatis et linearibus subtrinerviis acutis, spicâ etiam fructiferâ densâ ovatâ tandem oblongo-cylindricâ, floribus subsessilibus, calycis lanato-villosi laciniis ovato-oblongis, corollâ flavâ palato flavo-lanato, calcare tenui acuto dimidiâ corollâ sublongiore, capsulâ globosâ calyce subbreviore, seminibus marginatis disco tuberculatis ♀.

Hab. in Mesopotamiâ circa Suerek, Kochhassar, Kifri (Ky. 170! Haussk!).
Fl. Mai

Semipedalis et pedalidis, folia 9-10 lineas longa $1\frac{1}{2}$ -2 lineas lata, corolla cum calcare pollicaris. Inflorescentiâ et characteribus affinis *L. pyramidata* differt foliis multo minoribus angustis paucinerviis non semiamplexicaulibus.

18. L. fastigiata (Chav. Mon. p. 125, tab. 7, fig. a.) glabra glaucescens, caulis elatis ascendentibus superne sèpius fastigiatim ramosis, foliis crebris subamplexicaulibus ovato-vel oblongo-lanceolatis plurinerviis sèpius latis, spicis densis tandem elongatis, bracteis margine membranaceis, floribus medioeribus pedicello calyce subbreviore suffultis, calycis glabri laciniis subnerviis oblongis vel oblongo-linearibus margine membranaceis obtusis, corollâ flavâ palato lanato, calcare tenui eâ dimidio breviore, capsulâ ovatâ magnâ calyce longiore. seminibus late marginatis disco lèvibus ♀.

Hab. in graminosis et agris otiosis montanis Kurdistaniæ Persicæ et Persiæ austro-occidentalis, inter Bagdad et Kermanschah (Oliv! Noë 1108!), in montibus Pir Omar Gudrun, Zagros, Teng Nallî 5000'-8000' et in montibus Schahu et Juanro 8000'-10000' Haussk!). prope Schiraz (Ky. 546! 913! Auch. 5070!), Persiæ occid. (Auch. 1917!).

Planta 2-3-pedalis, folia caulina sèpe 3-4 pollices longa $1\frac{1}{2}$ -2 lata, interdum autem angustiora, corolla 9-10 lineas longa.

19. L. Kurdica (Boiss. et Hoh. Diagn. Ser. I, 4, p. 73) glabra glaucescens, caulis elatis superne sèpius fastigiatim ramosis, foliis crebris caulinis plus minus late oblongis vel oblongo-lanceolatis subquinq; quenerviis, raueis abbreviatis anguste lanceolatis vel linearibus omnibus acutis, spicis densis tandem elongatis, bracteis lanceolatis membranaceis, floribus breviter petiolatis, calycis sèpius glabri laciniis lanceolatis acutis margine membranaceis, corolle flavæ palato lanato, calcare recto tenui corolle equilongo, capsulâ ovatâ, seminibus late marginatis lèviusculis ♀.

Hab. in glareosis mont's Gara Kurdistaniæ (Ky. 591!), Cataoniae monte Barsuch Dagh supra Malatia (Haussk!), Armenia prope Erzerum (Huet! Tchiht!), agris otiosis ad Baibut (Bourg!). Armenia Rossicæ et prov. Karabagh (Buhse! Kowal! exs sub *L. macroura*), prov. Aderbijan Persie (Szov!).

Inflorescentiâ et foliis plerumque latis atlinis *L. fastigiata* a quâ differt floribus dimidio minoribus (6-7 lineas longis), calcare longiore, calycis laciniis acutis. An nihilominus ejus varietas Formæ adsunt foliis angustioribus facie *L. Italica* referentes que floribus longius pedicellatis, bracteis angustè linearibus, capsulâ parvâ longe differt.

20. *L. Aucheri* (Boiss. Diagn. Ser. I, 7, p. 44) glabra, caulis tenuibus erectis simplicibus vel fastigiatim ramosis, foliis crebris strictis lanceolato-linearibus utrinque attenuatis 4-3-nerviis, spicis tandem laxinsculis, bracteis oblongis submembranaceis, floribus breviter pedicellatis, calycis puberuli laciniis enerviis ovato-oblongis obtusis, corollæ flavæ palato hirsuto, calcare rectiusculo corollâ subbreviore, capsulâ globosâ, seminibus marginatis lœviusculis ♀. *L. fastigiata* β. *angustifolia* Bth. in n.c. Prodr. — *L. Cœlesyriaca* Boiss. Diagn. Ser. I, 12, p. 41.

Hab. in incultis et agris derelictis Libani (Auch. 4906!), inter segetes ad et supra Cedros (Ky. 291!), supra Dimam (Bl!), in Cœlesyria (Ehrenb! Boiss!) in Antilibano (Boiss!).

Pedalis et elatior, folia 3-4 lineas lata, corolla 8-9 lineas longa. Affinis *L. fastigiata*, saepius simplicior, folia angusta nervis paucioribus, flores tertia vel dimidiâ parte minores, calyx pubescens.

21. *L. pycnophylla* (Boiss. et Bal. Diagn. Ser. II, 6, p. 129) glabra glaucescens, caulis elatis superne fastigiatim ramosis rariis simplicibus, foliis sessiliibus subsemiamplexicaulibus late ovato-oblongis acuminatis 5-7-nerviis, rameis squamæformibus, spicis densis demum elongatis laxis, bracteis lanceolatis, floribus breviter pedicellatis, calycis crispule tomentelli laciniis enerviis ovatis obtusis, corollæ luteæ palato hirto, calcare recto corollâ subbreviore, capsulâ sphærica, seminibus marginatis lœvibus ♀.

Hab. in regione alpinâ montis Aslandach Cappadociæ (Bal. 1034!), in schistosis vallis Teng Armeniae australis (Ky. 514 sub *L. fastigiata*!).

♂. *eriocalyx*. — Folia angustiora, calyx dense albo-tomentosus. — *L. eriocalyx* Boiss. in Bourg. exs. 1860.

Hab. in agris ad Adessa prope Elmalu Lyciae (Bourg. 1841)

Folia in typo eis *L. fastigiata* similia 6-10 lineas lata, in varietate angustiora sed quinquenervia, flores 7 lineas longi. A *L. fastigiata* floribus duplo minoribus et calycis laciniis brevibus tomenteilis diversa. *L. confertiflora* foliis brevibus angustis, spicis etiam fructiferis densis etc, longius differt.

22. *L. vulgaris* (Mill. Dict. N° 1) inferne glabra inflorescentiâ pubescenti-glandulosâ, caulis crebre foliosis vel superne fastigiatim vel parce ramosis, foliis sparsis anguste lanceolato-linearibus acutis trinerviis nervo medio superne sulco impresso subtus prominentem, spicis brevibus densis fructiferis elongatis, bracteis breviter lanceolatis reflexis, floribus majusculis breviter pedicellatis, calycis parvi glabri laciniis parvis membranaceis triangulari-ovatis trinerviis, corollâ flavâ calcare tenui subæquilongo, capsulâ magnâ ovaâ vel ovato-cylindricâ calyce duplo longiore, seminibus marginatis disco saepius tuberculatis ♀. *Ant. Linaria* L. Sp. 858. Ic. Fl. Dan. tab. 982. — *L. speciosa* Ten.

Hab. in cultis Laconiae (Sibth!), Atticæ ad Ampelokypos (Heldr!), in Macedonia prope Bitolia (Orph!), faucibus Thraciæ prope Kalofor (Pichler!), Tauriæ et prov. Caucasicis (M.B! Stev!), Caucaso australi ad Darial '500' (Rupr!).

L. latifolia. — Folia latiora. — *L. Biebersteinii* Besser Enum., p. 25. — Rehb. Ic. Crit. tab. 434.

Hab. in Caucaso prope Lars (Rehmi!).

Ar. Geogr. Europa borealis et media a Scandinavia ad Rossiam medium et Gallia et Italiæ ad Rossiani meridionalem, Songaria.

23. *L. macroura* (M.B. Taur. Cauc. II, p. 75 sub *Antirrhino*) glabra, caulis erectis vel ascendentibus crebre foliosis simplicibus vel parce et breviter ramosis, foliis planis anguste linearibus elongatis, racemo denso brevi paucifloro tandem elongato, bracteis lanceolato-linearibus, floribus magnis breviter petiolatis, calycis glanduloso-hirsuti laciniis tenuiter plurinerviis ovato-oblongis obtusis, corollæ amplæ flavæ palato barbato, calcare recto acuto corollæ longiore, capsulâ ovatâ calyce sublongiore, seminibus levibus margine membranaceis ♀. — *L. macroura* Chav. Mon. Ic. — M.B. Cent. tab. 27. — Rehb. Ic. Crit. tab. 432. — *L. rupestris* C. A. M. Enum. p. 110!

Hab. in campestribus Tauriæ (M.B., Stev!), Caucasi septentrionalis (Lebed.), Ossetiæ in fauce Darial (C.A. M!) et distr. Alagir 3000' (Rupr!).

Facie similis *L. vulgari*, folia longiora et angustiora, flores majores, calcar manifeste longius, calyx hirtus. — *L. rupestris* C.A. M. ex specimine valde manco ejus herbarii nihil est nisi forma diffusa ramosissima ejusdem speciei.

24. *L. odora* (M.B. Taur. Cauc. II, p. 76) glabra glauca, caulis gracilis erectis a basi strictiusecula ramosis, foliis remotiuseculis linearis-subulatis semiteretibus subcanaliculatis, floribus paucis laxè et breviter racemosis breviter pedicellatis, calycis parvi glabri vel hirtuli laciniis subnerviis ellipticis obtusiusculis, corollæ luteæ palato barbato, calcare recto vel subincurvo corollæ breviore, capsulâ globosâ, seminibus late marginatis ♀. — *L. odora* Chav. Mon. Ic. — Rehb. tab. 64. — *L. Læselii* Schweigg. — *L. juncea* Rehb. pl. Crit. tab. 425 non B.C. — *L. venosa* Lindl. bot. Reg. 1841 Misc. p. 70. — *L. Altaica* Fisch.

Hab. in arenosis maritimis, ad Kila prope Byzantium (Duparq!), Tauriæ ad Sudak (Rehmi!), prov. Talysch alt. 4000' (ex Hohen.), ad radices montis Demavend Persiae borealis (Ky. 720! saltem ex incompleto specimine), prov. Khorassan (Bge!), littore orientali maris Caspii (Karel!), Affghania (Griff!).

Ar. Geogr. Germania borealis orientalis, Rossia media et australis, Songaria, Sibiria Uralensis et Altaica.

25. *L. Damascena* (Boiss. et Gaill. Diagn. Ser. II, 6, p. 130) glabra glaucescens, pluricaulis caulis ascendentibus foliosis in racemum simplicem post anthesin elongatum densiusculum glabrum abeuntibus, foliis alternis lanceolato-linearibus acutis subtrinerviis nervo medio superne non sulco impresso, pedicellis calyce brevioribus, bracteis oblongis acutis pedicello longioribus, floribus magnis, calycis breviter et crispule pubescens et ciliati laciniis plurinerviis oblongo-ellipticis acutiusculis basi attenuatis, corollæ flavæ glabré

palato concolor minute pruinosa, calcare recto acuto corollæ æquilongo, capsula sphæricâ calyce subbreviore, seminibus discoideis lœvibus ♀.

Hab. in collibus siccis Gebel el Assouad ad Meridiem Damasci sitis (Gaill!).
Fl. Apr.

Caulis denum cum racemo sesquipedale, folia 1-1½ pollicem longa sesqui-lineam lata, corolla 14-15 lineas longa. A *L. fastigiata*, Aucheri et affinis sepalis nervosis, corollâ magnâ, etc. distincta, magis affinis *L. vulgari* quæ discrepat foliis longioribus sulcatis, sepalis angustioribus, corollæ palato discolore barbato, capsula ovalâ calyce longiore. Planta e Damasco in D.C. Prodr. sub *L. Biebersteinianâ* citata huc spectat.

26. *L. Nurensis* (Boiss. et Haussk.) glabra glaucescens, caulis elatis erectis vel ascendentibus simplicibus basi demudatis dein crebre foliosis in racemum simplicem oblongum tandem elongatum abeuntibus, foliis crassiusculis lanceolatis acutiusculis supra subenerviis subtus elevatim 3-5-nerviis, bracteis oblongis, pedicellis calyce brevioribus, floribus magnis, calycis glabriuscili carnosuli lacinis elliptico-oblongis acutiusculis margine membranaceis basi elevatim 3-4-nerviis, corollâ flavâ concolor palato villosâ, calcare incurvo acuto corollâ breviore, capsula ovalâ calyci subæquilongâ, seminibus marginatis lœvibus ♀.

Hab. in regione alpinâ superiore montium Nur et Sawers Persiae australis 9600'-14300' (Haussk!). Fl. Jun.

Pedalis et bipedalis, racemus florifer 2-3-pollicaris, folia 1-1½-pollicem longa 2-3 lineas lata, corolla eâ *L. vulgaris* major fauce ampliori. Facies *L. Dalmaticæ* seminibus triquetris alienæ, affinis *L. Dainascene* que differt foliis angustioribus tenuioribus, calyce crispare pubescenti, calcare longiore recto, capsula sphærica breviore.

27. *L. Michauxii* (Chav. Mon. p. 124, tab. 7, fig. B) glabra glauca carnosula, caudiculis decumbentibus tenuibus nudis fragilibus in caules brevissimos ad apicem usque confertissime foliosos simplices procumbentes abeuntibus, foliis subsessilibus obovatis obtusis acutiusculis basi cuneatis, floribus magnis brevissime pedicellatis, bracteis oblongis pedicello longiocibus, calycis glabri lacinis elliptico-oblongis subspathulatis obtusis corollæ tubo duplo brevioribus, corollæ purpureo-violaceæ calcare recto sensim attenuato eæ æquilongo, palato barbato flavescente, capsula globosa maximâ calyce sublongiore, seminibus discoideis marginatis lœvibus nigris ♀. *L. Derderiana* Auch. MSS.

Hab. in Persiâ prope Teheran (Ky 36!), in districtu Veronine ad margines magni deserti salsi (Bode!), in montibus prope Ris-chm et Jezd (Buhse!), ad Ssertschah (Bge!), in monte Kohrud propè Kaschan (Bode!), circa Ispahan (Mich! Auch. 5072!), in monte Soi (Derderian!).

Caudiculi e radice plus minus elongati in caules 3-5-pollicares abeunt, surculi steriles pauci foliis subminoribus, folia caulinâ 4-6 lineas lata et longa saepè orbiculata, corolla sesquipollicem longa, capsula piso duplo major. Species insignis facie *L. glacialem* Boiss. quo-lammodo referens sed folia, etc., diversissima.

** Annua.

28. **L. Pelisseriana** (L. Sp. p. 855 sub *Antirrhino*) glabra, collo sureulos steriles foliosos edente, caule erecto gracili simplici vel parce ramuloso, foliis sureulorum oblongis verticillatis, caulinis remotis alternis longe et anguste linearibus, floribus paucis capitatis tandem laxe et breviter racemosis, pedicellis calyce sublongioribus strictis, bracteis calycinisque laciniis linearibus acutissimis, corollæ violaceæ lobis superioribus elongatis acutis, calcare recto tenuissimo corolla æquilongo, capsulâ globosâ calyce breviore, seminibus parvis discoideis radiationi ciliatis ⊖. *L. Pelisseriana* D.C. Fl. Fr. III., p. 589. — Ic. Fl. Græc., t. 591. — Rehb. Ic. tab. 62.

Hab. in cultis et herbidis regionis inferioris totius Graeciae (Sibth., Boiss! Heldr! Orph. Fl. exs. 713!), Macedonia et Thracia (Friv! Griseb!), Archipelagi (Urv!), Creta (Sieb! Raul!), totius Anatoliae occid. et australis!, Syrie maritimæ (Auch. Bl! Gaill!), prov. Transcaucasiarum (Wittm. ex Ledeb.).

Ar. Geogr. Gallia media et australis, Italia, Dalmatia.

29. **L. arvensis** (L. Sp. p. 855 sub *Antirrhino*) annua glaucescens inferne glabra, collo sureulos steriles breves foliosos edente, caule tenui erecto simplici vel parce ramoso, foliis linearibus, floribus parvis breviter pedicellatis capitatis, racemo fructifero elongato laxissimo, bracteis linearibus tandem reflexis, calycis glandulosi laciniis oblongo-linearibus obtusis basi subattennatis capsulâ globosâ subbrevioribus, corollæ tubo calyce breviore lobis superioribus patentibus, calcare subulato recto vel incurvo corollâ dimidio breviore, seminibus plano-orbiculatis marginatis disco levibus nec tuberculatis ⊖. *L. arvensis* Desf. Atl. — Ic. Rehb. tab. 62.

Hab. in arvis Peloponnesi (Sibth!).

β. flavidora. — Corolla lutea interdum violaceo-striata. — Ic. Fl. Græc. tab. 590. — *L. simplex* D.C. Fl. Fr. III., p. 588. — *Ant. parviflorum* Jaeq. Ic. rar. III, tab. 499 non Desf.

Hab. in cultis et arenosis totius ditionis in regione inferiore a Graeciâ! et Thraciâ! ad Anatoliam! Syriam borealem ad Aleppo et Mesopotamiam (Haussk!), Arabiam petreain (Schimp. 119!), Armeniam et Pontum Lazio (Bourg! Ball!), Tauriam (M.B.), Transcaucasiam (Hohen!), Persiam borealem (Auch! Bge!).

Ar. Geogr. Europa tota media et australis a Galliâ et Germaniâ ad regionem Danubialem, Africa borealis.

30. **L. micrantha** (Cavan. Ic. I. p. 51 sub *Antirrhino*) annua glauca, collo sureulos steriles breves foliosos edente, caule unico vel pluribus pumilis simplicibus, foliis brevibus surculorum linearibus caulinis oblongo-lanceolatis acutis, floribus capitatis brevissime pedicellatis, racemo fructifero laxe brevi, calycis glabri vel glandulosociliati laciniis oblongo-linearibus obtusis capsulâ ovatâ subbrevioribus, corollâ minimâ calycem parum excedente caruleâ, calcare brevissimo

conico, seminibus orbiculatis membranaceo-marginatis ⊖. *L. micrantha* Spreng. — Ic. Fl. Græc. tab. 587. — *L. parviflora* Desf. Atl. II, p. 44, tab. 137.

Hab. in cultis Atticæ et insulæ Syra (Heldr!), Eubeæ (Sprun!), Rhodi (Sibth.), Syriæ ad Damascum (Gaill!), Persiæ australis ad Gere (Ky!), Transcaucasiæ prope Baku (C.A. Mey!).

A *L. arvensi* formâ foliorum et calcare brevissimo gibbiformi distincta, flores minores.

Ar. Geogr. Gallia australis, Sardinia, Hispania, Africa borealis.

2 §. Oblongæ.

* Perennes.

31. *L. Dalmatica* (L. Sp. p. 857 sub *Antirrhino*) glabra glauca simplex vel sæpius paniculatim ramosa, radice repente, foliis alternis acutis ovatis oblongis vel lanceolatis semiamplexicaulibus, racemis tandem elongatis laxis, bracteis oblongo-lanceolatis acuminatis inferioribus pedicello longioribus, pedicellis strictis calyci æquilongis vel longioribus, calycis laciniis late lanceolatis acutissimis enerviis corollæ tubum subæquantibus, corollâ magnâ luteâ palato villoso, calcare recto corollâ breviore, capsulâ ovatâ calyce subbreviore, seminibus acute triquetris tuberculato-rugosis ♀. — *L. Dalmatica* Mill. № 13. — Ic. Rchb. Pl. Crit. tab. 438. — Ic. Germ. tab. 65. — *L. Smithii* Boiss. et Orph. in Fl. Græc. exs. № 711! (forma caulinibus elatis simpliciuseulis, racemo laxissimo.)

Hab. in regione inferiori montis Malevo Laconiæ (Orph!), monte Delphi Eubeæ (Sprun!), Macedoniâ in monte Orbelo (Friedr!).

β. grandiflora. — Flores maximi brevissime pedicellati, laciniæ calycinæ latiores longiores capsulam sæpe longe superantes. — *L. grandiflora* Desf. Cor. p. 30, tab. 21. — *L. calycina* Boiss et Bal. MSS.

Hab. in incultis Cariæ et Ponti (Tchih!), Syriæ borealis ad Aintab et in monte Sscaffagh (Haussk!), Cappadocia (Ball), Armenia (Tourn! Auch 1915!), ditionis Talysh et Armenia Rossica (C.A. Mey! Hoh!), Persiæ borealis (Buhse! Aucher!) et occidentalis ad Mendeli (Noël).

Valde variat caulinibus simplicibus vel paniculatis, foliis plus minus latis, floribus in typo pollicaribus, in varietate bipollicaribus, calycis laciniis plus minus longis et latis.

Ar. Geogr. Dalmatia, regio Danubialis.

32. *L. Macedonica* (Griseb. Spic. II, p. 19) glabra glauca, caule procero superne patule ramosissimo, ramis et ramulis tenuiter virgatis in racemos tenues laxos abeuntibus, foliis caulinis ovatis semiamplexicaulibus acutis, rameis parvis breviter lanceolatis acuminatis, bracteis minimis squamæformibus, pedicellis gracilibus erecto-patulis calyce et bracteâ 4-6-plo longioribus, calycis parvi

laciiniis triangularibus acuminatis corollæ tubo multo brevioribus, corollæ flavæ palato barbato, calcare tenuissimo recto corollæ æquilongo, capsulâ ovatâ retusâ calcare 2-3-plo longiore, seminibus acute triquetris tuberculato-rugosis ♂.

Hab. in Macedoniae occidentalis rupibus graniticis montis Babuna prope Perlepe 2500' (Orph!), prope Dichovo et Bitolia (Fried! Orph!).

An hæc quoque, cl. Fenzl (Tchih. As. Min. II, p. 25) indicante, formis *L. Dalmaticae* adnumeranda? Ex specimine valde differt foliis rameis bracteisque minimis, pedicellis tenuibus 4-6 lineas longis nec brevibus crassis, calyce minimo, calcare longissimo tenui, capsulâ minore retusâ calyce multo longiore; explorandum tamen an formæ intermediæ non occurrant.

33. *L. præalta* (Boiss. et Bal. Diagn. Ser. II, 3, p. 162) glaberrima, surculis sterilibus longis foliosis sub anthesi exsiccatis, caule recto procero superne stricte paniculato, ramis longis virgatis in racemos longos laxissimos abeuntibus, foliis caulinis semiamplexicaulibus ovato-lanceolatis acutis 5-7-nerviis, rameis valde diminutis oblongo-lanceolatis, pedicellis calyce et bracteâ squamiformi brevioribus, laciiniis calycinis oblongis acutis corollæ tubo brevioribus, corollæ flavæ latiusculæ brevis palato barbato, calcare subinfuso longitudine calycis, capsulâ sphæricâ calyce sublongiore, seminibus triquetris minute tuberculatis ♀ vel ②?

Hab. in fauce Guzel deré supra pagum Kechlick non procul a Mersina Ciliciæ littoralis (Bal!).

Surculi steriles pedales, caulis cum ramis pedalibus 4-5-pedalis, racemi pauciflori semipedales et longiores, folia caulinâ 1-1½-pollicaria 6-7 lineas lata. Monocarpica videtur sed ex specimine unico non dijudicare possum, a *L. genistifolia* differre videtur corollâ non majore (5 lin. longâ) sed quoad longitudinem latiore et in primis calcare brevissimo.

34. *L. genistifolia* (L. Sp. p. 858 sub *Antirrhino*) glabra glauca, caulis erectis paniculato-ramosis, radice repente, foliis strictis lanceolatis acutis 3-5-nerviis, rameis abbreviatis linearibus, racemis laxiusculis virgatis, pedicellis strictis calyce et bracteâ brevioribus, floribus medioeribus, calycis laciiniis lanceolatis acutis corollæ tubum æquantibus, corollæ luteæ palato barbato, calcare corollam subæquante rectiusculo vel eâ breviore, capsulâ ovatâ calyce subbreviore, seminibus acute trigonis rugulosis ♀. — *L. genistæfolia* Mill. 14. — Ic. Jacq. Austr. tab. 244. — Fl. Græc. tab. 596. — *L. silenifolia* Fisch. Cat. Gor. — Rehb. Crit. tab. 437. — *L. chloræfolia* Rehb. Crit. tab. 436. — *L. hians* Fenzl MSS. — *L. orgyalis* Reuter MSS.

Hab. in collibus, dumetis siccis, agris incultis Macedoniae (Heldr! Orph. Fl. Gr. exs. 711!), Byzantii (Boiss!), Anatoliæ borealis!, Armeniæ (Calv!), Tauriæ et Caucasi (M B! Rehm! etc.), Ciliciæ (Bal! Ky. 397!), montis Cassii Syriæ borealis (Boiss!).

β. confertiflora. — Racemi etiam fructiferi densiusculi, calyx capsulâ brevior. — *L. monochroma* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 41.)

Hab. in cumpis Isauriae et Pisidiæ (Heldr!), Cariæ (Pinard!), Lyciae ad Elmalu (Bourg!).

γ . *linifolia* — Folia anguste linear-lanceolata et linearia, flores dissiti. — *L. linifolia* auct. plur. non L. herb. quæ *L. vulgaris*. — *L. Syspirensis* G. Koch Linn. XXIII, p. 717. — *L. flavissima* Boiss. MSS. — *L. concolor* Griseb. Spic. II, p. 21. — *Antirrhinum supinum* Fl. Graec. VI, p. 75, tab. 595 non L. — Buxb. Cent. I, tab. 25.

Hab. in Macedonia (Griseb!), circa Byzantium (Sibth!), Broussam (Pichler! specimina iconi Sibthorpianae simillima), Tmolus Lydie (Boiss! Bal!), montibus Lyciae (Bourg. 186!), regione subalpinâ Argæi Cappadociae (Bal!), valle Djimil Ponti Lazici 6000' (Bal!), Armenia Turcica (C. Koch! Bourg!).

δ . *venosa*. — Foliis angustis similis varietati *linifolia* sed corolla sordide flava rufescenti-reticulata. — *L. Balansæ* Boiss. et Rent. Diagn. Ser. II, 3 p. 163. — *L. polyclada* Fenzl in Tchih. As. Min. II, p. 23.

Hab. in Tauro Cilicio circa Gülek et Bulgharmaaden (Ky! Bal!).

Species summopere polymorpha et cuius formas reiterato examine intermedii conjunctas video. Collum surculos steriles foliosos nullos sed interdum paucos edens; caules et rami plantæ junioris prostrati vel procumbentes, annorum subsequentium erecti elatiores foliis saepius latioribus, flores eis *L. Dalmatica* semper duplo vel triplo minores.

Ar. Geogr. Germania orientalis, Italia media, Dalmatia, regio Danubialis, Rossia media et australis, Sibiria.

35. *L. Sibthorpiana* (Boiss. et Heldr. in Heldr. exs.) pluricaulis, surculis sterilibus destituta, præter inflorescentiam glabra, caulis erectis virgatis crebre foliosis simplicibus vel paniculatis, racemis confertifloris oblongis fructiferis parum elongatis, foliis erecto-patulis anguste linearibus uninerviis elongatis, bracteis squamiformibus, pedicellis calyci aequilongis vel sublongioribus, calycis parvi lacinias margine membranaceis ovatis vel oblongis obtusis, corollæ luteæ palato barbato, calcare recto tenui aequilongo, capsula parvæ ovatæ ralyce 2-3-plo longiore, seminibus triquetris tuberculatis φ .

α . *Peloponnesiaca* — Racemi rachis et calyx glabri vel glabrescentes, laciniae calycinae ovatae. — *L. Peloponnesiaca* Boiss. et Heldr. Diagn. Ser. II, 3, p. 163.

Hab. in pascuis saxosis Taygeti Laconiae in regione alpinâ (Heldr. exs!), montis Malevo 4-5000' (Orph. Fl. Gr. exs. sub *L. genistefolia*).

β . *Parnassica*. — Racemi rachis, pedicelli et calyx pubescenti glandulosi, laciniae calycinae oblongo-lineares. — *L. Parnassica* Boiss. et Heldr. loc. cit. p. 164. — *Antirrhinum strictum* Fl. Graec. tab. 394 non *L. stricta* Guss. — *L. pallida* Sprun. exs.

Hab. in saxosis regionis abietinæ et alpine Parassi 3500'-7000' (Sprun! Heldr. Herb. Norm. 655!), in regione sylvaticâ Olympi Thessali 5000'-6000' (Heldr! Orph. Fl. exs. 715!), in Macedonia supra Vodena et in monte Peristeri (Orph!).

Planta 1-2-pedalis et in excelsioribus semipedalis, floribus magnitudinis *L. genistifolia*, sed ab hujus formis angustifoliis quibuscum stepe confusa fuit distincta foliis longis angustis fere *L. vulgaris*, et praesertim calycis laciniis obtusis membranaceo-marginatis ovatis vel oblongo-linearibus nec trianguli-lanceolatis acutis. Icon Flora Græcia nostram plantam exacte exhibit sed loci Caria et Cyprus erronei videtur, et planta Sienla indicata est species annua diversissima *L. aparinoides* Willd. = *L. stricta* Guss.

Ar. Geogr. Montenegro.

36. *L. lineolata* (Boiss. in Ky., exs. 1846. — Diagn. Ser. I, 12, p. 42) pluricanlis inflorescentia excepta glabra, canibus tenuiter virgatis valde foliosis erectis simplicibus vel parce ramulosis, foliis strictis filiformibus longissimis, racemo brevi densiusculo rachide glabra vel hirtula, pedicellis brevissimis, bracteis linearibus, calycis parvi glabri vel crispule hirti laciniis linearibus anguste membranaceis obtusiusculis, corolla lutea violaceo-lineolata, calcare tenui recto dimidiâ corollâ longiore, capsula. . . . ♀. *L. pulchella* Hohen. exs. — *L. striata* Ledeb. Fl. Ross. III, p. 210 quoad pl. Caucasicum! non D.C.

Hab. in fane Dudera montis Elbrus Persiae borealis (Ky. 265!), Persia boreali (Auch. 507!), in agris lapidosis ad Tatmi ditionis Talyisch (Hoh!).

♂. Elymaitica. — Glaberrima, folia lineam aut paulo amplius lata.

Hab. in regione alpinâ montium Nur, Sawers Persiae austro-occidentalis (Haussk!)!

Caules 1-1½-pedales, folia formæ typice 1½-2-pollicaria lineam dimidiâ vix lata in ♂ sublatiora, racemus densus, corolla 7-8 lineas cum calcare longa. Semina ignota sed hujus gregis omnino videtur, a formis angustifoliis *L. genistifolia* calycis laciniis elongatis angustis distincta. Facies *L. striatae* ♂ confertiflora.

37. *L. coridifolia* (Desf. Cor. Tourn. p. 32, tab. 22 (*corifolia*) glaberrima multicanalis glaucescens, caulis tenuissime virgalis erectis rigidulis simplicibus vel stricte ramulosis crenatim foliosis in racemos tenues laxifloros abeuntibus, foliis subulatis brevibus acutis, pedicellis strictis bracteâ lanceolata calyceque brevioribus, floribus parvis, calycis laciniis breviter lanceolatis acutis, corolla pallide griseo-violacea labio superiore elongato lobis acutiusculis, calcare conico tubo æquilongo vel breviore, capsula minimâ globosâ retusa calyce sublongiore, seminibus acute angulatis tuberculato-rugosis ♀. *Linaria Orientalis Coridis folio flore leucophovo* Tourn. Cor. Inst. 9. — *L. Dschlorochensis* C. Koch Linn. XXIII, p. 568! Semina errore quodam discoidea dicta.

Hab. in sterilibus lapidosis, collinis, agris derelictis Anatoliae in Cariâ (Bois!), Lycidâ ad Elmaliñ (Bourg. 185!), Pisidiâ et Lycaoniam (Heldr.), Ciliciâ ad Tauri radices (Ky! Bal!), Cappadociâ (Auch. 1912! Bad!), Catoniaâ (Haussk!), Paplagoniâ et Galutiâ (Wiedt!), Armenia circâ Karput, Erzerum, Gunnelslikhane (Tourn! Noe! Huët!), Galatia (Tchilt!).

Planta semipedalis vel pedalis, facies et flores minuti *L. striatae* a qua

differt foliis tenuissimis brevibus, racemis laxissimis, corollæ lobis superioribus elongatis acutis, capsulâ dimidio minore *Cannabeos* semen non æquante. Variat raro corollâ lutescente.

Obs. Inter plantas Sprunerianas habeo specimen *L. striatae* Gallicis similium et ex monte Delphi Eubeæ indicatum, sed vereor ne error in schedulâ irrepserit, nam *L. strata* est species occidentalis et planta Dalmatica hoc nomine salutata calycis laciniis obtusis certe differt.

38. L. Iconia (Boiss. et Heldr. Diagn. Ser. I, 12, p. 42) glaberrima, caulis pumilis ascendentibus a basi ramosissimis, foliis filiformibus acutiusculis, floribus laxe racemosis, pedicellis calyce brevioribus bracteas lineares æquantibus, calycis laciniis albo-marginatis linear-i-oblongis obtusiusculis, corollæ violaceæ palato aurantiaco-fulvo, calcare subulato incurvo corollæ æquilongo, capsulâ ovatâ calycem æquante, seminibus acute angulatis subincurvis reticulatis ♀.

Hab. in arenosis planitiei ad Iconium Lycaoniæ (Heldr!). Fl. Jun.

Semipedalis, facies *L. odoratæ*, foliis tenuibus et florum magnitudine affinis *L. coridifolia* a quâ præter caules a basi ramosissimos differt calcare longo subulato nec brevi conico.

39. L. purpurea (L. Sp. p. 853 sub *Antirrhino*) glaberrima, caulis erectis elatis crebre foliosis fastigiatis paniculatis, foliis linearibus elongatis inferioribus et surculorum linear-i-lanceolatis verticillatis cæteris alternis, racemis elongatis densiusculis, bracteis linearibus, pedicellis calyci æquilongis vel sublongioribus, floribus parvis, calycis laciniis anguste lanceolatis acutiusculis margine membranaceis, corollæ violaceæ calcare arcuato acuto tubo longiore, capsulâ parvâ globosâ retusâ calyce subtriplo longiore, seminibus triquetris rugosis ♀. *L. purpurea* Mill. N° 5. — Ic. Fl. Græc. tab. 589.

Hab. in Græciâ ex herb. Sibthorpiano, sed eam e ditione non vidi.

Ar. Geogr. Italia australis, Sicilia; ex Africâ boreali valde dubia.

* * Annuæ

40. L. triphylla (L. Sp. 852 sub *Antirrhino*) annua glabra, collo surculos steriles edente, caule crassiusculo simplici vel breviter et parce ramuloso, foliis carnosulis sessilibus ovatis vel ellipticis obtusis, surculorum oppositis, caulinis ternatis, spicâ terminali densâ tandem laxisculâ, floribus sessilibus, calycis laciniis ovato-oblongis obtusis, corollâ tricolore vel cæruleâ, calcare arcuato acutato corollæ subæquilengo, capsulâ magnâ ovatâ, seminibus triquetris profunde reticulatis ⊖. *L. triphylla* Mill. — Ic. Bot. Mag. tab. 324. — Rehb. Ic. tab. 63.

Hab. inter segetes Græcie (Sprun! Heldr!), Archipelagi (Urv.).

Ar. Geogr. Regio mediterranea, Europa tota, Africa borealis.

41. L. reflexa (L. Sp. 857 sub *Antirrhino*) glabra, caulis decurrentibus vel ascendentibus, foliis ovatis vel oblongis acutis

carnosalis glaucis sive rufis et inferioribus ternis supereribus alterois, floribus ad axillas racemum pauciflorum laxum foliatum formantibus, pedicellis rectis folio brevioribus vel aequilongis fructiferis arenato-retexis elongatis folio longioribus, calycis laciniis lanceolatis acutis, corollae cæruleæ vel albida lobis superioribus elongatis, calcare gracili rectiseculo corella subæquilongo, capsula globosâ calyce ampliata breviore, seminibus oblongis subincurvis profunde rugoso-scorbolatis ♂. *L. reflexa* Desf. Atl. II, p. 42. — Ie. Fl. Grec. tab. 593.

Hab. in cultis Argolidis (Sibth.). E ditione nondum vidi.

Ar. Geogr. Corsica, Sardinia, Sicilia, Italia australis, Africa borealis.

42. *L. Chaleensis* (L. Sp. 857 sub *Antirrhino*) glaberrima, collo surculos breves foliosos edente, caulis erectis in racemum spiciformem laxum abeuntibus, foliis surculorum et inferioribus oblongis et ellipticis oppositis cæteris linearibus alternis, pedicellis calyce bracteisque linearibus breviore, calycis laciniis patulis linearibus acatis corollam aequalibus aut eam superantibus, corollæ albae lobis superioribus lineari-oblongis, calcare lineari-subulato gracillimo arcuato corollâ sesqui vel duplo longiore, capsula ovatâ calyce aucto valde superatâ, seminibus angulatis grosse rugoso-serbiculatis ♂. *L. Chaleensis* Mill. Dict. № 12. — Ie. Fl. Grec. tab. 392. — Ie. Rehb. tab. 66.

Hab. in arenosis et cultis Peloponnesi (Boiss!), Attica et Eubœa (Sprun! Heldr!), Zacynthi (Marg!), Crete (Heldr!), Cycladum (Oliv!), totius Anatoliae occidentalis et australis! Syriae littoralis et circa Damascum (Gail!), Palestinae (Boiss!), prope Aleppo et in Mesopotamia (Ky! Hanssk!), Armeniâ ad Gu-muschkhâné (Bourg!). Persiae borealis in monte Elbrus et australis prope Persepolin (Ky!).

Ar. Geogr. Gallia australis, Italia, Dalmatia.

43. *L. Armeniaca* (Chav. Mon. p. 147) glaberrima, caule erecto simplici vel ramoso, racemis laxis paucifloris, foliis linearibus elongatis obtusis præter infima opposita alternis, pedicellis strictis calyce sublongioribus, bracteis minutissimis, calycis laciniis linearibus tubum corollæ subæquantibus, corollæ cæruleæ lobis superioribus oblongis, calcare subulato tenuissimo arcuato coroliâ 2-3-plo longiore, capsula globosâ calyce aequilongâ, seminibus triquetris scorbiulatis ♂. *L. segatalis* C. Koch Linn. XVII, p. 286!

Hab. in Armeniâ (Tourn.), Armeniâ Rossicâ (C. Koch!), in arenosis et saxosis Georgiae Caucasicæ (Hoh!), ditione Talysh (C.A. Mey!).

Valde affinis corolla formâ et magnitudine *L. Chaleensi* a qua differt tantum pedicellis longioribus, bracteis minimis nec linearibus elongatis, tandem calycis laciniis corollæ tubum tantum vel vix aequalibus. Impirendum tamen an formæ intermediae non ocurrant.

44. *L. Hælava* (Forsk. Eg. Arab. p. III sub *Antirrhino*) nana multicaulis inferne glabra ad inflorescentiam glanduloso-pubescentes,

caulibus diffusis tortuosis ramulosis, foliis linearibus crassiunculis inferioribus 3-4-nis vel oppositis cæteris alternis, racemis terminalibus spiciformibus paucifloris ovatis densiusculis fructiferis elongatis, pedicellis bracteolâ oblongâ calyceque brevioribus, calycis hirti laci-niis ovato-oblongis acutis, corollâ violacea, sulphureâ vel variegatâ, palato ampio, calcare recto acutato corollâ longiore, capsulâ globosâ retusa calyce æquilongâ, seminibus minutissimis reniformibus ten-uissime tuberculatis ⊙ *L. Hælava* Chav. Mon. p. 164.

Hab. in arenosis maritimis et desertorum Egypti, prope Alexandriam et Aboukir (Forsk., Ehrenb! Samar! Gaill!), in deserto ad Kahiram et ad Suez (Wiest! Boiss! Ky!). Arabiâ petreâ (Boiss!), in Palestinâ australi inter Gaza et Ascalon (Ky 542!).

Plantula 3-5-pollicaris collo ramosissima corollâ magnâ cum calcare 7-9 lineas longâ.

45. ***L. Ascalonica*** (Boiss. et Ky. Diagn. Ser. II, 3, p. 165) glaberrima pumila caule erecto flexuoso a basi ramoso, foliis suren-lorum oppositis breviter lineari-oblongis, caulinis et rameis sparsis longis subulato-filiformibus, racemis terminalibus paucifloris laxissi-mis, pedicellis strictis calyci æquilongis bracteâ brevi subulatâ suf-fultis, calycis glabri laci-niis linearibus membranaceo-marginatis acutiusculis, corollæ flavæ calcare recto eæ subæquilongo, capsulâ oblongo-cylindricâ calyce sesquitlongiore, seminibus minimis nigris transverse corrugatis.

Hab. in Palestinâ circâ Ascalon (Ky!).

Ex specamine unico cum specimenibus *L. Hælave* commixto mihi nota, 5-pollicaris. Affinis *L. Hælave* sed distinctissima. Folia multo angustiora pollicem longa, pedicelli elongati, flores dimidio minores, calycis glabri laci-niæ lineares, capsula imprimis valde elongata et seminum structura diversa. *L. albifrons* foliis latis, calcare brevi, seminibus profunde punctato-insculptis lon-gius differt.

46. ***L. albifrons*** (Sibth. et Sm. Prodr. I, p. 432 sub *Antirrhino*) annua glabra glaucescens pumila simplex vel a collo pluricaulis, cau-libus simplicibus vel stricte et parce ramosis, foliis oblongo-lanceolatis acutis basi attenuatis infinitis oppositis vel ternis cæteris alternis, floribus parvis brevissime pedicellatis paucis subcapitatis tandem breviter et laxe spicatis, bracteis linearibus calyces subæquatibus, calycis glaberrimi laci-niis linearî-oblongis obtusis corollam subæquantibus, corollâ albidâ tubo violascente palato concolore vel lutescente, calcare linearî corollâ triplo breviore, capsulâ obtongâ calyce aucto æquilongâ vel sublongiore, seminibus ovatis obtuse angulatis pro-funde punctato-impressis ⊙. Ic. Fl. Græc. tab. 588. — *L. modesta* Boiss. Diagn. Ser. I, 12, p. 43. — *L. glauca* Ehrenb. MSS. — *L. mi-nutiflora* C. A. M. Eruin. p. 109.

Hab. in agris inter segetes et arenosis, Rhodus (Sibth.), Egyptus ad Ale-xandrian (Ehr! Samar!), Arabia petrea (Boiss!), Palestina australis prope

Asdod et Gaza (Ky. 422!), ad Hierosolymam (Roth. 582!), insulæ Karak Persia australis (Ky!), Transcaucasia prope Baku ad Caspium (C.A. Mey!).

Plantula 3-5-pollicaris, folia 2-3 lineas lata, corolla cum calyce 3-3½ lin- longa, capsulæ magnæ. Huc prob. spectat quoque *L. tumens* Steud. non in tantum nota.

Ar. Geogr. Regnum Tunetanum.

SECT. IV. CHENORRHINUM.

47. **L. rubrifolia** (Rob. et Cast. in D.C. Fl. Fr. V, p. 410) annua humilis pubescenti-viscida simplex vel basi ramosa, foliis subitus purpurascensibus parvis intus breviter petiolatis ovatis oppositis cæteris alternis sessilibus oblongis et oblongo-linearibus, racemis laxissimis paucifloris flexuosis, pedunculis folio longioribus, calycis laciniis linearispathulatis obtusis capsulâ longioribus, corollâ pubescente purpurascente, calcare recto acuto tenui corolla duplo breviore, capsulâ ovata lobis inæqualibus, seminibus oblongis costatis ad costas tuberculatis ♂. Ic. Moris Fl. Sard. tab. 101 optima. — *L. rupestris* Guss. Fl. Sic. Syn. II, p. 420.

Hab. in sabulosis et saxosis montanis sœpe inter pineta, Attica in Pentelico (Sprun! Sart!), Hymetto meridionali et insulâ Salami (Heldr!), Persia australi in colle Kuh Kasser Ali prope Gere (Ky. 92!), Belutschia inferiore ad collum Gundava (Stocks exs. sub *Anticharide Arabicid.*).

Ar. Geogr. Hispania, Africa borealis, Gallia australis, Sardinia, Sicilia.

48. **L. cryptarnm** (Boiss. et Haussk. exs. 1865) annua, patule hispida viscidula tenera exilis simplex vel basi parce ramosa, ramis ascendentibus pumilis in racemum laxum abeuntibus, foliis teneris oblongis obtusis petiolatis, floralibus superioribus ellipticis diminutis, pedicellis calyce 3-4-plo longioribus, calycis hispidi laciniis linearibus corollâ dimidiâ sublongioribus, corollâ roseâ glabra, calcare recto latinculo dimidiâ corollâ longiore, capsulâ globosâ calyce breviore, seminibus minimis longitudinaliter costulatis costis remotiusculis subanastomosantibus ♂.

Hab. in speluncis montis Beglagh Cataoniae prope Malatia (Haussk!).

Plantula 3-5-pollicaris, folia inferiora 12-9 lineas longa, pedicelli inferiores folio axillari breviores, superiores ea valde diminuta superantes, corolla cum calcare 3½-4 lineas longa. Semina dimidio minora eis *L. minoris* a quâ foliis majoribus membranaceis, calcare elongato etc. differt.

49. **L. minor** (L. Sp. p. 852 sub *Antirrhino*) annua glandulosopilosa, foliis oblongis et lanceolato-linearibus obtusis in petiolum attenuatis fere omnibus alternis, racemis laxissimis paucifloris flexuosis, pedicellis calyce 2-3-plo longioribus, calycis laciniis linearispathulatis capsulâ sublongioribus, corollæ cæruleo-violaceæ hirtæ labio superiore divaricatum bilobo, calcare conico corollâ duplo breviore, capsulâ ovata glandulosâ. seminibus oblongis apice truncatis

longitudinaliter costatis costis leviusculis ⊖. *L. minor* Desf. Atl. II, 46 — Ic. Engl. bot. tab. 2014. — Reb. Ic. tab. 61.

Hab. in arenosis et cultis præsertim montanis, Græcia in regione media Parnassi (Sprun! Heldr!), Phrygia in arvis Ouchak (Bal!), Lycia ad Elmali (Bourg!), Cappadocia circa Cæsaream (Bal!), Taurus Cilicus 6000' (Ky!), Tauria et Caucasus sept. (M.B., C.A. Mey.).

Ar. Geogr. Europa omnis a Scandinaviâ ad Rossiam medium, Africa borealis.

50. *L. pterospora*. annua, glanduloso-pilosa inferne parce et subdivaricatum ramosa, foliis alternis oblongis et lanceolatis obtusis, racemis longis laxissimis flexuosis, pedicellis calyce 2-3-plo longioribus, calycis lacinis linearispathulatis capsulam superantibus, corollæ glandulosæ cæralescenti-roseæ labio superiore sinu angusto bilobo, calcare conico dimidiâ corollâ longiore, capsula ovata glandulosâ, seminibus oblongis apice truncatis longitudinaliter multialatis alis nucleo subdupo angustioribus ⊖ *L. littoralis* var. *pterospora* Fisch. et Mey. Ind. quartus Hort. Petrop.

Hab. in Anatoliâ prope Topicha et Tokat (Wiedem!), in arenosis regionis inferioris montis Sipyli Lydiæ supra Magnesiam (Boiss!), ir. alveis Tauri Cilicii 4000' (Ky. 299! Bal!), Persia inter Kerman et Yezd (Bge! ex spec. valde incompl.).

Interdum pedalis, *L. minori* et *littorali* intermedia, a priora statu majore, ramis crassioribus, seminibus duplo majoribus aliatis, a posteriori pedicellis elongatis et seminis majoris aliis latioribus distincta.

51. *L. Persica* (Chav. Mon. p. 174) annua tota plus minusve patule pubescens pumila erecta simplex vel inferne parce ramosa, caule ramisque crassiusculis, foliis crassiusculis infimis ovatis vel oblongis subpetiolatis superioribus sessilibus ellipticis et linearibus, florib. subsessilibus distantibus paucis laxe spicatis folio calycem subrequanti suffulti, calycis tubo crispuo lanato lacinis oblongo-linearibus basi angustatis obtusis corollam æquantibus fructiferis accretis capsula duplo longioribus, corollâ puberulâ roseo-cæralescenti limbo parvo, calcare corollâ 8-10-plo breviore conico incurvo, capsula velutinâ globosâ obtusâ longe mucronatâ indehiscente localis æqualibus. seminibus oblongis truncatis inter costas longitudinales transverse et profunde foveolatis ⊖. *L. rhytidosperma* Boiss. Diagn. Ser. I, 4, p. 72. — *Ant. rhytidospermum* F. et M. Ind. II, Petrop. p. 27. — *Ant. fugax* Boiss. Diagn. Ser. II, 3, p. 460.

Hab. in agris incultis et collibus, Pisidia prope Bouldour (Heldr!), Phrygia ad Boulghas Kieui (Bal!), Armenia ad Erzerum (Caiv!) et Ardana (Noët), Syria prope Aleppo (Ky. 194! Haussk!), ditio Transeaucasica Talysch prope Swant 4000'-5000' (Hohen!), Persia (Mich.), media ad Ssertschah (Bge!) et australis prope Kaserun (Haussk!), Turkestania (Lehm.).

Facies *Hclianthemi Niloticæ*, capsulae eis *L. littoralis* crassiores, ut videtur indehiscentes prope basin irregulariter ruptiles et nixum ad *Schoenfurthiam* probentes, calcar minimum et ideo facile prætervisum.

ANTIRRHINUM (L. Gen. p. 309).

Calyx quinquepartitus laciniis subæqualibus. Corolla personata tubo basi saccato, labio superiore erecto bifido vel bipartito inferiore patente trifido, palato ample prominente faucem claudente. Stamina 4 didynama. antherarum loculi bini oblongi Capsula ovata vel pyriformis obliqua loculis binis inæqualibus polyspermis superiore pero unico inferiore poris binis pluridentatis vel unico dehiscentibus. Semina aptera. — Herbae annuae vel perennes.

SECT. II. ORONTIUM Bth in p.c. Prodr. IV, p. 290. — Semina compressa dorso levia carinata, facie interiore concava marginata margine dentato.

1. **A. Orontium** (L. Sp., p. 860) glabriusculum vel superne patule glandulosum, caule erecto simplici vel parce ramoso, foliis subpetiolatis oblongo-linearibus et lanceolatis obtusis superioribus anguste linearibus, floribus axillaribus in racemum spiciformem laxissimum dispositis, pedicellis calyce brevioribus, calycis glanduloso-hirti laciniis anguste linearibus corollam purpuream æquantibus vel superantibus, capsula ovatâ pilosâ ♂. Rehb. Ic. tab. 57.

Hab. in cultis fere totius regionis in regione inferiori a Græciâ! et Thraciâ! ad Tauriam (Ledebt!), Syriam littoralem! Egyptum!, Arabiam Mascatensem (Aueb. 5073!), Persiam australem (Ky. 93! Haussk!).

Ar. Geogr. Europa tota ab Angliâ et Sueciâ meridionali ad Rossiam medianam, Sibiria Uralensis, jugum Himalaicum, Africa borealis, Abyssinia, Insulæ Canarienses et Madera.

SECT. II. ANTIRRINASTRUM Chav. Mon. p. 81 excl. spec. — Semina obovata foveolata-cristata.

2. **A. majus** (L. Sp. p. 859) basi tortuoso-ramosum saepe lignescens, ramis annotinis elongatis rectis crebre foliosis, racemi spiciformis elongati rachide glanduloso-pilosâ, foliis oblongo-lanceolatis linearibusque utrinque attenuatis glabris, pedicellis crassis calyce brevioribus, calycis dense glanduloso-pilosi laciniis late ovatis obtusis, corollâ purpureâ amplâ calyce pluries longiore, capsula pubescenti-glandulosâ ♀.

β. angustifolium Chav. Mon. p. 86. — Folia anguste linearia margine interdum revoluta. — *A Rhodium* Boiss. in Pinard exs.

Hab. ad rupes, muros vetustos Græciae (Sibth.), insulæ Andros (Sart!), Cretæ in faucibus montis Icarî prope Candia (Heldr!), Thraciæ ad moenia Byzantii (Janka!), Cypri (Ky. 385 sub *A. Siculo!*), Syriâ littorali prope Tyrum (Gaill!) et interiori circa Rascheya (Boiss!).

Foliis angustissimis a formâ typicâ Gallicâ et Hispanicâ discrepat et in Italia australi, Sicilia, quoque crescit. *A. angustifolium* Urv. Enum. in insulâ Melitâ nec in Cycladibus occurrit et ad *A. Sicum* Ucr. referendum est.

Ar. Geogr. Europa australis et in Europâ media saepe efferatum, Africa borealis.

3. **A. tortuosum** (Bosc ex Chav. Mon. p. 88) glaberrimum basi tortuoso-ramosum saepe lignescens, ramis annotinis elongatis rectis cebre foliosis, racemii abbreviati rachiide glabrâ, foliis lanceolatis vel linearibus utrinque attenuatis, pedicellis calyce brevioribus, calyels glaberrimi laciniis ovato-oblongis obtusis, corollâ purpureâ amplâ calyce pluries longiore, capsulâ glaberrimâ 2.

Hab. ad rupes et muros, Byzantii (Boiss!), insula Chios (Orphan!), ad ruinas castelli Kalaa Beled Schekif Syrie littoralis (Gaill!).

Valde affine praecedenti a quo racemis brevioribus, glabritie, fere tantum differt. *A. Sicum* Ucr. in Archipelago erronee indicatur.

Ar. Geogr. Hispania australis, Gallia austro-orientalis, Africa borealis.

**SCHWEINFURTHIA (A. Braun Bericht. Acad. Wissensch.
Berlin 1866, p. 872).**

Calyx quinquepartitus laciniis inaequalibus, posticâ majore. Corolla personata tubo basi obscure saccato labio superiore porrecto bifido inferiore reflexo trilobo, palato bigibbo fanceum claudente. Stamina 4 fertilia didynama cum rudimento staminis quinque. antheræ biloculares post anthesin divergentes. Capsula globosa loeno superiore angustissimo oligospermo vel vacuo indehiscente, inferiore polyspermo prope basin vel irregulariter ruptili vel foramine dentibus cincto dehiscente. Semina obconica longitudinaliter alato-costata. — Herbae facie *Antirrhini* et *Linaria*.

1. **S. pterosperma** (A. Braun Bericht. Ac. Wiss. Berlin 1866 p. 872, tab. I) annua glabrata erecta pumila a basi fastigiatim ramosa, foliis carnosis oblongo-spathulatis in petiolum longe attenuatis supremis linearibus, racemis terminalibus laxis foliosis, pedicellis filiformibus folium axillare subæquantibus calyce 3-5-plto longioribus, calycis laciniis oblongo-lanceolatis superioribus præsertim supremo paulo longioribus et latioribus, corollâ albâ roseo-tintâ calyce triplo longiore basi vix saccatâ, capsulâ globosâ mucronatâ calyce longiore antice versus basin foramine dentibus cincto dehiscente, seminibus obconicis truncatis 5-6-alatis alis undulato-plicatis ②. *Orontium Aratum* Ehr. MSS. — *Ant. pterospermum* Rich. Tent. Abyss. II, p. 115.

Hab. prob. in littore Arabico Egypti superioris unde tamen eam nondum vidi.

Facies *Linaria minoris* sed multo ramosior, pedalis vel longior, folia inferiora cum petiolo 1-2-pollicaria 3-4 lineas lata, flores eis *L. minoris* vix maiores.

2. S. papilionacea (Burm. Fl. Ind. p. 121, tab. 39, fig. 2 sub *Antirrhino*) annua forsitan perennans radice indurata, glauca breviter puberula vel glabrata, caulis a collo procumbentibus vel ascendebus brevibus crassis, foliis earnosis ovatis penninerviis basi cuneatis breviter petiolatis superioribus anguste ellipticis, floribus axillaribus majusculis pedicello brevissimo suffultis, calycis demum accreti laeñia superiore aliis longiore et qua ruplo latiore cordato-ovata aenata cæteris lanceolatis inferioribus binis angustioribus, corollæ calyce sesquiloniore sordide albidæ purpureo-venosa basi fere obsolete saccata, capsula magna globosa calyce breviore longe et abrupte micronata antice ad basin loculi inferioris irregulariter ruptili, seminibus obconicis truncatis quinquealatis alis transverse sulcatis ♂. *Linaria sphaerocarpa* Bth. in p.c. Prodr. X, p. 287. — *Schur sphaerocarpa* A. Br. loc. cit. p. 875. — *Antirrhinum glaucum* Wight Ic. tab. 1459.

Hab. in regno Cabulico (Griff.), Belutschia inferiore (Stocks!).

Rami semipedales rarius pedales, folia inferiora 1-1½-pollim longa, corolla 7-8 lineas longa, capsula pisiformis major; facies *Ant. sempervirens*.

Ar. Geogr. Ditio Seinde Indiae.

TR. III. CHELONE.E Bth. in p.c. Prodr.

Corolla campanulata vel tubulosa basi non saccata nec calcarata. Antheræ uniloculares. Fructus (in nostris) capsularis valvatum dehisens. Inflorescentia saepius composita, partiali centrifugâ. Folia opposita vel superiora alterna, rarius omnia (in *Anticharide*) alterna.

SCROPHULARIA (L. Gen. 312).

Calyx quinquepartitus. Corolla tubo ventricoso-urecolato rarius breviter cylindrico, limbi bilabiati lobis quatuor superioribus erexit, inferiore patente. Stamina 4 didynama declinata antherarum loculis in unum transversum confluentibus, appendix (rudimentum staminis quinti) squamæforme ad tubi apicem inter lobos binos superiores insertum, interdum nullum. Capsula bilocularis demum septicida valvis integris vel apice bifidis, placentis polyspermis demum liberis. Semina ovoidea vel oblonga transverse rugosa. — Herbae — Generis difficilimum, folia valde variant. Stamina in speciebus pluribus manifeste inclusa vel exserta sunt, sed in aliis haec nota incerta est; duratio tandem in multis manifesta, in nonnullis, observante quoque Cl. Bentham dubia, et tamen in generis dispositione usus sum sed interdum timido animo.

Specierum Orientalium distributio.

SECT. I. CERAMANTHE Rehb. Fl. exs. I, p. 376. — *Venilia* G. Don
Gard. Dict. I, p. 507. — Appendix (staminis quinti rudimen-
tum) nulla. Corollæ lobi subæquales. — Inflorescentia
thyrsoidæ, floribus pedicellatis. Herbae foliis indivisis
rarissime basi lobulo auctis.

* Biennes. Thyrus foliatus.

S. vernalis, chrysantha, lunariaefolia, Kotschyana, cryptophila, Claussii.

** Perennes.

+ Thyrus foliatus.

S. lateriflora.

+ + Thyrus aphyllus.

S. Orientalis, nervosa, polybotrys, Boissieriana.

SECT. II. PYCNANTHUM. — Appendix nulla. Corollæ labium superius longius. — Inflorescentia capitata floribus subsessili-
bus. Herba nana foliis indivisis.

S. minima.

SECT. III. MIMULOPSIS. — Appendix linear-i-oblonga toto dorso corollæ affixa. Corollæ labium superius longius. — Inflores-
centia spicata floribus pedicellatis. Folia indivisa.

S. amplexicaulis.

SECT. IV. SCORODONIA. Don. — Bth. in p.c. Prodr. X, p. 394. —
Appendix orbicularis vel transverse latior basi tantum corollæ adfixa. Corollæ lobi superiores longiores. — Inflo-
rescentia thyrsoidæ vel paniculata pedicellis sæpissime calyce longioribus. Folia saltem inferiora opposita (in nostris) indivisa venis anastomosantibus.

* Annuæ.

S. Ilvensis, arguta, peregrina.

** Biennes.

*S. Scopolii, crenophila, divaricata, aestivalis, capillaris, sylvatica, luridiflora,
oxysepala.*

*** Perennes.

S. nodosa, chlorantha, alata, macrophylla, Cretica.

SECT. V. *TOMIOPHYLLUM* Bth. loc. cit. p. 310. — Appendix formâ varia, basi tantum corollæ adfixa. Corollæ lobi superiores longiores. — Inflorescentia paniculata, thyrsoidæ vel racemosa pedicellis sæpiissime calyciæquilonigis vel brevioribus. Folia venis non vel vix anastomosantibus, sæpius lyrata, partita, vel secta, rarius indivisa et tunc species a sectione præcedenti pedicellis brevibus dignoscendæ. — Grex a *Scorodoniam* potius habitu quam characteribus firmis distincta.

§ I. **Oppositifoliæ.** — Folia saltem inferiora opposita.

* Appendix orbiculata, semiorbiculata vel reniformis transverse latior.

+ Biennes.

S. scariosa, tagetifolia, Mesopotamica, Michoniana, lucida, sphærocarpa, rutaefolia, laxa, myriophylla, versicolor.

++ Perennes.

S. lepidota, farinosa, xylorrhiza, Taygetea, pulverulenta, catariaefolia, Mesogitana, heterophylla, laciniata, Olympica, Ruprechti, Griffithii, depauperata, subaphylla, frigida, scabiosæfolia, rostrata, trichopoda, xanthoglossa, striata, deserti, marginata.

• • Appendix ovata, oblonga vel oblongo-linearis.

+ Biennes.

S. hæmatantha, Heldreichii, pruinosa.

++ Perennes.

S. glauca, prasiifolia, Pinardi, variegata, puberula, hyssopifolia, thesioides.

• • • Appendix anguste linearis. Perennes.

S. canina, Cabulica.

§ 2. **Sparsifoliæ.** — Folia omnia sparsa.

S. crassicaulis, Syriaca, Benthamiana, hypericifolia, leucoclada.

SECT. I. CERAMANTHE

* Biennes.

1. **S. vernalis** (L. Sp. 864) molliter papillari-lanata glandulosa, caule crasso fistuloso, foliis papyraceis orbiculato-cordatis duplicatim vel inciso-dentatis inferioribus longe petiolatis superioribus diminutis

sessilibus, cymis oppositis pedunculatis thyrsum longum laxissimum foliatum formantibus, bracteis linearibus, calycis glandulosi laciini ovato-oblongis obtusis non marginatis, corollâ viridi-lutescente urceolatâ ore contractâ, antheris tandem breviter exsertis, capsulâ oblongâ acutâ glabriuseulâ ②. Ic. Engl. bot. tab. 567. — *Ceramanthe vernalis* Rehb. Germ. tab. 55.

Hab. in rupestribus umbriosis Thraciae in montibus supra Kalofer (Pichl!).

Ar. Geogr. Gallia media et orientalis, Germania, Italia, regio Danubialis, Serbia.

2. S. chrysanthæ (Janb. et Sp. III. III, p. 26, tab. 220) erispule lanato-hirsuta insuper glandulosa, caulis crassiusculis fistulosis, foliis cordato-orbiculatis argute duplicato-serratis inferioribus longe petiolatis superioribus sessilibus, cymulis subsessilibus thyrsum brevenum densiusculum ovatum vel oblongum minute foliatum formantibus, calycis laciiniis oblongo-lanceolatis acutis ciliatis, corollâ flavâ urceolatâ, staminibus longe exsertis, capsulâ oblongâ glanduloso-hirtâ obtusiusculâ mucronatâ ②. *S. vernalis* M.B. Taur. Cauc. II, p. 76 non L. — *S. minima* Bth. d.c. Prodr. X, p. 303 non M.B. — *S. congesta* Stev! Enum. Taur. p. 267.

Hab. in rupestribus umbrosis, speluncis, Armeniae Turcicæ (Auch. 5103!), montis Karakaban (Huet!), circa Gumuschkhané (Bourg!), Transcaucasie monte Karagiöl (Szov!), Somchetia et Guriæ (Ledebe!), Iberiae monte Gutgora prope Tiflis 6600' (Rupr.), ditio Talysh (C.A. M!).

S. calycina. — Thyrsus ob cymulas dissitas inferiores pedunculatas elongatas laxus, calycis laciiniæ ampliate, corolla longior tubulosurceolata. — *S. calycina* Boiss. in Bal. exs. 1866.

Hab. ad muros umbrosos Ponti Lazici ad Djimil 6000' (Bal!).

Species affinis *S. vernali* a quâ differt laciinarum calycis formâ et antheris longe exsertis. Corolla insuper paulo major et thyrsus saltem in formâ typicâ brevis confertus.

3. S. lunariæfolia (Boiss. et Bal. exs. 1866) superne sub lente sparsim et minute pruinosa-glandulosa cæterum glabra, caule elato fistuloso, foliis tenuiter membranaceis profunde cordato-orbiculatis duplicati et argute dentatis inferioribus amplis longe petiolatis superioribus sessilibus diminutis, cymis in thyrsum laxum foliatum pyramidatum dispositis inferioribus longiusculè pedunculatis, bracteis oblongis acuminatis incisis, calycis laciiniis iate lanceolatis acutis, corollâ flavâ urceolatâ, staminibus longiusculè exsertis, capsulâ ovato-oblongâ glandulosâ acutâ ②.

Hab. in præemptis maritimis Kemer Bournou Ponti Lazici inter Rhizè et Athinæ (Bal!). Fl. Jun.

Bipedalis, folia inferiora diam. 4 poll. longa et lata. An varietas *S. chrysanthæ* a quâ glabritie præsertim distinguitur?

4. S. Kotschyana (Bth. in d.c. Prodr. X, p. 303) plus mi-

nusve lanato-hirsuta glandulosa, caulis flexuosis, foliis teneris sinu aperto cuneato cordato-orbiculatis acutiuscula inciso-dentatis inferioribus petiolatis superioribus diminutis sessilibus, cymis ex omnibus fere axillis pedunculatis laxis paucifloris thyrum foliatum laxissimum formantibus, pedicellis longis, calyeis glabri laciinis ovatis obtusissimis, corollâ roseâ urceolatâ, staminibus longe exsertis, capsula glanduloso-tomentosâ ovatâ obtusa mucronata (2) *S. viscosa* Boiss. in Bourg. pl. exs. (forma laxior), — *S. Byzantina* Bth. in D.C. Prodr. X. p. 303!

Hab. ad rupestris umbrosas, fances et speluncas Tauri Cilicici (Ky. 276! 241 Balt!), ditionis Kassan Oglu (Ky. 208!), montis Begdaghi Cataoniae (Haussk!), Armeniae Turcicae in niente Almuska prope Baibut (Bourg!), specioina tandem fructifera ad Ollismereti Ineretie lecta (Rupr!) huc quoque spectare videntur. Errore schedula prope Byzantium indicata.

A *S. vernali* et *chrysanthâ* florum colore statim dignoscenda. Staminis sterilis rudimentum nullum reperi.

5. ***S. cryptophila*** (Boiss. et Heldr. Diagn. Ser. I, 12, p. 31) crispule plus minusve lanata vel hispida, caulis erectis fragilibus, foliis teneris sinu angusto cordatis ovatis obtusis obtuse et duplicatim crenato-lobatis præter superiora valde diminuta sessilia acuminata longiuscula petiolatis, cymis ex axillis superioribus pedunculatis paniculam pyramidatam formantibus, bracteis linearibus minimis, pedicellis elongatis, calycis glabri laciinis ovato-oblongis obtusiusculis, corollâ virescenti-rubellâ ovatâ, antheris longe exsertis, capsula ovatâ obtusa mucronata glaberrimâ (2) ?

Hab. in fauibus et speluncis montium Lyciae (Pest! Bourg!), Isauriae in monte Gheidagh 5500' et in eacumine montis Karadagh Lycaoniae in ruinis ecclesie veteris (Heldr!), in montibus Dykmen ad Aneyram Galatiae (Wieden!).

Affinis *S. Kotschyanae* differt paniculâ patenti saepius abbreviatâ, foliis obtuse crenatis sinu angusto cordatis, bracteis angustis, corollâ majori, capsula glabra.

6. ***S. Claussii*** (Boiss. et Buhse Enum. p. 163) præter inflorescentiam glabrescens, caule rubello, foliis membranaceis cordato-ovatis duplicatim et acute lobato-dentatis inferioribus longe superioribus brevissime petiolatis, cymis ex axillis superioribus pedunculatis paucifloris paniculam brevem pyramidatam glandulosam formantibus, pedicellis elongatis, bracteis oblongo-triangularibus acuminatis, calycis glandulosi laciinis linearis-oblongis acutiusculis, corollâ (e sieco rubellâ) urceolato-campanulatâ calyce duplo longiore, antheris exsertis, appendice nullâ, capsula ovatâ acutâ calyce duplo longiore (2) ?

Hab. in sylvis umbrosis prov. Ghilan Persiae prope Langerud (Buhse!), prope Siaret Persiae bor. orient. (Bge!). Fl. Apr.

Bipedalis, affinis *S. cryptophila* differt glabritate, calycis laciinis longioribus, et aliis notis enumeratis.

•• Perennes.

7. S. lateriflora (Trautv. Mél. Biol. tom. VI.—Pl. Radd. p. 6) ad caulim partem inferiorem foliaque subtus ad nervos pubescens cæterum glabra. caulis teretibus gracilibus elatis simplicibus vel parce ramosis, foliis omnibus oppositis glaucescentibus brevissime petiolatis a basi cordata oblongo-lanceolatis acuminatis argute et breviter serrulatis, cynis ex omnibus axillis folio æquilongis vel brevioribus breviter pedunculatis laxe multifloris trifloris, pedunculis pedicellisque glaberrimis capillaribus hisce flore multoties longioribus, bracteis brevissimis setaceis, floribus parvis sæpe clandestinis corollâ destitutis, calycis laciniis triangulari-ovatis acutiusculis fere immarginatis, corollæ brevis pallide viridulæ labiis subæquilongis, antheris breviter exsertis, appendice nullâ, capsula globoso-ovatâ abrupte acutatâ calyce triplo longiore ♀. *S. clandestina* Rupr. exs.

Hab. in rupibus Caucasi, Suania ad Muri (Radde!). Tuschetia ad Dano, ditio Tindal, Pschawia prope Gubi supra fluvium Aragwa, Daghestania prope Dido et supra Kussur 5400'-6600' (Rupr!).

Facies *Veronicae urticifoliae*, caules e radice plures pedem et amplius longi, folia 4-2 poll. longa basi 10-14 lata valde acuminata, cynæ cum pedicellis rigidulis 2-2½ poll longæ, corolla sesquilineam vix longa, capsulae piso dimidio minores, semina ut in congeneribus tranverse rugulosa et præterea longitudinaliter sulcata. Species insignis.

8. S. Orientalis (L. Sp. 864) glabra vel ad petiolos et nervos minute pruinosa, caulis elatis angulatis in thyrsum elongatum nudum glanduloso-puberulum abeuntibus, foliis petiolatis strictis oblongo-lanceolatis acuminatis argute serratis inferioribus basi cuneatâ inciso-pinnatifidis vel segmento auctis, ramorum axillarium sterilium linearibus, calycis glabri laciniis ovato-orbiculatis marginatis, corolle subglobosæ luteo-viridis labio superiore subbreviore, staminibus exsertis ♀. Ic. Jaub. et Sp. Ill. 3, tab. 221. — *S. ebulifolia* M.B. Taur. Cauc. II. p. 77.

Hab. in Armeniâ (Tournef!) ad Baibut inter duneta ad fluv. Tchoruk ripas (Bourg!), ad Ispir (Huet!), in Caucaso occid. (C.A. Mey!), Iberiâ et Caucaso orientali (M.B.), in prov. Karabagh ad Basarthal (Buhsel!).

Ar. Geogr. Sibiria Uralensis et Altaica.

9. S. nervosa (Bth. in D.C. Prodr. X, p. 303) glabriuscula vel breviter pruinosa-puberula, caulis obtuse angulatis in thyrsum nudum glandulosum abeuntibus, foliis breviter petiolatis nervosis oblongo-lanceolatis subtiliter serrulatis vel subintegris, calycis plus minus glandulosi laciniis ovatis anguste marginatis, corolle subglobosæ livide violaceæ labio superiore subbreviore, staminibus exsertis ♀. *S. Olivieri* J. et Sp. Ill. 3 tab. 222 non Wydl.

Hab. in Persia monte Elvend prope Hamadan (Oliv., Auch. 280!) in valle

Liwan (Buhsel), in prov. Aderbidjan (Auch. 5059!), in rupestribus montium Avroman et Schahu Kurdistanae Persiae 7-800' (Haussk!).

Affinis *S. Orientali* differt foliis subintegris, floribus minoribus, calyce glanduloso.

10. *S. polybotrys* (Boiss. Diagn. Ser. I. 12, p. 32) caulis glabris elatis teretibus glabris in thyrum aphyllum glandulosum abeuntibus, foliis minute et parce pruinoso-puberulis oblongis et oblongo-lanceolatis acuminatis inferioribus breviter petiolatis argute denticulatis superioribus sessilibus integris, calycis fere eglandulosi laciniis ovato-oblongis anguste marginatis, corollæ subglobosæ violaceæ labio superiore subbreviore, staminibus exsertis ♀.

Hab. ad radices montis Demavend Persiae borealis prope Lar. (Ky. 315!).

Valde affinis *S. nervosæ* et an ab eâ satis distincta?, differt foliis non elevatim nervosis latioribus, inferioribus argute dentatis; calyces minuti *S. nervosæ*.

11. *S. Boissieriana* (Jaub. et Sp. Ill. Or. III, p. 29, tab. 223) caulis glabris virgatis obsolete angulatis in thyrum nudum glandulosum abeuntibus, foliis glabriusculis lucidis obsolete venosis subsessilibus inferioribus anguste lanceolatis parte inferiori argute denticulatis superioribus et ramorum axillarium sessilium anguste linearibus integris, calycis glandulosi laciniis ovatis obtusis anguste marginatis, corollæ subglobosæ luteo-viridis labio superiore subbreviore, staminibus exsertis ♀. *S. Orientalis* Boiss. in Ky. exs. non L.

Hab. in montibus Kuh Barfi et Kuh Delu Persiae australis ad nives (Ky. 529! et 330!).

Praecedentibus affinis foliorum formâ differre videtur. Flores minuti *S. nervosæ*.

SECT. II. PYCNANTHIUM.

12. *S. minima* (M.B. Taur. Cauc. II, p. 79) perennis glabra viscidula. caudiculis subterraneis fragilibus plus minus elongatis nudis, caulis nanis folia pauca capitulumque ovatum densiflorum foliis superatum edentibus, foliis longe petiolatis a basi cuneatâ rotundatâ vel truncatâ ovatis et oblongis simpliciter vel subduplicatim et obtusiuscule crenato-denticulatis, bracteis anguste lanceolato-linearibus scariosis calyce longioribus, floribus subsessilibus, calycis scariosi laciniis ovatis undulato-crispis erosulis, corollâ roseo-purpureâ tubulosâ calyce 2 1/2-plo longiore labio superiore longiore lobis rotundatis, capsulis ovatis abrupte acutis glabris striatis ♀.

Hab. in declivibus schistosis mobilibus regionis alpinæ Caucaasi, alpes Kasbek Caucaasi centralis (Adams), ditiones Tindal, Boghos, Tuschetia, Daghestania Caucaasi orientalis alt 8000'-10500' (Rupr!).

Species insignis facie a congeneribus aliena; caules 1-3-pollicares, foliorum majorum lamina 12-16 lineas longa petiolo apicilongo. capitulum unicum

(rarius bina breviter pedunculata) etiam fructu maturo densum 18-15 lineas longum diam. 9-10 latum, corolla 6 lineas longa, capsula piso dimidio minor. Appendix staminea nulla sed florem staminibus quinis fertilibus donatum semel observavi.

SECT. III. MIMULOPSIS.

13. S. amplexicaulis (Bth. in v.c. Prodr. X, p. 310) tota breviter pubescenti-viscidula canescens, caulis pluribus strictis teretinsculis simplicibus vel inferne parce ramosis, foliis sessilibus flabellatim plurinerviis angute et cibre denticulatis intimis minoribus obovato-oblongis obtusis basi breviter attenuatis, cæteris ovato-oblongis acutis vel acuminate basi cordato-plexicaulibus, cymis confertim 1-5-floris subsessilibus distantibus thyrsos terminales spiciformes aphylos laxos formantibus, bracteolis minimis lanceolato-setaceis membranaceis, pedicellis strictis calyci aequilongis vel brevioribus, calyeis glabri laciñis ovatis margine late membranaceis, corollæ flavidò-virentis ureolato-campanulatae calyce triplo longioris labio superiore longiore, staminibus inclusis, appendice oblongo-lineari subquadratâ sulcato-cymbiformi toto dorso corolle affixâ, capsulâ elongato-conicâ acutissimâ calyce sextuplo longiore ♀.

Hab. in regione alpinâ superiore ad nives Persiae, mons Ssavalan prov. Ghilan (Auch 5060!), montes Ssahend et Demavend (Buhset!), montes Sawers, Nur, Daëna Persiae australis 10-13000' (Haussk! Ky. 654!), montes Avroman et Schahu Kurdistanie Persiae (Hauss⁴).

Planta mucilaginosa 1-2-pedalis, folia intermedia 1¹/₂-2-pollicaria, spicæ laxe paucifloræ 6-9 poll. longæ vel breviores, corolla 4 lineas, capsula 5 lineas longa. Species foliis amplexicaulibus *Mimuli* vel *Centranthi*, capsulaque elongata insignis

SECT. IV. SCORODONIA.

* Annuæ.

14. S. Ilwensis (C. Koch Linn. XVII, p. 284) præter inflorescentiam glabra, caule simplici in thyrsum inferne foliatum vel aphyllum strictum longum abeante, foliis parvis subtus elevatim nervosis ovato-oblongis basi rotundatis vel cuneatis inferioribus confertis breviter petiolatis subdupliciter dentatis superioribus diminutis sessilibus oblongis acutis acutissime incisis interdum auriculato-semi-amplexicaulibus, cymis paucifloris breviter pedunculatis, bracteis minutis setaceis pedicellisque calyce brevioribus glandulosis, calyeis glabri laciñis orbiculatis late membranaceo-marginatis, corollâ purpureâ calyce duplo longiore, staminibus tandem breviter exsertis, appendice reniformi, capsulâ magnâ ovatâ acutâ calyce sesqui vel duplo longiore ♂. *S. Galverti* Boiss. Diagn. Ser. II, 3, p. 452.

Hab. in Sonchetie montibus Ilwensis prope Abotz (C. Koch!), Ponto Lazico supra Djimil 7500' (Bal!), Armenia Turcia circa Erzerum et Tortum

(Calv! Huett!), in incultis prope Baibut et rupestribus prope Gummisch Khané (Bourg!). Fl. Jun., Jul.

Folia radicaria rosulata 1-2-polices longa, caulis cum thyro saepe pedali 1-2-pedalis interdum antem semipedalis pauciflorus, unicus vel e collo 3-4 centralis elatior crassus, flores in cymā brevissime pedicellati glomerati, capsula pisum minutum aequans, corolla magna longitudini subaequilata. Species pulchra distinctissima.

15. *S. arguta* (Soland. in Ait. Kew Ed. I, Vol. 2, p. 242) plus minusve pubescens, caule simplici vel parce et stricte ramoso, foliis membranaceis petiolatis late ovatis subcordatis vel cuneatis argute et duplicitam dentatis praeter infima alternis; cymis axillaribus 2-5-floris breviter pedunculatis thyrum strictum foliosum interdum subaphyllum formantibus, pedicellis calyce vix longioribus bracteisque setaceis brevibus glandulosis, calycis glandulosi lacinii ovatis obtusis submarginatis, corollā rubrā urceolatā calycem parum exceedinge, staminibus inclusis, appendice orbiculari-obcordata, capsula glabrā a basi rotundatā longe rostrato-attennatā ⊖. Ic. Webb Fl. Canar. tab. 177. — *S. rostrata* Hochst. in Schinap. exs.

Hab. in regno Mascate Arabiae secūs torrentes (Auch. 5057!).

A *S. peregrina* capsulae formā, floribus duplo minoribus, etc. distinctissima.

Ar. Geogr. Insulae Canarienses, Regnum Maroccanum et Tunetanum, Nubia, Abyssinia.

16. *S. peregrina* (L. Sp. p. 866) glabra, caule subsimplici, foliis membranaceis petiolatis grosse dentatis cordato-ovatis superiores alternis sensim diminutis busi truncaitis sessilibus, cymis axillaribus pedunculatis laxe 2-5-floris paniculam longam laxam foliosam formantibus, pedicellis calyce 2-3-plo longioribus, bracteis setaceis, calycis glabri lacinii late et breviter lanceolatis acutis non marginatis, corollā sanguineā breviter campanulatā labio superiore multo longiore, staminibus inclusis, appendice orbiculari-ovatā, capsula glabrā obtusā mueronatā ⊖. Ic. Fl. Gr. tab. 597. — Rehb. Germ. tab. 55.

Hab. ad vias, sepes, muros regionis inferioris, Græcia et ejus insulæ ubique!, Creta (Raul!), Macedonia (Griseb.), Anatolia occidentalis et australis (Bourg! (Bal!)), Syria littoralis (Ehr! Bl! Gaill!).

Ar. Geogr. Regio mediterranea totius Europæ et Africæ borealis.

* * Biennes.

17. *S. Scopolii* (Hoppe Cent. exs. — Pers. Syn. II, p. 160) caule obtuse quadrangulo, foliis punctatis ovato-oblongis cordatis petiolatis plus minusve pubescentibus obtuse vel acute crenato-dentatis dentibus apiculatis, thyro oblongo vel elongato laxe aphylo vel basi subfoliato, bracteis linear-lanceolatis, pedicellis glandulosis calyce subdupo longioribus, calycis glabri lacinii ovato-orbiculatis

late albo-marginatis, corollâ virenti-purpurascenti, staminibus inclusis, appendice transverse latiore, capsulâ ovato-globosâ mucronatâ calyce subdupo longiore sœpe retusa (2). Ic. Rchb. Germ. tab. 54. — *S. glandulosa* W. K. III, p. 238, tab. 214. — *S. auriculata* Scop. non L. — *S. betonicæfolia* Wydl. Mon. — C. Koch Linnæa. — *S. Scorodonia* Host Austr. II, p. 214. — Griseb. Spic. II, p. 37 et Ledeb. Fl. Ross! non L. — *S. Castagneana* Wydl. Mon. p. 149. — *S. glandulifera* Clarke Trav. I, p. 660 ex diagnosi et loco. — *S. melissæfolia* Griseb. Spicil! 17, p. 37.

Hab. in sylvaticis a regione submontanâ et collinâ ad alpinam, Macedonia et Thracia (Griseb., Friv!), Byzantii (Cast! Noël), Bithynia in Olympo (Clem! Pichl!), Rhodus (Cadett!), Anatolia borealis et Pontus (Wied! Bourg!), Pontus Lazicus ad Djimit 6000' (Bal!), mons Amanus Syriae (Ky. 244!), Caucasus omnis (M.B., Rupr!), Tauria (Stev.), ditio Talysh (C.A. M.), Persia bor. (Bge! Haussk!).

$\beta.$ *grandidentata*. — Foliorum sœpe acutiorum et magis oblongo-ramum dentos acutiores, appendix non retusa. — *S. grandidentata* Ten. Fl. Nap. Suppl. II, p. 69.

Hab. in Graecia monte Malevo (Orph!), Velugo et Delphi Eubeæ (Heldr. 2744!) et herb. Norm. 683!, monte Korthiati Macedoniae (Heldr!).

$\gamma.$ *oligantha*. — Formæ præcedenti similis sed folia minora, planta humilior, thyrsus depauperatus. Forma alpina. — *S. oligantha* Boiss. et Heldr. Diagn. Ser. I, 12, p. 33. — *S. Galatica minor melissæfolio* Tourn. herb! et *S. melissæfolia* Urv. Mem. Soc. Linn. I, p. 331.

Hab. in summis Taygeti (Heldr!), in Ponto (Tournef!).

$\delta.$ *Smyrnæa*. — Folia profunde inciso-lobata lobis acute dentatis. — *S. Smyrnæa* Boiss. Diagn. Ser. I, 4, p. 66.

Hab. ad rivulos montium supra Bournabat prope Smyrnam (Boiss!).

$\varepsilon.$ *Tmolea*. — Pilosior, glandulosior, glaucescenti-canescens, calyx augustius marginatus. — *S. Tmolea* Boiss. Diagn. Ser. I, 4, p. 66. — *S. decumbens* F. et M. Ann. Sc. Nat. 3^{me} Ser. V, p. 180! — *S. fontana* Ky. pl. exs. 1859. — *S. puberula* Boiss. et Haussk. MSS.

Hab. in pratis humidis montis Tmoli Lydiæ ad Bozdagh (Boiss!), prope Tokat (Wied!), in monte Tchoschdag Cilicie orientalis (Ky. 90!), Berytdagh Cataonia 7000' (Haussk!).

Species indumento et foliis polymorpha, speciminum major copia nexus inter omnes formas præbuit.

Ar. Geogr. Germania orientalis, Italia media et australis, regio Danubialis, Rossia media et australis.

18. *S. crenophila* (Boiss. Diagn. Ser. I, 7, p. 41) tota crispule papilloso-hirsuta viscidula cinerasceus, caulinibus obtuse quadrangulis superne ramosis, foliis membranaceis ovato-oblongis obtusis obtuse dentatis inferioribus longiusculæ petiolatis basi cordatis superioribus sensim diminatis breviter petiolatis, floralibus trianguli-oblongis

acutis, paniculâ terminali laxâ foliolosâ glandulosâ, bracteis linearibus minimis, pedicellis calyce 3-4-plo longioribus divaricatis, calycis glabri laciniis orbiculatis anguste membranaceis, corollâ luridâ brevi, staminibus inclusis, appendice transverse latiore basi truncatâ apice subretusâ, capsulâ ovalâ acutâ calyce subdupo longiore ②.

Hab. in humidis montium Zagros, Avroman et Schahu Kurdistanie Persiae 5000'-8000' (Haussk!), montibus Sawers et Kuh Daëna Persiae australis 8000' (Haussk! Ky. 632! forma minus papillosa).

Bipedalis et elatior, folia *Melisse officinalis*, differt a *S. Scopolii* indumento papillari denso criso, inflorescentiae paniculatæ pedicellis elongatis, calyce angustius marginato. *S. divaricata* indumento et pedicellis longis divaricatis affinis differt cymis axillaribus.

19. ***S. divaricata*** (Ledeb. Fl. Alt. II, p. 240) secus caulem, petiolos, pedunculos crispule villosula, caule elato obtusangulo, foliis petiolatis membranaceis undulatis amplis pubescenti-papilloso cordato-ovatis acutis acete lobatis et dentatis, superioribus parum diuinutis oblongis acuminatis, cymis ex omnibus axillis divaricatum 3-multifloris, pedicellis calyce 3-4-plo longioribus, calycis glaberrimi laciniis ovatis obtusis margine membranaceis, corollâ fusco-viranti, staminibus inclusis, appendice reniformi transverse latiore, capsulâ majusculâ ovatâ glabrâ mucronatâ calyce 2 1/2-plo longiore ②. Ic. Ledeb. tab. 121. — *S. Georgica* Bth. in D. C. Prodr. X, pag. 305!

Hab. in sylvis Caucasi borealis ad Stavropol (Owerin!), centralis ad Kaischaur (Ovi!) et Darial (Rupr!), Suania (Radd!), Iberia prope Titlin (Ledeb.), in monte Sarial et prope Schuscha (Hoh!).

A *S. Scopolii* formis foliis membranaceis ea *Momordica claterii* interdum referentibus cymisque axillaribus divaricatis folio brevioribus optime distincta.

20. ***S. aestivalis*** (Griseb. Spic. II, p. 76) pallide virens viscidula pubescens, caulinis erectis simplicibus, foliis petiolatis a basi rotundatâ vel truncatâ oblongis inaequaliter duplicato-serratis, superioribus non diuinutis, cymis ad axillas supremas paucis 3-5-floris, pedunculis tenuibus petiolo sublongioribus, pedicellis tenuissimis corollæ subaequilongis, bracteis subulatis minimis, calycis parce et breviter glandulosi laciniis oblongis obtusis vix marginatis, corollâ flavidovirente, appendice parvâ obtiangulari obtusissimâ, staminibus vix exsertis, capsulâ . . . ȝ.

Hab. in monte Seardo Macedonia inter Kalkandele et montem Kobelitzu (Griseb!). Vid. in herb. cl. Griseb.

Caulis 1-1 1/2-pedalis, folia 1 1/2-2-pollices longa, facies *S. vernalis* a qua ceteris neglectis appendice aliena est, ex specimine juniori incomplete nota, ob folia floralia non diuinuta prope *S. divaricata* Ledeb. collocanda, hæc notâ, foliis non cordatis, a *S. sylvatica* sat differre videtur.

21. ***S. capillaris*** (Boiss. et Bal.) glabriuscula, caule elato acute

quadrangulo, foliis petiolatis cordatis ovatis obtusis obtuse et duplicitim crenato-lebatis crenis apiculatis, thyrsso longissimo laxo minute foliato, foliis cuneato-oblongis acutis acute dentatis, pedicellis capillaribus divaricatis calyce multolies longioribus minute et parce glandulosis, calycis glaberrimi laciniis orbiculatis late membranaceis. corollæ lurido-flavescenti, staminibus inclusis, appendice transverse reniformi retusa, capsula globosa magnâ apiculata calyce triplo longiore ②.

Hab. in regione inferiori Ponti Lazici prope Rhizé (Bal. exs! sub *S. divaricata* var.).

Folia inferiora 3 poll. longa 2½ lata, thyrsus pedem et amplius longus cymis breviter pedunculatis divaricatissime 5 multifloris constans. Species a *S. divaricata* caule acutangulo, glabritie, pedicellis tenuissimis saepe pollicem longis distinctissima. Corolla fere magnitudinis *S. Scopolii*.

22. ***S. sylvatica*** (Boiss. et Heldr. Diagn. Ser. II, 3, p. 153) viridis glanduloso-viscosa, caule elato ramoso obtuse quadrangulo, foliis petiolatis amplis membranaceis ovatis basi profunde cordatis grosse et obtusiuscula lobatis lobis parce dentatis, cymis breviter pedunculatis paniculas thyrsoides foliatas formantibus, foliis floralibus diminutis subsessilibus triangularibus acutis acute incisis, bracteis linearibus integris, pedicellis calyci subaequilongis fructiferis ramosis, calycis glanduloso-papillaris laciniis ovatis vix marginatis, corollæ flavidio-virenti, staminibus inclusis filamentis glandulosis, appendice obtriangulari minutâ, capsula ovata acutissimâ calyce sub-duplo longiore ②.

Hab. in regione sylvaticâ Olympi Thessali inter Mavroslonghos et Hagios Dionysios 5000'-7000' (Heldr. 2503!, Orph. Fl. Gr. exs. 730!).

Folia inferiora saepe 4-5-pollices longa. Species pulchra a *Sc. Scopolii* et *divaricata* caule ramoso, calyce hirsuto non marginato, pedicellis fructiferis recurvis, etc. discedens.

23. ***S. luridiflora*** (F. et M. Ind. VI, Hort. Petrop. p. 40.) molliter arachnoideo-lanuginosa, caule elato simplici obtuse angulato, foliis breviter petiolatis a basi truncata vel subcordata ovato-oblongis obtusis obtuse et subduplicatim crenatis, thyrsso terminali parte inferiori folioso cymis breviter et crassiuscula pedunculatis 5-7-floris constante, bracteis setaceis pedicellisque calyce vix longioribus hirsutissimis, calycis lanuginosi laciniis oblongis vix marginatis, corollæ brevi lurida, staminibus inclusis, appendice obcordata transverse latiore, capsula ovata acuta calyce vix sesquiloniore ② ?

Hab. in Anatoliâ boreali prope Tosia et Mersiwan (Wied!).

Species distinctissima a *S. Scopolii* formis indumento lanato, thyrsso inferne foliato, calyce lanuginoso, a *S. sylvaticâ* indumento, caule simplici, ab utrâque calyce ample corollâ et capsula vix breviore discedens.

24. ***S. oxysepala***, papilloso-lanuginosa glaucescens, caule obtuse angulato, foliis petiolatis a basi truncata vel cuneata ovatis vel

oblongis acentis acutiuscule lobulato-dentatis superioribus sensim diminutis sessilibus argute dentatis, cymis ex omnibus axillis pedunculatis 5-7-floris, pedicellis calyce duplo longioribus bracteisque linear-setaceis glanduloso-hispidis, calycis hirsuti laciniis obtongis acuminatis membranaceo-marginatis, corolla lurida labio superiore vix longiore, staminibus inclusis, appendice parvâ suborbiculariâ, capsulâ ovatâ acutissimâ calyce subdupo longiore (2).

Hab. in monte Ssavalan Persiae borealis (Auch. 5058!).

Ad *S. divaricatum* Ledeb. in D.C. Prodr. relata sed ab eis foliis non cordatis, pedicellis brevibus, et in primis calycis hirsuti laciniis cuspidatis omnino aliena et inter affines distinctissima.

*** Perennes

25. ***S. nodosa*** (L. Spec. p. 863) glabriuscula, radice tuberoso-nodosâ, caule elato argute quadrangulo, foliis petiolatis ovato-oblongis basi cordatis et subcordatis subduplicatim et argute serratis superioribus sessilibus lanceolatis, paniculâ aphyllâ vel inferne foliis linearibus obsitâ, calycis laciniis ovatis obtusis obsolete marginatis, corollâ brunneo-virenti, staminibus inclusis, appendice transverse latiore truncaâ vel retusa, capsulâ ovatâ acutâ calyce 3-4-plo longiore. ♀. Ic. Rehb. Germ. tab. 53. — *S. Hemschinica* C. Koch Linn. XXII, p. 558. — *S. macrobotrys* Led. Fl. Ross. III, p. 217!

Hab. in sylvaticis humidis ad rivos in Peloponneso (ex Sibth.), Ponto Lazico ad Rhizé (Balt!), Guriâ (Nordn!), Tauriâ (Stev!), Iberiâ (M.B.), ditione Talyesch (C.A. M.).

Ar. Geogr. Europa septentrionalis et media ad Hispaniam et Italiam centrales, Dalmatiâ et Rossiam australem usque, Sibiria Uralensis et Altaica, America borealis.

26. ***S. chlorantha*** (Ky. et Boiss. in Pl. exs. 1859) pubescens, caule elato crasso obtuse angulato, foliis amplis longe petiolatis ovatis profunde cordatis acuminatis acutiuscule dentatis, cymis pedunculatis paniculam pyramidatam inferne foliosam formantibus, foliis floralibus valde diminutis oblongo-lanceolatis longe acuminatis, bracteis setaceis pedicellisque calyce 2-3-plo longioribus valde glandulosis, calycis parvi pruinoso-glandulosi laciniis ovatis obtusis anguste marginatis, corolla lurida, staminibus inclusis, appendice parvâ obtiangulari transverse sublatiore, capsulis parvis ovatis acutis calyce triplo longioribus ♀.

Hab. in arenosis ad fontes pagi Goschkar prov. Musch Armenia-Turcicæ alt. 500' (Ky. 346!). Fl. Aug.

Folia *Salviae glutinosæ*, inferiora 4-5-pollices longa. Affinis *S. nodosa* differt pubescentiâ, caule obtusangulo, foliis profunde cordatis, calyce glandulosopubescenti, capsulâ duplo minore.

S. alata 27. (Gilib. Fl. Lithuan. II, p. 127) glabriuscula, radice fibrosâ, caule quadrangulo cum petiolis alato, foliis petiolatis ovato-

oblongis basi rotundatis vel obscure cordatis serratis, paniculâ elongatâ subaphyllâ, pedicellis calyce longioribus parce glandulosis, calycis glabri lacinii orbiculatis late membranaceis, corollâ viridi-brunneâ, staminibus inclusis, appendice transverse multo latiore obocordato-bilobâ lobis diivaricatis, capsulâ subglobosâ acutâ ♀. Ic. Rehb. Germ. tab. 51. — *S. Ehrharti* Stev. ex Bab. Man. 218. — *S. Neesii* Wirtg. — *S. cymigera* C. Koch Linn. XIX, p. 22 nec Linn. XXII. — *S. Samaranitiae* Boiss. et Heldr. MSS. — *S. aquatica* Koch Syn. El. I. et Auct. plur. non L.

Hab. ad aquas præsertim montanas, Græcia in montibus Malevo (Orph!), Parnasso (Heldr! Orph. Fl. exs. 443!), Thraciâ (Urv.), Bithyniâ (Thirke!), Anatoliâ boreali (Wied!), Armeniâ (Bourg!), regione Caucasicâ (Ledeb.), Persiâ (Buhse!), regno Cabulico (Griff!).

β. cordata. — Caules angustins alati, folia inferiora cordata, flores sæpius minores, appendix vel ut in formâ typicâ biloba vel tantum retusa. — *S. Pisidica* Boiss. et Heldr. Diagn. Ser. I, 12, p. 33. — *S. cymigera* C. Koch Linn. XXII, p. 557 non Linn. XIX.

Hab. in Pisidiâ ad Aglansun et Isauriâ ad lacum Beychehr (Heldr!), Lyciâ prope Elmalu (Bourg. 174!), Armeniâ Rossicâ (C. Koch!), Mesopotamiâ et Syriâ boreali ad Orfa, Aintab, Marasch (Haussk!), Libâno (Ky. 763!, Bl!), ad Hasrun, Mar serkis, Bescharré, Cœlesyriâ ad Heliopolin (Ehrenb!).

Species foliis acute vel obtusius dentatis basi rotundatis vel cordatis, florum et capsularum magnitudine et etiam appendicis (tamen semper transverse latoris) formâ valde polymorpha, a *S. aquaticâ* L. = *Balbisii* Horn. in Oriente non observatâ non tantum appendix sed inflorescentiâ paniculatâ nec anguste thyrsoidæ discedens.

Ar. Geogr. Europa præsertim borealis et media, Sibiria Altaica.

28. *S. macrophylla* (Boiss. Diagn. Ser. I, 12, p. 32) glabra, radice fibrosâ, caule elato acute quadrangulo fere aptero simplici vel parce ramoso, foliis amplis petiolatis basi cordatis grosse crenato-dentatis et prope basin interdum sublobatis, cymis brevissimis pendunculatis multifloris oppositis remotiusculis thyrso longos angustos interruptos aphyllos formantibus, pedicellis calyce longioribus glabriusculis, calycis glabri lacinii ovatis acutis albo-marginatis, corollâ luridâ, appendix transverse semilunari rotundatâ, capsulis ovatis acutis ♀. *S. Balbisii* var *macrophylla* Boiss. Diagn. Ser. II, 3, p. 154.

Hab. ad aquas Syriæ ad Tripolin (Bl!), ad Sidonem, Tortosam et circa Damascum (Gaill!), in Palestinâ ad lacum Merom et prope Jericho (Boiss!).

Planta pluripedalis, folia inferiora sœpe 6-8-pollices longa. Afinis *S. alata* var. *cordata* a quâ differt inflorescentiâ thyrsoidæ pseudoverticillatâ nec paniculatâ, appendicisque formâ. Floribus thyrsoides refert *S. aquaticam* L. = *S. Balbisii* Hornem. ad quam eam olim retuli, sed haec in Europâ occidentali ad Italiam usque tantum obvia omnino differt petiolis alatis, foliis obtuse crenatis, corollâ brunneâ, appendix transverse non latiori ovata vel rotundatâ.

29. *S. Cretica* (Boiss. et Heldr.) caulibus acute angulatis apte-

ris. foliis utrinque sed subtus magis pubescentibus petiolatis cordatis oblongis obtusis obtuse dentatis, floralibus minimis acuminatis, cymis breviter pedunculatis remotis thyrsum longum valde interrumpum formantibus, bracteis linearibus pedicellisque strictis calyce longioribus glandulosis, calycis papilloso-pubescentis laciniis ovatis obtusis membranaceo-marginatis, corollâ fuscâ, staminibus inclusis, appendice ovato-orbiculari, capsulâ globosâ mucronulatâ calyce vix sesquilociore ♀.

Hab. in fossis subhumidis circa hortos ad Cydoniam Cretæ (Heldr. exs. sub *S. auriculata*).

Caules pedem et amplius elati, thyrsus pedem sœpe longus, folia inferiora pollices duo non excedentia, flores magnit. *S. aquatica* L. = *S. Balbisii* Horn. a quā differt caule aptero, foliis calyceque pubescentibus. *S. auriculata* L. differt foliis lyratis, pedicellis brevissimis, calyce glabro.

SECT. V. TOMIOPHYLLUM.

§ 1. Oppositifoliæ.

* Appendix orbiculata, semiorbiculata, vel reniformis transverse latior.

+ Biennes.

30. ***S. scariosa*** (Boiss. Diagn. Ser. I, 4, p. 67) glaberrima, caule elato simplici folioso in thyrsum elongatum angustum abeunti, foliis amplis in laciniis anguste lanceolatas acutas acute incisas bipinnatisectis, foliorum superiorum minorum sessilium laciniis angustioribus, cynis brevibus bifidis 5-7-floris, pedicellis glandulosis calyce subbrevioribus, bracteis et bracteolis ovato-oblongis latissime albo-scariosis, calycis glabri laciniis ovatis membranâ albo-scariosâ eis latiore undulato-bullatâ marginatis, corollâ atropurpureâ, antheris inclusis, appendice orbiculatâ, capsulâ ovato conicâ longe attenuato-acuminatâ calyce duplo longiore ②.

Hab. in Syriâ boreali inter segetes, ad Aleppo (Montbret., Haussk!), ad Aintab (Auch. 1765!).

Planta 2-3-pedalis thyrsi pedem et amplius lengo, folia *Datisca canina*, flores magni. Species membranâ albâ bullatâ calycem marginante insignis et pulcherrima.

31. ***S. tagetifolia*** (Boiss. et Haussk. pl. exs. 1865) inferne glaberrima, caule elato subangulato simplici et ramoso in thyrsum longum laxum abeunte, foliis in laciniis anguste et breviter lanceolatas argute inciso-dentatas basi attenuatas vel petiolulatas pinnatim et bipinnatim sectis rachide interdum lobuliferâ, cymis longiuscule pedunculatis bifidis laxe 7-15 floris, pedicellis glanduliferis calyce 3-4 plo longioribus, bracteolis squamiformibus non scariosis, calycis

glabri laciniis orbiculatis membranâ scariosâ albâ denticulatâ eis angustiore marginatis, corollâ purpureâ, staminibus inclusis, appendice reniformi, capsulis parvis globosis mucronatis calyce vix longioribus ȝ.

Hab. ad segetes Syriæ borealis ad Samandara inter Aleppo et Aintab (Haussk!).

Facie et foliorum laciniis acutis affinis *S. scariosæ* differt cymis laxifloris, bracteis et bracteolis herbaceis, calyce anguste nec latissime albo-scarioso, capsulis dimidio minoribus calyce vix longioribus obtusis breviter mucronatis.

32. S. Mesopotamica (Boiss. Diagn. Ser. I, 12, p. 34) glabra superne parce et minute glandulosa, caule elato obtuse angulato in thyrum longum angustum interrupte spiciformem abeunte, foliis in laciniis longas late linear-lanceolatas argute dentatas pinnatum vel subbipinnatum sectis, cymis brevissime pedunculatis confertim 7-9-floris, pedicellis calyce brevioribus, bracteis et bracteolis oblongo-spathulatis late membranâ rufâ crispulâ eis subaequilatâ marginatis, corollâ purpureâ, staminibus inclusis, appendice ovato-rotundâ, capsula breviter ovato-conicâ acutâ calyce duplo longiore ②.

Hab. in Mesopotamiâ inter Orfa et Suerek (Ky. 57!).

Bi vel tripedalis, habitus et folia *S. scariosæ* a quâ præter foliorum laciniis sublatiores differt cymis brevius pedunculatis densius floriferis, bracteis et calyce membranâ non niveâ sed rufo-purpureâ et dimidio angustiore marginatis, capsulis minoribus acutis nec in rostrum subaequilongum attenuatis.

33. S. Michoniana (Coll. et Kral. in Cat. pl. Palest. p. 13) præter inflorescentiam glaberrima, caule elato crasso rubro angulato simplici vel inferne parce ramoso in thyrum laxum angustum longissimum abeunte, foliis amplis ambitu ovato-oblongis pinnatisectis segmentis ovato-oblongis obtuse dentatis vel lyratis inferioribus petiolulatis ad basin petioluli lacinulâ ovatâ sæpe auctis, superioribus secus rachidem decurrentibus, cymis brevissime pedunculatis glandulosis confertim 6-7-floris, bracteis linearibus longis bracteolisque setaceis herbaceis non marginatis, pedicellis calyce brevioribus, calycis glabri laciniis ovatis membranâ rubrâ æquilitâ valde undulato-crispâ marginatis, corollâ flavescens labio superiore multo majore, staminibus inclusis, appendice parvâ orbiculatâ, capsula subsplenicâ abrupte et longinsecule mucronata calyce duplo longiore ②. *S. rubricaulis* Boiss. Diagn. Ser. I, 12, p. 34.

Hab. in Galileâ ad Jenin et inter Banias et Tiberiadem atque Colesyriâ inter Hasbeya et Rascheya (Boiss!), in Libano ad Ghasir et Syriâ littorali ad Sidonem in alveis et ad sepes (Gaill!), inter Tripoli et Hama (Bl!), in Pales-tinâ ad Hierosolymam (Roth).

S. tenuisecta. — Foliorum segmenta et laciniæ diminutæ, folia magis decomposita, thyrsus abbreviatus. — *S. Hierochuntina* Boiss. loc. cit. p. 35.

Hab. in Palestinâ ad meridiem Gaza et circa Jericho in humidis (Boiss!), in Colesyriâ inter Merdj et Leitani (Gaill!).

Tri-quadripedalis, thyrsus in formâ typicâ saepe bipedalis, flores et capsulae majusculi. Affinis *S. Mesopotamicae* a quâ egregie differt foliorum dissectione, bracteolis non scariosis.

34 S. lucida (L. Sp. p. 865) glaberrima lucida, caule obtuse tetragono in thyrum longum minute glandulosum abeunte, foliis crassisculis imis parvis ovatis lobatis cæteris bis pinnatisectis segmentis oblongis inciso-dentatis, foliorum superiorum angustioribus acutis, cymis bifidis 3-7-floris floribus brevissime pedicellatis, bracteis linearibus brevissimis, calycis laciniis orbiculatis latiuscule scarioso -marginatis, corollæ lobis superioribus majoribus orbiculatis atropurpureis, staminibus majoribus exsertis, appendice late semi-orbiculari, capsulis majusculis globosis abrupte acutatis vel mucronatis calyce duplo longioribus (2).

α. genuina. — Humilior, foliorum laciniæ interdum angustiores Guss. Pl. rar. t. 43. — *S. glauca* Fl. Græc. VI, p. 78, tab. 599.

Hab. in rupestribus Cycladum (Sibth., Urv!), Tenos et Naxos (Wied!).

Hæc est forma in Italiâ circa Hydruntum crescens et a Limneo citata.

β. filicifolia. — Major 2-3-pedalis, foliorum laciniæ latiores. — *S. filicifolia* Fl. Græc. VI, pl. 79, tab. 600. — *S. glauco folio in amplas laciniias d'viso* Tourn. Voy. I, tab. 221.

Hab. ad rupes, muros Græciæ in Hymetto, Parnes et Pentelico (Sprunt Boiss! Heldr!), Parnassi in regione mediâ (Guic!), Cretæ (Sieb!), in monte Idæ et ad Neochori (Heldr!), insulis Andros, Hydra et Naxos (Heldr!), Rhodi (Bourg!), Asiæ minoris (Auch. 2011), Pisidiae in fauce Tsimboukkhan (Heldr!).

S. lucida Gren. et Godr. e Galliâ australi specifice differre videtur.

Ar. Geogr. Regnum Neapolitanum orientale.

35. S. sphærocarpa (Boiss. et Reut. Diagn. Ser. II, 3 p. 158 in notâ) glaberrima lucida, caule rubro obtuse tetragono in thyrum elongatum subpyramidalatum subglandulosum abeunti, foliis crassisculis imis minoribus ovatis lyrato-pinnatifidis cæteris bipinnatisectis segmentis divaricatis oblongis obtusis obtuse dentatis, foliorum superiorum angustioribus acutis, cymis bifidis 5-7 floris floribus subsessilibus, bracteis brevissime linearibus, calycis laciniis orbiculatis latiuscule scariosis, corollæ purpureâ, staminibus vix exsertis, appendice transverse reniformi, capsulis parvis globosis obtusis apiculatis calyce sesquilocularibus (2). *S. canina* Fl. Græc. VI, p. 78, tab. 598 non L. — *S. filicifolia* var *minor* Boiss. Diagn. Ser. I, 4. p. 70. — *S. livida* Fl. Græc. VII, p. I, tab. 601 ? (forma corollæ labio pallide purpureo).

Hab. in regione media Anatoliæ occidentalis, montes Mesogis, Tmolus, Cadmus, Lydiae et Cariæ (Boiss! Ball!), Lycia ad Elmalu (Bourg. 1781), Cilicia Trachea (Péron!), Cilicia littoralis in monte Torbol prope Tripoli (Boiss!), inter Ghazir et Beïtkachbo Libani (Gaill. 2756!).

Affinis *S. lucidæ filicifoliae* a quâ sat differe videtur foliorum minorum laciniis divaricatis obtusioribus, calyce latius scarioso, floribus et capsula

obtusiore minoribus, cymæ fructiferae ramis saepe magis elongatis. *S. livida* Fl. Græc. ex specimine juniori descripta corolla pallidæ variatione tantum differre videtur. Icon. *S. canine* Fl. Græcæ floribus magnis et staminibus inclusis toto cœlo a specie Linneanâ differt et huc potius quam ad *S. lucidam* referenda est.

36. *S. rutæfolia*, glaberrima, lucida, caule rubello elato ramoso elevatum quadrilineato, ramis longis in thyrsos longos angustos nudos abeuntibus, foliis carnosulis petiolatis in segmenta inæqualia oblonga cuneata cum mucronulo obtuse lobata vel dentata secus rachidem lobuliferam saepe decurrentia pinnatim vel bipinnatim sectis, bracteis bracteolisque breviter linearibus, cymis breviter et crassiusculæ pedunculatis rigide bisidis 13-5-floris, calycis laciniis orbiculatis membranâ fuscescente dentatâ subæquilatâ marginatis, corollæ lobis superioribus sublongioribus orbiculatis, antheris subexsertis, appendice magnâ transverse reniformi, capsulâ sphærico-depressâ mucronatâ calyce sesquilociore ②. *S. lucida* M.B. Taur. Cauc. II, p. 78! non L.

Hab. in Iberiâ prope Titlin (Steven! Rupr!), in Transcaucasiâ ad Borschom 3000' (Radde!).

Pluripedalis, affinis *S. lucideæ* sed præsertim *S. sphaerocarpæ*, ab utrâque distincta foliorum segmentis valde inæqualibus decurrentibus abbreviatis, rachide lobulatâ, floribus minoribus.

37. *S. laxa* (Boiss. et Heldr. Diagn. Ser. II, 3, p. 154) glaberrima a collo saepius pluricaulis, caulibus simplicibus vel basi parce ramosis obtuse quadrangulis parte tantum inferiori foliatis, foliis ambitu oblongis tenuiter bipinnatis sectis pellucido-punctatis, laciniis minimis oblongis acute dentatis, cymis crasse et breviter pedunculatis inferioribus 5-floris superioribus 3-1-floris thyrsos pyramidatos angustos laxos formantibus, pedicellis calyce brevioribus minute glandulosis, bracteis squamiformibus, calycis laciniis orbiculatis membranâ scariosâ latâ denticulato-lacerâ marginatis, corollæ labio superiore purpureo, filamentis glandulosis apice breviter exsertis, appendice semicirculari, capsulâ majuscûlâ sphæricâ abrupte et longe mucronatâ ②.

Hab. in regione alpinâ Peloponnesi inter lapides, mons Kyllene supra Livadi (Heldr! Orph. Fl. Gr. exs. 243!), regio superior Taygeti (Heldr. pl. exs. sub *S. lucideæ* et *felicifoliae* var!).

Pedalis vel elatior, foliis tenuiter sectis *S. Hoppii* similis sed ab eâ præter alias notas appendicis formâ aliena. A formis *S. lucideæ* pro quâ eam olim habuimus laciniis tenuibus foliorum, caulis subaphyllis, calycis marginâ toto scarioso dentato, etc. diversa.

38. *S. myriophylla* (Boiss. et Heldr. Diagn. Ser. I, 22, p. 39) glaberrima siecitate nigricans mulicaulis, caulis crassiusculis pu-milis parte inferiore crebre foliosis in thyrsum elongatum angustum inferne tantum foliosum abeuntibus, foliis breviter petiolatis ambitu oblongis in lacinulas confertas minutissimas breviter oblongo-lineares

neutas divaricatas tripinnasectis, cymis breviter pedunculatis 3-5-floris, bracteis bracteolisque breviter linearibus, pedicellis pruinosis calyce brevioribus, floribus parvis, calycis laciniis latiusculæ albo-marginatis denticulatis, corollæ urceolatæ calyce 2 ½-plo longioris lobis superioribus purpureis rotundatis, antheris subexsertis, appendice reniformi transverse latiore subretusâ, capsulâ.... (2).

Hab. in regione superiori montis Gheidagh Tauri Isaurie 6000' (Heldr.).

Semipedalis vel fere pedalis, facies et folia *S. Hoppii* a quâ foliis tenuius dissectis, antheris breviter exsertis et appendice late reniformi nec linearilanceolata differt. *S. laxa* magis affinis differt foliorum laciniis multo crassioribus, floribus duplo majoribus.

39. ***S. versicolor*** (Boiss. Diagn. Ser. II, 3, p. 156) tota minute glanduloso-pruinosa, caulinis pumilis ascendentibus simplicibus crebre foliosis in thyrsus oblongum undum abeuntibus, foliis subsessilibus ambitu ovatis parvis in segmenta conferta anguste lanceolata argute serrata vel incisa pinnatisectis, bracteis bracteolisque linear-setaceis, cymis breviter et crassiuseule pedunculatis 7-3-floris glandulosis, pedicellis calyce brevioribus, calycis glabri laciniis orbiculatis late albo et fusco-membranaceis, corollâ calyce duplo longiori lurido-rubellâ, antheris inclusis, appendice semicirculari transverse latiore basi truncatâ, capsulâ globosâ calycem parum excedente longe mucronatâ (2) ?

Hab. in montibus prope Erzerum Armeniæ (Calv. pl. exs. 984!).

Ex unico specimine mihi nota et ulterius observanda, florifera cum thyrsi 10-pollicaris, folia inferiora sesquipollicem vix longa eis *S. Hoppii* a quo appendice non linearis differt fere tenuius dissecta. Affinis *S. myriophyllæ* a quâ differt indumento pruinoso-glanduloso, foliorum laciniis non minutissimis, calyce multo majore etc.

+ + Perennes.

40. ***S. lepidota*** (Boiss. Diagn. Ser. I, 4, p. 71) tota squamulis peltatis adpressis margine stellatim ciliatis lepidoto-inæana cespitosa, caulinis teretibus simplicibus in thyrsus abbreviatum abeuntibus, foliis carnosulis oblongo vel linearispathulatis in petiolum longe attenuatis, cespitis confertis obtusis integris vel apice truncatis aut obsolete 3-5-dentatis, caulinis ad caulis partem inferiorem paucis trifidis, cymis brevibus 3-2-floris, pedicellis calyci aequilongis, bracteolis minimis linearibus, calycis subglandulosi laciniis rotundatis membranâ latâ denticulatâ marginatis, corollâ..., staminibus..., capsulâ ovatâ acutâ parvâ calyce subdupo longiore ♀.

Hab. in Armenia et Cappa loci orientali (Auch 2437).

Species indumento Hippophaës, Eleagni vel Alyssi distinctissima, caules cum thyrsi 10-12-pollicares, folia cum petiolo sesquipollicaria apice 1 ½-2-lineas tantum lata, capsule eis *S. caerulea* vix majores.

41. ***S. farinosa*** (Boiss. in Ky. Pers. exs. 1845. — Diagn. Ser.

I, 7, p. 40) tota tuberculis albis mimimis papillæformibus crebris farinosa cana, caulis e rhizomate crasso subcarnoso crassis simplicibus foliosis in thyrum brevem oblongum foliolosnm abeuntibus, foliis carnosulis longiuscule petiolatis ovatis vel oblongis obtuse crenato-lobatis vel plus minusve profunde incisis vel pinnatipartitis, bracteis ellipticis integris vel dentatis cymarum 3-1-florum pedunculos æquantibus vel superantibus, pedicellis calyce vix longioribus, calycis glandulosi laciniis ovatis anguste membranaceo-marginatis, corollâ cylindraceâ flavidâ calyce et latitudine suâ triplo longiore fauce constrictâ labio superiore longiore, antheris superioribus sub-exsertis, appendice minutâ reniformi-orbiculatâ, capsulis globosis minute papillosis acutis calyce duplo longioribus ♀. *S. longiflora* Bth. in n.c. Prodr. X, p. 313!

Hab. in fissuris rupium Persiae australis, mons Sabst Buschom ad Schiraz (Ky. 427!), prope Schahpur, Persepolin et in monte Parrow prope Kermanchah (Haussk!). Fl. Apr., Mai.

Caulis cum thyro 2 3-pollicares, foliorum lamina 6-9 lineas longa, corolla longitudine inter 2¹/₂-4 lineas varians. Species fragilis, carnosa, distinctissima.

42. *S. xylorrhiza* (Boiss. et Haussk.) e rhizomate brevi lignoso multicaulis minute glandulosa vel glabriuscula, caulis pumilis fragilibus simplicibus subangulatis in thyrum angustum parte inferiori foliatum abeuntibns, foliis ovato-oblongis acutiuscule inciso-dentatis basi cuneatis vel indivisis vel lyratis vel pinnatisectis laciniis oblongis dentatis superioribus confluentibus, foliis floralibus diminutis integris, cymis breviter pedunculatis 5-1-floris, bracteolis oblongis, pedicellis glandulosis calyce vix longioribus, calycis laciniis ovatis anguste marginatis corollâ campanulatâ calyce triplo longiore, antheris exsertis, appendice ovato-rotundâ parvâ, capsulâ globosâ sub-didymâ obtusâ apiculatâ calyce duplo longiore.

Hab. in fissuris rupium Syriae borealis et Mesopotamiae ad Aleppo, Aintab, Orfa, Gebel Taktak Mesopotamiae (Ky. 421! Haussk!), Palestinæ ad Tufileh (Roth 386!), Cilicie ad rupes Gulek Boghas (Ky. 400 sub *S. Pinardi* et Bal. sub *S. heterophylla*, specimina tantum fructifera sed quæ hue spectare videntur).

Caules cum thyro 4-7-pollicares, folia parva 6-8 lineas longa incisa vel divisa. Species saepe relata ad *S. heterophyllam* a quâ præter appendicis multo minoris formam differt corollâ proportione calycis multo longiore campanulatâ ore expansâ nec urceolata.

43. *S. Taygetea* (Boiss. Diagn. Ser. I, 4, p. 68) tota pubescens crispa glandulosâ obsita e rhizomate indurato multicaulis, caulis fragilibus pumilis simplicibus teretiusculis foliosis in thyrum abbreviatum glandulosissimum abeuntibus, foliis petiolatis triangulari-ovatis cuneato-truncatis vel subcordatis acutis acute et subduplicatim dentatis saepe lobatis, cymis breviter pedunculatis 7-3-floris supremis unifloris, bracteis oblongis, bracteolis breviter oblongo-linearibus, pedicellis calyci subæquilongis, calycis glabri laciniis orbiculatis late membranaceo-marginatis, corollâ virescenti-flavidâ, antheris inclusis,

appendice reniformi transverse latiore, capsula globosâ subdidymâ obtusissimâ apiculatâ calyce subdupo longiore ♀.

Hab. in fissuris rupium umbrosarum regionis inferioris Taygeti prope Mistra (Boiss! Heldr!).

Semipedalis vel paulo procerior, folia inferiora 1 $\frac{1}{2}$ poll. longa. Flores magnit. S. *Scopolii* ejus formas pumilas quodammodo refert, sed ob fragilitatem, rhizoma crassum sublignosum pereume, pedicellos breves, capsulum obtusissimam gregi S. *heterophylla* adsoiciata et ab omnibus affinibus indumento viscido diversa

44. S. pulverulenta (Boiss. et Noë Diagn. Ser. II, 3, p. 155) glandulis subsessilibus crebris cinerascenti-pruinosa, caulinibus obtuse quadrangulis foliosis simplicibus vel parce ramosis, foliis petiolatis enneatis vel basi truncatis oblongis obtusiusculè dentatis, thyro brevi aphylo, bracteis bracteolisque minutis linearibus, cymis breviter et crassinervie pedunculatis 3-7-floris, floribus majusculis, pedicellis calyce subbrevioribus, calycis glabri lacinias membranâ subglabrâ latâ marginatis, corollâ ureolatâ, antheris vix exsertis appendice transverse reniformi, capsula majuscula ovafo-globosâ acutâ ♀.

Hab. in subalpinis et alpinis ad Ardana Armeniae australis (Noë 4065!), ad radices montis Argai Cappadociae (Ky. exs. 244! spec. fructiferam).

Facies S. *catariaefolia* et ut ea floribus magnis, antheris subincisis, appendice, a formis pruinosis S. *variegata* aliena, ab eâ discrepat floribus brevis pedicellatis, foliis non cordatis, calyce non glanduloso.

45. S. catariæfolia (Boiss. et Heldr. Diagn. Ser. I, 12, p. 36) minute glanduloso-pubescentes, pluricaulis, caulinibus elatis parce ramulosis quadrangulis in thyrsum brevem aphyllum glandulosum abeuntibus, foliis petiolatis oblongis acutiusculè et duplicatim dentatis basi subcordatis, cymis 3-7-floris, bracteis bracteolisque linearibus, pedicellis calyce vix longioribus, floribus majusculis, calycis glanduloso-puberuli lacinias membranâ albâ latiusculâ denticulatâ marginatis, corollâ ureolatâ flavidâ, antheris non vel vix exsertis, appendice transverse reniformi basi truncatâ, capsula.... ♀. S. *nepetaefolia* Boiss. et Heldr. exs. non Presl.

Hab. in cacumine montis Karadagh Lycaonite in ruinis veteris ecclesie (Heldr!).

Sæpe sesquipedalis, flores magnitudinis S. *Scopolii* ejus formas foliis minoribus donatas refert, ab eâ floribus breviter pedicellatis statim distinguenda; a formis pubescentibus S. *variegata* floribus fere duplo majoribus, antheris non longe exsertis, appendicis formâ distincta, inter affines calyce glanduloso insignis.

46. S. Mesogitana, breviter glanduloso-pubescentis pluricaulis, caulinibus obtuse quadrangulis subsimplicibus in thyrsum subpaniculatum inferne minute foliosum abeuntibus, foliis ovato-oblongis basi truncatis et subcordatis grosse subuplicatim et obtusiusculè dentatis basi interdum pinnatisecto-lyratis inferioribus petiolatis superio-

ribus subsessilibus, bracteis ellipticis, bracteolis squamæformibus, cymis breviter pedicellatis divaricatum 5-7-floris, pedicellis filiformibus calyce sublongioribus, calycis glabri laciniis albo-marginatis, corollâ luridâ, antheris subexsertis, appendice parvâ transverse reniformi, capsulâ globosâ obtusâ apiculatâ calyce duplo longiore ♀.

Hab. in rupestribus calcareis regionis mediæ montis Mesogis Lydiæ (Boiss!).

Species subdubitanter proposita *S. catariefoliae* indumento et florum magnitudine affinis sed thyrso subfoliato, calyce glabro, antheris exsertis distincta. *S. variegatae* var. *rupestris* quoque proxima differt floribus majoribus, pedicellis gracilioribus, appendice. A formis *S. heterophylloë* cui adnumerata fuit aliena est indumento, caulis non fragilibus, capsulâ minore, etc.

47. *S. heterophylla* (Willd. Sp. III, p. 274) glabra sœpe glaucescens, caulis e rhizomate lignescenti pluribus ascendentibus fragilibus sœpius abbreviatis in thyrum aphyllum pyramidatum multiflorum abeuntibus, foliis carnosulis petiolatis ovatis obtusis basi subcuneatis truncatis vel subcordatis obtuse inciso-crenatis indivisis vel lyrato-pinnatifidis vel sectis segmentis lateralibus 1-2 multo minoribus, cymis laxe 5-9 floris, bracteolis squamæformibus, pedicellis calyci subæqualib; calycis laciniis ovatis latiusculæ albo-membranaceis, corollâ majuscûlâ lurida purpureâ calyce duplo longiore, antheris subexsertis, appendice reniformi transverse sublatiori interdum subretusâ, capsulâ calyce subtriplo longiore globosâ obtusâ apiculatâ sulcato-didymâ ♀. *S. Urvillæana*, Wydl. Mon. p. 160, tab. 2! et *S. Oliveriana* Wydl. p. 159! — *S. micrautha* Urv. Enum. p. 75. — *S. cæsia* Fl. Græc. VII, p. 3, tab. 604.

Hab. ad rupes Cretæ (Willd.) et ad prom. Grabousa (Raul!), insularum Naxos, Melos, Ferasia (Urv!), Chios (Heldr! Orph. Fl. exs. 728!), Thera (Sart!), Syra (Boiss!), Baotia supra Livadia (Heldr! Orph!), Argolidis ad Naupliam (Orph!), Atticæ ad Lycabettum (Heldr!).

S. pinnatisecta. — Folia profundius divisa pinnatisecta segmento terminali non majore, caules sœpe proceriores. Forma ad typum manifeste transiens. — *S. laciniata* var. *obtusiloba* Rent. et Marg. Fl. Zaczynth. pag. 70. — *S. chrysanthemifolia* Fl. Pelop. non Willd.

Hab. in rupibus regionis inferioris, Attica ad Acropolin. Lycabettum cum typo (Boiss! Heldr!), Peloponnesus ad Calymata et Pylos (Bory!), Cephalonia (Heldr!), Zaczynthus (Marg!), Acarnania (Nied!).

48. *S. laciniata* (W. K. Pl. rar. Hung. II, p. 185, tab. 170) glabra viridis pluricaulis, caulis rubellis tetragonis subsimplicibus foliosis in thyrum subaphyllum subpyramidatum abeuntibus, foliis petiolatis lyratis vel pinnatipartitis sectisve segmentis ovato-oblongis vel ligulatis acute dentatis vel incisis inferioribus plurifloris, bracteis linearibus, bracteolis setaceis, pedicellis calyci æquilongis vel brevioribus, calycis laciniis orbiculatis membranâ albâ integrâ eis angustiore marginatis, corollâ luride purpurascente, antheris inclusis, appendice late orbiculato-reniformi interdum retusâ, capsulâ calyce

vix duplo longiore globosâ apiculatâ ♂. *S. heterophylla* Fl. Græc. tab. 603 non Willd. — *S. frutescens* Sm. Prod. non L. — *S. Sibthorpiana* Spreng. Syst. 2, p. 786.

Hab. in rupestribus regionis subalpinæ et alpinæ Græciae 5000'-6000', montes Olenos, Kyllene et supra Megaspileon Achææ (Heldr!), mons Arachneon Argolidis (Sprun!), Olympus Thessalus (Heldr! Orph. Fl. Gr. exs. 729!).

S. multifida. — Folia in lacinias angustas lanceolato-lineares acute dentatas decomposita. Forma interdum puberula. — *S. multifida* W. Hort. Berol. tab. 58 et herb! — *S. casia* Griseb. Spie. II, p. 38 non Sibth

Hab. in rupestribus regionis subalpinæ et alpinæ Peloponnesus in regione superiore Taygeti et montis Chelmos (Heldr!), mons Velugo Ætoliae et Parnassus (Heldr. herb. Norm. 682!), mons Scardus Macedonie (Griseb!).

Valde affinis formis pinnatisectis *S. heterophylle* pro cuius stirpe subalpinâ et alpinâ haberri posset, differt tamen colore lat. viridi, foliis tenuius membranaceis nec carnosis lobis et dentibus siepius acutis nec obtusis, caulinibus non adeo fragilibus, capsula calycis proportione breviori. Nondum vidi editio formas hujus speciei foliis indivisis lobatis quæ in Dalmatiâ et Serbia sœpe observantur.

Ar Geogr. Regio Danubialis, Serbia, Dalmatia.

49. *S. Olympica* (Boiss. Diagn. Ser. I, 4, p. 69) parte inferiori glaberrima multicaulis, caulinibus nigricanti-rubellis nitidis subsimplicibus in racemum laxissimum pauciflororum aphyllum abeuntibus, foliis intense viridibus flavidulis præter nonnulla interdum indivisa oblonga inciso-dentata lyrato-pinnatifidis partitisve lacinias late linear-oblengis acute et duplicatim dentato-incisis, cymis breviter pedunculatis parce glandulosis inferioribus 3-2 floris ceteris unifloris, bracteis bracteolisque linearibus brevibus, floribus majusculis in dichotomiâ et lateralibus supra bracteolam sessilibus, calycis glabri lacinias orbiculatis membranâ rufescente vel purpureâ eis subæquilaterali undulato-crispâ marginatis, corolla flavidæ urceolatæ lobis binis superioribus purpureis, antheris inclusis, appendice transverse latiore seniorieculatâ basi late cordatâ, capsula sphaericâ obtusa mucronata calyce duplo longiore ♀. *S. pyrrotopha* Boiss. in Ky. exs. 1859. — *S. Oliveriana* C. Koch Linn. XXIII, p. 709 non Wydl.

Hab. ad rupes frigidas regionis alpinæ, Olympus Bithynus supra Kirkbouar (Boiss! Pichler!), mons Argeus Cappadociae (Bal. exs! sub *S. depauperata*), Armenia australis in monte Bingöldagh 9000' (Ky. 321 et 459! sub *S. laciniata* var.), Caucasus in Abchasiâ (Radde!), ad rupes montis Kasbek (Rehm!) et ad Baidera inter Vladikankas et Tiflis 6000' (Rupr!), in ditionibus Alagir, Rad-scha, Tindal Caucasi orientalis 3-9000' (Rupr!).

S. Lazica. — Flores e maximis (5 lineas longi). — *S. Lazica* Boiss. et Bal. MSS.

Hab. in valle Djimil Ponti Lazici 6000' (Bal!).

Caules e rhizomate indurato pedales aut sublongiores, racemus angustus brevis laxus cymis fere omnibus ad florem unicum reductis, folia 1½-3-pollices longa ea *Chrysanthemi frutescentis* fere referentia, flores magnitudinis

S. Scopoliæ. Species a *S. laciniatâ* præter inflorestiam racemosam calyce late et undulatim membranaceo et appendice magnâ basi cordatâ disticta.

50. *S. Ruprechtii*, præter inflorescentiam glaberrima, caulis e radice cylindricâ carnosâ longâ apice longe fibrosâ ascendentibus pumilis vel nanis simplicibus foliosis, foliis petiolatis ambitu oblongis lyratim pinnatipartitis vel sectis segmentis oblongis obtuse et irregulariter incisis, racemo terminali simplici oblongo tandem laxo glanduloso, cymis omnibus ad pedunculum uniflorum flori aequilongum reductis, bracteis linearibus pedunculo longioribus, bracteolis ad floris basin calyce aequilongis, floribus magnis, calycis ampli glabri laciinis orbiculatis late purpureo-membranaceis, corollæ ochroleucæ calyce duplo longioris lobis superioribus amplis sordide purpureis, staminibus inclusis, appendice transverse reniformi basi cordatâ, capsulâ magnâ ovalâ aentâ calyce vix longiore ♀.

Hab. inter lapillos regionis alpineæ superioris Caucasi orientalis, ditio Alagir ad glacies Zei 6600'. Tuschetia ad Didigwerdi et Samkuri Zweti 9000'-40000' (Rupr.).

Caules cum racemo spiciformi 2-3-pollicari 3-5-pollicares imâ basi nudi, folia 1½-2-pollices longa longisuscule petiolata, flores et capsule e maximis. Species insignis ab affini *S. Olympicâ* caulis nanis, radice carnosulâ in rhizoma induratum non abeunte, racemo conferto spiciformi, sepalis amplis capsulam subequalibus disticta.

51. *S. Griffithii* (Bth. in D.C. Prodr. X. p. 312) glabra vel minute glanduloso-pruinosa viridis vel canescens, caulis pluribus simpliciensulis inferne foliosis superne in thyrum virgatum angustum longum strictum abeuntibus, foliis breviter petiolatis oblongis profunde pinnatifidis vel pinnatisectis laciinis acute inciso-dentatis foliorum superiorum lanceolatis acuminatis, cymis minute glandulosis 3-4-floris breviter et stricte pedunculatis, floribus majusculis, calycis laciinis orbiculatis latiusculæ marginatis, antheris breviter exsertis, appendice reniformi-orbiculatâ, capsulâ ovalâ aentâ ♀.

Hab. in regno Cabulico (Griff. 616! et 619!).
Insignis thyro stipe sesquipedali angustissimo, folia inferiora 1-2-pollicaria.

Ar. Geogr. Tibetia (Hook. et Thoms. sub *S. Kotschyi*).

52. *S. depauperata* (Boiss. Diagn. Ser. I, 4, p. 68) tota minute glandulosa viridis multicaulis, caulis virgatis sepius opposite et stricte ramosis inferne foliosis superne in thyrum vel sepius paniculam laxam depauperatam abeuntibus, foliis breviter petiolatis et sessilibus teneris ambitu oblongis in segmenta parva oblongo-lanceolata aenta incisa basilaria minora superiora confluentia pinnatisectis, cymis remotis longisuscule pedunculatis imis 3-3-floris cæteris unifloris, bracteis bracteolisque linearibus, pedicellis glandulosis calyce subbrevioribus, floribus majusculis, calycis glabri laciinis orbiculatis angustæ albo-marginatis, corollæ virescenti-luteæ urceolatae

labio superiori purpureo, antheris inclusis, appendice spatulatâ transverse latiore, capsulâ sphæricâ apiculatâ calyce subdupo longiore ♀.

Hab. in glareosis et rupestribus regionis alpine Anatolie occidentalis, montes Sipylus, Mesogis et Tmolus Lyiae, Cadmus Curiae (Boiss!), mons Akdaghi Lyciae (Bourg!).

Caules 1-2-pedales, folia pro parte magnitudine parva anguste laciniata. A cl. Bth. in n.c. Prodr. *S. variegata* ad sociata a quâ floribus duplo majoribus, filamentis inclusis etc aliena, sed prope *S. Olympica* a quâ caulinibus ramosis, foliis parvis tenuiter laciniatis, appendice non cordatâ differt collocanda.

53. *S. subaphylla* (Boiss. Diagn. Ser. I, 7, p. 41) glabra multicanlis, caulinibus tenuiter virgatis junceis a parte inferiori opposite et striete ramosis in thyrsos racemiformes longos laxissimos nudos abeuntibus, foliis ad caulinum partem inferiorem panicis parvis subsessilibus oblongo-lanceolatis in laciniis parvas tenues pinnatifidis vel subbipinnatis et, bracteis bracteolisque squamiformibus, cymis breviter pedunculatis praeter inferiores trifloras unifloris, pedicellis glandulosis calyce subbrevioribus, calycis laeuiis orbiculatis latiuscula albo-marginatis, corollâ areolatâ rubellâ calyce duplo longiore, staminibus inclusis, appendice parvâ crassiusculâ transverse sublatiore margine obsolete denticulatâ, capsulâ parvis sphæricis apiculatis calyce duplo longioribus ♀. *S. Persica* Bth. Prodr. n.c. X, p. 314 ex parte!

Hab. ad rupes montium Persiae australis et Kurdistaniae conterminae 9000-12000', montes Schahm, Kellal, Ssebsseknh, Sawers, Kuh Nur, (Haussk!), Kuh Daëna et Salst Buschom prope Schiras (Ky. 429! 798! et 751! sub *S. depauperata*), in Persiâ loco non notato (Auch. 3169!).

Pedalis et sesquipedalis, insignis caulinibus tenuiter virgatis praeter partem inferiori subaphyllis, foliis parvis (6-10 lin. longis) tenuiter laciniatis. Facies *S. depauperata*, caules tenuiores, flores minores labio superiore vix longiore.

54. *S. frigida* (Roiss. Diagn. Ser. I, 7, p. 42) glabra glauco-cæsia, caulinibus elatis virgatis crassiusculis striete et longe ramosis in thyrsos longissimos angustos laxos saepe ramoso-paniculatos parte inferiori foliosos abeuntibus, foliis crassiusculis subsessilibus lanceolatis elevatim nervosis remote et acute inciso-dentatis, superioribus anguste linearibus, cymis breviter pedunculatis divaricatum 3-3-floris, pedicellis calyci æquilongis vel brevioribus, bracteolis squamiformibus aenatis, calycis laeuiis orbiculatis late membranaceis, corollâ areolatâ calyce duplo longiore rubellâ, antheris inclusis, appendice minutâ obtiangulari-orbiculatâ, capsulâ parvâ ovato-depressâ aenâ calyce duplo longiore ♀.

Hab. in rupestribus regionis alpine Persiae austro-occidentalis, in districtibus Bachtiaris et Kuh Kilouyeh, montes Tehinar, Sakawa, Kuh Nur, Kellal 8000' (Haussk!), in montibus Kuh Delu et Kuh Barli prope Schiras 329 et 411 (Haussk!). Spec. incompl. ex Tichmedesch Persiae bôr. (Bge!) hic quoque spectare videtur. Mammam nigrum copiose præbet.

Canes 2-3-pedales thyrso saepe paniculato et pedali, folia inferiora bipollicaria 5-6 lineas lata superiore valde diminuta. Facie affinis *S. subaphyllæ* sed glauca, folia carnosa lanceolata indivisa majora, inflorescentia multiflora, etc.

55. *S. scabiosæfolia* (Bth. in D.C. Prodr. X, p. 311) glabra glaucescens saepe pruinoso-punctata, caulinis numerosis erectis simplicibus vel parce ramosis in thyrsos aphyllos abeuntibus, foliis insimis interdum indivisis oblongis dentatis, cæteris vel omniibus lyratopinnatifidis vel sectis laciniis incisis vel crenatis oblongis lateralibus parvis terminali majore, cymis bifidis 3-3-floris, bracteis minutis breviter linearibus, floribus parvis, calycis laciniis ovato-orbiculatis anguste scariosis, antheris breviter exsertis, appendice orbiculata, capsulis parvis globosis ♂.

Hab. in regno Cabulico (Griff. 614! 822!) in Belutschia (Stocks 991!).

Pedalis et se-quipedalis, facies *S. variegatæ* sed appendicis forma diversa.

Ar. Geogr. India boreali-occidentalis..

56. *S. rostrata* (Boiss. et Buhse Aufz. p. 163) glabra pluricanlis, caulinis subsimplicibus obtuse quadrangulis in thyrsum laxum abeuntibus, foliis radicalibus petiolatis lyratis segmentis lateralibus 2-4 parvis terminali ovato obtuse lobato lobis crenatis, foliis caulinis diminutis subsessilibus interrupte pinnatipartitis vel sectis laciniis subpetiolatis oblongo-lanceolatis acute dentatis terminalibus saepe confluentibus, cymæ longiusculæ pedunculatae bifidae ramis terminalibus laxe paucifloris, bracteis bracteolisque linearis-setaceis, pedicellis parce glandulosis, calycis laciniis ovatis anguste marginatis, corollæ luridæ, antheris breviter exsertis, appendice reniformi, capsula ovata elevatim reticulata calyce duplo longiore in rostrum pungens eæ æquilongum abrupte abeunte ♀.

Hab. in prov. Ghilan Persie prope Imam Sadeh Haschim et prope Rudbar in valle Sebrud (Buhse!) Specimen floriferum e ditione Talysh prope Nou (Buhse!) hue quoque spectare videtur.

Pedalis et bipedalis, folia 2 pollices longa, facies *S. laciniatae* a qua et ab affinibus differt rostro tenuiter pyramidato 2 lineas longo capsulae fere æquilongo.

57. *S. trichopoda* (Boiss. et Bal. Diagn. Ser. II. 3. p. 456) glabra, caulinis proceris obtuse angulatis gracilibus in paniculas amplas longas foliolosas abeuntibus, foliis ambitu oblongis lyratis vel pinnatisectis 1-2 jugis segmentis grosse incisis vel pinnatifidis lateralibus minoribus petiolulatis, foliis superioribus diminutis subsessilibus floralibus elliptico lanceolatis integerrimis, cymis longe pedunculatis divaricatissime bis bifidis 3-9-floris, ramis cymæ pedicellisque minute glandulosis calyce multoties longioribus filiformibus, bracteolis setaceis brevibus, calycis glabri laciniis ovato-oblongis latinscule albo-membranaceis, corollæ purpurascens calyce 2½-plo longioris lobis superioribus magnis, staminibus inclusis,

appendice transverse latiore ampliâ semilunari basi subconcavâ, capsulâ sphærico-subdepressâ mucronatâ calyce duplo longiore ♂ ?

Hab. in fauce Guzelderé supra Sedichig in Ciliciâ littorali (Bal!). Fl. Mai.

Caules 4-5-pedales, panicula saepe sesquipedalis ob pedunculos ramos et edicellos longos filiformes divaricatos effusissima, folia inferiora 3-4-pollicaria, flores magn. *S. laciniata*. Species in sectione florum pedicellis 7-9 lineas longis distinctissima, sed ob habitum et folia dissecta *Tomiophyllum* nec *Scorodoniae* adnumeranda.

58. *S. xanthoglossa* (Boiss. Diagn. Ser. I, 12, p. 38) glabra glaucescens, caulis erectis rigidis crassiuseulis a parte inferiori vel dimidiâ in paniculam longam multifloram thyrsoideam tandem saepe ramosissimam abeuntibus, foliis firmis carnosulis intimis oblongo-cuneatis obtuse crenatis vel flabellatim incisis cæteris pinnatim sectis lacinij dentatis inferiorum minutis oblongis obtusis superiorum angustatis anguste lanceolatis vel linearibus denticulis acutis, cymis breviter pedunculatis bifidis, ramis tandem elongatis rigidis laxissime plurifloris, bracteis bracteolisque linearibus hisce calyces subsessiles æquantibus, floribus medioeribus, calycis minuti lacinis orbiculatis late albo-marginatis, corollâ calyce triplo majore lobis superioribus basi strangulatis orbiculatis magnis, staminibus plus minus exsertis, appendice magnâ flavidâ orbiculatâ margine crenulatâ, capsulâ parvâ ovatâ mucronatâ calyce sesquiloniore ♂.

α. genuina. — Flores submajores, corollæ lobi superiores amplissimi tubo latieres et sublongiores, filamenta longe exserta.

Hab. circa Hierosolymam frequens (Boiss!), circa Hebron (Ky. 862!), ad stagna Salomonis (Ky. 792! sub *S. laciniata*!).

β. decipiens. — Flores paulo minores lobis superioribus corollæ paulo minus amplis, filamenta brevius exserta. — *S. decipiens* Boiss. et Ky. Diagn. Ser. II, 3, p. 156. — *S. Aintabensis* Boiss. et Haussk. MSS. — *S. expansa* Reut. MSS. — *S. Orientalis* Ehrenb. exs. — *S. Persica* Bth. in D.C. Prodr. ex parte !

Hab. in Libano a radicibus prope Tripolin et Sidonem ad regionem alpinam usque (Ehr! Bl! Gaill!), Antilibano (Gaill! Ky. 149!), Mesopotamiâ (Oliv! Ky. 361!), Syria boreali ad Aleppo (Ky. 961!), ad Aintab (Haussk!), in deserto Chabor Assyriæ (Haussk!), Ciliciâ in ditione Kassanoglu et ad radices montis Argai (Ky. 68! et 215! sub *S. floribundâ*). Ciliciâ ad Bouloukli (Bal. exs. *S. canina*!) regione alpinâ Tauri (Ky. 42! sub *S. bicolor*!).

γ. hispidula — Tota superne papillaris et glandulosa. — *S. hispida* Boiss. et Bal. Diagn. Ser. II, 6, p. 157.

Hab. in schistosis faucis Guzelderé prope Mersina Cilicie littoralis (Bal!).

Species lobis superioribus amplis corollæ et appendice magnâ lobis saepius vix minore insignis, caulis plus minus elatis, foliis varie laciñiatis valde variabilis, interdum ramosissima et tunc facie similis *S. caninae* pro quâ eam ipse interdum habui, tamen characteribus aliena.

59. *S. striata*, glabra, multicaulis, caulis tenuissime virgatis angulatis striatis fere a parte inferiori in thyrsos longissimos angustos

multifloros parte inferiore foliatos abeuntibus, foliis in laciniis paucissimas lateralibus breviter triangulari-lanceolatis decurrentibus terminali trifidâ omnibus acutis pinnatipartitis, superioribus sessilibus basi attenuatis, cymis breviter pedunculatis laxe 9-5-floris ramis gracilibus, floribus minimis subsessilibus, calycis laciniis ovatis late albo-marginatis, corollæ calyce subdupo longioris lobis superioribus magnis orbiculatis, appendice orbiculari transverse sublatiore lobos superiores corollæ aequante, staminibus longe exsertis, capsulâ.... ♀.

Hab. in montibus prope Jezd Persiae orientalis ad pagum Dehballo (Buhse 1839!), in Persia loco non notato sub *S. bicolori* (Ky. 871!, Derderian!), prope Ichredabad, Kerman, Ispahan (Bge!).

Caulis cum thyro pedali 1½-2-pedalis, gracilitate affinis *S. thesioidi* pro qua specimine imperfecto tunc tantum noto eam h̄:bui differt foliorum laciniis paucis brevissimis latoribus et præsertim appendicis formâ et magnitudine.

60. ***S. deserti*** (Del. Fl. Eg. p. 96, tab. 33, fig. I) præter inflorescentiam obsolete et minutissime glandulosam glaberrima, caulis e rhizomate indurato virgatis rigidis inferne parce foliosis in thyrsum elongatum saepe paniculatum ramosum abeuntibus, foliis parvis crassiusculis margine cartilagineis inferioribus petiolatis ovatis vel lyritis superioribus sessilibus pinnatipartitis laciniis oblongo vel lineari-spathulatis obtusissimis obtuse crenatis vel dentatis, cymis breviter pedunculatis divaricatum bifidis 9-3-floris, bracteis bracteolique minute squamiformibus, floribus minutis, pedicello calyce breviore, calycis parvi laciniis carnosis orbiculatis anguste albo-marginatis, corollæ luride calyce duplo longioris lobis superioribus parvis rotundatis, antheris subinclusis, appendice orbiculata, capsula parva calyce duplo longiore subglobosa acuta ♀ *S. Sinaica* Benth. in D.C. Prodr. X, p. 314 non differt.

Hab. in desertis arenosis Egypti (Del.) circâ Suez (Bové!), ad Ouadi el Hof prope Heluan (Schweinf!), in Arabia Petreea regione Bestau (Schimp. 329!), Wadi Scheick (Boiss!), in Persia australis insulis Karak et Korgo (Ky. 46!), ad Bonchyr (Auch. 5051!), circa Kaserun et Schahpur (Haussk!), in Persia orient. (Bge!).

Sesquipedalis vel humilior, saepe tandem ramosissima cymarum ramis rigidis, foliorum laciniis obtusissimis margine plus minus cartilagineis recognoscenda.

61. ***S. marginata*** (Boiss. Diagn. Ser. 1, 4, p. 72) glaberrima a celo pluricaulis, caulis erectis et ascendentibus virgatis rigidis duris albo-corticosis simplicibus vel rugosis in thyrsu laxum rigidum abeuntibus, foliis crassis rigidis albo-marginatis obtusissime incisis vel crenatis radicalibus rosulatis ovatis basi cuneatis petiolo eis subæquilongo suffultis, caulinis diminutis breviter petiolatis saepe fasciculatis, cymis breviter pedunculatis bifidis vel abortu ad ramum reductis remote 5-1-floris, bracteolis squamæformibus minimis, floribus parvis subsessilibus, calycis laciniis ovatis anguste albo-mar-

ginatis, corollâ parvâ sanguineâ, staminibus... appendice obovato-orbiculatâ, capsulâ parvâ globosâ obtusissimâ apiculatâ (2).

Hab. in monte Elvend Persiae (Auch. 2898!), monte Avroman Kurdistaniæ Persice in schistosis 5000' (Haussk!), Assyria ad Beschabur juxta trajectum Tigridis (Ky. 390!).

Planta coriacea sed biennis nec suffrutescens, caules cum thyrso pedales, foliorum radicalium lamina 12-15 lineas longa, flores eis *S. caninæ* subminores. Species foliis crassis marginatis insignis facie gregem *S. caninæ* referens sed appendice ab eâ remota.

** Appendix ovata, oblonga, vel oblongo-linearis.

+ Biennes.

62. ***S. hæmatantha*** (Boiss. et Heldr.) glaberrima a collo pluricaulis, caulis tereti-subangulatis opposite et stricte ramulosis parle inferiori foliosis superne in thyrsos laxos pyramidatos aphyllos abeuntibus, foliis oblongis acute dentatis obtusis basi cuneatis inferioribus longiusculæ petiolatis superioribus sessilibus, cymis pedunculatis bifidis divaricatis laxe 3-7-floris, bracteolis brevissimis subulatis, pedicellis crassis parce glandulosis calyce subduplicite longioribus, calycis glabri laciniis ovatis membranâ latâ undulato-lacerâ marginatis, corollâ sanguineâ, labii superioris longioris lobis orbiculatis transverse latioribus, staminibus inclusis, appendice parvâ linearî oblongâ, capsulâ calyce sesquilongiore globosâ obtusâ longiusculæ mucronatâ (2).

Hab. in schistosis montium Avroman et Schahu 5000' (Haussk!).

Pedalis vel sesquipedalis, foliorum inferiorum lamina 1-1½-pollicaris 7-9 lineas lata, thyrsus floribundus, capsula mediocris. Inter species appendice linearî-oblongâ donatas radice bienni et foliis indivisis insignis.

63. ***S. Heldreichii*** (Boiss. Diagn. Ser. II, 3, p. 158) glabra, caulis rubris obtuse quadrangulis simplicibus in thyrsum paniculatum multiflororum abeuntibus, foliis præter primaria 1-2 ovata parva indivisa cito evanida ambitu ovato-oblongis in laciniulas conflertas parvas oblongas inferiorum obtusiusculas superiorum angustas acutas bipinnatisectis, inferioribus breviter petiolatis, cymis breviter pedunculatis bifidis laxe 5-9-floris vix glandulosis, pedicellis calyce brevioribus, bracteis et bracteolis squamiformibus, calycis laciniis ovatis vel oblongis membranâ albâ denticulatâ anguste marginatis, corollâ parvâ purpureâ calyce vix duplo longiore lobis superioribus rotundatis, antheris exsertis, appendice parvâ anguste oblongâ, capsulâ minutâ globosâ calyce duplo longiore apiculatâ (2).

Hab. in regione inferiore montis Davrosdagh Pisidie (Heldr. exs. *S. flicifolia* var. *minor*!), in regione alpinâ montium Elmada Lycie (Bourg!).

Semipedalis vel pedalis caulis a tertia parte floriferis, cymæ rigidule ramoscæ, flores parvi. Foliis temiter sectis affinis *S. laxe* et *S. myriophylla* a quibus ceteris neglectis appendice statim distinguitur. Facie quoque affinis *S. Hoppii* floribus capsulaque duplo majoribus, corollâ calyce triplo longiori, etc., distinctæ.

64. **S. pruinosa** (Boiss. Diagn. Ser. I. 12, p. 38) biennis, radice verticali subsimplici. caule unico vel 2-3 erectis inferne foliosis a parte inferiori in thyrum simplicem vel in paniculam ramosam abeuntibus, foliis carnosis punctis minimis calcareis elevatis crebris pruinoso-farinosis glaucis petiolatis subtus elevatim nervosis ovatis grosse crenato-lobatis basi interdum laciniulis 1-2 auctis lobis saepius minute et crispule denticulatis, cymis petiolatis bifidis laxifloris ramis divaricatis rigidulis, bracteis oblongo-linearibus, bracteolis squamiformibus, pedicellis glabris calyce 1 1/2-2 plo longioribus, calycis laciniis ovatis membranâ albâ æquilateralâ undulatâ denticulatâ cinctis, corollâ calycem parum excedente urceolatâ, antheris vix exsertis, appendice ovato-oblongâ, capsula globosâ obtusâ longe mucronata calycem parum excedente ②.

Hab. in schistosis montis Elbrus Persie borealis prope Derbend (Ky. 438!), in montibus Elvend et Kaliser supra Masibin 7000' (Haussk!).

Semipedalis vel pedalis, folia ad caulis partem inferiorem subcongesta 1-2-pollices longa, flores et capsulae mediocres; species indumento et calyce late membranaceo insignis.

+ + Perennes.

65. **S. glauca** (Decaisne in Anch. exs. — p.c., Prodr. X, p. 310) glaberrima glauca e rhizomate lignescenti pluricaulis, caulis simplicibus fragillimis teretiusculis, foliis carnosis ovatis obtusis grossè et obtusissime paucicrenato-incisis, inferioribus petiolatis cæteris oppositis subsessilibus et sessilibus cordatis, thyro brevi pyramidalo glaberrimo, bracteis ellipticis acutiusculis integris, cymis pedunculatis laxe 5-9-floris, bracteolis linearibus minimis, pedicellis calyce subduplo brevioribus, floribus parvis, calycis laciniis ovatis latiuscule albo-marginatis, corollâ calyce vix duplo longiore, antheris subexsertis, appendice parvâ ovatâ acutiusculâ, capsula globosâ mucronata ȝ.

Hab. in fissuris rupium Persie australis, inter Tarun et Darap (Anch. 5064!), in monte Sabst Buschom prope Schiraz (Ky. 373!), ad rupes Teng Machmnd ditionis Bachtiaris et ad Kotel Kumaredj 3000' (Haussk!). Fl. Mart., Apr.

Caules cum thyro 6-10-pollicares, folia *Veronicae urticæfolia* sed obtusiora intermedia 1-1 1/2-pollicem longa, flores et capsula parvi. A formis integrifoliis *S. heterophyllæ* foliis caulinis sessilibus cordatis, floribus parvis, appendice, etc. distineta.

66. **S. prasiifolia** (Boiss. et Haussk.) glaberrima pallide virens, caulis e rhizomate lignescenti pluribus ascendentibus simplicibus in thyrum parte inferiori foliosum elongatum multiflorum abeuntibus, foliis ovato-oblongis basi rotundatis vel subcuneatis obtusis obtuse et late inciso-crenatis inferioribus breviter petiolatis superioribus sessilibus, floralibus ellipticis integris acutis, cymis crebris pedunculatis laxe 5-9-floris, bracteolis squamæforinibus acutis, pedicellis calyce duplo longioribus, calycis laciniis ovatis latiuscule sca-

riosis, corollâ calyce duplo longiore, antheris exsertis, appendice parvâ ovalâ acutiusculâ, capsulâ.... ♀.

Hab. in rupibus calcareis praeeruptis ad Schahpur et Kotel Mallu Persiae australis (Haussk!). Fl. Apr.

Glabritie, floribus parvis affinis *S. glauca*, differt foliis oblongis non cordatis, thyro inferne folioso densiore longiore multifloro

67. S. Pinardi (Boiss. Diagn. Ser. I, 4, p. 70) e rhizomate lignoso multicaulis praeter inflorescentiam parce glandulosam glabra, caulis elongatis fragillimis tenuibus simplicibus vel ramosis in paniculas breves multifloras basi foliolosas abeuntibus, foliis membranaceis petiolatis oblongis grosse pinnatipartitis vel pinnatisectis segmentis oblongis obtuse lobulatis, foliis supremis parvis subsessilibus 2-3-dentatis, cymis pedunculatis 7-3 floris, ramis pedicellisque calyce 2-3-plo longioribus filiformibus, calyeis laevis ovatis anguste membranaceis, corollâ purpureâ calyce quadruplo longiore, antheris exsertis, appendice parvâ ovato-oblongâ, capsulâ globosâ subdidymâ calyce duplo longiore mucronulatâ ♀.

Hab. in rupestribus Cariæ (Pinard!), Pamphyliæ ad rupes maritimæ Adalia (Bourg. 177!).

Semipedalis vel pedalis fragilitate formas pinnatisectas *S. heterophylle* referens sed panicula ob ramos cymarum et pedicellis tenues elongatos multo gracilior, corolla calyce dimidio minore anguste membranaceo 3-4-plo nec duplo longior, antheræ longius exsertæ, appendix non reniformis. A *S. trichopoda* quoque affini pedicellis brevioribus, appendice, etc. distincta.

68. S. variegata (M.B. Taur. Cauc. II, p. 78) e rhizomate indurato multicaulis tota pruinosa-punctata vel pubescens rarius glabra, caulis rubellis simplicibus vel parce ramosis in thyrsi laxum aphyllum abeuntibus, foliis ambitu oblongis profunde pinnatisidis-partitisve interdum pinnatisectis laevis lanceolatis vel linearibus inciso-dentatis, bracteis elliptico-linearibus, bracteolis minimis, cymis breviter pedunculatis 5-1-floris, pedicellis calyce subbrevioribus, floribus parvis, calyeis laevis orbiculatis albo-marginalis, corollâ variegatâ urceolatâ, antheris valde exsertis, appendice ovato-spathulatâ rarius oblongâ, capsulâ parvâ globoso-depressâ apiculatâ calyce triplo longiore ♀. *S. laciniata* C. Koch Linn. XXIII, p. 710 non W. K.

Hab. in saxosis totius Caucasi 2500'-7000' (M.B., C.A. Mey! Rupr!), Iberia et prov. Talysh (Hoh!), Persia bor. in prov. Aderbidjan (Szov!).

S. cinerascens. — Tota crebrius pruinosa canescens, folia profunde incisa vel pinnatipartita. — *S. cinerascens* Boiss. in Tehih. As. Min. II, p. 46. — *S. laciniata* var. *adenophora* C. Koch Linn. XXIII, p. 710.

Hab. in monte Paryadres Ponti (Tchiht!), Armeniâ Turcicâ (C. Koch!), prope Baibut (Bourg!).

S. rupestris. — Pruinosa vel puberula, folia dentata vel incisa

sed indivisa. — *S. rupestris* M.B. Taur. Cauc. II, p. 79. — *S. Ani* C. Koch Linn. XVII, p. 285! — *S. chamædrifolia* Boiss. et Haussk. MSS.

Hab. in rupestribus Tauriae meridionalis (Stev!), Anatoliae borealis ad Tokat (Auch. 503!), Armeniae Turcicae (C. Koch! Huet!), ad Baibut et Gummusch Khané (Bourgt!), Cappadociae in monte Alidagh (Bal!), Cataoniæ montibus Baltandagh et Berytdagh (Haussk!).

♂. *Libanotica*. — Procerior, præter thyrsum elongatum parce glandulosum glabra. Folia dentata, rarius incisa. *S. multifida* β *densiflora* Bth in D.C. Prodr! — *S. Libanotica* Boiss. Diagn. Ser. I. 12, p. 36. — *S. Urvilleana* Decaisn. Fl. Sinait. non Wydl. — *S. Persica* Bth. ex parte. — *S. incisa* C. Koch non Weinm.

Hab. in monte Sinaitico Stæ Catherinæ (Bové! Schimp. 350!), in Libano (Ehrenb!), supra Cedros (Boiss! Ky. 310!), Antilibano supra Zebdani (Boiss! Ky. 20!), monte Cassio (Boiss!), Tauro Cilicico (sub *S. melissæfoliæ*! Ky! Bal!), monte Ssoffdagh Syrie borealis (Haussk!), monte Gara Kurdistaniae (Ky. 316!), montibus Pir Omar Gudrun, Avroman, Schahu (Haussk!), montibus Persie borealis (Ky. 2981 324!, Buhsel, Auch. 5064!).

Species foliis indivisis et varie partitis vel sectis valde polymorpha, hujus formæ foliis partitis cum *S. laciniatæ* saepè confusæ ab eâ floribus fere duplo minoribus, filamentis longe exsertis et appendice facile distinguuntur ut et formæ foliis indivisis eisdem notis a *S. pulverulentæ*, *Mesogitanæ* et affinibus.

69. **S. puberula**, tota sub lente minute pruinosa-puberula glaucescens pluricaulis, caulinis erectis virgatis parte inferiori foliosis in thyrsum elongatum angustum rigidum abeuntibus, foliis subsessilibus anguste linear-lanceolatis acute pinnatifidis, cymis brevissime pedunculatis breviter et rigide bisidis 7-3-floris, bracteis bracteolisque squamiformibus, floribus parvis, calycis minuti laciniis orbiculatis albo-marginalis, corollæ purpurascens calyce triplo majoris lobis superioribus rotundatis, antheris exsertis, appendice parvâ oblongâ acutâ, capsulâ.... ♀.

Hab. in Persia boreali (Auch. 5066!).

A cl. Bth. in D.C. Prodr. adnumerata *S. multifidae* Willd. quæ *S. laciniatae* W. K. varietatem sistit! Affinis *S. variegatae* a cuius formis differt foliis angustis acute incisis et præsertim appendice acutâ.

70. **S. hyssopifolia** (Boiss. et Haussk.) tota minute pubescenti-glandulosa, caulinis ascendentibus pumilis a basi stricte ramulosis cerebre foliosis in thyrsum stricte paniculatum brevem foliosum abeuntibus, foliis carnosulis subsessilibus basi attenuatis acutis inferioribus oblongo-lanceolatis basi utrinque interdum dente crasso triangulari auctis superioribus linearibus integris, cymis stricte pedunculatis inferioribus laxe 2-3-floris cæteris unifloris, bracteis elongatis linearibus bracteolis brevibus, pedicello supra bracteolas subnullo, floribus mediocribus, calycis glabri laciniis oblongis membranâ angustâ lacerâ marginatis, corollâ e seco rubrâ calyce 2 1/2-plo longiore, antheris inclusis, appendice oblongo-lineari canaliculatâ, capsulâ crassiusculâ ovato-globosâ mucronatâ calyce subtriplo majore ♀.

Hab. ad rupes montis Berydaghi Cataoniae 7000' (Haussk!), in Karduchiâ ad hospitium Chana Putkie (Ky. Suppl. 714!).

Caules cum thyrso 4-pollicari 6-10-pollicares, folia inferiora fere pollicaria 3 lineas lata, superiora minora et angustiora; species inflorescentia stricta, bracteis elongatis intermixta, foliis subintegris, appendice notabilis.

71. *S. thesioides* (Boiss. et Buhse Anz. p. 164) glabra multicaulis caulis elatis tenuiter virgatis subquadrangulis ramosis in paniculam longam gracilem foliosam abeuntibus, foliis in laciniis angustissime lineares 1-3 jugas pinnatiseptis terminali longissimâ, foliis floralibus indivisis cymarum pedunculos æquantibus, cymis bifidis laxe 3-5-floris rami gracillimi. pedicellis calyce brevioribus, calycis laciniis orbiculatis membranâ latâ lacerâ marginatis, corollæ purpureæ calyce subdupo longioris lobis superioribus rotundatis, antheris exsertis, appendice oblongâ, capsulâ.... ♀.

Hab. in fauibus salsuginosis Armeniæ Rossicæ prope Nackitschewan (Buhsei), prope Karababa (Szovt).

Caules erecti pedales et sesquipedales gracillimi rigiduli, foliorum laciniæ linea $\frac{3}{4}$ vix latæ terminalis pollicem vel amplius longa, flores parvi. Species distinctissima.

*** Appendix anguste linearis. Perennes.

72. *S. canina* (L. Sp. p. 865) glabra multicaulis caulis virgatis rigidis elongatis simplicibus vel ramosis, foliis pinnatiseptis inferioribus petiolatis segmentis inferiorum oblongis vel obovatis saepe confluentibus dentatis vel incisis superiorum sessilium angustis lanceolatis vel linearibus acutis lobatis, thyrso longe multifloro saepe paniculato, cymis pedunculatis semel vel pluries bifidis laxe 9-3 floris, bracteis bracteolisque anguste linearibus, pedicellis calyce brevioribus, calycis parvi laciniis orbiculatis late scariosis, corollæ purpureæ lobis superioribus margine albidis, staminibus exsertis, appendice anguste linearis rarius nullâ, capsulâ globosâ apiculatâ calyce duplo longiore ♀. — *S. bicolor* Fl. Gr. VI, p. I, tab. 602. — *S. chrysanthemifolia* Willd. Hort. Berol. tab. 59 et M.B. II, p. 78. — *S. multifida* Lam. Fl. Fr. non Willd. — *S. pyramidalis* Wydl. Mon. p. 165. — *S. ramosissima* Urv. Enum, p. 75 non D.C.!

Hab. in siccis, collibus, ad muros regionis inferioris et montanae, Graecia in Peloponneso et Atticâ (Sprun! Boiss! Heldr!), Cephaloniâ (Heldr!), Zacyntho (Marg!), Macedonia (Friv!), Thraciâ (Griseb.), Byzantii (Auch. 1760!), insula Chio (Orph! Fl. exs. 727!), Seyros (Urv!), Cariâ (Pinard!), Lyciâ (Bourg!), Bithyniâ et Paphlagoniâ (Griseb!), Tauriâ (Stev.).

β. *floribunda*. — Sub lente minute pruinosa, folia in laciniulas minimas confertas bipinnatipartita vel secta, calycis laciniæ ovato-oblongæ latius scariosæ laceræ, corollæ lobis superiores magis porrecti et staminia longius exserta. — *S. floribunda* Boiss. et Bal. Diagn. Ser. II, 3, p. 158.

Hab. in Anatoliâ circa Smyrnam (Auch. 1777! Boiss! Bal! 334!).

Species foliorum divisione et laciniarum formâ valde variabilis sed inter omnes formas transitus observantur. Hujus area ad Orientem ultra Tauriam et Anatoliæ occidentalem et borealem non progredi videtur et specimina ex Caucaso et Syriâ huc relata ad species diversas spectant

Ar. Geogr. Europa media et australis tota a Belgio, Germaniâ australi ad reginem Danubialem et Rossiam australiæ, Africa borealis.

73. S. Cabulica (Bth. in D.C. Prodr. X, p. 316) glaberrima multicaulis glaucescens, caulis erectis rigidis parte inferiori parce foliosis in thyrsos elongatos rigidos abeuntibus, foliis oblongo-lanceolatis obsolete repando-dentatis, cymis 5-7-floris inferioribus pedunculatis divaricatim bifidis supremis abbreviatis subsessilibus, floribus parvis remotis sessilibus, calycis laciniis obtongis obtusis vix marginatis, appendice linearie ♂.

Hab. ad Bamean regni Cabulici in alveis torrentiam (Griff. 3874! in herb. Kew!).

Ex speciminibus valde imperfectis nota, affinis *S. caninæ* sed folia fere integra.

§ 2. Sparsifoliæ. — Folia omnia sparsa.

74. S. crassicaulis (Boiss. Diagn. Ser. I, 42, p. 37) glaberrima pluricaulis, caulis crassis medullâ farctis erectis parce foliatis a parte inferiori ex axillis ramosis ramis strictis elongatis sterilibus, foliis omnibus alternis integerrimis obovato-spathulatis et linearibus obtusis, ramorum linearibus angustatis brevibus, thyro terminali laxo paucifloro, cymis inferioribus subtrifloris cæteris unifloris omnibus pedunculis longis crassis suffultis, pedicellis calyce brevioribus, bracteis linearibus acutis, bracteolis brevibus margine scariosis, floribus magnis, calycis laciniis ovatis latissime albo-marginatis, corollâ purpureâ, staminibus inclusis, appendice late reniformi, capsula magnâ globosâ obtusissimâ calyce sesquilongiore mucronata ♂.

Hab. in valle Talagon montis Elbrus Persiæ borealis (Ky. 523!).

Species insignis facie *Centranthi*. sesquipedalis, folia intermedia sesquiplicaria $2\frac{1}{2}$ lineas lata, infima multo breviora, rami laterales apice squamulosi, capsula pisum æquans vel superans.

75. S. Syriaca (Bth. in D.C. Prodr. X, p. 316) glabra basi suffrutescens pluricaulis, caulis præter folia rara parva squamiformia oblonga integra vel dentata aphyllis dicotome ramosis ramis in thyrsos valde depauperatos abeuntibus, cymis breviter pedunculatis bifidis 3-5-floris, bracteis bracteolisque minatis triangularibus, floribus sessilibus parvis, calycis laciniis orbiculariatis late albo-marginatis, staminibus inclusis, appendice minimâ oblongo-lineari, capsula ♂.

Hab. in Palestinâ circa Nazareth (Bové), loco non indicato (Auch. 1776!).

Pedalis, foliis parvis (4-5 lineas longis) integris insignis.

76. S. Benthamiana (Boiss. Diagn. Ser. I. 12, p. 39) basi suffrutescens pluricaulis glaberrima glauca, caulinis quadrangulis rigidis flexuosis fere a basi dichotome ramosissimis, foliis omnibus sparsis carnosis paucis parvis oblongis vel ellipticis inferioribus basi sensim in petiolum brevem attenuatis remote denticulato-serrulatis dentibus ascendentibus, superioribus diminutis sessilibus integris, cymis rigide crasse et breviter pedunculatis inferioribus trifloris ceteris unifloris, bracteis et bracteolis squamiformibus, floribus parvis sessilibus, calycis laciniis orbiculatis anguste albo-marginatis, corollæ brevis calyce sesquilocularis lobis superioribus parvis rotundatis, anthers subexsertis, appendice parvâ oblongâ, capsulâ ovato-sphaericâ acutiusculâ mucronatâ calyce duplo longiore ♀.

Hab. in Persiâ, in monte Sofi prope Ispahan (Bode! Derderian! Auch. 5031 et 5032!), ad radices montis Demavend (Ky. 6301), in prov. Khorassan et inter Kerman et Yezd (Bge!).

Pedalis rigida ramosissima, folia inferiora cum petiolo vix pollicaria apice 3-4 lineas lata. Facies *S. caninæ*, folia eis *S. Syriacæ* triplo majora et serrulato-dentata.

77. S. hypericifolia (Wydl. Mon. p. 166, tab. 5) frutescens glaberrima ramosissima ramis rigidis erecto-patulis basi foliolosis in racemum simplicem abeuntibus tandem persistentibus subspinosis, foliis sparsis parvis carnosulis integris margine interdum paucicallous inferioribus obovatis vel oblongis subpetiolatis ramis dtminutis, pedicellis calyce sublongioribus minutissime subbibracteolatis, floribus minutis, calycis laciniis orbiculatis latiuscule scariosis, corollâ calyce duplo longiore, staminibus exsertis, appendice linearis acutâ, capsulâ globosâ calyce subtriplo longiore ʒ.

Hab. in Mesopotamiâ inter Aleppo et Bagdad (Oliv!).

Frutex pedalis vel elatior, rami inferne pennâ anserinâ subtenuiores, folia 4-6 lineas longa, racemi 3-4-pollicares interdum semipedales. Species ramis fruticosis albis et cymis ad pedicellos unifloros racemosos dispositos reductis insignis.

78. S. leucoclada (Bge pl. Lehm. p. 424) glabra fruticosa ramosa ramis velutinis albo-corticosis, horuotinis inferne foliosis simplicibus vel ramosis in thyrsos elongatos multifloros abeuntibus, foliis omnibus sparsis oblongo-linearibus basi angustatis subsessilibus integerrimis, cymis numerosis brevissimis petiolatis bis vel ter dichotomis 11-3-floris, pedicellis calyci subaequilongis, calycis laciniis orbiculatis latiuscule membranaceo-marginatis, corollâ . . . capsulâ globosâ enspidatâ calyce duplo longiore ʒ.

Hab. ad rupestris graniticis Bakali deserti Kisylkum Turkestaniæ (Lehm!). Vidi frustul. sterile.

Frutex humilis ramosus ramis horuotinis cum thyrsos plus quam pedalibus, folia inferiora 9-13 lineas longa 1½-2 lata, capsula sesquilineam diametro lata. Caulibus fruticosis albis insignis, appendix ignota sed prob. affinis *S. hypericifolia* a qua differt foliorum formâ et inflorescentiâ.

SUTERA (Roth Bot. Bemerk. p. 172).

Calyx quinquepartitus laciniis linearibus. Corollæ deciduae tubus subæqualis, limbus parvus erecto-patulus laciniis integris superioribus subminoribus. Stamina 4 fertilia inclusa, antheræ subglobosæ confluentim uniloculares. Stylus apice breviter bilobus. Capsulæ valvæ apice bifidæ, placenta demum libera tereti-compressa. Semina numerosa minima foveolato-punctata. — Herba annua viscidula parviflora, foliis præter infima opposita alternis.

1. **S. glandulosa** (Roth loc. cit.) tota breviter glandulosohirta viscidula a collo ramosissima ramis præter centrale diffusis pumilis in racemos foliatos laxos abenntibns, foliis breviter petiolatis ambitu ovatis vel oblongis inciso-pinnatifidis vel dissectis laciniis incisis dentieulatis, foliis floralibus diminutis, pedicellis axillaribus strictis calycei æquilonigis folio subbrevioribus, corollâ glabrâ calyce tertia parte longiore, capsulâ oblongâ calycem parum excedente ♂. Ie. Wight III. tab. 856. — *Capraria dissecta* Del. descr. II, p. 239, tab. 32, fig. 2. — *Sutera dissecta* Walp.

Hab. ad ripas arenosas Nili in Ægypto superiore (Del., Sieb!, Auch. 1887! Boiss!).

Ar. Geogr. Nubia, India australis.

ANTICHRIS (Endl. Gen. p. 682. — Ascherson Ac. Wiss. Berl. 1866. — *Meissarhena* R. Br. — *Doratanthera* Bth. in Endl. Gen. 682. — *Distemon* Ehrenb. et Hemprich MSS.).

Calyx quinquepartitus. Corollæ deciduae tubus subcurvatus, faux elongata ampla, limbi quinqueffdi erecto-patentis lobi subæquales rotundati. Stamina bina, antica, inclusa, corollæ supra baseos constrictiōnem iuxta, antheræ magnæ loculis apice confluentibus. Stylus apice brevissime retuso-bilobus. Capsula ovato-oblonga acuminata valvis bipartitis, columna placentaris demum libera lineari-cylindrica. Semina numerosa oblonga longitudinaliter multicostata. — Herbae annuae viscidæ foliis alternis.

i. **A. Arabica** (Steud. et Hochst. in Schimp. Arab. exs. 1837 sub *Capraria*) breviter glandulosohirta, caule erecto pumilo a basi parce ramoso, foliis oblongo-linearibus acutiuseculis in petiolum brevem attenuatis, floribus in omnibus axillis solitariis pedicello filiformi versus medium bracteolato calyce sublongiore folio multo breviore suffultis, bracteolis setaceis, calycis laciniis elliptico-linearibus, corollâ roseâ viscidulâ calyce triplo longiore, antheræ lobis basalibus obtusis inæqualibus, capsulâ in rostrum abrupte attenuatâ calyce

duplo longiore ⊙ *Ant. Arabica* Endl., Icon. tab. 93. — *Meissarhena tomentosa* B. Br. in Salt Abyss. — *Distemou campanularis* Ehr. et Hempr. MSS.

Hab. in collibus Belutschiae (Stocks!), verosimiliter quoque ad littus Arabicum Egypti superioris.

Ar. Geogr. Nubia littoralis, Abyssinia, Arabia felix.

2. A. glandulosa (Aschers. Bericht. Ac. Wissensch. Berlin 1866, p. 880) patule longe et copiose glanduloso-hispida, caule pumilo a basi ramoso flexuoso, foliis ovatis et oblongis obtusis in petiolum brevem attenuatis, floribus in omnibus axillis solitariis pedicello filiformi versus medium bracteolato calyce sublongiore folio breviore suffultis, bracteolis lanceolatis basi attenuatis, calycis laciniis lineari-lanceolatis acutis, culis basi attenuatis, corollâ roseâ glaberrimâ calyce 2¹,₂-plo longiore, antherae lobis basalibus obtusis inaequalibus, capsula sensim acutatâ calyce sesquilociore ⊙. *Distemou glandulosus* Ehr. et Hempr. MSS.

Hab. in Egypto superiore littorali seens Mare Rubrum inter Kosseir et Ras Benass (Schweinf.), in deserto Arabicu Egypti (Husson!).

Affinis praecedenti, flores paulo maiores, indumento, foliorum formâ, etc diversa.

Ar. Geogr. Arabia tropica, regio Scinde Indiae.

3. A. linearis (Bth. in p.c. Prodr. X. p. 347 sub *Doratanthera*) siccitate nigricans tota brevissime hirtella viscida pumila erecta a basi ramosa, foliis longe et anguste linearibus uninerviis acutis, floribus in omnibus axillis solitariis pedicello filiformi calyce 2-4-plo longiori folio breviore suffultis, bracteolis in pedicelli parte superiori sub flore sitis setaceis, calycis laciniis lineari-lanceolatis acutis, corollâ roseâ hirtâ calyce triplo longiore, antherae lobis basalibus acuminatis subaequalibus, capsula oblongâ sensim acutatâ calyce sesquilociore ⊙. *A. linearis* Hochst. in Schimp. pl. Nub. — *Distemou angustifolius* Ehr. et Hempr. MSS.

Hab. in Egypto superiore (Bth. in p.c. Prodr.), Belutschia (Stocks!).

Anticharidi Arabicae et glandulosae omnino congener, antherarum forma eadem.

Ar. Geogr. Insulae Capitis viridis, Senegalalia, Nubia, Abyssinia, Arabia tropica, India borealis.

Tn. IV. GRATIOLEÆ (Bth. in p.c. Prodr.). — Corolla tubulosa vel campanulata non saccata. Antheræ biloculares. Capsula valvis 2-4 dehiscentia. Inflorescentia centripeta. Folia opposita.

MAZUS (Lour. Fl. Cechinch. p. 385).

Calyx campanulatus quinquesfidus. Corolla bilabiata labio superiori erecto breviter bifido, inferiori multo majori patenti trifido fice

bigibboso. Stamina fertilia 4, antherarum loculi divergentes. Stylus apice bilamellatus lobis ovatis æqualibus. Capsula ovata obtusa compressa valvulis dinabus integris. Placenta demum libera ovata. Semina numerosa lœvia. — Herbæ foliis infimis oppositis cæteris alternis.

M. rugosus (Lour. loc. cit. p. 385) pumilus glabriusculus sèpe stolonosus collo ramosus ramis erectis vel diffusis in racemos simplices aphyllos demum flexuosos laxos abeuntibus, foliis obovatis vel oblongo-œcatis grosse et obtuse dentatis radicalibus rosulatis, pedicello calyci subæquilongo, calycis aperte campanulati laciniis late lanceolatis tubo sublongioribus, corollâ calyce subtriplo longiore ♀. *Hornemannia bicolor* Willd. Enum. p. 654. — Rehb. Exot. t. 37.

Hab. in regno Cabulico (Griffith).

Ar. Geogr. India orientalis, China, Japonia, Sibiria Uralensis.

DODARTIA (L. Gen. p. 322).

Calyx campanulatus breviter quinquedentatus. Corolla elongata tubo sensim dilatato, labium superius ovatum retusum, inferius multo majns obovatum trifidum fave bigibbosâ. Stamina fertilia 4, antherarum loculi contigui divergentes. Stylus apice bilamellatus lobis ovalis æqualibus. Capsula globosa obtusa subindehiscens bilocularis, placentæ crassæ carnesæ. Semina numerosa lœvia. — Herba pereennis scoparia subaphylla foliis inferioribus oppositis.

D. Orientalis (L. Sp. p. 883) glaberrima, caulinis e rhizomate indurato ramoso numerosis erectis opposite et alternatim ramosissimis ramisque junceis subaphyllis, foliis infimis oppositis oblongis parvis sessilibus cito evanidis cæteris ad axillas breviter et anguste linearibus squamiformibus, racemis terminalibus laxe 3-7-floris, pedicellis bracteisque brevissimis, calycis dentibus triangularibus acutis, corollâ purpureâ extus glabrâ calyce quintuplo longiore, labio inferiore ad faveum intus borbato, stigmate exerto, capsulâ apienlatâ ♀. Ic. Gærtn. Fr. I, tab. 53 — Bot. Mag. tab. 2199. — Jaub. et Sp. III, tab. 410.

Hab. in campestribus subsalsis Caucaso bor-ali et australi adjacentibus (M.B., C.A. Mey!), Iberiâ Caucasicâ et prov. Karabagh (Hohen!), Armeniâ (Tourn., C. Koch), Persiâ (Jenisch!), Turkestaniâ (Lehm!), prov. Aderbidjan (Szov! Bge!).

Ar. Geogr. Rossia media et australis, Songaria, Sibiria Uralensis et Altaica.

LINDENBERGIA (Lehm. in Link et Otto Abbild. tab. 95. — *Bovea* Decaisne Fl. Sinaica).

Calyx tubulosus vel campanulatus quinquedentatus dentibus subinæqualibus. Corolla elongata, labium superius oblongum concavum

erectum retusum, inferius majus trilobum fauce prominulâ biplicatâ. Stamina fertilia 4, antherarum loculi disjuncti stipitati. Stylus apice subelevatus truncato-capitatus stigmatosus. Capsula oblonga vel ovoidea bisulca loculicide bivalvis, valvae integræ medio septiteræ, placenta centralis demum libera 4-partibilis, semina numerosa minima. — Suffrutices vel herbae foliis oppositis.

1. **L. Sinaica** (Decaisn. Fl. Sinaic. Ann. Sc. Nat. Ser. II, 2, p. 253 sub *Boved*) patule glanduloso-hispida basi fruticosa opposite ramosa ramis hornotinis virgatis inferne foliosis in racemos longos simplices laxos abeuntibus, foliis breviter petiolatis a basi rotundata vel truncatâ ovatis obtusis grosse crenato-dentatis, floralibus subsessilibus calyce minoribus, floribus oppositis secundis brevissime pedicellatis, bracteolis nullis, calycis hispidi viscidè dentibus triangularibus acutis, corollâ violacea hispidulâ calyce quadruplo longiore, fauce intus barbatâ, capsulâ hirtâ acutâ calyce sublongiore 5. *L. Sinaica* Bth. Scroph. Ind. p. 22.

Hab. ad rupes in Egypto littorali Arabico inter Kosseir et Ras Benass (Schweinf.), in valle Sancti Pauli ad scaturigines (Husson!), in Arabiae iugno Sinaitico ad Ouadi Hebran (Ehrenb! Bové! Schimp. 183!).

Frutex bipedalis racemis $\frac{1}{2}$ -4-pedalibus, folia ramorum novellorum pollucem longa superiora multo minora, corolla 7-8 lin. longa. Facies *Scutellariae* vel *Melissæ*.

2. **L. fruticosa** (Bth in D.C. Prodr. X, p. 377) breviter velutino-hirtella basi suffrutescens ramosa, rsmis tenuibus fragilibus totâ longitudine racemosis foliosis, foliis parvis ovatis paucicrenatis floralibus vix diminutis omnibus petiolatis, floribus axillaribus oppositis pedicello petiolo aequilongo suffultis, calycis parce papilloso-hispiduli dentibus triangularibus acutiusculis, corollâ roseâ glabrâ calyce triplo longiore fauce intus barbatâ, capsulâ acutâ hirtâ calyce sesquiongiore 5.

Hab ad rupes regni Mascate Arabiae (Auch. 5104!).

Afinis *L. Sinaicae* a quâ forsitan non sat differt, indumentum brevius minus viscidum, folia minora, floralia petiolata 2-2 $\frac{1}{2}$ lin. longa, racemus densior, flores paulo longius pedicellati, corolla extus glabra.

3. **L. urticæfolia** (Lehm. in Link et Otto Abbild. tab. 95) annua patule glanduloso-hirta erecta inferne ramosa, ramis fere a basi floriferis laxiuscule racemosis, foliis longe petiolatis ovatis obtusis obtuse crenatis floralibus conformibus subdiminutis, floribus oppositis axillaribus brevissime pedicellatis, calycis campanulati papilloso-hirsuti dentibus ovatis, corollâ glabrâ calyce 2 $\frac{1}{2}$ -plo longiore, capsulâ ovatâ acutâ hirsutâ calycem non excedente ⊖. *Stemodia littoralis* Vahl Symb. II, p. 69. — *Stemodia muraria* Roxb.

Hab. in Belutschia (Stocks ex Aitch. Cat. p. 107).

Ar Geogr. India

HERPESTIS (Gærtn. Fil. Suppl. p. 186).

Calyx quinquepartitus laciniis inæqualibus posticâ latiore. Corolla bilabiata labio superiore retuso vel bilobo inferiore trilobo, vel quinqueloba lobis subæqualibus. Stamina quatuor. antheræ loculis contignis parallelis vel divaricatis. Stylus breviter bilobus vel apice dilatatus capitatus. Capsula bisulca, valvulae 2 bipartitæ rarius integræ, placenta columnaris tandem libera. Semina numerosa parva. — Herbae foliis oppositis.

I. **H. Monnieria** (L. Sp. p. 24 sub *Gratiola*) glabra carnosula glaucescens, caulis repentibus elongatis ad axillas radicantibus, foliis subintegris obovato-cuneiformibus subsessilibus subnerviis, floribus ad axillas solitariis pedicello folio æquilongo vel longiore sub flore bibracteolato suffultis, calycis segmento postico ovato cæteris angustioribus, corollæ calyce sesquilonioris lobis ovatis subæqualibus, antheris subæqualibus sagittatis, stigmate capitato, capsulâ ovatâ acutâ calyce subbreviore ♂. *H. Monnieria* H. B. K. Nov. Gen. et Sp. II, p. 366. — Ic. Bot. Mag. tab. 2337. — *Limosella calycina* Forsk. Fl. Eg. Arab. p. 412.

Hab. in humidis et ad rivulos, regni Cabulici in paludibus Bussolah (Griff!), Arabiae Mascateensis (Auch. 5074!), Babyloniae ad Bagdad (Noë!).

Ar. Geogr. Arabia tropica, Africa, Asia et America tropicae.

GRATIOLI (L. Gen. p. 33).

Calyx subæqualiter quinquepartitus. Corollæ tubulosæ labio superiore integro vel bifido inferiore trifido pâlato non promiulo. Stamina 2 postica fertilia, 2 antica sterilia tiliiformia vel nulla, antherarum loculi distincti paralleli (in nostrâ) in connectivo dilatato peltiformi sessiles. Stylus apice bilamellatus. Capsula quadrivalvis, placenta columnaris demum libera. Semina numerosa reticulatofoveolata. — Herbae foliis oppositis.

I. **G. officinalis** (L. Sp. p. 24) glaberrima, rhizomate repente squamoso, caulis erectis crebre foliosis superne tetragonis, foliis semiamplexicanilibus lanceolatis subtrinerviis integris vel serrulatis, floribus axillaribus longe pedicellatis pedicello apice bibracteolato, calycis glanduloso-punctati laciniis lanceolatis, corollæ calyce quinque longioris tubo curvulo luteolo limbo roseo, fauce intus barbatâ, filamentis sterilibus elongatis, capsulâ ovatâ acutâ calyci æquilongâ ♀. Ic. Fl. Dan. tab. 363. — Rehb. Ic. tab. 56.

Hab. ad fossas et in paludosis, Macedonia et Thracia (Orph! Pichler!), Anatolia in Bithyniâ, Lyciâ, Isauriâ Lydiâ (Thirke! Bal! Bourg!), Transcaucasiâ (Szov!).

Ar. Geogr. Europa media a Daniā et Belgio ad Rossiam medium et ab Hispaniā boreali et Italiā ad regionem Danubialem et Dalmatiam, Sibiria Uralensis et Altaica, America borealis.

VANDELLIA (L. Mant. p. 89. — *Lindernia* All.).

Calyx quinquepartitus laciniis subaequalibus. Corolla tubulosa limbi labio superiore bifido inferiore majore trifido. Stamina 4 fertilia filamentis brevibus anticis basi appendiculatis. Antherae per paria cohaerentes vel libere loculis ovatis divergentibus. Stylus apice breviter bilamellatus. Capsula ovata bivalvis, valvulis indivisis, placenta columnaris tandem libera. Semina numerosa parva. — Herba foliis oppositis.

I. **V. pyxidaria** (All. Mise. Taor. III p. 178, tab. 5 sub *Lindernia*) glabra pumila basi sepe ramosa, foliis sessilibus ovatis vel oblongis integris obtusis trinerviis, pedunculis filiformibus ex axillis superioribus oppositis folio longioribus, laciniis calycinis linearibus, corollae labio superiore purpureo inferiore flavido, filamentis anticis basi obsolete glanduloso-appendiculatis, antheris non coherentibus, capsula calyce breviore, seminibus transverse foveolatis ⊖. *V. pyxidaria* Maxim. Pl. Japon. — *V. erecta* Blh. Seroph. Ind. p. 36.

β. grundiflora. — Maxim Prim. Fl. Am. p. 305. — Corolla calyce sesquilonigior aperta.

Hab. in fossis et locis hyeme inundatis, Pontus Lazicus prope Rhizé (Bal!), Iberia et ditio Talysh (M B., Hohen! Buhse!), Persia australis ad Teng Machmud et oryzeta Susiana (Haussk!).

Haec forma in Rossia australi, Serbia quoque occurrit, forma typica ex Europa, regione Amurensi, India differt corollae calyce breviore brevissimo tempore vel vix aperta.

Ar. Geogr. Gallia, Germania, Italia borealis, Serbia, Rossia australis, Sibiria Amurensis, India Orientalis.

PEPLIDIUM (Delile Fl. Eg. p. 423).

Calyx tubulosus quinqueangulatus breviter quinquedentatus. Corolla tubo brevissimo laciniis quinis subaequalibus. Stamina bina inclusa, filamentis brevibus basi subdilatatis, antherae uniloculares. Stylus brevis apice spathulato-dilatatus. Capsula ovata indehiscentis irregulariter ruptilis tenuiter membranacea. Columna placentaria libera. Semina numerosa angulata. — Herba procumbens foliis oppositis facie *Peplidis*.

I. **P. humifusum** (Del. loc. cit. tab. 4, fig. 2) glabrum ramosissimum ramis prostratis radicantibus, foliis parv's carnosulis ob-

ovatis integris enerviis breviter petiolatis, floribus in axillis oppositis subsessilibus minimis, corollâ calycem vix excedente ⊖. *Microcarpaea cochlearifolia* Sm. Hook. Bot. Misc. III, p. 95. Suppl tab. 29.

Hab. in hyeme inundatis, oryzetis Ægypti inferioris (Del., Sieb!), prope Alexandriam (Letourneux!).

Ar. Geogr. Africa tropica, India Orientalis, Nova Hollandia.

LIMOSELLA (L. Gen. p. 320).

Calyx campanulatus dentibus quatuor triangularibus. Corolla rotato-campanulata tubo brevissimo, limbo quinquepartito subbilabiato. Stamina in medio tubo inserta subæqualia, antheræ loculis confluentibus uniloculares. Ovarium inæquale, stylus obliquus, stigma capitatum. Capsula bivalvis basi tautum incomplete bilocularis. Columna placentifera libera. Semina numerosa costata. — Herbæ pusillæ cespitosæ vel reptantes glabrae.

1. **L. aquatica** (L. Sp. p. 881) foliis longe petiolatis anguste oblongis vel linearis-spathulatis, corollæ laciniis ovati-oblongis calyce subduplo longioribus, capsula calyce sublongiore ⊖. Ic. Fl. Dan. tab. 60. — Rehb. Ic. tab. 401.

Hab. in humidis, Egyptus (Schweinf. Beitr.).

Ar. Geogr. Europa fere tota, Sibiria, Abyssinia, America borealis, Australia.

Tr. V. DIGITALE E Bth. in n. c. Prodri. — Corolla tubulosa vel campanulata lobis planis. Antherarum loculi saepius apice confluentes. Capsula bivalvis. Inflorescentia centripeta racemosa. Folia alterna.

SIBTHORPIA (L. Gen. 320. — *Disandra* L. Fil.).

Calyx 4-8-fidus. Corolla subrotata tubo brevissimo, limbi laciniis tot quot calycis vel unâ additâ. Stamina numero laciniis corollæ aequalia vel uno deficiente, antheræ sagittatæ biloculares loculis apice contignis. Stylus apice capitatus. Capsula membranacea a latere compressa loculicide bivalvis. Semina dorso convexa facie umbilicata concava vel plana. — Herbæ prostratae radicantes teneræ foliis alternis vel fasciculatis.

1. **L. Africana** (L. Spec. p. 880) hispidula cespitosa ramis teneris filiformibus intricatis prostratis radicantibus, foliis longe petiolatis orbiculatis 7-9 crenatis crenis latis retusis, pedicellis axillaribus petiolos æquantibus vel brevioribus, calycis hirti laciniis triangula-

ribus acutis, corollæ luteæ laciniis ovatis calyce subduplo longioribus, antheris corollam subæquantibus, stylo glabro. φ . *Disandra Africana* Camb. Enum. Bal. p. 117, tab. 9.

Hab. ad fontes regionis montanæ, Laconia ad Hagios Petros (Boiss!), Creta (Sibth.), in castanetis humidis Euneachoria 1500' (Heldr!).

Valde affinis *S. Europeæ* quæ differt floribus minoribus, corollâ carneoâ calycem vix exceedente, ovario superne cum stylî basi barbatis. Cf. Cainbesedes in *S. Europaæ* discum quinquelobum a me nullibi visum indicat. Species distributione geographicâ insignis.

Ar. Geogr. Insulae Azoricae, Balearicæ, Abyssinia.

DIGITALIS (L. Gen. 313).

Calyx quinquepartitus. Corolla declinata tubo ventricoso campanulato vel cylindrico, limbi labium superius brevissimum retusum vel bifidum, inferioris lobus medius lateralibus latior et saepè longior. Stamina 4 didynama e corollæ fundo ascendentia inclusa, antherarum loculi divergentes apice confluentes. Stylus breviter bilobus. Capsula ovata septicide bivalvis, valvulis integris marginibus intreflexis placentam columnarem superne nudantibus. Semina numerosa minuta oblonga. — Herbæ perennæ.

§ 1. **Globifloræ** Bth. — Corollæ tubus abrupte inflatus globosus vel obconicus, labii inferioris lobus medius elongatus.

1. **D. ferruginea** (L. Sp. 867) caule elato folioso in racemum longum subsimplicem densiflorum vel laxiusculum abeunti, foliis glabris vel subtus et ad margines parce puberulis lanceolatis obscure repandis inferioribus in petiolum attenuatis, rachide glabrâ, floribus brevissime pedicellatis, bracteis lanceolatis deflexis calyce saepius brevioribus, ealycis glabri laciniis obovato-ellipticis obtusis scarioso-marginatis, corollæ glanduloso-pubescentis fulvae ferrugineo-reticulatae tubo subito inflato subgloboso, lobis superioribus abbreviatis rotundatis, lateralibus brevibus acutis inferiore porrecto tobum subæquante ovato integro vel obscure trilobo intus piloso. Ic. Fl. Græc. tab. 606. — Lindl. Mon. tab. 12 — Bot. Mag. tab. 1828. — *D. aurea* Linol. Mon. tab. 13. — *D. brachyantha* Griseb. Spic. II, p. 513.

Hab. in collinis et sylvaticis præsertim regionis montanæ et subalpinæ, Græcia in Laconia monte Malevo (Orph. Fl. Gr. exs. 40!), Taygeto et Parnasso (Heldr!), Macedoniâ (Frider.), Byzantii in monte Gigantis (Boiss!), Olympo Bithyno (Boiss!). Thnolo Lydiae (Boiss!, Bal. 337!), montibus Lyciae (Bourg!), Lyconiae (Heldr!), Anatoliâ boreali (Wied!), Ponto supra Trapezuntem (Bourg!) et Ponto Lazico ad Djimil (Bal!), Imeretiâ (Szov!), Armenia Turcicâ (Szov!), Libano (Bl!). In Tauriâ indicata sed ex Steven erroneâ.

Flores, præsertim in speciminibus ex Anatoliâ boreali, Ponto et Armenia variant iniores.

Ar. Geogr. Italia media et australis, Carnia, Dalmatia, regio Danubialis, Serbia.

2. *D. nervosa* (Steud. ex Hochst. in herb. Un. Itin. — Bth. in d. c. Prodr. X. p. 450) glabra, caule elato in racemum longum densiusculum abeunti, foliis oblongo-lanceolatis integris inferioribus petiolatis, floribus breviter pedicellatis, bracteis deflexis lanceolatis calyce brevioribus, calycis lacinis oblongis acutis vel acuminatis margine vix scariosis, corollæ parvæ lutescentis pubescentis tubo brevi subito ampliato lobis superioribus et intermediis abbreviatis inferiore porrecto ovato intus barbato tubo breviore ♀. ***D. laevigata*** C. A. Mey. Enum. p. 110 non W. K.

Hab. in Georgiâ Caucasicâ et ditione Karabagh, prope Lenkoran et in tractu Swant ditionis Talysh (Hohen!, Szov!), Persiae bor. prov. Ghilan (Buhse!).

Valde affinis formis parvifloris ***D. ferruginea*** a quibus lobo inferiori corollæ breviore et in primis lacinis calycinis acutis differt.

Ar. Geogr. Turcomania.

3. *D. laevigata* (W. K. Pl. rar. Hung. II, p. 171, tab. 158) glabra, caule elato in racemum sæpius simplicem elongatum laxiusculum abeunti, foliis. integris inferioribus oblongo vel obovato-lanceolatis in petiolum attenuatis, caulinis lanceolatis, floribus breviter pedicellatis, bracteis anguste lanceolatis calyce sublongioribus, calycis lacinis lanceolatis acutis et acuminatis margine subscariosis. corollæ flavidæ purpureo-venosæ extus puberulæ tubo subito inflato subgloboso inferne gibbo lobis superioribus et intermediis brevibus inferiore porrecto tubum subæquante oblongo repando intus piloso ♀. Lindl. Mon. tab. 10. — Rchb. Germ. tab. 71.

Hab. in sylvaticis regionis montanæ Græciæ, mons Malevo Laconia (Orph. Fl. Gr. exs. 41!), Taygetus supra conventum Gola (Heldr!), regio media montis Kyllenes supra Trikala (Heldr!), Parnassus et Olympus Thessalus (Heldr!), mons Korthiati Macedoniæ (Heldr!). Variat ut ***D. ferruginea*** floribus minoribus.

Ar. Geogr. Carnia, Dalmatia, regio Danubialis.

4. *D. lanata* (Ehrh. Beitr. 7, p. 23) caule glabro rubello, foliis glabriusculis vel ciliatis inferioribus oblongis petiolatis subrepandis cæteris lanceolatis integris, racemi densi multiflori rachide pubescente. floribus brevissime pedicellatis bracteam ciliato-lanatam lanceolatam subæquantibus, calycis lanati lacinis lanceolatis, corollæ villosæ albidae vel pallide ochroleucæ violaceo-reticulatae tubo subito inflato subgloboso lobis superioribus et intermediis brevissimis, inferiore porrecto ovato utrinque pubescente tubo subæquilongo ♀. Ic. W. K. Hung, tab. 74. — Lindl. Mon. tab. 15.

Hab. in regione sylvatica Parnassi (Sprun!), montis Korthiati Macedoniæ (Heldr! Orph. Fl. Gr. exs. 726!), Thraciâ circa Kalofor (Friv! Pichler!).

Ar. Geogr. Regio Danubialis.

5. **D. Orientalis** (Lam. Dict. II, p. 280) caule glabro in spicam laxam interruptam abeunti, foliis glabris sessilibus anguste linearibus subgramineis acuminatis in curvo-subflexuosis, rachide glandulosovallosa, floribus remotis subsessilibus, bracteis lanceolatis acuminatis inferioribus calycis longitu line crerteris brevioribus, calycis lanuginosi laciniis ellipticis obtusis, corollæ albidae pilosæ tubo subito inflato obconico subtus valde gibbo lobis superioribus et intermediis obsoletis inferiore porrecto oblongo-spathulato obtuso tubo æquilongo ♀. Ic. Bot. Mag. tab. 2253. Lindl. Mon. tab. 16 mala (forma hybrida?) — Jaub. et Sp. Ill. tab. 409.

Hab. in monte Hæmo Thraciae (Friv. ex Griseb.), Bithynia prope Bolu (Pest!), Galatia et Paphlagonia (Wiedem!), Ponto (Tchih!), Armenia ad Gumusch Khané (Bourgl.).

Ab affini *D. lanatâ* foliis angustis et floribus dissitis distincta.

6. **D. Cariensis** (Boiss in Pinard exs. 1843 — Diagn. Ser. II, 3, p. 159) caule glabro in spicam brevem densiusculam abeunti, foliis glabris linear-lanceolatis sessilibus acuminatis, rachide brevissime pruinoso-puberula. bracteis velutinis lanceolatis acuminatis calyces æquantibus, floribus breviter pedicellatis, calycis breviter velutini laciniis lanceolatis acutis longe mucronatis, corollæ parvæ pubescentis e sicco ochroleucæ vel albidae tubo subito ampliato obconico, lobis superioribus et intermediis brevissimis inferiori ovato-spathulato tubo subbreviore ♀. *D. Heldreichii* Jaub. et Sp. Ill. Or. Vol. V. (pagina nulla notata!)

Hab. in Cariâ (Pinard!) in montibus Pisidiæ prope Egirdir (Heldr. pl. exs. sub *D. Orientali*).

Hanc speciem olim *D. Orientali* formis adnumeravi sed præter spicam brevem densiusculam et flores dimidiò minores ab ea et afflinibus certe differt calycis laciniis lanceolatis et longe mucronato-acuminatis.

7. **D. leucophæa** (Sibth. et Sm. Fl. Græc. VII, p. 6, tab. 607) caule elato in spicam dense cylindricam multifloram abeunti, foliis glabris vel ciliolatis oblongo-lanceolatis lanceolatisque subintegris, rachide glandulosovalnatâ, bracteis linearibus deflexis lanuginosis flores subsessiles æquantibus vel superantibus, corollæ pubescentis parvæ tubo subito inflato ovato-globoso fulvo-reticulato lobis superioribus et intermediis brevissimis, inferiore albo purpurascens-venoso tubo æquilongo basi attenuato ungniculato apice in laminam lunato-rotundam expanso ♀. Ic. Lindl. Mon. tab. 14.

Hab. in monte Athone Macedoniæ (Sibth. Auch!), insulâ Tassos et in monte Scardo (Griseb!).

Specimen imperfectum tantum vidi hujus stirpis pulchrae *D. lanatæ* affinis sed spicâ longiori magis compactâ, bracteis angustis elongatis, floribus dimidiò minoribus et præsertim structurâ lobi corollini inferioris distinctæ.

§ 2. **Grandifloræ** Bth. — Corollæ tubus campanulatus, limbi lobi breves.

8. **D. grandiflora** (All. Ped. I, p. 70) pubescens, caule elato in racemum laxum secundum abeunte, folijs minute denticulatis oblongo-lanceolatis inferioribus breviter petiolatis superioribus semi-amplexicaulibus, rachide glandulosâ, floribus horizontalibus magnis pedicellis calyei æquilongis suffultis, bracteis minutis, calycis tomentelli laciniis elliptico-lanceolatis acutiusculis, corollâ ochroleucâ pubescente campanulatâ sensim ampliatâ ventricosâ lobis latitudine suâ brevioribus inferiore sublatiore ♀. *D. ochroleuca* Jacq. Hort. Vind. tab. 57. — *D. ambigua* Murray. — Rehb. Germ. tab. 69.

Hab. in regione sylvaticâ Olympi Thessali (Heldr! Orph. Fl. exs. 724!), Macedonia occidentali prope Bitolia (Griseb.), Olympos Bithyno (Noë!).

Ar. Geogr. Belgium, Gallia centralis, orientalis et in Pyreneis et alpibus, Helvetia, Germania, Italia, regio Danubialis, Dalmatia, Rossia media, Sibiria Uralensis.

9. **D. ciliata** (Trantv. Mél. Biolog. tom. VI, p. 7) tota crispule pubescens, caule crebre folioso in racemum brevem laxiusculum abeunti, foliis lanceolatis sessilibus acuminatis a medio præsertim argute serratis dentibus ascendentibus inæqualibus, rachide puberulâ, bracteis lanceolatis inferioribus pedicello longioribus superioribus eo æquilongis, pedicellis strictis calyce longioribus vel æquilongis, calycis laciniis ovato-ellipticis glbris obtusis ciliatis, corollæ (in siccâ ochraceâ) breviter glanduloso-pubescentis margine ciliatae brevis tubo campanulato, lobis ovatis obtusis parum inæqualibus inferiore lateralibus sublongiore ♀.

Hab. in ñtione Suaniâ Caucasi meridionalis prope Muschali (Radde!).

Caulis in specimine descripto sesquipedalis, folia caulina intermedia 2½-pollices longa 4 lineas lata, corolla è *D. grandiflora* manifeste brevior 9-10 lineas longa. Species foliis argute incisis *D. laciniata* Lindl. ceterum diversissimam referens

§ 3. **Tubifloræ** Bth. — Corollæ tubus tenuis cylindricus, lobi breves.

10. **D. viridiflora** (Lindl. Mon. 21, tab. 18) glanduloso-canescens, caule folioso in racemum tenuem longum secundum laxiusculum abeunte, foliis tenuiter membranaceis denticulatis oblongis et ovato-lanceolatis inferioribus subpetiolatis ceteris semiamplexicaulibus, superioribus sensim in bracteas lanceolatas floribus longiores transversibus, pedicellis brevissimis, calycis laciniis lanceolatis acuminatis, corollæ viscoso-pubescentis lutescenti-virentis tubo tenui elongato incurvo non ventricoso lobis parvis patentibus obtusis ♀. *D. lutea* Sm. Prodr. I, p. 439? non L.

Hab. in Fagetis et Castanetis Macedoniae, mons Korthiati (Heldr! Orph. Fl.

exs. 725!), in Athone (Sibth., Auch. 1892!, Pichler!), in monte Haemo Thracie (Friv!).

Species foliis tenuiter membranaceis insignis, corolla magn. *D. lutea* sed longior.

CAMPYLANTHUS (Roth Nov. pl. Sp. p. 4).

Calyx quinquepartitus laciniis aequalibus. Corolla infundibuliformis tubo elongato infra medium incurvo, limbo patente quinquepartito subæquali. Stamina bina prope basin inserta inclusa, antherarum loculi divaricati apice in unicum connati. Stylus brevis apice incrassato-dilatatus. Capsula ovato-compressa bilocularis septifrage dehiscens valvis bipartitis columnam placentiferam nudantibus. Semina numerosa orbiculato-compressa hyalino-alata (ex. cl. Webb) campylotropa hilo et chalazâ ad medium marginis approximatis. — Frutices foliis alternis subulatis.

I. C. ramosissimus (Wight Ic. IV, tab. 4416) a basi divaricatum ramosissimus ramis tortuosis, foliis villosis brevibus linearispathulatis, floribus paucis terminalibus subcorymbosis, pedicellis calyce brevioribus basi bracteolatis, calycis laciniis lanceolatis acutis, corollæ tubo calyce triplo longiore limbi laciniis oblongis acutis. *C. salsolooides* Stocks exs. non Roth.

Hab. in collibus siccis calcareis Belutschiae (Aitch. Cat. p. 408. Non vidi).

Ar. Geogr. Ditio Scinde.

TR. VI. VERONICEÆ (Bth. in D.C. Prodr.) Corolle tubus brevissimus, lobii patentes plani non vel obscure bilabiati. Stamina distantia in nostris bina. Antheræ muticæ biloculares vel confertim uniloculares. Fructus capsularis. Inflorescentia centripeta racemosa. Folia saltem inferiora opposita.

WULFENIA (Jacq. Coll. II, p. 66).

Calyx quinquepartitus. Corolla tubuloso-campanulata, limbi brevis erecto-patentis subbilabiati labio superiore latiore bifido inferiore trilobo. Stamina bina filamentis ad basin sinuum labia sejungentium insertis, antheræ reniformes loculis divergentibus apice confluentibus. Stylus capitatus. Capsula oblonga septicide et loculicide dehiscens. Semina ovata dorso convexa facie concava. — Herbae perennes foliis subradicalibus, pedunculo scapiformi.

I. W. Orientalis (Boiss. Diagn. Ser. I, 4, p. 75) glaberrima, foliis rosulatis cartilagineis oblongo-spathulatis in petiolum brevem

attenuatis duplicitim et obtuse crenato-sublobatis, scapo nudo foliis longiore in spicam tandem elongatam laxam abeunte, bracteis oblongo-lanceolatis acutis, pedicellis hirtulis bracteam superantibus calyce subdupo longioribus, calycis lacinis lanceolatis obtusis, corollae tubo calyce duplo longiore angusto basi æquali, capsula oblonga calyci æquilongæ ♂.

Hab. in rupestribus montium Syrie borealis ad Seleuciam (Auch. 4918!), inter Suadię et Antiochiam (Moubré in Gay. herb!).

Folia W. *Carinthiacæ* sed magis coriacea et glaberrima nec ad nervum medium lanutala, pedicelli calyce longiores nec multo breviores; calycis laciniae in W. *Carinthiacæ* apice longe attenuatae acutiusculæ corollino tubo latiore basi subgibbo vix breviores.

PÆDEROTA (L. Gen. 26).

Calyx quinquepartitus. Corolla tubulosa ad medium rarius fere ad basin usque quadripartita subbilabiata labio inferiore tripartito. Stamina bina tubo inserta, antheræ rimis binis longitudinalibus dehiscentes. Stigma capitato-subbilobum. Capsula turgida acuta loculicida. Semina plano-convexa facie internâ affixa. — Herbae perennes multicaules.

I. **P. Pontica** (Ruprecht. MSS.) crispule et minnte pubescenti-glandulosa, canibus e rhizomate suffrutescenti numerosis simplicibus pumilis inferne undis induratis, foliis subsessilibus a basi rotundata vel subcuneata oblongis obtusis vel acutiusculis pauciserratis, floribus axillaribus solitariis oppositis in racemum foliatum dispositis, pedicellis calyce longioribus folio axillari brevioribus, calycis corollæ dimidio brevioris laciinis oblongo-linearibus obtusis quinto minore, corollæ flavâ tubo brevissimo laciinis oblongo-spathulatis subæqualibus tubulose conniventibus, filamentis inclusis basi glabris, capsula pedicello apice recurvo cernuâ ovatâ acutâ lacinis calycinis patulis auctis breviore, seminibus minimis ovatis subcompressis umbilico minuto ♀.

Hab. in fissuris rupium Imeretiae inter Kutais et Achalzich paulo supra ruinas ecclesie 5100' (Rupr!). Fl. Jun.

Facies P. *Ageriae*, caules 5-6-pollicares, folia plantæ fructiferae 12-15 longa, florifera minor, corolla 6 lineas longa, capsula piso minor. A congeneribus corollæ fere ad basin usque quadripartitæ, tubulosa quidem et ad staminum insertionem non pilosa distincta, quæ notæ characterem genericum *Paderote* infirmant.

VERONICA (L. Gen. p. 14).

Calyx 4-5-partitus. Corollæ tubus brevis vel brevissimus, limbus patens quadrifidus laciñia superiore latiore. Stamina bina tubo in-

serta exserta, antheræ riniis binis longitudinalibus dehiseentes. Stigma integrum. Capsula turgida saepius compressa bilocularis obtusa saepius retusa. Semina planiuscula vel concava facie internâ affixa. — Herbae vel suffrutices.

Specierum Orientalium distributio.

SERIES I. — Perennes.

SECT. I. BECCABUNGA Griseb. Spic. II, p. 31. — Racemi axillares oppositi. Corollæ tubus brevissimus. Capsula loculicide dehiseens valvulis demum bipartitis alterâ vel utrâque a columna placentiferâ secedente. Semina ovata vel oblonga vix compressa.

V. Anagallis, anagaloides, Hayssknechtii, lysimachioides, oxycarpa, Beccabunga, Michauxii.

SECT. II. CHAMEDRYS Griseb. Spic. II, p. 28. — Racemi axillares oppositi rarius alterni. Corollæ tubus brevissimus. Capsula lateraliter compressa saepius marginata valvulis columnæ placentiferæ arce cohaerentibus. Semina compressa.

* Semina cymbiformia.

V. peduncularis, petraea, Caucasica, Armena, microcarpa, Aucheri.

** Semina plano-biconvexa facie interiori umbilico saepius margine elevato donato instructa, plus minus distinete rugulosa.

+ Capsula basi cuneata.

V. pectinata, multifida, Orientalis, Kurdica, thymifolia, polifolia, Aleppica, cuncifolia, surculosa, fragilis, einerea, Chamedrys.

++ Capsula basi rotundata.

V. melissifolia, stenobotrys, Galathica, urticifolia, Teucrium, latifolia, Austriaca, telephœfalia, euphrasiæfolia, aphylla.

*** Semina plano-convexa non umbilicata.

V. officinalis, minuta.

SECT. III. VERONICASTRUM Bth. in D.C. Prodr. X, p. 479. — Racemi terminales. Corollæ tubus brevissimus. Capsulae valvulae columnæ placentiferæ arce cohaerentes. Semina non cymbiformia.

V. gentianoides, monticola, bellidioïdes, Kotschyana, Thessalica, serpyllifolia, cespitosa, bombycina.

SECT. IV. PÆDEROTOIDES Blh. d.c. Prodr. X. p. 479. — Racemi axillares. Corollæ tubus elongatus. Facies Pæderotæ sed capsula Veronicaæ.

V. Pæderotæ.

SECT. V. PSEUDOLYSIMACHIJUM Koch Syn. p. 605. — Racemi terminales. Corollæ tubus latitudine suâ sublongior. Semina ovata vel oblonga vix compressa.

V. spuria, longifolia, spicata, orchidea, incana.

SERIES II. — Annuæ.

SECT. V. ALSINEBE Griseb. Spic. II, p. 23. — Flores in racemum terminali dispositi vel axillares solitarii. Capsula lateraliter compressa valvulis columnæ placentiferæ arcte cohærentibus.

* Semina plano-biconvexa facie plus minus manifeste umbilicata.

+ Pedicelli calyce breviores vel eo vix longiores.

1. Folia pinnatipartita.

V. chamæpithys, verna.

2. Folia indivisa integra vel crenata.

V. arvensis, Sartoriana, viscosa.

++ Pedicelli calyce manifeste et pluries longiores.

1. Capsula sinu angusto biloba.

V. acinifolia, nudicaulis, ixodes, conferta.

2. Capsulæ loculi sinu lato recto vel obtuso divergentes.

V. Reuteriana, hispidula, Syriaca, ceratocarpa, divaricata.

** Semina cymbiformia facie concava dorso convexa
(Sect. *Omphalospora* Besser).

+ Calycis laciniæ fere ad basin usque liberæ.

1. Folia floralia saltem superiora bracteiformia.

V. glauca, Chaubardi, Peloponnesiaca, amena, grandiflora, præcox,
Bungei, triphyllus, biloba, campylopoda, rubrifolia, macropoda.

2. Folia floralia omnis caulinis subconformia.

V. Buxbaumii, filiformis, didyma, agrestis, opaca, Cymbalaria, cymbalioides, hederæfolia.

++ Calycis laciniæ per paria connatæ.

V. Cristagalli, Griffithii.

SERIES I — Perennes.

SECT. I. BECCABUNGA.

1. V. Anagallis (L. Sp. p. 16) glabriuscula superne interdum parce glandulosa, rhizomate radicante, caulis erectis crassis fistulosis, foliis sessilibus serratis et integris a basi saepius cordato-amplexicauli ovatis et lanceolatis, racemis laxis axillaribus paniculatis, bracteis linearibus, pedicellis erecto-patulis calyce braeteisque longioribus, calycis laciniis 4 inaequalibus ellipticis, corollâ pallidâ calyce sublongiore, capsula glabrâ orbiculata subretusa calyce subbreviore, stylo capsulae subaequilongo ♀. Ic. Fl. Dan. tab. 903. — Rehb. Germ. tab. 81

Hab. in humidis ad fossas et rivos totius ditionis a regione inferiori ad alpinam a Graeciâ! ad Arabiam petream! Egyptum in Oasi magnâ (Schweinf!), Persiam australem! Affghanianâ!

β. montioides. — Caules (ex loco steriliore sicciore vel ex disseminatione gregariâ) teneri pusilli simplices saepe nani, racemus unicus vel pauci pauciflori, folia breviter ovata. — *V. montioides* Boiss. Diagn. Ser. I, 7, p. 43. — *V. pusilla* Bth. in D.C. Prodr. 10, p. 468 (saltem quoad specimina Afghana!).

Hab. in Caucaso prope Kasbek (Rehm!), humidis alpis Kuh Daëna Persiae australis (Ky. 63! et 679!), in fauce Dudera montis Elbrus Persiae borealis (Ky. 270!), regno Cabulico ad Akrobot (Griff!).

Forma forsitan interdum annua et cuius transitus ad typum manifesti videntur (verbi gratia specimina Kotschiana ex monte Hermon Syriae № 225! sub *V. anagalloide*), varietas parallela in *V. anagalloide* quoque occurrit.

Ar. Geogr. Europa tota, Sibiria, Japonia, China, India, Abyssinia, Africa borealis, Insulæ Canarienses, America borealis.

2. V. anagalloides (Guss. Pl. rar. p. 5. tab. 3) glabra vel puberula, rhizomate radicante, caulis erectis fistulosis, foliis a basi saepius amplexicauli linear-lanceolatis acutis subintegris, racemis laxis axillaribus paniculatis, pedicellis erecto-patulis calyce bracteisque linearibus longioribus, calycis laciniis inaequalibus ellipticis, corollâ albida striata calyce non longiore, capsula glabrâ ellipticâ obtusa vix retusa ♀.

Hab. in humidis et ad rivos Graeciae in Bœotia (Heldr!), Anatoliae in Pamphyliâ et Lyciâ (Bourg!), Syriae littoralis (Bl! Gaill!), Egypti (Ehrenb. exs. sub *V. Egyptiacâ* Ehr!), Arabiae Petreæ (Bové!), Transcaucasiae (C.A. M., Hoh.), Affghaniae (Griff!).

β. tenuis. — Forma macra caulis tenuibus simplicibus qui annui videntur. — *V. tenuis* Ledeb. Fl. Alt. I, p. 38! — Ic. tab. 217.

Hab. in prov. Talysh prope Lenkoran (Hansent!), in Persia boreali.

Valde affinis *V. Anagalli* pro enjus varietate notanici plures eam habent,

eam tamen cum formæ hujus sectionis plures in Oriente crescent seorsum describo.

Ar. Geogr. Hispania, Gallia, Italia, Songaria.

3. V. Haussknechtii, glabra, rhizomate repente radicante, caule erecto fistuloso, foliis sessilibus a basi amplexicanli linear-lanceolatis obtusiusculis integris summis angustissimis acutis, racemis axillaribus tenuibus laxis longe paniculatis, pedicellis tenuibus erecto-patulis bracteâ linear-setacea calyceque longioribus, corollâ albidâ laciniis calycinis oblongis obtusis vix longiore, capsula minimâ caiyce longiore ellipticâ retusâ stylo eâ breviore superatâ ȝ.

Hab. in pratis humidis et uliginosis Mesopotamiae ad Surug et secûs fluvium Hassawi (Haussk!).

Pedalis et elatior, folia inferiora 8-10 lineas longa tres lata. Facies et capsulae ellipticæ *V. anagalloidis* sed dimidio minores et insuper valde retuse. Ulterius investiganda.

4. V. lysimachioides (Boiss. Diagn. Ser. II, 3, p. 165) glabra, caule elato fistuloso, foliis a basi amplexicauli lanceolatis acutis elongatis subintegris, racemis axillaribus longis ample paniculatis densifloris. pedicellis strictis bracteâ setacea calyceque vix longioribus, calycis laciniis lanceolatis acutis, corollâ niveâ calyce longiore, capsula glabrâ ovatâ retusâ calyce sublongiore stylo æquilongo superatâ ȝ.

Hab. ad aquas circa Laodiceam Lydiam (Boiss!).

Planta 2-3-pedalis foliis inferioribus 2-3-pollicaribus. Videtur a *V. Anagalli* et *V. anagalloide* pedicellis non elongatis subpatulis sed abbreviatis strictis distincta.

5. V. oxyacarpa (Boiss. Diagn. Ser. I, 7, p. 44) inferne glabra superne minute glandulosa, rhizomate radicante, caule fistuloso, foliis sessilibus a basi amplexicauli lanceolatis acutiusculis obsolete crenatis, racemis axillaribus densiusculis elongatis paniculatis, pedicellis erectis bracteâ linear calyceque duplo longioribus, calycis laciniis ovatis acutiusculis corollâ rubellâ subbrevioribus, capsula glabrâ calyce longiore oblongâ apice attenuatâ acutiusculâ stylo subbreviore apiculatâ ȝ.

Hab. ad fontes alpinas Persiae australis, mons Kuh Daëna (Ky. 639!). ad Desgird ditionis Bachtiaris (Haussk!), in regno Cabulico ad Arghandah prope Candahar (Griff. exs. 3918!).

Facie *V. lysimachioide* et *anagalloide* refert ab utrâque corollâ rubellâ, a priori capsulis acutiusculis, a posteriore pedicellis strictis differre videtur. An forma *V. Michauxii* capsulis acutis quõque donata? sed differt formâ foliorum et tomenti defectu.

6. V. Beccabunga (L. Sp. 16) glabra rarius superne glandulos-puberula, caulinis basi radicantibus crassis fistulosis, foliis carnosulis petiolatis oblongo-ellipticis obtusis crenato-serratis, race-

mis axillaribus laxis, pedicellis patulis bracteâ calyceque longioribus, calycis laciniis ovato-lanceolatis acutis corollâ cœruleâ subbrevioribus, capsulâ suborbiculatâ turgidâ subretusâ stylo eâ subbreviore terminatâ ♀. Ic. Engl. Bot. tab. 655 — Rehb. Germ. tab. 80. — *V. Scardica* Griseb. Spic. II, p. 31 ex specimine mili nou diversa videtur. — *V. punctata* Hamilt. (saltē quoad specimina Griffithiana!).

Hab. ad aquas totius ditionis a Græciâ!, ad Arabiam petræam! Persiam orientalem (Bgel!) et regnum Cabulicum (Griff!) a regione inferiori ad alpinam crescent.

Occurrunt quoque in hâc specie forme pumilæ et nanæ eis *V. Anagallis* analogæ.

Ar Geogr. Europa omnis, Sibiria, Japonia, regio Himalaica, Abyssinia, Africa borealis.

7. V. Michauxii (Lam III. I, p. 44) papillis crispulis undique pubescens viscidula, caulis ascendentibus basi radicantibus, foliis ovatis crenato-dentatis inferioribus et ramorum steriliū petiolatis cœteris amplexicaulibus, racemis paucis ad axillas superiores densiusculis, pedicellis erectis bracteâ et calyce sublongioribus, corollâ rubellâ calycis laciniis oblongis acutis longiore, capsulâ calyce sublongiore glandulosâ ovatâ turgidâ acutiusculâ non retusâ stylo longo apiculatâ ♀. Ic. Jauth. et Sp. III. tab. 424.

Hab. ad aquas montis Elvend Persiae 7-9000' (Mich! Auch! Haussk!), in regno Cabulico (Griff. ex Bth).

Folia et habitus *V. Beccabungæ* sed indumentum velutinum papillosum et capsulae acuta.

SFCT. II. CHAMÆDRYS.

* Semina cymbiformia.

8. V. peduncularis (M.B. Taur. Cauc. I, p. 11) caulis numerosis ascendentibus vel procumbentibus subbifariam hirtis rigidulis, foliis breviter petiolatis et sessilibus glabris vel adpresso hirtis ovatis vel oblongis grosse dentatis vel incisis dentibus ascendentibus, racemis ex axillis superioribus longis laxis foliolosis, bracteis inferioribus oblongis subdentalis cœteris elliptico-linearibus integris, pedicellis filiformibus patulis calyce 4-5-plo longioribus, calycis laciniis elliptico-lanceolatis subæquilongis, corollâ albâ ad fæcum rubro-striatâ calyce longiore, capsulâ hirtulâ calyce subæquilongâ primum compressâ transverse latiore didymâ basi truncatâ angulo lato retusâ, seminibus cymbiformibus magnis in loculo 2-3-nis ♀. *V. secundiflora* C. Koch Linn. XVII, p. 288! — *V. nigricans* C. Koch L. XXIII, pag. 693! — *V. phanicaantha* C. Koch Linn. XXIII, p. 283! — *V. Beuthumi* C. Koch.

Hab. in dumosis et sylvaticis regionis inferioris et alpinæ, Pontus prope Trapezuntiem (Bourg!), et Rhizé (Bal!), circu Djimil 6000' (Bal!), Armenia

Turcica ad Kassuklu (Huet!) et in monte Bingöldagh 900' (Ky!), Armenia Rossica (Szov!), Imeretia 40°0'-500' (Szov! Rupr!), Caucasus omnis (C.A. Mey!), Georgia Caucasica prope Tiflis (Hohen!), Caucasus orientalis 6000'-800' (Rupr!).

V. umbrosa. — Folia minora angustiora, calycis laciniae capsula longiores. — *V. umbrosa* M.B. Taur. Cae. I, p. 11! — Ic. Fl. Ross. tab. 7.

Hab. in sylvis umbrosis Tauriae (Stev! Godet! Beaupré!).

Caules formae typicae semipedales vel ped.les, pedicelli 4-6 lineas longi, capsula transverse 2 lineas lata. Dubitatum sub *V. anisophylla* cl. Bth. in D.C. Prodr. p. 469 describit plantam Armenam (Auch. № 2531 in D.C. herb.) quæ a *V. anisophylla* C. Koch = *V. Orientalis* Mill. omnino differt et ad *V. pedunculare* referenda videtur, capsula tamen apice non retusa sed junior est et ex specimine pessimo et incompleto non rite dijudicandum.

9. ***V. petraea*** (Steven Mem. Mosq. III, p. 250) minute et crispu-pubescentes cespitosa, caulis pumilis diffusus et ascendentibus foliosis, foliis a basi subsessili cuneatæ oblongis vel ellipticis paucidentatis superioribus interdum integris, racemis axillaribus solitariis pedunculis longiusculis insidentibus brevibus laxiusculis, pedicellis bracteis oblongis calyceque 2-3-plo longioribus, calycis laciinis late oblongis capsulam æquantibus vel superantibus, corollæ cœruleæ ampliæ, capsula glanduloso-hirta bilobâ transverse latiore basi rotundata apice retusâ, seminibus cymbiformibus

Hab. in regione subalpinâ et alpinâ, Caucasus occidentalis (C.A. Mey!), australis ad Wladikawkas, Kobi, Lars, Kasbek 4000'-6000' (Stev! C.A. Mey! Rupr!), Tuschetia, Daghestania a 3000' ad 9500' (Rupr!).

In humilioribus 4-5-pollicaris foliis 6-10 lineas longis, facie *V. Alpinam* referens, in regione alpinâ superiori rami tenuiores abbreviati, folia minima 3-4-lineas longa Species capsula et seminibus affinis *V. pedunculari* a quâ differt foliis minutis, pedunculis brevioribus preter bracteas non foliolosis, pedicellis brevioribus non patulis

10 ***V. Caucasică*** (M.B. Taur. Cae. I, p. 13) undique crispu-pubescentes subglandulosa, caulis erectis vel ascendentibus, foliis subsessilibus ambitu ovatis vel oblongis semel vel bis pinnatisectis laciinis oblongis vel lineari-cuneatis basi saepè angustatis obtusis rarius acutatis, racemis ex axillis superioribus oppositis longe pedunculatis patentibus laxifloris, bracteis inferioribus pinnatisectis superioribus ellipticis integris, pedicellis filiformibus calyce triplo longioribus fructiferis patentí-recurvís, calycis laciinis late lanceolatis capsula longioribus, corollæ calyce longiore albâ lilacino-striatâ, capsula hirtâ valde compressâ didymâ transverse latiore basi truncatâ angulo lato retusâ, seminibus cymbiformibus ♀. Ic. Lodd. Bot. Cab. tab. 1369. — *V. Ossetica* Stev. Mem. Mosq. 2, p. 180.

Hab. in Caucaso occidentali (C.A. Mey!), australi 2400'-5000' (M.B., Rupr!), orientali 6000'-7500' (Rupr!).

Planta 5-6-pollicaris affinis *V. pedunculari* a quâ differt indumento copioiore, foliis *V. Austriaca*, capsula calycinis laciinis superatâ.

11. V. Armena (Boiss. et Huet Diagn. Ser. II, 3, p. 166) siccatate nigricans cespitoso-multicaulis, rhizomate indurato, caulis punnulis decumbentibus et ascendentibus temibus rigidis duris sublente crispule velutiniis, foliis sessilibus brevissimis in lacinias revolutas tenuissime lineares pinnatisectis, racemis ex axillis superioribus 2-4 pedunculis folia superantibus suffultis brevibus laxis, pedicellis tenuibus bracteis elliptico-linearibus calyceque 2-3-plo longioribus fructiferis patentibus, calycis laciniis quinis oblongis quintâ breviore, corollâ intense cæruleâ calyce plus duplo longiore, capsulâ glabratâ parvâ obcordatâ transverse latiore late retusâ calyce triplo longiore, seminibus cymbiformibus ♀.

Hab. in monte Techdag Armenia 7-8000' (Huet! Calvert!).

Canales 3-4 pollicares, foliorum laciniæ 3-5 lineas longæ, capsula fere *V. acinifoliae* apice $1\frac{3}{4}$ lin. lata. A *V. multifidae* formis aliis neglectis seminibus cymbiformibus aliena, a *V. Caucasica* laciniis foliorum tenuissimis revolutis, capsulâ minore calycem longe superante.

12. V. microcarpa (Boiss. Diagn. Ser. I, 4 p. 76) crispule et brevissime velutino-eanescens sicca nigricans multicaulis caulis punnulis diffusis vel ascendentibus parte inferiori induratis et nudis, foliis parvis petiolatis obovato-orbiculatis breviter cuneatis regulariter obtuse et interdum subduplicatim inciso-pectinatis, racemis 2-6 in axillis supremis oppositis longiuscule pedunculatis densifloris, pedicellis strictis bracteæ linearis calyceque subæquilongis, corollâ laciniis calycinis oblongo-linearibus obtusis inæqualibus longiore, capsulâ minimâ tomentellâ calyce breviore vix compressâ orbiculatâ transverse sublatiore emarginatâ, seminibus cymbiformibus.

Hab. in salsuginosis prov. Aderbidjan Persiae (Auch. 5096! Szov!) ad Bayazid Armenia Turcicæ (Auch. 5097!), ad Nackitschewan Armenie Rossicæ (Buhse!).

Planta 4-5-pollicaris, racemi fructiferi 2-3-pollicares, folia 3-5 lineas diam. lata, capsula linea vix longior et latior. Species foliorum formâ *V. cuneifoliae* affinis sed distinctissima indumento, ramis duris non radicantibus, capsulae minitie et formâ, præsertim antem seminibus cymbiformibus qua in *V. cuneifolia* biconvexa umbilicata sunt.

13. V. Aucheri (Boiss. Diagn. Ser. I, p. 80) tota crispule albo-velutina, dense cespitosa basi denum indurata ramis intricatis filiformibus fragillimis foliosis, foliis minimis petiolatis ovato-rotundatis obtuse 5-7 inciso-crenatis, floralibus imis basi cuneatis 2-3-dentatis seminis linearis-spathulatis integris, pedicellis axillaribus unifloris capillaribus flexuosis folio 2-3-plo longioribus, calycis tomentelli laciniis oblongis obtusis, corollâ albo-cæruleâ calyce duplo longiore, capsulâ calycem subæquante hirtâ transverse latiore breviter obcordatâ lobis rotundatis. stylo sinu multo longiore, seminibus minutis oblongis cymbiformibus brevibus ♀.

Hab. in saxosis regionis alpinæ Persiae borealis in monte Demavend 6300' (Auch. 5093! Ky. 342!), in valle Loura montis Elbrus (Ky!), ad Rudbar (Buhse!).

Ex speciminibus Kotschyani certe perennis cespites densos parvos formans, rami intricatissimi flexuosi 2-3-pollicares, folia infima $1\frac{1}{2}$ -2 lineas diam. lata, flores pro planta minutie majusculi. A praecedentibus floribus axillaribus solitariis nec in racemos axillares pedunculatos dispositis diversa.

• Semina plano-biconvexa facie interiori umbilico saepius margine elevato donato instructa, plus minus distinete rugulosa.

+ Capsula basi cuneata.

14. **V. pectinata** (L. Mant. p. 24) velutina canescens, caulis prostratis radicantibus, ramis ascendentibus racemum lateralem saepius unicum elongatum parte inferiori foliosum ramulumque terninalem brevem dense foliosum edentibus, foliis ramulorum obovatis vel elliptico-linearibus basi attenuatis subpetiolatis regulariter et pectinalim crenatis vel incisis, foliolis ad raceini partem inferiorem parvis oblongis crenatis in bracteas lineares sensim abeuntibus, pedicellis strictis calyce vix longioribus, calycis lacinias quaternis linearibus inaequalibus, corollâ magnâ pallide cæruleâ, capsulâ pubescente basi cuneatâ obtriangulari late et breviter retusâ vel truncatâ ♀. Ic. Buxb. Cent. I, tab. 39 fig. I.

Hab. in collibus et umbrosis Thraciae circâ Byzantium (Sibth., Orph!), Bithyniae ad Ghemlek et Moudania (Griseb., Barbe!), ad Broussam (Pichler!), Lydiae in monte Tartali ad Smyrnam et in Sipylo (Boiss!), Phrygiæ monte Bulghasdaghi (Ball!), Cappadociâ nec in Syria (Auch. 1956!) et in monte Ali-dagh (Bal!).

Planta sepinpedalis vel subprocerior, folia 7-9 lin. longa, racemi fructiferi 4-6-pollicares. Species racemis inferne foliolosis nec nudis, in ramo solitariis et fere nunquam oppositis inter affines insignis.

15. **V. multifida** (L. Sp. 17) breviter et erispule pubescentes, caulis basi induratis decumbentibus vel diffusis, foliis sessilibus in lacinias lineares integras vel dentatas pinnatisectis, racemis 2-4 ex axillis subquinis, pedicellis calyce vix longioribus fructiferis strictis, calycis lacinias subquinis valde inaequalibus, corollâ carneâ vel pallide cæruleâ calyce longiore, capsulâ calyci aequilongâ vel sublongiore glabrâ vel breviter glandulosâ transverse latiore basi cuneatâ obsoletissime obcordata vel truncata ♀.

Hab. in dumosis siecis Lydiæ in monte Sipylo (Auch. 1957!), Phrygiæ prope Ouchak (Ball!), Cariae in monte Cadmo (Boiss!), Pisidiae (Heldr!), Cilicie Tracheæ (Per!), Libani prope Bscherre (Ky. 74!), Anatoliæ bor. ad Tokat (Auch. 5098!), Tauriæ (Stev!).

β. tenuifolia. — Caules tenuiores, foliorum laciniae tenuissimæ abbreviate. — *V. tenuifolia* M. B. Gauc. I, p. 43. *V. parvifolia* Vahl Enum. I, p. 72.

Hab. in regione subalpinâ et alpinâ, Cadmi et Sipyli cacumen (Boiss!), Anatolia borealis (Wied!), Argens Cappadociae (Ky! Ball!), Armeniâ in montibus Gumusch (Bourg!), in valle Kassuklu (Huett!), supra Karput (Noë!), Georgia Caucasicâ (Hoht!), Tauriæ prope Laspi (Rehm!).

Species capsulâ *V. Orientali* similis et ab eâ foliis pinnatisectis (forsitan non

satis) distincta, transitus manifestos tamen nondum vidi. Varietas *tenifolia* minor, alpina, ad typum manifeste transit.

Ar. Geogr. Rossia australis, Songaria, Siberia Uralensis.

16. V. Orientalis (Mill. Soland in Ait. Kew I, p. 23) breviter et erispule pubescens rarius glabrata, caulis e rhizomate ligneo-menti decumbentibus vel prostratis, foliis sessilibus brevibus inferioribus cuneato-oblongis vel lanceolatis dentato-incisis rarius integris, superioribus saepius angustioribus lanceolatis et supremis saepe integris, racemis 2-4 ex axillis supremis brevibus laxis fructiferis secundis, pedicellis erecto-patulis calyce vix longioribus, calyeis laciniis 4 vel 5 linear-lanceolatis valde inaequalibus, corolla carneâ vel pallide cœruleâ calyce longiore, capsula glanduloso-pubescenti transverse latiori basi cuneatâ obtusissime obcordata vel truncatâ calyce sublongiore vel aequilongâ, seminibus ovatis ♀. *V. parviflora* (sphalmitate *parvifolia* Vahl! Enum. I, p. 72 — *V. Noëna* Boiss. Diagn. Ser. II, 3, p. 168 : forma foliis subintegratis) — *V. anisophylla* C. Koch Linn. XVII, p. 287 ! non Bth. in p.c. Prodr. — *V. Billardieri* Vahl Enum. I, p. 70 ex specimine archetypo herb. Desf !! forma foliis subintegratis.

Hab. in rupestribus siccis presertim montanis et alpinis Anatolie Orientalis in TAURO CILICICO 8-9000' (Ky! Bai exs. sub *V. pectinata* var.), ARMENIA Turcica (C. Kocht), in Libano et Antilibano frequentissima (Boiss! Gaill. 2170!, Bl. exs. 761 et 448!), circa Aintab (Auch! Haussk!), in montibus Mesopotamiae et Assyriae (Noël Ky. 62 !), montibus Avroman, Schahm, Pir Omar Gudrun Kurdistaniae Persiae 7000-8000' (Haussk!), Persia boreali (Bge!) et australi in monte Kuh Delu (Ky. 486!), ditione Bachtiaris Haussk!), montibus circa Ispahan (Auch!), Tauria (ex Stev!), prov. Talysch (C.A. Mey!), Armenia Rossica (Szovt).

β. tenuifolia (Boiss. Diagn. Ser. II, 3, p. 167). — Folia anguste linearia margine interdum subrevoluta integra rarius acute paucidentata. — *V. Taurica* Stev., Lond. Bot. Cab., tab. 911.

Hab. in Armenia australi ad Bakker Maaden (Noël), circa Baibut (Bourg!), ad Diarbekir Mesopotamiae (Ky. 187!), Persia boreali (Buhsel!), in monte Tchetyrdagh Taurie (Rehm!).

Species foliis formâ et magnitudine variis irregulariter dentatis vel integris valde varians, a *V. Tenuiro. latifoliâ*, *Austriacâ* capsula basi cuneatâ transverse latiore facile distinguenda. *V. parviflora* Vahl ex antopsis *Veronicae Cappadocice minime filiis laciniatis* Tourn. herb! huc adducenda est.

17. V. Kurdica (Bth. p.c. Prodr. X, p. 473) minule et erispule pubescenti-velutina cespitosa, ramis diffusis brevibus tenuibus intumatis, foliis parvis oblongis vel linear-lanceolatis margine saepius revolutis integris rarius paucidentatis, racemis pedunculatis brevissimis laxis, pedicellis bracteolas minutis ellipticas calycesque 2-3-plo superantibus fructiferis erecto-patulis, calycis laciniis linear-lanceolatis quaternis inaequalibus, corolla cœruleâ calyce quadruplo longiore, capsula parvâ calyce longiore glabra vel minute hirtâ obcordata bilobâ basi cuneatâ transverse latiore, seminibus ovatis ♀. *V. Syriensis* C. Koch Linn. XXIII, p. 698 !

Hab. in montibus Armeniæ Kurdicæ (Brant!), circa Artahan (C. Kochi!), in devexis australibus Bingöldagh 6000' (Ky. Suppl. 717 sub *V. Orientali*), Persiæ boreali in monte Elburz (Ky. 221!), monte Elamut (Auch. 5089!), montibus Ssahend, Ssawalan, Demavend (Buhsei!), Persiæ australis cacumine montis Kuh Daëna (Ky. 774! forma foliis minoribus brevioribus).

Caules 2-3-pollicares, folia 2-3 lineas longa, racemi fructiferi 2-3-pollicares. Valde affinis *V. Orientali* var. *tenuifolia* a quâ pedicellis longioribus (2-3 lineas longis) patentibus fere tantum differt. An species legitima? Speciminum e Persiæ semina matura non vidi, huc tamen ommino spectare videntur.

18. ***V. thymifolia*** (Sibth. et Sm. Fl. Græc. I. p. 5, tab. 6) breviter et crispile velutina cespitosa, candiculis prostratis radicantibus ramis brevibus ascendentibus crebre foliosis, foliis oblongo vel linearispathulatis basi attenuatâ sessilibus margine revolutis integris vel superne minute paucicrenatis, racemis brevibus paucifloris breviter pedunculatis in axillis supremis 1-2 interdum gemmæ terminalis abortu terminalibus, pedicellis calyce et bracteâ ovato-oblonga brevioribus, calycis laciniis 4 oblongis subinæqualibus, corollâ cœruleâ calyce longiore, capsulâ hirtâ transverse latiore obcordatâ basi cuneatâ lobis divaricatis. seminibus orbiculato-ellipticis ♀. *V. Cretira* Pall. in Link Jahrb. I, p. 41 et Willd. herb! *V. Tymphrestea* Boiss. et Sprun. Diagn. Ser. I, 4, p. 77. — *V. teucrioides* Boiss. et Heldr. Diagn. Ser. II, 3, p. 169.

Hab. in regione alpinâ Cretæ in montibus Sphacioticis (Sibth.), in montibus Idâ et Lassiti (Heldr!), in montibus Velugo Ætoliae (Tymphresto), in cacumine Parnassi (Sprunt), in Olympo Thessalo ad nives (Heldr!).

Rami bipollicares, folia 3-5 lineas longa 1-2½ lata. Capsulae formâ (quani non vidi sed e descript. et icon. Fl. Græca descripti) affinis videtur *V. Orientali* et *Kurdicae* sed spicâ densâ subcapitâtâ, pedicellis brevissimis, distincta. Racemi terminales interdum sed rarins sunt, abortu gemmæ terminalis.

19. ***V. polifolia*** (Bth. in D. C. Prodr. X, p. 473) tomentoso-cana, caulinibus e rhizomate indurato cespitosis brevibus ascendentibus, foliis sessilibus oblongo-vel linearispathulatis basi attenuatis margine revolutis obtuse et regulariter crenatis, racemis ex axillis supremis oppositis brevibus densis, pedicellis rectis calyce subbrevioribus, calycis laciniis quaternis obovato-oblongis, corollâ roseâ calyce longiore, capsulis minutis tomentellis basi cuneatis transverse latioribus late obcordatis calyce longioribus ♀. *V. macrostachya* Vahl Enum. p. 71 ex spec. archetypo herb. Desf! sed nomen improprium nam spica non longissima!

Hab. in regione subalpinâ et alpinâ Libani supra Eden (Labill! Boiss! Bl. 447! et 3625!), in Antilibano circâ Rascheya (Boiss!) et in alpe Bludan supra Zebdani 6000' (Ky. 41!), in monte Tauro? (Auch. 1964!). In Olympo Bithyno non crescit ubi ex schedulâ Aucherianâ commutatione indicata fuit, et planta Armena in D.C. citata ad *V. cinereum* referenda est.

Caules 3-4-pollicares, folia parva ea varietatum minorum *Teucrii Polii* referentia in calidioribus cana et anguste revoluta. Species capsulâ affinis *V. Orientali* a cuius formis foliis regulariter et obtuse pectinato-crenatis et laciniis calycis obovatis differt. *V. Aleppica* est planta multo major laciniis caly-

cinis lanceolatis capsulam superantibus. Folia margine convoluta cl. Vahl observatione incautâ integra dixit.

20. V. Aleppica (Boiss. Diagn. Ser. II, 2, p. 169) hirsuta vel tomentoso-caueens, caulis basi induratis decumbentibus vel diffusis elongatis, foliis subsessilibus basi cuneato-attennatis oblongis vel oblongo-linearibus regulariter et pectinatim inciso-dentatis, racemis ex axillis supremis 2 rarius 4 fructiferis longis unilateralibus, pedicellis strictis calyce vix longioribus, calycis laciniis quaternis rarius quinque lanceolatis inaequalibus capsula longioribus, corollâ carneâ amplâ, capsula hirsutâ obtriangulari latitudini æquilongâ apice truncatâ vix retusâ ♂.

Hab. in sylvaticis circa Aleppo Syriæ (cl. Pat. Johannes!), in monte Cassio (Boiss!). Amano (Ky. 86! et 206!), in ditione Kassan Oglu Cilicie orientalis (Ky. 69!), in Tauro Cilico prope Gülek Boghas (Bal. exs. sub *V. Orientali* var!), in Mesopotamiâ prope Mardin et Assyriæ monte Pir Omar Gudrun (Haussk!).

Caules cum racemis semipedales vel demum pedales, folia 7-12 lineas longa 3-5 lata. Affinis *V. multifida* et *Orientali* formâ capsula laevis cuneata, sed hæc obtriangularis et truncata nec transverse latior, calycisque laciniis manifeste superata, folia insuper eis *V. Orientalis* majora et regulariter pectinata, nunquam ut in *V. multifida* pinnatisecta. Specimina e Mesopotamiâ inducendo magis cano ad *V. polifoliam* accedunt sed ob tolla latiora et laciniis calycinis lanceolatas capsula longiores huc spectant.

21. V. cuneifolia (Don Ann. Nat. Hist. 7, p. 457) breviter pubescens cespitosa, caudiculis tentibus prostratis repentibus radicantibus, ramis brevibus ascendentibus, foliis parvis cuneato-ovobovatis utrinque regulariter et obtuse 3-5-crenatis, inferioribus subpetiolatis, racemis axillaribus saepius solitaris rarius oppositis binis tandem longisculis, bracteis inferioribus ovato-oblongis paucidentatis superioribus ellipticis integris, pedicellis calyci æquilongis vel vix longioribus rectis, calycis laciniis quaternis oblongo-lanceolatis subinæqualibus, corollâ cœruleâ margine albida calyce longiore, capsula hirtâ cuneato-obtriangulari retusa calyci subæquilongâ ♀. *V. Cariensis* Boiss. Diagn. Ser. I, IV, p. 75.

Hab. in Cariæ monte Cadmo supra Gheyra et in monte Tmolo Lydiæ ad Bozdagh (Boiss!), in montibus Lyciæ (Heldr! Bourg! sub *V. polifolia*) et monte Gheidagh Isauriæ (Heldr!).

β. villosa. — Magis villosa, folia interdum oblonga. — *V. dichrus* Schott et Ky. Oest. Woch. VII, № 26!

Folia 2-4 lineas longa ea *Teucrii chamædrys* referentia sed minora, capsula sesquilineam longa et lata.

22. V. surculosa (Boiss. et Bal. Diagn. Ser. II, 3, p. 170) glanduloso-hirta, caudiculis nudis tenuibus elongatis prostratis vel subterraneis, ramis brevissimis fragilibus confertim foliosis racemos saepius binos brevissimos etiam fructiferos oblongos densos ex axilla superiore edentibus, foliis sessilibus vel breviter petiolatis basi longe

enneatis oblongo-spathulatis utrinque obtuse 2-3-crenato-dentatis, pedicellis strictis bracteâ oblongo-lineari integrâ calyceque brevioribus, corollâ azureâ laciniis calycinis oblongis obtusis duplo longiore, capsulâ hirtâ basi enneatâ transverse latiore obcordata φ .

Hab. inter lapides in regione alpinâ Tauri Cilicici supra Bulgharmaaden (Bal!).

Rami floriferi pollicares, folia 6-8 lineas longa 3-3 $\frac{1}{2}$ lata, capsula eâ *V. cuneifoliae* major 2 lineas lata. Huic speciei affinis sed characteribus indicatis distincta.

23. *V. fragilis* (Boiss. et Haussk.) velutino-cinerea glandulosa e rhizomate indurato ramosissima fragilis, ramis erectis tenuibus ramulis filiformibus strictis, foliis parvis remotis sessilibus ovato-oblongis utriusque obtuse 5-7-crenato-incisis, racemis brevibus paucifloris ex axillis superioribus binis interdum terminalibus solitariis, pedicellis erectis calyce bracteisque ellipticis brevioribus rectis, corollâ albâ, capsulâ calycis laciniis oblongo-ellipticis breviore glandulosohirtâ transverse latiore angulo aperto obcordato-bilobâ φ .

Hab. in dumosis montium Sawers et Eschker Persiae austro-occidentalis 8000' (Haussk!).

Affinis *V. cuneifoliae* sed ramuli tenuissimi erecti fragiles, racemi breves, capsula forme diversæ vix lineam longa lin. 4 $\frac{1}{4}$ lata subdivaricatum biloba. Racemi normaliter axillares oppositi sed in ramulis maerioribus solitarii omnino terminales ut in speciebus nonnullis aliis transitum ad *Veronicastrum* præbentes.

24. *V. cinerea* (Boiss. et Bal. Diagn. Ser. II, 6. p 131) brevissime velutina canescens vel cinerea sicca nigricans cespitosa, caulinis pumilis inferne prostratis radicantibus deinde ascendentibus, foliis oblongis et oblongo-linearibus basi attenuatis subpetiolatis superne crenato-dentatis margine convolutis, racemis axillaribus breviter pedunculatis laxiusculis, pedicellis calyce bracteâ oblongo-lineari 1 $\frac{1}{2}$ -plo longioribus fructiferis strictis, corollâ carneâ calyce longiore, capsulâ calycinis laciniis æquilongis oblongo-spathulatis sublongiore velutinâ obcordato-triangulari transverse latiore, seminibus subrugosis φ .

Hab. in regione alpinâ Tauri Cilicici (Bal. sub *V. Kurdicâ!*, Ky. exs. 46! sub *V. pectinatâ*), ditionis Kassan Oglu (Ky. 149!), montis Argæi Cappadociae (Ky. 194! et Bal. sub *V. Orientali!*), montis Karamasdagh (Bal!), Berytdagh Cataoniae (Haussk!), Armeniae (Aucht!).

Planta 3-4-pollicaris affinis *V. Kurdicæ* et *V. Orientali* a quibus differt indumento cinereo, pedicellis strictis, calycis laciniis subæqualibus et caulinis inferne valde radicantibus.

25. *V. Chamædrys* (L. Sp. 17) caulinis ascendentibus basi prostratâ radicantibus superne bifariam pilosis, foliis pubescentibus subsessilibus ovatis inciso-crenatis basi rotundatis vel subcordatis, racemis ex axillis supremis 2-4 laxis, pedicellis calyce bracteisque sublongioribus fructiferis erectis, calycis laciniis 4 lanceolatis capsu-

lam superantibus, corollâ calyce longiore cæruleâ lobi inferiori albo, capsulâ compressâ hirtâ transverse sublatiore basi cuneatâ obcordato-triangulari ♀. Ic. Engl. bot. tab. 623 — Rehb. Germ. tab. 83, fig. 2.

Hab. in dumosis præsertim subalpinis, Olympus Thessalus (Heldr!), Macedonia et Thracia (Griseb.), Byzantii (Noe!), Olympo Bithynio et Bithynia orientali (Boiss! Noe!), Ponto (Tchih!), Armeniâ Rossicâ (Calv!), Imeretia (Szyv!), Caucaso centrali (M.B., Bupe!), Tauria (Stev!), Syriâ orientali (Auch. 4942!).

V. pilosa Bth. in n.c. Prodr. — Caules undique pilosi. — *V. chamaedryoides* F. Pelop., p. 15, t. I.

Hab. in Messeniâ (Bory), in Taygeti parte inferiore (Heldr!), montibus Achaei (Heldr!, Orph. Fl. Gr. exs. 717!), in regione abietinâ Parnassi (Sam. et Guicci! Heldr. Herb. Norm. 287!).

Ar. Geogr. Europa borealis et media a Scandinaviâ ad Rossiam, Sibiria, insulae Canarienses.

++ Capsula basi rotundata.

26. **V. melissæfolia** (Desf. ex Poir. Diet. 8, p. 256) caule ascendentî valde elato villis patulis hirsuto, foliis parœ villosulis vel glabratâs ovatis subinciso-crenatis basi rotundatis vel subcordatis inferioribus sessilibus obtusis superioribus acutiusculis interdum breviissime petiolatis, racemis axillaribus breviter pedunculatis longissimis laxiusculis villosulis paniculatis, pedicellis bracteâ linearî calyceque brevioribus rectis, corollâ pallide cæruleâ calycem parum excedente, capsulâ calycis laciñiis quaternis late lanceolatis æqualibus breviore compressâ ciliatâ transverse sublatiore sinn lato obcordata ♀. *V. Vestiana* R. et Sch. I, Mant. p. 11. — *V. maxima* Stev. Mem. Mosq. 2, p. 179, tab. II, fig. 8. — Buxb. Cent. I, tab. 34.

Hab. in umbrosis Phrygiae (Auch. 859!), Bithyniae (Buxb.), circa Toptcha (Wied!), Ponto inter Samsun et Tekke (Tchih!), Ponto Lazico in sylva Calia peope Trapezuntum (Bourg!), circa Rhizé (Ball!), Caucaso centrali (Owerin!), Iberiâ et Kachetiâ (Stev! Rupr!), Guriâ (C. Koch!).

Planta sape 2-3-pedalis, racemi eis *V. Chamaedrys* multo longiores et saepius numerosi, folia superiora vix diminuta subpetiolata, pedicelli multo breviores, capsula major basi non cuneata.

27. **V. stenobotrys** (Boiss. et Bl. Diagn. Ser. II, 3, p. 166) siccatate nigricans, caulis ascendentibus elati flexuosis erispule hirtis, foliis subtus ad nervos breviter et parœ superne sparsissime hirtis vel glabris oblongis acutiusculis in dentes ascendentes aculos regulariter inciso-serratis insimis subpetiolatis intermediis sessilibus subcordatis, racemis 2-4 ex axillis supremis longissimis angustis pubescensibus, pedicellis bracteâ linearî calycique subæquilongis fructiferis strictis, calycis hirti laciñiis anguste linearibns trinerviis obtusis quaternis quintâ minimâ rarius additâ, corollâ e sicco carneâ ampliâ extus hirtâ vel glabrâ, capsulâ nervulosâ hirtâ ovatâ breviter

obcordatâ compressâ durâ margine subincurvassatâ calyce subbrevisiore ♀.

Hab. in Ciliciâ prope Bouloukli (Bal. exs! sub *V. leiocarpâ*), in regione montanâ montis Amani Syriae prope Apisch Gaya 2500' (Ky. 265!), in sylvaticis Libani ad Ain Berdê et prope Bezulmar (Bl!), ad Beit Kaschbo et Khan el Birket (Gaill!).

β. leiocarpa. — Racemi, calyces, capsulaque giaberrimi. — *V. leiocarpa* Boiss. Diagn. Ser. I, 12, p. 45.

Hab. in sylvaticis regionis inferioris montis Cassii suprâ Kassab (Boiss!).

Planta pedalis et elatior, folia pollicem longâ firma, racemi fructiferi semi-pedales, capsula 1 $\frac{3}{4}$ lin. longa pauloque angustior. Affinis *V. melissæfoliæ* a quâ differt foliis minoribus angustioribus, calycis laciniis angustis inæqualibus et capsulâ ovatâ durâ elevatim nervulosâ.

28. V. Galathica, caulis a basi suffrutescenti denudatâ procumbenti ascendentibus crebre ramulosis firmis crispule pubescenti-ramis crebre foliosis, foliis firmis sessilibus oblongis vel oblongo-lanceolatis basi cuneatis obtusiusculis margine subrevolutis ramulorum inferiorum dentatis cæteris integris omnibus utrinque pubēcispulâ cinerascentibus facieque superiori eximie et elevatim pellucido-punctatis, racemis ex axillis superioribus longis multifloris fructiferis strictis, floribus , pedicellis strictis capsulae subæquilonigis bracteâ linearî eis breviore suffultis, sepalis lanceolatis obtusis capsulâ tertiatâ parte brevioribus quinto minore, capsulâ pubescente ovato-oblongâ basi rotundatâ apice retusâ, seminibus plano-biconvexis obsolete rugulosis ♀.

Hab. ad rupes prope Galathi Abchasie Caucasica non procul a Kutais (Rupr.).

Caules 6-9-pollicares, spica fructifera sœpe semipedalis, capsula 2 $\frac{1}{2}$ lineas longa, folia majora 8-9 lineas longa 2 $\frac{1}{2}$ lata. Species indumento et caulinibus basi induratis fere suffruticosis insignis.

29. V. urticifolia (L. fil. Suppl. 83) parce et crispule papilio-hirta, caulinibus erectis proceris, foliis sessilibus ovatis argute serratis superioribus acuminatis basi subcordatis, racemis axillaribus oppositis paniculatis laxis, bracteis linearibus minutis, pedicellis calyce 3-4-plo longioribus erecto-patulis fructiferis apice sursum recurvis, calycis laciniis quaternis minutis lanceolatis obtusiusculis, corollâ griseo-caeruleâ calyce longiore, capsulâ calycè duplo longiore orbiculari breviter retusâ, seminibus lœvibus ♀. Ic. Jacq. Austr. tab. 59. — Rehb. Germ. tab. 82. — *V. latifolia* Lam.

Hab. in sylvaticis subalpinis, Olympus Thessalus (Heldr!), Macedonia prope Bitolia (Orph!), Thracia (Friv!).

Ar. Geogr. Gallia, Helvetia, Alpes Germaniae, Italia borealis, Dalmatia, regio Danubialis.

30. V. Teucrium (L. Sp. p. 16) pubescens, rhizomate caules numerosos ascendentibus pumilos ramosque steriles edente; foliis

oblongo-lanceolatis vel lanceolato-linearibus obtuse crenato-incisis interdum remote dentatis intimis subpetiolatis cæteris sessilibus, racemis 2-4 ex axillis supremis tandem elongatis, pedicellis erectis calyei æquilonis vel longioribus, calycis laciniis 5 oblongo-linearibus obtusis superiore breviore, corollæ magnâ cæruleâ, capsulâ calyei subaequilongâ hirtâ obovato-obcordatâ basi rotundatâ ♀. *V. latifolia* ♂ minor Koch Syn. — *V. prostrata* Smith Prodr I, p. 7 et M.B. Taur. Gaue, non L. — *V. dentata* Schm.

Hab. in siecis præsertim alpinis, Peloponnesus in Messeniâ (herb. Fauché!), regione alpinâ Parnassi (Guicci!) et montis Velugo (Etolie (Samar!), Macedonia (ex Sibth.), Byzantii (Sibth! Noë!), Armeniâ Turcicâ circa Erzerum, Baibut, Tortum (C. Koch! Calvt Bourg!), Caucaso boreali (Ower), Tauriâ (Stev!).

Ar. Geogr. Belgium, Gallia, Germania, regio Danubialis, Lusitania, Hispania borealis, Italia, Rossia, Songaria.

31. ***V. latifolia*** (L. Sp. 18) pubescens, caulis erectis elatis, foliis sessilibus ovatis subcordato-triangularibus crenato-incisis, racemis 2-4 ex axillis supremis ortis tandem elongatis, calycis laciniis 5 lanceolatis superiore breviore, corollæ magnâ cæruleâ, capsulâ puberulâ calyce vix longiore obovato-obcordatâ basi rotundatâ ♀. *V. Teucrium* var. *major* Schrad. — Rehb. Germ. tab. 88. — *V. pseudochamaedrys* Jacq. Austr. tab. 60.

Hab. in regione montanâ Macedoniæ et Thraciæ (Griseb.), Peloponneso (herb! Fauché!), Tauriâ!, Caucaso (M.B.)

Ar. Geogr. Gallia, Germania, regio Danubialis, Rossia.

32. ***V. Austriaca*** (L. Sp. p. 17. — Bth. in n.c. Prodr. 10, p. 470) pubescens, caulis erectis rarius ascendentibus, foliis sessilibus ambitu ovatis in laciniis oblongas basi angustatas vel lineares integras vel incisas pinnatisectis, racemis 2-4 ex axillis supremis tandem elongatis, pedicellis strictis calyce sèpius longioribus, calycis laciniis quinque rarius quaternis linearibus valde inæqualibus, corollæ magnâ cæruleâ, capsulâ calyci æquilonis vel breviore hirtâ obovato-obcordatâ basi rotundatâ ♀. Ic. Rehb. Germ. tab. 89. fig. 3. — *V. multifida* Jacq. Austr. t. 327 non L. — *V. Jacquinii* R. et Sch. I, p. 106.

Hab. in monte Velugo (Etolie 3000' (Sprun!), Macedonia in pratis montanis Scardi et sylvis Athonis (Griseb.), monte Corinthi (Orph!), Thraciâ in monte Hæmo (Frivt!), Iberiâ Caucasicâ (Stev!), Georgia in montosis prope Tiflis (Hon. exs. sub *V. multifidâ!*), Armeniâ Rossicâ (Szov!).

β. canescens. — *V. recta* Bth. in n.c. Prodr. 10, p. 474.

Hab. in monte Tmolo Lydiae supra Philadelphiam (Boiss!), supra Sardes (Balt!).

Species capsulae formâ similis *V. latifolia* a quâ foliis pinnatisectis fero tantum differt. Varietas β indumento tantum a typo distinguitur, laciniæ calycine etenim in eâ quoque inæquales et quinta minima interdum adest.

Ar. Geogr. Austria, Dalmatia, regio Danubialis, Rossia media et australis.

33. V. telephiifolia (Vahl Enum. I, p. 63) glabra glaucescens, ramis diffusis radicantibus fragilibus, foliis carnosis minimis subimbricatis obovato-spathulatis vel oblongis breviter petiolatis obtuse panicrenatis, pedunculis axillaribus solitariis folia superantibus nudis in racemum brevem ovatum abeuntibus, pedicellis calyce 2-3-plo longioribus erectis, bracteis laciniosque calycinis elliptico-oblongis, corollâ cæruleâ calyce longiore, capsulâ glabrâ calyce subduplo longiore bitubâ transverse latiore basi rotundatâ late retusâ. seminibus ovato-orbiculatis margine radiatim subrugosis ♀.

Hab. in Armeniâ (Anch. 1966!), in monte Ararat (Helm!).

$\beta.$ *pilosula*. — Rami sub lente et folia secus nervos birtula, folia magis dissita paulo majora crenis utrinque 4-5 nec 2-3. — *V. orbicularis* Fisch. ex Trautv. Opusc. 1866, p. 8. — *V. Calcerti* Boiss. MSS. — *V. Livanensis* C. Koch Fl. XXIII, p. 698!

Hab. circa Erzernm (Calv!), in Transcaucasiâ prope Borschom (Radde!) ditione Liwaneh Transcaucasiae (C. Koch!).

Rami 2-3-pollicares, filiformes sed duriusculi sicci fragillimi, folia 2-3 lineas longa, pedunculus racemi 8-12 lineas longus, capsula sesquilineam longa 2 $\frac{1}{2}$ lata.

34 V. euphrasiæfolia (Link Jahrb. I, p. 41) glabriuscula cespitosa candiculis tenuiter filiformibus intricatis, ramis filiformibus ascendentibus brevibus laxe foliatis, foliis minimis breviter petiolatis a basi cuneato-rotundatâ oblongis obtusis denticulis utrinque 3-4 ascendentibus, pedunculis ex axillis supremis panicis longis 2-3-floris, pedicellis capillaribus calyce 3-4-plo longioribus, calycis glandulosi segmentis ovalis, capsulâ glanduloso-birtulâ obovatâ obtusissimâ basi cuneatâ calyce tertiatâ parte longiore, seminibus.... ♀. *V. thymifolia* Willd. herb! non Sibth.

Hab. in alpibus prov. Ghilan Persiæ (herb. Willd!).

Rami pusilli 2-3-pollicares, folia 2-3 lineas tantum longa. Valde affinis *V. telephiifolia* var. β a quâ ex unicâ capsulâ a me observatâ differt eâ obovatâ obtusâ nec valde retusâ transverse latiore.

35. V. aphylla (L. Sp. 14) pubescens subglandulosa cespitosa candiculis tenuibus radicantibus, caulinis brevissimis rosulatim foliosis, foliis breviter petiolatis obovato-ellipticis obtusis integris vel crenatis, racemis ex axillâ superiore longe pedunculatis corymbose 3-5-floris, pedicellis calyce et bracteâ 3-4-plo longioribus, calycis lacinios oblongis obtusis corollâ cæruleâ triplo brevioribus, capsulâ magnâ glandulosâ calyce duplo longiore compressâ ovato-obcordatâ, seminibus numerosis orbiculatis ♀. Ic. Rehb. Germ. tab. 86, flg. 2. — *V. depauperata* W. K. Hung. p. 272, tab. 245.

Hab. in fissuris rupium regionis alpinæ montis Parnassi (Heldr!).

Ar. Géogr. Hispania in montibus Castellæ veteris!, Pyrenæi, Jurassus, Alpes Galliae, Helvetia et Germaniae. Aprutium Neapolitanum, montes regionis Danubialis, Serbiae, Bosniae.

... Semina plano-convexa non umbilicata.

36. **V. officinalis** (L. Lp. p. 14) pubescens, caulinis basi repentibus et radicantibus apice ascendentibus, foliis basi attenuatâ breviter petiolatis obovatis et ellipticis serratis, racemis ex axillis superioribus pedunculis crassis insidentibus densis, pedicellis rectis calyce brevioribus, calycis lacinis lanceolatis capsulâ dimidio brevioribus, corollâ pallidâ calyce sublongiore, capsulâ glandulosâ compressâ obtriangulari truncatâ vel retusâ ♀. Ic. Fl. Dan. tab. 243. — Rehb. Germ. tab. 83.

Hab. in sylvaticis praesertim montanis et subalpinis, Macedonia et Thracia 4500'-4000' (Griseb.), Byzantii ad Buyukdere (Goumany!), Tauria et Caucasus (M.B., Radde!, Owerin!), Transcaucasia (Szov!), ditio Talyssch (C.A. Mey.).

Ar. Geogr. Europa borealis et media a Scandinavia ad Rossiam et Hispaniam boreali ad Dalmatiam, Sibiria Uralensis.

37. **V. minuta** (C. C. Mey. Enum. p. 105) pilosula caudiculis tenuiter filiformibus prostratis radicantibus, ramis brevibus ascendentibus, foliis teneris minimis breviter petiolatis ovatis vel orbiculari-latis integerrimis vel obsoletissime paucierenatis, pedunculis in axillis superioribus subsolitariis foliis longioribus foliolosis in racenium brevem pauciflorum abeuntibus, racemi foliis bracteisque ovato-oblongis ellipticisque, pedicellis erectis capillaribus bracteâ calyceque 3-4 plo longioribus, calycis laciinis oblongo-ellipticis pilosis corollâ cernueâ duplo brevioribus, capsulâ calyce sublongiore hirtâ bilobâ transverse latiore basi apiceque retusâ ♀.

Hab. in lapidosis regionis alpinae summae Caucaasi 8000'-11000', Caucasus occidentalis (C.A. Mey!), orientalis in ditionibus Tindal, Boghos, Tuschetia, Chewsuria, Daghestanië (Rupr!).

Facies *V. repens* D.C. Rami 1-2-pollicares, folia 2-3 lineas longa et lata, capsula sesquilineam lata. Afinis *V. telephifolia* sed multo tenerior, non fragilis nec carnosa, folia integra, capsula hirta minor, semina praesertim dimidiò minora oblonga nec orbiculata nec radiatum subuplicata.

SECT. III. VERONICASTRUM

38. **V. gentianoides** (Vahl Symb. I. p. 1) caulinis e rhizomate oblique præmorso subtus fibrillifero erectis elatis in racenium spiciformem longum multiflorum laxum pubescenti-glandulosum abeuntibus, foliis crassiusculis nitidis glabris vel glanduloso-pilosulis integris vel paucierenatis inferioribus rosularum caulinisque intimis approximatis oblongo-lanceolatis vel lanceolatis aenatis basi longe attenuatis, superioribus sensim valde diminutis linear-lanceolatis strictis, pedicellis erecto-patulis bracteâ linearis calyceque 3-4-plo longioribus, calycis glandulosi laciinis oblongis obtusis, corollâ cern-

leâ calyce duplo longiore, capsulâ calyce duplo longiore glandulosâ subcompressâ orbiculari-obcordataâ, seminibus concaviusculis umbilicatis ♀. Ic. Fl. Græc. tab. 5. — Vent. Malm. tab. 86. — *V. Olympica* Pall. — *V. Buxbaumiana* Pall. — *V. pallida* Hörn Hafn. p. 17.

Hab. in pratis alpinis humidis, Olympus Bithynus (Sibth., Auch. 1954! Boiss! etc.), mons Ilgasdag Anatoliæ borealis (Wied!), Argæus Cappadociaæ 9000' (Bal. 1028!), Pontus occid. (Tchih!) et Lazicus supra Djimil 7800' (Balt.), Armenia Turcica (Huet! Bourg!), Tauriæ alpes (Stev.), Caucasus occidentalis et centralis 2500'-9000' (M.B., Rupr!, Rehm!), Caucasus orientalis 7500'-9600' (Rupr!), Abchasia, Mingrelia, Guria (Eichw. Radde!), Georgia Caucasica (Hohén!).

β. latifolia. — Folia abbreviata radicalia ovato-vel oblongo-spatulata obtusa. — Ic. Bot. Mag. tab. 1002.

Hab. in Caucaso (Stev!), Armeniâ Rossicâ (Szov!), Alpibus Ghilani (Auch. 5009!).

39. *V. monticola* (Trautv. Pl. Nov. Dec. 1866, p. 8) pluricaulis caulis sub lente sparsim puberulis a basi procumbente radicante ascendentibus elatis simplicibus in racemum laxum abeuntibus, foliis glabris oblongis et oblongo-lanceolatis acutis acute et ascendentim dentato-serratis basi cuneatis inferioribus breviter petiolatis, pedicellis tenuibus erecto-patulis calyce 3-4 plo longioribus, bracteis oblongo-linearibus integris praeter infimas pedicello brevioribus, calycis laciniis 4 ellipsoideo-oblongis obtusis quintâ minore, corollâ calyce duplo longiore, capsulâ calyce duplo longiore minute puberulâ ellipsoideo-ovatâ obtusâ non retusâ, seminibus numerosis minimis ovatis. ♀.

Hab. ad latus meridionale montis Nachar Abchasiæ 6-7000' (Radde!).

Fere pedalis, folia intermedia 1 $\frac{1}{2}$ -2 pollices longa. Facies *V. Ponæ* capsulâ obcordata, foliis obtusis etc. distinctæ

40. *V. bellidioides* (L. Sp. 12) glanduloso-hirta, caulis brevibus cespitosis basi prostratâ radicantibus ascendentibus, foliis obovatis obtusis apice crenulatis ad basin caulis confertis, racemo terminali longe pedunculato confertum paucifloro subcapitato villoso, foliis secus pedunculum 2-4 oppositis remotis diminutis, pedicellis strictis calyci vix æquilongis, corollâ cærulescente laciniis calycinis oblongis obtusis longiore, capsulâ calyce duplo longiore magnâ ovalâ vix retusâ glandulosâ, seminibus numerosis ovatis umbilicatis ♀. Ic. Rchb. Germ. tab. 95.

Hab. in regione alpinâ Macedoniæ in Scardo (Griseb!), Thraciâ in alpinis Kalofer (Friv!).

Ar. Geogr. Pyrenei, Alpes, Sudeti, Italia media, regio Danubialis.

41. *V. Kotschyana* (Bth. in D.C. Prodr. 10, p. 480) cespitosa rhizomate sublignoso multicauli, caulis brevibus diffusis et ascendentibus crassiusculis duris inferne denudatis squamulosis dein

crebre foliosis pilosis, foliis subimbricatis oblongo-linearibus ciliatis et margine revolutis acutiusculis carnosulis strictis, spicâ terminali densâ ovato-oblongâ villosâ, pedicellis brevissimis, calycis lanatuli laciniis oblongo-linearibus obtusis subaequalibus, corollâ roseâ calyce longiore, capsulâ glanduloso-hirtâ obovatâ breviter et anguste retusâ subcompressâ calyce longiore, seminibus minimis numerosis oblongis biconvexis subumbilicatis ♀.

Hab. in fissuris rupium regionis alpine Tauri Cilicici 1000' (Ky. 117!) et 390 sub *V. Billardieri!*, Bal!.

Caules 2-3-pollicares, folia 3-4 lineas longa 4 lata, spica densa 1½-pollicaris vel brevior, capsula 2 lineas vel paulo amplius longa. Suffruticulus facie *Thymbrae capitatae*. Species ex affinitate *V. satureioides* Vis. Dalmaticæ a quâ tamen distinctissima.

42. ***V. Thessalica*** Bth. in d.c. Prodr. X, p. 480 pilis crispulis brevibus parce velutina cespitosa, caudiculis prostratis radicantibus, ramis diffusis simplicibus crebre foliosis, foliis parvis sessilibus basi attenuatis ovato-rhombeis acutiuscule dentatis, floribus 3-5 ad ramorum apicem approximatis, pedicellis bracteâ oblongâ integrâ calyceque subbrevioribus, calycis glandulosi laciniis oblongo-spathulatis obtusis subaequilongis, corollâ cœruleâ vel carueâ calyce longiore, capsulâ.... ♀. *V. erinoides* Boiss. Diagn. Ser. I, 4, p. 78.

Hab. in regione alpinâ montis Kyllenes Peloponnesi ad nives 7000' (Heldr!), in monte Parnasso 6500' (Sprnrl!), Olympo Thessalo (Auch. 1969!).

Plantula ex speciminibus paucis et tantum floriferis mihi nota ulterius investiganda, facie *V. thymifoliae* affinis sed ab eâ certe distincta foliis (2-3 lineas longis) rhombeis acutiuscule et crebre dentatis, floribusque terminatis nec in pedunculo axillari racemosis.

43. ***V. serpyllifolia*** (L. Sp 15) caulis minute puberulis basi prostratis radicantibus dein ascendentibus inferne saepè ramulosis in racemum laxum tandem elongatum abeuntibus, foliis glabris lucidis integris vel obsolete crenatis inferioribus et ramorum steriliū petiolatis ovatis intermediis cuneatis superioribus in bracteas oblongo-lanceolatas transeuntibus, pedicellis erectis calyce sublongioribus bracteas saltem superiores excedentibus, calycis laciniis ovato-oblongis subaequalibus, corollâ albâ vel pallide cœruleâ calyce parum longiore, capsulâ compressâ suborbiculari-obcordatâ transverse latiore calyce breviore, seminibus alterâ facie convexis umbilicatis ♀. Ic. Fl. Dan. tab. 492. — Rehb. Germ. tab. 97. — *V. Bungabecca* Janka Brev. II, p. 7 (forma submoustrosa).

Hab. in pratis humidis et umbrosis præsertim montanis et subalpinis, Macedonia et Thracia montes 2-6000' (Griseb., Orph! Friv!), Olympus Bithynus et Tmolus (Boiss!), Anatolia borealis (Wied!), Pontus prope Trapezuntum et Rhizé (Bourg! Bal!). Tauria (Stev!), Caucasus omnis (M.B.), Transeaucasia (Szov!), Persia borealis (Buhse! Bge!).

Ar. Geogr. Europa fere tota, Sibiria, jugum Himalaicum, Africa borealis, America borealis et australis.

44. **V. cespitosa** (Boiss. Dagn. Ser. I, 4, p. 79) cespitosa pulvinata, caudiculis intricatis filiformibus nudis prostratis, ramis nanis confertissime foliosis, foliis plus minus lanatis margine valde revolutis anguste linearibus obtusis, floribus 1-4 terminalibus exscapa vix exsertis, pedicellis ealyxi et bracteæ oblongo-lineari sub-aequilongis, calycis longe lanati lacinis oblongo-linearibus obtusiusculis inaequalibus, corollæ magnâ roseâ, capsulâ glabrâ calyci subaequilongâ orbiculari-obcordata transverse sublatiore, seminibus rugulosis alterâ facie umbilico elevato ovato obsoitis ♀.

Hab. in cacumine Olympi Bithym (Auch 1970! Boiss! Clem! Pichler!), Cadmi supra Ghyra in Cariâ (Boiss.), montis Davrosdagh Pisidie 7000' (Heidr!), montis Masmeneudagh Cappadocie 9000' (Ball!).

β teiophylla. — Folia glabra, inflorescentia hirsuta.

Hab. in cacuminibus Libani supra Eden (Ky. 301! Bl!).

Rami parte foliatâ 6-10 lineas tantum longi, folia 3-4 lineas longa vix lineam lata, capsula vix sesquilinea longa

45. **V. bombycina** (Boiss. et Ky. Diagn. Ser. II, 3, p. 171) adpresso bombycino-nivea cespitosa pulvinaris, caudiculis fragilibus nudis ramisque confertim foliosis brevissimis, foliis minimis sessilibus ovato-vel oblongo-spathulatis, floribus 1-3 terminibus pedicellis calyce brevioribus suffultis, bracteis minimis ovatis, calycis lacinis oblongis obtusis, corollæ rubellæ calyce plus duplo longiore, capsulâ parvâ tomentosâ calyce subbreviore orbiculari basi subcuneatâ apice obtusâ vix retusa, semiinibus ovatis convexis ♀.

Hab. in fissuris rupium summi montis Makmel Libani 8500' (Ky. 301!). Fl. Jul.

Nana, folia 2 lineas circiter longa, capsula 1½ lin longa et lata. Facie affinis *V. cespitosa* sed characteribus distinctissima

SECT. IV. PÆDEROTOIDES.

46. **V. Pæderotæ** (Boiss. Diagn. Ser. I, 4, p. 78) breviter et parce puberula cespitosa, caudiculis fragilibus longis nudis remote squamulosis prostratis, ramis brevissimis foliosis, foliis breviter petiolatis carnosulis ovatis obtusis obtuse crenulatis, racemis pedunculatis subcapitatis villosis axillaribus vel gemme terminalis brevis abortu terminalibus, pedicellis calyce brevioribus bracteas lineares aspernitibus, calycis lacinis longe linearis-spathulatis subaequalibus, corollæ carminea infunditiflora tubo calyce breviore lobis oblongis acutiusculis aeque longo, capsula (juniore) calyce duplo breviore puberula compresso-planâ apicib; lobis subdivaricatis ♀.

Hab. in sehistosis regionis alpina montium Pæstæ borealis, mons Elamout (Auch. 5088!), alpes Hasartschel regi Fibrusens (Ky. 500!).

Caudiculi bipolticares, folia conferta 3-4 lineas longa et lata, calyx lineas tres, corolla 4 longa, tacies *Limaria crassifolia* vel *Veronica macrostemon* Bge.

SECT. V. PSEUDGLASIMACHUM.

47. V. spuria (L. Sp. 14) velutina-pubescent, caulis clavis strictis, racemis terminalibus paniculatis, foliis oppositis vel 3-4 verticillatis a basi angustata vel cuneata oblongo-lanceolatis, argute serratis, bracteis linear-lanceolatis petiello brevioribus, pedicellis calyce sublongioribus, calycis laciniis oblongo-triangularibus, capsula ovalo-subrotundata turgida vix retusa glabra ♀. Ic. Rehbe. Germ. tab. 94. — *V. foliosa* W. K. Hung, tab. 102. — *V. brevifolia* M.B. I, p. 6. — *V. paniculata* L. Sp. 18.

Hab. in Hunnelicâ (Nocht), in Caucaiso (ex M.B.).

Ar. Geogr. Germania, regio Danubialis, Rossia merid et austri et Sibiria.

48. V. longifolia (L. Sp. p. 13) breviter pubescens vel subvelutina, caulis clavis strictis, racemis terminalibus paniculatis densiusculis, foliis oppositis vel 3-4-nis a basi ovata vel subcordata lanceolatis acutissime serratis, bracteis subauantatis petiellum calyci subaequilongum aequantibus, calycis laciniis acutis, capsula ovato-rotunda turgida retusa glabra ♀. Rehbe. Germ. tab. 93. — *V. maritima* et *V. chitica* M.B., — *V. spuria* Sm. Prodri. I, p. 6 non L.

Hab. in Arcadia (herb. Fauché), Hunnelicâ (Nocht), Armenia (Göstrem), 5000' (Tchihly) et ad lacum Gestemert prov. Musch 5200' (Ky. 295 sur le Jolma var.), Caucaiso (ex M.B.).

Ar. Geogr. Scandinavia, Germania, regio Danubialis, Dalmatia, Rossia, Songuria, Sibiria.

49. V. spicata (L. Sp. 14) tonitrella canescens, caule ascendenti in racemum subsolitarium spiciformem censim denuo, foliis oppositis petiolatis crenatis apice integris inferioribus ovatis obtusis superioribus lanceolatis brevibus, calycis pedicello duplo longioris laciniis oblongis obtusis, corolla laciniis oblongis tubo duplo longioribus, corollâ globosa turgida glanduloso-hirta vix retusa ♀.

Hab. in pascuis rie s. Laonia (ex Smith Prodri.), Tauria (M.B.), Caucasus (M.B.).

Ar. Geogr. Europa borealis et media a Scandinavia ad Rossiam, Hispaniam borealem, Italiam austriadem, Dalmatiam, Sibiria Urdensis et Altide.

50. V. Orchidea (Crantz Stirp. IV, p. 333) parce hirta siccitate nigricans, caule elato firme, racemo denso terminali solitari vel 3-5 paniculatis, foliis oppositis obtuse crenatis inferioribus ovatis petiolatis superioribus oblongis et ellipticis sessilibus, floribus subsessilibus, calycis laciniis ovatis glandulosis, corolla laciniis fine cri-acuminatis tubo quadruplo longioribus dum torto-conniventibus, fince barbata, capsula globosa turgida valde glanduloso-hirta subretusa ♀. Ic. Bochel. Baum. fig. 42. — *V. hybrida* M.B. Taur. Gane. I, p. 8. — *V. spicata var. cristata* Koch Rehbe. Germ. tab. 92, fig. 2.

Hab. in saxosis subalpinis montis Rhodopes Thraciae (Jankal!), Caucaiso boreali (Owerint!).

Ar. Geogr. Regio Danubialis.

51. **V. incana** (L. Sp. p. 14) adpressissime incano-tomentosa, caule a basi ascendentē erecto racemum simplicem densum vel racemes 3-5 paniculatos edente, foliis oppositis minute crenatis inferioribus oblongis petiolatis superioribus oblongo-linearibus integris, pedicellis brevissimis, calycis laciuiis canis elliptico-linearibus obtusis, capsula pubescente retusa calyce breviore ♂.

Hab. in rupestribus regionis subalpinæ et alpinæ montium Tauriae (M.B., Rehm!).

Ar. Geogr. Rossia australis, Sibiria, Mongholia Sinensis.

SERIES II. — Annuæ.

SECT. VI. ALSINEBE.

* Semina plano-biconvexa facie plus minus manifeste umbilicata.

+ Pedicelli calyce breviores vel eo vix longiores.

52. **V. Chamæpitys** (Griseb. Spic. II, p. 25) pumila a collo saepius ramosa caulinis simpliciuseulis crispule hirtis lateralibus ascendentibus, foliis imis ovatis petiolatis obtuse paucidentatis cæteris in 3-5 lacinias lineares pinnatisectis omnibus ciliatis, floribus sessilibus spicas densiusculas strictas foliatas rigidas formantibus, bracteis linearibus flores superantibus præter infimas indivisis, calycis laciuiis linearibus subinæqualibus obtusis, corollæ albâ cœruleo-striatâ calyce ampliore, capsula asperâ calyce longe supærata deltoideâ apice truncatâ vel late subretusâ durâ stylo longe superatâ, seminibus 12-15 ovato-subquadratis ⊖. *V. digitata* Auct. quoad pl. Orientalem non Vahl.

Hab. in collibus, agris sterilibus montanis et alpinis, Thracia prope Ruskoï (Griseb!), Byzantii et montes prope Angora (Wiedem!), Smyrnæ montes inter frutices (Boiss! Bal. 409!), Phrygia prope Ouchak (Bal!), Pisidia ad Baldur (Heldr!), Pamphylia supra Adalia (Bourg!), Cappadociae mons Alidagh (Bal. 1026!), Caria ad Mog'ah (Auch. 1952!).

Species a cl. Grisebach recie a *V. digitatâ* Vahl Hispanicâ distincta quæ differt foliis omnibus pinnatisectis, bracteis tripartitis, corollâ calyce duplo breviora minutâ, capsula calycem subæquante, stylo brevissimo.

53. **V. verna** (L. Sp. 19) pubescens pumila simplex vel a basi ramosa, foliis subpetiolatis imis ovatis integris cæteris in 5-7 lobos obtusos pinnatifidis, floralibus lanceolato-linearibus integris, racemis spiciformibus tandem longis laxiusculis, pedunculis strictis calyce brevioribus, calycis laciuiis lanceolato-linearibus corollam

pallide cæruleam superantibus, capsulâ glanduloso-ciliatâ compressâ obcordatâ longitu-line sublatiore angulo aperto bilobâ lobis rotundatis stylo sinu breviore, seminibus 12-15 ovatis minute umbilicatis ⊖. Engl. bot. tab. 25. — Rehb. Germ. t. 99. — *V. succulenta* All. *V. pinnatifida* Lam. № 194.

Hab. in cultis arenosis, sterilibus regionis præsertim montanæ et alpinæ, Laconia mons Malevo in cacumine (Orph!), Macedonia mons Korthiatî (Orph!), Pontus Lazicus ad Djimil in tectis (Ball!), Armenia (Huet!), Cappadocia in cultis montis Argæi 6000' (Ball!), Tauria, Caucasus et Transeaucasia (Ledeb.), Affghania ad Otipore (Griff!).

Ar. Geogr. Europa omnis a Scandinaviâ ad Rossiam, Sibiria Altaica.

54. *V. arvensis* (L. Sp. 18) pilosiuscula simplex vel diffuse ramosa pumila, foliis subpetiolatis cordato-ovatis crenatis, floralibus sessilibus breviter lanceolatis integerrimis, racemis spiciformibus tandem longis laxis, pedicellis rectis calyce brevioribus, corollâ minutâ pallide cæruleâ, capsulâ calyce breviore glanduloso-hirsutâ compressâ obcordatâ ad tertiam partem bilobâ lobis rotundatis, stylo fissuram subæquante, seminibus 15-20 ovatis minute umbilicatis ⊖. Ic. Engl. bot. tab. 734. — Rehb. Germ. tab. 99.

Hab. in cultis et saxosis montanis, Peloponnesus (herb Fauché!) Parnes Atticæ et Parnassus prope Rachova (Heldr. Herb. Norm. 746!), Macedonia et Thracia (Griseb.), Pontus Lazicus ad Djimil 6000' (Ball!), Syria ad Aintab (Haussk!) et Berythum (Gaill!), Cyprus (Sibth.), Tauria et prov. Caucasicis (M.B.), Talysh (Hohen.).

Ar. Geogr. Europa omnis a Scandinaviâ ad Rossiam, Songaria, Sibiria, Africa borealis, insulæ Canarienses.

55. *V. Sartoriana* (Boiss. et Heldr. Diagn. Ser. II, 3, p. 171) viscidulo-hispidula simpliciuscula pumila, foliis subpetiolatis ovatis crenatis subcordatis, floralibus sessilibus linearibus brevissimis, racemo brevi laxo paucifloro, pedicellis rectis calyce sublongioribus, corollâ minutâ, capsulâ obcordatâ ciliatâ compressâ calycem subæquante ad quartam partem bilobâ lobis rotundatis, stylo fissurâ subbreviore, seminibus 8-10 orbiculatis ⊖.

Hab. in pascuis saxosis montis Parnethos Atticæ et Parnassi prope Rachova in consortio *V. arvensis* (Heldr. Herb. Norm. 748!, Guiceardi!).

Valde affinis *V. arvensi* a quâ tamen distincta videtur racemis paucifloris, foliis floralibus brevioribus, pedicellis longioribus, capsulâ maiore quoad calycem longiore et præsertim seminibus duplo majoribus orbiculatis paucioribus.

56. *V. viscosa* (Boiss. Diagn. Ser. I, 12, p. 47) valde glanduloso-pubescent siccitate nigricans nana a collo ramosa, foliis brevisime petiolatis ovato-subcordatis obtuse crenatis, floralibus sensim diminutis inferioribus oblongis lobato-dentatis supremis ellipticis integris basi attenuatis, racemis densiusculis hispidis, pedicellis deum horizontaliter patentibus apice recurvis calyce et bracteâ sub-

longioribus, corollâ caeruleâ lacinis calycinis ovato-ellipticis acutis non longiore, capsulâ erectâ glabrâ obcordatâ compressâ ad medium divaricatum bilobâ lobis reticulatis obtusis, stylo fissu et aequilongo, semiinibus 4 orbiculatis ♂.

Hab. in arenosis regionis a fine Libani (Boiss!), in desertis Mesopotamiae circa Orfa, Ma-Ara, Kotschhae (Hanssk!).

Demum tri officinalis, cupri et eis *V. arvensis* triplo maiores, ab ea et *V. Sartoriana* pedicellis longis patentibus, capsule lobis divaricatis, etc., distincta.

⊕ ⊕ Pedicelli calice manifeste et plures longi res.

1. Capsula sinu angusto biloba

37. *V. acinifolia* (L. Sp. 19) glanduloso-hirta unila simplex vel a basi ramosa caulinibus lateralibus ascendentibus foliis breviter petiolatis ovalis suberectis, floralibus secundum dimiditatis elliptico-lanceolatis ut inque aitematis integri, ramo lanceolato elongato laxissimo, pedicellis fructiferis tenuibus erecto-palulis calyce 3-4-plo longioribus corollie intense caeruleae fave flavidæ calyce sublongioris lobis superioribus transverse latioribus inferiore minore, capsulâ glanduloso-ciliata transverse latiori compressâ calyce submajore ad medium in lobos orbiculatos divisâ sinu angusto, stylo sinu aequilongo, loculis 7-8-spermis, semiinibus ovalis labiis ♂. Ie. Vaill. Bot. tab. 43. — Relib. Germ. tab. 38, fig. 2. — *V. Romana* All. — *V. Gorum-sensis* Boiss. et Ky. in Sched.

Hab. in celtis, herbidis humidis presertim montanis et alpinis, Peloponnesus (Herb. Fanehæ), Attica in Parnethos et amioe (Heldr.), Creta (Heldr.), Thrace (Fris.), Lydia ad Smyram (Boiss!) et in monte Metopæ (Boiss!), Cilicia in alpini Tauri (Batt.) et in divisione Kotsangor (Ky. 50!), Serbia borealis (Hanssk!), Mesopotamia in huncidis (Hanssk!), Cyprus (Ky.), Armenia prope Guinu-chkhanæ (Bourgl.).

Ar. Geogr. Belgium, Gallia, Germania occid., Helvetia, Italia, regio Danubialis, Dalmatia, Madera.

38. *V. nudicaulis* (Kar. et Kir. Bull. Mosq. 1842, p. 413) puberula vel glabra, caule pusillo filiformi nano subsimplici in racemum paniciformem brevem laxum abeunti, foliis minimis oblongis integris basi angustatis sessilibus floralibus linearis-oblongis pedicellis recurvatis, pedunculis calyce sesquilocigibis patentibus, corollâ... capsulâ glabrâ vel glanduloso-ciliata calyce sublongo et compressâ transverse sublatiore ad $\frac{2}{3}$ sinu anguste bilobâ, loculis ovalis 3-5-spermis, stylo sinu duplo breviore ♂. *V. perfoliata* Boiss. Diagn. Ser. I. 7, p. 43. — *V. pusilla* Ky. exs. non Blh. — *V. acinifolia* var. *glabra* et var. *Karelinii* Trautv.

Hab. in celtis, alpinis Persie australis in alpe Kub Deh (Ky. 53!) et ad eius membris S. Casp. (Ky. 130!) (Hanssk!), in monte Totschal prope Teheran et Cilicia Persie borealis (Ky.).

Species erroneè *Omphalospora* adnumerata, seminibus plano-biconvexis donata et valde affinis *V. acinifoliae* a qua tamen foliis mininis oblongis integris, pedicellis fructiferis patentibus, stylo brevissimo differt. Specimina nostra Songaricis similia sed maiores.

Ar. Geogr. Songaria.

59. V. ixodes (Boiss. et Bal. Diagn. Ser. II, 3, p. 172) glanduloso-hispida a basi pluricaulis punica, foliis integer mis deltoideis in petiolum brevissimum attenuatis, floralibus ellipticis et linearibus pedicellos subaequantibus racemis longis tenuibus pedicellis fructiferis erecto-patulis arcuatis calyce vix longioribus, corollâ albâ parvâ calycem vix excedente, capsulâ ciliatâ calyce sublongi re transverse latiore compressâ ad medi um in lobos obovatos sinu angusto bipartitâ, stylo loculis 7-8-spermis dimidio breviore.

Hab. in regione alpinâ mo iitis Maesmenendagh Cappadociae (Bal.), V. sicc et viv. cult.

Affinis *V. acinifoliae* differt foliis integrerrimis deltoideo-circumatis, pedicellis brevioribus, corollâ albâ minntâ, styl. brevi. *V. undulatilis* Kar. et Kir. nostrae stylo abbreviato affinis differt foliis minimis ellipticis, pedicellis calyce longioribus valde patentibus apice recurvis.

60. V. conferta. nana glaberrima raris breviter et crispule hirtula simplex vel collo ramulosa, foliis minimiis oblongis integris basi attenuatis, floralibus conformibus pedicellos aequantibus vel superantibus, racemis fere radicalibus brevibus densius utis, pedunculis fructiferis patentibus arenatis calyce 1½-2-plo longioribus, capsulâ parvâ glabrâ turgidulâ calyci subaequilonigâ transverse subtaliore ad ³/₄ in lobos rotundatos sinu angustissimo partitâ, stylo loculis 3-4-spermis tripto breviore ⊖. *V. glaberrima* Boiss. et Bal. Diagn. Ser. II, 3, p. 172.

Hab. in pratis humidis regionis alpinei Tauri Ciliciei supra Bulgharmaaden (Bal. 689), in excelsis Libani (Bl. exs. 472!), in Ponto Lazio supra Djimi (8000') (Bal. 4).

Pollinosis, affinis *V. acinifoliae* et *ixodi*, ab utrâque racemo abbreviato distiore, foliis angustis, capsulis minoribus turgidulis oligospermis distincta et hoc motu a *V. undulatilis*. Nomen speciemini de specimina quedam hirta mut. redidit.

2. Capsulae loculi sinu lato recto vel obtuso divergentes.

61. V. Reuteriana (Boiss. Diagn. Ser. I, 12, p. 46) glandulosohirta nana simplex vel basi ramosa, foliis subpetiolatis parvis obovatis integris, floralibus linearispathulatis pedicello aequilongis, racemo etiam fructifero brevi, pedicellis fructiferis horizontalibus tice sursum flexis calyce 2-3½ longioribus, corollâ, capsulâ marginis glandulosâ ad ³/₄ birectilobâ calycem superante lobis angulo recto divergentibus obovalis, stylo loculis 4-5-spermis duplo breviore ⊖.

Hab. circa Diarbekir Mesopotamiæ (Ky. 240!), in argillosis subhumidis montis Mandschura Antilibani 5500' (Ky. 52!).

Plantula 4-1½-pollicaris facie *V. acinifoliae* a quâ præter folia integra differt capsulæ loculis oligospermis et angulo lato recto nec angusto divergentibus.

62. ***V. hispidula*** (Boiss. et Huet Diagn. Ser. II, 3, p. 172) pumila superne glanduloso-hirta, caule filiformi inferne parce folioso supra medium subcorymbose 3-5-racemoso, foliis minutis integris interioribus oblongis in petiolum brevissimum attenuatis, floralibus linearibus pedicellis subbrevioribus, racemis densiusculis, pedunculis patentibus sursum recurvis capsulâ vix longioribus, corollâ azureâ minutâ calycem vix excedente, capsulâ glandulosâ calyce sublongiore transverse latiore ad ¾ longitudinis in loculos obovatos angulo aperto divergentes partiâ, stylo sinu quabruplo breviore, loculis 3-4-spermis ⊖.

Hab. in arenosis Armeniæ prope Erzerum (Huet!), Phrygiæ ad Yachamicklar Kieui prope Ouchak (Bal!).

β. minima. — Specimina pusilla 3-4-flora. — *V. minima* C. Koch Linn. XVII, p. 288.

Hab. in Armeniâ Rossicâ ad radices montis Alagös (C. Koch!).

Planta 2-3-pollicaris, racemi fructiferi 1-1½-pollicares, capsulæ eis *V. acinifoliae* fere duplo minores; ab eâ præter folia differt loculis capsulæ sinu fere recto sejunctis divaricatis styloque brevi. *V. Reuteriana* magis affinis differt foliis latiusculis, capsulâ duplo majore, stylo longiore.

63. ***V. Syriaca*** (Ræm. et Sch. Syst. I, p. 116) glanduloso-hirta basi sæpius ramosa ramis brevibus, racemis demum longis laxis, foliis ovatis crenatis vel dentatis imis subpetiolatis, bracteis pedicello multo brevioribus linear-ellipticis integris, pedicellis capillaribus calyce multoties longioribus fructiferis erecto-patulis, corollâ azureâ magnâ sepalis oblongis duplo longiore, capsulâ compressâ glandulosociliatâ calyci æquilongâ transverse latiore ad mediani partem in loculos ovatos obtusos angulo lato bifidâ, stylo sinu duplo longiore, loculis 8-10-spermis, seminibus ovalis rugulosis ⊖. *V. pedunculata* Labill. Dec. 5, p. 8, tab. 5.

Hab. in pratis et cultis regionis inferioris Pamphyllæ prope Adalia (Heldr!), Ciliciæ (Auch. 1938!), Aleppi (Ky. 31! Haussk!), Damasci (Lab.), Syriae littoralis ad Berythum (Bl!), Palestinae (Boiss!).

64. ***V. ceratocarpa*** (C. A. Meyer Enum. p. 106) crispule papillosa, caulibus elongatis diffusis in racemos longos laxos abeuntibus, pedicellis filiformibus calyce 4-5-plo longioribus demum patentibus, calycis laciniis lanceolatis acutis capsulâ sublongioribus, corollâ pallide erubescâ calyce sublongiore, capsulâ subiunctante compressâ hirtâ elevatum reticulatâ transverse multo latiore in loculos oblongo-lanceolatos acutos angulo obtuso valde divergentes ultra medium bipartitâ, style longo, loculis 2-3-spermis, seminibus ovatis

radiatim subrugosis umbilico elevatim marginato ⊖. *V. reticulata* C. Koch Linn. XXIII, p. 552!

Hab. in pratis Transcaucasie inter Sallian et Lenkoran ad Caspium (G.A. Mey!).

Rami cum racemo $\frac{1}{2}$ -1-pedales, folia inferiora 4-5 lineas longa, pedicelli fructiferi 9-10 lineas longi, capsula 4 lineas lata 2 longa.

65. V. divaricata (Boiss. et Bal. Diagn. Ser. II, 3, p. 173) glanduloso-hirta pumila basi siepius ramosa, foliis minutis oblongis crenatis subpetiolatis, floralibus anguste linearibus pedicellis brevioribus, racemis laxis, pedicellis tenuibus calyce duplo longioribus fructiferis patentibus apice sursum curvatis, corollā azureā calyce duplo longiore, capsulis glandulosis in loculos 3-4-spermis ellipticos apice attenuatos angulo obtuso valde divergentes ad $\frac{3}{4}$ bipartitis, stylo longissimo sinum superante, seminibus oblongis minimis brevibus non umbilicatis ⊖.

Hab. in Ciliciā littorali ad pagum Alladagh non procul a Mersina (Bad. 688!).

Plantula 2-3-pollicaris, folia inferiora 4-5 lineas longa, racemi fructiferi 2- $\frac{1}{2}$ -pollicares. Corolla fere tres lineas diametro lata. Species loculis angustis et valde divaricatis capsule ea *V. ceratocarpa* similis sed triplo minoris et non elevatim reticulatae insignis. Semina insuper plano-convexa oblonga laevia.

** Semina cyathiformia facie concava dorso convexa

+ Calycis laciniæ fere ad basin usque liberae.

I. Folia floralia saltem superiora bracteiformia.

66. V. glauca (Sibth. et Sm. Fl. Gr. I, p. 6, tab. 7) glauca crispula et brevissime pubescens opposite et patentim ramosa, foliis ovatis dentatis imis cuneatis subpetiolatis caeteris subcordatis, floralibus parvis imis dentatis caeteris breviter linearibus, racemis laxis, pedicellis filiformibus calyce et bracteâ 3-5-plo longioribus fructiferis patentibus apice recurvis, corollā ceruleâ amplâ calycis laciniis ellipticis triplo longiore, capsulâ orbiculatâ glabrâ truncato-subretusa calycis laciniis subbreviore, loculis 6-7-spermis, stylo longo ⊖. *V. Graeca* Sprun. exs.

Hab. in cultis arenosis Atticae frequens (Sibth., Sprun! Boiss! Heldr!). Fl. Mart., Apr.

Specimina majora 6-7-pollicaria, folia 6-8 lineas longa 6 basi lata, corolla 5 lin. diam. lata, calycis laciniæ numquam dentatas vidi ut in Florâ Graecâ describuntur sed ultra insertionem breviter calcaratæ. *V. pumila* Clarke species male nota ex Bulgariâ extra fines ditionis nostræ oriunda *V. glauca* affinis videtur sed viscositate diversa.

67. V. Chaubardi (Boiss. et Reut. Diagn. Ser. II, 3, p. 174) pilis brevibus crispulis hirta patentim ramosa, foliis ovatis inferioribus petiolatis crenatis intermediis sessilibus subcordatis dentatis vel

subincisæ, floralibus inferioribus foliis similibus superioribus integris anguste et breviter oblongo-linearibus, racemis tandem elongatis laxis, pedicellis filiformibus tandem longissimis arcuato-deflexis, corollâ ceruleâ amplâ calyce triplo longiore, capsulâ compressâ glabrâ calyci æquilonigâ transverse latiore fere ad medium bilobâ loculis obovatis obtusissimis, stylo longo, seminibus magnis sphæricis cyathiformibus lævibus ⊖ *V. peduncularis* Fl. Pelop. p. I, non M.B.

Hab. in Messeniâ prope Methonem, Calamata, Armyros, Pylos, Phigaleam (Bory!).

Facie *V. glauca* affinis sed non glauca et capsula biloba.

68. **V. Peloponnesiaca** (Roiss. et Orph. in Fl. Graec. exs. 716) patule glanduloso-hirta basi opposite et patule ramosa, foliis ovatis dentatis imis subpetiolatis cæteris sessilibus subcordatis, floralibus minutis elliptico-linearibus integris, racemis laxissimis, pedicellis filiformibus calyce 2-3-plo longioribus fructiferis patentissimis apice recurvis, capsulâ glandulosâ rarius glabrâ orbiculatâ compressâ truncato-subretusâ calyce subbreviore, loculis 7-8-spermis, stylo longo, seminibus lævibus cyathiformibus ⊖.

Hab. in Argolide (Sprun!) monte Malevo Laconiaæ supra Hagios Joannis 300'-4000' (Orph!), in regione superiori montis Helicon Bœotiaæ (Orph!), in monte Cénos Cephaloniae 4000'-5500' (Heldr!), insula Tinos (Wiedem!).

Similis *V. glauca* et *Chaubardi* a priori indumento glanduloso, a posteriori indumento et capsulâ subretusâ nee profunde bilobâ differt.

69. **V. amœna** (Steven in M.B. Taur. Cauc. I, p. 14) crebre pubescenti-canescens glandulosa pumila simplex vel inferne ramosa, foliis parvis petiolatis ovatis crenatis, floralibus elliptico-linearibus integris pedicelle brevioribus, racemis demum laxis, pedicellis calyci æquilonigis vel brevioribus fructiferis strictis, calycis laciñiis lanceolatis acutis, corollâ cyaneâ fundo albida magnâ calycem superante, capsulâ hirsutâ calyce breviore turgidulâ orbiculari-obcordatâ sinu longiore, seminibus oblongis cyathiformibus lævibus ⊖. Ic. M.B. Cent. tab. 18! — *V. Albanica* C. Koch Linn. XXIII, p. 701!

Hab. in Transcaucasiâ, Iberia (Stev!), prope Baku ad Caspium (C.A. Mey! Seidl! C. Koch!), ad Schemacha (Bayern!).

Planta 2-3-pollicaris floribus magnis insignis, sæpe confusa cum *V. glauca*, *V. Chaubardi* et *V. Peloponnesiaca* pedicellis longis patentibus statim distinguendis. Specimen imperfectum huic in Boiss. et Buhse Enum. relatum huc non spectat.

70. **V. grandiflora** (Don Ann. Nat. Hist. Ser. I, VII, p. 458) glanduloso-pubescentis, caule pumilo erecto simplici vel ramoso, foliis inferioribus brevissime petiolatis ovatis obtusis crenatis, superioribus sessilibus pinnatis vel tripartitis segmentis linearibus obtusis terminali majore subspathulato, floribus racemosis pedicellis capillaribus glandulosis folio axillari triplo longioribus suffultis, calycis glandulosi

laciiniis quaternis linearibus obtusis anticis submajoribus, corollæ maxime cyaneæ laciiniis rhomboeo-ovatis anticâ minore, ovario subrotundo glabro integro ♂.

Hab. in Cariâ ad Meandrum et prope Mylassa (Fellows). Non vidi.

Bipollicularis, corolla diametro sex lineas lata. Capsula et semina ignota. Ex cl. auctore affinis *V. amurensis*.

71. *V. præcox* (All. Auct. 3, tab. I) glanduloso-pilosa pumila simplex vel basi ramosa, foliis subpetiolatis ovato-subcordatis dentatis, floralibus sessilibus parvis oblongis inferioribus crenatis cæteris integris, racemis demum elongatis tenuibus laxis, pedicellis erectis glandulosis calyce brevioribus vel vix longioribus, calycis laciiniis oblongis corollâ cæruleâ brevioribus, capsulâ subglandulosâ calyce sublongiore oblongâ turgidâ breviter retusa, stylo sinu longiore, seminibus globosis cyathiformibus levibus ♂. Ic. Rehb. Germ. tab. 100, fig. I.

Hab. in arenosis montanis et alpinis, Phrygia ad Boulghas Keui 3000' (Bal!), Crete montes Lassiti 5000' (Heldr!), mons Akdaghi Lyciae (Bourg!), Sonchetia Caucasică (C. Koch), Tauria (Stev.).

Ar. Geogr. Europa media et australis a Belgio, Galliâ, Germaniâ ad regionem Danubialem et Rossiam medium, Africa borealis.

72. *V. Bungei*, tota papillis albis obsita viscidula pumila simplex vel collo ramosa, foliis minatis subsessilibus ovatis crenato-dentatis, floralibus minoribus oblongis calyces subaequantibus integris, racemo brevi densiusculo, pedicellis fructiferis crassiunculis recurvis calyce brevioribus, sepalis oblongo-lanceolatis obtusis capsulâ longioribus, capsulâ turgidâ pubescente breviter et angulo obtuso bilobâ lobis rotundatis stylo sinu sublongiore, seminibus ovatis cyathiformibus minimis sub lente tuberculatis ♂.

Hab. in alpe Ssiachaneh Persiae boreali-orientalis inter Siaret et Schahrud (Bge!).

Bipollicularis, folia inferiora tres lineas longa, racemi 7-10 lineas longi basi laxi. Facies *V. præcocis* a quâ pedicellis fructiferis recurvis statim distinguitur.

73. *V. triphyllos* (L. Sp. 49) patule pubescens glandulosa basi ramosa sape diffusa pumila, foliis imis ovatis petiolatis indivisis, intermediis in 3-5 laciinas oblongo-spathulatas palmatisectis, floralibus sessilibus triseptis et summis indivisis, racemis laxis, pedicellis calyce et bracteis æquilongis vel longioribus erecto-patulis, calycis laciiniis oblongis obtusis corollâ cæruleâ longioribus, capsulâ magnâ suborbiculatâ glandulosâ calycei subæquilongâ obcordata turgidâ, stylo sinu longiore, seminibus levibus concavis ♂. Ic. Fl. Græc. tab. 40. — Rehb. Germ. tab. 40, fig. 2 et 3. — *V. digitata* Lam. Fl. Fr. II, p. 445 non Vahl.

Hab. in cultis arenosis Græciae (Boisst.), Thracie (Sibth.), Lydie (Bal!), Phrygiae in monte Almadagh (Bal!), Cappadocie (Bal!), Syrie ad Aleppo

(Ky! Haussk!), Meopoiamiae (Chesney), Tauris, Caucasi et Transcaucasias (Ledeb.).

Ar. Geogr. Europa media et australis a Daniâ et Gothia ad Rossiam meridiam, Turcomania, Africa borealis.

74. V. biloba (L. Mant. 172) breviter et parce glandulosos puberula, pumila simplex vel interne ramosa, foliis oblongis subpetiolatis serrato-dentalis, floralibus elliptico-lanceolatis acutis integris pedicello subbrevioribus, racemis tandem laxis, pedicellis filiformibus fructiferis erecto-patulis capsulae aequilengis vel sublongioribus calycis segmentis elevatim trinerviis ovato-lanceolatis acuminatis basi per paria breviter connatis, corollâ pallida minitâ calyre breviore, capsula glandulosâ compressâ fere ad basin in loculos obovatos divergentes 2-3-spermis paritâ, stylo sinu multo breviore, seminibus obsolete rugulosis cyathiformibus (Rehb. Icon. VII. tab. 645. — *V. bartsiifolia* Boiss. in Bal. exs. (forma major) — *V. exilis* Schott in Ky. exs. (forma pusilla).

Hab. in rupestribus et arenosis montanis et alpinis, Anatolia in Olympo Bithyno (Auch. 1944), caumine Mesogis (Boiss!), Isauria monte Gheidalagh 6000' (Heldr!), Tauro Cilicio (Bal!), Libani regione alpina (Bl!), Armenia Turcica et Ponto Lazico 6-7000' (Huet! Bal!), Armeniâ Rossica et prov. Talysch 5000' (C.A. M!), Caucaso orientali 7-9000' (Stev!), Persia bor. in monte Elbrus et ad Teheran (Ky!), australi in alpe Kuh Delu (Ky!), Turkestaniâ (Lehm.), Affghania (Griff.).

Ar. Geogr. Songaria, Sibiria, regio Himalaica.

75. V. campylopoda (Boiss. Diagn. Ser. I, 4, p. 80) papillari-hirtula pumila simplex vel basi ramosa, foliis oblongo-lanceolatis inferioribus obovatoe crenatis ceteris integris, floralibus anguste linearie-lanceolatis pedicello aequantibus, racemis laxis tandem elongatis, pedicellis filiformibus calyci aequilongis vel 1 1/2-plo longioribus fructiferis recurvis, calycis lacinias per paria breviter connatis anguste lanceolatis acuminatis trinerviis, corollâ calyce minore, capsula glandulosâ compressâ calyce subbreviore fere ad basin in loculos obovatos divergentes apice subangustatos bipartitâ, stylo sinu duplo breviore, seminibus cyathiformibus va'de rugulosis (Rehb.).

Hab. in cultis et arenosis praesertim montanis et alpinis, Phrygia prope Ouchak (Bal!), Lycae montes Elmalu (Bourg. 1821 sub *V. bilobæ*), Taurus Cilicus (Bal!), Libanus (Bl!), Antilibanus (Boiss!), caumen montis Sinai (Schimp. 118! Boiss!), Mesopotamia (Haussk!), Cappadocia (Bal!), Armenia Turcica (Bourg!) et Rossica (C. Koch!), Assyria (Auch. 1945!), Persia australis (Ky! Haussk!) et boreali, (Auch. 5090! Buhse!), Affghania (Griff!), Belutschia (Stocks!).

β. microtheca. — Bracteæ breviores, capsula dimidio minor, semina minora. — *V. microtheca* Boiss. et Bal. Diagn. Ser. II, 6, p. 131.

Hab. in cultis ad Ouchak Phrygiae (Bal!), ad Aleppo Syriæ (Ky. 145! Haussk!), Persia bor. (Buhse!), Affghania (Griff!).

Species valde affinis *V. bilobæ* a qua tamen foliis angustioribus subintegris,

pedicellis recurvis, sepalis angustis, seminibus valde rugulosis, stylo longiore differre videtur. Var. *microtheca* intermediis ad typum transit.

Ar. Geogr. Songaria.

76. V. rubrifolia (Boiss. Diagn. Ser. I, 12, p. 46) crispule et brevissime puberula rubella siccitate nigricans, collo ramosissima ramis decumbentibus filiformibus rigidulis brevissimis in racemos breves laxos abeuntibus, foliis ad collum rosulatis parvis ovatis crenulatis vel integris subpetiolatis, floralibus oblongo-linearibus integris basi attenuatis pedicello brevioribus, pedicellis capillaribus calyci aequilongis fructiferis patulis apice sursum incurvis, calycis lacinii ellipticis obtusis capsulam aequantibus, corollâ carneâ calyceum aequante, capsulâ subglandulosâ transverse sublatiore ad tertiam partem in loculos obovatos obtusos trispermos fissâ, stylo sinu subdupo breviore, semiibus ovalis cyathiformibus leviusculis ⊖.

Hab. in Persiâ mediâ (Derderian!), in monte Kohrud prope Kaschan (Kapherr!), in monte Elbrus prope Derbend 7-8000' (Ky. 224!), in montibus prope Rischun (Buhsel!), Persiæ australis (Ky!).

Rami cum racemis fructiferis vix bipollicares, capsula sesquilineam lata, facies *V. acinifolia* a quâ ramis decumbentibus rigidis seminisque chartere longe distat.

77. V. macropoda (Boiss. Diagn. Ser. I, 4, p. 79) basi ramosa ramis glabris filiformibus rigidis in racemos longos laxissimos eito abeuntibus, foliis radicalibus rosulatis oblongo-spathulatis sessilibus sub lente hirtulis, pedicellis capillaribus glabris patentibus longissimis basi bracteis minimis oblongo-linearibus suffultis, corollâ caeruleâ calyce vix longiore, capsulâ parce glandulosâ sepalis oblongis aequilongâ transverse latiore ad tertiam partem in loculos ovalis obtusos dispermos lobatâ, stylo sinu angusto duplo longiore, semiibus cyathiformibus rugulosis. ⊖.

Hab. in Persiâ circa Ispahan (Auch. 5094! Derderian!), circa Chabbise, Kerman, Ssertschah (Bge!).

Folia fere *Androsace maximowii*, racemi fere totâ rami longitudine fructiferi 4-6 pollices longi. Species insignis pedicellis capillaribus patentissimis 8-10 lineas longis basi bracteolâ minimâ stipatis. Capsula parva lineam longa sesquilineam lata. Semina que prius ex specimine immaturo plana esse autu-maveram cyathiformia sunt basi constricta. Facies *V. Syriacw*.

2. Folia floralia omnia caulinis subconformia.

78. V. Buxbaumii (Ten. Nap. I p. 7, tab. I) papillari-pubescentes, caulis elongatis prostratis inferne ramosis et saepe radicantibus, foliis ovatis subcordatis grosse crenato-serratis inferioribus oppositis petiolatis floralibus alternis vix diminutis, pedicellis axillaribus unifloris filiformibus erecto-patulis folio plerunque longioribus, calycis lacinii ovato-oblongis fructiferis valde divaricatis elevati nervosis capsulâ longioribus, corollâ cyaneo-venosâ calycem

superante, capsulâ ciliatâ elevatim reticulatâ longitudine duplo latiore angulo obtuso bilobâ loculis divaricatis obtuse triangularibus margine carinatis 5-8-spermis, stylo lobis capsulae sublongiore, seminibus oblongis rugosis cyathiformibus ⊖. Ic. Fl. Dan. tab. 1962. *V. Persica* Poir. Dict. 8, p. 542. Rehb. Germ. tab 78. — *V. filiformis* D.C. Fl. Fr. non Smith. — *V. Tournefortii* Gmel. — *V. agrestis* var. *Byzantina* Fl. Græc. I, tab. 8. — *V. hospita* M.K.

Hab. in cultis totius ditionis a Græciâ! Byzantio! ad Syriam!, Transcaucasiâ! Persiam borealem! (Bge!), regnum Cabulicum (Griff!), Turkestania (Lehm!).

Ar. Geogr. Europa media et australis a Daniâ et Gothiâ ad Rossiam medium, Hispaniâque boreali ad Dalmatiam.

79. *V. filiformis* (Smith Trans. Soc. Linn. 1, p. 195) parce papilloso-pubescentis ramis numerosis filiformibus elongatis prostratis radicantibus, foliis parvis orbiculatis obtuse subcrenatis subcordatis subpetiolatis floralibus subdiminutis, pedicellis tenuiter filiformibus axillaris erecto-patulis folio multoties longioribus, calycis laciniis ellipticis capsulâ brevioribus, corollâ cœruleâ calyce ampliore, capsulâ ciliatâ tenuiter reticulatâ orbiculari obcordata angulo acuto ad medium in loculos ovatos 8-10-ovulatos bilobâ, stylo sinu duplo longiore, seminibus oblongis tuberculatis cyathiformibus ⊖ vel perennans. — Ic. Rehb. Icon. tab. 644.

Hab. in cultis derelictis et humidis Asie minoris borealis (Auch. 5092!), ad Samisun (Tchih!), in Ponto Lazico ad Rhizé et Djimil (Bal!), Imeretia (Stev!), Armeniâ ad Ispir (Huet!), Caucaso centrali 4200'-6500' (M.B! Rupr!), ad Lenkoran (C.A. Mey!), in monte Avroman Kurdistaniæ (Haussk!).

Difert a *V. Buxbaumii* quâcum sæpe commutatur caulis tenuissimis, foliis 2-4 lineas tantum diam. latis, capsulâ suborbiculatâ vix transverse latiore, stylo longo. Semina triplo minora sunt et cyathiformia nec ut cl. Ledeb erronee dicit plano-convexa.

80. *V. didyma* (Ten. Nap. Prodr. 6) parce pubescens, caulis filiformibus ascendentibus et decumbentibus, foliis subrotundo-ovatis subcordatis petiolatis serrato-crenatis superioribus sensim diminutis ovatis, pedicellis axillaris unifloris folium subæquantibus vel brevioribus fructiferis reflexis, calycis laciniis late ovatis nervosis acutiusculis capsulâ sublongioribus, corollâ intense cœruleâ calycem æquante, capsulâ breviter et dense tomentellâ transverse latiore emarginata loculis inflatis dorso rotundatis 8-16-spermis, stylo sinu longiore, seminibus ovatis cymbiformibus rugulosis ⊖. Ic. Rehb. Germ. t. 77, fig. 1 et 2. — *V. polita* Fries. — Fl. Dan. tab 449.

Hab. in cultis totius ditionis a Græciâ! ad Syriam! Palestinam! Mesopotamiam et Babyloniam!, regionem Caucasicam!, Persiam borealem (Bge!) et australem (Ky. 194!), Affghaniam (Griff!), Belutschiam (Stocks!).

Ar. Geogr. Europa omnis a Scandinaviâ ad Rossiam. Africa borealis.

81. *V. agrestis* (L. Sp. 18) parce et breviter papillari-hirta,

caulibus filiformibus ascendentibus vel decumbentibus, foliis ovatis subpetiolatis crenato-dentatis superioribus sensim diminutis cuneatis, pedicellis axillaribus unifloris folio aequilongis vel brevioribus fructiferis recurvis, calycis laciis oblongis obtusis corollam pallidam vel albidastrum striatam superantibus, capsula calyce subbrevis glanduloso-hirta transverse sublatiore sinn angusto et brevi obcordata lobis obtusis carinatis tenuiter reticulatis, stylo sinu subbrevis, seminibus in loculo 4-5 ovato-globosis cymbiformibus rugulosis ⊖. Ic. Rehb. Crit. III, fig. 440. — Rehb. Germ. tab. 79, fig. 2 et 3. — *V. pulchella* Bast. et Guss.

Hab. in cultis Bithyniae (Sibth.), Anatoliae borealis (Anch. 5091!), Cretae ad Cydoniam (Haussk!), Tauriae (Stev.), regionis Caucasicae (M.B.).

Ar. Geogr. Europa omnis a Scandinaviâ ad Rossiam mediam, Songaria, Sibiria Altaica, Africa borealis.

82. **V. opaca** (Fries Nov. Ed. II, p. 3) parce pubescens, caulis filiformibus ascendebatibus et decumbentibus, foliis subrotundovalvis serrato-crenatis, pedicellis axillaribus solitariis folium subaequantibus vel brevioribus fructiferis reflexis, calycis laciis subspathulatis capsula longioribus, corolla cernuea calycem subaequante, capsula intertexte villosula transverse valde latiore acute emarginata loculis turgidis margine carinatis 4-5-spermis stylo sinu sublongiore, seminibus ovatis cyathiformibus rugulosis ⊖. Ic. Rehb. Crit. III, fig. 441. — Rehb. Germ. tab. 79, fig. I.

Hab. in Thraciâ (Griseb.), Iberiâ (Owerint), ditione Talysch (ex Hohen. Enum.).

Praeter notas indicatas a binis precedentibus staminibus fangi nec margini inferiori tubi insertis differre dicitur.

Ar. Geogr. Germania, Scandinavia, Italia borealis, Dalmatia, regio Danubialis.

83. **V. cymbalaria** (Bod. Diss. p. 3) papillis elongatis hirsuta, caulis flaccidis elongatis decumbentibus, foliis subcordato-semiorbiculatis vel cuneatis petiolatis 5-9-lobis, pedunculis folio longioribus tandem patentibus, calycis lacinis ovatis vel ellipticis obtusis basi attenuatis fructiferis patentissimis ciliatis, corolla albâ rotata calyce vix majore, capsula inflata didymâ 2-4-spermâ papillis hirsuta vel glabra, stylo longo, seminibus globosis urceolatis rugosis ⊖. Ic. Fl. Græc. tab. 9. — *V. cymbalariaefolia* Vahl Enum. I, p. 81. — *V. Paranitana* Tin. (forma capsula glabra).

Hab. in cultis regionis inferioris in Græciâ omni (Orph. Fl. exs. 718!), Archipelago!, Macedonia et Thraciâ (Griseb!). Anatolia occidentali (Bal!) et australi (Bourg!), Ponto Lazico ad Trapezunteum (Bourg!). Cypro (Ky!), Syriâ boreali (Bl!), Libano (Gaill!), Palestina (Boiss!), Tauriâ meridionali (Stev.). Forma leiocarpa et lasiocarpa mixta, haec frequentior.

Ar. Geogr. Regio mediterranea Europa ab Hispaniâ ad Dalmatiam, Africa borealis.

84. **V. cymbalariaoides** (Blanche in litt.) glabriuscula, canibus flaccidis elongatis decumbentibus, foliis trianguli-orbiculatis subcordatis vel cuneatis petiolatis 5-7-lobatis, pedunculis axillaribus folium subaequantibus tandem deflexis, calycis lacinii ovatis ciliatis saepe lobulo laterali auctis, corollâ cyathiformi lacteâ calyce breviore, capsulâ inflatâ didymâ glabrâ tetraspermâ, stylo sinu vix longiore, seminibus globosis ureolatis subrugulosis ⊖.

Hab. in cultis Syriae littoralis in consortio precedentis ad Sidonem et Tripolim (Bl!), ad Aleppo et Orfa et in monte Sindjar Mesopotamiae (Haussk!).

Facie et characteribus valde affinis *V. cymbalariae* a qua cl. Blanche et Haussknecht qui vivam observaverunt distinctissimam esse contendunt. Ego dubito.

85. **V. hederæfolia** (L. Sp. 19) pubescens, caulis elongatis decumbentibus, foliis petiolatis subcordatis suborbiculatis trilobis vel quinquelobis lobo terminali majore, pedunculis axillaribus solitariis folium subaequantibus fructiferis patenti-deflexis, calycis lacinii cordatis triangularibus fructiferi erectis, corollâ albâ vel pallide cœruleâ calyce breviore, capsulâ glabrâ globoso-quadrilobâ, loculis monospermis, stylo brevi, seminibus globosis cyathiformibus rugulosis ⊖. Ic. Rehb. Germ. tab. 77.

Hab. in cultis fere totius ditionis a Græciâ! et Thraciâ! ad Syriam (Gail!), Persiam prope Teheran (Ky. 8!).

Ar. Geogr. Europa fere omnis, Africa borealis.

+ + Calycis laciniae per paria connatae.

86. **V. Crista-Galli** (Stev. Trans. Soc. Linn. II, p. 408, tab. 31) papillosa, caulis ascendentibus vel decumbentibus flaccidis, foliis ovatis crenato-serratis inferioribus breviter petiolatis cæteris subcordatis, superioribus vix diminutis, pedicellis axillaribus 1 vel 2-3-floris filiformibus folio longioribus, calycis lacinii 4 in 2 phylla ovata cordata biloba dentataque concretis fructifero valde ampliato compresso, corolla pallide cœruleâ calyce duplo minore, capsulâ calyce subminore subciliata transverse ellipticâ subretusa loculis obtusissimis abortu monospermis, stylo brevissimo, semine magno ovato-orbiculari rugosissimo umbilicato ⊖ *Diplophyllum veronicaeforme* Lehmann. Berl. Mag. 7-4-2.

Hab. in sylvis umbrosis Caucasi orientalis 2500'-3000' (Stev., C.A. Mey!), in Grusiâ (Wilh.) in ditionis Talysh umbrosis prope Lenkoran (Hohen!), Persia bor. orientalis ad Siaret (Bge!).

Folia 8-10 lineas longa, calyx fructifer compressus 5-6 lineas longus et latus, semen 1 $\frac{1}{4}$ lineæ longum.

87. **V. Griffithii** (Bth. in p.c. Prodr. IV, p. 485) brevissime et crispule asperula nana basi ramosa, foliis oblongis subcrenatis, floribus lanceolatis acutis integerrimis pedicelli longitudine, racemis

brevibus laxis, pedicellis erecto-patulis filiformibus calyce brevioribus, calycis laciinis amplis late ovatis acutis trinerviis reticulatis ad medium per paria connatis, capsula asperula ultra medium in loculos obovatos obtusos calyce breviores partita, stylo brevissimo, semini-
bus leviusculis ♂.

Hab. in Affghaniâ prope Otipore (Griff. 642!).

Facie *V. biloba* similis differt calycis laciinis majoribus retinatis et inter se longe connatis. *V. cardiocarpa* Walp. Songarica eadem nota insignis et proxima sat tamen differre videtur pedicellis calyce longioribus et capsula breviter obcordata

Species male nota.

V. pumila (Clarke Trav. Ed. 4^a VIII, p. 223, Ic.) annua viscoso-
pubescens, caule nano subsimplici, foliis ovatis vel subrhomboides
profunde lobatis lobis inferioribus subdentatis, floribus pedicellis
patulis elongatis suffultis, corolla diametro semipollucaris laciinis
ovatis ♂.

Hab. in monte Hæmo Thraciae ad Chaligh Kavaek in viâ inter Kirkilissi et
Rutschuk.

Ex speciminiis junioribus incomplete descripta, ex affinitate *V. glauca*
et *amœna*.

TR. VII. GERARDIEÆ (Bth. in D.C. Prodr.) Corolla bilabiata lobis
planis, posticis binae saepissime interioribus. Stamina per
paria approximata, antherarum loculi distincti basi saepe
mucronati, alterum interdum deficiens. Fruetus capsula-
ris. Inflorescentia racemosa centripeta. Folia saltēm infe-
riore opposita.

STRIGA (Lour. Cochinch. p. 22).

Calyx tubulosus quinquefidus. Corolla tubus tenuis ad vel supra
medium abrupte incurvus, limbus patens obliquus bilabiatus labio
superiore integro vel bilobo inferiore trifido. Stamina inclusa fila-
mentis brevissimis, antheræ sagittatae loculo unico. Stylus apice bifidus. Capsula recta valvulis coriaceis tandem elasticæ dehiscentibus ♀.
Herbe seabra siecitatem nigricantes, parasiticae, floribus spicatis.

4. **S. Hermonthica** (Del. Fl. Eg. p. 309. tab. 34 sub *Buchneria*)
intense virens pilis e tuberculo ortis sebra, caule erecto elato stricte
ramoso, foliis linear-lanceolatis plicatis subcurvatis, spicis longis,
bracteis linear-subulatis ciliatis calycem subæquantibus, calycis
costis 5 sebris dentibus acutis, corolla purpureæ tubo calyce duplo
longiore ad medium infraucto, limbi ampli labio superiore breviore
retuso, capsula obtusa erlyce breviore ♀. *S. Hermonthica* Bth.
Seroph. p. 305.

Hab. in cultis Egypti mediae et superioris ad *Sorghum* et *Andropogonem Aleppensem* parasitica (Del! Sieb., Ehrenb! Boiss! etc.).

Ar. Geogr. Nubia, Abyssinia, Africa tropica.

LEPTORHABDOS (Schrenk. pl. Schr. fasc. I, p. 23).

Calyx quinquedentatus. Corolla tubus brevis in limbum erecto-patentem ampliatus, lobi 5 bifidi. Stamina didynama corollâ breviora, antheræ liberae loculis aequalibus nuncicis. Stylus apice breviter dilatatus obtusus. Capsula superne compressa obtusa loculicide delicens valvulis integris elastice deliscentibus. Semina in loculis bina vel abortu solitaria lateraliter infra medium affixa testâ rugosâ adhaerente. — Herba annua erecta vrgata strictæ facie *Odontitis*.

T. L. virgata (Bth. in p.c. Prodr. X, p. 510) erecta sub lente tenuiter pruinosa, caule tenuiter vrgato inferne folioso superne stricte breviterque ramoso, foliis in lacinias anguste lineares pinnatisectis, spicis terminalibus tenuibus laxissimis, pedicellis strictis calyce bracteaque linearí setaceâ subbrevioribus, calycis glandulosi dentibus lanceolatis mucronatis tubo duplo brevioribus, corollâ roseâ calyce-duplo longiore, capsula calyci æquilongâ vel longiore ♂,

Hab. in moute Elamont Persie borealis (Auch. 5005!), in prov. Khorassan inter Schahrud et Mesched (Bge!), in regno Cabulico (Griff.).

Planta 4-2-pedalis *L. micranthâ* tenuior et foliis multo angustioribus, ei ceterum floribus similissima et hujus forsitan varietas.

Ar. Geogr. Jugum Himalaicum.

TR. VIII. EUPHRASIEÆ (Bth. in p.c. Prodr.) Corolla bilabiata labio postico erecto concavo vel galeato erecto aestivatione inferiore. Stamina per paria approximata, antherarum loculi bini distincti basi saepius mucronati. Inflorescentia centripeta racemosa. Folia opposita vel alterna.

SIPHONOSTEGIA (Bth. in Hook et Arn. Beech. p. 203). — *Lesquereria* Boiss. Diagn. Ser. I, 12, p. 43. — Ser. II, 6, p. 432.

Calyx bilabiatus longe tubulosus elevatum plurinervius bilabiatus labiis ad basin in lacinias oblongo-lineares subæquales superiori tripartito inferiore bipartito. Corolla longe tubulosa bilabiata galea retusa, labio inferiore subæquilongo breviter trilobo, palatum non gibbosum sed longituinaliter plus minus biplicatum. Stamina 4 subæqualia sub galea, antheræ erectæ sagittatae loculis nuncicis. Capsula lanceolata aenta vix compressa calyce inclusa loculicide delicens. Semina numerosissima scobiformi-subiriquetra testâ laxiusculâ sed

ultra nucleus vix productâ. — Herba perennis caulis elatis duris. — Genus *Cymbaria* et *Bungea* characteribus affine, sed habitu alienum, a priori corollâ non campanulatâ, a posteriore galeâ retusâ et calyce quinquedentato, ab utrâque palato non gibboso, antheris non mucronatis et præsertim seminibus minimis numerosis distinctum, a *Schizanthus* calycis lobis subæqualib; capsulâ angustâ, seminis testâ non longiusculæ linearî-elongatâ diversum.

1. **S. Syriaca** (Boiss. et Reut. loc. cit. sub *Lesquerenoid*) breviter et patule hirsuta siccitate nigricans, caulis elongatis virgatis duris cæbre foliatis simplicibus vel parce ramosis in spicam longam laxiusculam foliosam abeuntibus, foliis subsessilibus oblongo-lanceolatis integris basi attenuatis, floralibus elliptico-linearibus calyces æquantibus vel longioribus, floribus oppositis sessilibus erectis, bracteolis calycinis linearibus, calycis tomentelli 11-nervoso-costati et profunde sulcatis labiis tubo quadruplo brevioribus, corolle purpurea tubo breviter exserto linibo parvo ♀.

Hab. in rupestribus dumosis Cilicie littoralis inter Mersina et Aladagh 2000' (Bal!), in valle Orontis inter Antiochiam et Saadieh (Boiss!), in Amano versus Tolosdagh 1000' (Ky 406!). Fl. Jul., Aug.

Caules 2-pedes et amplius longi, folia inferiora 1-2-pollicaria 2-6 lineas lata, calyx 8-9 lineas longus angustus, corolla 12-14 lineas longa, capsula valde pungens calyce inclusa 6-7 lineas longa vix sesquilinea diametro latior Semina eis *S. Chinensis* subduplo majora.

BUNGEA (C.A. Mey. Enn. p. 108).

Calyx bibracteatus breviter tubulosus laciniis 4 foliaceis longis, corolla tubulosa galeâ acuminatâ utrinque dente aucta, labio inferiore acute trifido, palatum biconvexum. Siamina sub galeâ di-dynamia, antheræ glabrae transversæ loculis æqualibus mucronatis. Ovarii placentæ bifidae. Capsula ovoidea acuminata loculicide breviter apice dehiscentis valvulis coriaceis dissepimenta diu coherentia bifida valde dilatata loculos fere dividentia medio fermentibus. Semina pauca maiuscula prope basin lateraliter affixa ascendentia oblongo-triquetra margine incrassata. — Herba cespitosa facie *Ajuga chamœpithys*. — Genus calyce bibracteato distinctum.

1. **B. trifida** (Vahl Symb. I. p. 44, sub *Rhinantho*) hirsuta caulis e rhizomate indurato pluribus erectis vel decumbentibus simplicibus brevibus foliatis cito in spicam densam oblongam abeuntibus, foliis inferioribus oppositis cæteris linearibus infimis brevibus integris cræteris in laciniis lineares tripartitis, floribus longis subsessilibus, bracteis linearibus calyci æquialongis, calycis laciniis uniner viis longe linearibus acutis rigidulis tubo multoties longioribus corollam superantibus, corollæ flavescentis extus intusque velutinæ

tubo longo, capsula longe rostrata φ . *B. trifida* C. A. Mey. l. cit. — *Bartsia trifida* Spr. Syst. II, p. 773. — *Pedicularis Orientalis supina folio trifido* Tourn. Cor. 9. — *Fedic. chamæpityos facie* Buxb. Cent. I, tab. 8.

Hab. in collibus aridis Phrygiae ad Ouchak (Bal!), Galatiae (Wiedem!), Cappadociae (Ky. 224! Bal. 432!), Armeniae Turcicæ (Anch! Bourg!) et Rossicae (Szov!), ditionis Talysch 3500'-4800' (C.A.M. Hoh!), Persiae borealis et mediae (Bge!), australis ad Persepolin (Ky 264!). Fl. Jun.

Caules cum spicis 4-7-pollicares, calycis laciniae pollicem et amplius longæ basi lineam latæ acutæ, capsula cum rostro 9 lineas longa.

EUPHRASIA (L. Gen. p. 30).

Calyx quadrifidus tubuloso-campanulatus interdum dente postico parvo auctus. Corollæ galea lata concava apice biloba, labium inferius patens trifidum lobis obtusis vel retusis, palatum non plicatum. Stamina didynama, antheræ sub galea, loculis mucronatis, spinulæ loculi inferioris staminum breviorum longiore. Capsula oblonga compressa. Semina plurima fusiformia pendula longitudinaliter sulcata. — Herbae annuae.

1. ***E. officinalis*** (L. Sp. 841) pubescens rarius glabrata, foliis parvis ovatis utrinque obtusiuscute 3-5-crenato-dentatis, foliorum superiorum acutis, galeæ lobis patentibus denticulatis, labii inferioris lobis emarginatis \odot . Species valde variabilis cajus varietates in Oriente frequentiores numero.

α . *pratensis*. — Koch Syn. — Glanduloso-pilosa, folia ovata vel oblonga, flores plerumque majusculi.

Hab. in pratis regionis inferioris Ponti prope Rhizé (Bal!), in regione alpinâ prope Djimil 7500' (Bal!), in promontorio totius Caucasi a regione montanâ ad alpinam (C. A. Mey.).

β . *micrautha*. — Crispule pubescens, folia ovato-oblonga. Flores minuti — *E. gracilis* Fries.

Hab. in Caucaso (Rupr!), Affghaniae alpibus (Griff!).

γ . *Tatarica*. — Elatior, pubescens vel glabrata, caules rigidiores, folia latiora fere orbiculata, etiam superiorum dentes obtusiusculi. Corolla majuscula — *E. Tatarica* Fisch. in Spr. Syst. II, p. 777. — Ledeb. Ic. tab. 435. — *Odontites Caucasica* C. Koch Linn. XVII. p. 289 ex specimine typico! — Forma affinis *E. officinali* var. *hir-tellei* Jord.

Hab. in pratis alpinis Tauriae (Rehm!), Ponti Lazici ad Djimil (Bal!), Armeniae Turcicæ (Ky! Huet exs. sub *E. Armenâ*), Persiae bor. in monte Kuhdaéna (Ky. 633!).

Ar. Geogr. Europa ombris borealis et media a Scandinaviâ ad Rossiam et Hispaniâ Italiamque borealibus ad regnum Danubiale, Sibiria, regio Himalaica, America borealis.

2. **E. Salisburgensis** (Funk in Hoppe Tasch. p. 190) pubescens vel glabrata, caulinis ramisque strictis rigidulis, foliis oblongis basi cuneatis utrinque 2-3-dentatis dentibus acutis, foliorum superiorum cuspidato-aristatis, galeæ lobis patentibus denticulatis, labii inferioris lobis emarginatis ♂.

Hab. in regione alpinâ, Parnasso (Heldr! Guicci!), Athone et monte Korathiæ Macedoniae (Auch. 1696! Pichler!), monte Anemas Lycaoniæ (Heldr!), Tauro Cilicico (Ky! Ball!), Anatoliâ in montibus Tokat (Wiedem!).

Planta ex Alpibus plerumque rigidior, folia saepè latiora, similis *E. pectinatae* Ten. ex Italâ australi. Utramque ob dentes foliorum superiorum cuspidatos *E. Salisburgensi* ad numero sed hæc forsitan ab *E. officinali* non sat differt.

Ar. Geogr. Alpes, Pyrenæi, Italia, Dalmatia, regio Danubialis.

3. **E. minima** (Schl. in D. C. Fl. Fr. III, p. 473) nana pusilla, foliis minimis ovalis paucidentatis, superiorum dentibus mucronatis, floribus minimis cœruleis vel cupreis, galeæ lobis connivētibus, labii inferioris lobis emarginatis ♂. Ic. Rehb. Germ. t. III, fig. 8. — *E. pygmaea* C. Koch Linn. XXIII, p. 686!

Hab. in alpibus Thracie (Frivl), Ponti (C. Koch!), Tauri Cilicici (Ky. 286!), Caucasi orientalis 6000'-9000' (Rupr!), Persiæ borealis (Ky! Buhse!).

Ar. Geogr. Alpes, Pyrenæi, montes Hispaniæ, Carpathi.

EUFRAGIA (Griseb. Spicil. II, p. 43).

Calyx tubulosus quadrisfidus. Corollæ tubus cylindricus, galea concava lobis non replicatis, palatum convexum, labium inferius patens trilobum. Stamina didynama, antherarum loculi aequales mucronati. Capsula oblonga vel lanceolata compressiuscula. Semina numerosa minuta non sulcata. — Herbæ annuae viscosæ.

1. **E. latifolia** (L. Sp. 841 sub *Euphrasia*) humilis glandulos-viscosa, foliis parvis subsessilibus breviter ovatis inferioribus crenatis superioribus subpalmatilobis, spicis ovatis demum elongatis et inferne interrupptis, calycis dentibus oblongis tubo brevioribus, corollæ parvæ purpureæ vel albo-suffusæ rarius albæ labio galeam subæquante, antheris glabris ♂. — *E. latifolia* Griseb. Spic. loc. cit. Rehb. Germ. tab. 404. — *Bartsia latifolia* Fl. Græc. VI, p. 69, tab. 386. — *Trixago purpurea* Stev.

Hab. in collibus et pratis regionis maritimæ totius Græciae! et insularum!, Macedoniae et Thraciae (Griseb! Janka!), Anatoliæ maritimæ et Ponti!, Syrie littoralis et interioris ad Aleppo (Kyt!) et in Cælesyriâ (Boiss!), Transcaucasie (Ledeb.), prov. Talysh ad Caspium (Hohen!), prov. Ghilan Persiæ (Buhse!, Bge!).

β. flaviflora. — Corolla sulphurea.

Hab. in rupestribus graminosis Persiæ australis prope Gere (Ky. 43!). Nomen specificum ineptum sed antiquius servavi.

Ar. Geogr. Gallia occidentalis, regio mediterranea totius Europæ et Africæ borealis.

2. **E. viscosa** (L. Sp. 839 sub *Bartsia*) elatior glanduloso-viscosa, foliis oblongis crenato-dentatis, spicis elongatis laxis, calycis lobis linearibus tubo vix brevioribus, corollæ flavæ labio inferiore galeâ longiore, antheris pilosis ⊖ *E. viscosa* Bth. in D.C. Prodr. 10, p. 543. — Rehb. Germ. tab. 103. — *Trixago viscosa* Rehb. exs. 360. — *Bhinanthus maxima* Lam. Dict. et Desf. Atl. non Willd.

Hab. in pratis in primis maritimis totius Graeciae (Sprun! Heldr! Orph!), Macedoniae et Thracie (Griseb.), Anatoliae occidentalis et australis (Auch!, Ky!), Syriae littoralis (Bl!), Egypti inferioris (Ehrenb! Wiest!), Transcaucasiae ad mare Nigrum et ad Caspium (Ledeb!), prov. Ghilan Persiae (Buhse!).

Ar. Geogr. Gallia occidentalis, regio mediterranea Europæ et Africæ borealis, Madera, insulae Canarienses.

TRIXAGO (Stev. Mém. Mosq. Vol. VI, p. 4).

Calyx ventricoso-campanulatus breviter quadrifidus. Corollæ tubus cylindricus, palatum bigibbosum, galea concava marginibus non replicatis. labiem inferius pateus trilobum. Stamina didynama, antherarum loculi aequales mucronati. Capsula ovata turgida placentis crassis bifidis. Semina minuta numerosissima longitudinaliter et transversiter costata. Herba annua.

1. **T. Apula** (Stev. loc. cit. p. 4) superne glanduloso-pilosa, foliis a basi sessili vel subamplexicauli oblongo-lanceolatis vel linearibus obtuse serrato-dentatis, spicâ densâ brevi tandem oblongâ, calycis dentibus ovatis tubo 4-5-plo brevioribus, corollæ annuae purpureæ vel versicoloris labio inferiori galeâ longiore, antheris pilosulis, capsula ovato-globosâ ⊖. *Bartsia Trixago* L. Sp. I, p. 602. — Rehb. Germ. p. 103. — *Trixago carnea* Griseb. Spic. p. 12. — *Bartsia versicolor* Pers. Syn. II, p. 151. — *Bartsia bicolor* D.C. Ic. rar. tab. 10.

β. *flaviflora*. — Corolla lutea. — *Bartsia Trixago* Fl. Graec. tab. 585. — *Trixago rhinanthina* Link. — *Rhinanthus maximus* Willd. III, p. 189.

Hab. cum varietate minus frequenti in pratis praesertim maritimis totius fere ditionis, Graeciæ!, Macedoniae et Thraciæ!, Anatoliæ littoralis, Syriae littoralis (Bl!), Palestinæ (Boiss!), Mesopotamiæ (Ky! Haussk!), Transcaucasiæ ad Euxinum et Caspium (Ledeb!), Lenkoran (C.A. Mey!).

Ar. Geogr. Gallia occidentalis, regio mediterranea totius Europæ et Africa borealis, P. bonæ spei.

ODONTITES (Haller Pers. Syn. II, p. 450).

Calyx quadrifidus tubulosus vel campanulatus. Corollæ galea concava integra vel retusa lateribus non replicatis, labium inferius tri-

tidum lobis integris, palatum non plicatum. Stamina didynama sub galea, antherarum loculi aequaliter mucronati rarius mutici. Capsula oblonga compressa. Semina plurima pendula longitudinaliter suteata. — Herba annua vel perennies suffrutescentes foliis oppositis,

§ 1. Annuæ.

* Antheræ glaberrime.

1. O. Aucheri (Boiss. Diagn. Ser. I, 4, p. 75) annua adpresso seabripto-canescens tuberulata glandulisque sessilibus plus minus viscida, caulis tenuibus rigidis, ramis filiformibus erecto-patulis, foliis strictis convolutis tenuiter linearibus, floralibus calyces subæquantibus, calycis tubuloso-campanulati dentibus lanceolatis strictis tubo subæquilonigis, corollæ rubella tubo vix exerto limbo ciliato galea retusa, antheris glabris galea inclusis muticis, capsula hirta calyxi subæquilonigra truncata retusa apiculata ♂. *Euph. rigidifolia* et *E. leianthera* Fenzl. MSS. — *Amblygauthera chrysanthia* Fisch. in Stev. exs.

Hab in siccis montanis et stepnis alpiniis, Phrygia prope Ouchak (Bal!), circa Tokat (Wied!), Tauris Cilicieus (Ky! Bal!), Cappadocia (Balt!). Cataonia in montibus Begdaghi et Beryt-dagh 7-8000' (Haussk!), Libanus (Bl!), Armenia (Ky. 1600!), ad Baibut (Bourg!), in prov. Musch (Ky. 3200!), Armenia Rossica (Szov!). Kurdistanie Persie mons Awiheng 7000' (Haussk!), mons Totschal Persie prope Teheran (Ky. 577!).

In excelsioribus 3-5-pollicaris spicis abbreviatis, in demissioribus interdum pedalis, folia subulata brevia, corolla 4-5 lineas longa.

2. O. lutea (L. Sp. 842 sub *Euphrasiid*) annua minute et crissipule pubescens, caule rigido ramis brevibus erecto-patulis, foliis lanceolato-linearibus vel linearibus integris vel inferioribus remete dentatis siæpe deflexis, floralibus flore subbrevioribus, spicis unilate-ralibus tandem elongatis, calyce campanulato adpresso pubescente ad tertiam partem in dentes triangulares fiso, corollæ luteæ pubescentis tubo e calyce non exerto labiis divaricatis aequilongis, antheris exsertis glabris liberis luteis mucronatis, capsula ovato-elliptica obtusissimâ pubescente ♂. *O. lutea* Rehb. exc. p. 339. — Jacq. Austr. tab. 398. — Rehb. Germ. tab. 108. — *Euph. linifolia* L. var. z.

Hab in collibus aridis, Cilicia in montibus prope Mersina (Bal!), Tauria et Caucasi latus boreale (M.B., C.A. Mey.). Fl. Aug., Sept.

β. hispidula. — Bractæ et calyces patule et longius hispiduli.

Hab. in sylvaticis Beit Kachbo Libani (Gail!).

Ar. Geogr. Hispania orientalis, Gallia, Germania, Italia, regio Danubialis, Dalmatia, Rossia australis.

** Antheræ hirtæ vel lanatae.

3. O. glutinosa (M.B. Taur. Cauc. II, p. 70 sub *Euphrasiid*) annua breviter glutinoso-pubescentis, caule rigido simplici vel stricte

ramoso ramis lateralibus spicâ terminali brevioribus, foliis strictis longe linearibus convolutis integris, floralibus abbreviatis calyce 1 $\frac{1}{2}$ -2-plo longioribus, spicis angustis strictis demum laxiusculis, calycis subsessilis tubulosi glandulosi dentibus strictis linearibus tubo sublongioribus, corollæ flavae tubo glanduloso subincurvo calyce 2-2 $\frac{1}{2}$ -plo longiore, limbo margine glabro, labii inferioris sublongioris lobis obtusissimis. antheris transversis inclusis villo connexis mucronatis, capsulâ calyce subbreviore apice truncatâ apiculatâ ⊙ *O. glutinosa* Bth. in D.C. Prod. — *O. ixodes* Boiss. et Heldr. Diagn. Ser. I, 7, p. 45.

Hab. in siccis regionis alpinæ, Taygetus (Heldr!), Olympus Bithynus (Clem! Pichler!), montes Davros et Boudroun Pisidiæ (Heldr!), Alidagh Capadociæ 5000' et Taurus Cilicicus (Bal!), mons Jildisdagh Galatiæ (Wied!), Pontus Lazicus ad Djimil 6000' (Bal!). Armenia prope Erzerum (Calv!), Tauriæ mons Tchatyrdagh (M.B.), Iberia Caucasicæ (M.B!).

Planta 4-8-pollicaris, folia 1-4 $\frac{1}{2}$ -pollicem longa, calyx 3 lineas longus, corolla 6-7 lineas longa. Contuli specimina herb. Biebersteiniani!

Ar. Geogr. Serbia.

4. ***O. serotina*** (Lam. Fl. Fr. II, p. 350 sub *Euphrasiâ*) pubescens annua, caule procero ramosissimo, foliis lanceolatis acuminatis basi attenuatis remotiuscule serratis, bracteis oblongo-linearibus flore brevioribus integris, spicis demum tenuibus elongatis laxis, calycis campanulati dentibus lanceolatis tubo æquilongis, corollæ rubræ pubescentis tubo non exerto labiis divergentibus inferiore breviore, antheris subexsertis transversis apiculatis basi barbatulis, capsulâ oblongâ obtusâ apiculatâ calyci subæquilonlâ ⊙. Ic. Rehb. Germ. tab. 106. — *Euphrasia Odontites* L. Sp. 841 et *Od. rubra* Pers. Syn. II, p. 150 ex parte. — *O. divergens* Jord.

Hab. in cultis et humidis totius ditionis a Graecia! et Macedonia! ad Persiam hor. (Ky. 633!), Kurdistaniam Persicam (Haussk!).

Ar. Geogr. Europa media et australis a Belgio et Germania ad Rossiam mediam, Sibiria.

§ 2. Suffruticosæ.

5. ***O. Linkii*** (Heldr. et Sart. in Boiss. Diagn. Ser. II, 3, p. 177) basi suffrutescens glabra siccitate nigrescens, caudice ramos abbreviatos dense foliosos caulesque floriferos elatos ramosos edente, ramis floralibus virgatis arcuatis erectis elongatis in spicas laxiusculas secundas abeuntibus, foliis anguste linearibus acutis uninerviis margine scabridis, floralibus calyce subæquilongis. calycis glabri tubulosi dentibus triangularibus acutis tubo quadruplo brevioribus, corollæ luteæ glabri tubo calyce sublongiore labiis divergentibus superiore recto obtuso longiore, antheris subexsertis mucronatis ad insertionem parce lanatis, capsulâ glaberrimâ oblongo-lineari truncato-subretusa calyce sublongiore 3. *G. rigidifolia* Link MSS. non Biv.

Hab. in regione inferiore et abietinā Parnassi 2500'-4000' (Sart! Heldr. Herb. Norm 67!). Fl. Aug., Sept.

Folia ramorum steriliū hornotiuorum 1½-pollīcīaria 2 lineas lata, caules floriferi 1-2-pedales rāni sīpe semipēdālibus, folia ramealīa brevia. Facies *O. purpurea* corollā purpureā, antheris submuticis etc. diversae et *O. Bocconi*. Guss. foliis latioribus, floribus confertis, bracteis ciliatis apice caudato-squarrosis, antheris glabris diversissimā.

6. O. Cypria. basi suffrutescens glabra siccitate nigricans, caudice ramosissimo ramos steriles breves breviter foliosos caulesque floriferos elatos virgalos parce ramosos edente, foliis etiam ramorum steriliū brevibus carnosulis anguste linearibus obtusis, spicis terminalibus brevibus densiusculis secundis, bracteis oblongo-lanceolatis calyce subbrevioribus, calycis campanulati glabri dentibus triangularibus tubo duplo brevioribus, corolla lutescentis glabrae tubo inclusō labiis divergentibus subaequilongis, antheris subexsertis apiculatis margine ciliato-lanatis, capsula puberula emarginatā calyce sublongiore ♂. *O. Bocconi* Kotschy Cypr. p. 292 non Guss.

Hab. in insulā Cypro prope Prodromo, Evrico, ad prom. Capo Gallo et ad radices montis Troodos (Kyrl.). Fl. Aut.

Folia ramorum steriliū 4-5 lineas tantum longa, caules sesquipedales, spicas etiam fructiferae pollīcīares. Ab *O. Bocconi* foliis, antheris hirsutis longe distat; magis affinis *O. Linkii* foliis et spicis elongatis, dentium calycis proportione, corolla labiis valde inaequalibus, capsula glabra, etc., diverse.

7. O. Cretica. basi suffrutescens glaberrima siccitate nigrescens, caudice ramos abbreviatos dense foliosos floriferosque sublongiores edente, foliis anguste linearī-lanceolatis acuminatis uninerviis margine laevibus, bracteis flores non superantibus, spicis solitariis vel 4-5 breviter paniculatis brevibus, calycis glabri dentibus margine scabridiis triangularibus tubo triplo brevioribus, antheris vix exsertis glabris brevissime mucronatis, capsula glabra ellipticā obtusa mucronatā calyce sublongiore ♀. *E. frutescens* Sieb. exs.

Hab. in Cretā ad Omalo districtū Sphakia (Sieb!). Vid. in herb. Kew.

Semipedalis, affinis *O. Linkii* caulinis elatis, spicis laxis tenuibus 3-4-pollīcīribus, calycis dentibus brevioribus, antheris sublanatis distinctae. *O. Bocconi* ad quam a el. Beuth. adducitur magis differt bracteis apice recurvis flores superantibus, capsulis puberulis.

RHYNCHOCORYS (Griseb. Spic. II, p. 42. — *Elephas* Guss.).

Calyx a latere compressus bilabiatus labio superiore bidentato vel retuso inferiore bifido. Corolla tubus brevissimus cuneato-campanulatus, galea in rostrum longum tenue proboscidiforme marginibus conduplicatis stylum vaginans a basi vel a medio abrupte attenuata, labium inferius latum patens trilobatum. Stamina didynama filamentis brevibus, antheris glabris obliquis, loculis aequalibus muticis. Capsula orbicularia plus minus compressa valvulis mediante dissepimento

cohaerentibus. Semina numero definita striata vel costata. — Herbæ oppositifoliæ perennes vel annuæ.

1. R. Elephas (L. Sp. 840 sub *Rhinantho*) glanduloso-pilosa vel glabra, radice dentato-squamosâ, caulinis elatis erectis vel ascendentibus flaccidis laxe ramosis, foliis brevissime petiolatis ovatis obtusis obtuse dentato-crenatis basi rotundatis vel subcordatis, floralibus sensim diminutis ellipticis acutius dentatis, floribus in axillis solitariis oppositis secundis breviter pedunculatis racemos longos laxos formantibus, calycis labiis inæqualibus inferiore longiore profundius bifido, corolla flavæ galeæ linearî infra medium abrupte inflexâ et superne gibbâ in rostrum subulatum rectum vel ascendens supra basin dentibus binis dependentibus subtus instructum apice non appendiculatum abeunti, labio inferiori amplio galeam aequante vel superante orbiculato obtuse trilobo, capsulâ globosâ hirtâ calyce breviore, seminibus 15-20 subglobosis lineato-striatis ♀. *Rhynchocorys Elephas* Griseb. Spic. II, p. 12. — *Elephas Columnæ* Guss. Sic. Prodr. II, p. 133. — *Elephas Campoclaresium* Col. Eephr. I, t. b. 188. — *E. recta* Don.

Hab. in sylvaticis, pratis humidis umbrosis regionis montanæ et subalpinæ, Thracia (Friv!), Pontus Lazicus in monte Kalonoros prope Rhizé 4500' et circâ Djimil 6000' (Bal!), Imaretia et Guria (Ledebr.), Georgia Caucasica et ditio Talyesch (Hohen!), prov. Ghilan Persiae (Auch. 5085! Buhse! Bode!), inter segetes ad radices montis Demavend (Ky. 722!), in monte Elbrus prope Passgala (Ky. 422!), prope Siaret (Bge!).

β. stricta. — Villoso, caulis simplicior abbreviatus minus flaccidus. Forma ex locis editioribus siccioribus. *R. strictus* C. Koch Linn. XIII, p. 684.

Hab. in Ponto Lazico supra Djimil 6000'-9000' (C. Koch! Bal!), Guriâ (Szov!).

Planta indumento, caule debili vel stricto, foliis, floribus et calycis magnitudine variabilis et in varietates agre dividenda.

Ar. Geogr. Sicilia, Italia meridionalis.

2. R. Orientalis (L. Sp. 840 sub *Rhinantho*) pubescens, radice brevi fibrosâ, caulinis erectis vel ascendentibus simplicibus vel opposite ramosis, foliis a basi subsessili subcordatâ triangulari-ovatis acutiusculis obtusinscule crenato-dentatis, floralibus sensim diminutis, floribus in axillis solitariis oppositiis secundis longiusculi pedunculatis laxe et longe racemosis, calycis labio superiore sublatiore lobis obtusis, corollæ flavæ galeæ a basi utrinque lobulo auctâ in rostrum subulatum elongatum valde incurvum edentulum apice in laminam parvam orbiculatam expansum abeunti, labio inferiori amplio ovalo-orbiculato obtuse trilobo, capsulâ villosâ orbiculari subcompressâ, seminibus 10-12 oblongis subineurvis profunde sulcatis et corticoso-costatis ♂. *Rhynchocorys Orientalis* Bth. in D.C. Prodr. — Rehb. Icon. tab. 730. — *Elephas* Tourn. Voy. tab. 299. — *Elephas incurva* Don.

Hab. in pratis montosis Caucasus (M.B., C.A. M., Owerint!), Caucasus orientalis 6-7000' (Rupr!), Georgiae Caucasicæ (Hohen!), Guriae (Nordm.), Armeniae Rossicae (Szov!).

Radice annua, rostri formâ, seminibus paucioribus et majoribus a praecedenti distinctissima.

RHINANTHUS (L. Gen. 304 ex parte. — *Alectorolophus* M.B.
Taur. Cauc. II, p. 68).

Calyx ventricoso-compressus quadridentalis. Corolla tubus ex indecens, galea ovata obtusa compressa antice sub apice bidentata, labium inferius brevius patens trilobum. Stamina didynama, anthæcae transversæ pilosæ obtuse bilobæ muticæ. Capsula compressa orbicularis valvulis membranaceis medio septiferis, placentis tenuibus. Semina pauca magna suborbiculata compressa alâ circumdata. — Herbe annue.

1. **R. minor** (Ehrh. Beitr. 6, p. 144) glabriusculus erectus, foliis sessilibus asperis oblongo-lanceolatis crenato-dentatis, spicis ovatis densis tandem elongatis laxis, bracteis concoloribus viridibus in dentes triangulares mucronatos incisis, corollæ flavæ tubo recto, galeæ dentibus ovatis aquilonis et latiſ \odot . Rehb. Pl. Crit. fig. 974. — Rehb. Germ. tab. 117. — *Alectorolophus parviflorus* Wallr. — *Rh. Crista galli* L. ex parte.

Hab. in pratis subalpinis Igasdaghi Anatoliae (Wiedem!) vallis Djimil Ponti Lazici 6000' (Balt!), Transcaucasia (Auch. 5101'!), Persia hor. ad Khoi prov. Aderbidjan (Szov!). Tauria (M.B.), Caucasus orientalis (C.A. Mey!), Caucasus orientalis 6000'-8000' (Rupr!).

Ar. Geogr. Europa borealis et media a Lapponiâ ad Lusitaniam, Hispaniam et Italiam medias, Dalmatiâ, Sibiria, America borealis.

2. **R. major** (Ehrh. Beitr. 6, 144) glabriusculus vel hirsutus, foliis sessilibus asperis oblongo-lanceolatis, spicis ovatis densis tandem elongatis laxis, bracteis decoloribus pallidis in dentes triangulares mucronatos incisis, corollæ flavæ tubo subcurvato, galeæ dentibus ovatis \odot , i.e. Rehb. Crit. fig. 975. — Rehb. Germ. tab. 118. — *Al. grandiflorus* Wallr. — *Rh. Cristagalli* L. ex parte.

Hab. in pratis subalpinis, Armenia prope Baibut (Bourgt!) et in monte Techlagh (Calv!), Cappadocia (Ky. 231!), Caucasus omnis (C.A. Mey.).

β . *Parnassicus*. — Minor, pubescens, facies *R. Alpini* sed bracteæ latiores et dentes galeæ abbreviati *R. minoris*. — *R. pubescens* Boiss. et Heldr. in Orph. Fl. Gr. exs.

Hab. in regione superiori montis Parmassi 6000'-7000' (Heldr! Orph. Fl. exs. 412!).

Ar. Geogr. Europa omnis presertim borealis et media a Scandinavia ad Lusitaniam et Hispaniam boreales, Italianamque medium, Sibiria.

3. R. Alpinus (Baumg. Enum. Transylv. II, p. 194) humilis, hirtus, foliis sessilibus asperis oblongo-lanceolatis crenato-dentatis, spicis brevibus tandem basi interruptis, bracteis decoloribus pallidis in dentes triangulari-lanceolatos incisis, corollæ flavæ tubo ascendente, galeæ dentibus triangularibus acutiusculis ♂. Rehb. Germ. tab. 119, fig. 2. — *R. alpestris* Wahlenb. — *R. pulcher* Schum.

Hab. in Rumeliâ orientali (Frivl.).

A quibusdam auctoribus cum *R. angustifolio* confusus a quo tamen forma et brevitate foliorum bractearumque dentibus non longe aristatis differt.

Ar. Geogr. Carpathi, Sudeti, Alpes Transylvaniae, Tyrolis, Banatûs, Bosniæ.

MELAMPYRUM (L. Gen. 305).

Calyx tubuloso-campanulatus quadrifidus. Corollæ tubus cylindricus superne latior, galea brevis compressa obtusa marginibus anguste replicatis, labium inferius inflato-bigibbosum tridentatum. Anthæ hirtellæ subverticales loculis æqualibus mucronatis. Capsula compressa ovata obliqua loculis 1-2-spermis. Semina lœvia strophiolata. — Herbæ annuæ.

* Flores spicati.

1. M. cristatum (L. Sp. 842) superne pubescens, caule erecto rigidulo divaricatum ramoso, foliis sessilibus lanceolato-linearibus patulis vel deflexis, spicæ densâ tetragonâ angulis cristæformibus, bracteis tetrastiche imbricatis cordato-orbiculatis complicatis argute duplicato-serratis acumine recurvo, calycis dentibus lanceolatis tubo brevioribus, corollæ flavidæ tubo brevi ♂. Ic. Fl. Dan. tab. 1104. — Rehb. Germ. tab. 116.

Hab. in pratis et sylvis siccis Byzantii (Sibth.), Tauriæ (ex M.B.). E ditione nondum vidi.

Ar. Geogr. Europa omnis præsertim borealis et media, Sibiria Uralensis et Altaica.

2. M. arvense (L. Spec. 842) asperulum, caule erecto rigidulo patule ramoso, foliis sessilibus lanceolatis vel linearibus, spicæ laxe cylindracea, bracteis lutescentibus ovato-oblongis longe et pectinatum subulato-dentatis apice subintegris acuminatis subtus saepius bifarium punctatis, calycis pubescenti-scabri dentibus a basi latiore longe setaceis inæqualibus tubo suo longioribus tubum corollæ æquantibus, corollæ purpureæ ad faucem flavo-maculatâ, capsula cuneatâ obovatâ mucronatâ ♂. Ic. Fl. Dan. tab. 911. — Rehb. Germ. tab. 115, fig. I.

Hab. in cultis argillosis, Thracia (Friv.), Byzantium (Coumany!), Anatolia borealis (Wiedem!), Armenia Turcica (Calv! Bourg!), Pontus Lazicus ad Djimil (Bali), Tauria (M.B.), Caucasus et Transcaucasia (C.A. M., Hohen.).

β. elatius. — Elatius, folia angustiora, spicæ laxiores, corollæ la-

bium superius densius barbatum. — *M. elatius* Reut. in Bourg. exs. 1862.

Hab. in dumosis supra Trapezuntem (Bourg! Ky. Suppl. 712!), Anatoliâ bor. (Wiedem!).

Ar. Geogr. Europa borealis et media a Sueciâ meridionali ad Rossiam medianam et Hispaniâ boreali atque Italiâ ad Dalmatiam.

3. **M. barbatum** (W. K. Hung. I, p. 89, tab. 86) àdpresso asperulo-hirtum, caule erecto rigidulo patule ramoso, foliis sessilibus lanceolatis vel anguste linearibus, spicâ laxe cylindricâ, bracteis a basi ovatâ lanceolatis in laciniis subulatas ciliatas parte inferiori pectinatim dentatis, calycis tubo glabriusculo dentibus setaceis ciliatis parum breviore, corollæ luteæ tubo calycinis laciniis multo longiore, capsulâ stipitatâ obovatâ mucronatâ ♂. Ic. Rehb. Germ. tab. 115, fig. 2. — *M. ciliatum* Boiss. et Heldr. Diagn. Ser. II, 3, p. 176 (forma foliis subangustioribus et spicâ laxiore).

Hab. in Thessaliâ (Auch. 1672!) inter frutices ad radices Olympi Thessali (Heldr.).

Ar. Geogr. Italia, Austria, regio Danubialis, Dalmatia.

4. **M. Caucasicum** (Bge pl. Abich. p. 16) breviter hirtum, caule erecto rigidulo patule ramoso, foliis sessilibus lanceolatis, floribus viridibus ovatis in laciniis lanceolatas acuminatas pectinato-incisis parte superiori subintegrâ, calycis dentibus tubum glabriuscum aequalibus corollæ tubo multo brevioribus, capsulâ subsessili elliptico-oblongâ subfalcata oblique mucronatâ ♂. *M. barbatum* Led. Fl. Ross. III, p. 305 non W. K. — *M. arvense* var. *chlorostachys* Hohen. Talyisch En. p. 81!

Hab. in Guriâ Caucasicâ (Nordm!), Armeniâ Rossicâ (Abich), ditione Rad-scha et Daghestaniâ prope Gimri 600'-1200' (Rupr!), Armeniâ Rossicâ (Szov!), Georgiâ Caucasicâ et ditione Talyisch (Hon!).

Facies *M. barbati* sed bractearum laciniæ latiores et capsulæ formâ præsertim ab eo distinctum videtur.

* * Flores in racemos laxos unilaterales dispositi.

5. **M. nemorosum** (L. Sp. 843) pubescens, caule erecto patule ramoso, foliis petiolatis ovatis et ovato-lanceolatis, floribus in axillis bractearum solitariis oppositis secundis racemos laxos formantibus, bracteis petiolatis ovato-triangularibus subcordatis setaceo-pectinatis summis violaceis, calycis villosi dentibus a basi lanceolatâ setaceis tubo sublongioribus, corollæ flavæ tubo calyce longiore, capsulâ oblongâ obliquâ oblique mucronatâ ♂. Ic. Fl. Dan. tab. 305. — Rehb. Germ. tab. 114.

Hab. in sylvaticis, Caucasus et Iberia (M.B.), Georgia Caucasicâ (Hohent.). E ditione nondum vidi.

Ar. Geogr. Europa borealis et media a Daniâ et Gothiâ ad Rossiam medium et australiem et a Galliâ et Italiâ mediâ ad Dalmatiam, Sibiria Uralensis.

6. *M. pratense* (L. Sp. 843) glabriusculum, caule erecto ramis patulis elongatis, foliis breviter petiolatis a basi rotundatâ lanceolatis, floribus solitariis axillaribus oppositis unilaterialibus racemos laxissimos foliatos formantibus, bracteis inferioribus foliis similibus superioribus basi dentibus paucis subulatis instructis, calycis glabri dentibus linearis-subulatis tubo sublongioribus quartam tubi corollini partem aequantibus, capsulâ breviter lanceolatâ acutâ calyce longiore ⊖. Ic. Engl. bot. tab. 113. — Rchb. Germ. tab. 113, fig. 11.

Hab. in pratis, ericetis et sylvaticis, prov. Caucasicae (Guldenst. ex Ledeb.).

Ar. Geogr. Gallia borealis et media a Scandinavia ad Rossiam borealem et ab Hispania centrali ad regionem Daunialem, Sibiria Uralensis et Altaica.

7. *M. Heracleoticum* (Boiss. et Orph.) pubescenti-asperulum caule erecto longe et patule ramoso, foliis basi attenuatis subsessilibus lanceolatis, floribus solitariis axillaribus oppositis unilaterialibus racemos laxos foliatos formantibus, bracteis petiolulatis supra basin minutam rhombream longe pectinato-setaceam longe et anguste linearibus integerrimis, calycis dentibus longe setaceis tubo longioribus tubum corolle eitrinae subequantibus, capsulâ breviter triangulari-lanceolatâ acutâ calyce longiore ⊖.

Hab. inter pterides in Macedonia supra Bucovo prope Bitolia (Heracleam) (Orph!) Fl. Jun., Julio.

Pedalis, facies *M. pratensis* a quo dentibus ciliiformibus longissimis bractearum et calycis differt, bractae inferiores pollicem et amplius longe basi vix sesquilineam late cuspide longâ linea $\frac{2}{3}$ latâ. Capsula *M. pratensis*. Species insignis ex paucis speciminibus nota.

8. *M. stenophyllum*, totum sub lente asperulum, caule erecto patule ramoso, foliis sessilibus a basi angustissime linearibus, floribus solitariis axillaribus oppositis paucis racemum breve in capitatum foliatum formantibus, bracteis in laevis paucas setaceas pinnatisectis, calycis asperuti dentibus a basi lanceolatâ setaceis tubo suo subaequilongis corolle tubum dimidium aequantibus, capsulâ.... ♀. *M. sylvaticum* C. Koch Linn. XXIII. p. 679 non L.

Hab. in sylvaticis vallis Fortuna Ponti Lazici (C. Koch!).

Itenosissimum semipedale vel elatius, a *M. sylvatico* calycis forma alienum, ut *M. pratense* magis accedens sed foliis basi non dilatatis lineam dimidiam vix longis, calycis dentibus setaceis tubi corollini proportione longioribus diversum. Ex unico specimine florifero notum et ulterius observandum.

9. *M. sylvaticum* (L. Sp. 843) glabriusculum, caule erecto pumilo patule et parce ramoso, foliis breviter petiolatis a basi eu-neatâ lanceolatis, floribus solitariis axillaribus oppositis unilaterialibus racemos laxissimos foliatos formantibus, bracteis longe lanceolatis, patentibus corolle parvae late tubo longioribus, capsula ovato-oblongâ aenniatâ rectâ calyce longiore ⊖. Ic. Engl. bot. tab. 804 — Rchb. Germ. tab. 113.

Hab. in sylvaticis in Caucaſi monte Beschtaw (Guldenst.). Editione non vidi.

Ar. Geogr. Europa praesertim borealis et media a Scandinaviâ ad Rossiam et Hispaniâ boreali ad Italiam et regionem Danubialem, Sibiria.

PEDICULARIS (L. Gen. 746).

Calyx tubulosus vel inflatus antice et interdum postice fisis varius 2-5-dentatus. Corollæ tubus cylindraceus, galea rostrata vel erosa, labium inferius trilidum. Stamina didynama sub galeâ ascendentia, antherarum transversarum loculi paralleli basi aequales vel aristati. Capsula compressa obliqua valvis membranaceis ad medium septiferis. Semina numerosa reticulata, foveolata vel multicostata. — Herbae perennes foliis alternis vel verticillatis.

Specierum Orientalium dispositio.

§ 1. **Erostres.** — Galea non rostrata nec vel obsolete dentata.

* Folia caulina verticillata.

P. Caucasia, pycnantha, Cadmea, verticillata, Pontica

** Folia caulina sparsa.

P. orthantha, Olympica, atropurpurea, Wilhelmsiana, condensata.

§ 2. **Rostratæ.** — Galea in rostrum attenuata edentula.

* Folia caulina verticillata.

P. subrostrata, crassirostris.

** Folia caulina sparsa.

P. Nordmanniana, Elephas, labellata.

§ 3. **Bicuspidatæ.** — Galea incurva brevissime vel vix rostrata apice oblique truncato bidentata.

P. leucodon, Graeca, Friderici-Angusti, brachyodonta, comosa.

§ 1. **Erostres.**

* Folia caulina verticillata.

1. **P. Caucasia** (M.B. Taur. Cauc. II, p. 72) radicis fibris crassis longis attenuatis, caulis pumilis simplicibus subbifariam

puberulis, foliis oblongo-linearibus pinnatisectis segmentis breviter oblongis incisis vel pinnatifidis, foliis caulinis verticillatis, spicâ ovatâ vel interruptâ oblongâ, bracteis crenulatis calycibusque hirsutis hisce cylindricis dentibus breviter lanceolatis acutis crenulatis, corollæ purpureæ tubo cylindrico ad medium infracto calyce sublongiore, galeâ linearî antice rectilineâ apice subattenuatâ obtusissimâ labium inferius subæquante, filamentis duobus plus minus barbatis antheris contiguis extus prominulî, capsulâ oblongo-lanceolatâ acutâ obliquâ calyce duplo longiore ♀. Steven Mon. tab. 8. *P. Persica* Willd. herb. — *P. nudicaulis* C. Koch Linn. XVII, p. 289.

Hab. in regione alpinâ, Armenia Turcica in montibus supra Gumuschkhané (Bourg!), Somchetia (C. Koch), Caucasus australis 7000'-9000' (M.B.).

β. albiflora. — Humilior, spica compacta, corolla albida. — *P. Armena* Boiss. et Huet Diagn. Ser. II, p. 175.

Hab. in regione alpinâ Ponti Lazici supra Djimil (Bal!), montis Fecelem prope Baikut (Bourg!) et Techdag supra Erzerum 8000' (Huet!), Caucaso in ditionibus Salataviâ et Tuschetiâ 8500'-10000' (Owerin! Ruprecht!).

γ. glabrescens. — Flores parvi, corolla albida galeâ apice purpureâ, calyx glabrescens intense nigro-lineatus dentibus parvis subintegris. Forma a var. *β* vix distincta.

Hab. in editissimis Caucasi Orientalis in ditionibus Tindal, Bogos, Tuschetiâ, Salataviâ (Rupr!), Persia boreali in montibus Ssahend, Demavend (Buhse!), Elbrus et Totschal (Ky. 166!).

δ. apiculata. — Typo præter galeam apiculo oblique terminatam similis.

Hab. in monte Alagös Armeniæ Rossicæ (Buhse!), in Adschariâ (Owerin!), in monte Areklidruk (Szov!), in Alpibus prov. Ghilan (Hablitzl ex herb. Pallas!).

Species facie *P. verticillatae* affinis, ab ea calycis cylindrici nec urceolati dentibus longioribus, antheris contignis, galeæ formâ, etc., eximie distincta.

2. **P. pycnantha** (Boiss. Diagn. Ser. I, 12, p. 45). radice.... caule pumilo ascendentî parce et crispule lanatulo, foliis oblongo-lanceolatis radicalibus in lacinias lanceolatas acutissimas pinnatilobatas pinnatisectis rachide latâ, caulinis verticillatis longis acuminatis denticulatis vel lobulatis, spicâ densiflorâ oblongâ, bracteis lanceolatis acutis subintegris calyces æquantibns ciliatis, calyce parce lanato ovato subinflato nigro nervoso dentibus breviter triangularibus acutis, corollæ e sicco pallide carneâ tubo cylindrico calyce sublongiore labio superiori linearî recto obtusissimo basi subangustato, inferioris subbrevioris lobo intermedio basi valde strangulato, filamentis glaberrimis, antheris contiguis subexsertis, capsulâ.... ♀.

Hab. in monte Totschal prope Teheran Persia ad nives (Ky. 166!).

Folia radicalia 3-4-pollicies longa medium versus pollicem fere lata, caulinâ sesquipollicaria 3-4 lineas lata, caules tripollicares, spica 1½-2-pollicaris. Corolla 8 lineas longa; labii superioris formâ *P. Caucasicam* referens sed foliorum, bractearum etc. characteribus species distinctissima.

2. P. Cadmea (Boiss. Diagn. Ser. I, 4, p. 82) cespitosa, radieis fibris crassiusculis elongatis apice attenuatis, foliis parvis ambitu oblongo-linearibus in lacinulas minimas breves divaricatas bipinnatisectis, caulinis verticillatis, spicâ densâ ovatâ arachnoideo-lanatâ, bracteis ovato-rhombeis pectinato-incisis superioribus apice 2-3-fidis, calycis tubulosi dentibus brevissimis oblongis mucronulatis parce dentieulatis, corollæ roseæ tubo calyce duplo longiore versus medium infracto in limbum superne ampliato, labio superiore linearis recto apice rotundato obtuso erostri, inferioris sublongioris lobo intermedio basi angustato, filamentis glabris, antheris contiguis subinclusis, capsulâ oblongâ obliquâ falcata acutâ calyce sesquilongiore ♀.

Hab. in regione superiore Cadmi supra Gheyra in Cariâ 6500' (Boiss!), Cappadociâ orientali (Auch. 1922!).

3. longiflora. — Corollæ tubus calyce triplo longior. Forma ad typum transiens.

Hab. in monte Akdag Lyciae (Bourg. 480!), Davrosdagh Pisidiae 6-7000' (Heldr!), Berytdagh Cataoniæ 8-10000' (Haussk!), Tauro Cilicio 8'000' (Ky. 59! sub. *P. Caucasicâ*! Bal!).

Species a cl. Bth. cum *P. Caucasicâ* combinata a quâ differt foliis tenuius et magis divisis, corollæ tubo elongato, limbo ampliore, filamentis glabris, capsulâ curvata.

4. P. verticillata (L. Sp. 846) caulis pluribus pumilis simplicibus 2-4-fariam puberulis, foliis linearibus in lacinias oblongas obtusas denticulatas pinnatipartitis, caulinis diminutis verticillatis, floribus verticillatis in spicam brevem ovatam basi saepe interruptam congestis, bracteis foliiformibus pinnatifidis, calycis rubelli hirti inflati fauce coarctati urceolati superne fissi dentibus brevissimis, corollæ purpureæ tubo calyce duplo longiore imâ basi infracto, galea linearis extremitate subarcuatâ rectangulâ apice acutiusculâ, labii inferioris æquilongi lobo intermedio suborbiculato, filamentis longioribus barbatis, antherarum paribus distantibus, capsulâ lanceolatâ acutâ subobliquâ calyce sesquilongiore ♀. Jacq. Austr. tab. 206. — Rehb. Germ. tab. 141.

Hab. in pratis regionis alpinæ Macedoniae et Thraciae, Scardus (Griseb!) Rhodope (Friv!).

At Geogr. Montes Hispaniæ australis, Italiae mediae, Pyrenæi, Alpes, regio Danubialis, Lapponia orientalis, Rossia arctica, Sibiria, America arctica.

5. P. Pontica, caulis pluribus pumilis simplicibus glabratiss vel parce et plurifariam puberulis, foliis ambitu oblongo-lanceolatis in lacinulas oblongas acutas acute denticulatas pinnatipartitis, caulinis verticillatis, floribus verticillatis in spicam brevem ovatam basi interdum interruptam congestis, bracteis lanatulis in lacinias lineares 3-4-fidis, calycis lanatuli tubuloso-campanulati dentibus lanceolatis obsolete denticulatis, corollæ roseæ tubo calyce sesquilongiore versus medium infracto, galea linearis rectâ extremitate subarcuatâ erostri

oblique truncatâ, labii inferioris sublongioris lobis ovatis, filamentis glabris, antheris contiguis non prominulis, capsulâ lanceolatâ acutâ rectiusculâ calyce sesquilongiore ♀. *P. verticillata* Tchih. As. Min. II, p. 46 non L.

Hab. in alpibus Ponti inter Trapezuntem et Baibut 6000' (Ky. 512!) et in Ponto Lazico supra Djimil 7500' (Bal. sub *P. Caucasicā*), in monte Bingöldagh Armeniae (Tchih!).

A *P. Caucasicā* formis galeâ non apice rotundatâ sed angulo recto subfalcato-curvatâ oblique truncatâ discedens, hâc corollæ formâ affinis *P. verticillata* sed distincta lacinias foliorum acutis, calyce non urceolato-gibbo, filamentis glabris, antherarum paribus non distantibus. *P. amena* filamentis barbatis, capsulâ ovatâ, etc. differt.

* * Folia caulinâ sparsâ.

6. *P. orthantha* (Griseb. Spicil. II, p. 15) radicis fibris cylindricis longis, caulis pumilis crispule hirtis vel glabratis, foliis ambitu oblongis in lacinulas angustas lineares brevissimas subpinnatisectis, caulinis sparsis, bracteis petiolatis anguste linearibus pinuatifidis calyces et sœpe corollas superantibus, floribus paucis breviter spicatis inferioribus pedicellatis, calycis lanatuli tubuloso-campanulati dentibus lanceolatis integris obtusinsculis tubo subbrevioribus, corollæ rubrae tubo calyce sesquilongiore subcurvato, galeâ linearis rectâ apice rotundatâ erostri, labii inferioris galeâ vix brevioris lobis rotundatis, filamentis superne parce pilosis, antheris contiguis extus subprominulis, capsulâ ovato-oblongâ acutâ subobliquâ calycem parum excedente ♀.

Hab. in regione alpinâ Macedoniae in monte Scardo (Peristeri) 5200'-6500' (Griseb!), in Thracie monte Rhodope supra Karlova (Friv. sub *P. rosea*!) et supra Kalofer (Jankat!).

Caulis 2-6-pollicaris, corolla 6 lineas longa. Affinis *P. rosea* Jacq. foliis ambitu multo angustioribus lacinulis acutis et galeâ subfalcata distinctissime.

7. *P. Olympica* (Boiss. Diagn. Ser. I, 4, p. 83) cespitosa, radicis fibris tenuibus brevibus, caulis nanis glabris, foliis ambitu anguste linearibus in lacinias breves oblongas obtusas calcareo-crenulatas pinnatipartitis vel fidis, foliis caulinis sessilibus diminutis sparsis, bracteis oblongo-lanceolatis calyces subsuperantibus inferioribus crenulatis cæteris integris, spicâ ovatâ densâ, floribus parvis sessilibus, calycis patule hirti campanulati dentibus lanceolatis acutis integris tubo triplo brevioribus, corollæ tubo subincurve calyce sublongiore labio superiore recte linearis apice rotundato erostri, labii inferioris duplo brevioris lobis rotundatis, filamentis glabris, antheris contiguis extus non prominentis, capsulâ... ♀.

Hab. in pratis humidissimis regionis alpine inferioris Olympi Bithyni (Auch. 1919! Boiss! Pichler! etc.). Floret Maio.

Planta 2-4-pollicaris, folia *P. flammæ* vel *versicoloris* sed angustiora, corolla 5-6 lineas tantum longa. A *P. orthantha* foliis angustis multo minus divisis, caulinis bracteisque sessilibus, floribus sessilibus, corollæ labio infe-

riore duplo breviore, filamentis glabris etc. distincta. *P. flammea* et *versicolor* galeā antice subfalcatā, etc., differunt.

S. P. atropurpurea (Nordm. Bull. Acad. Petrop. II, p. 313), radicis ramis crassis attenuatis, caule elato glabrinculo folioso foliis radicalibus longe petiolatis caulinis sessilibus omnibus ambitu late ovatis bipinnatisectis lacinias lanceolatis acuminatis acute pinnatifidis, spicā densā lanatā, bracteis flore brevioribus anguste linearibus apice parce serrulatis, calycis ovali membranacei lanati dentibus brevibus triangulari-ovatis acutis, corolla atropurpurea glabra tubo vix curvato calyce duplo longiore, galeā rectā linearī apice rotundatā sub apice edentulā vel obsolete utrinque unidentatā, labii inferioris brevioris lobis oblongis lateralibus angustioribus, filamentis duobus superne villosis, capsulā compressā ovatā rectā acutā calycem non excedente ♂. *P. rillobractea* C. Koch Linn. XXII, p. 682

Hab. in regione alpinā Ponti (C. Koch!), Ponti Lazici supra Djimil 7000' (Bal!), Gurie prope Achalzik (Nordm.), Armeniae Rossiae (Szov!).

Bipedalis, foliorum radicalium laciniae primariae bipinnatae, spica florifera 2 polli., fructifera 4-5-pollicaris, corolla angusta 9 lineas longa. A *P. condensata* longe differt corollae purpureae tubo recto, galeā angustiore subdentata, capsulā acutā, etc.

9. P. Wilhelmsiana (Fisch. in M.B. Taur. Cauc. III, p. 412) lanatula, radicis ramis elongatis, caule collum versus folioso ceterum nudo, foliis ambitu anguste lanceolatis in lacinias oblongo-triangulares acutas acute denticulatas pinnatisectis, spicā densā villosā bracteis angustis linearī-lanceolatis pectinatum pinnatifidis flores longe superantibus foliosā et comosā, calycis lanati nrceolati inflati dentibus linearibus elongatis inaequalibus, corollae parvae glabrae purpureae tubo recto calycem vix excedente galeā rectā linearī apice rotundatā erostri basi subangustatā, labii inferioris brevioris lobis lateralibus latioribus retusis, antheris omnibus barbatis, capsulā oblongā oblique cuspidatā calyce sesquilociore ② esse videtur. Ic. Stev. Mon. I. 16.

Hab. in regione subalpinā et alpinā Ponti Lazici supra Djimil 7500' (Bal!), Caucasi bor. ad Nartzanum et in monte Beschtan (M.B., Hoh!), monte Sarial Georgiae Caucasicæ 4500'-5000' (Hohen!), prov. Karabagh (Schmidt).

Palmaris vel pedalis, spicā longe foliosā, corollis pro plante magnitudine parvis insignis.

10. P. condensata (M.B. Taur. Cauc. II, p. 72) tota pubescens, radicis fibris crassis attenuatis, caule elato, foliis amplis ambitu late oblongis in segmenta lanceolata subbipinnatifida acute calcareo-denticulata pinnatisectis, spicā densā lanatā, bracteis dentatis intimis exceptis flore brevioribus, calycis campanulati lanati dentibus breviter triangularibus, corollae flavae glaberrimae tubo incurvo calyce duplo longiore, galeā gibbā obtusissimā antice subarcuatā, labio inferiore subbreviore lobis rotundatis intermedio minore, filamentis omnibus barbatis, capsulā obovatā equali obtusā calyce inclusā ♀.

Ic. Stev. Mon. tab. 17. — *P. campylosiphon* C. Koch Linn. XXIII, p. 682. — *P. Hacquetii* C. Koch Linn. XVII. p. 42 non Graf

Hab. in pratis alpinis et subalpinis montis Ylgasdagh Anatoliae borealis (Wied!), Ponti Lazici supra Trapezuntum (Ky. Suppl. 707!), supra Khabackar (Bal!), in Guriâ (Nordm.), Caucaso occid. (C.A. Meyer), centrali (Rehm!), orientali in Tuschetia, Salataviâ, Daghestaniâ 7000'-8500' (Rupr!), prov. Karabagh (Szov!).

β. glabrescens. — Canlis et folia glabra.

Hab. in pratis uidis Ponti Lazici supra Djimil (Bal!).

Sæpius bipedalis sed in excelsioribus occurruunt etiam formæ 3-4-pollicares. Species a *P. foliosâ* et *P. Hacquetii* affinibus bracteis flores non superantibus, corollâ glabrâ, capsulâ non obliquâ mucronata nec inæqualiter biloculari distincta.

§ 2. Rostratæ.

* Folia caulina verticillata.

11. **P. subrostrata** (C. A. Mey. Enum. p. 108) caule nano bifarium puberulo, foliis anguste linearibus in lobulos ovatos pinna-tisectis glabris, caulinis diminutis verticillatis, spicâ brevi densâ ovato-oblongâ, bracteis et calycibus præter pilos lanatulos paucos secus marginem glabris illis oblongis apice dentatis hujus laciniis subdentatis, corollæ purpureæ tubo sensim curvato calyce duplo longiore, galeâ rectâ brevissime rostratâ truncatâ et denticulo auctâ ♀.

Hab. in pratis alpinis Caucasi occidentalis 7-8500' (C.A. Mey!).

Descripti specimen unicum et radice orbatum herbarii Meyeriani; plantula tripollicaris, capitulum 9 lineas longum. Species inter erostres et rostratas v. gr. inter *P. Ponticam* et *crassirostrem* intermedia, huic tamen rostro etsi brevi tamen inanifesto magis accedens et ab eâ bracteis et calyce glabriusculis, corollæ tubo longiore distincta.

12. **P. crassirostris** (Bge' Bull. Acad. Petersb. VIII, pag. 248) radicis fibris tenuibus, caulis sæpius pluribus ascendéntibus pumilis seriatim pubescentibus, foliis anguste oblongo-linearibus in laciniis minimis ovatis bipinnatisectis, caulinis verticillatis diminutis, spicis ovatis et oblongis sæpius interruptis, bracteis modice lanatulis palmatisidis, calycis sublanati anguste tubuloso-campanulati laciniis oblongis acute denticulatis, corollæ purpureæ tubo sensim curvato calyce sublongiore, galeâ arcuatâ sensim attenuatâ in rostrum descendens concavum apice truncatum, labii inferioris sublongioris lobis oblongis, filamentis glabris, capsulâ oblongâ acutâ subenervata calyce sesquiongiore ♀. *P. Araratica* Bge Mem. Ac. Petersb. Ser. 6, tome VII, p. 594.

Hab. in pratis Caucasi, Imeretia ad Tchkméri (Guld.), Suania (Radde!), ad glacies Kasbek 10000' (Rehm!), Tuschetia supra Dano et Diklo 8000', Daghestania in montibus Saridagh, Gudurdagh, Artchikala, etc. 8000'-10000' (Rupr!), mons Ararat Armeniae (Abich!) et Alagös (Frick!).

Planta 2-5-pollicaris facie *P. rostrata* referens sed folia caulinis verticillata, flores subminores et rostrum brevius. Plantæ ex Caucaso orientali ob stationem sicciorum minores evadunt corollæ limbo subminore. Planta ex monte Ararat nisi rostro subbreviore differre videtur.

.. Folia caulinis sparsa.

13. **P. Nordmanniana** (Bge in Ledeb. Fl. Ross. III, p. 277) præter caules interdum lineatum et brevissime puberulos glabra, radicis fibris tenuibus, caulis pumilis vel mediocribus, foliis ambitu oblongo-lanceolatis angustis in segmenta oblonga acute et saepè duplicatim dentata pinnatisectis, caulinis diminutis sparsis, floribus paucis capitatis tandem racemosis inferioribus stricte pedicellatis, bracteis inferioribus trifidis dentatis, calyce glabro tubuloso ad tertiam partem in dentes linearis-oblungos inæquales denticulatos fiso, corollæ roseæ tubo calyce sublongiore, galea semicirculariter curvata in rostrum porrectum conico-lineare tenue apice truncatum abruptinuscule angustatâ, labii inferioris lobis latis ovatis galeam æquantibus, filamentis duobus superne parce pilosis, capsula oblonga acutâ subobliqua calyce tertia parte longiore ♀. *P. rostrata* C. A. Mey. Enum. p. 108 non L.

Hab. in pratis alpinis Ponti Lazici supra Djimil 7500' (Balt!), ditione Artaban (C. Koch!), Guriâ (Nordm!), Abchasiâ (Radde!), Caucaso occidentali (C.A. Mey.), ditionibus Alagir, Radscha, Chewsuriâ et Tuschetia 8000'-9500' (Ruprecht!), Armenia Rossica (Szov!).

Caules variant 2-7-pollicares; facies *P. crassirostris* foliis caulinis verticillatis et rostro breviore deorum magis flexo distinctæ. *P. rostrata* L. et *P. Jacquinii* Koch caulis pumilis ascendentibus, calyce plus minus hirsuto, rostro deorsum curvato nec horizontaliter porrecto, radicis fibris crassioribus differunt.

14. **P. Elephas** (Boiss. Diagn. Ser. I, 4, p. 81) glaberrima, caule nano, foliis parvis ambitu anguste linearibus in lacinulas ovatas subimbricatas ovataso btusas acute calcareo-denticulatas pinnatisectis calyces æquantibus, calyce glabro tubuloso ad tertiam partem in dentes linearis-spathulatos calcareo-denticulatos fiso dente inferiore breviore, corollæ roseæ tubo calyce sesquiongiore, limbi antorsum valde curvati labio superiore in rostrum sensim attenuatum longissimum circinnatum apice truncatum bifidum abeunti. labii inferioris latissimi lobis subæqualibus rotundato-subretusis, filamentis glabris, capsula oblongo-lineari obtusa oblique mucronata calyce sublongiore ♀.

Hab. in alpe Elamout Persiae borealis (Auch. 5083!), in valle Hajiguk Afghanistanæ ad nives 10500' (Griff. Journ. 1060!) eadem videtur.

Bipollicaris, folia 2-3 lineas lata, calyx 4½ lineas longus. Affinis *P. rhinanthoides* Schrenck et *P. Hookerianæ* Wall.; ambo sunt plantæ multo majores, prior differt insuper tubo corollino calyce plus duplo longiore et calyce lanatulo, posterior calycis hirsuti dentibus late ovatis, corollâ hirtella, etc.

15. **P. labellata** (Jacquem. in Decaisn. Voy. Jacq. p. 118, tab. 123) glabra, caulis humilibus ascendentibus, foliis parvis oblongis

pinnatilobatis laciniis parvis ovatis inciso-dentatis, floribus 3-4 terminalibus approximatis pedicellatis, bracteis foliis similibus minoribus margine arachnoideo-pilosus, calycis tubo arachnoideo-piloso tubuloso ad sextam partem in dentes glabros fisso dente postico breviore integro lanceolato cæteris ovatis argute serratis, corollæ roseæ tubo calyce subduplo longiore, limbi antrorsum valde curvati labio superiore in rostrum sensim attenuatum longum circinnatum abeunte, labii inferioris latissimi lobis obtusis lateralibus subquadratis subretusis intermedio rotundato angustiore, filamentis pilosis, capsulâ... ♀.

Hab. in regni Cabulici alpibus (Griff. 636). Non vidi.

Ar. Geogr. India boreali-occidentalis.

§ 3. **Bicuspidatæ.**

16. P. leucodon (Griseb. Spic. II, p. 17) radicis fibris a basi tenuiore incrassatis napuliformibus, caule lanatulo folia radicalia parum superante, foliis oblongo-lanceolatis pilosis in laciniis oblongo-lanceolatas pinnatipartitas pinnatisectis, spicæ densiusculâ, bracteis supremis subintegris calycem æquantibus, calycis campanulati pilosuli dentibus linearispathulatis apice denticulatis tubo triplo longioribus, corollæ ochroleucae (9 lin. longæ) glabræ galeâ sensim curvatâ juxta apicem truncatum dente triangulari-subulato utrinque auctâ, labii inferioris glabri lobis obtusissimis lateralibus latioribus, crista binis longis parallelis, filamentis superne pilosis ♀. *P. occulta* Janka Brev. 2, p. 7!

Hab. in regione alpinâ montis Kobelitza (Scardi) Macedoniae 5-7000' (Griseb!), in monte Rhodope Thracie supra Kalofe (Janka!).

Semipedalis, in sectione calycis dentium formâ insignis.

17. P. Græca (Bge Bull. Phys. Math. Petersb. I, p. 10) arachnoideo-lanatula, foliis ambitu ovato-oblongis in laciniis angustissime et breviter lineares divaricatas bipinnatisectis, spicæ floriferâ ovata dein elongatâ, bracteis subpetiolatis linearibus pinnatifidis inferioribus calyces superantibus, calycis lanati campanulati dentibus linearibus subdenticulatis dimidium tubum æquantibus, corollæ ochroleucae (12 lin. longæ) glabræ galeâ incurvâ erostri sub apice truncato longe subulato-bidentatâ. labii inferioris lobis denticulatis glabris, crista binis longis parallelis, filamentis duobus puberulis, capsulâ oblongâ acutâ subobliquâ calycem parum excedeute ♀. *P. rupestris* Boiss. et Orph. Diagn. Ser. II, 3, p. 175.

Hab. in regione alpinâ montis Velugo Ætoliae (Fraas), Parnassi 5500'-6000' (Guice! Heldr. Herb. Norm. 669! Orph!), montium Chelmos et Kyllene (Orph!).

Radice et facie affinis *P. Friderici* differt foliis tenuius sectis magis abbreviatis, bracteis elongatis pinnatifidis, calycis laciniis linearibus obtusis denticulatis. *P. leucodon* affinis quoque differt foliis angustis, radice napuliferâ, corollâ brevi, etc.

18. P. Friderici-Augusti (Tommasini Linn. XIII, p. 69, t. 2) radicis fibris funiformibus aequalibus, caule erecto glabriusculo, foliis glabris ambitu oblongo-lanceolatis in lacinias anguste lineares pinnatifidas pinnatisectis, spicula floriferà brevi densa, bracteis flores non aequalibus lanceolatis superioribus subulatis integris, calycis lanati campanulati dentibus lanceolatis acutis integris latitudine triplo longioribus, corollae ochroleucæ (12 lin. longæ) extus saepe parce hirsute galea sensim curvata justa apicem truncatum dente subulato utrinque aucta, labii lobis denticulatis glabris cristis binis longis parallelis, filamentis parce pilosis, capsula ovata acentra calyce quartâ parle longiore ♀. Ic. Rehb. Germ. tab. 130 et 141.

Hab. in regione alpinâ montis Kobelitz Scardi in Macedonia (Friedr. ex Griseb.).

Ar. Geogr. Illyria, Dalmatia, Serbia, Bosnia, Montenegro.

19. P. brachydonta (Schlosser et Vučatinovich Fl. Croatica p. 632) minute pubescens vel glabrata secca nigricans, radicis fibris napuliformibus, foliis ambitu oblongo lanceolatis in lacinias lanceolatas pinnatipartitas calcareo-denticulatas pinnatisectis, spicula oblonga, bracteis lanceolatis pinnatifidis et integris calyces aequalibus, calycis parce hirti dentibus triangularibus latitudini aequilongis, corolla ochroleucæ (8-10 lin. longæ) tubo parum curvato galea incurvâ subrostrata apice oblique truncata et bidentata, labii inferioris lobis obtusis denticulatis glabris, cristis ad gibbos ovatos binos sub lobo intermedio reductis, filamentis binis superne hirtis, capsula oblongo-lanceolata falcato-rostrata compressa calyce subduplo longiore ♀. *P. ochroleuca* Schlosser in Rehb. Germ. XX, p. 71, tab. 134). — *P. leucodon* Rehb. fil. Germ. p. 70, tab. 133 non Griseb. — *P. comosa* Griseb. Spic. II, p. 17 ex parte quoad plantam Macedonicam!

Hab. in montibus Ljubatrin et Peristeri (Scardus) Macedoniae (Griseb! Orph!).

Species a *P. comosa* et affinibus pyrami crista non elongatis sed ad gibbos ovatos reductis distincta, a *P. leucodonte* formâ dentium calvæ aliena. Planta Bosnica et Macedoniae (eius fructum non vidi) nihil similes videntur, in utriusque labium dentielatum quidem sed non ciliatum vidi.

Ar. Geogr. Croatia, Bosnia.

20. P. comosa (L. Sp. 847) pubescens vel glabrata, radicis fibris napuliformibus, foliis ambitu lanceolatis in lacinias lanceolatas pinnatipartitas cartilagineo-dentatas pinnatisectis, spicula ovata vel oblonga densa, bracteis inferioribus pinnatifidis superioribus integris, calycis campanulato-cylindrici membranacei dentibus brevissimis ovatis obtusis latitudine suâ brevioribus, corolla ochroleucæ 10-12 lineas longe galea incurvâ subrostrata apice oblique truncata et in dentes binos triangulari-subulatos abeunte, labii inferioris lobis obtusis vel retusis margine ciliolatis vel nudis, faucis cristis binis lon-

gis parallelis, filamentis duobus pilosis, capsula oblonga apice incurva acuta calyce tertia parte longiore ♀. Rehb. Germ. tab. 136.

Hab. in pratis subalpinis et alpinis, mons Paryadres Ponti (Tchih!), Caucasus centralis et septentrionalis (Rupr! Ower!), orientalis 8000'-9000' (Rupr!), Iberia 4200' (Rupr!), Persia borealis (Buhse!), inter Siaret et Schahrud (Bge!).

β . *Sibthorpii*. — Calycis criso-lanati dentes acuti vel apiculati, corollæ labium inferius glabrum. — *P. Sibthorpii* Boiss Diagn. Ser. I, 4. p. 83. — *P. flava* Sm. Prodr. I. p. 429 non Pall. — *P. bicuspidata* Griseb. It. Hung. p. 325. — *P. achilleæfolia* Hoh. exs. non Steph.

Hab. in Olympo Bithyno (Sibth! Boiss! Auch!), montibus prope Boli (Wied!), Tmolo supra Philadelphiam (Boiss!). Armeniæ (Huett!). Georgia Caucasicæ (Hohenst!), Persiæ boreali (Buhse!), montibus Tauriae (Rehm!).

γ . *acmodonta*. — Plantæ glabriuscensæ siccitate valde nigricans, calycis glabriuscensili vel glabri dentes muerone calcareo apiculati interdum æquilongi et lati, labium inferius glaberrimum. — *P. acmodonta* Boiss. loc. cit. p. 84. — *P. jueunda* Schott et Ky. in Sched.

Hab. in Pisidiâ prope Bouldour (Heldr!), Tauro Cilicio 8000' (Ky! Bal!), Cappadociâ (Auch. 1921! Ky. 482! Bal!), Cataoniae monte Berytdagh 8000' (Haussk!), Armeniæ Turcicâ supra Gumuschkhané (Bourg!), Ponto Lazico (Bal!), Persiæ boreali monte Savalan (Auch. 5082!), Demavend (Ky. 387!).

Formæ hic enumeratae altera in alteram transire videntur, character labii inferioris corollæ glabri vel ciliati non sat firmus. Varietates β et γ in Oriente formâ typicâ multo frequentiores sunt.

Ar. Geogr. Alpes Hispaniæ, Galliæ australis, Italiae, Dalmatiæ, regionis Danubialis, Rossia media et australis, Sibiria tota.

ORD. LXXXIX. OROBANCHACEÆ.

(Lindl. Introd. Ed. 2^a p. 287.)

Flores hermaplroditi irregulares. Calyx liber persistens 4-5-sepalus, sepalis in calycem gamosepalum 4-5-sidum vel in segmenta bina lateralia coalitis. Corolla gamopetala hypogyna tubulosa vel campanulata limbo obliquo plus minusve bilabiata. Stamina 4 didynama tubo corollæ inserta, antheræ biloculares. Ovarium liberum uniloculare placentis parietalibus per paria contiguis vel distinctis, stylus simplex, stigma capitato-bilobum. Ovula plurima anatropa. Capsula anilocularis in valvas binas ad medium placentiferas plus minus profunde dehiscentes. Semina plurima subglobosa vel oblonga minima testâ foveolata vel striata. Albumen carnosum, embryo prope hilum minutus, cotyledones indistinctæ. — Herbæ fibrillis radiciformibus e candice ortis aliarum plantarum radicibus adnatae, nunquam virides, pro foliis squamas cauli concolores gerentes, saepius perennes.

LATHRÆA (L. Gen. 743).

Flores hermaphroditi basi bracteæ fulti ebracteolati. Calyx campa-nulatus quadrifidus. Corolla bilabiata. Stamina 4 didynama corollæ tubo inserta, antherarum loculi basi discreti et mucronati. Ovarium antice glandulæ liberæ fultum, stylus crassus, stigma capitatum. Capsula unilocularis bivalvis valvis juxta axiu placentas quatuor geminatim confluentes ferentibus. Semina plurima vel pauca. — Herbae parasiticæ caulis parte subterranea rhizomatiformi.

* Placentæ dilatatae. Semina numerosa parva favosa.

1. **L. Squamaria** (L. Sp. 848) rhizomate squamis carnosis imbricatis vestito ramoso, caule brevi erecto paucisquamato, racemo terminali denso bracteato secundo ante anthesin cernuo, bracteis ovatis vel obtuse rhombis, floribus pedicellatis subpendulis, calyce campanulato ad medium in 4 lobos triangulari-ovatos obtusos fuso, corollæ albido-roseæ calyce sesquilongiore labio superiore subfornicato integro inferiore subbreviore tridentato, antheris villosis ♀. Ic. Fl. Dan. tab. 136. — Rehb. Germ. 143.

Hab. ad arbores et frutices parasitica.

Hab. in sylvis montanis, Macedoniae in quergetis Athonis 3000' (Griseb!), Bithynia (Griseb., Auch. 5081!, Wiedem!), in monte Amano Syriae (Auch. 2853!), Tauria (M B.).

E. erecta. — C. Koch Linn. XVII, p. 291. — Flores horizontales, corollæ labium superius sordide roseum, antheræ valde lanatae.

Hab. in Somchetia (C. Koch), Iberia Caucasicæ (Rupr!).

Ar. Geogr. Europa media et australis a Scandinaviâ ad Rossiam medium, Sibiria Uralensis, regio Himalaica.

** Placentæ versus medium vix dilatatae. Semina minuscula 3-5 subglobosa circa umbilicum præsertim costato-radiata.

2. **L. Rhodopea** (Dingler in Bot. Zeit. 1877, p. 74) rhizomate squamis carnosis imbricatis vestito ramoso, caule erecto gracili sparsim squamato, racemo terminali elongato cylindraceo denso, bracteis cuneato-lanceolatis obtusis, floribus breviter pedicellatis, calyce glandulifero campanulato ad medium bilabiato dentibus acutiusculis anticis linear-lanceolatis posticis oblique triangularibus, corollæ intus glabrae tubo calycem subsuperante limbo duplo longiore, labio superiore galeato mutico, autico subbreviore breviter triangulari acuminato apice breviter tridentato, ovario pauciovulato stigmate subcapitato-incrassato, capsula calyce breviore obcordata, antheris albo-villosis ♀.

Hab. in fagetis umbrosis montis Rhodopes meridionalis Thraciae (Karlyk-dagh) prope Gumeldschinam 3500'-3800' (Dingler!) Floret Maio

Cl. detectoris benevolentia specimen plantæ rarissimæ accepi sed ætate ninius provectum. corollæ et staminibus jam delapsi. hæc omnia igitur ex accuratâ auctoris descriptione depropensi. Caulis cum racemo denso fere semi-pollicari pedalis, calyx fructifer 5 lineas longus. Facie a *L. squamaria* et *clandestinâ* aliena sed *L. Japonica* Miq. affinis. Habitus omnino *Neottie nivalis* ovis.

ANOPLANTHUS (Endl. Gen. 727. — *Anoplon* Don).

Flores hermaphroditæ ebracteati. Calyx campanulatus quinquefidus. Corolla (in nostro) breviter campanulata limbo patentí subbilabiato subæqualiter quinquelobo. Stamina 4 didynama corollæ tubo inserta, antherarum loculi basi discreti mucronati. Ovarium uniloculare, stigma capitato-subbilobum. Capsula unilocularis bivalvis valvis intra marginem utrinque placentiferis placentis convergentibus. Semina plurima reticulata. — Herba parasitica caule subuniflora.

1. **A. coccineus** (Marsch. Casp. p. 58 sub *Orobanche*) glabra vel superne sub lente minnte glandulosa, caule tereti striato squamis ovato-oblongis semiamplexicilibus vaginantibus remotis obsito unifloro rarius bifloro, flore maximo, calyce campanulato corollæ tubum æquante fere ad medium in lacinias late triangulari-lineares obtusas tisso, corollæ coccineæ tubo breviter campanulato, limbo tubo sublongiore patentí subbilabiato lobis ovatis obtusis subæqualibus ♀. *A. coccineus* Walp. Rep. III. p. 481. — *A. Biebersteinii* Reut. in D.C. Prodr. II. p. 42. — *Anoplon Biebersteinii* C. A. Mey. Enum. p. 104. — *Lathraea Phœliepa* var. β Linn. Sp. 844. — *Phelipaea foliata* Lamb. Trans. Linn. Soc. X. 260, tab. 7. — Ic. M.B. Cent. t. 56. — Rehb. pl. Crit. fig. 609, 930.

Hab. in regione montanâ et alpinâ ad radices *Centaurearum* plurium parasitica, mons Cadmus Cariae supra Denisleh (Boiss! Pinard!), Ciliciae ditio Kassan Oghlu 4000' in quercentis (Ky. 44!), montis Cassii Syrie borealis regio superior (Boiss!), Tauria et Caucasus (M.B., Stev.).

z. peduncularis. — Squamæ ad basin caulis superne longe nudi magis congestæ. vix varietas. — *Phelipaea Tournefortii* Desf. Cor. tab. 10. — *Anoplanthus Tournefortii* Reut. in D.C. Prodr. II. p. 42. — *Phel. Orientalis* flore coccineo Tourn. Inst. tab. 479.

Hab. in Armeniâ Turcieâ circa Erzerum (Tourn! Huet! Calv!), in Kurdistanâ (Barré de Lancey!), Armeniâ Rossicâ (Buhse!), in pratis montosis Georgie Caucasicæ (Hoh!).

z. nigrovittatus. — Corolla vittâ latâ nigra obsita.

Hab. in graminosis montis Pir Omar Gudrun 5-6000' (Haussk!).

Scapus variat 4-5-pollicaris, pedalis et longior, corollæ limbus diametro 1-1½ pollicaris.

PHELIPÆA (Tourn. Cor. 47. — Desf. Atl. II, p. 60).

Flores hermaphroditæ basi bracteâ fulti lateraliter bibracteolati. Calyx 4-5-fidus. Corolla bilabiata labio superiore bifido inferiore

trifido. Stamina 4 didynama corollæ tubo inserta, antherarum loculi basi mucronati, connectivus muticus vel mucronatus. Ovarium uniloculare placentis parietalibus 4 geminatis, stigma capitato-bilobum. Capsula unilocularis bivalvis valvulis apice dehiscentibus. Semina plurima reticulata. — Herbae parasitæ facie *Orobanches*.

Sect. I. TRIONYCHION. — Wallr. Orob. p. 58. — Calyx 4-5-tidus lobis elongatis inæqualibus acuminatis. Antheræ superficie glabrae loculis secus dehiscentiae lineam glabris vel plus minus villosis. Capsula styli decidui basi coronata.

* Caulis semper simplex.

+ Corollæ lobi acuti vel acutiusculi.

1. **P. cærulea** (Vill. Dauph. II, p. 406 sub *Orobanche*) caule simplici saepè elato, spicâ laxinseculâ totâ pubescenti-glandulosâ, bracteâ mediâ ovato-lanceolatâ calyce breviore lateralibus subulatis, calycis laciniis 5 rarins 4 lanceolatis acuminatis, corollâ calyce $2\frac{1}{2}$ plo longiore pubescenti-glandulosâ amethystinâ tubulosâ autrorsum curvatâ tubo medio angustato laciniis labiorum acutis planis, antheris glabris vel apice parce barbatis, stigmate albido bigloboso \neq . *P. cærulea* C. A. Mey. En. 104. — Ie. Rehb. Crit. 7, fig. 928. Ie. Germ. tab. 141. — *P. Cedreti* Reut. in Noë exs. — *P. Syssirensis* C. Koch Linn. XXIII.

Hab. in *Anthemide Austriacâ* ad cacumen Pentelici Atticae (Sart!), Macedonia in Athone (Griseb.), collibus Lycae prope Elmali (Bourg!), Anatolia boreali (Wied!), Libano (Boiss! Noë!), Tauri in *Brassici polymorphâ* (ex Stev.), Armeniâ (Koch), Iberia (M.B., Hoh.), Persia boreali (Bge!).

Ar. Geogr. Europa media et australis ab Anglia et Belgio ad Rossiam medium, Sibiria.

2. **P. longiflora** (Pers. Enc. II, p. 182 sub *Orobanche*) caule simplici gracili, spicâ laxinseculâ pubescenti-glandulosâ, bracteâ mediâ ovatâ attenuato-acuminatâ calyce breviore lateralibus anguste linearilanceolatis, calycis laciniis 4-5-nis brevibus subulato-acuminatis, corollâ calyce triplo longiore pubescenti-glandulosâ anguste tubulosâ curvatâ cæruleo-violaceâ lobi oblongis acutiusculis, antheris lanatis, stigmate bilobo \neq . *P. longiflora* C. A. M. Eu. p. 104. — Ie. Rehb. Crit. 7, fig. 932.

Hab. Byzantii (Noë exs. 74!), in Ponto (Tchih!), ad Baibut Armeniae (Bourg!), in monte Ssodagh Cataonie (Haussk!), ad Helenendorf Transcaucasiae (Hohen!).

Affinis *P. arenarie* a qua differre videtur corollâ tenuiore longiore magis curvatâ, lobi minus rotundatis.

Ar. Geogr. Rossia australis.

++ Corollæ lobi obtusissimi.

3. **P. arenaria** (Borkh. in Rœm. Mag. p. 6 sub *Orobanche*),

caule simplici, spicâ laxiusculâ pubescenti-glandulosâ, bracteâ mediâ lanceolatâ calyceem æquante vel superante lateralibus linearis-subulatis, calycis laciniis 5 rarius 4-lanceolatis acuminatis, corollâ calyce 2 ½-plo longiore pubescenti-glandulosâ cærneleo-violaceâ tubulosâ subrectâ rarius curvatâ superne inflatâ lobis obtusis margine reflexis, antheris ad suturam valde lanatis, stylo stipitato-glanduloso, stigmate albo-flavido bigloboso ♀. *P. arenaria* Walp. III, p. 459. — Ic. Rehb. Crit. 7, fig. 929-931. — Ic. Germ. tab. 145. — *P. obtusiloba* Reut. in Haussk. exs.

Hab. *Artemisiarum* parasitica in Tauriâ (ex Ledeb.), in Iberiâ prope Tiflis (ex cl. Rehb. fil.), Caucaso orientali (Rupr!), in monte Berytdagh Cataoniae 8000' (Haussk!).

Ar. Geogr. Europa media et australis a Belgio et Germania ad Rossiam australem, Africa borealis.

4. **P. lanuginosa** (C. A. Meyer in Ledeb. Fl. Alt. II, p. 450) tota adpressiuscule albido-lanuginosa, caule simplici, spicâ brevi densâ, bracteâ mediâ ovato-lanceolatâ laterales lanceolatas calycemque æquante vel superante, calycis laciniis 5 rarius 4 breviter lanceolatis, corollâ lanuginosâ brevi calyce duplo longiore subcurvatâ cæruleâ lobis parvis obtusis denticulatis, antheris glabris, stylo hirtulo, stigmate bilobo ♀. Ic. Ledeb. tab. 377. — *Orobanche cæsia* Rehb. Crit. 7 p. 48, fig. 936 nec *Phel. cæsia* Rent. in D.C. Prodr. — *O. borealis* Turez. — *Ph. araneosa* Reut. MSS.

Hab. *Artemisie*, *Teucrii* sp. parasitica in Armeniâ circâ Erzerum (Calv!) et Baibut (Bourg!), in valle Djimil Ponti Lazici 6000' (Bal!), in Caucaso prope Stavropol (C.A. Mey!), ditione Talysh (Hohen!), Persiâ boreali ad Siaret (Bge!).

Corolla 7-8 lineas tantum longa. Differt a *P. cæruleâ* lanagine, bracteis longioribus in spicæ apice comosis, etc.

Ar. Geogr. Rossia australis, Songaria, Sibiria.

5. **P. cælestis** (Reut. in D.C. Prodr. II, p. 5) sæpe pluricaulis caulis pumilis, spicâ densâ brevi ovatâ breviter et crispuo lanuginosâ, bracteâ mediâ late lanceolatâ plurinervi calyce breviore lateralibus anguste lanceolatis, calycis laciniis 4 subulato-attenuatis, corollâ cyaneâ glanduloso-hirtâ calyce 2 ½-plo longiore tubulosâ infundibuliformi valde curvato-falcataâ lobis rotundatis obtusissimis, antheris ad suturas parce lanuginosis, stylo glabriuscule, stigmate bilobo ♀.

Hab. in cacumine montis Sipyli Lydiæ (Boiss!), in montibus Cariae (Pinard!), in ditione Kassan Oglu Cilicie (Ky. Suppl. 167!), in collibus ad Gumuschkhané Armeniæ (Bourg!), ad *Cirsia*, *Cousinias*, *Anthemideas*, in monte Berytdagh Cataoniæ, Gebel Taktak Mesopotamiae (Haussk!).

Caules 2-4-pollicares, corolla 10 lineas longa. Spicæ densæ *P. lanuginosæ* sed corolla longior multo magis curvata, antheræ lanatae, bracteæ breviores.

6. **P. pulchella** (C. A. Mey. in Eichw. Casp. p. 17, tab. 18) tota albo-lanata, caule simplici humili tenuissimo, spicâ pauciflorâ

brevi ovatâ, bracteâ mediâ ovatâ acutâ calyce subbreviore lateralibus linearibus calyceem subæquantibus, colycis lobis 4 subulato-attenuatis, corollâ calyce triplo longiore violaceo-cæruleâ albo-lanatâ subinervia sensim ampliatâ lobis orbiculatis denticulatis, antheris lanatis, stylo pilis longis albis copiosis hirsuto, stigmate flavo obliquo bilobâ ♀.

Hab. in Inneretiâ (Eichw.).

Indumento ad *P. lannigerosam* accedens sed ab omnibus caule 3-4-pollici filo emporetico vix crassiore, stylo piloso et stigmate obliquo nec nutans distincta videtur.

++ Caulis sœpe ramosus.

+ Corollæ lobi acuti.

7. ***P. oxyloba*** (Reut. in p.c. Prodr. II, p. 9) parce et brevissime puberula vel glabrescens, caule simplici vel ramoso cæbre squamioso, spicâ densiusculâ brevi, bracteâ mediâ oblongo-acuminatâ lateralibusque anguste lanceolatis calyce brevioribus, calycis lobis 4 a basi breviter lanceolatâ longe acuminatis, corollâ hirtulâ pallide cœruleâ calyce triplo longiore anguste tubuloso-infundibuliformi antice curvatâ lobis ovatis acutis ciliatis, filamentis brevibus, antheris parce lanatulis, stylo hirtulo, stigmate pallide cœruleo retuso-bilobo ♀.

Hab. in vineis, ad muros Anatoliæ, Smyrniæ (Bal. 299!), ad Elmaliu Lycia (Bourg!), ad Alya Pamphyliæ (Heldr!), ad Mudania Bithyniæ (Pichler!).

Semipedalis rarius pedalis, gracilis, corolla 9 lineas longa pro longitudine angusta lobis acutis.

8. ***P. hirtiflora*** (Reut. in p.c. Prodr. II, p. 10) valde glanduloso-papillosa, caule humili crasso simplici et ramoso, spicâ densâ strictâ, bracteis nervosis mediâ late lateralibus anguste lanceolatis acuminatis calyce parum brevioribus, calycis elevatim nervosi laciniis 4 a basi latiore longe subulatis, corollâ calyce 2 ½ plo longiore tubuloso-infundibuliformi lobis ovatis acutis, antheris valde lanatis, stylo subglanduloso, stigmate retuso ♀.

Hab. ad varios *Astragalos* praesertim spinosos parasitica in montibus Persia australis, prope ruinas Persepolis (Ky. 920! et 408! ex parte), radices montium Kuh Sowers et Eschker (Haussk!), Belutschia (Stocks!).

Insignis bracteis et lobis calycinis longe subulatis strictis, a formis *P. laran-* *dulacea* lobis corollæ acutis, a *P. oxyloba* corollis majoribus et antheris valde lanatis distincta.

++ Corollæ lobi obtusi

9. ***P. lavandulacea*** (Rehb. Crit. 7, p. 48, tab. 935 sub *Orobanchæ*, breviter papillosa, caule elato simplici vel inferne parce ramoso, spicâ longâ densiusculâ multiflorâ, bracteâ mediâ ovato-lanceolatâ acuminatâ calyce subbreviore lateralibus linearisubulatis, calycis coriacæ lobis 5 subulato-attenuatis, corollâ calyce triplo longiore cœruleâ a basi tenui glabriuscûla sensim ampliatâ breviter

pulverulentâ arcuatini antrorsum curvatâ lobis ovato-orbiculatis denticulatis ciliatis, antherarum barbâ inæquali. stigmate subquadrato ♂. — *P. lavandulacea* Reut. in D.C. Prodr. Ic. Rehb. Germ. tab. 147. — *P. Schultzii* Mut. Fl. Fr. II. p. 352. — *P. Fraasii* Walp. Rep. bot. 3, p. 460. — *P. tricholoba* Reut. in D.C. Prodr. II, p. 10 — *P. comosa* Reut. in Noë exs.

Hab. ad varias plantas parasitica, Peloponnesus (Despreaux, Fraas), Zacynthus (Marg!), Salamis (Heldr!), Cilicia ad Mersina (Bal!) et ditione Kassan Oglu (Ky. Suppl. 468! sub *P. Heldreichii*), Syria in monte Cassio (Boiss!). Libano (Bl!) et prope Damascum (Gaill!), in monte Hermone et Ithabure Palestine (Boiss!), montibus Berytida Cataoniae, Aroman et Schahu Kurdistanae Persiae et monte Sawers Persie occidentalis (Haussk!), montibus Persiae australis (Ky. 408! ex parte. Auch. 5076!), Belutschia (Stocks exs. 4087!).

Erronee schedulæ commutatione Reuterus *P. tricholobam* suam ad *Tragacanthas* parasiticam indicat, hic crescendi modus ad *P. hirtifloram* spectat

Ar. Geogr. Regio mediterranea Europæ, Africa borealis, Insulæ Canarienses.

10. *P. pyramidalis* (Reut. in D.C. Prodr. II, p. 7) breviter papillosa, caule simplici rarius ad spicæ basin ramoso, spicæ longâ densâ floribus pedicellatis, bracteâ mediâ lanceolatâ acuminatâ calyce multo breviore lateralibus linearis-subulatis, calycis membranaceis laciniis subquinis subulato-acuminatis tubo longioribus, corollâ cœruleâ glanduloso-hirsutâ calyce 2 1/2-plo longiore a basi sensim ampliatâ curvatâ lobis ovatis obtusis patentibus, antheris ad commissuram longe lanatis, stylo hirtulo stigmate retuso-bilobo ♀.

Hab. in monte Kyllene Peloponnesi supra Trikala (Orph!), Tmolo Lydiæ supra Philadelphiæ (Boiss!), monte Ssoffdag Syrie borealis 3000' in consortio *P. longiflora* (Hauffsk!).

Valde affinis *P. lavandulacea* a quâ forsitan differt spicis magis compactis, floribus distinctius pedicellatis, calyce minus coriaceo. corollæ lobis tubi proportione submajoribus.

11. *P. Heldreichii* (Reut. in D.C. Prodr. II, p. 8) breviter et parœ papillari-hirta, caule crassiusculo simplici vel ad spicæ basin ramoso, spicæ longâ densâ strictâ apice ante anthesin comosâ, bracteâ mediâ lanceolatâ bracteolisque subulatis calyce brevioribus. calycis lobis 4 longe lanceolato-subulatis, corollâ pallide cœruleâ calyce duplo longiore tubuloso-infundibuliformi subcurvatâ lobis ovatis obtusis intus hirsutis, antheris glaberrimis, stylo glanduloso, stigmate retuso-bilobo albido ♀.

Hab. in arenosis Lycaoniae circa Iconium (Heldr!).

Semipedalis et pedalis, corolla 41 lineas longa, facies *P. lavandulacea* et *P. pyramidalis* a quibus antheris glabris statim dignoscitur.

12. *P. ramosa* (L. Sp. 882 sub *Orobanche*) glanduloso-pubescentia flava, caulis simplicibus vel ramosis, spicis laxiusculis, bracteâ mediâ oblongo-lanceolatâ lateralibusque anguste lanceolatis

acuminatis calyce subbrevioribus, calyce ad medium in lobos 4 a basi trianguli subulatos diviso, corollà flavidâ vel violaceâ calyce 2¹/₂ triplo longiore supra ovarium strangulatâ superne curvata lobis ovatis obtusis, filamentis basi puberulis, antheris glabris vel ad suturas parce lanatulis, stigmate flavidio retuso ♀. *P. ramosa* C.A. M. Enum. p. 104. — Ie. Rehb. Crit. 933-934. — Rehb. Germ. t. 152.

Hab. in cultis, *Cannabis*, *Fabac.*, *Nicotianæ* etc. parasitica, Macedonia (Friv!), Syria in Libano (Bl!), Mesopotamia variis locis (Haussk!), Egyptus (Gaill! Schweint!), Tauria et prov. Caucasicæ (Ledeb.).

β. Muteli. — Spicæ breviores densiores, corolla major magis violascens supra ovarium magis constricta et superne paulo magis dilatata. — *O. Muteli* Schultz in Mut. Fl. Fr. 2, p. 353, tab. 43. — *P. Muteli* Reut. in D.C. Prodr. p. 8. Ie. Rehb. Germ. tab. 150. — *P. emarginata* Reut. loc. cit. p. 9 et *P. Hohenackeri* Reut. loc. cit. p. 10. — *P. rufescens* Griseb. Spicil. p. 59! et *P. casia* Griseb. ibid. — *P. serotina* Haussk. Pl. exs. — *P. macrantha* C. Koch Linn. XVII, p. 290.

Hab. ad *Compositas*, *Leguminosas* et alias species ex ordinibus diversissimis parasitica typo multo frequentior in ditione totâ a Græciâ et insulis, Macedonia! ad Syriam! Egyptum! Arabiam petream!, regiones Caucasicæ! Persiam!

γ. nana. — Caulis simplex brevis, spica brevis densa pauciflora. — *O. nana* Noë exs.

Hab. in arenosis Syriae prope Sidonem (Gaill!).

Species florum magnitudine et spicis plus minus densis polymorpha sed specimina etiam inter varietates propositas non semper rite distribuenda.

Ar. Geogr. Europa omnis ab Angliâ et Belgio ad Rossiam medium. Africa borealis, insulae Canarienses, Abyssinia, P. bon. Spei.

13. *P. Aegyptiaca* (Pers. Encycl. II, p. 181 sub *Orobanche*) breviter glanduloso-papillosa, caulis elatis simplicibus vel sèpius ramosis, spicis longis laxis, floribus sessilibus vel pedunculatis, bracteâ mediâ oblongâ lateralibus anguste lanceolatis calyce brevioribus, calycis laciniis a basi trianguli acuminatis, corollâ calyce triplo longiore cœruleâ tubalesco-infundibuliformi arcuatâ lobis ovatis obtusis, antheris ad suturas lanatis, stigmate retuso ♀. *P. Aegyptiaca* Walp. Rep. III, p. 463. — *O. Delilei* Decaisn. Ann. 24, p. 201 et *P. Delilei* Walp. — *O. ramosa* Del. Eg. non L. — *O. pedunculata* Viv. Æg. Dec 22. — *O. Indica* Roxb. Fl. Ind. III, p. 27 et *P. Indica* Reut. in D.C. Prodr. ex parte. — *P. Aleppensis* Reut. MSS.

Hab. ad *Cucurbitaceas*, *Gossypium*, *Nicotianam*, *Solanum*, *Melongenam*, *Phlomides* et varias alias parasitica, Byzantium (Noë!), Cyprus (Ky. sub *P. longiflorâ*!), Egyptus (Boiss! Husson! Del!), Palestina (Boiss!), Syria borealis, Mesopotamia, Babylonia (Haussk! Noë!), Persia (Haussk!) et borealis (Buhse! sub *P. hirtiflorâ*), prov. Khorassan (Bge!), Afghanistan (Griff!), Belutschia (Stocks!).

Valde affinis *P. ramosa* a quâ tamen distincta videtur corollis duplo majoribus et longioribus, antheris lanatis. Hec est vera *O. Indica* Roxb. ex descrip-

tione et speciminibus, sed *P. ramosa* typica in Indiâ quoque crescit. Planta Georgica in d.c. Prodr. ad *P. Indicam* citata est *Ph. ramosa*!

Ar. Geogr. India orientalis, Algeria interior.

SECT. II. CISTANCHE Link. et Hoffm. Fl. Portug. I, p. 319. — *Hæmodorion* Wallr. Gen. Orob. p. 69. — Calyx semiquinquefidus rarius fissus lobis obtusis sepius æqualibus. Anthereæ undique lanatæ. Capsula stylo persistente terminata. Caulis simplicissimus.

Glabræ.

14. *P. lutea* (Desf. Atl. II, p. 60, tab. 146) glabra, caule crasso elato squamis lanceolatis crebre obsito, spicâ longâ densâ, bracteâ mediâ late ovatâ obtusâ calycem æquante vel sublongiore lateralibus oblongis minoribus calyce brevioribus omnibus carnosis margine membranaceis, calycis campanulati lobis ovatis obtusis margine scariosis erosulis, corollâ luteâ calyce $2\frac{1}{2}$ -plo longiore e basi angustatâ antrorum arcuatâ valde ampliatâ lobis ovato-orbiculariatis æqualibus patentibus, antheris albo-lanatis connectivo apiculatis, filamentis basi hirsutis, stigmate emarginato ♂.

Hab. *Chenopodiacearum* frutescentium parasitica, in Egypto (Sieb! Schweinf!), deserto Tib. Arabie Petreæ (Boiss!), Palestinaustrali prope Hebron (Roth!).

Ar. Geogr. Hispania australis, Africa borealis.

15. *P. tubulosa* (Schenk Plant. Spec. p. 23) glabra, caule crasso elato squamis lanceolatis crebre obsito, spicâ longâ densâ, bracteâ mediâ late lanceolatâ obtusiusculâ calycem subsuperante lateralibus brevioribus anguste linear-lanceolatis, calycis tubulosi lobis oblongis obtusis æqualibus, corollæ (e sicco violaceentis) tubo calyce duplo longiore apice incurvo in limbum modice dilatato lobis rotundatis obtusis, filamentis inferne valde hirsutis, antheris albo-hirsutis connectivo non apiculatis ♂.

Hab. inter Suez et Tor Arabie Petreæ (Roth et Erdl.), in valle Ouadi Useit ad *Tamarices* parasitica (Boiss!), in desertis salsis Persiae ad *Salsolaceas* parasitica prope Rischm et Kaschan (Buhse!), prope Ssertschah (Bge!). Floret Fehr, Mart.

Procerior *P. lutea*, siepe 2-3-pedalis, spica 1-1 $\frac{1}{2}$ -pedalis, corolla 15-16 lineas longa. Distinguitur a *P. lutea* formâ bractearum, corollæ tubo calycis proportione longiore et limbo minus dilatato, antheris muticis, colore. Stigma e corollâ exsertum ut in omnibus sectionis, sed tantum anthesi peractâ et corollâ jam subemarcidâ.

16. *P. flava* (C.A. Mey. Enum. p. 104) glabra, caule crasso squamis linear-lanceolatis obsito, spicâ compactâ densâ, bracteâ mediâ a basi oblongo-lanceolatâ in ligulam linearem corollâ multo longiore abeunte, lateralibus linear-lanceolatis calycem æquantibus, calycis tubulosi lobis rotundatis, corollâ flavâ calyce duplo longiore

tubuloso-campanulatâ incurvebâ lobis patentibus subæqualibus ovatis, filamentis basi hirsutis, antheris lanatis connectivo apiculatis ♀. *P. gigantea* Kar. Enum. in Bull. Mosq. 1839, p. 165.

Hab. in arenâ mobili ad littus Caspium ad Erzek inter Baku et Sallian (C.A. Mey!).

Corolla sesquipolligem longa, insignis bracteis mediis cum ligula 2 lineam latâ 2 pollices longis flores multum superantibus.

Ar. Geogr. Turcomania ad Caspium.

** Tomentosæ vel ciliatae.

17. **P. salsa** (C.A. Mey. Fl. Alt. II, p. 461) tomentoso-lanata, caule crasso inferne squamis ovato-oblongis obtusis crebre obsito, spicâ longâ densâ, bracteis tomentosis vel glabris margine ciliatis mediâ ovato-lanceolatâ obtusâ calyce sublongiore, lateralibus linearioribus calycem aequantibus, calycis glabri campanulati seniquinquefidi lobis ovatis obtusis margine sepe tomentellis, corollâ glabrâ violacea rectiusculâ calyce subtriplo longiore tubuloso-campanulatâ lobis subæqualibus patentibus semiorbiculatis, filamentis basi lanatis, antheris lanatis connectivo breviter vel obsolete apiculatis, stigmate suborbiculato ♀. Ledeb. Icon. tab. 376. — *P. incana* Payne Expl. Pal. p. 116! — *P. Armena* C. Koch Linn. XVII, p. 291.

Hab. in argilloso-salsis *Chenopodiacearum* parasita, mons Gebel Abiad deserti Syriae ad Palmyram (Bl!), Palestina trans Jordanum (Payne!), Persia prope Ispahan (Arch. 50°0!), inter Kerman et Yezd, Ispahan et Teheran (Bge!), ad Sson inter Ispahan et Kaschan (Bulise!), prov. Aderbidjan (Szov!), Armenia ad Araxem (C. Koch!). Turcomania ad mare Caspium (Karel!), Turkestania (Lehm!).

Ar. Geogr. Songaria, Sibiria Uralensis.

18. **P. fissa** (C.A. Mey. Enum. p. 104) tomentoso-lanata, caule crasso inferne squamis ovato-oblongis crebris dense obsito, spicâ longâ densâ, bracteâ mediâ lanceolatâ obtusâ calyce sesquilongiore lateralibus anguste linearibus calyci aequilongis, calycis tomentosi fissi lacinia superiore et posteriore linearis, inferiore et anteriore spathaeformi obtuse quadridentata, corollæ calyce 2½-plo longioris rectiusculæ tubuloso-campanulatæ lobis patentibus subæqualibus semiorbiculatis margine lanatis, filamentis basi barbatis, antheris connectivo apiculatis, stigmate orbiculari ♀.

Hab. prope Baku (C.A. Mey.). Hac refero quoque specimina a cl. Bayern in campis salsis ad iluv. Kara ditionis Elisabethpol in *Chenopodiaceum* parasitica locata.

Valde affinis *P. salsa* à quâ bracteâ mediâ lanceolatâ elongatâ et calycis structurâ (an normali?) differre videtur.

19. **P. ambigua** (Bge pl. Lehm. p. 429) caule glabrinculo squamis sparsis lanceolatis obsito, spicâ densâ, bracteis ciliatis intermedia lanceolatâ calycem excedente lateralibus linearibus obtusis

calyceum æquantibus, calycis hyalini campanulato-tubulosi subinæqualiter quinquefidi dentibus ciliatis subinæqualibus ovato-lanceolatis obtusis postico subbreviore acutiusculo, corollæ glabræ violaceæ tubulosæ rectiusculæ calyce triplo longioris lobis æqualibus rotundatis patulis, filamentis basi antherisque lanatis, connectivo apiculato, stigmate capitato ♀.

Hab. in desertis argillosis Kisilkum Turkestanice (Lehm!). Non vidi.

Ex specimine unico descripta, caulis quadripollicaris, spica tripollicaris, calyx vix dimidium pollicem longus, corolla 14-15 lineas longa; affinis *P. fissæ et salse*.

OROBANCHE (L. Gen. 779).

Flores hermaphroditi bracteâ fulti bracteolati. Calyx saepius disepalus sepalis integris vel bifidis, rarius sepalis basi inter se concretis gamosepalus quadrifidus. Corolla bilabiata ringens rarius recta. Stamina 4 corollæ tubo inserta didynama antherarum loculis basi mucronatis. Ovarium uniloculare placentis parietalibus 4 geminatis. Stigma retusum vel bilobum. Capsula unilocularis bivalvis valvulis apice et basi coalitis lateraliter dehiscentibus. Semina plurima reticulata. — Herbe parasiticæ. — Hujus generis ut et *Phelipeæa* species notis certis difficile limitantur; characteres enim graviores, filamentorum insertio et indumentum, antheræ glabræ vel hirtæ, etc., non semper sat firmi; alii v. gr. plantæ et stigmatis color, corollæ forma, exsiccatione plus minusve evanescunt. Notæ ex plantâ vivâ in loco natali depromptæ quam maxime necessariæ.

Specierum Orientalium distributio.

§ 1. Sepala antica et postica in calycem 4-5-fidum connata.

O. gamosepala, quadrifida, connata.

§ 2. Sepala libera rarius basi anticâ vel posticâ unilateraliter et obsolete connata.

* Filamenta versus corollæ basin vel usque ad quartam partem ejus tubi inserta.

+ Filamenta basi vel parte inferiori hirta vel villosa.

1. Stigma (an in omnibus?) luteum vel flavidum.

O. Anatolica, leucopogon, Stockii, Kurdica, cruenta, Sprunerii, Palestina, alpigena

2. Stigma (an in omnibus?) violaceum, purpureum, rarius albidum.

O. speciosa, leucantha, alba, pubescens, hyalina, Teuerii, Galii, Epithymum, punctata, cuprea, fuliginosa.

+ + Filamenta glaberrima

O. euglossa, Wiedemannii, glabrata.

++ Filamenta versus tertiam aut dimidiari tui corollini partem vel superius inserta.

+ Filamenta ad basin vel inferne hirta vel villosa.

O. rubens, loricata, amethystea, minor, Hederae, Cypria, curvipesca Kotschy, spectabilis.

+ + Filamenta glabra

O. cernua, poyonanthera, Cumana, camptolepis.

§ 1. Sepala antica et postica in calycem 4-5-fidum connata.

1. **O. gamosepala** (Reut. in D.C. Prodr. II, p. 15) caule rubro papilloso, spicæ brevi laxa, bracteis multinerviis obtuse acuminatis margine præsertim lanatulis corollâ dimidio brevioribus, sepalis plurinerviis latis antice et postice alte connatis subinaequaliter et breviter bitidis corollæ tubo duplo brevioribus, corollâ atrosanguineâ extus minute glandulosâ incurvo-areuatâ labiis ciliatis superiore multo majore galeato obtuso vix retuso inferioris lobis minutis rotundatis intermedio submajore, filamentis ad corollæ basin insertis fere glabris antheris magnis ad suturas lanatulis, stigmate cernuo bilobo e sicco flavido ♀.

Hab. in pratis humidis prope Karaes et ad radices montis Matschuka Georgie Caucasicæ (Hohen!), in pratis alpinis montis Antschalala ditionis Tindal Caucasi 7000'-8000' (Rupr!), in monte Karagiol (Szov. exs. N° 289!).

Caulis 3-5-pollicaris, corolla magnitudinis *O. Galii*. Species distinctissima calyce brevi monosepalo dentibus brevibus triangularibus, corollâ atrosanguineâ labio superiore magno galeato, filamentis glabris, antheris lanatis.

2. **O. quadrifida** (C. Koch Linn. XXIII, p. 665) crispule furfuraceo-hirsuta, spicæ laxa, bracteis lanceolatis plurinerviis corollâ subbrevioribus, sepalis latis connatis plurinerviis quadrifidis laciniis subelatis mediis corollæ tubum æquantibus lateralibus brevioribus, corollâ (e sicco albida) tubuloso-subcampanulatâ subrectâ apice subcurvatâ labio superiore submajore bilobo utriusque lobis denticulatis, filamentis paulo supra basin insertis parte inferiori villosis et fere totâ longitudine hirtis, stigmate.... ♀.

Hab. in ditione Caucasicæ Grusiæ (Schmidt!).

Mihi ex unico exemplari vermisibus fere destructo herb. cl. C. Koch tantum nota, spica semipedalis, corolla 8 lineas longa. Ex cl. auctore affinis *O. Hedera* et *O. Galii*; sepalis quadrifidis insignis.

3. *O. connata* (C. Koch Linn. XXIII, p. 669) caule glabrescente, spicâ breviter puberulâ laxâ, bracteis plurinerviis ovato-oblongis aoutis corollæ tubo longioribus, sepalis latis bifidis laciiniis triangularibus in unicum antice longe postice breviter concretis, corollâ minute glandulosâ e siccо fusco-rubellâ strictâ tubulosâ rectâ labiis parvis denticulatis superiori bilobo lateraliter reflexo inferiore obtuse trilobo, filamentis versus tubi tertiam partem insertis basi crebre et breviter villosis, antheris glabriusculis, stigmate bilobo ♀. *O. cerulescens* C. Koch Linn. XVII, 291 non Steph.

Hab. in Armenia Rossica prope Eriwan in planitie secus Araxem (C. Koch!).

Ex specimine unico valdeque incompleto descripta tamen ut videtur species distincta, corolla 7 lineas longa stricta fere exacte tubulosa. Inter species sepalis connatis donatas *O. quadrifida* differt indumento longo furfuraceo, calycis laciiniis subulatis longioribus; filamentis totâ longitudine hirsutis; *O. gamosepala* corollæ incurvo-arcuatâ, antheris lanatis etc.

§ 2. Sepala libera, rarius basi antice vel postice unilateraliter subconnata.

• Filamenta versus basin corollæ aut paulo superius inserta.

+ Filamenta basi vel inferne hirta vel villosa.

4. ***O. luteum*** (Boiss. et Rent. in Pinard pl. exs. p c. Prodr. II, p. 17) tota ad scapum squamas bracteas et calyces tomento albo-araneoso plus minus vestita, caule crasso squamis imbricatis ovatis acutis obsito, spicâ densissimâ oblongo-cylindricâ, bracteis late oblongo-lanceolatis acutis corollam subæquantibus, sepalis plurinerviis ovatis ad medium vel tertiam partem bilobis antice coalitis corollâ duplo brevioribus, corollâ coriacea breviter campanulatâ parum arcuatâ intense purpureâ glandulis aspersâ labio superiore galeato inferioris lobis brevibus obtusis, filamentis ad corollæ basin insertis inferne hirtis rarius glabratis, stigmate luteo ♀. *O. colorata* C. Koch Linn. XVII, p. 291.

Hab. *Salvia Cedronelle* et congenorum parasitica, Anatolia borealis ad Kadikios (Wied!), mons Cadmus Carie supra Colossam (Boiss!), Lycia supra Elmalu (Bourgl.), Pisidia in collibus Buldur (Heldr!), Cappadocia ad Argæum (Ky!), Cilicie littoralis faux Guzeleré (Bal!), mons Anmanus Syriae borealis (Ky. 291), Armenia borealis (C. Koch!) et australis ad Bakker Maaden (Noët!), Pontus (Tehih!), mons Sariel Georgiae Caucasicæ (Hohen!), mons Pir Omar Gindruan Kurdistaniæ Persicæ (Haussk!).

Species insignis lanugine arachnoideâ, caulis crassus crebre squamosus, corolla coriacea 9-10 lineas longa lata.

5. *O. leucopogon* (Boiss. et Hausskn. exs. 1865) griseo-lanuginosa, spicâ densiflorâ, bracteis plurinerviis lanceolatis acutis sepalisque oblongis plurinerviis bifidis subulatis corollam subæquantibus, corollâ rubente extus parce glandulosâ tubuloso-infundibuli-

formi superne vix incurvâ, labiis ciliatis intus albo-lanuginosis superiore retuso inferioris lobis obtusis intermedio submajore, filamentis infrâ quartam tubi partem insertis ibique dilatatis et hirtis, stylo glanduloso, stigmate magno subexerto ♀.

Hab. ad *Carduos* parasitica in monte Ssodagh Cataoniæ 400' (Haussk!).

Spica semipedalis, indumentum lanuginosum *O. Anatolice*, insignis corollæ labiis intus dense albo-lanuginosis.

6. O. Stocksii, minute pruinosa, caule squamis breviter oblongis obsito, spicâ densâ ovatâ brevi, floribus magnis, bracteis multinerviis ovalis acentiuseulis corollâ quadruplo brevioribus, sepalis oblongis breviter bifidis bracteæ aequilongis, corollâ e secco albâ minute pulverulento-pruinosa late campanulatâ rectâ obtuse bilabiata lobis integris late semiorbiculatis, filamentis versus tubi quartam partem insertis basi valde dilatatis et hirtis, antheris barbatis, stigmate magno bilobo ♀.

Hab. in *Eryngii* sp. parasitica ad Douboud Affghaniae (Stocks!).

Caules semipedales, spica junior capitata demum bipollucaris; insignis corollâ sesquipollucari latâ a basin sensim et modice dilatata e secco obsolete labiatâ lobis subaequalibus margine integris, antheris barbatis. Species inter omnes congeneres corollâ maximâ donata.

7. O. Kurdica (Boiss. et Haussk) parce et breviter papilloso-pruinosa, caule squamis longis oblongo-linearibus obtusiusculis obsito, spicâ elongatâ densiusculâ, floribus majusculis, bracteis a basi latiori linear-lanceolatis obtusis corollas subaequantibus, sepalis 4-5-nerviis lanceolatis bifidis tubo corollæ longioribus, corollâ minute pruinosa pallide carneâ late tubuloso-campanulatâ apice vix curvatâ labiis repandis non ciliatis superiore bilobo inferiore trilobo lobis brevibus obtusis, filamentis paulo supra basin insertis inferne dilatatis breviter hirtis, stigmate luteo ♀.

Hab. ad *Phlomides*, *Tanacetum*, *Carduaceas* parasitica in montibus Avroman et Schahu Kurdistaniæ Persicæ 8-1000' (Haussk!).

Caules crassiusculi, spica semipedalis, corolla fere pollicaris ut et bracteæ et calyces exsiccata membranacea.

8. O. cruenta (Bertol. rar. Ital. Dec. III, p. 56) pubescenti-glandulosa, spicâ laxiusculâ, bracteis acutis corollæ tubum æquantibus vel superantibus, sepalis plurinerviis subaequaliter bifidis corollæ tubum superantibus, corollâ campanulatâ arcuatâ brunneo-flavâ intus eruentâ basi ventricosa labio superiore galeato integro vel subretuso inferioris lobis denticulatis glanduloso-fimbriatis, filamentis basi corolle insertis basi hirtis et superne glandulosis, antheris glabris, stigmate bilobo luteo sëpe purpureo-marginato ♀ Rehb. Crit. VII, fig. 896. — Rehb. Ic. Germ. t. 159. — *O. gracilis* Sm. Trans. Linn. 4, p. 172

Hab. ad Leguminosas parasitica in Argolide circâ Mycenæ (Heldr!), mon-

tibus Macedonie (Griseb.), Bithyniâ circâ Brussam (Fritsch!), Transcaucasia in Kachetiâ (Hohen. ex Ledeb.).

Ar. Geogr. Europa tota media et australis a Galliâ australi et Germaniâ mediâ et australi ad regionem Danubialem.

9. **O. Spruneri** (F. W. Schultz Flora 26, p. 130) papillari-pubescentia, spicâ densiusculâ, bracteis late lanceolatis calyce longioribus, sepalis plurinerviis late lanceolato-acuminatis indivisis vel bifidis corollæ tubum æquilibus, corollâ breviter et late campanulatâ flavo-rubrâ intus et limbo purpureâ basi ventricosâ arcuatâ, labio superiore majore galeato retuso inferioris lobis patentibus denticulatis, filamentis ad basin corollæ insertis basi villosis superne glandulosis, antheris glabris, stigmate bilobo sordide flavo ♀. *O. crassistyla* Schultz loc. cit. p. 131.

Hab. parasitica ad *Cistos* in Græciâ ad Naupliam (Sprun!), in monte Pentelicô Atticae (Heldr! Boiss!), monte Parnes (Heldr!), ad Kumi Eubeæ (Heldr!).

Affinis *O. cruentæ*, corolla major.

10. **O. Palestina** (Reut d.c. Prodr. XI, p. 718) parce papillosa, spicâ longissimâ laxiusculâ, bracteâ a basi oblongâ sensim linearisubulata corollam superante, sepalis paucinerviis late oblongis integris vel bifidis laciniis linearisubulatis corollæ tubo longioribus, corollâ glabriuscûlâ e sicco albidiâ tubulosâ brevi arcuatâ labiis parvis subæqualibus superiore subulato inferioris lobo medio submajore, filamentis versus quartam tubi partem insertis inferne latis et basi hirtis, stigmate (e sicco) flavescente ♀.

Hab. in Palestinâ circâ Sichem et Banias (Boiss!). Hierosolymæ (Roth exs. 494!), ad *Physocaulum nodosam* et *Anthriscum macrospermam* prope Marasch Syriæ (Haussk!).

Spica interdum pedalis, flores eis *O. minoris* submajores. Species spicis longis et bracteis elongatis peculiaris, facie Rehb. Germ tab. 168. *O. pallidifloram* Wimm. referens, sed haec differt bracteis et sepalis brevioribus, filamentis paulo supra basin insertis, stigmate rubro.

11. **O. alpigena** (C. Koch Linn. XXIII, p. 668) minute et breviter glanduloso-hirta, spicâ oblongâ densiusculâ, bracteis plurinerviis ovato-oblongis attenuato-acuminatis corollæ tubo subbrevioribus, sepalis paucinerviis indivisis corollæ tubo æquiblongis. corollâ minute glandulosâ tubulosâ apice vix subincurvâ labiis parvis denticulatis superiori retuso inferiore breviter trilobo, filamentis versus quartam tubi partem insertis basi hirtis, antheris barbulatis, stigmate flavidō ♀.

Hab. in Ponto Lazico prope Djimil (C. Koch!), ibidem *Betonicæ grandifloræ* parasitica (Bal!).

Spicæ 1½-2½-pollicares, corolla 7-8 lineas tantum longa. Affinis *O. rubenti* a quâ sepalis indivisis, antheris barbatis, corollâ vix curvatâ differt.

2. Stigma violaceum, purpureum vel albidum.

12. **O. speciosa** (d.c. Fl. Fr. Suppl. p. 393) crispule papillosa

caule saepe procero, spicâ longâ inferne laxiusculâ, bracteis lanceolatis subulato-acuminatis corollæ tubo longioribus, sepalis quinque-nerviis divergentibus in lacinias acuminatas corollæ tubo aequilongas saepius bifidis, corollâ campanulatâ arcuatâ albidâ cæruleo-striatâ siccâ papyraceâ pallide fusca labio superiore bilobo inferioris lobo medio majore omnibus denticulatis ciliatis rotundatis, filamentis paulo supra basin corollæ insertis basi pubescentibus superne glandulosis, stigmate pallide violaceo ♂. Rehb. Ic. Germ. tab. 461. — *O. pruinosa* Lap. Suppl. 87. Rehb. Crit. fig. 911. — *O. grandiflora* Bor. et Ch. Fl. Pel. № 983, tab. 23. — *O. segatum* Sprun., C. Koch Linnae. XIX, p. 26.

Hab. in *Vicia*!, *Ervo*, *Fabâ*, aliisque plantis parasitica in totâ ditione vulgaris a Greciâ ad Syria!-, Mesopotamiam!, Egyptum!, Transeaucasiam!

Ar. Geogr. Regio mediterranea totius Europa et Africæ borealis.

13. *O. leucantha* (Griseb. Spieil. II, p. 57) palmaris, pubescens, spicâ brevi laxiusculâ, bracteis lanceolatis corollæ tubum æquantibus, sepalis 1-2-nerviis subulatis vel bifidis, corollæ albæ tubo rectiusculo limbo patente, filamentis paulo supra basin sitis ad punctum insertionis breviter luteo-lanatis cæterum glaberrimis, stigmate violaceo ♀. *O. alba* Rehb. Crit. fig. 912 ex Grisebach excl. stigmatis colore.

Hab. in peninsula Athoâ prope Karaes in Castanetis umbrosis 2000' (Griseb.). Non vidi.

Affinis videtur *O. alba* et formis parvifloris *O. speciosæ*. Planta Bannatæ Rehb. Germ. tab. 109, filamentis hirsutis glandulosisque, etc., diversa videtur.

14. *O. alba* (Steph. in Willd. Sp. III, p. 350) breviter glanduloso-hirta, spicâ brevi laxiusculâ, bracteis lanceolatis corollæ tubo brevioribus, sepalis integris acuminatis bracteâ brevioribus vel aequilongis, corollâ pallidâ albâ purpureo-suffusâ campanulatâ dorso breviter curvatâ labio superiore transverse latiore retuso inferioris lobis subæqualibus obtusis, filamentis paulo supra basin insertis basi hirsutis superne glanduliferis, stigmate purpurascente-bilobo ♀. Ic. M.B. Cent. tab. 52. — Rehb. Germ. 215.

Hab. in graminosis Armenie circa Erzerum (Calv!) et Gumuschkhané (Bourg!), Tauria et Caucasi (M.B., C.A. M.), Georgiae Caucasicæ (Hohen!), tractus Swant (C.A. M.).

Species mihi e sicco obseura, flores illis *O. speciosæ* minores. Icon Rehb. Crit. fig. 912 ex cl. Rehb. fil. huc non spectat et ex cl. Grisebach suam *O. leucantham* sistit.

Ar. Geogr. Transylvania, Turkestania, Songaria, Sibiria Altaica.

15. *O. pubescens* (Urv. Enum. p. 76) tota crispula albo-lanuginosa, spicis longis densiuseulis, floribus parvis, bracteâ ovato-lanceolatâ acuminatâ corollam æquante, sepalis bi vel obsolete plurinerviis integris vel bifidis corollæ tubum superantibus, corollâ sublanu-

ginosâ flavidò-rubellâ vel lilacinâ breviter tubuloso-campanulatâ rectiusculâ vel arcuatâ labiis irregulariter denticulatis superiore subintegro inferioris lobis rotundatis, filamentis paulo suprâ tubi basin insertis inferne lanuginoso-pilosus, stigmate lilacino ♂? Ic. Rehb. Germ. tab. 180. — *O. versicolor* Schultz Fl. 28, p. 129. — *O. villosa* et *O. arachnoidea* Schultz in litt. — *O. fragrans* Griseb. Spic. II, 58 non Koch.

Hab. ad *Cichoraceas*, *Chrysanthemum*, *Orlayam*, *Tordylium Apulum*, *Calendulus*, etc. vulgatissima, in Græciâ et ejus insulis (Urv! Boiss!, Heldr! Orph. Fl. exs 21!), Macedonia (Griseb.), Byzantii (Cast!), Anatoliâ occid. et australi maritimis (Pin! Heldr!), Cretâ (Heldr!), Syriâ et Mesopotamiâ (Haussk!), Palestinâ boreali (Boiss!).

Species elegans facie *O. minorem* referens sed albo-lanuginosa, spicæ elongatæ, labium superius non bilobum, filamentorum insertio diversa, etc.

Ar. Geogr. Gallia australis.

16. *O. hyalina* (Sprun. in D.C. Prodr. XI, p. 24) crispule albo-papillosa, caule nano squamis apice subulatis crebre obsito, spicâ brevissimâ pauciflorâ ovatâ, bracteâ lanceolatâ flores æquantî, sepalis subuninerviis simplicibus a basi lanceolatâ longe subulatis tubo corollæ longioribus, corollâ minutâ in sicco hyalinâ tubuloso-campanulata subrectâ labiis parvis irregulariter denticulatis superiore integro inferioris lobis rotundatis æqualibus, filamentis paulo supra baain insertis totâ parte inferiori dilatatis hirtis, stylo parce glanduloso, stigmate violascente? ♂.

Hab. in arenosis maritimis insulæ Æginæ Atticæ rarissime (Sprun!), insulâ Cephalonîa prope Taphious (Heldr!).

Planta 4-5-pollicaris facie formam pumilam *O. minoris*, *O. pumilam* Noë, omnino referens, ab eâ filamentis prope basin corollæ insertis discedens; an ab *O. pubescens* spicâ brevi pauciflorâ, filamentis basi breviter hirtis sat distincta? *O. Salisii* Req. e Corsicâ hûc ab auctoribus Gallicis et Rehb. Ic. Germ. tab. 165 relata ob filamenta altius inserta est potius *O. minoris* forma.

17. *O. Teucrii* (F. W. Schultz Fl. 1835, p. 200) crispule glanduloso-hirta, spicâ laxiusculâ, bracteis lanceolatis acutis corollâ sub-brevioribus, sepalis subæqualiter bifidis apice subulatis antice subconnatis corollæ tubum dimidium æquantibus, corollâ fusco-rubrâ glandulosâ tubuloso-campanulatâ dorso rectiusculâ labio superiore galeato declivi integro vel retuso inferiori deorsum flexo trilobo utroque denticulato, filamentis supra corollæ basin insertis ad medium usque pilosis, stigmate fuscescenti sanguineo ♂. Ic. Rehb. Germ. tab. 169. — *O. atrorubens* Schultz Flora 1840, p. 128. — *O. marginata*, *O. hians* et *O. Lazica* Reut. in Bal. et Huet exs.

Hab. *Teucrii chamaedrys* parasitica in valle Djimil Ponti Lazici (Bal!), Imeretiâ (Szov!).

Ar. Geogr. Gallia, Helvetia, Germania.

18. *O. Galii* (Duby Bot. Gall. p. 349) glanduloso-pubescent, spicâ laxâ, bracteis lanceolatis acuminatis flores æquantibus vel su-

perantibus, sepalis plurinerviis antice coalitis bifidis corollæ dimidiatum tubum æquantibus, corollæ albo-carneæ glanduloso-pubescente campanulatâ arcuatâ a basi ad apicem sensim ampliatâ, labii superioris lateribus porrectis inferioris lobis subæqualibus denticulatis, filamentis paulo supra basin insertis pilosis superne ut et stylus glanduloso-pilosus, stigmate bilobo purpureo non marginato ♀. *O. caryophyllacea* Sm. Act. Linn. IV, p. 169. Rehb. Crit. fig. 890, 891. — Ie. Germ. tab. 162. — *O. vulgaris* D.C. Fl. Fr. III, 489. Rehb. Crit. 808-810. — *O. strobiligena* Rehb. Crit. fig. 905-905.

Hab. ad *Galia* parasitica in Tauriâ et prov. Caucasicis (M.B.), Georgia Caucasicâ (Hohen.), probab. in Macedoniâ.

Ar. Geogr. Europa media et australis ab Anglia, Belgio et Germaniâ ad Rossiam mediâ, Sibiria omnis.

19. *O. Epithymum* (D.C. Fl. Fr. III, p. 490) glanduloso-pubescentes, spicâ brevi laxâ, bracteis lanceolatis acuminatis corollæ labium inferius æquantibus vel superantibus, sepalis lanceolatis subulato-acuminatis tubo corollæ longioribus subindivisis, corollæ lutescenti rubro-suffusâ pilis glandulosis e tuberculo ortis hirtâ campanulato-tubulosâ subarcuatâ labiis inæqualibus acute dentalis margine crispatis superioris biloibi lobis patentibus inferioris lobo intermedio longiore, filamentis prope basin insertis inferne parcissime hirtis superne glandulosis, stigmate atropurpureo ♀. Rehb. Crit. f 885-888. — Rehb. Germ. tab. 163. — *O. cruenta* var. *leiosemon* Griseb. Spic. II, p. 55 ex Rent. — *O. ferruginea* C. Koch Linn. XVII, p. 291. — *O. psilandra* C. Koch Linn. XXIII, p. 668 !

Hab. ad *Thymos*, *Zizyphoram*, *Satureiam* parasitica, Lycabetus Atticæ (Sart!), Thraciâ (Griseb!), Zacinthus (Marg!), prope Byzantium (Courmany!). Armenia Turcica (Huet! Bourg!), prov. Transcaucasica Karabagh (Hoh. ex Ledeb.).

♂? *Buhsei*. — Sepala bifida. — *O. Buhsei* Reut. in Boiss. et Buhse Aufz. p. 169.

Hab. in Armeniâ Rossicâ prope Nackitschewan *Thymi serpylli* parasitica (Buhse!).

Ar. Geogr. Europa media et australis a Belgio et Germaniâ mediâ ad Rossiam mediâ.

20. *O. punctata* (F. W. Schultz loc. cit. p. 131) papillari-hirta, spicâ densiusculâ, bracteâ late lanceolatâ acuminatâ calyce longiore, sepalis paucinerviis anguste lanceolato-subulatis corollæ tubum æquantibus, corollæ ochrolencâ nervoso et punctato-violaceâ campanulatâ arcuatâ labiis denticulatis inferioris lobo intermedio longiore, filamentis prope basin insertis et ibi sparsim pilosis apice subglandulosis, stigmate pallide violaceo ♀.

Hab. in monte Lycabeto Atticæ ad *Cistos* parasitica (Sprun., Heldr!).

Affinis *C. Epithymo* spica densior, corolla major.

21. **O. cuprea** (Boiss. et Bal. exs. 1855) breviter glanduloso-papillosa tota fuscescenti-rubra, caule squamis oblongo-lanceolatis obtusis obsito, spicâ longâ laxâ, bracteis a basi late lanceolatâ longe attenuatis subulatis corollam æquantibus, sepalis a basi latiori longe subulatis indivisis corollæ subæquilongis, corollâ tubuloso-campanulatâ dorso medice curvatâ labiis subæquilongis acute denticulato-glandulosis superiore retuso inferioris lobis obtusis intermedio submajore, filamentis paulo supra basin insertis inferne dilatatis et valde hirtis, stigmate albo ♀.

Hab. *Origani Onitis* et *Satureiae Thymbræ* parasitica ad Bouloukli et in fauce Guzel Dere supra Sedichig Ciliciæ littoralis (Bal!).

Spica semipedalis vel fere pedalis, corolla 9 lineas longa, facies *O. hederæ* floribus minoribus et characteribus ceterum diversissimæ. Affinis *O. Epithymo* spicâ brevi, stigmate atropurpureo, etc. distinctæ.

22. **O. fuliginosa** (Rent. in D.C. Prodr. XI. p. 23) caule glabrescente minute squamoso, spicâ laxiusculâ. bracteis breviter pubescenti-glandulosis a basi ovatâ subulato-attenuatis corollâ subbrevirib. sepalis a basi ovato-oblongâ longe subulatis integris vel bifidis plurinerviis corollæ tubo æquilongis vel longioribus, corollâ breviter glanduloso-hirtâ violaceo-brunneâ tubuloso-campanulatâ superne galacto-fornicatâ labiis undulato-plicatis obtuse denticulatis superioris bifidi lobis porrectis inferioris æqualibus ovalis acutis. filamentis supra basin corollæ insertis inferne villosis superne parce glandulosis, stigmate violaceo ? ♀. Ic. Jord. Fragni. 3, tab. 9, fig. B.

Hab. in monte Hymetto Atticæ *Orobi* et *Cephalariae* parasitica (Sprunt! Sart!) insulâ Rhodo prope Bastida (Bourg!).

Valde affinis *O. Scabiosæ* Koch a quâ labii inferioris lobis acutis differre videtur.

Ar. Geogr. Galloprovincia.

++ Filamenta glaberrima.

23. **O. euglossa** (Rehb. fil. Ic. Fl. Germ. XX, p. 117, tab. 217) pumila, caule tenui squamis breviter lanceolatis obsito, spicâ brevisimâ pauciflorâ, bracteis lanceolatis acuminatis flores æquantibus, sepalis 4-3-nerviis lanceolatis acuminatis curvulis indivisis tubo corollino longioribus, corollâ parvâ parum curvatâ labiis parvis denticulatis superiore bilobo inferiore subæqualiter trilobo, filamentis glaberrimis supra basin insertis, stigmate violascente ? ♀.

Hab. *Hieracii Piloselle* parasitica Byzantii (Noë). Non vidi.

Ex iconè valde affinis videtur *O. hyalinae* a quâ labio superiore bilobo et filamentis glabris an satis differt.

24. **O. Wiedemannii**. glanduloso-hirta, caule pumilo tenui angulato squamis numerosis linearibus obsito, spicâ ovato-oblongâ densiusculâ pauciflorâ, floribus parvis bracteis sepalisque lanceolatis

longe subulato-acuminatis plurinerviis corollæ tubo longioribus hisce præter infima interdum bifida indivisis, corollâ intense purpureâ parce glanduloso-hirtâ anguste tubuloso-campanulatâ dorso parum curvatâ labiis parvis subæqualibus superiore bilobo porrecto inferioris intermedio submajore omnibus rotundatis denticulatis non ciliatis, filamentis paulo supra basin insertis glaberrimis, antheris glabris, stigmate bilobo e sicco flavido.

Hab. in Anatoliâ boreali prope Kastamonii (Wiedem!), in Daghestaniâ mediâ prope Tindi 55°0'-60°0' (Rupr!).

Planta 4-5-pollicaris spicâ 4-4½-pollicari, corolla 7 lineas longa. Species insignis colore intense purpureo, filamentis glaberrimis, sepalis fere semper indivisis.

25. O. glabrata (C. A. Mey. in Eichw. Casp. p. 16, tab. 17) glanduloso-hirta, caule humili squamis linearibus obsito, spicâ densâ brevi, bracteis linear-lanceolatis acutis corollam æquantibus, sepalis subintegris lanceolatis longe acuminatis uninerviis corollam dimidiâm æquantibus, corollâ glandulâ raris brevissime stipitatis adpersâ in sicco obscure rufâ campanulatâ rectâ labiis crispato-denticulatis superiore obsoiete et latissime bilobo inferioris subbrevioris lobis orbiculatis obtusis intermedio sublongiore, filamentis paulo supra basin insertis inferne glaberrimis, antheris glabris, stigmate retuso ♀.

Hab. in Transcaucasiæ ditione Kachetiâ (Eichw.), prov. Ghilan Persiæ (Jenisch!), ad *Cousinia* radices in monte Kellal ditionis Bachtiaris Persiæ (Haussk!).

Caulis crassitie pennæ corvinæ 4-5-pollicaris, corolla 7-8 lineas longa. Species ad *C. Galii* facie accedens, corollâ fere rectâ et filamentis basi glaberrimis a congeneribus distincta.

• Filamenta versus ¼-½ partem tubi corollini vel superius inserta.

+ Filamenta basi vel inferne hirta vel villosa.

26. O. rubens (Wallr. Orob. 46) glanduloso-pubescentia elata, spicâ longâ laxiusculâ, bracteis lanceolatis acuminatis plurinerviis corollam æquantibus vel subbrevioribus, sepalis plurinerviis ovatis inæqualiter bifidis corollæ tubum dimidium æquantibus antice contiguis vel coailitis, corollâ lutescenti-fuscâ vel rubellâ e basi curvatâ tubuloso-campanulatâ dorso rectâ labio superiore galeato declivi bilobo lobis patentibus, inferiore subæqualiter trilobo, filamentis versus tertiam tubi partem insertis a basi ad medium pilosis, stigmate cernino-flavo ♀. Ic. Rehb. Germ. t. 171. — *O. elatior* Rehb. Pl. Crit. fig. 901-902.

Specimen a cl. Szovits ex Transcaucasiâ loco non indicato in herb. Hort. Petrop. hinc referto. In radice *Medicaginum* parasitica.

Ar. Geogr. Europa media a Galliâ et Germaniâ ad Dalmatiam, Rossiam medianam et australem.

27. **O. loricata** (Rchb. Fl. exc. p. 335 excl. synon.) breviter pubescens, spicâ demum laxâ, bracteis oblongo-lanceolatis corollæ æquilongis, sepalis 3-5-nerviis in lacinias subulatas bipartitis corollæ tubum æquantibus, corollâ flavidâ rubello-striatâ tubuloso-campanulatâ inferne rectâ superne tantum arcuatâ labio superiore patule bilobo inferioris lobis æqualibus obtuse denticulatis, staminibus sub medio tubo insertis parte inferiori pilosis, stigmate violascenti ♀. Ic. Rchb. Crit. fig. 917. — *O. Artemisiæ* Vauch. — *O. caryophyllacea* C. A. Mey. Enum. ex Ledeb.

Hab. in ditione Transcaucasica Talyisch (C.A. Mey.). Non vidi et mihi valde dubia.

Ar. Geogr. Gallia australis, Helvetia, Germania, Italia borealis, Hungaria.

28. **O. amethystea** (Thuill. Fl. Par. Ed. II, 1, p. 317) glanduloso-pubescentia, caule viotaceo squamis anguste lanceolatis obsito, spicâ densiusculâ longâ, bracteis pubescentibus lanceolato-linearibus corollam æquantibus, sepalis 5-7-nerviis integris et bifidis a basi ovatâ subito subulato-angustatis, corollâ albida lilacino-suffusâ glanduloso-hirtâ tubulosâ a tertiatâ parte inferiore subito in genu curvatâ, labiis undulato-crispis acute denticulatis superiore porrecto retuso vel subquadriolo inferioris lobis subbifidis intermedio majore, filamentis paulo sub medio tubo insertis inferne parce pilosis, stigmate fusco-purpureo ♀. Ic. Rchb. Germ. tab. 185. — Rchb. Crit. fig. 920-921. — *O. Eryngii* Duby. — *O. Attica* Reut. in D.C. Prodr. pag. 28.

Hab. *Eryngii* specierum parasitica, Græcia ad littus prope Amphissa (Heldr!), in collibus maritimis Ponti ad Cerasuntem (Tchich!).

Facies *O. minoris*, bracteæ basi angustiores, corolla longior subito in genu flexa, filamentis altius insertis.

Ar. Geogr. Belgium, Gallia, Germania occid. et australis, Anglia, Hispania, Africa borealis.

29. **O. minor** (Sutt. Trans. Linn. IV, p. 178) tenuiter pubescenti-glandulosa, caule gracili, spicâ densâ vel laxâ, bracteis pubescenti-lanuginosis a basi ovatâ acuminatis corollam æquantibus vel subsuperantibus, sepalis plurinerviis integris vel bifidis a basi ovatâ subito subulato-contractis tubum corollæ æquantibus vel superantibus, corollâ parvâ albida lilaceo-suffusâ tubulosâ arcuatâ labiis undulatis obtuse denticulatis superiore bilobo porrecto inferioris lobis subrotundis subæqualibus, filamentis paulo sub medio tubo insertis basi parce pilosis, stigmate purpureo-violaceo ♂. Ic. Rchb. Germ. tab. 183. — Rchb. Crit. fig. 876-879. — *O. Grisebachii!* — *O. striata!* Reut. in D.C. Prodr. II, p. 28. — *O. condensata* Griseb. Spicil. 2, p. 56 non Moris.

Hab. ad *Trifolium*, *Coronillas*, *Cichoraceas* parasitica ad littora et in interioribus, Cephalonia (Heldr!), Attica (Orph!), Byzantium (Coum!), Pamphylia et Pisidia (Heldr!), Abchasia (Nordm. ex Ledeb.).

Ar. Geogr. Europa media et australis ab Angliâ et Dania ad regionem Danubialem, Dalmatiam, Africa borealis.

30. **O. Hederæ** (Duby Bot. Gall. 350) breviter glandulosopubescent, spicâ longâ gracili laxiusculâ, bracteis ovato-lanceolatis acuminatis enerviis corollam aequantibus violaceis, sepalis violaceis subuninerviis antice coalitis lanceolatis subulatis integris vel bifidis corollæ tubum aequantibus vel longioribus, corollâ flavâ violaceo-suffusâ glabriuscûlâ tubulosâ dorso arcuatâ labiis denticulatis non ciliatis supericie retuso inferioris lobo medio majore, filamentis paucis supra tertiam tubi partem insertis basi puberulis, stigmate flavo ♀. Ic. Vauch. Mon. tab. 8. — Engl. Bot. tab. 2859. — Rehb. Germ. tab. 182. — *O. Medicaginis* Rehb. Pl. Crit. fig. 919 et Griseb. Spieil. p. 57 non Duby.

Hab. ad *Hederæ* radices parasitica, Peninsula Athoa prope Karaes (Griseb.), Pontus Lazicus in valle Djimil (Bal!).

Facies *O. minoris* a quâ sepalis uninerviis, stigmatis colore, etc. differt.

Ar. Geogr. Europa media et australis ab Angliâ et Belgio, Germaniâ ad Dalmatiam.

31. **O. Cypria** (Reut. in Kotschy Cypr. p. 204) papilloso-viscida, caule nano squamis crebris membranaceis oblongo-triangularibus obsito, spicâ densâ oblongâ pauciflorâ, floribus parvis, bracteis ovatis acutis corollæ tubum aequantibus, sepalis oblongis plurinerviis apice breviter bifidis tubum corollæ superantibus, corollâ e sicco rubellâ tubuloso-cylindraceâ apice tantum curvatâ parce glandulosâ labiis parvis denticulato-fimbriatis superioris lobis binis latis retusis inferioris brevibus ovatis, filamentis basi dilatatis versus tubi tertiam partem insertis inferne hirtulis, stylo apice glanduloso, stigmate bilo ♀.

Hab. ad radices *Pterocephali multiflori* et *Scutellarie hirtæ* in insula Cypro ad Prodromo 4000' (Ky. exs. 854). Fl. Maio.

Plantula cum spicâ pollicari 2-3-pollicaris, flores e minimis, corolla 5 lineas longa superne non ampliata. Facie refert *O. camptolepidem* sed bracteæ nec coriaceæ nec hamatæ, corolla apice curvata nec recta, non ampliata, lobis valde denticulato-fimbriatis.

32. **O. cærulescens** (Steph. in Willd. Sp. III, p. 349) albo-lanuginosa, caule crasso, spicâ densâ compactâ, bracteis oblongo-lanceolatis acuminatis corollæ tubo longioribus, sepalis indivisis vel ad tertiam partem divisis tubo corollæ longioribus, corollâ cærulescente laxe lanuginosâ tubulosâ subbincurvâ labiis denticulatis superiore retuso inferioris lobis rotundatis concavis, filamentis supra medium tubum insertis basi parce pilosis, stigmate albido-cærulescente ♀. Ic. Rehb. Crit. fig. 925. — Rehb. Germ. tab. 188. — *C. ammophila* C. A. M. Fl. Alt. II, p. 454 Ledeb. Ic. tab. 389 non differre videtur.

Hab. *Artemisiarum*, *Medicaginiis* parasitica, Anatolia borealis ad Safranbol (Wied!), Transcaucasiæ ditio Guria (Nordm. ex Ledeb.), Iberia prope Tiflis (Rupr!).

Lanâ adpressâ griseo-albâ insignis, facies *Phelipeæ*.

Ar. Geogr. Germania, Rossia australis, Songaria, Sibiria Altaica.

33. O. Kotschyî (Reut. in D.C. Prodr. XI, p. 33) crispule piloso-araneosa, caule crassiusculo squamis linearibus obsito, spicâ densiflorâ, bracteis a basi latiore anguste linear-i-lanceolatis corollam æquantibus vel superantibus, sepalis nervosis ad medium vel basin in laciniis linear-i-subulatas dimidiâ corollam æquantes bipartitis, corollâ hirtulâ pallide flavâ tubuloso-infundibuliformi subincurvâ extus parce papillosâ labiis parvis subrepandis non ciliatis superiore bilobo inferioris lobis ovatis obtusis, filamentis supra medium tubum insertis basi hirtulis, antheris ad suturam lanatis, stigmate bilobo ♀.

Hab. in Persiâ australi (Ky. 891!), ad *Umbelliferas* in regione alpinâ montis Sawers 12000' (Haussk!).

Semipedalis et pedalis, corolla 8-9 lineas longa. Ex affinitate *O. pogonantheræ* a quâ differt bracteis angustioribus et multo longioribus, corollâ majori hirtâ superne magis ampliatâ.

34. O. spectabilis (Reut. in Boiss. et Buhse Aufz. p. 166) tota pilis albis crassis lanato-canescens, caule robusto squamis lanceolatis carnosulis sparsim obsito, spicâ longâ densâ cylindricâ, bracteis lanceolato-acuminatis corollam subæquantibus, sepalis latis plurinerviis ad vel ultra medium bipartitis lobis linear-i-lanceolatis corollæ tubum subæquantibus, corollâ pallide violascenti extus parce lanuginosâ tubo nocti sculo fauce ampliato labii superioris lobis binis subretusis extus revolutis, inferioris ternis subæqualibus ovato-rotundatis omnibus denticulatis, filamentis ad medium tubum insertis basi villosulis, antheris ad suturas ciliatis, stylo apice parce glanduloso, stigmate divaricatum bilobo ♀.

Hab. *Artemisiæ* parasitica in valle Derrehgoum montium prope Jezd Persiæ orientalis (Buhse). Fl. Apr.

Pedalis, spica 7-8-pollicaris, corolla pollicaris. Species pulchra et distinctissima.

++ Filamenta glabra.

35. O. cernua (Löfl. It. p. 452. — Linn. Sp. p. 882) minute pruinosa vel glabrescens, caule crassiusculo breviter squamoso, spicâ densâ cærulescenti-violaceâ, bracteâ elevatim et parallele plurinervi ovato-lanceolatâ acutâ corollâ dimidio breviore, sepalis plurinerviis ovatis integris vel bisfidis incurvis corollam dimidiâ æquantibus, corollâ parvâ glabrâ tubulosâ brevi ad medium valde curvatâ cernuâ cæruleo-violaceâ labiis parvis crenatis non ciliatis superiore bilobo inferioris lobis subæqualibus, filamentis supra me-

dium tubum insertis, glabris, stigmatis albidi lobis velutino-granulatis ♀. Ic. Rehb. Germ. t. 187. — *O. Hispanica* Boiss. Voy. Bot. Esp. p. 478. — *O. curviflora* Viv. Eg. tab. 2, fig. 17. — *O. Gallica* Gren. Bot. Zeit. 22, p. 576. — *C. Grenieri* Schultz Fl. 1845, pag. 739. — *O. cumana* Mut. non Wallr.

Hab. ad *Achilleas*, *Artemisiæ*, *Cynareae* parasitica, in Egypto inferiore ad Kahiran in *Hyoscyamo mutico* (Schweinf!), inter Alexandriam et Rosette (Ehr!), Arabiâ petreâ et Palestinâ (Boiss!), Mesopotamiâ (Haussk!), Persiâ ad lacum Urmiah (Auch. 5078!), prope Jezd (Buhsel!), prope Aghda (Bge!), Tauriâ (Stev!).

Species insignis corolla valde curvata limbo brevi tubo dilatato angustiore. *O. bicolor* C.A. Mey. — Ledeb. Ic. tab. 390 est hujus varietas sepalis postice basi subcoalitis. Area geographicâ valde extensa.

Ar. Geogr. Hispania, Gallia australis, Arabia felix, Songaria, Algeria interior, Nova Hollandia.

36. **O. pogonanthera** (Reut. in D.C. Prodr. XI, p. 33) parce et crispule papillosa, caule crassiusculo squamis linearibus obsito, spicâ elongatâ densâ compactâ, bracteis oblongo-lanceolatis minute glanduloso-pruinosis oblongo-lanceolatis nervosis bifidis corollâ 2 $\frac{1}{2}$ plo brevioribus, corollâ præter glandulas paucas glabrâ pallidâ anguste tubulosâ reetâ labiis parvis obsolete denticulatis non ciliatis superiore subretuso inferioris lobis ovato-oblongis obtusis, filamentis medio tubo insertis glabris, antheris apice barbatis vel nudis, stigmate bilobo flavo ♀.

Hab. *Lycii*, *Tamaricis* etc. parasitica, in Persiâ australi (Ky. 849!), ad ripas fluvii Dalaki prope Kotel Malo (Haussk!).

Spica semipedalis et longior, corolla 6 lineas longa. Facie, bractearum et florum characteribus valde affinis *O. cernuae* a quâ non charaktere ut videtur incerto antherarum barbatarum sed corollâ rectâ nec infracto-curvatâ differt.

37. **O. Cumana** (Wallr. Orob. p. 58) glabrescens, caule gracili breviter squamoso, spicâ longâ laxâ, bracteis oblongo-lanceolatis acutis nervosis dimidiâ corollam æquantibus, sepalis pluriuerviis oblongis breviter bifidis corollâ brevioribus, corollâ glabrâ cœruleâ anguste tubulosâ fauce non ampliatâ horizontaliter valde curvatâ labiis parvis subæqualibus margine subintegris superiore retuso inferiore trilobo, filamentis supra medium tubum insertis glabris vel basi puberulis, stigmata albido retuso-bilobo ♀. Ic. Rehb. Crit. fig. 926-927. Rehb. Germ. tab. 189. — *O. arenaria* M.B. Taur. Cauc. II, p. 83.

Hab. *Artemisiarum*, *Xanthii* parasitica, Tauria (M.B.), Caucasus ad Stavropol (C.A. Mey!), Turcomania (Eichw.).

Species floribus horizontalibus valde incurvis tubulosis insignis, characteribus affinis *O. cernuae* a quâ limbo corollæ tubo duplo longiore nec æquilongo statim dignoscitur.

Ar. Geogr. Podolia, Rossia australis

38. **O. camptolepis** (Boiss. et Reut.) brevissime pruinosa-glandulosa, caulis nanis crebre squamatis in spicam oblongam

brevem densifloram abeuntibus, squamis bracteisque coriaceis late ovalis in mucronem brevem hamato-recurvum abrupte attenuatis hisce corollæ tubum subsuperantibus, sepalis oblongo-quadratis plurinerviis ad tertiam partem in lacinias rigidulas subrecurvas corollæ tubo sublongiores bifidis, corollâ e sicco carueâ puberulo-glandulosâ anguste tubuloso-infundibuliformi apice parum ampliatâ rectâ labiis parvis repandis non ciliatis superiore bilobo inferioris submajoris lobis rotundatis, filamentis medio tubo insertis antherisque glaberrimis, stigmate retuso rosello ♀. *O. latisquama* Reut. in Bal. exs. nou Schnitz. — *O. platylepis* Reut. MSS.

Hab. *Polygoni maritimi* parasitica ad ruinas Pompeiopolis in Ciliciâ (Bal!). Fl. Apr.

Cum spicâ bipollicari 2-3-pollicaris, bracteæ 5-6 lineas longæ 4-5 latæ, corolla parva sex lineas tantum longa. Species bracteis in cuspidem hamatam in sicco rigidam abeuntibus corollâque rectâ insignis.

Species ab auctoribus insufficienter et plerumque tantum ex speciminiis siccis descriptæ, probabilissime omnes specierum supra descriptarum synonyma; oblivioni quam merentur omni jure tradendæ.

O. grandisepala (F. W. Schultz Flora 1843 p. 128) sepalis plurinerviis tridentatis corollam æquantibus dentibus subulatis superiore 2-4 plo longiore, corollâ tubuloso-campanulatâ subarcuatâ denticulatâ, filamentis infra medium tubi insertis inferne parce pilosis.

Hab. in Græciâ ad Naupliam (Fraas).

Bipedalis et elatior, flores magn. *O. minoris*, facies *O. rubensis*.

O. Lycica (Schultz loc. cit.) sepalis plurinerviis ovatis bifidis corollæ tubum æquantibus vel longioribus, corollâ tubuloso-campanulatâ arcuatâ fave ampliatâ acute densiusculâ labio superiore bilobo, filamentis sub medio tubo insertis inferne subvillosis.

Hab. ad Lycos Atticæ (Fraas).

Facies *O. rubensis* vel *Epithymi*.

O. parviflora (Schultz loc. cit.) sepalis obsolete 3-6-nerviis a basi ovatâ lanceolato-subulatis sèpius bifidis corollæ tubum superantibus, corollâ tubulosa subarcuatâ laciñis parvis obsolete denticulatis, filamentis infra medium tubum insertis sparsim villosis.

Hab. ad Naupliam (Fraas).

Planta albido-subvillosa, exsiccata griseo-ochroleuca. Facies *O. Salvie*.

O. subverticillata (Schultz l. cit.), sepalis obsolete 5-nerviis ovatis inæqualiter bifidis vel indivisis tubum corollæ æquantibus, corollæ tubuloso-campanulatae arcuatæ labiis denticulatis superiore subbilobo, filamentis ad curvaturam corollæ insertis inferne subvillosis.

Hab. ad Thebas *Vicie* parasitica (Fraas).

An *O. rubens*? sed flores subminores.

O. lilacina (Schultz loc. cit. p. 129) sepalis obsolete 1-3-nerviis a basi ovatâ subito subulatis indivisis vel dente auctis corollæ tubo æquilongis vel longioribus, corollæ tubuloso-campanulatae breviter arcuatae labiis obsolete denticulatis lobis subæqualibus rotundato-truncatis, filamentis sub medio tubo insertis inferne pilosis.

Hab. in Atticâ (Fraas).

Facies *O. rubensis* sed minor.

O. arcuata (Schultz loc. cit.) sepalis obsolete 2-3-nerviis a basi ovatâ lanceolatis integris vel dente divaricato auctis, corollæ tubulosæ arcuatae horizontaliter patentis lobis denticulatis rotundatis, labio superiore multo majore, filamentis supra basin insertis ultra medium villosis.

Hab. in Atticâ (Fraas).

Parva sparsim albido-lanuginosa, flores eis *O. minoris* breviores.

O. diaphana (Schultz loc. cit. p. 130) sepalis trinerviis diaphanis lanceolatis indivisis corollæ tubum vix æquantibus, corollæ tubuloso-campanulatae vix arcuatae labiis denticulatis superiore paulo sursum curvato, filamentis supra basin insertis basi subvillosis.

Hab. in Phliotide (Fraas).

An varietas *O. Galii* quam facie refert?

O. angustisepala (Schultz loc. cit.) sepalis obsolete 1-4-nerviis indivisis et bifidis a basi ovatâ anguste subulatis corollæ tubum æquantibus vel brevioribus, corollæ campanulata fave ampliata dorso subcurvatâ labiis expensis undulato-plicatis obsolete denticulatis superiore subbilobo inferioris lobo medio duplo majore, filamentis supra basin corollæ insertis inferne villosis.

Hab. in Hymetto *Orobi digitati* parasitica (Fraas).

An var. *O. pruinosa*? Crescit quoque prope Fiume Illyriæ (Noë).

O. breviflora (Schultz loc. cit.) sepalis 3-5-nerviis a basi ovatâ subulatis integris tubum corollam æquantibus vel superantibus, corollæ a basi amplissimâ campanulata antice basi ventricosâ dorso curvatâ labiis denticulatis glanduloso-fimbriatis superiore retuso, staminibus basi corollæ insertis basi villosis.

Hab. in Atticâ (Fraas).

Genitalia exserta, facies *O. cruentæ*.

O. hians (Stev. Taur. p. 273) sepalis ovato-lanceolatis integris vel bifidis dimidium tubum subæquantibus, corollæ sensim ampliata fave late hiante labiis apice reflexis vix denticulatis imberibus, filamentis supra basin insertis ad medium usque pilosis, stigmate magno bifido margine elevato.

Hab. in Tauriâ, Iberiâ, Rossiâ australi (Stev.).

Facies *O. alba*.

O. squalida (Stev. Taur. p. 273) sepalis subulato-acuminatis in divisis vel bifidis corollam subæquantibus, corollæ glanduloso-villosæ tubulose leviter arcuatae labiis acute denticulatis parce ciliatis, filamentis ad basin styloque pubescentibus.

Hab. in Tauriâ et Iberiâ (Stev.).

An *O. Epithymum* Ledeb.? a quâ specie differt spicâ longiore, pube densiore, corollâ sordidè cæruleo-rufescens. An *O. pubescens* (Urv? Stev.).

Genus dubium.

Anblatum Tourn. Coroll. p. 48, tab. 481.

Planta aphylla facie *Orobanches* vel *Lathree*, corollâ breviter bilabiata labiis indivisis. Probabilissime *Anoplanthi* Icon erronea. *Anblatum Orientale* flore purpurascente Tourn. loc. cit. (nihil amplius).

ORD. XC. ACANTHACEÆ.

(Juss. Gen. p. 102.)

Flores hermaphroditi sæpius irregulares. Calyx persistens sepalis vel segmentis quinque vel abortu quatuor liberis vel sæpius plus minus alte connatis. Corolla gamopetala limbo nunc æqualiter quinquelobo nunc bilabiato nunc unilabiato. Stamina 4 didynama vel bina antica fauci inserta, antheræ biloculares loculis nunc parallelis nunc altero connectivo altius affixo. Ovarium superum biloculare ovulis anatropis in loculo binis vel numerosis. Capsula globosa oblonga vel linearis ab apice ad basin loculicide et sæpe elastice dehiscens, valvis medio semisepta seminifera ferentibus. Semina (in nostris) funiculo acreto indurato incurvo ascendentे (*retinaculo*) suffulta. Embryo rectus plano-compressus, radiculâ inferâ. Albumen nullum. — Herbae suffrutices et arbores. Præter genus *Acanthum* species infra enumeratæ tropicæ vel subtropicæ sunt areæ geographicæ majorem partem extra ditionem nostram habentes.

TR. I. RUELLIEÆ. — Corollæ lobi contorti. Semina plano-compressa hilo prope basin marginali, funiculo sæpissime in retinaculum incurvum indurato.

HYGROPHILA (R. Br. Prodr. 479. — *Hemiadelphis* Nees in D.C. Prodr. XI, 80).

Calycis segmenta 5 angusta infra medium plus minus connata parum inæqualia. Corolla ringens bilabiata labio superiore bidentato

inferiore trilobo. Stamina perfecta 4 didynama, vel bina cum stamnodes totidem minimis, filamenta basi dilatata infra medium tubum affixa. Antheræ biloculares loculis parallelis. Stylus apice linearis recurvus. Capsula oblongo-lanceolata compressa bilocularis polysperma. — Herbæ.

1. **H. polysperma** (Roxb. Fl. Ind. I, p. 120 sub *Justicia*) annua, caulinibus repentebus tenuibus ad nodos radieferis, foliis elliptico-oblongis glabris vel hirtulis, spicis terminalibus tenuibus cylindricis, basi laxis, floribus ad bractearum foliis conformium axillas sessilibus subsolitariis, bracteolis lanceolatis, calyce villoso ad $\frac{3}{4}$ in lacinias lanceolatas partito, corollæ purpurascens tubo subexerto, staminibus fertilibus binis, capsulâ oblongo-lineari ⊖. *H. polysperma* Anders. in Journ. Proc. Linn. Soc. IX, p. 456. — *Hemiadelphus polysperma* Nees. Wight. Ic. tab. 1462.

Hab. in regno Cabulico (Griff!).

Ar. Geogr. India tropica.

RUELLIA (L. Gen. 784).

Calyx quinquefidus vel partitus segmentis linearibus vel lanceolatis acutis. Corollæ infundibuliformis tubus rectus incurvus vel inflexus in faucem campanulatam dilatatus, limbus patens lobis 5 ovatis subæqualibus. Stamina 4 didynama sub fauce affixa filamentis basi dilatatis, antheræ biloculares loculis parallelis. Stylus apice recurvus. Capsula oblongo-linearis vel clavata plus minus basi slijata et sterilis compressa, apice tumida polysperma vel oligosperma.

1. **R. longifolia** (Stocks in Kew. Journ. IV, p. 177 sub *Dipteracantho*) basi suffrutescens, ramulis herbaceis erectis stricte ramulosis, foliis elongatis linearibus integris utrinque attenuatis adpresso hirtulis pallide virentibus, floribus axillaribus solitariis vel binis sessilibus remotis, bracteis linear-lanceolatis florem æquantibus et superantibus, calycis segmentis subulatis ciliatis dimidiata corollam æquantibus, corollæ lilacinæ tubo brevi in faucem amplissimam dilatato limbi lobis brevibus obtusis, ovarii loculis sexovulatis, capsulâ linearis calyce duplo longiore basi aspermâ brevi 5. *R. longifolia* Anders. Kew. Journ. 14, p. 177.

Hab. in Belutschia inferiore ad Schah Bilawul (Stocks!).

Rami novelli 6-10 pollicares, folia 2-3-pollicaria 2-3 lineas lata, corolla 9-10 lineas longa sesquilineam lata.

TR. II. ACANTHEÆ. — Corollæ tubus brevis, limbus superne fissus inferne in labium unicum obovatum expansus. Semina plano-compressa, hilo prope basiu marginali, funiculo saepius in retinaculum indurato.

BLEPHARIS (Juss. Gen. 103).

Calyx quadripartitus segmentis superiore et inferiore exterioribus majoribns, lateralibus interioribus parvis. Corolla unilabiata tubo brevi ovoideo apice constricto, limbo a basi breviter cupulatâ in labium planum breviter 3-5-lobum antice expanso. Stamina quatuor didynama fauci affixa labio breviora filamentis incrassatis incurvis, posticis apice attenuatis, anticis ultrâ antherarum insertionem breviter productis. Antheræ per paria conniventes ciliatæ uniloculares. Stylus apice attenuatus bifidus. Capsula bilocularis oblonga quadrivel abortu disperma. — Herbæ vel fruticuli.

1. **B. edulis** (Forsk. Eg. Arab. p. 114 sub *Acantho*) minute aspera et ciliatula glauca collo spicam sessilem interdum unicam et ramos breves crassiusculos decumbentes 1-3-stachyos edens, foliis radicalibus cito evanidis integris margine spinulosis, rameis ad soicarum basin oblongis et lanceolatis spinoso-dentatis, spicis oblongis quadrifariis demum elongatis columnaribus radicali et dichotomia-libus sessilibus, lateralibus breviter pedunculatis, bracteâ durâ late lanceolatâ valide nervosâ canaliculatâ squarroso-patenti spinoso-acuminatâ et utrinque valide 3-5-spinosâ flores superante, bracteolis brevioribus linearis-subulatis villosis, calycis adpresso hirsuti membranacei lacinis externis oblongis superiore longiore inferiore tridentatâ lateralibus quadruplo minoribus oblongo-linearibus. corollæ cærulescentis labio obtuse trilobo cum denticulo utrinque externo minuto, capsulâ compressâ ovatâ acuminatâ ♀. *B. edulis* Pers. Syn. II, p. 180. — *Ruellia ciliaris* L. Mant. p. 89 non alior. — *Ruellia Persica* Burm. Ind. p. 135. tab. 42, fig. 1. — *Acanthodium spicatum* Del. Fl. Eg. p. 97, tab. 33, fig. 3.

Hab. in desertis Ægyptiacis inter Cahiram et Suez (Del! Husson!), Arabiae Petree (Schimp. 367!), Persiae australis (Garcin), regni Mascate (Auch. 5221!), Belutschiae (Stocks!). Arabice *Schok el Dabb*. Folia erada edulia sapida.

Ar. Geogr. Nubia, Arabia tropica, India boreali-occidentalis.

ACANTHUS (L. Gen. 793).

Calyx quadripartitus laciinis superiore et inferiore exterioribus amplis elongatis venosis, lateralibus interioribus parvis obovatis vel orbiculatis. Corolla unilabiata tubo annulari brevissimo cartilagineo limbo basi attenuatâ superne fisco inferne in labium amplum obovatum planum late trilobum expanso. Stamina 4 fauci affixa subdidynama, filamenta crassa rigida apice subincurva, antheræ uniloculares longitudinaliter barbatæ. Stylus breviter bifidus. Capsula bilocularis oblonga coriacea nitida a basi ad medium tetrasperma. — Herbæ

perennes foliis *Cardui* vel *Morinæ*, floribus in spicam magnam terminali congestis magnis speciosis, basi bracteâ dorsali et bracteolis binis lateralibus instructis.

1. **A. mollis** (L. Sp. 891) foliis amplis membranaceis ad nervos et margines parce ciliolatis pinnatipartitis laciñiis oblongis grosse sinuatis et dentatis præter intimas basi angustatas distinctas basi confluentibus dentibus triangularibus inermibus apiculatis, spicâ peduncule longo folia saepius superante sussultâ longâ laxiusculâ pubescente, bracteâ exteriore oblongâ spinulose et profunde inciso-dentata ealyce subbreviore, bracteolis linearis-lanceolatis acutis, calycis membranacei bipartiti laciñiâ superiore obovato-oblongâ concavâ apice subtridentatâ inferiore subbreviore linearis-spathulatâ apice bidentatâ, corollâ albida labio obtuse trilobo inferne angustato ♂.

Hab. in pinguibus regionis inferioris Thraciae et Macedoniæ (Frider., Griseb.). E ditione nondum vidi.

Forsan ex patriâ et area geographicâ specimina citata ad *A. longifolium* Host referenda sunt speciem affinem Dalmaticam foliorum laciñiis late ovatis omnibus distinctis et basi abrupte attenuatis, rachide alata, bracteolis subulatis, calycis labio inferiore integro distinctam. Planta a cl. Griseb. ex Albaniâ boreali (extra ditionem nostram) lecta est *A. longifolius*.

Ar Geogr. Lusitania, Hispania, Gallia australis, Italia, Dalmatia, Africa borealis.

2. **A. Dioscoridis** (L. Sp. Ed. Willd. III, p. 398) caule tomentello elongato, foliis sub lente papillaribus late viridibus membranaceis rigidulis radicalibus longe oblongo-lanceolatis in petiolum longe attenuatis integerrimis vel inferne utrinque 2-3 lobulatis lobis triangularibus saepè mucronato-spinnlosis, caulinis diminutis sessilibus spinnoso-dentatis, spicâ longâ crassâ densâ, bracteis late oblongo-acuminatis profunde spinoso-incisis calycem subæquantibus, bracteolis subulatis ciliatis, calycis hirsuti laciñiâ superiore oblongo-linearis integrâ corollâ vix breviore, inferiore anguste linearis apice dilatata integrâ, corollâ purpureâ labio lato trilobo basi angustato lobis latissimis subretusis ♀. *A. Raddei* Trautv. Aliq. sp. novæ p. 3.

Hab. in dumosis subalpinis Armeniæ in prov. Musch (Ky. Suppl. 722!) et in prov. Erzerum (Badde!), Mesopotamie inter Suerek et Diarbekir (Ky. 471!), Kurdistanie Persicæ ad Mendeli (Noë 1157!), inter Masibin et Awiheing et ad Nansud 7-8000' (Haussk.).

Folia inferiora cum petiolo saepè pedalia 1½-2-pollices lata, spica crassa 7-10-pollicaris. Ad hanc unicam speciem accommodatur phrasis Linnei « *foliis lanceolatis integerrimis margine spinosis* » quamvis spinæ paucissimæ sint et saepè omnino deficiant. Locus « *Libanus* » quoque non convenit, sed loci plantarum Orientalium a botanicis antiquioribus indicati saepius erronei. Specimen in herbario Linnei nullum.

3. **A. hirsutus** (Boiss. Diagn. Ser. I, 4, p. 86) totus pilis brevibus crispuulis deflexis hirsutus, caule saepius humili tollis radicalibus breviore, foliis radicalibus longis anguste lanceolatis petiolatis in

lacinias triangulares acutas triangulari-dentatas breviter et molliter spinulosas basi confluentes inferiores saepe retrorsas pinnatipartitis superioribus subbinis conformibus sessilibus, spicâ ovatâ vel oblongâ compactâ, bracteâ amplâ ovato-oblongâ cucullatâ hirtâ nervosâ in spinas tenues latitudine suâ breviores pinnatifidâ, bracteolis hirsutissimis subulato-linearibus apice spinulosis bracteâ vix brevioribus, calycis longe hirsuti laciñiâ superiore oblongo-spathulatâ subintegrâ corollam æquante, inferiore late linearî subbreviore dentibus tribus spinulosis, corollæ albidæ lutescentis inferne hirsutæ labio obtuse et breviter trilobo ♀. *A. spinosus* ex parte Nees in D.C. Prodr. XI, p. 271 nec L. nec Auct.

Hab. in regione montanâ Lydiâ et Cariâ in parte inferiori montium Tmolus, Mesogis, Cadmus (Boiss!), in Pisidiâ, Isauriâ et Lycaoniâ (Heldr!), Cappadociâ (Ky. 184! Bal!), Galatiâ ad Tokat et Angora (Noë! Wied!), insulâ Rhodo (Auch. 1699!).

Planta 9-10-pollicaris rarius pedalis, folia inferiora 9-15-pollicaria 2-pollices lata, spica 4-6-pollicaris 2-polices lata. Species foliis teneris minime coriaceis, spinulis tenellis brevibus non pungentibus ab omnibus *Acanthis spinosis* statim distinguenda.

4. **A. Syriacus** (Boiss. Diagn. Ser. I, 11, p. 135) puberulus, caule folia parum superante, foliis elongatis oblongo-lanceolatis basi in petiolum brevem attenuatis in lacinias lanceolatas iterum lobato-spinosas basi confluentes vel infimas distinctas pinnatipartitis spinis tenuibus pungentibus, foliis caulinis subbinis diminutis sessilibus, spicâ ovato-cylindricâ compactâ, bracteâ amplâ coriaceâ elevatum nervosâ et reticulatâ margine ciliatulâ dorso glabriuscûlâ ovato-oblongâ utrinque in 4-5 spinas validas pinnatifidâ flores subæquante, bracteolis brevioribus anguste linearibus hirtulis apice spinulosis, calycis glabriuscûli laciñiis majoribus herbaceis superiore oblongo-spathulatâ apice minute tri-denticulato-repandâ inferiore angustiore et breviore linearî apice dilatatâ bifidâ dentibus spinulosis, corollæ albidæ glaberrimæ labio trilobo lobis rotundatis ♀. *A. spinosus* ex parte Nees in D.C. Prodr. non L.

Hab. inter segetes, in siccis regionis inferioris et montanæ, Cilicia orientalis ad Sis (Ky. Suppl. 57!), Mesopotamia circa Orfa, Suerek, Harran (Ky. 63! Haussk!), Syria littoralis in regione inferiore Libani (Bl! Gaill!), Caesarea (Ehr! Boiss!), Palestina circa Panias, Nazareth, Hierosolymam (Boiss! Ky. 150!).

β. dentatus. — Folia breviora breviter et irregulariter spinoso-dentata.

Hab. in Syria boreali prope Aintab (Haussk!). Ciliciâ (Tchih!), Cappadociâ ad Euphratem (Auch. 2698!).

Planta saepius humilis, *A. hirsuto* tamen saepius major, huic affinis, distinctus spinis foliorum et bractearum multo validioribus pungentibus, irdumento glabrescenti, corollâ glabrâ.

5. **A. spinosus** (L. Sp. p. 891) glabriusculus, foliis ambitu late ovato-oblongis nitidis elevatum nervosis pinnatum et bipinnati-

partitis laciniis primariis divaricatis longe lanceolatis secundariis triangulari-lanceolatis apice et lateraliter in spinas longas abeuntibus, spicæ crassæ elongatæ glabriusculæ, bractæ oblongo-lanceolatæ coriaceæ rigidæ elevatim nervosæ lateraliter longe spinosæ in spinam subrecurvam canaliculatam abeunte, bracteolis lanceolato-subulatis acerosis bracteæ equilongis, calycis hirtuli laciniæ superiore oblongo-spatulatæ apice 3-5-deuticulatæ inferiore breviore et angustiore bidentatæ denticulis spinulosis, corollæ glabrae albae labio apice obtuse et late trilobo ♀. Ic. Fl. Græc. tab. 611. — *A. spinosissimus* Desf. Hort. Par. — Rehb. Ic. tab. 194.

Hab. in collibus, in cultis derelictis regionis inferioris, Pelopennesus (Bory!), Attica (Sprun! Heldr!), Zacynthus (Marg!), Macedonia (Friv!), Lydia (Boiss!), Rhodus (Bourg!), Creta (Sieb! Raul!).

Ar. Geogr. Dalmatia.

TR. III. JUSTICIE.E. — Corollæ limbus subæqualis vel bilabiatus lobis varie imbricatis. Semina plano-compressa hilo marginali, rarius ovoideæ hilo basilari, funiculo in retinaculum incurvum indurato.

BARLERIA (L. Gen. 785).

Calyx quadripartitus segmento antico et postico majoribus exterioribus, lateralibus interioribus angustis. Corolla infundibuliformis tubo sèpius longo, fauce dilatata, limbi lobis quinque subæqualibus. Stamina 4 didynama prope corollæ basin inserta, linea antica longiora, 2 postica interdum ananthera, autherae loculis parallelis. Stylus apice integer. Capsula ovoidea vel oblonga tetrasperma vel abortu disperma. — Herbæ vel frutices.

1. B. Hochstetteri (Nees in p.c. Prodr. XI, p. 231) suffrutescens ramis vetustis ramosissimis tortuosis novellis herbaceis opposite ramulosis breviter canescenti-velutinis, foliis viridibus pube adpresso bipartitæ hirtulis pedunculatis parvis ovatis vel ellipticis integris obtusis sèpe mucronulatis, ramulis trifloris, floribus pedunculatis pedunculo medio ebracteolato lateralibus bibracteolatis, calycis laciniis anticæ et posticæ ovato-ellipticis acutiusculis, corollæ tubo calyce 1½-duplo longiore, limbi lobis subæqualibus. — Hook. Ic. tab. 803 — *B. diandra* Hochst. et St. in Schimp. pl. Arab. fel.

Hab. in collibus Belutschiae (Stocks!).

Rami novelli semipedales, folia majora pollicaria, corolla pollicaris.

Ar. Geogr. Nubia, Abyssinia, Arabia felix, ditio Scindæ Indiae.

2. B. Aucheriana (Nees in p.c. Prodr. XI, p. 234) fruticulosa ramosissima ramis tortuosis adpresso canis, foliis parvis carnosulis

elliptico-oblongis obtusis glabrescentibus, ramulis trifloris demum subspinescentibus, floribus pedunculatis pedunculo medio ebracteato lateralibus 2-bracteolatis, calycis laciniis antica et posticæ ovato-lanceolatis acuminatis, corollæ tubo calyce subdupo longiore limbi lobis subæqualibus ♂.

Hab. in monte Chebek ditionis Mascate Arabiae (Auch. 5222!).

Fruticulus 3-4-pollicaris, folia 4-6 lineas, calyces $2\frac{1}{2}$ lineas longi, corolla 8 lineas longa. Ex cl. Anderson forma *B. Hochstetteri* a qua differre videtur pedunculus demum subspinescentibus, foliis parvis angustioribus, laciniis calycis acuminatis, sed specimina manca.

3. ***B. acanthoides*** (Vahl Syrib. I, p. 47) fruticosa ramosa pubescens, foliis obovato-ellipticis obtusis mucronulatis petiolatis, floribus axillaribus sessilibus subinde solitariis saepius 2-5 in spicam unilateralem patentem congestis, bracteis spinosis inferioribus pinnatis herbaceo-dilatatis superioribus spicæ subintegris, calycis laciniis majoribus amplis membranaceis reticulato-nervosis apice spinulosis margine nervulis excurrentibus spinuloso-ciliatis vel nudis, corollæ tubo longissimo ♂.

Hab. in Egypto superiore (Ky. ex Nees), Belutschia (Stocksl.).

4. ***B. candida*** (Nees in p.c. Prodr. XI, p. 240) fruticulosa adpresso cana ramosissima erinacea ramis brevibus intricatis, foliis minimis ovato-orbiculatis petiolatis obtusis, floribus axillaribus solitariis, bracteis in spinam longam subulatam patentem versus medium utrinque saepe unispinulosam mutatis, calycis laciniis majoribus oblongis spinoso-acuminatis reticulatis pubescenti-scabridis, corollâ.... ♂.

Hab. in aridis regni Mascate Arabiae (Auch. 5223!).

Ex cl. Anderson *B. acanthoidi* adnumerata sed e specimine cæterum valde incompleto omnino diversa videtur, folia cum petiolo 2-3 lineas tantum longa, spinæ bracteales non dilatatae longiores 6-8 lineas longæ, calycis laciniæ 3-6 lineas longæ 2 lineas latæ.

LEPIDAGATHIS (Willd. Sp. III, p. 400).

Calyx subquinquepartitus segmentis saepius scariosis acutis inæqualibus, postico latiore anticis binis saepe plus minus connatis. Corolle tubus cylindraceus in faucem campanulatam dilatatus, limbus bilabiatus labio superiore erecto-patente retuso vel bifido inferiore inæqualiter trilobo. Stamina quatuor didynama basi faucis affixa inclusa, antheræ biloculares loculis discretis muticis subæqualibus altero altius affixo. Stylus apice recurvus. Capsula oblonga acuta a basi bilocularis tetrasperma, semina orbiculata retinaculis validis suffulta. — Herbæ vel suffrutices inermes vel bracteis spinescentibus donati, floribus in spicas densas bracteatas saepius axillares dispositis.

1. **L. calycina** (Hochst. in Schimp. exs. — D.C. Prodr. XI p. 252) fruticulosa virens adpressiuscula hirta, foliis parvis breviter petiolatis oblongis et oblongo-lanceolatis saepe subobliquis, spicis axillaribus confertis strobiliformibus oblongis, bracteis seariosis trinerviis obovatis acutis apiculatis ciliatulis ad nervos puberulis, bracteolis subulatis ciliatis, calycis seariosi albidi laciniis anticus in unicam basi coalitis, posteriore trinervi latiore, lateralibus multo minoribus subulatis, corollæ albidae tubo inclusu limbo parvo ♂. *L. discolor* Stocks exs. — *L. strobilifera* Stocks in Hook. Journ. et Kew Misc. IV, p. 177.

Hab. ad Sehah Bilawul in Belutschia inferiore (Stocks!).

Fruticulus $\frac{1}{2}$ -1-pedalis, folia 7-10 lineas longa, spicæ pollicares vel breviore, calyx 4 lineas, corolla 7 longa.

Ar. Geogr. Abyssinia, regio Scinde Indie.

JUSTICIA (L. Gen. 27. — *Rostellaria* Nees in Wall. As. III, 76).

Calyx in segmenta 4-5 angusta æqualia partitus. Corollæ tubus limbo brevi sublongior superne ampliatus, limbus bilabiatus labio superiore concavo integro vel breviter bifido, inferiore patente trilobo. Stamina bina fauci affixa labio superiore breviora, antheræ bi-loculares loculis discretis, altero altius affixo mutico, altero inferiore basi calcare parvo albo appendiculato. Stylus filiformis integer vel minute bidentatus. Capsula ovata vel oblonga basi in stipitem solidum contracta tetrasperma, semina rugosa vel tuberculata. — Herbae, rarius frutices.

1. **J. peploides** (Nees in D.C. Prodr. XI, p. 375 sub *Rostellaria*) annua, ramosa ramis erectis vel procumbentibus ramulosis, foliis viridibus sparsim papillari-asperulis vel glabris parvis petiolatis ovato-oblongis vel oblongo-lanceolatis, spicis terminalibus densis tenuibus cylindricis basi foliis binis minoribus subsessilibus saepius suffultis, bracteis calycisque laciniis subæqualibus linear-lanceolatis acutis asperis et ciliatis rarius glabrinusculis, corollæ albidae tubo inclusu limbo parvo ♂. *J. Vahlii* Roth Nov. Sp. p. 14. — *J. quinangularis* Roxb. Fl. Ind. p. 123 ex parte..

Hab. in oryzetis, locis humidis, regnum Cabulicum (Griff!), Belutschia (Stocks!).

Ar. Geogr. Abyssinia, India.

ECBOLIUM (Kurz Journ. Asiat. Beng. 40, p. 75).

Calyx profunde quinquepartitus segmentis lanceolatis subæqualibus. Corollæ tubus longus tenuiter cylindraceus in fauces breviter ampliatus, limbus bilabiatus labio superiore linear breviter biden-

tato inferiore patente profunde trilobo. Stamina bina sub fauce affixa, antheræ subexsertæ biloculares loculis parallelis æqualibus muticis. Stylus apice bilobus. Capsula ovata septo parallele compressa in stipitem longum contracta abortu disperma. Semina plana tuberculata. — Frutices.

1. **E. Linneanum** (Kurz loc. cit.) fruticosum vel arboreum, foliis late flavo-virentibus sub lente puberulis vel glabrescentibus elliptico-oblongis acutiusculis in petiolum brevem breviter attenuatis, floribus in spicas terminales tetragonas congestis, bracteis ovatis herbaceis acutis vel obtusis calyces superantibus, calycis hirsuti laciniis anguste lanceolatis, corollæ tubo bracteis duplo vel triplo longiore, limbo tubo $2\frac{1}{2}$ -plo breviore 5. *Justicia Ebolium* Linn. Fl. Zeyl. p. 47. — Ic. Burm. Zeyl. tab. 4, fig. I. — *Eranthemum Ebolium* Anders. in Thw. Ceylan, p. 235.

Hab. in regno Mascate Arabiæ (Auch. 5224!).

Ar. Geogr. Nubia, Abyssinia, Arabia tropica, India.

DICLIPTERA (Juss. in Ann. Mus. Par. IX, p. 450).

Bracteæ binæ basi subconnatae florem unicum vel plures includentes. Calyx profunde in laciniæ subæquales quinquefidus. Corollæ tubus tenuis ad faucem plus minus ampliatus, limbus longe bilabiatus labio superiore integro vel retuso inferiore patente integro vel brevissime trilobo. Stamina bina fauci affixa, antheræ loculis binis discretis superpositis sæpius muticis. Stylus apice integer vel bidentatus. Capsula ovata vel orbiculata subcompressa subsessilis vel stipitata, placentæ sub dehiscentiâ basin versus a valvis elastice solutæ et sursum curvatæ. Semina quatuor vel abortu bina orbiculata plana retinaculis latiusculis acutis suffulta. — Herbæ.

1. **D. Roxburghiana** (Nees in Wall. Pl. As. rar. III, p. 111) pilis rarís centro affixis adpresso et parce strigulosa viridis, caulinis decumbentibus tenuibus elongatis ramosis, foliis ovato-oblongis acuminatis subrepandis in petiolum brevem attenuatis, cymis ad axillas 4-5-nis aliis brevius aliis longe pedunculatis laxè 3-5-floris, bracteis involucralibus unifloris a basi cuneatâ obovato-ellipticis inæqualibus mucronulatis ciliatis venoso-triangulariis, corollæ tubo breviter exerto, capsulâ ovato-subrotundâ compressâ unguiculatâ hirsutâ ♀. — *D. bupleuroides* Nees. — *D. hirtula* Nees. — *Justicia Chinensis* Roxb. Fl. Ind. I, p. 125 non L.

Hab. in regno Cabulico (Griff. ex Anders.).

Ar. Geogr. India Orientalis.

ORD. XCI. SELAGINACE.E.

(Lindl. Introd. p. 279.)

Flores hermaphroditici irregulares. Calyx inferus 3 rarius 3 partitus vel fidus, interdum spathaceus bipartitus. Corolla gamopetala tubo tenui rariis in fauce ampliato, limbo quinquefido rarius bilabiato. Stamina quatuor corollae tubo inserta rarius bina. Anthere loculis confluentibus uniloculares. Discus hypogynus annularis. Ovarium fiberam biloculare ovulis solitariis pendulis anatropis. Caryopses binæ coherentes vel tandem secedentes alterà siepe abortivæ. Embryo teres in axi albuminis carnosí rectus, radiculâ superâ.

GYMNANDRA (Pall. It. III. — *Layotis* Gaertn. Comm. Acad. Petrop. XIV, p. 533.).

Calyx spathaceus antice tissus postice integer vel 2-3-fidus. Corolla tubus brevis fauce ampliata, limbus bilabiatus labio superiore integro vel bifido inferiore trifido rarius bifido. Stamina bina fauci affixa, anthere versatiles confluentia uniloculares. Ovarium biloculare loculis uniovulatis. Stylus apice subdilatatus capitatus vel subbilebus. Fructus parvus exocarpio tenui subearnoso, endocarpio crustaceo. Semen pendulum, embryo teres rectus in albumine carnosò. — Herbae perennes facie *Pudrorot*. — Genus inter *Selaginaceus* anomalum, ab eis patriâ, corollæ tubo superne ampliato, limbo bilabiato, stigmate capitato discedens sed cæteris notis congruens, *Serophulariacearum* tribu *Veronicaeis* valle affine sed ovulis iu loculis solitariis ab eis discrepans.

1. **G. stolonifera** (C. Koch Linn. XVII, p. 289) glabra aequalis, rhizomate crasso brevissimo subobliquo præmorso fibras longas tenuiter cylindricas emitente, collo dense fibroso folia scapos breves et stolones edente, foliis omnibus radicalibus rosulatis late lanceolatis uninerviis integris vel subcrenatis acutiusculis in petiolum brevem complanatum attenuatis, stolonibus longis tenuibus decumbentibus squamuulis parvis linear-oblängis obsitis apice radicantibus et fasciculo foliorum parvorum terminatis, scapis radicalibus foliis brevioribus, spicâ ovato-oblängâ densiflorâ, floribus subsessilibus bracteâ oblongâ acutiusculâ æquilonâ suffultis, calyce tubuloso-spathaceo submembranaceo antice ad basin usque fisco postice in dentes 2 triangulares obtusos ad tertiam partem fisco, corolla cæruleæ calyce duplo longioris tubo subexerto labiis æquilonis superiori obovato cuneillato integro inferiore patente in lobos binos ovalos profunde fisco, staminibus breviter exsertis, stigmate capitato, fructu ovato-

oblongo glabro breviter rostrato et superne longitudinaliter nervoso calyce longiore ♀. Ic. Jaub. et Sp. Ill. tab. 254. — *G. Armena* Boiss. Diagn. Ser. I, 4, p. 75.

Hab. in pascuis alpinis, Anatolia prope Sivas (Botta), Cappadocia circa Karahissar (Auch. 50861), Armenia Turcica prope Erzerum et Zazalarhané (Huet! Calv!), supra Gumuschkhané (Bourg!), Kurdistania circa Van (Noë!), Somchetia (C. Koch), prov. Aderbidjan Persie (Szov!).

Collum crassiusculum, folia cum petiolo $1\frac{1}{2}$ -3-pollicaria 4-6 lineas lata, spicæ pollicares. Species stolonibus, corollæ labio superiore integro inferiore bi nec trifido a congeneribus distinctissima, fructus non bipartibilis videtur.

ORD. XII. GLOBULARIEÆ.

(D.C. Fl. Fr. III, p. 427.)

Flores hermaphroditi irregulares. Calyx quinquefidus vel partitus æqualis vel bilabiatus persistens. Corolla hypogyna gamopetala oblique bilabiata interdum unilabiata. Stamina 4 didynama tubo corollæ inserta, antheræ confluentia uniloculares. Stigma bifidum. Ovarium liberum uniloculare, ovulo unico pendulo anatropo. Caryopsis utricularis indehiscens. Embryo teres in axi albuminis rectus, radicula supera. — Ordo a cl. Bentham et Hooker Gen. II, p. 1127 *Selaginaceis* forsitan recte conjunctum ab eis tamen inflorescentia et ovario uniloculari discedens.

LOBULARIA (L. Gen. 442).

Calyx basi turbinatus vel campanulatus plus minus profunde 5-fidus vel partitus, æqualis vel bilabiatus. Corolla tubo brevi limbo obliquo bilabiato labio superiore bipartito breviore interdum fere obsoleto vel nullo, inferiore tripartito vel tridentato. Stamina 4 didynamæ fanci affixa, antheræ versatiles confluentia uniloculares. Ovarium uniloculare uniovulatum, stylus apice minute bidentatus. Fructus parvus calyce inclusus oblongus indehiscens membranaceus, semen pendulum, embryo in axi albuminis carnosus rectus. — Herbae, suffrutes vel frutices, floribus in capitula globosa involucrata dense congestis.

1. **G. vulgaris** (L. Sp. 139) glabra, rhizomate cespitoso multicipiti, caulibus herbaceis foliosis felia radicalia multo superantibus, foliis radicalibus rosulatis obovatis in petiolum attenuatis quinque-nerviis apice retusis rarius tridenticulatis, caulinis parvis sessilibus elliptico-lanceolatis acutis, capitulo terminali, involueri phyllis ovato-lanceolatis acutis trinerviis ciliatis, receptaculi paleis lanceolatis uni-nerviis a medio sursum setosis, calycis setoso-hispidi laciniis sub-

æqualibus lanceolatis acutis erectis tubo duplo longioribus, corollæ cœruleæ calyce $2\frac{1}{2}$ -plo longiore labio superiore brevi bipartito inferioris triplo longioris tripartiti segmentis late linearibus basi trinerviis superne uninerviis 2. Ic. Relib. tab. 196. — Willk. Glob. tab. I, fig. 4. — *G. Willkommii* Nym. Syll. p. 140.

Hab. in monte Haemo Thraciae (Friv.), Caucaso et Iberia (M.B., Willk.) E ditione nondum vidi.

Loci Anatolici ab auctoribus indicati ad speciem sequentem spectant.

Ar. Geogr. Lusitania, Hispania borealis, Gallia, Belgium, Helvetia, Germania, Italia, Dalmatia, regio Danubialis, Serbia.

2. *G. trichosantha* (Fisch. et Mey. Ind. Petrop. V, pag. 36) glabra glaucescens, rhizomate multicipiti stolonifero, caulis herbaeis foliosis folia radicalia superantibus, foliis radicalibus rosulatis obovatis et obovato-oblongis in petiolum longe attenuatis obtusis vel tridenticulatis subnunerviis venis secundariis obscuris, caulinis parvis sessilibus ellipticis et elliptico-linearibus mucronatis, capitulo terminali, involueri phyllis oblongo-lanceolatis longe acuminatis setosis, receptaculi paleis lanceolatis acuminatis trinerviis superne margine ciliatis, calycis setoso-hispidi laciñiis subæqualibus lanceolato-subulatis tubo 3-4-plo longioribus, corollæ cœruleæ sub-duplo longioris labio superiore duplo breviore in segmenta capillaria bipartito, inferioris segmentis ternis anguste linearibus usque ad medium trinerviis superne uninerviis 2. Ic. Jaub. et Sp. Illustr. tab. 259 B. — Willk. Glob. tab. I, fig. 4. — *G. vulgaris* var. *Bithynica* Griseb. Spic. II, p. 294. — *G. macrantha* C. Koch in Walp. Rep. IV, p. 175. — *G. pallida* C. Koch Linn. XXII, p. 634.

Hab. in siccis dumosis regionis montanæ, Byzantii (Goumany!), Bithynia circa Moudania, Broussa, (Thirke! Pichl! Barbey!), Caria in Cadmo supra Denisleh (Boiss! Pin!), Pamphylia (Heldr!), Cilicia prope Gülek (Ball!), Anatolia Borealis (Auch. 523!), et circa Thorbal et Ancyram (Wied!), Armenia Turcica (Calv! C. Koch) et Rossica prope Erivan (Buhsei!), Cappadocia (Ball!) Syria borealis prope Aintab (Haussk!).

Facies *G. vulgaris* a quâ rhizomate stolonifero, foliis subminerviis, capitulis et floribus majoribus, calycis laciñiis longioribus acuminatis, corolla laciñiis angustis eximie differt.

3. *G. cordifolia* (L. Sp. 139) glabra, rhizomate suffrutescenti ramosissimo ramis prostratis et radicantibus, foliis obovatis uninerviis in petiolum longe attenuatis obtusis integris retusis vel tridenticulatis, pedunculis elongatis nudis vel squamis 1-2 minimis oblongis vel linearibus obsitis, capitulo terminali cernuo, involueri phyllis obovato-oblongis mucronatis margine ciliatis, receptaculi paleis lanceolatis basi attenuatis acuminatis uninerviis margine breviter ciliatis, calycis bilabiati laciñiis tubo setuloso sublongioribus lanceolatis acutis labii superioris paulo angustioribus, corollæ cœruleæ calyce duplo longioris labio superiore bipartito laciñiis linearibus superne dilatatis, inferiore sublongiore trifido laciñiis linearibus usque ad tertiam partem trinerviis cæterum uninerviis 3.

Hab. in rupestribus, regio alpina Olympi Thessali (Heldr! Orph!), Scardus Macedoniae et Haemus Thraciae (Friv., Griseb.), in Tauriæ (Pallas ex Willd. herb.).

G. nana Lam. fide schedulæ mendacis Gittardianæ in Peloponneso erroneè indicata est species propria nec *G. cordifolia* varietas! in Pyrenæis, Hispaniâ boreali, Galliâ australi, Alpibus Liguriæ crescens.

Ar Geogr. Hispania borealis, Gallia orientalis et australis, Germania in Alpibus, Italia, regio Danubialis, Dalmatia.

4. **G. Stygia** (Orph. in Boiss. Diagn. Ser. II, 4, p. 69) glabra, rhizomate in caudiculus tennes fragiles denudatos prostratos et radicantes diviso, foliis ad collum surculos breves foliosos edens contortis uninerviis ovato-orbiculatis basi cuneatis petiolatis obtusissimis, capitulo terminali ex foliis vix exerto pedunculo eo vix longiore nudo vel squamâ oblongâ obsito suffulto, involuci phyllis oblongo-lanceolatis acuminatis margine ciliatis cæterum glabris, receptaculi paleis anguste linear-lanceolatis acutis glabris uninerviis, calyce fere ad basin in lacinias subæquales anguste lanceolatus apice subulatas undique longe albo-ciliatas partito, corollæ caruleæ calyce sesqui-longioris labio superiore in lacinias anguste lineares bipartito inferiore tertiatâ parte longiore tripartito laciñis linearibus ad $\frac{2}{3}$ trinerviis apice uninerviis 2.

Hab. in regionis alpinae fissuris rupium, mons Chelmos Peloponnesi supra Stygem rarissima (Orph.).

Folia fere *Carradoriu* *incanescens*, capitulum eo *G. cordifolia* paulo magius: ab ea caudiculis tenuibus fragillimis non sufractiosis, pedunculis brevissimis, involuci squamis angustioribus longioribus, calyce non bilabiato, quinquepartito nec quinquefido etc., a *Carradoriu* quam facie refert corollæ structurâ etc. statim distinguenda.

5. **G. Alypum** (L. Sp. 139) glabra glaucescens, fruticosa, ramosissima ramis erectis ramulosis, foliis sparsis coriaceis uninerviis basi attenuatis subpetiolatis apice mucronatis vel tridentatis, superioribus usque ad capitulum sensim diminutis, capitulo terminali, involuci phyllis numerosis imbricatis ovatis mucronatis margine ciliatis, receptaculi subglobosi basi in stipitem angustati paleis linearis-subulatis hirtis caducis, calyce longe barbato ad $\frac{2}{3}$ in lacinias subæquales linear-acuminatas partito, corollæ caruleæ calyce subdupo longioris labio superiore rudimentario bipartito segmentis dentiformibus, inferiore longissimo breviter tridentato dente medio trimervi lateralibus binerviis 3. Ic. Rehb. tab. 197.

Hab. in collibus regionis inferioris, Græcia et ejus insulae ubi *Staurias* vel *Sideri* audit (Sibth., Sprun! Heldr! Orph!), Anatolia occidentalis (Boiss.), Lisanus (Runwolf ex A. D.C.). Fl. Hyeme et vere.

Ar Geogr. Hispania orientalis et australis, Gallia australis, Italia, Africa borealis.

6. **G. Arabica** (Jaub. et Sp. III. Or. III. tab. 260) glabra glaucescens fruticosa ramosissima, ramis brevibus rigidis divergentibus,

foliis sparsis spathulato-oblongis in petiolum brevem attenuatis integris vel subtridentatis uninerviis, superioribus sensim diminutis, capitulo terminali, involuci phyllis imbricatis ovatis coriaceis setosis, receptaculi conico-cylindrici non stipitati acutiusculi paleis lanceolatis et linearibus cuspidato-acuminatis villosis, calyce longe barbato ad $\frac{3}{4}$ in lacinias subæquales lanceolato-subulatas partito, corollæ cœruleæ calyce sesquilongiore unilabiata apice breviter trilobâ lobis uninerviis 3. — *G. trichocalyx* Steud. in Schimp. Pl. Arab. — *G. Alypum* Del. Eg. p. 5 non L.

Hab. in montibus Sinaï et Serbal Arabie petrea (Schimp. 406! Botta), ad rupes circa Alexandriam Egypti (Delile, Samart), Arabice *Handeguk*.

Facies *G. Alypi*, rami breviores crassiores divaricati; ab ea receptaculi formâ, corollæ labio superiore prorsus nullo, etc., distincta.

7. G. Orientalis (L. Sp. p. 140) glabra, caudice lignoso ramoso ascendentí vel diffuso tortuoso. caulis floriferis fragilibus parce foliosis in racemum spiciformem basi laxissimum abeuntibus, foliis uninerviis radicalibus fasciculatis oblongo-vel obovato-spathulatis in petiolum attenuatis acutiusculis, caulinis paucis sparsis sessilibus oblongo-lanceolatis et linearibus acuminatis, capitulis 5-12 parvis inferioribus remotis pedunculis strictis eis saepe longioribus suffultis superioribus sessilibus congestis, involuci phyllis ovatis acuminatis concavis uninerviis hispidis, receptaculi paleis obovatis acuminatis trinerviis hispido, calycis membranacei hispido brevissime bilabiati dentibus breviter triangularibus labii inferioris profundiis fisis, corollæ cœrulecentis calyce subtriplo longioris labiis subæquilongis tubo subduplo longioribus superiore bipartito laciinis anguste linearibus superiore trifido laciinis trinerviis 3. Ic. Jaub. et Sp. Ill. tab. 259 A.

Hab. in argillosis Anatoliæ interioris (Tournef.), in monte Mesogi prope Derbent (Boiss!), Pisidiâ inter Ghelindost et Karagatch (Heldr!), circa Kadi-kios et in montibus prope Ancyram (Wiedem!), Cappadocia (Tchih! Bal!), Syria borealis prope Aintab (Haussk!).

ORD. XCIII. VERBENACEÆ.

(Juss. Ann. Mus. VII, p. 63.)

Flores hermaphroditæ rarius polygamo-dioici regulares vel irregulares. Calyx persistens gamosepalus 4-5-dentatus vel partitus. Corolla gamopetala hypogyna tubulosa vel hypocrateriformis limbo saepius obliquo et inæquali. Stamina 4 didynama, antheris bilocularibus. Ovarium liberum integrum disco insidens 2-4-loculare loculis uniovulatis, ovulis erectis vel pendulis. Fructus drupaceus siccus 2-4 pyrenis constans, rarius capsularis abortu monospermus. Embryo rectus radiculâ inferâ. Albumen nullum vel parcissimum.

TR. I. VERBENEÆ Schauer.

Inflorescentia indefinita spicata vel racemosa undique centripeta. Ovula e basi loculi erecta anatropa. Semina exalbuminosa.

LANTANA (L. Gen. 765).

Calyx minimus membranaceus truncatus vel sinuato-dentatus. Corolle tubus cylindricus aequalis vel ad semina subampliatus. limbus patens obliquus subbilabiatus in lobos 4-5 latos obtusos fissns. Stamina 4 didynama inclusa, antheræ loculi paralleli. Ovarium bilobulare loculis uniovulatis, stylus brevis stigmate crassiusculo obliquo. Fructus drupaceus, exocarpio plus minus succoso, putamine dno in pyrenas binas secedente. — Frutices foliis oppositis, floribus in spicas axillares pedunculatas saepe in capitula contractas dispositis.

1. **L. alba** (Mill. ex Link Enum. Berol. II, p. 126) fruticosa striguloso-aspera canescens, foliis petiolatis rugosis a basi truncatâ vel subcordatâ oblongis acutiusculis crenato-dentatis subtus magis canis, pedunculis siue folio longioribus, floribus capitatis, bracteis obovatis vel ellipticis tubum corollæ equantibus exterioribus foliaceis, corollâ velutinâ albâ, drupâ subbaccatâ ȝ. *L. Indica* Roxb. Fl. Ind. III, p. 89. — *L. dubia* Royle Illustr. p. 300, tab. 73. — *L. collina* Decaisne in Jacq. Voy. p. 136, t. 141.

Hab. in collibus Belutschiae (Stocks 635!).

Ar. Geogr. India, Abyssinia.

LIPPIA (L. Gen. 784. — *Zappania* Scop.).

Calyx parvus tubulosus membranaceus 2-4-dentatus. Corolla infundibuliformis tubo cylindrico, limbo obliquo subbilabiato labio superiore integro vel bilobo inferiore trifido. Stamina 4 didynama medio tubo affixa saepius inclusa, antherarum loculi paralleli. Ovarium bilobulare loculis uniovulatis, stylus brevis stigmate obliquo vel recurvo. Fructus siccus parvus calyce inclusus in pyrenas binas secedens. — Frutices (in nostris herbae) foliis oppositis, floribus spicatis vel capitatis.

1. **L. nodiflora** (L. Sp. 28 sub *Verbena*) herbacea, tota pilis adpressis centro affixis strigillosa virens vel canescens, caulis procumbentibus ad axillas radicantibus superne ascendentibus, foliis cuneato-spathulatis in petiolum brevem attenuatis parte inferiori integris superiore argute serratis obtusis vel acutiusculis, pedunculis

axillaribus solitariis folio longioribus, floribus in capitula ovata demum oblongo-cylindrica congestis, bracteis imbricatis obovatis concavis setulosis mucronatis vel muticis margine membranaceis erosulis tubum corollæ aequantibus, calyce bipartito leviter bicarinato ad carinas puberulo, corolla roseæ limbo subinæquali, fructu ovato subcompresso transverse sublatiore levi ȝ. Ic. Fl. Graec. tab. 333. — *Zappania nodiflora* Lam. III. p. 248, tab. 17. — *Lippia repens* Spr. — *Zappania repens* Bertol.

Hab. in arenosis et cultis uliginosis et humidis saepe maritimis regionis inferioris, Creta (Sibth! Sieb!), Thracia (Clarke), Pamphylia (Heldr!), Syria littoralis (Haussk! Gaill!), Egyptus inferior (Ehrenb! Auch! Schimp!), Babylonica (Noë!), Persia australis (Haussk!), bor. alis (Buhsel!), Transcaucasia in prov. Karabagh et Talysh (Mey! Hoht!), Turkomania (Kar.).

Ar. Geogr. Hispanie regnum Valentiniun, Italia media et australis, Sicilia, Africa borealis, insule Canarienses, Arabia tropica, Abyssinia, India Orientalis, China, P. Bonae Spei, America borealis, tropica et australis.

PRIVA (Adans. Fam. p. 303).

Calyx tubulosus costis 5 in dentes productis, fructifer ampliatus fructum arete amplectens ore contractus. Corolla hypocrateriformia tubo cylindrico recto vel incurvo, limbo patente obliquo subinæquali quinquelobo. Stamina 4 didynama medio tubo affixa inclusa, antherarum loculi paralleli vel divergentes. Ovarium bilocularre loculis septorum spuriorum ope mox bilocellatis, loculis uniovulatis. Stylus stigmate laterali antico subclavato. Fructus calyce aucto inclusus in pyrenas binas biloculares vel abortu uniloculares duras leves rugosas vel muricatas secedens. — Herbae foliis oppositis, floribus longe et tenuiter spicatis.

J. P. Abyssinica (Jaub. et Sp. Ill. Orient. tab. 453 et 454) basi suffrutescens pubescens, ramis herbaceis quadrangulis in spicas longas tenues abeuntibus, foliis petiolatis ovatis dentatis aentis inferioribus subcordatis caeleris basi rotundatis vel cuneatis, floribus laxis pedicello sublongioribus, corollæ calyce subduplo longiore segmentis binis posticis submajoribus, fructu calyce inflato submembranaceo pubescente et setuloso laxe inclusa tetragono-compresso transverse latiore coecis binis dorso valde rugulosis et echinalis bilocularibus indistinctibus dispermis tandem solutis constante, coecorum commissurâ biante marginibus hand intreflexis ȝ. *P. leptostachya* Aitches. Cat. p. 119 non Juss. — *P. dentata* Juss. ex parte. — *Steptium asperum* Roxb. Coron. II. p. 25, tab. 146.

Hab. in Belutschia (Stocks 571!).

Ex cl. Jaub. et Sp. loc. cit. a *P. leptostachya* Juss. et *P. Forskahlei* (Vahl sub *Verbenâ*) coecorum marginibus non valde intreflexis distincta, sed haec nota forsitan non sat firma et tunc omnes ille ad unicam speciem reducendae essent.

Ar. Geogr. Nubia, Abyssinia, India.

VERBENA (L. Gen. 32).

Calyx tubulosus quinquecostatus quinquedentatus, fructifer immutatus. Corolla infundibuliformis tubo recto vel incurvo, limbo patentí subbilabiato inæqualiter quinquelobo. Stamina 4 didynama inclusa, antherae ovatae loculis subparallelis, connectivo (in nostris) non appendiculato. Ovarium quadriloculare loculis uniovulatis, stylus apice breviter bifidus. Fructus calyce inclusus siccus in pyrenas quatuor crustaceas angustas secedens. Herbae (nostræ) floribus spicatis.

1. V. officinalis (L. Sp. p. 29) perennis, caule erecto elato tenui superne paniculato ad angulos scabridulo, foliis oblongis et oblongo-lanceolatis grosse dentatis incisis vel laciniatis vel lyratopinnatidis in petiolum brevem latum contractis, spicis longis laxis filiformibus, floribus minimis subsessilibus strictis, bracteis oblongis acuminatis calyce brevioribus, calycis adpresso setuloso dentibus brevibus acutis, corollæ lilacinae tubo sensim ampliato exerto ♀. Ic. tab. 91.

Hab. in ruderatis, ad vias, in ditione totâ a Græcia! Thracia! Cretâ (Raoul!), ad Syriam littoralem (Bl! Gaill!) et interiorem ad Damascum (Gaill!), Persiam borealem (Buhse! Bge!), Caucasum et Transcaucasiam (Ledeb.), regnum Cabulicum (Griff!).

Ar. Geogr. Europa omnis ab Angliâ et Gothiâ ad Rossiam medium, Japonia, Africa borealis, Abyssinia, India borealis, P. Bonæ spei, insulæ Canarienses, America.

2. V. supina (L. Sp. p. 29) annua hispidula canescens a basi ramosissima ramis ascendentibus vel decumbentibus, foliis petiolatis infimis oblongis incisis cito exsiccati cæteris in lacinias ovato-oblongas obtusiusculas integras vel dentatas bipinnatisectis, spicis alaribus et terminalibus solitariis breviter pedunculatis fructiferis elongatis laxis, floribus parvis subsessilibus erectis bracteâ lineari-lanceolatâ calyce breviore suffultis, calyce setuloso ad tertiam partem in dentes triangulari-ovatos fisso, corollæ caerulecentis tubo subexerto ⊕. Fl. Græc. tab. 554, — Rehb. Ic. tab. 94. — **V. procumbens** Forsk.

Hab. in argillosis et arenosis hyeme inundatis, Græcia in regione abietinâ Parnassi prope Livadia (Heldr!), Creta in ditione Sphakia (Heldr!), Anatolia occidentalis (Sibth.), Palestina in monte Carmel (Gaill!), Aegyptus inferior (Anch! Boiss! etc.), Mesopotamia et Babylonia (Ky! Noë! Haussk!), Transcaucasia (ex Stev.).

Ar. Geogr. Lusitania, Hispania, Italia australis, Hungaria, Dalmatia, Africa borealis, Nubia, insulæ Canarienses.

Tr. II. VITICEÆ Schauer.

Inflorescentia definita di-vel trichotome cymosa, cymis axillaribus vel composite paniculatis centrifuge floriferis. Ovula supra basin loculi inserta amphitropa vel subanatropa. Semina exalbuminosa.

VITEX (L. Gen. 790).

Calyx brevis campanulatus quinquedentatus. Corolle tubus brevis, limbus valde obliquus subbilabiatus quinquesfidus lobo antico maiore. Stamina quatuor distantia exserta, antheræ loculi distincti subdivergentes. Ovarium quadrilocellatum locellis uniovulatis, stylus breviter bifidus. Drupa siccata quadrilocularis. — Arbores vel frutices foliis oppositis digitatis.

1. **V. Agnus Castus** (L. Sp. 890) fruticosa vel arborea ramis obtuse quadragulis tomentosis in paniculam amplam abeuntibus, foliis longe petiolatis digitatim 5 rarius 7-foliolatis supra tomentellis viridibus subtus canescensibus foliolis lanceolatis acuminatis basi attenuatâ petiolulatis subintegris, cymis compactis subsessilibus in verticillastra remotiuscula longe spicata confertis, bracteis minutissimis vel nullis, calyce tomentoso campanulato dentibus triangularibus obtusis tubo triplo brevioribus, corollâ lilacinâ calyce triplo longiore tubo subexerto fauce inflatâ ore glabrâ, drupâ sphæricâ obtusâ calycem parum superante 5. Ic. Fl. Græca tab. 541. — Rehb. Ic. tab. 92.

Hab. secus rivos et torrentes regionis inferioris 0-1200! Græcia et ejus insulae! Macedonia et Thracia! Anatolia tota maritima! Syria littoralis (Bl! Gaill!) et interior ad Marasch (Haussk!), Palestina trans Jordanum (Weltst!), Egyptus (Auch! etc.), Tauria meridionalis (Stev!), Transcaucasia (Ledeb.), prov. Khorassan Persiae borealis (Bge!). Græcis *Ligaria*, Arabis *El Gar* audit.

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, Dalmatia, Africa borealis.

2. **V. Negundo** (L. Sp. 890) fruticosa ramis obtuse quadragulis tomentellis apice paniculatis, foliis utrinque adpresso sed subtus intensius canescensibus longe petiolatis digitatim quinquefoliolatis foliolis lanceolatis acuminatissimis integris basi attenuatis præter lateralia subsessilia pedicellatis, cymis laxifloris breviter pedunculatis oppositis subdistantibus longe racemosis, bracteis linearibus pedicellos æquantibus, calyce tomentoso oboconico sulcato dentibus triangularibus acutiusculis tubo triplo longioribus, corollâ calyce duplo longiore fauce barbatâ, drupâ globosâ calycem non superante. Wight. Ic. tab. 519.

Hab. in regno Cabulico (Griff!).

Foliola eis speciminum ex Indiâ boreali angustiora superne magis canescens, corolle tubus paulo brevior, sed characteres convenient.

Ar. Geogr. Ditio Scinde, India.

CLERODENDRON (L. Gen. 789. — *Volkameria* L. Gen. 788).

Calyx campanulatus quinquedentatus vel quinquesfidus fructifer sæpius auctus. Corollæ hypocraterimorphæ tubus sæpius longus

tenuiter cylindraceus, limbus patens quinquelobus lobis aequalibus vel antice prolatione. Stamina 4 longe exserta in alabastro involuta, antheræ loculi paralleli. Ovarium incomplete quadriloculare loculis uniovulatis, stylus longus bifidus. Drupa globosa exsucca quadripyrena vel pyrenis binatim coælitis dipyrrena, pericarpio osseo. — Fruites foliis simplicibns.

1. **C. Acerbianum** (Visian. Æg. Arab. № 95, tab. 4, fig. I sub *Volkamerid*) fruticosum inerme, ramis tomentellis, foliis viridibus tomentellis breviter petiolatis oppositis vel ternis ovatis vel oblongis acuminatis integris basi rotundatis vel cordatis, cymis axillaribus capitatis pedunculis petiolo longioribus suffultis bracteis linearibus inferioribus calyces æquantibus, calyeis hirsuti dentibus linear-lanceolatis acentis tubum dimidium æquantibus, corollæ albæ tubo tenui calyce quadruplo longiore limbi laciniis ovato-oblongis, genitalibus longissime exsertis, drupâ magnâ dipyrénâ calyce cupuliformi suffulta spongioso-rugosissimâ 5.

Hab. in Ægypto superiore Nubiæ contermino ad Nilum (Husson!).

Folia 2-3-pollicaria, corolla pollicaris glandulosa, drupa fructum *Rubi* similans Species tropica vix ditionis nostræ.

Ar. Geogr. Nubia.

TR. III. AVICENNLE Meisu.

Inflorescentia ultima centrifuge cymosa cymis parvis capitatis ad axillas pedunculatis vel corymbosis terminalibus. Ovula pendula amphitropa. Semen parce albuminosum.

AVICENNIA (L. Gen. App. № 4237).

Calyx brevis quinqnepartitus segmentis ovatis imbricatis. Corollæ tubus brevis cylindraceus rectus, limbus patens subæqualiter quadrifidus. Filamenta 4 brevia, antheræ subexsertæ loculis parallelis. Ovarium columnæ centrali quadrialatæ imperfecte quadriloculare, loculis uniovulatis, stylus brevis bifidus. Capsula magna subcarnosa valvis binis crassiusculis deliiscens. Semen abortu unicum, embryo maximus testâ destitutus intra pericarpium jam germinans cotyledonibus magnis longitudinaliter plicatis, radiculâ inferâ, plumulâ jam fructu nondum deciduo accretâ. — Arbores sempervirentes foliis oppositis coriacieis.

1. **A. officinalis** (L. Sp. p. 440) fruticosa vel arborea foliis oblongis ellipticis vel obovatis integris in petiolum brevem attenuatis supra viridibus subtus niveis, cymis capitatis panicæ et densifloris ad

axillas superiores et terminalibus subternis erasse et breviter pedunculatis, floribus sessilibus singulis bracteâ et bracteolis binis ovatis dorso sericeis calyce brevioribus suffultis, corollæ lobis e calyce semi-exsertis erecto-patulis ovatis erectiusculis supra glabris subtus sericeis, stigmatibus subsessilibus 5. *A. tomentosa* Wall. pl. Asiat. tab. 271 non Jacq.

Hab. ad littora maritima Ægypti superioris ad Ouadi Lechuma et Ouadi Gemul inter Kosseir et Ras Benass (Schweinf!). in Arabiâ petreâ ad Mare Rubrum prope Esmain (Schimp. 736!). Arabie Schora.

Ar. Geogr. Africa orientalis tropica, Natalia, Arabia tropica, India, China australis, Nova Hollandia.

ORD. XCIV. LABIATÆ.

(Juss. Gen. p. 140.)

Flores hermaphroditæ saepius irregulares; Calyx inferus persistens gamosepales quinquedentatus vel lobatus saepius bilabiatus. Corolla gamopetala hypogyna tubulosa limbo quinquesido vel lobis posticis connatis quadrifido bilabiato rarissime aequali. Stamina corollæ inserta quatuor saepius didynama interdum bina, antheræ bi vel uniloculares. Ovarium disco hypogyno insertum superum ad basin usque (rarissime ad medium) quadripartitum lobis erectis unilocularibus uniovulatis. Stylus e centro ovarii inter lobos erectus filiformis. Ovulum erectum anatropum. Fructus nuculis quatuor vel abortu paucioribus calyce inclusis liberis indehiscentibus crustaceis rarius subcarnosis constans. Semen erectum. Albumen nullum vel subnulum. Embryo rectus radiculâ inferâ. — Herbæ vel frutices foliis oppositis vel verticillatis exstipulatis.

Labiatarum Tribuum et generum dispositio.

Tr. I. OCYMOIDE.E Bth. — Stamina perfecta quatuor declinata saepius exserta, antheræ loculorum confluentia uniloculares. Corollæ labium superius in nostris quadrilobum inferius integrum.

Ocimum, Orthosiphon.

Tr. II. LAVANDULE.E — Stamina quatuor perfecta declinata inclusa. Antheræ loculorum confluentia uniloculares, Corollæ labium superius bifidum inferius trilobum.

Lavandula.

TR. III. SATUREINEÆ Bth. — Stamina perfecta quatuor rarius bina aequalia vel antica longiora, distantia, divergentia vel ascendentia.. Corollæ lobi sæpius plani, antherarum loculi (in nostris) bini paralleli vel divergentes.

Mentha, Lycopus, Origanum, Thymus, Thymbra, Zataria, Satureia, Micromeria, Calamintha, Melissa, Hyssopus, Zizyphora.

TR. IV. MONARDEÆ Bth. — Stamina perfecta bina antica. Antheræ biloculares connectivo in duobus generibus prioribus minimo, in posterioribus elongato.

Perovskia, Dorystæchas, Salvia, Rosmarinus.

TR. V. NEPETÆ Bth. — Stamina perfecta quatuor ascendentia, posticis antica superantibus, antherarum loculi divaricati, calyx sæpius 15-nervis.

Nepeta, Glechoma, Dracocephalum, Lallemantia. Hymenocrater, Hypogomphia.

TR. VI. STACHYDEÆ Bth. — Stamina perfecta quatuor sub labio superiore ascendentia parallela. antica longiora vel omnia tubo inclusa. Calyx sæpius 5-10-nervus.

SUBTR. I. SCUTELLARINEÆ Bth. — Calyx fructifer labiis incumbentibus complanato-clausus.

Scutellaria, Prunella.

SUBTR. II. MARRUBIEÆ Bth. — Calyx fructifer non clausus. Genitalia tubo corollæ inclusa.

Sideritis, Marrubium, Tapeinanthus.

SUBTR. III. LAMIEÆ Bth. — Calyx fructifer apertus. Stamina e tubo plus minusve exserta.

Chamophacos, Stachys, Betonica, Galeopsis, Leonurus, Melittis, Lamium, Wiedemannia, Molucella, Lagochilus, Ballota, Otostegia, Leucas, Phlomis, Eremostachys.

TR. VII. PRASIEÆ Bth. — Calyx, corolla et stamina *Stachydearum* sed nuculæ carnosæ nec siccae.

Prasium.

TR. VIII. AJUGOIDEÆ Bth. — Stamina (in nostris) perfecta quatuor ascendentia, antherarum loculi divaricati. Corolla ob labium superius brevissimum fissum subunilabiata. Nuculæ siccae favosæ vel reticulatae.

Ajuga, Teucrium.

TR. I. OCYMOIDEÆ.

OCYUM (L. Gen. 473).

Calyx ovatus vel campanulatus quinquedentatus dentis superioris orbiculati vel obovati membranacei marginibus decurrentibus alatus. Corollæ tubus exannulatus calyceae sæpius brevior, limbus bilabiatus labio superiore quadrisido, inferiore vix longiore declinato integro plano vel subconcavo. Stamina 3. Stylus breviter bifidus. Discus glandulas tumidas ferens. Nuculae ovoideæ laeves. Herbae vel suffrutescences, verticillastris sæpius sexfloris.

1. **O. Basilicum** (L. Sp. 833) annum glabrum, caulis erectis vel ascendentibus, foliis petiolatis utrinque angustatis ovatis et oblongis integris vel subdentatis, verticillastris intervallo calycibus longiore sejunetis, ealyce fructifero reflexo dentibus ciliatis supremo orbiculato, corollâ calyce subduplo longiore, filamentis superioribus basi denticulo auctis ♂.

Hab. in insulis fl. Dialæ in Babylonia (Noë 1034!), in arvis prope Mesched prov. Khorassan (Bge!), sed prob. introductum, ubique præsertim in australioribus cultum.

Ar. Geogr. Asia, Africa et America tropicæ.

2. **O. sanctum** (L. Mant. p. 85) perenne vel perennans pubescens, caulis ramosis inferne induratis, foliis petiolatis ovatis obtusis obtuse dentatis, verticillastris intervallo calycibus æquilongo vel breviore sejunetis, calycibus pedicello brevioribus nutantibus dente supremo obovato concavo breviter decurrente, corollâ calyceem vix superante, filamentis superioribus basi fasciculo pilorum auctis ♀.

Hab. ad Schah Bilawul Belutschiaæ (Stocks!), in regno Mascate Arabie (Auch. 5181!) forma villosa.

Ar. Geogr. India

ORTHOSIPHON (Bth. Lab. p. 25).

Calyx ovatus quinquedentatus dente superiore ovato membranaceo marginibus sæpe decurrente. Corollæ tubus exsertus, limbi labium superius 3 4-dentatum, inferius integerrimum concavum. Stamina 4 edentula. Stylus apice clavato-capitatus. Nuculae minutissimæ punctulatae. — Herbae vel suffrutescences.

1. **O. pallidus** (Bth. Lab. p. 708) annus humilis pubescens, foliis petiolatis ovatis obtuse dentatis, calycis pedicellati tandem deflexi fance nudâ dente supremo orbiculato breviter decurrente inferioribus binis longe subulato-attenuatis, staminibus corollâ brevioribus ♂.

Hab. in regno Cabulico (Griff!).

Facies et corolla brevis fere *Ocymi* sed labium inferius et calyx *Orthosiphonis*.

Ar. Geogr. India.

Obs. *Coleum Arabicum* Bth. sub nomine *Tencrui Mazziarii* ab amic. Heldreich accepi in insulâ Zaczyntho ad radices meridionales montis Xerocastello secus rivulum a cl. Mazziari lectum, sed certe non spontaneum.

TR. II. LAVANDULEÆ.

LAVANDULA (Tourn. Inst. I, p. 198).

Calyx ovato-tubulosus 13-rarius 15-nervius dentibus 5 brevibus supremo saepe apice dilatato-appendiculato, omnibus in fructu conniventibus. Corollæ labium superius submajus bilobum inferius trilobum. Stamina 4 declinata corollæ tubo inclusa, antheræ ovato-reniformes loculis confluentibus rimâ communi semicirculari dehiscentibus. Stylus bilamellatus. Discus squamis 4 carnosis nuculis antepositis auctus. Nuculæ glabrae laeves. — Herbæ vel suffrutices.

SECT. I. STÆCHAS Bth. — Bracteæ in spicam densam imbricatae, in axillis 3-5-floræ, supremæ steriles comosæ. Calycis dens superior appendiculatus.

1. L. Stæchas (L. Sp. 800) fruticosa tomentoso-canescens ramosissima ramis erectis, foliis linear-oblongis vel linearibus obtusis margine revolutis, axillis inferioribus fasciculigeris, spicis ovato-oblongis densis breviter pedunculatis, bracteis sterilibus obovatis violaceo-purpureis, corollæ violaceæ tubo calyce longiore, calycibus ovatis incanis, stigmatibus brevibus retusis 5. Ic. Fl. Græc. t. 549. — Rchb. Ic. tab. 26. — *Stæchas officinarum* Mill.

Hab. in collibus calidis regionis inferioris Græciae (Sprun! Heldr! Orph. Fl. exs. 273!), Zaczynthi (Marg!), Archipelagi (Sibth.), Cretæ (Sieb!), Byzantii et Bithyniæ (Griseb!), Lycie (Forb.), Libani et Antilibani in reg. inferiori (Gail!), Syriae ad Antiochiam (Bl!), Palestina borealis (Boiss!) et australis prope Gaza (Ky!). Fl. vere.

Ar. Geogr. Insulæ Canarienses et Madera, Lusitania et Hispania maritimæ, Gallia australis, Italia australis, Africa borealis.

2. L. Cariensis (Boiss. Diagn. Ser. I. 5, p. 3) fruticosa tenuiter velutina virens ramosissima ramis erectis, foliis linearibus margine revolutis acutiusculis, axillis inferioribus fasciculigeris, spicis longissime pedunculatis oblongis, bracteis sterilibus oblongo-lanceolatis pallidis nervosis, calycis cylindrici longe hirsuti appendice substipitata transverse latiore iobulata, corollæ violaceæ tubo calycem

non excedente, stigmatibus brevibus retusis 5. *L. pedunculata* β *Carriensis* Bth. in p.c. Prodr. 12, p. 144. — *L. spectabilis* C. Koch Linn. 21, p. 646.

Hab. in collibus Byzantii et in Bithynia (Thuret! Clem! Thirkel!), Lydia (Auch. 1763!), in monte Tmolo supra Birgui (Bal. 326!), Cariä (Pinard!).

Spicis longe pedunculatis et habitu valde affinis *L. pedunculata* Cav. Hispanicae a qua tamen specifice differre videtur indumento parco, calycis appendice lobulata nec integrâ, comâ pallidâ nec intense violacea.

SECT. II. PTEROSTÆCHAS Ging. — Bracteæ spicatae oppositæ in axillis nūfloræ, supremæ non comosæ. Calycis dens superior non appendiculatus.

3. *L. atriplicifolia* (Benth. p.c. Prodr. 12, p. 146) suffruticosa breviter cano-tomentosa ramis erectis, foliis integerrimis linearioribus, spicis subternis paniculatis densis, floribus solitariis oppositis 5.

Hab. in Ægypto superiore (Figari herb. Centr. Ital.). Non vidi.

Spicæ nutantes 1-2-pollices longæ, calyx et corolla fere *L. pinnatae*, corollæ lobi angusti.

4. *L. multifida* (L. Spec. p. 800) pubescens canescens multi-caulis, caulis flexuosis inferne ramosis, foliis in laciniis breves cuneatas obtusas trifidas vel lobulatas pinnatisectis, spicis longiusculæ pedunculatis saepius solitariis oblongis densis, bracteis hirsutis ovatis acutis nervosis calyce subaequilongis, calycis oblongo-cylindrici dentibus villosis triangulari-lanceolatis subæqualibus, corollæ tubo calyce longiore φ . Ic. Barrel 798.

Hab. in Ægypto superiore (Figari ex Bth.). Habeo specimen in horto Kahirino ex desertis Ægypti cultum.

Ar. Geogr. Lusitania, Hispania, Africa borealis

5. *L. pubescens* (Decaisne Flor. Sin. p. 8) patule pubescens viscidula, caulis erectis opposite ramosis, foliis in laciniis breves oblongo-lineares acutas bipinnatisectis, spicis solitariis vel paniculatis linearibus densifloris, bracteis breviter puberulis oblongis acuminatis nervosis calyce subbrevioribus, calycis puberuli oblongo-cylindrici dentibus triangularibus acutis supremo sublatiore, corollæ tubo calyce duplo longiore, stigmatibus ovatis φ . Ic. Jaub. et Sp. Ill. tab. 375.

Hab. in desertis Ægypti inter Keneh et Kosseir (Schweinf!), ad Ouadi Askar deserti Ægyptiaco-Arabicis (Schweinf!), Arabia petræa in rupestribus Sinaiticis (Bové! Schimp. 231!).

Affinis *L. multifidae*, differt spicis longioribus paniculatis breviter pedunculatis, bracteis calyceque breviter puberalis nec villosis, calycis dentibus inæqualibus.

Ar. Geogr. Arabia tropica.

6. **L. coronopifolia** (Poir. Dict. II, p. 398) breviter puberula vel glabra multicaulis caulibus longe stricte et opposite ramosis superne subnudis, foliis parvis in lacinias breviter et anguste lineares pinnatisectis, spicis solitariis vel paniculatis strictis longis gracillimis laxis, floribus oppositis, bracteis calyce multo brevioribus breviter lanceolatis, calyce minute puberulo cylindrico dentibus lanceolatis acutis æqualibus, corollæ tubo calyce longiore, stigmatibus ovatis ♀. *L. multifida* Burm. Fl. Ind. tab. 38, fig. I. — *L. striata* Del. Fl. Eg. I, p. 94, tab. 32.

Hab. in desertis Egypti prope Suez (Del., Ehrenb!), in Thebaide (Husson!), Arabiâ petreâ ad Ouadi Hebran (Schimp. 141!), ad Ouadi Scheik (Boiss!), Palestinâ australi prope Hebron (Roth!), Persiâ australi (Burm.). Arabice *Dictaë* vel *Siedé*.

Spicis laxis tenuissimis a *L. pubescente* et *L. multifidâ* distincta.

Ar. Geogr. Nubia, Arabia tropica.

Obs. *L. Spica* L. et *L. dentata* L. ex Prodr. Fl. Gr. in Græciâ indicatæ ibi spontaneæ non crescunt.

SECT. III. CHÆTOSTACHYS Bth. — Bracteæ spicatae alternæ in axillæ unifloræ. supremæ non comosæ. Calycis dens superior non appendiculatus.

7. **L. subnuda** (Bth. in D.C. Prod. 12 p. 148) tenuiter pubescens, caulibus virgatis opposite et stricte ramosis subaphyllis. foliis paucis parvis petiolatis oblongis paucilobatis lobis brevibus obtusis, spicis gracilibus elongatis densiusculis solitariis vel subternis breviter pedunculatis, bracteis triangulari-lanceolatis acuminatissimis, calycis cylindrici minute velutini dentibus lanceolato-subulatis æqualibus, corollæ tubo tenui calyce subdupo longiore.

Hab. in regno Mascate Arabiæ (Auch. 5220!).

Folia semipollicem longa vel minora, spicæ 1- $\frac{1}{2}$ -pollicares eas *L. coronopifoliae* referentes sed minus laxæ et flores alterni.

TR. III. SATUREINEÆ.

MENTHA (L. Gen. p. 291).

Calyx campanulatus quinquedentatus. Corolla infundibuliformis tubo inclusu sensim in limbum ampliato, limbus quadrifidus subæqualis laciniâ superiore sublatiore. Stamina 4 æqualia erecta distantia, filamenta glabra nuda, antheræ loculi paralleli rimis parallelis debiscentes. Stylus breviter bifidus, nuculæ ovatæ læves. — Herbæ perennes verticillastris in spicam aut capitalum terminale congestis, vel remotis. Species plures polygamо-dioicæ corollæ magnitudine et staminibus inclusis aut exsertis variantes.

SECT. I. EUMENTHA. — Calycis faua nuda, dentes subæquales.

1. **M. rotundifolia** (L. Sp. p. 825) caulis erectis, foliis sessilibus ovatis saepe cordatis rugosis crenatis supra minus pubescens, subtus canescentibus vel lanatis, spicis densis conico-cylindricis saepius paniculatis, bracteis lanceolatis, calycibus obsolete striatis hirtis fructiferis ovato-globosis dentibus brevibus acutis ♀. Ic. Rehb. tab. 81. — *M. rugosa* Lam. — *M. neglecta* et *M. macrostachya* Ten.

Hab. ad aquas in regione præsertim inferiore, Græcia (Sprun! Heldr!), Thracia (Friv!), Bithynia (Pichl!).

Ar. Geogr. Europa media et australis ab Angliâ et Germaniâ mediâ ad Rossiam medium, Madera, Africa borealis.

2. **M. sylvestris** (L. Sp. p. 804) caulis erectis, foliis subtus vel utrinque canescentibus vel canis rarius glabratis subsessilibus ovatis oblongis vel lanceolatis plus minusve denticulatis interdum undulato-crispis, verticillastris multifloris spicas conicas vel cylindraceas plus minus elongatas densas rarius inferne interruptas formantibus, bracteis linear-setaceis, calycibus tomentosis striatis fructiferis ventricosis superne coarctatis dentibus subulatis demum subconniventibus ♀. Rehb. Ic. tab. 82. — *M. mollissima* Bork. — *M. nemorosa* W. — *M. candicans* Mill. et multa alia synonyma quæ citare fastidiosum. — *M. nigrescens* et *M. microphylla* C. Koch Linn. 21 p. 648.

Hab. ad ripas et fossas totius ditionis a regione inferiori ad alpinam, a Græciâ! et Macedonia! ad Persiam borealem! et australem!, Turkestaniam! Egyptum!

Species quoad indumentum, foliorum formam et dentes, spicas breves vel longas admodum variabilis. Non omnes varietates sed quædam magis memorabiles numero; omnes transitus innumeros præbent.

β. *stenostachya*. — Spicæ longæ tenuiter cylindricæ inferne saepe interruptæ, flores minores, folia tomentosa vel lanata saepe plicata et margine undulata. — *M. tomentosa* Urv. Enum. p. 67. — *M. canescens* Sieb. — *M. Sieberi* C. Koch Linn. 21 p. 649.

Hab. in Græciâ!, Macedonia! et Thraciâ!, Anatoliâ occid. (Boiss!) et interiore (Wied!), Cretâ! Syriâ!, Arabiâ petreâ (Schimp. 815!), Persiâ australi (Bth.), prov. Khorassan (Bge!).

γ. *lavandulacea*. — Canescens vel cana, spicæ longæ angustæ et flores parvi præcedentis, sed folia longe et anguste linear-lanceolata plana saepe integra. — *M. lavandulacea* Willd. Enum. p. 609.

Hab. ad scaturigines Sinaiticas (Ehr! Schimp. 419!).

δ. *petiolata* — Folia petiolata ovata vel lanceolata subdentata vel serrata, spinæ variant breves vel elongatæ, planta saepius canescens vel cana. — *M. incana* Willd. En. p. 609. — *M. Royleana* Bth. in Wall. As. I, p. 29. — *M. Noëana* Boiss. in Noë exs.

Hab. in Græciâ (Sibth. ex Bth.), Libano (Gaill!), Kurdistaniâ (Ky. 395!), prope Bagdad (Noël), Persiâ boreali (Auch. 5105!), Turkestaniâ (Lehml!), regno Cabulico (Griff!).

ε. glabrata. — Viridis breviter et crispule puberula, folia subsessilia latiuscula sœpius argute dentata, spicæ crassiusculæ. — *M. Rosani* Ten. — *M. ambigua* Guss. — *M. sylvestris* var. *Kotschyana* Boiss. Diagn. Ser. II, 4 p. 5.

Hab. in Cataoniâ (Haussk!), alpibus Musch Armeniæ australis (Ky. 454!), valde Khabackar Ponti Lazici (Bal!), jago Koschan Tauri Cilicici 7800' (Ky 307!).

η. viridis. — Glabra viridis, folia breviter petiolata oblongo-lanceolata acute serrata, spicæ tenues interruptæ. — *M. viridis* L. Sp. p. 804.

Hab. in Macedonia (Friv.), prope Vodena (Orph!), ad Diarbekir Mesopotamiae (Haussk! sed an spontanea?).

Ω. suavis. — Molliter pubescens vel villosa, folia longiuscule petiolata ovato-oblonga serrata, spicæ laxæ basi interruptæ, bracteæ elongatae lanceolate. — *M. suavis* Guss. pl. rar. p. 367, tab. 66.

Hab. in humidis insulæ Tassos Macedonia (Frieder. ex Griseb.).

Videtur forma inter *M. sylvestrem* et *aquaticam* hybrida.

Ar. Geogr. Europa tota, Sibiria, insulæ Canarienses, Africa borealis, Abyssinia, India borealis, P. B. Spei.

3. *M. aquatica* (L. Sp. p. 805) caule retrorsum hirsuto, foliis petiolatis ovatis serratis basi rotundatis vel subcordatis glabratissimis vel pubescentibus, verticillastris 2-3 terminalibus remotiusculis vel in capitulum terminale congestis foliis floralibus superatis, bracteis lanceolatis, calyce oblongo sulcato dentibus a basi triangulare subulatis ♀. Ic. Rchb. tab. a.5. — *M. hirsuta* Sm.

Hab. in umbrosis et hamidis, Peloponnesus in Elide (Heldr!), Thracia, et Bithynia (Griseb.), Anatolia occidentalis (Sibth.), Syria borealis (Haussk!), Tauria, Caucasus et Transcaucasia (Ledeb.), Persia borealis (Szov.).

β. incana. — Tota lanato-tomentosa. — *M. eriantha* C. Koch Linn. 21 p. 650.

Hab. in agro Byzantino (Wied!), in Tauriâ (herb. C. Koch!).

Ar. Geogr. Europa omnis a Sueciâ ad Rossiam borealem, Sibiria Altaica et Baicalensis, Africa borealis, Madera, P. B. Spei, America borealis et tropica.

4. *M. arvensis* (L. Sp. 806), villosula vel glabriuscula, caulis ascendentibus retrorsum pilosis vel glabriusculis, foliis petiolatis ovatis dentatis basi cuneatis, verticillastris globosis fere secus totum caulem axillaribus remotis foliorum petiolo brevioribus, bracteis lanceolatis, calycis tubuloso-campanulati dentibus triangulare-laneeolatis demum porrectis ♀. Rchb. Ic. tab. 88. — *M. sativa* L.

Hab. in cultis humidis et ad fossas in ditione ut videtur rara, Byzantium (Sestini), Tauria (Stev.), Caucasus (M.B., C.A. M.).

Ar. Geogr. Europa præsertim septentrionalis et media, Sibiria, regio Himalaica, Madera.

SECT. II. PULEGUM Coss. — Calycis faux annulo pilorum clausa, dentes subbilabiati inferioribus binis angustioribus.

5. M. **Palegium** (L. Sp. 307) villosa, hirta vel glabrescens, caulis basi prostratis radicantibus, foliis ovatis vel oblongis breviter petiolatis minute et obscure denticulatis, verticillastris globosis omnibus remotis, floribus roseis, calyce hispido interdum dentibus accessoriis parvis aucto ♀. Ic. Rehb. tab. 89.

Hab. ad ripas et in inundatis totius ditionis in regione inferiori et montana a Græciâ et Macedonia! ad Persiam borealem (Szov!), Syriam! Egyptum inferiorem (Husson! Gaill!).

Ar. Geogr. Europa omnis ab Anglia et Sueciâ ad Rossiam medium, Sibiria Uralensis, insulae Canarienses, Africa borealis, Abyssinia, America septentrionalis et australis.

LYCOPUS (Tourn. Inst. p. 190).

Calyx campanulatus æqualis 4-5-dentatus fauce nudâ. Corolla campanulata subæqualis quadrifida calycem vix superans. Stamina distantia recta divergentia bina fertilia exserta antheris parallele bilocularibus, bina sterilia inclusa filiformia capitellata interdum nulla. Stylus apice bifidus. Nuculæ margine calloso incrassato trigonæ apice truncatae basi attenuatæ. — Herbe perennes verticillastris axillaribus densifloris globosis remotis.

1. L. **Europæus** (L. Sp. p. 30) glaber vel minute pubescens, caule rigido tetragono et quadrisulcato simplici vel opposite ramoso, oliis oblongo-rhombeis sinuato-dentatis in petiolum attenuatis, verticillastris multifloris, calycis dentibus triangularibus mucronatis, filamentis sterilibus minutissimis vel nullis ♀. Ic. Fl. Dan. tab. 1081. — Rehb. Ic. tab. 90. — *L. decrescens* C. Koch Linn. 21 p. 646!

Hab. ad aquas regionis inferioris et montanæ, Peloponnesus (Heldr!), Attica (Orph!), Cephalonia (Heldr!), Macedonia (Orph!), Byzantium et Bithynia (Wied!), Lydia (Heldr!), Galatia (Wied!), Armenia et Pontus Lazicus (Calv' Bal!) et interior ad Damascum (Gaill!).

Ar. Geogr. Europa omnis ab Anglia et Sueciâ australi ad Rossiam, Sibiria Altaica et Baicalensis, Africa borealis.

2. L. **exaltatus** (L. fil. Suppl. p. 87) pubescens, caule elato rigido tetragono ramoso, foliis oblongis pinnatifidis partitisque in petiolum brevem attenuatis, verticillastris parvis, calyce acute quinque-dentato, filamentis sterilibus acute capitatis ♀. Ic. Fl. Græc. tab. 12. — *L. pinnatifidus* Pall. Fl. Ross. II. tab. I.

Hab. in paludosis Thraciae (Friv.), Bithyniae (Sibth., Boiss!), Paphlagoniae (Wied!), Tauriae et regionis Caucasicæ (M.B., Rehm!).

Ar. Geogr. Regio Danubialis, Germania, Italia borealis, Dalmatia, Rossia, Sibiria Uralensis et Altaica.

ORIGANUM (L. Gen. 726).

Calyx campanulatus vel turbinatus bilabiatus antice fissus vel subæqualiter quinquedentatus. Corollæ tubus inclusus vel exsertus, limbi labium superius rectum retusum vel bifidum. Stamina 4 distantia superne patula, antheræ biloculares loculis distinctis divergentibus, Stylus bifidus. Nuculæ ovatae vel oblongæ lœves. — Suffrutices vel herbæ perennes floribus bracteatis plus minus dense in spicas ovatas vel cylindricas imbricatis.

Specierum Orientalium dispositio.

SECT. I. AMARACUS Bth. — *Amaracus* et *Anatolicon* Bth. in B.C.
Prodr. — Bracteæ amplæ sëpe coloratae calyces occultantes. Calyx bilabiatus.

* Calycis labium superius integrum.

O. Tournefortii, Dictamnus.

** Calycis labium superius plus minus profunde tridentatum.

+ Folia basi non cordata

O. pulchellum, Sipyleum, Libanoticum.

++ Folia basi plus minus cordata.

SECT. II. EUORIGANUM Bth. — Bracteæ mediocres Calyx subæqualiter quinquedentatus.

* Corollæ tubus longe exsertus.

O. levigatum, Ehrenbergia.

** Corollæ tubus subinclusus.

O. vulgare, hirtum

SECT. III. MAJORANA Bth. — Bracteæ floresque minimi dense imbricati.
Calyx turbinatus.

* Calyx bilabiatus.

O. micranthum.

** Calyx truncatus.

O. microphyllum.

*** Calyx antice fissus unilabiatus.

O. Maru, duvium, Onites.

SECT. I. AMARACUS.

1. **O. Tournefortii** (Sibth. in Ait Kew Ed. I, 5 p. 311) glabrum vel patule lanatum, rhizomate suffrutescenti multicauli, caulinis ascendentibus pumilis subimalicibus tota longitudine crebre foliatis, foliis sessilibus glaucis orbiculatis subcordatis integerrimis intimis minoribus ovatis, spicis terminalibus foliis summis involucratis solitariis vel 2-3-nis erectis oblongis, bracteis purpureis late ellipticis calyces superantibus, calycis labio superiore integerrimo obovato inferiore parvo bidentato, corollæ roseæ calyce triplo longioris tubo antice medium versus gibbo ♀. Ic. Fl. Græc. tab. 369. — *Amaracus Tournefortii* Bth. Lab. p. 334. — *Or. calcaratum* Juss. — *Origanum Dictamni Cretici* facie Tourn. Voy. p. 240.

Hab. in rupibus verticalibus insulæ Amorgos Cycladum ad Mariae canonium (Tourn., Sibth., Orphant).

Caulis spithamei, folia intermedia 8-9 lineas longa. Differt ab *O. Dictamno* caulinis omnino foliatis, corollæ tubo longiore ecalcarato, etc.

2. **O. Dictamnus** (L. Sp. 823) albo-lanatum, rhizomate suffrutescenti multicauli, caulinis ascendentibus pumilis inferne foliosis superne subnudis simplicibus vel parce panicleato-ramosis, foliis pannosis orbiculatis inferioribus breviter petiolatis cæteris sessilibus omnibus integris, spicis erectiusculis ovatis vel oblongis pedunculatis, bracteis coloratis calyces superantibus, calycis labio superiore integerrimo ovato inferiore obsoleto subretuso, corollæ roseæ calyce duplo longioris tubo non gibbo ♀. Ic. Bot. Mag. tab. 298. — *Amaracus Dictamnus* Bth.

Hab. ad rupes umbrosas faucium Cretæ in regione inferiore et montana ad 4500' usque fere ubique (Sibth., Sieb!, Heldr! Raul! etc.). Vulgo *Dictamnos*, *Stamatochorton*, *Airontas*.

3. **O. pulchellum** (Boiss. Diagn. Ser. II, 4 p. 9) pilosum cinerascens, caulinis ascendentibus longe anguste et unilateraliter racemoso-paniculatis, foliis ovatis obtusis integris inferioribus breviter petiolatis cæteris sessilibus, spicis nutantibus ovatis vel oblongo-prismaticis pedunculis remote foliolosis suffultis, bracteis rubellis ovatis obtusis glabris vel ciliatis calyces occultantibus, calycis glabri fauce barbati labio superiore ovato repando-tridentato inferiore brevissimo late et breviter bilobo, corollæ roseæ tubo calyce duplo longiore non gibbo ♀.

Hab. ex Oriente in Horto Parisiensi sub nomine *O. Dictamni* vel *pseudodictamni* cultum.

Ab *O. Dictamno* paniculâ secundâ polystachyâ, spicis nutantibus, foliis minoribus non orbiculatis, bracteis minoribus 2 $\frac{1}{2}$ -3 lineas longis distinctum.

4. **O. Sipyleum** (Linn. Sp. p. 823) inferne patule pilosulum cæterum glabrum, caulinis virgatis flexuosis ascendentibus elatis longe

paniculatis, foliis parvis ovatis glaucis integris turionum villosulis subpetiolatis caulinis sessilibus acutiusculis, paniculæ ramis foliolosis saepe deflexis, spicis pedunculatis ovato-oblongis, bracteis glabris elliptico-ovatis obtusiusculis calyces occultantibus, calycis labio superiore dentibus ternis brevibus triangularibus inferiore breviore in dentes angustiores profunde bifido, fauce pilosâ, corollæ tubo calyce duplo longiore obsolete gibbo ȝ. Ic. Fl. Græc. tab. 570.

Hab. in rupestribus calcareis Anatoliæ, mons Sipylus Lydiæ (Sibth!, Boiss! Bal!), Phrygia ad Argani (Mitch) et ad Ouchak (Bal!), Caria (Pinard!), Pisidia (Heldr!). Loci Græci et Syriaci hic citati ad alias spectant species.

5. **O. Libanoticum** (Boiss. Diagn. Ser. I, 5, p. 14) basi pilosulum cæterum glabrum, rhizomate suffrutescenti multicauli, caulis ascendentibus foliosis saepius elatis paniculatis, foliis pallide virantibus ovatis obtusis breviter petiolatis subdissitis, floralibus valde diminutis sessilibus, spicis pedunculis tenuibus nulis suffultis subnutantibus ovatis tandem laxiusculis, bracteis amplis rubellis ovatis acutiusculis calyces occultantibus, calycis labio superiore in dentes ovatos obtusos ciliatulos ad medium fisco inferiore breviore in dentes lanceolatos acutos partito, corollæ roseæ tubo calyce plus duplo longiore non gibbo ȝ.

Hab. in siccis regionis montanæ et subalpinæ Libani circa Eden, Becharré, Marserkis et in ditione Dauniæ (Ehr! Auch. 1656 bis! Boiss! Bl!, Ky. 250!).

Pedale vel bipedale, folia majora 7-8 lineas longa, spicæ tandem pollicares. Ab affinibus *O. Sipyleo* et *cordifolio* foliis pedunculatis, labii calycini superioris dentibus ovatis obtusis nec breviter triangularibus acutis, etc., distinctum.

6. **O. ciliatum** (Boiss. et Ky. Diagn. Ser. II, 4 p. 10) patule hirtum pallide virens, caulis numerosis tenuibus erectis vel ascendentibus crebre foliosis simplicibus vel breviter paniculatis oligostachyis, foliis ciliatis sessilibus ovatis subcordatis obtusis, spicis breviter et tenuiter pedunculatis ovatis laxiusculis junioribus subnutantibus, bracteis coloratis ovato-spathulatis acutiusculis ciliatis calyces occultantibus, calycis labio superiore elongato oblongo apice ciliato breviter tridentato, inferiore quadruplo breviore dentibus binis lanceolatis, fauce ciliatâ, corollæ roseæ tubo inclusu limbo basi gibbo ȝ.

Hab. in rupestribus umbrosis regionis montanæ superioris Tauri Cilicici inter Gulek Boghas et Gulek Maaden (Bal. 542!), in fauibus Agatsh Kisse 6000' (Ky. 238!).

Spithameum, folia majora 8-9 lineas longa, ab omnibus affinibus corollæ tubo inclusu distinctum.

7. **O. cordifolium** (Month. et Auch. Ann. Sc. Nat. 1836, 6 p. 43 sub *Amaraco*) glabrum glaucum, caulis ascendentibus rubellis subsimplicibus remotiuscule foliosis, foliis sessilibus cordatis ovatis acutiusculis integris vel subrepandis margine lævibus superioribus sensim diminutis, spicis ovatis, bracteis rubellis magnis orbiculari-rhombeis cuneatis truncato-subapiculatis calyces longe supe-

rantibus, calycis labio superiore breviter tridentato inferiore angustiore et subbreviore in dentes binos lanceolatos obtusos profunde bifido, corollæ roseæ calyce $2\frac{1}{2}$ -plo longioris tubo apice subgibbo ♀.

Hab. in Cypro ex cl. Bth. sed in Catalogo Kotschyano floræ Cypriæ non citatur, vel in Syriâ (Auch. 1616!).

Caules rubri pedales et longiores tenues, folia majora 6-7 lineas longa, spicae eis precedentium paulo laxiores, bractæ fere 5 lineas longæ et late. Calyx fere *O. Sipylei* sed folia et bractæ diversissimæ.

8. O. leptocladum, glaberrimum glaucescens, caulis ascendentibus inferne crebre foliatis superne paniculatis paniculae depauperatae ramis patentibus gracillimis, foliis parvis sessilibus coriaceis cordato-ovatis obtusis nervosis integris punctatissimis, superioribus minimis remotis, ramealibus squamiformibus acutis, spicis anguste linear-oblängis laxiusculis, bracteis rubellis glabris elliptico-linearibus acutis calyce subbrevioribus, calycis glabri fauce villes à labio superiore triplo majore ad tertiam partem inferioreque fere ad basin in dentes triangulares acutos fissis, corollæ roseæ tubo calyce sesqui-longiore ♀.

Hab. in monte Yamourdabadagh et circa Ermenek Cilicie Tracheæ 3500-4200! (Peronin!).

Pedale vel sesquipedale, folia majora 6 lineas longa, spicae 6-10 lineas longæ vix 3 lineas lata, bractæ 3 lineas longæ $4\frac{1}{2}$ latæ. Calyx $2\frac{1}{2}$ lineas, corolla $4\frac{1}{2}$ lineas longi, facies *O. levigati* sed *Amaracus*. In sectione spicis angustis et bracteis parvis insignis.

9. O. scabrum (Boiss. et Heldr. Diagn. Ser. I, 7, p. 48) inferne sub lente pruinosum ceterum glabrum pallide virens, caulis ascendentibus erectis apice breviter racemosis, foliis sessilibus nervosis coriaceis punctatis cordato-ovatis acutiusculis margine sebris. spicis ovatis pedunculis brevissimis nutantibus foliolosis suffultis, bracteis coloratis ovatis acutiusculis calyces occultantibus, calycis glabriuscili punctati labio superiore ad tertiam partem obtuse tridentato, inferiore triplo breviore in dentes anguste lanceolatos acutos bipartito, corollæ tubo calyce $2\frac{1}{4}$ -plo longiore non gibbo ♀.

Hab. in rupestribus regionis sylvaticæ Taygeti circa Hagia Varvara (Heldr!), in monte Malevo 4000' (Orph. Fl. Gr. exs. 42!).

Sesquipedale, folia majora fere pollicaria.

10. O. pulchrum (Boiss. et Heldr. Diagn. Ser. II, 4, p. 11) glaberrimum, rhizomate pluricauli, caulis ascendentibus crebre foliosis pumilis parce paniculato-ramosis, foliis sessilibus cordatis ovato-orbiculatis acutiusculis coriaceis nervosis punctatis margine leviusculis, spicis ovatis pedunculatis nutantibus, bracteis purpureis ovato-ellipticis obtusis flores occultantibus, calycis glanduloso-punctati labio superiore ad medium in dentes ovatos fisso, inferiore vix breviore in dentes lanceolatos acuminatos bipartito, corollæ roseæ tubo calyce duplo longiore non gibbo ♀.

Hab. in lapidosis mobilibus partis superioris montis Delphi Eubeæ 4600'-5500' (Heldr. Herb. Norm. 784!), in monte Condyli Eubeæ 4000' Orph!).

Affine *O. scabro* a quo differt foliis margine lœvibus, spicis longius pedunculatis, calycis enerviæ nec elevatim nervosi labiis subæquilongis nec valde inæqualibus. Hæc species in loco natali formas cum *O. hirta* hybridas gignit.

11. *O. rotundifolium* (Boiss. Diagn. Ser. II, 4, 11) basi birtulim cæterum glabrum pallide virens, caulis ascendentibus pumilis dense foliosis simplicibus monostachyis vel apice oligostachyis, foliis sessilibus elevatim nervosis subcoriaceis ovato-orbulatis basi anguste cordatis obtusissimis margine lœvibus, spicis sessilibus vel brevissime pedunculatis ovatis, bracteis maximis membranaceis albis glabris orbiculatis basi cuneatis, calycis glabri labio superiore in dentes ovatos acutiusculos ad tertiam partem fiso, inferiore subbreviore in dentes lanceolatos obtusiusculos ad basin partito, corollæ albidae tubo calyce duplo longiore non gibbo limbo apice e bracteis subexerto ♂.

Hab. in regione montanâ Armeniæ Turcicæ, in prov. Batoum (Calv!), circa Artwin (C. Koch!), inter Gumuschkhanè et Kerekli (Tchih!).

Semipedale, folia 9-10 lineas longa, bracteæ foliis fere maiores diametro 9 lineas latæ.

12. *O. Haussknechtii*, glabriuscum, caulis e rhizomate indurato ascendentibus rigidis paniculatis remotiusculæ foliosis, foliis parvis coriaceis punctatis nervosis cordato-ovatis obtusis margine sebris, floralibus sensim diminutis, spicis oblongis, bracteis coloratis glabris ovatis obtusis calyces occultantibus, calycis glanduloso-punctati labio superiore ad medium in 3 dentes ovatos obtusos fiso, inferiore subbreviore in dentes binos lanceolatos obtusos ad basin partito, corollæ roseæ tubo non gibbo calyce duplo longiore ♂.

Hab. in monte Akdag Cataoniae supra Adiaman 5000' (Haussk!).

Hæc species quam prius pro *O. rotundifolii* varietate habuimus quod formâ calycis refert ab eo certe differt foliis multo minoribus (5-6 lineas longis) disstis et margine scabris, præsertim bracteis triplo minoribus ovato-spathulatis 4-5 lineas longitudine non excedentibus.

SECT. II. EUORIGANUM.

13. *O. lœvigatum* (Boiss. in Ann. Sc. Nat. 54, 2, p. 252) glaberrimum, caulis basi procumbentibus vel ascendentibus elongatis gracilibus rigidulis elatis paniculatis, foliis parvis pallide virentibus sessilibus elliptico-oblongis acutiusculis integris vel obscure repandis, spicis breviter pedunculatis brevibus paucifloris laxiusculis, bracteis virentibus anguste ellipticis acutis subcarinatis calycei subæquilongis, calycis glabri imipunctati fauce villosi dentibus æquilongis triangulari-lanceolatis obtusiusculis tubo triplo brevioribus, corollæ purpureæ calyce triplo longioris tubo in limbum parvum sensim ampliato ♂.

Hab. in Antitauro Cataoniae meridionalis (Tchih!), in dumosis inter Beilan et Alexandrette Syriae borealis (Haussk!).

Bipedale, folia majora 9 lineas longa, spiculae terminales fasciculiformes, calyx $2\frac{1}{2}$ lineas, corolla 7 lineas longi. Species distinctissima facie *Amaraci* sed bracteis minutis et calyce ab eis remota, ex *O. compacti* affinitate sed spicae multo breviores laxae, etc. etc.

14. O. Ehrenbergii, inferne patule et parce hirtulum cæterum glabrum pallide virens, caulis elatis simplicibus, foliis ovatis obtusis glandulosso-punctatis deflexis inferioribus brevibus petiolatis cæteris sessilibus, spicis subsessilibus ad axillas oppositis inter se remotis oblongis laxiusculis interdum ramosis longe et anguste racemosis, bracteis spathulato-oblongis acutis glaberrimis glandulosso-punctatis calyce sublongioribus, calycis glabri crebre rubro-punctati fauce villosi dentibus aequilongis lanceolatis tubo duplo brevioribus, corollæ purpureæ calyce triplo longioris tubo in limbum sensim ampliato ♀.

Hab. in Libani monte Sanin (Ehr!), in arenosis ferrugineis inter Zachle et Beckfaya (Gaill!), ad Solima (Bl!).

Caules sesquipedales a tertia parte inferiori ramosi, spicæ angusta 6-8 lineas longæ, calyx sesquilineam, corolla $4\frac{1}{2}$ lineas longa. Species inflorescentiæ peculiaris et corollæ tubo longe exerto ab omnibus *O. vulgaris* et *hirti* formis discedens, corollæ forma *O. herigato* affinis.

15. O. vulgare (L. Sp. p. 824) viride plus minus hirsutum foliis petiolatis ovatis vel oblongis repandis, spicis densifloris oblongis vel elongatis prismatico-cylindricis in paniculam plus minus compositam sœpe corymbiformem dispositis, bracteis sœpius coloratis eglandulosis ovato-ellipticis acutis calyci aequilongis vel sublongioribus, calycis dentibus brevibus ovatis aequilongis acutis, corollæ tubo breviter exerto ♀. Ic. Rehb. tab. 61. — *O. Heracleoticum* Rehb. exc. 313 et prob. Linu. — *O. macrostachyum* et *megastachyum* Link (forma specis elongatis). — *O. Creticum* L. Sp. 823 et herb.

Hab. in montanis siccis et sylvaticis regionis montanae et subalpinæ, Graecia (Sibth.), Macedonia (Griseb.), Anatolia borealis, Libanus circa Eden (Bl!), Tauria, Caucasus et prov. Transcaucasicae (Led., Hohen!). Persia borealis (Bge!).

β . *viride*. — Bracteæ virides sœpe diminutæ glabrae vel pilosulae. — *O. parviflorum* Urv En. — *O. virens* C. A. Mey et Auct. mult. non Hoffm. et Link. — *O. hirtum* Auct. plur. non Vogel. — *O. pruinatum*, *angustifolium* et *gracile*. C. Koch Linn. XXI, p. 661. — *O. normale* Don.

Varietas typo in Oriente multo copiosior, Olympus Thessaliae (Heldr!), montes Macedoniae (Heldr! Orph!), Byzantium (Noë!), Anatolia bor!, Armenia et Pontus Lazicus (Bourg! Ky!), Cilicia (Bal!), Cataoniae Berydaghi (Haussk!), mons Avromen Kurdistaniae (Haussk!), Caucasus et Transcaucasia!, Persia borealis (Szov! Buhse!).

γ . *magnilimbis*. — Omnia var. β , sed corollæ limbus paulo major. — *O. albiflorum* C. Koch loc. cit.

Hab. in Bithynia (C. Koch!), Ponto Lazico circa Djimil (Bal!).

Varietas β saepe confunditur cum *O. virenti* L. et *H. Lusitanico* et *Hispanico* paniculā in thyrsū constrictā, bracteis calycis proportionē majoribus acutioribus magis membranaceis distincto.

Ar. Geogr. Europa omnis a Scandinaviā ad Rossiam septentrionalem, Sibiria, regio Himalaica.

16. ***O. hirtum*** (Vogel Linn. XV, p. 80 au Link ?) pallide virens plus minusve hirtum, foliis ovatis integris vel subrepandis inferioribus breviter petiolatis cæteris sessilibus, spicis densifloris ovatis vel oblongis glomeratis racemoso-paniculatis, bracteis viridibus parvis ovatis aentis glabris vel sæpius hirtis facie interiore glanduloso-punctatis calyce sublongioribus, calycis glanduloso-punctati dentibus aequalibus lanceolatis acutis, corollæ tubo subincluso φ . *O. Heracleoticum* Bth. Lab. p. 336. — *O. Smyrneum* Sibth. et Sm. Fl. Græc. VI, p. 57, tab. 571 non L. (hūc ob bracteas glanduliferas potius quam ad *O. vulgare viride* spectare videtur). — Griseb. Spic. II, p. 115. — *O. neglectum* Vogel Linn. 45 p. 81.

Hab. in siccis regionis inferioris et montanæ, Laconia in Taygeti regione infer. (Heldr!) et Malevo (Orph. Fl. exs. 528!), Messenia (Bory!), Arcadia (Heldr!), mons Delphi Eubœa (Heldr!), Cephalonia (Heldr!), Macedonia ad Athonis radices (Friv! Pichler!), Creta (Sieb! Heldr!), Cyclades (Orph!), Lydia (Sibth., Bal. 319!), Cilicia (Ky. 472!).

Valde affine *O. vulgare* β *viridi* quocum a nonnullis botanicis conjungitur et a quo tantum differt bracteis minoribus latioribus intus glandulosis, calyce magis glanduloso. An tamen hujus varietas? *O. glandulosum* Det. hūc sæpe relatum differt bracteis calyce brevioribus, calycis dentibus brevissimis.

SECT. III. MAJORANA.

17. ***O. micranthum*** (Vogel Linn. XV. p. 77) totum adpresse tomentosum canum, caulibus ascendentibus crassiusculis paniculatis, foliis minimis ovatis vel orbiculatis obtusis integris inferioribus breviter petiolatis cæteris sessilibus, spicis numerosis congestis minimis ovato-oblongis densis, bracteis canis orbiculatis obtusissimis eucullatis marginibus inflexis calycem subæquilonum includentibus, calycis breviter turbinati hirti ore villosi labio superiore sublongiore obsolete tridenticulato denticulis obtusissimis inferiore breviter bidentato dentibus triangularibus, corollæ roseæ tubo inclusō φ .

Hab. in Tauro Cilicie prope Gülekboghaz 4500'-5000' (Ky. 261! et 471! Bal!).

Planta semipedalis rarius pedalis, folia majora 5 lineas longa, spicule demum 4-5 lineas longæ, corolla 1 $\frac{1}{4}$ lineas longa.

18. ***O. microphyllum*** (Sieb. exs. — Bth. Lab. 338 sub *Majorana*) basi procumbenti radicans, ramis basi ranulosis erectis tenuiter virgatis rubellis glabriusculis anguste paniculatis, foliis minimis albo-tomentosis remetis breviter petiolatis ovatis et oblongis integris, spicis paucis subglobosis terminalibus solitariis vel ternis

breviter pedunculatis albo-lanatis, bracteis spathulato-ovatis obtusis calyce longioribus, calyce glabro glanduloso-punctato turbinato oblique truncato edentulo fauce barbatâ, corollæ roseæ calyce vix duplo longioris tubo vix exerto ♀. *O. Maru* Fl. Græc. VI. p. 59. t. 573. — Bot. Mag. tab. 2603 non L.

Hab. in saxosis apriis regionis montanae Cretæ, montes Sphaciotici prope Omalο 3500' (Sieb! Sibth., Heldr! Raul!).

Pedale vel sesquipedale, folia majora vix tres lineas longa, capitula piso minora, calyx vix $1\frac{1}{4}$ lineam, corolla $2\frac{1}{2}$ lineas longi.

19. O. Maru (L. Sp. p. 826) basi suffruticosum, ramis erectis breviter et patule hirtis, foliis subsessilibus ovatis obtusis integris crassis subtus elevatim nervosis, spicis oblongis et demum anguste cylindricis elongatis in paniculæ ramis glomeratis, bracteis tomentosis quadrifariis ovato-spathulatis obtusis calyci aequilongis, calyce ovato-spathulato antice fisso unilabiato obtusissimo integro vel obsolete retuso, corollæ albæ tubo inclusu ♂. *O. Syriacum* Sieb. exs. et prob. Linnei. — *Majorana crassifolia* Bth. Lab. p. 339. — *O. restitum* Clarke?

Hab. in regione inferiore et montanâ Syriae, Amano (Ky. 48 sub *O. Onite*!), Libano circa Sidonem, Tripolin, Eden, Ghazir, Dimam, Bescherre (Bl! Gaill! Ky. 260!), in Syria (Auch. 4658!), circa Antiochiam (Boiss!), in Galileâ (Gaill!), circa Hierosolymam (Sieb! Roth!), in ditione Golan (Wettstein!). Vulgo *Satar*.

♂ *Sinaicum*. — Indumentum minus canum magis patulum. — *O. nerrosum* Vogel loc. cit. p. 78. — *O. Egyptiacum* L. saltem ex parte.

Hab. in montibus Sinaiticis (Ehrenb! Bové! Schimp. 385!).

Ab *O. Onite* distinctum foliis canis integris subtus nervosis, inflorescentiâ paniculata nec complanato-corimbosa, spiculae longiores sunt et flores sub-majores.

20. O. dubium, basi suffruticosum, ramis erectis foliosis glabriusculis vel tomentellis, foliis petiolatis adpresso canescensibus ovatis obtusissimis integris nervis subtus non elevatis, spicis ovatis glomeratis paniculatis, bracteis tomentellis ovato-spathulatis obtusis calyce sublongioribus, calyce ovato-spathulato antice fisso unilabiato obtuso, corollæ albæ tubo inclusu ♂.

Hab. in insulâ Naxos Archipelagi (Lenormand!), argillosis Cypri inter Panteleimon et Paleomilo (Ky. 937! sub *O. Majorana*!), Cilicia Trachea (Peronin!). faucibus Guzulderé prope Mersina (Bal. exs. sub *O. Maru*!).

Simile *O. Majorana* L. (cujus specimina culta tantum vidi) calyce semper bilabiato donati cuius labium inferius parvum quidem sed manifestum atque bidentatum est. *O. dubium* ab *O. Maru* foliis petiolatis obtusissimis subtus non nervosis differt. ab *O. Onite* inflorescentiâ paniculata, foliis integris.

21. O. Orites (L. Sp. p. 824) basi suffruticosum ramis erectis crassis patule papillari-hirsutis et minute glandulosis, foliis tomentellis canescensibus ovatis subserratis insimis brevissime petiolatis

cæteris sessilibus acutis, spicis ovatis compactis in corymbum planiusculum compositum congestis, foliis floralibus oblongo-lanceolatis pedunculos patentibus non æquantibus, bracteis quadrifariis ovato-orbiculatis acutiuseculis nervosis hirtis eucullatis calyce sublongioribus, calyce ovato-spathulato antice fisso unilabiato obsoletissime tridenticulato, corollæ albæ tubo inclusu 5. Ic. Fl. Græc. tab. 572.

Hab. in collibus regionis calidæ Peloponnesi (Sprn! etc.), Cycladum (Urv! Orph!), Rhodi (Bourg!), Cretæ (Sieb! Raul! Heldr!), Lydiæ ad Smyrnam (Fleisch! Boiss!), Lyciæ (Bourg!), Pamphyliæ (Heldr!), Syriæ (Auch. ex Bth.).

Ar. Geogr. Sicilia nec Dalmatia ubi indicatur.

THYMUS (L. Gen. 727).

Calyx ovoideus 10-18-nervis bilabiatus fauce villosus, labium posticum latum tridentatum dentibus triangularibus vel lanceolatis, anticum dentibus binis subulatis constans. Corollæ tubus inclusus vel exsertus intus nudus, limbi subbilabiati labium posticum subplanum retusum, anticum patens trifidum. Stamina 4 recta distantia, antheræ biloculares loculis parallelis vel divergentibus. Stylus bifidus. Nuculæ ovatae vel oblongæ læves. — Herbae perennes vel suffrutices.

SECT. I. SERPYLLUM. — Calyx teres, corollæ tubus inclusus vel vix exsertus.

1. T. teucroides (Boiss. et Spr. Diagn. Ser. I. p. 15) suffrutescens ramosissimus, ramis breviter puberulis basi induratis procumbentibus dein erectis pumilis, foliis minutis petiolatis ovato-rhombeis obtusiusculis glanduloso-punctatis subtus obsolete puberulis, floralibus conformibus paulo minoribus, verticillastris axillaribus 2-6-floris distinctis raxe spicatis, pedicellis calyci æquilongis, bracteis oblongis pedicello brevioribus, calycis breviter campanulati ad medium bilabiati glabri vel pruinosi basi subgibbi labio superiore subbreviore dentibus tribus breviter et late triangularibus mucronulatis inferiore in dentes a basi lanceolata subulatos ciliatos ad basin fisso, corollæ roseæ amplæ tubo breviter exerto 2. *T. graveolens* Fl. Græc. VI, p. 61, tab. 576 non M.B.

Hab. in regione alpinâ Parnassi (Sibth., Heldr! Orph!), monte Æta et prope Zeitun (Sprn!), monte Chelmos Achaiae 7000' (Heldr!).

Rami cum racemo 6-8-pollicares, folia vix tres lineas longa lataque, corolla fere 5 lineas longa. Facies *Calam. Alpinae* sed stamina *Thymi*; affine *T. Pipe-relle* Hispanico ramis crassis rigidis, floribus floralibus bracteisque minimis, calycis labio superiore inferius superante, etc., distincto.

2. T. Serpyllum (L. Sp. 482) cespitosus, ramis inferne prostratis et radicantibus vel ascendent-erectis seriatim vel undique puberulis, foliis ellipticis oblongis vel linearibus basi plus minus

attenuatis vel cuneatis raro rotundatis breviter petiolatis, verticillastris sessilibus in capitulum congestis vel inferioribus plus minus distantibus racemosis, foliis floralibus subconformibus, pedicellis calyce subbrevioribus, calycis plus minus hirti ad medium bilabiati labio superiore sublongiore dentibus triangularibus glabris vel ciliatis inferiore in dentes ciliatos lanceolato-subulatos bipartito, corollæ roseæ tubo subincluso ♀. Species valde polymorpha cuius formæ mirum in modum confluent, frequentiores numero.

α. genuinus. — Rami tenues elongati, folia parva plana linear-elliptica glabriuscula, flores capitati. — Ic. Rehb. 65, 1 et 2.

Hab. in pratis præsertim regionis montanæ et alpinæ, Macedonia in Scardo (Griseb.), mons Delphi Eubeæ et cacumen Athonis (Orph!), Olympi Bithyni regio superior!, Pontus Lazicus supra Trapezuntum (Calv!), circa Djimil (Bal!), Caucasus (Ledeb.).

β. latifolius. — Rami erectiores elongati, folia majora ovato-oblonga, verticillastra inferiora interdum dissita. — *T. Chamædrys* Fries. Nov. 196. — *T. montanus* W. K. Hung. I, p. 72, tab. 71. — *T. collinus* M.B. III, p. 401. — *T. pulchellus* C. A. M. Enum. p. 87 ex specimine archetypo! lusus foliis angustioribus ad var. *angustifolium* vergens.

Hab. in Thraciâ et Macedoniâ (Griseb.), Caucaso et Transcaucasiâ (Led., C.A. Mey., Szovl).

γ. nummularius. — Rami a basi prostratae erecti fragiles patule hirsuti, folia ovata vel orbiculata basi saepè rotundata vel etiam subcordata, capitula globosa foliis summis involucrata, corolla speciosa limbo majusculo, staminibus exsertis. — *T. nummularius* M.B. Taur. Cauc. II, p. 58. Bot. Mag. tab. 2666!

Hab. in regione alpinâ Ponti Lazici supra Djimil 6000' (Bal!), Alpibus Caucasicis (M.B.), Daghestania 6000'-9000' (Rupr!).

Rami 4-5-pollicares, specimina quædam foliis subcordato-orbiculatis propriam sistere speciem videntur sed intermediae formæ occurunt.

δ. Marschallianus. — Rami elongati hirtelli vel villosi, folia saepius elongata oblonga et linear-elliptica ciliata vel undique hirsuta. — Rehb. tab. 6, № 2. *T. Marschallianus* Willd. Sp. 141. — *T. Pannonicus* Bth. in D.C. Prodr. et All. Ped. I, p. 20 (forma foliis brevioribus). — *T. Raegneri* C. Koch Linn. XXI, p. 666.

Hab. in Græciâ (Auch. 1642!), Macedonia monte Peristeri (Orph!) et Kor thiati (Heldr!), Tauriâ, Caucaso et Transcaucasiâ (C. Koch! Hoh!), Persia australi in Alpe Kuh Delu (Ky. 560!).

δ. bis. Ocheus. — Totus breviter et patentim canescens-setulosus, rami abbreviati, folia parva ovata et oblongo-spathulata, flores terminales capitati foliis summis involucrati. — *T. Ocheus* Heldr. et Sart. Diagn. Ser. II, 4, p. 4.

Hab. in summo monte Hagios Elias (Oches) supra Karisto Eubeæ (Sart. exs. sub *T. integræ*).

ε. Chaubardi Boiss. et Heldr. Diagn. Ser. II, 4, p. 6. — Rami

elongati varietatis δ , sed glabri vel retrorsum tenuiter puberuli. Folia elongata versus basin ciliis paucis instructa vel omnino glabra, verticillastra congesta vel inferiora subdistantia. Dentes calycis saepius omnes ciliati. — *T. lanceolatus* Sm. Prodr. 419 non Desf. *T. Sibthorpii* Bth. Lab. 345. — *T. glabratus* Bory et Chaub. Pelop. non Link. — *T. Chaubardi* Boiss. et Heldr. in Orph. exs. — *T. heterotrichus* Griseb. Sp. II. p. 416.

Hab. in pascuis siccis montanis et alpinis, Græcia in Taygeto (Heldr!), Messeniâ (Bory!), montibus Parnes et Ozia Atticae (Heldr! Orph. Fl. exs 396!), Citheron Bæotiae (Boiss!), Onion et Olympus Thessalus (Heldr!), Parnassus (Guicci!), montes Korthiati et Athos Macedoniae (Griseb., Orph! Janka! Pichler!), Byzantii (Counany!).

ζ . *Kotschyannus*. — Fruticulosus rami lignosis saeppe crassis saepius erectis, folia nervosa rotundata vel cuneato-ovata vel elliptica aut lanceolata subtus elevatim nervosa, flores in capitula interdum basi interrupta congesti, calycis dentes omnes saepius ciliati et superiores acuti vel acuminati, corolla saepius alba. — *T. Kotschyannus* Boiss. et Hoh. Diagn. Ser. I, 5, p. 16 et quoad formas stenophyllas *T. lanceolatus* Bth. in n.c. Prodr. quoad Persiam non Desf. — *T. xylorrhizus* et *T. Balansæ* Boiss. et Ky in Sched.

Hab. in siccis montium Berytdagh et Akkerdaghi Cataoniae (Haussk!), Alidagh Cappadociae (Bal!), Bingöldagh et Alischeri Armeniae (Ky 510! et 677!), montis Gara Assyriae (Ky. 327!), Kurdistanie Persicæ et Persie australis in montibus Avroman, Pir Omar Gudrum, Kellal, Kuh Daëna, Eschker, Kellal (Ky. 629! Haussk!).

A varietatibus β , ε et ζ differt ramis vetustis crassis saeppe fruticosis Diu eum pro specie propriâ habui sed formæ maiifeste intermediae adlunt.

η . *angustifolius*. — Rami saepius decumbentes glabri vel pubescentes, folia parva oblongo-linearia brevia glabra basi ciliata vel tomentella, verticillastra saepius in capitulum aggregata. — *T. angustifolius* Pers. Syn. II, 130. — *T. parvifolius* et *T. rariflorus* C. Koch Linn. XXI, p. 666. — *T. fallax* F. et M. Ann. Sc. Nat. 1834 p. 23. — *T. Borei* Bth. Lab. p. 342. — *T. Argæus* Boiss. et Bal. Diagn. Ser. II, 4 p. 7 (forma alpina microphylla velutina). — *Satureia Argæa* F. et M. Ann. Nat. 1834. p. 33. omnino *Thymus!* nec *Satureia* est, certe forma *T. Serpylli angustifolii*!

Hab. in monte Delphi Eubœa (Heldr!), ad Thessalonicam et in monte Korthiati (Orph. Fl. exs. 524!), in totâ Anatoliâ sept. (Wied!), Armeniâ (C. Koch!), Argæo Cappadociae in regione alpinâ (Bal!), Tauriâ et prov. Caucasicis (M.B., Rupr!), in deserto inter Suez et Gaza Palestinae (Bové!), ad Ouadi Askar deserti Egyptiaco-Arabici (Schweinf!).

Huc quoque spectare videtur *T. acicularis* W. K. Pl. rar. tab. 447 ex cl. Visiani ad *T. striatum* relatus, sed qui ob flores et capitula parva, calycis dentes superiores glabros ab eo differt.

ϑ . *squarrosum* — Rami breves saepius velutini, folia minuta rigida nervosa elliptica vel ovata patentia saeppe subrecurva et canaliculata, floralia subconformia, capitula ovata vel oblonga, corolla alba. — *T. Sipyrensis* Boiss. Diagn. Ser. I, 3 p. 46. — *T. punctatus* Visian.

III. p. 7 tab. 2. — *T. squarrosum* F. et M. Ann. Sc. Nat. 1854 p. 32. — *T. rigidus* Schott. et Ky.

Hab. in siecis circa Aneyram, Hamamli, Karahissar (Wied!), in monte Sipylo Lydie (Boiss!), in sylvaticis circa Ouchak (Bal!), ad Elmali Lycæ (Bourg. 223!), Cappadociæ (Tchib! Ky. Suppl. 195!), Armeniæ australi ad Gungun (Ky. Suppl. 670!), Tauro Cilicie in alpinis (Ky. 71! Bal!). Amano (Ky!).

Haec varietas, præsertim forma ex monte Sipylo, primâ fronte a *T. Serpylo* diversissima videtur sed intermediis in eum sensim transit.

Ar. Geogr. Europa tota, Sibiria et Dalmatia, Regio Himalaica, Abyssinia, Groenlandia.

3. **T. striatus** (Vahl Symb. p. 78!) rhizomate suffruticoso cespitoso, ramis lignosis procumbentibus, ramulis floriferis erectis strictis brevibus crenatis foliosis retrosum hirtulis, foliis rigidis sessilibus linearibus subspathulatis glabris plus minus ciliatis subtus nervo valido nervulisque prominentibus instructis, floribus in capitula ovata densa foliis summis involuerata congestis, bracteis oblongis nervosis apice saepe attenuatis, pedicellis brevissimis, calycis ad medium bilabiati labio superiore sublongiore, dentibus omnibus lanceolatis ciliatis labii inferioris angustioribus profundius fissis, corolle albæ tubo subincluso 5. *T. Zygis* Fl. Græc. VI, p. 60 tab. 374 et Linn. herb. ex parte non L. Spec. — *T. zygoides* Griseb. Spic. II, p. 418!

Hab. in siecis regionis inferioris et montanæ, Cephaloniæ mons Enos (Heldr!), Attica in Hymetto (Sprun!), Thracia (Griseb!), Anatolia (Auec. 1626! et 1627!), Lydia ad colles Smyrnæ (Boiss! Bal. 304!) et in monte Tmolo (Bal. 302!), Lycæa (Bourg. 298!). Lycæonia (Heldr!).

Planta *T. Serpylo* rigidior, rami densius tolosi, flores majores et calycis dentes superiores lanceolati nec triangulares; haec est certe cl. Gussone et Visiani observantibus *T. Zygis* herbarii Linneani ex parte, sed descriptio Sp. pl. 826 et synonymon Löllingii citatum ad *T. tenuifolium* Mill. referenda sunt An hic quoque a *T. Serpylo* satis differt?

Ar. Geogr. Italia australis, Sicilia.

4 **T. hirsutus** (M.B. Taur. Gaue. II, p. 39) cespitoso-pulvinatus, caudiculis ex rhizomate indurato procumbentibus, ramis nivellis filiformibus tenuibus dense foliatis axillis fasciculiferis, foliis impiunctatis sessilibus linearis-subulatis obtusis margine subrevolutis ciliatis, floribus in capitula oblonga foliis floralibus sublatioribus intermixta approximatis, bracteis subulatis, calycis rubelli hirsuti ultra medium bilabiati dentibus labii superioris lanceolatis acuminatis inferioris subulatis, corolle roseæ tubo inclusa 2. *T. vulgaris* Sm. Prôdr. I, p. 419 ex cl. Bth. non L. — *T. cherleroides* Visiani. III. p. 8.

Hab. in regione alpinâ Graeciae, mons Pateras Attice, Taygetus, Chelinos, Kyllene (Heldr!), Velinga Etolie (Samar!), Parnassus (Heldr!), Olympus Thessalus (Heldr! Orph!), mons Ida Anatolie (Parol., Schmidt!), mons Gheidagh Tauri Isaurici 6000' (Heldr!). Taurus Cilicieus 6-8° 00' (Ky! Bal!). Tauri montes (Stev!).

Facies *Crucianella capitate* vel *Galii Olympici*, folia tenuia 2-3 lineas longa-

in axillis fasciculata et strictissima, calycis dentes superiores sœpius acutiores et profundius fissi quam in *T. Serpylli* formis. In locis diversissimis faciem et characteres servat et species propria videtur.

5. ***T. odoratissimus*** (M.B. Taur. Cauc. III, p. 405) dumulosus, ramis junioribus tenuibus erectis foliosis, foliis glandulosopunctatis linearisubulatis margine subrevolutis obtusis ciliatis, axillis fasciculiferis, floribus capitatis foliis floralibus conformibus intermixtis, calycis fere ad medium æqualiter bilabiati dentibus superioribus lanceolatis acutiusculis inferioribus subulatis ciliatis, corollæ purpureæ tubo inclusa ♀. *T. pectinatus* F. et M. Ann. Sc. Nat. 1854 p. 32. — *T. Zygis* Marsch. Casp. p. 210.

Hab. in campestribus Ciscaucasicis (C. Koch), Cataoniâ (Tchih!).

Specimen Anatolicum Rossicis simile. An hæc planta quoque varietatibus *T. Serpylli*, cl. Ledebour proponente, adnumeranda sit nescio, facies propria, et hucusque formas transitorias non vidi.

Ar. Geogr. Rossia australis.

6. ***T. comptus*** (Friv. Flora 1836, p. 439) ramis tenuibus ascendentibus et decumbentibus, foliis anguste linearisubulatis margine convolutis obtusiusculis basi ciliatis, axillis fasciculiferis, floralibus brevibus sublatioribus, verticillastris distantibus laxe racemosis, calycis hirti dentibus ciliatis labii superioris triangularibus acutis inferioris subulatis, corollæ purpurascens tubo inclusa ♀. *T. glaucus* Friv. exs. (forma foliis sublatioribus).

Hab. in montibus Thraciæ (Friv!), in arenosis prope Kalofer (Pichler!), in Macedonia monte Korthiati (Friv! Orph!).

Folia convoluta anguste linearia *T. odoratissimi* et *hirsuti*, sed racemus longus verticillastris 8-12 distinctis constans.

7. ***T. Cappadocicus*** (Boiss. Diagn. Ser. I, 5 p. 17) basi suffructicosus prostratus ramis floriferis erectis brevissimis dense dumosis crebre et stricte foliosis, foliis linearisetaceis margine convolutis obtusiusculis ciliatis, floribus paucis in capitulum foliatum parvum foliis suffultum congestis, foliis floralibus vix latioribus flores superantibus, floribus minimis, calycis glabri labio superiore ad medium in dentes triangulares fisso, inferioris æquiloni dentibus binis subulatis ciliatis, corollâ albâ calyce vix longiore ♀.

Hab. in Cappadociâ ad Euphratem (Auch. 1624!).

♂. *pruinosus*. — Indumentum pruinosum canescens, folia floralia subbreviora. — *T. pruinosus* Boiss. Ann. Sc. Nat. 1854 p. 253.

Hab. in planicie lapidosa Cappadociæ Orientalis inter pagos Gurum et Mendjalick (Tchih!).

Itami novelli bipollicares, folia 2-3 lineas longa lineam dimidiad vix lata, flores lineam longi. An intermediis cum *T. Serpyllo angustifolio* confluant haec et quatuor species præcedentes ulterius investigandum.

8. **T. decussatus** (Bth. Lab. p. 482) suffruticosus dumulosus pumilus ramis tenuibus albo-tomentellis erectis rigidis opposite et stricte ramulosis vetustis spinescentibus, foliis parvis tomentellis anguste linearibus margine revolutis obtusis strictis ramulorum sterilium dense decussatis, floralibus conformibus, floribus paucis dense capitellatis, pedicellis strictis calyce brevioribus, calycis glanduloso-punctati et hispiduli vix ad medium bilabiati labio superiore brevissime et obtusiuscule tridentato, inferiore æquilongo in dentes subulatos ciliatos fisco. corollæ albidae tubo breviter exerto 5. *T. quadrangulus* Ehr. MSS!

Hab. in rupestribus Arabiae petreæ in monte Sinai (Ehr! Bové!) et prope Bestan (Schimp. 378!). Arabice *Sætheran*.

Dumus semipedalis, folia 2-3 lineas longa, corolla vix tres lineas longa. Ab omnibus *T. Serpylli* et *striati* formis ramis opposite ramulosis et tandem subspinescentibus diversus.

9. **T. Syriacus** (Boiss. Diagn. Ser. I, 12 p. 47) suffruticosus dumulosus a parte inferiore ramosissimus ramis numerosis erectis simplicibus virgatis remotiuscule foliosis albidis sub lente hirtis, foliis sessilibus rigidis erecto-patulis linear-lanceolatis acutiusculis crebre punctatis glabris basi parcissime ciliatulis, capitulo terminali ovato denso, bracteis ovato-oblongis elevatum 5-7-nerviis hirtelle calyces subæquantibus, calyce breviter hirtello punctato labiis subæquilongis superiore lato oblongo apice minutissime tridenticulato inferiore in dentes setaceos ciliatos bipartito, corollæ cernuae tubo breviter exerto 5. *Th. lanceolatus* var. *angustifolius* Bth. in D.C. Prodr. XII, p. 203.

Hab. in Cælesyriâ ad radices montis Makmel prope Balbeck (Boiss!), ad Ouadi el Karn et Kalaa Djendel (Gaill!), prope Hasbeya (Ky. 250!), in monte Amano prope Beilan (Ky. 59!) forma bracteis majoribus calycis dentibus longioribus inter Homs et Tripoli (Post!), circa Aleppo (Boiss!), inter Aintab et Arablar et in arenosis prope Erbil et Sulimanieh (Haussk!), inter Baïard et Semel (Ky. 428!).

Rami novelli 6-10-pollicares, folia 5-8 lineas longa lineam lata, flores 4 lineas longi. Foliis augustis acutis, calycis formâ a *T. lanceolatus* Desf. Africano omnino alienus. Species calycis labio superiore lato tridenticulato, foliis firmis acutis insignis, cum sequenti ob habitum et bracteas latas transiunt ad *Pseudothymbram* præbens.

10. **T. Cilicicus** (Boiss. et Bal. Diagn. Ser. II, 4 p. 8) suffruticosus a parte inferiori ramosissimus ramis erectis brevibus crebre foliosis hirtulis, foliis sessilibus strictis subulato-subtriquetris acutis pruinosis et basi ciliatis, capitulo terminali denso ovato, bracteis villosulis ovatis abruptiuscule acuminatis calyces subsuperantibus, calycis hirti rubelli labio superiore ad quartam partem acute tridentato inferiore in dentes subulatos bipartito dentibus omnibus ciliatis, corollæ purpureæ tubo inclusa 5.

Hab. in collibus calcareis Ciliciae ad septentrionem urbis Tarsus (Bal!); prope Anamour (Peron!), montibus Kassan Oghlu Ciliciae orientalis (Ky!).

Vix semipedalis, facies *T. capitati* a quo calyce, etc., alienus. Affinis *T. Syriaco* a quo differt foliis crebris 4-5 lineas longis multo angustioribus, capitulis minoribus avellanam subæquantibus, bracteis abrupte acuminatis.

SECT. II. PSEUDOTHYMBRA Bth. — Calyx teres, corollæ tubus longe exsertus

11. **T. Billardieri** (Boiss. Diagn. Ser. II. 4, p. 8) suffruticosus cespitosus, ramis procumbentibus radicantibus, ramulis floralibus brevibus erectis crispuis hirtis dense foliosis, foliis linear-ellipticis margine subconvolutis albo-ciliatis obtusiusculis, axillis fasciculiferis, capitulo paucifloro ovato-globoso, foliis floralibus oblongis vel lanceolatis ciliatis apice repando-denticulatis calyces subæquantibus, calycis rubri labio superiore lato breviter tridentato inferiore in dentes subulatos partito dentibus omnibus ciliatis, corollæ roseæ tubo ealyce 3-4-plo longiore 5. *T. villosus* Fl. Græc. VI, p. 62, tab. 578 non Linn. — *T. integer* Griseb. Spicil. II, p. 116 (nomen antiquius sed minus aptum nam bracteæ dentatae).

Hab. in insulis Archipelagi (ex Sibth.), ego eum tantum ex insula Cypro vidi unde (nec e Libano) hunc olim Billardierius retulit, crescit in regione montana ad Prodromo, S^a Croce (Ky. 832 et Suppl. 397!).

Palmaris, folia 3-4 lineas longa lineam lata, calyx tres lineas, corolla 11-12 longa. *T. villosus* L. est planta Lusitanica bracteis profunde lobatis, etc., diversissima. Vide Boiss. Voy. Bot. Esp.

SECT. III. CORIDOTHYMUS Rehb. — Calyx a dorso compressus anceps. — Calyx *Thymbrae* sed stamina *Thymi*.

12. **T. capitatus** (L. Sp. 793 sub *Satureia*) suffruticosus dumulosus pumilus ramis crassis erectis ramulosis rigidis canescensibus, ramulis sterilibus demum spinescentibus, foliis rigidis patentibus linearibus brevissimis subtriquetris acutiusculis punctatis basi ciliatis, axillis breviter fasciculiferis, capitulo ovato denso strobili-formi, bracteis ovatis et oblongis cucullatis ciliatis dense imbricatis calyces obtentibus, calycis compressi ad margines et dentes ciliati labio superiore dentibus brevissimis conniventibus, inferiore sublongiore in dentes subulatos bipartito, corollæ roseæ tubo breviter exerto, antheris exsertis 5. Ic. Fl. Gr. tab. 544. — *T. capitatus* Link et Hoffm. — *Thymus Creticus* Brot. Phyt. p. 27, tab. 12. — *Thymbra capitata* Griseb. Spic. II, p. 127. — *Coridothymus cupifatus* Rehb. Ic. 18 p. 40.

Hab. in collibus regionis inferioris totius Græciae et ejus insularum!, Macedoniae (Friv!), Lydiae (Bal. 300!), Cariae! Cypri (Ky.), Syriae littoralis (Ehr! Gaill!) et interioris in ditione Golani (Wettst!), Palestinae (Sieb!), Egypti ad Alexandriam (Samar!), Græcis Thumari, Arabis Zatar Farisi audit.

Ar. Geogr. Regio mediterranea totius Europæ et Africæ borealis.

THYMBRA (L. Gen. 708).

Calyx oblongus 13-nervis a dorso plano compressus subanceps bilabiatus labio superiore tridentato inferiore bipartito. Corolla tubus exsertus intus nudus, limbi labium posticum erectum retusum subplanum, anticum trifidum. Stamina 4 sub labio superiore ascendentia didynama, antherae per paria approximate biloculares loculis parallelis. Stylus bifidus. — Nuculae ovatae leves. — Suffrutex facie *Thymi*.

1. **T. spicata** (L. Sp. 795) suffruticosa dumulosa, ramis novellis numerosis erectis rigidis simplicibus bifariam puberulis, foliis punctatis linearibus rigidis obtusis uninerviis planis vel complicatis ciliatis vel glabris, ramorum sterilium confertis, axillis saepe fasciculigeris, spicis densis oblongo-cylindricis basi rarius subinterruptis, foliis floralibus bracteisque numerosis imbricatis anguste lanceolatis rubellis albo-ciliatis calyces obtegentibus, calycis compressi glabri ad angulos ciliati labio superiore dentibus breviter triangularibus, inferioris aequilangi dentibus linearis-subulatis omnibus ciliatis, corollae purpureae tubo calyce subduplo longiore 5. Ic. Fl. Graec. tab. 546. — *T. verticillata* L. loc. cit. forma monstrosa. — *T. ambigua* Clarke.

Hab. in collibus regionis inferioris Graeciae in Achaiâ (Sibth.), Atticâ (Spr!), Bœotia (Orph. Fl. exs. 529!), Creta (Sibth.), Archipelagi (Urv.), Thracia (Griseb.), Bithynia (Pichler!), Caria (Boiss!), Lydia (Boiss!), Cilicia (Heldr!), Balt., Syriae littoralis (Ehrenb! Boiss!) et interioris in monte Ssotdagh (Haussk!), Palestinae (Sieb! Auch!), Kurdistaniae Assyriacae in monte Gara (Ky. 3 6!). Anatoliae borealis ad Tokat et Sojut (Wied!).

ZATARIA (Boiss. Diagn. Ser. I, 5, p. 48).

Calyx ovatus membranaceus quinquenervis regularis quinquedentatus ore barbatus. Corolla tubo subincluso, limbo bilabiato labio superiore subbreviore breviter bilobo inferiore aequaliter trilobo. Stamina 4 didynama brevia distantia inclusa, antherarum loculi distincti paralleli demum divergentes. Stylus breviter bifidus. Nuculae ovatae leves. — Fruticulus facie *Thymi*, floribus minutissimis verticillastra densa distincta laxe spicata formantibus. — Genus ab affinis calyce quinque nervi nec 10-13-nervi distinctum.

1. **Z. multiflora** (Boiss. loc. cit.) fruticulosa ramosissima, ramis rigidulis tenuibus puberulis albidis in spicam interraptam vel paniculam racemosam aheunibus, foliis minimis brevissime petiolatis orbiculatis myrtiformibus crebre punctatis glabriusculis, cymulis globosis densis vel omnibus ad verticillastra sessilia reductis vel

inferioribus pedunculatis, bracteis oblongis calycem aequantibus vel eo brevioribus, calyce membranaceo pentagono ad angulos ciliato dentibus triangularibus brevibus, corollâ albida calyce sublongiore 5.

Hab. in Persia australi (Auch. 5192!), prope Kerman et Ispahan (Bunge!), inter Djendack et Yezd (Buhse!), in Belutschia (Stocks!).

Fruticulus intricatus palmaris, folia 3 lineas diam. lata. Flores vix lineam unam longi.

β. elatior. — Pedalis, folia submajora 3-6 lin. diam. lata, spiculæ longiores inferiores pedunculatae et sic paniculatae, bracteæ calyce sublongiores. — *Z. bracteata* Boiss. Diagn. Ser. II, 4 p. 12.

Hab. in regno Cabulico prope Otipore (Griffith!).

Cl. Bentham speciminibus numerosis collatis hanc formam intermediis ad typum transire observavit.

SATUREIA (L. Gen. 707).

Calyx campanulatus vel obscure 10-13-nervius quinquedentatus aequalis vel obscure bilabiatus dentibus tunc subimæqualibus. Corollæ tubus inclusus vel exsertus intns nudus. limbi labium superius erectum planum integrum vel retusum anticum trifidum. Stamina 4 distantia sub labio postico conniventia vel exserta subdivaricata, antheræ bilocularis loculis parallelis divergentibus. — Herbæ rarius annuae vel suffrutices. Genus a *Thymo* calyce regulari vel subregulari et staminibus subascendentibus diversum, *Micromeria* magis affine et ab ea potius habitu, foliis carnosulis exsiccatis fragilibus (Conf. Bge Lab. Pers. pag. 37) quam calycis structurâ distinctum.

§ 1. Annuæ. — Calyx regularis.

1. *S. hortensis* (L. Sp. 795) annua, pube curvula seabraida basi ramosissima ramis filiformibus erectis vel patulis remote foliosis, foliis anguste et longe oblongo-linearibus linearibusque acutiusculis basi in petiolum brevem attenuatis, florum minimorum rarius solitarium glomerulis axillaribus 3-5-floris remotis longe spicatis inferioribus breviter pedunculatis superioribus sessilibus, bracteis setaceis brevibus, calycis seabridi dentibus anguste lanceolatis æquibus rigidulis tubo sublongioribus, corollæ roseæ limbo parvo calycem vix excedente ⊖. Ic. Rehb. tab. 71. — *S. pachyphylla* C. Koch Linn. XVII, p. 295! — *S. filicaulis* Schott Hort. Vindob.

Hab. in sterilibus Anatoliae borealis ad Samsun et Tokat (Wied!), Armeniæ (C. Koch!). Cappadociæ et Cataoniae (Haussk!), Cilicie ad Tauri radices (Ky! Bal!), Tauriae (M.B.), Caucasi usque ad 4800' in Daghestaniæ et Transcaucasia (Rupr! Szov!), Persiae bor.-occid. (Seidl!).

β. grandiflora. — Variatio corollæ majoris tubo breviter exerto. — *S. laxiflora* C. Koch Linn. XXI, p. 668.

Hab. in Grasiā (Wilh!), prope Titlis (Rupr!), circa Erzerum Armeniae (Calv!).

Ar. Geogr. Hispania, Gallia australis, Italia borealis, Dalmatia, Sibiria.

§ 2. **Eusatureiæ.** — Suffrutescentes. Calyx regularis.

2. **S. montana** (L. Sp. 794) suffruticosa glabra vel seabrida ramis virgatis pubescens foliosis in racemos spiciformes laxiusculos foliosos abeuntibus, foliis coriaceis lucidis linear-lanceolatis acutis basi longe attenuatis sessilibus ciliatis, floralibus flores superantibus, cymis 3-5-floris breviter pedunculatis vel subsessilibus, bracteis linearibus, calycis subæqualis dentibus lanceolatis acutis tubo subæquilongis, corollæ roseæ tubo subineluso 5. Ic. Rehb. tab. 72. 1.

Hab. in Tauriā et Ossetiā Caucasicā (M.B.), in ditione Gumbet Caucasi orient. (Ower.).

β. stenophylla. — Folia angustiora linearia basi brevissime attenuata. — Ic. Fl. Graeca tab. 543. — *S. montana* var. *fasciculata* Friv.

Hab. in Rumeлиā (Friv!), in declivitate montis Balkan Thracie sed boreali (Janka exs sub *S. pygmæa*!), in monte Athone Macedoniae (Sibth.).

Specimina e Balkan ramis abbreviatis subsimplicibus *S. pygmæam* Sieb. referunt (quæ cæterum ex multorum opinione *S. montana* var. est) sed ab eis ramulis pubescens nec glabris differunt.

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, regio Danubialis Dalmatia, Rossia australis.

3. **S. Parnassica** (Heldr. et Sert.) rhizomate crasso lignoso ramos breves puberulos cespitosos procumbentes novellos brevissimos foliosos edente, foliis parvulis obovatis vel spathulato-linearibus acutis basi attenuatis hirtis subtus carinatis, floribus in axillis supremis subsessilibus binis aut solitariis, bracteis elliptico-linearibus acuminatis, calyeis hirti dentibus æqualibus lanceolatis tubo subæquilongis, corollæ roseæ tubo inclusō 5. *S. approximata* Friv. Fl. 1836, p. 438.

Hab. in rupestribus Saraka dictis regionis summae Taygeti (Heldr!), Par-nasso in regione abietinæ et superiore (Guice!), in cacumine Delphi Eubœe 5000' (Heldr!), Athone (Friv!), regione superiori montis Sipyli Lydie supra Magnesiam (Bal. exs. 309 sub *S. spinosa*!).

Facies *S. spinosa* pro cuius varietate eam diū habui sed ramuli vetust multo tenuiores tandem subpersistentes nec valide spinosi, folia hirta. Ulterius observandum an ad *S. montanam* aut forsū potius ad *S. cuneifoliam* non transeat.

4. **S. spinosa** (L. Sp. 795!) rhizomate crasso lignoso ramos puberulos rigidos brevissimos crassos densissime pulvinari-fastigiatos opposite ramulosos apice spinescentes edente, foliis minimis obovato-lanceolatis acutis basi attenuatis carinato-plicatis, floribus paucis axillaribus solitariis subsessilibus, calycis sebriuscili dentibus

æquilongis lanceolatis carinatis tubo subbrevioribus, corollæ albidæ tubo inclusa 5. Ic. Fl. Græc. tab. 545.

Hab. in fissuris rupium regionis subalpinæ et alpinæ Cretæ 4000'-6000' in montibus Sphakia, Ida, Lassiti (Tourn! Sibth., Sieb! Raul! Heldr!).

Dumulus erinaceus palmaris spinis brevibus validis albis, folia vix 2 lineas longa.

5. *S. cuneifolia* (Ten. Fl. Nap. 5, p. 3, tab. 155) suffruticosa undique vel superne tautum seabra vel hispidula, ramis erectis tenuiter virgatis rigidis subsimpleibus in spicas longas tenues interruptas abeuntibus, foliis inferioribus obovatis vel oblongo-spathulatis basi longe attenuatis superioribus linearibus omnibus acutis plus minusve canaliculatis et subrecurvis floralibus saepius abbreviatis, verticillastris numerosis plurifloris breviter pedunculatis et sessilibus distinctis et saepius remotis, bracteis lanceolatis minutis, calycis seabrido-hirsuti dentibus æqualibus lanceolatis acutis tubo subæquilongis, corollæ roseæ tubo calyce non longiore 5. Rehb. Ic. tab. 71. — *S. virgata* Visian. Fl. Dalm. tab. 18.

Hab. in siccis montanis, Cephalonia in monte Enos 2400' (Wiest! Heldr!) forma diffusa, Cilicia Trachea (Per!), Pontus ad Trapezuntem (Auch. 1623!), mons Gara Kürdistaniæ (Ky. 259!), Libani regio subalpina ad Eden! Bescherre, Dimam (Bl! Gailli! Ky. 259!).

S. Wiedemanniana. — Lallem. Suppl. ad Ind. Petrop. p. 64. — Omnia formæ typicæ præter folia floralia non abbreviata sed ut in *S. montana* flores superantia, verticillastra magis approximata.

Hab. in monte Budrun Pisidiae (Heldr!), ad Amasia Paphlagoniæ (Heldr!), in monte Kechchdag (Wied!).

γ. tenuis. — Folia etiam inferiora anguste lanceolato-linearia, cymæ laxiores longius pedunculatae.

Hab. in Tauro Cilicio inter Gulek Boghas et Gulek Maaden (Auch. 1632! Ky. 477! Bal!).

S. oborata Lag. Hispanica differt indumento glanduloso, foliis obtusis, corollæ labio superiore elongato.

Ar. Geogr. Italia australis, Dalmatia.

6. *S. intermedia* (C. A. Mey. Enum. p. 91) undique patule pubescenti-canescens basi suffrutescens ramis tenuibus diffusis, foliis obovatis obtusis in petiolum attenuatis, floralibus oblongo-spathulatis flores paulo superantibus, cymis breviter pedunculatis 3-5-floris laxe racemosis, bracteis linearibus curvatis calyce brevioribus, calycis hirti dentibus subulatis subrecurvis acutis tubo æquilongis, corollæ tubo inclusa 5.

Hab. in ditione Talysh prope Swant 4000' (C.A. Mey!), prope Orant (Seidl!).

Rami novelli diffusi flexuosi semipedales, folia inferiora 3-4 lineas longa. Ab affini *S. cuneifolia* differt foliis magis rotundatis muticis, calycis dentibus subulatis recurvis.

§ 3. **Subbilabiatae**. — Suffrutescentes vel perennies, calyx subbilabiatus.

7. **S. Boissieri** (Haussk. exs.) basi suffrutescens pluriceaulis caulis ascendent-erectis simplicibus velutinis crebre foliosis, foliis punctatissimis parce scabrido-hirtis elliptico-linearibus basi attenuatis obtusis subplicatis, floralibus calyces subaequantibus, verticillastris paucifloris brevissime pedunculatis et sessilibus in spicam terminalem densiusculam approximatis, calycis patule hirti subbilabiati dentibus superioribus triangularibus tubo triplo brevioribus, inferioribus lanceolatis profundius fisis omnibus obtusiusculis, corollæ calyce 2-plo longioris tubo subexerto 5.

Hab. in monte Akdaghi Cataoniae inter Malatia et Adiaman (Haussk!).

Caules cum spicâ 2-3-pollicari vix pedales, folia inferiora 9-10 linea longa 2 lata, calyx 2 lineas, corolla 4 lata. A *S. cuneifoliae* formis calycis subbilabiati dentibus brevibus etc. distincta.

8. **S. subdentata**, ex rhizomate lignoso crasso ramosissima ramis ascendentibus tenuibus simplicibus vel paree ramosis crebre foliosis, foliis viridibus lucidis glabris oblongo-et linearispathulatis basi longe attenuatis carinato-plicatis obtusiusculis parte superiori repando-denticulatis, axillis fasciculiferis, floralibus coniformibus, cymis axillaribus subtrifloris tenuiter et stricte pedunculatis, bracteis linearispathulatis canalicularis acuminatis calyces aequantibus, calycis basi hirtuli subbilabiati dentibus superioribus triangularis-lanceolatis brevibus inferioribus profundius fisis lanceolatis calyce tubo brevioribus omnibus obtusiusculis, corollæ calyce 2 $\frac{1}{2}$ -plo longioris tubo subincluso 5.

Hab. ad Arzuch supra fluvium Awar Koissa Daghestaniae superioris 3500'-4000' (Rupr!).

Rami novelli 6-8-pollicares, folia majora 5 linea longa 2 lata, flores pauci, cymæ laxæ foliosæ, corolla 3 $\frac{1}{2}$ lin. longa. Affinis *S. intermedia* differt foliis angustis, cymis paucis paucifloris foliosis, calyce.

9. **S. mutica** (Fisch. et Mey. Ind. Petrop. 1833, p. 49) caulis præter basin induratum herbaceis tenuiter virgatis erectis a parte inferiori longe et anguste paniculato-ramosis breviter hirtis, foliis glabris pallidis punctatis anguste oblongo-linearibus linearibusque basi attenuatis muticis, floralibus inferioribus cymas aequantibus, cymis breviter pedunculatis 3-7-floris laxiuseule racemosis, bracteis linearibus brevissimis, calycis subbilabiati pilis crispatulis hirti dentibus linearis-lanceolatis obtusis tubo subbrevioribus inferioribus sublongioribus, corollæ tubo subincluso, antheris subinclusis 2. *S. intermedia* Bth. in D.C. Prodr. ex parte non C. A. M.

Hab. in siccis lapidosis ditionis Talyesch (C.A. Mey! Hoh. exs. sub *S. intermedia*! Buhse!), Tataris condimento inservit et Marsa audit. Fl. Jul., Aug.

Pedalis et bipedalis, folia inferiora fere pollicem longa 2 lineas lata, corollæ tres lineas longa. Sæpe commutata cum *S. intermediâ* a quâ longe distat caulis herbaceis rectis, foliis glabris longe linearibus, etc.

10. *S. spicigera* (C. Koch Linn. XVII, p. 295 sub *Micromeridâ*)
a basi ramosa ramis herbaceis diffusis bifariam vel undique puberulis
ramulosis creberrime foliosis, foliis intense viridibus anguste oblanceolato-linearibus vel lineari-setaceis acutiusculis muticis subflexuosis
sæpius patentí-reflexis, floralibus flores subæquantibus. cymis
subtrifloris subsessilibus in spicas unilaterales confertis, floribus
pedicellatis, calycis glabriuscui subbilabiati dentibus lanceolatis ob-
tusiusculis inferioribus paulo angustioribus et longioribus, corollæ
albæ tubo subincluso limbo ampio, staminibus exsertis 2. *S. diffusa*
Bth. olim in D.C. herb. — *S. intermedia* β laxior Bth. in D.C. Prodr. —
S. alternipilosa C. Koch Linn. XIX, p. 25 et *M. alternipilosa* C.
Koch Linn. XXI, p. 668.

Hab. in regione inferiori Ponti (Thirke!), circa Trapezuntum (Auch. 1822!,
Calv! Bal!), Colchidis et Imeretiae (Rupr!), ditionis Radscha Caucasi (C. Koch!
Rupr!). V. sicc. et viv. cult. Fl. Oct., Nov.

Facies *Odontitis luteæ* sed diffusa, siccitate nigrescit. Affinis *S. muticæ* a quâ
differt ramis diffusis, foliis creberrimis angustis acutioribus flexuoso-reflexis,
spicis densioribus secundis, corollæ majore, staminibus exsertis.

11. *S. macrantha* (C. A. Mey. Suppl. Ind. XI, Petrop. p. 67)
basi lignosa crebre multicaulis seabraida canescens, caulis virgatis
ſtenuibus elongatis erectis simplicibus vel parce paniculato-ramosis,
a ſoliis crassiusculis lineari-spathulatis brevibus obtusis subpatentibus,
ſxillis fasciculiferis, floralibus brevissimis, verticillastris paucifloris
ubsessilibus remotiusculis longe laxe et tenuiter spicatis, calycis
parce hirti tubuloso-campanulati subbilabiati dentibus a basi trian-
gulari subulatis muticis tubo triplo brevioribus inferioribus longio-
ribus, corollæ tubo calyce 2 1/2-plo longiore 2.

Hab. in siccis prov. Transcaucasicæ Karabagh (Szov! Hoh.), prope Ssautsch
bulak ad lacum Urmiah (Seidl!).

Rami 1-1 1/2-pedales tenues sed illis *S. muticæ* duriores, folia 3-4 lineas longa
sat approximata, calyx 2 lineas, corolla 5-6 longa. Species corollæ longitudine
notabilis.

12. *S. longiflora* (Boiss. et Haussk) basi lignosa multicaulis
punctatissima minute puberula canescens, caulis tenuiter virgatis
elongatis densiusculis erectis flexuosis simplicibus vel parce panicu-
lato-ramosis, foliis obovato-oblongis et spathulato-linearibus sessilibus
obtusis, floralibus ovatis vel oblongis brevissimis, verticillastris 4-6-
floris breviter pedunculatis vel sessilibus approximatis longe et te-
nuiter spicatis, calycis glabri sub lente pruinosi obsolete bilabiati
dentibus triangularibus obtusis tubo quadruplo brevioribus inferiori-
bus lanceolatis sublongioribus, corollæ violaceæ tubo calyce duplo
longiore 2.

Hab. in fissuris rupium calcarearum montis Parrow prope Kermanchah Persiae occidentalis 4-8000' (Haussk!).

Caulis cum spicis sesquipedalis; valde affinis *S. macrantha* a qua distincta videtur foliis latioribus, verticillastris approximatis, dentibus abbreviatis non subulatis.

13. S. Atropatana (Bge Lab. Pers. p. 36) basi lignosa multi-caulis minutissime papilloso-asperula, caulis tenuiter virgatis duriusculis erectis subsimplicibus, foliis parvis remotinsculis obovato-oblongis superioribus linearibus canaliculatis omnibus obtusis floribus brevissimis, verticillastris plurifloris breviter pedunculatis vel sessilibus remotinsculis tenuiter et laxe spicatis, calycis sparsim papilloso-hirti dentibus superioribus brevissime deltoideo-triangularibus, inferioribus sublongioribus lanceolatis obtusis tubo quadruplo brevioribus, corollæ tubo exerto ♀.

Hab. in rupestribus montanis Mutschetdagh prope Nehmedabad prov. Aderbidjan Persiae (Szov!).

Affinis *S. longiflora*, indumentam papillare, cymæ distantes, calycis papillaris dentes brevissimi. An hujus varietas?

§ 4. Zatarioideæ. — Suffruticosæ, calyx ob nervos suturales vix conspicuos subquinquenervius.

14. S. Bachtiarica (Bge Lab. Pers. p. 37) basi lignosa multi-caulis minute puberula canescens punctata, ramis erectis tenuiter virgatis simplicibus vel parce ramulosis, foliis crebris minutis oblongo-spathulatis canaliculato-plicatis recurvis obtusis, axillis fasciculigeris, foliis floralibus minutissimis oblongis, verticillastris plurifloris subsessilibus distinctis parvis distantibus longe et tenuissime spicatis, floribus minimis, calycis breviter campanulati hirsuti dentibus trianguli-lanceolatis tubo aequilongis, corollæ albidae vel pallide rubellæ tubo subincluso ♂.

Hab. ad rupes calcareas prope Abergun ditionis Bachtiaricae Persiae austro-occidentalis (Haussk!), circa Kirrind (Noë 10381). Fl. Sept.

Rami cum spicis pedales et sesquipedales, folia 2-3 lineas longa, calyx lineam, corolla $2\frac{1}{2}$ lineas longi. Ab omnibus calyce ob nervos secundarios parum conspicuos subquinquenervi distincta, hac notâ *Zatarice* accedens sed facie et characteribus *Satureia* vera.

§ 5. Tragorigana — Suffruticosæ, calyx regularis bracteis longis suffultus.

15. S. Thymbra (L. Sp. p. 794) suffruticosa seabrido-canescens ramosissima ramis lignosæ tortuosis junioribus brevibus erectis, foliis obovato-cuneatis plicatis acutissimis basi longiuscule attenuatis subsessilibus, axillis breviter ramulosis, verticillastris multifloris sessilibus globosis valde distantibus racemosis vel in paniculam pa-

rum ramosam dispositis, foliis floralibus minntis deflexis bracteisque numerosis oblongis acuminatis calyces æquantibus, calycis patule albo-hirsuti dentibus subæqualibus a basi lanceolatâ subulatis tubo subbrevioribus, corollæ roseæ tubo subincluso 5. Ic. Fl. Græc. tab. 541. — *Thymus Tragoriganum* L. Mant. 84. — *Thymbra hirsuta* Pers. Syn. II, p. 114.

Hab. in calidis siecis regionis inferioris Græcæ et Cycladum (Sibth., Boiss! Heldr!), Zæcynthi (Margt!), Cretæ (Sieb!), Rhodi!, Anatolice occid. in Lydia (Anch 1631! Boiss!), Pamphyliæ (Heldr!, Bourgt!), Ciliciâ (Bal!), Syriâ littorali (Bl!), Palestinâ usque ad Hebron (Boiss! Ky!), Cypro (Ky!).

Ar. Geogr. Sardinia.

MICROMERIA (Bth. Bot. Reg. 4282).

Calyx tubulosus vel obconico-campanulatus 13-nervius subæquilater quinquedentatus. Corollæ tubus rectus intus nudus exsertus vel inclusus, limbi labium superius erectum subplanum inferius tribolum. Stamina 4 didynama ascendenta apice arcuata, antheræ bilobulares connectivo vario. Nuculæ lœves. — Suffrutices vel herbæ. — Genus hinc ad *Satureiam*, illine ad *Calamintham* transiens.

Specierum Orientalium distributio.

SECT. I. CYMULARIA. — Annuæ, antheræ loculi divergentes connectivo punctiformi apice conjuncti.

M. cymuligera.

SECT. II. EUMICROMERIA. — Perennes, antheræ loculi ad extremitates connectivi elongati inserti.

* Nuculæ acutæ vel mucronatæ.

M. nervosa, Julianæ, cristata, cremnophila, elliptica

** Nuculæ obtusiusculæ.

M. Græca, hispida, Sinaica, microphylla, Persica.

SECT. III. PSEUDOMELISSA Bth. — Perennes. Antheræ loculi divergentes connectivo punctiformi apice conjuncti. Sectio ad *Eucalamintham* accedens sed calyx non bilabiatus.

M. nummularifolia, Libanotica, mollis, serpyllifolia, congesta, origanifolia.

SECT. I. CYMULARIA.

1. M. cymuligera (Boiss. et Haussk. exs. 1865) annua, collo ramosa ramis ascendentibus vel procumbentibus crassiuseculis simplicibus a basi crebre foliosis et floriferis retrorsum et crispuis hirtis. foliis nervulosis punctatis tenuibus asperulis elliptico-rhombeis basi in petiolum brevem longe attenuatis apice attenuato-acuminatis, floralibus conformibus cymas superantibus, cymis ad axillas breviter pedunculatis dense multifloris subglobosis, bracteis numerosis flores æquantiibus externis ovatis et oblongis longe acuminatis, internis subulatis, floribus minimis pedicellatis, calycis asperuli obconici dentibus tubo subæquilongis a basi lanceolatis subulatis ciliato-plumosis corollæ carneâ longioribus inferioribus profundius fisis, seminibus ovato-globosis ⊖.

Hab. in glareosis torrentium ad radices montis Berydaghi Cataoniae 3000' (Haussk!).

Rami 6-10-pollicares a basi crebre cymigeri, folia inferiora 3-5 lineas longa, cymæ densifloræ piso subminores, calyx sesquilineam longus exquisite 13-nervis, corolla lineam longa tubulosa, labio superiore integro inferius vix superante. Stamina longa sed e fauce non exserta, antherarum loculis non ut in *Eumicromeria* connectivo elongato sejunctis sed apice contiguis; superiorum antheras diminutas substeriles vidi, an semper?

SECT. II. EUMICROMERIA.

2. M. nervosa (Desf. Atl. 2, p. 9, tab. 121 sub *Satureia*) inferne breviter pubescens superne patule et molliter villosa. caulinibus ex rhizomate suffrutescenti virgatis ascendentibus elongatis simplicibus vel parce ramosis, foliis subsessilibus margine revolutis subtus nervulosis subcordato-ovatis acutiusculis, floralibus flores æquantiibus, cymis hispidissimis multifloris axillaribus breviter pedunculatis spicatis inferioribus distantibus, bracteis subulatis, calycis plumosi fauce hispiduli dentibus setaceis erecto-patentibus tubo subbreviribus, corollæ roseæ tubo inclusa, nuculis ovatis acutiusculis ♀. *M. nerrosa* Bth. Lab. p. 376. — *Thymus filiformis* Sieb. Cret. non L. — *Th. Mastichina* Sm. Prodri. non L. — *Satureia plumosa* Hampe Flora 1842, 1, 127.

Hab. in collibus siccis regionis inferioris Achaiæ (Bory!), Atticæ et Bœotiae (Sibth., Spr!, Boiss! Heldr!), Crete (Sieb!), Cycladum (Ürv.), Rhoëli (Bourg!), Zæynthi (Margt), Syriae littoralis ad Sidonem (Bl!), Cypri (Ky. 217!), Pæstina ad Samariam (Boiss!) et prope Hebron (Ky!).

Planta e Balearibus et ex Siciliâ eadem est, boreali-Africanam non vidi.

Ar Geogr. Baleares, Sicilia, regnum Neapolitanum, Africa borealis.

3. M. Julianæ (L. Sp. b93 sub *Satureia*) pubescens vel villosa, ex rhizomate duro ramosissima ramis erectis strictis simplicibus,

foliis sessilibus margine revolutis obtusis inferioribus ovato-oblongis superioribus vel omnibus linearibus, cymis axillaribus subsessilibus densifloris fastigiatis longe et laxiuscule spicatis, foliis floralibus flores æquantibus, bracteolis setaceis strictis calyces subæquantibus, calycis fauce nudi dentibus subulatis strictis conniventibus tubo subbrevioribus, corollæ tubo inclusio, nuculis oblongis mucronatis ♀. Ic. Fl. Græc. tab. 540. — *M. Julianæ* Bth. Lab. p. 378.

Hab. in rupestribus regionis inferioris et præsertim montanæ et subalpinæ totius Græciæ (Heldr! Orph. Fl. exs. 530!) usque ad 4500!, Zacynthi (Marg!), Cycladum (Urv.), Macedonia (Griseb!), Creta in prov. Sphakia 3500' (Heldr!), Rhodi (Bourg!), Lydiæ in monte Sipylo (Bal. 321!), Syriæ littoralis a collibus inferioribus ad Libanum medium (Bl! Gaill!). In montibus elatis forma ramis pumilis, Parnassus! Delphi Eubeæ, Libanus.

β. myrtifolia. — Folia omnia ovata vel oblonga, inferiora subpetiolata, floralia verticillastris sëpe sublongiora. Hæc forma ad typum intermediis transit. — *M. myrtifolia* Boiss. et Hoh. Diagn. Ser. I, 5, p. 19. — *M. Reinholdi* Heldr. MSS.

Hab. in Peloponnesi insulâ Poros (Heldr!), Lyciâ ad Elmali (Bourg!), Smyrnæ (Bal. 325!), Pamphyliâ (Heldr!), Cappadociâ australi (Bal!), Syriâ dor. prope Marasch (Haussk!), Syriâ littorali (Ehr! Bl! Gaill!). Kurdistaniâ Assyriacâ prope pagum Gara (Ky. 305!), montibus Avroman, Schahu, Pir Omar Gudrun 6-9000' (Haussk!).

Ar. Geogr. Lusitania (Welw!), Gallia australis, Italia australis et Sicilia, Dalmatia, Africa borealis.

4. *M. cristata* (Griseb. Spic. II, p. 122) breviter pubescens canescens, ramis ex rhizomate lignoso sëpe crasso creberrime cespitosis nanis filiformibus rectis basi nudis cæterum dense foliosis simplicibus, foliis minutis sessilibus strictis inferioribus oblongis cæteris lanceolatis omnibus obtusiusculis subtus concavis trinerviis nervis lateralibus marginantibus elevatis, ramorum steriliū densius imbricatis, cymis subtrifloris breviter et stricte pedunculatis folio axillari subsuperatis in spicam tenuem non interruptam approximatis, bracteis calyce multo brevioribus, calycis dentibus lanceolatis rectis tubo triplo brevioribus, fauce villosâ, corollæ purpureæ tubo inclusio, nuculis oblongis mucronatis ♀. *Thymus inodorus* Friv. non Desf. — *M. xylophila* Boiss. et Heldr. Diagn. Ser. I, 12, p. 49.

Hab. in fissuris rupium Macedonia australis (Friv!), montis Bulghasdagh Phrygiae (Bal!), montium Budrun et Anemas Pisidiæ 6000' (Heldr!), Anatoliâ boreali ad Safranbol (Wied!), Masmeneudagh Cappadocie (Bal!), Libanus prope Eden (Bl!).

Rami creberrimi 1-3-pollicares, folia 2-3 lineas longa, a *M. Julianæ* formis foliorum formâ, cymis pedicellatis, calycis fauce villosâ, a *M. Græcâ* nuculis distincta.

Ar. Geogr. Serbia.

5. *M. cremnophila* (Boiss. et Heldr.) sub lente minute pubescens e rhizomate indurato ramosissima, ramis pumilis simplicibus filiformibus ascendentibus, foliis parvis subsessilibus basi atte-

nuatis obtusis margine revolutis subtus nervulosis inferioribus obovatis floralibus linearibus flores subæquantibus, cymis 3-5-floris breviter pedunculatis distantibus laxe et tenuiter spicatis, calycis hirtelli fauce villosâ dentibus triangulari-lanceolatis rectis tubo quadruplo brevioribus, corollæ tubo inclusa, nuculis cylindrico-ellipticis acutis ♀.

Hab. ad rupes verticales regionis abietinæ Parnassi ad Gourna 4000'-4500' (Heldr.).

Rami cum spicâ 3-5-pollicares, folia majora tres lineas longa, superiora breviora et multo angustiora. A *M. ellipticâ* cui nuculis affinis calyce duplo breviore aliisque notis diversa, a *M. cristatâ* foliis subtus nervulosis nec trinerviis.

6. ***M. elliptica*** (C. Koch Linu. XXI, p. 669) viridis sib lente scabrida, ramis ex rhizomate lignoso numerosis filiformibus ascendentibus vel erectis simplicibus, foliis parvis subsessilibus margine revolutis subtus elevatim nervulosis obtusiuseculis ovato-oblongis, superioribus elliptico-linearibus flores superantibus, cymis 3-4-floris breviter et stricte pedunculatis breviter et laxiuscule spicatis, calycis fauce barbati dentibus lanceolatis obtusiuseculis rectis tubo quadruplo brevioribus, corollæ tubo breviter exerto, nuculis cylindrico-ellipticis acutis ♀. *M. Calverti* Boiss. MSS.

Hab. in Armenia Turcicâ prope Erzerum (Calv!), in ditione Pertakrek (C. Koch!).

Caules 3-4-pollicares, folia majora 4 lineas longa. Distinguitur a formis pumilis *M. Græcæ*, cymis paucis et paucifloris, calycis dentibus brevibus strictis, nuculis elongatis. Facies *M. cristatæ* foliis trinerviis non nervulosis, nuculis, etc., diversissimæ.

β. pubescens. — Patule pubescens. — *M. pubescens* Boiss. et Ky. exs. 1859.

Hab. in rupibus verticalibus vallis Teng prov. Musch Armeniæ 6500' (Ky. 458!).

7. ***M. Græca*** (L. Sp. 794 sub *Satureid*) pubescens vel superne presertim villosula, ex rhizomate suffrutescenti ramosissima ramis ascendentibus vel erectis saepius elongatis tenuiter virgatis subsimplicibus, feliis sessilibus firmis margine subrevolutis subtus elevatim nervulosis inferioribus ovatis vel ovato-oblongis superioribus lanceolatis vel linearibus, floralibus inferioribus cymas subsuperantibus, cymis axillaribus breviter pedunculatis 3-7-floris secundis, bracteis subulatis brevibus, calycis fauce villosi dentibus subulatis erecto-patentibus tubo subduplo brevioribus, corollæ roseæ tubo breviter exerto, nuculis oblongis non mucronatis ♀. Ic. Fl. Græc. tab. 542. — *M. Cypria* Ky. Cypr. p. 270!

Hab. in siccis Græciæ ut videtur rarius, in Peloponneso (Sibth.), Eubœa (Sprn!), Macedoniâ in Athone (Auch. 1644!), Cretâ et Cycladibus (Sibth., Urv.), Byzantium (Noë!), Anatoliâ (Auch. 1083!) et boreali ad Aniasia (Wied!), Ciliciâ (Balt!), Rhodo (Bourg!), Cypro (Ky. 338!), Ciliciâ littorali et Libani

regione mediâ (Bl! Gaill!), Palestinâ ad Gaza (Boiss!), Egypto ad Alexandriam (Ehr!).

Ar. Geogr. Europæ regio mediterranea a Lusitaniâ ad Dalmatiam, Africa borealis.

8. M. hispida (Boiss. et Heldr. Diagn. Ser. I, 12, p. 48) patule et longe hispida grisea basi suffrutescens, ramis novellis filiformibus brevibus diffusis simplicibus, foliis parvis subsessilibus subtus nervulosis acutiusculis inferioribus ovatis superioribus oblongo-ellipticis calyces subsuperantibus, cymis 3-4-floris pedicellatis, bracteis subulatis, calycis cylindrici hispidi fauce hirti dentibus lanceolato-subulatis erecto-patulis tubum dimidium æquantibus inferioribus binis angustioribus a basi subulatis, corollæ roseæ tubo subincluso, nuculis..... ♀.

Hab. in rupestribus submontosis Cretæ inter Maglia et Kainourochorio Eparchiæ Mirabello (Heldr!) ad Aradena Eparchiæ Sphakia (Raul!).

Rami 3-4-pollicares, calyces 2 lineas longi. Affinis *M. Græcæ* et præsertim *M. canescens* Bth. cuius indumentum patule villosum habet et a quâ differre videtur rami brevibus diffusis et calycis dentibus subinæquilatis. An ambo tamen ulterius observatæ ad *M. Græcæ* varietates non referendæ?

9. M. Sinaica (Bth. Lab. pag. 380) adpresso pubescens canescens suffruticosa ramosissima ramis novellis erectis crassiusculæ virgatis elongatis rigidis simplicibus et parce ramulosis, foliis minutissimis paucis remotis subsessilibus ovatis margine subrevolutis subtus elevatim nervulosis inferioribus ovatis superioribus oblongis flores æquantibus vel subsuperantibus, verticillastris sessilibus multifloris parvis remotis longe et laxissime spicatis. floribus minimis breviter pedicellatis, calycis seabridulo-pruinosi ovati dentibus lanceolatis acutiusculis rectis tubo plus dimidio brevioribus, nuculis ellipticis obtusiusculis ♂.

Hab. ad rupes Sinaiticas (Bové! Schimp. 265!). Arabice *Sileslē*.

Rami cum spicæ 8-12-pollicares, folia 2 lineas longa. calyx lineæ $\frac{3}{4}$ longus. Congeneribus magis suffruticosa, insignis foliis minimis et calyce ovato.

Ar. Geogr. Nubie littoralis montes (Schweinf!).

10. M. microphylla (Urv. Enum. p. 71 sub *Thymo*) ramis ex rhizomate indurato filiformibus sœpius decumbentibus simplicibus glabris vel patule hirtis, foliis minutis subsessilibus ovatis vel oblongis acutiusculis margine subrevolutis subtus nervulosis glabris vel viscidulo-hirtis, cymis axillaribus breviter pedunculatis laxis laxe racemosis, foliis floralibus conformibus cimas subæquautibus, bracteis subulatis minimis, calycis patule hispiduli dentibus a basi latiore subulatis rectis tubo triplo brevioribus inferioribus subangustioribus, fauce villosa, corollæ roseæ hirtæ tubo brevissime exerto, nuculis ellipticis minutissimis ♀. *M. microphylla* Bth. Lab. pag. 377. — *M. Sphaerotica* Boiss. et Heldr. Ser. I, XII. pag. 220 et Ser. II, 4, p. 13.

Hab. ad rupes regionis sempervirentis in fauce Aretana prov. Sphakia (Heldr!).

Rami tenuissimi flexuosi intricati 3-4-pollicares, folia 2-2 $\frac{1}{2}$ lineas longa calyx lineam longus; formam hispidulam e Cretâ tantum vidi. *Satvicia filiformis* Desf. Atl. II. p. 8, tab. 121 ramis elongatis rectis, cymis multi- et densifloris etc. mihi species diversa.

Ar. Geogr. Sicilia, Melita.

11. M. Persica (Boiss. Diagn. Ser. I, 12, p. 48) adpresse hirtula vel cinerascenti-velutina, basi suffrutescens ramosissima ramis filiformibus elongatis rigidulis, foliis parvis sessilibus margine subrevolutis obtusis subtus nervulosis inferioribus ovatis superioribus oblongo-linearibus flores æquantibus, cymis subpedicellatis 2-5-floris axillaribus remotiusculis longe et tenuiter spicatis, bracteis subulatis calyce duplo brevioribus, calycis hirtelli dentibus tubo 4-5-plo brevioribus rectis superioribus lanceolatis inferioribus paulo profundius fassis, corollæ albidae tubo inclusa, nuculis.... ♀.

Hab. in Persiâ australi (Auch. 5190! Ky. 882!), in rupibus calcareis ad Schahpour prope Kaserun 3000' (Haussk!). Fl. Apr.

Rami semipedales ut in *M. microphylla* tenues sed erecti rigiduli, cymulæ magis densiflora et vix pedicellatae, calycis laciniae breviores minus tenues. Affinis quoque *M. biflora* Bth. Indicæ foliis myrtiformibus subcordatis margine eximie incrassatis diversæ.

SECT. III. PSEUDOMELISSA.

12. M. nummularifolia (Boiss. Diagn. Ser. I, 12, pag. 49) palli le virens tota pilis longis patulis superne copiosioribus molliter hirsuta, caulinis e rhizomate ramoso suffrutescenti procumbentibus nudis, ramis novellis filiformibus brevibus foliosis ascendentibus, foliis sessilibus ovatis subcordatis nervulosis superioribus acutiusculis, floralibus diminutis calyces æquantibus, floribus solitariis rarius binis axillaribus breviter pedicellatis in racemum terminalem ovato-oblongum basi laxiuseulum approximatis, calycis longe et molliter hispidae dentibus tubo subduplo longioribus a basi lanceolatâ setaceis erecto-patulis tandem subrecurvis, corollæ purpureæ tubo calyce vix longiore ♀.

Hab. in fissuris rupium calcarearum Libani supra Eden (Boiss!), ad Hasrun (Bl!), in monte Hermon (Pest!).

Rami 2-4-pollicares, folia 3-4 lineas longa, calyx lineas 2 $\frac{1}{4}$, corolla tres lineas longa. Facies *Calaminthæ Corsice*, sed calyx basi æqualis, etc.

13. M. Libanotica (Boiss. Diagn. Ser. I, 12, p. 50) dense tomentosa cana, ramis e rhizomate lignoso crasso numerosis erectis simplicibus pumilis fragilibus, foliis sessilibus ovatis obtusis parvis, cymis axillaribus breviter pedunculatis 7-5-floris remotis laxe racemosis rarius ob inferiores in ramulos evolutas subpaniculatis, foliis

floralibus diminutis, bracteis breviter setaceis, calycis breviter hirti rubelli dentibus æqualibus triangularibus acutis tubo 4-5-plo brevioribus, corollæ roseæ calyce 2 ½-plo longioris tubo exerto ♂.

Hab. in Libano supra Cedros in fissuris rupium (Boiss!), supra Eden (Lab! Bl!), inter Dimam et Yamouny (Bl!), in latere septentrionali montis Makmel 8000' (Ky. 298!).

Rami 3-5-pollicares, folia intermedia majora 4-5 lineas longa, racemi 3-4-pollicares, calyx sesquilineam, corolla 2 ½ lineas longa. A formis *M. serpyllifolia* ramis pumilis, indumento non adpresso, cymis paucifloris, calyce longiore et longius dentato diversa.

14. *M. mollis* (Bth. in D.C. Prodr. 42, p. 225) brevissime tomentella canescens, ramis tenellis intricatis filiformibus fragillimis, foliis minutis petiolatis ovato-oblongis obtusiusculis, cymis axillariibus paucis laxe dichotomis 2-5-floris pedunculis folio longioribus suffultis, calycis tomentelli dentibus lanceolatis obtusiusculis erectopatulis tubo 3-4-plo brevioribus, corollâ calyce duplo longiore ♀.

Hab. in Kurdistaniā Mesopotamiā (Ky. 522 a!).

Rami 2-3-pollicares, folia 2-2 ½ lineas longa. Ex speciminibus valde incompletis nota sed a *M. serpyllifolia* et *M. Libanotica* distincta ramis non rectis sed intricatis tenuissimis gracillimis, cymis laxis, paucifloris, longius pedunculatis, racemos vel paniculam non formantibus sed ad axillas paucis remotis.

15. *M. serpyllifolia* (M.B. Taur. Cauc. II, p. 40 sub *Nepeta*) adpressissime cana, caulinis ex rhizomate indurato ramoso erectis elongatis longe paniculatis, foliis breviter petiolatis cuneatis ovato-oblongis integris vel subrepandis superioribus acutiusculis, cymis breviter pedunculatis multifloris plures dichotomis, bracteis linearibus minutissimis, calycis breviter obconici dentibus triangularibus rectis tubo sextuplo brevioribus, villis fauēs dentes æquantibus, corollæ calyce subtriplo longioris labio inferiore sublongiore, nuculis oblongis acutis ♀. Ic. M.B. Cent. Ross. tab. 28 — *M. marifolia* Bth. Lab. p. 382 ex parte quoad Orientem. — *Melissa marifolia* Griseb. Spic. II, p. 125. — *Calamintha spicigera* G. Koch Linn. XXI, pag. 674.

Hab. in rupestribus regionis inferioris et montanae, Tauria (M.B.), Armenia Turcica (G. Koch! Calv!), Taurus Cilicus (Ky. 94! Bal!), Syria borealis circa Aintab et Marasch (Haussk!), rupes ad Seleuciam (Boiss!) circa Sidonem (Gail!).

S. barbata — Calycis sublongioris fauce densius barbati dentes paulo magis elongati tubi quartam partem aequantes. — *M. barbata* B. et Ky. Diagn. Ser. II, 4, p. 14.

Hab. in regione subalpinâ Libani (Ehr!), supra Eden (Bl!), in valle Barghiuté (Gail!), in districtu Danie 5500' (Ky. 342!), in Palestinae monte Ithabure (Gail!).

M. serpyllifolia hucusque confusa fuit cum *M. marifolia* (*Nepeta*) Cavan. Ic. 567. Hispanicâ, indumento tenui cinerascente, cymis paucifloris anguste racemosis, calyce tenuiter cylindrico, corollæ labiis æquilongis, nuculis vix acutis distinctâ.

Ar. Geogr. Albania.

16. **M. congesta** (Boiss. et Haussk.) adpressissime canescens, caulis ex rhizomate indurato numerosis ascendentibus vel erectis simplicibus, foliis petiolatis ovatis obtusis, floralibus calyce multo brevioribus, cymis brevissimè pedunculatis densifloris in spicas terminales oblongas basi saepe interruptas congestis, bracteis squamaeformibus minimis, calycis adpresso tomentelli sulcati dentibus triangularibus obtusis brevissimis, fauce nuda, corollæ albidae tubo vix exerto, nuculæ oblongis acutiusculis ♀.

Hab. in rupestribus calcareis montis Akdaghs Cataoniae inter Adiaman et Malatia (Haussk.).

Subpedalis, folia inferiora 7-8 lineas longa, spicæ terminales pollicares. Affinis *M. serpyllifoliae* a quâ praeter inflorescentiam congestam differt calyce fauce nudo nec barbato.

17. **M. origanifolia** (Vis. Dalm. II, p. 199 sub *Calamintha* non Host) adpresso pubescens punctata pallide virens, caulis ex rhizomate duro basi induratis ascendentibus ramosis, foliis breviter petiolatis ovatis repando-subdentatis, floralibus sensim diminutis, cymis breviter pedunculatis pluri- et densifloris longe et laxiusculæ in racemum spiciformem angustum dispositis, calycis tomentelli sulcati dentibus lanceolato-subulatis tubo duplo brevioribus inferioribus sublongioribus, corollæ albidae tubo breviter exerto ♀. *M. Dalmatica* Bth. in v.c. Prodr. XII, p. 225. — *Melissa Pulegium* Griseb. Spic. II, p. 125 non Rochele.

Hab. in Thraciâ prope Philippopolin (Friv!).

A *M. serpyllifolia* et *congesta* dentibus calycis subulatis subinaequalibus distincta, magis affinis *M. Pulegio* plantæ Bannaticæ que differt foliis dentatis et calycis fauce hirsutâ.

CALAMINTHA (Mœnch. Meth. 408. — Bth.).

Calyx tubulosus 13-nervius bilabiatus (interdum obscure) labio superiori tridentato inferiore in dentes binos subulatos fiso. Corollæ tubus rectus, labium superius planum vel leviter concavum inferius trifidum. Stamina 4 didynama ascendentia per paria plus minus conniventia, antheræ loculi paralleli vel divergentes. Stylus bifidus. nuculæ lœves. — Herbae annuae vel perennes basi interdum suffrutescentes. — Genus speciebus nonnullis obscure bilabiatis, a cæteris tamen non dissociandis, *Micromeria* affine.

Specierum Orientalium dispositio.

SECT. I. EU CALAMINTHA Bth. — Verticillastra cymis compositis constantia. Corollæ tubus intus non annulatus.

C. grandiflora, betulifolia, Pamphylica, officinalis, Nepeta, incana, Cretica, umbrosa, Clinopodium.

SECT. II. CYCLOTRICHUM. — Verticillastra cymis compositis constantia. Calyx saepe vix bilabiatus. Corollæ tubus intus sub staminum insertione pilorum annulo instructus.

C. organifolia, florida, Haussknechtii, staminea, depauperata, nivea.

SECT. III. ACINOS Bth. — Verticillastra pedicellis rigidis indivisis constantia. Calyx basi gibbus saepius incurvus. Corollæ tubus intus non annulatus.

• • Perennes vel perennantes

C. Alpina, suaveolens, Patavina.

• • Annuæ.

C. Acinos, graveolens, maritima.

SECT. I. EUCLAMINTHA.

1. **C. grandiflora** (L. Sp. 827 sub *Melissâ*) sparsim patule et longe papillari-hirtella, caulis e rhizomate repeute ascendentibus elongatis simplicibus, foliis petiolatis magnis ovatis vel oblongis acutiusculis profunde serratis infimis subcordatis cæteris basi rotundatis et cuneatis, floralibus ellipticis sensim diminutis, cymis axillaribus pedunculatis 3-7-floris, bracteis linearis-subulatis, calycis sparsim hirti fauce subnudi labiis patentibus dentibus ciliatis superioribus ovatis breviter acuminatis inferioribus sublongioribus lanceolato-subulatis, corollâ roseâ calyce 2-3-plo longiore, nuculis atris ♀. *C. grandiflora* Mænch Meth. — Ic. Bot. Mag. tab. 208.

Hab. in sylvis regionis montanæ et subalpinæ Græciæ in Taygeto (Heldr!), montibus Kyllene, Parnasso, Olympo Thessalo (Heldr! Orph!), montibus Macedoniae et Thraciae (Griseb!), Olympo Bithyno (Boiss!), Anatoliâ boreali (Wied!), Ponto Lazico (Tchih! Bal!), Tauriâ (M.B.), Caucaso occidentali et centrali (Rupr!), Amano Syriæ (Ky. 83!).

Ar. Geogr. Gallia, Helvetia et Germania australes, Italia, regio Danubialis, Dalmatia, Africa borealis.

2. **C. betulifolia** (Boiss. et Bal. Diagn. Ser. II, 4, p. 14) caulis ascendentibus simplicibus vel inferne parce ramulosis elongatis superne breviter hirtellis, foliis pallide virentibus subtus pallidioribus minute puberulis petiolatis a basi truncatâ vel rotundatâ ovatis acutiusculis acutiuscule dentatis, floralibus diminutis acutioribus, cymis 5-7-floris breviter pedunculatis, bracteis oblongo-rhombeis acutis calyce brevioribus, calycis sub lente pruinosis vix bilabiati dentibus sub-

æqualibus rectis ciliatis ovato-triangularibus breviter acuminatis, fauce subnudâ, corollâ roseâ calyce 2 1/2-plo longiore ♀.

Hab. in rupestribus regionis inferioris in valle Guzelderé prope Sedichig Cilicie (Bal. 496!). Fl. Apr.

Pedalis, facies *C. grandifloræ*, folia subminora et breviora, corolla 10-12 lineas longa. Differt præterea a *C. grandifloræ* bractearum formâ, calycis vix bibrati dentibus rectis.

3. **C. Pamphylica** (Boiss. et Heldr. Diagn. Ser. I, 12, p. 52) patule et molliter villosa, ramis ex rhizomate ramoso tenui procumbentibus vel ascendentibus fragillimis brevibus simplicibus, foliis parvis petiolatis triangulari-ovatis basi subcordatis vel truncatis acutiusculis denticulatis, cymis in axillis summis subtrifloris pedunculo petiolo subæquilongo suffultis, bracteolis linearibus brevibus, calycis tubulosi parce hispidi subbilabiati dentibus æquilongis a basi triangulari acuminatis tubo multoties brevioribus, inferioribus subcurvalis, corollæ tubo calyce duplo longiore ♀.

Hab. ad rupes calcareas montis Ghibelleis supra Alaya Pamphyliæ orientalis 2000' (Heldr!).

Rami 3-6-pollicares, folia majora 4 1/2-5 lin. lata 4 longa petiolo subæquilongo, calyx 5 lineas longus, corolla 9 lineas longa eâ *C. grandifloræ* tenuior.

4. **C. officinalis** (Mænch Meth. p. 409) patule hirta, caulis e rhizomate repente erectis simplicibus vel parce ramosis, foliis majusculis petiolatis ovatis dentato-subserratis, cymis axillaribus pedunculatis laxifloris unilateralibus inferioribus folia æquantibus, bracteis subulatis subulatis parvis, pedicellis calyces subæquantibus, calycis dentibus superioribus lanceolatis recurvis inferioribus sublongioribus rectis linearis-subulatis, fauce parce pilosâ, corollâ roseâ calyce 2-3-plo longiore, nuculis subrotundis ♀. Ic. Rehb. tab. 75, fig. 2. — *Melissa Calamintha* L. Sp. 358.

Hab. in sylvaticis regionis montanae, Macedoniae in monte Korthiati (Orph!), Athone (Sibth., Orph!), Thraciae prope Kalofer (Pichl!), Byzantii (Sibth.), Ponto Lazico ad Trapezuntum (Kyl!) et Rhizé (Bal!), Transcaucasiæ (Rupr!), Persia boreali (Szov!).

Ar. Geogr. Europa media et australis a Bataviâ et Germaniâ mediâ ad regionem Danubialem et Dalmatiam.

5 **C. Nepeta** (Linn. Sp. 828 sub *Melissa*) molliter pubescenticanescens. caulis a basi ascendentè erectis simplicibus vel ramosis paniculatis, foliis parvis petiolatis ovato-rhombeis obtusis crenatis, cymis axillaribus unilateralibus densiusculis pedunculatis folia floralia superantibus laxe racemosis, bracteis subulatis minimis, pedicellis calyce subbrevioribus, calycis fauce longius pilosi dentibus superioribus lanceolatis recurvis inferioribus longioribus subulato-acuminatis, corollâ calyce subdupo longiore, nuculis oblongis ♀. *C. Nepeta* Clairv. Man. 197. — *C. rotundifolia* Host. — *C. Spruneri* Boiss. Diagn. Ser. I, XII, p. 53 (forma foliis majoribus). — *C. nepetoides*

Jord. Obs. 4, p. 46, tab. 2. — *C. canescens* et *C. Byzantina* C. Koch Linn. XXI, p. 672!

Hab. in umbrosis, ad sepes, in humidiusculis regionis inferioris et montanae a Graciā et ejus insulis!, Macedoniā! et Thraciā! Cretā (Heldr!), ad Anatolian occid. (Bal!), meridionalem in Ciliciā (Bal!). Syriam borealem ad Alexandrette (Haussk!), Anatolian borealem et Armeniam (Wied! Calv!), Tauriam et Transcaucasiam (M.B., Hoh!), Persiam bor. in pr. Ghilan (Buhse!).

Ar. Geogr. Europa media et australis ab Anglia et Germaniā mediā ad regionem Danubialem, Africa borealis.

6. *C. incana* (Sibth. et Sm. Fl. Græc. VI, p. 62, tab. 577 sub *Thymo*) tota dense tomentosa cana, caulinibus decumbentibus elongatis ramosis, foliis parvis petiolatis orbiculatis subintegris floralibus diminatis deflexis flores non æquantibus, cymis axillaribus sessilibus 6-2-floris laxissime racemoso-spicatis, pedicellis calyce longioribus, calycis obsolete gibbi fauce villosâ labio superiore lato brevissime tridenticulato inferioris æquiflori dentibus lanceolatis, corollæ carneæ calyce duplo longioris tubo vix exerto, nuculis ovato-globosis ♀.

Hab. in campis Attice et Archipelagi (Sibth., Sprn! Heldr!), Byzantii (C. Koch! Wied!), Cypri ad Larnaca (Ky. 9-0 sub *C. Creticā*!), Aleppo (Russ.), Syriæ littorali et Libani regione inferiore (Bl! Gail!), Palestinæ (Roth!).

Variat tomento in parte superiori plantæ rarescenti. Circa Athenas formæ inter hanc et *C. Nepetam* hybridae interdum occurunt. *Acinos incanus* Griseb. Spic. est species diversa, *C. maritima* Bth.

7. *C. Cretica* (L. Sp. 828 sub *Melissā*) longe patule et flexuose piloso-hispida canescens basi ramosissima suffruticosens, ramis novellis tenuiter virgatis ascendenti-erectis simpliciuseulis, foliis parvis breviter petiolatis ovatis obtusis subintegris, floralibus subdiminutis calyces subæquantibus, cymis subtrifloris breviter pedunculatis distantibus laxissime et anguste spicatis, pedicellis brevissimis, calycis patule hirsuti subbilabiati fance villosâ dentibus lanceolatis rectiusculis inferioribus sublongioribus et profundius fisis, corollæ albidâ calyce duplo longiore ♂. *C. Cretica* Bth. in B.C. Prodr. XII, p. 227. — *Thymus hirtus* Sieb

Hab. in Cretæ regione montanâ superiore in torrentium alveis et glareosis 3500'-4000' ad Omalο, Askyphous etc. prov. Sphakia (Sieb! Heldr!). Vulgo *Libanochoro*.

Pedalis, folia majora 3-4 lineas longa et lata, spicæ semipedales. A *C. incanā* pilis patulis, ramis inferne suffruticosis, calycis dentibus omnibus elongatis etc. distinctissima.

8. *C. umbrosa* (M.B. Taur. Cauc. II, p. 63 sub *Melissā*) viridis patule pubescens, caulinibus elongatis diffusis vel ascendentibus inferne radicantibus, foliis petiolatis ovato-oblongis basi rotundatis vel cuneatis repando-crenatis, floralibus sensim diminutis, verticillastris subsessilibus remotis globosis densiusculis, bracteis subulatis pedicellis vix longioribus, calycis patule hirsuti submeurvi bilabiati

dentibus superioribus lanceolatis inferioribus longioribus profundius fassis lanceolato-subulatis ciliatis, corollà parvâ calyce sesquiloniore ♀. — *C. umbrosa* Bth. in p.c. Prodr. XII, p. 232. — *Arinos multiflora* C. Koch MSS.

Hab. in sylvaticis regionis inferioris Ponti Lazici ad Rhizò (Bal!), Imeretiae et Gurie (C. Koch!), Iberiae (M.B.), Daghestanicae (Bupr!), prov. Talysh (G.A. M!), Persia; hor. in prov. Asterabad (Bge!), regni Cabulici (Griff.).

Ab eâ vix differre videtur *C. repens* Bth. loc. cit. p. 233.

Ar. Geogr. Jugum Himalaicum, montes Peninsulae Indiae.

9. **C. Clinopodium** Bth. in p.c. Prodr. XII, p. 233) paule pubescens vel villosa, caulis erectis elatis simplicibus vel ramosis, foliis breviter petiolatis ovatis obtusis suberenatis basi rotundatis, floralibus oblongis acutiusculis sape deflexis, verticillastris sessilibus remotis multifloris densis, bracteis setaceis plumosis ad basin verticillastris numerosis et ad florum basin calycem aequantibus, calycis incurvi patulae hirsuti bilabiati dentibus plus minus copiose et longe ciliato-plumosis lanceolatis labii inferioris multo profundius fassis, corollà purpureâ calyce 2-3-plo longiore ♀. Ic. Rehb. t. 73, fig. 1. — *Clinopodium vulgare* L. Sp. 821. — *Clin. Egyptiacum* Lam. — *Cl. plumosum* Sieb. Bot. Zeit. 1822, p. 242.

Hab. in umbrosis siccis praesertim regionis montane a Græciâ et Macedonia ad Anatoliam borealem et australem, Taurum Cilicicum (Bal!), Cyprum (Ky!), Libanum (Ehr! Bl! Gaill!), Cataeniam (Haussk!), Tauriam Caucasum et Transcaucasiam (M.B., Hol!), Persiam borealem (Szov!), Kurdistaniam Persicam in monte Avroman (Haussk!).

Ar. Geogr. Europa omnis a Britanniâ et Scandinaviâ ad Rossiam, Sibiria, Africa borealis.

SECT. II. CYCLOTRICHUM.

10. **C. origanifolia** (Lab. Dec. 4, p. 14, tab. 9 sub *Clinopodio*) glabriuseula sub lente pruinosa vel adpresso velutina grisea, ramosissima ramis e rhizomate lignoso ascendentibus flexuosis simplicibus rarius parce ramosis, foliis remotis parvis sessilibus suborbiculatis obtusis integris punetatis, floralibus ellipticis verticillastro brevioribus, verticillastris 3-4 sessilibus multifloris remotissimis, bracteis linear-lanceolatis calyce subbrevioribus, calycis glanduloso-punctatâ subincurvi subbilabiati fauce barbatâ labio superiore ad medium in dentes triangulares acuminatos fisso inferiore ad basin usque in dentes lanceolato-subulatos partito, corolle tubo calyce vix longiore, filamentis subexsertis ♀. — *Micromeria? origanifolia* Bth. in p.c. Prodr. XII, p. 214. — *C. glomerulosa* Boiss. et Heldr. in p.c. Prodr. XII, p. 284.

Hab. inter lapides montis Davrosdagh Pisidiæ 6900' (Heldr!), in Libano supra Eden, Ain el Karn, in montibus Sanin et Makmel (Labill! Ehrenb! Bl!).

Rami 6-10-pollicares, folia 3-4 lineas longa lataque, verticillastra 1-2-pollices

distantia, calycis dentes in iconæ Billardieri erroneè mutici depicti; forma glabrescens velutinæ rarior videtur.

11. **C. florida** (Boiss. Diagn. Ser. I, 12, p. 51) basi suffruticosa ramosissima, ramis elongatis erectis flexuosis subsimplicibus pubescentibus foliis subpetiolatis ovatis obtusis integris minute puberulis, floralibus inferioribus verticillastræ æquantibus, verticillastris sessilibus multifloris remotis, bracteis linear-lanceolatis acuminatis calyce brevioribus, calycis patule scabrido-hirti subcurvati bilabiati labio superiore ad medium in dentes lanceolatos fiso, fauce barbatâ, inferiore in dentes subulatos ciliatos bipartito, corollæ violaceæ tubo exerto, filamentis longe exsertis ♀.

Hab. in regione subalpinâ et alpinâ Tauri Cilicici supra Gulek Maaden 5500'-6000' (Ky. 478! et 143! Ball!).

Semipedalis et pedalîs, folia majora 9-10 lineas longa. Differt a *C. origanifolia* foliis majoribus, corollæ tubo staminibusque longius exsertis, a *C. staminea* indumento, calycis vix bilabiati dentibus æqualibus rectis.

12. **C. Haussknechti** (Bunge Lab. Pers. pag. 36) patule glauculos-pubescentes canescens, ex rhizomate indurato ramosissima ramis erectis pumilis simplicibus et parce ramosis, foliis parvis subsessilibus ovatis acutuseulis integris, verticillastris subsessilibus sessilibusque dense multifloris distantibus breviter spicatis, bracteis lanceolatis calyce subbrevioribus, calycis recti glabri vix bilabiati dentibus æquilongis erectis lanceolatis ciliatis dimidium tubum æquantibus anterioribus profundius fissis, corollæ tubo subexerto, filamentis exsertis ♀.

Hab. in fissuris rupium calcarearum Teng Tokab prope Behehan Persiae australis 3000' (Haussk!).

Rami 4-6-pollicares, folia majora 6 lineas longa, calyx tres lineas longus; *C. staminea* calycis tubo et dentibus rectis affinis sed glandulosa, dentes tubi proportione longiores, corollæ tubus breviter exsertus, etc.

13. **C. staminea** (Boiss. Diagn. Ser. I, 12, p. 51) glabra sub lente minute pruinosa basi suffruticosa et ramosa, ramis erectis virgatis simplicibus vel stricte ramulosis, foliis subpetiolatis ovatis basi truncatis vel subcordatis integris superioribus acutuseulis floralibus minimis ellipticis acuminatis, verticillastris sessilibus 8-12-floris distantibus interrupte spicatis, bracteis lanceolato-linearibus calyce brevioribus, calycis tubulosi subcurvati obscure bilabiati fauce villosâ dentibus lanceolato-subulatis strictis subæquilongis tubo quintuplo brevioribus inferioribus profundius fissis, corollæ roseæ calyce $2\frac{1}{2}$ -plo longioris tubo exerto, filamentis longe exsertis ♀. *Micromeria staminea* Boiss. et Hoh. Diagn. Ser. I, 5, p. 19. — *C. glabrescens* Boiss. in Ky. exs.

Hab. in rupestribus montis Gara Kurdistaniæ (Ky. 511!), in subalpinis supra Musch Armeniae 6500' (Ky. Suppl. 688!).

Rami cum spicâ 3-4-pollicari $\frac{1}{2}$ -1-pedales, folia majora 7-9 lineas longa,

verticillastra 6-10, calyx 3 lineas, corolla 6-7 longi. Quamvis calyx obscure bilabiatus sit ob affinitatem cum *C. origanifolia* et *C. floridæ* *Calaminthis* certe adnumeranda.

14. *C. depauperata* (Bge Lub. Pers. p. 38) crispule et breviter pubescens suffruticosa ramosissima, ramis novellis virgatis erectis gracilibus subsimplicibus, foliis parvis subpetiolatis ovato-orbiculatis inferioribus integris obtusissimis floralibus diminutis ovatis acutis acute paucidenticulatis, cymis axillaribus 3-1-floris subpedunculatis laxissime et tenuiter spicatis, bracteis lanceolatis pedicello æquilongis, calycis tubulosi recti glabri ore villosi vix bilabiati dentibus rectis a basi lanceolatâ longe subulatis dimidio tubo longioribus inferioribus paulo profundius fisis, dentium sinibus margine subrevolutis, corollæ tubo inclusu, staminibus exsertis ♀.

Hab. in rupibus calcareis prope Bors in valle fluvii Chyrsan Susianæ (Haussk!).

Sesquipedalis, folia majora 4-5 lineas longa et lata, calyx 4 lineas fere excdens; inter aflies cymis depauperatis, calycis dentibus longis insignis

15. *C. nivea* (Boiss. Ann. Sc. Nat. 1854. II. p. 253) dense tomentosa nivea basi suffrutescenti ramosissima, ramis erectis virgatis simplicibus vel inferne ramosis, foliis parvis subsessilibus ovatis obtusis integris, floralibus diminutis verticillastro brevioribus, verticillastris plurifloris densis sessilibus remotis laxè spicatis, bracteis subulatis calyce triplo brevioribus, calycis tomentosi tubulosi recti vix bilabiati fauce villosâ dentibus æquilongis brevissimis superioribus triangularibus inferioribus subangustioribus lanceolatis profundius fisis, corollæ tube subincluso, filamentis longe exsertis ♀. Ic. Tchih. As. Min. tab. 24.

Hab. in lapidosis regionis montane Cappadociae inter Albistan et Gurum (Tchih!), in monte Akdaghs Cataoniae inter Adiaman et Malatia 6000' (Haussk!).

Fere pedalis, folia majora 6 lineas longa, verticillastra 5-7 pollicem vel amplius distantia, calyx 3 lineas, corolla vix 5 longi. Species a precedentibus indumento panno nivo calyceque brevissime dentato distincta.

SECT. III. ACINOS.

16. *C. Alpina* (L. Sp. 826 sub *Thymo*) perennis, brevissime puberula vel patule pubescens, ramis ex rhizomate tenui indurato a basi diffusâ radicanti ascendentibus flexuosis brevibus, foliis parvis nervosis petiolatis ovatis vel elliptico-rhombeis acutis parte superiori subserratis, floralibus calyces subæquantibus, floribus ad axillas superiores geminis vel ternis, calycis nutantis tubo incurvo basi antice gibbo labiis tubo duplo brevioribus superiore ad medium in dentes triangulares fisso inferiore in dentes subulatos bipartito dentibus omnibus ciliatis fructiteris erecto-patulis, corollæ violaceæ calyce

$2\frac{1}{2}$ -plo longioris tubo in faucom valde ampliato φ . *C. Alpina* Bth. — *Acinos Alpinus* Mænch Ic. — Rehb. t. 74, 1.

Hab. in lapidosis regionis subalpinæ et alpine, Græcia (Auch. 1645!), mons Malevo Laconia (Orph. Fl. Gr. exs. 532!), Chelmos supra Stygem et Olympus Thessalus (Heldr!), Macedonia in Scardo (Griseb., Orph!), Thracia in Rhodope (Friv.), Olympus Bithynus (Boiss! Thirke!), Cephalonia in monte Oenos (Heldr!), Cretæ montes (ex Heldr.).

Ar. Geogr. Montes Hispaniæ borealis, Galliæ mediae et australis, Helvetiæ, Germaniæ australis, Italiae et Siciliæ, regionis Danubialis, Dalmatiæ, Africæ borealis.

17. *C. suaveolens* (Sm. Prodr. I, p. 420 sub *Thymo*) perennis molliter et patule canescens, ramis ex rhizomate indurato ascendentibus, foliis breviter petiolatis ellipticis vel elliptico-linearibus rigidis acutis parce et remote denticulatis, floralibus calyces vix superantibus, verticillastris 6-9-floris spicas longas laxiusculas formantibus, calycis nutantis patule hirsuti vix incurvi basi gibbi dentibus subæquilongis a basi sublatiore longe subulatis rectis conniventibus tubo quadruplo brevioribus, corollæ roseæ calyce duplo longioris tubo exerto superne subinflato φ . *A. acuminatus* Friv. Flora 1835, p. 332. — *C. Patarina* β *acuminata* Griseb. Spic. II, 123.

Hab. in regione montana et subalpinâ Græcia, Taygeto (Heldr!), Kyllene (Heldr! Orph. Fl. exs. 283! et 550!), Parnasso (Sibth., Sprunt!, Heldr. Herb. Norm. 221!), Olympo Thessaliae (Auch. 1641!), Macedonia supra Thessalonicanam et in Scardo (Orph!), Thraciâ (Friv!).

A nonnullis pro *C. Acini* formâ habita sed multo major, calycis structura diversa, corolla multo major longius exserta. Magis affinis *C. Patavinae* a quâ distinguitur rhizomate suffrutescenti, foliis angustioribus acutioribus, calycis dentibus subæquilatis.

18. *C. Patavina* (Jacq. Obs. 4, p. 7, tab. 87) perennans pubescens, ramis herbaceis erectis vel ascendentibus, foliis petiolatis ovato-rhombeis vel orbiculatis remote denticulatis superioribus acutis, floralibus calyces superantibus, floribus ad axillas superiores subternis laxe racemoso-spicatis, calycis subinecurvi basi gibbi dentibus subæquilongis labii superioris lanceolatis sursum subinclinati inferioris subulatis, corollæ calyce duplo longioris tubo subincluso ②. *C. Patarina* Host — *Acinos rotundifolius* Friv. vix Pers. — *C. rotundifolia* Rehb. Ic. tab. 74, fig. 2. — *A. alpinus* β *elatior* Griseb. Spic. II, p. 122. — *Acinos ascendens* Mænch.

Hab. in regione superiore Macedonie in monte Corthiatî (Orph!), Thracie (Friv!).

Planta saepè pedalis foliis majusculis latis, corollæ limbo magno.

Ar. Geogr. Dalmatia, Croatia, Banatus, Transylvania.

19. *C. Acinos* (L. Sp. 826 sub *Thymo*) annua pubescens vel villosula, caule a basi ramoso ramis ascendentibus vel erectis, foliis petiolatis ovatis vel rhombeis nervosis integris vel subserratis venis subtus distantibus subrectis, floralibus conformibus acutis calyces

superantibus, floribus ad axillas superiores geminatis aut ternis, calycis nutantis tubo incurvo basi antice gibbo labiis tubo duplo brevioribus superiore medium versus subtruncato in dentes ternos subulatos abrupte abeunte inferiore in dentes subulatos bipartito dentibus omnibus ciliatis fructiferis erectis subcommixtibus, corollæ purpureæ calyce subdupo longiore tubo exerto ⊖. *C. Acinos* Bth. in d.c. Prodr. — Engl. bot. tab. 411. — *Acinos thymoides* Mœch.

Hab. in cultis Peloponnesi (Sibth.), Thraciae et Macedoniae (Friv!), Anatolia borealis, Tauriae, Caucasi et Transcaucasiae (Led.).

Ar. Geogr. Europa media et australis a Britannia et Scandinavia ad Rossiam.

20. *C. graveolens* (M.B. Taur. Cane. II, p. 60 *Thymo*) annua canescens-villosa, saepius basi ramosa ramis ascendentibus vel erectis pumilis, foliis parvis petiolatis ovatis subserratis acutis subtus elevatim arcuato-nervulosis floralibus angustioribus flores superantibus, verticillastris axillaribus 6-4-floris remotis vel confertis, calycis subsessilis basi subgibbi valde incurvi dentibus dimidio tubo sublongioribus sursum tandem curvatis superioribus a basi lanceolata longe subulatis inferioribus sabbrevioribus a basi subulatis, corolle tubo angusto inclusa limbo minuto ⊖. Ic. M.B. Cent. tab. 38. — *C. graveolens* Bth. in d.c. Prodr. — *Thymus exiguis* Sibth. et Sm. Fl. Gr. VI, p. 61, tab. 575 (forma minor foliis integris). — *Acinos erectus* Friv. exs.

Hab. in siecis regionis montanae, Græcia in monte Malevo (Spr!), Pentelico (Boiss!). Parnassi regione inferiore (Heldr. herb. Norm. 629!), Thracia (Friv! Jank!), Cycladibus (Oliv!), Cretâ (Heldr!), Anatoliâ boreali!, occid. ad Smyru-nam (Bal!), in Pisidiâ (Heldr!), Ciliciâ et Cappadociâ (Bal!), Cypro (Ky. 762!), Armenia (Huet!), Tauro et Transcaucasiâ (M.B., Hoh!). Persia ad Teheran (Bge!) et australi (Ky. 985!), Mesopotamiâ (Oliv!), Syriâ ad Aleppo (Ky. 119!).

Species Hispanica valde affinis dentibus calycis brevioribus vix sat distincta est *C. purpurascens* Pers. cuius *Acinos rotundifolius* Pers. non Friv. est probab. synonymon.

21. *C. maritima* (Bth. in d.c. Prodr. XII, p. 231) annua, colo ramosa villosula, ramis prostrato-ascendentibus pumilis, foliis minimis petiolatis transverse ovatis obtusissimis vel retusis a medio obsolete crenatis supra viridibus subtus canis nervis prominulis fere rectis nitrinque hirsutis, floralibus rotundato-spathulatis mucronatis calyces superantibus, floribus in axillis supremis 2-3-nis brevissime pedicellatis confertis, calycis hispida rectiusculi basi subgibbi labio superiore in dentes a basi late triangulari subulatos sursum curvatos inferiore in dentes subulatos sabbreviores divisis, corollæ calyce subdupo longiore tubo inclusa ⊖. *Acinos iucanus* Griseb. Spie. II, p. 123 excluso Syn. Fl. Græcae.

Hab. in arenosis maritimis Propontidis in Thracia prope Hersek (Griseb!).

Rami in specimine bipollicares, spicæ oblongo-capitate vix pollicares, folia 2-2 $\frac{1}{2}$ lin. lata 1 $\frac{1}{2}$ -2 longa, calyx vix tres lineas longus. Foliorum formæ et minutie insignis.

MELISSA (L. Gen. 728).

Calyx campanulatus 13-costatus bilabiatus tubo postice plano labio superiore subplano tridentato inferiore bipartito. Corollæ tubus exsertus recurvato-ascendens intus nudus, labio superiore erecto retuso inferiore patente trifido. Stamina 4 didynama sub labio superiore arcuato-conniventia, antherarum loculi divaricati. Stylus bifidus. Nuculæ ovatae lœves. — Herba perennis.

1. M. officinalis (L. Sp. 827) hirsuta vel villosa, erecta ramosa saepè elata, foliis ovatis saepè cordatis obtusis vel acutiusculis serrato-dentatis, floralibus sensim diminutis flores superantibus, cymis axillaribus subsessilibus 3-5-floris secundis, calyce in pedicello demum nutante hirsuto labio superiore lato reticulato-venoso dentibus brevissimis, inferioris dentibus lanceolatis aristatis, corollæ albâ calyce subdupo longiore ♀. Ic. Rehb. tab. 60. — *M. graveolens* Host == *M. cordifolia* Pers. — *M. altissima* Fl. Gr. VI, p. 62, tab. 579.

Hab. in umbrosis, dumosis, ad aquas regionis inferioris sed saepius montanæ a Græciâ! et insulis! Macedonia et Thracia! ad Tauriam, Caucasum' et Transcaucasiam, Syriam littoralem! et Libanum!, Persiam borealem! et orientalem (Bge!).

Ar. Geogr. Europa australis, Lusitania et Hispania, Corsica, Italia, Dalmatia, regio Danubialis, Africa borealis.

HYSSOPUS (L. Gen. 709).

Calyx tubuloso-campanulatus 15-nervius ore subobliquo dentibus 5 aequalibus fauce nudâ. Corollæ tubus rectus calycem aequans, limbi labium superius planum retusum, inferius trilobum lobo medio latissimo bilobo. Stamina 4 didynama exserta divergentia, antheræ loculi divaricati. Stylus bifidus. Nuculæ ovoideæ triquetrae. — Suffrutescens foliis angustis integris.

1. H. officinalis (L. Sp. 796) basi suffrutescens multicaulis, caulis erectis simplicibus vel parce paniculato-corymbosis, foliis punctatis sessilibus lanceolatis vel linearibus planis vel margine subrevolutis, axillis fasciculigeris, floralibus abbreviatis calyces aequalibus, verticillastris subsessilibus in spicas terminales basi saepè interruptas approximatis, calycis dentibus lanceolatis acuminatis ♀.

β. angustifolius. — Rami breviter retrorsum et crispule tomentelli, folia anguste linearia margine saepè revoluta infirma tomentella, calycis dentes dimidio tubo aequilongi vel sublongiores. — *H. angustifolius* M.B. Taur. Cauc. II, p. 38. — *H. Orientalis* Willd. En. Berol. II, p. 600. *H. Caucasicus* Spr.

Hab. in siccis Tauriæ, Caucasi et Transcaucasie (M.B., Rupr! Hoh!), Persiæ borealis (Szov!), in monte Djulfek (Auch. 5145!).

Varietas a typo præter folia angustiora calycis dentibus tubi proportione longioribus distincta sed formæ intermediae occurrere videntur.

Ar. Geogr. Hispania, Gallia australis, Italia, Dalmatia, regio Danubialis, Sibiria Uralensis et Altaica.

ZIZYPHORA (L. Gen. 36).

Calyx anguste tubulosus 13-nervis obsoletissime bilabiatus labio superiore 3-inferiore 2-dentato dentibus brevibus conniventibus, fauce villosa. Corollæ tubus non annulatus in faucem dilatatus, limbi labium superius erectum integrum inferius patens trifidum. Stamina perfecta bina antica sub labio superiore ascendentia, antheræ margine connatae uniloculares vel loculo parvo casso stipitato appendiculatae vel et in eadem specie! exappendiculatae. Filamenta sterilia brevia antheris bilocularibus cassi! vel nulla. Stylus bilobus. Nuculæ leves. — Herbæ perennes facie *Thymi Serpylli* vel amnæ facie *Aenos*. Genus, jam. am. a Bunge animadvertente a *Monardeis* alienum et facie, calycis structurâ, etc. *Satureineis* adsociandum a quibus statim superioribus sterilibus interdum nullis tantum differt

* Perennis.

1. Z. clinopodioides (M.B. Taur. Cae. I. p. 17) ramosissima basi plus minusve suffruticosa ramosissima, foliis glabris minute velutini vel pilosis ovatis oblongis vel lanceolatis, floralibus conformibus minoribus vel sublatioribus, verticillastris in capitulum terminale ovato-oblongum vel globosum congestis, calycis tenuiter cylindrici minute velutini vel villosi dentibus linear-lanceolatis brevissimis obtusiusculis conniventibus superioribus sublongioribus, corollæ tubo breviter exerto, antheris non appendiculatis ♂. vel ♀. Species more *Thymi Serpylli* adnodium variabilis et eius formæ manifeste altera in alteram transeunt. Frequentiores numero.

2. serpyllacea — Humilis, rami saepius tenues saepè decumbentes, folia lanceolata acuta glabra vel hirtula, floralia saepè sublatiora, calyces brevissime hirtuli vel velutini. — *Cunila capitata* L. Sp. 30. — *Thymus punctatus* Willd. Phyt. I, p. 8. — *Z. serpyllacea* M.B. I, Taur. Cae. p. 17.

Hab. in Armeniâ Turcicâ ad Baibut, Maimansur, Tortum, etc (Huet! Bourg!), in prov. Musch (Ky. 350!), Caueaso (M.B., C.A. Mey!), Georgia Caucasicâ ad Titlis etc. (Hohen!).

3. canescens. — Humilis suffruticosa, folia latiora ovato-oblonga canescens vel glabra, floralia saepè latiora, calyces adpresso cani vel molliter pubescentes. — *Z. canescens* Bth. Lab. p. 321. Forma

sæpius australior. — *Z. Gundelsheimeri* C. Koch Linn. XVII, 294.
— *Z. Brantii* C. Koch Linn. 47, p. 294!

Hab. in regione alpinâ Armeniæ (Gund. herb!), Tauri Lycaonici prope Turchalar (Heldr!) et Ciliciei (Ky. 469! Bal!), montis Akdaghs Cataoniac et Berytida (Haussk!), Libani 5-6000' (Ehr! Boiss! Bl! Ky. 272!), Hermonis (Boiss! Ky! Gaill!), Kurdistanie in monte Gara (Ky. 523!), montium Avroman, Schahu, Dalechani, Pir Omar Gudrun, Kurdistanie Persicæ 6-900' (Haussk!).

Z. rigida. — Magis fruticosa, ramæ rigidæ sæpius elongati interdum pedales, folia lanceolata acuta, calyces adpresso pubescentes interdum præterea patule villosuli. — *Z. fasciculata* C. Koch MSS.

Hab. in Armeniæ Rossicæ et Persiæ boreali (Szov!), monte Elbrus Persicæ bor. (Ky. 408!), Persiæ mediæ et australi (Auch. 1639! et 5115!), montis Schahu, Sawers, Eschker, Kuh Nur 8-13000' (Haussk!), Daëna (Ky. 698!), in regno Cabulico (Griff!), monte Chehel Tun Belutschiae (Stocks!).

Z. dasyantha. — Rami tenuiores decumbentes magis herbacei, folia orbiculata ovata vel ovato-oblonga obtusa glabra vel villosula, calyces hispidissimi vel lanati interdum tamen parce hirsuti et ad var. *z* transeuntes. — *Z. dasyantha* M.B. Taur. Cauc. I, p. 18. — *Z. nummularia* F. et M. Ann. Sc. Nat. 1854, p. 33!

Hab. in regione alpinâ Caucasii (M.B., C.A. Mey!) in Salataviâ et Daghستانiâ 7000' (Rupr!), in ditione Talysh prope Swant (Hoh!) et monte Totschal Persicæ borealis (Ky. 574!). (Formæ ad var. *serpyllaceam* transeuntes.

Ar. Geogr. Songaria, Sibiria Altaica.

** Annuæ.

2. *Z. capitata* (L. Sp. 31) annua adpresso canescens pumila erecta a basi subcorymbose ramosa, foliis lanceolatis et linearibus acutis, floralibus dilatatis arcuatim nervosis rhomboeocvatis attenuato-acuminatis capitula involuerantibus et superantibus, floribus dense globoso-capitalis, calycis longe et tenuiter cylindrici piloso-hispidi dentibus linearilanceolatis brevibus conniventibus, corolle tubo tenui breviter exerto, antheris non appendiculatis ♂. Ic. Fl. Græc. tab. 13. — *Z. compacta* Friv. Flora 1835, p. 336.

Hab. in cultis et collibus saxosis, Græcia in monte Malevo Laconiae (Orph!), Argolide, Thessaliâ, Etoliâ (Sprun!), Atticâ (Heldr. herb. Norm. 600!), Thraciâ (Friv!), Byzantii (Cast!), Anatoliâ boreali (Wied!), Bithyniâ, Lydia in monte Mesogi. Cariæ in Cadmo (Boiss!), Lyciâ (Heldr!), Ciliciâ (Bal!), Syriâ littorali (Gaill!) et interiori ad Aleppo (Boiss! Haussk!), Cypro (Sibth!), Palestinâ et Arabia petrieâ conterminâ (Boiss!), Armeniâ (Bourg!), Tauriâ (M.B!). Transcaucasia (Hoh!), Persiæ boreali (Buhse! Bge!) et australi ad Persepolin (Ky. 833!).

Ar. Geogr. Hispania interior, Italia anstralis, Etruria, Serbia, Africa borealis.

3. *Z. Hispanica* (L. Sp. p. 31) annua breviter pubescens pumila a basi ramosa ramis erectis et ascendentibus, foliis brevissime petiolatis ovatis acutis arcuatim nervosis, floralibus conformibus calyces subsuperantibus, verticillastris paucifloris spicas longas densiusculas stricte foliosas formantibus, floribus crasse et brevissime

pedicellatis, calycis anguste cylindrici pubescens vel breviter hirti dentibus lanceolatis conniventibus brevibus, corollæ calyce sesqui-longioris tubo tenuissimo breviter exerto, antheris non appendiculatis ⊖.

Hab. in cultis derelictis planitiei Syriæ inter Edlîp et Aleppo site (Boisst).

Speciminibus Hispanicis simillima. *Z. acinoides* L. est probab. forma hujus speciei verticillastris subdistantibus, calyce longius hirto quæ in Hispaniâ sœpe occurrit.

Ar. Geogr. Hispania, Africa borealis interior.

4. *Z. acutifolia* (Mouth. et Auch. Ann. Sc. Nat. 1836, I, p. 43) annua minute pubescens pumila basi ramosa erecta, foliis oblongo vel elliptico-lanceolatis utrinque attenuatis breviter petiolatis acutiusculis, floralibus abbreviatis ovatis vel oblongis acuminatis acutis calyces superantibus, verticillastris subdistantibus vel contiguis spicas longas laxiusculas vel oblongas densifloras formantibus, floribus subsessilibus, calycis anguste cylindrici breviter puberuli dentibus lanceolato-subulatis brevibus conniventibus, corollæ tubo tenui breviter exerto, antheris non appendiculatis ⊖.

Hab. in agris collinis prope Aleppo Syriæ (Auch., Ky. 203!) inter Aintab et Nisib (Haussk!).

Planta 4-7-pollicaris, facie *Z. Hispanicæ* et *tenuioris*, prior differt foliis inferioribus ovatis nec lanceolatis, calyce breviore, posterior foliis angustioribus seabris valde ciliatis et antheris appendiculatis.

5. *Z. tenuior* (L. Sp. p. 31) annua canescens, pumila a collo ramosa ramis erectis vel ascendentibus, foliis nervosis seabridis rigide ciliatis anguste lanceolatis acutis, floralibus conformibus flores valde superantibus, verticillastris axillaribus sœpe secus totam ramum longitudinem in spicas laxiusculas vel densas foliosas oblongas vel longe cylindricas dispositis, calyce longe et tenuiter cylindrico hispido dentibus ovato-triangularibus obtusis brevissimis conniventibus, corollæ tubo tenui breviter exerto, antheris loculo casso minimo clavæformi appendiculatis rarius non appendiculatis ⊖. *Fuldermannia parviflora* Trautv. Bull. Petersb. VII, p. 21. — *Z. Persica* Bge Lab. Pers. p. 39 forma antheris non appendiculatis.

Hab. in cultis et collibus Cariae in planicie Cybirensi (Boisst), Phrygiae (Bal!), Pisidie (Heldr!), Lycie (Bourg. 204!), Syrie ad Damascum (Ky! Bal!) et in deserto Syriæ (Bl!). Armeniæ Turcicæ (Huet! Bourg!), Assyriæ (Ky! Haussk!), totius Persie australis et borealis (Auch! Ky! Bge! etc.), Afghanistanæ (Griff!), Belutschiae (Stocks!).

Ar. Geogr. Songaria, Sibiria Altaica.

6. *Z. Taurica* (M.B. Tanr. Cane. I, p. 48) annua canescens, pumila basi ramosa ramis ascendentibus vel erectis, foliis linearilanceolatis basi angustatis, floralibus conformibus nervosis seabris ciliatis calyces superantibus, verticillastris axillaribus fere secus totum ramum laxiuscule vel dense spicatis, calycis subsessilis longe

et tenuiter cylindrici hispidi dentibus triangularibus brevissimis conniventibus, corollæ calyce duplo longioris tubo subexerto in limbum magnum ampliato, antheris loculo sterili oblongo stipitato appendiculatis vel non appendiculatis, staminodiis binis sterilibus brevibus saepe obviis ♂. *Faldermannia Taurica* Trautv. — *Dracocephalum odoratissimum* Poir. Suppl. II, p. 521. — *Z. lanceolata* Boiss. exs.

Hab. in arenosis Lydiæ in montibus prope Smyrnam (Bal. 389!), prope Tralles (Boiss!), in planicie ad Philadelphiam (Boiss!), in Cariâ (Auch. 1633!), Paphlagoniâ et Galatiâ (Wied!), Cappadociâ (Ky. Suppl. 292!) et ad Talasse (Bal!), Armeniâ australi (Noë 861!), Syria bor. et Mesopotamiâ ad Orfa et Aintab (Haussk!), Tauriâ meridionali in arenosis maritimis et montibus (Stev! Rehm! etc.), Persiâ (ex cl. Bth.).

Inter specimina Taurica ex eodem loco orta et simillima alia antheris appendiculatis alia antheris absque omni appendice reperio, hæc nota igitur nullius momenti. *Z. Taurica* foliis et facie *Z. tenuiori* simillima ab ea corollæ longius exsertæ limbo multo majore differt.

7. ***Z. cleoniooides*** (Boiss. Diagn. Ser. I, 5, p. 14) annua erecta simplex vel basi ramosa, foliis præter pubem minutam seabridam longe ciliatis nervosis sessilibus a basi rotundatâ ovatis acutis, floralibus conformibus sed majoribus nervosissimis dense imbricatis longe attenuato-acuminatis saepe rubellis calyces occultantibus, spicâ elongatâ densiflorâ, calycis longe tubulosi brevissime hirtuli glanduloso-punctati dentibus lanceolatis brevibus conniventibus, corollæ roseæ calyce duplo longioris limbo amplio, antheris loculo sterili oblongo stipitato appendiculatis, staminodiis binis sterilibus brevibus ♂.

Hab. in collibus siccis ad basin montis Mesogis supra Sardes (Boiss!).

Planta 4-9-pollicaris, calyces 4½ lineas, corollæ 8-9 longæ. Affinis *Z. Taurica* differt foliis ovatis basi rotundatis, floralibus 4 lineas fere latis, calyce et corolla longioribus. Species elegantissima.

TR. IV. MONARDE.E.

PEROWSKIA (Kar. Bull. Mosq. 1841 p. 43. — Bge Pl. Lehm. p. 432 et Labiat. Persicæ p. 40. — Stocks in Boiss. Diagn. Ser. II, 4, p. 16).

Calyx tubuloso-campanulatus bilabiatus labio superiore integro vel tridenticulato, inferiore bidentato, fructifer auctus apertus. Corolla tubo subampliato subexerto intus annulato, limbi patentis subbilabiati labium superius quadrilobum lobis subæqualibus vel intermediis paulo minoribus et lateralibus reflexis, labium inferius integrum rotundatum. Stamina fertilia bina exserta divergentia, duo sterilia minima sub lobis labii superioris inserta. Antheræ biloculares loculis linearibus parallelis, connectivo minimo. Discus æqualis vel antice in glandulam tumens. Stylus bifidus lobis latis complanatus. Nuculae obovatae. — Herbæ perennes dichogamæ habitu *Lavandule*

sed characteribus alienæ, corolla *Ocymoidearum* sed antheræ bi-nec uniloculares. Affinitas in Ordine subincerta.

1. P. abrotanoides (Karel. Bull. Mosq. 1841, p. 15, tab. 4) caulis erectis elatis paniculatis sicut folia glabris vel indumento brevissimo floccoso-pruinoso secedenti obsitis, foliis petiolatis ambitu ovato-oblongis rarius incisis stepius in segmenta linearia decurrentia integra vel incisa obtusa bipinnatisectis, verticillastris sessilibus 6-2-floris distinctis in spicas longas nudas laxas dispositis, bracteis oblongis calyce brevioribus, calycis patule longe et plus minus copiose lanato-hispidi labio superiore subintegro ♀.

Hab. in Persiâ boreali-orientali (Buhse!), in ditione Khorassan prope Schahrud et Meschhed (Bge!) in littoribus Turcomaniæ (Karel!).

Bi-tripedalis, variat multum foliorum lacinis 1-2 lineas latis vel tenuissimis, amic. a Bge formas foliis tantum incisis observavit.

Ar. Geogr. Thetia.

2. P. scrophulariæfolia (Bge pl. Lehm. p. 433) caulis elatis superne paniculatis pube brevissimâ ramosâ secedenti canescens, foliis glabrinculis resinoso-punctatis petiolatis ovato-oblongis obtusis inaequaliter crenato-dentatis, verticillastris 6-2-floris laxiuseule et longe spicatis, bracteis minimis, calycis albo-villosi labio superiore obsoletissime tridenticulato ♀.

Hab. in rupestribus Turkestaniæ occidentalis in monte Karatau (Lehm!).

Hujus plantæ specimina incompleta tantum nota, an satis differt a *P. abrotanoides* quam indumento refert?

3. P. artemisioides (Boiss. Diagn. Ser. II, 4 p. 15) tota indumento stellato non secedente adpressissime cana, caulis erexit elatis crebre et stricte foliosis paniculatis, foliis petiolatis ambitu oblongis in lacinias oblongas vel lineares obtusas decurrentes bipinnatisectis, verticillastris 6-2-floris sessilibus distinctis in spicas longas graciles nudas laxas dispositis, bracteis lanceolato-linearibus minimis, calycis dense bombycino-lanati labio superiore tridenticulato ♀.

Hab. in Belutschia frequenter (Stocks!).

A *P. abrotanoides* differre videtur non foliorum divisione et laciniarum latitudine variabili sed indumento brevissimo stellato non secedenti. An potius *P. atriplicifolia* varietas?

4. P. atriplicifolia (Bth. in D.C. Prodr. XII, p. 261) tota indumento leproso squamulis stellatis constante adpresse et dense cana, caulis elatis simplicibns et stricte paniculatis, foliis breviter petiolatis oblongis vel lanceolatis repandis denticulatis vel incisis rugosis, verticillastris 6-2-floris remotis laxe spicatis, calycis hispidi labio superiore tridenticulato ♀.

Hab. in regno Cabulico inter Ghazni et Cabul (Griff!), in Belutschia superiore (Stocks!).

DORYSTÆCHAS (Boiss. et Heldr. in D.C. Prodr. XII, p. 264. —
Diagn. Ser. I, 12, p. 54).

Calyx breviter campanulatus bilabiatus labio superiore indiviso inferiore bipartito, fructifer valde auctus elliptico-cylindraceus ore clausus. Corollæ tubus tenuis breviter exsertus intus nudus, limbi patentis labium superius subbrevis retusum, inferius trilobum. Stamina bina antica vix exserta parallela subdistantia. antheræ biloculares loculis linearibus parallelis subcurvis apice connectivo vix perspicuo connexis. Staminodia nulla. Stylus apice breviter bifidus lobis subcomplanatis. Nuculæ oblongo-lineares acuminato-subrostratae. —Suffrutex facie *Salvie* vel *Meriandrae*. Genus ab eis connectivo punctiformi nec elongato distinctum, a *Perovskia* calyce fructifero clauso, staminodiis nullis, nuculis acuminatis et habitu diversissimum!

1. **D. hastata** (Boiss. et Heldr. loc. cit. p. 55) suffruticosa a basi ramosa, ramis vetustis erectis lignosis flexuosis nudis, novellis adpresso pubescentibus simplicibus, foliis breviter tomentellis superne griseis rugulosis subtus canescensibus lanceolato-hastatis obtusis et acutiusculis crenulatis, verticillastris numerosis sessilibus 10-12-floris foliis floralibus lanceolatis brevibus suffultis in spicas longas tennes cylindricas congestis. floribus minimis sessilibus, calycis albo-hirsuti valde glandulosi labio superiore sublongiore, inferioris lobis triangularibus acutis ♂.

Hab. in Lyciâ prope Elmalu (Pestal), in glareosis ad rivulos montis Climax prope Kourmalu in Pamphyliâ Lyciae finitimâ (Heldr.). Fl. Mai.

Suffrutex $\frac{1}{2}$ -4-pedalis, folia $1\frac{1}{2}$ - $2\frac{1}{2}$ -pollicaria basi 8-15 lineas lata, spicæ 2-4-pollicis longæ tenues, calyx florifer sesquilineam, fructifer 4-5 lineas longus labiis conniventibus clausus, corolla $2\frac{1}{2}$ lineas longa & siccō carnea.

SALVIA (L. Gen. 39).

Calyx campanulatus vel ovoideus rarius tubulosus plus minusve distinete bilabiatus labio postico breviter tridentato rarius integro, antico profundius bifido. Corollæ tubus inclusus vel exsertus intus nudus vel piloso-annulatus vel squamulâ penicillatâ auctus. limbus bilabiatus labio superiore concavo recto vel falcato, inferiore patente trifido lobo medio latiore. Stamina perfecta bina filamentis brevibus cum connectivo elongato transverse articulatis, connectivum lineare postice ascendens et loculum fertilem ferens, antice porrectum vel deflexum loculum cassum rarius polliniferum ferens vel nudum. Staminodia postica bina minima vel nulla. Stylus bifidus. Nuculæ ovoideæ laeves interdum subtriquetræ. — Suffrutices vel herbæ. Genus vastissimum.

Specierum Orientalium coordinatio.

SECT. I. EUSPHACE Bth. — Corollæ tubus intus piloso-annulatus, labium superius subrectum, antherarum connectiva antice porrecta loculum sæpius cassum ferentia, calycis fructiferi vix aucti dentes acuti.

• Simplicifoliæ.

S. divaricata, Aucheri, grandiflora, Libanotica, Cabulica, Kurdica, macrochlamys.

• • Pinnatæ.

+ Fruticosæ.

1. Folia floralia (infimis exceptis) floribus multo minora.

S. triloba, Cypria, Cedronella, roseifolia, suffruticosa, spiræifolia, potentillæfolia, Pisidica, Tchihatcheffii, Wiedemannii.

2. Folia floralia floribus subæquilonga.

S. cespitosa, Russeggeri.

++ Caules (ex rhizomate sæpe indurato) herbacei.

1. Folia floralia (infimis exceptis) parva flores non æquantia.

S. ringens, scabiosæfolia, pinnata, recognita, rubifolia, Cataonica, pilifera.

2. Folia floralia magna calyces superantia.

S. bracteata, trichoclada.

SECT. II. HYMENOSPHACE Bth. (ex parte speciebus Capensisibus exclusis). — Corollæ tubus intus piloso-annulatus, labium superius subrectum, antherarum connectiva antice porrecta loculum sæpius cassum ferentia, calyx fructifer auctus labiis membranaceo-dilatatis venosis superiore integro, repando, vel brevissime tridentato. Species binæ priores transitum ad *Eusphacem* præbent.

S. Smyrnæa, Cadmica, pomifera, calycina, Euphratica, Haussknechtii, Bodeana, dracocephaloïdes, Hydrangea, Sheilii, Szovitsiana, Rascheyana, acetabulosa, cryptantha, Pinardi.

SECT. III. DRYMOSPHACE Bth. — Corollæ tubus intus piloso-annulatus, labiuni superius falcatum compressum, antherarum connectiva antice porrecta et loculum cassum ferentia. Calyx fructifer vix auctus dentibus acutis.

S. glutinosa, Forskahlei.

SECT. IV. *ÆTHIOPIS* Bth. — Corollæ tubus intus exannulatus nudus vel squamulâ penicillatâ auctus, labium superius plus minusve falcatum compressum, antherarum connectiva antice deflexa abrupte dilatata et extremitate callosâ connexa. Calyx campanulatus labio superiore convexo tridentato. Species ultima transitum ad *Plethiosphacem* præbet.

§ 1. **Homalosphaceæ.** — Sect. *Homalosphace* Bge Lab. pag. 49. — Corollæ tubus sensim in faucem ampliatus intus squamulâ destitutus.

S. Syriaca, rhytidia, compressa, eriophora, Montbretii, canescens, hypochionea, phlyctidea, lericifolia, spinosa, Reuteriana, Persepolitana, Palwestina, macrosiphon.

§ 2. **Gongrosphaceæ.** — Sect. *Gongrosphace* Bge Lab. pag. 44. — Corollæ tubus abrupte in faucem saccatam intus squamulâ penicillatâ auctam ampliatus.

S. graveolens, Scarea, Æthiopis, aristata, ceratophylla, chionantha, argentea, Tmolea, verbascifolia, Atropatana, Urmensis, xanthocheila, frigida, modesta, candidissima, crassifolia, cyanescens, oligophylla, hypoleica, limbata, Beckeri, Sahendica, Chorassanica, brachycalyx.

SECT. V. **PLETHIOSPHACE** Bth. — Corollæ tubus intus exannulatus nudus vel squamulâ obsoletâ instructus, labium superius erectum vel falcato-compressum. Antherarum connectiva antice deflexa et abrupte dilatata extremitate callosâ connata. Calyx ovatus labio superiore concavo-bisulcate dentibus ternis brevissimis conniventibus.

S. Austriaca, staminea, pratensis, Hierosolymitana, virgata, viscosa, sylvestris, nutans, Verbenaca, controversa.

SECT. VI. *HORMINUM* Bth. — Corollæ (parvæ) tubus intus exannulatus nudus, labium superius erectum concavum. Connectiva antice deflexa abrupte dilatata extremitate callosâ connexa. Calyx tubulosus labio superiore truncato remote bidentato inferiore tridentato.

S. viridis, Horminum.

SECT. VII. **EREMOSPACHE** Bge Lab. p. 51. — (*Notiosphace* Bth. ex parte). — Corollæ (parvæ) tubus intus exannulatus nudus, labium superius rectum. Connectiva antice distantia et deflexa loculum cassum dilatatum ferentia. Calyx ovatus vel tubuloso-campanulatus labio superiore tridentato (interdum ut in *Plethiosphace* dorso bisulcato), inferiore tridentato.

S. Egyptiaca, deserti, santolinifolia, eremophila, trichocalycina, Tebesana, macilenta.

SECT. VIII HEMISPHACE Bth. — Corollæ tubus intus piloso-annulatus labium superius erecto-patens fornicatum. Connectiva parte anticâ breviter subulata.

S. verticillata, Peloponnesiaca, napifolia, Judaica, Russelii.

SECT. I. EUSPHACE.

* Simplicifoliæ.

1. **S. divaricata** (Montb. et Auch. Ann. Sc. Nat. 1836, 2, pag. 37) basi suffruticosa, caulis adpresso et breviter pubescentibus in paniculam nudam abenntibus, foliis oblongis minute crenatis rugosis supra glabris subtus pubescentibus canescens basi attenuatis petiolatis petiolis interdum ciliatis, paniculæ glabrae ramis divaricatis, bracteis nullis, verticillastris 5-2-floris, pedicellis erecto-patulis rigidis calyce triplo longioribus, calycis sparsim et breviter glanduloso-hirti viscidii labiis æquilongis angulo recto distantibus superiore obsolete tridenticulato inferiore bidentato, corollæ violacea calyce duplo longiore 5.

Hab. in Cappadociâ orientali (Auch. 1528!).

Folia 2-2 $\frac{1}{2}$ -pollicaria 6-8 linea lata, panicula pedalis pedicellis 1-2-pollices longis, calyx 7 linea, corolla 15 longi.

2. **S. Aucheri** (Bth. Ann. Sc. Nat. 1836, 2, p. 38) basi indurata, caulis a basi procumbenti radicanti erectis pubescentibus in paniculam nudam abeuntibus, foliis oblongis minute crenulatis tenuiter rugosis supra glabriusculis vel minute puberulis pallidis subtus canescens vel canis basi rotundatis petiolatis, paniculæ ramis elongatis erecto-patulis, verticillastris 4-2-floris, bracteis squamæformibus deciduis, floribus subsessilibus, calyce glanduloso-hirtulo dentibus 5 lanceolatis subæqualibus tubo quadruplo brevioribus, corollæ violacea tubo triplo longiore 5.

Hab. in regione montanâ Tauri Cilicie prope Gülek 2500'-4000' (Auch. 1529!), Ky. 435!, Heldr! Bal!.

β. canescens. Boiss. et Heldr. — Folia minora utrinque cana, flores minores, calycis dentes abbreviati late triangulares.

Hab. ad Boumarbachi et in valle Djoksu inter Karaman et Ermenek I-aureæ (Heldr!). Planta 2-3-pedalis, folia in typo 2-3-pollicaria pollicem lata, paniculæ 1-1 $\frac{1}{2}$ -pedales, corolla in typo 12-13 linea, in varietate 9 linea longa.

3. **S. grandiflora** (Ettling Salv. N° 2) basi indurata, caulis ascendenti-erectis elatis pubescenti-tomentellis, foliis petiolatis a

basi rotundatâ vel cordatâ ovatis et oblongis minute crenulatis rugulosis pubescentibus subtus pallidioribus, superioribus diminutis acuminateis, floribus racemosis rarius paniculatis, verticillastris 10-6-floris distinctis, bracteis parvis submembranaceis, calycis pallidi viscidulo-hirti subbilabiati dentibus triangulari-lanceolatis acutis intermedio interdum minore, labii inferioris profundius fisis, corollâ cæruleâ calyce subtriplo longiore 5. Ic. Jacq. Eclog. tab. 36.

Hab. in collibus regionis inferioris Anatoliae occid. in Bithynia (Auch-Thirke!), Phrygiâ ad Ouchak (Bal!), Lydiâ ad Smyrnam (Boiss! Bal!), Lyciâ (Bourg. 228!). Galatiâ et Paphlagoniâ (Wied!), Ciliciâ (Bal!). Syriâ boreali (Boiss!), Libano (Ehr! Bl!), Cypro (Ky!), Cataoniae monte Berydagh 7000' (Hausk!), Tauriâ (Rehm!).

β? rotundifolia. — Folia obtusiora ovato-orbiculata, dens superior calycis lateralibus manifeste brevior, corollæ tubus paulo longior. — *S. rotundifolia* Visian. Illustr. pag. 6. tab. I.

Hab. in monte Idâ Bithyniae (Parolini). Non vidi et huc ex auctoritate cl. Webb in Giorn. Ital. bot. Ann. I, pars 2, p. 51 adduco.

Affinis *S. officinali*, folia majora minus cana, ovato-oblonga nec oblongo-lanceolata, basi rotundata vel cordata nec attenuata, calycis dentes acuti quidem sed non subulato-acuminati; planta major foliis majoribus. — *S. Cretica* L. Sp. p. 33 est forma hortensis submonstrosa foliis oblongo-linearibus *S. officinalis* vel *S. lavandulæfoliæ*, in Cretâ minime obvia.

4. ***S. Libanotica*** (Boiss. et Gaill. Diagn. Ser. II, 4, pag. 16) fruticosa ramosa tota indumento criso adpreso viscidulo grisea, ramis floriferis erectis simplicibus vel parce ramulosis, foliis parvis petiolatis a basi rotundatâ vel truncatâ ovato-oblongis obtusis rugulosis miuite crenulatis, verticillastris 6-4-floris distinctis longe racemosis, foliis floralibus breviter lanceolatis flores non æquantibus, pedicellis calyce brevioribus, calycis patule hispida valde glandulosi vix bilabiati dentibus triangularibus acutis tubo quadruplo brevioribus, corollâ calyce vix duplo longiore 5.

Hab. in regione inferiore Libani inter Hallalie et Baramie supra Sidonem (Gaill!), in monte Torbol supra Tripoli (Bl!), Cypro in rupestribus Buffavento 1000' (Ky. 483!). Folia majora vix pollicaria, racemi semipedales, calyx 4 lineas, corolla 7-8 lineas longa.

An varietas *S. trilobæ?*, folia minora semper simplicia, corollæ minores pulchre cæruleæ albo-punctatae.

5. ***S. Cabulica*** (Bth. in D.C. Prodr. XII, p. 268) fruticosa ramosissima molliter villosa canescens, ramis novellis brevibus rigidulis subsimplicibus, foliis parvis cordatis ovato-orbiculatis crenatis rugulosis petiolatis, floralibus parvis oblongo-lanceolatis, verticillastris 2-3 terminalibus 4-2-floris approximatis, calyce pedicellato campanulato ad tertiam partem bilabiato labio superiore breviter tridentato inferiore bitido dentibus omnibus mucronatis, corollâ calyce 2 1/2-plo longiore 5.

Hab. in regno Cabulico ad Bharowul et Pusehut (Griff!), in Belutschia superiore (Stocks 881!).

Rami vetusti cortice albâ vestiti, folia majora 6-7 lineas longa, racemi pollicem circiter longi, calyces 5 lineas longi fructiferi vix aucti, corolla 11 lineas longa e sicco cærulea. Facies *S. nivea* Capensis.

6. S. Kurdica (Boiss. et Hoh. in p. c. Prodr. XII, p. 268) basi sulfrutescens pallide virens papillari-pubescentis viscida, ramis erectis in panienlam foliatam abenutibus, foliis longe petiolatis magnis late cordato-ovatis crenatis rugulosis, verticillastris bisforis in racemos breves laxos axillares folio breviores dispositis, pedicellis calyce brevioribus bracteâ lanceolatâ subæquilonigâ suffultis, calycis molliter hirsuti viscidii divaricatum ad tertiam partem bilabiati labio superiore sublongiore minute tridenticulato, inferiore ad medium in dentes late lanceolatos fiso, corollâ calyce subduplo longiore ♀.

Hab. ad rupes prope mænia urbis Amadieh Kurdistanæ Assyriacæ (Ky. 547!).

Ex frustulis imperfectis descripta, folia ea *S. macrochlamydis* referentia laminâ triplicari 2-pollices latâ, inflorescentia ob racemos breves axillares peculiaris, calyx fructifer 9-10 lineas longus, corolla 10 lin. longa intus annulata.

7. S. macrochlamys (Boiss. et Ky. pl. exs. 1859) patule papillari-hirsuta viscidula pallide virens, caulibus erectis crassis elongatis crebre foliosis, foliis longe et stricte foliosis amplis ovatis valde cordatis obtusis crenulatis non rugosis utrinque parce papillari-hirtis concoloribus, panienlâ terminali elongatâ strictâ patule et longe hirsutâ viscosâ, foliis floralibus amplis sessilibus membranaceis albidis late oblongo-lanceolatis, verticillastris bisforis secus paniculæ ramos breves approximatis, bracteis membranaceis albidis late oblongo-lanceolatis acuminatis cymbiformibus calyces superantibus, calycis membranacei hispidi ad medium bilabiati labio superiore tridentato-inferiore bifido dentibus lanceolato-subulatis, corollâ calyce vix sesquiloniore ♀.

Hab. in schistosis alpium prope Musch Armeniæ 5000' (Ky. 508!).

Specimina valde imperfecta ramis sterilibus foliatis pedem et amplius longis et paniculis seorsim lectis quinquepedalibus constantia. Foliorum lamina 2-4-pollices longa 2-2 $\frac{1}{2}$ lata, bracteæ 1 $\frac{1}{2}$ -pollicares, calyx vix pollicaris. Bracteis magnis persistentibus affinis *S. bracteata*, hic artificialiter ob folia indivisa collocata.

** Pinnatæ.

8. S. triloba (L. fil. Suppl. p. 88) fruticosa adpresso albo-tomentosa, foliis petiolatis rugulosis crenulatis subtus intensius canis a basi truncatâ vel subcordatâ oblongis vel lanceolatis obtusis saepius ad laminæ basin segmentis binis parvis ovatis vel ellipticis auctis, superioribus diminutis triangulari-auriculatis, floralibus parvis acuminatis, floribus racemosis vel saepius paniculatis paniculæ viscosissimæ ramis brevibus, verticillastris 6-4-floris, bracteis parvis membranaceis deciduis, calycis hirti viscosi subbilabiati dentibus triangularibus acutis labii superioris majoribus, corollâ violacea calyce

triplo longiore ȝ. Ic. Fl. Græc. tab. 17. — *S. auriculata* et *S. fruticosa* Mill. Dict. — *S. Sipylea* Lam. Ill. I, 68 — *S. incarnata* Ettl. Salv. № 15 (saltem ex parte et ex specimine herb. Vahl !).

Hab. in collibus regionis calidæ totius Græciae et Archipelagi (Silth! Spr! Boiss! Orph. Fl. exs. 548!). Thraciæ et Bithyniæ (Griseb.), Anatoliæ occidentalis! Cretæ (Rauh!), Syriæ littoralis (Bl! Gaill!), Palestinæ (Auch. 1541! Boiss! Ky. 448!), Cypri (Ky!).

Frutex 3-5-pedalis, folia 1½-2-pollicaria, panicula stricta semipedalis, calyx 4 lineas longus, corolla fere pollicaris.

Ar. Geogr. Sicilia, Italia meridionalis.

9. S. Cypria (Ky. Cypr. p. 266) fruticosa adpresso canescens ramis erectis opposite ramulosis, foliis parvis petiolatis minute rugulosis crenulatis obtusis a basi subcuneatâ truncatâ vel subcordatâ ovatis vel oblongis interdum (ex cl. Ky) ad laminæ basin lobulis binis auctis, racemis glandulosissimis, foliis floralibus parvis obovato-cuneatis, bracteis squamiformibus membranaceis deciduis, calycis papillari-glandulosi vix bilabiati dentibus breviter ovato-triangulibus obtusis, corollæ calyce triplo longioris violaceæ fauce flavomaculatâ ȝ.

Hab. in valle montanâ versus Demitlen prope Prodromo (Ky. 724!). Fl. Mai.

Frutex 3-5-pedalis, foliorum lamina 6-9 lineas longa, racemi 3-4-pollicares laxiusculi verticillæ paucifloris, calyx 3 lineas, corolla 9 longi.

An sit varietas *S. trilobæ* ulterius inquirendum, folia multo minora breviora, flores fere duplo minores, calycis dentes multo breviores obtusi.

10. S. Cedronella (Boiss. Diagn. Ser. I, 5, p. 3) fruticosa ramosissima dumosa ramis rigidis glaberrimis simplicibus foliosis; foliis glaucis brevissime petiolatis carnosulis glabrescentibus brevibus pinnatisectis segmentis 1-2-jugis confertis cuneato-oblongis obtuse dentatis terminali majore, verticillastris 4-2-floris distantibus racemosis foliis floralibus præter infima secta ovatis acuminatis parvis suffultis, pedicellis calyce brevioribus, calycis glabri vel breviter papilloso labio superiore tridentinato inferiore bifido dentibus omnibus acuminato-spinulosis, corollâ flavâ calyce duplo longiore ȝ.

Hab. in collibus argillosis Cariæ ad orientem urbis Denisleh (Boiss!), Phrygiæ ad Ezilair Kivi prope Ouchak (Bal!).

Frutex bipedalis, foliorum petiolus cum rachide 3-6 lineas tantum longus, segmentum superius 6-12 lineas longum, calyx 5 lineas, corolla 9-12 longa.

11. S. rosæfolia (Smith Icon. I, p. 5 tab. 5) basi suffrutescens breviter et retrorsum pubescenti-canescens prætereaque ad ramos juniores et petiolos sœpe setulosa, ramis erectis foliosis simplicibus vel parce ramosis, foliis longe et tenuiter petiolatis non rugosis pinnatisectis segmentis 2-3-jugis ellipticis vel oblongis lateralibus sœpe linearibus petiolulatis vel decurrentibus terminali longiore omnibus obtusis vel acutiusculis remote dentatis, floribus ovato-lanceolatis

acutis inferioribus calyces æquantibus, verticillastris 5-2-floris distantibus breviter racemosis, pedicellis calyce brevioribus, calycis adpresso pubescentis colorati bilabiati labio superiore sublongiore dentibus ternis brevibus acutis, inferiore in dentes ovato-lanceolatos acuminiatos fisso, corollâ purpureâ calyce subduplo longiore 5.

Hab. in Armeniâ Turcicâ (Tourn.), circa Erzerum (Calvi), Ispir (C. Koch !), Baibut (Bourg !).

Rami cum racemo 7-10-pollicares, verticillastra 3-5, calyx fere 5 lineas, corolla 9 lineas longa.

12. *S. suffruticosa* (Montb. et Auch. Ann. Sc. Nat. 1836, 2, pag. 391) suffruticosa basi ramosa, ramis herbaceis erectis virgatis gracilibus glaberrimis albidis foliosis, foliis petiolo stricto albido hirtulo suffultis pinnatisectis segmentis breviter pubescentibus interdum canescens 1-2 jugis oblongo-lanceolatis obtusiusculis sub-integris terminali 2-3-plo majore basi attenuato, foliis summis parvis in lacinias paucas lineares breves fisis, verticillastris 6-4-floris dissitis racemosis foliis floralibus ovatis acuminatis calyce brevioribus suffultis, pedicellis brevibus, calycis papillosi saepe glandulosi labio superiore breviter tridentato inferiore bifido dentibus mucrone lutescente subpungenti subincurvo terminatis, corollâ luteâ calyce duplo longiore 5. *S. Vanensis* Boiss. et Noë Diagn. Ser. II, 4 p. 17.

Hab. in monte Akdaghan Cataoniæ (Auch. 1518!), in faucibus Karakoi Cappadociæ ad Argæum 4500' (Ky. 185!), in Kurdistaniâ (Brant), inter Van et Ischri (Noë 153!), circa Ancyram et Kadikiöi (Wied!).

Rami 4-1½-pedales, calyx 4-5 lineas, corolla 7-8 lineas longa. Facie et foliis valde affinis *S. roseæfoliæ* a qua florum colore, calycis dentibus magis pungentibus differre videtur.

13. *S. spireæfolia* (Boiss. et Hoh. Diagn. Ser. I, 5 p. 5) suffruticosa basi ramosa, ramis erectis virgatis subsimplicibus glabris foliosis, foliis petiolatis pinnatisectis 2-3-jugis, segmentis supra parce subtus crebre papillari-hirtis lineatis acute dentatis lateralibus elliptico-lanceolatis acutis terminali majore elliptico-oblongo, foliis summis in lacinias lineares partitis, verticillastris 6-4-floris dissitis racemosis foliis floralibus ovatis acuminatis calyce brevioribus suffultis, calycis breviter pedunculati adpresso pubescentis labio superiore breviter tridentato inferiore bifido dentibus acutis vel mucronatis, corollâ dilute cœruleâ calyce 2-2 ½-plo longiore 5. *S. suffruticosa* ex parte Bth. in p.c. Prodr.

Hab. in Syria circa Aleppo (Ky. 256!), in agris argillosis Mesopotamiae inter Tcharmelik et Surug. inter Bir et Orfa et in deserto Dara (Haussk!).

Ab affinibus *S. roseæfoliâ* et *suffruticosâ* foliorum segmentis majoribus acute dentatis subtus lineatis acutis, florum colore, etc. distincta.

14. *S. potentillæfolia* (Boiss. et Heldr. p.c. Prodr. XII, p. 270) suffruticosa adpresso canescens, ramis virgatis simplicibus

vel parce ramulosis, foliis parvis petiolatis triseptis segmentis subsessilibus oblongo-cuneatis superne denticulatis intermedio submajore, verticillastris 4-2-floris distantibus racemosis foliis floralibus lanceolatis minutis inferioribus 2-3-partitis suffultis, pedicellis calyce brévioribus, calycis glanduloso-hirsuti ad tertiam partem bilabiati labio superiore tridenticulato dente intermedio minimo lateralibus mucronatis, labio inferiore in dentes lanceolatos longe acuminatos fisso, corollâ luteâ, cærulecente, vel bicolore calyce $2\frac{1}{2}$ -plo longiore 5.

Hab. in collibus rupestribus Lyciae circâ Elmalu (Bourg. 232-234!), in valle speluncarum supra Ermenek Ciliciæ Tracheæ (Heldr!).

Rami novelli pedales, foliorum segmenta 4-6 lineas longa, corollæ magnæ 1- $1\frac{1}{2}$ -pollicares. Facies *S. suffruticosa* et *roseofolia*, sed folia ternata nec pinnata, segmento terminali sessili nec petiolulata.

S. microphylla. — Foliola minora. Huc ob ramos erectos et verticillastra dissita spectare videtur.

Hab. in rupestribus ad Duden prope Elmalu Syriæ (Bourg. exs 235! sub *S. Pisidicâ*).

15. ***S. Pisidica*** (Boiss. et Heldr. in D.C. Prodr. XII, p. 269) suffruticosa minute pruinosa cauescens prætereaque ad ramos et petiolos ciliata, ramis procumbentibus saepe radicantibus novellis erectis pumilis simplicibus crebre foliosis, foliis triseptis rarissime bijugis segmentis parvis sessilibus lateralibus lineari-spathulatis terminali oblongo-cuneato, floralibus lanceolatis pedicellos vix superantibus, verticillastris 6-4-floris in racemos laxiusculos approximatis, calycis glandulosi et patule hirsuti fere ad medium bilabiati labio superiore truncato bidentato cum dente intermedio fere obsoleto, inferiore in dentes lanceolato-acuminatos bisido, corollâ violaceâ calyce $2\frac{1}{2}$ -plo longiore 5.

Hab. in collibus tophaceis Pisidiæ prope Baldur (Heldr!), Phrygiæ ad Yaparlar Kieui prope Ouchak (Bal. 1191!).

Affinis *S. potentillefolia* a qua præter folia angustiora et saepius minora differt ramis primariis prostratis radicantibus, verticillastrisque approximatis, an tamen sit hujus varietas in locis natalibus investigandum.

16. ***S. Tchihatcheffii***, suffruticosa ramosa primariis procumbentibus ramulos steriles breves è fertiles sublongiores omnes crebre foliatos edentibus, foliis petiolo brevissimo ciliato suffultis glabris in lacinias ternas anguste lineares margine revolutas triseptis, floralibus ovatis acuminatis calycem aequantibus, verticillastris in spicam densam $1\frac{1}{2}$ -2 pollicarem confertis, floribus parvis, calycis pruinosi ciliati ad medium bilabiati labio superiore in dentes ternos triangulari-lanceolatos apice fisso, inferioris dentibus binis lanceolatis omnibus acutis, corollâ calyce vix duplo longiore 5. *Dracocephalum Tchihatcheffii* F. et M. Ann. Sc. Nat. 1854, I, p. 34!

Hab. in vallibus herbo-is sylvaticis montis Kuredagh Galatiæ 3600' (Tchih!).

Rami primarii semipedales, ramuli floriferi erecti 2-3-pollicares, folia 6-8 lineas longa lacinis tenuibus acutis 4-5 lineas longis, calyx 4-5 lineas, corolla 7-8 longa. Nihil *Dracocephali* habet sed *Salvia* genuina *S. cespitosa* affinis!, ab ea et a finibus foliorum lacinis angustis floribusque parvis distincta.

17. S. Wiedemannii. suffrutescens parce et crispule papillosa pallida, ramis abbreviatis erectis rigidulis simplicibus, foliis depresso-punctatis petiolo brevissimo complanato longe ciliato suffultis brevibus in lacinias 3-5 anguste linearis-oblongas integras elongatas pinnatisectis, floralibus inferioribus conformibus flores superantibus, superioribus indivisis oblongis parvis, verticillastris 6-4-floris subdistantibus breviter racemosis, floribus parvis breviter pedunculatis, calyeis costati papillari-ciliati et glandulosis sessilibus adspersi labio superiore breviter tridentato dentibus triangularibus intermedio latiore, labio inferiore in dentes lanceolatos acutos profunde fisco, corollâ calyce sesquilociore 5.

Hab. in Anatoliâ prope Kadikiöi (Wied!).

Rami floriferi cum racemo semipedales, foliorum laciniae 6-12 lineas longe 1-1½ late, terminalis ceteris longior. Calyx 4 lineas longus, corolla vix 6 lineas longa. Affinis *S. Tchihatcheffii* ad quam forsitan confundit sed utriusque specimen unicum tantum vidi, differre videtur foliis crispulis et punctatis lacinis plurijugis et latioribus, racemo laxo, calycis dentibus superioribus brevissimis.

18. S. cæspitosa (Month. et Auch. Ann. Sc. Nat. 1836, p. 39) suffruticosa ramosissima cespitosa ramis punnulis ad apicem usque dense foliosis, foliis pubescentibus vel tomentellis petiolis ciliatis basi dilatatis suffultis ambitu angustis pinnatisectis segmentis 3-4-jugis oblongis crenato-dentatis lateralibus minimis saepe decurrentibus terminali submajore, floribus paucis in racenum brevissimum terminalem e foliis caulinis vix exsertum confertis, foliis floralibus membranaceis reticulatis ovalis calyces superantibus inferioribus apice in lobulos herbaceos fissis ceteris indivisis herbaceo-acuminatis, calycis patule hirsuti ad tertiam partem bilabiati labio superiore tridenticulato dente intermedio submajore, inferiore in lentes ovatos fisco omnibus mucronatis, corollâ violaceâ calyce triplo magiore 5. *S. peettinifolia* F. et M. Ann. Sc. Nat. 1854, I, p. 33! — *S. pa-chystachys* Trautv. Pl. Sp. nov. № 4!

Hab. in alpinis Anatoliæ, mons Beréketdaghi Lycie (Ball!), mons Akdaghi (Auch. 1519!) et Berydaghi Cataonie 7-8000' (Haussk!), Armenia Rossica (Radde!).

Rami 3-4-pollicares, folia cum petiolo 1-2-pollicaria segmentis 3-4 lineis longis. Flores subcapitati, corolla 14-16 lineas longa.

19. S. Russeggeri (Fenzl Del. Sem. Vindob 1851, — Tchih. As. Min. 2, p. 138) fruticosa ramosissima ramis novellis erectis simplicibus crebre foliatis, foliis petiolis strictis longe ciliatis inferne latiusculis suffultis pinnatisectis unijugis rarius bijugis segmentis

viridibus glanduloso-asperulis elevatim lineatis crenulatis lateribus anguste linearibus terminali cuneato ovato vel oblongo multo majore, verticillastris 6-4-floris in capitulum ovatum vel oblongum densum approximatis, foliis floralibus ovatis et oblongis acuminatis calyces superantibus et aequantibus, calycis glanduloso-asperuli ciliatiali ad quartam partem bilabiati labio superiore breviter et acute tridentato dente intermedio pargo, inferioris dentibus lanceolatis acuminatis, corollâ calyce duplo longiore ♂. *S. Kotschyi* Bth. MSS. non Boiss. — *S. Benthamiana* Boiss. in D.C. Prodr. 12, p. 209 non Field. et Gardn. — *S. Heldreichiana* Boiss. in D.C. Prodr. XIII Errata.

Hab. in collibus ad radices Tauri prope Gülek Boghas (Ky. 437! Bal!), prope Karaman in Isauriâ (Heldr!).

Suffrutex 1-2-pedalis, rami novelli 6-8-pollicares, foliorum segmentum superius $\frac{1}{2}$ -1-pollicare saepe complicatum, racemus 2-4-pollicaris densus quâ notâ a *S. suffruticosâ, rosæfoliâ* etc. differt, corolla 14 lineas longa.

20. ***S. ringens*** (Sibth. et Sm. Fl. Græc. I. p. 14, tab. 18), rhizomatis indurati ramis prostratis saepe radicantibus, caulis foliferis erectis glabris basi crebre foliosis superne subnudis in racemum vel in paniculam parce ramosam abeuntibus, foliis pallide virentibus rugulosis inferne præserlim pubescenti-viscidulis irregulariter et saepe interrupte pinnatisectis segmentis 3-7 lateralibus ovato-rotundatis vel ellipticis saepe subpetiolulatis terminali multo majore ovato vel oblongo saepe acutiusculo, verticillastris 6-2-floris remotis, bracteis parvis membranaceis deciduis, calycis in pedicello brevi nutantis glanduloso-hirsuti subincurvi labio superiore breviter tridentato inferiore bifido, corollæ magnæ cœruleæ calyce quadruplo longioris tubo decurvo limbo ampliato recurvo-ascendente ♀.

Hab. in regione montanâ et sylvatica Græciae, Achaia ad Megaspilæon (Sibth.), Kyllene regio interior 3500'-4000' (Heldr! Orph. Fl. exs. 100!), Parnes Attice, Parnassus, Olympos Thessalus (Sprun! Heldr!), Macedonia inter Istib et Kaßadar (Griseb.), Thracia in monte Rhodope (Janka!).

Pedalis et bipedalis, folia segmentorum numero et formâ valde variantia, corolla saepe sesquipollicaris. Species pulchra.

21. ***S. scabiosæfolia*** (Lam. Journ. Hist. Nat. N° 14, p. 44, tab. 27) pluricaulis breviter hirta et patule hispidula virens, caulis ascendentibus vix erectis simplicibus crebre foliosis, foliis pinnatisectis plurijugis segmentis indivisis vel iterum sectis integris oblongo vel lineari-lanceolatis basi et apice attenuatis acutis terminali saepe submajore, verticillastris 10-6-floris in racemum simplicem rarius basi 2 ramulis auctum approximatis, foliis floralibus ovatis acuminatis inferioribus calyces aequantibus, calycis ample campanulati membranacei hirti ultra tertiam partem bilabiati labio superiore breviter tridentato inferiore bifido dentibus omnibus mucronatis, corollâ violacea calyce plus duplo longiore ♀. Ic. Jacq. Eclog. I,

tab. 3. *S. pinifolia* Pall. — *S. Taurica* Habl. — *S. Hablitziana* Willd. in Schrad. Journ. I, 289, tab. 2.

Hab. in collibus cretaceis Tauriæ circâ Bachtchisarai et Karassubazar (Stev! Rehm!).

β. latifolia. — Foliola lateralia pauciora (2-4) omnia latiora breviora. — *S. vulnerariaefolia* Willd. Sp. I, p. 149 et herb!

Hab. in Oriente (Willd.) sed hanc formam tantum cultam vidi.

22. **S. pinnata** (L. Sp. p. 39) tota patule glanduloso-hirsuta valde viscida pallide virens, caulinus ascendentibus elatis simplicibus vel parce ramosis totâ longitudine foliosis, foliis petiolatis interrupte pinnatisectis segmentis utrinque 3-4 petiolulatis saepius alternis ovatis vel oblongis inciso-crenatis obtusis terminali saepius majore oblongo, foliis floralibus parvis membranaceis ovatis cito deciduis, verticillastris 6-4-floris remotis longe et laxe racemosis, pedicellis calyci aequilongis vel brevioribus, calycis nigricantis piloso-lanati urceolati truncati labiis fere obsoletis superiore tri-inferiore bidentato dentibus latis vix prominentibus mucronulatis, corollæ purpureæ calyce 2 1/2-plo longioris tubo recto lato ♀.

Hab. in cultis derelictis, ad vias, Bithynia prope Brussa (Pichl!), prope Smyrnam (Bal. 315!), Cilicia littoralis (Bal!), Syria borealis inter Aintab et Nisib (Haussk!), Palestina (Boiss! Ky!).

Caules saepe bipedales, segmenta foliorum magnitudine valde variantia, corollæ pollicem et amplius longæ.

23. **S. recognita** (F. et M. Ann. Sc. Nat. 1854, 2 p. 33) caulinus floriferis e rhizomate indurato erectis proceris inferne hispidis foliosis superne in racemum elongatum vel paniculam pauciramosam longe glanduloso-villosam abeuntibus, foliis pallide virentibus præter pubem brevem subtus ad nervos papillari-hirsutis rugosis pinnatisectis segmentis lateralibus 2-4 oblongis vel ellipticis parvis terminali magno ovato vel oblongo interdum cordato omnibus crenulatis, foliis superioribus subnullis vel binis laciinis parvis lanceolatis, floralibus parvis ovatis deciduis, verticillastris 6-4-floris distautibus, floribus subsessilibus, calycis glanduloso-hispidissimi labiis ovatis superiore minute tridentato inferiore in dentes binos ovatos acates fisso, corollæ roseæ calyce quadruplo longioris tubo subexerto fauce inflatâ sursum recurvâ ♀. *S. Argata* Boiss. et Bal. MSS. — *S. orgyalis* Fenzl in Tchih. As. Min. II, p. 136. — *S. incarnata* Ettl. N° 45 ex parte (Specimina culta in variis herbariis!).

Hab. in Cappadociæ monte Alidagh prope Cæsaream (Tchih! Bal!), valie Agatsch Kisse montis Bulghardagh Cilicie et in ditione Kassan Oghlu 4500' (Ky. 140!).

Planta 4-6-pedalis, caules florales versus medium glabri glauco-pruinosi, segmentum superius foliorum 2-3-pollicare, racemi interdum pedales, corolla 15-18 lineas longa. Huc spectant specimina herbariorum (semper culta) *S.*

incarnatae Ettl., sed hæc species ex specimine archetypo herbarii Vahliani est *S. triloba*. Icones Morisonii et Phukenetii ad *S. incarnatam* citatae prior ad *S. pinnatam* posterior ad *S. bracteatum* referenda videtur.

24. **S. rubifolia** (Boiss. Diagn. Ser. I, 12 p. 56) viridis tota patule papilloso-hispida viscida, caulis a basi indurata procumbenti erectis elongatis simplicibus, foliis petiolatis triseptis rarius bijugis membranaceis, segmento terminali duplo majori cordato-ovato lateralibus oblongis basi obliquâ saepe petiolulatis omnibus obtusis crenatis, verticillastris 6-4-floris distantibus racemosis, foliis floralibus parvis triangulari-lanceolatis acuminatis calyce multo brevioribus, pedicellis brevibus, calycis patule-hispidi et glandulosi ad quartam partem bilabiati dentibus a basi triangulari subulato-aristatis subpungentibus, corollâ cœruleo-roseo duplo longiore ♀. *S. incarnata* β? *Libanotica* Bth in p.c. Prodri. 12, p. 267.

Hab. in rupestribus umbrosis Antilibani circa Rascheya et supra Zebdani (Boiss!), in monte Garbi 4800' (Ky. 21!), in valle Ouadi Benka inter Dimar et Jantha (Gail!), in Libano (Aueh. 4543!).

Folia *Ononidis rotundifoliae* sed segmenta majora, superius 1-1½-pollicare, caules saepe bipedales, racemus semipedalis, calyx 10-12 lineas, corolla sesquipollim longa. Affinis *S. pilifera* ramis glabris, verticillastris paucifloris, calycis dentibus acutis nec aristatis, etc., distinctæ.

25. **S. Cataonica** (Boiss. et Haussk) ramis erectis elongatis induratis simplicibus inferne subnudis aliis annotinis superne foliosis sterilibus, aliis subaphyllis in racemum longum laxum abeuntibus, foliis petiolatis pinnatis crassiusculis crenulatis superne sericeo-tomentosis valde rugoso-lineatis subtus tomentosis opacis segmentis lateralibus binis parvis obovatis terminali magno ovalo basi rotundato vel subcordato, verticillastris 6-2-floris remotis, foliis floralibus oblongis acuminatis præter infima calyce brevioribus, floribus pedicellatis, calycis campanulati parce et patule hirti viridis elevatim striati et nervoso-reticulati labii superioris dentibus tribus late et breviter triangularibus mucronatis, inferioris bifidi dentibus late lanceolatis acutissimis, corollâ..... ♀.

Hab. in regione inferiore montis Berydagh Catonie 3-4000' (Haussk!).

Rami pedales, folia tres pollices longa segmento terminali 1-1½-pollicari, racemus 7-8-pollicaris verticillastris 5-6 remotis, calyx fructifer 7-8 lineas longus. Affinis *S. recognite* et *rubifoliae* differt foliis crassis sericeo-lineatis, etc.; fructifera tantum nota et ulterius investiganda.

26. **S. pilifera** (Moutb. et Auch. Ann. Sc. Nat. 1836, 2 p. 40) caulis floriferis a basi procumbenti erectis virgatis flexuosis ramosis glaberrimis, foliis petiolo gracili longe et sparsim ciliato suffultis pinnatisectis 1-2-jugis segmentis glabriusculis membranaceis ovato-oblongis obtusis crenatis terminali triplo majore basi subcordato, verticillastris 4-2-floris subdistantibus laxe et breviter racemosis foliis floralibus oblongis acuminatis calyce brevioribus sufful-

tis, pedicellis calyce duplo brevioribus, calycis hirtuli ad tertiam partem bilabiati dente superiore sublongiore dentibus tribus triangularibus, inferioris dentibus late lanceolatis omnibus acutissimis, corollâ calyce duplo longiore ♀.

Hab. in monte Akdagl Cappadocia (Auch. 4527!).

Sesquipedalis, foliorum segmentum terminale sesquipollicare, racemi 3-4-pollicares, calyx 7-8 lineas longus.

27. S. bracteata (Russ. Alepp. 2 p. 242) tota patule glanduloso-hirsuta viscida a rhizomate suffrutescenti, pluricaulis, caulis ascendentibus herbaceis paniculatim ramosis, foliis petiolatis pinnatisectis segmentis crenatis lateralibus sapius binis rarius 4-5 ovatis vel oblongis terminali majore ovato-oblongo, floralibus sessilibus persistentibus cymbiformibus basi submembranaceis coloratis apice herbaceis acuminato-attenuatis subrecurvis flores amplectentibus calyce longioribus, verticillastris 8-6-floris remotis, calycis sessilis hispidissimi viscidii labio superiore sublongiore oblongo indiviso vel subtridentato inferioris dentibus ovatis acutis, corollâ purpurascente calyce duplo longiore ♀.

Hab. in collibus et cultis derelictis Cariae in planicie Cybirensi (Boiss!), Phrygiae ad Ouchak (Bal!), Syriæ ad Aleppo (Russ., Auch. 4732! Ky!) et Aintab (Haussk!), ad Damascum (Ky!), Armeniæ Kurdistæ (Ky!), Armenia Turcicæ ad Gumusebkhane (Tehih!), et Ispir (Bourg!).

28. S. trichoclada (Bth. in D.C. Prodr. XII, p. 267) tota patentim pilosa superne præserit valde viscida pallide virens, ramis herbaceis elongatis, foliis membranaceis longe petiolatis pinnatisectis segmento superiore magno ovato vel oblongo grosse dentato basi rotundato vel cuneato lateralibus unijugis parvis oblongis saepe petiolatis, verticillastris 6-10-floris remotis foliis floralibus sessilibus oblongo-lanceolatis acuminatis integris calyces superantibus suffultis, calycis sessilis campanulati pubescenti-viscidi labio superiore apice truncato minute tridentato inferiore bifido, corollâ rubrâ calyce plus duplo longiore ♀.

Hab. in Mesopotamiæ (Auch. 4566!), in monte Pir Omar Gudrun 3500' (Haussk!).

Rami herbacei laxi saepe bipedales, foliorum segmentum superius 1¹-2²-pollicare, calyx 6-7 lineas, corolla 12-14 lineas longa. Affinis *S. bracteata*, pili multo longiores, foliorum caulinorum segmentum majus, corolla longius exserta, planta laxior.

SECT. II. HYMENOSPHIACE.

29. S. Smyrnæa (Boiss. Diagn. Ser. I, 5 p. 5) pluricaulis caulis ascendentibus simplicibus pumilis parte inferiori foliatis patule glanduloso-hispidis, foliis pallide virentibus pubescenti-viscidulis longuscule petiolatis crenulatis a basi rotundata vel subcordata

oblongis vel ellipticis obtusis, floralibus submembranaceis sessilibus ovato-rhombeis breviter acuminatis calyce brevioribus, racemo glanduloso-hispido verticillastris 6-4-floris distinctis subdistantibus constante, calycis membranacei patule hispidi fructiferi aucti ample campanulati divaricatum bilabiati labio superiore sublongiore dentibus tribus breviter triangularibus mucronatis, inferiore in dentes ovatos mucronatos fisso, corollæ cærulescenti-violaceæ calyce triplo longioris tubo recto sensim ampliato ♀.

Hab. in cacumine montis Tartali prope Smyrnam (Boiss!), in monte Bulghasdagh Phrygiæ (Bal!). Fl. Mai.

Semipedalis rarius pedalis, folia fera pollicem longa, racemus semipedalis, corolla sesquipollicularis. Specimiibus fructiferis nunc notis inter *Hymenosphaces* certe militat.

30. S. Cadmica (Boiss. Diagn. Ser. I, 5, p. 6) e rhizomate indurato pluricaulis patule glanduloso-hispida, caulis pumilis ascendenter basi foliosis, foliis longe petiolatis rugulosis crenulatis a basi rotundata vel retusa oblongis ellipticis obtusis sæpius indivisis interdum 2-4-segmentis petiolulatis minimis ovatis ad laminæ basin auctis, racemo subradicali verticillastris 6-2-floris distantibus constante valde hispido, foliis floralibus membranaceis sessilibus cordato-ovatis breviter acuminatis, calycis membranacei longe hispidi tandem ample campanulati divaricatum bilabiati labio superiore truncato interdum obsolete tridentato, inferioris lobis ovatis acutiusculis, corollæ albæ calyce duplo longioris tubo non exerto ♀.

Hab. in regione alpinâ montis Cadmi supra Colossam in Cariâ (Boiss!), Davrosdagh Pisidiæ 4500'-5000' (Heldr!), Bulghasdagh Phrygiæ 4500' (Bal!), Anatoliâ bor. ad Kadikios et Therakli (Wied!).

Semipedalis vel pedalis, folia 1-1 $\frac{1}{2}$ -pollicaria, calyx fructifer 10 lineas longus pollicem apice latus, corolla pollicaris. Affinis *S. Smyrnæa* differt foliis sæpe appendiculatis et præsertim corollâ abbreviatâ albâ calycisque dentibus superioribus fere obsoleteis.

31. S. pomifera (L. Sp. p. 34) fruticosa adpressissime cana, ramis simplicibus, foliis petiolatis rugulosis margine crenulatis et undulato-erispis a basi rotundata vel subcordata oblongis et oblongo-linearibus obtusis, verticillastris 4-2-floris in racemum ovatum vel oblongum approximatis inferiore sæpe subdistante, bracteis parvis membranaceis coloratis deciduis, calycis sessilis late campanulati hirti labiis membranaceis purpureis latis superiore subintegro inferiore bilobo lobis ovatis obtusiusculis mucronatis, corollæ cæruleo-violacea calyce duplo longiore ♂. Ic. Fl. Græc. tab. 45. — *S. crispa* Rivin. — *S. fragifera* Ettl.

Hab. in collibus et fruticetis regionis sempervirentis Cretæ (Tourn! Sibth., Sieb! Heldr! Raul!). In Græciâ et Palestinâ indicata sed ut videtur erroneè.

Insectorum icta gallas fert ad ramulorum extremitatem globosas, cerasi magnitudine, foliis abortivis coronatas, sapore dulci gratas et edules tandem lignosas.

32. S. calycina (Sibth. et Sm. Fl. Græc. I. p. 13, tab 16) fruticosa adpressissime cana ramis elongatis simplicibus vel apice anguste paniculatis, foliis petiolatis rugulosis planis a basi rotundatâ vel subcordatâ ovatis obtusis, verticillastris 4-2-floris racemos breves constituentibus inferioribus remotis, bracteis membranaceis coloratis deciduis, calycis sessilis late campanulati hirti labiis membranaceis dilatatis margine purpureis superiore integro vel repando inferiore bilobo lobis obtusis vel retusis, corollâ roseo violacea calyce plus duplo longiore 5.

Hab. in collibus regionis calide, Atticae in Hymetto et ad prom. Sunium (Sibth. Sprunt!), Lycabeto (Boiss! Heldr!), insulâ Poros (Wied!), Bœotia (Auch. 1525!), insulâ Naxos (Urv!) et Melos (Bory!).

Affinitas *S. pomiferae* differt foliis brevioribus et latioribus non undulatis, rami longiores, racemi saepè subpaniculati, calycis lobi obtusiores.

33. S. Euphratica (Month. et Auch. in Ann. Sc. Nat. 1836, 2, pag. 40) fruticosa ramosa, ramis villosis parte tioriferâ interdum glabratris, foliis rugosissimis crenatis tomento lanato tandem saepè detersili obsitis inferioribus longinseule ceteris breviter petiolatis a basi rotundatâ vel subcordatâ ovatis vel oblongis obtusis, floralibus membranaceis præter intima ovatis calyces non æquantibus, verticillastris 4-2-floris paneis subsecundis laxiusculæ et breviter racemosis, calyce magno tomentello campanulato labiis divergentibus superiore sublongiore subrecurvo acuminato subintegro inferiore obtuse et rotundatim bilobo, corollâ e sicco albidâ calyce duplo longiore tubo amplio 5.

Hab. in Cappadociâ ad Euphratem (Auch. 1516!), in Armeniaustrali (Auch. 1524!).

Variat multum indumento, foliis 1-4½-pollicem longis 8-18-latis. Calyces floriferi 9 lineas fructiferi saepè pollicares et apice pollicem lati, corolla 19 lineas longa.

34. S. Haussknechtii, suffruticosa ramosa viridis pilis brevibus sparsis papillaribus asperula, ramis pumilis ascendentibus ad apicem usque foliosis, foliis petiolis longe ciliatis insidentibus 2-3-jugis segmentis basi attenuatis obtuse denticulatis lateralibus ellipticis vel linearibus interdum breviter decurrentibus terminali paulo majore oblongo-elliptico, verticillastris in capitulum oblongum approximatis foliis floralibus ovatis acuminatis calyce brevioribus suffultis, floribus subsessilibus, calyce fructifero ample membranaceo reticulato-venoso ad nervos et margines ciliato labiis divergentibus superiore sublongiore rotundato retuso inferiore in lobos binos late ovatos mucronatos fiso, corollâ..... 5.

Hab. in saxosis montis Berydagħ Cataoniae 7-8000' (Haussk!).

Fructifera tantum nota, facies et foliis *S. Russeggeri* similis sed calyce diversissima, ramis cum racemo 2-3-pollicari vix semipedales, folii segmentum terminale 6-9 lineas longum, calyx fructifer 9-10 lineas longus labiis apice pollicem fere remotis.

35. **S. Bodeana** (Bge Lab. p. 42) adpresso cinereo-tomentella ramis simplicibus abbrevialis, foliis denticulatis crebre reticulato-rugosis pinnatisectis unijugis segmentis lateralibus ovato-oblongis subsessilibus terminali oblongo multo majore, floralibus oblongo-lanceolatis deflexis acutis floribus brevioribus, verticillastris densis 10-12-floris 3-4 subdistantibus, calycis pubescentis demum valde ampliati purpurascens ad tertiam partem bilabiali labio superiore submajore obtuso obtusissime subsinuato-trilobo inferioris lobis ovatis obtusis, corollæ tubo inclusa ♀.

Hab. prope Djehan Nameh in declivitate meridionali jugi Elbrusensis, ad orientem Teheran (Bode in herb. Hort. Petrop!).

Specimina incompleta 4-5-pollicaria, folia cum petiolo 2-4-pollicaria segmento superiore saepe sesquipollicari, calyx fructifer pollicem longus transverse 14 lineas latus labiis divergentibus, corolla pollicaris. Facies *S. Euphratice*.

36. **S. dracocephaloides** (Boiss. Diagn. Ser. I, 5 p. 4) fruticosa erispula tomentello-papillosa subcanescens, ramis erectis virgatis simplicibus, foliis pinnatisectis 2-3-jugis segmentis subbenerviis foliorum inferiorum obovalis superiorum oblongo-lanceolatis omnibus in petiolulum attenuatis terminali saepius submajore, verticillastris 3-5 distantibus 6-4-floris, foliis floralibus oblongis acuminatis inferioribus ealyces aequantibus, calycis campanulati adpresso canescentis tandem membranaceo-dilatati rabi labio superiore subrecurvo integro vel subdenticulato inferiore in dentes ovatos acutiusculos bifido, corollæ ealyce vix duplo longiore ♂.

Hab. in Persiae borealis deserto Seidkhodji (Auch. 5112! Szov!), in salsis Armenie circa Nackitschewan (Buhse!), circa Tabris et Marand (Bge!).

Rami floriferi $\frac{1}{2}$ -4-pedales, racemus laxus 3-4-pollicaris, folia 1 $\frac{1}{2}$ -2 $\frac{1}{2}$ -pollicaria, flores mediocres, calyx florifer 5 lineas, corolla 9 lineas longa. Absque dubio *Hymenophyllum*.

37. **S. Hydrangea** (p.c. herb. — Bth. Lab. p. 717) basi fruticosa adpresso canescens ramis strictis erectis simplicibus albidis, foliis petiolo brevi stricto insidentibus minime rugosis pinnatisectis rachide brevi, segmentis 3-4-jugis integerrimis acutiusculis lateralibus oblongo-lanceolatis vel anguste linearibus terminali longiore et latiore elliptico-lanceolato, foliis floralibus sessilibus inferioribus pinnatisectis flores superantibus superioribus oblongo-lanceolatis indivisis brevibus, verticillastris pluribus 10-6-floris remotis laxe racemosis supremis abortivis, floribus breviter pedicellatis, calycis magni tenuiter membranacei rosei praeter nervos hirtulos glabri labiis amplis superiore lato obtuso angulato-sinuato inferiore in lobos ovatos acutiusculos bifido, corollæ ealycem parum superante ♂.

Hab. in Persia (Oliv! Auch. 5201!), in plantis Teheran (Jenisch), ad radices montis Sahend prope Tabris et in jago Elbrus (Buhse!), prope Sultanieh (Bge!) in Kurdistania inter Van et Ischri (Noë 154!).

Rami cum racemo $\frac{1}{2}$ -4-pedales, folia inferiora bipollicaria, calyx florifer 9

lineas longus fructifer pollicaris. Species calycibus amplis intense roseis teneris elegantissima.

38. **S. Sheilei** (Boiss. Diagn. Ser. I, 7, p. 45) basi fruticosa adpresso et crispule canescens, ramis erectis rigidulis simplicibus, foliis breviter petiolatis minime rugosis pinnatisectis 2-3-jugis rachide longinseculâ segmentis oblongo-lanceolatis acutiusculis repandis superioribus subdecurrentibus terminali submajore, floralibus membranaceis reticulato-venosis ovatis acuminatis brevibus, verticillastris 6-4-floris remotis laxe racemosis supremis abortivis, floribus subsessilibus, calycis magui glabri tenuiter membranacei rosei labiis amplis subequalibus superiore lato obsolete angulato trilobato inferiore in lobos binos ovatos aequos fissos, corollâ calycem parum superante ♂.

Hab. in Persia australi (Auch. 5201), ad Persepolim et in regionibus altioribus montis Sabst Buschom prope Schiras (Ky. 3771).

Affinis *S. Hydrangea* eni a cl. Bth. conjungitur, differt foliorum segmentis minoribus minus inaequalibus eroso-denticulatis, floralibus membranaceis, flore fere dimidio minores, calycis labia vix inaequalia. Pilorum annulus tubi corollæ ex cl. Bth. in *S. Hydrangea* manifestus in *S. Sheili* obsoletus.

39. **S. Szovitsiana** (Bge Lab. Pers. p. 43) basi sicciflavae pubescentia stellata canescens, ramis numerosis ascendentibus simplicibus pumilis, foliis ad ramorum floriferorum partem inferiorem confertis parvis longe petiolatis rugosis crenatis ovatis obtusis basi rotundatis vel subcordatis lobulis binis interdum auctis, floralibus parvis membranaceis ovatis acuminatis, verticillastris remotis 8-4-floris laxe racemosis terminali abortivo, racemi axi glabri, floribus parvis breviter pedicellatis, calycis hirtuli tubuloso-campanulati purpurascens fructiferi ampliati labiis brevibus aequilongis superiore acutiusculæ tridentato inferiore late bilobo, corollâ calyce vix duplo longiore ♂.

Hab. in Persia boreali prope Urinah (Szovt) in monte Shahkuh ad occasum urbis Schahrud (Derderian).

Rami cum racemo 3-4-pollicari vix semipedales, foliorum laniæ 5-10 lineas longæ, calyx sub anthesi 4 lineas tandem sex lineas longus, corolla fere 8 lineas longa. Facies *S. acetabulosæ* et *cryptanthæ* sed flores multo minores.

40. **S. Rascheyana** (Boiss. Diagn. Ser. I, 12 p. 58) patula villosula viscida pallide virens, ramis simplicibus ascendentibus pumilis, foliis petiolatis pinnatisectis unijugis foliotis rugosis crenulatis oblongis obtusis lateralibus minoribus vel minimis, verticillastris 6-4-floris remotiuseculis longe racemosis foliis floralibus ovatis parvis basi suffultis, calyce parce glanduloso-piloso breviter bilabiato labio superiore sublongiore breviter et late tridentato inferiore bidentato dentibus omnibus mucronulatis, fructifero modice aucto non expanso elevatum reticulato-nervoso, corollâ cærulescente calyce sesquilociore ♀.

Hab. in Antibano circa Rascheya (Boiss!).

Caules semipedales, foliorum segmentum superius vix pollicare, verticillastri 5-7, calyx fructifer 8-9 lineas longus. Affinis *S. acetabulosa* et *S. cryptantha* a quibus præter folia floralia minora differt calyce modice acreto campanulato nec valde expanso.

41. *S. acetabulosa* (Vahl Enum. I, p. 227 et herb!) caulis pluribus ascendentibus simplicibus præter pubem brevem viscidulam patule hispidulis, foliis longe petiolatis rugosis adpresso cinereis aliis simplicibus a basi rotundata vel subcordata ovatis obtusis, aliis segmentis binis parvis a terminali magno subdistantibus auctis, verticillastris pluribus 8-4-floris distantibus longe et laxe racemosis, foliis floralibus inferioribus sessilibus lanceolatis flores superantibus superioribus brevioribus ovato-acuminatis submembranaceis, calyce hispido sub anthesi campanulato breviter bilabiato labio superiore sublongiore breviter tridentato vel subintegro inferiore bidentato dentibus triangularibus acutiusculis, fructifero valde aucto limbo ampio expanso membranaceo-reticulato sæpe colorato, corollæ calyce sesquilongiore ♀. — *S. Tauricola* Schott et Ky in Sched.

Hab. in collibus siccis Orientis (Vahl), ad Tokat (Wied!), Armeniæ Turcicæ ad Erzerum (Huett! Calv!), circa Gumuschkhané (Bourg!), Cataoniæ Beryt-dagh (Haussk!), Cappadociæ ad Argæum (Ky. 206 et Suppl. 242!), Tauro Ciliciæ (Ky.), ditione Kassanoglu Ciliciæ orientalis (Ky. 88!), Mesopotamiæ ad Mardin (Haussk!).

β. simplicifolia. — Folia omnia simplicia, vix varietas. — *S. Molucellæ* Bth. Lab. p. 316. — *S. multicaulis* Vahl Enum. p. 227 ex spec. archetypo herb. Desf!

Hab. in Syriâ circa Aleppo (Auch. 1561! Ky. 69!), in deserto Kotch Hassar Mesopotamiæ (Haussk!), Amano Syrie (Ky. 79!), Libano circa Eden (Bl!), Antilibano (Gaill! Ky. 10!) et circa Rascheya abunde (Boiss!), Armeniæ prope Van (Deyrolle!) et prope Karput (Noël!).

Caules cum racemo 1-4-pedales, calyx fructifer *Molucellæ* diametro sæpe pollicaris.

42. *S. cryptantha* (Montb. et Auch. Ann. Sc. Nat. 1836, 2, p. 40) pilis ramosis brevissimis canescens vel cana, caulis herbaceis ascendentibus simplicibus pumilis sæpe ut et petioli patule hispidis, foliis longe petiolatis rugosis minute crenulatis a basi cuneatâ rotundata rarius subcordata oblongis ellipticis vel elliptico-lanceolatis obtusis, floralibus sessilibus inferioribus flores subæquantibus superioribus membranaceis ovatis parvis, verticillastris 8-4-floris in racemum approximatis inferioribus rarius subdistantibus, calyce basi parce hispido cæterum glabro membranaceo viridi ad medium bilabiato labio superiore duplo longiore recurvo apice obtuso repando, inferioris lobis binis ovatis mucronatis, fructifero valde aucto limbo ampio expanso albo reticulato, corollæ albâ calyce vix sesquilongiore ♀.

Hab. in collibus argillosis Pisidiæ inter Karagatsch et Ghelindost (Heldr!),

Anatoliae bor. ad Kadikios (Wied!), Cappadociæ (Auch. 4531!), in montibus Alidagh et Masineneudagh (Bal!).

Afinis *S. acetabulosæ*, caules breviores cum racemo 6-8-pollicares, folia longiora angustiora, verticillastri sæpissime approximati, calyx præsertim albo-membranaceus labiis valde inæqualibus.

43. S. Pinardi (Boiss. Diagn Ser. I, 12 p 59) basi suffrutescens, ramis erectis inferne tomentellis parte floriferâ glabris, foliis longe petiolatis pube stellatâ a lpressâ subtus copiosiore canescentibus rugulosis crenulatis ovatis vel oblongis basi cordatis interdum segmentis binis lateralibus minutis auctis, verticillastris 6-4-floris distantibus in racemum longum laxissimum basi interdum ramulosum dispositis, foliis floralibus inferioribus oblongo-spathulatis flores æquantibus cæteris membranaceis **acuminatis** brevibus, calycè membranaceo albido fere glabro labio superiore majore obsolete repando-tridentato inferioris lobis ovatis mucronatis, fructifero aucto limbo expanso subreticulato, corollâ calyce sesquilongiore cærulecente ♂.

Hab. in Antilibano inter Rascheya et Damascum (Boiss!), ad Ouadi el Karn et circa Damascum (Gaill!), in monte Schergi prope Zebdani (Ky!), in monte Gebel Abiad deserti Syriaci (Bl!).

Afinis *S. acetabulosæ* et *cryptanthæ* differt basi suffrutescens, racemo laxissimo sæpe pedali et ramuloso, calyce glabro fructifero obsolete nec elevatim reticulato.

SECT. III. DRYMOSPHACE.

44. S. glutinosa (L. Sp. p. 37) caule herbaceo erecto superne præsertim cum inflorescentiâ racemosâ vel paniculatâ glutinoso-piloso, foliis magnis petiolatis ovato-oblongis grosse serratis acuminateis acute cordato-hastatis pubescentibus, floralibus oblongis acutis minutis, verticillastris distinctis 6-2-floris, floribus pedicellatis, calycis tubuloso-campanulati viscidì hirti labio superiore acuto integro vel minute tridentato inferioris lobis binis ovalo-triangularibus acutis, corollæ flavæ calyce triplo longioris tubo exerto intus hirto fauce ampliatâ ♀. Ic. Rehb. tab 45.

Hab. in umbrosis et sylvaticis, Macedonia (Orph!), Thracia in Hæmo (Friv!), mons Amanus Syriae bor. 4000' (Ky. 465!), Pontus Lazicus (Bal!), Armenia Turcica (Huet!), Tauria, Caucasus et Transcaucasia (Led! Rupr!), Persia borealis (Auch. 5196! Buhse!).

Ar. Geogr. Hispania borealis, Gallia media et australis, Helvetia, Germania meridionalis, Italia borealis et media, regio Danubialis, Dalmatia, Rossia australis.

45. S. Forskahlei (L. Mant. p. 26) viridis birta viscidula, caule erecto herbaceo inferne folioso superne nudiusculo in racemum longum simplicem vel paniculam parce ramosam abeunte, foliis magnis tenuiter membranaceis petiolo longo hispido insidentibus grosse et obtuse repando-lobatis ovatis sæpius cordatis sæpissime

basi auriculato-lobatis vel lyratis lobis lateralibus parvis inæqualibus saepè deflexis, floralibus parvis ovatis acuminatis, verticillastris 6-2-floris distantibus, floribus pedicellatis, calycis hirti viscidii labio superiore subtruncato dentibus tribus minutis mucronatis, inferioris dentibus binis lanceolatis subpungentibus, corollæ calyce triplo longioris violaceo-cæruleæ ad labium inferius medium albidae tubo intus annulato ♀. Ic. Fl. Gr. tab. 21. Bot. Mag. tab. 988. — *S. bifida* Forsk Eg. Arab. p. 202. — *S. longepetiolata* C. Koch Linn. XXI, pag. 657!.

Hab. in sylvis circa Byzantium (Forsk., Sibth., Boiss!), circa Bronssam et in regione inferiore Olympi Bithyni (Boiss! Pichl!), Bithyniæ et Galatiæ (Wied!). Ponto prope Niksar (Tehih!) et Trapezuntum (Bourg!), Armeniæ prope Köprubachu (Huet!).

SECT. IV. ETHIOPIS.

§ 1. Homalosphaceæ

46. **S. Syriaca** (L. Sp. p. 36) pluricaulis caulinibus herbaceis erectis saepè elatis glabrescentibus vel puberulis, foliis rugosissimis ininute puberulis flavidо-virentibus cordatis ovato-oblongis crenulatis et inciso-dentatis præter supraem petiolatis, paniculae saepè amplissime et laxæ ramis erecto-patulis elongatis, verticillastris 6-2-floris distantibus, foliis floralibus ovato-oblongis acuminatis florum pedicellos vix superantibus, floribus parvis, calycis tomentelli campanulati obscure bilabiati dentibus brevibus subæqualibus lanceolatis acutis spinulosis, corollæ calyce subduplo longiore albâ ♀. *S. parviflora* Vahl En. I, p. 268. — Labill. Ic. Dec. IV, tab. 7. — *S. varia* Vahl En. p. 273 ex Bth.

Hab. in cultis derelictis, Anatolia interior in Phrygiæ prope Ouchak (Ball!), Cariæ (Boiss!), Pisidiæ et Isauriæ (Heldr!), Lyciæ ad Elmalu (Bourg!), Cappadociæ (Bal!), Galatiæ (Wied!), Ponto australi et Armeniæ (Tehih! Bourg!), Syria ad Aleppo et Damascum (Ky! Gaill!), Mesopotamiæ ad Orfa (Hanssk!), Cœlesyriæ et Antilibano (Boiss!), ditione Talysh (Hoh!), Persiæ boreali (Buhse!) et mediæ circa Ispahan (Bge!).

Sæpe bipedalis et elatior, valde floribunda, calyx florifer $3\frac{1}{2}$ lin. corolla 6 lineas longi. Species floribus parvis ab omnibus sectionis distincta.

47. **S. rhytidæa** (Bth. in D.C. Prodr. XII, p. 280) breviter pubescens, caulinibus erectis elatis superne laxe paniculatis, foliis petiolatis rugulosis subcordato-ovatis erosio-dentatis vel sublobatis subtus saepius canescentibus, superioribus diminutis subsessilibus oblongo-lanceolatis acentis, verticillastris 2-6-floris numerosis inter se remotis, bracteis orbiculatis mucronatis calyce brevioribus, calycis virentis papillari-asperi ad tertiam partem bilabiati labiis divergentibus superiore breviter tridentato inferiore bifido denti us breviter aristatis, corollæ albidae tubo vix exerto ♀.

Hab. in regno Cabulico (Griff. 474 in herb. Kew!).

Planta bipedalis, folia inferiora 3-5-pollicis longa, calyx fructifer $3\frac{1}{2}$ lineas longus et latus. Ex cl. Bth. affinis *S. Syriaca* sed flores multo maiores. Tubi corollini structuram non vidi, affinitas idem subdubia.

48. *S. compressa* (Vahl Enum. I, p. 273) crispuè pubescens, caulis erectis ramosissime paniculatis panicula ramis elongatis patentibus, foliis inferioribus petiolatis a basi rotundatâ vel subcordatâ late ovatis acutiuscelis eroso-dentatis subrungosis superne parce et adpresso arachnoideis, superioribus cordato-amplexicaulinibus, verticillastris 6-4-floris remotinusculis foliis floralibus membranaceis orbiculatis breviter acuminatis calyce subbrevioribus suffultis supremis ahorlivis, calycis tomentelli membranacei fructiferi valde acreti saepe colorati labiis aquilongis in lobos ovatos obtusos bifidis, superioris dente tercio supremo minimo mucroniformi, corolla rosea calyce vix duplo longioris tubo inclusa recto intus nuda ♀. Ic. Vent. Gels. tab. 59. — *S. polyclonos* Boiss. Diagn. Ser. I, p. 47.

Hab. in Mesopotamia inter Mossul et Bagdad (Olivy), in Kurdistania inter Van et Ischri (No. 457!), Persia australi ad Dalechi (Ky. 425!) et prope Kasernu (Haussk!), ad Dair Hadj Afghanistan (Griff!).

Planta 1- $1\frac{1}{2}$ -pedalis, panicula $\frac{1}{2}$ -4-pedem lata, folia 3-4 poll. longa, calyx fructifer $3\frac{1}{2}$ lineas longus fructifer 6 lineas longus et apice latus. Ex amic. a Bge ab calycem fere *Hymenosphaerae* sectionis propriae *Tonosphaera* typus, sed fere eadem calycis structura in aliis *Ethiopitis* speciebus v. gr. in *S. hypoleuca* observatur.

49. *S. eriophora* (Boiss. et Ky. exs. 1859) pluricanulis, caulis erectis pumilis crebre foliosis in paniculam brevem pyramidatam abeuntibus, foliis viridibus adpresso-roseis hirsutis rugosissimis anguste lanceolato-linearibus aenatis inferioribus in petiolum attenuatis superioribus sessilibus, verticillastris 4-2-floris in racemos tenues albo-lanatos approximatis, foliis floralibus membranaceis cærulescentibus ovatis mucronulatis calyces subæquantibus, floribus parvis sessilibus, calycis albo-lanati cærulescentis tubuloso-campanulati dentibus longe subulato-spinosis rectis superioris intermedio duplo breviore, corolla cærulea calyce sesquilongoris tubo recto inclusa intus nuda labio superiore rectinuscule ♀.

Hab. ab Jool Baatch ad radices montis Allahdagh Cilicie 5000' (Ky. 229!).

Planta semipedalis, folia 2- $2\frac{1}{2}$ -pollicaria 2-3 lineas lata, racemi tripollieares tenues ut in *Ajuga* cærulescentes lanati, calyx fructifer 4 lineas, fructifer 5 $\frac{1}{2}$, corolla 6 lineas longa. Species characteribus insignis, nulli affinis.

50. *S. Montbretii* (Bth. Ann. Sc. Nat. 1836, 2 p. 42) caulis ex rhizomate incrassato pluribus erectis elongatis inferne canescens-tibus in racemum viscidulum simplicem vel basi parce et striete ramosum abeuntibus, foliis radicalibus confertis supra virentibus subtus adpresso canis oblongo-spathulatis vel obverse lanceolatis obtusis in petiolum longe attenuatis integris vel acute paucidentatis,

intermediis linearibus acutis, verticillastris 10-6-floris numerosis distantibus longe racemosis foliis floralibus transverse latioribus breviter acuminato-recurvis extus canis et lineatis dimidiis calyces æquantibus suffultis, calycis cani striati labiis æquifloris superiore breviter tridentato inferioris dentibus lanceolatis omnibus mucronatis, corollæ violaceæ calyce plus triplo longiore ♀. *Salvia hypargyeia* F. et M. Ann. Sc. Nat. 1854, I, p. 34. Tchih. As. Min. tab. 22.

Hab. in monte Karadagh Lycaoniæ (Heldr!), Cappadociâ (Tchih! Ball!), Galatiâ (Wiedt!), Tauro Cilicio et ditione Kassan Oghlu (Ky! Ball!), Syria boreali ad Aintab et Nisib (Auch. 1534! Ball!).

Sesquipedalis, folia inferiora cum petiolo 3-5-pollicaria 5-8 lineas lata, racemi 6-9-pollicares, calyx 4 $\frac{1}{2}$ lineas, corolla 12-15 lineas longa. A *S. phlomoide* Asso Hispaniæ et Algeriæ cui eam prius adnumeravi certe distincta bracteis obtusioribus calyces non subæquantibus, calycis labio superiore non valde dilatato et truncato. Ambarum tubum corollæ intus glaberrimum video.

51. ***S. canescens*** (C. A. M. Enum. p. 86) rhizomatis ramis prostratis induratis radiculiferis, foliis omnibus fere radicalibus rosulatis cespitosis utrinque adpresso albo-lanatis a basi sensim attenuata oblongis vel lanceolatis acutiusculis sinuato-lobatis vel sublyratis rarius subintegris, ramis floralibus pumilis erectis subnudis inferne albo-lanatis in racemum simplicem vel basi parce ramulosum patule viscosissimum abeuntibus. foliis floralibus transverse ovatis abrupte et breviter acuminatis minimis, verticillastris 6-4-floris remotis, floribus parvis subsessilibus, calycis viscidii costati labio superiore subbreviore truncato dentibus breviter triangulari-ovatis mucronulatis inferioris lanceolatis acuminatis, corollæ violaceæ calyce duplo longiore ♀. Ic. Bot. Reg. 1838, tab. 36 (mediocris, ex specimeni culto).

Hab. in Caucaso boreali ad Nartzana (C.A. Mey.), in monte Elbrus (Radde!), Ossetiâ et districtu Alagir 2500'-3000' (Ruprl!), Caucaso orientali in Daghستانâ 4800'-5500' (Ruprl!).

Sæpius *S. Montbretii* minor, ramis 4-8-pollicaribus, folia cum petiolo brevi 1-3-pollicaria, verticillastri 3-6 parvi, calyx 4 lineas, corolla 8 lineas longa. Species foliis albo-panniosis elegans foliis abbreviatis acutis, floribus fere duplo minoribus, etc., a *S. Montbretii* distincta.

52. ***S. hypochionæa*** (Boiss. Diagn. Ser. I, 5 p. 8) caulis ex rhizomatis ramis horizontalibus ascendentibus pumilis simplicibus, foliis ad caulum basin confertis vix rugosis supra viridibus parce araneosis subtus adpresso canis petiolatis a basi truncato-subcordata ovatis et oblongis obtusis irregulariter eroso-sublobatis, verticillastris 6-4-floris subdistantibus racemosis, foliis floralibus persistantibus concavis ovato-orbiculatis breviter acuminatis lineatis calyces subæquantibus, calycis visciduli tubuloso-campanulati breviter bilabiati dentibus superioribus brevibus subrecurvis intermedio breviore, inferioribus lanceolatis omnibus acutis, corollæ calyce duplo longiore ♀.

Hab. in argilloso montis Seidkhodji prov. Aderbidjan Persie borealis (Auch. 5207! Szov!).

Caulis cum racemo interdum semipedali 6-12-pollicares, folia radicalia $2\frac{1}{2}$ -4 poll longa $1\frac{1}{2}$ lata, calyx florifer 7 lineas fructifer 9 longus. Ex affinitate *S. Montbretii* a qua foliis latis et bracteis multo amplioribus differt

53. *S. phlyctidea* (C. Koch Linn. XXI. p. 655) caulis inferne arachnoideo-canescens in racemum basi striete et breviter ramulosum abeuntibus, foliis superne glabrescentibus ramosissimis subtus adpressissime niveis a basi rotundata ovatis obtusissimis crenato-dentatis inferioribus longe petiolatis superioribus sessilibus basi breviter cuneatis, floralibus ovatis cuspidatis membranaceo-violaceis cito deciduis calyce brevioribus, verticillastris 6-4-floris subdistantibus, calycis papilloso-hirti visciduli campanulati labiis divergentibus dentibus superioris breviter subulatis conniventibus, inferioris lanceolatis apice subulatis, corollæ calyce duplo longioris tubo recto intus nudo, labio superiore falcato ♂.

Hab. in Grnsiâ (Schmidt et Wilhelms ex C. Koch!).

Folia 2-3-pollices longa, racemus subramosus $\frac{1}{2}$ -1-pedalis, calyx florifer 5 lineas, fructifer subdilatatus 6 lineas longus. Folia et facies *S. hypoleuca* bracteis latis concavis persistentibus, calyce fructifero tubuloso 9 lineas longo etc. alienæ. Tantum corollam emarcidam observare potui.

54. *S. leriae-folia* (Btb. in n.c. Prodr. XII. p. 287) tota dense et adpresso albo-pannosa, caulis humilis crassis erectis in racemum oblongum simplicem vel basi ramulo auctum abenntibus, foliis crassis radicalibus amplis oblongis irregulariter sinuato-lobatis et basi saepe incisis in petiolum attenuatis, caulinis paucis subsessilibus, floralibus oblongis acutiusculis præter infima calyce multo brevioribus, floribus magnis subpedicellatis 10-5-verticillatis, calycis adpresso arachnoidei ad medium fere bilabiati labio superiore brevissime tridentato inferioris dentibus lanceolatis omnibus mucronatis, corollâ albida calyce vix sesquiloniore ♀.

Hab. in regno Cabulico (Griff!), in planicie arida circa Nichapur prov. Khorassan Persie boreali-orientalis (Bge!).

Folia radicalia cum petiolo 6-9-pollicaria 2- $2\frac{1}{2}$ pollices lata racemum saepe æquantia, racemus 4-6-pollicaris præter verticillastros inferiores densiusculus, calyx florifer 9 lineas longus. Species pulchra.

55. *S. spinosa* (L. Mant. p. 511) superne præsertim pubescenti-viscida, caule erecto elato in paniculam amplam pyramidatam abennte, foliis adpresso pubescenti-canescens in vel pallide virgentibus amplis ovatis subcordatis vel basi rotundatis erosio-dentatis saepe sublobatis inferioribus petiolatis superioribus amplexicanibus acutis, floralibus late ovatis acuminatis submembranaceis calyce subbrevioribus, verticillastris remotiusculis 4-2-floris, calycis viscidohirti tubulosi fructiferi aucti labiis brevibus in dentes superioris ternos subæquilongos inferioris binos trianguli-lanceolatos spino-

sos fissis, corollæ albæ calyce sesquilonigioris tubo inclusu recto intus nudo, labio superiore subfalcato abbreviato ♀. Jacq. Ic. rar. tab. 7! *S. præcox* Vahl Enum. I, p. 274 excl. Tournef. ex Bth.

Hab. in Arabiâ petræa (Schimp. 343!), Syriâ ad Aleppo (Ky. 199!, Boiss!), desertis Mesopotamiæ (Oliv!) et ad fl. Chabur et in monte Sindjar (Haussk!), in Persiâ indicata sed hic locus valde dubius.

Pedalis et elatior, panicula æque longa ac lata, calyx fructifer fere pollicaris, corollæ labium superius inferiori viz. longius.

56. *S. Reuteriana* (Boiss. Diagn. Ser. I, 5 p. 10) caule erecto inferne papilloso superne in paniculam viscidulam abeunti, foliis rugulosis adpresso et subtus densius canescensibus ad nervos hirsutis petiolatis a basi rotundata vel cordatâ ovatis obtusis crenulatis, floralibus amplissimis hyalino-membranaceis albido ovato-orbiculatis breviter acuminatis calyces æquantibus, verticillastris 6-2-floris approximatis, calyce membranaceo albido longe tubuloso-campanulato ad venas glanduloso-puberulo dentibus rectis lanceolatis apice subulatis non pungentibus superiore brevissimo, corollæ calyce vix sesquilonigioris tubo inclusu labio superiore abbreviato ♀.

Hab. in Persiæ borealis monte Elbrus prope Passgala (Ky. 191!), in ditione Khorassan (Bge!), circa Ispahan (Auch. 5205!), et in ambitu deserti salsi (Bge!).

Affinis *S. spinosæ* differt bracteis amplioribus, calycis hyalino-membranacei dentibus teneris non pungentibus superiore brevissimo; planta humilior, panicula confertiflora.

57. *S. Persepolitana* (Boiss. Diagn. Ser. I, 12 p. 60) caulis papilloso-asperis crebre foliosis mox in paniculam pyramidatam papillari-subviscidam abeuntibus, foliis rugulosis viridibus ad nervos papillosis ovatis vel oblongis erosio-dentatis sublobatis basi rotundatis vel subcordatis, floralibus membranaceis albido ciliatulis cucullatis orbiculatis breviter acuminatis calyces æquantibus, verticillastris 6-4-floris approximatis, calycis late campanulati papillosi fructiferi aucti dentibus late ovato-triangularibus spinulosis superiore breviore, corolle lilacinae calyce sesquilonigioris tubo subexerto labio superiore brevi rectiusculo ♀.

Hab. ad rupes prope ruinas Persepolis (Ky. 279! sub *S. Reuterianæ*).

Pedalis, odorem suavissimum spirans, folia 1 $\frac{1}{2}$ -2-pollicaria, calyx florifer 5 lin. fructifer 6-7 longus et apice latus, corolla 7 $\frac{1}{2}$ lin. longa. A *S. Reuterianæ* quæcum olim combinavi, *S. Palestinæ*, calyce campanulato nec tubuloso statim dignoscenda, a *S. Selareæ* cæteris omisssis corollæ tubo sensim in faucem ampliato intus nudo et labio superiore brevi diversa

58. *S. Palæstina* (Bth. Lab. p. 718) caule erecto patule et longe papilloso strigilloso in paniculam strictam abeunte, foliis rugosis adpresso et subtus canescensibus vel canis inferioribus petiolis strigosis suffultis oblongis basi saepius pinnatilobatis vel lyratis segmentis lateralibus saepè confluentibus terminali multo majore, floralibus membranaceis saepè coloratis orbiculatis concavis cuspidatis

calyce subbrevioribus, verticillastris distantibus 6-4-floris longe ramosis, calycis tubulose-campanulati glanduloso-punctati papilloso labiis brevibus in dentes lanceolatos rectos longe spinulosos superiore trifido inferiore bifido abeuntibus, calyce fractifero aucto tubuloso, corollae albæ calyce subtriplo longioris tubo recto intus nudo, labio superiore elongato rectiusculo ♀. *S. Sinaica* Delile Mss. — *S. Sieberi* Presl Bot. Beuerk. p. 100. — *S. Rassami* Boiss. olim in Ky. exs.

Hab. in collibus siccis Ægypti in valle Ouadi Askar deserti Arabie (Schweinf!), Arabiae Petree et jugi Sinaitici (Boiss! Auch. 4539! Schimp. 321! sub *S. Selareæ*), Palestine (Bové, Sieb.), Syria ad Damascum (Boiss! Gaill!), ad Zebdani (Ky. 680!), in Libano (Ehr!), ad Aleppo et Aintab (Ky. 200! Boiss! Haussk!). Mesopotamia (Haussk! Noë!). Persie australis ad Gere et Kaserun (Ky. 406! et 980! Haussk!), inter Yezd et Ispahan (Bge!).

Pedalis et elatior, indumentum et bractæ membranaceæ. *S. Selareæ* calyce fructifer breviter campanulato, corolla tubo brevi intus appenicalato abrupte ampliata et labio superiore valde falcato diversissimæ. Calyx *S. Palestinae* florifer 6 lineas fructifer 8-9 longus, corolla 14 lin. longa. *S. spinosa* L. affinis quoque differt foliis ovatis indivisis, paniculâ late pyramidatâ, calyce longiore saepè pollicari, præsertim corollæ brevioris labio superiore brevi nec longo.

59. *S. macrosiphon* (Boiss. Diagn. Ser. I, 5, p. 41) caule gracili elato superne subundo in paniculam longe ramosam breviter viscidulam abeunte, foliis breviter petiolatis parvis rugosis ubi inque canis a basi rotundatâ rarius 4-2 segmentis minutis auctâ ovato-oblongis obtusis crenatis. floralibus subnembranaceis ovato-acuminatis calyce multo brevioribus, verticillastris 4-2-floris remotissimis, calycis papilloso-asperi longe tubulosi dentibus rectis lanceolatis acutis vel apice subulatis superiore breviore, corollæ albidae calyce subduplo longioris tubo longe exerto labio superiore elongato rectiusculo ♀.

Hab. in Persia australi inter Fasa et Schiraz (Auch. 5197!).

β. *Kotschyi*. — Calyx et corolla subbreviores. — *S. Kotschyi* Boiss. Diagn. Ser. I, 7, p. 46.

Hab. inter segetes prope Schiraz (Ky. 291!), prope Ispahan (Bodé!), prope Kerman et in prov. Khorassan (Bge!), in Belutschia (Stocks N° 709!), regno Cabulico (Griff. 475).

Planta 1½-2-pedalis, folia 1½-2-pollices longa, panicula rami longi remotiflori. Calyx fructifer in typo pollicaris, in varietate paulo brevior, corolla in typo sesquipollicaris in varietate pollicaris. Minime *S. spinosa* L. sed *S. Palestinae* affinis a qua corollæ tubo longe exerto statim dignoscitur.

§ 2. Gongrosphaceæ.

60. *S. graveolens* (Vahl Enum. I, p. 273) basi suffruticosa ramosa ad ramos et petiolos patule papillari-hirsuta ad folia adpresso canescens, ramis erectis vel ascendentibus simplicibus vel saepius stricte paniculatis, foliis rugosissimis margine crenatis et saepè undulato-lobatis a basi truncata vel cordata triangulari-oblongis oblulis

inferioribus breviter petiolatis cæteris sessilibus, floralibus parvis ovatis acuminatis, verticillastris 6-4-floris præter infimos subdistantes approximatis longiuscule racemosis, calycis hispidissimi campanulati labiis patentibus superioris dentibus ternis brevibus inferioris binis late lanceolatis omnibus acutis breviter mucronatis, corollâ albâ calyce subdupo longiore tubo brevissimo intus appendice pilosâ aucto labio superiore elongato ♂. *S commutata* Bth. Lab. p 222. — *S. Dominica* L Sp. 35 non Swartz.

Hab. in Palestinâ circa Hierosolymam et Ramleh (Boiss!), in Syriâ circa Damascus et in Cælesyriâ (Auch. 1558!, Gaill!, Socint!).

Pedalis et sesquipedalis, folia inferiora bipinnicaria, racemi hirsutissimi, calyx florifer 4½-5 lin. longus fructifer subauctus. Ab omnibus speciebus sectionis *Aethiopis* suffrutescentiâ et habitu proprio distincta.

61. ***S. Sclarea*** (L. Sp. 38) caule herbaceo crasso erecto inferne hirsuto superne in paniculam viscidulam ramosissimam strictam abeunte, foliis viridibus pubescentibus inferioribus petiolatis sæpius amplis ovatis plerumque cordatis crenatis vel dentatis, floralibus magnis ovatis concavis tenuiter acuminatis membranaceis coloratis calyces superantibus tandem deflexis, verticillastris distantibus 6-2-floris supremis abortivis, calycis pedicellati papilloso-asperuli fructiferi modice aucti breviter et late campanulati labio superiore tridentato inferiore bifido dentibus omnibus subspinoso-subulatis, corollæ roseo-caeruleæ calyce triplo longioris tubo inclusa intus penicillo appendiculato ♀. Ic. Fl. Græc. tab. 23. — Rehb. Ic. tab. 48. — *S. bracteata* Bot. Mag. tab. 2329 nec Russ.

Hab. in umbrosis siccis regionis præsertim montanae, Græcia in monte Kyllene 3-4000' (Heldr!), Macedonia et Thracia (Friv!), insula Cos (Urv.), Anatolia tota! Libanus (Bl! Gaill!), Armenia (Huet! Bal!), Tauria, Caucasus et regio Transcaucasica (Hoh! Mey!), Kurdistania Persica (Haussk!), montes Persici australis (Haussk!), et borealis (Auch! Kyl!), ditio Khorasan (Bge!).

Ar. Geogr. Europa media et australis a Belgio et Gallia ad Germaniam australem, regionem Danubialem et Rossiam australem, Africa borealis.

62. ***S. Aethiopis*** (L. Sp. p. 39) albo-lanata non viscosa, caule crasso erecto a parte inferiori ample et pyramidatim paniculato, foliis fere omnibus ad collum confertis expansis breviter petiolatis rugosis ovato-rhombeis eroso-crenatis et irregulariter plus minusve incisio-lobatis, caulinis paucis cordato-amplexicaulibus, floralibus latissimis striatis sœpe glabrescentibus subrotundo-ovatis cuspidatis calyce subbrevioribus, verticillastris 10-6-floris distinctis secus paniculæ ramos laxe racemosis, calycis lanati fructiferi accreti dentibus lanceolatis apice subulato-spinosis superiore breviore, corollæ albæ calyce duplo longioris tubo inclusa in faucom abrupte dilatato intus appendiculo penicillato aucto ♀. Ic. Jacq. Anstr. tab. 211. — Rehb. tab. 47. — *S. Kochiana* Kunze Sem. Hort. Lips. 1847! ex specimine herbarii G. Koch!

Hab. in siccis et cultis otiosis, Graecia (Sibth.), Macedonia (Fryi), Anatolia omnis interior et maritima a Bithynia ad Lycaiam (Wied!), Boiss! Heldr! Bourg!), Armenia Turcica (Huet!), Tauria Caucasus et Transcasica (M. B., Ledeb., Ruprecht!), Persia borealis (Buhse!, Bge!).

S. leuconeura. — Folia subtus secus nervos cerebrini pilosa albo-lanata. Flores subminores. — *S. leuconeura* Boiss. Diagn. Ser. II, 4, p. 20.

Hab. in Assyria prope Mardin (Aueh. 1543!).

Ad *S. verbascofolium* ex cl. Bth. refuta sed inflorescentiae indumento non viscido hue spectat.

Ar. Geogr. Hispania, Gallia et Germania australes, Italia media, Dalmatia, regio Danubialis, Rossia australis.

63. *S. aristata* (Aueh. ex Bth. n.c. Prodr. XII, p. 270) praeter canes inferne glabriusculos tota patula pubescenti-glandulosa, caulinibus crassis brevibus patulo supra basin in paniculam pyramidaliter abenitibus, paniculae ramis rigidis erecto-patulis, foliis ambitu oblongis petiolatis in segmenta anguste linearia secus rachidem decurrentia indivisa vel incisa pinnatisectis, floralibus sessilibus indivisis longe linear-i-lanceolatis attenuato-acuminatis, floribus pedicellis calyce saepius longioribus suffultis in verticillastra 4-2-flora laxa secus paniculae ramos dispositis supremis abortivis, calycis hispido dentibus a basi late lanceolata longe mucronato-aristatis tubo vix brevioribus superiore minimo mucroniformi, corollae albæ calyce triplo longioris tubo recto breviter exerto ♀. *S. Overini* Trautv. Stirp. nov. v. № 42!

Hab. in monte-is Persie occidentalis (Aueh. 1563!), inter Sengar et Sultanich (Bge!), prope Mendeli (№ 1183!), ad lacum Urniyah (Overin!).

Plantæ vix pedalis, folia bipinnicaria lacinii 1-1½ lineam lati, paniculæ rami 4-6-pollicares, calyx 7 lineas longus fructifer subdilatatus, corolla 15-16 lineas longa, connectiva postice deflexa loculis ut videtur fertilibus nec inter se connexis, lineæ binae oblique pilorum in corollæ tubo. Ex cl. Bunge sectionis proprie *Physophace* typus sed mihi omnino inter *Ethiopes* militare videtur.

64. *S. ceratophylla* (L. Sp. 39) inferne eum foliis laxe albo-lanata, caule crasso erecto protero paulo supra basin longe et ample paniculato, paniculæ hispido-viscidæ ramis elongatis erecto-patulis, foliis inferioribus humili adpressis ambitu late lanceolatis in lacinias elongatas late lineares integras vel pinnatisidas secus rachidem alatum decurrentes sectis, foliis floralibus ovatis acuminatis calycei aquilon-gis vel brevioribus, verticillastris 6-4-floris distantibus superioribus abortivis, calycis papilloso-hispidi fructiferi aucti labiis demum divaricatis superiore dentibus ternis intermedio breviore, inferioris dentibus binis lanceolatis omnibus spinulosis, corollæ albæ calyce sub-duplo longiore ♀.

Hab. in collibus aridis Lydie ad radices Mesogis (Boiss!), Pisidia (Heldr!), Paphlagoniae (Wied!), Cappadociae (Aueh. 1549!, Bal!), Armenia Turcica (Huet!), Syriae ad Aleppo (Haussk!) et deserti Syriaci (Bl!), Mesopotamia et

Assyriæ (Ky. 281! Haussk!), ditione Talysch (C.A. M! Hoh!), Persiæ boreali et mediæ (Oliv! Auch. 5200! Bge!).

Planta $\frac{1}{2}$ - $\frac{1}{2}$ -pedalis, folia magnitudine valde variabilia interdum pedalia, calyx florifer vix 5 lineas, fructifer 9 lineas longus et apice latus. Corollæ tubus intus squamâ fimbriatâ auctus.

65. *S. chionantha* (Boiss. Diagn. Ser. I, 5 p. 8) caule crasso valde elato superne amplissime paniculato ut panicula tota hirto-viscosissimo, foliis utrinque laxe albo-lanatis lenge linearis-lanceolatis minutissime crenulatis vel integris obtusiusculis basi attenuatis inferioribus breviter petiolatis, superioribus anguste lanceolatis acuminate, floralibus triangularibus mucronatis iniunctis florum pedicellos vix æquantibus, verticillastris 8-4-floris secus paniculæ ramos erecto-patulos elongatos remotis, calycis parce glanduloso-hirti labiis divergentibus superiore late et breviter tridentato dente medio minimo inferioris dentibus lanceolatis omnibus breviter spinuloso-mucronatis, corollæ niveæ calyce $2\frac{1}{2}$ -plo longioris tubo inclusa labio superiore lato valde falcato ♂.

Hab. inter segetes planitiei elatae Cariæ ad meridiem montis Cadmi (Boiss!) in Lyciâ prope Elmalu (Bourg. 224!).

Species pulchra foliis *S. Montbretii*, floribus *S. argenteam* referens, folia 4-9 poll. longa pollicem lata vel angustiora, caulis tripodalis, panicula $1\frac{1}{2}$ -pœdem longa et basi lata, calyx 6-7 lineas, corolla 12-15 lineas longa.

66. *S. argentea* (L. Sp. p. 38) caule erecto ample paniculato, paniculæ hirto-viscidæ ramis erecto-patulis elongatis, foliis ad caulis basin confertis breviter et grasse petiolatis ovatis et oblongis basi rotundatis vel cuneatis utrinque sed subtus densius arachnoideo-canescens sinuato-lobatis erosis, floralibus membranaceis viscidulis viridibus ovatis cuspidatis calyce duplo brevioribus, verticillastris 6-4-floris distantibus racemosis, calycis glanduloso-hirti labio superiore tridentato dente intermedio brevi, inferiore bidentato dentibus omnibus apice subulato-spinulosis, corollæ albæ calyce triplo longioris tubo inclusa appendiculâ hirtâ aucto superiore valde incurvo falcato ♂. Ic. Fl. Græc. tab. 27. — Rehb. Ic. tab. 46.

Hab. in cultis derelictis prope Athenas (Spr! Orph. Fl. exs. 547!), in regione alpina Parnassi (Heldr!), Macedoniâ (Friv!), agro Trojano (Schmidt!).

Planta saepe bipedalis, folia radicalia saepe 4-7 poll. longa, calyx florifer 6 lineas longus fructifer subauctus, corolla 16 lineas longa. In Cretâ a Linnæo indicata sed a recentioribus ibi non reperta. *S. Gussouii* Boiss. Italica et Sicula et *S. patula* Desf. boreali-Africana hujus forsitan varietates sunt sed characteribus nonnullis diversæ.

Ar. Geogr. Serbia, Dalmatia.

67. *S. Tmolea* (Boiss. Diagn. Ser. I, 5, p. 9) inferne laxe arachnoideo-canescens superne hispidula viscosissima, caulis proceris superne paniculatis, foliis inferioribus longe petiolatis a basi attenuatâ oblongo-lanceolatis vel subrhombatis profunde et irregula-

riter lobatis eroso dentatis, superioribus sessilibus acute dentatis acuminatis, floralibus ovato-orbiculatis cuspidatis calyce subbrevioribus, verticillastris 6-4-floris remotis, calycis viscosi labio superiore tridentato dente intermedio brevi nuncio inferiore hirsuto dentibus omnibus spinuloso-mucronatis, corolle calyce triplo longioris labio superiore caeruleo incurvo-falcato inferiore albido, tubo inclusu ♀.

Hab. in rupib. alpinis Tmoli occidentali supra Philadelphiam (Boiss!).

Bipedalis, folia inferiora longe petiolata 4-6-pollices longa, calyx 5-6 lineas, corolla 14-15 longa ea *S. argentea* fere major. Huic et *S. verbascifolia* affinis differt foliis profunde lobatis, corollae colore.

68. *S. verbascifolia* (M.B. Taur. Cauc. 3 p. 24) inferne canescens vel laxe cana partibus floralibus breviter viscido-hirta, canle erecto in racemum simplicem vel paniculam parum ramosam abenente, foliis inferioribus petiolatis rugosis ovatis vel oblongis inciso-lobatis basi cuneato-truncatis vel subcordatis, superioribus paucis sessilibus oblongis acutis vel acuminatis, verticillastris 6-4-floris distinctis racemosis foliis floralibus subrotundis cuspidatis membranaceis calyce triplo rarius duplo brevioribus siccatis, calycis papilloso-hirtelli visciduli labio superiore tridentato dente intermedio nimirum, inferiore bidentato dentibus apice longiusculo subulato-spinulosis, corolle calyce subdupo longioris labio superiore albido inferiore ochroleuco, tubo inclusu intus appendicula piloso aucto ♀. *S. candidissima* M.B. Taur. Cauc. I, p. 22 non Vahl.

Hab. in ditione Cubensi Iberia et prope Tiflis (Ower!), circa Elisabethpol (Hoh!), in monte Beschbarmak (Karel!), in prov. Aderbidjan Persie borealis (Buhse! Szov.).

β. cana. — Albo-lanata, ramosior elatior, folia ovata vel oblonga basi subcordata vel truncata aut breviter cuneata. — *S. microstegia* Boiss. et Bal. Diagn. Ser. II-4, p. 19. — *S. pseudoathiopis* Boiss. in Ky. Sched. — *S. nicea* Ehrenb. MSS. — *S. atomaria* Boiss. Diagn. Ser. II, 4, p. 18 (forma calycis dentibus brevissime spinulosis).

Hab. in alpinis Teng prov. Musch Armenie 7000' (Ky. 667!), monte Alidagh Cappadociae 5000' (Bal!), Berytdagh Cataonia 70° 0' (Hanssk!). Tauro Cilicieo supra Gülek Boghas (Bal. sub *S. Tmolea*), Syria borealis monte Amano (Ky. 227! sub *S. Tmolea*), Libano ad Becherre et Cedros (Auch. 1544! Ehr! Boiss! Bl! Ky. 289! sub *S. Tmolea*), Hermone (Ky. 927!).

Valde variabilis foliorum forma et indumento, valde affinis *S. argentea*; flores et praesertim corolle minores 9-10 lineas tantum longe, folia basi saepius truncato-subcordata, inflorescentia in formâ typicâ simplicior sed panicula in var. β saepe ampla.

69. *S. Atropatana* (Bge Lab. Pers. p. 47) inferne arachnoideo-canescens partibus floralibus viridis glanduloso-hirta, caule elato ample paniculato, paniculae ramis divergentibus elongatis, foliis inferioribus petiolatis a basi cuneatâ rarius truncatâ oblongis vel linearilanceolatis obtusis erosulis et sinuato-lobatis, floralibus viridibus ovato-orbiculatis cuspidatis pungentibus saepe recurvis dimidiis

calyces aequantibus, verticillastris 6-4-floris remotissimis longe racemosis, corollæ calyce duplo longioris tubo inclusu intus squamuâ obsoito ♀. *S. argentea* β? *angustifolia* Bth. in d.c. Prodr. XII. p. 285. — *S. hypochionea* Buhse exs. non Diagn. — *S. verbascifolia* var. Boiss. herb. — *S. Bachtiarica* Bge Lab. Pers. p. 47? ex descript.

Hab. in prov. Aderbidjan Persiae prope Tikmedesch (Bge!), in monte Elbrus Persiae borealis (Ky. 254!), circa Schiraz et in monte Kuh Delu Persiae australis (Ky. 535! et 982!), in monte Sawers, Nur, Eschker, Kellal ad nives 9-12'000' (Haussk!), Avroman et Schahu (Haussk!).

Affinis *S. verbascifoliae* et *argenteae* sed folia angustiora et minora, panículæ rami elongati divergentes, bractæ longius recurvato-cuspidatae. Flores interdum illis *S. verbascifoliae* minores interdum majores sed inter omnes formas transitus manifesti. An tamen *S. verbascifoliae* varietas quodam modo geographica dicenda ulterius observandum. *S. Bachtiarica* Bge ex specimine ab eodem in schedula speciminis Bodeani in herb. Petrop. servati hoc nomine inscripto est *S. ceratophylla* ipsa, sed ex descriptione ad *S. Atropatanam* referenda videtur.

70. S. Urmensis (Bge Lab. Pers. p. 48) tenuiter floccosovillosa virens. caule elato in paniculam paucirameam abeunte, foliis radicalibus amplissimis longe petiolatis cordato-ovatis grosse incisodentatis lœvibns. caulinis sensim diminutis sessilibus suborbiculatis abrupte acuminatis omnibus subtus præter nervos subglabris. verticillastris numerosis sexfloris superioribus approximatis, bracteis canescensibus late ovatis calycem superantibus. calycis campanulato-infundibuliformis tenuissime 47-18 nervii extus villosuli dentibus ovatis subulato-mucronatis, corollæ tubo fanceque inclusis, galea falcatâ glandulosâ ♀.

Hab. in elatis montiam ad viam pone pagum Kuschra districtus Urmiah prov. Aderbidjan (Szovt!). Non vidi.

Ex claro auctore *S. Atropatana* proxima, calyce tenuiter multistriato distincta. Ex unico specimine descripta et ulterius investiganda.

71. S. xanthocheila (Boiss. Diagn. Ser. 1, 12, p. 59) inferne laxe lanata canescens vel cana superne patule hispida glandulosa, caule brevi erecto mox in paniculam brevem pyramidatam densifloram abeunte, foliis ovatis subcordatis lobat-dentatis inferioribus petiolatis superioribus sessilibus acute dentatis, floralibus membranaceis late ovatis cuspidatis calyces aequantibus vel subbrevioribus, verticillastris 6-8-floris approximatis, calycis hispida campanulati dentibus latis triangularibus spinuloso-aristatis superiore brevi submutico. corollæ calyce 2½-plo longioris tubo inclusu labio superiore albo inferiore sulphureo ♀. *S. frigida* Boiss. in Buhse Aufz. non Diagn.

Hab. in regione alpinâ Armeniae in monte Techdag (Huet!), Armeniæ Rossicae in montibus Erivan et in ditione Talysh (Buhse!), in districtu Airum (Kolen!), Persiae bor. prov. Aderbidjan (Auch. 5215!), in monte Elbrus prope Passgala (Ky. 428!).

Planta semipedalis vel vix procerior, folia radicalia 1½-3-pollices longa, paniculæ rami 2-3-pollicares. Characteribus et corollæ colore affinis *S. verbasci-*

foliæ, an ejus formis quoque adnumeranda sit ulterius observandum; differt canle pumilo, panicula brevis verticillis approximatis, bracteis majoribus.

72. **S. frigida** (Boiss. Diagn. Ser. I, p. 10) inferne arachnoideo-canescens partibus floralibus hirta viscosissima, canle pumilo mox in paniculam abeunte, foliis petiolatis parvis rugosis ovatis basi truncatis vel subcordatis obtuse lobato-dentatis et denticulatis, floralibus ovato-orbiculatis obtusis breviter mucronatis calyces subæquantibus vel brevioribus, verticillastris 6-4-floris remotis, floribus parvis, calycis breviter campanulati papilloso-hirti visciduli labiis aquilongis dentibus triangularibus apice spinulosis intermedio labii superioris breviore, corollæ cœruleo-violaceæ calyce duplo longioris tubo inclusa, labio superiore angusto apice attenuato modice curvato ♀. *S. oreades* Schott et Ky in Sched.

Hab. in regione alpinâ excelsiori Anatoliae occid. et australis, Cariae Cadmus supra Khonas et supra Geyra (Boiss!), Lycia in montibus supra Elmalu (Bourg!), Pisidiae mons Davrosdaghi 6000' (Heldr!), Taurus Cilicus 80° 0' (Ky. 54! Bal!), Cappadocia australis mons Masmeneudagh (Bal!).

β. oblongifolia. — Folia viridiora oblonga basi rotundata. — *S. spinulosa* Montb. et Auch. ex Bth. in D.C. Prodr. XII, p. 283!

Hab. in Tauro orientali (Auch. 1523! in D.C. Prodr.), in rupes tribus ad marginem sylvarum supra Gorumse 6000' in ditione Kassan Oghlu Tauri Orientali (Ky. 113!), Plantam Mesopotanicam (Ky. 23!) huc citatam non vidi.

Planta saepius 4-6-pollicaris sed interdum major panicula ramis elongatis 6-10-pollicaribus, folia 1 $\frac{1}{2}$ -2-pollices longa atque lata, calyx 4 lineas, corolla 7-8 longa. A formis *S. verbascifoliae* floribus saltem duplo minoribus, corollæ colore et labio superiore angusto acutinuscuso et minus falcato distincta.

73. **S. modesta** (Boiss. in Bal. exs. 1856) inferne adpresso araneoso-canescens, ad inflorescentiam patule hirta vix viscosa, caulis erectis gracilibus superne paniculatis, foliis inferioribus rugosis petiolatis a basi rotundata vel cuneata oblongis vel ovatis obtusis eroso-denticulatis interdum sublobatis, floralibus membranaceis violaceouscentibus ovatis acuminatis calyces subæquantibus, verticillastris 6-4-floris secus paniculae ramos graciles remotis summis abortivis, floribus parvis, calycis patule papillari-hirti saepe violaceouscentis labiis subdivergentibus dentibus a basi ovata longe subulato-spinulosis superiore subbreviore, corollæ violaceæ calyce vix duplo longioris tubo inclusa labio superiori abbreviato angusto vix falcato ♀.

Hab. in Cappadociæ regione subalpinâ montis Argæi in valle Kameehly Tchæf (Bal!), in Armeniâ prope Erzerum et Ispir (Huet! Calv!).

Pedalis, gracilis, folia 1 $\frac{1}{2}$ -2-pollices longa, calyx tres lineas corolla 6 lineas tantum longi. Affinis *S. frigida* sed verticillastri remoti, paniculae indumentum vix viscosum, calycis dentes longius spinulosi et superior minus brevis, præsertim insignis corollæ labio superiore brevi vix falcato.

74. **S. candidissima** (Vahl Enn. I, p. 278) caulis erectis inferne tomentellis superne in paniculam strictam minute glandulosam

abeuntibus, foliis ad partem inferiorem caulum congestis petiolatis utrinque pannosis canis rarius canescentibus a basi rotundata vel cordata ovalis oblongis vel oblongo-lanceolatis repando-crenatis dentatisve, floralibus membranaceis saepius lanatis a basi ovalata vel triangulari setaceo-acuminatis parvis, verticillastris 6-4-floris remotis supremis abortivis, calycis campanulati papilloso-asperi striati labiis divergentibus superiore dentibus apice subulatis subconniventibus tandem subrecurvo, inferioris dentibus bicus lanceolatis longe subnatis, corollæ albâ calyce triplo longiore labio superiore valde falcato inferiore flavicante, genitalibus exsertis ♀. *S. odorata* Willd. Enum. Berol. I, 43. — Jacq. fil. Eclog. fasc. II, p. 26, tab. 15! — *S. albida* Jacq. Obs. p. 10. — *S. Armena* C. Koch Linn. XXI, p. 654!

Hab. in collibus siccis, in Græcia monte Geranion supra Isthmum Corinthiacum (Stuart Mill!) et ad radices montis Veluchi (Etolia) (Nicolaidis!) rara, Anatolia in Phrygiâ prope Onchak (Ball!), Cariâ (Boiss! Pinard!), Lyciâ ad Elmalu (Bourg. 225!), Isauriâ ad Koniah (Heldr!), Galatiâ (Wied!), Cappadociâ (Auch. 4515!) et prope Talasse (Ball!), Cataonia (Haussk!), Armeniâ (Tourn!), prope Erzerum (Huet!) et Baibut (Bourg!), Assyriâ inter Mossul et Bagdad (Oliv!).

Planta 1½-2-pedalis, folia radicalia 2-3-pollices longa longe petiolata, calyx 4½ lin. longus, corolla 12-14 lin. longa, *S. odorata* Willd. specimina prorsus eamdem speciem sicut floribus tantillum minoribus.

75. ***S. crassifolia*** Sibth. (t Sm. Fl. Græc. I, 19, tab. 26) caulinibus erectis inferne tomentellis superne in paniculam brevissime glandulosam ramis paucis virgatis constantem abeuntibus, foliis utrinque canis vel canescentibus breviter petiolatis cordato-orbiculari obtusissimis acutinsculis crenulatis, floralibus parvis herbaceis sublanatis ovato-orbiculari mucronatis, verticillastris 6-4-floris dis-sitis, calycis papillosi campanulati labio superiore subbreviore subrecurvo minute tridenticulato dentibus conniventibus intermedio mutico lateralibus acutis, labii inferioris dentibus binis ovato-lanceolatis apice subnatis, corollæ violaceæ calyce triplo longioris labio superiore valde falcato, genitalibus vix exsertis ♀. *S. Cilicia* Boiss. Diagn. Ser. II, 4, p. 19.

Hab. in siccis insulae Cypri (Sibth.) et prope Kithrea (Ky. 333!), in vinetis Cilicie prope Amatcha 4000' (Ky. 355!), in Cappadociâ prope Bosante Tchâf (Ball!) et Cataoniae monte Berydag (Haussk!).

Icon Flora Græca specimen preter solitum humile exhibit, planta saepius bipedalis paniculæ ramis temibus longis, folia 1½-2½ lineas longa et lata. Ab affini *S. candidissimâ* quâcum commixta fuit egregie distincta foliorum formâ, floralibus herbaceis obtusis, calycis minoris (3½ lin. vix longi) dentibus superioribus brevissimis intermedio mutico, corollæ colore.

76. ***S. cyanescens*** (Boiss. et Bal. Diagn. Ser. II, 4, p. 19) caulinibus erectis inferne tomentosis superne in paniculam breviter papillari-glandulosam abeuntibus, foliis cano-tomentosis longe petiolatis a basi subcordata ovalis obtusiusculis crenulatis, floralibus membranaceis violaceis ovatis abrupte et breviter cuspidatis parvis

tandem reflexis, verticillastris 6-4-floris parum distantibus racemosis, calycis breviter papillosi campanulati violacei labio superiore subbreviore sursum subcurvato denticulis subulatis intermedio subtaliore, inferioris dentibus a basi late lanceolatâ subulatis, corollâ pallide violacea calyce triplo longiore labio superiore falcato, genitalibus exsertis ♀.

Hab. in collibus secus flumen Kamechly Tchâï ad radices Masmeneulagh Cappadociae (Bal!), Anatoliâ boreali prope Tokat et Mersivan (Wied!).

Planta 1-1½-pedalis, foliorum lamina 8-14 lineas longa, calyx florifer 4 lineas longus, corolla fere pollicaris; affinis *S. candidissima* et præsertim *S. crassifolia*, ab ea foliis non obtusissimis, bracteis et calyce cyaneis, calycis dente supremo acuto lateralibus latiore, etc., distincta videtur.

77. *S. oligophylla* (Auch. MSS. ex Bth. in p.c. Prodr. XII, p. 279. — Bge Lab. p. 44) glabra, caulinis a basi in ramos tenuiter virgatos fere nudos strictos iterum atque iterum trichotomos scoparie abenuntibus, foliis inferioribus cito deciduis, ramealibus paucis minutis sessilibus triangulari-ovatis crenulatis, floralibus squamiformibus, floribus secus racemos breves solitariis cernuis paucis distantibus, calycis glabri glandulosò-punctati campanulati labiis brevibus divergentibus subcrenatis, superioris dentibus ternis brevibus inferioris binis lanceolatis acuminatis, corollæ tubo inclusa ♀.

Hab. in collibus aridis prov. Ghilan (Auch. 5029!), in valle fluvii Sefirud (Buhse!).

Bipedalis ramosissima scoparia ramis numerosis tenuibus, calyx fructifer 4½ lineas longus. Species ex speciminiibus incompletis foliis inferioribus et corollâ destitutis tantum nota. *S. hypoleuca* calyce quidem affinis sed vegetazione, inflorescentia etc. diversissima.

78. *S. hypoleuca* (Bth. in p.c. Prodr. 12, p. 279) caule glabro crasso elato folioso in paniculam ramosam elongatam glabram abeunte, foliis amplis coriaceis supra viridibus subtus rugosis adpressissime canescens in orbiculariatis obtusissimis crenulatis inferioribus petiolatis cæteris cordato-amplexicaulibus, floralibus parvis membranaceis ovatis, verticillastris 4-2-floris distantibus superioribus abortivis, calycis membranacei breviter campanulati pubescentis et glandulosò-punctati labiis valde divergentibus subcrenatis superioris dentibus ternis breviter ovatis submuticis, inferioris binis ovatis muticis vel ovato-lanceolatis mucronatis, corollæ allve calyce subtriplo longioris tubo inclusa, labio superiore falcato, genitalibus subexsertis ♀.

Hab. in Persia borealis prov. Lar et prope Teheran (Auch. 456!) et 5021!, in fauce Schirdere montis Elbrus (Ky. 409!), prope Ask (Buhse!).

Mixta cum paniculâ saepè 1-2-pedali tripodalis, folia diam. 2½-4-pollicaria, apicula rotundata, calyx 3½-lineas, corolla fere pollicaris. Species foliis ampliis coriaceis et calycis labiis obtusis divergentibus fere *S. compressa* 3½-reticulata.

79. *S. limbata* (C. A. Mey. Enum. p. 86) caule glabrescente

ecto subnudo paniculato, foliis viridibus ad caulis partem inferiorem confertis rugosissimis subtus glanduloso-punctatis et sub lente asperulis superne glabris late ovatis basi cordatis erosodentatis saepe sublobatis, floralibus ovatis aeuminatis minimis deciduis, verticillastris 6-2-floris secus paniculae ramos virgatos elongatos distantibus, calycis campanulati striati nitidi papillis sparsis asperi labiis divaricatis superiore apice recurvo breviter et acute tridentalate dente medio subbreviore, inferioris dentibus lanceolatis acuminatis, corollae calyce triplo longioris albidæ labio superiore valde falcato dorso papilloso, labii inferioris lobo medie lutescente fuso-marginato tubo inclusu brevi ♀. *S. polyadenia* Boiss. Diagn. Ser. I, 5 p. 7.

Hab. in aridis saxosis ditionis Talysch (C.A. Mey! Hoh!), Persiae bor. in prov. Aderbidjan (Auch. 5206!, Szov!), inter Teheran et Tabris (Bge!), inter Isfahan et Kaschan (Buhse!), in monte Elburs (Ky. 257!) forma dentibus calycis longius subulatis), prope Kohrud (Bge!), in Armenia ad Erzerum (Huet! Calv!).

Planta 1½-2½-pedalis, folia radicalia 4-5-polices longa 3-4 lata, calyces 3 lineas, corolla 9 lineas longa. Insignis pilis muriciformibus calycis et hujus dentibus superioribus horizontaliter curvatis, facies *S. pratensis* sed calyx *Ethiopis* nec *Plethosphacez.*

80. ***S. Beckeri*** (Trautv. Act. Hort. Petrop. III, p. 276) caulinis erexit pruinoso-canescensibus in racemum elongatum simplicem vel basi ramulis paucis brevibus auctum abeuntibus, foliis a basi subcordata ovatis obtusis repandis superne glabris rugosissimis subtus adpresso niveis inferioribus petiolatis summis sessilibus, floralibus minimis membranaceis cito evanidis, verticillastris 4,2-floris remotiusculis, calycis cyanei breviter papillosi et glandulis sessilibus adspersi labio superiore recurvo dentibus binis a basi triangulari subulatis intermedio minimo, inferioris dentibus binis a basi lanceolata subulatis, corollae violaceæ calyce triplo longioris labio superiore longo falcato ♀. *S. Ruprechti* Boiss. in Sched.

Hab. ad pagos Kankro, Chäbischu et Chondodo inter Karasa et Chuschtada Daghestanie mediae 540'-6000' (Rupr!), ad Achty (Becker!).

Cum racemo semipedali sesquipedalis, foliorum majorum lamina 2½-polices longa et lata, calyx 4½ lineas corolla 12-14 longi. Ab affini *S. limbata* foliis hypoleucis, calycis colorati dente supremo minutissimo, a *S. cyanescens* quam facie refert foliis superne glabris, bracteis obsoletis etc., discrepans.

81. ***S. Sahendica*** (Boiss. et Buhse Aufz. p. 172) minute pubescens pallide virens, caulinis ascendentibus simplicibus vel inferne ramulos foliosos edentibus in racemos subsimplices abeuntibus, foliis supra glabrescentibus subtus glandulis/adspersis crenulatis inferioribus petiolatis a basi rotundata vel truncata oblongo-lanceolatis obtusis superioribus sessilibus acutis, floralibus ovato-acuminatis submembranaceis calyces subequalibus et brevioribus, verticillastris 6-4-floris inferioribus distantibus, calycis papilloso-asperi et glanduloso-punctati labio superiore subrecurvo dentibus brevibus

acutis inferiore sublongiore dentibus lanceolatis acuminatis, corolla e sicco albido tubo inclusa, labio superiore falcato, genitalibus exsertis ♀.

Hab. gregaria in prov. Aderbidjan (Auch. 5199!), ad radices montis Sahend (Bnhse!) inter Teheran et Tabris (Bge!).

Pedalis et bipedalis, folia fere *S. officinalis* sed virentia 2-2 $\frac{1}{2}$ pollicaria basi 7-9 lineas lata, racemi semipedales apice densiusculi, calyx 5 lineas, corolla 10-11 longa. Affinis *S. limbata* a qua foliis angustis et racemo simplici differt.

82. S. Chorassanica (Bge Lab. Pers. p. 45) pallide virens minute puberula et resinoso-punctata, caulinis erectis simplicibus vel superne striete et parce ramosis, foliis anguste linear-lanceolatis suberectis acutis vix rugosis basi attenuatis inferioribus breviter petiolatis ceteris sessilibus, floralibus triangularibus acuminatis calyces vix aequantibus, verticillastris 6-2-floris remotis, calycis campanulati breviter papillosi et glandulis adspersi labio superiore i dentes tres breves conniventes fisso, inferiore bifido dentibus omnibus lanceolatis subulato-spinosis, corollae calyce triplo longioris tubo brevi inclusa labio superiore falcato ♀.

Hab. in subalpinis jugi elati in ditione Khorassan inter Nichapur et Mesched siti (Bge!).

Canes sesquipollicares, folia inferiora 2-3-pollicis longa 4-6 lineas lata, verticillastra 3-6, calyx 5 $\frac{1}{4}$ lineas longus. Affinis *S. Sahendica*, foliis, verticillastris remotis, etc., diversa.

83. S. brachycalyx, caule valde clato pilosuscule superne nudo paniculato, paniculae breviter papilloso-hirtae viscidulae ramis longissimis, foliis inferioribus petiolo longo crispule hirsuto suffultis magnis tenuiter membranaceis subtus ad nervos pubescentibus ovalis basi truncatis vel late cordatis erosio-dentatis et sèpius grosse lobatis, floralibus triangularibus acuminatis deflexis inferioribus calyces aequantibus, verticillastris 6-4-floris valde distantibus, calycibus breviter pedicellatis hirsuto-viscidis breviter campanulatis labiis brevissimis latis subhorizontaliter truncatis superiore dorso bisulcato tridenticulato inferiore bidentato dentibus omnibus breviter spinulesis, fructifero nutante aucto subinflato, corollae calyce quadruplo longioris cærulescentis tubo brevissimo sub fauce abrupte ampliata squamâ latâ penicillatâ obsito cum labio inferiore purpureo et nigro maculato, labio superiore longo valde falcato, genitalibus exsertis ♀. *S. Indica* L. Sp. p. 37. — Ic. Jacq. Vindob. I, tab. 78. — Bot. Mag. tab. 395!

Hab. in rupestribus umbrosis Palestinae ad Hierosolymam (Ky. 449!), ad Beeroth (Boiss!). Antilibano (Boiss! Schub! Gaill!), Syriâ prope Aleppo (Auch 4557!), Mesopotamia ad Mossul (Auch!), Kurdistania inter Van et Ischri (Noë 155!).

Planta 3-4-pedalis, folia radicalia 4-6-pollices longa et lata, paniculae rami internum sesquipedales, flores magni cito et facile decidui, calyx florifer 3 $\frac{1}{2}$ lineas longus fructus sex lineas longus et apice latus dorso concave et late

bisulcatus quā notā ad *Plethiosphacem* accedit, corolla pulcherrima 12-15 lineas longa fauce abrupte dilatato-gibba et intus squamā latā auctā quā notā *Gongrosphaceis* recte adnumeratur. Species errore botanicorum antiquorum ex Indiā indicata et cuius ideo nomen specificum mutandum erat.

SECT. V. PLETHIOSPHACE.

84. **S. Austriaca** (L. Syst. p. 69) caule erecto breviter pubescente a basi foliosā subnudū in racemum simplicem vel paniculam parēt et breviter ramosam longe et patule hispidam abeunte, foliis fere omnibus radicalibus rosulatis expansis breviter et late petiolatis a basi truncatā vel subcordatā late ovatis duplicato-crenatis sāpē lobatis supra glabris subtus pubescentibus, floralibus ovatis breviter acuminatis herbaccis calyces aequantibus vel brevioribus, verticillastris 6-4-floris inferioribus subdistantibus, calycis subsessilis pilosissimi campanulati labiis brevibus ovatis superiore breviter tridentato inferiore in dentes ovatos mucronatos bisido, corollā ochroleucā calyce triplo longiore labiis aequilongis superiore subfalcato, staminibus corollā duplo longioribus ♀. Ic. Jacq. Austr. tab. 112. — Rehb. Ic. tab. 50.

Hab. in montosis demissioribus Tauriae (M B., Stev.).

Ar. Geogr. Austria, regio Danubialis, Rossia australis.

85. **S. staminea** (Montb. et Auch. Ann. Sc. Nat. 1836, 2, p. 41) breviter papillari-pubescentis vel araneosa pallide virens, caule erecto crassiusculo folioso in racemum simplicem aut in paniculam striete et breviter ramulosam abeunte, foliis rugosis a basi rotundatā vel cordatā oblongis obtuse dentato-lobatis inferioribus longe petiolatis obtusis supremis sessilibus diminutis acutis, floralibus minimis ovato-orbiculatis muticis vel mucronatis, verticillastris 8-10-floris in racemos approximatis, floribus parvis pedicellatis, calycis campanulati hirsuti labio superiore subbreviore subrecurvo denticulis ternis brevibus conniventibus acutiusculis intermedio minimo, inferioris dentibus binis lanceolatis acutis, corollae cærulescentis calyce 2½-plo longioris labio superiore subaequilongo angusto recto, staminibus corollā duplo longioribus ♀. *S. Austriaca* C. Koch Linn. XXI, p. 636 non L. — *S. Calverti* Boiss. MSS.

Hab. in argillosis et cunctis derelictis, Cappadocia Orientalis (Auch. 4526!), prov. Musch Armeniae 5-7000' (Ky. 366!), inter Baibut et Ispir (Huet! Bourg!), prope Erzerum (Calv!).

Pedalis et sesquipedalis, folia intermedia 4-5 poll. longa 4-1½ lata, racemi 5-8-pollicares. Species calyce postice concavo bisulcato et omni indole a sectione *Ethiopide* aliena et *Plethiosphaci* adnumeranda, affinis *S. Austriaca* a quā caule folioso, foliis elongatis angustis, racemis non lanatis, calycis dentibus muticis, corollae labio superiore angusto recto egregie differt.

86. **S. pratensis** (L. Sp. p. 35) pubescens viridis superne

glandulos-pilosa, caule erecto parce folioso in paniculam parce ramosam abeunte, foliis rugulosis superne glabris radicalibus petiolatis cordato-ovatis duplicato-crenatis interdum lobatis, superioribus sessilibus acutis acute dentatis, floribus parvis subreflexis cordato-ovatis crenipatatis, verticillastris subsexfloris disjunctis, calyeis subsessilibus pubescenti-viscidii campanulati labio superiore breviter anguste et acute tridentinato, inferioris dentibus binis a basi ovali mucronulatis, corollæ caruleæ rarius alba vel roseæ calyce triplo longioris tubo calycem aquante labio superiore longiore falcato ♀. Ic. Engl. bot. tab. 153. — Rehb. Ic. tab. 51. — *S. Barrelieri* Ten. — *S. Tenorii* Spr. — *S. dubia* C. Koch Linn. XXI, p. 637!

Hab. in pratis regionis montanae, Graecia in monte Ithome Messeniae (Heldr!), montibus Atticae (Orph!) et monte Hagios Elias Æginæ (Heldr!), Bithynia in Olympo (Auch. 1557!), Pontus (C. Koch!), Caucasus borealis (C.A. Mey.).

Ar. Geogr. Europa borealis et media a Suecia meridionali ad Rossiam medium et australem, Hispaniam borealem et medium, Italiam, regionem Danubialem.

87. *S. Hierosolymitana* (Boiss. Diagn. Ser. I, 12, p. 61) caule elato ad angulos retrorsum spinuloso-aspero cæterum glabro superne in paniculam papillosum viscidulam abeunte, foliis preter hinc caulinam sessilia radicalibus petiolo longo papilloso suffultis membranaceis viridibus parce papilloso-asperis a basi cordata late ovatis obtusis grosse lobatis lobis acutiusculæ denticulatis, floralibus minimis oblongis acuminatis, verticillastris subsexfloris distinctis, calyeis subsessilibus rubelli papilloso-glandulosi labio superiore sublongiore subrecurvo dentibus tribus subulatis subaequilongis, inferioris dentibus binis a basi triangulari-ovata longe subulatis, corollæ atrorubentis calyce triplo longioris labio superiore sublongiore incurvo ♀.

Hab. in ropestribus ad septentrionem versis Palestine (Ky. 642!), circa Hierosolymam frequens (Boiss!), Syria littoralis in Libano supra Sidonem (Gail! Bl!) et in valle Nahr el Kelb (Gail!), in Antilibano ad Ainhata (Gail!).

Plauta 1½-2-pedalis facie et florum magnit. *S. pratensis*, folia 3-4-pollicaria. Differt a *S. pratensi* foliis latioribus, in lumento aspero partis inferioris plantæ, calyeis dentibus longius aristatis, corollæ labio superiore minus falcato; a *S. virgata* et *S. viscosa* floribus magnis, etc., longius distat.

88. *S. virgata* (Ait. Kew I, p. 39) papillari-pubescentis pallide viridis, caule elato superne in paniculam longe virgato-ramosam pubescentem vel villosam vix viscidam abeunte, foliis rugulosis obtuse crenatis et lobatis inferioribus petiolatis cordatis ovato-oblongis, superioribus paucis diminutis breviter petiolatis et sessilibus acutis, floralibus minimis ovato-oblongis acuminatis saepè deflexis et coloratis, verticillastris 6-4-floris distantibus longe racemosis, floribus parvis, calyeis breviter pedicellati campanulati patule hirti labio superiore subrecurvo dentibus tribus parvis conniventibus, inferioris binis ovatis subulato-attenuatis, corollæ calyce

2-2½-plo longioris cernueæ vel albidae labio superiore falcato ♀.
Ic. Jacq. Hort. Vind. t. 37. — *S. Garganica* Tea. — *S. Sibthorpii*
Fl. Gr. I, p. 17, tab. 22. — *S. campestris* M.B. — Rehb. Icon. bot.
t. 709. — *S. nudicaulis* C. Koch Linn. XIX, p. 24 non Vahl (speci-
mina parva caule simplici!). — *S. oblonga* C. Koch Linn. XIX, p. 24!

Hab. in herbidis siccis fere totius regionis, Graecia in Laconie monte Malevo
(Orph. Fl. exs. 164!), Aetolia (Heldr. Herb. Norm. 679!), Bœotia (Sprun!),
Macedonia et Thracia (Friv! Griseb!), Anatolia omni septentrionali, occidentali
et australi (Boiss! Bourg! Pichl!) ad Ciliciam usque (Ball!), Rhodo (Bourg!),
Anatolia interiore (Wi-d!), Armeniæ (Kyt!), Syriæ boreali in Amaro (Ky. 447!),
Tauriæ (M.B.), Georgiæ Caucasica (Hohen!), Persiæ boreali (Szov., Buhse!
Ky!) et australi prope Schiraz (Ky. 667! et 593!), montibus Sawers, Eschker
et in ditione Bachtiaris (Haussk!).

Indumento polymorpha; species racemis elongatis et floribus fere triplo
minoribus a *S. prateasi* distincta.

Ar. Geogr. Italia australis, regio Danubialis, Rossia australis.

S. viscosa (Jacq. Misc. II, p. 328 Ic. rar. tab. 5), caule
i... e breviter superne densius viscoso-pubescente superne sim-
plici longe racemoso vel paniculato ramis paucis elongatis, foliis
adpresso supra parce subtus densius papillari-pubescentibus mem-
branaceis rugulosis late cordatis ovato-oblongis sinuato-repandis
infimis petiolatis obtusis intermediis et superioribus breviter petio-
latis et sessilibus acuminatis, floralibus minimis ovatis acutis, verti-
cillastris 6-4 floris valde remotis, floribus parvis, calycis subsessilis
valde papillari-glandulosi labio superiore subbreviore obtuso den-
tibus ternis minimis conniventibus sursum curvalis inferioris binis
ovatis mucronatis, corollæ roseæ vel rubræ viscosæ calyce 2½-plo
longioris labio superiore longiore subfalcato ♀. — *S. Gaillardotii*
Boiss. olim.

Hab. in regione inferiore et mediâ Libani (Ehrenb!) Auch. 4542! Bl! Gaill!).

Affinis *S. virgatae* a quâ differre videtur foliis tenuioribus præter infima
magis acuminatis, partium floralium viscositate, dentibus calycis brevius
mucronatis. Corolla eâ paulo major. Locî Italici et Hispanici ab auctoribus
indicati certe erronei.

90. **S. sylvestris** (L. Sp. 34) caulis erectis adpresso lomenti-
tellis crebre foliatis rariis in racemum simplicem sèpius in pan-
iculam strictam breviter pubescentem vel tomentellam abeuntibus,
foliis supra minute puberulis subtus caulis vel canescentibus a basi
rotundatâ oblongo-lanceolatis crenatis inferioribus petiolatis obtusis
superioribus sessilibus cordatis acute dentatis acuminatis, floralibus
parvis ovato-orbiculatis acuminatis sèpe coloratis, verticillastris
10-6-floris inferioribus vel omnibus distinctis longe racemosis, flor-
ibus parvis subsessilibus, calycis tubuloso-campanulati patule papil-
losi et glanduloso-punctati labio superiore subbreviore subrecurvo
tridentato inferiore bidentato dentibus subulato-attenuatis, corollæ
purpureo-violaceæ calyce duplo longioris labio superiore longiore

inecurvo ♀. Ie. Jaq. Austr. tab. 212. — Rehb. Icon. bot. tab. 707.
— *S. nemorosa* L. Sp. p. 35. — Rehb. Icon. bot. tab. 708.

Hab. in grammosis ad cultorum margines, Graecia in Olympo Thessalo (Auch. 4559!), Macedonia et Thracia (Friv! Orph! Hebr!), Lycia ad Elmali (Bourg 227!), Cappadocia (Bal!), Armenia (Auch! Bourg!), Tauria, Caucasus et Transcaucasia (M.B., Stev! Hohen!), Persia borealis (Buhse! Bge!) et occidentalibus prope Sihna (Haussk!).

Amiens Bge ex appendiculi pilosi tubi corollae absentia *S. nemorosa* et *S. syvestri* se pupigit sed hec nota non mihi nec sat gravis nec sat constans ex ea videtur.

Ar. Geogr. Germania orientalis, regio Danubialis, Rossia media et australis, Songaria, Sibiria Uralensis et Altaica.

91. **S. nutans** (L. Sp. p. 39) caule herbaceo pubescente submudo, paniculae racemis longe pedunculatis abbreviatis densioris ante anthesin nutantibus, foliis subradicalibus longe pedunculatis firmis rugosis subtus adpresso canescensibus a basi cordata ovato-oblongis obtusis duplicito-crenatis, floralibus minimis ovato-orbiculatis mineronatis, verticillastris 6-4-floris approximatis, calycis hirsuti labio superiore subbreviore subincurvo minute tridentato, inferiore in dentes ovatos apice subulatos bilido, corollae intense violacea calyce triplo longioris labio superiore sursum patente recto longiore ♀. Ie. W. Kit. Hung. tab. 62. — Rehb. Ic. tab. 49.

Hab. in sylvis Macedonie prope Vodena (Griseb.), in collibus Tauriae frequens (Stev!), in ditione Talysh ad Siwirs (ex G.A. Meyer).

Ar. Geogr. Regio Danubialis, Rossia media et australis.

92. **S. Verbenaca** (L. Sp. p. 35) breviter pubescens superne saepe villosula, canibus ascendentibus simplicibus vel parce ramosis, foliis ovatis et oblongis crenatis rugosis glabriusculis basi rotundatis vel cordatis indivisis lobatis vel pinnatifidis inferioribus petiolatis superioribus sessilibus amplexicaulibus cuspidatis, floralibus orbiculari-cordatis acuminatis tandem reflexis, verticillastris remotis 6-4-floris, calycis ovato-campanulati hirsuti siccis colorati labio superiore subbreviore semiorbiculato brevissime denticulata, inferioris dentibus a basi ovato-lanceolata subulatis, corollae caeruleae labio superiore latiusculo subincurvo ♀.

z. serotina Boiss. Voy. Bot. Esp. p. 484. — Caules elatiiores foliosi, folia crenata vel lobata nec pinnatifida, verticillastri remoti, corolla limbus parvus calycem saepe parum excedens. — Rehb. Icon. bot. tab. 703. — *S. oblongata* Vahl. — Rehb. Icon. bot. tab. 702. — *S. Sibthorpii* Fl. Pelop. № 37 non Fl. Græc. — *S. Spiehmanniana* M.B. Taur. Cauc. I, p. 21. — *S. disermas* Sm. Prod. non L. Floret aestate.

β. vernalis Boiss. loc. cit. — Caules humiles suaphylli, folia fere omnia radicalia profunde incisa vel pinnatifida, racemus densior, corolla major calyce duplo longior. — *S. clandestina* L. Sp. 36 non

herb. nec Fl. Græc. — Rehb. Icon. bot. tab. 704 et 705. — *S. polymorpha* Hoffm. et Link Fl. Port. tab. 19. — *S. hyemalis* Brot. Phyt. tab. 23. — *S. praecox* Savi. — *S. multifida* Fl. Gr. I, p. 17, tab. 23. — *S. horminoidea* Pourr. — *S. laciniata* Willd. Enum. Ber. Floret hyemic et vere.

Hab. in herbidis regionis inferioris utraque forma, in Græciâ et insulis vulgaris!, Macedonia (Friv!), Byzantii (Sibth.), Anatolia orientalis (Müll.), Cyprus (Ky!), Syria ad Aleppo (Haussk!) et littoralis (Bl! Gaill!), Palestina (Roth!), Ægyptus inferior (Sam!), Tauria meridionalis (Stev!), Transcaucasia ad Caspium (C.A. Mey!).

Ar. Geogr. Britannia et Belgium, Gallia occidentalis et australis, Lusitania, Hispania, Italia, Dalmatia.

93. **S. controversa** (Teñ. Syll. p. 18) ex rhizomate indurato pluricaulis pumila adpressissime canescens et pilis longis crispis plus minus hispida, caulis simplicibus vel a basi ramosis, foliis ambitu oblongis in lacinias lineares bullato-rugosissimas margine revolutas obtusissimas crenulatas divaricatas sectis lateralibus divaricatis remotis abbreviatis terminati elongatâ, floralibus parvis ovato-orbiculatis acutis, verticillastris 8-6-floris remotiusculis vel in racemos patule hispidissimos approximatis, calycis longe albo-lanati subsessilis labio superiore subrecurvo orbiculato minutissime tridentato, inferioris dentibus binis lanceolatis mucronatis., corolla rubellæ calyce duplo longioris labio superiore angusto subfalcato ♀. *S. clandestina* L. herb. nou Sp. — Fl. Græc. tab. 24. — *S. lanigera* Poir. Dict. Suppl. 5, p. 48. — *S. rugosissima* Zuce. Acad. Monac. III, p. 244, tab. 7.

Hab. in arenosis salsis Egypti ad Alexandriam (Ehr! Auch. 1535! Samar!), in valle Ouâdi Dngra deserti Arabici (Schweinf!), Arabiae petreæ (Bové! Boiss!), Cypri (Sibth., Ky!), Syriae ad Damascum (Gaill!), Mesopotamiae (Haussk!).

Adest quoque ut in *S. Verbenacâ* forma corollâ clandestinâ calycem vix vel parum superante.

Ar. Geogr. Africa borealis.

Obs. *S. ceratophylloides* L. Mant. p. 26 in Egypso a Vahllo indicata ibi non crescit.

SECT. VI. HORMINUM.

94. **S. viridis** (L. Sp. 34) annua villosa pallide virens, a collo simplex vel pluricaulis caulis erectis parum elongatis simplicibus vel 1-2 ramis auctis, foliis petiolatis a basi rotundatâ vel subcordatâ ovatis vel oblongis crenulatis, superioribus sessilibus, floribus ovato-rhombeis latis acutis persistentibus calyces subæquantibus basi utrinque bracteâ subulatâ auctis viridibus terminalibus diminutis, racemis simplicibus, verticillastris 6-4-floris subdistantibus, calyce tubuloso aspero elevatim striato labio superiore oblique truncato

dentibus binis parvis remotis, inferioris dentibus ternis a basi lanceolatâ spinulosis, fructifero aucto plicato reflexo, corollæ pallide violaceæ calyce vix duplo longioris tubo subexerto ♀. Ic. Fl. Graec. tab. 49. — Desf. Atl. tab. I. — *S. truncata* Willd. Enum. I, p. 34.

Hab. in collibus graminosis siccis regionis inferioris a Græciâ et ejus insulis!, Thraciâ (Griseb!), Byzantio (Wied!), Anatoliâ maritimâ boreali occidentali et australi! ad Syriam prope Aleppo (Ky!), Trauscanasiam (Hoh! Szov!), Persiam boreali-occid. Szov! Buhsei!.

Ar. Geogr. Hispania, Italia, Sicilia, Dalmatia, Africa borealis.

95. S. Horminum (L. Sp. p. 34) annua villosa pallide virens a collo simplex vel pluricaulis, caulinis erectis simplicibus vel 1-2 ramis auctis, foliis petiolatis a basi cuneatâ vel rotundatâ ovato-oblongis obtusis crenatis superioribus sessilibus, floralibus ovato-rhombeis latis acutis persistentibus calyces subequalibus basi utrinque bracteâ subulatâ auctis, terminalibus sterilibus membranaceis dilatatis violaceis comosis, racemis simplicibus, verticillastris 6-4-floris distantibus, calyce tubuloso aspero elevatim striato labio superiore oblique truncato dentibus binis remotis inferioris dentibus ternis a basi lanceolatâ spinulosis, fructifero aucto reflexo plicato, corollæ rubello-violaceæ calyce subduplo longioris tubo subexerto ♀. Ic. Fl. Gr. tab. 20.

Hab. in cultis et collinis regionis inferioris a Græciâ et ejus insulis! Macedonia et Thraciâ (Gris!), Byzantio!, Anatoliâ occidentali et australi! ad Syriam littoralem (Bl!), Palestinam (Boiss!), Tauriam (M.B.).

β. angustifolia. — Folia oblongo vel linearis-spathulata.

Hab. in arenosis mobilibus Syriae circa Tripoli (Bl!).

S. Horminum et *S. viridis* formis intermediis forsan conflunt.

Ar. Geogr. Italia, Dalmatia, Africa borealis.

SECT. VII. EREMOSPHACE.

96. S. Aegyptiaca (L. Sp. p. 33) brevissime canescens-pubescentia suffruticosa pumila a basi ramosissima dumulosa, ramis decumbentibus et erectis patule et rigide ramulosis intricatis, foliis paucis parvis inferioribus oblongis in petiolum brevem attenuatis superioribus linearibus sessilibus omnibus obtuse serrato-crenatis bullatis obtusis, verticillastris 4-2-floris remotis secus ramulos floribus deciduis persistentes racemosis supremis abortivis, foliis floralibus parvis persistentibus oblongis acutiusculis, floribus in pedicello brevi erecto nutantibus parvis, calycis ovati pubescentis vel hirsuti labio superiore subbreviore dorso concavo bisulcate apice subincurvo dentibus 3 minimis conniventibus, inferioris dentibus a basi trianguli-lanceolatâ subulatis, corollæ glabræ calyce vix sesquilonioris labio superiore subbreviore recto, connectivis antice ascendentibus

liberis ♂, Ic. Jacq. Vind. tab. 108. — *S. pumila* Bth. Lab. p. 726. — Jacquem. Voy. tab. 133 (nullo modo distincta). — *Melissa perennis* Forsk. Fl. p. 108. — *Thymus hirtus* Viv. Lib. 30, tab. 44. — *Thym. Syrticus* Spr. Syst. II, p. 697.

Hab. in desertis Egypti (Forsk! Auch! etc.), Arabiae petreeæ (Boiss!), Persia australis et insulæ Karak ad sinum Persicum (Ky! Auch! Haussk!), regni Cabulieci (Griff!).

Dumulus semipedalis, folia 4-6 lineas longa, racemi 2-3-pollicares, calyx florifer $1\frac{1}{4}$ lin. longus fructifer fere 3 lin. longus.

Ar. Geogr. Insulæ Canarienses, insulæ Capitis viridis, Africa borealis, Nubia littoralis, Arabia felix, India boreali-occidentalis.

97. *S. deserti* (Decaisne Flor. Sinaic p. 10) cano-tomentosa suffruticosa ramosissima dumulosa, ramis annotinis erectis pinnatis simplicibus vel parce 1-2-ramulosis, foliis parvis ovatis vel oblongis petiolatis obtusis margine revolutis crenato-bullatis, floralibus ellipticis acutiusculis, verticillastris 8-4-floris subremotis longe racemosis, calycis longe et patule hirsuto-lanati ovati labiis subæqualibus, corollæ glabrae calyce vix sesquilonioris labio superiore subbreviore erecto ♂.

Hab. in valle Ouadi Dachl deserti Aegyptiaco-Arabici (Schweinf!), in deserto Sinaitico (Bové! Auch. 415!), in rupibus montis Sinai (Schimp. 484!), in valle Ouadi Scheick (Boiss!).

Dumulus 6-9-pollicaris, foliorum majorum lamina 7-8 lineas longa $2\frac{1}{2}$ -3 lata; species affinis *S. Aegyptiacæ* a qua distinguitur tomento densi, verticillastris crebrioribus plurifloris magis approximatis et praesertim calycis longe albo-hispidi labio superiore non recurvo et dentibus omnibus præter supremum æquilongis.

98. *S. santolinæfolia* (Boiss. Diagn. Ser. I, 5, p. 43) suffruticosa ramosissima breviter et patule canescens, ramis tortuosis novellis brevibus erectis, foliis minimis subpetiolatis anguste et breviter linearibus pectinatim lobatis lobulis minimis subcontignis ovatis margine revolutis, floralibus inferioribus conformibus superioribus minimis ovatis integris, verticillastris subbifloris in racemum tenuem approximatis, floribus parvis, calycis ovati longe plumoso-hispidi tandem nutantis dentibus æquilongis triangularibus acutis labii superioris minus profunde fissis, corollæ calyce vix quartâ parte longiore ♂.

Hab. in desertis ad sinum Persicum (Auch. 5214!), in arenosis circa Dalaki (Haussk!).

Dumus semipedalis, folia cum petiolo vix tres lineas longa linea $\frac{3}{4}$ lata, calyx 2 lineas longus. Calyce plumoso affinis *S. deserti* sed distincta foliis angustissimis pinnatilobatis, verticillastris paucifloris, floribus minoribus.

99. *S. eremophila* (Boiss. Diagn. Ser. I, 5, p. 12) suffruticosa ramosissima tota patule hispida canescens, ramis novellis crebris erectis brevibus subsimplicibus crebre foliosis, foliis petiolatis minutis oblongo-linearibus brevibus in lobulos ovatos obtuses margine

revolutos pectinatim pinnatisectis, floralibus minutissimis oblongo-linearibus integris, verticillastris 6-4-floris in racemos oblongos approximatis, calycis longe albo-hispidi campanulati pedicello setaceo aequilongo suffulti labii brevibus aequilongis divergentibus obtusissimis brevissime et obtusiuscula superiore tri inferiore bidentato, corollā.... 5.

Hab. in Persia australi (Auch. 5194!), in montosis inter Yezd et Mehris et inter Kerman et Yezd, in rupibus ad septentrionem urbis Tebbes ad margines orientales deserti regni (Bge!).

Damus vix semipedalis, folia petiolo 3-5 lineas longo suffulta 3 lineas longa lineam lata lobulis valde complicato-crispis, racemi sesquipollulares plumosi, calyx 2 lineas longus et apice latus subcompressus. Calyce a praecedentibus distinctissima.

100. S. trichocalycina (Bth. in D.C. Prodr. XII. p. 336) suffruticosa ramosissima patule hispidula canescens, ramis novellis brevibus rigidulis tenuibus saepe patule ramulosis tandem persistentibus, foliis petiolatis parvis anguste linearibus in lobulos contiguos obtusos margine revolutos sinuato-pinnatisidis, floralibus minimis linear-oblengis, verticillastris paucis 2-1-floris breviter et laxe racemosis, calycis hispidi pedicello erecto-patulo dimidio breviore suffulti tubuloso-campanulati dentibus erectis a basi lanceolata longe subulatis labii superioris subbrevioribus 5.

Hab. ad rupes calcareas Jurunek regni Cabulici (Griff. 528!).

Damus 6-9-pollicaris, folia 3-6 lineas longa lineam lata, calyx fructifer 4-4¹/₂ lineas longus. Folia fere *S. santolinifolia* sed calyces tubulosi plus duplo maiores dentibus longe subulatis.

101. S. Tebesana (Bge Lab. Pers. p. 32) suffruticosa ramosissima patule et breviter pubescentis ramis crebris erectis pumilis stricte et opposite ramulosis, foliis minimis petiolatis ovato-rhombeis utrinque 2-3-dentatis suberispato-undulatis, floralibus subsessilibus integris, verticillastris bifloris laxe racemosis, calyce pedicello breviore tandem recurvo suffulto longe et patentim lanato-hispido tubuloso-campanulato dentibus ovatis breviter mucronatis labii superioris brevioribus, corollae calycem parum excedentis glabrae labio superiore subbreviore 5.

Hab. in præcipuis calcareis Persie orientalis supra Tebbes et prope Meibut inter Jezd et Ispahan (Bge!), ad Tacht i Rustam prope Ispahan (Bodel).

Damus 6-9-pollicaris, foliorum majorum lamina 3-4 lineas longa 2¹/₂ lata, calyx sub anthesi 2 lineas fructifer 3 lin. longus. Foliorum formâ et calycis dentibus breviter mucronatis a *S. trichocalycina* diversa.

102. S. macilenta (Boiss. Diagn. Ser. I, 3 p. 13) suffruticosa ramosissima glabrescens punctis resinosis crebris adspersa, ramis crebris subundis erectis strictis tenuiter virgatis, foliis paucis minutissimis petiolatis oblongo-linearibus utrinque obtuse 2-3-denticulatis, floralibus oblongis acutiusculis integris, verticillastris paucis bifloris

terminalibus remotiusculis, calycis pedicello setaceo æquilongo tandem recurvo suffulti breviter campanulati longe plumoso-hispidi labio superiore bi inferiore tridentato dentibus late et breviter triangularibus mucronulatis labii inferioris sublongioribus, corollæ glabra calyce sesquiloniore ♂.

Hab. ad radices montium regni Mascate Arabie (Auch. 5210!).

Rami tenuiter juncei 3-6-pollicares, foliorum petiolus 3-4 lineas limbis vix lineam latus, calyx sesquilineam longus Facies *Stachydis glutinosæ*; a precedentibus glabritie (calyce excepto), ramis junceis nudis, etc., distincta.

SECT. VIII. HEMISPHERE.

103. *S. verticillata* (L. Sp. p. 37) viridis breviter papillari-pubescentes, caulis elatis foliosis superne simplicibus vel paniculatis, foliis late sinuato-crenatis inferioribus petiolatis ovatis basi late cordatis saepius lyratis unijugis segmentis lateralibus auriculiformibus parvis oblongis, foliis superioribus a basi sessili cordato-amplexicauli ovatis acuminatis, floralibus parvis ellipticis caudatis deflexis, verticillastris 20-40-floris distantibus longe racemosis, pedicellis calycei æquilongis, calycis nigricantis subbilabiati hirti labio superiore breviter tridentato inferiore in dentes binos acutos profundius fiso, corollæ cœruleæ calyce duplo longioris tubo subexerto ♀. [c. Rehb. tab. 54.]

Hab. in sterilibus ad vias, Bœotia (Bth.), Macedonia et Thracia (Griseb.), Byzantium et Anatolia borealis (Wied!), Pisidia ad lacum Egirdir (Heldr!), Cilicia ad Gülek Boghaz (Ky! Bal!), Armenia Turcica (Bourg! Ky. 332!), mons Gara Kurdistanæ (Ky. 355!), Tauria, Caucasus et Transcaucasia (Led., Rupr! etc.), Persia borealis (Buhse! Bge!).

Ar. Geogr. Hispania media et Gallia (ibi forsitan non spontanea), Germania, Italia, Dalmatia, regio Danubialis, Rossia media et australis.

104. *S. Peloponnesiaca* (Boiss. et Heldr. Ser. I, 7, p. 47) indumento adpresso sericeo-canescens, caulis elatis foliosis superne paniculatis, foliis crenato-dentatis inferioribus petiolatis pinnatipartite-lyratis lobis utrinque 2-3 triangulari-ovatis basi confluentibus terminali submajore cuneato-oblongo, foliis superioribus sessilibus triangulari-oblongis acutis basi saepe lobatis, floralibus parvis ovato-acuminatis, verticillastris 40-30-floris distantibus longe racemosis, pedicellis calycei æquilongis, calycis hirti bilabiati labio superiore sublongiore subrecurvo tridentato Inferiore bidentato dentibus omnibus acutis, corollæ violaceæ calyce 2 1/2-plo longioris tubo breviter exerto ♀.

Hab. in Messeniâ prope Pylos et in regione inferiori montis Ithomes in fruticetis (Bory! Heldr!), ad Kalamata (Heldr!), in Ætolia ad Missolunghi (Heldr!).

A *S. verticillata* indumento adpresso sericeo-lanato, foliis pinnatisectis segmentis minus inæqualibus et terminali non cordato, calyce evidenter bilabiato distincta.

105. **S. napifolia** (Jacq. Hort. Vindob. II, p. 51, tab. 152) viridis pilis papillaribus longis patulis viscidis ad caulem et petiolos crebre ad folia sparsim obsita, caulis erectis superne paniculatis, foliis suberenatis inferioribus petiolatis lyratis segmento superiore ovato-subcordato obtuso lateralibus utrinque 1-2 parvis auricula-formibus saepe conniventibus, superioribus a basi sessili cordato-amplexicauli ovato-oblongis acutiusculis, floralibus oblongis acutis deflexis minutis, verticillastris 12-6-floris remotis, pedicellis calyce brevioribus, calycis bilabiati hirti labio superiore subrecurvo dentibus ternis brevibus intermedio minimo, inferioris dentibus binis lanceolatis acutis, corollæ violaceæ calyce vix sesquilonioris tubo inclusu φ .

Hab. in insulâ Cos (Urv!), Lydiâ ad Trianda (Boiss!), et ad Smyrnam (Boiss! Bal. 313!), circa Tokat et Safranbol (Wied!), Ciliciâ littorali ad Mersina (Bal.).

Valde affinis *S. verticillatae* a qua distinta videtur pilis longioribus viscidis caulis et petiolorum, foliis obtusioribus, verticillastris paucifloris et præsertim calyce evidenter bilabiato labio superiore subrecurvo, dentibus acutioribus. Flores quidquam majores.

106. **S. Judaica** (Boiss. Diagn. Ser. I, 42 p. 61) viridis brevissime et sparsim papilloso-asperula, caulis erectis superne stricte paniculatis, foliis rugoso-bullatis repando-crenatis radicalibus petiolatis lyrato-pinnatipartitis segmentis ovatis obtusis lateralibus utrinque 4-2 interdum confluentibus terminali majore vix cordato, foliis superioribus sessilibus cordato-ovatis, floralibus membranaceis oblongis acuminatis, verticillastris 12-6-floris distantibus longe racemosis, pedicellis calyce brevioribus, calycis rubelli scabridi bilabiati labio superiore sublongiore tridentato inferioris dentibus binis tenuioribus subulatis, corollæ violaceæ calyce sesquilonioris tubo inclusu φ .

Hab. in Palestinæ montibus circa Hierosolymam (Boiss!), ad Hebron (Ky. 794!), in Syriâ littorali circa Sidonem et Amrith (Gailli), Tripolin (Bl!), Laodiceam (Post!).

A *S. verticillata* et *napifolia* foliis rugoso-bullatis, indumento brevi scabro distincta.

107.. **S. Russelii** (Bth. in D.C. Prodr. XII, p. 357) adpressissime canescens, caulis erectis virgatis simplicibus vel stricte paniculatis, foliis tenuiter rugosissimis crenulatis inferioribus petiolatis a basi cuneatâ oblongo-lanceolatis obtusis indivisis vel petiolo rarius lobulis paucis aucto, superioribus sessilibus, floralibus parvis deflexis oblongo-cuspidatis, verticillastris 30-20-floris remotis longe racemosis, pedicellis calyci subæquilonigis, calycis breviter bilabiati dentibus omnibus breviter triangularibus acutis, corollæ violaceæ calyce vix sesquilonioris tubo inclusu φ . *S. sinapifolia* G. Koch Linn. XXI, p. 659.

Hab. in Mesopotamiâ inter Orfa et Suerek (Ky. 402!), Syriâ ad Aleppo (Russ.), in montibus Ssöffdag et Berytdagh Cataonie 3500'-6000' (Haussk!),

Cappadociæ ad Euphratem (Auch!), valle Teng prov. Musch Armeniae (Ky. 1684!).

A præcedentibus foliis angustis rugosis, calycis dentibus brevibus aenatis sed non mucronatis distincta.

Species non sat note prob. præcedentium synonyma.

S. Lorentei (Hochst. Flora 1846, p. 28. Sect. *Ethiopis*) lanatovillosa, foliis inferioribus petiolatis ovato-lanceolatis basi lobatis superne dentatis, paniculâ divaricata, verticillastris remotis sexfloris, foliis floralibus cordatis acuminatis coloratis, calycis glanduloso-vilosoi bilabiati dentibus subspinosis, corollæ calyce subdupo longioris labio superiore angusto parum incurvo ♀.

Hab. prope Diarbekir Mesopotamiae (Lorent).

Ex loco *S. spinosa* vel *S. Palestina*.

S. Linkiana (Ræm. et Sch. Syst. Mant. I, p. 182) Planta minime Orientalis ex speciminibus cultis tantum descripta, probabiliter proles hybrida specierum e sect. *Eusphace*. == *S. pilantha* Link Enum. I, p. 40, Vid. sp. cult. in herb. Kew.

ROSMARINUS (L. Gen. 38).

Calyx campanulatus bilabiatus labio superiore concavo minute tridentato inferiore bilido, fauce nudâ. Corolle tubus brevis exseritus intus nudus in fauē ampliatus, limbi labium superius erectum angustum bifidum inferius patens trilobum lobo medio maximo concavo pendulo. Stamina fertilia bina antica sub labio superiore ascendentia exserta recurva filamentis cum connectivo continuis infra medium dente tenui parvo reflexo instructis, antheræ inter se cohærentes loculo linearie ad apicem connectivi dorsifixo. Stylus bifidus. Nuculae ovatae lœves. — Frutices.

1. R. officinalis (L. Sp. 33) fruticosus remosissimus ramis erectis crebre foliosis, foliis persistentibus coriaceis linearibus margine revolutis viridibus subtus canescensibus, floribus sessilibus paucis oppositis approximatis in racemos breves axillares dispositis, bracteis minutis tomentellis lanceolatis caducis, calyce pubescente albido, corollâ cæruleâ ♂. Ic. Fl. Gr. tab. 14.

Hab. in collibus calidis regionis inferioris in Oriente rarius, Zacynthus (Marg.), Bœotia (ex Sibth.), Macedonia et Thracia (ex Griseb.), insula Melos (Sibth.), Cilicia littoralis prope Mersina in pinetis (Bal!).

Ar. Geogr. Europæ regio mediterranea a Gallia australi ad Dalmatiam, Africa borealis.

T. V. NEPETÆ.

NEPETA (L. Gen. 710).

Calyx tubulosus rectus vel incurvus ore recto vel obliquo et tunc dentibus superioribus sublongioribus minus profunde fisis. Corollæ tubus exannulatus in fauce ampliatus inclusus vel exsertus, labium superius planum rectum bifidum, inferioris lacinia media rotundata valde concava. Stamina 4 sub labio superiore parallela antrorsum curvata inferiora breviora, antheræ biloculares loculis divergentibus vel divaricatis, nuculæ læves, malleatæ, rugulosæ vel tuberculatæ. — Herbae perennes rarius annuae. — Genus ob specierum ingentem numerum et characterum essentialium deficientiam totius ordinis difficillimum et nimis artificialiter in series dispositum.

Specierum Orientalium dispositio.

SECT. I. EUNEPETA. — Flores omnes hermaphroditi. Corollæ labia parum inæqualia.

SERIES I. — Perennes.

+ Nuculæ læves.

§ 1. **Spicatæ** Bth. — Verticillastra in spicas basi interdum interruptas condensata. Calycis fauce glabri os non obliquum, dentes recti subulati. Nuculæ læves.

N. podostachys, subincisa, pubescens, raphanorrhiza, Maracandia.

§ 2. **Capituliferæ** Bth. — Verticillastra in capitula terminalia ovata vel globosa condensata. Calycis fauce glabri os obliquum, dentibus anticis profundius fisis. Nuculæ læves. — Herbae villosæ ramis fragilibus.

N. layopsis, cephalotes, prostrata, floccosa, lasiocephala.

§ 3. **Catariæ.** — Verticillastra spicas sepius interruptas formantia, calycis fauce glabri os obliquum dentibus anticis profundius fisis. Corollæ tubus inclusus vel breviter exsertus. Nuculæ læves.

N. supina, crispa, menthoides, chionophila, Cataria, Bodeana, ruderalis, rugosa, crinita.

§ 4. **Longifloræ.** — Cymæ paucifloræ saepius laxæ. Calycis fauce glabri os obliquum. Corollæ tubus calyce 2-4-plo longior. Nuculæ læves.

N. lamiifolia, longiflora, teucriifolia, Calverti, gracilescens, Trautvetteri, brevifolia, macrosiphon, leptantha, leptoclada.

§ 5. **Psilonepetæ** Bth. — Cymæ paucifloræ laxæ. Calyx fauce pilosus. Corollæ tubus longe exsertus. Nuculæ læves.

1. Calyx ore rectus dentibus æqualibus.

N. sessilifolia, laxiflora, oxydonta, depauperata.

2. Calyx ore subobliquus dentibus inferioribus profundius fissis.

N. denudata.

+ + Nuculæ malleatæ, tuberculatæ vel serobiculatæ.

§ 6. **Macrostegiæ.** — Verticillastra densiflora remota. Bractæ ovatæ vel oblongæ. Calyx fauce glaber.

1. Calyx ore obliquus.

N. glomerulosa, juncea.

2. Calyx ore rectus.

N. Scordotis, Sibthorpii, leucostegia.

§ 7. **Stenostegiæ.** — Verticillastra densiflora vel laxiuscula. Bractæ lanceolatæ vel linearisubulatæ. Calyx fauce glaber.

1. Calyx cylindricus ore rectus dentibus a basi lanceolatæ subulatis.

N. Cadmea, Isaurica, camphorata, Parnassica, Spruneri.

2. Calyx ore subobliquus.

a. Corollæ tubus calyce non longior.

N. Orientalis, Cæsarea, viscida, Tmolea, cyanea.

b. Corollæ tubus calyce 1 $\frac{1}{4}$ -2-plo longior.

N. aristata, septemcernata, Persica, Kotschyii, betonicæfolia, glabrescens concolor, grandiflora, speciosa, melissæfolia, racemosa, Mussini, marifolia stenantha, Ruprechtii.

c. Corollæ tubus calyce triplo longior.

N. Cilicica, glomerata, curviflora.

3. Calyx ovatus ore rectus dentibus linearis-lanceolatis.

N. nuda, marrubioïdes.

SERIES II. — Annuæ.

§ 8. **Micranthæ.** — Calycis os obliquum. Affines N. *Mussini* et *cyanæa*.

N. humilis, *micrantha*, *saccharata*.

§ 9. **Micronepetæ.** — Calyx ore rectus.

N. petrava, *hymenodonta*, *Ispahanica*, *Schiraziana*, *pungens*, *satureioides*, *Daenensis*, *bracteata*, *globiflora*, *spathulifera*.

SECT. II. OXYNEPETA Bth. — Gen. *Oxynepeta* Bge Lab. pag. 58. — Flores dimorphi, centrales cymularum præcociores forminei antheris effetis, laterales paulo maiores submasculi. Calyx ore æqualis. Corollæ labium superius abbreviatum. — Herbae perennes paniculatae cymis 2-3-tomis.

N. Ucranica, *heliotropifolia*, *curvidens*, *calycinæa*, *cryptantha*.

SECT. I. EUNEPETA.

§ 1. **Spicatæ.**

1. **N. podostachys** (Bth. in D.C. Prodr. 12 p. 372) glabrescens vel puberula, caulis a basi decumbenti erectis elatis stricte ramosis, foliis parvis breviter petiolatis oblongo-lanceolatis acutis serratis basi rotundatis, spicis longe pedunculatis oblongo-cylindricis basi interdum interruptis, bracteis linearisubulatis calyces æquantibus, calycis hispidi dentibus subulato-lanceolatis tubo sublongioribus corollæ tubum æquantibus, corollæ limbo minuto, nuculis levibus ♀.

Hab. in regno Cabulico ad Kalou et in ascensu ad jugum Onnnye (Griff.)

Caules 1¹-2-2-pedales, folia inferiora vix pollicaria, spicæ 4-2-pollicares.

2. **N. subincisa** (Bth. in D.C. Prodr. XII. p. 373) breviter asperulo-pubescent multicaulis, caulis erectis elatis stricte et parce ramosis parce et minute foliosis, foliis breviter petiolatis ovato-lanceolatis basi breviter cuneatis subinciso-dentatis, spicis pedunculatis breviter ovalis densifloris basi saepe interruptis, bracteis lanceolatis et subulatis calyces æquantibus, calycis glabriuscui dentibus plumoso-hispidis subæqualibus tubo subæquilonis corollæ tubo subbrevioribus ♀.

Hab. in regni Cabulici valle alpina Erak (Griff. 514!). Vid. in herb. Kew.

Pedalis et elatior, folia remota vix pollicaria.

3. N. pubescens (Bth. in D.C. Prodr. XII, p. 373) breviter et crispule puberula canescens, caulis clatis ascendentibus ramulosis, foliis petiolatis ovato-oblongis basi obtusis regulariter dentatis, spicæ supra folia suprema diminuta brevissime pedunculata oblonga, bracteis lanceolatis calyce subbrevioribus, calycis molliter pilosi ore valde obliqui dentibus valde inæqualibus triangulari-lanceolatis tubo brevioribus, nuculis levibus ♀.

Hab. in montibus regni Cabulici prope Puschut (Griff!).

A *N. podostachyde* foliis majoribus longe petiolatis, indumento, calyce distincta.

4. N. raphanorrhiza (Bth. Lab. p. 734) rhizomate tuberoso oblongo, caulis pluribus ascendentibus clatis pilis papillaribus crassis sparsis obsitis inferne parce ramosis, foliis breviter papillari-hispidulis petiolatis a basi cuneatâ ovato-oblongis obtusiusculis grosse crenato-serratis, spicis breviter pedunculatis tenuibus basi stepe interruptis foliis supremis diminutis sussultis, bracteis oblongo-lanceolatis et lanceolatis calyces subæquantibus, calycis breviter pubescentis dentibus lanceolato-subulatis subinæqualibus tubo parte dimidiâ brevioribus, corollæ tubo breviter exerto limbo minuto ♀.

Hab. in regno Cabulico (Griff!).

Pedalis et sesquipedalis, rhizoma magnitudine nucis.

Ar. Geogr. Regio Himalaica.

5. N. Maracandia (Bge Pl. Lehm. 434) glabriuscula ascendens ramosa, foliis caulinis anguste oblongis basi cuneatis utrinque acutatis acute et grosse pauciserratis, ramealibus linearis-oblongis utrinque 1-2-dentatis, spicis ovato-oblongis densissimis pubescentibus, bracteis oblongo-lanceolatis pungenti-acuminatis calycem æquantibus, bracteolis subulatis, calycis ore æqualis dentibus tenuiter subulato-acuminatis ciliolatis tubum suum æquantibus, corollæ tubo inclusu, nuculis levibus nitidis ♀.

Hab. in cacumine montis Karatau supra Samarkand Turkestaniæ (Lehm!). Non vidi.

Pedalis, folia caulina vix pollicaria 3-4 lineas lata, spica vix pollicaris, calyx $3\frac{1}{4}$ lineas longus, corolla glabriuscula 5 lineas longa lobo inferiore latissimo bilobo. Ab aequalibus ex cl. Bge foliis basi cuneato-angustatis nec cordatis vel truncatis distincta.

§ 2. Capituliferæ.

6. N. lagopsis (Bth. in D.C. Prodr. XII, p. 379) tota pilis longis flexuosis albo-hispida a collo ramosissima ramis fragilibus brevibus erectis et diffusis, foliis parvis breviter petiolatis ovatis obtusis grosse et obtusiuscula dentatis, capitulis ovatis albo-hispidissimis foliis supremis diminutis involucratis, bracteis subulatis

calyces aequantibus, calycis dentibus longe lanceolato-subulatis tubo longioribus corollae subaequifloris ♀.

Hab. in regno Cabulico (Griff. N° 494 in herb. Kew!).

Rami semipedales foliati, folia 4-6 lineas longa, spicæ vix pollicares, facies *N. prostrata*.

7. *N. cephalotes* (Boiss. Diagn. Ser. I, 3 p. 20) albo-tomentosa pluricaulis caulis basi induratis ascendentibus parte inferiori foliosis superiori nudis simplicibus, foliis supra tomentosis subtus niveis parvis crassiunculis a basi truncatâ ovato-orbiculatis crenatis rugulosis inferioribus petiolo aequilongo suffultis superioribus parvis subsessilibus, floribus in capitulum longe pedunculatum globosum aggregatis, bracteis linearis-lanceolatis acuminatis calyceem aequantibus, calycis hirsuti recti dentibus lanceolato-subulatis tubo subbrevioribus inferioribus multo profundius fisis, corollæ calyce subduplo longioris tubo exerto, nuculis oblongis subcompressis levibus ♀.

Hab. in argillosis Persiae (Auch. 5141!).

Fere pedalis, folia majora sex lineas longa lataque, capitulum pedunculo longo minute bibracteolato suffultum nuce subminutus, calyx 3½ lineas, corolla 5-6 lineas longa. Species elegans foliis parvis crassis subtus niveis, capitulo solitario globoso insignis.

8. *N. prostrata* (Bth. in L.C. Prodr. XII, p. 380) laxe lanata, caulis e rhizomate crasso corticoso tenuibus prostratis fragilibus brevibus simplicibus, foliis petiolatis minutis inferioribus orbiculatis cito evanidis, floralibus ovatis verticillastri multo brevioribus, verticillastris globosis densis præter summa bina approximata distinctis, bracteis linearibus lanatis exterioribus calyces subaequantibus, calycis pedicellati sublanati subincunyi dentibus superioribus breviter lanceolatis tubo quadruplo brevioribus inferioribus profundius fisis, corollæ calyce vix sesquifloris tubo vix exerto limbo parvo, nuculis minimis ovatis laevibus ♀.

Hab. in Persia prope Ispahan (Auch. 4741!), prope Aghda inter Yezd et Ispahan (Bge.).

Caules 4-7-pollicares, folia inferiora diametro 4-6 lineas lata, verticillastri avellanæ sæpius minores, calyx tres lineas, corolla 4 lin. longa.

9. *N. floccosa* (Bth. Lab. p. 736) tota lanâ albâ floccosâ vestita caulis decumbentibus vel ascendentibus divaricatim ramosis, foliis inferioribus longiusculæ caeteris breviter petiolatis cordatis suborbiculatis obtuse repando-crenatis, floralibus squamiformibus breviter linearibus, verticillastris capituliformibus globosis oppositis inferioribus longe superioribus brevius pedunculatis, bracteis linearibus hirsutis calyces aequantibus, calycis lanati ore obliquo dentibus breviter lanceolatis inæqualibus tubo 4-5-plo brevioribus, corollæ tubo non exerto limbo parvo, nuculis anguste ellipticis subcompressis levibus ♀.

Hab. in regno Cabulico (Griff. 470 ex Bth.).

Caules nunc humiles nunc 1-2-pedales, indumentum fere et folia *N. cephalotis* sed tenuius et subtus minus nivea, capitula in squamarum axillis opposita pedunculata nec unicum solitarium terminale, calycis (3 lin. longi) os obliquus. Specimina Tibetica descripti.

Ar. Geogr. India boreali-occidentalis, Tibetia occidentalis.

10. *N. lasiocephala* (Bth. in D.C. Prodr. XII, p. 380) laxe lanata pallide virens, ramis e rhizomate corticoso indurato numerosis fragilibus tenuibus nanis procumbentibus cespitosis basi foliatis, foliis parvis petiolatis ovato-orbiculatis crenatis, floralibus sessilibus capitulo brevioribus, floribus in capitulum terminale globosum pendunculatum conglobatis, bracteis numerosis lanatis linearis-setaceis calyces æquantibus, calycis longe tubulosi crispo-lanatuli striati dentibus lanceolato-subulatis tubo quintuplo brevioribus inferioribus binis multo profundius fissis, corollæ hispidulæ calyce sesquialongioris tubo calyce vix longiore, nuculis levibus ♀. *Melissa teucrioides* Boiss. Diagn. Ser. I, 7, p. 49. — *Dracocephalum teucrioides* Bge Lab. Pers. p. 59.

Hab. ad fontem Dcheschme Pias et in cacumine montis Kuh Daëna Persiae australis (Ky. 757!).

Rami tenuis 2-3-pollicares, folia 4-5 lineas longa, capitulum avellanæ submajus, calyx 5 lineas corolla 7 longa. Facies *Calamintha origanifolia* vel *Dracocephali Aucheri* et a cl. Bunge hinc generi adnumerata sed ob corollæ labium superius non fornicatum omnino *Nepeta*.

§ 3. Catariæ.

11. *N. supina* (Stev. Mem. Soc. Nat. Mosq. III, p. 2) adpresso papillari-pubescentis, caulis humilibus ascendentibus basi saepius ramosis, foliis subsessilibus parvis ovatis basi breviter cuneatis crenatis, spicæ paucifloræ ovatæ verticillastro inferiore 2-3-floro interdum subdistante, bracteis lanceolatis calyce subbrevioribus, calycis hirsuti subbilabiati ore obliquo dentibus lanceolato-subulatis inferioribus subbrevioribus tubo 2¹/2-plo brevioribus superioribus altius connatis, corollæ tubo tenui curvato breviter exerto, nuculis oblongis levibus ♀.

Hab. in regione alpina Caucasi orientalis, ad torrentem Chodja (Stev.), in alpe Tufandagh 9000' (C.A. Mey!), in Tuschetia, ditione Tindal et Daghestaniæ australi 6000'-10000' (Rupr!).

Caules plures semipedales vel sublongiores, folio majora 7-8 lineas longa, spica brevis, calyx 4 lineas, corolla 10 lin. longa ea *N. betonicifolia* et *grandiflora* major.

12. *N. crispa* (Willd. Sp. III, p. 50) tota papillis crispulis ramosis canescens, caulis ascendentibus flexuosis ramosis, foliis petiolatis a basi subcordatæ vel truncatæ ovatis obtusis rugulosis

margine crispulis et aeniuseule inciso-crenatis, floralibus inferioribus cuneatis flores æquantibus, verticillastris sessilibus densifloris inferioribus 1-2 dissitis cæteris oblongo-capitatis, bracteis subulatis calycinum tubum subaequantibus, calycis pilis ramosis hirti recti ore obliquo dentibus a basi lanceolatâ setaceis tubo quadruplo brevioribus superioribus longioribus, corolla cœrulea tubo breviter exerto, nuculis lœvibus ♀.

Hab. in saxosis graniticis montis Elvend Persiae occidentalis (André, Auch-1750!, Haussk!).

Pedalis, folia majora pollicaria, calyx $3\frac{1}{2}$ lineas, corolla 6 lineas longa Species indumento pilis ramosis constante insignis.

13. N. menthoïdes (Boiss. et Buhse Aufz. p. 174) tota dense et adpresso tomentosa cana, caulinibus ascendentí-erectis simplicibus et ramosis, foliis breviter petiolatis et sessilibus cuneatis oblongis acutiuscula denticulatis secus nervum subuplicatis et sepe subrecurvis, floralibus oblongo-linearibus acutis inferioribus flores superantibus, verticillastris sessilibus paucifloris inferioribus 1-2 remotis cæteris in spicam angustam oblongam approximatis, bracteis linearilanceolatis flores æquantibus, calycis rubelli parce lanati brevis subincurvi ore obliqui dentibus triangulari-lanceolatis acuminatis tubo triplo brevioribus, corolla calyce vix duplo longioris tubo inclusa in limbum horizontaliter curvata, nuculis ovato-trigonis levibus ♀.

Hab. in regione alpinâ montis Ssavalan Persiae borealis usque ad 11000 (Bubse! Seidl!), in alpinis prov. Khorassan inter Nichapur et Meschhed (Bge!).

Caules fere pedales, folia majora pollicem longa 4-5 lineas lata, calyx 4 lineas longus; facies *Mentha sylvestris* Species distinctissima facie quodammodo affinis *N. crispa* indumento pulverulento, foliis cordatis, corolla tubo exerto, etc. diversissime.

14. N. chionophila (Boiss. et Haussk) multicaulis tota pubescens crispulâ brevissimâ adpresso cinerascens, caulinibus erectis firmis inferne ramosis, foliis ovato-oblongis breviter petiolatis basi cuneatis vel rotundatis obtusis obtuse crenato-repandis, floralibus linearilanceolatis sessilibus integris subrecurvis, verticillastris distantibus racemoso-spicatis, cymis densifloris inferioribus pedunculatis, bracteis linearibus calyce duplo brevioribus, calycis pilis crispulis ramosis brevibus obsiti ore obliquo dentibus triangulari-lanceolatis tubo triplo brevioribus, corolla hirtæ tubo valde curvata breviter exerto, nuculis oblongis trigonis levissimis ♀.

Hab. gregatim ad nives montium Kellal et Ssebsekul Persiae austro-occidentalis (Haussk!).

Bipedalis, folia inferiora 8-10 lineas longa, calyx circiter 3 lineas, corolla 6 longa. Species pulchra affinis *N. menthoïdi* a qua differt indumento tenuissimo, corolla tubo exerto.

15. N. Cataria (L. Sp. p. 796) breviter pubescenti-cauescens, caulinibus elatis ramosis, foliis petiolatis ovatis cordatis acutis profunde

et acutiuscula dentatis subtus magis canescens, racemis densifloris oblongis basi saepe interruptis et subramosis, bracteis subulatis calycis tubum subequantibus, calycis pubescentis ovali ore obliqui dentibus subincurvis a basi lanceolatâ subulatis tubo subbrevioribus superioribus sublongioribus, corolla hirtæ albidae rubello-punctatæ tubo inclusa, nuculis ovatis subeompressis levibus ♀. Ic. Fl. Dan. tab. 580. — Rehb. tab. 41.

Hab. in dumosis, ad saepes et in underatis regionis praesertim montane et subalpine, Græcia in montibus Kyllene et Olympo Thessalo 3-4000' (Heldr!), Parma 2.000' (Orph. Fl. Gr. exs. 551!), Macedonia et Thracia (Griseb.), Byzantii (Sibth.), regio Taurico-Caucasica (Ledeb., Rupr!). Persia borealis (Buhse, Bunge!), Kurdistanie Persiae montes Avroman et Awiheng (Haussk!), Libanus (Auch!), regnum Cabulicum (Griff!).

Ar. Geogr. Europa fere tota a Scandinavia ad Rossiam, India boreali-occidentalis.

16. N. Bodeana (Bunge Lab. Pers. p. 53) breviter pilosa pallide virens, caulinis erectis acute quadrangulis stricte foliosis, foliis sessilibus cordato-oblongis crebre dentatis superioribus acutiusculis, verticillastris confertis interrupte spicatis spicâ terminali stricte ramulosa, bracteis oblongo-linearibus subulato-acuminatis calycem aequalibus, calycis rectiuseuli villosuli dentibus subaequalibus tubum aequalibus, corolla tubo inclusa, nuculis levibus ♀.

Hab. prope Djehanuameh in declivitate australi montis Elbrus ad orientem urbis Teheran (Bode!).

Pedalis vel elatior, folia inferiora sesquipolligaria, spica racemis brevibus breviter pedunculatis strictis constans. Ex affinitate *N. Catariae*, ex specimine pessimo descripta et ulterius indaganda.

17. N. ruderale (Hamilt. ex Bth. in Wall. Pl. As. rar. I, p. 64), adpresso et breviter pubescenti-canescens pluricaulis, caulinis elatis ramulosis, foliis petiolatis ovatis obtusis crenatis basi truncatis vel subcordatis racemis simplicibus subsecundis, cymis multifloris densiusculis inferioribus pedunculatis, bracteis subulatis calyce brevioribus, calycis asperuli costati ore subobliqui dentibus lanceolatis acutis rectis superioribus longioribus sed tubo brevioribus corollæ albidae tubum aequalibus, limbo minuto ♀. *Glechoma erecta* Boxb. Fl. Ind. 3, p. 7. — *Thymus nepetoides* Don Prod. Nep. pag. 113. — *N. clinopodioides* et *N. calamithoides* Bth. in D.C. Prodr. XII, p. 382.

Hab. in demissioribus regni Cabulici (Griff. in herb. Kew!).

Ar. Geogr. India.

18. N. rugosa (Bth. in D.C. Prodr. XII, p. 382) adpresso canescens e rhizomate indurato multicaulis, caulinis tenuiter virgatis elatis simplicibus vel inferne parce et stricte ramosis et foliatis in racemos interruptos tenues longos abeuntibus, foliis minutis rugosissimis minute crenulatis petiolatis ovatis vel oblongis basi cunea-

tis vel subcordatis, superioribus oblongo-lanceolatis, cymis paucifloris inferioribus subpedunculatis, bracteis oblongis membranaceis acutiusculis imbricatis calyce triplo brevioribus, calycis breviter tomentelli elevatim costati subobliqui subincurvi dentibus triangularibus acutiusculis tubo quintuplo brevioribus, corollæ tubo breviter exerto, nuculis levibus ♀.

Hab. in regno Cabulico prope Yonutt 11000' (Griff!).

Bipedalis, folia 4-6 lineas tantum longa, racemi tenues semipedales, facies et characteres *N. juncea* sed ob semina laevia inter *Cataria* aptius collocanda.

Ar. Geogr. Arabia felix (Botta ex Bth.).

19. *N. crinita* (Montb. et Auch. Ann. Sc. Nat. 1836, p. 46) breviter pubescenti-canescens, caulis erectis gracilibus subsimplicibus, foliis breviter petiolatis parvis subcordatis ovatis obtusis erenatis, floralibus oblongis et lanceolatis aristato-acuminatis calyces æquantibus, verticillastris paniculatis inferioribus remotissimis superioribus approximatis, bracteis subulatis strigosulis calyces æquantibus, calycis strigulosi ore subobliqui dentibus a basi lanceolata longe subulato-criniferis tubo longioribus inferioribus profundius fllsis, corollæ hirtæ tubo calyce non longiore ♀.

Hab. in monte Akdaghi Cataoniae (Auch. 1726!).

Pedalis, folia majora 7-9 lineas longa, verticillastri inferiores 2-3-pollices distantes, calyx 3½ lineas, corolla 5 longa. Planta bracteis et dentibus calycis eriniformi-attenuatissimis insignis. Nuenæ juniores tantum notæ laeves esse videntur.

§ 4. Longifloræ.

20. *N. lamiifolia* (Willd. Enum. p. 602) tota pilis albis adpressis papillaribus cinerascens, caulis e rhizomate corticoso prostrato-ascendentibus brevibus parce et patule ramulosis, foliis parvis breviter petiolatis ovato-orbiculatis basi truncato-subcordatis obtusis dentibus ovatis obtusis. floralibus minimis subsessilibus. cymis breviter pedunculatis 3-5-floris insimis distantibus cæteris in capitulum brevem ovatum approximatis, bracteis longe setaceo-acuminatis tubum calycis subæquantibus, calycis longe et patule albo-papillosi ore valde obliquo dentibus a basi lanceolata longe acuminatis tubo quadruplo longioribus, corollæ tubo incurvo calyce subtriplo longiore sensim ampliato, nuculis.... ♀. *Cataria Orientalis* *ninima* *lamiifolio flore longissimo* Tourn. Cor.

Hab. in Armeniâ (Tourn!). Vid in herb. Willd!

Caules cum inflorescentiâ vix semipedales, folia majora 4-5 lineas longa, calyx 3 lineas, corolla 10-11 longi. Nulla cum *N. brevifoliâ* C.A. M. affinitas. Inflorescentiâ subcapitata a cæteris *Longifloris* distincta et facie *N. prostratanam* vel *crispam* referens.

21. *N. longiflora* (Vent. Hort. Cels. p. 66, tab. 66) adpres-

sissime canescens, caulis ascendentibus gracilibus virgatis paniculatim et laxe ramosis superne denudatis, foliis crassiusculis breviter petiolatis ovatis subcordatis obtusis obtuse cordatis, superioribus minutis sessilibus oblongis floralibus bracteisque subulatis minimis, cymis longe et tenuiter pedunculatis 3-5-floris, floralibus pedicellatis, calycis longe tubulosi asperuli incurvi ore obliquo dentibus lanceolatis acutis tubo 4-5-plo brevioribus, superioribus latioribus anterioribus antice subcurvatis profundius fassis, corollæ cœruleæ tubo calyce subduplo longiore, nuculis trigenis laevibus 2.

Hab. in Persia monte Elbrus et inter Ispahan et Teheran (Oliv! Auch. 4749!), in prov. Aderbidjan distr. Khoi ad torrentem Lavin et in collibus sterilissimis circa Pere (Szov!), circa Marand (Bge!), ad nives montium Avroman et Schahu (Haussk!).

Bi- et tripedalis, paniculae rami nudi gracillimi inferiores pedales, folia inferiora pollicem longa et basi lata, calyx 4 lin., corolla 9 longi.

22. N. teucriifolia (Willd. Enum. Berol. pag. 602 et herb! glaucescens glabra vel sub lente adpresso puberula, caulis ascendentibus elatis parte inferiori excepta subaphyllis, foliis crassiusculis petiolatis subcordatis ovato-triangularibus obtusis obtuse dentatis, superioribus basi truncatis vel cuneatis, floralibus paucis minimis, cymis dichotomis 7-3-floris inferioribus laxis longe superioribus brevius pedunculatis paniculam laxissimam elongatam formantibus, bracteis lanceolatis pedicello sublongioribus, calycis breviter papillosi striati tubulosi incurvi ore valde obliquo dentibus oblongis acutis tubo quintuplo brevioribus inferioribus antice subcurvatis profundius fassis, corollæ cœruleæ tubo tenui incurvo calyce triplo longiore, nuculis oblongis trigonis laevibus 2. *N. fissa* C. A. M. Enum. pag. 93. — *N. laxa* Bth. Lab. p. 483. — *N. glabra* et var β *Persica* Bth. Lab. pag. 483! — *N. rarifolia* et *N. virgata* C. Koch Linn. XXI, p. 676!

Hab. in Cappadocia monte Akdag (Auch. 4730!), Armenia Kurdiæ (Ky. 473!), Armenia boreali (C. Koch!), ditione Talyssch (C.A. M! Hoh!), Persia boreali (Auch. 4754!), in monte Elbrus prope Derbend (Ky. 592!) et prope Radkann (Buhse!).

Affinis *N. longijfora* a qua differt glabritie, corollæ tubo longiore. Bracteæ non ovatae sed lanceolatae!

23. N. Calverti (Boiss. Diagn. Ser. II, 4 p. 25) sub lente adpresso et minutissime pubescens, caulis ex rhizomate incrassato fibroso ascendentibus simplicibus, foliis minutis petiolatis cordatis ovatis obtusis obtuse dentatis sursum sensim diminutis, floralibus minimis oblongo-lanceolatis, cymis 7-3-floris anguste paniculatis inferioribus longe superioribus breviter pedunculatis pedunculis gracilibus strictis, bracteis setaceo-linearibus acuminatis tubo calycino brevioribus, calycis secus nervos præsentim asperuli incurvi ore obliquo dentibus ovatis acuminatis tubo sextuplo brevioribus,

corollæ violaceæ tubo calyce $2\frac{1}{2}$ longiore, nuculis lœvibus ovatis subcompressis ♀. *N. obtusicrena* Boiss. in Ky exs.

Hab. in Armeniâ circa Erzerum (Calvi!), inter Mamahatun et Jeniköi prope Erzerum (Tchih. exs. 202! ad *N. marifoliam* erroneâ relata), ad Müküs 6500' (Ky. Suppl. 685!).

Pedalis, panicula racemosa 4-6-pollices longa. Species a *N. teucrifoliâ* affini foliis minoribus (6-7 lineas longis), corolle tubo duplo breviore, etc., distincta. Calyx 3 lineas, corolla 7 longi. Affinis quoque *N. marifoliae* sed non cana, folia cordata, calycis dentes ovati, nuculae lœves.

24. N. gracilescens (Boiss. in Bourg. pl. Arm. exs. 4862) sub lente adpresso et minutissime pubescens canescens, caulinis ascendentibus simplicibus tenuiter virgatis, foliis parvis petiolatis cuneato-oblongis obtusis obtuse crenatis, floralibus minutis linearilanceolatis integris acutis, cymis laxè 3-3-floris pedunculatis angustè subpanienlati-racemosi, cyme ramulis pedicellisque divaricatis, bracteis linearilanceolatis apice subulatis, calycis papilloso-hirti incurvi ore obliquo dentibus oblongo-lanceolatis membranaceo-marginatis acuminatis tubo 4-5-plo brevioribus inferioribus profundius fissis, corollæ tubo calyce subduplo longiore, nuculis lœvibus ♀.

Hab. in collibus Armeniæ Turcice prope Ispir (Bourg.).

Pedalis, racemus depauperatus 4-5-pollicaris, calyx fructifer $3\frac{1}{2}$ lineas longus. Differt a *N. teucrifoliâ* et *Calverti* foliis non cordatis sed cuneato-oblongis (8-9 lineas longis 4 latis), calycis dentibus margine scariosis, a *N. marifoliâ* calycis dentibus margine membranaceis et setaceo-acuminatis distincta.

25. N. Trautvetteri (Boiss. et Buhse Aufz. p. 775) caulinis elatis basi tantum foliosis sub lente pruinosis inferne interdum hirtellis, foliis parvis longe petiolatis basi subcordatis rotundatis crenatis superne breviter subtus longius tomentellis canescensibus, floralibus minimis subsessilibus oblongis, cymis longe pedunculatis dichotomis laxifloris paniculam longam formantibus, bracteis minimis mucronatis, floribus unilateralibus subsessilibus, calycis subincurvi valde striati asperuli ore valde obliquo dentibus brevissimis triangulari-oblongis margine membranaceis acutis inferioribus brevioribus, corollæ hirtulae tubo calyce subduplo longiore, nuculis lœvibus ♀.

Hab. in ditione Transcaucasiciâ, Karadagh prope Alliar (Buhse!) in lapidosis montanis vallis Koschadara (Szov!).

Planta 1-1½-pedalis, folia 5-7 lineas longa et lata, panicula ½-1-pedalis, calyx 3½ lineas longus. Ab affini *N. longiflorâ* differt foliis parvis, cymis bifidis laxifloris et imprimis dentibus calycinis latis.

26. N. brevifolia (C. A. Mey. Enum. p. 93) virens sub lente tenuissime pubescens, caulinis assurgentibus tenuibus ramosis, foliis minutis late cordatis triangulari-orbiculatis grosse paucierenatis inferioribus longe petiolatis, floralibus minutis subsessilibus oblongis

acutis acute dentatis, cymis paucifloris inferioribus breviter pedunculatis verticillastra 3-4 dissita formantibus, bracteis lanceolatis setaceo-acuminatis tubo calycino subbrevioribus, calycis striati minute glandulosi et patule papilloso incurvi ore obliquo dentibus ovato-lanceolatis acuminatis tubo 8-4-plo brevioribus, corollæ cæruleæ calyce triplo longioris tubo tenai incurvo longe exerto, nuculis.... ♀. *N. lamiifolia* Bth. in n.c. Prodri. non Willd!

Hab. in schistosis montis Tufandagh Caucasi orientalis 9000' (C.A. Mey!), in Armenia (Willd).

Pedalis, folia inferiora pollicem dimidium longa et lata dentibus 5-7 obtusissimis, corolla 7-8 lineas longa. Afinis *N. longiflora*, humilior, indumentum tenuius et parceus, calycis dentes non acuti sed spinuloso-acuminati.

27. ***N. macrosiphon*** (Boiss. Diagn. Ser. I, 7 p. 51) patule et brevissime pubescens subcanescens, caulis e rhizomate indurato ascendentibus elongatis laxe et anguste paniculatis, foliis cordatis ovatis grosse et obtuse dentatis vel sublobatis petiolatis superioribus sessilibus valde diminutis, cymis confertim 5-7-floris inferioribus longe superioribus breviter pedunculatis, bracteis minimis lanceolatis breviter ciliatis, calycis incurvi nigricantis ad nervos prominentes sparsim papilloso-asperi ore valde obliquo, dentibus tubo sextuplo brevioribus acutis superioribus oblongis inferioribus profundius fisis lanceolatis, corollæ cæruleæ hirtæ tubo valde incurvo calyce 2 1/2-plo longiore, nuculis levibus ♀.

Hab. in fissuris rupium montis Kuh Daëna Persiae australis (Ky. 588! et 986!) ad nives montis Schahu supra Juanro 12000' et in monte Dalechani supra Sungur 9000' (Haussk!). Odor gratus.

Cum paniculâ angustâ sœpe pedali pedes duo longa, folia inferiora 1-2-pollices longa, calyx fere 5 lineas, corolla 12-14 lineas longi. Species pulchra corolla longissima inter affines insignis.

28. ***N. leptantha*** (Boiss. et Haussk.) breviter pubescens canescens, caulis a basi indurata ascendenter simplicibus vel parce ramosis, foliis breviter petiolatis cordatis ovatis obtusis grosse et obtuse dentatis, floralibus oblongo-lanceolatis integris acutis, verticillastris paucifloris sessilibus parvis inferioribus præsertim longe remotis, bracteis subulatis ciliatis calycinum tubum æquantibus, calycis rubelli recti sparsim et patule albo-papilloso ore subobliquo dentibus lanceolatis acuminatis albo-ciliatis tubo 2 1/2-plo brevioribus inferioribus angustioribus profundius fisis, corollæ hirtæ tubo angustissimo subincurvo calyce 2 1/2-plo longiore, nuculis.... ♀. *N. graciliflora* Boiss. et Haussk exs. non Bth.

Hab. in rupestribus montis Berytdagh Cataoniae 7000' (Haussk!).

Sesquipedalis, folia majora 12-15 lineas longa, verticillastra 5-9 avellanæ multo minora, calyx florifer 4 lineas, corolla 9 longa. Species bracteis et calycibus pectinatim ciliatis insignis. An semina lævia?

29. ***N. leptoclada*** (Trautv. Stirp. Nov. N, № 13) caule elato

gracili inferne longe molliter et patule hirsuto superne glabresecti anguste et longe paniculato, foliis supra breviter pubescentibus subtus patule hirsutis cinerascentibus cordato-orbiculatis grosse et obtuse crenato-dentatis inferioribus longiusculæ petiolatis cæteris subsessilibus, paniculae ramis brevibus tenuibus 3-5-floris, bracteis minimis oblongis acutis, pedicellis calyce subbrevioribus, calycis anguste tubulosi rectiusculi papillis albis asperi ore valde obliquo dentibus ovatis acutis vel micronatis inferioribus subrecurvis, corollæ carneaæ tubo tenuissimo calyce 2 $\frac{1}{4}$ -plo longiore, nuculis levibus ♀.

Hab. in Armenia Rossicâ ad pagum Belaw (V. sp. in herb. cl. Trautv.)

Herba bipedalis, folia inferiora diam. 45 lineas lata interdum transverse sublatiōna, panicula angusta pauciflora semi-pedalis, calyx 2 $\frac{1}{2}$ lineas longus. Inter affines foliis orbiculatis et indumento molliter et longe hirsuto cinereo notabilis.

§ 5. **Psilonepetæ.**

30. **N. sessilifolia** (Bge Lab. Pers. p. 54) viridis sub lente glandulis minimis subsessilibus pruinosa, caulis erectis elatis superne striete ramosis paniculatis, foliis majusculis sessilibus a basi truncatâ late ovatis grosse et acutiusculæ dentatis floralibus diminutis oblongis, cymis longe pedunculatis confertim 5-3-floris, bracteis subulatis pedicello vix longioribus, calycis recti valde striati glanduloso-asperi ore æqualis dentibus breviter triangularibus acutis tubo quintuplo brevioribus, corollæ glabrae tubo calyce subduplo longiore, nuculis obovato-oblongis levibus ♀.

Hab. prope Kohrud inter Ispahan et Kaschan (Bge!).

Bipedalis, folia fere *Mimuli* 1 $\frac{1}{2}$ -2-pollicaria, calyx fructifer 3 $\frac{1}{2}$ lineas longus

31. **N. laxiflora** (Bth. in n.c. Prodr. XII. p. 393) glaberrima, caulis elatis gracilibus striete paniculato-ramosis, foliis petiolatis cordatis oblongis grosse et obtuse dentatis dentibus latis ascendentibus, superioribus subsessilibus acutis subintegris, cymis 7-3-floris longe et graciliter pedunculatis, bracteis subulatis minutis, calycis valde striati ad strias minute asperi vel glabri recti dentibus triangularibus acutis æquilonis tubo 5-6-plo brevioribus, corollæ glabrae tubo calyce 2 $\frac{1}{2}$ -plo longiore, nuculis levibus ♀.

Hab. in Persiaustrali (Auch. 1745!), in summis montis Schahu Kurdistaniæ Persice ad niveis 12000' (Haussk!).

Caules 2 $\frac{1}{2}$ -pedales, folia inferiora 1 $\frac{1}{2}$ -pollicem longa, calyx 2 $\frac{1}{2}$ lineas, corolla 7 lineas longi. Species formâ dentium calycis affinis *N. sessilifolia* sed hujus folia sessilia formâ omnino dissimilia, indumentum pruinoso-glandulosum, etc.

32. **N. oxyodonta** (Boiss. Diagn. Ser. I. 7 p. 51), caulis numerosis sub lente tomentellis vel glabratis a basi indurata erexit

tenuiter virgatis elongatis simplicibus vel stricte ramosis superne subnudis, foliis adpresso canescenti-tomentosis subcordatis ovato-orbiculatis subinciso-dentatis inferioribus breviter petiolatis, superioribus paucis sessilibus oblongo-linearibus minimis, cymis 5-7-floris inferioribus breviter pedunculatis superioribus subsessilibus saepe approximatis, bracteis linear-lanceolatis calycis tubo brevioribus, calycis asperuli elongati recti dentibus lanceolatis apice subulatis spinulosis subæqualibus tubo quadruplo brevioribus, corollæ carneæ tubo calyce subduplo longiore, nuculis oblongis subcompressis levibus ♀.

Hab. in Persiâ australi inter frutices faecium alpis Kuhdehu et in caeruleum montis Sabst Buschom prope Schiraz (Ky. 507! et 415!), in montibus Sawers, Nur. Eschker districtu Kuh Kilonyeh 8-10000' et alpe Kellal ditionis Bachtiaris (Haussk!).

Bipedalis basi fere suffrutescens, folia 8-9 lineas longa lataque. calyx demum 5 lineas, corolla 9 longi. ·

33. N. depauperata (Bth. in p.c. Prod. 12 p. 392) sub lente minute glandulosa suffruticosa ramosissima ramis simplicibus erectis rigidis tenuiter virgatis, foliis minimis breviter petiolatis ovatis subcordatis plicato-rugosissimis margine crispis et dentatis, cymis strict pedunculatis 3-1-floris anguste et laxè racemosis, bracteis lanceolatis longitudine pedicelli, calycis striati sub lente pruinosi recti tubo cylindrico in limbum sublatiorem et paulo breviorem in dentes late lanceolatos acutos ad $\frac{2}{3}$ tissum expanso, corollæ tubo calyce 2 $\frac{1}{2}$ -plo longiore, nuculis glabris ♀. *N. septemcrenata* var. *depauperata* Boiss. Diagn. Ser. I, 7 p. 41.

Hab. ad rupes faecium montis Sabst Buschom prope Schiras (Ky. 374!), in Persiâ australi (Auch. 5158! ex Bth.).

Semipedalis, folia 2-3 lineas longa, insignis suffrutescentia, foliis plicatis minimis. Calyx florifer $2\frac{1}{2}$ lineas longus in limbum abruptiuscula dilatatus. Corolla 7-8 lineas longa.

34. N. denudata (Bth. in p.c. Prodr. XII, p. 390) caulis elatis tenuiter virgatis junceis stricte ramosis parte inferiori exceptâ aphyllis sub lente minute tomentellis vel glabratis, foliis utrinque adpresso aubo-lanatis petiolatis ovatis obtusis obtuse crenatis subrugosis, floralibus bracteisque linear-subulatis, cymis 2-3-floris inferioribus breviter pedunculatis superioribus sessilibus, calycis tomentelli elongati recti ore subobliquo dentibus acute lanceolatis tubo quadruplo brevioribus anterioribus subbrevioribus profundius fisis, corollæ tubo tenui calyce subduplo longiore, nuculis levibus ♀.

Hab. in argillosis Persiae borealis (Auch. 5129!), in schistosis montis Elbrus prope Derbend (Ky. 410!).

Bipedalis, racemi laxissimi tenuissimi, folia pauca majora pollicem longa, calyx tres lineas, corolla sex longi.

§ 6. **Macrostegiæ.**

35. N. glomerulosa (Boiss. Diagn. Ser. I, 5 p. 21) adpressissime lanata nivea, ex rhizomate lignoso multicaulis caulinis tenuiter virgatis elongatis subsimplicibus superne subnudis, foliis parvis ovatis crenatis lineato-rugosis inferioribus breviter petiolatis cæteris sessilibus, floralibus ovatis submembranaceis, verticillastris parvis densitioris inferioribus 1-2 remotis cæteris in spicam oblongam approximatis, bracteis membranaceis ovato-oblongis acutis eucullatis calyces æquantibus, calycis longe hirsuti membranacei striati ore subobliqui dentibus lanceolato-subulatis tubo subbrevioribus, corollæ roseæ tubo inclusa, nuculis oblongis minute tuberculatis ♀.

Hab. in lapidosis montis Elbrus et prope Tchesme Ali Persiae borealis (Buhse!), in montibus prov. Khorassan (Bge!), prope Isphahan (Auch. 1748! et 5139!), in calcareis ad nives Kuh Nur, Sawers, Maregun, Eschker 11-13000' (Haussk!).

Pedalis vel sesquipedalis, folia inferiora 4-8 lineas longa, racemi 2-3-plicares, calyx 2½-3 lineas longus. Ex affinitate *N. Sibthorpii* et *leucostegia*, flores minores et praesertim bracteis ovatis distinctissima.

36. N. juncea (Blh. p.c. Prodr. XII, p. 379) adpressissime tomentoso-canescens a basi suffrutescenti ramosissima, ramis erectis tenuiter virgatis basi foliosis superne nudis inferne ranulosis, foliis minimis paucis breviter petiolatis ovatis obtusis crenatis basi truncatis rugosis, verticillastris parvis paucifloris inferioribus valde distantibus superioribus interrupte spicatis, bracteis scariosis ovatis acuminate calyces æquantibus, calycis ore obliqui hirsuti dentibus breviter lanceolatis tubo duplo brevioribus inferioribus profundius fassis, corollæ calyce sesquialongiore, nuculis minutissime tuberculatis ♀.

Hab. ad Karabagh et Schusgao regni Cabulici (Griff!), ad Chehel Tun Belutschia (Stocks 1009!).

Pedalis et sesquipedalis, valde affinis *N. glomerulosa*, folia rariora minora 3-5 lineas longa, verticillastris pauciflori remotissimi et multo minores pisum vix æquantes, calyces 2 lineas parum excedentes brevius dentati.

37. N. Scordotis (L. Sp. p. 698) caulinis ascendentibus elatis foliosis simplicibus vel inferne ramosis papillis longis crispis hirsutis, foliis adpresso papilloso-lanatis subtus præserit canescens cætibus majusculis ovatis obtusis grosse et obtuse crenatis basi anguste cordatis inferioribus petiolatis superioribus subsessilibus, floralibus bracteisque ovatis subacuminatis calyces æquantibus, verticillastris 5-10 spicatis inferioribus distinctis superioribus confluentibus, calycis albo-lanati dentibus lanceolato-subulatis æquialongis vix incurvis tubo æquialongis, corollæ puberulae cærulecentis tubo calyce sesquialongiore tenui sensim ampliato, nuculis subcompressis rugulosotuberculatis ♀. *S. tomentosa* Sieb. exs.

Hab. ad vias et in siccis regionis inferioris Cretæ (Sieb., Heldr! Weiss!).

Bipedalis, folia inferiora 2-2 $\frac{1}{2}$ -pollices longa, spica 4-5-pollicaris basi valde interrupta, calyx florifer 3 lineas, corolla 6 lin. longa. Facies *Stachydis circumdata*.

38. N. Sibthorpii (Bth. Lab. p. 474) adpresso tomentosa cana, caulis erectis vel ascendentibus simplicibus vel parce ramosis, foliis breviter petiolatis rugosis ovato-oblongis cordatis crenatis, verticillastris paucis densifloris inferioribus distantibus superioribus in spicam oblongam condensatis, bracteis numerosis membranaceis albidis oblongo-lanceolatis subulato-acuminatis calyces æquantibus, calycis recti dentibus subæqualibus longe lanceolato-subulatis tubo æquilongis, corollæ albæ pubescens tubo calycem non excedente, nuculis tuberculatis ♀. *N. Italica* Sibth. et Sm. Fl. Græc. 6, t. 548 non L. — *N. Argolica* Bory et Chaub. Exp. Mor. № 889, tab. XVII (mala).

Hab. in siccis regionis inferioris Græciæ rarius, saepius in montanâ et subalpinâ, ad Naupliam et Argos (Bory, Sprun!), mons Kyllene 3-4000' (Heldr! Orph. Fl. Gr. exs. 210!), regio abietina montis Parnes Atticæ (Sibth., Sprun! Heldr!), circa Patras (Link).

β. Dirphyæ. — Planta minus cana, flores paulo minores. — *N. Dirphyæ* Heldr. herb. Norm. 777 !

Hab. in lapidosis montis Dirphys Eubeæ 4500' (Heldr!).

Species affinis *N. Orientali*, magis cana, spicæ breviores magis condensatæ, bracteæ latiores albo-membranaceæ, calyx profundius fissus.

39. N. leucostegia (Boiss. et Heldr. Diagn. Ser. I, 12 p. 63) adpressissime cana multicaulis, caulis ascendentibus gracilibus simplicibus vel ramosis, foliis parvis petiolatis cordatis ovato-oblongis obtuse crenatis, verticillastris numerosis parvis multifloris distantibus longe et laxe racemosis. bracteis numerosis nervosis albo-marginatis oblongo-lanceolatis mucronatis calyce vix brevioribus, calycis tomentelli recti dentibus æqualibus lanceolatis tubo æquilongis, corollæ albæ pubescens tubo subincurvo subexerto nuculis tuberculatis ♀. *N. Sibthorpii* Bth. ex parte quoad plantam ex Anatoliâ et Syriâ.

Hab. in Lyciâ prope Elmaju (Bourg. 213!), Pisidiâ ad Bouldur et Egirdir (Heldr!), ad Amasia, Tokat, Safranbol, Samsun, Mersivan (Wied!), Ciliciâ ad Gülek (Ky. 458! Bal!), monte Ssöffdaghs Cataoniae et prope Orfa Mesopotamiae (Haussk!), in monte Pir Omar Gudrun Kurdistaniæ 3-4000' (Haussk!), Syria prope Zebdani et Antilibano 4-6000' (Ehr! Boiss! Ky. 4!).

Affinis *N. Sibthorpii* a qua differt indumento brevissimo, verticillastris numerosis distantibus et præsertim ut et a *N. Orientali* floribus minoribus, bracteis et calyce breviter nec longe aristatis, corollâ subexsertâ.

§ 7. Stenostegiæ

40. N. Cadmea (Boiss. Diagn. Ser. I, 5 p. 21) adpresso et brevissime pruinoso-tomentella pallide virens, caulis erectis elatis

strictè ramosis, foliis petiolatis ovato-oblongis cordatis crenatis, floralibus sensim diminutis oblongis et linearibus spinuloso-acuminatis calyces æquantibus, verticillastris multifloris subsessilibus fere omnibus distinctis inferioribus remotis, bracteis subulatis viscidò-pruinosis calyces superantibus, calycis viscidò-pruinosi recti dentibus lanceolato-subulatis tubo æquilongis subinaequalibus, corolla albæ calyce duplo longioris tubo vix exerto, nuculis tenuissime tuberculatis.

Hab. in dumosis regionis montane ad basin montis Cadmi Cariæ supra Colossam (Boiss.), in rupestribus pinorum Pamphylæ inter Marla et Adalia 4000' (Heldr.).

Bipedalis, affinis *N. Orientali* a qua differt bracteis longissime subulatis nec lanceolatis, calycis dentibus longe subulatis, verticillastris minoribus. *N. crinita* affinis quoque differt indumento calycis et bractearum striguloso nec brevissime pruinoso-glanduloso, bracteis et calycis dentibus tenuius et longius subulatis.

41. *N. Isaurica* (Boiss. et Heldr. in p.c. Prodr. 12 p. 378) brevissime velutina cauæseens vel cana e rhizomate indurato ramosissima, ramis tenuibus ascendentibus pumilis, foliis parvis rugulosis petiolatis ovato-triangularibus truncatis vel cordatis, verticillastris in spicam oblongam basi interruptam approximatis, bracteis linearisubulatis calyce vix brevioribus, calycis hirsuti recti dentibus lanceolatis tubo subbrevioribus, corollæ calyce subdupo longioris tubo subexerto, nuculis tuberculatis ♀.

Hab. in monte Gheidagh Isauriæ 5200' (Heldr.).

β. Scorodonia. — Paulo elatior et magis cana, verticillastri numerosiores dissiti. — *N. Scorodonia* Boiss. Diagn. Ser. I, 12 p. 63.

Hab. cum typo.

Caules in typo 4-5-pollicares, folia 4-5 lineas longa, calyx 3-3½ lineas longus, varietas β paulo elatior. Affinis *N. Orientali* differt statu minima, floribus dimidio minoribus, calycibus acutis nec longe aristato-mucronatis.

42. *N. camphorata* (Boiss. et Heldr. Ser. I, 7 p. 49) Iota pilis patulis crispis viscidulis lanatula cauæseens multicaulis basi indurata, caulis erectis vel ascendentibus elongatis ramosis, foliis parvis rugosis breviter petiolatis cordatis trianguli-ovatis margine crispule crenatis, floralibus minimis, verticillastris numerosis parvis remotis longe racemosis, bracteis subulatis calyce triplo brevioribus, calycis viridis patule hirti recti dentibus lanceolato-sabulatis tubum dimidium æquantibus, corollæ albæ tomentellæ tubo calyce sesqui-longiore, nuculis tuberculatis ♀.

Hab. in alveo torrentis Pentaplon in regione abietinæ Taygeti (Heldr.).

Tota planta odorem gravissimum camphoratum spirat; caules 1½-pedales, folia majora 6-8 lineas longa, verticillastri avellanæ minores, calyx florifer 2½ lineas, corolla fere 5 lineas longa. Foliis margine crispule crenatis notabilis, a *N. Orientali* floribus duplo brevioribus, a *N. Sibthorpii* bracteis subulatis, etc., distincta.

43. **N. Parnassica** (Heldr. et Sart. in Boiss. Diagn. Ser. II, 4 p. 22) breviter velutina viscidula canescens multicaulis, caulinis ascendentibus elongatis ramosis, foliis a basi truncato-subcordatâ ovato-oblongis crenatis rugulosis breviter petiolatis, verticillastris numerosis inferioribus distantibus cæteris in spicam approximatis, bracteis anguste linearibus calycis tubo subbrevioribus, calycis valde striati viridis tomentello-hirti dentibus aequalibus tubo subæquilon-gis, corollæ albidæ hirtæ tubo curvato breviter exerto, nuculis tuberculatis ♀.

Hab. in regione abietinâ et superiore montis Parnassi (Heldr! Sart! Fl. Grac. exs. 1872!), (Orph. Fl. exs. 433!).

β. Orphanidea — Adpresso cana. — *N. Orphanidea* Boiss. loc. cit. p. 22.

Hab. in regione media et superiore montis Malevo Laconiae supra Kastanitza (Orph. Fl. Grac. exs. N° 84!).

Tota valde camphorata, characteribus et bracteis angustissimis affinis *N. camphorata* a quâ floribus majoribus, calycis 3½ lineas longi dentibus tubo aequilongis differe videtur.

44. **N. Spruneri** (Boiss. Diagn. Ser. II, 4 p. 23) adpresso cano-pubescente velutino-canescens, caulinis ascendentibus inferne stricte ramosissimis, foliis parvis breviter petiolatis a basi truncato-subcordatâ oblongis subrugosis crenulatis concoloribus, verticillastris inferioribus valde dissitis superioribus confluentibus, bracteis subulatis calycino tubo vix longioribus, calycis adpresso-canis valde costatis recti dentibus tubo duplo brevioribus, corollæ pubescentis albae tubo calyce vix longiore, nuculis tuberculatis ♀. *N. Tymphrestea* Heldr. et Sart. Herb. Norm. 663!

Hab. in glareosis torrentium (Etoliae (Sprun!)), in regione abietinâ montis Veluchi (Sunar. et Guiee!).

Valde affinis *N. camphorata* et *Parnassica*, indumentum totius plantæ et præsertim calycis multo brevius et adpresso, calyx magnitudinis *N. Parnassica* minus profunde fissus.

45. **N. Orientalis** (Mill. Dict. N° 9) tenuiter cano-pubescente, caulinis erectis elatis simplicibus vel parce ramosis, foliis mediocribus petiolatis ovatis cordatis obtuse crenatis vix rugosis, floralibus sensim diminutis verticillastro brevioribus, verticillastris numerosis sessilibus racemosis distinctis inferioribus remotis superioribus approximatis, bracteis numerosis rigidis nervosis lanceolatis subspinosecenti-micronotatis calyces subæquantibus, calycis brevissime hirti subineurvi vix obliqui dentibus lanceolatis acuminatis pungentibus tubo subbrevioribus superiore longiore, corollæ calyce vix duplo longioris tubo calycem non superante, nuculis minute tuberculatis ♀. *N. Italica* L. Sp. p. 798. — Jacq. Hort. Vind. tab. 412. — Rehb. Ic. tab. 483. — *N. marrubioïdes* Willd. Enum. Berol. II, p. 603.

Hab. in rupe tribus regionis montane, Lydia supra Koukouludja prope

Smyrnam (Bal!), Isauria ad Ermenek (Heldr!), Cilicia ad Kechlik (Bal!), Syria borealis ad Aintab (Auch. 1729! Hassk!), mons Cassius in regione inferiore (Boiss!), Libanus (Bl! Gaill!), mons Gara Kurdistani (Ky. 408!).

Planta 2-3-pedalis, folia majora pollicaria, calyx fructifer 5 linea longus, Nec in Hispaniâ nec in Italiâ crescit, nomen *N. Italica* igitur omnino rejiciendum.

46. N. Cæsarea, breviter tomentosa canescens, caulis elatis firmis stricte paniculatis, foliis subsessilibus oblongo-lanceolatis obtusiusculis subintegratis, floralibus diminutis acuminatis, verticillastris in spicas terminales oblongo-cylindricas densifloras basi interdum subinterruptas approximatis, bracteis setaceis plumosis calyci æquilongis, calycis ore subobliqui dentibus a basi latiore longe sevaceis tubo suo longioribus superioribus sublongioribus, corollæ albidae tubo dentibus calycis vix longiore, nuculis (junioribus) tuberculatis ♀.

Hab. ad Cæsaream Cappadocie (Wiedemann exs. 318!).

Bipedalis et claviger, spicæ sesquipolligares, calyx fere 6 linea, corolla 7 longi. Species a *N. Orientali* et *cunctâ* inflorescentiâ dense spicato-capitata, calycis laciniis longissime setaceis distincta.

47. N. viscidæ (Boiss. Diagn. Ser. I, 5 p. 20) tota pilis patulis crassis papillosa viscida viridis, caulis erectis subsimplicibus, foliis sessilibus subcordatis ovato-oblongis crenatis, floralibus sensim diminutis lanceolatis calyces subaequantibus, verticillastris multifloris numerosis distinctis viscosis, bracteis linearibus calyce subbrevioribus, calycis recti ore subobliqui dentibus rectis lanceolatis acutinsculis tubo subduplo brevioribus superioribus sublongioribus, inferioribus profundius fissis, corollæ cœruleæ viscidulae tubo gracili subineluso, nuculis tuberculatis ♀.

Hab. in regione superiore montis Tmoli supra Philadelphiam (Boiss!), in monte Bulghasdag Phrygiae 4500' (Bal!).

Pedalis, folia inferiora fere bipolligares, verticillastri ampli. calyx striatus tandem fere 5 linea longus.

48. N. Tmolea (Boiss. Diagn. Ser. I, 5 p. 22) indumento brevi crispulo hirsuta, caulis elatis simplicibus vel stricte ramesis, foliis præter infima subpetiolata sessilibus ovato-oblongis cordatis crenatis, verticillastris numerosis violaceis hirsutis multifloris præter summa distinctis inferiorum cymis subpedunculatis, bracteis subulatis ineurvis calyce subduplo brevioribus, calycis hirsuti striati recti ore subobliqui dentibus subæqualibus linear-lanceolatis acutinsculis tubo subæquilongis, corollæ cœruleæ hirtæ tubo vix exerto, nuculis tuberculatis ♀.

Hab. in regione superiori montis Tmoli supra Philadelphiam (Boiss!).

β. laxior. — Cymæ præsertim inferiores laxiores longius pedunculatae, bractearum et calycis indumentum brevius.

Hab. in parte superiori vallis Yachamichlar ad sept. urbis Ouchak Phrygiae (Bal!).

Affinis *N. viscidæ* sed indumentum vix viscidum, folia inferiora subpetiolata, cymæ inferiores sublaxiores, bractæ breves, dentium calyeis cum tubo proportio diversa. In monte Tmolo formæ inter *N. viscidam*, *Tmolcam* et *nedam* hybridae frequentes occurunt.

49. ***N. cyanea*** (Stev. Mém. Soc. Nat. Mosq. III, p. 265) adpressissime canescens, caulinibus ascendentibus humilibus a basi saepius paniculato-ramosis, foliis parvis petiolatis rugosis oblongo-cuneatis crenulatis, floralibus elliptico-linearibus brevibus, cymis axillaribus breviter pedunculatis dichotome laxifloris secundis, bracteis subulatis brevissimis, calyeis cano-tomentosi incurvi cyanei ore obliquo dentibus breviter lanceolatis tubo 4-5-plo brevioribus superiore longiore, corollæ albido-purpureæ calyce parum longioris tubo inclusa, nuculis oblongis malleato-insculptis ♀. *N. incana* M.B. II, p. 40 et Rehb. Ic. bot. IV, tab. 303 non Pl. Crit. — *N. acinifolia* Spreng. Pug. p. 41.

Hab. in schistosis Caucaesi ad fluvium Terek et prope Nartzana (C.A. M.), ad torrentem Ghodjal (Stev.), ad Dzerani (Rehnl.), in monte Elbrus 4000' (Radde!), ditione Alagir 2500' et Caucaso orientali 4500'-6900' (Ruprl.).

Semipedalis patentim ramosissima, prætrea differt a *N. Mussini* foliis non cordatis, cymis laxifloris secundis, corollæ tubo inclusa, nuculis malleatis. Ramificatio sectionis *Oxynepetæ*.

50. ***N. aristata*** (Boiss. et Ky in Ky. exs. 1859) glabra viridis, caulinibus elatis crassiuseulis simplicibus vel parce et stricte ramosis, foliis magnis a basi truncato-subcordatâ late ovato-oblongis grosse inciso-dentatis inferioribus brevissime petiolatis cæteris sessilibus, floralibus anguste linear-lanceolatis spinoso-acuminatis integris calyeis aequantibus, verticillastris multifloris densis subsessilibus inferioribus remotis supremis confluentibus. bracteis numerosis subulatis calycinum tubum aequantibus, calyeis sub lente pruinoso-asperuli recti ore subobliquo dentibus a basi lanceolatâ longe subulatis tubo subtriplo brevioribus inferioribus profundius fisis, corollæ tubo tenui subincurvo calyce subduplo longiore, nuculis ovatis compresis malleato-tuberculatis ♀. *N. trichocalyx* Boiss. et Haussk. MSS.

Hab. in rupestribus vallis Teng prov. Musch Armenie 6500' (Ky. exs. 458!), in monte Berytdagh Cataoniae 7000' (Haussk!).

Bi- et tripedalis, folia inferiora 2-3-polices longa, racemus basi laxissimus 1-1-pedalis, calyx florifer 5 lineas, corolla 10 lineas longa; dentibus calycinis longe aristatis et corollæ magnæ tubo elongato inter affines insignis.

51. ***N. septemcrenata*** (Ehrenb. in Bth. Lab. p. 484) minute glanduloso-viscida prætereaque patule papilloso-hirta, caulinibus virgatis elongatis simplicibus vel 1-2 ramulis strictis instructis, foliis pallide viridibus breviter ovatis profunde cordatis grosse et obtuse inciso-dentatis intimis breviter petiolatis cæteris sessilibus, cymis paniculosis intimis breviter et stricte pedunculatis cæteris sessilibus verticillastra longe et laxe racemosa formantibus, bracteis lanceolato-

subulatis florum pedicellis aequilongis, calyeis sub lente pruinosis vix incurvi ore oblique dentibus lanceolatis acutis tubo quintuplo brevioribus, corollae tubo incurvo calyce duplo longiore, nuculis tuberculato-muricatis ♀.

Hab. in desertis Egypti (Ehrenb.), in fissuris rupium montium Sinaiticorum (Bové! Schimp. 137! Auch. 1731!).

Pedalis, racemi semipedales, folia inferiora fere pollicem longa et lata, floralia infima ceteris conformia cetera minuta oblonga acuta integra, calyx $3\frac{1}{2}$ lineas, corolla 8 longa.

52. N. Persica (Boiss. Diagn. Ser. I, p. 66) minute glandulos-viscida et praeterea patule papilloso-hispida pallide virens ex rhizomate indurato multicaulis, caulinibus elongatis virgatis foliosis simplicibus vel inferne ramosis, foliis cordatis ovato-orbiculatis obtuse et grosse crenatis inferioribus petiolatis superioribus sensim diminutis sessilibus, floralibus inferioribus flores aequantibus ceteris minutis oblongo-linearibus, verticillastris plurifloris laxe et longe racemosis inferioribus distantibus breviter et striete pedunculatis ceteris subsessilibus, bracteis linear-oblengis pedicello sublongioribus, calycis crispule hirtuli recti ore vix obliquo dentibus lanceolatis acutis rigidis dimidium tubum aequantibus, corolla rubella albo-striata tubo calyce vix duplo longiore, nuculis ovato-subtrigonis muricatis ♀. *N. septemcrenata* Bth. in n.c. Prodr. ex parte quoad locos Persicos.

Hab. in Persia borealis monte Elbrus supra Derbend (Ky. 258!), circa Schahrud et Kerman (Bge!), in montibus circa Isphahan (Bode!), in Persiaustrali (Auch. 5157!), in collibus saxosis ad Schiras et Persepolim (Ky. 398! et 401!).

Pedalis et sesquipedalis, folia inferiora diametro 9-10 lineas lata. A *N. septemcrenata* eximie differt foliis multierenatis, verticillastris plurifloris, calycis obconici nec cylindrici vix $2\frac{1}{2}$ lineas longi dentibus dimidio tubo aequilongis. Corolla $5\frac{1}{2}$ lineas tantum longa.

53. N. Kotschy (Boiss. Diagn. Ser. I, 7 p. 50) adpresse canescens-tomentosa ex rhizomate indurato suffrutescenti multicaulis, caulinibus ascendenti-erectis virgatis foliosis simplicibus et striete ramosis, foliis ovato-orbiculatis subcordatis obtuse et grosse crenatis inferioribus breviter petiolatis, verticillastris breviter et striete pedunculatis dissitis longe et laxe racemosis, bracteis oblongis minimis, calyeis tomentosi fructiferi oblongi recti ore vix obliquo dentibus breviter lanceolatis aentiusculis tubo subtriplo brevioribus, corolla carneae tomentellae tubo calyce duplo longiore, nuculis muricatis ♀.

Hab. in Persia australi ad Perezend (Auch. 5149!), in collibus prope Schiras usque ad regiones altiores alpis Kuh Delu (Ky. 320! 495! et 507!).

Pedalis et sesquipedalis, folia inferiora 8-9 lineas diam. lata, calyx $2\frac{1}{2}$ lineas corolla 7 lineas longi. Ob nuculas muriculatas *N. septemcrenata* et *Persica* appropinquanda, ceteris neglectis indumento statim distinguenda.

54. N. betonicæfolia (C. A. Mey. Enum. p. 92) brevissime canescens-pubescentis, caulibus ex radice repente simplicibus erectis, foliis subsessilibus ovato-oblongis profunde cordatis crenato-dentatis superioribus subdiminutis acutiusculis, spicæ oblongæ basi interdum interruptæ, bracteis villosis linearis-subulatis calyce dimidiò brevioribus, corollæ pubescentis tubo tenui incurvo calyce sesquilociore, nuculis.... φ .

Hab. in graminosis alpinis ditionis Transcaucasicae Talysh 6000' (C.A. Mey!).

Vix pedalis, folia majora 12-15 lineas longa eorum paria approximata, spica densa bipollicaris diametro 12-15 lineas lata, calyx 4 lineas, corolla 7 longa. Affinis *N. racemosæ* differt spicæ compactâ, calycis dentibus acuminatis, etc.

55. N. glabrescens (Boiss. in Bourg. exs. Armen. 1862) viridis sub lente brevissime et crispule puberula, caulibus crassiusculis simplicibus vel stricte ramosis, foliis cordatis oblongis obtuse crenatis inferioribus breviter petiolatis superioribus sessilibus, floralibus oblongo-lanceolatis dentatis parvis, cymis inferioribus breviter pedunculatis laxiusculis remotis superioribus approximatis, bracteis linearibus pubescentibus calyce duplo brevioribus, calycis breviter pubescentis ore subobliquo dentibus rectis æquilongis lanceolatis tubo 2-4 plo brevioribus, corollæ hirtæ calyce sesquilocioris tubo subexerto, nuculis tuberculatis φ . *N. cordifolia* C. Koch MSS.

Hab. in collibus prope Baibut Armeniae (Bourg!), in Cappadociâ (Auch. 1728!), in valle Merga Sauk prov. Musch (Ky. 367!).

Sesqui vel bipedalis, folia inferiora 1-1 $\frac{1}{2}$ -pollicaria, calyx 3-3 $\frac{1}{2}$ lineas longus. Affinis *N. betonicæfolia* cui cl. Bth. plantam Aucherianam adnumeravit sed *N. betonicæfolia* differt cynis in spicam oblongam et villosam condensatis, floralibus majoribus, calycis dentibus abbreviatis. Magis affinis *N. Tmoleæ* et præsertim hujus varietati β calycis hispida dentibus sublinearibus tubo æquilongis diversæ. Proxima quoque *N. racemosæ* calyce molliter villoso, etc., diversæ.

56. N. concolor (Boiss. et Heldr. Diagn. Ser. I, 12 p. 65), caulibus glabris elatis subsimplicibus, foliis papillis minimis sparsis sub lente obsitis concoloribus viridibus cordatis evatis grosse et obtuse crenato-lobatis inferioribus breviter petiolatis cæteris sessilibus, floralibus parvis acutis acute dentatis superioribusque integris, verticillastris paucifloris inferioribus remotissimis supremis magis approximatis, bracteis subulatis calycis tubo brevioribus, calycis pilis papillaribus patentibus rigidulis hirsuti striati ore obliqui dentibus lanceolatis acutis subpatentibus tubo duplo brevioribus, corollæ valde hirtæ tubo calyce subduplo longiore, nuculis tuberculatis φ .

Hab. in cacumine occidentali montis Gheidagh Tauri Isaurici 6000' (Heldr!).

Planta 1 $\frac{1}{2}$ -2-pedalis, folia inferiora bipollicaria, racemi laxissimi semipedales, corolla sex lineas longa. Affinis *N. glabrescens* quæ sat differre videtur foliis floralibus acute dentatis, verticillastris inferioribus valde remotis, calycis indumento, dentibusque subpatentibus. Nuculae prius laeves descriptæ insecti iectu monstrosæ erant, normales juniores tuberculatæ videntur.

57. **N. grandiflora** (M.B. Taur. Cane. II, p. 42) glabriuscula viridis, caulis elatis ramulosis, foliis breviter petiolatis magnis oblongis cordatis crenato-dentatis, floralibus lanceolatis inferioribus cymas æquantibus, cymis multifloris laxe racemosis inferioribus pedunculatis, bracteis subulatis calyce multo brevioribus, calycis violacei breviter asperulo-hirti striati elongati incurvi ore obliquo, dentibus lanceolatis acuminatissimis tubo subtriplo brevioribus superioribus sublongioribus, corollæ cœruleæ tubo exerto, nuculis impresso-subtuberulatis ♀. Ic. Rehb. Icon. bot. 3 tab. 279. — *N. melissæfolia* Pers. Syn. 2 p. 115 non Lam. — *N. colorata* Willd. Enum. 2 p. 602.

Hab. in graminosis sylvaticis et subalpinis totius Caucasi 45°00'-60°00' (M.B., G.A. Mey.), Daghestanie 6-7000' (Rupr!), Iberie et prov. Karabagh (Hoh!).

Bipedalis et elatior, folia inferiora 2-2 $\frac{1}{2}$ -pollicaria, racemi interdum basi ramoso-subpaniculati, calyx 4 $\frac{1}{2}$ lineas, corolla 8 longa.

58. **N. speciosa** (Boiss. et Noë Diagn. Ser. II, 4 p.¹/34) viridis sub lente parce et brevissime hirtella, caulis elatis simplicibus, foliis majusculis ovatis truncato-subcordatis grosse et obtuse crenato-dentatis inferioribus brevissime petiolatis cæteris sessilibus, floralibus inferioribus oblongo-lanceolatis acute dentatis flores subæquantibus cæteris linear-lanceolatis parvis, verticillastris numerosis multifloris racemosis inferioribus distantibus subpedunculatis, bracteis subulatis calyce triplo brevioribus, calycis pubescentis recti striati ore subobliquo, dentibus lanceolatis subæqualibus tubo 3-4-plo brevioribus, corollæ cœruleæ incurvæ calyce 2 $\frac{1}{2}$ -plo longioris tubo exerto limbo ampio, nuculis.... ♀.

Hab. in subalpinis Kurdistaniæ Persicae prope Mendeli (Noë 1186!), in rupestribus montium Avroman et Schahu 10000' (Haussk!).

Bipedalis, folia inferiora fere bipollicaria, racemi saepe semipedales verticillastris 5-9, superioribus approximatis, flores 7 lineas longi. Facies omnino et flores magni *N. grandiflora* a qua differt caule simplici, cymis inferioribus non longe subpaniculato-pedunculatis, calyce rectiusculo nec longe incurvo, etc.

59. **N. melissæfolia** (Lam. Dict. I, p. 711) patule papilloso-pubescentes, pallide virens vel canescens, radice fasciculato-fusiformi, caulis ascendentibus simplicibus, foliis ovatis cordatis grosse crenatis inferioribus longiusculæ petiolatis, floralibus sessilibus oblongis et superioribus breviter lanceolatis integris, cymis laxis laxe racemosis inferioribus longius cæteris breviter pedunculatis, bracteis subulatis calyce multo brevioribus, calycis breviter papilloso-hirti viridis striati incurvi ore obliquo dentibus triangulari-lanceolatis acutis tubo quadruplo brevioribus superioribus submajoribus, corollæ rubro-punctatae calyce duplo longioris tubo breviter exerto, nuculis minute tuberculatis ♀. Ic. Desf. Cor. tab. 12.

Hab. inter fruticos regionis sempervirentis Cretæ (Tourn., Heldr!), insula Melos (Urv., Armenis!).

Pedalis et bipedalis, folia inferiora 1-2-pollicaria, facies et flores magni *N.*

grandiflorae, racemus brevior, calyces pro longitudine latiores dentibus brevioribus et latioribus, corolla tubus brevior. Ab ea et affinibus insuper radice fibris fusiformibus 1½-2-pollices longis fasciculatis distinctissima.

60. **N. racemosa** (Lam. Dict. I. p. 711) pubescens pallide virens. caulis crassiusculis ascendent-erectis strictis elatis simplicibus, foliis cordatis ovatis vel oblongis crenatis inferioribus breviter petiolatis cæteris sessilibus, floralibus linear-lanceolatis parvis, cymis omnibus sessilibus vel inferioribus breviter pedunculatis laxe racemosis, bracteis linear-subulatis calyce multo brevioribus, calycis cyanæ molliter tomentosi vix curvati ore subobliquo dentibus breviter lanceolatis tubo duplo brevioribus, corollæ cœruleæ incurvæ tubo calyce sesquilongiore, nuculis subcompressis ruguloso-tuberculatis ♀. Ic. Rehb. Ic. bot. 6, tab. 585. — *N. Reichenbachiana* F. et M. Ind. 8 Petrop. p. 68.

Hab. in fæcibus et pascuis montanis Armeniæ circa Erzerum (Huett!), Baibut (Bourg!), in prov. Musch (Ky. 686!), circa Bayazid (Auch. 5148!), Persiæ boreali (Auch. 5131! et 5142!) in montibus Ssahend et Elbras (Buhse! Ky. 172!), inter Teheran et Tabris (Bge.).

Sesqui vel bipedalis, affinis *N. Mussini* differt caulis crassioribus erectis, calycis curvati dentibus longioribus.

61. **N. Mussini** (Henk. Adumb. p. 15) caulis numerosis diffusis vel ascendentibus crispule tomentoso-sublanatis, foliis parvis petiolatis cordatis ovatis obtusis crebre crenatis, floralibus sessilibus oblongis sensim diminutis cymas non æquantibus, cymis paucifloris racemosis inferioribus pedunculatis dissitis, bracteis paucis linear-lanceolatis calycis tubo brevioribus, calycis molliter tomentosi cyanæ incurvi ore obliquo dentibus triangulari-lanceolatis calyce quadruplo brevioribus superioribus longioribus, corollæ cœruleæ tubo calyce sesquilongiore, nuculis ruguloso-serobiculatis ♀. *N. longiflora* C. A. Mey. Ind. Cauc. p. 92 non Vent.

Hab. in collibus saxosis Iberiae circa Tiflis (MLB., Rupr!), Georgiae ad Elisabethpol (Hohen!), Daghhestaniæ et Tuschetiae 5500' (Rupr!), prov. Talysh (C. A. Mey.), Persiæ borealis (Szov!) et in alpe Demavend 5-10000' (Buhse!), prope Musch Armeniae Turcice australis 4500' (Ky. Suppl. 687!), Armenia (Auch. 5146!).

β. crassifolia. — Adpresse nivea pannosa, calycis dentes subbreviiores. — *N. crassifolia* Boiss. et Bulise Aufz. p. 175.

Hab. in fæce Tchehordeh montis Elbrus Persiæ bor. (Buhse!), in montosis inter Asterabad et Schahrud (Bge!).

Humilior et magis diffusa *N. grandiflora*, calycis 3½ lineas longi molliter tomentosi nec asperuli dentes acuti quidem sed minime acuminato-aristati.

62. **N. marifolia** (Boiss. et Huet Diagn. Ser. II, p. 24) adpressissimè cinereo-canescens multicaulis caulis tenuibus elongatis subsimplicibus, foliis breviter petiolatis rugulosis crenatis truncato-subcordatis oblongis obtusis, floralibus parvis ellipticis et linearibus,

cymis paucifloris dissitis longe et laxe racemosis breviter pedunculatis, bracteis linearibus brevissimis, calycis violacei tomentello-sublanati incurvi ore obliquo dentibus breviter lanceolatis tubo quintuplo brevioribus superioribus longioribus, corollæ calyce $2\frac{1}{4}$ -plo longioris tubo breviter exerto, nuculis sub lente scrobiculatis ♀. *N. tenuicantha* Boiss. in Tchih. exs.

Hab. ad radices montis Techdaghi prope Erzerum Armeniae 6-700' (Huet!), in collibus prope Baibut (Bourg!).

Affinis *N. Mussini* sed caules tenuiores erecti 4-4½-pedales, folia magis elongata vix cordata, calycis dentes breviores obtusiores, nuculae obsoletius scrobiculatae.

63. N. stenantha (Ky. et Boiss. in Sched.) minutissime puberula pallide virens, caulinibus erectis tenuibus inferne striete ramulosis, foliis oblongis subcordatis obtuse crenatis inferioribus breviter petiolatis, cymis laxiuseule paucifloris strictissime racemoso-paniculatis inferioribus longe superioribus breviter pedunculatis, bracteis linearibus brevissimis, calycis costato-striati brevissime et crispule tomentelli subincurvi ore obliquo dentibus triangulari-lanceolatis acutiusculis tubo triplo brevioribus superioribus longioribus, corollæ tubo breviter exerto, nuculis ruguloso-tuberulatis ♀.

Hab. in rupestribus praeruptis vallis Teng prov. Musch Armeniae Turcicae 6000' (Ky. 510 a!).

Pedalis, folia inferiora sesquipollicaria, racemus semipedalis angustus. Affinis *N. marifolia* differt foliis majoribus cordatis, calyce adpresso tomentello, racemis angustis, nuculis rugulosis.

64. N. Ruprechti, pallida sub lente brevissime pubescens, caulinibus elongatis basi prostrata vel ascendenti ramosis, foliis pellucido-punctatis subtus canescentibus breviter petiolatis a basi truncatâ triangulari-oblongis acutiusculis profunde dentatis, cymis pauci et laxifloris breviter pedunculatis subpaniculatis racemosis, bracteis linearibus brevissimis, calycis eburnescens patule hirsuti valde incurvi ore obliquo dentibus lanceolatis acuminatis tubo quadruplo brevioribus superiore longiore, corollæ tubo breviter exerto, nuculis sub lente tenuissime malleatis ♀.

Hab. in Caucasi orientalis Daghestaniam mediâ prope Golotl et supra Anarskoi Koissu 2100'-3000' (Rupr!).

Pedalis et sesquipedalis, folia majora 9-10 lineas longa, calyx 4 lineas longus. Ex affinitate *N. Mussini* a qua ramis elongatis, foliis profunde dentatis non cordatis, nuculis minutis malleatis, etc., differt.

65. N. Cilicica (Boiss. p.c. Prodr. XII, p. 388. Diagn. Ser. I, 12, p. 64) viridis glabra vel sub lente minute puberula, caulinibus elatis simplicibus vel ramosis, foliis petiolatis cordatis ovatis grosse dentatis, superioribus et floralibus sessilibus acutius dentatis, verticillastris pauci vel multifloris sessilibus racemos terminales basi interruptos formantibus, bracteis numerosis subulatis hispidis calycis

tubo brevioribus, calycis glandulis sessilibus adpersi et hispiduli recti ore obliquo dentibus lanceolatis acutis tubo triplo brevioribus inferioribus profundius fisis, corollæ tubo tenui subincurvo calyce triplo longiore, nuculis tuberculatis apice vel undique muricatis ♀. *N. pyrenantha* Bth. in p.c. Prodr. loc. cit. forma multiflora. — *N. Andrica* Boiss. et Heldr. Diagn. Ser. I, 12 p. 64.

Hab. in regione subalpinâ et alpinâ montis Gheidagh Isauriae 6000' (Heldr!), Tauri Cilicici 6-7000' (Ky. 400! Balt!), monte Aslandach Cappadociae (Bal!), monte Cassio Syriae borealis usque ad cacumen (Boiss!), in Libano (Labiilt!), supra Bischerre (Ehrt!), supra Eden et ad Cedros (Boiss!), inter Yamouny et Dimam (Bl!), monte Garbi Antilibani (Ky. 20!), Hermione ad cacumina usque (Boiss!).

3. Pisidica. — Folia minora subcordata, verticillastri paniciflori, bracteæ abbreviatæ. — *N. Pisidica* Boiss. et Heldr. Diagn. Ser. I, 12 p. 65.

Hab. in saxosis montis Boudroun Pisidiae (Heldr!).

Species pulchra foliis 7-8 lineas vel sesquipollicem longis, verticillastris pauci vel multifloris, calycis dentibus acutis vel acuminatis variis (caules 1-2-pedales, calyx 4-5 lineas, corolla 12-13 lineas longi. *N. Andrica* nucule quedam submonstrose auctæ leves sunt, sed normales tuberculato-muricatas esse observo.

66. *N. glomerata* (Montb. et Auch. Ann. Sc. Nat. 1836, p. 46) minute glandulosa et papillari-pubescentia, caulis ascendentibus ramosis, foliis inferioribus breviter petiolatis cordatis ovatis obtuse crenatis superioribus sessilibus subauriculatis, verticillastris densis parvis sessilibus remotis longe spicatis, bracteis lanceolato-subulatis calyce dimidio brevioribus, calycis subulato-striati ad costas glanduloso-asperi recti ore obliquo dentibus lanceolatis acutis tubo triplo brevioribus anticis binis profundius fisis subbrevioribus, corollæ tubo curvato calyce subduplo longiore, nuculis obtuse tuberculatis ♀.

Hab. in Cappadociâ ad Euphratrem (Auch. 1727!), in montibus Ssosdag et Karabiyukle Cataonie (Hanssk!), in Libano (Ehrt!), supra Eden et Dimam (Bl!), in monte Hermon (Gail! Ky. 219! sub *N. septemcrenata*).

Valde affinis *N. septemcrenata* a qua sat differre videtur calyce ad costas aspero nec undique minute pruinoso, nuculis obtuse tuberculatis nec muricatis. Calyx vix tres lineas, corolla 9 lineas longi.

67. *N. curviflora* (Boiss. Diagn. Ser. I, 5 p. 22) tota adpresissime tomentella canescens, caulis elatis paniculatis, foliis petiolatis a basi truncatâ vel subcordatâ oblongis obtusis obtuse crenatis, floralibus minutis oblongo-lanceolatis brevissime petiolatis, cymis multifloris brevissime pedunculatis et sessilibus pluries dichotomis densis racemos longos interruptos formantibus, bracteis subulatis curvatis brevissimis, floribus parvis, calycis canescens breviter tubulosi fructiferi oblongi ore subobliquo dentibus oblongis obtusis tubo quadruplo brevioribus superiore sublongiore, corollæ cœruleæ pubescentis tubo calyce triplo longiore valde incurvo, nuculis ovatis tuberculatis ♀.

Hab. in siccis regionis montanae Syriæ littoralis in monte Carmel (Labill!), Libano circa Eden (Boiss! Bl!), in viâ inter Berythum et Damascum (Ehr! Gaill!), Palestinā in collibus prope Samariam (Boiss!) et Hierosolymam (Roth 322!).

Planta 2-3-pedalis, folia inferiora pollice parum longiora, panicula multiflora interdum pedalis, calyx vix 2 lineas, corolla 6-7 lineas longa. Facies *Micromeria* e sect. *Pseudomelissa*, floribus minutis in genere insignis.

68. N. nuda (L. Sp. 797) minute pubescens vel glabriuscula, caulis elatis simplicibus vel saepius striete paniculatis, foliis viridibus oblongis et oblongo-lanceolatis basi cordatis crenatis obtusis inferioribus breviter petiolatis ceteris sessilibus, verticillastris secus paniculae ramos laxe racemosis breviter pedunculatis laxinusculis, bracteis subulatis brevissimis, calycis breviter pubescentis fructiferi ovali recti valde striati dentibus anguste lanceolatis tubo subbrevioribus, corollæ pallide violaceæ tubo subexerto, nuculis crebre tuberculatis apice pilosiusculis ♀. Ic. Fl. Græc. tab. 547. — *N. violacea* Will. Dauph. II, p. 367 an Linn? — *N. Pannonica* Jacq. Austr. tab. 129. — *N. paniculata* Crantz.

Hab. in graminosis et sylvaticis praesertim regionis montanae et interdum alpinæ, Græcia in Parnasso (Heldr! Orph. Fl. Gr. exs. 432!), Macedonia in monte Korthiati (Orph! Heldr!), Thraciâ (Friv!), Asiâ minore (Auch!), Armeniâ Rossicâ (Seidl!), Tauriâ et Caucaso (Led!).

β. albiflora. — Calycis dentes sublatiores evidentius lanceolati et pallide marginati. Corolla albida. — *N. nuda* β *parviflora* Bth. in D.C. Prodr. (sed flores eis typi non minores). — Ic. Jaeq. Austr. t. 24. — *N. Ucranica* M.B. II, p. 41 non L. — *N. alba* Desf. — *N. Pannonica* L. Sp. p. 797? — *N. sulphurea*! et *N. Pontica*! G. Koch Linn. XXI, pag. 677.

Hab. in montanis et alpinis formâ typicâ fere vulgator, Macedonia mons Korthiati (Orph!), Scardus (Griseb.), Lydie mons Tmolus (Boiss! Ball!), Lycia supra Elmalu (Bourg. 212!), Isauria (Heldr!), Cappadocia (Ky. Suppl. 207!), Armenia (Bourg! Ky!) et Pontus Lazicus ad Djimil 6000' (Ball), Tauria et regio Caucasica (Led. etc.), Persia borealis (Szov!), Mesopotamia (Noë 170!), Libanus 5-6000' (Ehr! Ky. 294!).

Ambo formæ omnino intermediis confluere videntur.

Ar. Geogr. Gallia australis, Hispania borealis, Germania australis et orientalis, Italia, Dalmatia, regio Danubialis, Rossia media et australis, Sibiria.

69. N. marrubioïdes (Boiss. et Heldr. Diagn. Ser. I, 12, pag. 67) tota molliter velutina canescens, caulis elatis paniculatum ramosis, foliis sessilibus ovatis cordatis, verticillastris dense multifloris longe et laxe racemosis inferioribus stricte et breviter pedunculatis, bracteis hirsutis linear-setaceis brevibus, calycis albo-hirsuti recti fructiferi oblongi dentibus strictis anguste lanceolatis tubo æquilongis acutis, corollæ albidae tubo incluso, achenis glabris minute tuberculatis ♀.

Hab. in regione inferiori montis Karadagh Lycaonie supra ruinas Larenda (Heldr!), in regione alpinâ Olympi Bithyni (Boiss! Pichl!).

Sesquipedalis, valde a finis *N. nudr* a qua specifice differre videtur foliis brevioribus latioribus, indumento velutino molli, corollæ tubo inclusa.

§ 8. Micranthæ.

70. **N. humilis** (Bth. in D.C. Predr. XII, p. 382) annua minutiſſime glandulosa et papillari-puberula, ascendens vel erecta basi ramosa, foliis inferioribus petiolatis ovatis basi cordatis vel truncatis grosse dentatis vel repandis, floralibus sessilibus elliptico-lanceolatis integris acuminatis, cymis multifloris densis inferioribus pedunculatis superioribus sessilibus racenos longos paniculæformes formantibus, bracteis lanceolatis et linearibus acuminatissimis calyce subbrevioribus, calycis longe tubulosi glandulosi et hirtuli subincurvi striati ore obliquo dentibus lanceolatis rectis longe acuminatis tubo triplo brevioribus superioribus latioribus et sublongioribus, corollæ tubo apice vix exerto; nuculis oblongo-trigonis lœvibus punctulatis ⊖.

Hab. in Persiâ (an potius Assyrâ?) (Auch. 1743!), montibus Avroman, Schahu, Pir Omar Gudrun Kurdistaniæ Persicæ 5-9000' (Haussk!), in subalpinis prope Mendeli (Noë 1160!).

Semipedalis et pedalis, folia inferiora pollicem longa, calyx florifer $3\frac{1}{2}$ lineas fructifer fere 5 lineas longus. A *N. micrantha* præter flores submajores nuculis, dentibus calycinis longe acuminatis distincta.

71. **N. micrantha** (Bge in Ledeb. Fl. Alt. II, p. 401) annua tenuiter pubescens, caule erecto simplici vel sæpius inferne ramoso, foliis petiolatis ovatis obtusis crenatis basi rotundatis vel subcordatis, floralibus sessilibus oblongis vel lanceolatis, cymis oppositis multifloris densis rarius paucifloris et laxiusculis inferioribus longiusculæ pedunculatis superioribus sessilibus racemum paniculæformem formantibus, bracteis linear-lanceolatis calyce triplo brevioribus, calycis patule pubescentis striati subincurvi ore obliquo dentibus lanceolatis tubo quadruplo brevioribus superioribus sublongioribus inferioribus antice inclinatis, corollæ hirtæ tubo exerto, nuculis fuscis obsolete malleatis vel subscrabiculatis ⊖. Led. Ic. tab. 412. — *N. Meyeri* Bth. Lab. p. 478. — *N. pallida* C. Koch Linn. XXI, p. 675!

Hab. in lapidosis et arenosis interdum salsis Armenie Rossicæ (C. Koch!), prov. Talyisch (C.A. M! Hohen!), totius Persiae borealis (Szov! Auch. 1746! et 5132! Buhse!), in toto ambitu deserti magni prope Schahrud, Chabbis, Kerman, Jezd, Téheran, Ispahan (Bge!), in Turkestaniâ (Lehm!), Belutschia (Stocks 1023!).

Variat semipedalis vel pedalis et elatior, calyx circ. tres lineas longus. Optime observante amic. Bunge *N. Meyeri* non specifice differt a *N. micrantha* cuius est forma magis australis cymis magis multifloris condensatis.

Ar. Geogr. Songaria, Sibiria Altaica.

72. **N. saccharata** (Bge Lab. Pers. 56) annua tenuissime pubescens, caule erecto ramoso, foliis inferioribus petiolatis ovatis

subcordatis grosse dentatis intermediis subsessilibus basi cuneatis, floralibus linearis-oblongis subintegris, cymis oppositis pedunculatis laxe dichotomis ample et laxissime paniculatis, bracteis linearis-lanceolatis calyce subtriplo brevioribus, calycis papilloso striati subincervi ore obliquo dentibus lanceolatis acentis tubo quadruplo brevioribus superioribus latioribus sublongioribus, corollæ tubo inclusu, nuculis pallidis grosse verrucosis verrucis lucidis ⊙ *N. Meyeri* var. *laxiflora* Ky exs. 1846.

Hab. in vallibus arenosis Persiae borealis in monte Elbrus prope Passgala (Ky 175!), in montibus Siokuh et Kohrud (Bode), in monte Sefi prope Ispahan (Kapherr), prope Ssof et Teheran (Bge!).

Pedalis et sesquipedalis, calyx tres lineas longus. Simillima *N. micrantha* sed inflorescentia laxissima et nuculæ majores sculpturâ dissimiles.

§ 9. Micronepetæ.

73. *N. petræa* (Bth. D.C. Prodr. XII, p. 394) annua, caule saepius elato ramoso ample paniculato, foliis adpresso lanatulis albidis inferioribus petiolatis ovalo-orbiculatis crenatis, intermediis oblongo-lanceolatis, floralibus subsessilibus oblongis cuneatis, cymis pedunculis longis filiformibus insidentibus confertim 3-5-floris, bracteis lanceolatis obtusis calyce brevioribus, floribus pedicellatis, calycis recti ad nervos asperuli dentibus rectis linearis-lanceolatis obtusinseculis tubo sesquibrevioribus superiore sublongiore, corollæ albidae puberulae tubo calyce quartâ parte longiore, nuculis oblongo-trigonis levibus ⊙.

Hab. in Persia lapidosis loco non notato (Auch. 1747!), in schistosis montis Avroman 5000 (Haussk!).

Specimina Ancheraiana semipedalia sed planta a cl. Haussknecht lecta pedes duo et amplius longa paniculâ ramosissimâ sesquipedali, folia intermedia pollicem et amplius longa, inferiora minora, calyx tres lineas longus, cymarum pedunculi bipollicares et flores pedicellis calyce vix brevioribus suffulti.

74. *N. hymenodonta*, annua adpresso tomentella canescens, caule erecto simplici vel inferne ramuloso, foliis parvis longe petiolatis ovalis crenatis, superioribus diminutis sessilibus, verticillastris 4-5 subsessilibus in spicam oblongam basi interruptam approximitis bracteis linearis-subulatis ciliatis spinuloso-acuminatis curvatis calyce vix brevioribus, calycis recti ad nervos papillari-hirti ore valde obliquo dentibus membranaceis superioribus longioribus ovatis acutiusculis, inferioribus profundius fissis triangulari-lanceolatis acutis, corollæ albidae hirtæ tubo subexserto, nuculis ovato-trigonis levibus sub lente punctulatis ⊙.

Hab. in Persia inter Yezd et Ispahan (Bge!).

Planta 8-10-pollicaris ab amic. a Bunge *N. petreæ* cuius specimina typica non viderat adnumerata sed distincta verticillastris sessilibus nec longe petiolatis, bracteis subulatis spinosis, calycis apice membranacei rubelli dentibus superioribus latissimis.

75. **N. Ispahanica** (Boiss. Diagn. Ser. I. 5 p. 23) annua breviter latulata, canule pumilo a basi ramuloso, foliis petiolatis ovatis obtusis obtuse crenatis insimis saepe subcordatis, floralibus ovato-oblongis acutis flores subæquantibus, floribus in capitula multiflora terminalia confectis, bracteis linearibus calyces æquantibus, calycis recti patule hirti dentibus subæquilonigis lanceolatis acutis inferioribus angustioribus et profundius fisis, corollæ parvæ tubo inclusa, nuculis minime punctulatis ♂.

Hab. in Persiâ prope Ispahan (Auch. 5143!) prope Teheran et Kohrud (Bode, Kapherr), in toto ambitu magni deserti salsi a Shahrud et Djendak ad Jezd et Kerman (Buhse! Bge!).

Planta 3-5-pollicaris, folia 6-7 lineas longa, capitula fructifera magnit. avellanae, calyces vix tres lineas longi.

76. **N. Schiraziana** (Boiss. Diagn. Ser. I. 7 p. 52) annua breviter canescens-velutina pumila basi ramosa, foliis petiolatis ovatis basi rotundatis vel cuneatis obtuse et late crenatis, floralibus oblongis sessilibus flores non æquantibus, floribus in capitula terminalia ovata longe pedunculata aggregatis, bracteis linearibus calycem æquaتابibus, calycis lanatuli recti dentibus lanceolatis æquilonigis et æquilateralis tubo subdupo brevioribus, corollæ roseæ tubo exserlo, nuculis grosse albe-tuberculatis ♂.

Hab. in argillos ad pedem montis Sabst Buschom prope Schiraz (Ky. 451! sub *N. Ispahanicâ*).

Planta 6-10-pollicaris, capitula magnitudine avellanae, corolla 4-5 lineas longa tubo angusto. Species valde affinis *N. Ispahanicæ* a quâ calyce et nuculis differt.

77. **N. pungens** (Bth. Lab. p. 487) annua asperulo-pubescentis erecta pumila simplex vel basi stricte ramosa, foliis inferioribus longe petiolatis minime ovatis crenatis intermediis subsessilibus longe cuneatis oblongis dentatis, floralibus lanceolatis acuminatis subintegris flores æquantibus, verticillastris panicis multifloris deusis subsessilibus laxè et breviter racemosis, bracteis nervosis lanceolatis apice curvatis spinoso-acuminatis calyce longicribus, calycis recti ad nervos pubescentis ore æqualis dentibus lanceolato-subulatis rigidis æqualibus tubo duplo brevioribus, corollæ albo-cærulescentis tubo subincluso, nuculis minimis trigonis laevibus nitidis ♂. *Zizyphora pungens* Bge Fl. Alt. I, p. 23. — Ledeb. Ic. tab 124. — *N. pusilla* Bth. Lab. p. 487.

Hab. in Persiâ inter Bagdad et Kermanchah (Oliv!), in argillosis montis Elbrus prope Derbend (Ky. 462!), prope Schahrud et Sof (Bge!), in convallibus regionis superioris montis Kuh Barfi prope Schiraz (Ky. 340!).

Planta 4-6-pollicaris, calyx $3\frac{1}{2}$ lineas longus. Species bracteis longis rigidis apice recurvis insignis. Cl. Bunge observante nullum inter plantam Persicam et Songaricam discriben.

Ar. Geogr. Songaria.

78. N. satureioides (Boiss. Diagn. Ser. I, 3 p. 24) annua puberula canescens, caule erecto a basi patule ramoso ramisque tenuibus rigidulis, foliis radicalibus parvis petiolatis ovatis ceteris anguste lanceolatis utrinque attenuatis subdentatis, floralibus linearis-subulatis acutissimis verticillastri longioribus, verticillastris sessilibus densifloris distantibus, bracteis linearis-lanceolatis acuminalis calvees et quantibus vel superantibus, calycis pubescentis recti equalis dentibus lanceolatis acuminato-spinulosis tubo aequilongis, corollae tubo breviter exerto, nuculis ovatis compresse trigonis levibus nitidis ♂.

Hab. in Persia prope Isfahan (Auch. 5147!), in montibus Sefā et Siokah (Boile), in apricis montium prope Schahrul, Taur, et alibi in regno Cabulico (Griff. 497!).

Semipedalis et pedalis, facies *Satureiæ hortensis*, folia intermedia pollicem et amplius longa $1\frac{1}{2}$ -2 lineas lata, verticillastri piso paulo maiores, calyx 2 $\frac{1}{2}$ lineas, corolla 4 lineas longa

79. N. Daenensis (Boiss. Diagn. Ser. I, 7 p. 53) annua sublente asperula, caule nano filiformi simplici vel ramuloso, foliis parvis lanceolatis vel linearibus subpetiolatis integris vel remote denticulatis, verticillastris in spicam terminalem ovatam condensatis, bracteis lanceolatis hispidulis calycem subæqualibus, calycis asperulo-pruinosi dentibus æqualibus subulatis tubo aequilongis, corollæ carneæ tubo inclusa, nuculis ovatis minicinis levibus ♂.

Hab. in lapidosis montis Elbrus prope Derbend et montis Totschal prope Teheran Persie borealis (Ky. 179! et 628!), in regionibus altioribus montis Kuh Daena Persie australis (Ky. 915!).

Plantula 2-4-pollicaris facie *Ziziphora*, folia majora 4-5 lineas longa sesquilineam lata, capitula avellana minora. Aflinis *N. pungeant* et *satureioides* differt inflorescentia capituliformi, etc.

80. N. bracteata (Bth. in p.c. Prod. XII, p. 393) annua sublente pruinosa subcanescens, caule pumilo filiformi a basi patule ramuloso ramisque rigidulis, foliis petiolatis enucleatis oblongis vel rhombicis remote dentalis, floralibus sessilibus capitula involuerantibus et saepe superantibus, bracteis numerosis nervosis oblongis vel ovatis mucronulatis subintegris calyces superantibus, verticillastris in capitula ovala congestis, calycis dentibus rigidis subulatis ciliatis tubo duplo brevioribus, corollæ tubo inclusa, nuculis oblongis levibus nitidis ♂. *Zaturia? humilis* Bth. in p.c. Prod. XII, p. 183.

Hab. in Persia prope Isfahan (Auch. 5136!), in monte Elbrus prope Derbend (Ky. 179!), prope Jezd (Buh-sel!), in aridis totius ambitus deserti magni salsti prope Mehr, Nichapour, Kerman, etc. (Bge!), Belutschia (Stocks 936! et 503!).

Facies *Ziziphora*, 3-6-pollicaris, folia floralia 5-8 lineas longa, capitula magnitudine avellanae. A *N. Daenensis* bracteis magnis latis, nuculis angustioribus, etc., distincta.

81. N. globiflora (Bge Del. Sem. Darp. 1859 et Lat. Pers. pag. 58) annua, canescens, caule pumilo filiformi divaricatum paniculoso, foliis caulinis paucis parvis petiolatis ovatis paucide lati-

floralibus majoribus sessilibus numerosis orbiculatis integris nervosis capitula involuerantibus et superantibus, floribus in capitula globosa congestis, bracteis oblongo-lanceolatis intimis submembranaceis calycis tubo brevioribus, calycis hirti dentibus subulatis rigidis tubo triplo brevioribus, corollae tubo tenui inclusa, nuculis ovatis levibus lucidis ⊖.

Hab. in aridis ad montem Shahkuh Persiae (Derderian), in prov. Khorassan prope Schahrud (Bge!).

Valde affinis *N. bracteatae* a qua differt foliis floralibus orbiculatis integris, bracteis membranaceis, calycis dentibus tubi ratione brevioribus.

82. *N. spathulifera* (Bth. in p.c. Prodr. XII, p. 380) annua! a collo ramosa ramis pumilis diffusa vel ascendentibus hispidulis parte inferiori foliatis, foliis minimis canescensibus longe petiolatis ovatis vel orbiculatis inciso-crenatis, spicis globose-capitatis lanatis longe pedunculatis basi interdum verticillastro remotiusculo auctis, bracteis longe stipitatis calyces aequalibus vel superantibus exterioribus ovatis ceteris elliptico-lanceolatis omnibus acuminatis, calycis membranacei lanati dentibus aequalibus lanceolato-subulatis rectis tubo suo subbrevioribus corollam subaequantibus, nuculis levibus ⊖.

Hab. in regno Cabulico (Griff. 500 in herb. Kew!).

Rami 6-8-pollicares, folia 4-6 lineas longa et lata, capitula avellanæ vix majora. Species bractearum formæ insignis.

SECT. II. OXYNEPETA.

83. *N. Ucranica* (L. Sp. p. 797 non M.B. et auct. plur.) tomentella vel glabrescens, caulis erectis patule paniculatis, foliis breviter petiolatis oblongis subcordatis crenatis, superioribus sensim diminutis subsessilibus lanceolatis, cymis pedunculatis laxe dichotomis, bracteis linearibus calyces aequalibus, calycis pubescentis obconico-tubulosi ore aequalis dentibus lanceolatis acutissimis rectis tubo aequilongis intus parce pilosis corollam cœruleam aequalibus vel subsuperantibus, nuculis apice subtuberculatis ♀. Ic. Rehb. tab. 42. — *Teucrium Sibiricum* L. Sp. 788. — *N. Sibirica* M.B. II, p. 41. — *N. parviflora* M.B. II, p. 41.

Hab. in campis et collibus siccis præsertim regionis montanæ, Tauria (Stev!), Caucasus (M.B. etc.), Karadagh (Szov!), ditio Talysch 4-500' (Hoht!), Persia borealis (Buhsel!), Armenia in neglectis Erzerum (Huet! Calv!), Capadoccia (Auch. 1545!).

Species ab auctoribus propositæ non nisi indumento valde varianti interdum subullo differunt, nuculae am. a de Bunge observante nunquam apice pubescentes.

Ar. Geogr. Transylvania, Rossia australis, Soongaria, Sibiria Uralensis et Altaica.

84. *N. heliotropifolia* (Lam. Dict. I, p. 711) inferne et ad

folia breviter et adpresso canescens ad inflorescentiam patule vel crispule pubescent., caulis erectis patule paniculatis, foliis breviter petiolatis oblongis basi angustatis subintegris, cymis pedunculatis lance dichotomis, bracteis linearibus calyces æquantibus patulo-recurvatis, calycis hispidae ore æqualis dentibus a basi lanceolata setaceis tubo longioribus areuato-recurvatis corollâ subbrevioribus, nuculis.... ♀.
N. trichodonta Boiss. et Noë Diagn. Ser. II, 4, p. 25. — *Satureja stricta* Russ. Aleppo II, p. 255.

Hab. in Oriente (Lam), in Mesopotamiâ prope Diarbekir (Noë 1941), in Kurdistanâ (Brant) et inter Van et Bitlis (Barré de Lancy!), Caeteri loci a cl. Bth. citati mihi ad *N. Ucranicam* vel species affines spectare videntur.

♂? major. — Calyces ut videtur majores sed planta fructifera tantum visa.

Hab. in Persiâ boreali prope Sengan inter Teheran et Tabris (Bge!).

Species a *N. Ucranicâ* ut am. a Bge animadvertisit foliis integris, bracteis calycisque laciniis patulo-recurvatis distincta. Corollæ majores.

85. ***N. curvidens*** (Boiss. et Bal. Diagn. Ser. II, 4, p. 26) brevissime et crispule tomentella, caulis patule paniculatis, foliis breviter petiolatis ovato-oblongis obtuse crenatis, superioribus et floralibus lanceolato-linearibus integris acutis mucronatis, cymis confertim 5-7-floris, bracteis linear-lanceolatis acerosis squarroso-patentibus calyces æquantibus vel superantibus, calycis dentibus lanceolatis subpungentibus subrecurvis tubo brevioribus corollam parvam violaceam superantibus, calyce fructifero ovato, nuculis apice obsolete tuberculatis ♀. *N. congesta* F. et M. Ann. Sc. Nat. 1854, p. 34.

Hab. in collibus et agris otiosis Cappadociae prope Karahissar (Bal!), ad Dundarli (Ky. 218!), ad Yaparlar Kieui prope Ouchak Phrygiae (Bal!), in Lycania (Tchih!).

Differet a *N. Ucranicâ* et *heliotropifolia* cymis confertifloris, a priori insuper bracteis et dentibus calycis recurvis, a posteriore calycis dentibus abbreviatis.

86. ***N. calycina*** (Fenzl Flora 1843, p. 400) adpresso canescens, caulis erectis rigidis fere a basi patule paniculatis, foliis brevissime petiolatis oblongis obtusis subintegris, rameis anguste lanceolatis acutis, cymis laxis dichotome 5-7-floris, bracteis linearibus calyces æquantibus, calycis canescens striati dentibus lanceolatis acutis rectis innocuis tubo duplo brevioribus corollâ parvâ carnâ subbrevioribus, nuculis malleatis ♀.

Hab. in agris argillosis Mesopotamiae inter Suerek et Diarbekir (Ky. 56! et 145!), inter Orta et Kharran (Haussk!), in collibus Armeniae australis inter Karput et Siwas (Noë 841!), in incultis prope Baibut (Bourg!).

Vix pedalis, facies *Marrubii parviflori*, a congeneribus differt calycis dentibus brevibus basi latis acutiusculis quidem sed non acuminatis.

87. ***N. cryptantha*** (Boiss. et Haussk. in Haussk. exs. 1865) patule et longiuscule papilloso-hispida, caulis erectis paniculato-

ramosis, foliis inferioribus breviter petiolatis ovatis et oblongis grosse et obtusiusculæ dentatis, floralibus sessilibus lanceolatis acutis integris, cymis dichotomis confertifloris, bracteis lanceolatis acuminatis calyces involucratis et aequalibus, calycis hispiduli striati elongati dentibus lanceolatis acuminatis rectis subpungentibus tubo æquilongis corollam parvam carnem superantibus, muculis malleatis apice tuberculatis. — ♀ *Oxynepetea inrotularu* Bge Lab. Pers. p. 59.

Hab. in agris argilloso Mesopotamiae (Auch. 4792!), circa Snerek et in deserto Chabur (Hausskt.), in Syria interiore circa Honus (Dr. Post!), in Persia prope Urmiah (Szov.), inter Ispahan et Teheran (Bge!).

Indumento ad caules et ramos praesertim atque foliorum margines pilis longis basi dilatatis aliis parvis viscidulis intermixtis constante, foliis profunde dentatis, bracteis, calycisque laciniis latis a precedentibus distincta.

Species delenda.

N. Vaillantii Bth. Lab. p. 471. Non in herbario Vahlii sed in herbario Jussiaci adest! Ex Oriente erronee indicata, est forma foliis subtalioribus *N. tuberosa*.

GLECHOMA (L. Gen. 714).

Calyx tubuloso-campanulatus rectus ore subæqualis quinquedentatus. Corolla tubus exannulatus sensim ampliatus, labium superius planum rectum bifidum, inferioris lacinia media plana obcordata. Stamina 4 approximata sub labio superiori parallela, antheræ per paria in cruce dispositæ rimis dehiscentes, muculae lœves. — Herba procumbens. — Genus *Nepeta* valde affine sed inflorescentia peculiaris et quædam in corollæ formâ discrimina genus Linnaeanum servare suadent.

1 **G. hederacea** (L. sp. 807) molliter hirta, caulis repentinibus radiculiferis ramos floriferos erectos sterilesque prostratos saepe elongatos edentibus, foliis petiolatis reniformi-orbiculatis cordatis grosse crenatis, glomerulis 2-3-floris brevissime pedunculatis axillaribus, bracteis brevibus setaceis, calycis hirsuti dentibus triangulari-lanceolatis acuminatis tubo tri-quadruplo brevioribus, corollæ tubo calyce subtriplo longiore ♀ Ic. Lam. III. tab. 505. — Relib. Ic. tab. 40.

Hab. in umbrosis praesertim regionis montane et in parte magis septentrionali ditionis, Laconia (Sibth.), Byzantii (Tchih.), Anatolia borealis (Thirke! Wied!), Pontus Laziensis ad Rhizé (Bal!), Tauria, Caucasus et Transcaucasia (M.B., Rupr!).

β. hirsuta. — Villosa, calycis dentes longiores. — *G. hirsuta* W. K. Pl. rar. p. 124 tab. 119.

Hab. in Rumelia (Friv!).

Ar. Geogr. Europa tota a Scotiâ et Scandinaviâ ad Rossiam, Sibiria, Japonia.

DRACOCEPHALUM (L. Gen. 729).

Calyx tubulosso-campanulatus rectus vel subincurvus saepius bilabiatus labio superiore tridentato inferiore bifido, dentibus labio superioris et praesertim supremo latioribus. Corollæ tubus exannulatus in fauce ampliatus, labium superius fornicatum retusum intus plieis destitutum, inferius patens trifidum lobo medio latiore. Stamina approximata sub labio superiore parallela apice antrosum curvata, inferiora breviora, antheræ biloculares loculis divaricatis. Stylus apice subaequilateris bifidus. Nuculae leves. — Herbae annuae vel perennes.

* Antheræ glabrae.

1. **D. botryoides** (Stev. Mém. Soc. Nat. Mosq. 3, p. 266) velutino-hispidulum cinereum, caulis proambientibus fragilibus, foliis parvis petiolatis ovato-roundatis profunde in lobos 5-7 ovatos vel oblongos obtusos margine subrevolutos pinnatifidis, verticillastris in capitulum ovatum confertis, bracteis basi longe cneatidis trifidis calyces equantibus, calycis pubescentis nervosi dentibus acuminatis inferioribus oblongo-lanceolatis superiore sublatiore ovato, corollæ pubescentis tubo subincurvo calyce vix longiore labio superiore brevi recto ♀.

Hab. in glareosis torrentium Caucasi orientalis, in alpe Tyfandagh (G.A. Meyt.), ad torrentem Chodjal (M.B.).

Caules 4-6-pollicares, foliorum lamina 3-4 lineas diam. lata, calyx demum 5 lineas longus. Species rara videtur a Ruprechtio non observata.

2. **D. Aucheri** (Boiss. Diagn. Ser. I, 5, p. 25) velutino-tomentellum viscidulum cinereum a collo ramosissimum, caulis brevibus prostratis interne nudis, foliis minutis petiolatis triangulari-orbiculatis in lobos 5-7 ovatos pinnatipartitis, verticillastris in capitulum densum ovatum confertis, bracteis oblongo-lanceolatis acutis basi attenuatis infinitis trilobis superioribus indivisis, eaiyeis crispile hirti nervosi subbilabiati labio superiore ultra medium trifido dentibus attenuato-subulatis superiore duplo latiore ovato, labio inferiore in dentes lanceolato-lineares subulato-attenuatos partito, corollæ pubescentis tubo brevissime exerto labiis subaequilongis ♀. *Dr. botryoides* Boiss. et Buhse. — Ky. exs. Pers. non Stev.

Hab. in summo monte Demavend Persiae borealis 11-1200' (Auch. 5134! Buhse!) in planitiis Besmischal ejusdem junci 9000' (Ky. 3531!)

Foliis parvis et habitu simile *D. botryoides* sed calyx fere dimidio minor evidentius bilabiatus, corollæ labia subaequalia.

3. **D. thymiflorum** (L. Sp. p. 831) sub lente breviter pubescens, caulis erectis tenuibus simplicibus vel parce et stricte ramosis, foliis petiolatis radicalibus cordatis ovato-orbiculatis flores superan-

tibus minute denticulatis et integris, verticillastris distinctis plurifloris longe racemosis calycis nervosi tubulosi subincurvi hirto-asperuli dente supremo ovato cuspidato cæteris lanceolatis subulato-attenuatis, corollæ tenuis purpurascens calycem vix superantis limbo nimuto ☽.

Hab. in montibus Ssamam, Elbrus et in valle Jolu Persiæ borealis (Buhse!), in prov. Asterabad prope Siaret (Bge!).

Ar. Geogr. Scandinavia, Rossia media et australis, Songaria, Turkestania borealis, Sibiria Uralensis et Altaica.

4. *D. multicaule* (Montb. et Auch. Ann. Sc. Nat. Ser. II, 6, p. 47) sub lente adpresso velutino-pubescentis, caulis e rhizomate lignoso pluribus pumilis simplicibus erectis rigidis foliosis, foliis parvis oblongo-lanceolatis integerrimis erectis basi in petiolum brevem attenuatis, floralibus sensim angustatis linearibus acutis calyce brevioribus, verticillastris bifloris racemum oblongum laxiusculum formantibus, bracteis lanceolatis integris aristatis, floribus magnis breviter pedunculatis, calyce hirtulo nervoso reticulato fere ad medium usque bilabiato, labii superioris dentibus ternis ovatis mucronatis, inferiore vix a medium bisido lobis lanceolatis apice subulatis, corollæ pubescentis calyce duplo longioris tubo exerto sensim ampliato ♀.

Hab. in Armeniæ monte Kechichdag (Montb.), prope Erzerum (Auch. 1675!), in alpinis Bimgöldagh (Ky. Suppl. 689!), in Persiæ borealis prov. Aderbidjan (Auch. 5217!), prope Tabris (Bge!).

Semipedale vel humilius, folia 5-9 lineas longa 2-3 lata, calyx 7 lineas, corolla 15 longi. Affine *D. peregrino* glabritie, foliis paucidentatis, distincto.

5. *D. Kotschy* (Boiss. Diagn. Ser. I, 7, p. 53) adpresso et crispule tomentellum cinereum, caulis e rhizomate duro pluribus pumilis simplicibus foliosis, foliis parvis breviter petiolatis obtusis inferioribus ovatis crenatis superioribus oblongis integris, verticillastris 2-4-floris in racemum laxiusculum ovato-oblongum aggregatis, bracteis oblongo-rhombeis incisis dentibus utrinque 2-4 longe spinulosis, calyce pubescente et glanduloso-punctato ad medium bilabiato labii superioris dentibus ternis ovatis mucronatis, inferiore profundius in lobos lanceolatos subulato-attenuatos fisso, corollæ flavidæ tomentellæ calyce duplo longioris tubo exerto sensim ampliato ♀.

Hab. in rupestribus Persiæ borealis in monte Totschal prope Teheran (Ky. 231!), in jugis Ssamam et Demavend usque ad 10000' (Buhse!), prope Schah-rud (Bge!), Persiae anstralis in monte Kuh Daëna (Ky. 577!). Persice *Alef Mast* vel *Palengemischk*, et cum lacte comeditur.

Semipedale, facies *D. multicaulis* a quo foliis crenatis et bracteis setigeris differt. Corolla 12-14 lineas longa.

6. *D. setigerum* (Boiss. et Huet Diagn. Ser. II, 4, p. 21) adpresso pubescens pallide virens, caulis e rhizomate duro pluribus pumilis simplicibus foliosis, foliis parvis inferioribus petiolatis ovatis obsolete repandis intermediis lanceolatis sessilibus, floralibus

et bracteis calyces aequalibus utrinque 2-3-dentatis dentibus longissime setaceo-aristatis, verticillastris 2-4-floris racemum ovato-oblongum laxiusculum formantibus, calycis nervosi hirtuli et glandulis sessilibus aspersi bilabiati labio superiore in dentes tres ovatos inferiore in lobos lanceolatos profundius fisso dentibus omnibus longissime setaceo-aristatis, corollae hirtae calyce duplo longioris tubo exerto sensim ampliato ♀.

Hab. in incultis prope Erzerum Armeniae (Huet!).

Semipedale, flores pollicares, ab affini *D. Kotschyi* foliis floralibus longo setuloso-dentatis et calycis dentibus multo longius cuspidatis diversum. Observandum tamen an transitus inter utrumque et *D. multicaule* non adsint.

** Antheræ villosæ.

7. D. Ruyschiana (L. Sp. 830) multicaule caulinibus erectis simplicibus foliosis tenuiter pubescentibus, foliis angustè linearilanceolatis integrerrimis muticis glabris margine subrevolutis, axillis fascieuligeris, verticillastris paucifloris in spicam ovatam basi interdum interruptam approxinatis, bracteis ovato-lanceolatis integris, calyce nervoso glabro vel puberulo vix ad medium bilabiati dentibus breviter mucronatis superiore latiore ovato cæteris lanceolatis, corollæ violaceæ pubescentis calyce plus duplo longioris tubo recto exerto ♀.

Hab. in pasenis lapilosis regionis alpinei Caucasi (Ledeb., C.A. Mey., Auch. 1758!) in Pschawiâ et Daghestaniâ Caucasi orientalis 6600' (Rupr!).

Ar. Geogr. Gallia in Pyreneis et Delphiniâ, Italia borealis, Norvegia et Suecia, Germania, Hungaria, Rossia septentrionalis, media et australis, Son-garia, Sibiria, Japonia.

8. D. Austriacum (L. Sp. 829) multicaule, caulinibus pubescentibus erectis simplicibus foliosis, foliis puberulis vel glabratibus in lacinias lineares revolutas mucronatas 3-5 pinnatipartitis, axillis fascieuliferis foliis interdum integris, verticillastris subsexfloris subdistinetis racemum ovatum vel oblongum formantibus, bracteis tritidis aristatis, calycis hirsuti nervosi transverse venosi subbilabiati dente supremo latiore ovato mucronato cæteris lanceolatis mucronatis, corollæ cæruleo-violaceæ puberulæ calyce triplo longioris tubo ampliato ♀. Jacq. Ic. rar. tab. 112.

Hab. in pratis alpinis montis Bousdonandagh Ponti Lazici 6600' (Bal!), in Caucaso (M.B., C.A. Mey., in Ossetia et Daghestaniâ 1800'-2500' (Rupr!).

Ar. Geogr. Gallia in Pyreneis et Galloprovinciâ, Helvetia et Germania in Alpibus, Bohemia et Gallicia, regio Danubialis, Rossia australis.

LALLEMANTIA (F. et M. Ind. VI. Petrop. p. 40).

Calyx tubulosus ore subobliquo dentibus quinis supremo latiore, inferioribus binis angustioribus. Corolla latium superius concavum subfornicatum intus plicis duabus arcuatis margini subparallelis

auctum, inferius patens trifidum. Stamina 4 approximata sublabio superiore parallela apice antrorsum curvata, inferiora breviora. Antheræ biloculares loculis divaricatis. Stylis apice subæqualiter bifidus. Nucula leves. — Genus a *Dracocephalo* galeæ plicis interioribus tantum distinctum.

1. **L. peltata** (L. Sp 831 sub *Dracocephalo*) annua glabra vel pubescenti-asperula, caule erecto simplici vel inferne stricte ramoso, foliis inferioribus petiolatis ovatis vel oblongis minute crenulatis, floralibus basi attenuatâ subsessilibus lanceolatis et linearibus sub-integris flores longe superantibus, verticillastris 6-10-floris spicas interdum longas et basi interruptas formantibus bracteis subpetiolatis reticulato-nervosis ovatis vel orbiculatis crenuncircâ crenâ et breviter aristato-dentatis calyce brevioribus, calyeis glabriuscali longitudinaliter et transverse nervosi dentibus mucronatis intimis lanceolatis, lateralibus oblongis, superiore lateralibus duplo inferioribus triplo latiore, corollæ cæruleæ calyce sublongioris tubo angusto ⊖. *L. peltata* F. et M. Ind. VI. Petrop. p. 53 — *D. ozymifolium* Mill. Dict. № 4 — *Zornia peltata* Mœch Meth 411.

Hab. in arenosis et arvis incultis, jugn o Parayadges Tonti (Tehilt!), Poutus Lazicus ad Djinoil (000') (Bal!). Armenia Turcica (Huett! Bourg!). Cappadocia (Auch. 1670!) Bal! Ky!). Kurdistania (Brant). Caucasus australis (1300m!) et orientalis 6-8000' (Rupr!), prov. Karabagh et Talysh (Hoh!), Persia borealis (Ky! Buhse!).

2. **L. Royleana** (Wall. in Bth. Lab. p. 497 sub *Dracocephalo*) annua breviter et crispare pubescens canescens simplex vel basi ramosa, caulis erectis rigidis sepius puniulis, foliis radicalibus longe petiolatis a basi truncatâ vel breviter cuneatâ ovatis obtuse multierenatis, floralibus subsessilibus oblongo-cuneatis paniculatatis flores superantibus, verticillastris 4-6-floris inferioribus subdistans stricte spicatis, bracteis petiolatis cuneati-flabellatis utrinque longiusculæ 2-3 aristato-dentatis, calycis patule hirti tubulosi nervosi dentibus oreibus submuticis tandem convointis et tubum claudentibus superioribus ovatis inferioribus binis subangustioribus, corolle cæruleæ tabo inclusa limbo minuto. ⊖ *L. Royleana* Bth. in dc. Prodr. 12, p. 404. — *D. Inderiense* Kar. et Kir. Bull. Mosq. 15, p. 423. — *Nepeta erodiifolia* Boiss. Diagn. Ser. I, t. 5, p. 24.

Hab. in arenosis apricis Persie prope Teheran (Ky. 42!), Ispahan (Auch. 5130!), Schiras (Haussk!), ubique circa desertum magnum (Bge!). Turkestania (Lehm!), Afghanistan (Griff.), Belutschia (Stocks!).

Facies *Ziziphore*, foliorum radicalium lamina 5-9 lineas longa, calyx ex *L. peltata* diuidio minor.

Ar. Geogr. India boreali-occidentalis.

3. **L. Iberica** (M. B. Taur. Caue. t. 2, p. 54 sub *Dracocephalo*) annua glabrinseula, caule erecto simplici vel inferne ramoso, foliis inferioribus breviter petiolatis oblongis obsolete repando-crenatis,

floralibus sessilibus linear-lanceolatis obtusis flores duplo superantibus, bracteis petiolatis nervosis cuneatis ovato-flabellatis apice longe 5-7-dentato aristatis, calycis nervosi aspernli vel glabrai dentibus macronatis superioribus ovalis supremo latiore inferioribus lanceolatis, corollae carunculae rarius lutescentis tubo angusto inclusa limbo parvo ♂. *L. Iberica* F. et M. loc. cit. — *D. aristatum* Bertol. Comm., Bononi. 6, p. 225. — *L. sulphurea* G. Koch Linna. XXII, p. 679.

Hab. in agris incultis haemosis et saxosis regionis inferioris et subalpinarum, Lycia ad Elmali (Bour., 211!), Almadagh Phrygiae (Bal!), Anatolia borealis ad Kastamouni et Angora (Wied!), Cappadocia (Bal!), Cilicia ad Pylas (Bal!), Armenia Turcica (Koch! Bourgl.) et Rossica (Koch!), ditio Talysh (e. A. M!), Persia borealis (Boiss! Bubse!) et australis ad Ispahan (Aueh 5133!), Kurdistania Persica in monte Pir Gudrun et Mesopolanaria (Hanssk!), Syria et Aleppo (Aueh! Ky! Hausk!), Palestina inter Hierosolymam et Jericho (Boiss!), Cœles-syria (Boiss!).

Diderit a *L. Roggweilii* calyces duplo longiore (6 lineas longo), caule crassiori magis protero, foliis longioribus angustioribus subintegris, a *L. peltata* bracteatum forma. Colitur in Per le prov. Kerman sub nomine *Gundschu Sighah* (Se amnum ingrimum), oleum semina præbent (Bge.).

4 L. canescens (L. sp. 831 sub *Draeocephalo*) biennis breviter tomentoso-canescens, caulis a radice indurata pluribus ascendentibus crassiuscantis simplicibus, foliis inferioribus petiolatis crenatis oblongo-lanceolatis, floralibus sessilibus linearibus calyces superantibus, verticillastris 4-6-floris laxiuscente spicatis, bracteis parvis subsessilibus nervosissimis cuneato-flabellatis 5-7-dentato-aristatis, calycis tomentelli dentibus numerosatis supremo latiore ovato-oblongo inferioribus lanceolatis, corolle calyce plus duplo longioris cerulea tubo in faucem sensim ampliato ♂. *L. canescens* F. et M. loc. cit.

Hab. in agris incultis regioni montane et alpine, Syria (Russell ex Bth.), Armenia Turcica (Tchih!), ad Baibut (Bourg!), in monte Techdaghi (~800) (Huet!), in Persia borealis monte Elbrus (Ky. 761).

Caules siepe pedates, calyx 7-8 lineas, corolla 14-15 longi. A precedentibus corolle tubo longe exerto et ampliato distincta.

L. brachystegia — Folia profundius crenato-dentata, bracteæ calyce quintuplo minores breviter aristatae. — *L. azurea* Boiss. et Huet olim.

Hab. circa Erzerum (Huet!), in monte Palanteuken (Ky. Suppl. 6751).

HYMENOCRATER (Fisch. et Mey. Ind. Secund. Petrop. p. 39).

Calyx tubo cylindrico quindecimnervi limbo fructifero valde aucto membranaceo rotatim expanso reticulato profunde quinquelobo. Corolle tubus superne in faucem ampliatus, labium superius erectum bifidum inferius patens trilobum lobo medio maiore retuso. Stamina 4 didynaria sub galea ascendentia, postica longiora antheræ loculis divaricatis. Stylus apice subæqualiter bifidus. Nuculae ovoideæ levées

vel tuberculatæ. Suffrutices vel herbæ perennes calycibus *Molucellæ*. — Genus valde affine *Nepetæ* a quâ calyce tantum differt.

1. *H. bituminosus* (Fisch. et Mey. loc. cit. p. 39) fruticosus præter papillas tuberculiformes sparsas glaber viridis, ramis simplicibus, foliis resinoso-punctatis petiolatis ovato-oblongis cordatis obtusis obtuse crenato-dentatis, floralibus diminutis, cymis subsessilibus in verticillastra 2-3 approximata oblongo-spicata dispositis, bracteis oblongo-lanceolatis utrinque attenuatis, calycis glabris tubo sulcato-striato, fructiferi limbo chartaceo reticulato-nervoso tubo longiore in lacinias pallidas late ovato-suborbicularias obtusas mucronulatas profunde partito, nuculis minute verrucosis ♂. *H. Aucheri* et *H. secundiflorus* Jaub. et Sp. Ill. 18. V. p. 62 et 93, tab. 456 et 457.

Hab. in collibus aridis prope pagum Perodse distr. Khoi prov. Aderbidjan Persie (Szov!), prope Ardebil (Auch 5121!), inter Teheran et Tabris, Jezd et Ispahan (Bge!).

Frutex pedalis vel humilior, folia inferiora 9-10 lineas longa, corolla 7-9 lineas longa, calycis fructiferi laciniae 6-7 lineas longæ 5-6 latæ. Recte observante am. Bunge *H. secundiflorus* est tantum status junior plantæ corollis florum præcociorum sublongioribus. Corolla glanduloso-hirtella.

2. *H. calycinus* (Boiss. Diagn. Ser. I, 5 p. 41 sub *Sestiniæ*) fruticosus glabriuscens viridis, ramis simplicibus, foliis resinoso-punctatis petiolatis a basi truncatæ vel cordatæ ovato-oblongis obtusis obtuse dentatis, cymis axillaribus paucifloris breviter pedunculatis in verticillastra plura distantia dispositis, bracteis linear-lanceolatis, calycis tubo minute glanduloso sulcato-striato, fructiferi limbo chartaceo reticulato-venoso tubo longiore violascente in lacinias oblongas acutas basi attenuatas ad basin partito, nuculis grosse verrucosis ♂. *H. calycinus* Bth. in D.C. Prodr. 12, p. 408. — Jaub. et Sp. Illustr. tab. 458. — *H. bituminosus* Boiss. et Buhse Aufz. p. 177 non F. et M.

Hab. in Persiæ boreali in jugo Ssamam prope Pokideh (Buhse!), in præruptis prope Schahrud (Bge!).

Valde affinis *H. bituminoso* a quo potissimum laciñis calycinis angustatis (6 lin. longis 3 lati) differre videtur. Corollæ tubus hirsutus.

3. *H. pallens* (Bge Lab. Pers. pag. 61) fruticosus glaber pallide virens, foliis vix resinoso-punctatis breviter petiolatis cordato-oblongis obtusissimis paucidentatis, cymis axillaribus subsessilibus paucifloris inferioribus remotis, bracteis ovato-lanceolatis membranaceis acuminatis, calycis tubo glabro limbi fructiferi profunde quintupartiti albo-virentis membranacei laciñis latissime ovatis acutis, nuculis tenuiter verruculosis ♂.

Hab. in alpe Schahkuh Persiæ borealis inter Asterabad et Schahrud (Derderian in herb. Hort Petrop!), prope Ispahan (Auch. 5121!).

A præcedentibus calycis fructiferi limbo tenerrime membranaceo albo laciñis latioribus distinctus; an tamen am. Bge indicante illæ tres species in unam polymorpham conjungendæ? sed formæ intermediæ adhuc desunt.

4. H. macrophyllus (Bge Lab. Pers. p. 62) frutienlosus glaber, ramis elongatis simplicibus, foliis majusculis breviter petiolatis subcordato-ovalis acutissimum grosse et acute pluridentatis, cymis pedunculatis subsecundis multifloris remotis, bracteis linearisubulatis, calycis tubo striato minute glanduloso, fructiferi limbo tubo longiore ad basin usque in lacinias chartaceas oblongas cuspidatas partito, nuculis nigris grosse verrucosis ♂.

Hab in jugo Elbrus prope Djehannameh ad orientem Teheran (Bode). Vidi in herb. Hort. Petrop.

Rami floriferi $1\frac{1}{2}$ -2-pedales, folia inferiora inter se distantia $1\frac{1}{2}$ -2-pollices longa basi pollicem lata, superiora sensim minora, pedunculi cymarum inferiorum pollicares, calycis laciniae demum semipollicares utrinque attenuate ciliolatae versus medium vix ultra duas lineas late. Corolla sex lineas vix excedens sub anthesi calycem non superans.

5. H. longiflorus (Bth. in B.C. Prodr. 42 p. 40) totus papillis sparsis hirtus viscidulus viridis, caelibus floriferis herbaceis elongatis erectis, foliis majusculis subtus elevatim nervosis cordatis ovalis obtusis obtuse et grosse dentatis inferioribus breviter petiolatis cæteris sessilibus, floralibus minutis acutis inlegris, cymis multifloris distantibus inferioribus longiuscula superioribus breviter pedunculatis longe et stricte subpanienlato-racemosis, bracteis herbaceis lanceolatis acutis, calycis floriferi tubo papillari-hirto viscidulo striato limbi lacinii rubellis lanceolatis acutis fructiferis valde auctis purpureis ovatis micronatis ultra medium partitis tubo multo longioribus, nuculis levibus ♀. *H. Haussknechtii* Boiss. et Rent. MSS.

Hab in montibus Avroman et Schahn Kurdistaniae Persiae 5-9000' (Haussk! Auch. 1534! undé? non ex Persia nec Armeniâ).

Caules 1-2-pedales, panicula stricta saepe pedalis, folia inferiora sesquipollicem longa et lata. Corolla caerulescens eis congenerum fauce et limbo amplior. Antheræ longe exsertæ! Calycis fructiferi limbus rotatus tandem diametro pollicaris et ultra.

6. H. paniculatus (Bge Lab. Pers. p. 62) basi fruticulosus, ramis floriferis herbaceis pubescentibus elongatis stricte paniculatis, foliis minute puberulis inferioribus petiolatis late cordato-ovatis subintegris superioribus subsessilibus oblongis grosse et irregulariter paucidentatis, cymis remotis inferioribus longe superioribus breviter pedunculatis, bracteis petiolatis oblongis scariosis, calycis tubo striato papilloso et glanduloso-hirto, fructiferi limbo rotato tubo longiore pellucido albo ultra medium in lacinias orbiculares obtusissimas partito, nuculis levibus dorso tenuissime nervosis ♂.

Hab in præruptis rupestribus vallium prope Schahrud Persie borealis (Bge).

Rami floriferi $1\frac{1}{2}$ -2-pedales, folia inferiora $1\frac{1}{2}$ -2-pollices longa et basi lata. Panicula pedalis et longior, cyme confertiflora inferiores pedunculis bipaniculatis suffulte, calycis fructiferi limbus diametro 10-12 lineas latus lacinii 5-6 lineas longis et latis. Corolla 6-7 lineas longa sub anthesi calyce longior hirtula.

7. *H. elegans* (Bge Lab. Pers. p. 63) basi fruticulosus ramis numerosis erectis herbaceis pubescentibus simplicibus, foliis breviter petiolatis pubescentibus resinoso-punctatis ovatis inciso-crenatis basi truncatis vel subcordatis summis diminutis sessilibus, cymis paucifloris inferioribus brevissime pedunculatis summis sessilibus verticillastra 2-4 valde remota formantibus, bracteis minutis subscariosis linearis-oblongis, calycis tubo hirsuto nervoso, fructiferi limbo rotato intense roseo-violaceo tubo duplo longiore in lacinias reticulatas ovalo-orbiculares obtusas ad $\frac{2}{3}$ partito, nuculis levibus nervosis ♂.

Hab. in declivitate meridionali montis Elbrus prope Djehannamch, Firuskuh et Schahkuh (Bodel), in subalpinis supra Tasch in via inter Asterabad et Schahrud (Bge!).

Rami hornotini $\frac{1}{2}$ -1-pedales, folia semipollicem tantum longa. Calycis fructiferi limbus valde auctus 14-16 lineas diametro latus. Corolla curvata 9 lineas longa, anthere subinclusa. Species calycibus etiam sub anthesi intense coloratis elegantissima.

8. *H. sessilifolius* (Bth. in D.C. Prodr. 12 p. 406) minute glanduloso-pubescentis, caulis herbaceis elongatis basi stricte ramulosis, foliis sessilibus cordatis ovatis rigidulis acute et subduplicatis dentatis, superioribus diminutis triangulati-lanceolatis, cymis multifloris verticillastra 3-5 remota formantibus inferioribus breviter pedunculatis ceteris sessilibus, bracteis scariosis late ovatis acutis, calycis tubo striato glanduloso-hirsuto fauce villosa, fructiferi limbo scariose reticulato purpurascente tubo longiore in lacinias ovato-oblongas acuminatas ad $\frac{2}{3}$ partito, nuculis obsoletae punctulatis ♀.

Hab. ad Kalou et in collo Erak regni Cabulici (Griff).

Saltem quoad ramos floriferos pedales herbaceus, folia inferiora 1-1 $\frac{1}{2}$ -pollicaria, calycis (fructiferi sed immaturi) laciniae 6-7 lineas longae. Inter affines foliis sessilibus insignis.

9. *H. incanus* (Bge Lab. Pers. p. 63) totus pube brevi crispata densa canescens multicaulis ramis erectis herbaceis basi ramulosis, foliis parvis cordato-oblongis crenato-dentatis inferioribus petiolatis summis oblongis subsessilibus, cymis 1-3-floris axillaribus oppositis remotis inferioribus pedunculatis summis subsessilibus, bracteis herbaceis oblongo-linearibus calycis tubum vix aequantibus, calycis floriferi tubo striato pubescente limbi laciniae linearis-lanceolatis demum ampliatis oblongo-lanceolatis coloratis acuminatis, nuculis.... ♀.

Hab. in Persia occidentali mediâ inter Ispahan et Kaschan prope Murtchehar, Ssof et Kohrud (Bge!).

Herbaceus nec basi suffruticosus, caules sub anthesi 7-10-pollicares, folia 6-7 lineas longa. Calyx sub anthesi 4 $\frac{1}{2}$ -5 lineas longus, laciniae post anthesin jam elongata 4-5 lineas longae 1 $\frac{1}{2}$ latæ roseæ. Corolla 7 lineas longa, anteræ juniores exsertæ. Species pube cinereâ, cymis paucifloris distincta, facie N.peter, quoad calycis fructiferos nondum nota.

HYPOGOMPHIA (Bunge in Bull. Acad. Pétersb. XVIII, p. 30).

Calyx campanulatus membranaceus obsolete quinque nervius sub-equaliter quinquedentalatus. Corolla tubus exannulatus leviter curvatus breviter exsertus in fauce vix dilatata, limbi labium superius elongatum angustum subincurvum, inferius patens trifidum lobis lateralibus oblongis intermedio majore obocordato. Staminia binas sub labio superiore arcuato-ascendentia filamentis ultra antheram breviter productis; antherae loculis confluentibus uniloculares. Staminodia binas antica apice clavata. Stylus breviter et inaequaliter bifidus. Nuculae obtuse tenuissime granulatae. — Herbae annuae nrae.

1. H. nana (Bth. in Bth. et Hook. Gen. Pl. II, p. 1201) annua nana simplex tota molliter et longe pilosa, foliis oblongo-spathulatis integris vel apice obtuse et obsolete crenatis, verticillastris 1-2. sub-distantibus, floribus in axillis solitariis subsessilibus, calyce florifero obconico-cylindrico fructifero campanulato aneto dentibus brevibus late triangularibus, corollae albide tubo breviter exerto ⊖.

Hab. in regno Cabulico prope Syghan (Griff!). Vid. in herb. Kew.

Plantula 1-2-pollicaris, folia 3-4 lineas longa. Differre videtur ab *H. Turkestanica* Bge extra fines nostros obviā que ramosa, cuius folia sesquipollucaria dicuntur et flores in axillis gemini.

Tu. VI. STACHYDEÆ.

TAPEINANTHUS (Boiss. Diagn. Ser. I, 42, p. 68).

Calyx campanulatus decemnervius obscure bilabiatus dentibus triangularibus vix inaequalibus superioribus subbrevioribus, fructifer subinflatus. Corolla tubo exerto exannulato, fance vix dilatata, limbi pauci labio superiore breviter lanceolato planiusculo subbifido inferioris trilobi lobo intermedio latiore. Staminia quatuor fertilia filamentis brevibus inclusis subparallelis inferioribus sublongioribus, antherae loculis confluentibus subuniloculares oblongae filamentis dorso medio inter loculos affixa. Stylus inaequaliter bifidus lobo superiore truncato, inferiore basi superiore amplectente. Nuculae magnae oblongae basi subangustatae facie interiori carinatae. — Herbae annuae pumilæ odoris ingrati facie *Odontitem vernam* referentes.

1. T Persicus (Boiss. Diagn. Ser. I, 42, p. 63) annus glanduloso-viscidus sordide virens simplex vel a collo pluricaulis, caule erecto pumilo, foliis uninerviis remote et ascendentib- denticulatis infe-

rioribus oblongis longius superioribus lanceolatis breviter petiolatis, floribus in axillis superioribus binis oppositis breviter pedunculatis folio longe superatis, calycis extus et intus breviter puberuli dentibus triangularibus acutis sub anthesi tubo sextuplo brevioribus, corollæ albidae roseo marginatae tubo calyce plus duplo longiore ⊖.

Hab. in siccis demissioribus totius Persiae, in arenosis prope Teheran (Ky. 48!), in arenosis salsis distr. Khoi prov. Aderbidjan (Szov.), in ambitu toto deserti magni ad Djendack (Buhse!), ad Schahrad, Kerman, Yezd, Ispahan, Kohrud (Bge), in Turkestani e deserto Kisilkum (Lehm!).

Planta 4-4-pollicaris, raro (ex Bge) subpedalis, folia 6-10 lineas longa 2-3 lata, superiora sensim angustata, calyx $2\frac{1}{2}$ lineas, corolla 5-6 lineas longa.

2. **T. Brahuicus** (Boiss Prodr. Ser. II, 4, p. 29) annus albo-hispida a collo saepius multicaulis caulinis pumilis erectis, foliis integris lanceolatis acutis et acuminatis in petiolum brevem attenuatis, floribus in axillis superioribus binis oppositis breviter pedunculatis folio longe superatis, calycis longe et palea extus et intus albo-hispida dentibus triangulari-lanceolatis subulato-acuminatis sub anthesi tubo triplo brevioribus, corollæ roseæ tubo breviter exerto ⊖.

Hab. in Persia austro-orientali prope Kerman (Bge), in Belutschia (Stocksl.).

Facies praecedentis a quo differt indumento, foliis integris, calycis dentibus longioribus, corollæ tubo breviore.

CHAMESPACOS (Schrenk Enum I, p. 27).

Calyx campanulatus deceinnervius in dentes aequales longe setaceo-attenuatos ultra medium fissus. Corollæ tubus filiformis exsertus in faucem subampliatus labio superiore plano emarginato inferiore brevi trifido. Stamina quatuor fertilia exserta ascendenter approximatæ subaequalia, antherarum loculi divaricati. Syllos apice subaequaliter bifidus. Nuculae elongatae oblongo-lineares laeves lateribus et praesertim apice anguste alatae — Herba annua pumila. — Genus affine *Tapeinanthe* quem facie refert et a quo staminibus exsertis et nuculis apice alatis differt.

1. **C. ilicifolius** (Schrenck loc. cit. p. 28) annus glabriusculus, caule erecto pumilo simplici vel inferne ramoso, foliis rigidulis breviter petiolatis oblongis et oblongo-lanceolatis longe spinuloso-acuminatis et spinuloso-dentatis, floribus binis axillaribus oppositis subsessilibus, calycis extus parce et ad faucem crebre hirsuti dentibus a basi latiore longe subulatis spinescentibus incurvo-patulis, corollæ violaceæ tubo dentibus calycinis sublongiore ⊖.

Hab. in sabulosis desertorum Karakum et Kisilkum Turkestaniæ (Lehm!).

Planta 4-5-pollicaris, folia inferiora 9-10 lineas longa, floralia diminuta, calyx demum 5 lineas longus, nuculae angustæ 2 lineas longæ.

Ar. Geogr. Songaria.

SCUTELLARIA (L. Gen. 734).

Calyx campanulatus bilabiatus labiis integris brevibus latis fructiferis clausis, superiore dorso squamâ rotundatâ concavâ (sepalo postico sepalis lateralibus ei introrsum adnatis quasi excluso Blh.) instructo deciduo. Corolla bilabiata tubo longe exerto. Stamina 4 approximata sub labio superiore parallela apice antrosum curvata, antheræ ciliatæ staminum superiorum divaricatum bilobulare inferiorum uniloculares loculo altero obsoleto. Styli lobus superior brevissimus. Nuculæ subglobosæ sœpius tuberculosæ. — Herbae basi interdum suffrutescentes.

Specierum Orientalium distributio.

SECT. I. LUPULINARIA Hamilt. — Flores non secundi, tetragonospati vel racemosi.

* Flores in spicam plus minus tetragonam congesti.

S. Orientalis, virens, pinnatifida, alpina, glchomoides, pectinata, Haussknechtii, Stockii.

** Flores plus minus laxe racemosi.

S. fruticosa, cretacea, multicaulis, Szovitsiana, Chorassanica.

SECT. II. VULGARES. Benth. in Benth. et Hook. Gen. II, p. 1202. — Flores oppositi racemosi secundi.

1. *Salviæfolia*. — Decumbentes sœpius humiles, facies præcedentium sed flores secundi.

S. salviæfolia, Pontica, diffusa, heterophylla.

2. *Peregrinæ*. — Sœpius majores erectæ, rarius decumbentes, folia floralia caulinis plus minus ve dissimilia.

S. Tournefortii, Columna, peregrina, altissima, a'bida, Sieberi, hirta, nutriculata, glutinosa.

3. *Galericulatæ*. — Erectæ, folia floralia caulinis subconformia.

S. galericulata, hastifolia.

SECT. I. LUPULINARIA.

1. S. Orientalis (L. Sp. p. 834) ramosissima ramis induratis ascendentibus abbreviatis pubescentibus vel canis. foliis parvis petiolatis ovatis vel oblongis inciso-dentatis supra virentibus glabris vel pubescentibus subtus adpresso canis, floribus in spicas oblongas tetragonas confertis foliis floralibus membranaceis ovato-oblongis integris imbricatis calyces occultantibus. corollæ longâ pubescente flavâ vel labio inferiore fuscescente ♀.

Variat multum foliorum magnitudine, divisione et indumento, foliis floribus plus minus dilatatis, molliter villosis vel glabratis. Formæ sequentes intermedii omnino confluent.

z. genuina — Folia majora crenato-dentata. — Ic. Fl. Græc. tab. 580. — Rehb. Ic. bot. tab. 8. — *S. Caucasica* Hamilt. — *S. bicolor* Hochst. in Lorent Reis. p. 338 ?

Hab. in saxosis regionis montane et subalpinæ, Græcia in Parnasso (Heldr. Herb. Norm. 684!), Anatolia in Lyciâ (Bourg!), Cariâ (Boiss!), Bithyniâ (Griseb!), monte Amano Syriæ borealis (Ky. 82!). Armeniâ!, Caucaso et Transcaucasiâ!, Persiâ boreali (Auch! Bunge!), Tauriâ!

β. pinnatifida. — Folia minora profunde pinnatifida et interdum usque ad nervum medium pinnatipartita. — Ic. Jaub. et Sp. Ill. tab. 412.

Hab. in saxosis regionis subalpinæ. Parnassus et Velugo (Etoliae (Heldr! Sprn!)), Thracia (Griseb!), Galatia (Wied!), Phrygia (Bal!), Armenia (Tchih! Bourg!), Transcaucasia, Persia borealis (Bge!), Kurdistaniæ mons Pir Omar Gudrun (Haussk!).

γ. atropina. — Rami tenuiores prostrati sœpe radicantes, folia parva utrinque grisea vel canescens, spica sœpius abbreviata pauciiflora.

Hab. in regione alpinâ, Olympus Bithynus (Sibth! Auch! Boiss! etc.), Tmol et Cadmî cacumina (Boiss! Bal!), mons Berydagh Cataoniac (Haussk!), Taurus Cilicus (Ky. 174! Bal!), Libani regio superior (Ehr! Auch. 1610! Boiss! Bl!), Persia borealis in Ghilan (Auch. 5155!) et prov. Khorassan (Bge!).

Ar. Geogr. Hispania australis, Dalmatia, Croatia, Sibiria Uralensis, Turcomania.

2. S. virens (Boiss. et Ky. in Sched. 1859) glabra ramosissima ramis induratis prostratis vel ascendentibus brevibus, foliis majusculis petiolatis viridibus concoloribus a basi truncatâ ovato-oblongis profunde et obtuse inciso-dentatis, floribus in spicas subtetragonas oblongas confertis, foliis floralibus membranaceis ovalis acutis imbricatis calyces occultantibus, corollæ rectâ pubescente flavâ ♀.

Hab. ad parietes umbrosas rupium alpinarum vallis Teng in provinciâ Musch Armeniæ australis (Ky. 457!). Fl. Sept.

Facies *S. Orientalis* sed omnino glabra, folia pollicem et amplius longa utrinque lata viridia, spicæ similes sed bractæ acutiores et corollæ tubus paulo brevior. A *S. Alpinâ flaviflorâ* foliis profunde incisis glabris longius petiolatis, ramis rigidis induratis, etc., magis discelens.

3. S. pinnatifida (Arth. Hamilt. Mon. pag. 16, tab. 2) basi indurata suffrutescens molliter et patule pubescens vel villosa grisea, ramis ascendentibus simplicibus brevibus, foliis breviter petiolatis ovatis vel oblongis profunde incisis vel pinnatifidis laciniis linearibus obtusis, spicis tetragonis oblongis tandem elongatis et basi subinterruptis, foliis floralibus sessilibus ovato-rhombeis præter supraemam integram utrinque 2-3 dentato-incisis, corollâ longâ purpurascenit-fulvâ pubescente ♀.

Hab. in Persiâ inter Kermanchah et Hamadan (Oliv!), prope Mendeli (Noë 1485!), circa Ispahan (Auch. 5152!), in face Murch Mahal montis Elbrus (Ky. 167!), in montibus Avroman et Schahu Kurdistanae Persice ad nives 7-9000' (Haussk!).

Folia eis *S. Orientalis* interdum majora saepius griseo-villosa et profunde incisa, facies et floram magnitudo hujus speciei sed ab eâ foliis floralibus inciso-dentatis statim distinguenda.

4. S. Alpina (L. Sp. p. 834) pubescens vel glabrata a basi ramosissima ramis decumbentibus, foliis viridibus ovatis vel oblongis minute serrato-crenatis inferioribus breviter petiolatis superioribus subsessilibus acutis, floribus in spicas ovatas tetragonas densas tandem elongatas laxas congestis, floralibus membranaceis purpurascen-tibus oblongo-lanceolatis calyees occultantibus, corollâ pubescente violacea labio inferiore albido ♀. Ic. W. K. Hung. tab. 137.

Hab. in regione alpinâ Græciæ raro, mons Olenos in cacumine 6600' (Heldr!), Macedonia in monte Scardo (Griseb!).

Varietateni flavifloram, *S. lupulinam* L., e ditione non vidi.

Ar. Geogr. Hispania, Gallia in Pyrenæis et Alpibus, Helvetia australis, Italiae alpes, regio Danubialis, Dalmatia, Rossia media et australis, Sibiria.

5. S. glechomoides (Boiss. Diagn. Ser. I, 12 p. 67) caudieulis filiformibus fragilibus nudis subterraneis in ramos floriferos brevissimos simplices abeuntibus, foliis parvis hirtis glandulosisque petiolatis ovatis obtusis basi truncatis repando-crenatis, spicâ terminali densâ 4-5-florâ, foliis floralibus submembranaceis purpurascen-tibus oblongis obtusis villosis calyce quadruplo longioribus, corollæ pubescentis violacea tubo elongato ♀. Ic. Jaub. et Sp. Illustr. tab. 377.

Hab. in alpinis Persiae borealis, in monte Elamout (Auch. 5158!), ad lacum congelatum Darja-i-fach montis Totschal prope Teheran (Ky. 568!).

Facies *Linarie pallide* Ten., caudiculi tenuissimi inter saxa repentes, rami floriferi vix pollicares 4-8-phyllici, folia 4-6 lineas longa, corolla frē pollicaris. Recete mōnente amici s. a Bunge *Lupulinariis* alnumeranda.

6. S. pectinata (Month. et Auch. Ann. Sc. Nat. 1836, p. 44) adpresso incano-tomentosa a basi inderatâ ramosa ramis erectis vel ascendentibus simplicibus pumilis, foliis parvis breviter petiolatis lanceolatis in laciniulas minimas margine revolutas obtusas confertas pectinatum pinuatifilis, floribus in spicae oblongam tetragonam cum-

fertis, foliis floralibus canis ovatis acutis integerrimis, corollâ purpureâ pubescente ♀.

Hab. in Cappadociâ ad Euphratem (Auch. 1609!). Errore schedulæ probab. ex Trapezunte indicata.

Caules semipedales, folia 4-7 lineas longa basi 2 lata, hâc notâ et floribus in spicas breves condensatis a *S. fruticosa* certe differt.

7. S. Haussknechti, breviter tomentosa cana, caulis elongatis prostratis opposite et patule ramosis subpaniculatis, foliis rugulosis petiolatis cordatis ovatis regulariter crenatis subtus intensius canis, floribus in spicas oblongas congestis, foliis floralibus submembranaceis oblongis acutis imbricatis parvis, corollâ pubescente rubro-lateritiâ labio inferiore pubescente ♀.

Hab. ad Karput Armeniae australis (Haussk! semina in Hort. Valeyres culta!), circa Diarbekir (Ky. 312!).

Caules demum pedales et sesquipedales spicis numerosis pedunculatis, folia inferiore pollicaria, corollæ pollicares. A *S. fruticosa* caulis prostratis subpaniculatim ramosis, indumento adpressiore vix glanduloso, spicis densifloris; ab omnibus *S. Orientalis* formis caulis elongatis prostratis ramosis, bracteis parvis, corollis purpureis etc. distinctissima. *S. pectinata* differt ramis brevibus simplicibus rectis, foliis anguste lanceolatis pinnatifidis vel partitis.

8. S. Stockssii (Boiss. Diagn. Ser. II, 4. p. 28) molliter griseolanata basi suffrutescens ramosissima ramis vetustis prostratis duris annotinis brevissimis fragilibus ascendentibus foliosis, foliis parvis elliptico-rhombeis integris acutiusculis in petiolati brevem attenuatis, floralibus conformibus subdiminutis calyce longioribus, spicis paucifloris capituliformibus terminalibus ex cespite vix exsertis, corollâ hirtâ calyce pluries longiore ♀.

Hab. ad Chehel Tan Belutschiae (Stocks!).

Facies *Ajuga salicifolia* var. *tomentosa*, rami annotini 2-3-pollicares, folia majora 6 lineas longa 3½ lata, capitula parva foliis fere occultata subquadri-flora, corolla 10-11 lineas longa. Species inter gregem *S. Orientalis* et *S. hirtæ* subdubia.

9. S. fruticosa (Desf. Cat. Par. p. 63) tomentosa et glandulos-viscidula basi indurata ramis numerosis erectis vel ascendentibus virgatis sæpius elongatis simplicibus vel opposite ramosis, foliis parvis rugosis oblongis obtuse inciso-crenatis basi truncatis utrinque tomentosis subtus canis, floralibus herbaceis ovatis acutis calycem parum superantibus, floribus oppositis laxe racemosis, corollâ longâ flavâ labio inferiore sæpe purpurascente ♀. *S. tomentosa* Bert. Nov. Comm. Bonon VI, p. 225.

Hab. in collibus calidis siccis regionis inferioris montanae et subalpineæ Syriae prope Aleppo (Russell, Ky. 237!), ad Birejjick et Aintab (Haussk!), in Colesyria (Ehr!), collibus circa Damascum (Boiss! Ky! Gaill!), Antilibano (Auch!), monte Hermon usque ad 6000' (Ky. 217!), ad Hierosolymum (Roth 333!), in Persiâ australi ad Persepolin (Ky. 391!), inter Fasa et Schiraz (Auch!).

Caules saepe 1-1½-pedales, folia inferiora 6-9 lineas longa, racemi tandem 4-6-pollicares. Corolla pollicaris. Nomen improprium nam pars inferior plantæ potius indurata quam fruticosa.

10. *S. cretacea* (Boiss. et Haussk.) tomentosa basi indurata et ramosissima ramis erectis simplicibus pumilis, foliis parvulis subtus densius canis reticulatisque petiolatis a basi truncatâ ovato-orbiculatis obtusissimis obtuse crenatis, floralibus oblongis acutis integris canis, floralibus paucis terminalibus laxiuscule oblongo-spicatis, corollâ longâ pubescente purpureâ ♀.

Hab. in cretaceis regionis superioris montis Sindjar Mesopotamie (Haussk!).

Rami numerosi 4-5-pollicares, folia 4-5 lineas longa lataque, corolla 13-14 lineas longa. Affinis *S. fruticosæ* a quâ differre videtur indumento non glandulosō, ramis brevibus, foliis minimis rotundatis, inflorescentiâ magis condensata.

11. *S. multicaulis* (Boiss. Diagn. N. Ser. I, 7, p. 61) basi suffruticosa ramosissima ramis tenuiter virgatis erectis simplicibus breviter puberulis, foliis parvulis cinerascenti-velutinis et subtus glandulis aspersis breviter petiolatis a basi truncatâ vel breviter cunealâ ovatis obtuse paucidentatis interdum repandis, floralibus paucis oppositis distantibus, foliis floralibus ovatis acutis viscidulis integris calyces subsuperantibus, corollâ pubescenti-viscidâ rectâ purpureâ vel flavo-rubellâ ♂. *S. nepetaefolia* Bth. in p.c. Prodr. 12, p. 414!

Hab. in regione alpinâ Persiae austro-occidentalis 8-10000' ubi cespites format, alpes Kuh Delu (Ky. 526!), montes Kuh Nur, Sawers, Eschker, Kellal et Ssebekuh (Haussk!), monte Elvend (Auch. 1931!), regio alpina Persie bor. orientalis inter Nischapur et Meschhed (Bge!).

Rami floriferi stricti 5-9-pollicares, folia majora 5-9 lineas longa, racemi 2-3-pollicares 6-10-flori, corolla eā *S. fruticosæ* multo brevior 8-9 lineas longa.

♂. *Cabalica* Bth. loc. cit. — Folia angustiora.

Hab. in regni Cabulici jugo Erak 9-12000' (Griffith!).

12. *S. Szovitziana* (Bge Lab. Pers. p. 65) basi suffruticosa ramis erectis tenuiter virgatis elongatis remote foliosis, foliis minutis sub-sessilibus firmulis obtuse crenatis vel subintegris supra glabris subtus elevatim nervosis et sub leute minute pulveralis oblongis acutiusculis, floralibus ovatis acuminatis calyce fructifero brevioribus, verticillastris subbifloris remotis, corollâ longâ purpureâ ♀.

Hab. in montibus Seidkhodji prov. Aderbidjan (Szov!). Vidi in herb. Hort. Petrop.

Caules cum spicâ pauciflorâ laxâ 12-15-pollicares, verticillastra 3-6 remota. Affinis *S. multicauli* sed corolla longior.

13. *S. Chorassanica* (Bge Lab. Pers. p. 65) canescens, caulis ascendentibus elongatis simplicibus, foliis villosulis et subtus glandulosō-punctatis petiolatis ovatis basi rotundatis crenulatis, floralibus ovatis acutatis integerrimis calycem æquantibus, floralibus

12-16 oppositis laxiuscole racemosis, corolla flavâ villosso-pubescente ♀.

Hab. in montibus province Khorassan Persiae ad austrum urbis Sebsewar sitis (Biepert). Non vidi.

Ex unico specimine nota, caules fere pedales, folia majora vix semipelticaria, Ex el. noctore affinis *S. multiflora*.

SECT. II. VULGARES.

1. *Salviæfolia*

14. *S. salviæfolia* (Bth. Lab. p. 433) caulis a collo indurato pluribus basi prostratis vel ascendentibus pumilis, foliis parvis griseis breviter velutino hirtis et glanduloso-adspersis supra lineatis subtus rugosis minute crenulatis longiuscole petiolatis a basi rotundato-cuneatâ vel truncatâ ovatis obtusis, floralibus breviter petiolatis oblongis obtusis subintegris calyce sublongioribus, floribus oppositis secundis densiuscole racemosis, corolla flava hispidulae tubo basi incurvo ♀.

Hab. in argillosis regionis montane et subalpine regio superior montis Davro dagh Pisidie (Heldr.), Galatia prope Tokat (Wied.). Armenia prope Baibut et Germeskhané (Bourg.). Cappadocia in argilosis Argei (Ky. Suppl. 220), Cataonia mons Akdaghi (Anch. 162!), Galicia in fancibus Gulek Boghaz (Ky. 462! et 357!) ad Allahdaghi prope Mersina (Bal!). Ex schedulâ herb. Sibthorpiani & Graciæ sed prob. erronee.

Caules cum racemo 3-4-pollicares, foliorum lamina pollice dimidio vix longior, racemus fractiter tripollicularis, corolla 8-9 lineas longa

15. *S. Pontica* (C. Koch Linn. XXI, p. 704) viridis, caudiculis ex collo numerosis tiliiformibus prostratis radicantibus nudis, ramis floriferis brevibus ascendentibus simplicibus minute puberulis, foliis prater ciliola glabris obovatis obtusis obsolete crenato-repandis basi subcuneatis inferioribus longiuscole petiolatis, floralibus subsessilibus calyces superantibus corollâ multo brevioribus, floribus paucis oppositis in racemum secundum brevem laxiusculum approximatis, corolle pubescenti-viscidæ purpureo-violaceæ tubo latiusculo basi incurvo ♀.

Hab. in lapidosis alpinis Ponti Lazici 6-1000' (C. Koch!), supra Djimil (Bal!).

Caudeculi filiformes sed duri 3-5-pollicares, rami floriferi 2-3 pollicares, foliorum majorum limbis 5-6 lineas longus, corolla fere pollicares. Facies *S. Alpina* a qua valde differt foliis longe petiolatis obtusissimis vix crenatis, floribus herbaceis obtusis, floribus secundis laxe racemosis nec dense et tetragone apicatis. Magis affinis *S. salviæfolia* a quâ distinguitur caudiculis elongatis nudis prostratis, corollis violaceis

16. *S. diffusa* (Bth. in p.c. Prodr. 12 p. 421) viridis breviter velutino-hirtella et crebre glanduloso-adspersa collo ramosissima ramis prostratis crassiuseulis brevibus radicantibus, foliis minutis

petiolatis ovatis suberanatis subtus elevatim nervosis, racemis brevibus, foliis floralibus petiolatis ovatis subintegris calyces fructiferos vix superantibus, corollā hispidā viscidā rubellā brevi ♀. *S. numularia* Fenzl in Ky. Scheld.

Hab. in regione alpina inferiore, Cappadocia in montibus Alidagh et Menendagh (Bal!), mons Berytdagh Cataonice 6000' (Haussk!), Taurus Ciliensis (Ky. 461!), mons Amanus Syria borealis in jago Tolo (Ky. 429!).

Ramii numerosi a basi cerebre foliosi simplices vel ramosi 2-4-pollicares, folia 2-4 lineas longa, racemi fructiferi 2-4-pollicares, corolla 5 lineas longa. Species a *S. salinifolia* affini ramis brevissimis crassiunculis, foliis minibus, corollā duplo breviore purpurascenti-fulvā egregie distincta.

17. *S. heterophylla* (Mouth. et Auch. Ann. Sc. Nat. 1836, pag. 45) caulinis e collo pluribus decumbentibus vel ascendentibus minute puberulis simplicibus vel parce ramosis, foliis parvis pallide virentibus breviter velutinis et glandulis adspersis supra lineatis subtus rugulosis minute crenulatis ruficulis petiolatis ovatis, caulinis oblongis et oblongo-lanceolatis lanceolatis subsessilibus, floralibus linear-lanceolatis corollā subbrevioribus, floribus oppositis breviter racemosis secundis, corollā hispido-viscidā flavidā ♀.

Hab. in montosis dumosis, Cilicia ad Tchaousli prope Mersina (Bdt!), Syria borealis prope Aintab (Auch. 461!), regio inferior montis Cassii et montes prope Antiochiam (Boiss!). Fl. ver.

Caules 1-1½-pedales virgati in racennum 2-3-pollicarem abentes, foliū radicalia breviora latiora 6-9 linea longa caulina interdum pollicaria 2-2½ lin. longa, corolla 7-8 lineas longa tubo fusi recurvo. Affinis *S. salinifolia* differt caulinis elongatis, foliis caulinis angustatis elongatis, floralibus calyces longe superantibus.

2. *Peregrinae.*

18. *S. Tournefortii* (Bth. Bot. Reg. XVIII ad calcem N° 1493) preter pubem brevissimam lente tantum perspicuum glabra, caulinis erectis elatis simplicibus vel parce ramosis, foliis petiolatis a basi truncata vix cordata ovato-triangularibus grosse crenatis, racemo unilaterali laxiunculo sub lente parce hirsuto, foliis floralibus ovato-rhombeis acutis calyce brevioribus, corollā e sicco purpureo-fusca hirtulā calyce hirsuta sextuplo longiore, seminibus sebris ♀. *S. altissima* var. *grandiflora* C. A. Mey. Ind. p. 89.

Hab. in sylvis et nemorosis Transcaucasiae ad Caspium prope Lenkoran et Astara (Höh! G.A. Mey!), Persia boreali maritimā prope Rescht (Buhse!), prope Asterabad (Kar!) et Siaret (Bge!).

Bi- et tripedalis, folia inferiora interdum 2-2½-pollucaria, corolla 10 lineas longa; species glabritie et corollis longis a sequentibus distincta.

19. *S. Columnæ* (All. Pelem 1. p. 40, tab. 84) caulinis elatis simplicibus vel parce ramosis undique pubescentibus, foliis breviter pubescentibus petiolatis a basi truncata vel subcordata ovato-oblongis grosse crenatis, racemo unilaterali vel secundo denum elongato laxo patule glanduloso-villoso, foliis floralibus subpetiolatis villosis ovatis

aculis calyce brevioribus, corollâ purpurascente pubescenti-viscidâ calyce sextuplo longiore, seminibus granulatis stellatim et minutissime puberulis ♀. *S. pallida* Guss. — *S. Gussonei* Ten.

Hab in umbrosis montanis, Latonia in monte Malevo (Orph. Fl. Gr. exs. 545!), Etolia prope Carpenisi (Sannar!), Eubea prope Kourbatsi (Wild!), Olympus Thessalus prope Hagios Dionysios (Orph!), mons Korthiatî Macedoniae (Orph!).

Bipedalis et elatior, folia inferiora 2-3-pollicaria, corolla 10-11 lineas longa. Folii majoribus, bracteis calyce brevioribus et corollis elongatis a *S. peregrina* var. *Sibthorpii* quâcum sâpe confusa distinguitur.

Ar. Geogr. Italia media et australis, Sicilia, Africa borealis.

20. *S. peregrina* (L. Sp. p. 836) caulinibus ascendentibus vel erectis glabriusculis paniculatim ramosis, foliis mediocribus petiolatis a basi truncatâ vel cordatâ ovatis grandidentatis, racemis elongatis laxis præter indumentum breve crispulum pilis brevibus glandulosis sub lente obsitis, foliis floralibus ovatis acutis calyce longioribus, corollâ intense violaceâ pubescente calyce quadruplo longiore, seminibus punctato-depressis pube stellatâ canescensibus ♀. *S. rubicunda* Hornem. — *S. adenotricha* Boiss. et Heldr. MSS.

Hab in regione montana et subalpinâ, regio inferior montis Parnes Atticæ 2500' (Orph!), Parnassi supra Rachova 3500'-4000' (Heldr! Orph. Fl. Gr. exs. 430!), montes Bœotiae (Sprun!).

β. *Parnassica* Boiss. Diagn. Ser. II, 4 p. 28. — Humilior semi-pedalis, caules tenuiores subsimplices, planta breviter puberula. — *S. Parnassica* Heldr. et Sart. MSS.

Hab. in regione superiore montium Parnes et Pateras Atticæ (Auch. 4613! Heldr!), Delphi Eubœa (Heldr!), Parnassi ad Karkaria (Heldr!).

γ. *Sibthorpii* Boiss. et Reut. loc. cit. — Caules et folia plus minusve velutino-hirta sâpe canescens, racemi præter pilos glandulosos eodem indumento criso sed longiore et copiosiore obsiti. — *S. Sibthorpii* Boiss. et Reut. MSS. — *S. Columnæ* var. *Sibthorpii* Bth. in d.c. Prodr. XII, p. 419. — *S. peregrina* Sibth. et Sm. Fl. Gr. tab. 582 (corollæ nimis longæ)

Hab. in umbrosis regionis inferioris et montanae, Peloponnesus in monte Malevo (Orph. Fl. exs. 661!) et Kyllene supra Trikala (Heldr!), insulâ Naxos (Sart!), Byzantii (Sibth., Auch. 1618!), Ponto Lazico ad Khabackar (Bal!), monte Aladagh Cappadocia et Ciliciâ littorali (Bal!), Syriâ littorali in monte Amano (Ky. 18! sub *S. albidâ*), ad Sidonem (Gaill!) et Libano ad Arissa (Ehr!) et Eden (Bl!), Cypro (Sibth., Ky. 48?!), ad Hierosolymam (Roth 337!), Mesopotamiâ ad Orfa (Haussk!).

Formæ enumeratae intermedii evidenter confluunt. Species a *S. altissimâ* ejus corollas abbreviatas (6-8 lineas longas) habet foliis minoribus, bracteis calyces superantibus et seminum charactere distincta. Var. *Sibthorpii* indumento ad *S. Columnæ* accedens ab ea bracteis calyces superantibus et corollis abbreviatis discedit.

Ar. Geogr. Sicilia.

21. *S. altissima* (L. Sp. Ed. II, p. 836) caulinibus elatis simplicibus vel ramosis præter angulos pubescentes glabris, foliis præter

margines et nervos subtus ciliatulos glabris petiolatis cordato ovatis grandidentatis. racemis elongatis pubescenti-glandulosis secundis, foliis floralibus subpetiolatis ovato-rhombeis acutis calyce subbrevioribus, corollā pubescente cæruleā calyce quadruplo longiore, seminibus scabris ♀. *S. commutata* Guss Prodr. II, p. 136 — *S. Columnæ* Host Fl. Austr. non All. — *S. peregrina* W. K. Hung. II, tab. 125 non L.

Hab. in nemorosis, Tauria et Caucasus (M.B., Hohent!), Iberia (Boschniak!).

Planta 2-3-pedalis, folia inferiora tres pollices longa, corolla 6-7 lineas longa. Specimen Aucheri ex Atticā № 1613 hūc citatum est *S. peregrina* var. *Parnassica*.

Ar. Geogr. Italia australis, Dalmatia, regio Danubialis, Bosnia, Rossia australis.

22. *S. albida* (L. Mant. p. 248) herbacea, inferne breviter et crispule pubescenti-tomentella superne et secus racemos patule hispido-viscidula, caulinis elatis ramosis, foliis petiolatis late ovatis obtusis grosse dentalis intimis cordatis, racemis elongatis densiusculis secundis, foliis floralibus magnis corollas subaequantibus petiolatis elliptico-oblongis utrinque acutis integerrimis, calyce hispido parum acreto, corollā hispidā albido-oehrōleucā calyce triplo longiore, seminibus granulato-scabris ♀. Ic. Fl. Græc. tab. 581. — *S. Cretica* Mill. Dict. — *S. nigrescens* Spr. Syst. II. p. 702 — *S. pallida* M.B. Taur. II, p. 65 non Guss.

Hab. in umbrōsis præsertim regionis montane et subalpinis, Macedoniae peninsula Athoa (Griseb., Pichler!), mons Korthiati (Orph!), Thracia propre Carlova (Janka!), prope Byzantium (Sibth!), in Bithynia et Olympo (Thirke! Sibth., Pichl!), Phrygiæ mons Bulghasdaghi (Bal!), Anatolia borealis prope Tokat (Wied!), Rhodi montes supra Maritza (Bourg. exs. sub *S. Sieberi*!), mons Avroman Kurdistaniæ Persice 4-6000' (Haussk!), Tauria (M.B., Stev!).

Corolle breves *S. peregrinæ* à quā differt totā inflorescentiā longius villos-viscidā, racemis densiusculis, bracteis majoribus latioribus, colore florum. Specimina ex Libano citata ad *S. peregrinam* var. *Sibthorpii* referenda sunt.

Ar. Geogr. Banatus.

23. *S. Sieberi* (Bth in D.C. Prodr. XII, p. 420) basi indurata suffrutescens adpresso et brevissime canescens, caulinis annotoninis herbaceis erectis paniculatis, foliis petiolatis a basi truncato-subcordata ovatis obtuse et grosse crenatis, racemis secundis brevibus confertifloris fructiferis laxiusculis, foliis floralibus petiolatis oblongo-rhombeis acuminatis calyces superantibus, corollā pubescente calyce quadruplo longiore, nuculis . . . ♀. *S. hirta* Sieb. exs. non Fl. Græca.

Hab. ad rupes montis Ida Cretæ (Sieb!), prope Critza prov. Mirabello et Anatoli prov. Girapetra (Heldr!), ad prom. Maleka et Aradana (Raul!), ad Cydoniam (Weiss!), in montibus insulae Rhodi supra Maritza (Bourg!).

Pedalis vel sesquipedalis, folia inferiora 2-2½-pollices longa. Affinis *S. albida* differt caulinis basi suffrutescentibus, indumento breviori, spicis condensatis, bracteis acuminatis fere imbricatis Corolla 7-8 lineas longa.

24. **S. hirta** (Sibth. et Sm. Fl. Græc VI, p. 66, tab. 583) indu-
mento ædpresso pilisque longis patulis hirta viscidula cinerea, ramis
e rhizomate indurato ramoso basi decumbentibus et subnudis fragi-
libus brevibus, foliis longe petiolatis ovato-subcordatis crenato-
dentatis, spicis secundis brevibus oblongis densiusculis, foliis flora-
libus subpetiolatis ovatis integris calyce sublongioribus, calyce villosa
fructifero modice aucto, corollâ pubescente calyce triplo longiore
ochroleucâ labio superiore rubello, nuculis granulatis ♀. *S. decum-
bens* Sieb. exs.

Hab. in rupestribus regionis alpineæ ditionis Sphakia Cretæ, montes Stra-
vopodia, Hagion Pneuma, Hellinoseli etc. 6-7000' (Sieb! Heldr! Raul!), mons
Ida (Heldr!), montes Cypri circa Prodromo (Ky. 699!).

β. brachystegia. — Bracteæ minores calycem floriferum vix supe-
rantes. — *S. rupestris* Boiss. et Heldr. Diagn. Ser. I, 7, p. 60.

Hab. in regione alpinâ Taygeti ad Kakochoni (Heldr!), specimina incompl.
e cacumine montis Ænos Cephaloniae (Heldr!) huc spectare videntur.

Caules vix spithamei, folia pollice plerumque breviora, corolla sex lineas
longa. Species a *S. utriculata* Labill. calyce fructifero modice aucto nec valde
inflato distincta.

25. **S. utriculata** (Labill. Ic. Syr. Dec. 4, p. 11, tab. 6) patule
hispido - villosa viscidula canescens, ramis e rhizomate indurato
ramoso decumbentibus fragilibus, foliis parvis petiolatis subcordato-
ovatis crenato-dentatis, racemis densiusculis demum elongatis sub-
laxis, foliis floralibus subpetiolatis ovatis integris obtusis calyce
florifero sublongioribus fructifero brevioribus, calyce hispidissimo
fructifero valde inflato, corollâ hispidâ calyce quadruplo longiore,
nuculis.... ♀.

Hab. in Libano (Labill!) supra Cedros (Boiss!), ad Marserkis et supra
Yamouny (Bl!), in Antiilibani fauibus Ouadi Barrada (Boiss!) et in valle
Manschura 5000' (Ky. 417!), in regione mediâ Hermonis (Boiss!).

Rami 6-10-pollicares parte inferiori nudi, folia 6-9 lineas longa, calyx fructi-
fer 6-7 lineas longus et latus, corolla 10 lineas longa. Species calyce fructi-
fero valde inflato et corollâ elongatâ a *S. hi tā* distincta

26. **S. glutinosa** (Bth. in p.c. Prodr. 12, p. 419) suffruticosa
ramosissima undique patule hispida glutinosa, ramis hornotinis
erectis brevibus ramulosis intricatis, foliis breviter petiolatis cuneatis
oblongo-rhombeis utrinque paucidentatis, racemis laxis, foliis flora-
libus breviter lanceolatis calyci suhæquilongis, corollâ viscoso-
pubescente calyce subtriplo longiore ♂.

Hab. in regno Cabulico (Griff!).

Ex speciminibus incompletis junioribus descripta, rami juniores 4-6-polli-
cares, folia 6-7 lineas longa Ex affinitate *S. Arabicæ* J. et Sp.

3. Galericulatæ.

27. **S. galericulata** (L. Sp. p. 835) parce et breviter puberula,
radice repente, caulis erectis patentim sæpius ramosis, foliis a basi

cordatâ oblongo-lanceolatis remote et obtuse dentatis acutatis sub-petiolatis, floralibus conformibns sensim diminutis, floribus axillari-bus oppositis secundis, calyce pilis simplicibus brevibus hirtulo, corollæ cœruleæ calyce multoties longioris tubo basi fere rectangule curvato, nuculis tuberculatis ♀. Ic. Engl. Bot. tab. 723. — Rehb. Ic. tab. 55.

Hab. in humidis, Macedonia prope Vodena (Orph!), Byzantium (Sibth.), Anatolia borealis ad Kadikiōi (Wied!). Pontus Lazicus ad Rhizé (Balt!), Caucasus et Transcaucasia (Hohen., Rupr!).

Ar. Geogr. Europa fere omnis sed in australioribus rarer, Sibiria omnis, India boreali-occidentalis, Africa borealis, America borealis.

28. **S. hastifolia** (L. Sp. 835) parce et breviter puberula, radice repente, caulis tenuibus erectis simplicibus vel inferne ramosis, foliis breviter petiolatis basi truncatis hastato-sublobatis integris inferioribus ovatis cœteris lanceolatis, floralibus conformibus sensim diminutis, floribus axillaribus oppositis in racemum terminalem secundum approximatis, calyce glanduloso-pubescenti, corollæ cœruleæ calyce multoties longioris tubo basi fere rectangule curvato, nuculis tuberculatis ♀. Ic. Rehb. t. 55, fig. I. — Fl. Dan. tab. 1822.

Hab. in humidis Thraciae (Friv!), Lydia in valle Caystri prope Trianda (Boiss!), prov. Caucasicæ (Hoh. ex Ledeb.).

Ar. Geogr. Scandinavia meridionalis, Gallia media, Germania, Italia borealis, regio Danubialis, Rossia.

BRUNELLA (L. Gen. 783).

Calyx tubuloso-campanulatus reticulato-venosus supra planus bilabiatus, fructifer compresso-clausus, labio superiore plano lato truncale breviter tridentato inferiore bifido. Corolla tubus subexsertus ascendens intus annulatus, limbus bilabiatus labio superiore galeato inferiore dependente obtuse trilobo. Stamina approximata sub labio corollæ parallelæ, superiora præsertim apice breviter bidentata dente inferiore antherifero, antheræ divaricatim biloculares. Stylus apice subulatim bifidus. Nuculæ oblongæ laeves. — Herbæ perennes.

1. **B. vulgaris** (L. Sp. 837) glabriuscula vel pubescens, radice repente, caulis ascendentibus pumilis, foliis petiolatis vel oblongis integris vel superioribus dentatis aut pinnatifidis, caulinis remotis, spicæ ovatâ vel oblongâ, braeteis membranaceis hirtis orbiculatis abrupte acuminatis, dentibus labii superioris calycinæ brevissimis truncatis mucronatis, inferioris lanceolatis, corollæ violacea breviter exsertâ, filamentorum longiorum mucrone subalato recto ♀. Ic. Fl. Dan. t. 910. — Engl. bot. tab. 961. — *B. luciniata* L. var. γ.

Hab. in pratis siccis, sylvis apertis regionis inferioris et subalpinæ totius

ditionis a Græciâ! ad Syriam littoralem et Libanum (Bl! Gaill!), regionem Caucasicam!, Persiam borealem (Auch! Buhse!).

B. Crelica. — Pusilla, nacra, calycis labium superius subintegrum.

Hab. in quercetis meridionalibus montis Ida supra Vorisa (Heldr!).

Ar. Geogr. Europa omnis, Sibiria, Japonia, China, India borealis, Africa borealis, Nova Hollandia, America borealis.

2. *B. alba* (Pall. ap. M.B. Taur. Cauc. II, p. 67) pubescens, radice repente, caulibus ascendentibus pumilis, foliis petiolatis ovatis vel oblongis inferioribus integris superioribus sæpius dentatis vel pinnatifidis, spicâ ovatâ vel oblongâ, bracteis membranaceis orbiculatis, dentibus labii superioris calycini late ovatis truncatis abrupte mucronatis inferioris lanceolatis, corollâ ochroleucâ breviter exsertâ, filamentorum majorum mucrone antrorsum arcuato ♀. *P. laciniata* L. Sp. 837 excl. var γ. — Ic. Jacq. Austr. tab. 378.

Hab. in siccis, ad ripas præsertim in regione montanâ et subalpinâ, Zacythus (Sibth., Marg!), mons Malevo Laconiae (Orph!), Olympus Thessalus (Heldr!), mons Korthiati Macedonia (Orph!), Byzantii (Sibth., Wiedem'), Anatolia borealis (Wied!), Tauria, Caucasus et Transcaucasia (Ledeb.), Persia borealis (Szov! Buhse!).

Ar. Geogr. Europa media et australis a Belgio et Germaniâ ad regionem et Rossiam meridionalem, Africa borealis.

3. *B. grandiflora* (Mænch Méth. p. 414) glabrescens vel parce et minute hirta, radice repente, caulibus ascendentibus, foliis petiolatis ovalo-oblongis integerrimis rarius dentatis vel pinnatifidis, spicâ ovatâ vel oblongâ, bracteis membranaceis orbiculatis ciliatis apiculatis, dentibus labii superioris calycini late et brevissime triangularibus mucronatis inferioris lanceolatis, corollâ violaceâ calycem duplo superante, filamentis muticis longioribus apice gibbo parvo instructis ♀. Ic. Jacq. Austr. tab. 377.

Hab. in pratis montanis siccis, Bithynia supra Broussam (Thirke ex Griseb.), Caucasus Orientalis 4800' (Rupr!).

Ar. Geogr. Europa media a Daniâ et Gothiâ ad Rossiam medium, Lusitania et Hispania borealis et media, Italia, Dalmatia, regio Danubialis.

Obs. *B. hyssopifolia* L., species occidentalis erronee ex Prodromo Fl. Græcæ in agro Byzantino indicatur.

MARRUBIUM (L. Gen. № 721).

Calyx tubulosus 5-10-dentatus æqualis. Corollæ tubus inclusus rarius breviter exsertus, limbus bilabiatus labio superiore erecto (in nostris) retuso vel bifido, inferiore patente trifido. Stamina 4 didynama recta parallela tubo inclusa, antheræ divaricatim biloculares. Stylus bifidus lobis brevibus obtusis. Nuculæ apice obtusæ. — Herbæ perennes rarius annuæ.

Specierum Orientalum dispositio.

SECT. I. EUMARRUBIUM Bth. — Corollæ tubus subannulatus labio superiore emarginato vel bifido. Bracteæ subulatæ vel setaceæ integræ.

* Calycis dentes quini.

× Calycini dentes recti tubo 2-5-plō breviores.

1. *Verticillastra capitato-congesta.*

M. lutescens, cephalanthum.

2. *Verticillastra saltem inferiora plura distantia.*

M. peregrinum, depauperatum, Astracanicum, purpureum, Kotschyi, Libanoticum, globosum, condensatum, micranthum, fauicidens, trachyticum, Bourgei, rotundifolium.

× × Calycini dentes recti tubo vix breviores.

M. heterodon, Thessalum.

× × × Calycini dentes demum stellatim patentes elongati.

M. catariifolium, procerum, Alysson, Persicum, leonuroides, propinquum, velutinum, Frivaldszkyanum.

** Calycis dentes subdeni (Confer *M. parviflorum* var.).

M. præcax, vulgare, crassidens, cuneatum, polyodon, parviflorum.

SECT. II. BALLOTOIDES. — Corollæ tubus exannulatus labio superiore bifido. Bracteæ oblongæ dentalæ.

M. ballotoides.

SECT. I. EUMARRUBIUM.

* Calycis dentes quini.

1. **M. lutescens** (Boiss. Diagn. Ser. I, 5 p. 34) indumento denso molli adpresso sericeo totum flavescens, ramis sterilibus brevibus crebre foliosis, floralibus ascendentibus elongatis superne opposite stricte et breviter ramulosis anguste paniculatis, foliis petiolatis ovatis lineatis suberenatis, floralibus minutis oblongo-cuneatis flores superantibus, verticillastris paucifloris in capitula terminalia saepe laxiuscula approximatis, bracteis panicis subulatis ealyces subsequantibus, calycis erispule hirsuti dentibus a basi lanceolata spinu-

losis rectis tubo quadruplo brevioribus, corollæ flavidæ tubo breviter exerto limbo parvo stellatim tomentello ♀.

Hab. in pratis alpinis montis Tmoli supra Philadelphiam et Cadmi supra Denisleh (Boiss!), in monte Mouraddagh Phrygiae 3300' (Bal!).

Caules cum paniculâ sesquipedales, folia majora 9-12 lineas longa, calyx 2 lineas longus. Species inflorescentiâ stricte paniculatâ et floribus in capitula congestis insignis.

2. *M. cephalanthum* (Boiss. et Noë Diagn. Ser. II, 4 p. 50) dense albo vel flavescenti-pannosum basi induratum cespitosum ramis sterilibus brevissimis crebre foliosis, floralibus elongatis ascendentibus parce et remotissime foliosis, foliis parvis petiolatis lineatis et rugosis oblongis crenulatis, floralibus diminutis flores subsuperantibus, verticillastris 2-3 paucifloris in capitula ovata vel oblonga condensatis, bracteis hirsutis tenuiter subulatis calycis tubum subæquantibus, calycis hirsuti dentibus rectis a basi lanceolatâ tenuissime subulatis inermibus tubo triplo brevioribus, corollæ flavidæ villosissimæ tubo breviter exerto ♀. *Stachys Armenia* Boiss. et Haussk. MSS.

Hab. in rupestribus regionis alpinæ Cappadociae in monte Masmeneudagh (Bal!). Berytlagh Cataouie 6-8000' (Haussk!), ad rupes Bakker Maaden Armeniae australis (Noë 851!).

Caules floriferi 6-10-pollicares subtetraphylli, capitula 4-1½-pollicaria, calyx 3 lineas longus. Facies *Stachydis*, inflorescentiâ et corollis flavidis insigne; ab affini *M. lutescens* indumento breviore non sericeo, caulis floriferis sessilibus, calycis dentibus tenuissime et longe subulatis inermibus discrepans.

3. *M. peregrinum* (L. Sp. p. 815) adpressissime canum, caule humili patentim ramosissimo, foliis supra linealis subtus elevatim rugulosis oblongo-rhombeis subpetiolatis cuneatis parte interiori integris superiori crenato-serratis, floralibus minutis lanceolatis flores superantibus, verticillastris 6-10 floris ad ramorum apicem congestis, bracteis adpressis tomentosis calyce brevioribus, calycis breviter et adpresso stellatim lepidoti tomentosi profunde sulcatis dentibus 5 lanceolato-subulatis tomentosis erectis tubo triplo brevioribus, corollæ albae limbo parvo ♀. Ic. Jacq. Austr. t. 160. — Rehb. Ic. t. 23. — *M. Crelicum* Mill. Diet.

Hab. in ruderatis et aridis regionis inferioris et montanæ. Græcia in Arcadiâ et Doride (Heldr!), Bœtiâ (Sprun!), Macedonia et Thraciâ (Gris! Orph Fl. exs. 521!), Bithyniâ et Lydiâ (Boiss! Heldr!), Anatoliâ boreali (Wied!), Tauriâ et Caucaso boreali (Stev., Led.).

Ar. Geogr. Germania, regio Danubialis, Rossia media et australis.

4. *M. depauperatum* (Boiss. et Bal. Diagn. Ser. II, 4 p. 49) totum adpressissime tomentosum cinereum pluricaule, caulis tenuiter virgatis superne breviter paniculatis, foliis parvis breviter rugulosis minute crenulatis inferioribus petiolatis oblongis, floralibus minimis sessilibus deflexis obtusis calyce longioribus, verticillastris

bifloris in spicas longas graciles approximatis, bracteis nullis, calycis adpresso tomentosi dentibus quinque triangularibus muticis brevissimis exterieribus binis sublongioribus, fauce albo-pilosâ, corollæ albae minute tubo subexerto ♀.

Hab. in collibus saxosis prope pagum Erek Cappadociae inter fluvium Zamante Sou et Caesaream 4500' (Bal!).

Inđumentum *M. peregrini*, sesquipedale, folia inferiora 6-8 lineas longa, spicæ 2-3-pollicares tenues, calyces vix 2 lineas longi. Species distinctissima.

5. M. Astracanicum (Jacq. Ic. Rar. I. p. 11, tab. 109!) caulis humilibus basi decumbentibus sublanatis superne simplicibus vel breviter ramulosis adpresso hirtis, foliis adpresso sericeo-griseis superne profunde lineatis subtus rugosissimis inferioribus obovato-orbiculatis cuneatis petiolatis apice crenatis, floralibus oblongis basi attenuatis apice dentatis verticillastra duplo superantibus, verticillastris 3-5 multifloris superioribus approximatis, bracteis subulatis pungentibus rigidis calyces aequantibus, calycis pilis stellatis et simpli-cibus tomentelli dentibus subulatis rectis tubo dimidiâ vel tertia parte brevioribus, corolla purpureæ calyce duplo longioris tubo inclusa ♀. Rohb. Ic. tab. 270

Hab. in pratis siecis subalpinis et alpinis, Olympus Bithynus (Pichl!), Anatolia borealis in monte Jylidisdagh (Wied!) et supra Tokat (Noë 852!), Bulghasdagh Phrygiae (Bal!), Tauris Ciliciens in valle Giusguta (Bal!), Cappadociae mons Aslandach (Bal!), Berytdagh 600' (Hanss-k!), Pontus (Tchih!), Armenia Turcica montes (Bourg! Huet! Ky. Suppl. 684!), Armenia Rossica (Bge), ditio Talysh (C.A. Mey! Hoh!), Persia borealis in montibus Ssahend, Elbrns (Buhse! Auch. 5108!).

β. glandulosum. — Elatins, calycis pili subglandulosi.

Hab. in Tauro Cilicio (Ky. 454! sub *M. leoniroide*).

Hujus speciei nomen improprium, in Rossiaustrali enim non crescit.

6. M. purpureum (Bge Lab. Pers. p. 67) caulis ascendenter simplicibus molliter et adpressiusculè lanatis, foliis molliter lanato-villosis supra lineatis subtus rugulosis petiolatis orbiculatis vel reniformibus subcordatis crenulatis, floralibus verticillatos superantibus, verticillastris globosis multifloris densis superioribus confluentibus, bracteis subulatis plumosis tubo calycis brevioribus, calycis superne præsertim longe plumoso-villoso dentibus lanceolato-subulatis erecto-patulis tubo quadruplo brevioribus, corolla violacea tubo inclusa, labio parvo, labio superiore profunde bifido ♀.

Hab. in subalpinis Persie boreali-occidentalis inter Siaret et Schahrud (Bge!).

Semipedale et pedale, folia inferiora transverse latiora 9-10 lineas lata. Facies et verticillastra lanata *M. globosi*, ab eo et affinis foliis subcor lat's, verticillastris superioribus confluentibus, etc., diversum.

7. M. Kotschyi (Boiss. et Hohen. Diagn. Ser. I. V. p. 33) ramoso-pluricaule caulis a basi procumbenti erectis elongatis simplicibus vel parce ramosis basi illo-lmatis superne minute tomentellis, foliis

rugosis petiolatis inferioribus albo-lanatis-superioribus virescentibus griseo-tomentellis ovatis crenatis, floralibus oblongo-cuneatis breviter petiolatis verticillatos superantibus, verticillastris remotis densis, bracteis subulatis incurvis calycis tubum æquantibus, calycis hirsuti et stellatim pilosi dentibus lanceolatis spinulosis rectis tubo 4-5 plo brevioribus, corollæ roseæ vel albæ tubo exerto labio superiore profunde bifido ♂. *M. Noëcanum* Boiss. MSS.

Hab. in rupestribus regio iis mediæ montis Gara Kurdistaniæ supra Mossul (Ky. 379!), in montibus circa Diarbekir (Noë 698!), montes Pir Omar Gudrun et Juanro Kurdistaniæ (Haussk!).

β. brachyodon. — Totius plantæ et calycis indumentum brevius fere tantum pilis stellatis constans. — *M. brachyodon* Boiss. Diagn. Ser. I, 7, p. 58.

Hab. in regione alpinâ montium Avroman et Schahu Kurdistaniæ Persicæ (Haussk!), Persia austro-occidentalis, Kuh Daëna (Ky. 596!) et Kuh Delu (465!), montes Nur, Sawers et Eschker (9-12000' (Haussk!).

Species a *M. Astracanico* indumento breviore, parciore, dentibus calycinis brevioribus, corollæ tubo exerto et labio superiore longe nec breviter bifido distincta.

8. ***M. Libanoticum*** (Boiss. Diagn. Ser. I, 12, p. 73) dense et breviter tomentosum flavescens ramosissimum ramis a basi procumbenti albo-lanata erectis simplicibus, foliis rugosissimis parvis ovatis vel orbiculatis subcrenulatis inferioribus longe petiolatis, floralibus breviter petiolatis ovato-oblongis verticillastro longioribus, verticillastris denses, bracteis subulatis calyce subbrevioribus, calycis brevissime et stellatim tomentelli sulcati dentibus lanceolatis spinulosis rectis tubo triplo brevioribus, corollæ carneæ limbo parvo calycem parum superante ♀.

Hab. in summis Libani supra Cedros (Ehrenb! Boiss! Ky. 302!).

γ. Hermonis. — Indumentum canum, corollæ limbus paulo major subexsertus. — *M. Hermonis* Boiss. Diagn. loc. cit. p. 74.

Hab. in regione mediâ et superiore montis Hermon supra Rascheya (Boiss! Dr. Post!).

Semipedale et pedale, folia majora 6-7 lineas longa, calyx 3 lineas tantum longus. Facies et folia parva *M. globosi* sed calycis indumentum pilis stellatis constans; hæc nota *M. Kotschy* var. *brachyodonti* affine sed folia parva suborbiculata nec ovato-oblonga dense subpannosa nec tomentello-virescentia, calyces sublongiores.

9. ***M. globosum*** (Montb. et Auch. Ann. Sc. Nat. Ser. II, 6 p. 53) ramosissimum ramis simplicibus a basi procumbenti indurata erectis albo-lanatis, foliis dense et adpresso sericeis superne lineatis parvis cuneato-ovatis vel orbiculatis apice obtuse crenatis inferioribus longe floralibus breviter petiolatis, hisce diminutis verticillastris longe superantibus, verticillastris parvis dissitis globosis, bracteis subulatis villosissimis adpressis calyce subbrevioribus, calycis villo-

sissimi dentibus lanceolato-subulatis spinulosis erectis brevissimis, corollæ albæ tubo subexerto limbo parvo ♀.

Hab. in pratis alpinis Tmoli supra Geyra in Cariá (Boiss.!) montis Akdagħ Lycæ (Bourg. exs. sub *M. micrantho*) montis Akdagħ Tauri orientalis in Cataoniā (Auch. 1787!), montis Argæi Cappadociæ radicibus (Ky. Suppl. 290! sub *M. micrantho*).

Rami $\frac{1}{2}$ -1-pedales, folia majora diam. vix pollicaria, verticillastra magn. avellanæ. Calyx 3 lineas longus dentibus tubo 5-6-plo brevioribus.

10. ***M. condensatum*** (Boiss. in Noë pl. exs.) ramosissimum breviter et parce albo-lanatum, ramis a basi procumbenti erectis, foliis parvis obovatis subcrenatis albo-lineatis in petiolam attenuatis, floralibus diminutis oblongis breviter petiolatis verticillastro longioribus, verticillastris parvis globosis compactis bombycinis distantibus, bracteis setaceis hispidissimis calyces æquantibus, calycis minimi turbinati inferne glabrescentis parte superiori valde lanati dentibus spinulosis erecto-patulis tubo triplo brevioribus. corollæ albæ limbo minimo ex lanâ calycis vix exerto ♀.

Hab. in montibus Mesopotamiae prope Diarbekir (Noë 850!).

Valde affine *M. globoso* pro cuius formâ in Diagn. Ser. II, 4, p. 51 habui sed distinctum tomento parciore breviore non sericeo, foliis etiam inferioribus vix semipollicaribus, calyce $2\frac{1}{2}$ lineas vix longo subturbinato basi glabrescente et apice valde lanato, dentibus longioribus.

11. ***M. micranthum*** (Boiss. et Heldr. Diagn. Ser. I, 12 p. 73) ramosissimum ramis simplicibus a basi procumbenti erectis adpresso-sime albo-tomentosis, foliis parvis adpresso tomentosis non sericeis supra flavidis subtus canis ovatis obtuse crenatis inferioribus longe petiolatis floralibus breviter petiolatis flores vix superantibus, verticillastris glomeratis binis oppositis densis breviter pedunculatis, verticillastris panicis inferioribus distantibus, bracteis subulatis calyce brevioribus, calycis flavidо-villosi dentibus subulatis rectis vix spinulosis tubo 4-6 plo brevioribus, corollæ carneæ tubo subincluso, limbo parvo ♀. *M. pseudoalyssum* F. et M. Ann. Sc. Nat. 1854, p. 34 non de Noë.

Hab. in pascuis montis Ghelipel Tauri Isaurici et cæteris montibus supra Ermenek 4000'—6000' (Heldr.!), in Cataoniā (Tchih!).

Cespitosum, rami $\frac{1}{2}$ -1-pedales, folia 5-6 lineas longa et lata. Valde affine *M. globoso* a quo differre videtur indumento breviore adpressiore non sericeo, foliis minoribus, verticillorum saltem inferiorum glomerulis distincte pedunculatis.

12. ***M. faucidens*** (Boiss. et Bal. Diagn. Ser. II, 4 pag. 50) basi ramosissimum totum densissime albo-vel lutescenti-pannosum, ramis simplicibus a basi procumbenti erectis, foliis petiolatis crenatis obovatis vel oblongis basi cuneatis, floralibus subsessilibus verticillois superantibus, verticillastris valde remotis, bracteis subulatis plumosiss calycino tubo subbrevioribus, calycis valde hirsuti dentibus su-

bulatis rectis apice spinulosis glabris tubo triplo brevioribus, fauce dense pilosâ gibbis (in sicco obsoletis) angustatâ, corollæ albidæ tubo subincluso limbo minuto ♀.

Hab. in regione alpinâ inferiore Tauri Cilicici ad septentrionem angustiarum Gülek Boghas (Bal. 531!), in monte Amano in rupestribus Apisch Gaya (Ky. 238 sub *M. globoso*), in Cataoniâ supra Seytun et in monte Akkerdagh 4000' (Haussk!).

Caules floriferi pedales, folia rainorum steriliū párva, indumentum eo *M. Kotschyi* et *M. Libanotici* densius, simplex nec stellatum. Magis affine *M. micrantho* a quo sat differre videtur florum glomerulis non pedunculatis, calycis fauce gibbosi dentibus longioribus.

13. *M. trachyticum*, multicaule adpresso et parce lanatum cinereum sericum, caulis elatæ ascendentibus parce ramosis, foliis supra lineatis subtus rugosissimis dentatis oblongis cuneatis inferioribus petiolatis floralibus oblongo-linearibus subsessilibus verticilos multo superantibus, verticillastris remotis, bracteis subulatis spinulosis adpresso calyce triplo brevioribus, calycis tubulosi adpresso stellatim albo-tomentosi dentibus subulatis rectis subspinulosis subinæqualibus tubo 4-5-plo brevioribus, corollæ parvæ albæ tubo inclusa ♀.

Hab. in monte trachytico Hussein Kazi prope Ancyram Anatoliæ (Wied!).

Sesquipedale, folia turionum minuta, caulinæ inferiora pollicem longa, calyx lineis 3 sublongior. Species indumento, foliis præsertim floralibus angustis elongatis, calyx breviter cano *M. parviflorum* *heterodon* referens, sed ab eo calycis dentibus brevibus rectis nec longis squarroso-patentibus alienum, juxta *M. micranthum* et *faucidens* collocandum.

14. *M. Bourgæi*, adpresso flavo-tomentosum e rhizomate duro multicaule caulis ascendentibus simplicissimis, foliis rugulosis crenulatis turionum parvis oblongis, caulinis inferioribus ovato-spatulatis basi cuneatis petiolatis, floralibus breviter petiolatis diminutis ovato-oblongis flores superantibus, verticillastris multilloris distantibus, bracteis subulatis hirtis patentibus incurvis calycino tubo æquialongis, calycis flavidо-plumosi striati dentibus lanceolato-subulatis hirtis subinæqualibus rectis tubo triplo brevioribus, corollæ carneæ vel flavidæ? tubo inclusa ♀.

Hab. in regione alpinâ montium supra Elmalu Lyciæ (Bourg! exs. sub *M. heterodon*).

Folia et facies *M. heterodon* a quo eximie differt verticillis ob flores minutos dimidio minoribus, calycis 3½ lineas tantum longi dentibus brevibus. Indumento flavo, foliorum multo minorum formæ, bracteis patentibus elongatis, etc., a *M. rotundifolio*, calycis flavidо-plumosi dentibus longioribus subinæqualibus a *M. Kotschyi* et *M. micrantho* diversum.

15. *M. rotundifolium* (Boiss. Diagn. Ser. I, 5 p. 33) multicaule totum albo-griseum dense pannosum vix sericeum ramis sterilibus subelongatis foliosis, caulis floriferis ascendentibus simplicissimis, foliis valde rugosis et superne lineatis crenatis orbiculatis

inferioribus petiolatis basi rotundatis vel subcordatis, floralibus subsessilibus flores longe superantibus, verticillastris multifloris remotis, bracteis subulatis tomentosis adpressis calyce multo brevioribus, calycis striati breviter et stellatim tomentelli dentibus rectis lanceolato-subulatis tubo quadruplo brevioribus, corollæ pallide violaceæ tubo inclusa ♀.

Hab. in pratis alpinis Tmoli orientalis supra Philadelphiam (Boiss!), in parte superiore Sipyli supra Magnesiam (Bal. 329!).

Pedale et sesquipedale, folia magna diam. 1-1 $\frac{1}{4}$ -pollicaria *M. velutini* cuius faciem habet et a quo ut et a *M. Frivaldszkyano* et *heterodon* egregie differt calycis 3 $\frac{1}{2}$ lineas tantum longi dentibus brevibus, bracteis brevissimis.

16. *M. heterodon* (Boiss. et Bal. Diagn. Ser. II, 4 p. 52) adpresso cinereo vel flavido-tomentosum ex rhizomate duro multicaule, caulis ascendenter simplicissimis, foliis rugosissimis crenulatis turionum parvis oblongis, caulinis inferioribus ovato-spathulatis in petiolam attenuatis, floralibus conformibus sessilibus inflexis verticilos superantibus, verticillastris multifloris distantibus, bracteis hirtis subulatis erectis calyces æquantibus, calycis stel. diu hirti striati dentibus breviter plumosis tenuiter subulatis rectis valde inæqualibus, binis longioribus tubum æquantibus, corollæ amplexibus dentes calycis subæquante ♀. *M. velutinum* var. *heterodon* B'g. I. d.c. Prodr. XII, p. 451.

Hab. in regione alpinâ Tauri Cilicii ad Gusguta et Gulek Mag. (Boiss! Ky. 455!), in valle alpinâ Kar Boghas (Bal!).

Semipedale et pedale, folia eis *M. velutini* minora angustiora secundum partem basi cuneato-attenuata, calyces quoque longissimi (6-7 lineas 1-1 $\frac{1}{2}$ pollicis) et rectibus valde inæqualibus et rectis nec patulis rigidis.

17. *M. Thessalum* (Boiss. et Heldr. Diagn. Ser. I, 3 p. 5 multicaule, caulis ascendenter simplicibus albo-lanatis non rugulosis supra adpresso cinereis sericeis lineatis subtus int. dist. canescens crenulatis inferioribus ovatis cuneatis petiolatis floralibus oblongis subsessilibus verticillo longioribus, verticillastris multifloris globosis distantibus lanatis, bracteis subulatis adpresso-cerato tubo calycino subbrevioribus, calycis inferne glabrescentis superioris plumosi dentibus rectis tenuiter subulatis plumosis apic. glauco-inermibus tubo subæquilongis, corollæ albæ tubo inclusa ♀.

Hab. in pasculis alpinis montis Olympi Thessali (Heldr. Orph. Fl. Gr. p. 520!).

Sesquipedale, folia fere *N. candidissimi* a quo verticillastris hirtis calycisque dentibus non squarroso-patentibus differt, eidem notâ calycis superne plumoso-lanati nec adpresso lepidotî, foliisque ovatis nec orbiculariis a *M. velutinum* et *rotundifolium* dignoscendum.

18. *M. catariæfolium* (Desr. in Lam. Diet. II, p. 717 annuum, caule erecto rubello sub lente parce et brevissime pubescente a basi opposite et breviter ramoso, foliis in-lumento tenuissimis

stellato canescens supra lineatis subtus elevatim nervosis petiolatis oblongis profunde et obtuse paucicrenatis, floralibus brevius petiolatis angustatis deflexis verticillo longioribus, verticillastris 3-4 distantibus multifloris, bracteis subulatis spinulosis rigidis patentibus calycis tubo æquilongis, calycis pilis stellatis adpressissime tomentelli valde sulcati dentibus lanceolato-subulatis patentissimis rigidis spinulosis dimidio tubo longioribus, corollæ roseæ vel carneæ tubo breviter exerto ⊖. *M. candidissimum* C. A. Mey. non L. — *M. Ibericum catariefolio* Tourn. herb !

Hab. in Caucaso ad ripas fluvii Terek prope Wladikaukas (Rehm!), Iberia Caucasica (Wilm! Hoh! Blum in Lœdeb. herbt!) Somchetia (Eichw.), Cappadocia (Tourn.).

Fere pedale, folia 6-10 lineas longa subtus magis canescens, calyx 3¹/₂ lineas longus. Species indumento brevissimo tenui, calycis valde nervosi et sulcati dentibus valde subulatis patentis subrecurvis insignis.

19. ***M. procerum*** (Bunge Lab. Pers. p. 66) totum indumento brevissimo stellato canescens, caule procero superne opposite ramoso, foliis grosse et acutiuscule dentatis inferioribus petiolatis obovato-suborbiculatis, floralibus subsessilibus oblongis acutis acutius dentatis deflexis verticillo longioribus, verticillastris multifloris magnis inferioribus distantibus, bracteis tenuiter subulatis patentibus hispidulis calyce subbrevioribus, calycis profunde sulcati tenuiter tomentelli et glandulosi dentibus lanceolato-subulatis pungentibus tandem stellatum patentibus tubo subbrevioribus, corollæ carneæ tubo breviter exerto ⊖ ?

Hab. in montosis prov. Khorassan Persiae inter pagos Derrud et Dschegar in viâ inter Nischapur et Mesched (Bge!).

Bipedale et altius, foliorum inferiorum lamina 2-2¹/₂-pollicaris ut in *M. leonuroide* grosse sed acutius dentata, verticillastra diametro pollicaria, calyx ut in *M. catariefolio* sulcatus dentibus stellatum patentibus sed major. Ex affinitate cum *M. catariefolio* annum esse suspicor.

20. ***M. Alysson*** (L. Sp. 815) annum, caulibus lanatis crassiunculis ascendentibus pumilis simplicibus vel parce ramulosis, foliis cinereis tomentosis rugosissimis petiolatis cuneato-flabelliformibus antice late crenato-lobatis, floralibus diminutis deflexis flores superantibus, verticillastris 10-15 floris distinctis laxiuscule et longe spicatis, bracteis obsoletis, calycis adpresso lanati dentibus trianguli-lanceolatis spinosis demum rigidis stellatum patentibus tubo duplo brevioribus, fauce demum lanâ clausâ, corollæ purpurascens limbo minuto labio superiore brevi ⊖. Ic Clus. Hist. p. 35. — *M. plicatum* Forsk. Eg. p. 213.

Hab. in ineulis et pascuis siccis, Egyptus ad Alexandriam (Forsk., Ehr! Samar!), Arabia petrea ad Palestine limites (Boiss!), Syria (Auch. 1789).

Ar. Geogr. Hispania australis, Africa borealis, Sardinia, Italia australis.

21. ***M. Persicum*** (C. A. Mey. Enum. p. 93) adpresso lanatum niveum caulibus elongatis inferne ramulosis, foliis rugosis ovato-

orbiculatis breviter petiolatis crenatis subtus intensius niveis, floralibus oblongis basi longe cuneatis deflexis verticillo longioribus, verticillastris remotis 6-8-floris, bracteis pancis brevissimis spinosis recurvis, calycis adpresso lanati cani dentibus patentissimis subulatis rigidis spinosis dimidium tubum æquantibus, corollæ violaceæ tubo inclusa ♀.

Hab. in aridis lapidosis et arenosis ad Swant et Tatuni ditionis Transcaucasicæ Talyesch (C.A. Mey! Hoh!).

Pedale, folia majora 9 lineas lata. Species bracteis brevissimis dentibusque calycis patent-recurvis rigidis spinosis insignis.

22. **M. leonuroides** (Desr. in Lam. Diet. III, p. 715) breviter tomentello-hirtum pallide virens, caulis crassiusculis, foliis subtus canescentibus petiolatis ovatis majusculis grosse et obtuse inciso-dentatis, floralibus oblongis breviter petiolatis dentatis verticillastra remota parva multiflora multo superantibus, bracteis subulatis villosis apice spinulosis glabratis calyce subbrevioribus patentibus, calycis albo-lanati dentibus subulatis apice spinulosis glabris recurvo-patentibus tubo duplo brevioribus, corollæ roseæ tubo subexerto ♀.
M. Astracanicum M. B. 11 p. 52 non Jacq. — Rehb. Ic. tab. 299.

Hab. in lapidosis Caucasi ad Narzana (M B.), Pätingorsk (Hohen!), ad Girgieti (Rehm!), in ditionibus Ossetiæ et Alagir 2500' et Daghestaniæ boreali 4500'-6000' (Rupr!).

β. plumosum. — Indumentum breviter et molliter cinereum, folia inferiora reniformia transverse sublatiora, floralia orbiculata, calycis dentes setacei plomosi tubo vix breviores. — *M. plumosum* C. A. Mey. Enum. p. 96.

Hab. in regione alpinâ Caucasi Orientalis 8-9000' (C.A. Mey!).

Pedale et bipedale sed interdum et præsertim in varietate humilius; species foliis profunde incisis et verticillis compactis remotis insignis. Erroneæ in Tauriæ et Rossia austro-orientali indicatum.

23. **M. propinquum** (F. et M. Index I Petrop. p. 33) villosocanescens, caulis erectis ramosis, foliis petiolatis subrugosis subtus intensius cinereis grosse crenatis inferioribus a basi cuneatâ orbiculatis, floralibus breviter petiolatis obovato-oblongis basi attenuatis verticillastra multo superantibus, verticillastris globosis distantibus, bracteis subulatis deflexis calyce subbrevioribus, calycis striati villosi dentibus subulatis spinulosis demum patentibus tubo dimidio æquilongis vel sublongioribus, corollæ roseæ tubo vix exerto ♀. *M. cærulescens* Bth. in D.C. Prodr. ex spec. Barclayano! an Desf. Cat. Ed. 3^a p. 395 et 433?

Hab. in agris incultis ditionis Talyesch prope Tatuni (C.A. Mey! Hoh!), prope Helenendorf (Hohen!), in Persiæ boreali prope Tabris (Bge!).

Valde et forsitan nimis affine *M. leonuroidi*, lana laxior, folia crenata nec incisa, etiam inferiora basi cuneata, flores subminores. *M. cærulescens* Desf. Hort Par. ex Catal. p. 433 est potius *M. leonuroides*.

24. *M. velutinum* (Sibth. et Sm. Fl. Gr. VI. p. 49, tab. 561) multicaule caulis a basi procumbenti erectis elongatis simplicibus vel parce ramosis adpresso tomentosis, foliis dense et adpresso tomentosis velutinis flavidio-nitentibus eximie lineatis orbiculatis subcrenulatis inferioribus petiolatis. floralibus parum diminutis subsessilibus flores superantibus, verticillastris multifloris magnis dissitis, bracteis numerosis tenuiter subulatis tomentellis a basi curvatae erectis calycis tubum æquantibus, calycis stellatim tomentosi striati dentibus a basi tenuiter subulatis tubo subbrevioribus patentibus. corollæ flavæ tubo inclusa ♀. *M. circinnatum* Desr. in Lam. Dict. 3 p. 717!

Hab. in regione alpinâ Parnassi 4-6000' frequentissimum et totum jugum sociâ *Daphne buxifolia* obtgens (Heldr!), (Sibth! Orph. Fl. exs. 431! Sart!), in monte Velugo (Etoliae) (Sprun!).

♂. *Cylleneum*. — Indumentum albo-cinereum, bracteæ et dentes calycini abbreviati dimidium tubum vix æquantes. — *M. Cylleneum* Boiss. et Heldr. Ser. II, 4 p. 51.

Hab. in regione alpinâ montium Chelmos et Kyllenes Peloponnesi 3-5000'-7000' (Heldr! Orph. Fl. exs. 299!).

Pedale et elatius, folia inferiora sesquipollucem longa, verticillastra magnitudine nucis. Species nitore flavidio in locis natalibus elegantissima.

25. *M. Friwaldskyianum* (Boiss. Diagn. Ser. I, 12 p. 74) adpresso tomentellum griseum. caulis a basi procumbenti erectis simplicibus, foliis supra lineatis subius reticulato-rugosis inferioribus petiolatis orbiculatis basi subcordatis grosse et obtuse dentatis. floralibus subsessilibus ovato-oblongis acute dentatis flores longe superantibus, verticillastris multifloris remotis, bracteis hirtis setaceis patenti-subreflexis incurvis calyci aquilongis, calycis breviter et stellatim hirtuli valde striati dentibus setaceis divergentibus tubo vix brevioribus, corollæ albidae tubo inclusa ♀. *M. candidissimum* Friv. exs. non L.

Hab. in Rumeliâ (Friv!), in saxosis borealibus montis Rhodopes Thraciae supra Stanimak (Janka!).

Bipedale, affine *M. velutino* a quo differre videtur foliis profundius et superioribus acute dentatis, calycis (6-7 lineas longi) dentibus longioribus, corolla ut e seco videtur albâ. *M. candidissimum* L. in Italiâ australi. Dalmatiâ, Montenegro (nec in Hispania et Persia) crescens longius differt indumento cano, foliis minoribus oblongis, etc.

• Calycis dentes subdeni.

26. *M. præcox* (Janka pl. Transylv. exs. in Sched.) adpresso canescens, caulis erectis superne parce paniculato-ramosis ramis brevibus erectis, foliis superne lineatis subtus magis canis elevatis reticulatis inferioribus petiolatis ovatis et ovato-oblongis crenatis, floralibus valde diminutis oblongo-lanceolatis flores superantibus, verticillastris 6-8-floris distinctis, bracteis subulatis tomentellis adpresso calycis tubum æquantibus, calycis stellatim tomentosi

dentibus lanceolato-subulatis erectis ♂ vel 8-10 inæqualibus tubo profunde sulcato triplo brevioribus, corollâ albâ ♀. *M. paniculatum* Desr. — *M. Pestalozzae* Boiss. Diagn. Ser. II, 4 p. 54!

Hab. in sterilibus Anatoliæ occidentalis, Bithynia prope Moudania (Pichler!), Ouchak Phrygiæ (Bal!). Lydia supra Magnesiam ad Sipyli radie (Bal. exs. sub *M. catariaefolio*!), Lycia (Pest!), Tauria ad Sebastopol (St-Sapery!), Caucasus borealis (ex C.A. Mey.).

A *M. peregrino* foliorum formâ, verticillastris remotis, calycis dentibus accessoriis sèpius obviis distinctum. An diversum a *M. Pannonicum* Clus. — *M. remoto* Kit. quod tamen forma inter *M. vulgare* et *peregrinum* hybrida videtur?

Ar. Geogr. Transylvaniam.

27. *M. vulgare* (L. Sp. p. 816) pluricaule caulis crassis albo-lanatis ascendentibus simplicibus vel parce ramosis, foliis rugulosis crenatis molliter villosis cinerascentibus vel subtus atbo-lanatis ovato-orbiculatis inferioribus petiolatis sèpè cordatis, floralibus diminutis subsessilibus deflexis cuneato-oblongis verticillo sublongioribus, verticillastris multifloris globosis distantibus, bracteis subulatis apice nudis uncinalis. calycis stellato-tomentelli dentibus 10 patentibus subulatis apice valde uncinalis et nudis alternatim brevioribus, quinis longioribus tubo triplo brevioribus, corollæ albæ limbo minute ♀. Ic. Fl. Dan. t. 1036. — Rebb. Ic. tab. 23. — *M. Anulum* Ten. — *M. anisodon* C. Koch Linn. XXI, p. 696.

Hab. in ruderatis regionis inferioris et montanea a Græciâ! et Macedonia (Griseb.), ad regiones Caucasicas. Persian borealem (Bge!) et australem (Ky. 312!), Syriam (Gail!) et Palestinam (Roth!), Belutschiam (Stocks!).

Ar. Geogr. Europa omnis a Sueciâ meridionali ad Rossiam medium, Africa borealis, Africa tropica. In Americam introductum.

28. *M. crassidens* (Boiss. Diagn. Ser. I, 5 p. 35) pluricaule, caulis elongatis simplicibus vel parce ramosis inferne dense albo-lanatis superne minute tomentellis, foliis adipesse ramosis sericeis flavidis supra lineatis subtus rugosis crenatis inferioribus ovatis breviter cuneatis petiolatis, floralibus oblongis basi valde attenuatis deflexis inferioribus flores valde superantibus, verticillastris dissitis, bracteis subulatis tubum calycis subæquantibus, calycis tomentosi sulcati dentibus 10-15 radiatim patentibus lanceolatis apice spinulosis et subuncinatis alternatim subbrevioribus dimidium tubum æquantibus vel superantibus, fance pilis clausâ, corollæ albæ non exsertæ limbo minuto ♀.

Hab. in monte Gara Kurdistaniæ (Ky. 324!), Persiâ australi in montibus Kuh Sawers (Hausskl.), et inter segetes prope Schiras (Ky. 312!). inter Yezd et Ispahan (Bge!).

Diffrerit a *M. vulgari* foliis minoribus ovato-cuneatis nec orbiculari-subcordatis, dentibus calycinis longioribus vix uncinalis. Huic olim erroneè retuli plantam Syriacam ad *M. cuneati* formas referendam.

29. *M. cuneatum* (Russ. Alepp. II, p. 255) pluricaule, cauli-

bus elatis inferne adpresso albo-lanatis superne stellatim tomentellis virentibus paniculatis vel rarius simplicibus, foliis parvis rugosissimis crenulatis stellatim supra virentibus subtus cano-tomentosis ovato-cuneatis, floralibus deflexis angustatis, verticillastris parvis spicatis inferioribus distantibus, bracteis paucis brevissimis linearibus, calycis stellatim tomentosi dentibus subdenis patentibus breviter lanceolatis muticis inæqualibus saepius alternatim trifidis tubo subtriplo brevioribus, fauce gibbis 10 clausa, corollæ albæ limbo minimo ♀.

Hab. in ruderatis circa Aleppo Syriæ (Russ., Auch. 1785! Ky!).

β spinulosum. — Calycis dentes sublongiores in spinulam brevem interdum subuncinatam abeuntes. — *M. radiatum* Delile in Bth. Lab. p. 571.

Hab. in siccis, Libanus (Labill! Bl!), mons Hermon 5500'-6000' (Ky. 167! sub *M. crassidente*) inter Antilibanum et Damascum (Boiss! Gaill!), inter Hama et Homs (Bl!) in Mesopotamia circa Mossul (Noë 23! Haussk!).

Sesqui- et bipedale, caules eis *M. vulgaris* tenuiores et calyces minores. A speciebus affinibus gibbis in fauce calycis distinguitur. Varietas typo vulgarior ad eum sensim transit. Saepè cum *M. crassidente* commutatum cujus dentes longiores sunt et nunquam trifidi.

30. ***M. polyodon*** (Boiss. Diagn. Ser. 1, 12 p. 75) pluricaule, caulibus elongatis simplicibus vel parce ramosis inferne densius albo-lanatis, foliis albo-panniosis crenulatis oblongo-cuneatis inferioribus breviter petiolatis floralibus flores superantibus, verticillastris dissitis, bracteis paucis subulatis brevissimis, calycis tomentosi sulcati dentibus subdenis late et breviter lanceolatis irregulariter 2-3 fidis spinulosis basi subcoalitis tubo vix brevioribus patentibus, fauce villosa, corollæ albæ limbo parvo ♀.

Hab. in aridis montis Elbrus Persiae prope Derbend (Ky. 244!), prope Teheran (Bode), prope Schahrud (Bge!).

Valde affine *M. cuneato*, ab eo calycis fauce pilosa nec gibba, dentibus latioribus in limbum basi coalitis an satis diversum.

31. ***M. parviflorum*** (Fisch. et Mey. Ind. Petrop. I, p. 83) totum breviter et dense canum, ex rhizomate indurato pluricaule, caulibus erectis tenuibus simplicibus vel superne parce ramosis, foliis parvis plus minus sericeis et lincatis subtus rugulosis oblongis longe cuneatis petiolatis crenatis, floralibus oblongo-linearibus subsessilibus deflexis apice saepè dentatis verticillo longioribus, verticillastris globosis distantibus spicatis, bracteis subulatis muticis tubo calycino triplo brevioribus, calycis breviter albo-tomentosi dentibus denis subulatis subspinulosis demum squarroso-patentibus tubo subbrevioribus, corollæ albæ tubo inclusa ♀. *M. peregrinum* C. A. Mey. Enum. non L.

Hab. in monte Mesogis Lydiæ (Boiss!) monte Olklatschi Pisidiæ (Heldr!), Cappadociæ (Bal!), Anatoliæ boreali ad Samsun, Safranbol, Mersivan (Wied!) Armenia ad Erzerum (Huet!), ditione Karadagh (Buhsel), ditione Talyssch (C.A. Mey!), Persia bor. (Szov!).

β. oligodon — Dentes calycini tantum quini. — *M. album* Boiss. et Bal. Diagn. Ser. II, 5 p. 52. — *M. candidissimum* C. Koch Linn. XXI, p. 696 non L.

Hab. in Cappadociâ ad Kamechli Khan et in monte Alidagh (Bal!), circa Baibut Armeniæ (Bourg!).

Planta 1-2-pedalis a *M. cuncato* var. *spinuloso* = *M. radiato* Del. quo cum confusum fuit foliis floralibus angustis, calycis dentibus angustioribus et longioribus nunquam trifidis nec uncinatis omnino distincta; typus ad varietatem eo rariorem intermediis transit.

SECT. II. BALLOTOIDES.

32. ***M. ballotoides*** (Boiss. et Bal. Diagn. Ser. II, 4 p. 53) pluricaule dense et breviter tomentosum cinereum, caulis erectis basi albo-lanatis superne parce et breviter paniculato-ramosis, foliis rugosissimis griseis subtus magis canis inferioribus ovato-orbiculatis petiolatis, floralibus parvis deflexis anguste oblongis flores vix aequantibus, verticillastris approximatis, bracteis oblongo-cuneatis obtuse 3-5-dentatis dimidium calycem vix aequantibus, calycis breviter et stellatim tomentosi limbo erecto-patulo tubo striato quintuplo breviore dentibus subquinis aliis trilobis lobis ovatis obscure mucronulatis aliis minoribus indivisis ovatis, corollæ albæ limbo minuto vix exerto ♀.

Hab. ad Balamout Kaivé inter Smyrnam et Magnesiam in consortio *M. vulgaris* (Bal. exs. 320!).

Facies et indumentum *M. vulgaris* sed ab omnibus congeneribus bracteis non setaceis et calyce fere *Ballotæ* discrepans. Stamina in fauce inclusa *Marrubii*. Hanc plantam pro prole hybrida *Marrubii* et *Ballote* habuisse sed nuculæ normales sunt, obtusæ trigonæ faciebus internis subconcavæ.

SIBERITIS (L. Gen. 742).

Calyx tubuloso-campanulatus interdum subbilabiatus dentibus 5 plerumque spinosis. Corolla tubo inclusa limbo bilabiato labio superiore erecto subplano integro vel retuso-bifido, inferiore patente trifido. Stamina 4 didynama tubo inclusa superiora brevissima antheris divaricatim bilocularibus, inferiora longiora antheris sæpius difformibus. Stylus tubo inclusus bifidus lobo superiore brevi truncato inferiore dilatato basi superiore amplectente. Nuculæ apice rotundato-obtusæ. — Herbae annuae et perennes vel suffrutices.

SECT. I. ***HESIODIA*** Bth. — Herbae annuae. Folia floralia inferioribus conformia sed minora integra vel crenata.

1. ***S. lanata*** (L. Sp. 804) annua molliter et patule villosa, collo sæpius pluricaulis caulis sæpius ramosis erectis, foliis crenatis

ovatis obtusis inferioribus petiolatis, floralibus verticillastro longioribus, verticillastris 6-4-floris distinctis sed in racemos basi interruptos approximatis, calycis molliter lanati subircurvi subbilabiati dentibus lanceolatis subulato-attenuatis spinulosis tubo subbrevioribus supremo sublatiore, corollæ calyce subbrevioris limbo parvo fuscescente ♂. *S. elegans* Murr. Comm. Gött. tab. 4. — *S. nigricans* Pers. Syn. 11, p. 117. — *Stachys paniculata* Bth. in D.C. Prodr. XII, pag. 485!

Hab. in cultis et arenosis Peloponnesi (Sart!), insulæ Leros Cycladum (Urv.), Cretæ (Sieb!), Hellesponti (Auch!), Lydiæ et Cariæ (Boiss! Bal. 307!), Galatiaæ (Wied!), Ciliciae (Ky!), Cappadociae ad Cæsaream (Bal!), Anatoliae loco non notato (Mitchell in herb. Bth!).

2. *S. Romana* (L. Sp. p. 802) annua molliter pilosa viridis pumila simplex vel saepius collo ramosa ramis ascendentibus vel diffusis, foliis ovatis et oblongis crenato-dentatis radicalibus in petio lum attenuatis, verticillastris 6-4-floris distantibus, foliis floralibus oblongis dentatis verticillastro sublongioribus, calycis viridis patule hirsuti basi valde gibbi subbilabiati dentibus spinuloso-aristatis superiore multo majore ovato inferioribus lanceolatis, corollæ albæ limbo, breviter exserto labio superiore latiusculo interdum roseo ♂. Ic. Fl. Græc. tab. 552. — Rchb. Ic. tab. 25. — *Burgsdorffia rigida* Meench.

Hab. in cultis totius Græciae (Sprun! Orph. Fl. eks. 537!), Archipelagi (Urv!), Cretæ (Sieb!), Smyrnæ (Boiss!), Cyprī (Ky!), Syriæ littoralis (Bl! Gaill!).

β. mutica. — Calycis dentes ovati vel oblongi acuti mutici superior duplo major. — *S. mutica* Boiss. MSS.

Hab. in collibus Pamphyliæ prope Adalia (Bourg!).

Ar. Geogr. Regio mediterranea Europæ, Africa borealis.

3. *S. purpurea* (Talbot in Bth. Lab. p. 742) annua, molliter villosa, collo ramosa ramis ascendentibus vel diffusis, foliis ovatis et oblongis crenato-dentatis, verticillastris sexfloris distantibus longe racemosis, foliis floralibus spinuloso-mucronatis verticillastro sublongioribus, calycis crebre et patule villosi basi subgibbi dentibus longe spinuloso-aristatis superiore maximo oblongo inferioribus a basi latiore subulatis, corollæ purpureæ rarius albæ calyce sesquilonioris labio superiore angusto ♂. Ic. Rchb. tab. 25.

Hab. in collinis et agris sterilibus Coreyrae (Talb.), Zacynthi (Marg!), Peloponnesi (Bory! Sprun! Heldr!), Thessaliæ (Orph. 538!), Eubeæ (Wild!), Etolie (Nieder!), Asiæ minoris (Auch. 1676).

Valde affinis *S. Romanae* cuius forsitan varietas, indumentum longius mollius et corollæ limbus major magis exsertus.

4. *S. montana* (L. Sp. p. 802) annua pumila laxe et longe pilosa viridis simplex vel a collo ramosa, foliis oblongo-lanceolatis superne remote denticulatis, verticillastris numerosis distinctis sed approximatis subsexfloris longe racemosis, foliis floralibus oblongis spinuloso-

mucronatis calyce longioribus, calycis patentim pilosi viridis subbilabiati dentibus omnibus æqualibus oblongis reticulatis spinuloso-mucronatis demum patentibus labii superioris ternis altius connatis, corollæ luteæ demum fuscescentis limbo parvo breviter exerto ⊖.
Ic. Jacq. Austr. tab. 434. — Rchb. Ic. tab. 25. — *Hesiodia bicolor* Mænch.

Hab. in cultis totius regionis a Græciâ! et Macedonia! ad Syriam! regiones Caucasicast, Persiam borealem! et australem!, Affghaniam (Griff!); a regione inferiori et montana ad (præsertim in australioribus, Libano!, Persiâ australi!) alpinam.

Variat *cryptantha*, corollâ diminutâ subobartivâ tubo calycis inclusâ, et *comosa*, foliis floralibus auctis flavidis ad apicem ramorum confertis comata.

Ar. Geogr. Gallia australis, Hispania, Italia, regio Danubialis, Dalmatia, Rossia media et australis, Africa borealis.

5. S. remota (Urv. Enum. p. 66) annua, albo-lanata sœpius a basi divaricatim ramosa ramis elongatis rigidis, foliis oblongo-spathulatis remote denticulatis, verticillastris distantibus subsexfloris longe et interrupte racemosis. foliis floralibus deflexis ovatis verticillastro æquilongis vel sublongioribus, calycis basi albo-lanati dentibus subæqualibus aristato-mucronatis demum patentibus, corollæ luteæ demum fuscescentis limbo exerto ⊖. *S. montana* Fl. Græc. tab. 55! non L.

Hab. in agris derelictis Atticæ et Eubeæ (Sprun! Boiss! Heldr!). Macedonia (Friv!). Ioniae ad Smyrnam (Boiss! Bal. 306!), insulae Samos (Urv!), Lycie (Bourg!), Pisidie (Heldr!), Cilicie littoralis (Bal!), Galatiæ et Ponti (Tchih!).

Affinis *S. montana* a quâ differt ramis elongatis crassioribus firmis verticillastris remotis, indumento lanato nec patule hirsuto calycis, calycinis dentibus elongatis sensim spinuloso-attenuatis, corollæ limbo in majore.

β. Lycaonica. — Flores subminores, folia parva, verticillastra magis approximata.

Hab. in tophaceis aridis Lycaoniæ inter Beychehr et Koniah (Heldreich!).

6. S. Balansæ (Boiss. Diagn. Ser. II, 4 p. 35) annua patule et longe hirsuta viscidula pallide virens a basi ramosa pumila ramis crassiusculis, foliis oblongo-lanceolatis suberenatis obtusiusculis, floralibus angustatis calyces æquantibus, verticillastris 6-4-floris subdistantibus vel approximatis racemosis, calycis patule hirsuti limbo reticulato subbilabiato, dentibus æquilongis ovato-oblongis longe spinulosis labii superioris ternis altius connatis, corollæ albæ vel purpureæ limbo calyce subbreviore ⊖.

Hab. ad Gülek Boghas Ciliciæ (Bal!), Lyciæ in collibus Elmali (Bourg!), in incultis Armeniae prope Gumuschkhané (Bourg!).

Calycis formâ affinis *S. montana* et *remote*, sed calyx duplo major fere quinque lineas longus, corolla purpurea vel alba nec flavo-fuscescens.

SECT. II. EMPEDOCLEA Bth. Lab. p. 374. — Herbæ perennes pluricaules basi induratae vel suffrutescentes. Folia floralia integerrima flores in verticillastris sessiles amplectentia. — Species habitu proprio donatae, sed characteribus gravioribus non distinctæ et interdum ægre limitandæ ut jam cl. Benth. animadvertisit.

7. **S. hololeuca** (Boiss. et Heldr. Diagn. Ser. I, 12 p. 69) basi indurata multicaulis dense et adpresso tota albo-pannosa, caulis superne trichotome corymbosis, verticillastris globosis intermedio solitario pedunculato lateralibus solitariis vel binis approximatis, foliis orbiculari-ovatis minute crenulatis obtusissimis inferioribus petiolatis, superioribus sessilibus oblongis, floralibus in urceolum brevem truncatum obsolete et brevissime bimucronatum calycibus dimidio breviorem connatis, calycis albo-pannosi dentibus breviter triangularibus obtusis, corollæ luteæ tubo brevissime exerto limbo minimo ♀. *S. rhytidia* Spr. — *S. rugosa* Poir.

Hab. in rupestribus calidissimis vallis Djoksu Tauri Isaurici inter Karaman et Ermenek 3000' (Heldr.).

Caules pedales, folia inferiora 9-12 lineas longa et lata, rami floriferi bis et trichotomi capitulo centrali brevius pedunculato, capitula avellanæ minora, calyx tres lineas longus. Species inflorescentia corybosæ insignis.

8. **S. Cretica** (Boiss. MSS. non L.) basi indurata multicaulis tota dense albo-lanata, caulis elongatis simplicibus vel rarius opposite ramosis, verticillastris numerosis distantibus longe spicatis, foliis oblongis et oblongo-lanceolatis basi attenuatis insimis obtusis subcrenatis intermediis acutiusculis, floralibus cordatis abrupte et breviter acuminatis calyces æquantibus, calycis lanati dentibus lanceolatis acutis tubo subduplo brevioribus, corollæ luteæ tubo subincluso ♀. *S. Syriaca* L. Spec. p. 801. — Ic. Fl. Græc. tab. 550. — Rehb. Ic. Exot tab. 57.

Hab. in siccis regionis montanae et subalpinæ Cretæ 2500'-5000' (Tourn., Sieb! Sibth., Heldr! Rauli!). Vulgo *Malotoxa*. Flores apibus gratissimi.

β. condensata. — Verticillastri superiores rarius omnes in spicam continuam condensati. Intermediis ad formam typicam transit.

Hab. in saxosis regionis superioris montis Delphi Eubeæ 4500'-5000' (Sprunt Heldr! Orph!). Vulgo *Tsai apo tou Delphi*.

Pedalis dense pannosa, capitula avellanæ minora. Hæc species in Syriâ non crescit et nomen Linneanum omni jure mutandum erat ut et illud *S. Cretica* Linn. quæ in Cretâ non habitat sed in insulis Canariensis, et est nunc *S. candicans* Ait.

9. **S. Sipylea** (Boiss. Diagn. Ser. I, 5 p. 32) basi indurata multicaulis adpresso cana, caulis erectis simplicibus vel stricte ramosis, verticillastris parvis spicatis inferioribus distantibus cæteris

approximatis, foliis integris lanceolatis inferioribus obtusis superioribus apice acutiusculis callosis, floralibus cordato-orbiculatis abrupte et brevissime apiculatis calyce subbrevioribus, calycis adpressiusculae hirsuti dentibus triangularibus acutis tubo duplo brevioribus, corollæ luteæ tubo breviter exerto ♂.

Hab. in arenosis calcareis regionis montanæ montis Sipyli prope Magnesiam Lydiæ (Boiss!).

Pedalis et sesquipedalis, capitula vix pisum æquantia, calyx 3 lineas longus. A *S. Creticā* indumento tenui adpresso, caulis ramosis, verticillastris minoribus valde distincta.

10. *S. erythrantha* (Boiss. et Heldr. Diagn. Ser. I, 12 p. 70) basi suffrutescens multicaulis adpressissime grisea, caulis erectis stricte ramulosis, verticillastris parvis distantibus laxe spicatis, foliis rugulosis crenato-denticulatis oblongis inferioribus breviter petiolatis obtusiusculis superioribus sessilibus acutiusculis, floralibus ovato-rotundis abrupte et breviter acuminatis calyce brevioribus, calycis griseo-hirti dentibus lanceolatis acentis tubo subbrevioribus, corollæ parvæ purpureæ tubo subexerto ♂.

Hab. in rupestribus Tauri Isaurici inter Marla et Adalia 4000' (Heldr!). Fl. Jul.

Planta sesquipedalis, folia pollicem et amplius longa 6-7 lineas lata, verticillastris 6-7 piso duplo maiores, calyx tres lineas longus. Species elegans florum colore, etc., a congeneribus distincta.

11. *S. Lycia* (Boiss. et Heldr. Diagn. Ser. I, 12 p. 70) basi suffrutescens pluricaulis, caulis crebre foliosis, ramis gracilibus virgatis subsimplicibus, verticillastris valde remotis laxissime spicatis, foliis parvis dense niveo-pannosis oblongo-spathulatis obtusis integris subpetiolato-attenuatis, axillis gemmiferis, floralibus oblongo-lanceolatis canaliculatis rigidis subrecurvis glabrescentibus acuminatis verticillastro longioribus, calycis inferne villosi dentibus strictis lanceolatis acutis multicis tubo sesquibrevioribus, corollæ luteæ tubo calyceem vix excedente ♂.

Hab. in saxosis pinetorum regionis inferioris Lyciae supra portum Tcherali (Heldr!). Fl. Mai.

Planta 1-1½-pedalis, folia 6-7 lineas longa apice 4 lata. Species foliis minutis, axillis gemmiferis, bracteis in cuspidem longam rigidam duram attenuatis insignis. Verticillastris avellana minores, calyx 4 lineas longus.

12. *S. Taurica* (M.B. Taur. Cauc. II, p. 43) basi suffrutescens pluricaulis tota adpresso tomentosa cana, foliis oblongo-lanceolatis inferioribus crenulatis obtusis spathulatis basi longe attenuatis superioribus sessilibus acutiusculis, verticillastris inferioribus (rarius omnibus) distantibus superioribus in spicam continuam approximatis, foliis floralibus membranaceis reticulatis virentibus cordatis abrupte et breviter acuminatis verticillastro vix longioribus, calycis dentibus lanceolatis tubo sesqui vel duplo brevioribus, corollæ luteæ tubo in-

cluso ꝑ. Ic. M.B. Cent. tab. 39. — *S. distans* Willd. Sp. III, p. 66 et ejus herb! specimen pessimum absque foliis! — *S. dura* Bth. Lab. p. 576! *S. Syriaca* Pall. Index non L. — *S. Georgica* C. Koch Linn. XXI, pag. 694.

Hab. in rupestribus calcareis Tauriae ad Karassubazar, Balaclava, Sebatopol et in alpibus (M.B., Stev., Rehm!), in præruptis Olympi Bithyni (Clem! Pichl!), in monte Tmolo Lydiæ supra Bozdag (Boiss!), Anatolia boreali inter Tokat et Gerida (Wied!), in Grusia Caucasica (C. Koch).

Indumentum eo *S. Creticæ* magis adpressum, bractæ longius acuminatæ membranaceæ nec herbaceæ tomentosæ; ab eâ certe differt. *S. distans* Willd. ad formam verticillastris omnibus distantibus spectat.

13. S. Ræseri (Boiss. et Heldr. Diagn. Ser. II, 4 p. 30) basi indurata multicaulis tota adpresso grisea, caulis virgatis simplicibus, verticillastris distinctis spicatis inferioribus distantibus, foliis mediocribus turionum magis canis elliptico-spathulatis obtusis petiolatis, intermediis linear-lanceolatis acutis crenulatis saepe subcurvis, floralibus viridibus membranaceis reticulatis abrupte et breviter acuminatis calyces non æquantibus, calycis flavi hirsuti dentibus lanceolatis acuminatis tubo subbrevioribus, corollæ luteæ tubo non exerto ꝑ.

Hab. in regione media montis Parnassi 4000'-5000' (Heldr! Guicci!, Orph. Fl. Gr. exs. 429!), montibus Arapokephalo et Velugo Eurytaniæ (Sprun!, Heldr. Herb. Norm. 490!).

Affinis *S. Tauricæ* et præsertim *S. Siculæ* Ucr. = *Bruticæ* Ten. indumento dense pannoso, foliis majoribus, calycis dentibus abbreviatis non acuminatis distinctæ. Folia inf. cum petiolo pollicaria 3-4 lineas lata. Capitula avellana submajora.

14. S. Scardica Griseb. Spicil. II, p. 144) basi indurata pluricaulis adpresso albo-lanata, foliis integris subtus rugulosis inferioribus oblongo-lanceolatis obtusis subspathulatis in petiolum brevem attenuatis superioribus linear-lanceolatis acutiusculis, verticillastris in spicam cylindricam densam solitariam vel duabus lateralibus brevioribus auctam confertis, foliis floralibus membranaceis reticulatis ciliatis ovato-deltoides in cuspidem patulam e spicâ exsertam attenuatis, calycis hirsuti dentibus lanceolatis acuminatis ciliatis tubo æquiforis, corollæ luteæ calycem parum excedente ꝑ. *S. florida* Boiss. et Heldr. Diagn. Ser. II, 4 p. 31.

Hab. in regione sylvatica Olympi Thessali prope œnobium Hagio Dionysi 4000' (Heldr! Orph. Fl. Gr. exs. 539!), in pratis montanis Scardi Macedoniae (Griseb!).

Sesquipedalis, folia inferiora 2-2½ poll. longa 4-5 lineas lata, spicæ 2-3-policares 9-10 lineas lata. Calyx 5-6 lineas longus.

15. S. theezans (Boiss. et Heldr. Diagn. Ser. I, 7 p. 58) basi indurata pluricaulis adpresso tomentella flavicans, caulis ascendentibus simplicibus vel superne opposite ramulosis, foliis rugulosis crassiœculis inferioribus oblongo-spathulatis obtusis petiolatis obso-

lete crenulatis vel integris superioribus late linear-lanceolatis, verticillastris inferioribus distantibus cæteris in spicam condensatis, foliis floralibus pubescentibus herbaceo-membranaceis subreticulatis a basi dilatata longe lanceolato-acuminatis saltem inferioribus verticillastro 2-3-plo longioribus, calycis adpresso pubescens dentibus lanceolatis acutis trinerviis tubo subbrevioribus, corollæ luteæ tubo inclusa ♀. *S. Syriaca* Bory et Ch. — *Phlomis clandestina* Bory et Chaub. Ed. prima. — *S. Cretica* Sm. Prodr. I, p. 400 non L.

Hab. in saxosis et alveis regionis superioris montium Malevo et Taygetus Laconiae (Heldr! Orph. Fl. Gr. exs. 14!), Odore et sapore gratissima sub nomine *Tzai* vel *Theæ Græcæ* colligitur et per totam Græciam mittitur.

6. *Cyllenea* Heldr. MSS. — Flavo-virescens, verticillastri magis dissiti, bractæ brevius acuminatæ. — *S. Peloponnesiaca* Boiss. et Heldr. Diagn. Ser. II, 4 p. 35.

Hab. in regione mediâ et superiore montium Kyllene et Olenos (Heldr!).

Semipedalis et pedalis, folia 1½-2-pollicaria 3-4 lineas lata, bractæ pollicem longæ. Affinis *S. Scardicea*, indumentum tenuius, spicæ laxiores, bractææ multo longius acuminatæ non longe ciliatæ, calyx adpresso pubescens nec longe hirsutus.

16. **S. Cilicica** (Boiss. et Bal. Diagn. Ser. II, 4 p. 29) basi vix indurata ad folia inferiora et ramos steriles adpresso canescenti-lanata superne patule viscido-hirta viridis, caulis a basi ascendentib; erectis simplicibus vel stricte ramosis in spicas elongatas laxas verticillastris distinctis constantes abeuntibus, foliis obscure crenulatis inferioribus oblongo et elliptico-spathulatis obtusis, floralibus tenuiter membranaceis reticulatis flavidis cuculiatis brevissime et obtuse acuminatis calyces obtentib;ns, calycis breviter glanduloso-hirti dentibus pilosis oblongo-linearibus trinerviis abrupte et brevissime mucronatis tubo sesquibrevioribus, corollæ luteæ tubo brevissime exerto ♀.

Hab. in Ciliciâ secus viam inter Bouloukli et Alladagh in regione montana inferiore (Bal. 504!) Fl. Jun.

Bipedalis, spicæ sœpe pedales, folia intermedia 1½-1-pollicem longa 4-5 lineas lata, verticillastri numerosi, calyx 4 lineas longus. Species dentium calycinorum structuræ inter affines insignis.

17. **S. phlomoïdes** (Boiss. et Bal. Diagn. Ser. II, 4 p. 33) basi suffrutescens ramosa procumbens tota longe et patule villosa flavo-virens, caulis erectis simplicibus, verticillastris paucis valde distantibus, foliis oblongo-spathulatis obtusis rugulosis obsolete crenatis radicalibus longe caulinis brevius petiolatis, floralibus membranaceis reticulatis ovato-cordatis longe attenuato-acuminatis calyces multo superantibus, calycis hirsuti et glandulæ aspersi dentibus a basi lanceolatâ trinervi longe subulatis tubo sesquibrevioribus corollam luteam subæquantibus ♀.

Hab. in monte Masmeneudagh Cappadociæ (Bal!).

Semipedalis, foliorum inferiorum lamina pollicaris 5-6 lineas lata, calyx 6 lineas longus; species foliis etiam caulinis longe petiolatis et calycis dentibus longe subulatis inter affines recognoscenda.

18. S. Pisidica (Boiss. et Heldr. Diagn. Ser. I, 12 p. 72) basi suffrutescens pluricaulis patule hirta patule virens, caulinis erectis simplicibus vel parce ramosis, verticillastris omnibus distantibus laxe spicatis, foliis rigidis valde nervosis integris vel rarius prope apicem paucidenticulatis late lanceolato-linearibus acutis, floralibus membranaceis a basi ovata breviter acuminatis, calycis pilosi et glandulis sessilibus obsiti dentibus lanceolatis acutis tubo sesqui-brevioribus, corollæ luteæ tubo inclusa ♀.

Hab. in rupestribus apricis montis Budrun Pisidiæ 5000' (Heldr!).

Pedalis, folia fere *Phlomidis herbaventri* 2-2½-pollices longa 6-8 lineas lata, calyx 4 lineas longus; affinis *S. Libanotica*, folia subintegra nec argute serrulata, bractæ majores membranaceæ.

19. S. Libanotica (Labill. Ic. Syr. 4, p. 13, tab. 8) basi suffruticosa pluricaulis ad ramos et folia novella albo-lanata superne plerumque virescens glabreseens, caulinis angulatis gracilibus elongatis simplicibus vel striete ramosis, verticillastris parvis longe et laxe spicatis inferioribus præsertim valde distantibus, foliis rigidis venoso-reticulatis duris oblongo-lanceolatis acutis mucronatis saepius argute serrulatis, floralibus submembranaceis orbiculato-cordatis abrupte et tenuiter mucronatis calyces æquantibus vel superantibus, calycis laciniis lanceolatis spinescenti-mucronatis tubo duplo brevioribus, corollæ luteæ tubo inclusa ♀. *S. arguta* Boiss. Diagn. Ser. II, 4 p. 32.

Hab. in Libani regione alpinâ (Labill! Bl!), Hermone (Post!), sylvaticis montis Cassii et ad rupes Antiochiae (Boiss!), monte Amano supra Arsus (Ky. 118!)!, valle Marla montis Gheidagh Isauriæ (Heldr!), Anatoliâ boreali inter Amasia et Toptcha (Wiedt!).

β. incana. — Tota etiam superne adpresso plus minus canescens vel cana. — *S. ambigua* Fenzl ex parte. — *S. Bourgæi* Boiss. MSS.

Hab. in Libani cedretis Hadet, ad Mar Serkis et Ain Zehalta (Bl!), in apricis Marasch Cataoniae (Haussk!), monte Gara Kurdistaniae (Ky. 575!), Tauro Cilicio circa Gülek (Ky. 280! Bal!), Armenia prope Baibut (Bourg!).

γ. linearis. Bth. in p.c. Prodr. — Virescens vel canescens, folia linear-lanceolata vel linearia sepe angustissima acutiora. — *S. microstegia* Boiss. et Haussk. MSS. — *S. ambigua* Fenzl in Sched. ex parte.

Hab. in Libano prope Bescherre (Ehrenb!), inter Aintab et Nisib (Haussk!), Tauro Cilicio 5000'-8000' (Ky. 448 et 239! Bal!), monte Aslandach Cappadociae (Bal!), et Akdaghi Catoniae (Haussk!), Ciliciæ Tracheæ prope Ermenek (Peron!), Lyciæ prope Elmali (Bourg. 209!).

Indumento, foliis plus minus angustis, etc., valde variabilis, sed verticillastris parvis avellanam æquantibus vel minoribus valde dissitis et foliis serratis semper recognoscenda.

20. *S. condensata* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 71) basi indurata patule et breviter lanatula sordide virens, caulis erectis simplicibus vel superne opposite ramosis paniculatis, foliis oblongis et oblongo-lanceolatis reticulato-nervosis argute serrulatis acutis, insimilis cito exsiccatis in petiolum attenuatis superioribus rigidis pungentibus, verticillastris omnibus in capitula oblonga condensatis, foliis floralibus cordato-orbiculatis ciliatis reticulatis in cuspidem tenuem calyces superantem abruptissime abeuntibus, calycis hirsuti dentibus lanceolatis acuminatis subpungentibus tubo sessilibrevioribus, corollæ luteæ tubo inclusæ ♂.

Hab. in regione inferiori Tauri Pamphyli inter Marla et Adalia 1000' (Heldr!) Fl. Julio.

Sesqui vel bipedalis, folia intermedia 1½-2-pollicaria 6-7 lata, capitula compacta 1-2-pollices longa. calyx 4 lineas longus. Species foliis argute dentatis distincta. *S. Libanotica* eodem gaudet charactere sed flores dimidio iniores et verticillastros dissitos habet.

β. procumbens. — Tota adpresso araneoso-grisea, caules breviores ascendentes vel basi procumbentes, spicæ basi interruptæ. — *S. Taurica* var. *procumbens* Diagn. loc. cit. p. 71.

Hab. in saxonis regionis subalpinae Tauri Pamphyli inter Maria et Adalia 4000' (Heldr!), in Cilicie Trachea prope Anamur (Per!).

Ob folia argute denticulata huic potius quam *S. Tauricea* adnumeranda.

21. *S. stricta* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 72) basi suffrutescens pluricaulis præter ramos steriles cum foliis villosos parce hirsuta late viridis. caulis erectis simplicibus, verticillastris præter infimum saepè subremotam in spicas longas approximatis, foliis ramorum sterilium ellipticis parvis longiusculæ petiolatis cæleris linearibus rigidis argute denticulatis spinuloso-mucronatis, foliis floralibus membranaceis concavis reticulatis virentibus ciliatis calyces occultantibus abruptæ et brevissime acuminatis, calycis glabriuscui dentibus ciliatis lanceolatis acutis tubo duplo brevioribus. corollæ luteæ tubo subincluso ♂.

Hab. in fruticetis apricis Tauri Pamphyli ad Ormana inter Marla et Adalia 3000' (Heldr!).

Pedalis et elatior, folia caulina 1-2-pollicaria 3-4 lineas lata, spicæ 2-6-pollicares. Foliis angustis serratis affinis *S. Libanotica*, sed præter caules simplices et verticillastros approximatos differt bracteis multo majoribus 8-9 lineas longis atque latis; a *S. condensata* foliorum formâ et bracteis magnis. Spicæ continuæ ut in *S. Tauricea* formis, sed in hac folia obsolete crenata nec serrata.

22. *S. pullulans* (Vent. Hort. Cels tab. 98) inferne griseo-sublanata superne pubescenti-viscida virens, caulis crassis elatis opposite et striete ramosissimis, verticillastris numerosis vel omnibus distantibus vel superioribus condensatis. foliis rigidis valde reticulatis crenato-denticulatis apice spinulosis oblongo-lanceolatis, floralibus nervosissimis membranaceis ciliatis suborbiculatis transverse

latrioribus in cuspidem patulam vel recurvam abrupte attenuatis flores superantibus, calycis viridis puberuli dentibus lanceolatis rigidis ciliatis corollam subæquantibus tubo subbrevioribus ȝ.

Hab. in siccis regionis inferioris et montanæ Libani supra Tripoli usque ad Eden (Bl!), supra Sidonem (Gaill!), ad Anubin 3500' (Ky. 362!), in Palestinâ prope Nazareth (Bové! Sieb! Auch! Gaill!).

Caules rigidi sœpe tripedales tetragoni, folia coriacea 2-4-pollices longa, verticillastri mediocres. Species ramositate et bracteis valde nervoso-exaratis insignis.

23. *S. perfoliata* (L. Sp. 892) partibus junioribus longe albo-hirsuta vel lanata demum glabrescens breviter viscidulo-hirta flavovirens, caulis crassis elatis rarius simplicibus sœpius stricte paniculatis, foliis magnis coriaceis valde nervosis infimis in petiolum attenuatis minoribus, cæleris semiamplexicaulibus oblongis et oblongolanceolatis denticulatis acutis vel acuminatis, verticillastris sœpius distinctis et inferioribus distantibus, foliis floralibus membranaceis viridibus reticulatis late cordatis longe acuminato-attenuatis flores superantibus, calycis dentibus lanceotatis ciliaus tubo glandulis asperso dimidio brevioribus, corollæ flavæ tubo inclusu ȝ.

Hab. in siccis, alveis regionis inferioris et montanæ, regio media montis Athos Macedoniæ (Janka!), montes Smyrnæ supra Bournabat (Boiss!), Taurus Pamphylius (Heldr!), Cilicia prope Mersina et Gülek (Bal! Ky!), Syria prope Antiochiam et in montibus Latakieh (Boiss!).

β. condensata. — Verticillastræ in spicas oblongas contractas condensati, corolla sœpe rosea vel purpurascens.

Hab. in dumetis prope Karabiyukle 4000' inter Aintab et Marasch (Haussk!), circa Beyrouth (Bl!).

Species erronee in Siciliâ et Hispaniâ indicata.

Species delenda.

S. incana L. *β tragoriganum* Blh. in D.C. Prodr. XII, p. 444 ex specimine herb. Benthamiani est omnino *S. incana* Hispanica probabilissime ex schedulæ commutatione erronee ex Cretâ indicata.

STACHYS (L. Gen. N° 719).

Calyx tubuloso-campanulatus 5-10-nervius ore æqualis vel obliquus rarius subbilabiatus. Corollæ tubus cylindricus æqualis inclusus rarius breviter exsertus intus transversim vel oblique annulatum pilosus rarissime nudus, limbus bilabiatus labio superiore erecto integro vel retuso rarissime bifido sœpe subfornicato, inferiore longiore patente trilobo laciniâ mediâ maximâ. Stamina quatuor didynamia exserta inferioribus longioribus defloratis sœpe ad latera faucis

dejectis; antheris bilocularibus loculis divergentibus vel divaricatis. Nuces rotundato-obtusæ. — Herbæ perennes et annuæ, vel suffrutescens.

Specierum Orientalium dispositio.

SECT. I. ERIOSTACHYS Blh. Lab. p. 534. — Perennes. Verticillastri sæpius pluri vel multiflori bracteis dimidium calycem æquantibus vel superantibus. — Sectio quoad specierum delimitationem difficilis et cujus nonnullæ e posterioribus transitus ad *Stachyotypum* præbent.

§ 1. **Germanicæ.** — Corolla majuscula (magn. *S. Germanicæ* vel major).

* Calyx non bilabiatus.

+ Corollæ labium superius integrum vel retusum.

1. Corolla ochroleuca vel flava.

S. Orientalis, Tmolea.

2. Corolla carnea rosea vel purpurea.

S. Libanotica, lanata, Tournefortii, alpina, Cretica, Thirkei, Germanica, Ehrenbergii, Cassia, Heldreichii, Huetii, Balansæ, pinetorum, spectabilis.

++ Corollæ labium superius bifidum.

S. Græca.

** Calyx bilabiatus.

S. Persica.

§ 2. **Micranthæ.** — Flores eis præcedentium fere triplo minores. Corolla carnea vel rosea labio superiore integro vel retuso.

S. viticina, setifera, longespicata, rosea.

SECT. II. STACHYOTYPUS. — Sectiones *Stachyotypus*, *Chilostachys*, *Chamaesideritis*, *Zietenia* et *Ambleia* Blh. in D.C. Prodr. — Perennes, verticillastri sæpius pauciflori, bracteæ sæpissime brevissimæ vel nullæ. Forma dentium calycis, corollæ color et ejus annulus pilosus plus minus manifestus notæ nimis variabiles sunt et ideo omnes sectiones superne citatas in unam conjungere debui.

§ 3. **Sylvaticæ.** — Caules herbacei et facies *Eriostachydis*.

S. sylvatica, hydrophila, patustris, viridis, mollissima, Messeniaca, canescens.

§ 4. **Fragiles.** — Caules ex fissuris rupium penduli fragiles.

S. Finardi, longiflora.

§ 5. **Rectæ.** — Caules erecti virgati rigiduli.

S. recta, Parolinii, nepetæfolia, Iberica, virgata, angustifolia, distans, bombycinæ, aleurites, diversifolia, subnuda, viscosa, Benthamiana.

§ 6. **Fruticulosæ.** — Inferne fruticulosæ ramosissimæ ramis erectis pumilis in nonnullis apice demum spinescentibus. Plantæ glabrae vel plus minusve pilosæ rarius cinereæ sed non albo-tomentosæ.

S. Swainsoni, Spruneræ, tetragona, spinosa, micronata, acerosa, fruticulosa, multicaulis, Aucheri, ixodes, pilifera.

§ 7. **Aambleiæ.** — Basi suffrutescentes ramosæ ramis erectis brevibus, undique dense albo-tomentosæ.

* Calycis dentes mutici.

S. obtusierena, nivea, inflata, parviflora, affinis.

** Calycis dentes spinulosi.

S. Palestina, Kotschyæ, tomentosa, laxa, candida, chrysanthæ, Iva.

§ 8. **Infrarosulares.** — A sectionibus prioribus ramis floriferis sub rosulæ vel sub ramo centrali sterili nascentibus distinctæ.

S. lavandulæfolia, pumila, rupestræ, citrina, maritima, pubescens.

SECT. III. OLISIA Dumort. — Species annuae.

S. amara, plumosa, spinulosa, Arabica, arvensis, ramosissima, satureioidæ, burgdorffioïdes, nevralcalycina, Persepolitana.

SECT. I. ERIOSTACHYS.

§ 1. GERMANICÆ.

1. **S. Orientalis** (Vahl Symb. II, p. 64, an L. Sp. p. 813 ?) perennis molliter villosa pallide virens, caule elato subsimplici, verticillastris multifloris 1-2 inferioribus valde distantibus cæteris in capitulum ovatum vel oblongum terminalem confertis, foliis crenulatis rugulosis inferioribus longe petiolatis oblongo-lan-

lanceolatis obtusis basi rotundatis rarius subcordatis, floralibus ovatis cordatis attenuato-acuminatis vel acutis flores superantibus, bracteis elliptico-rhombeis acutis, calycis hirsuti recti dentibus lanceolato-subulatis dimidium tubum æquilibus vel superantibus, corollæ ochroleucæ villosæ labio superiore integro vel retuso ♀. Ic. Fl. Græc. tab. 560. — *S. obliqua* W. K. Hung. II, p. 142 tab. 134! — *S. Montbretii* Bth. Ann. Sc. Nat. 6, p. 48. — *S. Heraclea* β lutea Bth. in D.C. Prodr. XII, p. 463. — *S pauciflora* Visian. Ill. pag. 10, tab. 2 ex descript. et iconæ hujus videtur forma pauciflora.

Hab. in fruticetis regionis inferioris et montanae, Thracia borealis (Jankat!), radices Olympi Bithyni (Sibth! Month! Auch! Pichler), Tmolus supra Philadelphiam et Sipylus supra Magnesiam (Boiss!).

Sesqui vel bipedalis, insignis foliis floralibus latis inferioribus praesertim cordatis, bracteis dilatatis, corollâ flava. *S. Heraclea* All. planta occidentalis verticillastris dissitis paucifloris, calycis abbreviati dentibus ovatis, corollâ purpureâ omnino aliena est

Ar. Geogr. Croatia, Dalmatia.

2. S. Tmolea (Boiss. Diagn. Ser. I, 5 p. 29) adpresse et molliter tomentosa sericea, caulinis simplicibus ascendentibus, verticillastris 3-4 inter se remotissimis multifloris, foliis suberenuvatis inferioribus anguste oblongo-spathulatis obtusis in petiolum attenuatis, floralibus a basi latiore ovato-lanceolatis vel lanceolatis acuminatis verticillastra superantibus, bracteis subulato-lanceolatis, calycis hirsuti vix obliqui dentibus lanceolatis acuminatis tubo subtriplo breviribus, corollæ ochroleucæ labio superiore retuso ♀.

Hab. in regione alpinâ montis Tmoli supra Bozdagh in Lydiâ (Boiss!), in faucibus Olympi supra Broussam (Pichler!).

Corollis flavis affinis *S. Orientali*, foliorum indumentum multo densius sericeo-pannosum, folia basi attenuata nunquam cordata, verticillastri omnes remotissimi, bracteæ multo angustiores, dentes calycini breviores.

3. S. Libanotica (Bth. in D.C. Prodr. XII, p. 462) caulinis elatis ramosis virgatis rubellis rigidis patule setulosis, verticillastris 6-8-floris omnibus valde remotis, foliis rugulosis crenatis molliter et adpresse villosis subsericeis inferioribus oblongis petiolatis basi rotundatis vel subtruncatis, floralibus lanceolatis acutissimis flores superantibus supremis glabratis rigidis, bracteis subulatis calyces subæquantibus, floribus magnis, calycis denum coriacei tubo elongato limbo infundibuliformi reticulato-nervoso glanduloso glabrato dentibus rigidis carinato-nervosis triangularibus pungenti-spinosis, corollæ valde hispidæ tubo subexerto labio superiore integro ♀.

Hab. in Libano (Auch. 1714!), ad Khan el Djedid inter Berythum et Damascum (Gaill!), in Antilibano ad margines vinorum prope Rascheya 4500' (Ky. 160!), ad Ouadi el Habesch (Gaill!), ad Ouadi el Harir (Gaill!).

Planta elata, folia 3-4-pollicaria, verticillastris 2-4 pollices remoti, calyx de num fere sex lineas longus valde coriaceus angulis valde nervosis dentibus reticulatis valde pungentibus. Corollæ color e sicco carneus. Species insignis.

β. minor. — Caules humiliores simplices, calyces breviores dentibus brevius spinosis. — *S. ciliaris* Boiss. Diagn. Ser. I, 12 p. 78.

Hab. in sylvaticis regionis inferioris montis Cassii Syriæ borealis (Boiss!).

4. *S. lanata* (Jacq. Ic. Rar. I. p. 44, tab. 107) perennis cespitosa densissime pannoso-lanata sericea, caulis a basi procumbenti radicanti erectis simplicibus vel parce ramosis, foliis subcrenulatis inferioribus oblongo-spathulatis in petiolum attenuatis superioribus sessilibus ellipticis, floralibus supremis verticillastro brevioribus, verticillastris multifloris supremis in spicam approximatis, calycis dense albo-lanati ore obliqui dentibus triangularibus acutis, corollæ roseæ tubo inclusa labio superiore integro lanato ♀. Ic. Rehb. tab. 9. — *S. Byzantina* C. Koch Linn. XXI, p. 686!

Hab. in siccis circa Byzantium (C. Koch!), Bithyniâ ad Boli et Toptcha (Wied!), Tauriâ ad Sebastopol (S^r-Supéry!), Caucaso 3500-6000' (M.B., C.A. Mey.), Daghestaniâ australi 4800' (Rupr!), Georgiâ Caucasicâ (Hohen!), Persiæ borealis prov. Ghilan (Auch. 5176! Buhse!), ad Asterabad et Siaret (Bge.).

β. alpina. — Minor, semipedalis rarius pedalis, folia diminuta radicalia 3-4 lineas lata in petiolum longe attenuata, verticillastris insimne sæpe excepto in capitulum oblongum approximati.

Hab. in siccis regionis alpinæ Olympi Bithyni (Boiss! Pichler!).

5. *S. Tournefortii* (Poiret Dict. V, p. 227) perennis dense pannosa albo-lanata, caulis elatis simplicibus vel parce ramosis, verticillastris multifloris omnibus distantibus, foliis crenulatis minute rugulosis inferioribus petiolatis ovatis cordatis, floralibus oblongo-lanceolatis obtusiusculis inferioribus verticillastro vix longioribus, calycis albo-lanati recti dentibus lanceolatis acutis spinosis tubo triplo brevioribus, corollæ purpureæ tubo inclusa labio superiore hirsuto integro ♀. *S. Sieberi* C. Koch Linn. XXI, p. 685. — *S. Cretica* Sieb. exs. non L. — *Stachys Cretica latifolia* Tourn. Inst. 186. Voy. I, 29, tab. 1.

Hab. in campis sterilibus Cretæ (Sieb!), prope Cydoniam (Tourn.), in prov. Messara prope Gortynam (Heldr!), ad Khania, Malaxa, Omalon (Raoul!).

Planta 2-3-pedalis, indumentum albo-pannosum sericeum *S. lanata*, sed folia cordata et calyx rectus nec obliquus quo charactere a formis *S. Germanicae* etiam discrepat.

6. *S. Alpina* (L. Sp. 812) perennis caule elato ramoso hirsuto superne glanduloso hirta, foliis viridibus hirtis membranaceis crenatis inferioribus ovato-oblongis cordatis petiolatis, verticillastris 10-12-floris distinctis inferioribus remotis, foliis floralibus lanceolatis acutis integris flores superantibus, calycis glanduloso-hirti subobliqui dentibus lanceolato-ovatis acutis spinosis tubo duplo brevioribus, corollæ rubro-fuscae hirtæ labio superiore integro ♀. Ic. Rehb. tab. 8

Hab. in sylvaticis et pratis, mons Scardus Macedoniæ in regione subalpinâ (Griseb.), Caucasus (Rehm!).

β. discolor. — Folia subtus canescentia, corolla roseo-purpurascens.

Hab. in regione superiori montis Peristeri Macedoniæ 6500' (Orph!).

Ar. Geogr. Hispania borealis, Gallia, Belgium, Germania, Italia, regio Danubialis.

7. **S. Cretica** (Sibth. et Sm. Fl. Græc. VI, p. 47, tab. 558 et prob. L. Sp. 812) perennis adpresso tomentosa canescens vel pallide virens, caulis erectis stipes simplicibus, verticillastris multifloris remotis, foliis tenuiter rugulosis crenulatis obtusis inferioribus oblongo vel linear-lanceolatis petiolatis basi attenuatis, floralibus oblongo-triangularibus vel breviter lanceolatis inferioribus verticillastro longioribus, calycis lanatuli subobliqui dentibus triangular-lanceolatis breviter spinosis tubo subtriplo brevioribus, corollæ roseæ tubo inclusa labio superiore recto lanato retuso ♀. *S. Italica* Bot. Græcorum vix Miller.

Hab. in collinis et lapidosis regionis inferioris montanae et in australioribus subalpinæ, Zacynthus (Marg!), Attica et Bœotia (Sprun! Heldr! Orph. Fl. exs. 77!), Cyclades (Orph! Urv!), Rhodus (Bourg. exs. sub *S. Græcæ*), Creta (Heldr!), Anatolia omnis occid. et australis a Lydiâ ad Ciliciam (Boiss! Bal! Bourg!), Paphlagonia (Wied!), Armenia Turcica ad Baibut (Bourg!), Tauria (ex Ledeb.), Cappadocia (Bal!), Cyprus (Ky!), Syria borealis inter Aintab et Nisib (Haussk!), et littoralis (Boiss!), Libanus usque ad Eden (Bl! Gaill!), Antilibanus (Ky. 476!), Palestina (Boiss!).

β. paniculata. — Inflorescentia ramosa saepe paniculata. — *S. Mersinæ* Boiss. Diagn. Ser. II, 4 p. 37.

Hab. in incultis Pisidiæ (Heldr!), Ciliciae littoralis ad Mersina (Bal!), Syria prope Aintab (Haussk!), Palestina ad Hierosolymam (Auch. 1740!).

γ. *Garana*. — Inflorescentia ramosa paniculata sed verticillastri omnes in spicas breves condensati. — *S. Garana* Boiss. Diagn. Ser. I, 12, p. 76.

Hab. in lapidosis montis Gara Kurdistaniæ (Ky. 413!).

Species in parte occidentali ditionis frequentissima et ab affinibus foliis angustioribus basi semper attenuatis nec truncatis vel cordatis distincta. Huic valde affinis est *S. Italica* Mill. = *S. salviaefolia* Ten. Italiam australi et Dalmatiæ frequens sed que foliis inferioribus basi subcordatis differre videtur. Nostra est certissime species in Fl. Græcæ descripta et depicta et prob. quoque Linnei, indumentum variat luride virens et canescens.

8. **S. Thirkei** (C. Koch Linn. XXI, p. 685) perennis, adpresso tomentosa canescens, caulis erectis simplicibus, verticillastris plurifloris omnibus præter supra distinctis et remotis, foliis parvis subcrenulatis subrugosis inferioribus oblongo-lineariibus obtusis in petiolum attenuatis, floralibus oblongo-triangularibus inferioribus verticillastro sublongioribus, bracteis lanceolatis spinosis calyce brevioribus, calycis longe et densissime albo-lanati dentibus demum patentissimis et recurvatis a basi lanceolata longe subulatis rubris dimidio tubo longioribus, corollæ roseæ extus valde hirsutæ labio superiore integro ♀.

Hab. in Bithyniâ (Thirke!).

β . *condensata*. — Caulis ramosus, verticillastra præter inferiora 1-2 interdum distantia in spicas oblonga condensata.

Hab. circâ Byzantium et in Anatoliâ boreali circa Sojut (Wiedem!).

Indumentum adpressum, caules rigidi 1-1 $\frac{1}{2}$ -pedales, folia angusta basi attenuata *S. Creticæ*. Species quodammodo etiam *S. Orientali* affinis sed ab omnibus calycis dentibus demum patenti-recurvatis longissime spinulosis distinctissima.

9. **S. Germanica** (L. Sp. p. 812) perennis dense et laxe lanato-villosa, caulis simplicibus vel parce ramosis, verticillastris multifloris inferioribus remotis superioribus approximatis, foliis rugulosis crenatis inferioribus petiolatis ovatis cordatis, floralibus lanceolatis verticillastro sublongioribus, calycis villosi subobliqui dentibus triangulari-ovatis breviter spinulosis tubo duplo brevioribus, corollæ purpureæ tubo subincluso labio superiore hirsuto integro vel retuso γ . Ic. Jacq. Austr. tab. 319. — Rchb. Ic. tab. 9. — *S. polystachya* Ten. Nap. tab. 53. — *S. Reinerti* Heldr. exs. — *S. Pisidica* Boiss. et Heldr. Diagn. Ser. I, 12, p. 75 (forma laxius villosa virens). — *S. obtusata* Boiss. in Bourg. exs.

Hab. in pratis siccis et in sylvaticis regionis montanæ et subalpinæ, Olympus Thessalus (Auch. 1720!, Heldr! Orph!), Eubæa (Wild!), mons Tymphratus (Etolia 50°) (Sam. et Guicc. in Heldr. herb. Norm. 743!), Macedonia et Thracia (Griseb.) mons Davros Pisidiæ (Heldr!), Anatolia borealis (Wied!), Pontus (Tchih!), Pontus Lazicus ad Djimil (Ball!), Caucasus (Ledeb., Rupr!), Tauriæ montes (Rehm!).

β . *Bithynica* — Pedalis, indumentum magnis floccosum, folia latiora magis cordata, verticillastræ pauci in spicam oblongam basi interruptam condensati, folia floralia ovato-subcordata. — *S. Bithynica* Boiss. Diagn. Ser. I, 5 p. 28.

Hab. in lapidosis regionis alpinæ Olympi Bithyni supra Kirkbounar (Boiss!), montis Kitirlidagh (Pichl!).

γ . *penicillata* — Humilior, folia minora inferiora obscurius cor data, floralia oblonga obtusa, verticillastræ pauciores. — *S. penicillata* Heldr. et Sart. in Boiss. Diagn. Ser. II, 4 p. 37. — *S. Italicæ* et *S. dasyanthes* var. Boiss. et Heldr. olim.

Hab. in saxosis regionis subalpinæ et alpinæ Taygeti, Chelmos, Olenos, Kyllenes, montis Delphi Eubæa (Heldr!), Parnassi (Heldr! Orph. Fl. Gr. exs. 439!).

Hæc forma in Aprutii monte Morrone quoque crescit.

δ . *intermedia*. — Elatior ramosa, verticillastræ magis dissiti. — *S. Alpina* var. *intermedia* Bth. in D.C. Prodr. — *S. intermedia* Ait. Kew 2 p. 201. — *S. Sibirica* Link Enum. II, p. 109. — *S. rosea* Hohen. Enum. Talysh. p. 240!

Hab. in Caucaso (C.A. Mey.), in herbosis montanis ditionis Elisabethpol Gerogiae Caucasicæ (Hohen!).

Indumento longe et patule lanato et glandularum inflorescentiæ defectu *S. Germanicae* ut cl. Bentham ipse suspicatur nec *S. Alpinae* adnumeranda.

Ar. Geogr. Europa media ab Anglia et Belgio ad Lusitaniam, Hispaniam centralem, Italiam, Dalmatiam, regionem Danubialem, Rossiam medium et australem.

10. S. Ehrenbergii. perennis, tota dense breviter et molliter tomentosa cana, caulis valde simplicibus vel parce ramosis, foliis rugosissimis crenulatis inferioribus petiolatis ovatis cordatis obtusissimis, floralibus oblongis praeter supra flores superantibus, verticillastris omnibus remotissimis, bracteis subulatis calyce brevioribus, calycis tomentoso-lanati subobliqui tandem accreti dentibus ovatis longe spinulosis inferioribus brevioribus et angustioribus, corollæ valde hirsutiæ labio superiore retuso ♀. *Leonurus mollis* Ehrenb. pl. exs.

Hab. in lapidosis et ad rupes regionis alpinae Libani, ad Kafrum prope Bischerre et in summo monte Makmel (Ehrenb!), circa Cedretum et in summis Djord Arasch (Ky. 307 sub *S. Pisidicā*), ad Star es Soud et in cacuminibus inter Yamouny et Dimam (Bl !), supra Khan el Djedid (Gaill. 2146!).

Pluripedalis, in summis humilior, insignis tomento denso albo absque ullo nitore, foliis valde reticulato-rugosis obtusissimis, verticillastris magnis tomentosis valde remotis. His notis et praesertim calyce inaequidentato et valde accrescenti a formis polymorphæ *S. Germanicæ* distincta videtur. Floris color e sicco incertus.

11. S. Cassia (Boiss. Diagn. Ser. I. 12 p. 76) perennis, caulis elatis parce et adpresso hirtis simplicibus vel parce ramosis, foliis rugulosis crenulatis superne viridibus subtus adpresso canescensibus inferioribus petiolatis oblongis truncato - subcordatis, floralibus oblongo-lanceolatis inferioribus flores superantibus, verticillastris multifloris inferioribus dissitis cæteris contiguis, calyeis hirsuto-lanati albi obliqui dentibus a basi latiore longe subulato-spinosis demum suhpantibus tubo brevioribus, corollæ roseæ tubo inclusa labio superiore hirsutissimo integro ♀. *S. Italica* var. *chondrostachys* Heldr!

Hab. in sylvaticis regionis inferioris montis Cassii Syrie borealis (Boiss!), in monte Korthiati Macedoniae (Heldr! Orph. Fl. Gr. exs. 535!).

Planta 2-3-pedalis a *S. Creticā* et *S. Italicā* foliis subcordatis, verticillastris superioribus condensatis, dentibus calycis longioribus patentibus longe subulato-spinosis, indumento parco adpresso distincta, a *S. Germanicæ* formis indumento adpresso canescens et dentibus calycis longe pilosis etiam aliena.

12. S. Heldreichii. perennis, adpresso et breviter tomentella canescens, caule elato ramoso vel paniculato, foliis tenuibus crenulatis subrugosis inferioribus majusculis petiolatis ovato-oblongis cordatis, verticillastris numerosis praeter inferiora remota in spicas longas condensatis, foliis floralibus oblongo-lanceolatis inferioribus flores subsuperantibus, floribus mediocribus, calycis hirsuti dentibus triangularibus spinulâ terminatis tubo triplo brevioribus, corollæ roseæ tubo inclusa labio superiore hispido integro ♀. *S. dasyanthes* Heldr. exs. non Rafin. — *S. penicillata* var. *ramosa* Boiss. olim.

Hab. in herbidis humidis et paludosis Bœotiae ad Lebadeam (Heldr!), Thessalia ad Katarine (Heldr! Orph!), Eubeæ prope Creos (Orph!), Byzantii, (Auch. 1707! Coumany!), Bithyniae ad Safranbol (Wied!).

Planta elata 2-3-pedalis, spicæ basi interruptæ sœpe pedales. Differt a formis *S. Germanice* indumento brevi adpresso, spicis longis condensatis, floribus paulo minoribus, foliis membranaceis, *S. dasyanthes* Raf. Sicula floribus minutis affinis differt foliis angustis acute dentatis, verticillastris omnibus valde distantibus. *S. Cassia* denique indumento adpresso et inflorescentia condensatâ quoque donata distinguitur floribus majoribus et calycis dentibus longe et patule spinoso-attenuatis.

13. S. Huetii, perennis. breviter lanata luride cinereo-virens, caulis herbaceis a basi procumbenti denudatâ erectis inferne striete ramulosis, foliis parvis crenulatis rugulosis oblongis breviter petiolatis obtusis basi truncatis vel subcordatis, verticillastris 6-10-floris parvis albo-lanatis remotissimis foliis floralibus breviter oblongo-lanceolatis acutis inferioribus eis sublongioribus suffultis, bracteis linearis-subulatis calyce brevioribus, calycis hirsuto-lanati subobliqui demum acreti dentibus dimidio tubo longioribus lanceolatis longe spinuloso-attenuatis, corollæ (e secco carneæ) labio superiori parce hirsuto integro ♀. *S. Brantii* Huet pl. exs. non Bth.

Hab. ad radices montis Techdagh Armeniae Turcicæ et circa Tortum (Huet!), prope Erzerum (Calv!).

Caules pedales inferne breviter ramulosi in spicam laxissimam verticillastris 6-10-inferioribus 2 pollices distantibus constantem abeunt, folia etiam inferiora vix pollicem longa, calyx fructifer sex lineas fere longus. Species insignis facie *Clinopodii* bracteis abbreviatis ad *Stachytypum* accedens.

14. S. Balansæ (Boiss. et Ky. in Ky. exs. 1859) crispule hirsuta cinerascens, caulis erectis simplicibus vel parce ramosis, verticillastris multifloris inferioribus distantibus supremis sœpe approximatis, foliis crenatis vix rugulosis a basi subcordatâ anguste et elongatim oblongis inferioribus longe petiolatis obtusis, floralibus oblongo-lanceolatis integris acutis inferioribus verticillastro sublongioribus, bracteis linearibus calyces subæquantibus. floribus pedicellatis, calycis longe albo-hirsuti subincurvi et obliqui dentibus lanceolatis spinosis tubo duplo brevioribus tandem patentí-subcurvatis, corollæ lilacinae tubo inclusò labio superiore albo-hirsutissimo integrò ♀.

Hab. in schistosis alpium prope Musch inter Astragalos 7500' (Ky. 441!) monte Jyldisdag Galatiae (Wied!), in graminosis Berydag Cataoniae 6000' (Haussk!), in collibus prope Baibut (Bourg!), in monte Aslandach Cappadociae (Bal.).

Sesquipedalis, folia angusta et longa fere *Betonice* (inferiora 2½-4-pollices, longa 6-15 lineas lata), verticillastris 3-5 hirsutissimi magn. nucis. Species inter affines floribus longius pedicellatis et calycis dentibus subcurvatis insignis.

15. S. pinetorum (Boiss. et Bal. Diana. Ser. II, 4 p. 36) perennis, caule elato adpresso tomentello subsimplici, foliis crenato-

dentatis amplis membranaceis vix rugosis ovatis vel ovato-oblongis cordatis petiolatis supra viridibus pubescentibus subtus albo-tomentellis, verticillastris plurifloris distantibus, foliis floralibus oblongis acuminatis inferioribus verticillastro longioribus, bracteis linearisubulatis calyce brevioribus, calycis virentis patule hirti et subglandulosi subincurvi subobliqui dentibus subæqualibus lanceolatis spinulosis, corollæ carneæ labio superiore integro longe hirsuto ♀.

Hab. in sylvis Pini Laricionis supra Gülek Boghaz Tauri Cilicie (Bal. 535!).

Planta 2-3-pedalis, foliis discoloribus membranaceis ad *S. spectabilem* accedens a quā tamen floribus duplo majoribus, calyce subincurvo ore obliquo nec recto, etc. recedit. Magis affinis *S. Balansæ* a quā foliis magis membranaceis discoloribus latis nec anguste oblongis, floribus subsessilibus, calycis non albo-hirsutissimi dentibus non patent-curvatis specificè differre videtur; an tamen hujus varietas? *S. spectabilis* Choisy foliis discoloribus quoque donata differt floribus dimidio minoribus, calyce recto nec obliquo, etc.

16. ***S. spectabilis*** (Choisy in Pl. rar. Hort. Gen. I, p. 27) perennis, caule elato tomentello simplici vel sèpius ramoso, foliis crenulatis non rugosis supra viridibus rarius canescensibus subtus cano-tomentosis petiolatis ovatis cordatis, floralibus oblongo-lanceolatis acutis integris inferioribus verticillastra superantibus, verticillastris 6-10-floris distinctis superioribus sape confluentibus, bracteis elliptico-rhombeis cuspidiatis calyce brevioribus, calycis recti dense tomentose-hirsuti dentibus lanceolatis longe spinoso-attenuatis dimidiis tubo longioribus, corollis purpureis vel carneis ochroleuco-variegatis extus hirtis labio superiore integro ♀. *S. hypoleuca* C. Koch Linn. XXI, p. 688 et *S. elata* C. Koch ibid. p. 687.

Hab. in dumosis humidis et præsertim ad rivulos regionis sub. inæ et alpinæ, Armenia ad Erzerum (Auch. 1705! Calv!), ad Baibut (Bourg!), Cataonia ad Marasch (Haussk!), Amanus Syriae bor. (Ky. 148!), mons Gara Kurdistanæ (Ky. 328!), montes Avroman et Schahu Kurdistanæ Persice 7000' (Haussk!), Persia austro-occid. in montibus districtus Kuh Kilonyeh 8000' (Haussk!) et Kuh Daëna (Ky. 640!), mons Elbrus prope Derbend (Ky. 412!).

Planta 3-4 et interdum 6-8-pedalis, foliis magnis discoloribus submembranaceis minime rugosis, etiam caulinis sapius cordatis, indumento adpresso, calycis recti nec subobliqui dentibus attenuato-spinulosis a formis *S. Germanicae* cui eam cl. Bth. adnumerat certe aliena.

17. ***S. Græca*** (Boiss et Heldr. Diagn. Ser. I, 12, p. 77) perennis, caulis ascendentibus simplicibus patule hispidis, verticillastris omnibus distantibus, foliis crenatis subrugulosis membranaceis adpresso hirsutis subsericeo-canescensibus vel virentibus, inferioribus oblongis obtusis basi rotundatis vel subcordatis, floralibus subcordatis oblongis verticillastris superantibus supremis acutis, calycis longe hirsuti recti dentibus triangulari-lanceolatis inæquilateris breviter spinuloso-attenuatis tubo duplo brevioribus, corollæ roseæ limbo ampio parce hirsuto labio superiore profunde bifido ♀. *S. Lusitanica* Bor. et Ch. Fl. Pelop. non Link et Hoffm. — *S. Heraclea*

Bot. Græc. non L. — *S. acutifolia* Link Linn. IX, p. 575 nomen omnino incongruum.

Hab. in fruticetis, vineis, agris Messeniæ (Bory! Heldr!), montis Malevo (Sprun!), insulae Poros (Wied!), Argolidis (Heldr!), montis Onii Bæotiae (Heldr!), montis Delphi Eubeæ (Sprun!).

Pedalis vel sesquipedalis, facies *S. Heracleæ* a quā ut et ab omnibus affinis differt corollā magnā fere sex lineas longā ejus labium superius non ut in omnibus gregis integrum vel retusum est sed profunde bifidum.

18. S. Persica (Gmel jun. ex C. A. Mey. Enum. p. 94) perennis, caule adpresso tomentello simplici vel ramoso, foliis membranaceis crenatis petiolatis ovatis cordatis supra viridibus subtus canescentibus vel canis, floralibus oblongo-lanceolatis inferioribus flores superantibus, verticillastris inferioribus distantibus superioribus congestis, bracteis lanceolatis calyce subbrevioribus, calyce hirsuto bilabiato dentibus triangulari-lanceolatis spinosis labii superioris ternis altius connatis, corollæ roseæ extus hirsutæ labio superiore integro ♀. *S. sericea* Ledeb. Fl Ross. 3, p. 412 non Wall.

Hab. in Persiaæ prov. Ghilan (Gmel.), in sylvis montis Djulfekkuh (Aucn. 5174!), ad jugum Schimlar (Buhse 803!), prope Asterabad (Karel. exs. sub *S. rosea*!), prope Siaret (Bunge!).

Bipedalis et elatior, habitu, foliis discoloribus membranaceis similis *S. spectabilis* a quā tamen egregie differt calycis distincte bilabiati dentibus superioribus brevioribus, inferioribus binis profundius fissis. *S. sericea* Wall. quācum in Bth. Lab. p. 537 conjungitur est species diversissima *S. setifera* affinis.

§ 2. Micranthæ.

19. S. viticina (Boiss. Diagn. Ser. I, 12, p. 77) perennis adpresso et breviter tomentosa dense canescens, caulibus erectis elatis ramosissimis pyramidatim paniculatis, foliis crenulatis subrugosis inferioribus breviter petiolatis a basi truncato-subcordata oblongis obtusis, floralibus parvis inferioribus flores æquantibus, verticillastris parvis numerosis tomentosis inferioribus distinctis supremis confluentibus, bracteis lanatis oblongis calyce subbrevioribus, floribus parvis, calycis lanati dentibus triangulari-ovatis acutis muticis tubo subdupo brevioribus, corolle roseæ limbo hirsuto labio superiore integro ♀.

Hab. in dumosis sæpe humidis Syriæ borealis (Auch. 528! et 4712!), prope Antiochiam, Suadich, Seleuciam (Boiss!), circa Sidonem (Gailli!), Tripolin et Libanum circa Eden (Bl!).

Planta tripedalis, folia inferiora 2-2½-pollices longa, racemi semipedales, calyces 2 lineas longi. Species floribus minutis, bracteis oblongis et calycis dentibus muticis insignis inflorescentia *Viticem agnum castum* referens.

20. S. setifera (C.A. Mey. Enum. p. 94) perennis, molliter pilosa pallide virens vel canescens, caulibus erectis elatis flexuosis ramosis, verticillastris 6-8-floris omnibus distantibus longe racemosis, foliis membranaceis oblongis ascendenti-denticulatis apice spinulosis infe-

rioribus subsessilibus, floralibus sessilibus gradatim minoribus verticillastra superantibus, bracteis lanceolatis aristatis, floribus parvis, calycis patule hirti recti tandem accreti dentibus triangularibus aristulâ subulatâ terminatis tandem subrecurvis, corollæ glabrescentis carneæ vel sordide rubræ labio superiore subretuso ♀. *S. lycopsiformis* C. Koch Linn. XXII. p. 692!

Hab. in dumosis humidis et ad fossas et rivulos, Lycia ad Elmalm (Bourg. 220!), Cappadocia ad Caesaream (Wied!), Armenia Turcica ad Baibut (Bourg!) et Erzerum (Calv!), ditio Transcaucasica Talysch (C. A. Mey! Hoh!), Persia borealis in monte Elbrus (Buhset) et prope Derbend (Ky. 403!), in prov. Khorassan (Bge!), prope Kermanschah et Hamadan 5000' (Haussk!), in montibus Sawers, Eschker 8000' (Haussk!), Kuh Daéna et prope Schiraz (Ky. 572! Auch. 5177!), regno Cabulico (Griff!).

21. *S. longespicata* (Boiss. et Ky. exs. 1859) perennis molliter tomentosa canescens, caulis erectis elatis flexuosis ramosissimis, verticillastris præter inferiora 1-2-remota dense congestis spicatis, foliis ascendentis-denticulatis oblongis inferioribus petiolatis cordatis, floralibus triangulari-lanceolatis integris inferioribus flores superantibus, bracteis subulatis, floribus parvis, calycis albo-hispidissimi recti dentibus a basi lanceolata longe spinuloso-attenuatis tubo vix brevioribus, corollæ roseæ extus hirtæ labio superiore retuso ♀.

Hab. in dumosis secus fluvium Aksu prope Marasch Cataonie (Haussk!), inter Noreb et Angag prov. Boglan Armeniæ australis (Ky. 429!), in Cœlestiâ inter Zaclre et Balbeck (Ehrenb!).

Bi-tripedalis, flores parvi et inflorescentia in spicas condensatas *S. viticina* dentibus brevibus muticis calycis distinctissime; a *S. setiferâ* foliis cordatis muticis petiolatis, verticillastris non distantibus, calycis dentibus rectis, etc., aliena.

22. *S. rosea* (Desf. Coroll. Tourn. pag. 20, tab. 13, sub *Sideritide*) patule hispidula siccitate nigrescens, caulis erectis vel ascendentibus abbreviatis ramosis virgatis, foliis sessilibus elliptico-oblongis obtusis remote crenatis erenis ascendentibus basi subcordatis, floribus axillaribus subsessilibus solitariis oppositis basi bracteis binis lanceolatis acutis calyci æquilongis suffultis, calyce campanulato minute glanduloso et patule hirsuto dentibus æqualibus triangulari-lanceolatis acutis tubo subdupo brevioribus, corollæ pallide roseæ hirsutæ tubo inclusa, limbi minuti labio superiore breviore subretuso inferioris lobo intermedio vix majore ♀. *Sideritis Aegyptiaca* Pers. Syn. II, p. 117.

Hab. non in Egypto ut erroneè Fontanesius sed in Oriente? loco non indicato (Dr. Granger in herb. Juss. sub *Sideritis Orientalis scordioides flore purpureo*). Olim in Hort. Par. culta.

Rami 6-12-pollicares, facies *Teucrii Scordium*, folia pollicem longa omnia ad axillas florifera, calyx 4 lineas corolla $5\frac{1}{2}$ longa tuho piloso-annulato. Stanina quatuor inter se conformia e fauce subexserta omnino *Stachydis* nec *Sideritidis*. Pars inferior plantæ non nota et ideo affinitas dubia, tamen ob bracteas magnas inter *Eriostachydes* prope *S. setiferam* militare videtur quamvis a cæteris sectionis verticillastris bifloris distincta sit.

SECT. II. STACHYOTYHUS.

§ 3. *Sylvaticæ.*

23. **S. sylvatica** (L. sp. 811) perennis parce hirsuta viridis, caulinis e rhizomate repenti erectis superne saepius ramosis et glanduloso-pilosis, foliis membranaceis petiolatis ovatis cordatis acuminateis crenatis vel serratis, verticillastris 2-6-floris foliis floralibus lanceolatis longe cuspidatis calyces superantibus suffultis distinctis interrupte spicatis, calycis glandulosi dentibus æqualibus a basi triangulari subulatis tandem divaricatis, corollæ brunneo-purpureæ tubo exerto labio superiore breviter papilloso ♀. Ic. Engl. bot. tab. 416. — *S. Trapezuntea* Boiss. Diagn. Ser. II. 4, p. 38.

Hab. in sylvaticis humidis, Laconia et Thracia ad Byzantium (Sibth.), Anatolia borealis ad Mersiwan (Wied!), Pontus ad Cerasuntem (Tchih!), Pontus Lazicus ad Trapzuntem (Calvi!) et Rhizé (Bal!), Tauria (Stev.), Caucasus (M B., Owerin!), Transcaucasia (Hoh., C.A. Mey.).

Ar. Geogr. Europa fere tota a Scandinaviâ ad Rossiam septentrionalem, Sibiria Uralensis.

24. **S. hydrophila** (Boiss. Diagn. Ser. I, 12, p. 81) perennis patule et tenuiter hispida viridis, caulinis teneris elongatis a basi decumbenti ascendentibus ramosis, foliis tenuiter membranaceis amplis petiolatis cordatis ovatis aculiuseculis obtuse crenatis, verticillastris 4-6-floris foliis ovato-oblongis subrhomboides ea superantibus vel aequantibus suffultis distinctis longe et laxe racemosis, calycis viridis hirti visciduli breviter campanulati dentibus triangulari-lanceolatis spinulosis tubo subbrevioribus, corollæ albido-carnææ tubo inclusa labio inferiore ampio ♀.

Hab. in umbrosis ad rivulos et sepes circa Tripolim Syriæ (Boiss! Bl!), ad Eden Libani (Bl!), circa Sidonem (Gail!).

Planta 2-3-pedalis, folia inferiora saepe 3-4-pollices longa 3 lata. Affinis *S. sylvatica* a qua egregie differt foliis non acuminateis, floralibus non angustis longeque subulato-cuspidatis, calycis dentibus non divaricato-subrecurvatis. Corolla insuper coloris diversi, tubo non exerto, labio inferiore majore lobo intermedio lato

25. **S. palustris** (L. sp. 811) perennis, caulinis a rhizomate repente simplicibus vel breviter ramulosis ad angulos retrorsum setulosis, foliis breviter pubescentibus viridibus interdum canescensibus a basi cordatâ lanceolatis acutis crenato-serratis inferioribus breviter petiolatis cæteris sessilibus, floralibus amplexicaulibus flores multum superantibus, verticillastris 6-8-floris superioribus approximatis, calycis glanduloso-villosi subbilabiati dentibus a basi lanceolatâ longe subulato-spinosis tubo vix brevioribus, corollâ purpureâ parce puberulâ calyce duplo longiore tubo non exerto labio inferiore elongato ♀. Ic. Engl. bot. tab. 24.

Hab. in paludosis et ad ripas, Bæotia ad Lebadeam (Orph!), Macedonia (Friv.), Byzantii (Sibth.), Anatolia borealis (Wied!), Armenia Turcica (Tchih!), Tauria et Caucasus (M.B.), Georgia Caucasica (Hoh.)

Ar. Geogr. Europa fere omnis a Scandinaviâ ad Rossiam, Sibiria, China, America borealis.

26. *S. viridis* (Boiss. et Heldr. Diagn. Ser. II, 4, p. 39) perennis, *viridis glandulosa* et setis patentibus sparsis hispida, caulis elatis paniculatis, foliis sessilibus oblongo-lanceolatis late et obtuse crenulatis inferioribus obtusis superioribus acutis, floralibus a basi latiore acuminate inferioribus calyces superantibus, verticillastris 6-8-floris in spicas elongatas congestis inferioribus subdistantibus, calycis campanulati glandulosi et sparsim setulosi dentibus lanceolatis longe spinuloso-attenuatis subæqualibus tubo æquilongis, corollæ carneæ purpureo-suffusæ tubo inclusa labio superiore extus piloso, inferiore longiore ♀.

Hab. in alveis torrentium montis Olympi Thessali supra Metochi et montis Korthiati supra Thessalonicam 2000' (Heldr!).

Bipedalis, flores magnitudinis *S. spinulose* et *S. menthaefolue* Vis, quasi habitu refert et que calycis bilabiati dentibus superioribus alte connatis, corollâ ochroleucâ, etc. longe differt.

27. *S. menthoides* (Ky. et Boiss. in Ky exs. 1859) *viridis*, caulis erectis simplicibus stricte foliosis glabris, foliis subsessilibus ad petiolorum insertionem margineque ciliolatis cæterum glabris oblongis acutis acutiusculæ et ascendentí-serrulatis, floralibus diminutis inferioribus flores superantibus, verticillastris 6-4 floris in racemos breves basi interdum subcompositos approximatis, floribus minutis, bracteolis subulatis calyce plus duplo brevioribus, calycis minute glandulosi dentibus lanceolatis spinulâ terminatis denum subrecurvis tubo subdupo brevioribus, corollæ parvæ tubo inclusa labio superiore breviter hirto ♀.

Hab. in graminosis fruticosis Armeniæ Turcicæ australis prope Gumgum ad pagum Gestemert 5200' (Ky. Suppl. 680!). Fl Aug.

Pedalis, folia pollicem longa, racemus 1-2-pollicaris. Affinis *S. viridi* differt glabritate, floribus duplo minoribus; calyx $2\frac{1}{2}$ lineas longus dentibus abruptius spinulosis et subrecurvis. Corolla eâ *S. menthaefolue* et *viridis* triplo minor coloris e sicco mihi incerti.

28. *S. mollissima* (Willd Hort. Berol. p. 60, tab. 60) perennis canescens molliter villosa, caulis ascendentibus humilibus fragilibus simplicibus aut inferne parce ramosis foliosis. foliis ovatis cordatis petiolatis obtusis obtuse crenatis. verticillastris plurifloris in capitulum ovatum foliis summis sessilibus calyce sublongioribus involucratum dense congestis, bracteis late linearí-spathulatis acutis calyci subæquilongis, calycis *viridis* villosi campanulati dentibus late lanceolatis spinulosis subæquilatis. corollæ pallide ochroleucæ extus hirtæ tubo calycem non excedente ♀. *S. decumbens* Pers. Syn. 2, p. 123, Willd. En. Suppl. (Vide Aschers. Bot. Zeit. 1872, p. 631).

Hab. in rupestribus insulae Coreyrae (herb. Willd! Dr Schrader ex Aschers!).

Planta ut videtur humilis $\frac{1}{2}$ -1-pedalis, foliis majoribus pollicem longis, capitulo pollicari. Facie affinis *S. circinnata* ex Hispania quâcum prob. confusa ex Hispania erroneè indicata fait. Species bracteis valde evolutis a sectionis charactere divergens, sed huic tamen facie et aliis notis adnumeranda nec inter *Eriostachydes* inserenda.

29 S. Messeniaca (Boiss. Diagn. Ser. I, 12 p. 81) perennis, viridis patule et molliter hispidula basi indurata et ramoso-pluricaulis, caulis a basi ascendentí erectis foliosis simplicibus vel breviter 1-2-ramnlosis. foliis ovato-oblongis cordatis obtusis obtuse crenatis inferioribus breviter petiolatis cæteris sessilibus, floralibus inferioribus flores subsuperantibus, verticillastris 6-4-floris in racemum brevem approximatis inferioribus distinctis, bracteis paucis minimis setaceis, calycis viridis hirti visciduli obscure bilabiati dentibus triangulari-lanceolatis breviter spinulosis inferioribus binis vix angustioribus. corollæ pallide ochroleucæ purpureo-maculatæ labio superiore glanduloso-hirto inferiore ampio 2-3-plo longiore tubo subexerto ♂. *S. Tournefortii* Bory et Ch. Fl. Pélop № 913 non Poiret.

Hab. in rupestribus Messeniae circa Pylos, Androuvista et ad Phygaleam (Guérin Mill!).

Facies *Teucrī Scorodonīce*, pedalis vel humilior, folia majora pollicem longa 7-9-lineas lata, racemus 3-4-pollicaris, calyx 4 lineas longus. Aflnis *S. mollissimæ* a quâ foliis sublongioribus sed præsertim verticillastris longe racemosis et bracteis setaceis brevissimis differt.

30. S. canescens (Bory et Ch. Fl. Pélop. № 916 sub 19) perennis tomentoso-canescens, caulis erectis, foliis petiolatis oblongo-lanceolatis acutis superioribus sessilibus, floralibus diminutis calyces superantibus et æquantibus, verticillastris subsexfloris in racemum brevem approximatis, calycis dentibus lanceolatis spinulosis rectis tubo æquilongis, corollæ albidae lobio superiore hirsnto, inferiore longiore tubo subexerto ♂.

Hab. in Peloponeso ad Tsimova et Vitylos, ad prom. Tenarum (Bory) Non vidi.

Ex iconē affinis *S. Messeniacea* a quâ differre videtur indumento tomentoso-canescente, foliis basi rotundatis, calycis dentibus tubi proportione longioribus. Specimina ad Lutraki Sinū Corinthiaci lecta dubitanter ab am. Heldreich hic referuntur sed foliis brevibus obtusiusculis nec apice attenuatis acutis ab iconē differunt et ad *S. Swainsoni* var. *Seyronicam* referenda sunt.

§ 4. Fragiles.

31. S. Pinardi (Boiss. Diagn. Ser. I, V, p. 29) brevissime papilloso-glandulosa et setis patulis pallucidis hispidula, rhizomate indurato ramoso, caulis humilius fragilis simplicibus, foliis pallide virentibus concoloribus longe petiolatis profunde cordatis ovatis obtusis crenatis, floralibus sessilibus a basi cuneatâ

rotundatis dentatis acutiusculis inferioribus flores æquantibus, verticillastris 6-4-floris distinctis in racemum terminalem breve in approximatis, bracteis nullis, calycis sparsim setosi campanulati dentibus subæqualibus triangularibus spinuloso-attenuatis tubo triplo brevioribus, corollæ albœ tubo calyce duplo longiore, labio superiore hirtulo, inferiore longiore ♀.

Hab. in rupestribus Cariæ littoralis (Pinard!).

Caules incurvi cum spicâ brevi 5-6-pollicares, folia 1 $\frac{1}{2}$ -2-pollicies longa et lata, calyx 3 lineas, corolla tubus 5 $\frac{1}{2}$ lineas longus. Facies et fragilitas *S. circinnatæ* L'her. indumento sericeo, foliis floralibus petiolatis, corollæ roseæ tubo subincluso, etc. distinctæ.

32. *S. longiflora* (Boiss. et Bal. Diagn. Ser. I 4 p. 39) minute glandulosa patuleque et molliter hispida, caulis fragillimus elongatis dependentibus, foliis petiotis eis æquilongis suffultis ovato-orbiculatis profunde cordatis circumcirent acute et grosse triangulari-dentatis, floralibus a basi petiolatæ cuneatæ ovato-flabellatis 3-5-cuspidato-dentatis calyces superantibus, verticillastris bifloris racemos breves laxiusculos terminalem et sœpe laterales axillares formantibus, calycis visciduli dentibus lanceolatis longe spinulosis tubum dimidium æquantibus, corolle viscidulae purpurascens tubo longissimo calycem ter superante labio superiore retuso inferiore duplo longiore ♀.

Hab. ex rupibus verticalibus speluncarum faucis Guzelderé supra Sedichig Cilicie littoralis pendula (Bal.).

Ex speciminibus autumnalibus rursum florentibus descripta, caules tennes pedales, folia majora diametro 1-1 $\frac{1}{2}$ -pollicaria, raceini sesquipollicares, calyx 4 $\frac{1}{2}$ lineas, corolla 15-16 longa. Affinis *S. circinnatæ* et præsertim *S. Pinardi* a quâ foliis acute et grosse dentatis, corollæ tubo longissimo egregie differt.

§ 5. Rectæ.

33. *S. recta* (L. Mant. 22) perennis hirsuta viridis, caulis ascendentibus elongatis simplicibus vel striete ramosis, foliis oblongis vel oblongo-lanceolatis crenatis inferioribus petiolatis superioribus sessilibus, floralibus ovato-oblongis integris aristulatis, verticillastris 4-6-floris præter superiora distinctis longe et interrupte spicatis, bracteolis setaceis minimis, calycis hirti ovato-campanulati dentibus triangularibus longe in spinulam glabram mucronato-aristatis tubo subduplo brevioribus, corollæ ochroleucæ vel albidæ fauce purpurascens pubescens tubo inclusa, staminibus labium superius subæquantibus ♀. Ic. Jacq. Austr. tab. 359. — *S. Sideritis* Vill. — *S. procumbens* Lam. Synonyma *S. annua* et *S. glutinosa* Smith Prodr. a. cl. Bth. citata ex eo Lab. p. 556 dubia sunt.

Hab. in declivibus herbidis Thraciae borealis prope Carlova (Janka!), Tauria et Caucaso (M.B., Ledeb.). Loc. Læonia (Sibth.) et Creta Sibth ex Bth. mihi dubii.

β. sideritoides. — Hirsutior, folia lanceolata et lanceolato-linearia acuta argutins denticulata rarius subintegra, calycis magis hirsuti dentes tubo subæquilongi. — *S. sideritoides*, *S. atherocalyx* et *S. linearifolia* C. Koch Linn. XXI, p. 691 et 692. — *S. patula* Griseb. Spic. II, p. 142!

Hab. in Macedonia (Griseb., Friv. exs. sub *S. recta angustifolia*). Armenia prope Erzerum (Tchih!), Caucaso (C. Koch), Daghestan 4800'-7800' (Rupr!).

γ. fragilis. — Calycibus exceptis glabrescens, folia subintegra vel integra lanceolata et ramealia linearia. — *S. fragilis* Vis. Fl. Dalm. tab. 16, fig. I.

Hab. circa Byzantium (Coumany).

δ. leucoglossa. — Facies præcedentis, hirtula, folia anguste lanceolata inferiora serrata superiora et ramealia integra linearia. Calycis papillis sparsis scabridi dentes tubo æquilongi. — *S. leucoglossa* Griseb. Spic. II, p. 140.

Hab. in Macedonia monte Athone (Auch. 1718! Griseb!).

Omnis formæ hic enumeratae ut et *S. subcrenata* Visiani inter se et cum typo intermediis evidenter confluent.

Ar. Geogr. Europa tota media et australis a Gallia et Germania ad Rossiam medium.

34. *S. Parolinii* (Visian. Ill. p. 10, tab. IV) perennis, caulis retrorsum hispidis virgatis inferne divaricatim ramulosis foliosis superne remote microphyllis, verticillastris paucis 6-4-floris in racemum terminalem ovatum basi sepe interruptum congestis, foliis viridibus hispidis inferioribus petiolatis oblongis obtusiusculis ascendenti-serrulatis superioribus remotis floralibusque anguste lanceolatis deflexis verticillastro longioribus, bracteis setaceis minimis, calycis breviter campanulati subbilabiati villosi dentibus lanceolatis longe spinulosis erecto-patulis tubo subbrevioribus, corollæ ochroleuce tubo vix exerto labio superiore breviore hirto ♀.

Hab. circa Lepanto (Naupactem) Etolie (Parol.), insulæ Cephallenia in monte Atro et inter Argostoli et Omalo (Heldr!).

Pedalis, folia inferiora 8-12 lineas tantum longa, caulina superiora valde dissita angusta, racemus 1-1½-pollicaris, calyx 3 lineas longus. Species valde affinis *S. recta* var. *leucoglossa* a qua tamen verticillastris in capitulum ovatum conglobatis calyceque viloso obscure bilabiato profundius fiso differre videtur.

35. *S. nepetæfolia* (Desf. Cat. Hort. Par. p. 58) perennis minute et adpressiuseule pubescens, caulis ascendentibus basi ramulosis, foliis petiolatis ovatis cordatis obtusis acutiusculæ crenato-dentatis, verticillastris distantibus subsexfloris, foliis floralibus valde diminutis sessilibus linear-lanceolatis subintegris apice inaequaliatis calyci subæquilongis deflexis, bracteis setaceis minimis, calycis adpresso canescens tubuloso-campanulati dentibus a basi triangulari spinulosis tubo triplo brevioribus, corollæ albide vel ochroleuce

adpresso pubescentis tubo subincurvo calyce sesquilongiore, labio superiore brevi subretuso, antheris non exsertis ♀. *Nepeta circinata* Willd. Enum. p. 602 et herb!

Hab. in Cappadociâ (ex Willd.).

Caules tenues cum racemo 1-1 $\frac{1}{2}$ -pedales, folia inferiora pollicaria cætera sensim diminuta, racemus laxus 6-8-pollicaris, calyx rectiusculus tres lineas longus. Species ob antheras e tubo non exsertas potius *Sideritidi* adnumeranda, sed perennis et facies *Stachydis*. Planta a collectoribus recentioribus non reperta ex Hispaniâ erroneâ et quâdam confusione indicata.

36. S. Iberica (M.B. Taur. Cauc. II, p. 54) perennis hirsuta viridis, caulis ascendensntibus elongatis virgatis simplicibus vel ramosis, foliis anguste oblongis basi longe attenuatis subpetiolatis superne obtuse serratis obtusiusculis, floralibus integris a basi latiore lanceolatis acutiusculis muticis inferioribus calyces superantibus, verticillastris 6-10-floris distinctis superioribus approximatis, bracteis subnullis, calycis hirsuti campanulati dentibus a basi triangulari-lanceolatâ longe mucronato-aristatis dimidio tubo æquiflora vel longioribus, corollæ purpureæ hirsutæ tubo exerto, staminibus e fauce vix subexsertis ♀. *S. congesta* Urv. Enum. p. 68. — *S. Caucasicâ* C. Koch Linn. XXI, p. 693.

Hab. in Thessalîæ monte Cœta supra Mustapha Bey (Sprun!), Ponto (Tchih!), Ponto Lazico ad Djimil (Bal!), Armeniâ ad Baibut (Huet!), in apricis Iberiæ (M.B., Hohen!).

β. pallidiflora — Calyces hirsuti vel glabrescentes, corolla carnea vel albido-ochroleuca. — *S. leucoglossa* var. *Anatolica* Boiss. Diagn. Ser. II, 4 p. 41.

Hab. in Anatoliâ boreali (Tchih! Wied!), Tauro Isaurico (Heldr!), Tauro Cilicio (Ky. 443! Bal!), Cappadociâ (Bal!), Armeniâ prope Baibut (Bourgl!), Tauriâ (M.B.).

γ. brachydonta. — Flores pallidi varietatis præcedentis sed calycis dentes triangulares brevius aristati. — *S. stenostachya* Boiss. Diagn. Ser. II, 4 p. 41.

Hab. in Syriae borealis monte Cassio (Boiss!), Amano (Ky!).

Ab omnibus *S. rectæ* formis quibus habitu, calyceque accedit antheris sub-inclusis nec labium superius subaequantibus, foliisque floralibus muticis nec aristatis egregie distincta. *S. arenaria* Vahl planta Sicula et boreali-Africana quâcum nostra sœpe confusa fuit differt foliis floralibus mucronatis, corollæ labio superiore longo valde bifido, antheris exsertis.

37. S. virgata (Bory et Chaub. Fl. Pélop. N° 911, tab. 18) perennis tota adpresso sericeo-lanata, caulis virgatis ciatis opposite et stricte ramosis sparsim foliosis, foliis integris linearibus inferioribus obtusiusculis basi sensim in petiolum attenuatis superioribus acutis, floralibus bracteisque calyce brevioribus, verticillastris bifloris remotis longe racemosis, floribus subsessilibus strictis, calycis tubuloso-campanulati dense sericei dentibus lanceolatis spinosis tubo quadruplo brevioribus, corollæ albae sericeæ tubo calyce breviore, staminibus exsertis ♀. *S. Zuccarinii* Bth. Lab. p. 535.

Hab. in Argolide ad Kranidi et Trezenam (Despréaux!), ad Acrocorinthum (Zuccar.).

Caules 2-3-pedales, folia inferiora pollicaria $2\frac{1}{2}$ lineas lata, calyx 5 lineas longus. Minime suffruticosa; indumento dense sericeo ad *S. spinosam* cæterum diversissimam accedens.

38. *S. angustifolia* (M.B. Taur. Cauc. II, p. 51) perennis glabra pluricaulis caulis tenuiter virgatis ascendentibus elongatis stricte ramosis, foliis sessilibus inferioribus anguste linear-lanceolatis in lacinias distantes lineares pinnatifidis partitisve. cæteris angustissime linearibus indivisis vel lacinulâ auctis, floralibus calyce brevioribus, verticillastris distinctis bifloris longe et anguste racemosis, floribus pedicellatis, calycis campanulati glabri vel parce hirtio-scabridi dentibus lanceolatis spinoso-attenuatis tubo subbrevioribus, corollæ purpurascens vel albidae rarius citrinæ adpresso pubescentis tubo breviter exerto, staminibus fauce vix longioribus ♀. *S. tenuifolia* Pall., Janka. — Rehb. Exot. tab. 28 non Willd. — *S. abrotanifolia* Boiss. et Heldr. olim.

Hab. in rupestribus Macedoniæ, mons Athos (Auch. 1716!), prope Bitolia et in monte Korthiati (Heldr! Orph. Fl. exs. 536!), Thraciæ prope Karlova (Janka!), Tauria (M.B., Rehm!).

Foliis inferioribus pinnatipartitis cæteris angustissimis, verticillastris bifloris, etc. distincta; suffruticosa erroneè dicitur.

Ar. Geogr. Rumelia, Rossia australis.

39. *S. distans* (Bth. in n.c. Prodr. XII, p. 472) perennis multicaulis caulis albo-lauatis virgatis flexuosis elatis ramicis a basi ascendentí erectis, foliis parvis remotiusculis utrinque adpresso canescensibus vel subtus canis inferioribus petiolatis cordatis ovatis obtusiusculæ crenato-dentatis superioribus sensim diuinutis sessilibus acutis, floralibus breviter oblongo-lanceolatis flores vix æquantibus, verticillastris multifloris valde distantibus vel 2-3 interdum in capitulum oblongum approximatis, bracteis lanceolatis calyce brevioribus, calycis nervosi breviter pubescenti-glandulosi dentibus lanceolatis longe spinuloso-attenuatis tandem divergenti-subpatentiis tubo æquilengis vel brevioribus, corollæ ochroleuce glanduloso-hirtæ tubo subexerto, limbo parvo ♀.

Hab. in Libano (Auch. 1753!), ad niuros pontis Nacher Beryti (Ehr! Ky. 463!), ad Khan Hussein Libani inferioris et circa Tripoli (Boiss!), ad Bezulmar (Bl!) et Sidonem (Gaill!).

β. oxyodonta. — Folia virescentia oblonga acutius dentata.

Hab. in ponte ad fluvium Jordanum infra Banias Galileæ (Boiss!).

γ. teucriifolia. — Rami parce et laxe sublanati interdum glabri, folia minora virentia, calycis dentes abbreviati triangulares acuti. — *S. teucriifolia* Boiss. et Ky. Diagn. Ser. II, 4, p. 40.

Hab. ad rupes regionis subalpinæ Libani, Eden (Ky. 587!), Bischerre (Ehr!) inter Diuan et Kanobin et ad Hasran (Bl!).

Sesqui vel tripedalis, folia inferiora 8-12 lineas longa, superiora remotissima et valde diminuta, calyx $2\frac{1}{2}$ lineas longus. Varietates enumeratae intermediis alia in alteram transeunt.

40. S. bombycina (Boiss. Diagn. Ser. I, 12, p. 79) perennis tota densissime albo-lanata, caulis numerosis a basi procumbenti erectis flexuosis elongatis simplicibus vel inferne ramosis, foliis parvis ovatis et oblongis obtusis crenato-dentatis inferioribus petiolatis superioribus panicis remotissimis diminutis integris acutiusculis, floralibus calyce brevioribus, verticillastris 1-3-multifloris distantibus, bracteis oblongis acuminatis calyce brevioribus, calycis dense et longe albo-lanati dentibus a basi lanceolatâ subulatis rectis tubo æquilongis, corollæ albidae limbo parvo hirtulo ♂.

Hab. in Syriâ loco non notato (Pestal.).

Pedalis et sesquipedalis, folia 6-7 lineas tantum longa, verticillastri nuce avellanâ submajores. Affinis *S. distanti*, an satis distincta sit ulterius explorandum; differt foliis dense albo-lanatis, calycibusque bombyce niveâ occultatis dentibus rectis nec subrecurvo-divergentibus.

41. S. aleurites Boiss. et Heldr. (Diagn. Ser. I, 12, p. 79) perennis ex rhizomate indurato pluricaulis, caulis elatis virgatis simplicibus vel basi ramosis inferne adpresso araneoso canis demum glabrescentibus, foliis paucis remotis petiolatis parce et adpressissime araneoso-canescensibus demum glabratris ovatis grosse et acutiuseule crenato-dentatis acutis inferioribus basi cordatis, superioribus remotissimis floralibusque lanceolatis acuminatis, verticillastris multifloris inferioribus præsertim valde distantibus foliis floralibus superatis superioribus approximatis, bracteis lanceolatis acuminatis calyce subbrevioribus, calycis araneosi cani dentibus a basi lanceolatâ longe subulatis rectis tubo subbrevioribus, corollæ e sicco ochroleucæ adpresso hirtæ tubo inclusa ♀.

Hab. ad rupes maritimæ ad Adalia Pamphyliæ (Heldr. Pestal.). Fl. Maio.

Caulis 2-3-pedales, folia eis ramorum sterilium *Lamii Galeobdolon* similia, majora sesquipolligaria, verticillastri 4-5, racemus 4-6-polligaris. Affinis *S. distanti* differt indumento adpresso araneoso, verticillastris plurifloris, calyce duplo majore $4\frac{1}{2}$ lin. longo, etc.

42. S. diversifolia (Boiss. Diagn. Ser. I, 12, p. 80) perennis basi suffruticosa ramosa prostrata, caulis hornotinis herbaceis virgatis glabris elongatis simplicibus vel superne parce et breviter ramulosis, foliis ramorum sterilium petiolis longis hirsutis insidentibus adpresso et parce setulosis viridibus a basi caneatâ ovato-oblongis obtusis mucronulatis integris vel apice obsolete crenulatis, caulinum floriferorum glabris sessilibus anguste lanceolato-linearibus acutis, verticillastris præter terminale sœpe 3-4-florum bifloris remotis, foliis floralibus brevissimis, bracteis linearis-setaceis calyce æquilongis, calycis glabri viridis visciduli dentibus lanceolatis punctatibus rectis tubo subbrevioribus, corollæ pallide carnea extus hirtæ tubo inclusa ♀.

Hab. in dumosis regionis montanæ in Syriâ inter Latakieh et Antiochiam sitæ (Boiss!). Fl. Junii fine.

Foliorum inferiorum limbus 10-12 lineas longus 6-8 latuſ, folia caulinæ 1-1½-pollicaria 2-3 lineas lata, caules floriferi sesquipedales in racemum laxè pauciflorum macrum abeuntes. Calyx florifer 2½ lineas longus. Ex *S. distantis* affinitate.

43. *S. subnuda* (Montb. et Auch. Ann. Sc. Nat. 2, VI, p. 50) perennis minutissime et parce puberula, caulis numerosis virgatis opposite et stricte ramosis, foliis parvis remotis infimis petiolatis ovatis superioribus oblongis subcrenatis, floralibus calyce brevioribus, verticillastris 6-2-floris valde remotis, bracteis subnullis, calycis viridis puberuli campanulati dentibus lanceolatis carinatis subpungentibus rectis tubo duplo brevioribus, corollæ ochroleucæ pubescentis tubo subexerto ♀.

Hab. in Cappadociâ ad Euphratem (Auch. 1702! et 1703!).

β. *Kurdica*. — Folia glabrescentia majora dentata, floralia inferioria flores interdum superantia. — *S. Kurdica* Boiss. et Hoh. Diagn. Ser. I, 5, p. 31.

Hab. in rupestribus montis Gara Kurdistaniæ (Ky. 390!), in montibus Avroman et Schahu 5-7000' (Haussk!).

Caules pedales et sesquipedales tetragoni juncei, calyces demum 4 lineas longi.

44. *S. viscosa* (Montb. et Auch. Ann. Sc. Nat. Ser. II, 6, p. 49) perennis, multicaulis viridis minute et plus minusve glandulosa prætereaque viscido-pubescentia. caulis herbaceis elongatis ascendentibus ramosis, foliis ovatis cordatis crenato-dentatis breviter petiolatis, floralibus subscissilibus triangularibus acutis calyces subsuperantibus, verticillastris 4-6-floris remotis, bracteis linearibus brevissimis, calycis viridis præter glandulas sessiles hirti dentibus lanceolatis longe subulatis spinescentibus carinatis dimidio tubo sublongioribus, corollæ flavidæ minute viscido-hirtulæ tubo inclusa ♀.

Hab. in Armeniae Turcicae rupestribus circa Gumuschkhané (Tchih! Bourg!), ad Euphratem superiorem (Auch. 1706!).

Caules graciles 1-1½-pedales, folia majora pollicaria, verticillastræ 1½-2-pollicies distantes, calyx 4 lineas longus.

γ. *elatior*. — Minus glandulosa, glabrescens, foliæ acutiora acutius dentata, verticillastræ superiores minus distantes, tota planta major. — *S. laetevirens* Ky. et Boiss. MSS.

Hab. in præruptis argillosis vallis Goschkar prov. Musch Armeniæ australis 6000' (Ky. 355!).

45. *S. Benthamiana* (Boiss. herb!) perennis e rhizomate indurato multicaulis patule hispidula viridis superne glandulos-viscidula, caulis tenuiter virgatis flexuosis ramosis, foliis parvis breviter petiolatis ovato-oblongis dentatis basi cordatis vel truncatis,

floralibus verticillastra superantibus vel æquantibus, verticillastris paucis 6-4-floris distantibus supremis interdum approximatis, bracteis subulatis brevissimis, calycis campanulati glandulosi dentibus lanceolatis acutis tubo duplo brevioribus, corollæ flavidæ viscido-hirtulæ tubo subexerto ♀. *S. viscosa* var. *Kotschyi* Bth. in D.C. Prodr. XII, p. 483.

Hab. ad rupeſ abſcissaſ Persie australis (Olivier! Auch. 2889!), prope Schiras et in monte Kuh Delu (Ky. 345! et 490!), ad ruinas Perſepolis et in monte Kuh Eschker (Haussk!).

β . *clinopodioides*. — Humilior semipedalis vix glandulosa sed tota patule vel retrorsum velutina canescens, verticillaſtri 1 vel 2 inferioribus interdum exceptis in capitulum ovatum approximati, calycis velutino-hispidi denteſ interdum acutiores sed mutici. — *S. ballotiformis* et *S. plebeia* Vatke Bot. Zeit. 1875, p. 448.

Hab. in rupiſ calcareiſ montium Schahu, Avroman, Pir Omar Gudrun Kurdistaniæ 6-8000' (Oliv! Haussk!). Fl. Jun.

Species a *S. subnudā* foliis cordatis, dentibus calycinis non carinatis nec pungentibus diversa, a cl. Bth. *S. viscosa* adnumerata quæ tamen calycis dentibus longe et anguste subulato-attenuatis differt.

§ 6. Fruticosæ.

46. **S. Swainsoni** (Bth. Lab. p. 535) tota adpreſſe et molliter cinerascenti-cinerea basi ſuffruticosa ramosiſſima cespitosa ramis pumiliſ herbaceiſ, foliis parviſ rugulosis ovatiſ obtuse crenatiſ obtuſiſſimiſ inferioribus in petiolum brevem ſtenuatiſ cæteriſ ſeſſiliſ cuneato-rhombeiſ acutiſ calyces ſuperantibus et æquantibus, verticillaſtri 4-2-floriſ in racemoſ oblongoſ interdum laxiusculoſ approximati, calyciſ tomentelli obscure bilabiati dentibus lanceolatiſ spinulosis tubo ſubduſpo brevioribuſ, corollæ carneæ tubo ſubincluso labio ſuperiore hirsuto inferiore duſpo longiore ♀. Ic. Visiani Ill. tab. 3 — *Prasium hirsutum* Poir. Dict. 5, 611 ex parte.

Hab. in fissuriſ rupiū Parnassi ſupra Rachova et Delphos 4-5000' (Swains., Sprun! Heldr. Herb. Norm. 901 Guicc! Orph. Fl. Gr. exs. 436!).

β . *Scyronica* — Magis sericea, rami abbreviati, folia minora. — *S. Scyronica* Boiss. Diagn. Ser. I, V, p. 31.

Hab. ad rupeſ Isthmi Corinthiaci ad Kakiscaſa et Lutraki et ad Scyronides petras (Boiss! Heldr! Orph. Fl. Gr. exs. 438!), insula Cephallenia (Heldr!), Ithaca (Webb).

γ . *Argolica*. — Viſescens, indumentum parcus minus adpreſſum, folia rugosiora. — *S. Argolica* Boiss. Diagn. Ser. I, 12, p. 83.

Hab. in rupiſ Peloponnesi ad Naupliam et Port Tolon (Sprun! Boiss! Heldr! Orph!), in insula Hydra (Heldr!).

Rami 4-6-pollicares, folia 5-10 lineaſ longa, calyx 4 lineaſ longus. Species ramis abbreviatis, foliis parviſ non cordatiſ ſæpius rugosiſ, corollæ carneæ, etc. a *S. molliſſimā* diſtincta, mediante var. β cuius indumentum sericeum ad *Ambicias* accedens.

47. S. Spruneri (Boiss. Diagn. Ser. I, 12, p. 82) viridis sparsim et patule hispida basi suffruticosa ramosa, ramis pumilis rigidis acute quadrangulis simplicibus vel parce ramulosis, foliis rigidis valde rugosis argute serratis infimis oblongo-ellipticis obtusis in petiolum brevissimum attenuatis cæteris sessilibus linear-lanceolatis acutis, floralibus subdilatatis oblongis acuminatis integris flores superantibus, verticillastris 6-4-floris in spicas ovatas congestis inferiore sèpe remotiusculo, bracteis nullis, calycis hirsuti viridis sublabiati dentibus lanceolatis rectis longe spinulosis tubo dimidio brevioribus, corollæ albidae tubo inclusa labio superiore longe hispido, inferiore longiore ♂.

Hab. in rupibus ad radices montis Parnes Atticæ supra Minidi et Metochi (Sprun! Boiss! Heldr! Orph. Fl. Gr. exs. 439!) alt. 500'-1500', monte Pateras (Heldr!), monte Kerata prope Eleusin (Heldr!), regione media montis Onion Bœotiae (Heldr!). Fl. Apr., Mai.

Rami semipedales, folia intermedia 1-2-pollices longa 3-4 lineas lata, calyx 3 lineas longus. A *S. Swainsoni* var. *Argolicâ* cui affinis distincta foliis rigidis argute serratis angustis, floralibus valde dilatatis acuminatis.

48. S. tetragona (Boiss. et Heldr. pl. exs. 1858) perennis ex rhizomate indurato ramosissima ramis pumilis viridibus parce setulosis tenuiter virgatis flexuosis tetragonis et sulcatis parce foliosis, foliis ramulorum basilarium sterili congestis oblongo-spathulatis ellipticis obtusis prope apicem obtuse crenatis in petiolum brevem attenuatis adpresso-circumflexuosis hirtis, superioribus et floralibus minimis squamiformibus oblongo-linearibus, verticillastris bifloris 2-6-remotissimis, calycis parce setulosi rigidi campanulati dentibus æqualibus triangularibus breviter spinulosis tubo triplo brevioribus, corollæ ochroleucæ tubo inclusa labio superiore hirto inferiore subduplo longiore ♀.

Hab. ad rupes regionis inferioris montis Delphi Eubeæ supra Steni (Heldr!).

Rami tenues intricati 3-6-pollicares subnudi, folia ad surculos steriles 3-5 lineas longa. rarius cum petiolo pollicaria, calyx vix tres lineas longus. Facies quodammodo *S. glutinosa* ramis oppositis spinescentibus, etc., distinctissimæ. Calyce affinis *S. angustifoliae*.

49. S. spinosa (L. Syst. Veg. 3 p. 58) suffruticosa humilis ramosissima dumuloso-erinacea tota adpresso sericeo-lanata, ramis rigidis opposite et plures ramulosis, ramulis apice spinosis sericeis, foliis sessilibus linearibus integerimis, verticillastris 1-2 sub spinâ terminali bifloris vel abortu unifloris foliis floralibus oblongis acutis calyce brevioribus suffultis, calycis sessilis sericei tubuloso campanulati dentibus triangulari-lanceolatis spinosis tubo triplo brevioribus, corollæ pilosæ albidae tubo inclusa 5. Ic. Fl. Græc. tab. 559.

Hab. in collibus aridis regionis inferioris et montanae Creta 0-1500' Amparo, Messara, Malevisi, etc. (Tourn., Sibth., Sieb! Raul! Heldr!). Fl. Mai., Jun.

Dumus semipedalis vel pedalis, folia pollicaria 2-3 lineas lata, spine terminales 1/2-1-pollicares, calyx 3 lineas longus.

Obs. *S. glutinosa* Linn. a Willd. et Smith Prodr. in Cretâ indicata ibi non

crescit, specimen herb. Sibth. ex cl. Bth. est *S. recta* vel *Iberica*; specimen herb. Billardierani ex Corsicâ nec ex Oriente oriundum est, cl. collector schedules Orientis et Corsicæ (v. gr. *Helich. frigidum*) sœpe commiscerit.

50. *S. mucronata* (Sieb. Cret. exs. — Spr. Syst. II, 733) suffruticosa ramosissima nana longe et laxe lanata pallide virens, ramis crassiusculis rigidis opposite ramulosis, foliis rugosis crenulatis oblongis in petiolum attenuatis, floralibus minutis sessilibus acutis calyce brevioribus, verticillastris 2-3 bi vel quadrifloris approximatis, floribus stricte et crassiuscule pedicellatis, calycis tubuloso-campanulati coriaceti sulcato-costati hirsuti virentis dentibus acerosis duris subdivergentibus tubo subbrevioribus, corollæ villosissimæ e sicco carneæ tubo inclusa ζ .

Hab. in saxosis regionis inferioris Cretæ, Capo Sidero, inter Palæokastron et Zakro (Sieb! Heldr!). Fl. Maio.

Planta 3-pollicaris ramis plus minus persistentibus sed minime spinescentibus, folia pollicaria 4-5 lineas lata, calyces 5 lineas longi.

51. *S. acerosa* (Boiss. Diagn. Ser. I, 7, 57) suffruticosa ramosissima humilis late erinacea viridis pruinosa parce hirtula, ramis rigidis stricte et opposite ramulosis apice tenuiter spinescentibus, foliis parvis sessilibus basi angustatis integris inferioribus oblongis muticis cæteris oblongo-linearibus apice spinulosis, floralibus flores æquantibus vel brevioribus, verticillastris 2-3 subdistantibus bifloris, floribus longiuscule pedicellatis strictis, calycis pruinosi visciduli tubuloso-campanulati dentibus lanceolatis spinosis tubo subæquilongis, corollæ albæ puberulæ tubo inclusa ζ .

Hab. in apricis regionis alpinæ Persiæ austro-occidentalis, mons Sabst Buschom prope Schiras (Ky. 418!), montes Kuh Nur, Sawers, Eschker, Maregun 10-12000' (Haussk!), montes Bachtarici (Bode!).

Semipedalis vel pedalis, folia majora 8-10 lineas longa 2-3 lata, calyx 5 lineas longus.

52. *S. fruticulosa* (M.B. Taur. Cauc. II, p. 51) fruticulosa ramosissima humilis viridis pilis sparsis brevibus crispulis obsita, ramis subsimplicibus, foliis parvis subsessilibus basi attenuatis oblongo-lanceolatis vel linearibus integris vel parce ascendentí-denticulatis inferioribus obtusis, floralibus linearibus acutis calyces superantibus et æquantibus, verticillastris paucis bifloris remotis laxe racemosis, floribus breviter pedicellatis, calycis tubuloso-campanulati breviter hirtuli dentibus a basi triangulari-lanceolatâ acutissimis subspinosis tubo subduplo brevioribus, corollæ pallide purpureæ adpresso pubescens tubo inclusa ζ .

Hab. in lapidosis montanis Iberiæ Caucasicæ et Somchetiæ (M.B., C.A. Mey!, Hoh!), prov. Aderbidjan Persiæ (Auch. 5182! Szov! Bge! Buhse!).

$\beta.$ *macrocheilos*. — Corollæ allidæ labium superius longius angustius. — *S. macrocheilos* Boiss. Diagn. 5 p. 30.

Hab. in collibus aridis prov. Aderbidjan Persiæ (Auch. 5185!).

Semipedalis, folia 6-10 lin. longa 2- $\frac{1}{2}$ lata, calyx 3 lineas longus.

53. S. multicaulis (Bth. in D.C. Prodr. XII, p. 486) fruticulosa ramosissima bimilis viridis sparsim pilosula ramis erectis tenuibus simplicibus rarius opposite ramulosis, foliis parvis crassiusculis sessilibus basi subattenuatis lanceolato-linearibus acutiusculis integrermis, floralibus brevissimis, verticillastris bifloris distinctis laxe spicatis, floribus breviter pedicellatis, calycis tubuloso-campanulati hirti dentibus lanceolatis carinatis nervosis tubo subbrevioribus, corollæ flavæ hirsutæ tubo inclusæ ♂.

Hab. in Persiâ loco non notato (Auch. 1809!).

β. brachyodonta. — Indumentum brevius pruinatum, calycis dentes breviores diimidium tubum æquantes.

Hab. in fissuris calcareis regionis alpinæ montium Dalechani prope Sungur et Parrow prope Kermanchah Persiae occidentalis 9-10000' (Haussk!).

Dumi 6-10-pollicares, folia 6-8 lineas longa 2 lata, calyx 4 lineas longus. Affinis *S. fruticulosæ*, rigidior, differt foliis integris acutis, calyce nervosiore, corollæ flavæ nec purpureæ labiis minoribus.

54. S. Aucheri (Bth. in D.C. Prodr. XII, p. 487) suffruticosa ramosissima tota glanduloso-viscida prætereaque plus minusve hispidula viridis, ramis erectis tenuibus rigidis simplicibus et ramosis apice subspinescentibus, foliis parvis sessilibus integris breviter oblongo-lanceolatis apice spinulosis, floralibus flores subsuperantibus, verticillastris pluribus distantibus bifloris, floribus brevissime pedicellatis, bracteolis subnullis, calycis glandulosi longeque pilosi late campanulati dentibus late lanceolatis longe spinosis elevatim uninerviis divergentibus tubo sublongioribus, corollâ albida pilosâ calycem parum superante ♂.

Hab. in Persiâ (Auch. 1807!), ad rupes calcareas montium Eschker, Sawers, Nur 8-10000' (Haussk!).

Semipedalis et pedalis. folia 10-12 lineas longa 3 lata, calyx 5 lineas longus.

55. S. ixodes (Boiss. et Haussk. in Haussk. Sched. 1868) suffruticosa ramosissima patule villosa viscosa, ramis erectis tenuibus simplicibus vel stricte ramulosis, foliis sessilibus integris linearilanceolatis basi attenuatis acutis mucronulatis, floralibus flores æquantes, verticillastris pluribus distinctis bifloris, bracteolis subulatis calyce duplo brevioribus interdum nullis, calycis patule hirsutissimi et viscidæ dentibus lanceolatis tenuiter spinosis divaricatis tubo longioribus, corollæ roseæ hirsutæ tubo calycis dentibus non longiore ♂.

Hab. ad rupes calcareas montis Teng Nalli Persiae australis 9000' et inter Dilegun et Maregun (Haussk!).

Valde affinis *S. Aucheri*, longius et densius pilosa, rami non spinescentes, flores saepius bracteolati, calycisque 6-7 lineas longi dentes angustiores longiores et valde divaricati.

56. S. pilifera (Bth. in D.C. Prodr. XII, p. 487) suffruticosa ramosissima pilis mollibus patulis villosa viscosa sordide viridis,

ramis erectis basi stricte ramulosis, foliis sessilibus oblongis micro-nulatis subintegris vel minute paucidentatis, floralibus flores aequantibus, verticillastris pluribus 4-2 floris distantibus, bracteis lanceolatis calyce brevioribus, floribus subsessilibus, calycis late campanulati villoso dentibus late lanceolatis spinulosis tubo aequilongis, corollæ albidae hirsutæ tubo calycis dentes non excedente ♀.

Hab. in Persia australi (Auch. 1808!), inter Ssor et Kohrnd (Bge!), in faucibus regionis mediæ alpis Kuh Daëna (Ky. 631!), in montibus Bachtiaricis Kellal et Ssebekuh (Haussk!).

Affinis *S. Aucheri* et *S. ixodi*, folia latiora breviora obtusiora denticulata, bracteolæ illis *S. ixodis* latiores.

§ 7. Ambleiæ.

57. **S. obtusicrenæ** (Boiss. Diagn. Ser. I, 7 p. 57) basi suffruticosa multicaulis tota densissime pannosa cana, caulis erectis crassiusculis simplicibus vel opposite et stricte ramulosis, foliis sessilibus late ovalis vel orbiculatis obtusissimis late et obtuse crenatis basi subcordatis, verticillastris 8-10-floris distantibus summis approximatis, foliis floralibus flores superantibus, bracteolis nullis, calycis albo-pannosi dentibus triangularibus obtusis brevissimis, corollæ roseæ striatæ tubo inclusæ ♀.

Hab. in rupestribus regionis alpinæ Persiae austro-occidentalæ, mons Kuh Daëna (Ky. 599!), mons Kuh Nur et montes Bachtiarici Kellal et Ssebekuh in valleculis frigidis 14000' (Hausekt!). Fl. Aug., Sept.

Caulæ 1-1½-pedales, folia intermedia diam 1-1½-pollicaria ea *Trichodesmatis mollis* referentia, calyx 5 lineas longus. Species pulcherrima.

58. **S. nivea** (Labill. Ic. Syr. III, p. 5 tab. 3) basi suffruticosa dense tomentoso-lanata ramosissima ramis crassis erectis simplicibus vel parce ramosis, foliis subsessilibus integerrimis subtus prominenter nervosis inferioribus oblongis obtusis basi angustatis, floralibus lanceolatis acutiusculis flores superantibus, verticillastris pluribus 6-4-floris subdistantibus vel approximalis longæ racemosis, bracteis subnullis, floribus sessilibus, calycis nivei tubuloso-campanulati dentibus triangulari-lanceolatis muticis tubo triplo brevioribus, corollæ albæ pubescentis tubo inclusæ ♀.

Hab. in collibus Syrieæ circa Damascum et inter Damascum et Antilibanum (Lab!) Boiss! Ky. 473! Gaill!), in Antilibano frequenter (Gaill!), circa Balbeck (Ehr!), in monte Sinai (Auch. 1784! An errore schedulae?).

Rami 1½-1-pedales, folia intermedia 1-1½-pollicaria, racemi sepe semi-pedales, calyces magni sex lineas longi.

59. **S. inflata** (Bth. Lab. p. 562) basi suffruticosa et ramosissima tota adpresso albo-tomentosa, ramis erectis subflexuosis simplicibus vel parce ramosis, foliis subsessilibus subtus sepe magis canis et elevatim nervosis integerrimis obtusis oblongo-ellipticis, superioribus

oblongo-linearibus, floralibus calyce multo brevioribus, verticillastris subsexfloris distantibus vel superioribus approximatis, bracteolis linearis-setaceis brevibus, calyce albo-tomentoso dentibus brevissimis ovatis acutiusculis, florifero oblongo-tubulosso, fructifero inflato valde aucto, corollæ purpureæ extus sericeæ tubo inclusa 5.

Hab. in cretaceis Mesopotamiae in monte Sindjar et in arenosis Derbent i Basian prope Sulimanieh (Haussk!), Persia australi ad Tengnalli 7000' (Haussk!), inter segetes prope Schiraz (Ky. 359!), in calcareis collium demissorum totius Persiae mediae et in ambitu deserti magni frequens (Auch! Bunge! etc.) in Persiae bor. prov. Aderbidjan (Auch!) monte Elbrus prope Derbend (Ky. 749!), circa Yezd (Buhse!), in Caucaso loco non indicato (Kowalewsky!).

Pedalis vel semipedalis, folia 8-12 lineas longa 3-4 lata, calyx florifer 6 lineas fructifer 8 lineas longus 4 latus. Species foliis minoribus angustioribus et calyce tandem inflato a *S. nivea* distincta.

60. ***S. parviflora*** (Bth. in D.C. Prodr. XII, p. 490) basi vix suffrutescens pluricaulis dense subfloccoso-lanata, caulinis erectis crassiusculis a basi opposite ramosissimis, foliis sessilibus superne lineatis integris inferioribus oblongo-ellipticis acutiusculis superioribus et floralibus lanceolatis acutis hisce parvis flores superantibus, verticillastris 4-2-floris remotiusculis longiusculis et anguste spicatis, floralibus parvis sessilibus, bracteolis nullis, calycis ovati dense albolanati dentibus conniventibus linearis-ovatis muticis tubo triplo brevioribus, corollæ albidae hirsutæ limbo minimo calycem vix excedente 2.

Hab. in vallis sterilibus regni Cabulici centralis (Griff!).

Pedalis, folia caulina 1-1½-pollicem longa 6-8 lineas lata, superiora et ramealia minora et angustiora. Inter affines floribus parvis insignis, calyx 2 lineas vix excedens, corolla vix exserta.

61. ***S. affinis*** (Fresen. Mus. Senkenb. p. 91) suffruticosa ramosissima adpresso tomentoso-lanata canescens vel cana, ramis virgatis flexuosis saepè ramulosis, foliis subpetiolatis basi attenuatis oblongis et oblongo-linearibus obtusis muticis rugulosis, floralibus angustatis calyce sublongioribus, verticillastris bifloris distinctis racemosis, calycis breviter pedicellati tomentelli tuboloso-campanulati dentibus triangularibus carinato-nervosis acutis muticis tubo triplo brevioribus, corollæ pubescentis carneæ tubo inclusa 3. *S. Orientalis* Forsk. Eg. Arab. 68 non L. — *S. Palæstina* Vahl Symb. 2, p. 64 non L. — *S. pauciflora* Bth. Lab. p. 560. — *S. Ægyptiaca* Pers. Syn. II, p. 124 (ex specimine herb. Juss!).

Hab. in Arabiâ Petreâ ad radices Sinai (Ehr! Bové! Schimp. 164!, Auch. 1793! Boiss!) in desertis Kahirinis Egypti (Ehr! Ky. 406!) ad Mogheta et Ouadi Hof deserti Ægyptiaco-Arabici (Schweinf! Arabice *Gerratum*).

Affinis *S. Palestinae* a qua differt foliis basi attenuatis nec amplexicaulibus, apice calycisque dentibus muticis, verticillastris særissime bifloris.

62. ***S. Palæstina*** (L. Sp. 1674) suffruticosa ramosissima humilis, ramis crassiusculis adpressissime canis rigidis simplicibus vel

ramosis, foliis sessilibus rugulosis adpresso supra canescentibus subtus canis elevatum reticulato-nervosis integris amplexicaulibus oblongis vel lanceolatis obtusis vel acutiusculis superioribus praesertim mucronatis, floralibus ovatis vel rhombis calyces superantibus, verticillastris 6-4-floris rarius bifloris praeter inferiora remotiuscula in racemos foliosos oblongos approximatis, bracteis miniinis setaceis, calycis sessilis tubuloso-campanulati tomentelli dentibus triangulari-ovatis spinulosis tubo triplo brevioribus, corollæ roseæ hirtulæ tubo subexerto 5. *S. lanata* Mœnch Meth. p. 397 non Jacq.

Hab. ad rupes Palestinæ prope Gaza (Bovè!), ad Hierosolymam! Sieb!, in monte Carmel (Boiss! Gaill!) in Libano prope Bescherre (Ky. 367!) et Dimam (Bl!), Antilibano (Gaill!), circa Damascum (Auch. 1805! Gaill!).

Variat foliis plus minusve angustis et elongatis interdum sesquipollicaribus, calyces 3 $\frac{1}{2}$ lineas longi.

63. *S. Kotschy* (Boiss Diagn. Ser. I, 5 p. 32) suffruticosa ramosissima adpressissime cana, ramis vetustis lignosis denudatis ascendentibus hornotinis erectis pumilis, foliis breviter petiolatis integris ovatis obtusiusculis subtus magis canis et elevatis nervosis, floralibus diminutis oblongis flores æquantibus, verticillastris 4-6-floris in racemum oblongum laxum approximatis, bracteolis setaceis brevissimis vel abortivis, floribus brevissime pedicellatis, calycis breviter tomentosi, campanulati dentibus lanceolatis acutis subspinescentibus divergentibus tubo duplo brevioribus, corollæ sericeæ roseæ tubo inclusa 5. *S. Haussknechtii* Vatke Bot. Zeit. 1875, p. 461.

Hab. in rupestribus prope cacumen montis Gara Kurdistanæ (Ky. 367!), in monte Pir Ömr Gudrun Kurdistanæ Persicæ (forma foliis paulo densius pannosis) (Hausskl.).

Rami novelli 3-5-pollicares, calyx 4 demum 6 lineas longus. Species inter affines dentibus calycis longe attenuatis spinescentibus divaricatis insignis. Rami steriles sub *S. Cydni* nomine ex Ciliciâ ad Cydni fontes a cl. Ky. exs. 329 relati ex foliorum formâ hûe forsitan spectant.

64. *S. tomentosa* (Bth. in p.c. Prodr. XII, p. 489 non Bunge) suffruticosa tomentoso-cana ramis novellis herbaceis simplicibus, foliis breviter petiolatis oblongo-lanceolatis obtusiusculis subtus non nervosis, floralibus diminutis supremis flores non æquantibus, verticillastris sexfloris remotiusculis, bracteis brevibus subulatis villosulis, calyce villoso anguste campanulato fere ad medium in lacinias lanceolatas acutissimas rectas partito, corollâ dentibus calycis breviore 5. *S. Bodeana* Bge Lab. Pers. p. 71.

Hab. in Persicæ prov. Ispahan (Auch. 5106!), in montibus Bachtiaricis (Bode!) Vid. spec. unicum in herb. Hort. Petrop.

Ramus cum racemo 9-10-pollicaris, folia inferiora 12-15 lin. longa 4 lata, verticillastri 6-7, calyx 7-8 lineas longus. Calyce acutissime dentato aſfinis *S. ixodi* sed ob indumentum albo-tomentosum *S. Kotschy* appropriquaanda a quâ foliis angustioribus, calycis dentibus rectis, discedit. *S. tomentosa* Bge Lab. Pers. est forma *S. lavandulæfoliæ*.

65. **S. laxa** (Boiss. et Buhse Aufz. p. 179) basi indurata ramosa adpressissime canescens, ramis erectis tenuiter virgatis subsimplicibus, foliis subsessilibus anguste elliptico-linearibus acutiusculis superne depresso-lineatis subtus elevatim nervulosis, floralibus calyces superantibus et aequantibus, verticillastris remotis 6-4-floris tenuiter et laxe racemosis, bractolis setaceis brevibus, calycis tubuloso-campanulati adpressissime cani dentibus triangulari-lanceolatis spinescentibus tubo quadruplo brevioribus, corollæ roseæ adpresso pubescentis tubo exerto ♂.

Hab. in Persiae borealis jugo Elbrus prope Radkann et in valle Nika (Buhse!), in rupestribus supra Siaret prov. Asterabad (Bgel!).

Facies *Tcuerii Cretici*, rami novelli tenues saepe pedales, folia intermedia pollicem longa lineas tres lata, calyx tres lineas longus; inter affines calycis parvi nervosi dentibus spinulosis brevibus, corollæ tubo exerto insignis.

66. **S. candida** (Bory et Chaub. Fl. Pélop. № 917, tab. 20) perennis pumila cespitoso-multicaulis tota dense et molliter sericeolanata, ramis ascendentibus vel basi prostratis subsimplicibus, foliis breviter petiolatis ovatis obtusis obscure crenulatis superioribus sensim diminutis, floralibus oblongis, verticillastris 6-4-floris præter infimum saepe distantem in spicam oblongam approximatis, bracteis subnullis, calycis lanato-tomentosi campanulati dentibus lanceolatis longe subulato-attenuatis tubo æquilongis demum divergentibus, corollæ carneæ purpureo-maculatae pubescentis tubo inclusa ♀.

Hab. ad rupes Laconiae et Messeniæ ad Mistra et in Taygeto usque ad regionem subalpinam (Bory! Boiss! Heldr!), in monte Diaforti (Bory!).

Rami novelli semipedales pannosi fragiles, folia 8-12 lineas longa, capitulum 1-2-pollicare laxiusculum, calyx 4 lineas longus.

67. **S. chrysantha** (Boiss. et Heldr. Diagn. Ser. I, 7 p. 56) perennis pumila cespitoso-multicaulis tota densissime pannosa cana, ramis a basi prostratae ascendentibus simplicibus et parce ramosis, foliis breviter petiolatis ovato-subrotundis subcrenatis superne sensim diminutis, floralibus supremis calyce brevioribus, verticillastris 6-4-floris in spicas laxiusculas approximatis, bracteis paucis minimis, calycis campanulati dense pannosi dentibus triangularibus spinulosis tubo dimidio æquilongis, corollæ ochroleuce hirtulae tubo inclusa ♀. *S. candida* var. *chrysantha* Bth. in D.C. Prodr. XII, p. 475.

Hab. ad rupes montis Malevo Laconiae prope Agriani 3000' (Heldr! Orph. Pl. exs. 76!). Fl. Jun., Jul.

Fragilis, folia diametro 6-7 lineas lata, valde affinis *S. candida* a qua differre videtur indumento cano absque nitore, calycis dentibus abbreviatis, corollæ colore.

68. **S. Iva** (Griseb. Spic. II, p. 143) safruticosa pumila tota densissime pannosa candida, ramis brevibus erectis, foliis subsessilibus oblongo-lanceolatis obtusis basi attenuatis subintegris, verticillastris paucis 4-6-floris in spicam brevem ovatam approximatis foliis

floralibus ellipticis calyces subæquantibus intermixtis, calycis campanulati lanati dentibus tubo duplo brevioribus in aristulam glabram teneram flexuosam abeuntibus, corollæ ochroleucæ tubo subexerto ♀.

Hab. in Macedonia australi inter Perlepe et Trojazchan 1400' (Griseb!). V. frustul. in herb. Cl. Griseb.

Rami palmares inferne ramulos steriles foliosos edentes, folia 8-12 lineas longa 2-4 lata longitudinaliter rugulosa, calyx 5-6 lineas longus. Aristula molli flexuosa dentium calycis insignis, accedens quoque ad *S. Scainsonii* var. *Scyronicam*.

§ 8. Infrarosulares.

69. *S. lavandulæfolia* (Vahl Symb. I, p. 42) suffruticosa ramosa pumila tota indumento adpresso brevi pilisque longis patentibus obsita, ramulis ascendentibus simplicibus aliis hornotiniis sterilibus brevibus fasciculatim foliosis aliis floriferis infra ramos hornotinos ortis, foliis parallele plus minusve sulculoso-lineatis oblongo-et linear-lanceolatis obtusis in petiolatum longe attenuatis, floralibus sessilibus ovato-ellipticis deflexis calycino tubo sublongioribus, verticillastris 6-4-floris remotiusculis vel in racemum oblongum approximatis, floribus subsessilibus, calycis laciniis plumosis teneris a basi anguste lanceolata longe subulatis patentibus tubo quadruplo longioribus, corollæ purpureæ hirtæ calyce breviore ♀ *S. calycantha* M.B. Taur. Cauc. 3, p. 393. — Rehb. Ic. I, tab. 15. — *Zietenia Orientalis* Gleditsch Act. Ber. 1866.

Hab. in rupestribus regionis subalpinæ et alpinæ ab Anatoliâ ad Persiam, Caria in monte Cadmo (Boiss!), Lyciae montes (Pest! Bourg!), Taurus Pamphylicus et Isauricus 4-6.000' (Heldr!), Galatia (Tchich!), Taurus Cilicius 8000' (Ky. 446!), Cappadocia et Cataonia mons Berydagh 7-9000' (Auch. 1672! Haussk!), Armenia Turcica (Auch. 5188!) ad Kassuklu (Huet!) et Baibut (Bourg!), mons Gara Kurdistaniæ (Ky!) Avroman et Schahu 6.000' (Haussk!), Persia australis mons Kuh Daëna (Ky. 764!) montes Bachtariæ (Auch. 1187!), Persia borealis Alpes (Ky. 457!), prov. Khorassan (Bge!), ditio Talysh (Hohen!), Iberia (M. B.).

β. brachyodon. — Dentes calycini brevius subulati. — *S. tomentosa* Bge Lab. Pers. p. 71, non Bth.

Hab. ad lacum Ourmiah (Seidl!).

Forsan tantum status autumnalis hebetatus speciei.

70. *S. pumila* (Russell Nat. Hist Alepp. II, p. 235 tab. 40!) perennis pumila basi suffruticosa reliquiis petiolorum obsita, ramis sterilibus abbreviatis crebre foliatis, foliis longe petiolatis rugulosis utrinque adpresso tomentellis subtus canescensibus ovalis cordatis crenulatis, ramis floriferis tenuibus ex axillis inferioribus ortis ramum sterilem superantibus foliis minoribus oblongis dentatis floralibus flores superantibus obsitis, verticillastris 6-10-floris dissitis vel superioribus confluentibus, bracteis paucis setaceis pedicellos florum vix æquantibus, calycis campanulati breviter tomentoso-pu-

bescantis dentibus lanceolatis subulatis tandem subrecurvis tubo brevioribus. corollæ flavæ adpresso pubescentis tubo inclusa ♀.

Hab. in Syriâ boreali ad Aleppo (Russ.), ad mœnia Antiochiae (Boiss!), in monte Cassio (Auch. 1701!).

β. brachyodontia. — Calycis dentes abbreviati dimidium tubum æquantes. — *S. floribunda* Montb. et Auch. loc. cit.

Hab. ad rupes Antiochiae cum typo (Auch!), ad Assy prope Aintab (Haussk!).

Semipedalis, rami parte denudata lignosâ procumbentes, folia majora 1 $\frac{1}{2}$ -pollicaria, verticillastri 3-5, calyx 4 lineas longus.

71. *S. rupestris* (Montb. et Auch. Ann. Sc. Nat. Ser. II, 6, pag. 49) perennis basi indurata ramosa pumila viridis patule glanduloso-pubescentia, ramis sterilibus abbreviatis crebre foliatis, foliis longe petiolatis rugulosis ovatis subcordatis obtusis crenatis, ramis floriferis tenuibus a basi rami sterilis ortis eumque sœpe superantibus simplicibus vel parce ramulosis foliis paucis minutis oblongis subpetiolatis obsitis, verticillastris paucis 6-10-floris dissitis vel superioribus confluentibus foliis floralibus subsuperatis, bracteis linearis-setaceis dimidium calycem subæquantibus, floribus pedicellatis, calycis tubuloso-campanulati breviter pubescenti-viscidi dentibus a basi lanceolatâ longe subulato-attenuatis tandem subincurvis tubo vix brevioribus, corollæ flavæ pubescentis tubo subexerto ♀.

Hab. ad rupes Cilicia (Auch. 1701!), in fave Guzelderé prope Mersina (Bal!), prope Gülek 4000' (Ky. Suppl. 334!), prope Alexandrette (Schlumb!).

Semipedalis, folia inferiora sœpe bipollicaria longe petiolata, verticillastri 2-4, calyx 5 lineas longus. Valde affinis *S. pumila* a qua sat differre videtur induimento longiore patulo glanduloso, bracteis sublongioribus, calycis sub-majoris dentibus elongatis.

72. *S. citrina* (Boiss. et Heldr. Diagn. Ser. I, 12 p. 83) perennis suffruticosa basi ramosa pumila tota tomento adpresso denso griseo-flavido pannosa, ramis sterilibus brevissimis foliatis, foliis rugulosis elliptico-oblongis obtusis obscure crenulatis petiolo subæquilongo suffultis, ramis floriferis infra ramum sterilem oriundis eo sublongioribus foliis paucis remotis parvis elliptico-linearibus obsitis, floralibus inferioribus verticillastro sublongioribus, verticillastris paucis in capitulum ovatum basi sœpe interruptum condensatis, bracteis paucis lanceolatis calyce duplo brevioribus, calycis tomentoso-lanati campanulati dentibus triangularibus acutis tubo quadruplo brevioribus, corollæ citrinæ hirsutæ tubo inclusa ♀.

Hab. in rupestribus regionis alpinæ montis Anemas Lycaoniæ et Gheidagh Isauriæ 6-7000' (Heldr!).

β. chamæsideritis — Tenuius et adpressissime tomentella, calycis dentes sublongiores. — *S. chamæsideritis* Boiss. et Bal. Diagn. Ser. II, 4 p. 41.

Hab. in regione alpinæ superiore montis Aslandach Cappadociæ (Bal!), monte Berydagha Cataoniæ 8000' (Haussk!).

Facies *Sideritidis hyssopifoliae*, semipedalis, folia inferiora 1-2-pollices longa petiolo aquilongo, calyces 4 lineas tandem 5-6 longi. A *S. pumila* et *rupestris* indumento, foliis basi cuneatis, calyce brevius dentato, etc., distincta.

73. S. maritima (L. Mant. 82) perennis breviter tomentella canescens-virens, ramis floriferis infra rosulas steriles ascendentibus subsimplicibus, foliis subrugosis tenuiter crenatis radicalibus ovato-spathulatis in petiolum attenuatis, floralibus oblongis obtusis præter insinua calyces non æquantibus, verticillastris 6-4-floris inferioribus distantibus superioribus in spicam approximatis, calycis tubulosocampanulati tomentoso-villosi dentibus lanceolatis acutis spinulosis ad apicem usque hirsutis tubo triplo brevioribus corollæ ochroleucæ tubum æquantibus ♀. Ic. Rehb. tab. 42. — Jacq. Hort. Vind. t. 70.

Hab. in rupestribus maritimis Peloponnesi (herb. Fanché!), insulae Hydra (Sibth.).

Ar. Geogr. Gallia et Hispania mediterraneæ, Italia, Dalmatia.

74. S. pubescens (Ten. Fl. Nap. Prodr. p. 34) perennis pubescens pallide virens, ramis floriferis infra rosulas steriles ascendentibus subsimplicibus, foliis tenuiter crenatis obtusis inferioribus a basi rotundata vel truncata oblongis petiolatis, floralibus ellipticis acutinsculis calyces æquantibus, verticillastris subsexfioris præter superiora distantibus, calycis tubulosocampanulati tomentoso-villosi dentibus lanceolatis longe spinuloso-attenuatis dimidio tubo longioribus ad apicem usque hirtis, corollæ ochroleucæ hirtæ tubo subexerto ♀. Ic. Ten. Nap. tab. 239. — Rehb. Ic. tab. 42. *S. decumbens* Willd. Enum. Suppl. non P. rs. — *S. maritima* Urv. Enum. p. 68 non L. — *S. Lazica* Böiss. in Bal. exs. (forma glabrescens floribus majoribus).

Hab. in agris incultis siccis atque collibus Thraciae ad Byzantium (Auch. 1704!), Anatolie borealis (Wied!), Lyciae (Bourg!), Cataoniae ad Mirasch (Haussk!), Cappadociæ (Auch.), Armeniae Turcice (Huet!), Ponti (Tchih!), Ponti Lazici in maritimis Of et in interioribus (Bal!), ad littora Colchidis et Alchasiæ (Urv., Rupr!), littora Caspii ad Lenkoran (Hoh!), Persia boreali in hortis, montosis (Ky. 69! Buhse! Bge!).

Affinis *S. maritima* differre videtur foliis inferioribus basi non attenuatis, foliis floralibus longioribus, calycis dentibus angustioribus longius spinulosis, corollæ tubo exerto.

A. S. annua radice perenni, calycis dentibus rectis nec curvatis distincta.

Ar. Geogr. Italia meridionalis, Dalmatia.

SECT. III. OLISIA.

75. S. annua (L. Sp. p. 813) annua inferne glabrescula superne pubescens simplex vel collo ramosa, caulis erectis opposite ramulosis, foliis petiolatis crenatis basi cuneatis inferioribus parvis ovati-oblongis superioribus subsessilibus oblongo-lanceolatis, florali-

bus linear-lanceolatis acutis integris calyces superantibus, verticillastris sexfloris distinctis supremis approximatis, calycis tubuloso-campanulati hirsuti glandulisque sessilibus adspersi dentibus lanceolatis acutissimis subspinosis sursum curvatis subaequilongis, corollae albido-ochroleucæ pubescentis tubo subexerto ⊖. Ic. Jacq. Austr. tab. 360. — Rehb. Ic. tab. II, fig. 2 — *S. adenocalyx* C. Koch Linn. XXI, p. 591! — *S. micrantha* C. Koch Linn. XXI, p. 690 (variatio corollâ submonstrose diminutâ calycem non excedente).

Hab. in cultis, Parnassi regio inferior (Guicci!), Thracia et Byzantium (Friv., Sibth.), Bithynia (Thirkel), Pontus Lazicus (Balt!), Armenia Turcica (Bourg!), Tauria, Caucasus et Transcaucasia (Ledebe., Hoh., C.A. Mey.).

β. ammophila Diagn. Ser. II, 4 p. 43. — Rami abbreviati sœpe prostrati, calyces breviores densius et mollis pilosi. — *S. brachyclada* Cosson! in Cat. Mich. Pl. Palest. p. 44 non Noë.

Hab. in arenosis Syriae littoralis ad Berythum et Tripoli (Bl! Gaill!), in Ciliciâ (Ky. It. 1859, N° 6).

γ. Cilicica. — Typo similis, elata, sed folia inferiora basi rotundata nec cuneata, calyceis spinulæ subbreviores. — *S. Cilicica* Boiss. et Bal. Diagn. Ser. II, 4 p. 42.

Hab. in herbidis faucis Guzelderé supra Sedichig (Bal!).

Ar. Geogr. Europa media ab Angliâ et Daniâ ad Rossiam medium, Hispaniam et Italiam centrales, regionem Danubialem.

76. *S. plumosa* (Griseb. Spic. II, p. 139) annua longe et patule hirsuta pallide virens, caule elato simplici, foliis sessilibus ellipticis acute et minute ascendent-serrulatis basi attenuatâ integris apice obtusis in mucronem abrupte attenuatis, verticillastris 6-10-floris in spicam oblongam basi subinterruptam foliis diminutis breviter lanceolatis integris superioribus alabastra subsuperantibus bracteatam approximatis, calyce hirsuto tubuloso-campanulato fere ad medium in dentes lanceolatos longe subulato-aristatos fisso aristis superioribus sublongioribus, corollæ ochroleucæ siccae luridæ tubo inclusa galea indivisa ⊖.

Hab. in Macedonia boreali inter Komanova et Strazia (Vid. sp. in herb. cl. Grisebach).

Pedalis, folia inferiora sesquipollucaria intermediis pollicaribus, spica bipollucaria, calyx cum aristis fere 2 lineas longis 6 lineas longus. Facies *S. hirtæ* a quâ differt indumento longiore, foliis non petiolatis nec cordatis ovatis, galea non bifida, etc.

77. *S. spinulosa* (Sibth. et Sm. Prodr. I, p. 410) annua viridis, caulis ad angulos seberrimis ascendentibus vel erectis elatis sœpe flexuosis simplicibus vel ramulosis, foliis parce hirtis ovatis cordatis crenatis longe petiolatis, floralibus sessilibus oblongis vel lanceolatis integris acutis flores superantibus, verticillastris 6-4-floris distinctis, calyce tubuloso-campanulato viridi glanduloso-hirto dentibus triangulari-ovatis subspinulosis tubo subtriplo brevioribus,

corollæ ample pubescenti-viscidæ albidò-ochroleuca tubo subexserto ⊖.

Hab. in dumosis regionis inferioris et montane totius Graecie (Sprunt Boiss! Heldr!), Thracie et Bithynia (Griseb!), Anatoliae occidentalis (Urv.), Zacynthi (Marg!), Creta (Heldr!).

Ar. Geogr. Dalmatia, Serbia.

78. S. Arabica (Hornem. Hafn. p. 554) annua patule papillari-hirsuta viscidula, caulis erectis elatis crassiuseculis simplicibus vel ramosis, foliis crenulatis imis petiolatis ovatis basi cordatis acutiusculis, superioribus sessilibus floralibusque acute serratis a basi latiori rotundata vel amplexicauli triangulari-lanceolatis acutis verticillastra superantibus, verticillastris numerosis 6-4-floris dissitis longe spicatis, calyce magno piloso parallela et cerebro nervuloso fructifero acreto supra tubum globosum subconstricto dentibus late triangulari-oblongis spinulosis tubo subaequilongis, corollæ violaceæ pubescentis tubo inclusa limbo amplio ⊖. *S. macrosperma* Boiss. Diagn. Ser. I, XII, p. 84.

Hab. in pinguisbus Palestine prope Ramla et in planitie Esdraclon (Boiss!), prope Sidonem et Tortosam (Gall!), in Syria prope Aleppo (Auch. 1697!), inter Tripoli et Hama (Bl!).

β minor. — Flores calyces et semina minora. — *S. obscura* Boiss. et Bal. Diagn. Ser. II, 4 p. 43.

Hab. in cultis derelictis Syrie littoralis prope Mersina (Bal. 512!).

Sæpe bipedalis, insignis calycibus fructiferis latis 6-8 lineas longis versus medium subconstrictis, dentibus latis.

79. S. arvensis (L. Sp. 814) annua patule papillari-pilosa simplex vel ramosa erecta vel decumbens pumila, foliis parvis petiolatis crenatis ovalo-cordatis obtusis, floralibus sessilibus oblongis vel lanceolatis aristulatis calyce sublongioribus, verticillastris 6-4-floris distantibus, calycis hispida tubuloso-campanulati dentibus lanceolatis mucronatis carinato-nervosis tubo subaequilongis, corollæ roseæ hirtulae limbo calycem vix excedente ⊖. Ic. Fl. Dan. tab. 587. — Rehb. Ic. II, fig. I.

Hab. in cultis arenosis Graecie (Sibth.), Peloponnesus (herb. Fauché!), Creta ad Cydoniam (Weiss!).

Ar. Geogr. Insula Canarienses, Maderenses et Azoriev, Europa a Scotiâ ad Rossiam medianam et ex Lusitanâ ad Dalmatiâ, Africa borealis. In Americam borealem et tropicam introducta.

80. S. ramosissima (Montb. et Auch. Ann. Sc. Nat. Ser. II, 6 p. 50) annua molliter canescenti-villosa pumila, caule a basi divaricatim ramoso, foliis inferioribus oblongis in petiolum attenuatis subdentalis, floralibus subsessilibus integris flores longe superantibus, verticillastris subsexfloris inferioribus distantibus superioribus approximatis, calycis tubulosi molliter villosi decennervii dentibus

a basi latiore longissime subulatis recurvis dimidio tubo sublongioribus, fauce dense pilosâ, corollæ purpureæ hirtæ tubo inclusa ⊖.

Hab. in Cappadociâ ad Euphratem (Auch. 1668!), in glareosis Mesopotamiae ad fluvium Zab Ala (Haussk!).

Planta 4-5-pollicaris, calyces fere 5 lineas longi, ex cl. Bth. interdum 10-dentati. Facies *Calamintha Acinos*.

81. *S. satureioides* (Montb. et Auch. Ann. Sc. Nat. Ser. II, 6 p. 51) annua viridis sub lente pruinosa-vel papilloso-puberula, caule humili a basi ramoso ramis tenuibus duriusculis rubellis, foliis integris vel obsolete 1-2-dentatis inferioribus elliptico-oblongis in petiolum longe attenuatis cæteris linearibus sessilibus acutiusculis, floralibus calyces superantibus ciliatis vel hirtulis, verticillastris 4-2-floris distantibus, floribus sessilibus, calycis tubulosi hirtuli subquinquenervii dentibus a basi lanceolatâ longe subulatis subspinosis demum recurvis tubo triplo brevioribus, fauce pilosâ, corollæ purpureæ pubescens tubo breviter exerto ⊖.

Hab. in agris argillosis Syriæ borealis ad Aintab (Auch. 1699!), prope Bir et in deserto Mesopotamiae prope Terek (Haussk!).

Planta 4-6-pollicaris erecta vel basi diffusa, folia intermedia 1½-1-pollicaria 1½-2 lineas lata, calyx 3½ lineas longus. A *S. ramosissimâ* cui affinis induimenti villosi defecta, verticillastris magis dissitis et præsertim calycis brevioris dentibus abbreviatis distincta.

82. *S. burgsdorffioides* (Boiss. Diagn. Ser. I, 12, p. 85) annua patule pubescenti-viscidula viridis, caule humili a basi saepius ramoso ramis rigidulis virgatis, foliis oblongo-linearibus basi longe attenuatis obtusis remote denticulatis, floralibus flores superantibus, verticillastris subquadrifloris remotis laxe spicatis, calyce papilloso tubuloso elevatim decemnervi dentibus a basi lanceolatâ subulato-spinosis demum incurvis tubo triplo brevioribus, fauce albo-pilosissimâ, corollæ purpureæ hirtulæ tubo dentibus calycis non longiore ⊖. *S. satureioides* var. *burgsdorffioides* Bth. in p.c. Prodr.

Hab. in Syriâ prope Aintab (Auch. 1698!), in Mesopotamia inter Suerek et Diarbekir (Ky. 401!), in vineis Armenie australis ad Karput (Noë 860!).

Valde affinis *S. satureioidi* a quâ differre videtur calyce 10-nervi subbreviore, corollæ tubo inclusa, totius plantæ pube copiosiore patulâ viscidulâ.

83. *S. neurocalycina* (Boiss. Diagn. Ser. I, 12 p. 85) annua pumila a basi ramosa glabella, ramis erectis vel diffusis tenuibus, foliis inferioribus oblongis obtusis crenatis in petiolum attenuatis, floralibus subsessilibus ellipticis flores superantibus et aequalibus, verticillastris quidrifloris distantibus longe et laxe racemosis, calycis parce et breviter papilloso viridis nitidi campanulati rigidi elevatim nervosi et transverse reticulati dentibus ovatis ciliatulis abrupte et breviter spinulosis tubo subtriplo brevioribus, fauce glabrâ, corollæ purpureæ hirtæ tubo inclusa ⊖.

Hab. in siccis Palestinæ circa Naplouse et in monte Carmel (Boiss !), in valle Nahr Aoulé Libani inferiori (Bl!), in cultis ad Ras Beyrouth (Bl!). Fl. vere.

Planta semipedalis vel paulo elatior, calyx 3 lineas longus. Facies *Sideritis purpurea* vel *St. satureioides* a quâ calyce eximie differt.

84. S ? Persepolitana (Boiss. Diagn. Ser. I, 7 p. 56) annua glabrescens, caulis pusillis ascendentibus pumilis simplicibus, foliis teneris reniformi-ovatis subcordatis obtuse paucierenatis omnibus etiam floralibus longe petiolatis, verticillastris 1-2 bifloris remotis. calycis sessilis breviter campanulati ore obliqui dentibus ciliatulis ovatis mucronulatis trinerviis et reticulato-nervosis dimidio tubo subbrevioribus, corollâ..... ⊖.

Hab. in rupestribus umbrosis montis Kuh Ajub prope Persepolim Persiae (Ky. 897! et 407 a!).

Planta 2-4-pollicaris, folia majora diam. 6-8 lineas lata, calyx tres lineas longus, facies *S. Corsicae* a quâ cæterum omnino differt sed et genus ob corollam ignotum mihi adhuc subdubium; an suadente cl. a Bunge potius *Lamii* species?

Species dubiæ male descriptæ.

S. Olympica (Poiret Dict. Suppl. V, 326) villosa, coule simplici recto stolonifero, foliis ellipticis argute serratis lanagine ciliatis, caulinis distantibus apice petioli recti patentibus, spicâ densâ lanuginosâ.

Hab. in monte Olympo (Toscan).

S. Macedonica (Don ex Griseb. Spic. II, p. 143) sordide albo-lanata, foliis subsessilibus oblongo-lanceolatis basi angustatis crenatis, verticillastris sexfloris, bracteolis minutis, calyce lanato pungente.

Hab. in Macedonia (herb. Lambert).

BETONICA (L. Gen. 718).

Calyx tubuloso-campanulatus æqualiter quinquedentatus dentibus aristatis. Corollæ labium superius concavum inferius patens trilobum lobio medio majore. Stamina quatuor sub galeâ parallele adscendentia exteriora post anthesin ad latera faucis vix dejecta, antheræ biloculares loculis parallelis vel subparallelis. Stylus apice subæqualiter bifidus. Nuculæ apice rotundato obtusæ. — Herbae perennes. — Genus valde affine *Stachydi*.

SECT. I. ALOPECUROS Bth. Lab. p. 531. — Corollæ tubus intus annulatus. Antherarum loculi paralleli.

1. **B. Jacquini** (Gren. et Godr. Fl. France, 2, p. 694) pubescens, caulis ascendentibus simplicibus elatis, foliis ovatis profunde et

acute dentatis basi valde cordatis radicalibus longe petiolatis, caulinis 2-4 subsessilibus, spicis ob verticillastra inferiora distincta laxe cylindricis, bracteis lanceolatis integris, calycis hirsuti striato-nervosi dentibus apice subulatis tubo duplo brevioribus, corollæ flavæ extus hirsuta tubo calycem vix excedente labio superiore angustato bilobo ♀. *Sideritis Alopecuros* Scop. Carn. tab. 8. — *B. Alopecuros* Jacq. Austr. p. 50, tab. 78. — *B. Orphanidea* Heldr. MSS.

Hab. in rupestribus regionis alpinæ et abietinae Græciae in montibus Kyllene et Chelmos (Orph. Fl. exs. 311 sub *B. Alopecuro*!), Parnasso et Olympo Thessalo (Heldr.). Macedonia in monte Scardo (Griseb. sub *B. Alopecuro*!).

Betonica Alopecurus L. vera ex Pyrenæis et Alpibus Delphinatūs specificē differt spicā crassiori ovatā densā, calyce reticulato-venoso, corollæ brevius hirtæ labio superiori obtuse retuso, etc.

Ar. Geogr. Alpes Helvetie, Italiæ, Germanie australis, Italiae, Dalmatia, regio Danubialis.

SECT. II. BETONICA Bth. Lab. p. 732. — Corollæ tubus intus nudus. Antherarum loculi subparalleli vel divergentes.

2. B. Brantii (Bth. in D.C. Prodr. XII, p. 463 sub *Stachyde*), moltiter et adpresso velutina, caule erecto simplici folioso in capitulum ovatum foliis involueratum et intermixtum abenante, foliis breviter petiolatis valde cordatis ovatis acutiusculis profunde et acute dentatis, floralibus acuminatis calyces superantibus, bracteis linearibus calyces æquantibus, calycis tomentelli dentibus a basi lanceolatâ subulatis flexuoso-subrecurvis tubo quadruplo brevioribus corollæ sulphureæ pilosæ tubum superantibus ♀.

Hab. in Kurdistanū (Brant!). Vidi in herb. Kew.

Pedalis, folia inferiora $1\frac{1}{2}$ -pollicaria pollicem lata, capitulum nuce duplo majus. Corollæ tubus annulo destitutus. Facies *B. Alopecurus* et *Jacquinii*. Omnibus notis *Betonica* nec *Stachys*.

3. B. Orientalis (L. Sp. 811) adpresso pubescenti-velutina, caulis elatis simplicibus vel parce ramosis, foliis subtus canescens-tibus a basi cordatâ lanceolatis obtusis obtuse multierenatis inferioribus longe petiolatis cantiis superioribus subsessilibus, spicis densis oblongis vel oblongo-cylindricis basi interdum interruptis, bracteis lanceolatis apice spinulosis, calycis stellatim pubescentis nervoso-striati dentibus subulatis tubo subduplo brevioribus laue valde hispidâ, corollæ purpureæ glabrae tubo subincurvo calyce sublongiore, staminibus exsertis ♀. *B. macrostachya* Wend. — *Stachys longifolia* Bth. Lab. p. 533. — *B. hirsuta* C.A. Mey. Enum. non L!

Hab. in Caucaso ubi rara videtur (Wilhelms ex Ledeb.), Armeniâ Turcicâ in montibus circa Erzerum (Calv! Tchih!), in pascuis prope Baibut (Bourg!).

4. B. nivea (Stev. McM. Mosq. III, p. 266) tomentella, caulis ascendentibus simplicibus, foliis superne virginalibus subtus canis

lanceolatis obtusis obtuse multierenatis radicalibus longe floralibus brevius petiolatis, verticillastris paucis distinctis foliis floralibus superatis, bracteis lanceolatis lanatis, calycis villosi membranacei reticulato-venosi dentibus lanceolatis inermibus tubo triplo brevioribus fance ciliata, corollae villosae roseae rarius ochrolentice tubo incurvo sensim dilatato calyce longiore, staminibus exsertis ♀.
Stachys nivea Bth. Lab. p. 533.

Hab. in lapidosis alpinis Caucaasi Orientalis (Stev!), in ditionibus Tindal et Daghestaniæ 6000'-7500' (Owerin! Rupr!), Persiaæ bor. monte Elbrus prope Warahosul (Buhset).

Semipedalis vel pedalis, folia angusta *B. Orientalis* sed subtus nivea, calyx 8-9 lineas longus et corolla fere sesquipollicularis incurva *B. grandifloræ* sed tubo breviore. Species elegans.

5. ***B. grandiflora*** (Willd. Sp. III, p. 96) pubescens vel villosa, caulinis ascendentibus simplicibus, foliis subtus subdiscoloribus a basi late cordata late ovatis obtusis late crenato-dentatis inferioribus longe petiolatis, floralibus sessilibus verticillastra superantibus, verticillastris paucis inferioribus 1-2 distinctis, bracteis late ovato-oblongis acutis, calycis membranacei reticulato-venosi superne hirsuti dentibus lanceolatis subulatis tubo duplo brevioribus, fance hirsutâ, corollæ pubescentis purpureæ calyce 3-4-plo longioris tubo incurvo sensim ampliato calyce multo longiore, staminibus subexsertis ♀. Ic. Bot. Mag. tab. 700. — Rehb. Icon. bot. tab. lab. 715. — *Stachys grandiflora* Bth. Lab. — *B. macrantha* G. Koch Linn. XXI, p. 683!

Hab. in pratis alpinis Armenie Turcicæ (G. Koch!), montis Paryadres et circa Gunnischkané (Tchih!), Ponti Lazici supra Djimil 6600' (Bal! Anch!), Caucaasi occid. et centralis 2500'-6000' (G.A. Mey! etc.), Somchetie et Ibericæ (Hoh!), prov. Karabagh (Stev.), Persiaæ borealis monte Karagöl (Szov!).

Species pulcherrima corollâ 12-15 lineas longâ interdum albâ.

6. ***B. Græca*** (Boiss. et Sprun. Diagn. Ser. I, 5 p. 27) pilis stellatis hirsuta caule simplici elato folioso, foliis subtus canescensibus cordatis oblongo-lanceolatis profunde crenato-dentatis inferioribus breviter petiolatis superioribus sessilibus, floralibus inferioribus deflexis verticillastro longioribus, verticillastris multifloris spicatis inferioribus remotis, bracteis oblongo-lanceolatis acuminatis, calycis stellatin hirti obsolete nervosi dentibus lanceolatis subulatis tubo duplo brevioribus, fance villosâ, corollæ carneæ pubescentis tubo calyce vix longiore, staminibus subexsertis ♀. *B. Scordica* Griseb. Spic. II, p. 136.

Hab. in regione sylvaticâ montis Velugo (Etolia), in regione sub-alpinâ montium Scardi et Bertisci Macedoniae 3-4500' (Griseb.).

Bipedalis, facies *Stachydis* vel *Nepeta* e sectione *Pycnonepetâ*, folia inferiora 2-3-pollicaria, calyx sex lineas longus. A *B. hirsutâ* foliis breviter petiolatis, verticillastris remotis, calyce non reticulato et corollæ tubo non exerto valde distincta.

7. **B. officinalis** (L. Sp. 810) pubescens vel glabrata, caule ascendentē tenui simplici, foliis a basi cordatâ ovato-oblongis vel ovatis obtusis crenatis inferioribus longe petiolatis, caulinis remotis sessilibus, floralibus inferioribus lanceolatis serratis acutis supremis linearibus integris, spicâ oblongo-cylindricâ interruptâ, bracteis ovatis mucronatis, floribus parvis, calyce avenio dentibus lanceolatis spinulosis tubo vix brevioribus, corollæ pubescentis purpureæ rarius albæ tubo calyce vix longiore, staminibus subinclusis ♀. Ic. Fl. Dan. tab. 726. — *Stachys Betonica* Bth. Lab. — Rchb. Ic. tab. 16. — *B. stricta* Crit. — *B. serotina* Host. — *B. glabrata* C. Koch Linn. XXI, p. 684!

Hab. in dumosis regionis montanæ, Laconia (ex Sibth.), Macedonia in monte Korthiati et supra Bitolia (Orph!), Bithynia et Armenia (ex C. Koch), Tauria Caucasus et Transcaucasia (Ledeb., Rupr!).

Ar. Geogr. Europa tota ab Angliâ et Gothiâ ad Rossiam.

GALEOPSIS (L. Gen. 717).

Calyx tubuloso-campanulatus quinque-nervius ore subæqualis dentibus quinis spinescentibus. Corolla tubo recto exerto intus nudo, fauce dilatatâ, limbi labio superiore ovato integro fornicate, inferiore patente trifido lobe medio obcordato basi cornubus binis conicis aucto. Stamina exserta sub labio superiore parallela, antheræ biloculares loculis oppositis transverse et bivalvatim dehiscentibus valvulis internis minoribus ciliatis. Stylus apice subæqualiter bifidus. Nuculæ ovoideæ apice rotundatæ. — Herbæ annuæ.

1. **G. Ladanum** (L. Sp. 810) caule erecto ramoso pilis deorsum adpressis mollibus pubescente aequali, foliis ovatis vel lanceolatis serratis, verticillastris inferioribus remotis, floribus sessilibus, bracteis linear-lanceolatis, calyce pubescente eglanduloso, corollæ purpureæ tubo rectiusculo superne dilatato calyce 1-2-plo longiore ⊖. *G. latifolia* Hoffm. Rchb. Ic. tab. 27. — *G. glandulosa* C. Koch Linn. XXI, p. 684!

Hab. in cultis derelictis ditionis septentrionalis, Pontus Lazicus ad Djimil (Bal!), Tauria (Koch!), Caucaso (Ledeb., Rupr!) et prob. in Macedonia et Thracia.

Ar. Geogr. Europa tota a Scandinavia ad Rossiam, Sibiria.

2. **G. Tetrahit** (L. Sp. 810), caule pilis rigidis hispido ramoso sub geniculis incrassato, foliis hispidulis petiolatis ovatis vel oblongis dentatis acuminatis, verticillastris inferioribus remotis floribus sessilibus, bracteis lanceolato-subulatis, calycis dentibus subulato-spinosis fauceque hispidis tubo longioribus, corollæ purpureæ vel albæ tubo calyce subbreviore ⊖. Ic. Rchb. tab. 30, fig. I.

Hab. in cultis, ruderatis ditionis septentrionalis, Macedonia supra Bitolia (Orph!), Pontus Lazicus supra Djimil (Bal!), Caucasus et Transcaucasia (Ledeb., Rupr!).

Ar. Geogr. Europa borealis et media usque ad Hispaniam et Italianam centrales, Sibiria.

3. G. versicolor (Curt. Lond. f. 6, tab. 36) caule hispido subgeniculis incrassato, foliis petiolatis oblongo-ovatis acuminatis serrato-dentatis, verticillastris superioribus approximatis, floribus sessilibus, bracteis lanceolato-subulatis, calycis dentibus subulato-spinosis fauceque hispidis, corollæ magnæ luteæ violaceo-variegatae vel sulphureæ tubo calyce longiore ♂. Ic. Rehb. 30, fig. 3. — *G. pubescens* Griseb. Spic. non Besser.

Hab. in cultis Thracie (Friv!), Caucasi (ex C: Koch).

Ar. Geogr. Europa borealis a Scandinaviâ ad Rossiam, Gallia, Helvetia, Germania, regio Danubialis.

LEONURUS (L. Gen. 722).

Calyx turbinatus subæqualis dentibus quinis apice subulatis spinescentibus demum patentibus. Corollæ tubus stepius inclusus, limbus bilabiatus labio superiore oblongo integro, inferiore patente trifido. Stamina exserta, antheræ per paria approximatæ biloculares loculis parallelis rarius divergentibus. Stylus apice subæqualiter bifidus. Nuculae læves triquetræ apice truncatæ angulis acutis. — Herbæ.

SECT. I. CARDIACA Bth. Lab. p. 517. — Corollæ tubus intus oblique annulatus supra annulum dilatatus, labium superius planiusculum basi angustatum, inferioris lobus medius integer.

1. L. Cardiaca (L. Sp. p. 817) adpresso pubescens, caule elato ramoso, foliis concoloribus vel discoloribus, foliis petiolatis ovatis palmatisidis lobis lanceolatis integris vel dentatis, floralibus sensim diminutis cuneatis oblongo-lanceolatis trifidis vel dentatis, verticillastris dissitis densifloris foliis floralibus longe superatis laxe et longe spicatis, floribus sessilibus, bracteis setaceis, calycis glabri dentibus a basi triangulari subulatis tubo vix brevioribus, corollæ roseæ galeæ hirsutissimæ ♀. Ic. Fl. Dan. tab. 727. — Rehb. Ic. t. 31 et 32. — *L. discolor* C. Koch Linn. XXI, p. 680! — *L. Tataricus* Tchich. As. Min. non L.

Hab. in collibus, ad vias et in ruderatis, Ætolia prope Carpenisi (Samar!), monte Korthiati prope Thessalonican (Heldr!), Armeniâ (Bourgt), Tauriâ, Caucaso et Transcaucasiâ (Led.).

B. villosus. Bth. — Caulis, folia et etiam calyx plus minus hirsuta vel villosa. — *L. villosus* Desf. Cat. Par. p. 75.

Hab. in Tauriâ (Urv.), Caucuso et Transcaucasiâ (Owerin! Ch. Koch!), Anatoliâ bor. (Wied!).

Ar. Geogr. Europa tota a Sueciâ ad Rossiam, Sibiria, regio Himalaica.

2. L. Persicus (Boiss. Diagn. Ser. I, 5 p. 27) breviter tomenitello-hirtus, caulis tenuiter virgatis stricte ramosis, foliis viridibus parvis breviter petiolatis ovatis in laciniis oblongo-lineares incisas palmatisidis vel partitis, floralibus anguste linear-lanceolatis laciniatis verticillastra remota vix superantibus, bracteis subulatis villosis, calycis villoso-lanati vel velutino-pubescentis dentibus a basi triangulari subulatis tubo subaequilongis, corollæ roseæ galea hirsutissimæ ♀.

Hab. in Persia boreali prope Teheran (Ky. 560!), in monte Totschal 7500' (Ky. 500!), in alpibus Elamout (Auch. 5113!), in alpe Demavend 9-1000' (Buhse!).

Habitu potius quam characteribus gravioribus a *L. Cordiacæ* formis diversus; caules multo graciliores stricte ramosi, folia floralia angustissima brevia subsessilia nec longiuscule petiolata.

SECT. II. CHAITURUS Blh. Lab. p. 520. — Corollæ tubus intus nudus æqualis, labium superius fornicateum inferius erectiusculum lobis subæqualibus medio subintegro.

3. L. Marrubiastrum (L. Sp. 817) adpresse pubescens, caule elato ramoso, foliis petiolatis oblongo-rhombeis acutis subinciso-dentatis, floralibus subangustatis verticillastra densa remota longe spicata multum superantibus, floribus sessilibus, bracteis subulatis, calycis breviter pubescentis dentibus a basi triangulari spinulosis tubo 2-3-plo brevioribus, corollæ minutæ roseæ breviter pubescentis tubo inclusa ②. *Chaiturus Marrubiastrum* Ehrh. — Ic. Rehb. tab. 32.

Hab. in dumosis humidiuseculis, Thessalia in planicie maritimæ prope Katarine (Heldr!), circa Byzantium (Audr.), Caucasus et Transcaucasia (Ledeb.).

Ar. Geogr. Gallia, Belgium, Dania, Germania, Hispania et Italia boreales, Dalmatia, regio Danubialis, Rossia media et australis, Sibiria Altaica.

MELITTIS (L. Gen. 731).

Calyx campanulatus amplius membranaceus subbilabiatus labio superiore lato rotundato subbilobo vel 2-3-dentato inferiore bifido lobis rotundatis. Corolla ampla tubo exerto exannulata labio superiore orbiculato subconcavo patente inferiore patente trilido. Stamina 4 sub labio superiore ascendentia non appendiculata, antheræ per paria approximatae biloculares loculis distinctis divergentibus. Stylus apice bifidus. Achenia trigona apice rotundata. — Herba perennis.

1. M. Melissophyllum (L. Sp. 832) pubescens vel hirsuta, caulis erectis subsimplicibus, foliis breviter petiolatis ovatis crenatis inferioribus saepe cordatis superioribus basi cuneatis subdimi-

nutis flores superantibns, verticillastris ad axillas superiores 2-6-floris unilateralibus, pedicellis calyce brevioribus, corolle magna purpurea vel roseo-alba g'abrae vel hirtulae tubo calyce 2-2 $\frac{1}{2}$ -plo longiore ♀. Jacq. Austr. tab. 26. — Rehb. Ic. tab. I. — *M. grandiflora* Smith.

Hab. in sylvaticis et faucibus umbriosis regionis montane et subalpinis, Laconia (Sibth.), Taygetus supra Anavryti (Heldr!), fauces montis Kyllenes 3-4000' (Orph! Heldr!), insula Poros (Wiedt), Macedonia in monte Athone (Griseb.), Olympus Thessaliae (Auch.).

Ar. Geogr. Europa omnis media et australis ab Anglia, Belgio, Germania ad Rossiam meridionalem.

LAMIUM (L. Gen. 746).

Galyx tubulosus vel campanulatus quinquedentatus ore æquali vel obliquo. Corolla tubo saepius exerto intus nudo vel annulato-piloso, limbo bilabiato labio superiore fornicato vel galeato integro retuso vel bifido, faux dilatata, inferioris lobi laterales truncati vel angulati interdum appendiculati, medius substipitatus emarginato-bilobus raro indivisus. Stamina 4 approximata sub labio superiore parallela, antheræ divaricatim biloculares. Stylus aequaliter bifidus. Nuculae acute triquetrae apice truncatæ. — Herbæ.

Specierum Orientalium distributio.

SECT. I. GALEOBIDOLON Blh.—Antheræ glabræ. Corollæ tubus oblique annulato-pilosus.

L. Galeobdolon.

SECT. II. LAMIOPSIS Dumort. Fl. Belg. p. 45 (charactere mutato). — Antheræ hirsutæ. Corollæ tubus intus exannulatus glaber.

* Perennia.

L. molle, Bithynicum, striatum, veronicaefolium, Armenum, microphyllum, cymbalariaefolium, tenuiflorum, stenosiphon.

** Annua.

L. bifidum, amplexicaule, Aleppicum, Ehrenbergii, macrodon, eriocephalum.

SECT. III. LAMIOTYPUS Dumort. (charactere mutato). — Antheræ hirsutæ. Corollæ tubus intus annulato-pilosus infra annulum saepè constrictus.

* Perennia.

L. Lycium, truncatum, maculatum, Gundelsheimeri, album, Robertsoni, tomentosum, crinitum.

** Annua.

L. moschatum, galactophyllum, Ponticum, purpureum.

SECT. I. GALEOBDOLON.

1. **L. Galeobdolon** (L. sp. p. 810 sub *Galeopsis*) viride parce pubescens, collo caules floriferos erectos stolonesque longos foliosos steriles siue radicantes edente, foliis membranaceis petiolatis inferioribus ovatis subcordatis obtuse crenatis superioribus et floralibus basi cuneatis breviter petiolatis oblongo-lanceolatis acuminatis acute dentatis flores superantibus, verticillastris pluribus distantibus, bracteis lanceolatis tubo calycis subbrevioribus, calycis breviter et adpresso pubescentis dentibus lanceolatis subulato-spinulosis tubo brevioribus demum patentibus, corollæ luteæ tubo brevi calyce non longiore intus oblique annulato, labio superiore valde elongato falcato fornicate ciliato, inferioris lobis tribus acutis ♀. *L. Galeobdolon* Crantz Stirp. 262. — *Galeobdolon luteum* Huds. Engl. bot. tab. 787.

Hab. in sylvaticis Thraciae et Byzantii (Sibth.), Ponti Lazici ad Rhizé (Ball), in Persie borealis prov. Ghilan (ex Buhse).

Ar. Geogr. Europa media et australis a Norvegia et Gothia ad Rossiam mediam, Sibiria Uralensis.

SECT. II. LAMIOPSIS.

* Perennia.

2. **L. molle** (Boiss. et Orph. non Willd.) caulis crassiusculis erectis adpresso et breviter hirtis inferne nudis superne simplicibus foliatis, foliis breviter tomentellis canescentibus rugulosis crenatis petiolatis cordatis ovalis, floralibus calyces superantibus, verticillastris multifloris approximatis dense hirsutis, calycis dentibus late-triangulari-lanceolatis breviter acuminatis tubo 2-3-plo brevioribus, corollæ roseæ tubo recto calyce duplo longiore fere a basi sensim ampliato intus nudo, labio superiore hirsutissimo retuso, lobis lateribus brevibus angulatis ♀.

Hab. in regione superiori montis Peristeri Macedoniæ 6800' (Orph.).

Pedale, valde affine *L. Garganicum* a quo foliis molliter tomentellis rugulosolineatis, calycis molliter tomentelli nec virentis hirtuli dentibus latioribus subbrevioribus atque distributione geographicâ diversum esse autumno. *L. Garganicum* in Græciâ et Olympo Bithyno erroneâ indicatum hucusque tantum ex Italiâ australi vidi.

3. **L. Bithynicum**. Bth. in D.C. Prodr. XII, p. 505) glabrescens, caulis ascendentibus simplicibus, foliis parvis a basi cordatâ trian-

gelari ovatis crenatis inferioribus longe petiolatis, floralibus sensim diminulis breviter petiolatis calyces superantibus, verticillastris approximatis 6-4-floris, calycis glabri dentibus a basi triangulari-lanceolata longe acuminatis tubo duplo brevioribus, corollæ roseæ tubo recto calyce subduplo longiore fauce ampliatâ, galea hirsuta sub-integrâ, lobis lateralibus breviter angulatis ♀. *L. Garganicum* var. *glabratum* Griseb. Spic. II, p. 133.

Hab. in saxosis Olympi Bithyni a regione inferiori prope Brussam (Griseb.), usque ad alpinam (Boiss! Clem! Pichl! etc.).

Affine *L. longifloro* Ten. ab eo dentibus calycinis non tantum acutis sed longiusculæ subulati-acuminatis ut videtur distinctum. Planta Serbica a cl. Pancie missa huc spectare videtur.

4. *L. striatum* (Sibth. et Sm. Fl. Gr. VI, p. 46, tab. 557) pubescens, caulis ascendentibus simplicibus, foliis cordatis ovatis obtusiusculis crenatis inferioribus longe petiolatis floralibus diminutis brevissime petiolatis calyces superantibus, verticillastris 8-6-floris inferioribus distantibus, calycis hirti dentibus a basi lanceolata subulatis dimidio tubo sublongioribus, corollæ calyce quadruplo longioris carneæ roseo-striatae tubo recto tenui intus glabro fauce dilatata, galea elongata bifida, lobis lateralibus angulatis obsolete denticulatis ♀. *L. glechomoides* Sm. in Rees Cycl. V, 20. — *Moldavica Orientalis hederæ terrestris folio* Tourn. herb! — *Dracocephalum lamiifolium* Desf. in Tourn. Cor. tab. 15.

Hab. in rupestribus regionis montanae et subalpinæ Græciae, montes Ænæ, Cephaloniae et Lassiti Cretæ (Heldr!), Hymettus et Parnes Atticae (Sibth., Auch. 1683! Sprn! Boiss! Orph. Fl. exs. 97! Heldr!), mons Helikon Bæotiae et Delphi Eubœa (Heldr! Orph!), Veluchi Ætolie (Samar!) et Æta Thessaliae (Sprn!), Macedonia in monte Corthiati (Orph!) et Atho (Griseb.), Laconia in monte Malevo (Orph! Heldr!), Anatoliæ occidentali in montibus Cadmo et Tmolô (Boiss!).

β. minus. — Pubescens, caules breviores magis decumbentes, folia diminuta, flores in verticillastris saepe paucioribus pauciores, flores minores, calycis dentes interdum brevius subulati et corollæ galea minus profunde bifida. — *L. Cytteneum* Boiss. et Heldr. MSS. — *L. nivale* Boiss. et Heldr. Ser. I, 7 p. 54. — *L. rectnm* Schenk Pl. Spec. p. 19.

Hab. in glareosis regionis alpinæ Taygeti (Boiss! Heldr!), montis Kyllene 3500' (Orph!), Parnassi regione abietinâ 3500'-4000' (Heldr. Herb. Norm. 727! Orph!), in subalpinis Amani Syriæ (Ky. 236!), in totius Libani et Antilibani regione subalpinâ et alpinâ (Ehr! Schub., Boiss! Bl! Gaill!).

γ. *nepetæfolium*. — Folia ovato-triangularia apice attenuata sed obtusa profundius dentata. — *L. nepetæfolium* Boiss. in Ky. Sched.

Hab. in Ciliciâ littorali ad Kechlick (Bal!), Tauro Cilicio (Ky. 47!), monte Djebel Nur Ciliciæ (Ky. 44 sub *L. nivali*).

δ. *reniforme*. — Caules abbreviati saepe decumbentes, verticillastris pauciflori, folia glabra vel puberula hirsuta et saepe minima obtuse inciso-crenata ovato-reniformia subcordata vel basi truncata, inferiora

longe petiolata, calycis parce hirtuli laciniae lanceolato-subulatae saepe tubo vix breviores, galea breviter bisida. — *L. reniforme* Montb. et Auch. Ann. Sc. Nat. Ser. II, 6, p. 47. — *L. plebeium* Boiss. et Huet MSS.

Hab. in regione alpinâ Ponti (Tchih!), Ponti Lazici (Ky. 1685!). Armeniae Turcicæ ad Zazalarhane (Huet!) et monte Fecelein prope Baibut (Bourg!), Kurdistaniâ (Barré de Lancy!), monte Berydagh Cataoniae 8-900' (Haussk!), regione alpinâ montis Argæi Cappadocie (Bal!), ditione Kassanoglu Ciliciae orientalis (Ky. 166!), Antilibato supra Bludan 6000' (Ky. 55!).

ε. pictum. Caules siliiformes breves diffusi et folia minuta formæ præcedentis, sed minus profunde lobato-crenata, reniformia vel ovato-orbiculata, flores quoque pauci, calycis parce et breviter hirtuli dentes apice brevis subulati. — *L. pictum* Boiss. et Heldr. Diagn. Ser. II, 4 p. 44.

Hab. in regione alpinâ Parnassi ad rupes umbrosas (Heldr! Orph. Fl. Gr. 440!).

Foliorum magnitudine et caulis elatis vel pumilis mutabile, corollæ subpollicares interdum 45-46 lineas longæ. Facies *L. longiflori* et *Bithynici* a quibus galea bisidâ distinguitur.

5. ***L. veronicæfolium*** (Bth. Lab. p. 510) glabriuscum cespitosum. caudiculis nudis fragilibus inter lapides occultatis in caules brevissimos simplices abenntibus, foliis parvis longe petiolatis reniformibus late cordatis grosse et obtuse crenato-lobatis, floralibus submajoribus calyces superantibus, verticillastris 2-3 bifloris approximatis, bracteis lanceolatis brevissimis, calycis glabri dentibus triangularibus acutis tubo triplo brevioribus, corollæ roseæ purpureo-maculatae tubo recto nudo calyce triplo longiore, fance ampliatâ, lobis lateralibus obtuse angulatis, galeâ hirsutâ truncatâ retusâ vel obsolete 3-4 denticulatâ ♀. *L. Manganottianum* Clem. Sert. Or. t. 4.

Hab. in lapidosis alpinis Olympi Bithyni (Sibth., Auch. 1683, Boiss! Pichler!).

Caudiculis cum caule florifero 4-5-pollicares, foliorum majorum lamina diam. 6 lineas lata, calyx 4 lineas, corolla 20 lineas longa. A *L. striato* var. *reniformi* et var. *picto* quibus facie affine est differt calycis dentibus brevibus, corollæ tubo multo magis elongato, galeâ truncato-retusâ nec bisidâ.

6. ***L. Armenum*** (Boiss. Diagn. Ser. I, 7 p. 55) glabriuscum cespitosum, caudiculis nudis fragilibus inter lapides occultatis in caules brevissimos simplices abenntibus, foliis parvis longe petiolatis ovatis cuneatis obtuse crenatis, floralibus submajoribus calyce longioribus, verticillastris paucis 6-4-floris terminalibus congestis, bracteis subulatis tubo calycinæ subbrevioribus, calycis dentibus hirtulis lanceolatis subulato-attenuatis tubo vix brevioribus, corollæ roseæ calyce quintuplo longioris tubo recto intus nudo in fance ampliato, galeâ tomentosâ integrâ ♀. *L. veronicæfolium* Bth. Ann. Sc. Nat. 1835, p. 47 non Bth. Lab.

Hab. in alpibus Kechichdag Armeniæ (Auch. 1682!).

Facie, foliis parvis, affine *L. veronicæfolio*; differt foliis basi cuneatis nec

cordatis, calycis dentibus elongatis subulatis. *L. striatum* var. *reniforme* affine quoque habet folia reniformi-cordata majora, galeam bisidam.

7. ***L. microphyllum*** (Boiss. Diagn. Ser. I, 5 p. 25) glabriusculum, caudiculis e rhizomate elongato suberoso tenuissimis subterraneis fragilibus in caules tenuissimos breves abeuntibus, foliis minimis longe petiolatis rotundato-reniformibus subcordatis obtuse paucicrenatis, floralibus non diminutis calyces superantibus, verticillastris 2-3 bifloris terminalibus confertis, bracteis lanceolato-linearibus calycis tubum æquantibus, calycis dentibus hirtulis triangulari-lanceolatis acuminatis tubo subdupo brevioribus, corollæ carneâ calyce 5-6 plo longiore tubo subcurvato intus nudo in faucom purpureo-maculatam sensim ampliato lobis lateralibus angulatis, galeâ valde elongatâ hirsutâ in lobos oblongos profunde divisâ ȝ. Ic. Tchih. As. Min. tab. 25.

Hab. in lapidosis calcareis mobilibus regionis alpine montis Cadmi supra Colossas in Cariâ (Boiss.).

Caules parte non subterraneâ pollicares, folia 2-4 lineas diam. lata, calyx 4-5 lineas longus. Corolla sesqui et fere bipaniculata, labium superius 8-9 lineas longum ultra tertiam partem bilobum quâ notâ, foliis minoribus, etc., haec species pulcherrima ab affini *L. veronicæfolio* differt.

8. ***L. cymbalariaefolium*** (Boiss. in Bourg. exs. Lyc. 1860) glabrum, caudiculis e rhizomate elongato numerosis tenuissimis fragilibus longis nudis in caules brevissimos abeuntibus, foliis longe petiolatis minutissimis reniformi-subcordatis 3-5-lobis etiam floralibus longe petiolatis calyces æquantibus, verticillastris 2-3 bifloris dense capitatis, bracteis lanceolatis tubum calycis æquantibus calycis dentibus longe lanceolato-acuminatis ciliatulis tubo sesquilonioribus, corollæ roseæ calyce quadruplo longioris tubo recto lobis lateralibus angulatis, galeâ profunde bilobâ ȝ.

Hab. in lapidosis regionis alpinæ montis Akdaghi Lycae (Bourg. exs. 215!).

Affine *L. microphyllum* a quo differt foliis 2-2½ lin. tantum latis 3-5-lobis, calycis majoris (demum 9 lineas longi) laciniis longissimis, corollæ tubo breviore.

9. ***L. tenuiflorum*** (Fisch. et Mey. Ann. Sc. Nat. 1854 p. 34) pubescenti-villosulum canescenti-virens multicaule caulibus pumilis ascendentibus simplicibus, foliis petiolatis ovato-triangularibus obtuse et grosse crenatis, floralibus calyces superantibus cuneatis subsessilibus, verticillastris 1-3 paucifloris, bracteis lanceolatis tubum calycinum subæquantibus, calycis dentibus hirsutis a basi lanceolata longe subulatis divergentibus tubo glabriusculo sublongioribus, corollæ hirsutæ tubo gracillimo recto calyce triplo longiore in faucom ampliato, lobis lateralibus angulatis, galeâ profunde bisidâ ȝ.

Hab. ad Guenzlu Lycaoniae 3000' (Tchih!), in rupestribus montis Tcharyklar Pamphyliæ supra Adalia (Bourg. exs! sub *L. recto*), in monte Davros Pisidiæ (Heldr. exs. 1845 sub *L. nivali* var!).

Caules semipedales, folia demum pollicaria, corolla fere pollicaris. Species a formis *L. striati* a quo eam prius non distinxii calyce multo longius fisco et corollæ tubo duplo angustiore et tenuiore distinctissima.

10. *L. stenosiphon* (Boiss. et Haussk) glabrum, caulis punnatis simplicibus, foliis minutis reniformibus paucicrenatis basi subtruncatis vel subcordatis omnibus et inferioribus longe petiolatis floralibus submajoribus flores superantibus, verticillastris 2-4 approximatis 2-4 floris, calycis glabri subinervi dentibus ciliatulis lanceolatis acuminatis tubo subdupo brevioribus, corolle roseæ tubo recto tenuissimo intus nudo calyce 2 1/2-plo longiore, fauce subampliatâ, lobis lateralibus angulatis, galeâ hirsutâ subretusâ ♀.

Hab. in rupestribus calcareis alpinis montium Avroman et Schahu Kurdistaniae Persicae 9100' (Haussk. 7981).

Caules 4-5-pollicares, folia *Veronicae hederaefoliae* 3-7 lineas diametro lata, calyx 3 1/2 lineas, corolla 13 lineas longa. Species foliis minutis reniformibus, calycis dentibus brevibus, corollæ galeâ indivisa ab affini *L. tenuifolio* distincta.

• • Annua.

11. *L. bifidum* (Cyrill. pl. rar. fasc. I, p. 22, tab. 7) annum pubescens vel glabratum, caulis crassiusculis ascendentibus, foliis inferioribus petiolatis cordato-ovalis crenatis vel subincisis, floralibus approximatis breviter petiolatis triangularibus acuminatis profunde incisis flores superantibus, verticillastris 2-3 capitatis vel inferioribus distantibus, bracteis linearibus brevibus, calycis hirti obliqui laciniis lanceolatis apice subulatis tubo brevioribus demum patentibus, corellæ albæ tubo recto non annulato calyce 3-4-plo longiore, fauce dilatatâ, lobis lateralibus dente brevi linearis auctis, galeâ bifida lobis obovatis saepe divaricatis, acheniis reticulatim albo-tuberculatis ♀. Ic. Ten. Fl. Nap. tab. 153.

Hab. in cultis Zaeynthi (Marg!), Eubeæ (Wied!), Peloponnesi (Sart!), Atticæ (Bots!), Thessaliae (Sprun!).

Ar. Geogr. Corsica, Sardinia, Sicilia, Italia australis, Dalmatia.

12. *L. amplexicaule* (L. Sp. 809) annum pumilum a collo saepius pluricaule pubescens, foliis inæqualiter et obtuse crenatis inferioribus petiolatis orbiculari-ovatis saepius cordatis, floralibus sessilibus reniformibus amplexicanilibus lobatis calyces superantibus, verticillastris 1-2 terminalibus dissitis, bracteis nullis, calycis tubulosi villosi dentibus lanceolato-subulatis conniventibus tubo subbrevioribus, corollæ purpureæ tubo recto tenui intus nudo! calyce subtriplo longiore, fauce dilatatâ, lobis lateralibus muticis, galeâ villosa ovata obtusa, acheniis laevibus ♂. Ic. Engl. bot. t. 770. — Rehb. tab. 3. — *L. Mesogæon* Heldr. MSS.

Hab. in cultis regionis inferioris et alpinæ in totâ fere ditione a Græciâ ad Macedoniam et Thraciam! regionem Caucasicam!, Persiam borealem! et australem (Ky! Bgl!), Syriam et Palestinam (Boiss!).

β. incisum. — Folia etiam inferiora profunde incisa imo trisecta segmentis petiolatis partitis.

Hab. in cultis prope Aleppo et in deserto Mesopotamiae ad Terek, Kotsch-hassar, etc. (Haussk!).

Ar. Geogr. Europa omnis a Britanniâ et Scandinaviâ ad Rossiam medium, Sibiria, Africa borealis.

13 L. Aleppicum (Boiss. et Haussk. in Haussk exs. 1865) annum pumilum a collo saepius pluricaule sub lente puberulum, foliis obtuse crenatis inferioribus petiolatis ovatis et ovato-oblongis, floralibus sessilibus amplexicaulibus oblongis profunde et obtuse incisis calyces superantibus, verticillastris 2-3 distantibus, calycis dentibus albo-villosissimis late lanceolatis tubo subbrevioribus, corollæ purpureæ tubo recto tenui intus nudo calyce subtriplo longiore, fance dilatata lobis lateralibus muticis, galea villosa ovata obtusa, acheniis tuberculatis albis confluentibus leprosis ⊖.

Hab. inter segetes prope Aleppo Syriæ ubi *L. amplexicauli* conumixtum crescit (Heldr!). Fl. Febr., Mart.

Valde affine *L. amplexicauli* a quo ex amic. Haussknecht qui ambo ex vivo comparavit specifice distinguitur foliis oblongis, calycis dentibus hirsutissimis latioribus, acheniis.

14. L. Ehrenbergii (Boiss. et Reut.) unum sub lente puberulum, foliis minimis ovatis paucicrenatis inferioribus petiolatis, floralibus sessilibus cuneatis calyces æquantibus, floribus in capitula terminalia ovata congestis, calycis pubescentis dentibus ovalis macro-natis tubo triplo brevioribus, corolla..., acheniis albo-vermiculatis ⊖.

Hab. in monte Makmel Libani prope nives (Ehrenb!), in summis moutibus Elmali Lyciae (Bourg!).

Plantula 1-2-pollicaris affinis *L. amplexicauli* et *L. Aleppico* a quibus dentibus calycis ovatis brevibus et achenii sculpturâ differt. Specimina quæ vidi cryptantha sunt quod in *L. amplexicauli* sæpe occurrit.

15. L. macrodon (Boiss. et Huet Diagn. Ser. II, 4. p. 45) annum adpresso pubescens collo pluricaule pumilum, foliis inferioribus petiolatis ovatis basi subtruncatis crenatis, floralibus majoribus sessilibus amplexicaulibus lobato-crenatis flores superantibus, verticillastris 1-3 approximatis densilloris, calycis hirsuti dentibus a basi lanceolata longissime et tenuissime setaceis tubo duplo longioribus tandem subpatulis, corollæ roseæ tubo recto intus nudo calyce parum longiore, fance subampliatâ, galea villosa obtusa, acheniis albo-reticulatis ⊖ *L. orthodon* et *L. plebeium* Boiss. olim.

Hab. in arvis incultis montanis Armeniæ Turcicæ prope Tortum (Huet!) et Gümüşkhané (Bourg!), montium Argei et Masmenenagh Cappadociæ 6000' (Bal!), in cedretis ditionis Kassanoglu Ciliciæ 6000' (Ky. 146!).

Planta 2-4-pollicaris, folia floralia majora eis *L. amplexicauli* a quo distinguitur calycis duplo majoris 6 lineas longi lacinias longissime setaceis tandem subpatulis tubo duplo longioribus, acheniis reticulatis.

16. *L. eriocephalum* (Bth. in D.C. Prodr. XII, p. 506) annum collo ramosum radice simplici, caulis crassis fragilibus foliosis, foliis inferioribus hirsute pubescentibus longe et tenuissime petiolatis minutissimis ovato-orbiculatis obtuse 5-9 crenatis, floribus in capitula globosa bombycina compacta dense congestis, foliis floralibus multo majoribus sessilibus breviter cuneatis subamplexicaulibus apice flabellatim in laciniis lineares obtusas incisis calyces æquantibus lanatis, calycis laciniis lanatis a basi late lanceolata subulatis tubo subbrevioribus, corollæ magnæ purpureæ tubo recto non annulato calyce $2\frac{1}{2}$ plo longiore in faucom valde ampliato, lobis lateralibus breviter dentatis, galea villosa oblonga integrâ, achenis longis laevibus ⊖.

Hab. in lapidosis mobilibus regionis alpinæ jugorum Gisyl Tepe, Koschan, Metdesis Tauri Cilicici 9000' (Ky. 1141, Bal 5031).

Planta 2-4-pollicaris, foliorum inferiorum lamina 2-4 lineas diam. lata, capitulo valde lanatum demum nuce majus, corolla 14-15 lineas longa, achenia 2 lineas longa. Alpium Cilicie eximum decus!

SECT. III. LAMIOTYPUS.

* Perennia.

17. *L. Lycium* (Boiss. in Bourg. exs. 1860) adpresse pubescens canescens, caulis numerosis ascendentibus pumilis simplicibus, foliis parvis triangulari-ovatis cordatis obtusiusculis crenatis petiolatis, floralibus calyces superantibus, verticillastris binis vel solitariis paucifloris, bracteis lanceolatis brevissimis, floribus parvis, calycis hirti dentibus late lanceolatis acutis tubo subduplo brevioribus, corollæ roseæ hirtæ tubo recto tenui intus piloso-annulato infra annulum non angustato calycem parum excedente, fauce subampliata, lobis lateralibus bidentatis, galea elongata angusta profunde bifida ♀.

Hab. in rupestribus regionis alpine montium supra Elmalu Lyciae (Bourg. 217!).

Caules semipedales, folia sub anthesi 6 lineas longa et lata demum subampliata, corolla 9-10 lineas longa tubo eo *L. striati* multo tenuiore. Facies *L. tenuiflori* sed tubus corollinus calycis proportione multo brevior intus piloso-annulatus nec nudus. A cæteris speciebus sectionis galea bifida alienum.

18. *L. truncatum* (Boiss. Diagn. Ser. I, 42, p. 86) viride sublente parce et adpresse setulosum, caulis diffusis elongatis, foliis teneris longe petiolatis ovato-oblongis obtusis basi truncato-subcordatis grosse et obtuse crenato-lobatis, floralibus breviter petiolatis flores superantibus, verticillastris 2-3 distinctis 6-8-floris, bracteis subulatis hirsutis brevissimis, calycis ad nervos adpresse setulosi dentibus breviter ciliatis a basi lanceolata longe setaceo-acuminatis tubo aequilongis vel longioribus, corollæ roseæ tubo calyce subduplo longiore paulo supra basin oblique piloso-annulato versus medium subcurvato fauce subampliata lobis lateralibus longe setaceo-acuminatis, galea valde hirsuta indivisa truncata ♀.

Hab. in rupestribus umbrosis vel humidis, Galilea circa Banias ((Boiss!)), Syria littoralis circa Tripoli (Bl!), Libanus circa Ghazir et Haramoun (Gail!), circa Hasrun, Eden et Bezulmar (Bl!), Amani regio inferior (Ky. 263!), circa Beilan (Haussk!).

Caulis 1-2-pedales, petioli inferiores sa-pe tripollicares, folia 2-3-poll. longa, calyx 7 lineas, corolla 14-15 lineas longa tubo infra annulum obconice angustato. Species corollæ tubo multo longiore vix curvato infra annulum obliquum nec transversum sensim nec abrupte attenuato, galea truncata a *L. maculato* distincta.

19. *L. maculatum* (L. Sp. 809) plus minus pubescens, caulis basi prostrata radicantibus erectis, foliis petiolatis flores superantibus, verticillastris 2-5 distinctis 6-10-floris, bracteis minimis, calycis dentibus lanceolatis longe subulato-acuminatis ciliatis tubo aequilongis vel longioribus tandem divaricatis, corollæ purpureæ tubo calycis dentibus sublongiore recurvo-porrecto supra basin transverse annulato et infra annulum abrupte constricto, fauce dilatata, lobis lateralis triangularibus sæpe appendice subulata auctis, galea oblonga integrâ ♀. Ic. Fl. Græc. tab. 556. — Rehb. tab. 4. — *L. rugosum* Ait. non Sibth. et Sm.

Hab. in umbrosis regionis montanae, mons Malevo Laconiae supra Hagios Petros (Orph. Fl. exs. 543!), Argolis (Sibth.), Thracia et Macedonia montes (Sibth., Orph! Friv!), Pontus supra Samsun (Tchih!), montes Ylgas Dagh et Aladagh Szeben Anatoliae bor. (Wied!), atque in montibus Gumusch Armeniae (Bourg!) (formæ alpinæ foliis minoribus), Caucaso (Led., Rupr.), Persia boreali in prov. Ghilan et Mazanderan (Bge! Buhse!).

Ar. Geogr. Europa borealis et media ab Angliâ et Sueciâ ad Rossiam medium, Lusitaniam, Hispaniam et Italiam medias, Dalmatiam.

20. *L. Gundelsheimeri* (C. Koch Linn. XVII, p. 297) caule elato striato parce rainoso setis deflexis sparsis obsito, foliis membranaceis teneris sparsim setulosis petiolatis ovatis acuminatis argute serrato-dentatis inferioribus basi rotundatis floralibus cuneatis oblongo-lanceolatis flores superantibus, verticillastris pluribus remotis, bracteis subulatis brevibus, calycis parce setosi dentibus præter basin latiorem subulatis ciliatis tubo sublongioribus, corollæ purpureæ tubo rectiusculo prope basin annulato et infra annulum subangustato calyce sesquiongiore, dentibus lateralibus 2-3 lacinulatis, galea oblonga hirsutâ ♀. *L. Cappadocicum* nunc moschatum nunc fatidum flore mayno Gundelsh. herb!

Hab. in Armeniâ Turcicâ (Gundelsh. in herb. Berol!), in tractu Daratchitachak (C. Koch).

Affine *L. maculato*, elatus sæpe tripedale, folia teneriora inferiora basi rotundata nec cordata, cætera cuneata, corolla 10-12 lineas longa. Ulterius comparandum.

21. *L. album* (L. Sp. 809) plus minus pubescens, caulis basi prostrata radicantibus erectis, foliis ovato-cordatis acuminatis inæqualiter serratis, floralibus breviter petiolatis flores superantibus, verticillastris 8-9 distinctis plurifloris, bracteis parvis ciliatis, calycis

laciinis lanceolatis longe subulato-acuminatis ciliatis demum patulis tubo longioribus, corollae albae tubo lato calyce sublongiore recurvo porrecto supra basin oblique annulato-piloso infra anulum valde constricto, fauce dilatata lobis lateralibus in 2-3 dentes altero saepe subulato ab initibus, galea obtusâ ♀. Ic. Engl. Bot. tab. 768. — Rehb. Ic. t. 4.

Hab. in subalpinis et alpinis humidis, Ponti Lazici ad Djimil (Bal!). Armenia ad Erzerum (Huet!), Baibut (Bourg!), in monte Bingöldagh (Ky!), ad fontes Djoschdag Cilicie orientalis (Ky. 92!), Caucaso et Transcaucasiâ (Ledeb., G.A. May! Rupr!), montibus Persiae borealis in prov. Ghilan et Mazanderan (Buhse, Bge), ad rivulos in monte Totschal prope Teheran (Ky. 49!) et in monte Ssahend (Buhse) sub *L. Robertsonii*.

Ar. Geogr. Europa borealis et media ab Anglia et Scandinaviâ ad Rossiam borealem, Hispaniam et Italiam boreales, Sibiria.

22 L. Robertsoni (Boiss. Diagn. Ser. I, 7, p. 54) glaucum siccitate nigricans, caulinis striato-sulealis glabris parce ramosis, foliis sub lente parce et minute puberulis membranaceis petiolatis obtuse et grosse crenato dentatis ovatis vel ovato-orbiculatis basi truncatis vel late subcordatis, floralibus breviter petiolatis acutiusculis flores superantibus, verticillastris panicis remotis, bracteis subulatis hirtis saepe deflexis, calycis breviter puberuli laciinis a basi lanceolatâ longe subulatis patentibus tubo sesquilonioribus, corollæ albae tubo recto calycis dentes non superante supra basin obliquo annulato-piloso infra anulum constricto, fauce dilatata, lobis lateralibus subulato-appendiculatis, galea albo-hirsutâ truncatâ denticulatâ ♀.

Hab. in glareosis alpis Kuh Daena Persiae australis (Ky. 604!), in rupestribus calcareis montium Avroman et Schahu (Haussk!).

Pedale vel sesquipedale, calyx 7 lineas longus, corolla pollicaris. *L. albo* affine differre videtur glaucidine, foliis obtusis obtuse crenatis, corollæ longioris tubo recto.

23 L. tomentosum (Willd. Sp. III, p. 90) molliter tomentosum, candens tenuibus elongatis subterraneis nudis, caulinis pulcherrimis, foliis petiolatis ovatis cordatis crenatis, floralibus flores superantibus, verticillastris 2-5-distinctis 6-8-floris, bracteis subulatis plumosis, calycis dense albo plumosi dentibus lanceolatis tubo subbrevioribus, corolle roseæ tubo subcurvato calycem parum superante supra basin oblique annulato et infra anulum abrupte coarctato, fauce subampliciatâ dentibus lateralibus dente interdum subulato elongato aequalis, galea hirsutissimâ obtusâ denticulatâ ♀. *L. vestitum* Blh. in p.c. Prodr. XII, p. 508. — *L. alpestre* Trantv. Stirp. Nov. deser. V, p. 14!

Hab. in glareosis regionis alpine, Armenia Turcica in Ponto Lazico supra Djimil 9'00' (Bal!), montibus Erzerum (Calv!), districtu Kars (Radde!), summis Agerowdaghi Armeniae Kurdiæ 4200' (Ky. Suppl. 672!), Caucaso omni 8'00'-9'50' (G.A. May!), Caucaso orientali usque ad 10500' (Rupr!), Persia boreali in monte Totschal prope Teheran (Ky. 565 a!), in alpibus Demavend (Auch. 5020! Buhse!).

β. filicaule. — Caules breves 1-2 pollicares, folia dinninta diametro 4-6 lineas lata, calycis laciniæ triangulares mucronatæ tubo plus duplo breviores. — *L. filicaule* Boiss. Diagn. Ser. I, 12 p. 86.

Hab. in glareosis montis Diklo Tuschetiae Caucasi orientali 90° 0' (Rupr!), monte Totschal Persiae ad lacum Derja-i-Jach (Ky. 295!).

Species a formis minoribus alpinis *L. maculati* præter indumentum molliter tomentosum verticillastris valde plumosum, foliis rotundatis obtuse crenatis, tubo corollæ rectiusculo oblique annulato et galea hirsutissimâ distincta. *L. Columnæ* Ten. huc saepe relatum est *L. maculati* forma

24. *L. crinitum* (Montb. et Auch. Ann. Sc. Nat. 1836 p. 48) breviter villosum, caulis ascendentibus, foliis breviter petiolatis ovatis cordatis acuminatis acute serratis, floralibus flores superantibus, verticillastris approximatis, bracteis tenuiter subulatis, calycis dentibus ciliatis a basi sublatiore angustissime subulatis divergentibus tubo hirtulo duplo longioribus, corollæ tubo calycis dentibus multo breviore rectiusculo supra basin annulato, fauce subampliatâ lobis lateralibus subulato-appendiculatis, galea villosâ oblongâ ♀. *L. Persicum* Boiss. et Buhse Enum. p. 180.

Hab. in Cappadociâ orientali (Auch. 1084!), in jugo Paryadres Ponti (Tchih!), in montibus Elbrus prope Warahosul et Demavend Persiae bor. (Buhse!).

Semipedale, valde affine *L. maculato* et *tomentoso* a quibus dentibus calycinis subulatis duplo longioribus differe videtur, a priore insuper indumento villosulo, a posteriore foliis acuminatis et acute dentatis.

" Annuæ.

25. *L. moschatum* (Mill. Dict. N° 4) annum, caulis crassis striatis ascendentibus saepe ramosis, foliis viridibus hirtulis petiolatis late cordatis ovatis crenatis, floralibus flores longe superantibus basi saepe albo-maculatis acutiusculis subacuminatis, verticillastris numerosis distinctis plurifloris, bracteis lanceolatis ciliatis tubo calycinio brevioribus, calycis glabriuscui fructiferi aucti dentibus triangulari-lanceolatis medio dorso et margine aspero-ciliatulis tubo subæquilongis superiore sublongiore omnibus demum stellatim patentibus, corollæ albæ tubo calyce breviore recto annulato-piloso, fauce amplâ lobis lateralibus triangularibus, galea hispi valde attenuatâ villosâ incrvâ oblongâ ♂. *L. calycinum* Urv. Enum. p. 67. — *L. ochroleucum* Link Linn. 9 p. 574.

Hab. in ruderatis et rupestribus umbrosis regionis inferioris Græciae, Peloponneso (Bory! Heldr! Orph!), circa Athenas copiosum (Sprun! Boiss!), Cretâ (Heldr!), Thraciâ littorali (Grisebt!), insulis Samos (Urv.), Chios (Auch. 1686!), Rhodos (Bourg!), agro Trojano (Schmidt!), Smyrnæ (Bal. 304!), Anatoliâ boreali ad Sinopem (Tchih!).

β. micranthum. — Flores minores, calycinæ laciniæ subtenuiores.

Hab. in cultis totius Syriæ littoralis (Bl! Gaill!), totius Palestinæ (Boiss! Roth!), Cypri (Ky. 295!).

Species tubo corollino brevissimo insignis, corolla in typo 8-9 lineas longa, in varietate tantum sex lineas.

26. *L. galactophyllum* (Boiss. et Reut. in Bourg. exs. 1862) annum, caulis glabriusculis crassis ascendentibus, foliis viridibus sub lente minute puberulis reniformi-rotundatis grosse et obtusissime crenatis inferioribus petiolatis subcordatis floralibus majoribus sessilibus breviter cuneatis flores longe superantibus basi saepius albo-maculatis, verticillastris pluribus multifloris distantibus, bracteis subulatis, calycis glabriusculi fructiferi aucti dentibus a basi latiore tenuiter subulatis ciliatulis tubo sesquilongioribus demum stellatim patentibus, corollæ albæ tubo calyce breviore subcurvato annulato-piloso, fauce amplâ, lobis lateralibus subulato-appendiculatis, galeâ hirsutâ basi attenuatâ incurvâ oblongâ ⊖.

Hab. in lapidosis mobilibus alpinis, Armenia Turcica supra Gumuschkhané (Bourg!), prope Erzerum (Calv!).

Semipedale et elatius, folia floralia saepè $1\frac{1}{2}$ -2-pollices lata, corolla vix 6 lineas longa. Affine *L. moschato* var. *micranthro* a quo eximie differt foliis reniformibus obtusissimis, dentibus calycinis longioribus eximie subulatis, corollæ lobis lateralibus setaceo-appendiculatis, etc.

27. *L. Ponticum* (Boiss. et Bal. MSS.) annum, caulis glabriusculis rectis saepè ramosis, foliis viridibus puberulis obtuse crenatis inferioribus petiolatis cordatis ovato-orbiculatis floralibus sub-majoribus breviter cuneatis triangularibus flores superantibus, verticillastris paucis distinctis, bracteis subulatis, calycis hirti laciniis a basi lanceolatâ longe subulatis tubo sublongioribus demum patentibus; corollæ albidae vel carneæ tubo recto dentibus calycis subbrevisiore oblique annulato, lobis lateralibus angulatis multicis, galeâ hirsutâ oblongâ subcurvatâ ⊖.

Hab. in Ponto Lazico maritimo circa Rhizé (Bal!).

Planta 6-9-pollicaris, folia floralia pollicem longa et lata, calyx 5 lineas, corolla 9 longa. Affine *L. galactophylo* differt foliis non reniformibus, calycis hirti dentibus brevioribus, corolla longiore rectiusculâ nec valde curvatâ, lobis lateralibus non appendiculatis. Ex unico specimine mihi notum.

28. *L. purpureum* (L. Sp. 809) annum pubescens, pluricaule, caulis erectis et ascendentibus, foliis viridibus omnibus petiolatis ovato-cordatis crenatis rarius incisis, floralibus confertis breviter petiolatis flores superantibus, verticillastris in capitulum confertis inferiore interdum subdistante, bracteis brevibus subulatis, calycis laciniis a basi lanceolatâ longe subulatis tubo longioribus, corollæ purpureæ rarius albæ tubo recto calyce $1\frac{1}{2}$ -plo longiore supra basin piloso-annulato, fauce dilatatâ, lobis lateralibus subhidentatis, galeâ ovatâ obtusâ integrâ, acheniis laevibus ⊖. Ic. Fl. Dan. tab. 523.

Hab. in cultis ditionis fere totius, Peloponnesu et Cyclades (Bory), Thracia et Byzantium!, Anatolia orientalis (Bal!) et borealis (Auch. 5180!), Pontus (Bourg!), Tauria, Caucasus et Iberia (Lödeb.), Syria interior ad Aleppo (Haussk!) et littoralis (Bl!).

Ar. Geogr. Europa omnis, Sibiria.

WIEDEMANNIA (F. et M. Ind. IV. Petrop. p. 51).

Calyx campanulatus bilabiatus labio superiore unidentato dente latiore breviore, inferiore quadridentato dentibus altius connatis angustioribus. Corolla tubo exserto intus annulato-piloso, limbo bilabiato labio superiore fornicato, faux dilatata, lobi laterales appendiculati vel acutati, medius bifidus. Stamina 4 sub labio superiore parallela, antheræ divaricatim biloculares hispidae. Stylus aequaliter bifidus. Nuculae acute triquetre apice truncatæ. — Herbe annuae. — Genus calyce tantum et vix a *Lamio* distinctum in quo interdum calyx obscure bilabiatus occurrit.

1. **W. multifida** (L. Sp. p. 809 sub *Lamio*) annua adpresso et parce puberula, caulis crassiuseulis erectis simplicibus vel fere a basi opposite racemosis, foliis obtuse crenato-dentatis inferioribus petiolatis ovatis intermediis incisis, floralibus sessilibus oblongis profunde incisis vel in lacinias obtusas pinnatifidis vel partitis verticillastra superantibus, verticillastris pluribus multifloris distinctis superioribus approximatis, bracteis adpresso hirsutis ovatis et oblongo-lanceolatis calyce brevioribus, calycis labiis hirsutis tubo glabrescenti duplo brevioribus superiore breviore unidentato late triangulari mucronato, inferioris quadridentati dentibus a basi lanceolata subnatis inferioris altius connatis, corollæ purpureæ tubo calyce sesquilociore, fauce subampliata, lobis lateralibus appendice brevi linearie auctis, galea villosa obovatâ basi attenuata ⊖. *W. multifida* Bth. in D.C. Prodr. XII, p. 503.

Hab. in Armeniâ Turcicâ (Tourn., Auch. 1881!), in valle Kassuklu (Huet!) agris otiosis ad Baibut (Bourg!), in monte Ilgasdag Anatoliæ borealis (Wied!).

Semipedale vel pedale, corolla pollicaris.

2. **W. Orientalis** (F. et M. Ind. IV. Petrop. p. 51) annua breviter pubescens viscidula, caulis crassiuseulis erectis simplicibus vel a basi ramosis, foliis ovato-oblongis inferioribus petiolatis crenatis superioribus sessilibus obtuse et grosse dentatis flores superantibus, verticillastris pluribus multifloris distinctis, bracteis ovatis acuminatis ciliatis appressis calyce brevioribus, calyce hirsuto fructifero aucto bilabiato labio superiore breviore lanceolato unidentato inferioris dentibus 4 erectis altius connatis lanceolato-subulatis, corollæ purpureæ tubo calyce vix longiore oblique annulato, lobis triangulari-acuminatis, galea purpureo-vilosissimâ ovatâ ⊖. *Lamium erythrotrichum* Boiss. Diagn. Ser. I, 5 p. 26. — *W. erythrotricha* Bth. in D.C. Prodr.

Hab. inter messes Anatoliæ borealis ad Hamâqli, Safranbol, Tosia, Mersivan, Angora (Wied!), ad Tokat (Auch. 5178!), Phrygiæ ad Ouchak (Bal!) Lydiæ ad radices Mesogis (Boiss!), Pisidiæ ad Buldur (Heldr!), Lyciæ ad Elmaliu (Bourg).

216!), Cappadociae (Ky. 183! Ball!), Armeniae australis ad Karput (Noë!), Syriae (Socin!).

Eximie differt a præcedente foliis floralibus indivisis, corollâ brevi 6 lineas tantum longâ. Calyx sub anthesi 3 lineas fructifer 6 lineas longus.

MOLUCCELLA (L. Gen. 296 ex parte).

Calyx oblique campanulatus limbo ampio membranaceo reticulato-venoso. Corolla tubo inclusò intus oblique annulato, labio superiore erecto inferioris trilobi lobo intermedio latiore emarginato. Stamina sub galeâ ascendentia, antheræ in apice filamenti lateraliter pedicellatæ loculis divaricalis. Stylus bifidus. Nuculæ acute triquetrae apice truncatae. — Herbe annuae.

1. **M. laevis** (L. Sp. p. 821) annua glaberrima laevis, caule crasso erecto simplici vel inferne parum ramoso fere a basi florifero, foliis longe petiolatis a basi rotundatâ vel subcordatâ rotundatis grosse et obtuse dentatis floralibus calyces æquantibus vel superantibus, verticillastris sexfloris subdistantibus longe spicatis, bracteis subulatis spinosis patenti-deflexis calycis tubo brevioribus, calycis tubo breviter obconico limbo amplissimo membranaceo subpentagono brevissime 5-mucronato, corollâ calyeis limbo breviore albidâ ⊖. Ic. Fl. Græc. tab. 566. — Bot. Mag. tab. 1852.

Hab. in cultis et ruderatis, Anatolia occidentalis (Sibth. Auch. 1793!), Galatia (Wied!) Cilicia littoralis (Bal!), Syria littoralis et interior prope Damascus (Gail!) usque ad Libani regionem medium prope Eden (Bl!), ad Aleppo (Ky!) et Aintab (Haussk!), Babyloniam ad Bagdad (Noë!).

2. **M. spinosa** (L. Sp. p. 821) annua glabriuscula, caule erecto procero quadrangulo sulcato simplici vel inferne parce ramoso, foliis petiolatis ovatis profunde inciso-lobatis floralibus verticillastra sub-superantibus, verticillastris 6-10-floris, bracteis tenuiuer subulato-spinosis deflexis calycis tubo brevioribus, calycis tubo obconico brevi papillari-glanduloso, limbo coriaceo bilabiato valde obliquo labio superiore sublongiore erecto lanceolato longe spinoso inferiore semiobiculari-pentagono spinis 7 tenuibus alternatim brevioribus, corollæ albidae limbo calyeem superanti ⊖. Ic. Fl. Græc. tab. 567.

Hab. in Græciâ ad Parnassi radices (Sibth.), Syriâ littorali (Ehrenb! Bl!), Palestina (Auch! Boiss!).

Ar. Geogr. Hispania australis, Italia australis et Sicilia.

LAGOCCHILUS (Bunge in Bth. Lab. p. 30).

Calyx tubuloso-campanulatus subquinquenervius dentibus 5 vel lacinia subæqualibus apice spinosis. Corollæ tubus subexsertus intus annulatus, galea oblonga apice bifida, labium trifidum lobis lateralibus

brevibus erectis acutis intermedio late bifido. Filamenta basi villo coherentia sub galeâ ascendentia, antheræ per paria approximatae loculis parallelis vel divergentibus margine ciliatis. Stylus subaequaliter bifidus. Nuculæ acute triquetrae apice truncatae. — Suffrutices rarius herbæ bracteis interdum in spinas axillares acreas et persistentibus. Rami exsiccati in articulos facile secedentes fragiles.

§ 1. Spinosi. — Bracteæ annotinæ in spinas quaternas axillares persistentes acreas.

1. L. Aucheri (Boiss. Diagn. Ser. I, 5 p. 38) fruticulosus, ramis glabris erectis albis inferne denudatis simplicibus sæpius parce ramosis, foliis sub lente scalrido-hirtulis cuneiformibus subpetiolatis in lacinias anguste lineares integras vel trifidas margine revolutas apice spinulosas ad basin usque tripartitis, spinis tenuiter acerosis calyce brevioribus, calycis glabri laciñis lanceolato-linearibus acutis spinulosis margine ciliolato scabris tubo plus duplo longioribus 5.

Hab. in Persiâ (Auch. 1979!), ad Sergendeh supra Teheran (Buhse!), in montosis Firozkuh (Bode), in montibus Zagros, Kuh Nur, Sawers Persiae austro-occidentalis 8-900' (Haussk!).

Rami $\frac{1}{2}$ -1-pedales, foliorum laciñiæ elongatae vix lineam latæ, calycis 10 lineas longi laciñiæ linea vix latiores.

2. L. Kotschyanus (Boiss. Diagn. Ser. I, 12, p. 87) suffruticosus multicaulis pumilus, caulis albis tomentellis subsimplicibus, foliis ovato-cuneiformibus ad basin usque in lacinias anguste lineares margine revolutas 2-3-fidas apice spinulosas, tripartitis, spinis tenuibus rectis puberulis folio subbrevioribus, calycis laciñis elliptico-linearibus basi attenuatis acutis spinulosis tomentellis tubo ovato lanato plus duplo longioribus 5.

Hab. ad pagum Asadbar montis Elbrus occidentalis (Ky. 459!), in montosis inter Teheran et Tabris (Bge!).

Affinis *L. Aucheri* et forte ut am. a Bunge suspicatur hujus varietas, præter indumentum laciñiæ calycis paulo breviores et latiores basi attenuatae.

3. L. hispidus (Bélanger Voy. Ic. sine nuncero sub *Yermoloffia*) suffruticosus humilis totus patule hispidus, foliis subpetiolatis circumscriptione ovalis in lacinias anguste lineares revolutas muticas triseptas tripartitis, spinis folio brevioribus, calycis glabri laciñis oblongo-linearibus ciliatulis oblongis mucronulatis tubo duplo longioribus 5.

Hab. in Persiâ (Belanger), in jugo Elbrus supra Nur (Buhse!).

A *L. Kotschyanus* calycis tubo glabro laciñisque mucronulatis nec spinulosis obtusis nec acutis distinctus.

4. L. Cabulicus (Bth. in D.C. Prodr. XII, p. 515) minute piloso-hirtellus fruticulosus humilis ramis erectis, foliis circumscripti-

tione ovatis sessilibus basi cuneatis palmatim 3-5-partitis laciniis oblongo linearibus integris vel incisis obtusis raro mucronulatis, spinis tenuibus acrosis calyce vix brevioribus, calycis glabri laciniis margine ciliatulo - scabridis oblongo - linearibus obtusis mucronatis tubo duplo longioribus ȝ.

Hab. ad Konrd Kabul et Huft Kotul Affghaniæ (Griff!), in montosis ad Kalaiti prope Nichapur prov. Khorassan (Bge!).

Semipedalis, folia majora diametro vix pollicaria, spinæ majores 9-10 lineas longæ, calyx 9 lineas longus laciniis sesquilineam latis. Valde affinis *L. hispido* cuius laciniæ calycinæ quoque obtuse sed indumento pruinoso sub lente tantum perspicuo et laciniis calycis apice spinulosis distinctum.

5. ***L. insignis*** (Bélanger Voy. absque numero sub *Yermoloffiâ*) fruticulosus, ramis albidis tenuiter pubescentibus, foliis parvis cuneiformibus trifidis laciniis brevibus spinulosis margine revolutis inferiorum breviter 2-3-fidis, foliis superioribus lanceolatis tridentatis vel integris, spinis validis glabris albis folia superantibus, calycis sub lente pruinosi campanulati coriacei laciniis lineari-lanceolatis acutis spinulosis tubo subæquilongis ȝ. *L. macracanthus* F. et M. in Schrenck Enum. p. 30.

Hab. in Persiâ prope Teheran (Jenisch, Bge), prope Asadbar (Ky!).

Spinæ 10 lineas longæ validæ, calyx 6-7 lineas longus.

6. ***L. glutaceus*** (Bunge Lab. Pers. p. 74) fruticosus glaberimus ramis elongatis simplicibus albis, spinis validis, foliis coriaceis inferioribus cuneatis trilobis lobis oblongis integris spinulosis superioribus ellipticis in spinulam abeuntibus, calycis glaberrimi laciniis ellipticis mucronatis tubo tertiatâ parte longioribus ȝ.

Hab. in prov. Khorassan occidentali ad limites deserti magni inter Meiomei et Miandescht (Bge!).

Affinis *L. insigni*, elatior 2-3-pedalis, calyx glaberrimus nec pruinosus, corolla galea emarginato biloba. Calyx eo *L. cuneati* duplo major 9-10 lineas longus.

7. ***L. cuneatus*** (Bth. in D.C. Prodr. 12 p. 515) fruticosus ramosus sub lente minute velutinus, ramis rigidis, foliis minimis coriaceis subpetiolatis obovato-cuneatis integris vel apice obtuse et obsolete tridentatis 1-3 spinulosis, spinis validis acrosis folia superantibus, calycis parvi sub lente puberuli laciniis elliptico-linearibus acutis mucronulatis tubo vix longioribus ȝ.

Hab. ad colluni Khyber regni Cabulici (Griff!).

Fruticosus videtur nec ut in congeneribus basi tantum fruticulosus, spinæ majores 10 lineas longæ, folia 4-5-lineas longa, calyx parvus fructifer 5 lineas longus.

§ 2 Inermes. — Bracteæ anno insequenti non persistentes.

8. ***L. inebrians*** (Bge. pl. Lelum. p. 438) herbaceus, caule patule

hirsuto divaricatum ramoso, foliis pubescentibus inferioribus a basi cuneatâ subpetiolatâ late obovato-orbiculatis grosse lobato-dentatis muticis, floralibus diminutis cuneato-obovatis obsolete trilobis flores non æquantibus, verticillastris numerosis subcontignis, bracteis subulatis puberulis calycis tubo brevioribus, calyce pubescenti tubuloso-campanulato coriaceo in limbum sensim ampliato, dentibus a basi ovalo-triangulari acrosis tubo pluries brevioribus ♂.

Hab. in collibus aridis Turkostaniae inter Bukara et Samarkand (Lehm!). Planta trita melle vel saccharo condita inebriat. Tatarice *Bangî Diwana*, Arabice *Basul baasch*.

Semi vel 1-pedalis, calyx 6-7 lineas longus. Affinis *L. hirta* foliis oblongis non coriaceis, calyce longe hispido, nuculis convexis nec planis distincto.

9. **L. acutilobus** (Ledeb. in Eichwald Pl. Nov. tab. 23 sub *Moluccella*) glaberrimus humilis, caulis e rhizomate indurato sub-herbaceis albidis sepe breviter ramosis, foliis coriaceis cuneiformi-ovatis petiolatis 3-5-fidis vel incisis undulatis acute spinoso-dentatis, bracteis subulatis spinosis patentibus calyce triplo brevioribus, calyce coriaceo glabro inferne tubuloso sursum in limbum pentagonum sub-ampliato dentibus a basi triangulari longe acrosis tubo 4-5-plo brevioribus ♂. *L. acutilobus* F. et M. Schrenk Enum. p. 31.

Hab. ad littus orientale Caspii in Turcomaniâ (Eichw! Karel! Lehmt!).

Calyx demum 6 lineas longus.

Ar. Geogr. Songaria.

10. **L. glaberrimus** (C. Koch Linn. XXI, p. 679) glaberrimus, caulis pumilis ascendentibus inferne parce ramosis, foliis minimis longe petiolatis ambitu orbiculatis in lacinias breves cuneatas obtusas 3-5-fidas triseptis, floralibus sessilibus cuneatis 3-5-fidis calyces æquantibus, bracteis linear-lanceolatis non pungentibus calycem dimidium æquantibus, verticillastris 4-6-floris in spicam ovato-oblongam approximatis, calycis glaberrimi lacinias lanceolatis setaceo-acuminatis tubo triplo brevioribus, corolla galea hirsuta elongata apice longe bifidâ ♀.

Hab. in Tauria (Rögner ex C. Koch!) sed an locus certus? nam nullibi hanc plantam e Tauria vidi

Caules ex speciminibus incompletis semipedales, foliorum lamina 3-4 lineas diam. lata, calyx 7 lineas longus, corolla rosea 12-14 lineas longa in specimine incompleta; species ulterius investiganda.

BALL TA (L. Gen. 720 circumscr. mutatâ).

Calyx infundibuliformis vel obconicus tubo 10-nervio saepius sulcato dentibus 5-19. Corolla tubus inclusus vel vix exsertus intus piloso-annulatus, labius superius erectum oblongum inferius patens trilobum lobo medio retuso, stamina sub galea ascendentia, antheræ

e tubo exsertæ per paria approximatae biloculares loculis divaricatis distinctis. Stylus bifidus. Nuculæ apice obtusæ. — Herbæ basi interdum suffrutescentes, rarius frutices.

SECT. I. BERENGERIA Bth. Lab. p. 594. — Calycis limbus 10-pluridentatus vel crenatus. Bracteæ non spineformes.

* Calyx limbus late membranaceo-expansus breviter crenatus vel dentatus

1. **B. pseudodictamnus** (L. Sp. p. 817 sub *Marrubio*) tota adpresso albo-lanata basi indurata suffruticosa et ramosa ramis erectis simplicibus, foliis parvis breviter petiolatis ovato-orbiculatis integris vel obscure crenatis subcordatis rugosis, floralibus conformibus flores superantibus, verticillastris multifloris globosis distantibus, bracteis oblongo-spathulatis obtusis exterioribus calycis tubum æquantibus, calycis dense et adpresso lanati limbo demum expanso tubo sublongiore obtusissime quinquelobo lobulis accessoriis interdum quinis minimis, corollæ albidae villosæ tubo breviter exerto, galea bifidâ ȝ. *B. pseudodictamnus* Benth. Lab. p. 594.

Hab. in rupestribus regionis inferioris Cretæ, Cydonia, Heracleion, Malaxa Sphakia, etc. (Sieb., Heldr! Raulin!)

Pedalis, folia majora 9 lineas longa, calycis limbus demum diametro tres lineas latus quâ notâ, foliis parvis subintegris et indumento densissime cano nec tenui canescens a *B. acetabulosa* statim distinguitur.

2. **B. Damascena** (Boiss. Diagn. Ser. I, 12 p. 87) multicaulis caulinis herbaceis crassiusculis erectis flexuosis densissime et adpresso albobombycinis, foliis paucis parvis breviter petiolatis griseo-lanatis orbiculatis rugoso-crispis obtuse crenatis, floralibus conformibus calyce brevioribus, verticillastris 2-3 remotissimis multifloris, bracteis anguste oblongo spathulatis calyce brevioribus, calycis dense albo-tomentosi limbo expanso tubo breviore brevissime 10-12-crenato crenis minute spinulosis, corolle roseæ valde hirsutæ tubo inclusa galea rectâ anguste linearis bifidâ ȝ.

Hab. in collibus apricis ad Salehieh prope Damascum (Boiss! Ky. 14!), in valle Ouadi Tin deserti Ægyptiaco-Arabici (Schweinf!) in Ægypto Asagharr dicta

Pedalis, folia diametro 6-7 lineas tantum lata, verticillastra magnit. nucis parvæ, calycis limbus diametro 2 ½-3 lineas latus. A *B. pseudodictamnus* caulinis herbaceis crassioribus, foliis minoribus undulato-crispis crenatis discrepans.

3. **B. acetabulosa** (L. Sp. p. 817 sub *Marrubio*) multicaulis, caulinis basi induratis ascendentibus simplicibus palea villosulis, foliis breviter petiolatis ovatis vel orbiculatis basi subcordatis grosse et obtuse crenatis utrinque adpresso sed subtus intensius canescens lanatis rugulosis, floralibus flores superantibus et æquantibus. verticillastris parum distantibus vel contiguis longe spicatis, bracteis exte-

rioribus petiolatis oblongo-spathulatis, calycis adpresso tomentelli limbo expanso membranaceo reticulato tubo multo longiore irregulatiter multicrenato crenis muticis vel brevissime mucronatis, corollæ tubo breviter exerto, galea albâ hirsutâ bifidâ labio purpurascente ♀.
B. acetabulosa Bth. Lab. p. 593. — *Marrubium pseudodictamnus* Fl. Græc. VI, p. 50, tab. 562 non L! — Græcis *Neropapakyria* vel *Manromargo*.

Hab. in collibus regionis inferioris Laconiae, Argolidis et Atticae (Sprun! Boiss! Orph. Fl. exs. 526!), Cycladum (Sibth., Urv.), Cretæ (Raulin!) Rhodi (Bourg!), Cariae et Lydiæ (Fleisch! Boiss! Ball!).

Planta bipedalis, folia 1-1 1/2 pollicem longa, calycis limbus 9 lineas diametro latus. Floribus et forsan foliis igniarii loco sub *Lychnitidis* nomine in Atticâ utuntur.

4. ***B. undulata*** (Fresen. Mus. Senck. p. 92 sub *Marrubio*) basi suffrutescens et ramosa multicaulis, caulinis patule hirsutis glandulosis, foliis utrinque villosis majusculis breviter petiolatis rugoso-crispis orbiculatis crenatis, floralibus conformibus preter infima verticillastro brevioribus, verticillastris numerosis multiorum remotis, bracteis brevibus linearis-spathulatis, calycis pubescenti-glandulosi limbo expanso membranaceo tubo subbreviore margine undulato-crispo mutice et obsolete pluricrenato, corollæ pallide roseæ tubo inclusa, galea hirsutâ rectâ bifidâ ♀. *B. undulata* Bth. Lab. p. 595. — *Marrub. crispum* Sieb. exs.

Hab. in apricis lapidosis Antilibani circa Rascheya (Ky. 445! Gaill!), circa Damascum (Gaill!), Palestinâ circa Banias, Tiberiadem, inter Hierosolymam et Jericho (Boiss!). Arabiâ petreâ in regione Sinaiticâ (Ehr! Bové! Schimp. 423!) Arabice *Rasa*

Sesquipedalis, folia majora diam. 1-1 1/2 pollicaria, calycis limbus diam. 6-7 lineas latns. Afinis *B. acetabulosa* differt indumento parciori patulo viscidulo, limbi calycinis cum tubo proportione, bracteis angustis. Speciminum Sinaiticum folia et calyces saepè minora.

5. ***B. Aucheri*** Boiss. Diagn. Ser. I, 5 p. 39) multicaulis, caulinis inferne ramosis molliter hispidis et glandulosis, foliis utrinque dense canescenti-tomentosis rugosissimis breviter petiolatis orbiculatis cordatis obtuse multicrenatis, floralibus conformibus verticillastris superantibus et æquantibus, verticillastris multifloris remotis, bracteis linearibus tubum calycinum æquantibus, calycis hirsuti tubo costato limbo patente dimidium tubum æquante in dentes triangulari-lanceolatos sub-16 acutos alternatim immores fiso, corollæ roseæ tubo breviter exerto galea rectâ hirsutâ bifidâ ♀.

Hab. in rupestribus calcareis regionis montanae et alpine, mons Pir Omár Gudrun Kurdistaniae Persiae 3-5000' et montes Sawers, Eschker, Nur Persiæ australis (Haussk!) prope Schiras (Ky. 445!) et ad Perezend (Auch 511!).

Sesqui et bipedalis, folia diam. pollicaria, calycis limbus diam. 4 lineas latns; refert *B. hirsutam* Hispanicam cuius dentes calycinis breviores latiores et denticulati.

** Calycis limbus brevis profunde vel ad basin usque dentatus.

6. **B. Larendana** (Boiss. Diagn. Ser. I, 12, p. 88) tota molliter et patule hirsuta caulibus opposite ramosis, foliis petiolatis supra laxe hirsutis subtus canescens vel canis orbiculatis cordatis grosse et rotundatim crenatis, floralibus conformibus sensim diminutis, verticillastris multifloris superioribus approximatis, bracteis linearibus acutis calyce triplo brevioribus, calycis parvi hirsuto-viscidi tubo costato limbo calyce quadruplo breviore patente in dentes sub 10 trianguli-lanceolatos in spinulam subulatam abeentes ad medium fisso, corollæ roseæ tubo inclusu galea rectâ hirsutissimâ bifidâ ♂.

Hab. ad radices montis Karadagh Lycaoniæ ad ruinas urbis Larenda (nunc Bin bir Klissa Heldr!), Cilicia prope Anamour ad sepes (Pér!) ad vias in monte Alidagh Cappadociae 4500' (Bal!).

Planta 1-2-pedalis, folia diam sesquipollucaria, calycis limbus diam. vix 2 lineas latus quā notā a *B. Aucheri* distinguitur.

7. **B. rotundifolia** (C. Koch Linn. XXI, p. 697) mollior hirsuta viscidula canescens, caulibus erectis inferne ramosis, foliis petiolatis subcordatis ovato-orbiculatis acutiusculae crenato-dentatis floralibus flores superantibus, verticillastris multifloris dissitis, bracteis subulatis calycis tubum æquantibus, calycis breviter hirti sulcati limbo palulo tubo triplo breviore ad basin fere usqne in dentes 10 a basi anguste lanceolatâ subulatos apice glabros alternatim minores partito, corollæ tubo inclusu galea hirsutissimâ rectâ bidentatâ ♀. *B. Armena* Boiss. Diagn. Ser. II, 4 p. 54 — *B. glandulifera* Trautv. Aliq. sp. nov. N° 2! inter Erzerum et Olti (Radde!).

Hab. in Armenia Turcicâ, in valle Tchabanz (C. Koch!), circa Tortum (Calv!).

Facies *B. Larendanae* a quā limbo calycis ad basin nec ad medium in dentes angustissimos sesquilineam longos partito differt.

8. **B. macrodonta** (Boiss. et Bal. Diagn. Ser. II, 4 p. 54) pluricaulis caulibus a basi ascendentib; erectis minute et viscidule hirtis inferne parce ramosis, foliis petiolatis parvis cordatis ovato-orbiculatis obtuse subangulatis supra viridibus subtus canis floralibus diminutis, verticillastris 2-4 multifloris superioribus sœpe confluentibus, bracteis linearis-subulatis rigidis calyci æquilongis, calycis patule hirti paucicostati limbo patenti tubo æquilongo in 10 dentes lanceolatos subulato-acuminatos patenti-recurvos fere ad basin partito, corollæ purpurascens tubo brevissime exerto, galea hispidâ bilobâ ♀.

Hab. in Cappadociæ regione montanâ superiore montis Masmeneudagh (Bal. 519!).

Caules fere pedales, folia majora diam. 7-10 lineas lata, verticillastra magnitudine nucis, calyx 7-8 lineas longus dentibus tres lineas longis eis *B. rotundifoliae* multo latioribus.

9. **B. saxatilis** (Sieb. in Bth. Lab. p. 596) basi suffrutescens, caulis numerosis ascendentibus basi denudatis ramosis parce et patule hirtis, foliis patule hirtis subtus canescens petiolatis valde rugulosis crenatis cordato-ovatis, floralibus diminutis, verticillastris laxiusculis saepe subpedunculatis paucifloris, bracteis subulatis calyce triplo brevioribus, calycis patule hispiduli visciduli tubo valde costato et striato limbo calyce quadruplo breviore subpatente in dentes denos triangulares breviter mucronato-spinosos subrecurvos alternatum minores fere ad basin usque partito, corollæ pallide roseæ tubo subexerto galea hirsuta bifida ♀. *B. rugosa!* (forma culta!), *B. obliqua!* et *B. Russeliana* Bth. Lab. p. 596 et 597.

Hab. ad rupes regionis inferioris, Cilicia Trachœa circa Ermenek (Pér!) Syria borealis ad Aleppo, Aintab (Haussk!), Mesopotamia inter Orfa et Suerek (Ky. 423!), Libanus in monte Torbol supra Tripoli (Boiss! Bl!), supra Berythum (Ehrenb!) et Sidonem (Gaill!), Antilibanus prope Dinar (Gaill!) et Hasbeya (Ky. 384!), Galilea circa Panias copiose (Boiss!), in monte Ithabure (Gaill!), circa Hierosolymam (Sieb! Roth!) Folia pollicem longa, saepe multo minora, calyx cum dentibus lineam longis quatuor lineas longus, variat indumento, foliorumque magnitudine.

B. brachydonta. — Indumentum adpresso canescens stellatum, folia minuta oblonga calycis pilis stellatis canescens dentes brevissimi.

Hab ad Mezerolou Chan inter Tarsus et Gulek in Ciliciâ (Bal!)

SECT. II. **BALLOTA** Bth. Lab. p. 597. — Calycis limbus quinque-dentatus. Bracteæ non spinæformes.

10. **B. nigra** (L. Sp. 814) herbacea hirsuta vel glabrata, foliis petiolatis ovatis grosse dentatis, floralibus inferioribus flores superantibus, verticillastris plurifloris remotis saepe subpedunculatis, bracteis linearis-subulatis calyce subbrevioribus, calycis costati sulcatis hirsuti dentibus rectis subæqualibus triangulari-lanceolatis in spinulam inermem attenuatis, corollæ roseæ tubo vix exerto ♀. Ic. Rehb. tab. 7. — *B. fatida* Lam. Diet. I, p. 357.

Hab. in ruderatis totius ditionis a Græciâ et Macedoniâ ad Caucasum! Syriam! Mesopotamiam Kurdicam (Ky. 354!) Persiam borealem (Auch. 510!) Buhsel Bgef!

Ar. Geogr. Europa omnis, Africa borealis.

SECT. III. **ACANTHOPRASUM** (Bth. Lab. p. 598). — Calycis limbus 5-10-dentatus. Bracteæ rigidæ spinæformes.

11. **B. integrifolia** (Bth. Lab. p. 599) fruticosa intricatim ramosissima glabra vel adpresso et parcissime hirtula viridis, foliis minutis petiolatis ovatis obtusis integris vel obsoletissime 1-2-crenatis, floralibus calyces æquantibus, bracteis quaternis spinæformibus crassiu-

seulis patentí-recurvis calyce brevioribus, floralibus solitariis oppositis breviter pedunculatis cernuis, calyce viridi pubescenti decem-sulcato limbo patente tubo duplo breviore fere ad basin usque in dentes quinos triangulares acutos partito, corollæ albæ fauce purpureo-lineatae tubo non exerto galea villosa. *Molucella frutescens* Fl. Græca VI. p. 55, tab. 568 non L.

Hab. in rupestribus Cypri inter Cerinia et Nicosia (Sibth.), supra Chrysostomo ascendendo ad Castello Regina (Ky. 3961). V. spec. sterilem.

Aflinis *B. frutescens* = *Moluc. frutescens* L. ex Alpibus maritimis, quæ differt villositate, foliis grosse dentatis, spinis tenuissimis rectis, calycis dentibus denis a basi ovatâ abrupte et longe cuspidatis.

Species dubia negligenda.

Ballota Orientalis Presl Del. Prag., p. 80.

OTOSTEGIA (Bth. Lab. p. 601).

Calycis tubus cylindricus 10 nervius, limbus membranaceus dilatatus bilabiatus labio superiore angustiore et minore. Corolla tubus inclusus intus annulatus, galea erecta hirsutissima labio trifido lobo intermedio retuso. Stamina sub galea ascendentia, antheræ per paria sub galea approximatæ loculis divaricatis. Stylus bifidus. Nuculae apice obtusæ — Frutices interdum bracteis persistentibus et accrescentibus spinosi.

§ I. Inermes.

1. **O. Schimperi** (Bth. in n.c. Prodr. 12, p. 519 sub *Ballotæ*) fruticosa ramis elongatis strictis virgatis minutissime velutinis, foliis minimis dense et breviter tomentosis breviter petiolatis crenatis subtus rugosis floralibus calyce brevioribus, verticillastris 2-4-floris, bracteis panicis subulatis brevibus, calycis subpatule et breviter tomentosi limbo reticulato valde obliquo subbilabiato obsolete sub-10-crenulato labio inferiore duplo longiore tubum fere aequante. *Molucella microphylla* Del. Fragm. pag. 10, fig. 2! — *B. microphylla* Bth. Lab. p. 596! — *Marrubium microphyllum* Desr. in Lam. Dict. 13. p. 720.

Hab. ad rupes regionis Ouadi Gurra Arabiæ petree (Schimp. 304! sub *B. saxatili*), ad Ouadi Else (Ehrenbl!).

Rami sepe pedales et longiores, folia diam. 4 lineas tantum lata, calycis labium inferius fere tres lineas longum. Synonymon Delileanum ad *B. saxatile* Sieb. in n.c. Prodr. relatum huc certe spectat. Haec planta transitum inter *Ballotam* et *Otostegiam* præbet, huic tamen ob ramos fruticosos, habitum et limbum calycinum obscurius tamen subbilabiatum adnumeranda. Planta herbarii Juss. a Desrousseaux descripta est *O. Schimperi* calyce vix evoluto!

2. O. moluccoides (Vahl Symb. I, p. 42, tab. 14, sub *Phlomide*) fruticosa ramis novellis herbaceis elongatis virgatis minute hirtis, foliis longiusculæ petiolatis a basi breviter cuneatâ vel truncatâ ovatis obtuse dentatis minute hirtis viridibus subtus canescensibus, cymis axillaribus paucifloris subpedunculatis, bracteis setaceis brevissimis, calycis adpresso hirti limbo reticulato labio inferiore ampio rhomboidato obsolete 5-7-angulato-dentato, superiore minimo ovato integro acuto 5. *O. moluccoides* J. et Sp. Ill. Or. IV, p. 117, tab. 378 — *O. scariosa* Bth. Lab. p. 602. — *Clinopodium fruticosum* Forsk. Eg. Arab. p. 107. — *Molucella Sinaiatica* Ehrenb. MSS.

Hab. ad Ouadi Hamme Arabiæ petreæ (Ehrenb!).

Foliorum lamina 1-2-pollicaris, calycis labium inferius 4 lineas inferius sesquilineam longum. A speciminibus Arabiæ tropicæ foliis majoribus subtus minus canescensibus et cynis 2-3 nec 5-6 floris tantum differt.

Ar. Geogr. Abyssinia.

§ 2. Spinosæ. — Spinæ axillares (bracteæ annotinæ persistentes accretæ subquaternæ) quaternæ. Bracteæ hornotinæ subulatæ pungentes.

3. O. Persica (Burm. Fl. Indic. pag. 128, tab. et herb! sub *Molucella*) fruticosa ramosa ramis strictis obtuse tetragonis velutinis, spinis albidis tomentellis patulis acerosis folia superantibus, foliis parvis rugosissimis utrinque adpresso tomentosis obovato-cuneatis apice spinulose 5-7-dentatis in petiolum brevem attenuatis, fasciculis subtrifloris axillaribus oppositis, bracteis subulatis calycis tubo brevioribus, calycis tubo velutino limbo reticulato parce hirtulo labio superiori oblongo-lanceolato integro apice 1-3-spinuloso tubum subæquante inferiore multo ampliore flabelliformi spinulose 3-7-lobulato 3. Ic. Jaub. et Sp. Ill. tab. 381. — *O. microphylla* Boiss. Diagn. Ser. I, 5, p. 39. — *Ballota Persica* Bth. Lab. p. 598.

Hab. in Persiaustrali (Auch. 5123!).

Spinæ axillares semipollicares, folia cum petiolo 4-6 lineas longa, calycis labium superius 3 lineas longum inferius $4\frac{1}{2}$ longum latumque.

4. O. Kotschy (Boiss. Diagn. Ser. I, 7, p. 59) fruticosa ramosissima rami obtuse tetragonis velutinis, spinis albidis tomentellis acerosis patulis folia subæquantibus, foliis utrinque tomentellis griseis rugulosis suborbiculatis basi cuneatis obsolete circumeirca crenulato-spinulosis breviter petiolatis, fasciculis bifloris axillaribus oppositis, bracteis subulatis calycis subbrevioribus, calycis labii subæquilongis tubum velutinum subæquantibus superiore anguste oblongo-spathulato basi attennato apice obsolete et spinulose tridentato, inferiore orbiculari circumpunctata multierenato brevissime spinuloso-ciliato 3.

Hab. in declivibus Persie australis ad Daleki (Ky. 1481), ad Kotel Kumaredj 300'-300' (Haussk!).

Frutex quadrangularis, folia fere *Marrubium Alysson* 7-8 lineas diam. lata.

5. **O. Aucheri** (Boiss. Diagn. Ser. I, 5, p. 40) fruticosa inferne ramosissima ramis elongatis tenuiter striatis glabriusculis, spinis tenuibus patulis foliis brevioribus, foliis parvis carnosulis subsessilibus oblongis vel ellipticis integerrimis apice spinulosis viridibus minute puberulis, fasciculis axillaribus oppositis substrifloris bracteis subulatis eis brevioribus sussultis, calycis glabri limbo bilabiato labio superiore herbaceo ovato-triangulari integro mucronato tubo 2-3-plo breviore, inferiore membranaceo reticulato trilobo lobo intermedio subquadrato retuso tubum æquante lateralibus multo minoribus ovatis mucronulatis ȝ. Ic. Jaub. et Sp. tab. 382.

Hab. in Persiâ australi (Auch. 5122!), in Belutschia (Stocks 713!).

Rami novelli graciles sæpe pedales, spinæ axillares 4 lin. longæ, folia 6-10 lineas longa 2-4 lata, lobus medius labii calycini inferioris 4-5 lineas latus. Affinis *O. limbatæ* = *Ballotæ limbatæ* Bth. calyce distinctæ.

LEUCAS (R. BR. Prodr. 504).

Calyx tubulosus vel campanulatus striatus apice rectus vel incurvus ore æqualis vel obliquus 8-10-dentatus. Corollæ tubus inclusus, galea erecta, labium inferius trifidum lobo medio majore. Stamina sub galea ascendentia, antheræ per paria approximatae subbiloculares loculis divaricatis confluentibus. Styli lobus superior brevissimus. Nuculæ triquetrae apice obtusæ. — Genus *Ballotæ* affine, stylo, calyce non infundibuliformi diversum. Species omnes subtropicæ vel tropicæ.

1. **L. urticæfolia** (Vahl Symb. 3, p. 76 sub *Phlomide*) annua tenuiter canescens-tomentosa, caule herbaceo simplici vel parce ramoso, foliis tenuibus petiolatis ovatis grosse crenatis basi breviter cuneatis, floralibus conformibus flores longe superantibus, verticillastris 2-3 remotis globosis dense multifloris, bracteis lanceolato-linearibus acutis ciliatis calyces subæquantibus, calycis villosuli subincurvi ore obliquo limbo supra fisso inferne producto breviter ovato breviter spinuloso-lobulato ȝ. *L. urticæfolia* R. Br. Prodr. p. 504. — *Ballota Arabica* Hochst. et Steud.

Hab. in regno Mascate Arabiæ (Auch. 5144!), in Belutschia (Stocks!).

Ar. Geogr. Nubia et Abyssinia, India Orientalis.

2. **L. inflata** (Bth. Lab. p. 744) fruticosa tomentosa canescens, foliis inferioribus breviter petiolatis obovatis cuneatis apice obtusissimo 3-4-denticulatis, floralibus sessilibus cordato-orbiculariatis flores superantibus et æquantibus, verticillastris 6-10-floris numerosis distinctis, bracteis obsoletis, calycis albo-hirsuti ovato-inflati striati ore contracto æquali dentibus 10 brevibus triangulari-lanceolatis acuminatis ȝ.

Hab. in Arabiâ Sinaiticâ (Bové!).

Ar. Geogr. Nubia, Arabia tropica.

3. L. cephalotes (Spreng. Syst. 2, p. 743) annua pubescenti-birta, foliis viridibus oblongis et oblongo-lanceolatis serratis breviter petiolatis, verticillastro terminali multifloro globoso foliis supremis eo longioribus involucrato, bracteis numerosis oblongis acuminatis ciliatis dense imbricatis calyces æquantibus, calyx longe tubuloso striato sparsim hirsuto apice antrorsum subincurvo ore obliquo dentibus denis lanceolato-subulatis brevissimis vix inæqualibus ♂.

Hab. in regno Cabulico (Griff. 501 ex Bth.).

Pedalis, folia $1\frac{1}{2}$ -2-pollicaria, capitulum nuce paulo minus, calyx 7 lineas longus.

Ar. Geogr. India Orientalis.

PHLOMIS (L. Sp. 295 ex parte).

Calyx tubulosus striatus saepe sulcatus ore æquali truncato dentibus aculis vel obtusis interdum retusis. Corolla tubo inclusa intus infra medium annulatim pilosa, galea compressa, labio patente trifido. Stamina sub galea ascendentia, filamenta superiora basi in appendiculam saepe producta. Antheræ sub galea per paria approximatæ loculis divaricatis confluentibus. Nuculae triquetrae apice obtusæ vel truncatae. — Herbae et frutices, plantæ særissime tomentosæ vel lanatae.

Specierum Orientalium dispositio.

SECT. I. EUPHLOMIS Bth. Lab. p. 622. — Corollæ galea arcuata tomentosa vel villosa, labii inferioris lobus mediis latus.

§ 1. **Gymnophlomides** Bth. loc. cit. — Caules herbacei interdum basi suffruticosi. Bracteæ saepius paucæ vel nullæ, lineares vel setaceæ, molles nec pangentes.

* Calycis dentes tubo æquilongi vel longiores.

P. Bruguieri, Aucheri.

** Calycis dentes tubo duplo breviores.

+ Folia saltem radicalia cordata.

P. Nissolii, Orientalis, brevilabris, cordata, capitata.

++ Folia basi cuneata vel in petiolum angustata.

P. Armeniaca, linearis, oppositiflora, Ghilanica.

2 §. **Dendrophlomides** Bth. loc. cit. p. 539. — Frutices, bracteæ numerosæ.

* Bracteæ non pungentes.

P. fruticosa, lunariæfolia, elliptica, lanata, Lycia, floccosa, aurea, ferruginea, Bourgvi.

** Bracteæ rigidæ subulatæ subpungentes.

P. chrysophylla, viscosa.

§ 3. **Oxyphlomides** Bth. loc. cit. p. 540. — Caules herbacei e rhizomate rarius suffrutescente, bracteæ subulatæ sæpe subpungentes sæpius numerosæ.

P. anisodonta, cancellata, Chorassanica, rigida, Samia, thapsoides, Herbaventi, Mesopotamica, Persica.

SECT. II. **PHLOMIDOPSIS** Link Handb. p. 479. — Corollæ galea recta pilosa intus et margine longe barbata, labii inferioris lobus medius lateralibus vix major.

P. tuberosa.

SECT. I. EUPHLOMIS.

§ 1. **Gymnophlomides.**

* Calycis dentes tubo æquilongi vel longiores.

1. **P. Bruguierii** (Desf. Mem. Ann. Mus. II, p. 9, tab. 5) caulis humilibus simplicibus vel parce ramulosis basi induratis, foliis tomentosis subtus magis canis elevatum reticulato-rugosis oblongis et lanceolatis basi attenuatis præter insima sessilibus, verticillastris subsexfloris in capitulum ovato-oblongum foliis lanceolatis flores superantibus intermixtum confertis, bracteis subulatis tomentosis calyce brevioribus, calycis tomentosi laciniis subulatis mollibus tubo cylindrico 1 ½-2-plo longioribus corollæ purpurascente longioribus, labio superiore e lentulo, filamentis basi non appendiculatis, nuculis glaberrimis ♀.

Hab. in Mesopotamiâ inter Diarbekir et Mardin et circa Beschabur ad Tigrim (Ky. 321! et 388!), prope Mossul (Noë!), in monte Pir Omar Gadrûn Kurdistanie (Haussk!), in Persiâ prope Hamadan (Oliv., Auch. 1819!), Persiâ australi inter Seytan et Bebehau (Haussk!).

Vix pedalis, folia 2-3-pollicaria 4-6 lineas lata, calyx florifer 9-10 lineas longus, fructiferi lacinae divergentes elongate 8-9 lineas longæ.

2. **P. Aucheri** (Boiss. Diagn. Ser. I, 5 p. 36), caulis simplicibus vel at basi in florescens 1-2 ramulis auctis, foliis tomentosis supra excavatum subtus elevatum reticulato-rugosis oblongis crenulatis inferioribus cordatis et longe petiolatis, verticillastris 3-6 remotis 6-10-

floris foliis floralibus oppositis sessilibus ovatis acutatis ea superantibus suffultis, bracteis paucis linearis-subulatis calyces subaequantibus, calycis molliter et longe plumosi dentibus linearis-lanceolatis subintegralibus tubo æquilongis, corollæ flavæ calycem parum excedente galea integrâ, filamentis basi non appendiculatis, nuculis glabris ♀.

Hab. in Persia australi (Auch. 5161!), in collibus ad lacum salsum Nemek Darja prope Schiraz (Ky. 454!), in montibus Bachtiarieis (Bode ex Bge).

Pedalis vel sesquipedalis, folia inferiora 3-4-pollicaria, spica laxa 6-8-pollicaris, calyx 8-10 lineas longus. A *P. Bruguieri* verticillastris dissitis, foliorum floralium formâ, dentibus calycinis brevioribus etc. distincta, facies *P. crinita* dentibus calycinis brevissimis, etc., aliena.

• Calycis dentes tubo multo breviores.

+ Folia saltem radicalia cordata.

3. ***P. Nissolii*** (L. Sp. 819) tota dense subfloccoso-pannosa, caule paniculato, foliis ovato-oblongis plus minus rugosis radicalibus longe petiolatis profunde cordatis, verticillastris subsexfloris foliis floralibus ellipticis subsessilibus ea subaequantibus suffultis remotis laxè spicatis, bracteis paucis subulatis brevibus, calycis floccoso-pannosi angulati dentibus brevibus ovatis obtuse carinato-plicatis, corollæ flavæ galea subproductâ obsolete tridentata, filamentis majoribus basi subappendiculatis, nuculis glabris ♀.

Hab. in collibus, Lydia prope Smyrnam (Fleisch! Bal. 314!), ad colles Laodiceæ (Boiss!), Phrygia prope Ouchak (Bal!), Galatia (Wied!), Cappadocia prope Caesaream (Bal!), Cilicia Trachea in valle Djoksu (Heldr!) et circa Ermenek (Peronin!), moute Amman Syriæ (Ky. 78! sub *P. Armeniacæ*).

β. leptorrhacos. — Indumentum tenuius adpressum non floccosum, folia siepe angustiora, calycis dentes acutius carinati acutiores. — *P. Syriaca* Boiss. Diagn. Ser. I, 12 p. 89. — *Ph. Tarsensis* Boiss. in Ky. exs.

Hab. in Syriâ circa Aleppo et Antiochiam (Boiss!), in Libano (Bl!), Anti-libano et ad Damascum (Gaill! Ky. 236!), Ciliciâ prope Tarsus (Ky. Suppl. 255!), Lyciâ ad Elmalu (Bourg. 202!).

Formæ transitoriae et distributio geographicæ docuerunt *P. Syriacam* potius ut varietatem *P. Nissolii* habendam esse.

4. ***P. Orientalis*** (Mill. Diet. N° 9) tota tomentosa vel subfloccoso-lanata, caulis basi induratis ascendentibus simplicibus vel ramiculis binis floralibus auctis, foliis supra saepè glabrescentibus rugulosis ovato-oblongis petiolatis cordatis, floralibus oblongo-lanceolatis cuneatis subpetiolatis flores superantibus, verticillastris 8-10-floris dissitis, bracteis linearis-setaceis brevibus paucis interdum nullis, calyce floccose rarius adpresso tomentoso cylindrico angulato dentibus lanceolatis acutis vel obtusiusculis tubo circ. triplo brevioribus, corollæ flavæ calyce duplo longiore limbo ampio galea obscure tridentata, filamentis non appendiculatis, nuculis glabris ♀. *P. Armeniaca* Blh. in D.C. Pradr. et Auct. plur. non Willd. — *P. Olivieri* Blh. Lab. p. 624.

Hab. in siccis regionis inferioris et montanæ, Caria prope Geyra (Boiss! Pint!), Galatia (Wied!), Armenia (Auch. 4814!) et in prov. Musch (Ky. 445!), Mesopotamia et Abyssinia (Haussk!), Kurdistania Persica et Persia australis montes ubi a 3000' ad 1000' ascendit (Ky! Haussk!), Persia borealis (Szov! Ky! Buhse! Bge! etc.), prov. Talysh (C.A. M., Hoh!).

β. brachyodon. — Dentes calycini breviores et obtusiores. — *P. Armeniaca* var. *brachyodon* Boiss. Diagn. Ser. I, 12 p. 88.

Hab. in Palestinâ inter Hierosolymam et Jericho (Boiss!).

Species a *P. Nissolii* inflorescentiâ simplici vel subsimplici nec paniculata calycisque dentibus longioribus angustioribus distincta. Varietas *brachyodon* inter utramque intermedia videtur sed caule humili simplici tamen huic adnumeranda. Willdenowius ex descriptione et herbario sub suâ *P. Armeniacâ* non hanc sed speciem foliis angustis basi cuneatis habuit.

5. *P. brevilabris* (Ehrenb. MSS.) humilis multicaulis basi suffrutescens et procumbens, caulis erectis simplicibus, foliis parvis adpressissime tomentosis utrinque sed subtiliter praesertim canis reticulato-rugosis oblongis subcordatis petiolatis, floralibus subsessilibus ellipticis sensim diminutis inferioribus flores superantibus, verticillastris paucifloris remotiuseulis, bracteis brevibus setaceis reflexis, calyce patule tomentoso-hirsuto angulato dentibus lanceolatis tubo subtriplo brevioribus, corollâ flavâ calyce subduplo longiore galeâ obscure tridentatâ, tubo intus parce piloso, filamentis non appendiculatis, nuculis glabris ♀. *P. Armeniaca* var. *microphylla* Boiss. herb.

Hab. in regione subalpinâ Libani ad Cedros (Boiss!), ad Dimam et Hasrun (Gaill!), ad Gisr el Hajar (Ehrenb!), ad El Toghra inter Sidonem et Damascum (Gaill!), Antilibano in regione inferiori Hermonis (Boiss! Gaill!).

Semipedalis rarius pedalis; a formis *P. Orientalis* differre videtur foliis parvis (1 rarius 2 poll.) adpressissime et tenuiter tomentosis rugosissimis nec floccosis, subcordatis vel truncatis nec profunde cordatis; bracteae numerosiores, folia floralia magis diminuta.

6. *P. cordata* (Boiss. et Ky.) adpresso tomentosa cano-flavida, caulis ascendentibus, foliis parvis ovatis etiam floralibus cordatis inferioribus longiusculæ cæteris breviter petiolatis, verticillastris 4-6-floris remotissimis foliis floralibus superatis, bracteis setaceis paucis minimis, calyce ovato dense lanato dentibus a basi lanceolatâ setaceis plumosis tubo subæquilongis, corollâ flavâ calyce 2½-plo longiore galeâ obscure tridentatâ, filamentis non appendiculatis, nuculis glabris ♀.

Hab. in graminosis porphyricis montis Tolos Amani supra Arsus Syriæ borealis 4500' (Ky. 400!).

Folia inferiora pollicem longa, verticillastri 2-4 remotissimi, calyx eis specierum affinum brevior 5 lineas longus valde lanatus, laciniis longis setaceo-attenuatis insignis, quibus notis si constantes sint a *P. brevilabri* differt.

7. *P. capitata* (Boiss. Diag. Ser. II, 4 p. 96) humilis multicaulis basi suffruticosa caulis erectis subsimplicibus, foliis adpressissime

albo vel flavido-tomentellis reliculato-rugosis parvis ovato-oblongis radicalibus cordatis longe petiolatis, caulinis subsessilibus cito deciduis, verticillastris in capitulum terminale ovatum densum confertis inferiore interdum remotiusculo, foliis floralibus ellipticis flores subaequantibus, bracteis paucis subulatis brevissimis, calycis adpresso tomentosi angulati dentibus lanceolatis obtusiuseculis tubo subtriplo brevioribus, corollæ luteæ calyce duplo longioris limbo ampio galea integrâ, nuculis glabris ♀. *P. Haussknechti* Bge Lab. Pers. p. 78.

Hab. in regione alpinâ Tauri Cilicici in valle Gusguta 650' (Ky. 451! Bal!), montibus Akkerdag et Berytdagh Cataoniæ (Haussk. exs. sub *P. Armeniaca microphylla*), monte Aslandach Cappadociae (Bal!).

Semipedalis rarius pedalis, folia parva et indumentum adpressum *P. brevilabris* sed folia profundius cordata et inflorescentia capituliformis.

++ Folia cuneata vel in petiolum angustata.

8. *P. Armeniaca* (Willd. Sp. 3 p. 19 et herb! non aliorum) basi suffruticosa multicaulis, caulinis saepius humilibus simplicibus rarius rameulo 1-2 florali brevi auctis, foliis adpresso tomentosis subtus præsertim canescensibus vel canis ellipticis vel lanceolatis obtusis vel acutiusculis basi cuneatis inferioribus longe petiolatis, floralibus subsessilibus flores superantibus et aequantibus, verticillastris 4-6 floris 2-4 inferioribus distantibus superioribus magis approximatis, bracteis paucis linearibus deflexis, calycis cylindrici tomentosi dentibus lanceolatis acutis tubo 3-4-plo brevioribus, corollâ flavâ calyce duplo longiore galea integrâ, nuculis glabris ♀. *P. lanceolata* Boiss. et Hoh. Diag. Ser. I, 5. p. 36.

Hab. in siccis regionis montanæ et alpinæ, Tmolus Lydiæ supra Philadelphiam (Boiss!), Bulghasdagh Phrygiæ (Bal!), Galatia (Wiedem!). Lycia prope Elmali (Bourg. 2001), Pisidia et Lycaonia (Heldri), Cilicia Trachea prope Ermenek (Perl), Karamasdagh Cappadociæ et Ssodagh Cataoniæ (Haussk!), mons Gara Kurdistaniæ (Ky. 383! 385!), Armenia ad Erzerum (Tourn., Calv!) et prope Baibut (Bourg!).

Species ab omnibus formis *P. Orientalis* foliis angustis elongatis basi cuneatis nec cordatis distincta; folia variant 5-12 lineas lata 2-6-pollicaria.

9. *P. linearis* (Boiss. et Bal. Diagn. Ser. II, 4 p. 46) multicaulis, rhizomatis ramis radicantibus, caulinis saepius humilibus simplicibus, foliis adpresso tomentellis pallide virentibus tenuiter rugulosis linearibus obtusis in petiolum longe angustatis, verticillastris 2-4 distantibus foliis floralibus subsessilibus linearibus longe superatis, bracteis linear-lanceolatis calyce brevioribus, calycis dense et patentim lanati dentibus lanceolatis acutis dimidio tubo sublongioribus, corollæ flavæ calyce plus duplo longioris labio integro, filamentis non appendiculatis, nuculis apice pilosis ♀. *P. angustifolia* Boiss. et Bal. exs. schedulae errore.

Hab. in regione alpinâ inferiore montium Masmeneu-lagh et Aslandach

Cappadociæ (Bal !), supra Seytun et in monte Berydagħ Cataoniæ 7-8000 (Haussk !), in valle Merga Sauk Armeniæ australis (Ky. 401 !).

Planta $\frac{1}{2}$ -1-pedalis, folia 3-6-polices longa 3-6 lineas lata. Affinis *P. Armeniacæ* a quā differt indumento adpressiori et tenuiori foliorum, dense lanato calycis, calycinis dentibus tubi proportione longioribus, nuculis apice pilosis

10. *P. oppositiflora* (Boiss. et Haussk.) caulibus tenuibus opposite et stricte ramosis glabriusculis, foliis tenuiter tomentellis canescentibus rugosis inferioribus anguste lanceolatis obtusiusculis in petiolum longe angustatis, verticillastris bifloris remotis laxissime et longe spicatis foliis anguste linearibus calyce longioribus suffultis, bracteis nullis, calyce obconico-cylindrico dence albo-panno vix angulato dentibus inæqualibus trianguli-lanceolatis tubo sextuplo brevioribus, corollâ flavâ calyce vix duplo longiore galeâ integrâ, nuculis apice pilosis ♀.

Hab. in Cataoniæ planicie Palanga owa et in monte Scherdagh supra Albistan (Haussk !). Fl. Aug.

Pedalis vel sesquipedalis, foliorum radicalium limbus bipollicaris 4-5 lineas latus, calyces 7 lineas longi. A formis stenophyllis *P. tinearis* caulibus paniculatis, verticillastris bifloris, calycis magni dentibus brevibus distincta.

11. *P. Ghilanica* (Koch Linn. XXI, p. 698) herbacea, caulis ascendentibus simplicibus, foliis rugulosis stellatim tomentellis superne virentibus subtus cinereis radicalibus elliptico-lanceolatis obtusiusculis obtuse crenatis in petiolum longum attenuatis, floralibus late linearibus acutis integris flores superantibus, verticillastris paucis distantibus 6-10-floris, bracteis linearis-setaceis plumosis calyces æquantibus, calycis longe plumosi dentibus trianguli-lanceolatis acutis tubo 4-5-plo brevioribus, corollâ flavâ calyce duplo longiore galeâ incurvâ retusâ, nuculis.... ♀.

Hab. in alpibus prov. Ghilan Persiæ bor. prope Gambu (Pallas in herb. Willd. sub *P. lychnitide*!).

Species ex unico specimine nota, folia radicalia 3 poll. longa 10-12 lineas lata petiolo bipollicari, caulis vix pedalis verticillastris tribus remotis, calyx sex lineas corolla 12 longi. Species bracteis longis plumosis a precedentibus diversa.

§ 2. **Dendrophlomides.**

* Bracteæ non pungentes.

12. *P. fruticosa* (L. Sp. p. 818) fruticosa, adpresso cano-tomentosa, foliis oblongis subtus rugosis et magis canis basi rotundato-cuneatis inferioribus breviter petiolatis, verticillastris binis remotis vel unico terminali dense multifloris foliis floralibus binis sessilibus fere superatis, bracteis tomentosis adpressis ovato-ellipticis vel lanceolatis acutis, calyce hispido angulato truncato dentibus brevissimis in acumen parvum subulatum patens vel subrecurvum abeuntibus, corollæ luteæ galeâ retusâ, filamentis binis basi appendiculatis, nuculis glabris ♂. Ic. Fl. Græc. tab. 563. Rchb. Ic. tab. 21.

Hab. in collibus regionis inferioris, Zacynthus (Marg !), Coreyra (Boiss !), Peloponnesus (herb. Fauché !), Attica (Sprun !), Thracia et Bithynia (Griseb., Noë !), Creta (Raul !).

Ar. Geogr. Sardinia, Italia australis.

13. *P. lunariæfolia* (Sibth. et Sm. Prodr. I, p. 414) fruticosa adpresso caescenti-virens, foliis oblongis obtusis inferioribus utriusque canis minoribus petiolatis basi truncatis, ramorum hornotinorum subsessilibus basi cuneatis supra virentibus subtus canescentibus rugulosis, capitulo sphaerico terminali multifloro foliis binis deflexis lanceolatis eo longioribus suffulto. bracteis imbricatis membranaceis ovatis mucronatis elevatim nervulosis margine tuberculato-ciliatis cæterum glabris, calyce sub lente minute et stellatim puberulo virenti truncato lobis 5 brevibus latis retusis, mucronibus brevibus subulatis rectis ex sinubus, corollâ flavâ calyce $2\frac{1}{2}$ -plo longiore galeâ retusa, filamentis longioribus appendiculatis, nuculis glabris ♂. *P. imbricata* Boiss. in Bourg. pl. Lyc. 1860.

Hab. in saxosis Duden prope Elmalu Lyciae (Bourg. 296!), circa Anamour Ciliciae (Peronin !), in insulâ Rhodo inter Chrysou et Slewra (Ky. 678 sub *P. lunariæfolia*!).

Rami pedales monocephali, capitula magna, calyx 8-9 lineas longus. Affinis *P. fruticosæ*, ab eâ indumento tenui, bracteis membranaceis ciliatis glabris calyceque distincta. Hæc ex charaktere bractearum lataram omnino videtur esse planta in Prodromo Floræ Græcæ descripta, sed loci (Peloponnesus et Athos mons) ibidem quâdam confusione indicati ad *P. Samiam* spectant.

14. *P. elliptica* (Bth. Lab. p. 626) fruticosa ramosa, ramis novellis brevibus simplicibus tomentosis, foliis parvis ellipticis obtusissimis rugosis breviter petiolatis suprà virentibus subtus canis, verticillastro multifloro capituliformi foliis floralibus conformibus flores æquantibus suffulto rarius binis subdissitis, bracteis linearibus rigidis tomentellis numerosis calyces æquantibus, calycis obconicæ tomentosi truncati dentibus brevissimis latis obtusis vel retusis mucronulatis, corollâ lilacinâ calyce duplo longiore, galeâ obsolete et obtuse tridentalâ, tubo infra faucem valde hirsuto, filamentis majoribus basi appendiculatis, nuculis glabris ♂.

Hab. in apricis faucium montis Sabst Buschom prope Schiraz (Ky. 372!), in Persiâ loco ignoto (d'Arcy ex Bth.). Fl. Maio.

Facies et folia parva (pollicaria) *P. lanatae*. Calyx 6 lineas longus, capitulum magn. nucis.

15. *P. lanata* (Willd. Enum. Suppl. 41) fruticosa divaricatim ramosa, ramis novellis ferrugineo subfloccoso-lanatis, foliis parvis ovatis et ellipticis obtusissimis rugosis superne virentibus subtus albo-lanatis, verticillastro unico terminali vel binis remotis 6-8-floris foliis floralibus binis conformibus ea subæquantibus suffultis, bracteis oblongo-lanceolatis acutis adpresso calyce brevioribus, calycis tomentoso-lanati truncati dentibus deceem brevissime late et obtuse trian-

gularibus, sinubus 5 brevissime mucronatis, corollâ flavâ calyce duplo longiore galeâ integrâ, filamentis longioribus basi appendiculatis nuculis apice hirsutis ♀. Ic. Janb. et Sp., Ill. Or. tab. 411 (mala). — *P. microphylla* Sieb. Reis. Cret. II, p. 319, tab. 8. — *P. parvifolia* Presl. Del. Prag. p. 86.

Hab. in rupestribus Cretæ (Tourn., Sieb!), ad sinum Spinalunga (Heldr!), in regione inferiore montium Sphacioticorum (Raul!).

Folia majora pollicaria, calyx 5 lineas longus ob dentes quinos profunde bifidos decemdentatus.

Ar. Geogr. Calabria.

16. *P. Lycia* (D. Don in Tayl. Ann. Nat. Hist. 7, p. 458) fruticosa ramosa adpressissime et stellatim tomentosa pallide virens, foliis parvis rugulosis subtus pallidioribus breviter petiolatis cordatis oblongis, floralibus subsessilibus verticillastra superantibus, verticillastris 1-2 parvis densissimis, bracteis margine longe albo-lanatis inter se lanâ connexis incurvis adpressissimis lanceolatis acuminatis subpungentibus calyce subbrevioribus, calyce stellato-tomentello truncato et ore longe luteo-barbato dentibus brevissimis latis truncatis retusis in spinulas brevissimas patentes subuncinatas abrupte abeuntibus, corollâ aureâ calyce duplo longiore, filamentis basi appendiculatis, nuculis glabris ♂.

Hab. in sylvis montanis, Lycia septentrionalis (Fellowes ex Don), in Anatoliâ occidentali loco speciali non indicato (Dr. Löw! ex herb. Koch).

Facies et folia parva (pollicem rarius sesquipollicem longa). *P. lanatae* sed basi eximie cordata et indumentum totius plantæ adpressius, verticillastra densiflora magnitudine inter nucem et avellanam media. Species insignis bracteis adpressissimis planis dorso tomentellis et virentibus margine albo-lanatis. Plantam cl. Don non vidi sed ejus descriptio ad eam herbarii cl. Koch optime quadrat.

17. *P. floccosa* (Don in Bct. Reg. 45, tab 1300) fruticosa floccoso-lanata, foliis rugosissimis ovato-oblongis basi cordatis petiolatis, floralibus breviter petiolatis basi truncatis verticillastro sublongioribus, verticillastris 3-5 remotis multifloris, bracteis linearibus albo-lanatis calyci aequilongis, calycis albo-lanati apice truncati dentibus brevibus latis truncatis in mucronem longum subulatum apice uncinatum glabriusculum abeuntibus, corollæ flavidæ ad labium inferius sœpe purpurascens calyce subdupo longioris galeâ retusâ, filamentis majoribus basi appendiculatis, nuculis g'abris ♀. *P. Samia* var. *bicolor* Viv. Lib. 30, tab. 45. — *P. bicolor* Bth. Lab. 629.

Hab. in Egypto prope Alexandriam versus Abusir (Ehrenb!), ad Mariut (Le-tourn! Planta!).

Folia inferiora 3-4-pollicaria, verticillastri 2 pollices distantes, calyx 8 lineas longus dentibus subulatis patentibus uncinatis insignis.

Ar. Geogr. Africa boreali-orientalis in Cyrenaica.

18. *P. aurea* (Decaisn. Pl. Sinaic. p. 13) fruticosa tota dense et

floccosa aureo-tomentosa, ramis novellis longis crassis, foliis oblongis obtusis inferioribus basi truncatis vix subcordatis breviter petiolatis, floralibus sessilibus oblongo-lanceolatis sensim diminutis, verticillastris multifloris valde remotis, bracteis oblongo-linearibus et linearibus obtusis adpressis calyce subbrevioribus, calyce aureo-tomentoso apice truncato dentibus 5 brevissime triangulari-subulatis, corollæ aureæ galeâ integrâ, filamentis longioribus basi appendiculatis, nuculis glabris ♀. *P. angustifolia* et *P. flavescentia* Mill. Dict. N° 2 et 10 ex cl. Bth.

Hab. in vallis Sinaiticis (Bové! Ehrenb! Schimp. 313! Boiss!). Arabis Aroa.

Frutex 3-4-pedalis gregarius, verticillastra nuce majora 3-4 pollices distantia, calyx 8 lineas longus. Synonyma Milleriana ob descriptiones mancas ex plantis cultis et patriam indicatam nullam mihi valde dubia videntur, ad varias species flavifloras æquo jure referenda et ideo nomen specificum cl. Decaisne præposui.

19. *P. ferruginea* (Ten. Fl. Nap. II, p. 36, tab. 57) fruticosa ramis novellis elongatis laxe floccoso-tomentosis, foliis rugosissimis supra velutinis virentibus subtus canescens vel dense lanatis canis inferioribus longe petiolatis cordatis, verticillastris 2-3 valde dissitis multifloris foliis binis subpetiolatis subrhombatis eis longioribus suffultis, bracteis lanceolatis vel subulatis acutis adpressis calyci subæquilongis, calycis viridis setis basi stellato-pubescentibus hispido dentibus brevissimis latis truncatis abrupte in mucrones subulatos dente ipso saepius longiores rarius breviores subinæquales abeuntibus, corollæ luteâ calyce duplo longiore, galeâ subintegrâ, filamentis longioribus breviter appendiculatis, nuculis glabris ♀. *P. Cretica* Presl Del. Prag. — *P. fruticosa* Sieb. exs! non I.

Hab. in regione inferiori Cretæ (Sieb!), prope Cydoniam (Wiss!), monte Malevo Laconiae 3000' (Orph. Fl. Gr. exs. 5231 sub *P. fruticosa* var.), in Insula Rhodi rupibus Filierimo prope Trianda (Bourg. exs. sub *P. viscosa* var!).

Facies *P. viscosa* sed folia minora crassiora rugosa, culyces breviores (6-7 lin. longi) mucronibus brevioribus, bracteæ non subpungentes. Variat in Creta ut in Italiâ indumento plus minus denso.

Ar. Geogr. Italia australis.

20. *P. Bourgæi*, fruticosa virens patule pubescens superne glanduloso-viscidula, foliis tenuibus vix rugosis ovato-oblongis cordatis inferioribus petiolatis superioribus subsessilibus verticillastro longioribus, verticillastris 1-3 valde remotis, floribus subpedicellatis, bracteis numerosis plumosis subulatis patentibus deflexis a medio sursum curvatis calyci subæquilongis, calycis hirsuti virentis apice truncati dentibus latis brevissimis truncatis in subulas longas patulas inæquales abeuntibus, corollæ flavæ galeâ subtridentata, filamentis subappendiculatis, nucibus glabris.

Hab. in Pamphyliæ rupibus maritimis prope Adalia (Bourg. exs. 2711 sub *P. viscosa*).

Affinis *P. viscosa* specifice differre videtur bracteis non adpressis rigidis subpungentibus sed setaceo-subulatis patentibus curvatis, calyx brevior dentibus tenuioribus patentibus.

** Bracteæ rigidæ subulatæ subpungentes.

21. **P. chrysophylla** (Boiss. Diagn. Ser. 1, 12, p. 89) fruticosa tota adpresso aureo-tomentosa, ramis novellis elongatis, foliis subtus rugosissimis petiolatis late et breviter cordatis obtusissimi cordatis, verticillastro terminali vel binis remotis multifloris foliis floralibus ea æquantibus vel superantibus suffultis, floribus subpedicellatis tribracteatis, bracteis anguste linearibus rigidis tomentosis adpresso calyce brevioribus, calycis aureo-pannosi angulati sulcati truncati dentibus latis brevissimis retusis angulis in spinulas 5 minitissimas excurrentibus, corollæ aureæ calyce plus duplo longioris galea obsoleta retusa, filamentis majoribus basi appendiculatis, nuculis glabris ♀.

Hab. in regione subalpinâ Libani circa Eden et Hasrun (Bl!), regione inferiori Antilibani in Hermone supra Rascheya et in montibus Zebdani (Boiss! Gaill!).

β. oblongifolia. — Folia oblonga basi subcuneata, verticillastri minores pauciflori.

Hab. in Amano supra Arsus (Ky!).

Facies et indumentum aureum *P. aurea* sed adpressum nec floccosum, ab eâ insuper floribus subpedicellatis et bracteis rigidis linearisubulatis *P. viscosa* discrepans. Varietas β est prob. tantum forma autumnalis mihi tantum fructifera nota et ejus folia ramorum sterilium minora et augustiora sola adsunt.

22. **P. viscosa** (Poir. Diet. 5. p. 271) fruticosa parce et breviter tomentella superne præsertim viscidula, ramis novellis elongatis, foliis majusculis tenuibus vix rugosis subtus canescentibus ovatis et oblongo-lanceolatis inferioribus petiolatis valde cordatis, floralibus subsessilibus cuneato-lanceolatis acuminatis flores longe superantibus, verticillastris multifloris 2-4 remotissimis, floribus subpedicellatis subtribracteatis, bracteis lanceolato-linearibus rigidis subpungentibus incurvis adpresso ciliatis calyce subbrevioribus, calycis viridis stellatim hispiduli visciduli dentibus a basi late ovatâ subulato-acuminatis valde inæqualibus binis elongatis, corollæ flavæ calyce duplo longiore galea bifidâ, filamentis binis basi appendiculatis, nuculis glabris ♂. *P. vires* D.C. Cat. Monsp. p. 132. — *P. glandulosa* Schenk Pl. Nov. p. 20. — *P. Herbaventi* var. Russell Aleppo, tab. 16. — *P. lunatifolia* var. *Russeliana* Bot. Mag. tab. 2842. — *P. Russeliana* Lagasca ex Blh?

Hab. in siccis regionis inferioris Syriæ bor. (Auch. 1812!), in monte Amano (Ky. 38!), ad radices Cassii (Boiss!), Syriæ littorali et regione infer. Libani (Bl! Gaill!), Palestinâ in Samariâ (Boiss!), ad Hierosolymam (Roth!), Cypro (Ky!).

β. angustifolia. — Folia lanceolata 7-10 lineas basi tantum lata,

verticillastra minora, calycis dentes minores minus inaequales. — *P. longifolia* Boiss. et Bl. Diagn. Ser. II, 4. p. 47.

Hab. in Libano circa Solima (Bl!), prope Beyruth (Ehrenb!).

Folia inferiora in typo 2-4-pollicaria sesquipollicem et amplius lata, calycis dentes iini inferiores $2\frac{1}{2}$ lineas longi. Species a *P. ferruginea* floribus subpedicellatis et bracteis rigidis basi incurvis distincta. Specimen *P. Russeliana* a cl. Lagasca in herb. Mus. Brit. descriptum ibi nunc deest.

§ 3. Oxyphlomides.

23. *P. anisodonta* (Boiss. Diagn. Ser. I, 5, p. 37) a basi suffrutescenti pluricipiti herbacea pluricaulis, caulis erectis tomentellis simplicibus, foliis supra virentibus subtus pallescentibus tomentosis inferioribus longe petiolatis a basi rotundata vel subcuneatâ ovatis vel oblongis obtusis crenatis, caulinis breviter petiolatis, floralibus subsessilibus oblongo-lanceolatis deflexis verticillastro longioribus, verticillastris 2-4 remotis, bracteis numerosis subulatis basi sursum curvatis ciliatis calyces æquantibus, calycis hirsuti dentibus brevibus ovato-triangularibus saepe retusis in aristulas subulatas ciliatas abeuntibus inferioribus longioribus dimidium tubum æquantibus, corollâ roseâ, filamentis longioribus basi appendiculatis, nuculis apice pilosis ♀.

Hab. in regione alpinâ Persiae borealis ad Djulfeck (Auch. 5159!), in jugo Elbrus (Ky. 483! Buhse!), australis in montibus Kuh Daëna et Kuh Delu (Ky. 519! et 797!), montibus Nur, Savers et alpibus Bachtiaricis (Hausk!), alpibus Avroman, Schahu, Dalechani Kurdistaniæ Persicae 9-12000' (Hausk!).

Variat $\frac{1}{2}$ - $1\frac{1}{2}$ -pedalis, foliorum inferiorum lamina 1-2-pollicaris, calycis tubus 5 lineas circ. longus. Variat quoque floribus albis et foliis subtus magis canis.

24. *P. cancellata* (Bge Lab. Pers. p. 76) herbacea caule humili simplici tomentello, foliis rugosis adpresso tomentosis subtus magis canis petiolatis oblongo-lanceolatis crenatis, floralibus breviter petiolatis acutis flores multoties superantibus, verticillastris paucis remotis 6-10-floris, bracteis subulatis hispidis calyce sublongioribus, calycis tubo stellato-tomentoso dentibus subulatis valde inaequalibus inferioribus recurvo-deflexis tubo longioribus cæteris triplo brevioribus, corollâ albida calyce duplo longiore tubo intus hirto, filamentis appendiculatis, nuculis apice stupposis ♀.

Hab. in rupestribus montium ad occidentem urbis Schahrud Persiae bor. (Bge!).

Semipedalis, folia inferiora tripollicaria. Valde affinis *P. anisodontæ* cuius forsitan varietas, calycis tubus subbrevior 4 lineas longus et dens anticus fere 5 lineas longus. Galea ex cl. Bge antice rectilinea nec dente aucta.

25. *P. Chorassanica* (Bge Lab. p. 76) herbacea adpresso stellato-tomentosa, caule humili simplici fere a basi florifero spiciformi, verticillastris numerosis approximatis 6-8-floris, foliis subtus magis

canis breviter petiolatis oblongo-lanceolatis obtusis crenato-dentatis floribus multoties longioribus, bracteis subulatis tomentosis calyces æquantibus, calycis subfloccoso-tomentosi dentibus subulatis inæqualibus anticis tubo vix brevioribus, corollâ albidâ calyce subdupo longiore, nuculis apice stupposis ♀.

Hab. in arvis prope Tas non procul a Meschhed prov. Khorassan (Bge !).

Semipedalis, folia tres pollices longa 7-9 lineas lata infimis exceptis in axillis florifera. Affinis binis præcedentibus sed ob verticillastra approximata inflorescentia spiciformis, calycis tubus 4 lineas longus, laciñæ anticæ 3 lineas longæ.

26. *P. rigida* (Labill. Dec. Syr. III, p. 15, tab. 10) herbacea, caule simplici patentim hispido, foliis rugosis tomentosis pallide virentibus oblongis et oblongo-lanceolatis inferioribus longe petiolatis, floralibus subsessilibus flores superantibus, verticillastris 3-5 magnis multifloris remotiusculis, bracteis patule hispidis a basi anguste lanceolatâ subulatis rigidis subpungentibus adpressis calyces æquantibus, calycis hirsuti angulato-sulcati dentibus a basi breviter ovalâ truncatâ subulatis rigidis inferioribus sublongioribus, corollâ rosâ calyce $2\frac{1}{2}$ -plo longiore labio inferiore longissimo, tubo fere toto intus hirta filamentis non appendiculatis, nuculis apice hirtis ♀.

Hab. in siccis arenosis regionis montanae, valles Tauri Isaurici inter Karaman et Ermenek 3500' (Heldr!), Gumgum Armeniæ australis (Ky. 679!), mons Akkerdaghi Cataoniae prope Marasch (Haussk!), Mesopotamia inter Orfa et Suerek (Ky. 81!) et ad Derbent i Basian (Haussk!), Antilbanus prope Damascum (Lab!), in monte Gebel Scheiek (Ehren! Gaill!), in monte Garbi (Ky. 94!).

Species splendida pedalis et sesquipedalis verticillastris ovo majoribus.

27. *P. Samia* (L. Sp. 819), caule herbaceo pubescente valde elato simplici vel opposite ramoso paniculato, foliis amplis ovato-oblongis valde cordatis acutiusculis subtus canescens radicalibus longe petiolatis, floralibus subsessilibus triangulari-rhombeis acutis vel acuminatis flores superantibus, verticillastris 3-7 multifloris distantibus, floribus subpedicellatis, bracteis ciliatis subulatis rigidis subpungentibus calyces æquantibus, calycis patule glanduloso-visciduli dentibus lanceolatis pungentibus tubo subtriplo brevioribus fructiferis patentibus, corollâ sordide flavidâ demum purpureâ tubo intus late piloso-annulato, filamentis appendiculatis, nuculis glabris ♀. Ic. Vent. Choix t. 4. — Fl. Græc. tab. 564. — *P. superba* C. Koch Linn. XXI, p. 699! — *P. lunariaefolia* Pr. Fl. Gr. ex loco nec ex descriptione.

Hab. in sylvaticis regionis montanae, Laconia in Taygeto (Heldr!) et Malevo (Orph., Fl. Gr. exs. 522!), Eubœa (Sibth.), Macedonia in monte Korthiati (Heldr!) et Atho (Griseb., Pichl!), Bithynia (Boiss!), Galatia (Wied!), Pontus Lazicus ad Khabaekar (Bal!), Lycia (Bourg!), Cilicia prope Gülek (Bal!).

Caulis 3-4-pedalis, folia radicalia sæpe semipedalia, calyx 9-10 lineas longus.

28. P. thapsoides (Bge pl. Lehm. p. 440) tota dense et crasse tomentosa, caule herbaceo erecto subramoso, foliis obsolete dentatis subrugosis supra parce subtus dense lanatis inferioribus longius petiolatis cordato-ovatis, floralibus oblongis subpetiolatis, verticillastris subsexfloris, bracteis paucis tomentosis erectis pungentibus exterioribus calyces subsuperantibus, calycis tomentosi angulati subtruncati dentibus brevissimis patulis pungentibus, corollâ purpureâ calyce duplo longiore tubo intus villoso, filamentis appendiculatis, nuculis glabris ♂.

Hab. in collibus aridis Turkestanie circa Samarkand (Lehm.). Non vidi.

Ex. cl. auctore *P. Cachemirianæ* Royle proxima.

29. P. Herba-Venti (L. Sp. 819) herbacea patula hirsuta vel adpresso canescenti-tomentosa, caulis erectis divaricatum et opposite paniculato-ramosis, foliis oblongo-lanceolatis crenatis vel subintegris inferioribus longius caulinis breviter petiolatis radicalibus basi subcordatis, verticillastris 2-5 remotis foliis floralibus sessilibus superatis, floribus subsessilibus, bracteis subulatis basi sursum curvatis calyce sublongioribus, calycis hirsuti vel tomentosi sulcati dentibus lanceolato-subulatis rigidis erecto-patulis subinæqualibus longioribus tubum dimidium æquantibus, corollâ roseâ, filamentis longioribus basi appendiculatis, nuculis glabris ♂.

α. typica. — Patula hirta vel hirsuta, viridior, folia crenata subtus pallidiora, calyx hirsutus.

Hab. in ditione rarior, Græcia prope Kalamata!, Galatia (Wiedem!), Tauria! (herb. Fauché).

β. tomentosa. — Adpresso tomentosa foliis subtus canis interdum præsertim superioribus subintegris, floralibus angustioribus, calyces adpresso tomentosi. — Ic. Fl. Græc. tab. 564. — *P. pungens* Willd. Sp. III, p. 121.

Hab. in agris incultis et in sterilibus regionis inferioris et montanæ. Attica (Sibth.), Pérotia (Heidr!), Laconia in regione inferiori montis Malevo (Orph. Fl. Gr. 65!), Macedonia et Thraciæ (Griseb., Friv!), totius Anatolie occidentalis!, Galatiae et Paphlagoniae (Wied!), Cappadociæ et Cataoniae (Haussk!), Syriae bor. (Auch! 1813!), Mesopotamiae (Ky. 30!), Armeniae Turciæ ad Bai-but (Bourgl!), Tauriae, Caucasi et Transcaucasiae (M.B., Hoh!), totius Persiae borealis (Auch! Buhse! etc!).

Ar. Geogr. Lusitania, Hispania, Gallia meridionalis, Sicilia et Italia australis, Rossia media et australis, Sibiria Uralensis, Africa borealis.

30. P. Mesopotamica (Boiss. Diagn. Ser. II, p. 47) herbacea adpresso subfloccosa tomentella pallide virens, caulis opposite ramosis, foliis subtus pallidioribus crenatis, radicalibus . . . caulinis breviter petiolatis basi cuneatis oblongo-lanceolatis, floralibus angustioribus verticillastris 2-3-plo longioribus, verticillastris paucifloris numerosis remotiusculis, bracteis tenuiter subulatis hirtis adpressis calyce triplo brevioribus, calycis adpresso tomentosi angulato-sulcati

dentibus a basi ovatâ linearis-subulatis rectis dimidium tubum æquans, corollâ e sicco flavidâ, filamentis appendiculatis, nuculis glabris ♀. — *P. pungens* β *srubutica* Bth. in D.C. Prodr. XII, p. 543.

Hab. in Mesopotamiâ inter Orfa et Suerek (Ky. 49).

A *P. Herbaventi* certe distincta bracteis paucis brevissimis, calycis dentibus rectis; corolla insuper flava videtur.

31. ***P. Persica*** (Boiss. Diagn. Ser. I, 5, p. 37) herbacea adpressissime tomentosa cana, caulinibus simplicibus vel sæpius opposite paniculatis, foliis lanceolatis integris in petiolum brevem attenuatis acutiusculis, floralibus superioribus linearibus verticillastri sublongioribus, verticillastris oppositis ob rachidem elongatam cymuliformibus unilaterali 3-5-floris, bracteis tomentosis subulatis calyci æquifloris in spinulam brevem abeuntibus, calycis sulcati subflocoso-tomentosi dentibus triangularibus brevibus in spinulas nudas breves patentibus inferiorum sublongiores abeuntibus, corollâ roseâ filamentis non appendiculatis, nuculis glabris ♀.

Hab. in Persiâ australi inter Fasa et Schiraz (Auch. 5160 A!), ad Tchinar prov. Lazistan (Haussk!).

β. *cymifera*. — Cymulæ ob rachidem sublongioram laxiores, bracteæ et calyces adpressius tomentosi spinulis terminalibus elongatis. — *P. cymifera* Boiss. Diagn. Ser. I, 7, p. 60.

Hab. prope Schiraz (Ky. 557!), inter Sihna et Kermanschah et prope Dinar-war (Haussk!).

Facies *P. Herbaventi* var. *tomentosæ* a quâ eximie differt verticillastris non condensatis sed in cimas unilaterales evolutis, calyce breviore (4 lin. longo) dentibus multo brevioribus.

SECT. II. PHLOMIDOPSIS.

32. ***P. tuberosa*** (L. Sp. p. 118) herbacea glabriuscula viridis, radicis fibris elongatis hinc inde in tubera ovata incrassatis, caule rubello elato paniculato rarius simplici, foliis petiolatis amplis ovatis obtusis crenatis basi profunde cordatis, floralibus oblongo-lanceolatis sessilibus inferioribus flores superantibus, verticillastris numerosis inferioribus remotis, bracteis numerosis subulatis calyces subæquantibus ciliolatis vel hirtis, calycis glabri vel hirsuti dentibus truncato-retusis costis excurrentibus breviter aristatis, corollâ roseâ calyce duplo longiore, filamentis appendiculatis, nucibus apice pilosis ♀. Ic. Bot. Mag. t. 1555. — Rehb. Ic. tab.

Hab. in collibus et pratis Rumeliæ (Friv.), Armeniæ Turcicæ ad Erzerum (Calv!), Bailut (Bourgl), Gumuschkhané (Tchih!), Tauriâ, Caucaso et Transcaucasiâ (Ledeb.), ditione Talysh (Hohen!), Persiæ prov. Aderbidjan (Buhse!).

P. discolor Led. Fl. Ross. III, p. 435 = *P. bicolor* Willd. herb. non Bth. est planta Sibirica nec Caucasica et varietas vel species aequalis *P. tuberosæ* foliis subtus canis.

Ar. Geogr. Austria, regio Danubialis, Rossia media et australis, Sibiria tota.

EREMOSTACHYS (Bunge in Ledeb. Fl. Alt. II, p. 444.
— *Phlomidis* L. Sp.)

Calyx tubuloso-campanulatus vel infundibuliformis, limbo interdum expanso membranaceo, dentibus 5 latis spinosis. Corolla tubo inclusa, galea compressa labio trifido. Stamina sub galea ascendentia filamentis superioribus basi appendiculatis. Antheræ *Phlomidis*. Nuculae apice densissime pilosæ. Herbae perennes. Genus habitu nec characteribus a *Phlomide* distinctum cui seclio prior fere æquo jure referenda esset.

Sect. I. PHLOMOIDES Bge Lab. Pers. p. 78. — Calycis cylindrici vel campanulati limbus rectus non ampliatus.

• *Verticillastri* dense 6-multiflori.

1. E. laciniata (L. Sp. p. 819 sub *Phlomide*) caule crasso elato in spicam saepius longam longe et densissime albo-lanatam abeunti, foliis virentibus hirtis rarius sublanatis radicalibus amplissimis petiolatis bipinnatisectis segmentis oblongo-lanceolatis vel linearibus inferioribus remotis, foliis caulinis diminutis subsessilibus, floralibus ovato-oblongis incisis flores subæquantibus, verticillastris numerosis multifloris inferioribus vel omnibus remotis, bracteis exterioribus lanceolato-linearibus, calyce campanulato lanâ densâ gossypinâ vestito truncato dentibus latis obtusis in spinulas brevissimas rectas deinceps patentibus abeuntibus, corollâ ochroleucâ vel purpurascenti-luridâ ♀. *E. laciniata* Bge in Led. Fl. Alt. 2 p. 416. — Ic. Bot. Reg. p. 52. — *P. Iberica* Vis. — *I. macrocheila* Jaub. et Sp. Ill. V. p. 13.

Hab. in regione inferiore et montana, Cilicia littoralis (Balt!), Syria (Schubert!), Libani regio inferior (BII Gaill!), Coëlesyria (Boiss!), Palestina ad meridiem Gaza (Boiss!), Transcaucasia et ditio Talyisch (Hohl!).

2. E. adenantha (Jaub. et Sp. Ill. Or. V. p. 13) collo dense lanato, caule erecto modice elato tomentoso-lanato in spicam breviter glanduloso-tomentellam basi subinterruptam abeunte, foliis virentibus supra hirtis subtus canescenti-sublanatis in segmenta oblongo-lanceolata pinnatisida vel partita pinnasectis inferioribus longe petiolatis, verticillastris multifloris numerosis inferioribus subdiscretis, foliis floralibus ovatis acutis subintegratis flores subæquantibus, bracteis linearis-lanceolatis calyce subbrevioribus calycibusque breviter tomentoso-glandulosis, calycis dentibus latis brevissimis truncatis retusis in spinulas brevissimas excurrentibus, corollâ lurido-flavida ♀.

Hab. in Persiae australis collibus prope Schiras et Persepolin (Ky. exs. 303! et 801! sub *E. laciniata*), in dumetis montium Sawers, Eschker, Kotel Mallu et prope Kaserun (Heussk!).

Facies *E. laciniatae* pro cuius formâ olim habui et a qua foliis floralibus

subintegris et spicâ non dense bombycino-lanatâ sed virenti tomentello-viscidâ differt.

3. E. pulvinaris (Jaub. et Sp. Ill. Or. V. p. 12) caule erecto modice elato adpresso tomentello sed ad basin et petiolos dilatatos foliorum radicalium dense gossypino, foliis parce et adpresso arachnoideis subvirentibus in segmenta anguste lanceolata decurrentia pinnatifida pinnatisectis, verticillastris 6-8-floris magno intervallo discretis, foliis floralibus lanceolatis pinnatifidis, bracteis subulatis pungentibus calyce dimidio brevioribus, calycis pube molli brevissimâ velutini dentibus brevissimis latâ in spinulas breves excurrentibus, corollâ lurido-flavidâ ♂.

Hab. in Persiâ prope Ispahan (Auch. 5163), inter Jezd et Ispahan (Bge.!), in montibus Bachtariacis (Bode!).

Sesquipedalis, affinis *E. Tournefortii* a quâ calycis pube breviter velutinâ differt.

4. E. lævigata (Bunge Lab. Pers. p. 81) caule erecto simplici inferne glabro superne adpresso lanatulo, collo dense pulvinato-lanato, foliis radicalibus glaberrimis carnosulis longe petiolatis ambitu oblongis in lacinias paucas lineares elongatas parce inciso-dentatas pinnatisectis, superiorum petiolo brevi lanatulo laciñisque abbreviatis, verticillastris 6-8-floris inferioribus remotiusculis cæteris dense spicatis, foliis floralibus oblongis indivisis spinuloso-cuspidatis, bracteis lanceolatis pungentibus calycis tubum æquantibus, calycis lanati apice truncati dentibus subulato-spinosis brevibus, corollâ lurido-flavidâ sericeâ ♀.

Hab. in monte Schah Kuh Persiæ (Derderian in herb. Hort. Petrop!).

Inter affines foliis glaberrimis paucilaciñatis insignis.

5. E. Tournefortii (Jaub. et Sp. Ill. Or. V. p. 12, tab. 412) caule erecto inferne glabro superne laxiuscule et adpresso lanatulo in spicam abeunti, foliis viridibus subtus canescens in radicalibus longe petiolatis ambitu oblongis in segmenta oblonga vel lanceolata obtuse pinnatipartita pinnatisectis, floralibus sessilibus oblongo-lanceolatis superne dentatis inferioribus flores superantibus, verticillastris multifloris discretis et inferioribus remotis, bracteis subulatis adpressis calyci subæquilongis, calyce tubuloso-campanulato parce lanatulo subglabrato dentibus brevissimis truncatis in spinulam brevem abeuntibus, corollâ albidiâ ♀.

Hab. in Armeniâ (Tourn.), in seccis prov. Aderbidjan (Auch. 5162!) circa Khoï et Seidkhodji (Szov!).

Spicæ indumento parvo laxo et calycibus brevioribus (7 lineas longis) ab *E. laciniata* discrepans. *E. gymnochalyx* Schrenk eni cl. Bth. specimen Aucherianum refert est planta caule multo tenuiori, laciñis foliorum angustissimis, calyce minore glabro.

6. E. glabra (Boiss. Diagn. Ser. I, 12 p. 90) glabra, caule tenui erecto in spicam simplicem abeunti, foliis indivisis oblongis et oblongo-

lanceolatis obtuse et grosse crenatis radicalibus longiuscule petiolatis, caulinis paucis sessilibus, floralibus cuneato-spathulatis paucidentatis apice spinulosis inferioribus flores superantibus, verticillastris numerosis valde dissitis subsexfloris, bracteis setaceis rigidis calyce duplo brevioribus, calyce glabro infra dentes membranaceos truncato-retusos ciliatos dorso in spinas rigidas dentibus longiores abeunte, corollæ albâ parvâ, galeâ glabriuscûlâ margine ciliatâ ♀.

Hab. in monte Elbrus Persiae borealis prope pagum Passgala (Ky. 290!).

Caulis bipedalis, foliorum radicalium limbus 3-4-pollicaris 12-15 lineas latus, calyx 5-6 lineas longus aristis 2 lineas longis.

7. **E. loasæfolia** (Bth. in D.C. Prodr. XII, p. 547) caule procero superne breviter velutino simplici vel parce ramoso, foliis supra glabrescentibus subtus papilloso-velutinis inferioribus petiolatis ambitu oblongis lyrato-pinnatifidis segmento terminali multo majore ovato-oblongo profunde lobato, caulinis sessilibus lanceolatis dentatis, floralibus ovatis subdentatis calyce subbrevioribus, verticillastris 6-8-floris remotis longissime spicatis, bracteolis subulatis paucis calyce quadruplo brevioribus, calycis breviter et dense tomentosi tubulosi dentibus breviter ovatis abrupte et breviter spinulosis, corollæ e secco luridâ ♀. *E. Stocksii* Boiss. Diagn. Ser. II, 4, p. 48.

Hab. in Belutschia 3000'-4000' (Stocks exs. 314! sub *E. laciniata*), in Affghania (Griff.).

Folia eis *E. laciniatae* minus dissecta, spicæ 1-1½-pedales, calyx sex lineas tantum longus. Species bracteis brevissimis insignis.

8. **E. acanthocalyx** (Boiss. Diagn. Ser. II, 4 p. 49) collo lanato, caule glabriusculo in spicam simplicem longam abeunti, foliis radicalibus elongatis petiolatis supra glabrescentibus subtus patule hispidulis in segmenta lanceolata incisa vel pinnatifida pinnatisectis, floralibus triangularibus sessilibus incisis inferioribus verticillastrum superantibus, verticillastris 6-8-floris inferioribus valde distantibus, bracteis numerosis lanceolatis et lineari-setaceis acerosis rigidis calyces subæquantibus, calycibus sub lente tuberculatis stellatim pilosis setulis terminatis obsitis, dentibus brevibus retusis costâ excurrente in spinam validam basi carinatam abeuntibus, corollæ albidâ? ♀.

Hab. in Belutschia prope Doulbund (Stocks exs. 1047!).

Foliorum radicalium limbus 6-8-pollicaris; species a congeneribus induamento minuto pruinosostellato calycis et ejus dentibus validioribus (2 lineas longis) dignoscenda.

9. **E. Lehmanniana** (Bge Lab. Pers. p. 81) caule procero glabro superne longe ramoso, foliis glaberrimis radicalibus lyrato-pinnatisectis segmentis lateralibus unijugis terminali maximo latissime obovalo inciso-lobato, floralibus calyces æquantibus, verticillastris remotis longe spicatis 10-14-floris, bracteolis oblongis calycem æquantibus pungenti-cuspidatis, calycis obconici laxe villosuli denti-

bus truncatis valde et squarrose spinoso-cuspidatis, corollâ calycem duplo excedente ♀. *E. superba* Bge pl. Lehm. p. 411 non Royle.

Hab. in regione alpinâ montis Karatau prope Samarkand Turkestaniae occidentalis (Lehn.). Non vidi.

Planta 3-4-pedalis, petiolus semipedalis, limbi lobus terminalis 8-pollices longus 6 latus. Spicæ pedem et amplius longæ, bracteolæ 8-9 lin. longæ 2 late. Calyx cum spinis 9 lineas longus, corolla fere bipollucaris. Foliis glaberrimis, verticillastris remotis nec in spicam densam condensatis, etc. ab *E. superba* Royle distincta.

• • Flores bini axillares oppositi.

10. ***E. hyoscyamooides*** (Boiss. et Buhse Aufz. p. 181) caule albido parce floccoso fere a basi opposite ramoso, foliis glabriusculis ambitu oblongis breviter petiolatis in lacinias lanceolatas lobulatas pinnatifidis, verticillastris bifloris spicas distichas basi laxiusculas formantibus, foliis floralibus oblongis subintegris apice spinulosis flores sessiles paulo superantibus, bracteolis sub calyce binis linearisetaceis dimidium calyce aequantibus, calyce adpresso tomentello truncato dentibus brevibus latis subtruncatis spinuloso-mucronatis, corollâ calyce duplo longiore galea tomentosa margine dense barbatâ ♀.

Hab. in montosis apricis Persiae ad limites septentrionales deserti magni, prope Semnam (Buhse!), prope Schahrud et Meiomei (Bunge!).

Planta subpedalis, folia pauca cum petiolo 4-5-pollicaria, calyx fructifer coriaceus tenuiter nervosus campanulatus 9 lineas longus. Verticillastris bifloris insignis.

11. ***E. Transoxana*** (Bge Pl. Lehm. p. 441) caule glabro ramoso, foliis petiolo ciliato suffultis supra pubescensibus subtus glabris in segmenta elongata pinnatisectis, floralibus oblongo-lanceolatis acuminateis cuspidatis integrerimis, verticillastris bifloris, floribus pedunculatis bibracteolatis, bracteolis linearibus pungentibus calyce infundibuliforme glaberrimo laevi pungenti-quinquedentato brevioribus ♀.

Hab. ad Jamam Kisilkum Turkestaniae occidentalis (Lehm.). Non vidi.

Ex descriptione ab *E. hyoscyamoide* floribus pedicellatis, calyce glaberrimo, etc. distincta.

12. ***E. Aralensis*** (Bunge Pl. Lehm. p. 442), caule erecto superne ramoso patentem hirsutissimo, foliis subtus glabris superne canescensibus ovatis grosse et acute dentatis, verticillastris bifloris, floribus sessilibus bracteis binis subulatis hirsutissimis pungentibus calyce subaequantibus suffultis, calyce tubuloso undique densissime et patule hirsutissimo dentibus subulatis patentissimis corollæ limbum dimidium aequantibus ♀.

Hab. ad Batkuk Kum Turkestaniae occidentalis (Lehm.). Non vidi.

Ex cl. auctore facies *E. phlomoidis*, ab *E. hyoscyamoide* calycis patule hirsutissimi dentibus valde elongatis, etc., distincta.

*** Verticillastri in cymas laxilloras evoluti.

13. E. thyrsiflora (Bth. in D.C. Prodr. XII, p. 048) crispule pubescenti-canescens, caule humili crassiusculo simplici nudo, foliis radicalibus oblongo-lanceolatis in petiolum longe attenuatis obtuse inciso-dentatis, floralibus sessilibus oblongis dentatis inferioribus flores aequantibus, verticillastris in cynas 5-7-floras evolutis racemum late ovatum formantibus, cymae flore centrali in racemi axi sessili cæteris secus cymæ ramos unilateralibus, bracteis binis linearisubulatis molliter pilosis calyceem aequantibus erecto-patulis, calycis canescenti-tomentosi tubuloso-infundibuliformis dentibus a basi latiore longe subulatis corollâ subbrevioribus ♀.

Hab. in regno Cabulico (Griff. 492), in Belutschia prope Ghurghina (Stocks 971!).

Planta 9-10-pollicaris, foliorum radicalium lamina tripollicaris, thyrsus semipedalis 3 pollices latus, bracteoleæ calycesque pollicares dentibus 3-4 lineas longis.

SECT. II. MOLUCELLOIDES (Bge loc. cit.) Calycis infundibuliformis limbus tandem membranaceo-dilatatus rotatus expansus.

14. E. macrophylla (Montb. et Auch. Ann. Sc. Nat. Ser. 2, VI, p. 54) plus minus dense lanâ adpressâ lanata, caule crasso parum elato simplici vel a basi opposite ramoso, foliis radicalibus amplis petiolatis ovatis vel orbiculatis grosse crenatis superne virentibus vel lanatis subtus canis, floralibus sessilibus cuneatis orbiculatis dentatis calyces aequantibus, verticillastris sexfloris in spicam oblongam vel cylindricam approximatis, bracteolis plumosis subulatofiliformibus rigidis calycis tubum aequantibus, calyce infundibuliformi tubo lanuginoso, limbo demum accrescente maximo pelviformi reticulato villosulo dentibus latissimis semiorbiculato-ovatis breviter mucronatis, corollâ calyci vix aquilongâ ♀. Ic. Jaub. et Sp. tab. 461.
— *E. pyramidalis* Jaub. et Sp. V. p. 69, tab 462 non differt.

Hab. in cultis derelictis, collibus lapidosis regionis inferioris et montane, Galatia (Wied!), Cappadocia ad Karahissar (Ball) et prope Malatia (Auch. 4975!), Armenia australis ad Bakker Maaden (Noë!), ad Erzerum (Auc. 5126!) et Baibut (Bourg!), Gebel Belas deserti ad Pahmyram (Bl!), desertum Khabur et mons Sindjar Mesopotamiae (Haussk!), Persia australis in montibus Bachtiarieis (Haussk!) et ad Persepolin (Ky!), media (Auch. 5127!) orientalis ad Kerman (Bge!), borealis a ditione Aderbidjan (Szov! Buhse!), ad Mazanderan et Khorassan (Bge!).

Pedalis vel elatior, calyx fructifer expansus diametro sesquipollicaris, ab *E. molucelloide* calycis tubo villosissimo nec breviter puberulo corollâque calyce non longiore distincta.

Ar. Geogr. Turkestania.

TR. VII. PRASIEÆ.

PRASIUM (L. Gen. 302).

Calyx campanulatus decemnervius irregulariter bilabiatus labio superiore breviter trifido inferiore profundius bipartito. Corolla tubo inclusa intus pilis squamosis annulata, labiis subæquilongis superiore ovato integro inferiore trifido lobo intermedio majore integro. Stamina sub galea ascendentia, antheræ loculi divaricati. Stylus æqualiter bifidus. Nuculae subbaccatæ. — Frutex.

1. **P. majus** (L. Sp. p. 838) fruticosum glabrum vel superne parce hispidulum, foliis petiolatis a basi truncatâ vel cordatâ ovatis crenatis vel serratis, floribus angustatis minus dentatis vel integris calyces superantibus, verticillastris bifloris plus minus approximatis, calyces breviter pedicellati viridis glabri dentibus ovatis aristatis inferioribus tubo sublongioribus, superioribus tribus altius subirregulariter conualis, corollâ albida glabra 5. Ic. Fl. Græc. tab. 584. — Rehb. Ic. tab. 2.

Hab. ad sepes et in rupestribus regionis inferioris, Græcia et ejus insulæ (Sprn! Heldr! Orph. Fl. exs. 527!), Creta (Sieb!), Rhodus (Bourg!), Chios (Athen. 1507!), Anatolia occidentalis (Sibth!) et australis in Ciliciâ (Per!), Syria littoralis (Gall!), Palestina (Roth! Ky!).

Ar. Geogr. Madera, Lusitania, Hispania, Italia australis, Corsica et Sardinia, Dalmatia, Africa borealis.

TR. VIII. AJUGOIDEÆ.

AJUGA (L. Gen. 705).

Calyx campanulatus vel obconicus 5-fidus vel dentatus. Corolla marcescens tubo inclusa vel exerto intus saepe annulata-pilosâ sparsa recto rarius spiraliter torta, limbus bilabiatus, labium superius brevissimum retusum rarius bipartitum, inferius elongatum patens trilobatum laciniis lateralibus parvis oblongis intermediâ latâ saepe bilobatum. Stamina 4 ascendentia sparsa exserta didynama, antherarum loculi divergentes vel divaricati demum confluentes. Stylus bifidus. Nuculae favosae vel corrugatae. — Herbae vel suffrutices.

SECT. I. BUGULA Bth. Lab. p. 692. — Verticillastri multiflori.

1. **A. reptans** (L. Sp. p. 785) perennis, radice truncatâ e collo stolones elongatos emittente, caule bifarium piloso, foliis ovatis vel

obovatis integris vel grosse crenatis vel dentatis glabriusculis radicalibus persistentibus expansis in petiolum longe attenuatis, caulinis paucis diminutis remotis, floralibus sessilibus ovatis inferioribus flores superantibus, verticillastris 6-8-floris in racemum elongatum basi interruptum approximatis, calycis laciniis lanceolatis villosis tubo æquilongis, corollæ cœruleæ tubo exerto recto, staminibus breviter exsertis ♀. Ic. Engl. bot. tab. 489. — Rchb. tab. 33.

Hab. in pascuis in australioribus montana et subalpina, Peloponnesus in Taygeto (Heldr!), Areadiâ (Heldr!), monte Kyllene 2500' (Orph!), Byzantii (Sibth.), Anatoliâ totâ boreali (Wied! Tchih!), Ponto ad Trapezuntum et Batonum (Bourg! Calv!), Caucaso et Transcaucasiâ (Ledeb., Ruprl.), prov. Talysch (C.A. M.), Persia borealis (Buhse!).

Ar. Geogr. Europa præsertim borealis et media ab Angliâ et Gothiâ ad Rossiam medium.

2. A. Genevensis (L. Sp. 785) perennis exstolonosa, caule undique villoso, foliis oblongo-ellipticis sinuatis vel crenatis plus minus hirsutis radicalibus petiolatis sub anthesi sæpius evanidis, floralibus cuneatis dentatis vel trilobis rarius integris superioribus flores non æquantibus, verticillastris in spicam longam basi interruptam approximatis, calycis lanatuli dentibus lanceolatis tubo longioribus, corollæ cœruleæ tubo exerto recto, staminibus exsertis ♀. Ic. Engl. bot. tab. 477. — Rchb. Ic. tab. 33. — *A. pyramidalis* M.B. 2, p. 32, non L.

Hab. in Macedonia peninsula Chalcidice (Griseb.), prope Bitolia (Orph!), Armenia Kurdiçâ (Brant ex Bth.), Tauriâ (Stev.), Caucaso et Transcaucasiâ (Ledeb.), Caucaso orientali 4500'-6000' (Ruprl.).

Ar. Geogr. Europa borealis et media a Daniâ et Gothiâ ad Rossiam medium, China.

3 A. pyramidalis (L. Sp. 785) perennis exstolonosa, caule undique villoso, foliis plus minus hirsutis radicalibus persistentibus magnis expansis obovatis obsolete crenatis vel integris, floralibus ovali-oblongis sœpe purpureis omnibus flores superantibus, verticillastris sexfloris in racemum tetragono-pyramidalatum sæpius continuum approximatis, calycis lanatuli dentibus lanceolatis tubo duplo longioribus, corollæ parvæ pallide cœruleæ tubo recto breviter exerto, filamentis exsertis ♀. Ic. Fl. Dan. t. 185. Engl. bot. tab. 1270. — Rchb. tab. 33.

Hab. in subalpinis Caucasi Orientalis 5500' (C.A. Meyer), in monte Hæmo Thracie (Sibth.). E ditione nondum vidi.

A precedentibus foliis radicalibus amplis, floralibus etiam supremis flores superantibus, floribus manifeste minoribus distincta. Flore Orientalis dubia civis et forsitan cum *A. Genevensi* commutata.

Ar. Geogr. Scandinavia, Scotia, Alpes, Gallia et Germania septentrionalis, regio Danubialis, Rossia septentrionalis.

4. A. bracteosa (Wall. pl. Asiat. rar. I, p. 29) perennis exstolonosa patule villoso-canescens, caulis diffusis, foliis radicalibus

ovatis vel oblongis sinuato-dentatis in petiolum angustatis, floralibus flores multo superantibus, verticillastris parvis remotis, floribus parvulis, calycis albo-hirsuti globoso-campanulati dentibus lanceolatis tubo æquilongis, corollæ purpurascens tubo recto vix exerto, staminibus inclusis ♀.

Hab. ad margines agrorum ad Chagur regni Cabulici (Griff. Journ. N° 30!).

A præcedentibus verticillastris distantibus, floribus fere triplo minoribus, staminibus inclusis distincta.

Ar. Geogr. India borealis.

5. A. Orientalis (L. Sp. 785) perennis exstolonosa villosolanata, foliis radicalibus persistentibus oblongis elongatis obtuse sinuatis vel inciso-lobatis in petiolum longe attenuatis, floralibus sessilibus oblongis 3-5-lobis flores superantibus, verticillastris distantibus vel supremis approximatis, calycis lanati laciniis lanceolatis tubo duplo longioribus, corollæ cœruleæ tubo breviter exerto intus nudo torto labio resupinato, staminibns subinclusis ♀. Ic. Dill. Elth. tab. 53.

Hab. in pratis pinguibus, sylvaticis, a regione inferiori ad montanam et in australioribus subalpinam, Cephalonia in monte Oeno (Heldr! forma bracteis violaceo-purpurascens), Attica, Bœotia, Parnassus (Sprun! Boiss! Heldr!), Thracia (Sibih.), Olympus Bithynus (Auch! Boiss!), Anatolia borealis omnis (Wied! Tchih!), Pamphylia ad Adalia (Heldr!), Cilicia (Bal!), Cyperi mons Troodos (Ky!), Syria in Libano et Antilibano (Boiss! Gaill!), Armenia mons Techdagh (Huet!), Georgia Caucasica (Hohen!).

β. condensata. — Humilior, racemus magis condensatus, folia floralia minora, superiora floribus breviora. — *A. Orientalis* var. *orthosiphon* G. Koch Linn. XVII, p. 301 (sed corollæ tubum etiam in speciminibus Kochianis tortum video!).

Hab. praesertim in alpinis, Armenia Turcica (C. Koch!) et supra Gumusch Khané (Bourg!), Pontus Lazicus 8000' (Bal!), Taurie montes (Stev!), Caucasus omnis 3000'-9000' (Hoh! Rupr!), Armenia Rossica (Buhse!).

Vix varietas, forma ex loco elatiore.

Ar. Geogr. Lusitania, Hispania, Italia australis.

SECT. II. CHAM.EPITHYS Tourn. — Verticillastris biflori raro quadri-flori. Corollæ labium superius non fissum.

6. A. salicifolia (L. Mant. p. 80 sub *Teucrios*) adpresso et brevissime canescens, caulis e rhizomate indurato subrepente numerosis ascendentibus vel basi diffusis simplicibus, foliis oblongis vel ellipticis obtusis integris in petiolum brevem angustatis, floralibus conformibus flores multoties superantibus, floribus binis oppositis pedicellatis, calycis adpresso tomentelli fructiferi subdilatati laciniis lanceolatis obtusis tubo subbrevioribus, corollæ luteæ calyce duplo longiore ♀. *A. salicifolia* Schreb. Un. p. 26. — Ic. Fl. Gr. tab. 626

Hab. in collibus siccis et sylvaticis, Thracia (Friv!), Phrygia (Sibth.) et prope Ouchak (Bal. 4173!), Galatia (Wied!), Pisidia et Lycaonia (Heldr!), Cilicia (Ky!), Cappadocia (Auch. 4606! Ball!), Armenia (Tourn.), prope Baibut (Bonrg!) et Tortum (Calv!), Tauria (Stev.), Caucasus (M.B.), Kurdistania (Auch. 1577!).

β. tomentella. — Breviter sed non adpresso tomentella. — *A. Chamæcistus* var. *β.* Bth. in p.c. Prodri.

Hab. ad Ispahan Persie (Auch. 5164!).

Huc nec ad *A. Chamæcistum* ob ramos herbaceos flexuosos nec fruticosos rigidos referenda videtur, corolla insuper e sicco flava.

7. A. oblongata (M.B. Taur. Cauc. III, p. 388), patula villosa, caulis crassis ascendentibus ramosis, foliis oblongis vel ellipticis integris obtusis basi attenuatis subsessilibus, floralibus sensim angustatis flores multo superantibus, floribus binis oppositis subpedicellatis, calycis longe hirsuti dente supremo minuto ceteris tubo longioribus, corolla flavâ calyce triplo longiore ♀. *A. salicifolia* Stev. Mem Mosq. III, p. 265.

Hab. in sabulosis Transcaucasiae ad fluvium Cyrum et prope Salliam (Stev!), in cultis Mesopotamiae prope Bagdad (Auch. 1575! Noe 19!).

Affinis *A. salicifoliae*, caules crassiores pedales ramosi magis herbacei, indumentum longum patulum nec adpressissimum, calycis dentes magis inaequales, corolla calycis proportione longior. Contuli specimen Stevenianum herb. M.B.

8. A. Chamæcistus (Ging. in Bth. Lab. p. 698) suffruticosa ramosissima ramis erectis adpresso velutinis albidis demum persistentibus aphyllis subspinescentibus, foliis sessilibus strictis oblongo-ellipticis et elliptico-linearibus obtusis integerrimis, floralibus conformatibus flores longe superantibus omnibus adpresso canescensibus, floribus binis oppositis pedicellatis, calycis adpresso canescens campanulato-tubuloso dentibus lanceolatis tubo æquilongis superiore breviore, corollâ purpureâ calyce 2½-plo longiore ♂.

Hab. in collibus Persie (Oliv.), in collibus argillosis ad basin montis Demavend Persie borealis (Ky. 395!), subalpinis prope Schahrud et circa Teheran (Bge), circa Ispahan (Bode!), Persia australi circa Schiras (Ky. 360!), in subalpinis Kurdistaniae Persie circa Mendeli (Noe 1162!).

β. stenophylla. — Folia longe et anguste linearia 1-1½-pollucaria linea unâ vix latiora margine subconvoluta. — *A. scoparia* Boiss. Diagn. Ser. I, 7, p. 63.

Hab. in saxosis ad ruinas Persepolis (Ky. 246!).

γ. euphrasioides. — Folia angusta varietatis sed dentibus seu lobis acutis utrinque 1-3 aucta. — *A. euphrasioides* Boiss. Diagn. Ser. I, 7, p. 63.

Hab. in collibus ad lacum Nemeck Derja prope Schiras (Ky. 458!).

Species ab *A. salicifolia* ramis fruticosis rigidis albidis corolla que purpurea facile distingueenda sed ramis plus minus spinescentibus, foliis (vix iam ½-1½-pollucies longa 1-5 lineas lata) integris vel dentatis valde variabilis. Tota planta moschum reololet.

9. **A. Iva** (L. Sp. p. 787 sub *Teucrio*) perennis canescens vel villosa basi ramosissima ramis basi induratis brevibus prostratis vel ascendentibus valde foliosis, foliis linearis-oblongis et linearibus marginé revolutis utrinque remote 1-3-dentatis vel integerrimis, verticillastris 2-4-floris foliis floralibus longe superalis secus totum ramum racemum basi interdum laxiorem formantibus, calycis albo-lanati dentibus lanceolatis obtusiusculis tubo brevioribus, corollâ purpureo-roseâ vel luteâ calyce triplo longiore interdum abortiva, seminibus foveolato-alveolatis ♀. *A. Iva* Schreb. Unil. p. 25. — Ic. Fl. Græc. tab. 525. — Cav. Ic. tab. 120. — Rchb. tab. 34. — *A. pseudo-iva* n. c. Fl. Fr. Suppl. 395.

Hab. in aridis et cultis Græciae (Sprun! Heldr! Barbey!), Archipelagi et Cretæ (Sibth.), Cypri (Kyl), Egypti ad Alexandriam (Ehrenb! Samar!).

Ar. Geogr. Insulæ Canarienses, Madera, Lusitania, Hispania, Gallia et Italia australes, Dalmatia, Africa borealis.

10. **A. Chamæpitys** (L. Sp. 787 sub *Teucrio*) annua pilosohispida caulis brevibus diffusis crebre foliosis, foliis viscidulis infimis minimis linearis-oblongis petiolatis cæteris in lacinias angustè lineares tripartitis, floralibus flores superantibus, floribus oppositis racemos foliosos præter basin laxiusculam densos secus totum ramum formantibus, calycis hirsuti dentibus inæqualibus lanceolatis tubo sublongioribus, corollâ luteâ calyce triplo longiore, nucularum areolis fere omnibus pentagonis ⊙. *A. Chamæpitys* Schr. Unil. p. 24. — Ic. Fl. Dan. tab. 733. — Engl. bot. tab. 77.

Hab. in cultis Græciae, Archipelagi et in Bithyniâ (ex Sibth.), Caucaso et Iberia Caucasicâ (Rupr!), Persiâ boreali (Bge!), Tauriâ (Stev.).

Ar. Geogr. Europa media et australis ab Angliâ et Germaniâ ad Rossiam mediam, Africa borealis.

11. **A. Chia** (Poiret Suppl. II, p. 772 sub *Teucrio*) perennis patule et longe albo-hispida, rhizomate ramoso saepè indurato, caulis prostratis vel ascendentibus, foliis crebris infimis in lacinias integras vel dentatas trifidis vel tripartitis cæteris in lacinias latiusculæ lineares trifidis, floralibus corollâ saepius subbrevioribus, verticillastris bifloris in racemos longos laxiusculos vel densos dispositis, calycis albo-hirsuti lacinias lanceolatis acutis subinæqualibus tubo brevioribus, corollâ calyce quintuplo longiore luteâ purpureo-striatâ rarius purpureâ, nuculis præter areolas paucas terminales pentagonas transverse corrugatis ♀. *A. Chia* Schreb. Unil. p. 25. — Ic. Fl. Græc. tab. 524. — *A. intermedia* Boiss. et Orph. MSS.

Hab. in sterilibus et cultis derelictis fere totius ditionis a Græciâ (Sprun! Boiss! Heldr! Orph. Fl. exs. 533!), Archipelago (Tourn., Sibth.), Byzantio et Bithyniâ (Griselb.), Anatoliâ boreali (Wied!), orientali (Bourg!) et australi in Ciliciâ (Heldr! Ball!) ad Cappadociam (Bal!), Armeniam Turcicam (Auch! Tchih!), Tauriam (Stev.), Transcaucasiam (Hohen!), Syriam littoralem (Bl! Gaill!).

Valde affinis *A. Chamæpithysi* et ab ea non semper facile dignoscenda,

eam specifice distinctam tamen automo, rhizoma perenne interdum indurato-suffrutescens, foliorum laciniæ (non semper) latiores, corolla longior 9-10 lineas longa calyx quintuplo nec triplo longior et in primis nucula transverse corrugatae nec (apice excepto) foveolatae. Formas sequentes ab auctoribus et me ipso quondam ut species proprias habitas, intermedii conjunctas, nunc ut varietates enumerare cogor.

β. latiloba. — Foliorum laciniæ latiores abbreviatæ, corolla magna lutea, labii lobus medijs basi rotundatus non cuneatus. — *A. Mesogitana* Boiss. Diagn. I, 7 p. 42.

Hab. in regione alpinâ montis Mesogis Lydie in terrâ mobili (Boiss!), Anatoliâ boreali prope Amasia et Toptcha (Wiedem!), in monte Bulghasdegh Phrygiae (Ball).

γ. suffrutescens. — Intensius canescens basi indurata suffrutescens, rami sæpius abbreviati, calycis dentes subbreviores, foliorum laciniæ sæpe abbreviatæ, corolla lutea interdum purpurascens. — *A. Palæstina* Boiss. Diagn. Ser. I, 12 p. 92.

Hab. in regione alpinâ, Olympus Bithynus in latere australi (Pichler!), mons Gheidagh Tauri Isauriei 7-8000' (Heldr!), Taurus Cilicus 3-9000' (Ky. 16! 54!), Libani et Hermonis cacumina 8-9000' (Boiss! Bl! Gaill!), et quoque in australioribus calidis, circa Hierosolymam (Boiss! Ky. 63!), Damascum (Boiss! Gaill!), Aleppo et monte Sindjar (Haussk!), mons Pir Omar Gudrun Assyriæ 4000' (Haussk!).

δ. tridactylites — Patule hispidissima, rhizoma sæpe induratum sed rami teneri et folia membranacea abbreviata cuneata in lobos latos ovato-oblongos trifida. Corolla purpurea. — *A. tridactylites* Gingins in Bth. Lab. p. 699.

Hab. in rupestribus umbrosis, mons Sinaiticus Ste Catherine (Bové Schimp. 447! Boiss!), Antilibanus ad Zebdani (Ky. 8!), Cyprus ad rupes Buffavento (Ky. 426!), Persia australis in fissuris calcareis, ad Gere (Ky. 34!), Perezend (Auch!), Shahpur, Kaserun, Daleki (Haussk!).

12. *A. vestita* (Boiss. Diagn. Ser. I, 7, p. 600) perennis albo-lanata, rhizomate crasso pluricipiti, caulis crassis fere a basi in racemum densum ovato-oblongum abeuntibus, foliis radicalibus petiolatis rhomboc-ovatis obtuse 3-4-crenatis, caulinis sessilibus cuneatis ad medium trilobis corollas æquantiibus, calycis albo-lanati laciniis lanceolatis æquilogis tubo sublongioribus, corolla albâ! calyce $2\frac{1}{2}$ -plo longiore, nuculis.... *γ. A. tridactylites* var. *lanata* Bth. Prodr. D.C. 12, p. 600, ex parte.

Hab. in fissuris rupium Assyriæ circa Mardin (Auch. 4576! Haussk!), inter Diarbekir et Mardin (Ky. 320!).

Vix bipanicarisi, ab omnibus *A. Chia* formis racemo brevi densissimo, floribus parvis sex lineas tantum longis, calycis laciniis tenuibus longioribus, corolla albâ distincta.

13. *A. bombycina*, annua? tota dense et molliter argyreobombycina, caulis prostratis vel ascendentibus brevibus, foliis radicalibus rosulatis oblongo-spathulatis integris vel paucicrenatis, floralibus cuneatis late brevissime et obtuse trilobis corollas sub-

superantibns, verticillastris bifloris in racemum inferne laxum dispositis, calycis lanati dentibus subulatis tubo æquilongis, corollâ luteâ calyce triplo longiore, nuculis omnibus transverse corrugatis ⊖.
A. tridactylites var. *lanata* Bth. ex parte.

Hab. in rupestribus promontorii Aiava Pamphyliæ (Heidr!), circa Keban Maaden Cataonæ (Haussk!).

Indumentum fere *A. vestue* sed mollius longius argyreum, radix annua vel saltem exilis nec rhizoma incrassatum, racemi laxi 3-4-policares, flores 6-7 lineas longi, nucule eis *A. Chiae* fere triplo minores.

14. ***A. laevigata*** (Russ. Alepp. II. p. 255 sub *Tenurio* non Vahl) annua herbacea glabra, caulis ex radice verticali demum indurata pluribus ascendentibus et erectis simplicibus vel corymbosis, foliis radicalibus linear-lanceolatis longe petiolatis integris vel paucidentatis, floralibus basi subdilatatis sessilibus in lacinias longas lineaes integras corollam æquantes tripartitis, verticillastris bifloris in spicas basi laxiusculas approximatis, calycis glabri laciñis lanceolatis acutis subæqualibus tubo sublongioribus apice interdum paniculosis, corollâ luteâ inferne purpureo-striatâ calyce quintuplo longiore, nuculis magnis transverse valde corrugatis apice foveolatis ⊖. *A. glabra* Bth. Lab. p. 700 vix Presl.

Hab. in saxosis et agris Cilicie ad Gulek et Goramse (Ky. 277! et 108!), ad Sedichig (Ball!), Cilicie ad Aleppo (Russ., Auch. 1564!, Ky. 206!), prope Nisib (Haussk!), in Armeniâ australi prope Musch (C. Koch!) et Warto (Ky. 406!).

Caules cum racemis semipedales, corollæ 9-10 lineas longæ, nuculae sesquilineam et amplius longæ. Ab *A. Chamœpithye* glabritie, omnibus partibus triplo majoribus, nuculis corrugatis distincta. *A. glabra* Presl (cujus nomen cæterum recentius) planta Sicula male nota videtur forma glabrescens *A. Chiae*.

SECT. III. PHLEBOANTHE Tausch. — Verticillastri biflori. Corollæ labium superius bipartitum.

15. ***A. Laxmanni*** (Linn. Syst. Veg. p. 439 sub *Tenurio*) perennis patula pilosa, rhizomate indurato ramoso, caulis pluribus crassis erectis simplicibus crebre foliosis, fo'liis ramorum steritum obovato-oblongis basi longe attenuatis apice paucidentatis, caulum fertilium sessilibus oblongis vel ellipticis integris, floratinus conformibus calyces multo superantibus, verticillastris bifloris, calyce pedicellato villosa laciñis oblongis obtususeculis tubo sublongioribus subinaequalibus, corollæ calyce duplo longioris flavidae purpureo-venose tubo superne fisso ♀. *A. Laxmanni* Blh. Lab. p. 697. — Ic. W. K. Hung. tab. 69. — Rehb. Ic. tab. 35. — *Phleboanthe Laxmanni* Tausch.

Hab. in regione inferiori et montana in sylvaticis Macedoniae et Thracie (Grisch!), Byzantii (Couniany!), Olympo Bithyno (Auch! Boiss!), Tauriâ (Stev!). Iberia Caucasicæ (M.B., etc.).

Ar. Geogr. Hungaria, Croatia, Transylvania, Rossia australis.

TEUCRIUM (L. Gen. 706).

Calyx tubuloso-campanulatus vel campanulatus quinquedentatus dentibus æqualibus vel inequalibus. Corolla decidua tubo brevi intus non annulato, limbi labio superiore brevissimo fisco inferiore quinquelobo lacinia infera maximæ rotundata vel oblonga saepius concavæ cæteris parvis oblongis (in nostris) declinatis. Stamina quatuor ascendentes saepius longe exserta didynama, antherarum loculi confluentes. Stylus apice subæqualiter bifidus. Nuculae basi obliquæ reticulatae vel rugosæ. — Frutices vel herbæ.

Specierum Orientalium distributio.

SECT. I. TEUCRIS Ging. — Perennia. Pedunculi oppositi axillares 1-3-flori racemosi vel paniculati, Calyx rectus æqualis.

* Folia integra.

T. rosmarinifolium, brevisolium, Pestalozzei, Aroanium.

** Folia dentata vel saepius partita.

+ Filamenta exserta.

T. multicaule, Orientale, pruinatum, Taylori, procerum, rigidum, Olivieri.

++ Filamenta subinclusa.

T. macrum, parviflorum.

SECT. II. STACHYOBOTRYS Bth. Lab. p. 672. — Perennia. Flores sparsi vel irregulariter subverticillati in spicam simplicem dispositi. Calyx campanulatus bilabiatus dente supremo latiore ovato.

T. Hyrcanicum, lamiifolium.

SECT. III. SCORODONIA Bth. Lab. 674. — Perennia. Pedunculi oppositi axillares uniflori secundi racemosi. Calyx campanulatus bilabiatus dente supremo ovato latiore.

T. Kotschyanum, Massiliense.

SECT. IV. SCORDIUM. — Perennia. Verticillastri 2-6-flori axillares distantes. Calyx basi obliquus subgibbus dentibus subæqualibus.

T. Scordium, scordioides, serratum, mellissoides.

SECT. V. SPINULARIA. — Annum, verticillastri 2-4-flori axillares distantes. Calycis basi gibbi dentes spinosi superior multo latior ovatus.

T. spinosum.

SECT. VI. ISOTRIODON. — Perennia. Verticillastri biflori in spicas ovato-oblongas vel racemos breves dispositi. Calycis dentes terni superiores ovati vel oblongi, inferiores bini angustiores.

T. pederotoides, Montbreti, Odontites, Persicum.

SECT. VII. CHAMÆDRYS Bth. Lab. p. 680. — Perennia. Verticillastri 2-4-flori in racemum laxum dispositi. Calyx basi subgibbus dentibus lanceolatis subæqualibus.

T. Chamædrys, flavum, divaricatum, microphyllum, leucophyllum, coniortodes, Socinianum.

SECT. VIII. POLIUM Bth. Lab. p. 684. — Perennia. Verticillastri in capitula globosa vel ovata terminalia condensati. Calyx tubuloso-campanulatus dentibus subæqualibus vel inferioribus subangustioribus et acutioribus.

T. montanum, Mascatense, alpestre, Cyprium, leucocladum, cuneifolium, Stocksonianum, Polium, Sinaicum.

SECT. I. TEUCRIS.

1. **T. rosmarinifolium** (Lam. Diet. II. p. 693) fruticosum, ramis virgatis elongatis internodiis distantibus, foliis supra viridibus subtus albōmentosis margine revolutis integris inferioribus crebris elliptico-linearibus inferne attenuatis superioribus anguste linearibus, floralibus calyce brevioribus, floribus oppositis pedicello calyce aequilongo suffultis spicas longas tenues laxiusculas formantibus, pedicellis infimis interdum 2-3-floris, calycis albōmentosi breviter campanulati dentibus triangulari-lanceolatis acutis tubo subbrevioribus, corollâ purpurascenti-cæruleâ calyce subduplo longiore 5. *T. Creticum* L. Sp. 788 sed in Cretâ non crescit! — Fl. Græc. tab. 529. — *T. hyssopifolium* Schreb. Unil. p. 28. — *T. Charamonense* Cavan.

Hab. in collibus regionis inferioris Ciliciae ad Bouloukli prope Mersina (Balt!). Cypri (Sibth! Ky. 334!), Syrie borealis in monte Torbol ad Tripolin (Boiss!). Palestine in Carmelo (Boiss! Cadet!), ad Arimathæam (Sieb!).

Frutex pluripedalis, folia omnino *Rosmarini*.

2. T. brevitolium (Schreb. Unilab. p. 27) fruticosum humile ramosissimum ramis brevibus saepe intricatis, foliis parvis adpressissime canescentibus subsericeis oblongis et linearibus obtusis integris margine revolutis, floralibus conformibus calyces superantibus, floribus oppositis pedicellatis racemos breves terminales formantibus, calyce adpresso tomentello in dentes oblongos obtusiusculos tubo subbreviores subæqualiter quinquefido, corollâ cœruleâ purpureostriatâ ♂. Ic. Fl. Græc. tab. 528. — *T. Creticum* Lam. Dict. II, p. 592 non L.

Hab. in rupibus maritimis Cretæ (Sibth! Sieb!), Melos (Urv., Armenis!), Poros et Naxos (Wied!), Atticæ ad promontorium Sunium et prope Laurium (Heldr!).

Frutex 1-2-pedalis, folia 3-4 linea longa sesquilineam lata, flores eis *T. fruticantis* duplo minores.

3. T. Pestalozzæ (Boiss. Diagn. Ser. I, 7, p. 90) fruticulosum humile rameolum rami novellis erectis crebre foliosis sub lente pruinosis, foliis parvis rigidulis sub lente adpresso puberulis oblongo-linearibus obtusis margine valde revolutis, floralibus linearibus acutis pedunculos superantibus, pedunculis unifloris oppositis calyci subæquilongis breviter racemosis, calycis birtuli elevatim quinque-nervii laciniis lanceolatis rigidis subpungentibus tubo æquilongis, corollâ cœruleâ subtus barbatâ ♂.

Hab. in Lyciâ prope Elmalu (Pestal.).

Rami 3-5-pollicares, folia majora 6-8 linea longa 1-1½ lata. Ab aſlini *T. brevitolio* indumento subnullo nec sericeo, calycis dentibus pungentibus nec obtusis, etc. distinctum.

4. T. Aroanium (Orph. in Boiss. Diagn. Ser. II, 4, p. 55) suffruticosum ramosissimum caudiculis nudis duris tenuibus radicantibus prostratis, ramis novellis brevissimis ascendentibus crebre foliosis canis, foliis parvis petiolatis obovatis et ovato-oblongis integris adpressissime supra canescentibus subtus canis. floribus oppositis pedicellatis magnis in racemum breviter ovatum 2-6-florum terminali approximatis, bracteis ellipticis acutis calyce brevioribus, calyce glanduloso campanulato viridi dentibus ovato-triangularibus acutissimis tubo æquilongis, corollâ cœrulecente calyce multo longiore laciniâ inferiore longe crispulo-papillosa, filamentis longissime exsertis ♂.

Hab. in regione alpinâ montis Chelmos (Aroanios) Achæa ad Stygem (4500'-5800') ubi rupes letis tapetibus vestit. (Orph. Fl. Græc. exs. 402!).

Rami floriferi 2-3-pollicares, folia 9-10 linea longa 2-3 lata. Flores eis *T. fruticantis* fere maiores. Insigne Floraë Græcæ decus.

5. T. multicaule (Month. et Auch. Ann. Sc. Nat. Ser. II, 6, p. 54) fruticulosum basi lignosum ramosissimum partibus novellis adpressiusculo hirtum pallide virens, ramis erectis strictis crebre

foliosis subsimplicibus in racemum angustum laxum abeuntibus, foliis in lacinias lineares longas margine subrevolutas integras strictas ad vel sæpius infra medium tripartitis, floralibus flores superantibus et æquantibus, summis indivisis, pedunculis solitariis oppositis strictis calyci subæquifloris, calycis tomentelli laciñiis subæqualibus lanceolatis acutissimis tubo duplo longioribus, corollæ extus hirtæ cœruleæ calyce subdupo longioris lobis obtusiusculis superioribns ellipticis, inferiore longiore, filamentis exsertis glabris, nuculis minute pruinoso-velutinis ♂.

Hab. in Cappadociâ ad Euphratem (Auch 4598!), in cretaceis Syriæ borealis inter Aïntab et montem Ssöfflagh (Haussk!), inter Orfa et Suerek (Ky. 420!).

Rami novelli cum spicâ semipedales, foliorum laciñiæ sepe pollicares, flores eis *T. Orientalis* subminores. Ab eo et sequentibus inflorescentiæ spicato-racemosâ et trunko fruticoso alienum. *T. pseudochamæpithys* caulinis herbaceis, pilis patentibus, filamentis longe exsertis, corollâ majore etc. differt.

6. *T. Orientale* (L. Sp. p. 786) perenne, rhizomate multicauli interdum indurato-suffrutescenti, caulinis erectis patule hispidulis superne opposite ramulosis breviter paniculato-thyrsoideis, foliis viridibus vel canescentibus patule hispidulis ambitu ovatis subpetiolatis in lacinias anguste lineares longas margine subrevolutas semel bisve pinnatisectis, pedunculis 1-3-floris, pedicellis calyce bracteâque longioribus, calycis hirtuli breviter et patentim campanulati dentibus lanceolatis acutissimis carinatis tubo hemisphaericâ sublongioribus, corollæ cœruleæ calyce 3-4-plo longioris laciñiâ inferiore elongatâ acutiusculâ concavâ subtus hirtâ, filamentis longe exsertis glabris, nuculis glabris minute pellucido-pruinosis ♀.

Hab. in collinis siccis, cultis derelictis Anatolie orientalis in interioribus Lydiæ et Cariæ (Boiss! Bal!), Galatiâ et Paphlagoniâ (Wied!), Armeniâ Turcicâ ubi frequens (Huet! Calv! Bourg!), Cappadociâ (Tchih! Bal!), Caucaso, Armeniâ Rossieâ et Georgiâ Caucasicâ (C.A. M., Hoh! Szov!), Persiâ boreali (Szov! Bge).

β. nivale. — Humilius semipedale, pedunculi sæpius uniflori. — *T. nivale* Boiss. Diagn. Ser. I, 12. p. 91. •

Hab. in regione alpinâ Libani (Ehrenb! Bl! Gaill!), Antilibani in Hermone et supra Orny 5000'-900' (Boiss! Gaill! Ky. 212!).

7. *T. pruinatum*, perenne totum adpresso canescens, e rhizomate indurato multicanle, caulinis superne opposite et stricte ramulosis paniculato-thyrsoideis, foliis ambitu ovatis in lacinias convolutas anguste lineares strictas breves pinnatum vel subbipinnatum sectis, pedunculis 1-3-floris, pedicellis calyce et bracteâ longioribus, calycis adpresso canescentis dentibus lanceolatis apice subuncinatis tubo elevatione quinquenervi sublongioribus, corollâ cœruleâ pruinosâ calyce triplo longiore lobo inferiore elongato acutiusculo, filamentis exsertis glabris, nuculis glabris pellucide pruinosis ♀. *T. Orientale* var. *pruinatum* Boiss. MSS.

Heb. in aridis subsalsis planitiei prope Karaman Lycaoniae ad radices montis Karadagh (Heldr!), ad Caesaream Cappadociae (Bal!), ad Aintab Syriæ borealis (Haussk!), Antilibano prope Dinar (Gaill!), ditione Golaa (Wettstein!).

A formis *T. Orientalis* specifice differre videtur indumento adpresso, dentibus calycis apice subuncinatis, foliorum laciñiis abbreviatis. Planta 1-1 $\frac{1}{2}$ -pedalis.

8. T. Taylori Boiss. Diagn. Ser. I, 7, p. 61) totum brevissime cano-tomentosum e rhizomate iuerassato multicaule, caulis rigidis superne breviter paniculatis, foliis circumscriptione ovato-orbiculatis in laciñias breves convolutas anguste lineares pinnatim vel subliplinatum sectis, pedicellis 1-3-floris calyce et bracteis 2-3-plo longioribus, calycis subglandulosi breviter campanulati dentibus a basi triangulari longe subulatis tubo aequilongis, corolle cœruleæ calyce triplo longioris lobo inferiori cœullato obtuso subtus hirsuto, filamentis longe exsertis parte inferiori pubescentibus, nuculis glabris pellucido-pruinosis ♀.

Hab. in Persiaustrali (Auch. 5167!) ad radices montis Sabst Buschom prope Schiraz et Kuh Ajub prope Persepolin (Ky. 370! et 393!), prope Dalechi et Kaserua (Haussk!), prope Rudbar (Buhse!).

Indumentum *T. pruinosi*, calycis dentes lengius et rigidius subulati, ab eo et *T. Orientali* filamentis inferne pubescentibus et labio inferiore obtuso distinetum.

9. T. procerum (Boiss. et Bl. Diagn. Ser. II, 4, p. 56) glabrum, caulis proceris valde quadrangulis a basi ramulosis in paniculam longissimam pyramidatam abeuntibus, foliis lucidis in laciñias longas lineares subconvolutas semel vel bis pinnatisectis, pedicellis hirtulis calyce et bracteis longioribus, calycis breviter campanulati hirtuli laciñiis lanceolatis acutissimis tubo aequilongis, corolle cœruleæ calyce quadruplo longioris extus patule glanduloso-hirsutæ lobo inferiore elongato acuto superioribus longe acuminatis, filamentis longe exsertis inferne hirtulis, nuculis glabris pellucido-pruinosis ♀. *T. pictum* Boiss. et Haussk. MSS.

Hab. in Syria (Auch. 459!), Libano prope Solima (Bl!), in arenosis ferrugineis Merdj Aliou supra Sidonem (Gaill!), in monte Makmel supra Ainete (Boiss!), monte Berydaghi Cataoniae et montibus Schaln et Avreman Kurdistanie Persie in demensis 6-9900' (Haussk!).

Sæpe 2-3-pedale in excelsioribus humilius; præter glabritiem ab affinis corolle glanduloso-hirsutæ laciñiis superioribus non tantum aëantis sed acuminatis differt.

10. T. rigidum (Bth. in o.c. Prodr. 12, p. 578) breviter et molliter cano-tomentellum, caulis crassis rigidis longe paniculatis, foliis in laciñias convolutas anguste lineares semel vel bis pinnatisectis, pedunculis 1-3-floris, pedicellis bracteis et calyce longioribus, calycis tomentosi dentibus lanceolatis acutis tubo longioribus, corolle cœruleæ hirtulæ calyce triplo longioris laciñiis superioribus binis

brevioribus acutis, inferioribus ellipticis obtusiuseulis intermediâ cymbiformi obtusiusculâ sublongiore, filamentis longe exscapa glabris, nuculis pilosissimis ♀.

Hab. in Assyriâ (Auch. 1604! et 1605!), ad Derbent i Basian (Haussk! mixtum cum *T. parvifloro*).

Stamina longe exserta *T. Orientalis* a quo laciniis tribus inferioribus corollae parum inaequilibus nuculisque lanatis differt. *T. parviflorum* et *macrum* staminibus inclusis, nuculis breviter hirtis distinguuntur.

11. T. Oliverianum (Gingins in Bth. Lab. p. 668) molliter canescenti-velutinum pluricaule caulinibus erectis rigidis crassiuseulis fere a basi opposite ramosis pyramidatis, foliis cuneatis apice breviter et obtuse 3-3-dentatis dente medio latiore infimis subpetiolatis, floribus sensim diminutis tridentatis et summis minimis elliptico-linearibus integris, pedunculis unifloris calyce longioribus, calycis tomentosi campanulati dentibus late lanceolatis acentiusculis tubo sublongioribus, corollâ hirtâ latâ calyce subtriplo longiore laciniis acutis vel acuminatis inferiore parum elongata, filamentis longe exsertis inferne hirsutis, nuculis magnis pilosissimis ♀.

Hab. in Assyriâ prope Bagdad (Oliv! Auch. 1603!), Persiaustrali prope Daleki (Haussk!).

Pedale et sesquipedale, folia inferiora cum petiolo sesquipollucaria 7-8 lineas lata, flores eis *T. Orientalis* vix minores. Foliorum forma, nuculis magnis, etc. a praecedentibus distinctissimum.

12. T. macrum (Boiss. et Haussk.) breviter tomentellum vel glabrescens, caulinibus gracilibus erectis superne breviter et anguste paniculatis paucifloris, foliis subsessilibus in laciniis anguste lineares convolutas breves confertas bipinnatisectis, pedunculis gracilibus 4-3-floris, pedicellis filiformibus bracteâ et calyce longioribus, calycis brevis tomentelli vel glabri dentibus triangularibus acutis tubo subbrevioribus, corollâ calyce 2½-plo longiore laciniis superioribus vix inaequilibus acuminatis intermediâ parum longiore acutâ, filamentis subinclusis, nuculis breviter pubescentibus ♀.

Hab. in graminosis et dumosis regionis alpinæ montium Sawers et Nur Persiae Australis 6-10000' (Haussk!).

Caulis pedales, flores minores eis *T. Orientalis* a quo corollæ lobis superioribus acuminatis, inferiori non elongato, filamentis subinclusis, nuculis hirtis differt *T. parviflorum* caulinibus crasse tetragonis et late patuleque pyramidatis, corollæ minoris laciniis obtusis distinguuntur.

13. T. parviflorum (Schreb. Unilab. p. 31 cum icono) breviter tomentellum canescens vel canum, e rhizomate indurato pluricaule, caulinibus crassiuseulis quadrangulis a medio patule et longe ramosis late et breviter pyramidatim paniculatis, foliis in laciniis anguste lineares margine revolutas bi-trternatim vel pinnatim sectis, pedunculis unifloris divaricatis bracteâ et calyce multo longioribus, floribus parvis, calycis tomentelli campanulati dentibus lanceolatis acutis tubo

subæquilongis, corollæ cœruleæ extus tomentellæ calyce sesquilongiore lobis obtusiuseulis, filamentis inclusis, nuculis brevit pilosis ♀.

Hab. in cultis, collibus, neglectis, Anatolia borealis ad Adikiöi (Wied!), Lycaonia ad Karaman et in planicie ad radices Karadagh (Heldr!). Armenia Turcica (Touru! Huet!), Syria borealis ad Aintab et Aleppo (Auch. 1597! Haussk! Ky. 225!), desertum Syriacum (Bld.), regio inferior montium prop. Latakieh (Boiss!), Antilibani radices (Ehrenb! Boiss!), Damascus (Ky!), Mesopotamia (Haussk!), in Persia prope Kasbin (Bge!).

Paniculæ latæ brevi divaricata insigne, flores eis *T. Orientalis* duplo minores.

SECT. II. STACHYOBOTRYS.

14. T. Hyrcanicum (L. Sp. 789) perenne, pubescens, caulis herbaceis elatis ramosis paniculatis rarius simplicibus, foliis subtus canescens petiolatis a basi cordata vel truncata ovato-oblongis obtuse multierenatis, spicis longis cylindricis gracilibus densis, bracteis linearis-subulatis calyce subbrevioribus, calyce pedicello breviori erecto-patulo suffulto nutanti patule villosa campanulata bilabiata fructiferò parum aucto dentibus breviter lanceolatis superiore latiore ovato membranaceo reticulato, corollæ purpureæ extus hirta calyce vix sesquilongiore, filamentis exsertis ♀. Ic. Bot. Mag. tab. 2013.

Hab. in sylvaticis et dumosis Ineretiae (Rupr!), Iberia Caucasicae (M.B. Holen!, C.A. Mey!), Persia borealis in prov. Ghilan et Mazanderan (Auch. 5175! Buhse!), ad Asterabad (Karelin!).

Spica 4-8-pollicaris.

15. T. lamiifolium (Urv. Enn. p. 64) perenne villosum griseum, caulis ascendentibus crassiusculis herbaceis simplicibus vel parce ramosis, foliis petiolatis ovatis basi cordatis vel truncatis obtuse dentatis, spicis densissimis oblongis demum breviter cylindricis, bracteis subulatis plumosis calyce longioribus, floribus horizontalibus demum subdeflexis breviter pedicellatis, calycis villosohispidi fructiferi aucti bilabiati dente supreino maximo ovato obtuso apiculato sursum replicato membranaceo-reticulato, internediis parvis ovatis obtusis inferioribus lanceolato-subulatis, corollæ albida calycem parum superante, staminibus inclusis ♀. *T. Arduini* Fl. Græc. VI, p. 26, tab. 581 et Griseb. Spicil. non L. — *Scutellaria Cretica* L. Sp. 836. — *Teucr. densillorum* Vis. Sem. II. Patav. 1847, p. 4.

Hab. in dumosis Thraciae (Urv.), circa Byzantium et ad radices Olympi Bitihui (Auch! Boiss!), Lydia prope Trianda et ad Sipylum (Boiss!), Caria (Pin!), Cilicia (Ky!), Syria in monte Cassio et in Libano (Ehrenb! Boiss! Bll. Gailli!).

Sæpe confusum cum *T. Arduini* e Dalmatiæ indumento breviori adp. essor, foliis non cordatis, bracteis flore brevioribus, calycis dente superiore plano oblongo spinuloso-acutissimo omnino alieno.

SECT. III. SCORODONIA.

16. **T. Kotschyanum** (Pœch in Flora 1844, p. 454) ad pressissime hirtellum pallide virens, caulis erectis elatis foliosis simplicibus vel tristachyis, foliis breviter petiolatis a basi rotundata vel cuneatâ oblongis obtusis minutissime crenulatis superioribus sessilibus, floribus oppositis pedicellis calyce brevioribus suffultis longe et laxiuscule spicato-racemosis, foliis floralibus oblongo-rhombeis attenuato-acuminatis inferioribus calyces superantibus, calycis patule hirti breviter campanulati bilabiati dentibus triangulari-ovatis longe spinulosis supremo duplo majore ovato mucronato, corollæ ochroleucae hirtæ tubo breviter exerto, filamentis exsertis ♀. *T. Smyrneum* Boiss. Diagn. Or. Ser. I, 5, p. 42. — *T. Scorodonia* Sm. Prodr. I, p. 392 non L. (ex loc.)

Hab. in sylvaticis montium Lydiæ prope Smyrnam (Boiss!), Ciliciâ Trachæa ad Ananur (Per!), Ciliciâ in montibus pone Mersina (Bal. 527!), monte Troodos Cypri (Ky. 921!), forsitan in Peloponneso (Sibth.).

Pedale et sesquipedale, affine *T. Scorodoniam* foliis crenatis cordatis, bracteis minimis ovatis atque *T. Baeticum* Boiss. et *Siculo* Guss. eisdem fere notis et calycis dentibus minus longe spinosis distinctis.

17. **T. Massiliense** (L. Sp. p. 789) patule tomentellum canescens ex rhizomate indurato pluricaule, caulis herbaceis ascendentibus simplicibus vel parce ramosis, foliis parvis rugulosis petiolatis ovato-oblongis crenatis basi rotundatis vel subcordatis, floribus oppositis pedicellis calyci subaequali longis suffultis racemum laxum secundum formantibus, bracteis ovato-rhombeis acutis petiolulatis calyce brevioribus, calycis breviter campanulati hispiduli dentibus ovatis spinulosis supremo majore ovato demum concavo, corollæ purpureæ tubo inclusa ♀. Ic. Jacq. Vindob. tab. 94.

Hab. in Cretæ rupibus et muris umbrosis inter Castaneta, Euneachoria, Lassiti, etc. (Sieb! Heldr! Raut!).

Speciminiibus Gallicis et Italiciis simile.

Ar. Geogr. Hispania et Gallia australes, Corsica, Sardinia.

SECT. IV. SCORDIUM.

18. **T. Scordium** (L. Sp. 790) perenne villosum pallide virens, radice repenti stolonifera, canibus ascendentibus cæbre foliosis, foliis sessilibus oblongo-ellipticis multicrenatis caulinis basi rotundatis, ramicis basi attenuatis, floralibus flores superantibus, ramulis floralibus axillaribus anguste paniculatis, floralibus 2-4 unilateralibus pedicellatis, calycis hispido-glandulosi campanulati basi obliqui gibbi dentibus lanceolatis acutis tubo duplo brevioribus, corolla purpuras-

cente calyce duplo longiore, antheris subexsertis ♀. Ic. Fl. Dan tab. 593. — *T. palustre* Lam. Fl. Fr. II. p. 411.

Hab. in humidis et uliginosis, Thracia (Friv!), prov. Caucasicæ (Ledeb.).

Ar. Geogr. Europa borealis et media ab Angliâ et Gothiâ ad Rossiam medium et Hispaniâ et Italiâ centralibus ad regionem Danubialem, Sibiria.

19. *T. scordioides* (Schreb. Unilab. p. 37) perenne lanuginosum, radice repenti, caulis sèpius proceris foliosis, foliis sessilibus ovato-oblongis crenatis canlinis basi cordato-amplexicanibus, rameis basi rotundatis, floralibus flores superantibus, paniculâ longâ angustâ, floribus 2-4 axillaribus pedicellatis, calycis hispido-glandulosi campanulati basi obliqui gibbi dentibus lanceolatis acutis tubo duplo brevioribus, corollâ purpurascente calyce duplo longiore, staminibus subexsertis ♀. *T. lanuginosum* Hoffm. et Link. Fl. Portug. p. 84, tab. 3.

Hab. in humidis totius regionis specie praecedenti vulgatius a Graeciâ ad Macedoniâ et Thraciam (Griseb.), Tauriam et prov. Caucasicas (Ledeb!), Persiam boreali occidentalem (Szov!), Susianam (Haussk!), Mesopotamiam (Kyl!), Syriam et Libanum (Bl! Gailli).

An varietas *T. Scordii* a quo differt præsertim foliorum formâ?; elatius est, plus minus lanatum.

Ar. Geogr. Anglia, Europæ regio mediterranea, Africa borealis.

20. *T. serratum* (Bl! in D.C. Prodr. 42, p. 586) perenne villosum viscidulum canescens, caulis proceris foliosis, foliis sessilibus a basi subcuneatâ oblongis argute serratis, floralibus breviter lanceolatis serratis acutissimis flores superantibus, ramis longis gracilibus paniculatis, verticillastris 2-4-floris remotiuseulis, pedicellis calyce sublongioribus, calycis hispidi breviter campanulati basi obliqui dentibus lanceolatis acutissimis dimidio tubo longioribus, corollâ purpurascenti calyce subdupo longiore, staminibus subexsertis ♀.

Hab. in regno Cabulico circa Jugdulluk et Khuggali (Griff!).

Planta 1-2-pedalis, species affinis *T. Scordio* a quo foliis argute serratis floralibus acutissimis præsertim differt.

21. *T. melisoïdes* (Boiss. et Haussk. exs. 1867) perenne adpressissime velutino-canescens vel canum, radice repente multicauli, caulis elatis longe paniculatis, foliis subsessilibus a basi brevissime cuneatâ vel rotundatâ elliptico-oblongis obtusis crenatis, floralibus sensim diminutis flores superantibus, verticillastris 2-4-floris remotiuseulis, pedicellis calyce longioribus, calycis cani basi obliqui campanulati dentibus lanceolatis acutis tubo subbrevioribus, corollâ albâ basi intescente calyce duplo longiore, staminibus exsertis ♀.

Hab. ad rivulos et in humidis regionis inferioris et montane in Persiâ anstro-occidentali, districtus Kuh Kilouyeh, prope Bebehan et in montibus Ayroman et Schahu Kurdistaniæ Persicæ (Haussk!).

Caules bipedales, inflorescentia sepe ramosissima. A *T. Scordio* eximie distinctum caulis rigidis, indumento adpressissime tomentoso-cano, pedicellis elongatis.

SECT. V. SPINULARIA.

22. **T. spinosum** (L. Sp. 793) annum patule pubescens pumilum, caule erecto crassiusculo fere a basi decussatim ramosissimo, ramis divaricatis pyramidatis apice demum spinescentibus, foliis inferioribus cito evanidis petiolatis oblongis in laciniis obtuse 3-5-fidias pinnatipartitis, superioribus sessilibus sensim diminutis dentatis, summis minimis integris ovatis, verticillastris remotis 2-4-floris, bracteis in spinas rigidas calyces subsuperantes mutatis, calyeibus pedicellis eis brevioribus suffultis demum deflexis hispido-glandulosis tubo ad basin subtus gibbo, dentibus spinosis superiore lato ovato extus concavo inferioribus quaternis subulatis, corollæ albidæ calyce vix duplo longioris tubo inclusa, antheris vix exsertis ♂. Ic. Fl. Græc. tab 539. — *T. mucronatum* L. Sp. p. 793. — *Scordium spinosum* Cav. Ic. I, p. 19, tab. 31.

Hab. in cultis, Anatoliâ occid. (Sibth.), ad Brussam (Pichler!). Ciliciâ littorali (Ball), Syriâ circâ Aleppo (Auch. 1580!), Galilæâ (Boiss!).

Ar. Geogr. Insulæ Canarienses, Lusitania et Hispania, Sicilia et Italia australis, Africa borealis.

SECT. VI. ISOTRIODON.

23. **T. pœderotoides** (Boiss. et Haussk.) patule velutino-tomentosum viscidulum griseum, ramis ex caudice indurato numerosis tenuibus erectis crassis foliosis in spicam ovato-oblongam densifloram abeuntibus, foliis subsessilibus cuneato-oblongis obsolete paucierenatis sublus elevatim nervosis inferioribus obtusis superioribus acutiusculis spicam involuerantibus, bracteis oblongis acuminate-attenuatis flores superantibus, floribus parvis brevissime pedicellatis, calycis hispiduli dentibus dimidio tubo sublongioribus superioribus tribus oblongis obtusiusculis inferioribus paulo angustioribus lanceolatis acutis, corollâ roseâ hirtâ calyce sesquiloniore, antheris exsertis ♀.

Hab. in fissuris rupium Syriæ borealis supra Assy non procul ab Aintab (Haussk!). Fl. fine Junii.

Collum reliquis ramorum persistentibus strictis horridum, rami cum spica 9-10 lineas longâ 2-3-pollicares, folia 6-9 lineas longa, calyx $1\frac{1}{4}$ lineas longus; facies quodammodo *Pœderotæ Bonarotæ* omnibus partibus diminutæ. Species distinctissima.

24. **T. Montbreti** (Bth. Ann. Sc. Nat. Ser. II, VI. p. 56) mollier velutino-tomentosum canescens, ramis e rhizomate indurato numerosissimis tenuibus fragilibus siexuosis intricatis, foliis breviter petiolatis a basi truncata ovatis vel oblongis obtuse et grosse crenatis, bracteis linear-lanceolatis acutis calyce brevioribus, spicis densis

ovalo-oblongis, calycis pedicello subæquilongo suffulti cernui basi gibbi campanulati hispidi dentibus ternis superioribus ovatis obtusis inferioribus subangustioribus lanceolatis acutis omnibus tubo duplo brevioribus, corollâ e sicco rubellâ calyce duplo longiore, antheris subexsertis ♂.

Hab. ad rupes verticales regionis inferioris Syriæ borealis, in monte Cassio (Auch. 4594!), ad Antiochiam, Seleuciam et in montibus inter Darkusch et Edlip (Boiss!), Tauro Cilicico (Ky. 440!), in Pamphyliâ circa Adalia (Bourg. sp. sterilia).

Rami 2-10-pollicares, folia 6-7 lineas longa sed saepe majora, spicæ pollicares. Planta fragillima.

25. T. Odontites (Boiss. et Bal. Diagn. Ser. II, 4, p. 57) perenne, ex collo indurato ramosissimum ramis filiformibus fragilibus hirtellis decumbentibus flexuosis simplicibus vel parce ramulosis, foliis hirsutis supra minute papilloso-pruinosis subtus velutinis brevissime petiolatis ovato-orbiculatis late et obtuse paucicrenatis basi rotundatis vel subcordatis sursum sensim diminutis, floralibus oblongo-linearibus integris pedicellis subbrevioribus, floribus binis oppositis pedicellis calyce longioribus suffultis racemos laxos paucifloros breves rarius ramoso-subpaniculatos formantibus, calycis breviter hirtuli campanulati nutantibas gibbi dentibus tubo duplo brevioribus superioribus tribus ovatis acutiusculis, inferioribus binis sublongioribus acutiusculis, corollâ purpureâ tubo subduplo longiore, staminibus exsertis ♂.

Hab. in fissuris rupium verticalium faucis Guzel Dere Ciliciæ littoralis supra Sedichig (Bal. exs. 525!). Fl. Junio.

Rami innumeri intricati tenuissimi 6-6-pollicares, folia majora 3-5 lineas diam. lata, racemi 1-2-pollicares, flores eis *T. Chamaedrys* dimidio minores. Crescendi modo et calyce *T. Montbreti* affine eximie differt ramis tenuissimis, foliis triplo minoribus, racemo laxissimo.

26. T. Persicum (Boiss. Diagn. Ser. I, 5, p. 43) totum patule et longe tomentoso-hispidum canum basi suffrutescens et ramosissimum, ramis erectis pumilis simplicibus vel divaricatum ramulosis valde fragilibus, foliis parvis in petiolatum brevissimum attenuatis inferioribus oblongis apice obtuse 5-7-dentatis superioribus integris, floribus oppositis pedicellis calyce brevioribus suffultis longiusculæ laxiusculæ et anguste racemosis, foliis floralibus elliptico-linearibus obtusiusculis calyces paulo superantibus basi longe attenuatis, calycis hispido-lanati dentibus calycis tubo brevioribus superioribus ternis triangulari-oblongis acutis inferioribus binis angustioribus lanceolatis sublongioribus, corollâ roseâ hispidulâ calyce subduplo longiore, staminibus exsertis ♂.

Hab. in Persiâ australi inter Tarun et Darap (Auch. 5170! et 5171!).

Vix semipedale, rami crassiusculi sed fragiles, folia 5-6 lineas longa, calyx tubuloso-campanulatus 2 $\frac{1}{2}$, lineas longus, facies *T. fragilis* Boiss. Hispanici,

SECT. VII. CHAMÆDRYS.

27. T. Chamædrys (L. Sp. p. 790) viride, glabrescens breviter hirtum vel patule hirsutum canescens, rhizomate stolonifero ramos plurimos basi radicantes ascendentibus herbaceos in racemum brevem oblongum folialum unilateralem abeuntibus, foliis parvis firmis petiolatis in lobos oblongos saeppe subdentatos incisis basi cuneatis, floralibus diminutis calyces superantibus ellipticis saepius dentatis, verticillastris 2-6-floris, pedicellis calyce subbrevioribus, calycis rubelli pubescentis basi subgibbi dentibus lanceolatis acutissimis subæqualibus tubo subbrevioribus, corollâ purpureâ calyce $2\frac{1}{2}$ -plo longiore, staminibus exsertis ♀. Ic. Engl. bot. tab. 780. — *T. Nuchense* C. Koch Linn. 21, p. 704.

Hab. in rupestribus, dumosis siccis totius fere regionis in regione montanâ et subalpinâ, montibus fere omnibus Graeciae et Peloponnesi!, Macedoniae! et Thracie!, Anatoliâ boreali (Wied!), Armeniâ (Huet!), et Ponto Lazico, a littore mari ad 6000' (Balt!), Olympo Bithyno (Boiss!), montibus Cappadocie (Balt!), Tauro Cilicie 3500' (Ky!), monte Amano Syriæ bor. (Ky!), Tauria Caucaso et Transcaucasiâ (M.B., Ledeb! Ruprl!), Persiâ boreali in provinciis Ghilan et Asterabad (Buhse! Bge.).

β. canum. — Folia utrinque vel subitus cano-tomentosa. — *T. canum* Fisch. et Mey. Ind. 1835, p. 40 — *T. Sysspirense* C. Koch Linn. XXI, p. 704.

Hab. in aridis saxosis Armeniæ (C. Koch!), Somchetiæ et Georgiae Caucasiæ (Hohen!), Anatoliâ boreal! in montibus Tokat, Amasia (Wieden!).

Ar. Geogr. Europa omnis a Britanniâ et Belgio ad Rossiam medianam, Sibiria Uralensis, Africa borealis.

28. T. flavum (L. Sp. 791) suffruticosum-basi ramosum ramis erectis viridibus crispulis et parce hirtis foliatis in racemum spiciformem abeuntibus, foliis crassiuseulis rigidis parvis petiolatis a basi truncata vel subcuneata ovatis utrinque 3-5-crenato-dentatis, floralibus ellipticis integris calyce subbrevioribus acutis, verticillastris 4-6-floris inferioribus subdistantibus, calycis pedicello eo subbreviore suffulti campanulati basi subgibbi hirsuti glanduloso-punctati dentibus lanceolatis acutis subæqualibus tubo subbrevioribus, corollâ flavâ hispidâ calyce $2\frac{1}{2}$ -plo longiore, staminibus exsertis ♀. Ic. Fl. Græc. tab. 533. — Rehb. Ic. 35.

Hab. in rupestribus regionis inferioris et montanæ Graeciae, Pentelicus (Heldr!), Malevo, Taygetus, Kyllene, Parnassus usque ad 3000' (Heldr! Orph. Fl. Gr. exs. 541!), Zacynthus (Marg!), Creta (Sieb!), Archipelagus (Sibth.).

Ar. Geogr. Gallia australis, Corsica, Sardinia, Sicilia, Italia australis, Dalmatia, Africa borealis.

29. T. divaricatum (Sieb. exs.) suffruticosum ramosissimum ramis pumilis erectis vel ascendentibus patule tomentosis in racemum saepius laxum vel basi interruptum abeuntibus, foliis utrinque hirtis

subtus canescens parvis petiolatis ovatis obtuse paucicrenatis basi rotundatis vel subcuneatis, floralibus integris ovato-oblongis acutiusculis calyces æquantibus vel brevioribus, verticillastris 4-6-floris subdistantibus, calycis pedicello eo breviore suffulti campanulati tomentosi basi subgibbi dentibus lanceolatis acutis tubo subbrevioribus, corollæ purpureâ calyce duplo longiore, staminibus exsertis 5. *T. flavum* var. *purpureum* Bth. — *T. regium* Auct. an Schreb? — *T. lucidum* Fl. Græc. tab. 532 non L.

Hab. in rupestribus regionis inferioris et montane, Græcia in Lycabetto, Hymetto, Parnasso, Eubœa (Heldr! Orph. exs. 5421), Creta (Sieb! Heldr!), Cyclades! et Rhodus! Anatolia occid. a Byzantio (Cast!) ad Pamphyliam (Heldr!), Cyprus (Ky!), Syria maritima in parte inferiori Libani (Boiss! Bl! Gaill!), Carmelus Palestine (Boiss!), ad Hierosolymam (Roth!).

Valde affine *T. flavo* et forsitan ejus varietas, praeter colorum florum tamē differre videtur ramis humilioribus tenuioribus, indumento sepius tomentello, distributione geographicā; a *T. Chamaedrys* basi suffruticosā, foliis minus profunde dentatis, floralibus integris.

30. *T. microphyllum* (Desf. Coroll. p. 18, tab. 11) humile basi ramosissimum suffrutescens, ramis divaricatis tenuibus flexuosis adpresso canis in racemum brevem unilateralē oblongum abeuntibus, foliis minimis supra viridibus subtus canis petiolulatis cuneato ovatis crenato-dentatis, floralibus subintegris flore brevioribus, floribus binis oppositis brevissime pedicellatis, calycis adpresso canescens campanulati basi subgibbi dentibus lanceolatis subæqualibus acutis, corollæ roseâ calyce duplo longiore, staminibus exsertis 5. *T. quadratum* Fl. Græc. VI. p. 25, tab. 530, nō: Schreb

Hab. in saxosis regionis inferioris Cretæ usque ad 3000' frequens (Tourn., Sieb!, Heldr! Raulin!).

Rami 2-5-pollicares, folia $1\frac{1}{2}$ - $2\frac{1}{2}$ lin. longa. A *T. divaricato* ramis tenuibus flexuosis, foliis minimis subtus canis, verticillastris bifloris longe alienus. ***T. Marum* L.** ramis fruticosis, foliis convolutis integris distinctum in Bory Fl. Pelop. fidè schedula mendacis erroneé indicatur. *T. canum* q. *latifolium* Bth. in D.C. Prodri. Auch. 1587 bis! est *T. Marum* L! a speciminiibus Telonensis minime diversum, sed specimen neutiquam in Asia minore lectum cederem.

31. *T. leucophyllum* (Montb. et Auch. Ann. Sc. Nat. Ser. II, 6, p. 55) basi suffrutescens et ramosissimum adpressissime canum, ramis tenuibus erectis simplicibus in racemum brevem oblongum abeuntibus, foliis parvis in petiolum brevissimum attenuatis inferioribus oblongo-linearibus obtuse paucidentatis cæteris strictis linearibus margine revolutis subintegris, floralibus ovatis acutiusculis calyce brevioribus, verticillastris paucis approximatis 2-4-floris, pedicellis calyce duplo brevioribus, calycis villosuli dentibus lanceolatis acutis tubo duplo brevioribus, corollæ roseâ calyce subduplo longiore, antheris exsertis 5. *T. canum* var. *augustifolium* Bth in D.C. Prodri. 12, p. 588.

Hab. in Cappadociâ ad Euphratem (Auch. 1893 in D.C. herb!).

Semipedale, racemus sesquipollucaris, folia 6-7 lineas longa inferiora planiuscula sesquilineam cætera lineam lata, flores fere magnit. *T. divaricati* a quo ramis tenuibus adpresso canis nec crassis tomentosis foliisque angustissimis fere ut *T. Webbianum* a formis *T. Chamaedrys* differunt.

32 T. coniortodes (Boiss. et Blanche) basi suffrutescens totum adpressissime canescens tomentosum canum, ramis procumbentibus vel ascendentibus tenuibus in racemos longos laxos tenues abeuntibus, foliis minimis utrinque canis cuneato-oblongis margine subrevolutis dentibus 2-3 utrinque ascendentibus obtusiusculis, floralibus integris flores subæquantibus, verticillastris bifloris remotiusculis, calyce adpresso cano pedicello eo breviore suffulto basi gibbo dentibus subæqualibus lanceolatis acutis calyce duplo brevioribus, corollâ purpureante calyce duplo longiore, antheris subexsertis 5.

Hab. ad radices orientales montis Gebel Abiad deserti Syriaci inter Hama et Palmyram (Bl.).

Rami cum racemo 4-6-pollicares, folia inferiora tres lineas longa, facies *T. microphylli* et *leucophylli*, a priore foliis utrinque canis, a posteriore foliorum formâ, ab utroque racemo longo laxe discedens.

33. T. Socinianum, basi suffrutescens ramosissimum ramis pumilis ascendentibus herbaceis canis in racemum oblongum laxiusculum abeuntibus, foliis parvls sessilibus tomentoso-cinerascentibns marginibus convolutis linearciylindricis in lobulos ovatos crispis utriusque marginis contiguos pinnatilobatis, floralibus ellipticis acutis planis calyce brevioribus, floribus binis oppositis pedicellis calyce brevioribus suffultis, calycis campanulati viscido-hispidi basi subgibbi dentibus lanceolatis acutissimis subæqualibus tubo duplo brevioribus, corollâ purpureâ calyce 2½-plo longiore, staminibus longe exsertis 5.

Hab. in elivo orientali Antilibani prope pagum Malula 9 leucas ad septentrionein Damasci situm (cl. Dr. Socin).

Rami novelli 3-4-pollicares, racemus tandem bipollucaris, folia more *Salviae santoinofoliae* convoluta et pectinatum lobulata 5-6 lineas longa lineam lata, calyces eis *T. divaricati* vix minores. Inter affinia foliis convolutis pectinatis insigne.

SECT. VIII. POLIUM.

34 T. montanum (L. Sp. 791) basi induratum ramosissimum ramis tenuibus inferne prostratis et nudis deis ascendentibus foliosis pumilis, foliis firmis oblongo-linearibus linearibusve basi in petiolum brevem attenuatis margine subrevolutis supra viridibus subtus canis summis capitulum involuerantibus, capitulis corymbosis densifloris, bracteis linearciylindricis flores non æquantibus, calycis minute puberuli vel glabri tubuloso-campanulati dentibus tubo duplo brevioribus superioribus ovatis acutis inferioribus tribus ovato-lanceolatis acuminate, corollâ albâ calyce subduplo longiore, antheris breviter exser-

tis ♀. Ic. Fl. Græc. tab. 534. — Rehb. Ic. tab. 37. — *T. supinum* L. Sp. 791 (forma foliis angustis).

Hab. in rupestribus regionis alpinæ, Græcia in Parnasso (Heldr!), monte Athos Macedonie (Auch. 4590!), Olympo Bithyno (Sibth., Boiss.), Tauria (Stev.), Taurus Cilicicus supra Bulgharmaaden (Bal!), Cilicia Trachea (Pér!).

β. hirsutum. — Rami, folia et calyces patule hispidi. — *T. Pannonicum* Kerner.

Hab. in Olympo Thessalo et monte Delphi Eubeæ (Heldr!), monte Sipylo supra Magnesiam (Bal!).

Ar. Geogr. Europa media et australis a Belgio et Germania mediâ ad Rossiam, Soongaria.

35. *T. Mascatense* (Boiss. Diagn. Ser. I, 5 p. 44) crispule et breviter pubescens griseum, ramis tenuibus filiformibus elongatis basi indurata procumbentibus simplicibus, foliis minutis elliptico-linearibus margine convolutis basi attenuatis subpetiolatis integris rarius apice obsolete 1-2-denticulatis, capitulo terminali ovato densifloro, bracteis linearibus calyce brevioribus, calycis sessilis obconici minute puberulo-glandulosi virentis dentibus tubo subtriplo brevioribus superioribus ovato-triangularibus acutis inferioribus binis sublongioribus acutioribus, corollâ albido-flavescente calyce sesquiloniore, staminibus longe exsertis ♀.

Hab. in monte Chebek Arabiæ Mascatensis (Auch. 5172!).

Rami novelli 3-4-pollicares simplicissimi flexuosi, folia 2-3 lineas longa lineam dimidiâ lata, capitulum magnitudine avellanæ, calyx 2½ lineas longus. Facies *T. montana* a quo calyce dimidio breviore, etc. differt.

36. *T. alpestre* (Sibth. et Sm. Fl. Græc. VI, p. 31, tab. 538) patule villosulum pallide virens humile suffruticosum ramosissimum ramis vetustis tortuosis lignosis procumbentibus novellis brevibus simplicibus vel saepius opposite breviter et divaricatum ramulosis, foliis minimis subsessilibus obovato-vel rotundato-cuneatis obtuse 3-7 crenato-lobulatis nervis supra impressis subtus prominulis, floribus brevissime pedicellatis oppositis in capitula terminalia laxa 2-5-flora foliis trilobis calyce brevioribus suffulta approximatis, calycis viridis hirsuti tubuloso-campanulati dentibus tubo duplo brevioribus lanceolatis acutiusculis inferioribus binis subangustioribus, corollâ albida calyce sesquiloniore, antheris vix exsertis ♀.

Hab. in Cretæ saxosis inter frutices regionis subalpineæ 4500'-5000' (Sibth., Sieb! Heldr! Rauli!). Fl. Jun.

Rami vetusti suffrutescentes breves valde intricati et tortuosi, hornotini 2-3-pollicares, folia 2-3 lineas longa concoloria pallide virentia, capitula avelanæ minora eis *T. Polii* et affl. laxiora. Calyx vix 2 lineas longus.

β. majus. — Rami crassiusculi firmiores 3-5-pollicares divaricatum ramulosiores, folia submajora obovato-oblonga niargine sepe revoluta, capitula majora 7-12-flora densiora.

Hab. in saxis apricis prope Norisa ad latus meridionale montis Idæ et in arenosis insulae Gaidaronisi (Heldr.).

Forma e locis demissioribus.

37. T. Cyprium (Boiss. Diagn. pl. Or. Ser. I, 5 p. 43) patule bipnidum griseum suffruticosum cespitosum ramosissimum humile ramis decumbentibus novellis brevibus flexuosis simplicibus vel parvœ ramulosis, foliis miutis subsessilibus obovato-cuncatis obtuse crenatis, capitulis terminalibus ovato-globosis, foliis floralibus ellipticis integris flores subæquantibus, calycis hispidi et glandulis adspersi tubuloso-campanulati dentibus tubo quintuplo brevioribus ovatis acutiusculis inferioribus binis subacutioribus, corollæ albida calycem parum excedente, staminibus subexsertis ♂.

Hab. in pinetis regionis superioris montis Troodos Cypri 600' (Auch. 155!, Ky. 783!).

Rami novelli 2-3-pollicares, folia majora 4 lineas longa, calyx 3 lineas longus. Affine *T. alpestri* a quo differt ramis subsimplicibus, capitulis majoribus, foliis floralibus integris calyces subsuperantibus, calycis dentibus abbreviatis.

38. T. leucocladum (Boiss. Ser. I, 5, p. 44) fruticulosum totum adpressissime canum humile a basi ramosissimum ramis brevibus crassis rigidis opposite et divaricatim ramulosus tandem indurato-subpinescentibus, foliis minimis subsessilibus cuneato-obovatis obtuse paucicrenatis, floralibus capitulo brevioribus ellipticis integris, capitulis parvis subsphaericis dense paniculatis breviter pedunculatis, bracteis ellipticis obtusis, floribus minimis sessilibus, calycis cani dentibus brevissimis ovatis obtusis, corollæ albida calyce sesquiloniore, antheris exsertis 5. *T. cuneifolium* Herb. Un. itin. non Fl. Graec.

Hab. ad rupes Quadi Hebræa Petreæ (Schimp. 145!).

Fruticulus semipedalis ramis vetustis lignosis tortuosus, rami novelli albi 2-3-pollicares ramulique divaricati rigidissimi, folia majora 4 lin. longa apice 3 lata, capitula piso vix majora, flores minimi. Species dentibus rotundatis calycis (vix ultra lineam longi) insignis a *T. cuneifolio* omnino aliena, magis affinis *T. alpestri* quod indumento crispo-lanato, ramis tenuibus non rigidis, floribus duplo majoribus, calycis dentibus cæterum longe differt.

39. T. cuneifolium (Sibth. et Sm. Fl. Graec. VI, p. 30, tab. 537) dense cano vel flavescenti-tomentosum ramosissimum ramis crassis inferne nudis procumbentibus superne ascendentibus ramulosus crebre foliosis, foliis crassiusculis cuneato-ovatis vel rotundis basi attenuatis obtusissime multicrenatis, verticillastris paucis in capitula terminalia ovata approximatis, foliis floralibus integris calyce brevioribus, calycis dense tomentosi tubulosi dentibus tubo quadruplo brevioribus subæqualibus obtusiusculis, corollæ albæ calyce vix duplo longiore, staminibus breviter exsertis ♂.

Hab. in glareosis et rupestribus maritimis regionis inferioris Cretæ (Sibth., Sib., Heldr. Raal!).

Rami semipedales et longiores, folia majora 7-9 lineas longa, capitula siepe 2-3 pauciflora in unicum laxum approximata. Inter affinita indumento dense tomentoso insigne.

40 T. Stocksianum (Boiss. Diagn. Ser. II, 4, p. 58) adpresiusculæ et breviter velutinum canescens basi suffruticosus ramosissimum, ramis novellis erectis pumilis fragilibus simplicibus vel opposite ramulosis, foliis parvis subsessilibus ovatis vel obovatis cuneato-attenuatis obtusis parte superiori obtusiusculæ 7-9-crenatis, capitulis terminalibus ovato-globosis, foliis floralibus ellipticis integris flores subsuperantibus, calycis campanulati crispule pubescentis dentibus tubo quadruplo brevioribus inferioribus sublongioribus acuminatis intermediis binis ovatis acutis superiore obtuso sappiis tridentato, corollæ e sicco albida calyce sesquiloniore, staminibus exsertis ♀.

Hab. ad collum Hurbab montium Hala Belutschiae 5000' (Stocks!).

Semipedale, folia inferiora 4-5 lineas longa, capitulum magn. avellanæ, calyx latiusculus fere 4 lineas longus.

41. T. Polium (L. Sp. p. 792) tomentosoc-anescens vel canaliculatum basi suffruticosus ramosissimum, ramis ascendentibus vel erectis apice breviter paniculatis vel sappiis corymbosis, foliis sessilibus oblongis vel linearibus obtusis crenatis, capitulis densilloris subsphaericis vel ovatis breviter pedunculatis, bracteis linearispatulatis flores non æquantibus, calycis subsessilis tomentosi breviter campanulati dentibus villo occultatis breviter triangularibus acutiuseulis vel obtusis subæqualibus, corollæ calycem parum superante, antheris vix exsertis ♀. Ic. Rehb. tab. 37. Species valde polymorpha.

α. vulgare. — Folia oblonga plana vel margine revoluta, capitula majora lanata. Corolla alba. — Ic. Fl. Græc. tab. 535. — *T. pseudohyssopus* Schreb. Unil. p. 45.

Hab. in rupestribus et collinis siccis regionis inferioris et præsertim montane in alpinam ascendens ubique a Græciâ! et ejus insulis! ad Macedoniam! et Thraciam!, Anatoliâ omnem maritimam! et interiorem in Paphlagoniâ, Galatiâ (Wied!), Cappadociâ (Balt.), Armeniâ (Calv!), Syriâ maritimâ! interiori ad Rascheya (Boiss!) et Damascum (Gaill!), in Libano usque ad Eden (Blt.), Palestinâ (Boiss! Roth!), Ägypto ad Alexandriam (Ehr!), et in planicie elata Galala (300') deserti Ägyptico-Arabici (Schweinf!), Mesopotamia, Assyriâ et Kurdistaniâ Persicâ (Ky! Haussk!), Persiâ boreali (Szov! Buhse, etc.), et australi in montibus Kuh Kilouyeh 8000' (Haussk!), Caucaso!, Transcaucasiâ! et Tauriâ.

β. roseum. — Simile var. α sud corollæ roseæ. — *T. capitatum* Fl. Gr. tab. 536.

Hab. in insula Zæyntho et alibi (Sibth.), Cephallenia insula (Heldr!), Cyprus (Ky. 925!).

γ. angustifolium Bth. — Folia angustiora magis convoluta linearia, rami graciliores, capitula minora. — *T. capitatum* L. Sp. 792.

Hab. in siccis, formâ vulgari in Oriente multo rarius, Macedonia (Griseb.), Bœotiam (Auct. 1591!), mons Sipylus supra Magnesiam Lydiæ (Boiss!), Aegæo Syriæ (Ky. 2361), mons Carmelus Palestinæ (Gaill!), Persia australis, monte Kuh Delu (Ky. 591!), Sawers et Eschker (Haussk!).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, Dalmatia, Rossia media et australis, Asia borealis.

42. T. Sinaicum (Boiss. Diagn. Ser. I, 12 p. 91) fruticosum ramosissimum patule et dense hirsutum griseum, ramis erectis tenuibus strictis simplicibus vel parce ramulosis, foliis sessilibus concoloribus oblongo-linearibus margine convolutis obtusis margine pectinatim obtuse et contigue lobulatis, capitiulis terminalibus globosis densis solitariis vel 2-3 approximatis foliis floralibus eis brevioribus suffultis, bracteis linearibus plumosis, calycis glanduloso-punctati et hirsuti dentibus brevibus superioribus triangulari-ovatis inferioribus binis sublongioribus paulo angustioribus, corollâ albâ calyce sesquio longiore, staminibus breviter exsertis *s. T. Polium* var. *pilosam* Decaisne Flor. Sin. p. 12.

Hab. in rupestribus montis Sinai Arabiæ petreæ (Bové! Schimp. 204! Boiss!). Arabice *Jaada*.

A fine *T. Polio* sed ab omnibus ejus formis ramis vetustis pedalibus vere fruticosis crassitie pennæ anserinæ, indumento totius plantæ non tomentoso nec lanato sed tenue natulo viscidulo distinctum; foliorum margines insuper regularius pectinatobrullati.

ORD. XCV. PLUMBAGINEÆ.

(Endl. Gen. p. 410.)

Flores regulares hermaphroditæ. Calyx gamosepalus rarissime pentaphyllus infundiduliformis limbo scarioso quinquelobo plicato, rarissime herbaceus tubulosus. Corolla hypogyna saepius tenera, vel quinquepetala petalis unguiculatis vel gamopetala infundibuliformis limbo quinquepartito æstiv. contorto. Stamina quinque corollæ lobis vel petalis opposita. Ovarium sessile liberum uniloculare uniovulatum ovulo ex apice funiculi liberi e fundo ovarii ascendentis pendulo. Styli à filiformes vel a basi liberi vel in unicum basin ad medium, vel omnino concreti. Stigmata filiformia vel capitata. Fructus utricularis calyce inclusus subcoriacens, vel tenuiter membranaceus subindehiscentia vel operculo aut valvis irregularibus dehiscens. Semen loculum ampleius inversum, testa membranacea albumini farinacco adnatâ, embryone recto axili. -- Ordo naturalissimus nulli arcte affinis, ne ad *Primulaceas* illinc ad *Plantagineas* quodammodo accedens.

TR. I. STATICEÆ Bartl. Ord. p. 127. — Stylo a basi vel saltem parte superiore liberi. Pericarpium utricularare vasi lacrum vel apice circumscissum.

ACANTHOLIMON (Bois. Diagn. Ser. I, 7, p. 69).

Calyx infundibuliformis tubo tenui, limbo scarioso quinque nervi 5-10-lobo. Petala quinque imâ basi annulatim connata longe unguiculata. Filamenta quinque libera glabra. Ovarium cylindrico-lucare in stylos attenuatum quinos glabros basi cohærentes. Stigmata capitato-depressa. Utriculus crustaceo-inembranaceus indehiscens vel ad angulos fissilis — Suffrutices erinacei foliis acerosis vel spinosis, floribus in spiculas uni vel 2-5-floras tri vel 4-5-bracteatas varie dispositas confertis, rachide fragillima. — Genus ex structurâ consimili et ideo characterum graviorum deficientiâ hand facile coordinandum, variabilis florum in unâquâque spiculâ numerus, variabilis quoque intra certos limites inflorescentia, &c. ceterum ex nimiâ ejus partium semper articulatarum fragilitate in speciminibus siccis non semper rite intelligi potest. Characteres primarios minus fugaces cum amic. a Bunge ex nervis calycis facie interiore glabris vel hirtis et cum cl. Spach et Bunge ex foliorum structurâ deprompsi; ea nempe vel *vernalia* vere jam evoluta ad partem inferiorem rami hornotini in rosulam sæpius contracti obvia, alia *aestivalia* serius evoluta in rosulâ superiora et interiora sunt; in specierum majore numero subconformia sed in sectione *Tragacanthiná* heteromorpha. Confer de *Acantholimone* accuratissimam monographiam cl. et am. a Bunge Petrop. 1872 quam paucis mutatis secutus sum.

Specierum Orientalium distributio.

SERIES I. — Calycini limbi nervi facie superiori (internâ) villosi.

SECT. I. CYMARIA Bunge. — Inflorescentia dichotome cymosa spiculis 2-3-floris dense et brevissime spicatis. — Folia aestivalia triquetra spinescentia margine lævia.

A. cymosum.

SECT. II. PTEROSTEGIA Bunge. — Inflorescentia capitulis 1-2 parvis concreta terminalibus constans, spiculæ 2-3-floræ. Folia omnia margine scabrida.

A. pterostegium.

SERIES II. — Calycini limbi nervi facie superiori (internâ) glabri.

Subseries I. — Folia homomorpha nempe vernalia aestivalibus similia quamvis breviora, omnia persistentia acerosa et margine acuto scabrida. Confer *A. diapensioides*.

SECT. III. ACMOSTEGIA Bunge. — Inflorescentia spicis brevibus distichis 3-4 in capitulum terminale confertis constans. Spiculæ inferiores bisporæ cæteræ unifloræ. Calycis tubulosi limbus minutus.

A. acmostegium.

SECT. IV. ARMERIOPSIS Boiss. — Inflorescentia spicâ terminali brevi capitulif. rmi constans. Spiculæ 2-5-floræ 4-5-bracteatæ bracteis exterioribus latissimis late membranaceis. Calyx infundibuliformis.

A. bracteatum, cephalotes, Cabulicum, distachyum.

SECT. V. GLUMARIA Boiss. — Spiculæ 2-3-floræ rarius abortivæ unifloræ in spicam plus minusve longam et laxam saltem basi interruptam dispositæ. Calyx infundibuliformis.

A. latifolium, Tataricum, Griffithianum, scirpinum, bromifolium, restiacum.

SECT. VI. PULVINARIA. — Spica 1-3-spiculata fasciculiformis in rosulâ sessilis, spiculæ 1-2-floræ 4-5-bracteatæ. Folia minima columnariter imbricata mutica nec ut in congeneribus omnibus acerosa

A. diapensioides.

SECT. VII. STATICOPSIS Bunge. — Inflorescentia spicata, spiculæ unifloræ tribracteatæ.

§ 1. **Rhodocalycina** Bge. — Calyx $4\frac{1}{2}$ lineas longus vel longior limbo purpureo.

* Scapus 2-pleiostachyus, spicæ densifloræ.

A. avenaceum, Calverti.

** Scapus monostachyus spicâ plus minus laxiflorâ.

A. venustum, laxiflorum, petraeum, Atropatanum, Senganense.

§ 2. **Caryophyllacea.** — *Caryophyllacea* et *Androsacea* Bge. — Calyx $4\frac{1}{2}$ lineas vel amplius longus limbo albo.

* Scapus sæpius pleiostachyus.

+ Spicæ laxifloræ elongatæ.

A. tenuiflorum, Araxanum.

+ + Spicæ breves dense et distiche imbricatæ.

A. Hohenackeri, Huetii, glumaceum, Sahendicum, Bodeanum, Lycaonicum, Cesareum.

• Scapus semper monostachyus.

A. Auganum, acerosum, caryophyllaceum, Baltanense, Armenum, Iconicum, Kotschyî, Libanoticum, Echinus, Wiedemannii, Peronini, Fausti.

§ 3. **Microcalycina** Bge. — Calyx tres ad summum quatuor lineas longus limbo fere semper (conf. *A. melananthum*) albo Hujus gregis species nonnullæ inter se valde affines ex speciminum majore copiâ et observatione in loco natali forsan reducendæ.

* Spicæ laxæ imbricatae.

A. lepturoides, Aspadanum, festucaceum, flexuosum, oliganthum, Eschke-rencse, brachystachyum.

• • Spicæ dense imbricatae.

+ Spicæ angustæ.

A. scabrellum, subulatum.

+ + Spicæ late breves.

A. melananthum, genistoides, brachyphyllum.

Subseries II. — Folia heteromorpha nempe vernalia (inferiora) carnosula sæpius margine scabrida cito fracta vel evanida, aestivalia (ramuli vel rosæ superiora) spinosa cylindrica vel superne planiuscula sed semper margine obtuso laevissima.

SECT. VIII. TRAGACANTHINA Bge.

§ 1. **Stenostomata** Bge. — Calycis subtubulosi limbus angustus plicatus.

* Spicæ secus scapum plures sessiles spiculis dense imbricatis.

A. polystachyum.

• • Spicæ e scapo sæpius plures laxissime plurispiculatæ.

A. Karelini, curviflorum, quinquelobum, Heratense, truncatum, longiflorum.

§ 2. **Eurytomata.** — *Eurystoma* et *Erythrostoma* Bge.

— Calycis infundibuliformis limbus ampliatus.

* Calycis limbus albus.

+ Spicæ laxæ vel laxiusculæ.

A. aristulatum, tragacanthinum, Scorpius, Stocksi.

+ + Spicæ breves distiche vel fasciculatim densifloræ.

1. Folia glabra.

A. fusciculare, horridum, viscidulum, Schirazianum.

2. Folia tomentella.

A. tomentellum, Talagonicum, Schahrudicum.

Calycis limbus purpureus vel roseus, spicæ densifloræ.

A. roseum, Gulistanum.

SECT. I. CYMARIA Bge.

1. **A. cymosum** (Bge l. c. p. 15) fruticulosum glaucum, ramis vaginis obsitis, foliis vernalibus brevibus planiusculis margine scabriusculis, æstivalibus e vaginâ latâ ciliatâ spinosis triquetris margine lœvibus, scapis ex axillis superioribus pancis folia parum superantibus hispidulis in cymam contractam pluries et breviter dichotomam tomentellam abeuntibus, spiculis 2-3-floris in spicas brevissimas distiche imbricatis, bracteis exterioribus tomentosis herbaceis albo-marginalis deltoideis transverse sublatioribus abrupte mucronatis, interioribus multo minoribus aristulatis, calycis tubo et limbo albo ad nervos extus et intus villoso ♂.

Hab. in rupestribus dolomitieis prov. Khorassan Persiæ prope Schahrud.

Rami in specimine 3-4-pollicares, folia æstivalia $1\frac{1}{2}$ -pollicaria, cyma densiflora diam. $1\frac{1}{2}$ -pollicaris, calyx $3\frac{1}{2}$ lineas longus.

SECT. II. PTEROSTEGIA Bge.

2. **A. pterostegium** (Bge loc. cit. p. 15) suffruticosum erinaceum glaucum ramis numerosissimis brevibus horridis, foliis primariis paucis brevibus cito emarcidis, æstivalibus rigidissimis acerosis supra planis subtus obtuse carinatis margine obsolete sebris, scapis folio subbrevioribus capitula parva 2-3 terminalia sessilia conferta ferentibus hirtulis, spiculis 2-3-floris in capitulo congestis, bracteis externis orbiculari-subcordatis centro coriaceo-herbaceis pubescentibus mucronatis margine late hyalinis, interioribus præter nervum hyalinis, calycis tubo dense hispidulo limbi nervis dilatatis longe ante marginem membranaceam albidum obsolete quinquelobum muticum evanidis intus hispidis ♂.

Hab. in rupestribus ad Tcheschine Ghilas ad orientem urbis Meschhed prov. Khorassan Persiæ (Bgel).

Specimina pauca fere arida tantum collecta, suffrutex humilis, folia longiora 1- $1\frac{1}{2}$ -pollicaria, capitula avellanâ minora, bractea exterior 4 lineas longa 5 lata, calyx 4 lineas longus.

SECT. III. ACNOSTEGIA Bge.

3. **A. acnostegium** (Boiss. et Buhse Aufz. p. 482) suffruticosum humile glabrum glaucum dense erinaceum, ramis brevibus

horridis, foliis crassis brevibus acerosis linearis-subtriquetris margine denticulato-scabris, scapis foliis brevioribus vel brevissimis, spicis 2-4 sessilibus 5-8-spiculatis brevibus distiche compressis capitulum confertis, spiculis inferioribus bifloris cæteris unifloris, bracteâ exteriore amplâ ovatâ dorso coriaceâ rufescente late albo-membranacea attenuato-acuminata subcarinata, intermediâ angustiore latius mucronata calyceum arete cingente, calycis tubulosi glabriuscui tubo in limbum angustum albido rectum quinquelobum et nervis excurrentibus mucronatum abeunte, corollâ amplâ anâne roseâ 5.

Hab. in montosis Persiæ orientalis prope Rischm ad margines deserti magni (Buhsei!), prope Damgan (Göbel), prope Schahrud (Bge!).

Folia $\frac{1}{2}$ -4-pollicaria, capitulum compositum nucis magnitudine, bractea inferior 4 lineas longa tres lata, interior 5 lineas longa. Corolla 8 lineas longa.

SECT. IV. ARMERIOPSIS Boiss.

4. A. bracteatum (Girard Ann. Sc. Nat. 1844, p. 330 sub *Statice*) cespitosum glaucescens vix suffrutescens ramis brevissimis foliis vetustis dense obtectis, foliis acerosis inferioribus brevibus planis cæteris elongatis angustatis omnibus margine seabridis tandem deflexis, scapis folio æquilonigis vel sublongioribus 1-4 squamatis, spicâ capituliformi compressâ 7-8 spiculis trifloris constante, bracteâ exteriore ovato-orbiculata præter basin subcoriaceam nervumque breviter excurrentem hyalinâ, bracteis interioribus cuneatis e medio dorso breviter aristatis, calycis tubo pilosulo limbo pallide roseo eroso-truncato mutico vel nervis breviter excurrentibus ♀. *A. bracteatum* Boiss. Diag. Ser. I, 7 p. 70.

Hab. in aridis Armeniæ Turcicæ prope Erzerum (Cały!), prope Ispir (Bourgl!), prov. Aderbidjan Persiæ ad Khoi (Szov!), ad Seidkhodji (Auch. 5242).

β. splendidum. — Calycis tubus glaber. — *A. splendidum* Bge loc. cit. p. 17.

Hab. in Persiæ borealis montosis infer Aghkent et Mianeh (Bge!).

Rami 3-4-pollicares, folia æstivalia 2-2 $\frac{1}{2}$ -pollicaria, scapi in typo et varietate longitudine (2-6 pollicari) et squamis 1 vel 4 variantes, capitula magnit. nucis, calyx semipollicaris. Var. *β. omnibus* notis præter calycis tubum glabrum similis est.

5. A. cephalotes (Boiss. in p.c. Prodr. I. c. p. 622) cespitosum suffrutescens glaucum breviter erinaceum, foliis brevibus patentissimis acerosis margine seabridis inferioribus latioribus planis cæteris plano-triquetris, scapis foliis 4-5-plo longioribus, spicâ ovato-oblongâ compressâ 5-7 spiculis distichis 2-5-floris constante, bracteâ exteriâ ovatâ naviculari præter dorsum late coriaceum in mucronatum excurrentis late hyalinâ, interioribus cuneatis refutis præter nervum e sinu in mucronatum omnino hyalinis, calycis tubo inter costas hirtulo, limbi tubo sesquibrevioris albidi eroso-truncati albidi mutici nervis roseis 5.

Hab. in regno Cabulico prope Kalou (Griff.).

Folia æstivalia 8-9 lineas longa lineam basi lata, calyx 5 lineas longus, capitula fere pollicem longa.

6. A. Cabulicum (Boiss. I. c. p. 623) cespitosum suffrutescens glaucum calcareo-tuberculatum ramis brevibus densissime foliosum, foliis annotinis acerosis longissimis plano-triquetris, scapis folia non superantibus, spicâ ovata compressâ dense et distiche 5-7-spiculatâ spiculis trifloris subpedicellatis, bractea externâ vix coriacea rufescente oblongâ concavâ calyces superante in mucronem tenuem acerosum longe attenuatâ interioribus præter nervum hyalinis obtusis vel truncatis muticis vel mucronulatis, calycis tubo inter costas parce puberulo limbo sordide albo truncato-eroso mutico 5.

Hab. in Affghania. (Vidi olim in herb. Hook sub N° 1581).

Affine præcedenti differre videtur foliis sœpe bipinnicaribus, scapis brevibus, bractea externâ longe cuspidatâ, internis apice non bilobis; spica et flores fere ampliores

7. A. distachyum (Boiss. loc. cit. p. 623) erinaceo-cespitosum glaucum calcareo-punctatum, ramis abbreviatis, foliis annotinis rigidis longis acerosis plano-triquetris, scapis folium subæquantibus, spiculis 2-3-floris binis terminalibus in capitulum approximatis inferiore sessili, bractea exteriore oblongâ basi coriacea late et sordide albo-marginatâ dorso excurrente tenuiter acerosâ interioribus dorso spathulatis aristulatis, calycis tubo in er costas hirtulo, limbi fusescenti-albidi eroso-truncati nervis pallidis mucronulatim excurrentibus 5.

Hab. in regno Cabulico (Griff. in herb. Hook sub N° 1582).

Folia 12-15 lineas longa, spiculæ 5 lineas, calyx 4 lineas longus.

SECT. V. GLUMARIA Boiss.

8. A. latifolium (Boiss. Diagn. Ser II, 4 p. 61) dense cespitosum glaucescens ramis brevissimis, foliis dense rosulatis patentibus planis brevibus late linearibus pungentibus albo-marginatis et margine seabridis, scapo foliis pluries longiore monostachyo, spicâ ovato-oblongâ spiculis 6-12 bi vel trifloris magnis inferioribus subdistans cæteris approximatis constante, bracteis pallidis late albo-marginatis exteriore subbreviore ovato-triangulari apice attenuatâ acutâ, cæteris latioribus ample obovatis latissime scariosis brevissime mucronatis, calycis tubo glaberrimo, limbo ampio rubro tubo breviore obtuse et breviter eroso-sublobato mutico 5.

Hab. in Kurdistaniæ Persicæ prope Mendeli (Noë 185!), in monte Avroman 600' (Haussk!).

Species per pulchra ad sectionem præcedentem spicâ ad capitulo formam accedente transiens, folia 12-15 lineas longa interdum bipinnicaria sesquilineam fere lata illis affiniunt minus dura, scapus 5-10-pollicaris, spica 2-3-pollicaris, calyces in genere maximi 10 lineas longi.

9. **A. Tataricum** (Boiss. loc. cit. p. 623) confertim cespitosum glaucescens ramis foliis reflexis horridis, foliis brevibus acerosis plano-subtriquetris, scapo brevissime hirtulo folia plures superante, spiculis 5-7 sessilibus alternis remotiusculis bifloris laxe et stricie spicatis, bracteâ exteriore breviore coriaceâ fuscescente margine anguste membranaceâ oblongâ breviter acerosâ, interioribus nervo dorsali apice mucronanti excepto omnino hyalinis, calycis tubo inter costas apice puberulo, limbo duplo breviore albo nervis atropurpureis ad marginem usque productis mutico 5.

Hab. in rupestribus regionis montanae et subalpinæ montis Karatau prope Samarkand Turkestaniae (Lehml.).

Folia hornotina 6-8 lineas tantum longa acerosissima, scapi 3-4-pollicares inferne 3-4-squamati, spica terminalis bipollicaris spiculis inferioribus magis remotis, calyx $4\frac{1}{2}$ lineas longus.

10. **A. Griffithianum** (Boiss. loc. cit. p. 623) erinaceum virens, ramis foliis vetustis horridis, foliis tenuibus subtriquetris acerosissimis, scapis virgatis tenuibus longis basi ramulo uno altero raro auctis superne spiculas plures 2-3-floras parvas sessiles alternas et distantes laxe spicatas ferentibus, bracteâ inferiore aliis dimidio breviore coriaceâ fusca ovato-triangulari mucronatâ superne albo-marginatâ, cæteris oblongis præter nervum medium in mucrone longiusculum excurrentem fusco-membranaceis carinatis calyceem æquantibus, calycis tubo glaberrimo limbo duplo breviore pallide rubello obsolete lobato-eroso expanso 5.

Hab. in Affghaniâ (Griff. N° 156 in herb. Hookt.).

Folia 9-10 lineas tantum longa, scapi cum spicâ laxâ fere pedales, calyx vix $3\frac{1}{2}$ lineas longus. Affine *A. polystachyo*, spiculis unifloris fasciculatis, etc. distincto.

11. **A. scirpinum** (Bge. loc. cit. p. 22) dense erinaceo-cespitosum pallide virens, ramis brevibus foliis reversis densissime obsitis foliis brevibus linear-lanceolatis acerosis margine tenuiter scabris scapis gracilibus virgatis longis remotissime 4-5-spiculatis interdum ramulo uno altero raro auctis, spiculis sessilibus 1-2-floris, bracteis muticis vel brevissime mucronatis exteruâ dimidio breviore ovalâ acutâ coriaceâ membranaceo-marginalâ, interioribus obovato-oblongis subcarinatis dorso herbaceis, calycis tubo inter costas hispidulo, limbi albi tubo subdupo brevioris mutici eroso-sublobati nervis purpureis 5.

Hab. in montosis lapidosis supra pagum Tehegar ad orientem urbis Mesched in prov. Khorassan Persie (Bge.).

Folia 9 lineas longa Lasi vix lineæ $\frac{3}{4}$ lata, scapi 5-6-pollicares, spiculae subquaternæ inferiores intervallo $1\frac{1}{2}$ -pollicari inter se distantes. Calyx fere 5 lineas longus limbo albo, quibus notis ab affini *A. Griffithiano* differt.

12. **A. bromifolium** (Boiss. et Haussk. in Sched.-Bge loc. cit. p. 23) glaucescens, ramis vetustis suffrutescentibus ascendentibus denudatis hornotinis brevibus dense foliosis, foliis longis planis mar-

gine albo-cartilagineis sub lente seabrido-velutinis vel glabris apice tenuiter acerosis, scapis crassis firmis teretibus folia pluries superantibus, spiculis numerosis distantibus 1-2-floris longe et laxe spicatis strictis rachide rectâ, bracteis herbaceo-coriaceis late hyalino-marginatis exteriore æquilongâ longe attenuatâ pungente, interioribus subplicatis brevissime mucronatis calycis tubum pubescentem superantibus, limbi pallide rosei mutici tubo $2\frac{1}{2}$ -plo brevioris eroso-quinquelobati nervis purpureis 5.

Hab. in schistosis alpinis montis Geminan Persiæ inter Sihna et Kerman-schah 7-9000', in monte Ataschke inter Sihna et Awiheng 9-10000' et in monte Dalechani Kurdistaniæ (Haussk!).

Rami vetusti elongati decumbentes non cespites vel saltem cespites laxissimos formant. Folia 2-3-pollicaria saepè inferne sesquilineam lata, scapi cum spicâ semipedali pedales et longiores. Calyx $5\frac{1}{2}$ -6 lineas longus. Foliis longis planis subtili xnosis insigne.

13. A. restiaceum (Bge loc cit. p. 22) cespitosum valde glaucum, ramis brevibus densissime foliosis, foliis rigidissimis patentibus tandem reversis planis acerosis basi ad marginem sebris, scapis firmis folia subsuperantibus, spicâ simplici elongatâ laxâ fragillimâ rachide valde anfractuosa, spiculis bifloris sessilibus distantibus patulis, bracteis coriaceis inferiore triangulari anguste albo-marginata longe aristata triplo breviore interiore areolissime convolutâ albo-marginata cuspidata calyces includente, interfloralibus abbreviatis, calycis glabri purpurascenti-fusei limbo brevissimo arce plicato lobis lanceolatis 5.

Hab. in Persiâ bor. orientali ad Abdulabad inter Meschhed et Turbet Scheik i Dehami (Bge).

Ex specimini bus paucis nimis vetustis tantum notum, folia calcareo-squamulosa 1-1½-pollicaria basi 4 ½ lin. lata, scapi absque spicâ 2 ½-3-pollicares, spicæ articuli semipollicares, spiculae 8 lineas, calyces 6 lineas longi. Floribus magnis et habitu *A. bromifolio* accedens, distinctum rachide anfractuosa, bractearum structurâ, etc.

SECT. VI. PULVINARIA.

14. A. diapensioides (Boiss. in D.C. Prodr. XII, p. 624) glaucum, ramis foliis minimis squarroso-patentibus persistentibus dense imbricatis columnaribus brevissimis in pulvinos densos congestis, foliis brevissimis crassiuseulis obtuse triquetris obtusis vel acutiseulis basi ad marginem sebridulis, spiculis 1-3 in centro rosulae sessilibus unifloris rarissime bifloris, bracteis 4-5 ovatis rufescensibus membranaceis mucronatis exteriore ovato-triangulari rigidula, superioribus duplo longioribus truncatis brevissime mucronatis, calycis tubo æquilongo eroso mutico 2.

Hab. in alpe Koh i Baba Affghanie 44-15000' (Griff!).

Facies *Saxifrage caesia*, folia ½-1 ½ lineas tantum longa, calyx vix 3

lineas longus. Am. a Bunge hanc speciem facie distinctissimam in sect. *Glu-*
mariæ medianibus *A. Ruprechti* et *lycopodiode* in ditione nostrâ non indi-
genis collocat, sed habitus omnino alienus et foliis muticis ab omnibus con-
generibus distincta.

SECT. VII. STATICOPSIS Bge.

§ 1. Rhodocalycina.

15. **A. avenaceum** (Bge loc. cit. p. 25) dense erinaceo-cespi-
tosum glaucescens, ramis brevibus foliis patentibus horridis, foliis
tenuibus rigidis acerosis plano-triquetris margine seabridis, scapis
gracilibus elongatis glabris teretibus saepe ramosis, spicis omnibus
brevibus dense et fasciculatim 3-5 spiculatis inferioribus pedunculatis,
superioribus unispiculatis, spiculis unifloris, bracteâ exteriore ovata
erioaceo-herbacea acuta fuscescenti-marginata, interioribus duplo
longioribus convolutis obtusis præter carinam virentem fuscō-mem-
branaceis, calycis tubo hirsuto in limbum purpurascentem subtrun-
catum obsolete erosō-quinquelobum submuticum glabrum sensim
ampliato ♂.

Hab. in lapidosis montanis ad pagos Tchegar et Gulistan ad occid. urbis
Meschled prov. Khorassan (Bge!).

Folia 7-10 lineas longa, scapi cum spicâ pedales, calyx 4 $\frac{1}{2}$ lineas longus.
Species facie *A. scirpino* simillima sed spiculae semper unifloræ tribracteatæ
et limbus purpurascens nec albus.

16. **A. Calverti** (Boiss. Diagn. Ser. II, 4 p. 65) cespitosum viride,
ramis tenuibus basi vestigiis foliorum circinnatim recurvis vestitis,
foliis longis angustissime linearibus planiusculis margine seabridulis
in mucronem vix pungentem abeuntibus non rigidis, scapis folia vix
superantibus puberulis 1-2-stachyis, spicis contiguis ovatis compressis
dense et distiche 5-7-spiculatis, bracteis subæquilongis herbaceis
fuscescenti-marginatis glabris exteriore attenuato-mucronata cæteris
præter nervum membranaceis abrupte et breviter mucronulatis,
calycis tubo obsolete puberulo, limbo subæquilongo obsolete lobato
roseo vel pallide rubello mutico nigro-nervoso ♀.

Hab. in Armeniæ monte Techdagh prope Erzerum (Calv. exs. 507!), ad
Palanteiken (Ky. Suppl. 584).

Ex unico specimen notum fere herbaceum, folia sesquipollucaria tenuissima
minime pungentia, spicae 9-10 lineas longæ, calyx 5 $\frac{1}{2}$ lineas longus. Omnia
facies et inflorescentia *A. glumacei* J. et Sp. sed folia minus rigida vix pun-
gentia, bracteæ glabræ, calycis limbus purpureus et interdum in pallide
rubellum vergens.

17. **A. venustum** (Boiss. Diagn. Ser. I, 7 p. 80) laxe cespito-
sum glaucescens calcareo-squamulatum, foliis latiusculis plano-cari-
natis præter infima subinermia pungentibus, scapis brevibus folia
superantibus superne cum rachide seabridis, spicâ simplici laxiusculâ

unilaterali vel distichâ erectâ vel subcurvatâ rachide vix flexuosa, bracteâ exteriore dimidio breviore late triangulâ acuminatâ carinatâ anguste membranaceo-marginatâ, interioribus oblongis mucronatis præter dorsum interdum obsolete puberulum scariosis, calycis tubo sparsim piloso limbo tubo subbreviori lato truncato plicato mutico fuscescenti-rubello ȝ. Ic. Tchih. Voy. tab. 72. — *Statice venusta* Fenzl MSS. in Ky. exs. — *S. dianthifolia* J. et Sp. Ill. I, p. 162 excl. syn. — *Limonium Orientale humilius Caryophylli folio angustiore flore spicato suave rubente* Tourn. Cor. p. 25.

Hab. in alpinis Tauri Cilicici prope Gülek Boghas 7000' (Ky. 441 Bal!), montis Argæi Cappadociae (Bal!), Ssöffdaghi Cataonie 3500' (Haussk!), Armenia (Tchich!), Anatoliæ bor. monte Yldizdaghi (Wied!).

β. Olivieri. — Folia longiora angustiora, spiculæ magis dissitæ. — *A. Olivieri* J. et Sp. loc. cit. tab. 93. — *A. laxiflorum* Boiss. in Bourg. exs. 295!

Hab. in Lycia prope Elmalu (Bourg. 295!), Tauri regione subalpinâ ad Kar Boghas (Bal!), monte Berytdagh Cataonie (Haussk!), Persia occid. inter Kermanchah et Hamadan (Oliv.).

γ. Assyriacum. — Spica laxa varietatis præcedentis sed scapus superne cum rachide et bracteis velutino-pubescentes. — *A. Assyriacum* Boiss. Diagn. Ser. I, 7 p. 81.

Hab. in Assyria inter Diarbekir et Mardin (Ky. 313!).

Ex speciminum comparatorum copiâ species in synonymia enumerate in *A. venustum* manifeste transierunt. Bractearum indumentum valde variabile, in speciminibus e Tauro Cilicico specimina alia bracteis velutinis aliis bracteis glaberrimis in eodem loco lectis simillima observavi. Species corollis amplis intense purpureis elegans, spica in typo 2 pollices, in formis laxifloris tres pollices longa, calyx 7-8 lineas longus.

18. ***A. laxiflorum*** (Boiss. in Ky. pl. Syr. bor. 1862) glaucum fruticulosum, ramis elongatis squamosis nigricantibus denudatis rosulâ paucifoliâ terminatis, foliis carnosulis brevibus triquetris enerviis in spinulam tenerrimam fuscam subito contractis, scapo foliis longiore in spicam longam laxam rectam unilateralem abeundi cum rachide lœvi glabro vel velutino, bracteis herbaceis glabris vel velutinis latiuscule albo-marginatis nervo longe excurrente rubro acuminato-cuspidatis exteriore tertia parte breviore, calycis tubo pilosulo, limbo pallissidime rubente tubo duplo breviore mutico obsolete quinque-lobo ȝ.

Hab. in glareosis fluvii Nahr Syad infra pagum Ursusa montis Amani Syriae bor. 200' (Ky. 428!).

Ramis parte denudata 2-3-pollices longis insigne, fruticulosum. Ab omnibus *A. venusti* formis rosulis oligophyllis, foliis carnosis fragilibus triquetris 6-8 lineas rarius pollicem longis, spicâ longâ rectâ semipedalî, spiculis ob bracteas internas arcte convolutas angustioribus et præsertim calycis (5 $\frac{1}{2}$ lineas longi) limbo tubi proportione multo breviore distinctum.

19. ***A. petraeum*** (Boiss. et Haussk Scied. pl. 1867. — Bge loc. cit. p. 28) dense cespitosum glaucum, ramis brevissimis confertis,

foliis brevissimis lanceolato-linearibus planis apice triquetris pungentibus acerosis, scapo tenui foliis parum longiore in spicam tenuem laxissimam brevem abeunte et cum ejus rachide subflexuosa glabro levigata spiculis 3-4 valde dissitis, bracteis glaberrimis externâ oblongo-lanceolatâ acutâ pallidâ albo-marginatâ quartâ parte breviore, cæteris præter costam in mucronem excurrentem hyalinis, calycis tubo adpresso puberulo limbo tubo subbreviore obsolete quinquelobo mutico pallide roseo ♂.

Hab. in monte Pir Omar Gudrun Kurdistaniae Persicae 3-6'000' (Haussk!). Planta Armena e prov. Musch (Ky. exs. № 379 huc a cl. a Bunge citata in herb. meo deest et eam non vidi).

Folia 6-8 lineas tantum longa, scapi cum spicâ 2-3-pollicares, rachidis articuli bracteis saepius longiores, calyx 6 lineas longus. A formis laxilloris *A. venusti* præter folia brevissima spicis paucifloris laxissimis, rachide tenui levii, bracteis minus inæqualibus distinctum.

20. **A. Atropatanum** (Bge loc. cit. p. 29) dense cespitosum valde glaucum ramis brevissimis confertis, foliis brevibus lanceolato-linearibus plano-triquetris rigide acerosis, scapo foliis parum longiore in spicam brevem laxiusculam disticham abeunti cum rachide flexuosa glaberrimo, bracteis subæqualibus glaberrimis late hyalino-marginalatis exteriore oblongo-lanceolatâ subcarinata cuspidato-attenuata intimâ obtusiusculâ mucronulatâ præter nervum hyalinâ, corolle tubo adpresso hirto, limbo tubo subbreviore roseo plicato truncato-subblobato mutico ♂.

Hab. in montosis aridis prov. Aderbidjan Persiae inter Turkmenchai et Tik-medesch (Bge!).

Folia 6-9 lineas tantum longa, scapi cum spicâ 7-8-spiculatâ 2-3-pollicares, rachidis articuli bracteis subbreviores, calyx 6½ lineas longus. Species a formis laxilloris *A. venusti* præter folia brevia valde glauca rigidiora inflorescentia glaberrimâ levii et presertim bracteis subæqualibus latius membranaceis distincta sed forsitan varietas *A. petrei* a quo differt tantum spicâ paniculatâ laxissimâ, bracteis non subæqualibus, exteriore non longe cuspidatâ. Observandum an formæ intermediae occurront.

21. **A. Senganense** (Bge loc. cit. p. 29) dense cespitosum valde glaucum, ramis brevissimis confertis dense foliosis, foliis anguste linearibus longe aceroso-pungentibus planiusculis apice subtrigonis, scapis gracilibus folia paulo superantibus in spicam simpliæ 3-6-spiculatam abeuntibus cum rachide reclinuscule scabridis, bracteis glabris exteriore quartâ parte breviore late et subauriculatim hyalino-marginalatâ acutâ, cæteris præter nervum breviter mucronantem hyalinis, calycis tubo hirtulo, limbo tubo breviore roseo truncato mutico ♂.

Hab. in montosis Persiae borealis inter Chorrumderreh, Sultanieh et Sengan (Bge!).

Folia tenuia 8-9 lineas tantum longa, scapi cum spicâ 2-4-pollicares, calyx 5 lineas longus. Species valde affinis *A. petreæ* et *Atropatanæ* a quibus foliis tenuioribus, bracteâ inferiore late marginatâ et basi subauriculata etc. differt.

§ 2. Caryophyllacea.

22. **A. tenuiflorum** (Boiss. Diagn. Ser. I, 7 p. 78) cespitosum glaucum ramis brevibus, foliis angustissime linearibus acerosis subulato-triquetris vetustis reversim imbricatis, scapis gracillimis folio multo longioribus paniculatum 2-5-stachyis, spicis laxissime remotiuscula 3-12-spiculatis rachide rectâ glaberrimâ, spiculis tenuibus strictis, bracteâ exteriore membranaceo-marginatâ cuspidato-attenuatâ interioribus præter nervum medium in mucronem excurrentem hyalinis subbreviore, calycis tubo parce hirtulo, limbo albido obsolete quinquelobo nervis fuscis glabris ante marginem evanidis percurso ȝ. *Statice acerosa* Hohen. Elisab. et Talysh Enum. non W. nec M.B. — *S. Echinus* Linn. ex parte. — *Limonium spicatum* foliis aculeatis Baxb. Cent. II, p. 18, tab. 10.

Hab. in Transcaucasiâ inter Gandcha et Schemacha (Buxb.), in apricis lapidosis ad Helenendorf et in tractu Suwant (Hohenl.).

Vix suffrutescens, folia 8-10 lineas longa tenuissima, scapi 3-5-pollicares, panicula sæpe semipedalis spicâ terminali 5-12-spiculatâ et spicis 1-3 lateraliibus longius pedunculatis paucius spiculatis constans, spicâ inter se sæpe plus quam earum longitudine remotâ, calyx 4½-5 lineas longus.

23. **A. Araxanum** (Bge loc. cit. p. 33) glaucescens calcareo-squamulatum ramis elongatis, foliis longissimis lineari-triquetris longe pungenti-acerosis vetustis reversis, scapis glaberrimis folia parum superantibus distachyis rarius monostachyis, spicis fragillimis alterâ longiore laxe 7-10-spiculatâ alterâ breviore paucispiculatâ, rachidis flexuosa glaberrimâ articulis bracteis æquilongis, bracteis glabris exteriore lanceolatâ acuminato-attenuatâ albo-marginatâ interioribus brevioribus obtusis præter nervum in mucronem excurrentem membranaceis, calycis tubo patule piloso, limbo subbreviore albo subtruncato nervis purpureis glabris ad marginem fere productis ȝ.

Hab. in apricis siccis ad Araxem in districtu Choï prov. Aderbidjan Persia (Szov!).

Folia superiora 2-pollices et amplius longa superne valde attenuata, scapi squamati squamis acuminatis. Calyx sex lineas longus. Differt ab *A. tenuifloro* foliis multo longioribus, floribus majoribus, bracteâ inferiore elongatâ, etc.

24. **A. Hohenackeri** (Jaub. et Sp. Ill. I, p. 162, tab. 92 sub *Statice*) cespitoso-erinaceum glaucescens calcareo-squamulosum et impresso-punctatum, ramis subelongatis foliis vetustis patentibus vel refractis creberrime obsitis, foliis brevibus anguste subulato-triquetris acerosis, scapis foliis subduplo longioribus scabro-puberulis 1-3-stachyis, spicis brevibus spiculis 5-8 densiuscula et distiche imbricatis, rachide hispidulâ, bracteis glabris herbaceis rubellis acutis albo-marginatis externâ tertiatâ parte breviore, calycis tubo undique adpresso hirsuto, limbi albi tubo brevioris eroso-truncati mutici nervis nigris extus puberulis ȝ. *A. Hohenackeri* Boiss. Diagn. 7, p. 75. — *St. Echi-*

nus Taur. Cauc. I, p. 251. — *Statice tenuifolia* et *S. aciphylla* J. et Sp. loc. cit. p. 162. — *S. horrida* Girard Ann. Sc. Nat. Ser. III, 2, p. 331 ex parte.

Hab. in Caucaso orientali (Seidl.), tractu Suwant ditionis Talysch (C.A. Mey! Hoh!), prov. Ghilan Persiae bor. (herb. Fisch! Auch. 5f41!), prov. Aderbidjan prope Nehmetabad (Bge!).

Folia tenuia 6-9 lineas longa, spicarum fasciculus vix pollicem longus, calyx 5 lineas longus.

25. **A. Huetii** (Boiss. Diagn. Ser. II, 4 p. 65) glaucescens totum scabrido-velutinum a collo ramosissimum ramis brevissimis denudatis subsquamosis, rosulis oligophyllis, foliis laxis carnosulis subtrigonis brevissimis recurvo-patnlis in mucronem brevem tenuissimum abrupte contractis, scapo foliis duplo longiore distachyo, spicis sessilibus ovatis dense distichis 5-9-spiculatis, bracteis amplis velutinis in mucronem attenuatis exteriore breviore latius herbaceâ, interioribus abruptius mucronatis, calycis tubo hirtulo, limbi subbrevioris albi sub 10-lobulati nervis puberulis rubris ad marginem usque productis ♀.

Hab. in Armeniâ Turcicâ supra Maimansour (Huet!).

Speciei singularis unicum specimen lectum, rami vix suffruticosi tennes, folia 6-7 lineas tantum longa herbaceo-carnosa fragilia ut in *A. Calverti* abrupte et tenuiter mucronata, spica dense imbricata vix pollicem longa, calyx 6½ lineas longus. Spiculam unicam bifloram vidi! Affine *A. glumaceum* et *A. Calverti* a quibus foliis, etc.; cæterum longe differt.

26. **A. glumaceum** (Jaub. et Sp. Ill. I, p. 162, tab. 91 sub *Statice*) vix suffrutescens viride cespitosum, ramis basi nudis superne foliis reflexis vestitis, foliis anguste subulatis tenuissime acerosis breviusculis, scapis foliis 2-3-plo longioribus 1-3-stachyis rachideque velutinis, spicis brevibus confertim 7-9-spiculatis, bracteis subvelutinis exteriore late ovata herbacea margine membranacea cuspidatâ, interioribus sublongioribus praeter dorsum viridem hyalinis obtusis mucronatis, calycis tubo ad costas hispidulo limbo albo nervis atroviolaceis latis ad marginem productis percurso ♀. *A. glumaceum* Boiss. Diagn. 7, p. 75 — *S. Hohenackeri* Led. Fl. Ross. ex parte quoad plantam Armenam. — *Limonium Orientale caryophylli* folio acutissimo, floribus in capitulum congestis acaulon et calyce albo Tourn. Cor. 25.

Hab. in Armeniâ Rossicâ, mons Ararat (Tourn.) et in valle Arguri (Parrot, Abich), ad rivulum Hamanlu (Frick!).

27. **A. Sahendicum** (Boiss. et Buhse Aufz. p. 183) viride glaberrimum dense cespitoso-erinaceum ramis brevibus foliis vetustis dense et squarrose obsitis, foliis anguste linearis-subulatis subtriquetris acerosis, scapis tenuibus folia subsuperantibus glaberrimis monostachyis, spicâ brevi laxiuseule et digitche 3-7-spiculatâ, rachide anfractuosâ glaberrimâ, bracteis glabris rubro nigricantibus ovatis

breviter mucronatis carinatis membranaceo-marginatis inferiore breviore, ealycis tubo inter nervos patule piloso, limbo albo tubo breviore truncato nervis atropurpureis usque ad apicem productis ♂.

Hab. in alpinis montis Ssahend prov. Aderbidjan Persiae 8-900' (Buhse! Seidl.).

Affl. e *A. glumaceo* sed scapi, rachis, bracteæ glaberrimæ tusco-rubræ, calyx 5 lineas longus. Ex specimine valde macro monostachyum sed probabilissime etiam ex maximâ cum *A. glumaceo* et *Bodeano* affinitate pleiostachyum.

28 ***A. Bodeanum*** (Bge loc. cit. p. 42) glaberrimum glaucescens erinaceo-cespitosum calcareo-squamulosum, ramis vetustis foliis velutinis squarrosis dense vestitis, foliis linear-lanceolatis trigonis rigide acerosis, scapis folio 2-3-plo longioribus 1-2-stachyis, spicis approximatis brevissimis distiche 5-7-spiculatis basi laxiusculis, rachide rectâ glaberrimâ, bracteis glabris exteriore triplo breviore breviter ovatâ anguste membranaceâ muticâ vel acutiusculâ, internis oblongis late fuscescens-membranaceis obtusis muticis, calycis tubo superne patentim piloso, limbo subæquilongo truncato albo nervis nigris glaberrimis usque ad apicem productis ♂.

Hab. ad Fullad Mahaleh Persiae borealis ad orientem urbis Teheran versus Schahrud eundo (Bode!).

Folia 7-9 lineas longa tenuia, scapi 2-2½-pollicares, calyx 5½ lineas longus. Facie, rachide scapoque glaberrimis affine *A. Sahendico* a quo foliis glaucescentibus, bracteis valde inæqualibus muticis abunde differt.

29. ***A. Lycaonicum*** (Boiss. et Heldr. in D.C. Prodr. XII, p. 627) dense cespitosum vix suffruticosum viride, foliis brevibus lanceolato-linearibus superne subtriquetris acerasis, scapis folia parum superantibus monostachyis vel rarius distachyis cum rachide velutinis, spicis sessilibus obovato-capitatis 5-8-spiculatis distiche et dense imbricatis interdum subscorpioideis. bracteis latis exteriore herbacea membranaceo-marginata mucronata interioribus longioribus præter nervum latum sub apice in mucronem excurrentem membranaceis, calycis tubo superne pilosulo, limbi albi submutici nervis atropurpureis ad apicem usque productis et ibi saepe dilatatis ♂.

Hab. in collibus Lycaoniae inter Beychehr et Koniah (Heldr!), in Phrygia inter Kiutaya et Arzani (Mitch!).

ξ. *Cappadocicum*. — Totum pubescenti-velutinum, folia sublatiora.

Hab. in regione alpinâ montis Aslaudach Cappadociae (Bal!).

γ. *pictum*. — Velutinum, iolia ut in formâ typicâ angustiora magis triquetra, bracteæ dorso velutinæ longiores mucronatae parte membranacea eleganter purpurascentes. — *A. Cataonicum* Bge loc. cit. p. 43 ex loco vix ex descriptione.

Hab. in monte Berytdagh 8-9000' (Haussk! sub *A. Tournefortii* var.).

Species foliis brevibus viridibus *A. Echinum* referens sed scapi folia superantes saepe distachyi et spicæ omnino dissimiles capitatae distiche compressæ, bracteæ latiores; ad *A. glumaceum* et *Calverti* cæterum distinctissima proprius

accedens. Varietas 7, sub nomine *A. Cataonici* ab. amic. a Bunge videtur descripta, sed specimina forsitan in schedula commixta fuerunt nam tota planta ex cl. Bge descriptione glaberrima dicta omnibus partibus velutina est. Folia 7-12 lineas longa, scapi foliis sesquiloculares sepius monostachyis, spicæ 7-9 lineas tantum longæ, calyx 6 lineas longus.

30. A. Cæsareum (Boiss. et Bal. Diagn. Ser. II, 4 p. 66) viride erinaceo-cespitosum ramis brevibus foliis vetustis patentibus horridis, foliis brevibus anguste linearis-trigonis subulato-attenuatis, scapis gracilibus foliis 4-5-plo longioribus superne cum rachide puberulis 1-2-stachyis, spicis oblongis distiche et 5-9-spiculatis basi-laxiusculis, bracteis glabris exteriore breviore herbacea fusco-marginata attenuato-mucronata, internis præter nervum in mucronem excurrentem omnino fusco-membranaceis obtusis, calycis tubo inter nervos hirtulo, limbo albo tubo subbreviore eroso-sublobato nervis atropurpureis ad marginem productis basi hirsutis §.

Hab. in regione subalpinâ montis Alidagh Cappadociae 4200' (Bal. exs. 9461).

Folia 9-12 lineas tantum longa, scapi tres pollicares, spicæ 1-4½-pollicem longæ, calyx 6 lineas longus. Proximum *A. glumaceo* differt spicis longioribus sublatiорibus, bracteis glabris fusco-membranaceis; ab *A. Lycaonico* scapis elongatis et spicâ longiore basi laxiusculâ longius distat.

31. A. Auganum (Bge. loc. cit. p. 39) glabrum viride, ramis tenuibus elongatis foliis vetustis flexuosis laxe obsitis, foliis brevissimis tenuiter subulato-triquetris acerosis inferne membranâ hyalinâ glaberrimâ marginatis, scapis gracilibus folia subsuperantibus monostachyis, spicâ capituliformi conferentia paucispiculata, rachide brevisimâ glabrâ, bracteâ exteriore ovata viridi margine late albo-membranacea acuminata, interioribus fere duplo longioribus oblongis præter nervum dorsalem mucronantem omnino hyalinis obtusis, calycis tubo hispidulo, limbi albidi tubo subbrevioris nervis atropurpureis glabris ad apicem usque productis ².

Hab. in alpinis summis regni Cabulici, jugum Unu 11300' et faux Yonutt 19800' (Griff!).

Specimen unicum vidi floribus spicæ jam exarticulatis, folia 6-7 lineas longa basi longe in membranam margine integerrimam non ciliatam dilatatâ margine tenuissime quidem sed evidenter seabridâ, bractea exterior 2 lineas interna 4 lineas longa, calyx 6 lineas longus. Annon sæpius pleiostachyum et inter species gregis præcedentis spiculis abbreviatis capituliformibus prædictis potius inserendum?

32. A. acerosum (Willd. Ges. Nat. Freund. III, p. 420 sub *Statice*) suffruticosum dumoso-erinaceum glaucescens impresso-punctatum, ramis elongatis, foliis crassis longis acerosissimis plano-triquetris margine seabridis, scapis folio longioribus in spicam laxam elongatum multispiculata abeuntibus, rachide flexuosa seabridâ, bracteis late scarioso-marginalis externâ triangulari longe spinuloso-acuminata interioribus brevius acuminatis carinatis calycis tubum superantibus tertia vel quartâ parte breviore, calycis tubo patule hirsuto, limbo

breviore albo truncato mutico nervis extus puberulis 5. Ic. Jaub. et Sp. III. tab. 80. — *A. acerosum* Boiss. Diagn. Ser. I, 7 p. 80. — *A. Phrygium* Boiss. Diagn. 7, p. 79. — *Limonium Orientale frutescens Caryophyli folio in aculeum rigidissimum ubeunte* Tourn. Cor. p. 25. — *A. caryophyllaceum* Boiss. in D.C. Prodr. ex parte.

Hab. in regione alpinâ Anatoliae occidentalis, Sipylus supra Magnesiam (Boiss! Bal. 284!) Cadmus (Jaub!), Tuolus supra Philadelphiam (Boiss!) Lycia ad Elmaliu (Bourg. 249! sub *A. Pinardi*).

B. brachystachyum. — *A. Pinardi* Boiss. Diagn. 7, p. 79. — *A. Listoniae* Boiss. in D.C. Prodr. XII, p. 631! — *A. Tchihatcheffii* F. et M. Ann. Sc. Nat. 1854, p. 30. — Tchihi. Voy. tab. 36.

Hab. in Anatoliâ bor. (Dom^{na} Liston!), ad Safranbol (Wied!), in Galatiâ (Sestini!), Cariâ (Pinard!), Phrygiâ inter Kntaya et Argani (Mitch!) regione inferiore Tauri Cilicieci ad castellum Gulek et in valle Gusgula (Ky. 99! sub *A. Pinardi*).

Domi lati 6-10-pollicies longi, folia 2-3-pollicaria basi sesquilineam lata, in specimine e Lyciâ breviora; spicæ in typo 4-8-pollicies in varietate 2-3-pollicies longæ, calyx 6½-7 lineas longus. Discrimen inter *A. acerosum* et *A. Pinardi* e corolle limbo calyce inclusa vel majore exerto depromptum ex culturâ diologamico variat.

33. *A. caryophyllaceum* (Boiss. in Ky. Alepp. Kurd. exs. 368 sub *Statice*) suffruticosum dumulosum erinaceum pallide virens, ramis subelongatis, foliis elongatis linearis-trigonis undique scabridulis, scapis folio sublongioribus in spicam rectam laxiusculam abeuntibus, rachide hispidulâ, bracteis dorso herbaceo plus minusve velutinis albo-marginatis breviter micronato-attenuatis exteriore subbreviore, tubo hispidulo limboque albo obsolete quinquelobo æquilongis nervis purpureis extus hirtis ad marginem usque productis 5. *A. caryophyllaceum* Boiss. Diagn. Ser. I, 7 p. 78.

Hab. in rupestribus montis Gara Kurdistaniae (Ky. exs. 368!), in districtu Warto prov. Musch Armenie australis ad Gestemert 5000' (Ky. 345!), in Kurdistanâ (No^o 207!).

Folia 4-4½-pollicaria, spicæ 3-4-pollicares internodiis bracteam inferiorem subaequantibus, calyx 6 lineas longus. Proximum *A. aceroso* var. *brachystachys* quoque in D.C. Prodr. coniuncti, differt tantum foliis viridibus undique scabridulis et bracteis parte herbacea plus minus velutinis. Hoe ut speciem propriam describendo am. a Bge sequor.

34. *A. Baltanense* (Boiss. et Hanssk.) dumoso erinaceum glaucescens, ramis longiusculis foliis vetustis retrorsis densissime obsitis, foliis tenuibus longissimis linearibus plano-subtriquetris sensim attenuato-pungentibus, scapis laevibus foliis longioribus crebre squamosis in spicam longam laxiusculam abeuntibus, rachidis flexuosæ intus scabriuscule articulis bracteâ externâ subbrevioribus, bracteis glabris late marginato-membranaceis exteriore lanceolatâ in cuspidem tenuem bracteas interiores breviter micronatas superante, calycis tubo paree hirto, limbo albo breviore mutico nervis nigricantibus fere ad apicem usque productis 5.

Hab. in montibus Baltandagh et Begdaghi Cataoniae 5000' (Haussk! exs. sub *A. Phrygio*).

Folia 1½-2-pollicaria, scapi parte nudâ 2-3-pollicares, spicæ tripollicares, calyx 7 lineas longus. Affine *A. aceroso* et *caryophyllaceo* differt foliis angustioribus minus rigidis et præsertim bracteâ externâ cæteras manifeste superante nec eis breviore.

35. ***A. Armenum*** (Boiss. et Huet Diagn. Ser. II, 4, p. 64) cespitosum erinaceum glaucum, ramis abbreviatis, foliis patentibus rigidis acerosis lanceolatis plano-triquetris magine seberrimis, scapis folia æquantibus in spicam rectam densiuscula et distiche 12-15-spiculatam abeuntibus, rachidis levis vel scabrinuscule recte vel subflexuose articulis bracteâ inferiore brevioribus, bracteis æquilongis glabris inferiore albo-marginatâ breviter aceroso-acuminatâ, interioribus præter dorsum membranaceis obtusis mucronatis, calycis tubo hirto, limbi albi subæquilongi erosio-truncati nervis rubris extus hirtis ante marginem evanidis 5. *A. Kotschyi* var. *Cataonicum* Bge loc. cit. p. 35.

Hab. in collibus aridis Armenie Tureicae prope Erzerum (Huet!), prope Baibut (Bourgt!), prope Gunuschikhané (Tchit!), in Cappadocie monte Ali-dagh (Bal. exs. sub *A. Pinardi*!) et in monte Berytdagh Cataoniae (Haussk!).

♂. *Balansæ* Boiss. loc. cit. — Folia subangustiora magis triquetra sublongiora, spicæ breviores, bractea exterior subbrevior intra marginem hyalinum fuscescenti-pieta — *A. Balansa* Boiss. in Bal. exs. et Bge. loc. cit. p. 36. — *A. Haussknechtii* Bge. loc. cit. p. 37.

Hab. in regione alpinâ montis Aslandach Cappadociae (Bal!), n. atus Akkerdagh Cataoniae (Haussk!).

Species ab affini *A. aceroso* foliis abbreviatis, spicis densioribus, floribus minoribus (calyces 5-6 lineas tantum longi) atque præsertim bracteis subæquilongis et limbi nervis longe ante marginem evanidis distincta. Varietas quam prius pro specie propriâ habui mediantibus specimenibus e monte Ali-dagh ad speciem transit. Planta Cataonica a cl. Bge ad *A. Kotschyi* conjuncta hoc meo sensu ob folia angusta longe attenuata 12-18 lineas longa spicamque elongatam spectat.

36. ***A. Iconicum*** (Boiss. et Heldr.) cespitoso-erinaceum valde glaucum, ramis abbreviatis, foliis brevibus linear-lanceolatis patentibus subcurvatis triquetris, scapis brevissimis in spicam multispiculatam laxiuscula imbricatam subincavam abeuntibus cum rachide subflexuose scabridulis, bracteis late albo-marginatis aceroso-attenuatis carinatis, inferiore cæteris breviore, calycis tubo hirsuto limbi albi tubo subbrevioris concoloris nervis pallidis ad apicem usque productis, petalis albidis 5. *A. Kotschyi* ♂ *Iconicum* Boiss. in p.c. Prodr. p. 628.

Hab. in collibus aridis prope Zilla ad occidentem urbis Ieonium (Heldr!).

Folia 9-11 lineas longa, spica 2-2½-pollicaris 45-48-spiculata, calyx 5 lineas longus. Planta prius ad *A. Kotschyi* adducta sed foliis angustioribus longioribus subtriquetris, spicis triplo longioribus subrecurvis, bracteis inaequalibus, petalis albis diversa.

37. **A. Kotschy**i (Jaub. et Sp. Ill. Or. I, p. 162 sub *Statice*) dense cespitosum erinaceum valde glaucum, ramis abbreviatis, foliis brevibus latiusculis lanceolatis rigide acerosis, scapis monostachyis brevissimis subnullis, spicis brevibus 5-7-floris densiuscule imbricatis distiche compressis rectis, bracteis subæquilongis late albo-membranaceis mucronatis, internis dorso acute carinato hirtulis, calycis tubo hirtulo, limbi albi concoloris tubo subbrevioris nervis pallidis ante marginem evanidis, corollâ roseâ 5. *A. Kotschy* Boiss. Diagn. Ser. I, 7, p. 74. — *A. breviscapum* Boiss. et Haussk. MSS.

Hab. in valle Karli Boghaz Tauri Cilicie (Ky. 4161 et 436!), in monte Berydagha Cataoniae (Haussk!).

Folia 6-8 lineas tantum longa inferne $1\frac{1}{4}$ lata. Spicæ pollicares. Calyx in planta Cilicie 4½-5 lineas in Cataonica ob limbum submajorem 5½ longus. Species spicis subsessilibus paucifloris densiuscule imbricatis, foliis brevissimis latiusculis rigidis glaucis, calycis limbo ob nervos pallidos subconcolore insignis.

38. **A. Libanoticum** (Boiss. in D.C. Prodr. XII, p. 630) subglaucescens dense cespitoso-erinaceum dumulosum ramis subelongatis foliis vetustis dense horridis, foliis rigidis triquetris acerosis margine scabris, scaro foliis breviore in spicam brevem laxiusculam rectam 4-7-spiculatam abeunte, rachide seabrido-velutinâ rectiusculâ, bracteis subæquilongis carinatis late membranaceis exteriore in cuspide attenuatâ interioribus abrupte cuspidatis, calycis tubo hispidulo limbo albo æquilongo obsolete quinquelobo nervis purpureis extus glabris ultra marginem non vel vix mucronantibus 5.

Hab. in Libani regione alpinâ et subalpinâ ad Cedros, in declivitate orientali montis Makmel et in Cœlesyriâ circa Ainhata (Boiss!), ad Djurd Hasrun (Bl!), in regione Amygdalorum montis Hermon Antilibani 5500'-6000' (Ky. exs. 163 sub *A. Kotschy* var!).

B. ulicinum. — Virescens, folia abbreviata, spica brevissima confertim et fasciculatim 3-5-spiculata sœpe viscidula, bractea exterior sœpe subbrevior. — *Statice ulicina* Willd. herb. Rém. et Sch. VI, p. 798. — *Statice hystrix* Jaub. et Sp. Ann. Sc. Nat. XX, p. 252.

Hab. in regione summi Libani inter Dimam et Yamouny (Bl!), in summis cacuminibus montis Makmel et in monte Gebel Baruck (Boiss!), in monte Hermone versus Orny 5000' (Ky. 243! Gaill!) et in cacumine ejusdem montis 9000' (Boiss!).

Folia in formâ typicâ 9-11 lineas, in varietate 6-8 lineas tantum longa eis praecedentiam manifestius et acutius triquetra. Ab eis præterea differt spicis paucifloris subsessilibus. Formæ inter typum et varietatem intermediae occurruunt. Varietas *B. ulicina* facie, spicis paucifloris fasciculiformibus ad *A. echinum* accedit, sed hoc differt foliis plano-triquetris, bracteâ exteriore multo breviore, calycis limbo aristulato, etc.

39. **A. Echinus** (L. Sp. 395 ex parte sub *Statice*) viride hre-viter erinaceo-cespitosum ramis brevissimis foliis vetustis horridis, foliis abbreviatis inferioribus planiusculis cæteris plus minusve sublato-triquetris acerosis sœpius glabris, scapis monostachyis pubes-

centibus subnullis vel brevissimis rarissime folia subæquantibus, spicæ abbreviatâ fasciculatim confertim et distiche 3-7-spiculata, rachide velutinâ, bracteis basi vel ad totam partem herbaceam velutinis acuminato-cuspidatis externâ breviore herbaceâ membranaceo-marginatâ cæteris nervo excepto membranaceis, calycis tubo hirtello, limbo albo tubo subbreviore nervis atropurpureis percurso mutico vel breviter mucronulato §. Ic. Fl. Græc. tab. 300 — *A. androsaceum* Boiss. Diagn. Ser. I, 7, p. 73. — *Statice androsacea* J. et Sp. Ill. I, tab. 89, et *S. pauciflora* J. et Sp. eod. loc. p. 162.

Hab. in regione alpinâ Græciae et Anatoliae, Peloponnesus in montibus Taygeto et Olenos (Heldr!), Malevo (Orph. Fl. exs. 80!), monte Velugo et Parnasso (Sprun! Heldr. Herb. Norm. 337!), Bithynia in regione alpinâ Olympi (Sibth! Auch! Boiss! etc.), Lydiâ in cacuminibus Tinoli (Boiss!), Pauro Cilicio 8000' (Ky! Bal!).

β. *Tournefortii*. — Folia subelongata magis attenuata erectiora lores æquantia. — *Statice Tournefortii* J. et Sp. tab. 90. — *Limonium Armenum Caryophylli folio acutissimo floribus veluti in capitulum congestis* Tourn. Cor. p. 25 et herb!

Hab. in montibus Armeniae (Tourn., Huet!), prope Baikut (Bourg!).

γ. *Creticum* Boiss. in D.C. Prodr. — Folia brevissima velutino-uberula. Spica sessilis 2-3-flora. — *Limonium Creticum juniperifolium* Tourn. Cor. 25.

Hab. in cacuminibus montium Sphacioticorum Cretæ (Alp., Tourn., Sieb! Sibth! Rau!), montis Ide (Heldr!).

δ. *puberulum*. — Folia patule pubescenti-velutina, spicæ fasciculiformes subsessiles, bracteæ internæ abrupte mucronatae. — *A. puberulum* Boiss. et Bal. Diagn. Ser. II, 4, p. 62.

Hab. in regione alpinâ montis Aslandach Cappadociæ (Bal!), montis Beryt-lagh 8-9000' (Haussk!).

Species foliis angustis vel latiusculis sed semper brevibus varians, emper habitu pulvinato, spicis subsessilibus vel brevissime pedunculatis paucifloris angustis fasciculiformibus recognoscenda.

40. A. **Wiedemannii** (Bge loc. cit. p. 45) glaberrimum iride confertissime pulvinari-cespitosum ramis brevissimis foliis retustis reversis confertim vestitis, foliis brevissimis tenuiter subutatis acrosis subtetragonis, scapis brevissimis folia non superanibus monostachyis, spicis confertim 2-5-spiculatis brevissimis distichis, rachide glaberrimâ viscidulâ, bracteis oblongo-lanceolatis carinatis acuminatis exteriore breviore anguste interioribus late rubello-membranaceis, calycis tubo piloso, tubi albi subæquilongi subdecemlobi nervis atropurpureis ultra marginem in mucronem productis ♀.

Hab. in monte Jyldisdagh Galatiae (Wiedem!).

Folia tenuia 6-8 lineas tantum longa, spica 9-10 lineas, calyx 5 lineas longi. Facies omnino varietatum minorum *A. Echini* a quo scapo rachideque glaberrimis, spicis magis compressis differt.

41. A. Peronini, glauquum dumuloso-erinaceum ramis elongatis foliis vetustis patulis horridis, foliis patule hispidulis tenuissime subulatis patulo-subcurvatis tenuiter acerosis, scapis velutinis folia non excedentibus monostachyis, spicæ 3-5-spiculatis fasciculiformibus brevibus distichis dense imbricatis, bracteis velutinis lanceolatis dorso herbaceo velutino carinatis exteriore subbreviore acuminate-attenuatâ cæteris subbidentatis mucronatis, corollæ tubo glabrinseculo limbi albi tubo vix brevioris obsolete quinquelobi nervis atropurpureis apice subexcurrente subdilatatis 5.

Hab. in montibus Cilicie Tracheæ inter Ermenek et Mout (Peronin!).

Rami 6-9-pollicares, folia e tenuissimis pollicem non excedentia, spicæ 9-10 lineas longæ, calyx sex lineas longus. Species insignis foliis tenuibus rigidulis tenuiter acerosis, indumento patenti. His notis, glaucedine, floribus majoribus et bracteis lanceolatis elongatis ab omnibus *A. Echini* formis distincta.

42. A. Fausti (Trantv. Stirp. Nov. Descr. V, p. 16) viride dense cespitosum, foliis linearis-subulatis trigonis brevissimis præter basin ciliatulam glabris vetustis deflexis, scapis tenuibus foliis 5-6-plo longioribus monostachyis apice 2-3-spiculatis spiculis contiguis subcapi-tatis unifloris, bracteis glabris exteriore herbacea late rubello dein albido-membranacea acutâ mucronatâ, cæteris triplo longioribus præter dorsum anguste herbaceum rubello et albo-membranaceis obtusis mucronulatis, calycis tubo hirtello, limbo tubum subæquante nervis purpureis glabris ultra marginem in mucronem productis.

Hab. in Persiâ bor.-orientali inter Asterabad et Schah Kuh (Faust!). Vidi in herb. Hort. Petrop.

Mihi ex unico et minimo specimine notum, insigne foliis 4-5 lineas tautum longis, scapi 1 $\frac{1}{2}$ -2-pollicares, calyx 5 lineas circ. longus. Ex affinitate *A. Echini*.

§ 3. *Microcalycina.*

43. A. lepturoides (J. et Sp. Ill. I, tab. 95 sub *Statice*) glau-cum calcareo-punctatum cespitosum, ramis abbreviatis foliis vetustis retrorsis dense vestitis, foliis lanceolato-linearibus basi planis superne triquetris rigide acerosis, scapis glaberrimis folia 3-4-plo superantibus in spicam unicam vel binas longas æquilongas laxissimas multispi-culatas abeuntibus, rachidis glaberrimæ flexuose articulis bracteis longioribus, bracteis glaberrimis exteriore lanceolatâ anguste margi-natâ longe pungenti-acuminatâ interiores et tubum calycis superante, interioribus præter nervum carinanum membranaceis obtusis mucro-nulatis, calycis tubo hispidulo, limbo breviore sordide albo truncato nervis nigricantibus basi hispidulis paulo ante marginem evanidis ♀. *S. accrosa* M.B. Taur. Cauc. I, 252 exel. syn.

Ha' in Iberiâ praesertim occidentali inter Tiflis et Mzheti (Stev., Wilh!), Kachetiâ et Souchetâ (Eichw!).

Folia 8-12 lineas longa, scapus 2-3-pollices, spicæ 4-5-pollices longæ, calyx nunquam lineis 4 longior, quâ notâ, bracteâ externâ longiore etc. ab *A. tenui-*

floro quocum sæpe confusam fuit et ad quod locus « ditio Talysh » sæpe indicatus spectat. Amic. a Bge observante *A. lepturoïdis* Boiss. Diagn. 7, p. 78 ex specimine herb. Fisch., in quo spicae *A. acerosi* rosulis *A. lepturoïdis* commixtae fuerant, descriptio erronea delenda.

44. A. Aspadanum (Bge. loc. cit. p. 50) valde glaucum crebre calcareo-squamatum, ramis abbreviatis dense cespitosum vix suffrutescens, foliis confertis longe et late lanceolato-linearibus longe euspidato-attenuatis acerosis planis, scapis folio brevioribus 1-3-stachyis cum rachide scabris, spicis flexuosis laxe 6-12-spiculatis, rachidis articulis saltem inferioribus spiculae subæquilongis, bracteæ inferiore breviore triangulari præter carinam albo-hyalinâ mucronatâ, interioribus obtusissimis vel bilobis muticis, calycis tubo hirsuto limbo subæquilongo albo nervis rubris ad apicem perductis glabris ♀.

Hab. in montosis Persiae mediae occidentalis prope Ispahan et Murtchehar (Bgel), prope Kohrud (Kapherr!).

Folia majora sesquipollucaria versus basin 1-1½ lin. lata, spicae angustè ½-1-pollicares, bractea exterior sesquilinea interiores vix 2 lineas longæ, calyx 3 lineas longus. Ab affini *A. oligantho* ramis brevissimis vix suffrutescens, foliis longioribus et multo latioribus, bracteis internis muticis, etc., distinctum.

45. A. festucaceum (Jaub. et Sp. Ann. 20 loc. cit. p. 254 sub *Statice*) glaucescens erinaceo-cespitosum ramis abbreviatis foliis vetustis deflexis crebre obsitis, foliis hornotinis linearis-triquetris acerosis elongatis, scapis folia 2-3-plo superantibus superne paniculatum 3-7-stachyis, spicis breviter pedunculatis inferioribus strictis distantibus exteris subfasciculatis omnibus laxiusculæ 6-12 spiculatis, bracteis scariosis fuscescentibus albo-marginatis carinatis inferiore subbreviore triangulari-mucronatâ, superioribus obtusis dorso excurrente mucronatis, calycis tubo puberulo limbi albi tubo subbrevioris nervis rubris basi puberulis usque ad apicem non productis 3. *A. festucaceum* Boiss. loc. cit. p. 76.

Hab. in Persiâ loco non notato (Auch. 2169 et 5239 A!), in Mesopotamia (potius Babylonîa?) (Auch. 2511).

Folia longiora tenuia sesquipollucaria, scapi 4-5-pollicares, spicae pollicem et amplius longæ, calyx 4 lineas longus.

46. A. flexuosum (Boiss. et Haussk. — Bge loc. cit. p. 51) glaucescens calcareo-lepidotum suffruticosum erinaceum, ramis foliis vetustis horridis, foliis rigidis acerosis linearis-lanceolatis superne planis, scapis elongatis teretibus flexuosis racemose pleiostachyis, spicis sessilibus remotis angustis laxiusculæ 5-12-spiculatis, rachidis intus velutinae rectiusculæ articulis bracteis æquilongis vel brevioribus, bracteis dorso rufescenti-glumiaceis late albo-marginatis inferiore ovato-triangulari acuminatâ duplo breviori interioribus nervo hispidulis obtusissimis muticis, calycis tubo hispidulo limbum album erosio-truncatum aequante, nervis atropurpureis basi puberulis ante marginem ovanidis 3.

Hab. in calcareis alpinis montium Kuh Kilouyeh prov. Louristan Persiae occidentalis, Tehinar, Dilegun, Maregun, Sakawa 7-10000' (Hau-sk!).

Folia hornotina 12 lineas longa, scapi parte nudâ 1-2-pollicares in racemum 4-10-pollicem longum spicis 5-10 constantem abeuntes. Spica angustæ 6-12 lineas longæ, calyx 4 lineas longus. Scapi interdum abbreviati unicam spicam laxam ferentes. Affine *A. festucaceo* differt foliis brevioribus validioribus, bracteis interioribus muticis, calycis limbo tubi proportione longiore. Autem hujus varietas?

47. A. oliganthum (Boiss. Diagn. Ser. I, 7, p. 77) glaucescens calcareo-punctatum suffruticosum dumoso-erinaceum ramis vetustis patule horridis, foliis erecto-patulis subulato-triquetris acrosis basi albo-marginatis, scapis foliis brevioribus 1-2-stachyis, spicis tenuibus brevibus laxe 3-5-spiculatis, rachidis tenuis glabræ flexuosa articulis bracteis vix brevioribus, bracteis glumaceis pallidis exteriore carinata mucronato-attenuata vix breviore interioribus obtusis nervo excurrente abrupte mucronulatis, calycis tubo parce et breviter puberulo, limbi tubo duplo brevioris sordide albi nervis nigris glabris ad apicem usque productis 3.

Hab. in declivitate occidentali cacuminis alpis Kuh Delu Persiae austro-occidentalis (Ky. 559!).

Dumus saepè pedem altus, folia superiora pollicem et amplius longa, scapus cum spicâ laxissimâ folia non superans, spica altera saepius brevior, calyx 3½ lineas vix longus.

48. A. Eschkerense (Boiss. et Haussk.), glaucescens calcareo-punctatum late et laxe cespitosum ramis parte inferiori laxe horridis, foliis tenuissime subulatis inferne longe albo-membranaceis strictis tenuiter acrosis, scapis folia vix superantibus glabris 1-3-stachyis spicis 5-12-spiculatis laxissimis saepè elongatis, rachidis glabræ flexuosa articulis spiculâ subbrevioribus, bracteis scariosis pallide fuscis albo-marginatis inferiore breviore acutâ mucronata interioribus obtusis nervo excurrente mucronatis, calycis tubo parce pilosulo, limbi tubo brevioris albi nervis rubris puberulis ad apicem usque productis 5.

Hab. in monte Kuh Eschker Persiae australis 10000' (Haussk!).

Folia superiora 12-14 lineas longa tenuissima, spica terminalis saepè 2-2½-pollicaris, inferiores breviores paucifloræ. Valde affine *A. oligantho*, an ejus varietas? differt tamen cespitosus laxis, spicis longis multifloris e cespite longe exsertis. Ambo ulterius observanda.

49. A. brachystachyum (Boiss. et Haussk. MSS. — Bge loc. loc. cit. p. 51) glaucescens calcareo-punctatum erinaceo-cespitosum, foliis subulatis acrosis basi latiusculæ membranaceis inferioribus abbreviatis latioribus, scapis folia æquantibus glabris 1-2-stachyis spicis brevibus 3-7-spiculatis parte præsertim inferiore laxiusculis, rachide rectiusculâ seabrâ, bracteis glabris externâ breviore præter basin herbaceam membranaceâ triangulari acutâ, interioribus basi purpurascensibus membranaceis obtusis muticis vel breviter mucro-

natis, calycis tubo inter costas hirto, limbi tubo æquilongi nervis latis purpureis ad apicem usque productis 2.

Hab. in regione alpinâ montium Dalechani et Schahu Kurdistanie Persicæ 9-12000' et in monte Sawers 8000' (Haussk!).

Folia fere pollicaria in plantâ e monte Dalechani tenuissima, spicæ pollicem non excedentes saepè breviores, bractæ marginè albo excepto nigro-purpurascentes. Aflne *A. oligantho* spicæ laxissimæ rachide flexuosa, bracteis pallidis, etc., distincto.

B. brachyphyllum. — Folia abbreviata infima 2-3 lineas superiora 7-8 lineas longa. — *A. Kurdicum* Bge loc. cit. p. 52.

Hab. in montibus calcareis Avroman et Schahu Kurdistanie Persicæ (Haussk!).

Hæc forma intermediis ad typum transire videtur et facie *A. melanantho* similis ab eo tanen spicâ non arctissime et distiche imbricatâ sed laxiusculâ differt.

50. *A. scabrellum* (Boiss. et Haussk.) late et dense erinaceo-cespitosum glaucescens calcareo-punctatum, foliis undique seabrido-puberulis tenuiter subulato-triquetris acrosis, scapis folio brevioribus monostachyis pubescentibus, spicâ 5-10-spiculatâ angustâ præter basin dense et distiche imbricatâ, rachide intus hirtulâ, bracteis basi exceptâ seariosis pallidis inferiore ovato-triangulari acutâ cæteris sublongioribus obtusis submuticis, tubo parcisssime hirto limbi albi tubo vix brevioris nervis latis nigro-purpureis puberulis ad apicem usque productis 5.

Hab. in grainiosis montium Ssebsekuh et Kellal Persicæ austro-occidentalis (Haussk!).

Folia pollicaria vel sublongiora, spicæ angustæ 12-15 lineas longe. Aflne *A. brachystachyo*, *Eschkerensi*, *oligantho*, differt foliis seabrido-velutinis, spicis præter basin dense imbricatis, ab *A. genistoidi* foliis non biformibus, spicâ longiore angustâ.

51. *A. subulatum* (Boiss. in D.C. Prodr. XII, p. 629) dense cespitoso-erinaceum virens glabrum, foliis subulato-triquetris tenuiter et longiusculæ spinuloso-acrosis, scapis seabridis folia 3-4-plauperantibus, spicis sessilibus angustis scapo adpressis dense et stricte 5-9-spiculatis inferioribus 1-2 remotiusculis superioribus terminabilis subfasciculatis, bractæa inferiore triangulari-ovatâ mucronatâ dorso herbacea albo-marginatâ superioribus obtusis mucronatis sublongioribus, corolle tubo glabro limbi dimidio minoris albi nervis rubris ante marginem eranidis 5.

Hab. in alpinis regni Cabulici ad septentrionem urbis Cabul (Griff!).

Folia superiora 8-9 lineas longa, scapi 3-4-pollicares 3-C-stachyi, spica angustæ 6-7 lineas longæ, calyx 3 $\frac{1}{2}$ lineas longus.

52. *A. melananthum* (Boiss. Diagn. Ser. I, 7, p. 74) glaucescens pulvinari-cespitosum ramis brevibus dense horridis, foliis

glabris insimis brevissimis cæteris validis brevibus acerosis latiuscule plano-triquetris, spicâ subsessili e foliis non exsertâ dense et distiche 6-8-spiculatâ, rachide puberulâ, bracteis viscidulis purpureo-nigricantibus exteriore triangulari mucronatâ, interioribus duplo longioribus hyalino membranaceis obtusis apiculatis, calycis tubo glabriuscule limbi albidi interdum rosei nervis nigris vix mueronulato-prominulis ♀.

Hab. in regionibus superioribus alpis Kuh Daëna prope Schiraz (Ky. 591!).

Folia 9-10 lineas longa lineam lata, spicæ foliis fere occultatæ 6-7 lineas tantum longe et totidem latae, calyx 4 lineas longus. Inter affinia foliis brevibus latis, spicis dense imbricatis sessilibus, bracteis purpureis insigne. Calycis limbus in speciminibus Kotschyianis variat violascens vel albus.

53. A. genistoides (Jaub. et Sp. Ann. loc. cit. XX, p. 252, sub *Statice*) glaucescens erinaceo-cespitosum, ramis elongatis foliis vetustis remotiuseculis horridis, foliis totâ superficie hispidulis inferioribus carnosulis brevioribus submuticis apice subdilatatis cæteris triquetris acerosis, scapis folia 2-3-plo superantibus parce puberulis 2-3-stachyis, spicis brevibus dense et distiche 6-9-spiculatis rachide velutinâ, bracteis glabris externâ ovato-triangulari dorso herbaceo-rubellâ albo-marginatâ acutâ, interioribus quartâ parte longioribus praeter nervum omnino albo-membranaceis obtusis submuticis, calycis tubo hirtulo limbi albi tubo brevioris nervis atropurpureis ad marginem usque productis 5. *A. genistoides* Boiss. loc. cit. p. 76.

Hab. in Persiâ loco non notato (Auch. 2024!).

Foliis biformibus hispidulis inferioribus carnosulis submuticis, cæteris acerosis 4-4 1/4-pollicaribus, rosulis paucifoliis insigne. Spicæ breves *A. glutinacei*, calyx 4 lineas tantum longus.

54. A. brachiphyllum (Boiss. in n.c. Prodr. XII, p. 628) cespitoso-pulvinatum viridi-atratum glabrum, ramis brevissimis foliis vetustis squarroso-patentibus horridis, foliis tenuissimis tenuiter subulato-triquetris in spinulanâ tenuem flavidam abrupte attenuatis, scapis viscido-puberulis tenuibus foliis 2-3-plo longioribus monostachyis, spicâ brevissimâ dense et distiche 5-8-spiculatâ, bracteis glabris inferiore ovato-triangulari herbacea nigricanti albo-marginatâ mucronatâ, interioribus tertiatâ parte longioribus praeter nervum nigricantem albo hyalinis obtusis muticis, calycis tubo glabro, limbi albi nervis nigris ante marginem evanidis 5.

Hab. in regni Cabulici jugo Ounou et fauce Yonutt 10°00'-113°0' (Griff!).

Folia 4-5 lineas tantum longa lineæ tertiam partem lata, scapi 4 1/2-pollicares, calyx 3 lineas longus, spica densa brevis *A. genistoides*. Ex incompleto specimine descriptum, an potius pleiostachyum? Ad cl. Bge cum *A. subulato* dubitanter conjunctum sed foliis brevissimis, bracteis muticis, floribus minoribus certe diversum.

SECT. VIII. TRAGACANTHINA Bunge.

§ 1. Stenostomata.

55. **A. polystachyum** (Boiss. in D.C. Prodr. XII, p. 624) duinosum glaucum, ramis paucifloris elongatis, foliis vernalibus carnosulis brevibus margine scabridis mox recurvis et fractis, aestivalibus longe spinosis brevibus, scapis rigidis longe virgatis tota longitudine remote spicigeris, spicis parvis ovatis sessilibus squamâ suffultis dense 3-5-spiculatis, bracteâ exteriore herbaceâ ovatâ anguste, membranaceâ obtusâ mucronulatâ, interioribus duplo longioribus oblongo-linearibus carinato-plicatis præter carinam serrulato-scabram hyalinis, calycis tubo puberulo, limbo breviter exerto anguste obsolete eroso-lobato rubello striato ad nervos puberulo ȝ.

Hab. in Affghaniâ circa Karabagh prope Kabul (Griff).

Folia longiora sesquipollucaria, rami 2-3-pollucares, scapi 1-4½-pedales, spicae 10-25 intervallis semipollucaribus remotæ, calyx 2½ lineas longus.

56. **A. Karelini** (Szegl. Bull. Mosq. 1851, 2, p. 475, tab. 13 sub *Statice*) glaucum calcareo-squamulatum fruticulosum laxe foliosum, foliis vernalibus carnosulis plano-subtriquetris margine scabris deciduis, aestivalibus spinosis elongatis basi late hyalinis superne planis, scapis folia vix superantibus paniculatum pleiostachyis, spicis breviter pedunculatis elongatis laxis flexuosis 3-12-spiculatis fragilimis, spiculis subcurvatis angustis, bracteâ exteriore ovatâ acuminatâ viridi late membranacea, interioribus sublongioribus præter nervum hyalinis obtusiusculis mucronatis calycis dimidium tubum glabrum vel subpuberulum aequantibus, calycis limbo angusto plicato lobis 5 ovatis nervo prominente mucronulatis ȝ. *A. Szovitsii* Boiss. et Buhse Aufz. p. 184.

Hab. in Armeniae Rossicæ salsuginosis inter Nachitchewan et Ordubad (Kowal! Szov!).

Spiae 12-17 lin. longæ, scapi (1-poll.) cum inflorescentiâ 3½-4-pollucares, rachidis articuli longitudine bractearum vel inferiores longiores. Calyx 4 lineas longus.

57. **A. curviflorum** (Bge loc. cit. p. 57) virens fruticosum ramis elongatis longe denudatis, foliis primariis carnosulis abbreviatis recurvis fugacibus, aestivalibus remotis subulatis gracilibus basi membranaceo-marginatis, scapis gracilibus longissimis fragilibus ramoso-pleiostachyis, spicis pedunculatis laxissimis inferioribus remotis paucifloris superiore 8-12-spiculata, rachidis flexuose articulis arcuatâ saltem inferioribus spiculâ longioribus, spiculis tenuibus, bracteâ exteriore subbreviore ovatâ acutâ interioribus præter nervum membranaceis acutiusculis vel mucronulatis tubi calycini glabri tertiam

partem æquantibus, calycis tubo in limbum angustum inter lobos 5 obtusum truncatum sensim abeunte 5.

Hab. in rupestribus Persie mediae occidentalis prope Kohrud et Schurab inter Ispahan et Kum (Bge!).

Fruticulus pedalis vel altior, spinæ tenues 1 $\frac{1}{2}$ -2-pollicares, scapi cum inflorescentiæ semipedales et longiores, spica terminalis longior 3-4-pollicaris, calyx 5 lineas longus. Scapis elongatis, bracteis calycis proportione brevioribus, calycis longioris (5 lineas longi) lobis obtusis muticis ab *A. Karelini* diversum.

58. ***A. quinquelobum*** (Bge loc. cit. p. 56) glaucescens dumosum horridum, foliis vernalibus carnosulis plano-trigonis margine scabris cito emarcidis, æstivalibus subulato-spinosis supra planiusculis, scapis folio sublongioribus gracilibus 2-3-stachyis, spicis laxissimis flexuosis 4-12-spiculatis, rachidis fragillimæ articulis inferioribus spiculae æquilongis, spiculis angustis rectis, bracteis herbaceis albo-marginatis exteriore dimidio breviore acutâ, interioribus obtusis muticis dimidium tubum calycis superantibus, tubo glaberrimo in limbum angustissimum inter nervos retractos quinquelobo sensim abeunte, lobis obtusis enerviis 5.

Hab. in rupibus dolomiticiis prope Schahrud Persiae bor.-orientalis (Bge!).

Spinæ 1-1 $\frac{1}{4}$ -pollicares, spica terminalis 1 $\frac{1}{2}$ -3-pollicaris, inferiores breviores paucifloræ, calyx 4 lineas longus. Species limbo calycino inter nervos abbreviatus nec secus nervos quinquelobo ab affinibus distincta.

59. ***A. Heratense*** (Bge loc. cit. p. 55) glaucum dumosum horridum, foliis vernalibus planis brevibus margine scabris deciduis, æstivalibus spinosis subulatis triquetris, scapis folio brevioribus ? monostachyis ?, spicis elongatis laxis flexuosis fragillimis, rachidis articulis inferioribus spiculâ brevioribus superioribus abbreviatis, spiculis angustis, bracteis pallide virentibus albo-marginatis exteriore breviter ovalâ acutâ, interioribus duplo longioribus tubi trientem æquantibus carinatis acutis, calycis tubo glabro rubro-striato vel punctato in limbum minutissimum decemlobum abeunte lobis lanceolatis subæqualibus acutis 5 enerviis, 5 nervo percursis.

Hab. in apricis rupestribus prope Herat Affghaniae occidentalis (Bge!).

Sero autumno pauca specimina incompleta lecta et ideo scapi et spicæ longitudi incerta, spinæ 1-1 $\frac{1}{4}$ -pollicares, rachidis articuli inferiores 4 lineas longi; calyx 5 lineas longus limbo æqualiter decemlobo hanc speciem ab affinibus sejungit.

60. ***A. truncatum*** (Bge loc. cit. p. 56) glaucum dumosum horridum, foliis vernalibus carnosulis brevibus plano-triquetris margine scabris, æstivalibus longis spinosis trigonis calcareo-verrucosis basi membranaceis, scapis folio longioribus 2-3-stachyis, spicis elongatis laxis fragillimis, rachidis rectiusculæ articulis spiculâ brevioribus, bracteâ exteriore ovatâ acutiusculâ, interioribus sublongioribus obtusissimis vel bilobis tubi calycini tertiam partem æquantibus, calycis

tubo glaberrimo concolore in limbum brevissimum angustum omnino truncatum muticum abeunte ȝ.

Hab. in rupestribus apricis Persiae mediae occid. prope Schurab (Bge), prope Siokuh ad meridiem Teheran prope limites deserti magni (Bode). Non vidi.

Facies *A. Heratensis*, spine biplicares, bracteæ interiores $1\frac{1}{2}$ lineam non excedentes, calyx $4\frac{1}{2}$ lineas longus limbo omnino truncato non lobato quâ notâ ab affinibus discriminatur.

61. A. longiflorum (Boiss. Diagn. Ser. II, 4, p. 63) glaucescens dumosum spinis horridum, foliis vernalibus herbaceis abbreviatâ subtrigonis, æstivalibus spinosis elongatis supra planis basi vix membranaceis, scapis foliis non longioribus subcorymbose 2-3-stachyis, spicis laxe 2-4-spiculatis rachide valde flexuosa, spiculis tenuibus subcurvatis longis, bracteâ exteriore triangulari acutâ herbacea albo-marginata, interioribus duplo longioribus præter dorsum carinantem membranaceis mucronulatis. calycis tubo glabro bracteis duplo longiore in limbum vix ampliatum abeunte, lobis ȝ erectis triangulari-lanceolatis mucronatis vel muticis ȝ.

Hab. in regno Cabulico circa Choky et Kuelah (Griff!), in Belutschia circa Doubund (Stocks!).

Rami 2-3-pollicares, spine 1-1 $\frac{1}{2}$ -pollicares tennes, spicæ fragillimæ, calyx 7-8 lineas longus quâ notâ hæc species ab affinibus egrégie differt.

§ 2. **Eurystomata.**

62. A. aristulatum (Bge loc. cit. p. 60) glaucum dumoso-fruticosum, ramis elongatis, foliis vernalibus carnosulis trigonis brevibus cito emarcidis recurvis, cæteris longe spinosis præter basin planiusculam teretibns patentibus, scapis folia superantibus pleiostachyis, spicis longe pedunculatis laxissime et flexuose 3-6-spiculatis, rachidis gracilis articulis spiculâ longioribus, bracteâ exteriore ovatâ herbacea late hyalino-marginalâ in cuspidem brevem attenuatâ, interioribus duplo longioribus calycis tubum æquantibus præter nervum in aristulam productum hyalinis obtusis, calycis tubo glabro in limbum infundibuliformem quinquelobum sensim ampliato, limbi lacinias late ovato-triangularibus purpureo-aristulatis ȝ.

Hab. ad fines austro-orientales Persiae prope Ssertschah, Serdani (Bge!).

Fruticulus semipedalis vel altior, spine sesquipollicares vel longiores, calyx 4 lineas longus. Inflorescentia omnino *Stenostomatum* a quibus distinguitur limbo calycino magis dilatato infundibuliformi *Eurystomatis*.

63. A. tragacanthinum (J. et Sp. Ann. Sc. loc. cit. p. 226 sub *Statice*) glaucescens dumoso-fruticosum, ramis elongatis, foliis vernalibus carnosulis flexuoso-recurvis trigonis margine sebris, spinis longis acerosis supra planiusculis subtus convexis basi puberulis et in vaginam membranaceo-marginalam brevem dilatatis, scapis folia subsuperantibus flexuosis paniculatim 2-5-stachyis, spicis bre-

viter pedunculatis laxe 5-10-spiculatis, rachidis valde flexuosæ articulis etiam inferioribus spiculâ uniflorâ brevioribus, bracteâ exteriore albo-marginatâ triangulare-lanceolatâ interiores præter dorsum hyalinæ obtusas submuticas subæquante, calycis tubo glabro bracteis æquilongo limbo brevi patente lobis 5 triangulari-ovatis acutis nervis rubris puberulis in mucronem productis 5. *A. tragacanthinum* Boiss. Diagn. 7, p. 84. — *Statice Spachii* Girard Ann. Ser. 3, vol. I, 332 ex parte.

Hab. in Persiâ boreali occid. ad lacum Urmiah (Auch. 5239 ex parte!), inter Chorrumdereh et Sultanieh (Bge!).

Spinæ 1-1 ½-pollicares, spicæ 1-1 ½-pollicares, calyx 5 lineas longus.

64. **A. Scorpious** (Boiss. Diagn. Ser. I, 7, p. 81) glaucum dumosofruticosum ramis elongatis, foliis vernalibus carnosulis lanceolato-trigonis margine scabris demum recurvis et deciduis, spinis longis validis basi glabris et breviter membranaceis, scapis folio brevioribus vel æquilonigis 2-5-stachyis, spicis densiuscule plurispiculatis, rachidis articulis brevissimis, bracteâ exteriore mucronulatâ vel muticâ, inferioribus duplo longioribus scarioso-hyalinis obtusis vel retusis muticis vel mucronulatis, calycis bracteis subæquilongi tubo glabro, limbo inter 5 lobos ovatos mucronulatos truncato 5. *St. Scorpious* et *Sc. ferox* Jaub. et Sp. loc. cit. p. 255. — *St. Spachii* Girard ex parte.

Hab. ad Sinum Persicum (Auch. 5239 ex parte), prope Kerman (Bge!).

β. leucacanthum. — Folia vernalia snbrigidiæ non recurva, spiculæ paulo minus confertæ, calycis tubus hispidulus. — *St. leucacantha* J. et Sp. loc. cit. — *A. leucacanthum* Boiss. loc. cit. p. 81.

Hab. ad Sinum Persicum (Auch. 5239! ex parte), prope Ispahan (Bode ex Bge).

γ. incomptum. — Folia vernalia ut in formâ typicâ flexuosa, spicæ laxiuscule spiculatae rachidis articulis brevissimis, calycis tubus glaber, limbus obsolete quinquelobus muticus. — *A. incomptum* Boiss. et Buhse Aufz. p. 183.

Hab. in rupestribus pagi Taft prope Yezd (Buhse!), prope Kermanschahi inter Yezd et Kerman (Bge!).

Ipsò indicante amiciss. (Bge 1. cit. p. C2) species ternas enumeratas unicam constituere persuasus sum ramis elongatis rigidis laxe valide et longe spinosis insignem et ab *A. tragacanthino* cui cæterum valde affinis spicis plus minus densis et bracteâ inferiore multo breviore distinctam.

65. **A. Stocksii** (Boiss. Diagn. Ser. II, 4, p. 63) glaucum glabrum fruticosum ramis etiam parte hornotinâ elongatis, foliis omnibus rigidis et spinosis erecto-patulis primariis multo brevioribus cæteris elongatim spinosis subulatis tereti-subcompressis brevissimis basi late membranaceo-vaginantibus ciliolatis, scapis folia subæquantibus 2-3-stachyis, spicis brevibus laxe 2-4-spiculatis, rachidis flexuose articulis spiculâ uniflorâ brevioribus, bracteâ inferiore herbacea

triangulari albo-marginatâ acutâ, interioribus duplo longioribus herbaceo-carinatis late hyalinis mucronulatis, calycis inferne glabri superne ad costas hirsuti tubo bracteas interiores æquante, limbo patente albo lobis 5 brevibus brevissime mucronatis 3.

Hab. in Belutschia prope Ispingli (Stocks exs. 1091!).

Diffr. ab affinis ramis elongatis laxe spinosis 6-7-pollicis longis et foliis primariis etiam spinosis sed abbreviatis, folia æstivalia 1 $\frac{1}{2}$ -2-pollicaria, spicæ vix pollicares, calyx 5 lineas longus.

66. A. fasciculare (Boiss. in D.C. Prodr. XII, p. 626) viridi-glaucescens glaberrimum erinaceo-cespitosum confertifolium, ramis brevibus horridis, foliis vernalibus brevioribus complanatis margine scabris tandem revolutis, spinis tenuiter subulatis basi membranaceo-dilatatis, scapis folia æquantibus vel subsuperantibus apice fasciculatim 2-3-stachyis, spicis sessilibus brevissimis confertim 3-5-spiculatis, spiculis bifloris, bracteâ inferiore viridi anguste marginatâ aceroso-mucronatâ, interioribus sublongioribus præter nervum viridem membranaceis obtusis mucronulatis calycis tubo parce hispidulo sublongioribus, limbi truncati mutici nervis nigris vix puberulis ad marginem usque productis 5.

Hab. in montibus Affghaniae ad collum Shusgao (Griff!).

Cespites humilis, spinæ horridæ 12-14 lineas longæ, calyx 4 lineas longus.

67. A. horridum (Bge loc. cit. p. 64) glabrum glaucum dumosum ramis elongatis, foliis vernalibus breviter lanceolatis, spinis patentissimis crassis brevibus rigidis, scapis monostachyis folio brevioribus, spicis dense et distiche 2-4-spiculatis, rachide viscidulâ glabra, spiculis 1-3 floris, bracteis glabris pallide virentibus margine membranaceis exteriore breviter ovatâ obtusiuseulâ obtuse apiculatâ interioribus duplo longioribus obovatis obtusis latius hyalinis, calycis tubo glabro limbo obsolete quinquelobo nervis excurrentibus breviter aristulato 5.

Hab. in aridis montosis ad Gerdekuh inter Kerman et Yezd Persicæ australis (Bge!).

Spinae crassæ rigidissimæ 8-10 lineas tantum longæ, scapi 4-6 lineas longi, spiculae sèpius trifloræ, calyx 5 $\frac{1}{2}$ lineas longus. Affine *A. fasciculare* sed foliis brevibus crassis, floribus majoribus, etc., distinctum.

68. A. viscidulum (Boiss. Diagn. Ser. II, 7, p. 71) lète viride cespitosum erinaceum, foliis subglanduloso-viscidulis confertissimis infimis brevioribus mucronatis planiusculis basi lœvibus ceteris tenuiter spinosis brevibus supra planis subtus obtuse carinatis, scapis brevissimis monostachyis, spicâ inter folia subsessili dense et inbrickatim 4-5-spiculatâ capitato-subscorpioideâ, spiculis unifloris, bracteis nitidis viscidulis exteriore triplo breviore viridi vix marginatâ ovatâ muticâ obtuse mucronatâ, interioribus oblongis latius marginatis

obtusis muticis tubo calycis glaberrimo rubello subbrevioribus, limbi albi truncati nervis rubris glabris ad marginem productis 5.

Hab. in Persia loco non notato (Auch. 2500!).

Rami tenues 4-5-pollicares inferne foliis vetustis patentibus vix rigidis laxè obsiti, folia 6-10 lineas longa lineæ tertiam partem diam. lata, spica foliis suboccultata piso subdupo major, calyx 5 lineas longus.

69. A. Schirazianum (Boiss. Diagn. 7 p. 72) glaucescens dense cespitoso-erinaceum, foliis viscidulis vernalibus brevissimis triangulare-lanceolatis planis mucronatis basi margine scabris, cæteris patentibus brevibus tereti-subulatis basi brevissime membranaceis, scapis brevissimis monostachyis, spicâ inter folia subsessili dense et imbricatim 5-9-spiculatâ capitato-scorpioideâ, spiculis subunifloris, bracteis viscidulis exteriore triplo breviore ovato-concavâ viridi dorso plurinervi angustissime marginatâ breviter et obtuse mucronatâ, interioribus præter dorsum carinantem late hyalinis obtusis mucronulatis calycis tubum aequantibus, tubo superne parce hispidulo limbi subquinquelobi nervis rubris in mucrones subproductis 3.

Hab. in rupestribus ad monumentum Nakschi Rustam prope Schiraz Persiae australis (Ky. 305!).

Affine *A. viscidulo* a quo differt glaucescentiâ, foliis tereti-subulatis nec facie superiore planis, bracteâ inferiore marginatâ et apice abruptius attenuatâ etc.

70. A. tomentellum (Boiss. Diagn. Ser. I, 7, p. 72) glaucescens dumoso-cespitosum totum viscidulo-tomentellum, ramis brevibus foliis confertis horridis, foliis vernalibus brevissimis triangulari-lanceolatis margine scabris, aestivalibus validis rigidis breviusculis acerosissimis supra planis subtus convexis, scapis brevissimis pubescentibus monostachyis, spicâ e foliis non exsertâ arcte et distiche 5-6-spiculatâ capitato-scorpioideâ, rachide glabrâ viscidulâ, spiculis mono-rarissime 2-3-floris, bracteâ exteriore dimidio breviore ovatâ obtusiusculâ mucronulatâ albo-marginatâ, interioribus dorso viscidulis latissime albo-membranaceis truncato-subretusis, calycis tubo adpresso puberulo viscidulo bracteis subbreviore, limbi subtruncati mutici nervis rubris puberulis ad marginem usque productis 3.

Hab. in lapidosis alpis Kuh Daïna Persiae australis supra Schiraz (Ky. 670!).

Dumi erinacei semipedem longi, spine validae 9-10 lineas longæ, calyx 4 lineas longus, spicæ piso subdupo maiores compressæ.

71. A. Talagonicum (Boiss. in D.C. Prodr. XII, p. 625) viride erinaceo-cespitosum confertifolium totum breviter pubescens, ramis subelongatis, foliis vernalibus brevissimis recurvis planis margine scabris aestivalibus subulatis longis spinosis supra planis subtus convexis basi breviter marginatis, scapis folio sublongioribus 2-3-stachyis, spicis brevissime pedunculatis approximatis arcte et imbricatim 6-8-spiculatis brevissimis subscorpioideis, spiculis 1 rarissime 2-3-floris,

rachide viscidulâ, bracteis glabris viridibus nitidis late albo-margina-tis inferiore breviter ovatâ submuticâ interioribus duplo longioribus truncatis muticis calycis tubum æquantibus, calycis tubo parce hispi-dulo, limbi erosi vix sinuati mutici nervis puberulis pallide ad margi-nem usque productis 5.

Hab. in valle Talagon jugi Elbrus Persiæ bor. ad Gattade (Ky. 514!).

Folia eis *A. tomentelli* tenuiora viridia longiora 1-1 $\frac{1}{4}$ -pollicaria, scapus longior pleiostachys, bracteæ tubi proportione breviores.

72. **A. Schahrudicum** (Bge loc. cit p. 65) glaucum breviter tomentellum dumoso-fruticosum ramis junioribus interdum subflexuosum, foliis vernalibus carnosulis brevibus trigonis margine sca-bris, spinis validis elongatis supra planis subtus convexis basi anguste marginatis, scapis pubescentibus folio duplo longioribus 3-5-stachyis, spicis cymosis inferioribus longius pedunculatis omnibus confertim et distiche 5-8-spiculatis, rachide viscidulâ glabrâ, spiculis 1-3-floris, bracteis omnibus pallide virentibus glabris nitidis late scariosis obtusis muticis exteriore dimidio breviore internis truncato-bilobis tubum calycis subglabrum æquantibus, limbi truncati mutici erosuli nervis pallidis ante marginem evanidis 5.

Hab. in aridis dolomiticis prope Schahrud (Bge!), ad Ssof inter Ispahan et Kaschan (Buhse! Aufz. sub A. sp. indetem.), prope Kum (Bode!).

Inflorescentiâ multiolorâ cymosâ ab *A. Talagonico* distinctum, folia crassi-ora longiora (1 $\frac{1}{2}$ -pollicaria), spiculae sæpius trifloræ, calyx 4 lineaæ longus.

73. **A. roseum** (Boiss. Diagn. Ser. 1, 7, p. 71) viride dense erinaceo-cespitosum ramis abbrevialis crebre foliosis, foliis præter vernalia pauca brevia cito decidua tenuiter spinosis erectis vel erecto-patulis supra planiusculis subtus obtuse carinatis pubescentibus rarius glabris, spicis terminalibus solitariis subsessilibus vel scapo brevissimo pubescenti suffultis fasciculatim 3-5-spiculatis rachide brevissimâ pubescente, spiculis unifloris, bracteis glabris vel velutinis exteriore breviore pungente lanceolatâ viridi anguste membranaceâ, interioribus præter carinam rufesceni - membranaceis acutis vel retusis mucronatis calycis tubo glabro brevioribus, limbo tubo bre-viore expanso roseo truncato mutico 5. *St. pungens* et *St. erinacea* J. et Sp. Ill. I, p. 163. — *St. Jauberti* Girard Ann. 1844, p. 331.

Hab. in jugo Elbrus Persiæ bor. (Hablitzl, Auch. 5249!), in alpinis prov. Khorassan et prov. Kuhistan Persiæ orientalis (Bge!), in montanis prope Teheran (Jenisch), in alpibus regni Cabulici (Griff!), in monte Dalechani Kurdistaniæ Persice 10000' (Haussk!).

Cespites densi latiusculi 3-5-pollices alti, folia tenuia 7-9 lineaæ longa. Calyx 4 $\frac{1}{2}$ lineaæ longus. Species calycibus purpureis elegans.

74. **A. Gulistanum** (Bge loc. cit. p. 67) glaucescens tenuiter puberulum fruticulosum ramis elongatis laxe et subverticillatim foliosis inter spinas denudatis, foliis vernalibus numerosis fugacibus,

spinis tenuibus elongatis curvato-patentibus, scapis tomentellis folia æquantibus vel brevioribus fasciculatim 2-3-stachyis, spicis confertim et distiche 4-7-spiculatis rachide pubescente, bracteis puberulis exteriore subbreviore a medio lanceolato-attenuatâ cuspidatâ basi membranaceo-marginatâ interioribus viridi-carinatis late fusco-marginatis acutis calycis lobo longioribus, calycis tubo glabriusculo in limbum tubo duplo breviorem obsolete quinquelobum muticum angustum pallide rubellum sensim dilatato 5.

Hab. in rupestribus aridis demissoribus prov. Khorassan Persiae prope Gulistan et in montibus Tabat Kuh et Kerat Kuh (Bge!).

Dumus semipedalis et elatior, spinæ ob innovationes distinctas subverticillatæ tenues 1-1½-pollicares, spicæ stricte et anguste distichæ 5-6 lineas longæ, calyx 5-6 lineas longus.

GONIOLIMON (Boiss. in D.C. Prodr. XII, p. 632).

Calyx infundibuliformis limbo scarioso quinquenervi plicato quinquelobo. Petala quinque imâ basi tantum annulatim connexa unguium marginibus in tubum contiguis. Filamenta basibus glandulosis subconnexa. Styli quinque liberi totâ longitudine papilloso-asperi. Stigma capitato-subdepressa. Utriculus membranaceus superne operculo brevissimo circulariter circumscisso dehiscens. — Herbæ scapis ramisque elatis vel angulatis.

1. **G. Tataricum** (L. Sp. I, p. 395 sub *Statice*) glaucescens, foliis obovatis vel oblongo-spathulatis mucronatis in petiolum brevem sensim attenuatis, scapo humili paulo supra basin divaricatim dichotomo late et patentim corymboso ramis anguste trialatis patentirecurvis, spiculis 1-2-floris in spicas terminales densiusculas distiche dispositis. bracteis herbaceis coriaceis anguste membranaceis pungentibus acute carinatis interiore tricuspidatâ cuspidibus rectis subæquilongis, calycis tubo undique breviter puberulo limbi æquilongi lobis oblongis obtusis ♀. Ic. Gmel. Sib. 2, tab. 92. — *Gon. Tataricum* Boiss. in D.C. Prodr. XII, p. 632. — Rchb. Ic. Germ. tab. 88.

Hab. in salsis prov. Caucasicarum (M.B., Hohen!), Tauriæ (Stev!).

β. angustifolium Boiss. loc. cit. — Folia angustiora lanceolata. rami angustissime alati, spiculæ sæpius unifloræ, calycis tubus sæpe glabrescens. — *S. incana* M.B. Taur. Cauc. I, p. 251 non alior. — *S. Besseriana* R. et Sch. VI, p. 789. — Rchb. Ic. Crit. tab. 270.

Hab. in Tauriæ (Urv!).

Ar. Geogr. Transylvania, Bulgaria, Rossia australis, Sibiria Uralensis.

2. **G. graminifolium** (Ait. Kew. p. 383 sub *Statice*) glaucescens, foliis anguste linearibus canaliculatis acutis, scapo superne divaricatim dichotomo patentim corymboso ramis tenuibus angulatis apte-

ris patent-recurvis, spiculis unifloris distantibus spicas tenues elongatas unilaterales formantibus, bracteis carinatis late membranaceo-marginatis interiore tridentata dentibus rectis intermediâ longiore lateralibus saepe rotundatis, calycis limbo in lobos lanceolatos acutiusculos profunde fisco ♀. *G. graminifolium* Boiss. loc. cit. — Ic. Rehb. Crit. fig. 963. — Trautv. Imag. tab. 9.

Hab. in salsis Tauriae (herb. Fauché!).

Foliis elongatis lineam latis, ramis aperis, spiculis dissitis, bracteis late membranaceo-marginatis, calycis dentibus elongatis a precedente distinctum videtur.

Ar. Geogr. Rossia australis.

3. **G. Dalmaticum** (Presl Bem. 105 sub *Statice*) glaucum, foliis parvis albo-tuberculatis margine anguste et crispule membranaceis, scapo brevi in paniculam brevem triangularem secundam abeunte, ramis brevibus angulatis fere aperis, spiculis subbifloris in spicas brevissimas distiche et dense imbricatis, bracteis coriaceis anguste membranaceis carinatis pungentibus interiore tricuspidatâ cuspidibus rectis intermediâ sublongiore, calycis tubo adpresso puberulo limbi lobis oblongo-lanceolatis obtusis ♀. *G. Dalmaticum* Rehb. Ic. Germ. XVII, tab. 149. — *S. incana* Visian. Fl. Dalm. II, p. 7 non L.

Hab. in herbidis collium ad Philippopolin Thraciae (Janka exs. sub *S. col-* *kind*!).

Affine *G. Tatarico* quocum hoc in Prodr. conjunxeram, folia multo minora calcareo-tuberculata, rami fere aperi, spicae breviores fasciculiformes, bracteae intimae dentes inaequales.

Ar. Geogr. Dalmatia, Croatia.

4. **G. collinum** (Griseb. Spicil. II, p. 300 sub *Statice*) glaucescens, foliis anguste oblongo-lanceolatis lanceolatisque aristato-mucronatis in petiolum longe attenuatis, scapo humili jam prope basin divaricatum dichotomo corymboso-paniculato ramis triquetris patuio-recurvis, spiculis unifloris paucis in spicas fasciculiformes breves crebras densiusculae imbricatis, bracteis viridibus coriaceis carinatis subrecurvis interiore inferne membranaceo-marginata a medio tricuspidatâ cuspidibus acutis intermediâ sublongiore subrecurvâ, calycis tubo glabro limbi tubo brevioris lobis oblongis obtusis ♀. *G. collinum* Boiss. loc. cit. p. 633. — *S. speciosa* Forsk. Cat. Constant. non L. — *S. incana* Vahl Symb! Linn. Mant? — *S. Tatarica* β. *trigonoides* Pois. Encycl!

Hab. in siecis Thraciae (Friv! sub *S. Besserianâ*), Byzantii (Forsk.), Bithyniae (Thirke!), Lydie (Auch. 2494! Boiss! Bal. 286!), Anatoliæ sept. ad Sojut (Wied!), Syriae interioris inter Tripolin et Hama (Bl!).

Species spicis brevibus fasciculiformibus et bracteis subrecurvis a praecedentibus distincta. *G. Serbicum* Vis. et Paucie Doc. I, tab. IV postea a cl. auctoribus in Dec. III, p. 24 ad *G. collinum* reductum ab eo bracteis pubescitibus, cuspidibus rectis differt.

5. **G. Sartorii** (Boiss. Diagn. Ser. II, 4, p. 67) glaucum, rhizomate lignoso cespitoso, foliis parvulis ovato-rhombeis acuminatis in petiolum breviter attenuatis, scapis brevissimis simplicibus angulatis in paniculam parvam brevem oblongam secundam abeuntibus, spiculis paucis unifloris fasciculatis, bracteis coriaceis basi late albo-marginatis recurvis inferiore deltae ad tertiam partem in cuspides ternas breves recurvas fissâ, calycis tubo obsolete glanduloso, limbi lobis breviter triangularibus 5.

Hab. in littoribus Atticæ prope Ergastiria in circuitu Laurion Atticæ (Heldr!), insula Mycone ad sinum Phtelia (Sart!).

Foliorum lamina 3-4 lineas longa, scapus bipollicaris, panicula pollicem longa spicis 5-7 tantum constans. Hic notis, floribus minoribus et præsertim rhizomate lignoso pluricipiti a precedentibus distinctum.

STATICE (L. Gen. 388 ex parte. — Willd. Enum. p. 335).

Calyx obconicus tubulosus vel sæpiissime infundibuliformis limbo scarioso quinquenervi, quinque vel rarius decemlobo vel partito. Corolla vel polypetala vel annulatim et basi tantum aut ad limbum usque gamopetala. Stamina basi rarius ad medium petalis adnata. Styli glabri vel liberi vel basi brevissime connati. Stigmata filiformi-cylindrica totâ superficie glandulosa. Utriculus parte superiori durior inferiore membranaceus vel apice operculatum vel prope basin irregulariter ruptilis. — Herbæ perennes vel annuæ aut suffrutices floribus in spiculas tribracteatas dispositis.

Specierum Orientalium distributio.

SERIES I. — Petala libera vel imâ basi tantum connexa.

SECT. I. PTEROCLADOS Boiss. in D.C. Prodr. — Ramuli florales ancipi-ti-complanati vel trialato-obpyramidati. Calyx insertione rectus limbo ampio plicatissimo.

S. sinuata, Thouini.

SECT. II. LIMONIUM Boiss. loc. cit. — Scapi et rami teretiusculi vel angulati. Calyx insertione subobliquus vel subobliquus. Petala integra vel retusa. Inflorescentia varia.

* Perennes rhizomate sæpe lignoso sed non suffruticosæ.

+ Bracteæ inferiores margine tantum membranaceæ.

S. Limonium, Gmelini, *effusa*, *pycnantha*, *globulariæfolia*, *Corinthiaca*, *ocymifolia*, Sieberi, *rorida*, *virgata*, *cancellata*, *Cosyrensis*, *nuda*.

++ Bracteæ inferiores fere omnino hyalinæ.

S. latifolia, Caspia, Iconia, pruinosa, otolepis, perfoliata.

++ Rami suffruticosi foliati.

S. obovata, suffruticosa, leptophylla, carnosa, axillaris, Arabica, Stocksu.

SECT. III. SPHEROSTACHYS Boiss. in D.C. Prodr. XII, p. 663. — Perennis. scapi et rami teretiusculi. Calyx insertione rectus, spiculæ in globulos pisiformes congestæ.

S. globulifera.

SECT. IV. SCHIZOPETALUM Boiss. Diagn. Ser. II, 4, p. 68. — Perennis, scapis et ramis teretibus. Petalorum limbus bipartitus. Spiculæ longe et laxe spicatæ.

S. macrorrhabdos.

SECT. V. SCHIZYMENTUM Boiss. in D.C. Prodr. XII, p. 665. — Spiculæ laxe spicatæ. Calycis tubus insertione rectus, limbus in 5-10 lacinias partitus.

S. echiooides, Oicerini, Cabulica.

SERIES II. — Corolla gamopetala.

SECT. VI. SIPHONANTHA. — Calycis tubus insertione rectus, limbus ad basin usque quinquepartitus. Filamenta fere usque ad apicem corollæ tubo adnata. Inflorescentia scorpioideo-corymbosa. Perennis.

S. tubiflora.

SECT. VII. PSYLLIOTACHYS Jaub. et Sp. — Calycis tubus insertione rectus infundibuliformis. Filamenta inferne tantum corollino tubo adnata. Inflorescentia spicata. Herbæ annuæ.

S. spicata, leptostachya.

SECT. I. PTEROCLADOS.

1. **S. sinuata** (L. Sp. p. 397) perennis pilis e tuberculo ortis hispido-seabra, foliis rosulatis sinuato vel pinnatifido-lobatis vel lyratis sinibus lobisque rotundatis terminali setifero. scapis dichotome corymbosis crispatum 3-5-alatis alis ad nodos in appendices

ternas verticillatas linear-lanceolatas acuminatas abeuntibus, spiculis 3-4-floris in spicas breves secundas summis ramulis trialatis apice triangulari-appendiculatis insidentes dense et distiche imbricatis, rachide facie inferiori squamulis linearibus ciliatis biseriatim pectinatâ, bracteis inferioribus membranaceis lanceolato-subulatis carinatis ciliatis, interiore viridi coriacea latiore dorso acute bicarinata apice truncata inaequaliter 2-3 dentata, calyx tubo cylindrico glabro limbo ampio cæruleo truncato crenulato, corollâ ochroleucâ ♀. Ic. Fl. Græc. tab. 301. — Bot. Mag. tab. 71. — *S. hirsuta* Presl Bot. Bem. p. 105.

Hab. in collibus regionis inferioris Peloponnesi! et Atticæ!, Macedonia in monte Athos (Griseb!), Thracia ad Byzantium (Noë), Cycladum (Urv! etc!), Cretæ (Sieb!), Rhodi (Bourg!), Cypri (Ky!), totius Anatoliæ occid. (Bal! Forbes!), Syriae littoralis (Bl! Gaill!), Palestine (Roth!).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Sicilia, Sardinia, Italia australis, Africa borealis.

2. S. Thouini (Viv. Cat. Hort. Negr. p. 34) annua glaucescens, foliis rosulatis sinuato vel pinnatifido-lyratis lobis sinibusque rotundatis margine breviter ciliatis, scapis superne dichotome corymbosis plus minus late trialatis alis rectis sub dichotomiis superioribus in appendices triangulares productis, spiculis 2-3-floris in spicas breves secundas summis ramulis trialatis insidentes distiche et dense imbricatis, ramulorum alis reticulatis a basi sursum valde dilatatis sub spicâ in appendices triangulares flores subæquantes productis, rachide subtus nudâ, bracteis inferioribus membranaceis cuspidis, interiore multo majore virenti coriacea dorso bicarinato-plicata carinis sub apice membranaceo trilobo in dentes binos duros patulo-recurvos inaequales abeuntibus, calycis tubo glabro, limbo albo tubo æquilongo ad medium in lobos triangulari-lanceolatos acutos et quinque setas cum lobis alternantes fiso, corollâ flavâ ♂. *S. Egyptiaca* Pers. Del. Fl. Eg. t. 25. — *S. tripteris* H. Par. -- *S. alata* W.

Hab. in siccis regionis inferioris, insula Ægina Græciae (herb. Fauché!), Palestina ad conventum Mar Saba et in valle Jordani (Boiss!), Arabia petrea (Auch! Boiss!), Egyptus inferior (Del! Ehr!), Persia australis ad Dalechi (Haussk!).

Ar. Geogr. Hispania australis, Africa borealis a regno Maroccano ad Cyrenaicam.

SECT. II. LIMONIUM.

3. S. Limonium (L. Sp. 394) glabra glaucescens, scapis elatis teretibus dichotome ramosissimis laxe paniculato-corymbosis, ramis elongatis patentibus, spicis secundis densis vel laxiusculis saepe scorpioides, foliis oblongis vel oblongo-lanceolatis in petiolum attenuatis obtusis, spiculis 2-3-floris, bracteis inferioribus herbaceis late albo-marginatis muticis vel dorso subcarinato excurrente submucronatis,

interiore triplo majore muticà late membranaceo-marginatâ, calycis tubo ad costas piloso, limbi tubo brevioris lobis triangularibus acutiusculis φ . — *S. Limonium* var. α . *genuina* et γ . *macrooclada* Boiss. in D.C. Prodr. — *S. serotina* Rehb. Ic. Crit. 8, tab. 998. — *S. Gmelini* Koch Syn. non W. — *S. Meyeri* Boiss. in D.C. Prodr. quoad specimina Graeca

Hab. ad littora maritima, Attica ad Phalerum (Sprun!), Peloponnesus et Cyclades (Bory!), Creta (Sieb!), Byzantii (Boiss!), Smyrnae (Bal. 232!), Syria littoralis ad Sidonem et Berythum (Bl! Gaill!).

Ar. Geogr. Littora totius Europae australis et Africæ borealis. Quoque ad littora Oceani et Baltica, *S. Limonium* β . *Behen* Boiss. in D.C. Prodr. = *S. pseudolimonium* Rehb. specifice non differre probatur.

4. **S. Gmelini** (Willd. Spec. I, p. 1524) glabra, scapis elatis gracilibus confertim corymboso-paniculatis, ramis patent-curvatis, spiculis subbitloris in spicas breves secundas subscorpioideas dense imbricatis, foliis ovatis vel oblongis obtusis in petiolum attenuatis interdum sessilibus, floribus parvis, bracteis duabus inferioribus ovato-rotundis fere omnino membranaceis exteriore interdum mucronulata, bracteâ interiore triplo majore late albo-marginatâ obtusâ, calycis tubo ad costas omnes vel nounullas piloso limbi lobis rotundatis erosio. φ . Ic. Rehb. Crit. tab. 391. — Ic. Germ. tab. 90.

Hab. in maritimis Eubœa sept. (Wied!), in peninsulâ Athoâ Macedonia (Orph!), Byzantii (Cast!), Tauriâ (M.B., Rehm!). Quoque in interioribus sal-suginosis, planities Suleovasti Ponti (Tchih!), in Armenia ad Euphratem prope Erzerum (Calv!).

β . *lilucina*. — Spiae densifloræ formæ typicæ sed folia minora acutiuscula et calycis tubus longius atque etiam inter costas sœpe pilosus. — *S. lilacina* Boiss. Diagn. Ser. II, 4, p. 68.

Hab. in salsis Cappadocie inter Indjesou et Cæsaream 3300' (Bal. 997!).

γ . *laxiflora* Boiss. in D.C. Prodr. — Panicula amplior magis divaricata, spiculae laxius spicatae. — *S. scoparia* M.B., C.A.M. non Willd. — *S. Meyeri* Boiss. in D.C. Prodr. XII, p. 645.

Hab. in maritimis Tauriæ (Stev!), ad littora Cis- et Transcaucasica mari Caspii (C.A. M! Hohen!), Persiæ borealis (Szov! Buhse!) et quoque interioribus prov. Khorassan inter Mesched et Herat (Bge!).

Species a *S. Limonio* floribus minoribus, bracteis externis fere omnino membranaceis et calycis lobis obtusis distincta, specimina quædam tamen (in secco) interdum subdubia.

Ar. Geogr. Banatus, Transylvania, Bulgaria, Rossia australis, regio Uralensis, Sibiria Altaica et Baicalensis.

5. **S. effusa** (Boiss. in D.C. Prodr. XII, p. 646) glabra glaucescens, scapis elatis dichotome ramosissimis in paniculam effusam abeuntibus, ramis elongatis tenuibus patulo-recurvis, foliis oblongis obtusis setaceo-mucronatis in petiolum brevem attenuatis, spiculis bifloris parvis valde dissitis unilateraliter spicatis, bracteis binis exterioribus

præter basin herbaceam hyalinis acutis interiore quadruplo majore late hyalino-marginatâ obtusâ, calycis tubo basi ad costas binis ciliato, limbi tubo duplo brevioris lobis 10 minatis subæqualibus erosulis ♀.

Hab. in Lydiâ interiore prope Hierapolin (Mitch!).

Folia cum petiolo triplicaria pollicem lata, scapi bipedales, spiculae 1½-2 lineas distantes. Facies *S. latifoliae* sed magis affinis *S. Gmelini laxiflorae* a qua spiculis valde dissitis et limbi calycini plicis lobos subæquantibus differt. Ex unico specimine mihi cæterum nota.

6. ***S. pycnantha*** (C. Koch Linn. XXI, p. 716) glabra glauca, foliis crebre rosulatis oblongis acutiusculis in petiolum attenuatis, scapis pluribus pumilis erectis corymboso-paniculatis spiculis parvis bifloris in spicas brevissimas scorpioideas dense imbricatis, bracteis binis externis ovato-triangularibus hyalinis acutiusculis, tertia triplo majore obtusa late membranace marginatâ, calycis tubo basi intus ad costam hirtulo, limbo tubo breviore fere usque ad basin in lobos triangulares acutos partito ♀. *S. Balansæ* Boiss. Diagn. Ser. II, 4, p. 69.

Hab. in salsuginosis ad occidentem urbis Cæsareæ in Cappadociâ (Bal!), ad Tuslachai prov. Musch Armeniæ (C. Koch!).

Species mihi ex paucis speciminibus nota; folia sesquipollucaria 4-5 lineas lata, scapi 4-5-pollicares, spicæ valde scorpioideæ subcapituliformes in corymbo subcontiguae, flores illis *S. Gmelini* minores, ab ea calycino limbo acute et profunde quinquepartito distincta

7. ***S. globulariæfolia*** (Desf. Fl. Atl. I, p. 274) glauca lœvis glabra, foliis rosulatis coriaceis obovatis acutiusculis vel obtusis macronatis in petiolum brevem attenuatis, scapis elatis rigidulis flexuosis dichotome ramosis effuse corymboso-paniculatis, ramis inferioribus paucis sterilibus ramulosis interdum nullis, spiculis 2-3-floris in spicas tenues subincurvas unilaterales sœpe elongatas laxè dispositis, bracteis rufis albo-marginatis inferiori ovato-triangulari carinatâ acutâ, superiore triplo longiore incurvâ dorso excurrente apiculatâ, calycis tubo ad costas parce et adpresso puberulo, limbo albo tubo subtroplo breviore lobis ovatis acutiusculis ♀. *S. Raddeana* Boiss. in d.c. Prodr. XII, p. 653.

Hab. in salsuginosis Egypti inferioris ad Alexandriam (Radde! Gaill! Dupar.!), ad Damiatam (Ehrenb!).

Scapi cum paniculâ 1-3-pedales, folia cum petiolo 2-3-pollicaria, spicæ 1-1½-pollicem longæ, calyx vix 2 lineas longus. Rami steriles in speciminiibus Ægyptiacis plus minus numerosi vel prorsus nulli variant et nullum igitur à plantâ Atlanticâ disserimen.

Ar. Geogr. Africa borealis.

8. ***S. Corinthiaca*** (Boiss. et Heldr. Diagn. Ser. II, 4, p. 69) glabra glaucescens, basi induratum cespitosa, foliis rosulatis oblongo-spatulatis obtusis in petiolum attenuatis, scapis erectis rigidulis in paniculam secundam abeuntibus, ramis omnibus fertilibus, spiculis

2-3-floris in spicas terminales erectas vel subineurvas secundas dense et distiche imbricatis, bracteis rufescentibus coriaceis anguste marginatis inferioribus ovato-triangularibus acutiusculis interiore triplo majore dorso convexâ obtusâ, calycis tubo inferne ad costas pilosulo, limbi minuti albidi lobis ovatis obtusis ♀.

Hab. in saxosis prem. Maero Nicolaos ad sinum Corinthiacum (Heldr!).

Folia sesquipollucaria vel breviora, scapi cum paniculâ 6-9-pollucares, affinis *S. oxylepidi* Boiss. = *densiflora* Guss. non Girard, differt floribus manifeste brevioribus et spicis angustioribus, bracteis inferioribus superioris proportione brevioribus et praesertim calycino limbo brevissimo nec valde evoluto albo-membranaceo. A *S. ocyminifolia* spicis densioribus, spiculis dimidio minoribus, calycis tubo basi tantum nec toto piloso, foliis minoribus angustioribus distinguitur.

9. *S. ocyminifolia* (Poiret Dict. V, p. 238) glaucescens leviuscula, rhizomate indurato, foliis carnosis obovato-spathulatis obtusissimis in petiolum æquilongum sensim attenuatis, scapis flexuosis dichotome ramosis paniculato-corymbosis, ramis sterilibus subnullis, spiculis 2-3-floris rectis in spicas tenues longas unilaterales distiche et densiuscule approximatis, bracteis herbaceis coriaceis anguste fuscescenti-membranaceis inferiore ovato-triangulari acutiusculâ superiore triplo longiore obovato-oblongâ obtusissimâ, calycis tubo tenui adpresso piloso, limbi brevissimi albidi lobis brevibus erectis ovatis obtusis ♀.

Hab. in insulis Archipelagi (Tourn. in Desf. herb!), in insula Audros (Santorit!), in insula Melos prope Clima (Orph. Fl. Gr. exs. 694!), ad seopulum Raphiti Attice (Urv!).

♂. *bellidifolia*. — Spiculae dissitae. — *S. bellidifolia* Sibth. et Sm. Fl. Græc. III. p. 90, tab. 295 excl. syn. — *S. Aucheri* Girard Ann. Sc. Nat. 3^{me} sér. 2, p. 38.

Hab. in maritimis insulae Rhodi (Sibth., Bourg!), insulae Thermia (Auch. 2506!).

Scapi cum paniculâ 1/2-1-pedales et interdum longiores, spicæ 1-3-pollucares densiusculæ et laxæ variant cum internodiis. Species affinis *S. spathulata* Desf. spiculis tertiatâ saltem parte majoribus (4 lineas longis), bracteis inferioribus omnino membranaceis inferiore late membranaceo-marginatâ distinctæ. *S. psiloclada* Boiss. affinis quoque praesertim varietati ♂. differt spiculis minoribus, bracteis inferioribus omnino membranaceis, superiore acutiusculâ etc.

10. *S. Sieberi* (Boiss. Voy. Esp. p. 530 in notâ) glaucescens leviuscula, rhizonate indurato ramoso, foliis obovato-spathulatis obtusis vel acutiusculis in petiolum longe attenuatis, scapis elatis erectis rigidulis flexuosis dichotome ramosissimis ramis inferioribus sterilibus paucis rigidis abbreviatis, spiculis 1-2-floris rectis valde dissitis spicas longas laxas unilaterales formantibus, bracteis herbaceis coriaceis anguste fuscescenti-membranaceis exteriore oblongâ acutiusculâ interiore triplo longiore marginibus involvâ, calycis tubo adpresso hirtulo limbi parvi lobis ovatis obtusis ♀. *S. spathulata* Sieb. herb.

Cret. non W. — *S. Græca* & *Sieberi* Boiss. in D.C. Prodr. XII, p. 650.

Hab. in maritimis Cretæ ad Retimo (Sieb!), ad Khania (Raul!), Atticæ ad Phalerum (Spr! Heldr! Orph!), Archipelagi (herb. Fauché!), insulæ Leros (Oliv!), Syriæ littoralis ad Berythum (Ehr! Bl! Gaill!).

Scapi $\frac{1}{2}$ -1-pedales, folia eis *S. ocytropifoliae* sæpius minora; præsertim ejus formis laxifloris valde affinis et forsitan ulterius conjugenda, differt tamen scapis et ramis rigidioribus, ramis inferioribus sæpe abortivis, spiculis oliganthis submajoribus (tres lineas longis) magis dissitis, braetea interna involuta. Proxima quoque *S. virgata* W. quâcum mixta sæpe crescit et formas transitiones (hybridas?) sæpe præbet, scapis et ramis robustioribus, spiculis tertia saltem parte longioribus magis dissitis rectis nec incurvis tantum distingua.

11. *S. rorida* (Sibth et Sm. Fl. Græc. III, p. 91, tab. 298) tuberculato-aspera glaucescens basi indurata cespitosa, foliis parvis confertis ovato vel oblongo-spathulatis basi longe attenuatis obtusis subtus carinatis, scapis a basi infraeto-ramosissimis breviter articulatis crassinervulatis, ramulis inferioribus sterilibus rigidis divergentibus brevibus superioribus in spicas laxas breves unilaterales aequalibus, spiculis 1-3-floris rectiuseculis, bracteis inferioribus submembranaceis acutis interiore triplo longiore coriaceâ anguste fusco-marginatâ obtusa, calycis tubo hirtulo limbi parvi lobis ovatis obtusis φ . *S. echioides* Smith Prodr. p. 213, non L. — *S. Græca* var. *microphylla* Boiss. in D.C. Prodr. p. 650.

Hab. in maritimis Atticæ prope Eleusin (Orph. Fl. Gr. exs. 265!), ad Phalerum (Heldr!), insulis Tenos et Thera (Sart!), Syra et Myconos (Orph!), Melos (Armenist!), Naxos (Bory!), in Syriâ prope Berythum (Bl! Gaill!). Indicatur præterea a Sibthorpio in Cretâ, littoribus Ioniæ et Cypro.

$\beta.$ *palmaris*. — Scapi nani, spicæ breves paucispiculatæ, spiculæ minores, folia lanceolato-spathulata vel obovata. — *S. Græca* Poir. Dict. Suppl. 5, p. 237, Jaub. et Sp. Ill. tab. 389. — *S. palmaris* Fl. Græc. III, p. 91, tab. 297. — *S. Tournefortii* Jaub. et Sp. Ill. tab. 388.

Hab. in maritimis Græciæ, insula Helena (Macronisi) Atticæ (Tournef.), littora Corinthiaca (Pichl!), insulæ Archipelagi (Tourn.), littora Asiae minoris (Sibth.).

$\gamma.$ *prolifera*. — Scapi elongati ad dichotomias fasciculos foliorum radicalibus similes edentes, ramuli steriles pauci, folia obovato-spathulata — *S. prolifera* Urv. Enum. p. 35. — Jaub. et Sp. Ill. tab. 390.

Hab. in littoribus Atticæ ad Eleusin, Megaram et ad Munychiam (Heldr!), insulis Ferusiâ, Therâ et Samo (Urv!).

$\delta.$ *hyssopifolia*. — Scapi elongati ad dichotomias fasciculos foliorum radicalibus similius sed minorum sæpe edentes, folia angustæ lanceolato-spathulata sæpe acutisecula, spicæ elongatæ. — *S. Oliviana* Andr. in D.C. herb. — *S. hyssopifolia* Girard loc. cit. p. 329.

Hab. in maritimis Cretæ prope Kissamos (Oliv! Heldr!), Rhodi (Bourgl!).

Species polymorpha a *S. ocytropifolia* et præsertim *S. Sieberi* quibus affinis foliorum et ramorum superficie tuberculato-aspera distincta.

12. S. virgata (Willd. Enum. Berol. I, p. 336) glabra lăviusculă basi indurata cespitosa, foliis oblongo vel lanceolato-spathulatis obtusis basi longe attenuatis, scapis numerosis saepius elatis rigidulis flexuosis stricte ramosissimis, ramis inferioribus rigidis sterilibus superioribus paniculatis, spiculis 2-4-floris angustis saepius incurvis spicas unilaterales saepius elongatas laxiusculas formantibus, bracteis anguste rufescenti-marginatis inferioribus ovatis acutiusculis interiori quadruplo longiori coriacea naviculari involuta dorso carinata obtusa, calycis tubo subincurvo acute costato pilosulo, limbi albidi tubo duplo brevioris lobis ovatis obtusis ♀. *S. oleifolia* Sm. Prodr. I, p. 213. — *S. riminea* Schrad. — *S. Smithii* Ten. Fl. Nap. tab. 223. — *S. cordata* Desf. Atl. non L.

Hab. in littoribus maritimis Graeciae in insulis Zacynthio (Marg!), Cephallenia (Heldr!), Melos (Armenis!), Cretâ (Sieb! Heldr!), Peloponneso (Bory!), Attica (Sprun! Heldr!), Macedonia (Janka!), Thracia ad Byzantium (Castl Auch. 2504!), Anatoliâ occid. ad Smyrnam (Boiss!). Syriâ littorali ad Sidonem (Bl! Gaill!).

Ar. Geogr. Regio mediterranea totius Europe et Africæ borealis.

13. S. cancellata (Bernh. in Bertol. Fl. It. 3, p. 525) cinereo-pubescentis rarins glabra, rhizomate indurato cespitoso, foliis parvis obovato-spathulatis obtusis basi longe attenuatis petiolatis uninerviis, scapis numerosis brevibus a basi articulatim ramosissimis cancellatim infractis, ramis divaricatissimis intricatis saepe refractis inferioribus sterilibus inter articulos breves strangulatis, spiculis parvis subtriangularis in spicas terminales laxas distichas dispositis, bracteâ inferiore trianguli minimâ acutâ anguste albo-marginata interiore quadruplo longiore acutiusculâ margine et apice hyalinâ, calycis tubo adpresso hirsuto, limbi tubo brevioris lobis oblongis acutiusculis ♀. *S. furfuracea* Rehb. Ic. Crit. fig. 327, non Lag. — *S. rorida* Auct. Fl. Dalm. non Fl. Græc.

z. glabra.

Hab. in maritimis Cephallenianis prope Kiniti Petra (Heldr!), in Peloponneso ad Navarin (herb. Fauché!).

β. laxa. — Glabra, folia submajora, scapi et rami elongati tenuiores fragillini articulis longioribus vix strangulatis, rami floriferi arcuato-recurvi spiculis laxissimis. — *Limonium articulatum supinum* Bocc. Ic. Sic. tab. 83 ex Melitâ hanc formam refert.

Hab. ad littora Cephallenianae ad Kipouria, Athera et Poros (Heldr!).

Specimina var. *α* glabritie exceptâ Dalmaticis simillima.

Ar. Geogr. Illyria, Dalmatia.

14. S. Cosyrensis (Guss. Prodr. Sic. Suppl. p. 90) glaberrima lăvis viridis, foliis parvis flaccidis obovato vel oblongo-spathulatis obtusis vel retusis in petiolum longe attenuatis, scapis numerosis a basi stricte ramosissimis articulatis flexuosis fragillimis, ramis steri-

libus, supremis spiciferis, spiculis subunifloris tenuibus cylindricis rectis vel subincurvis in spicas tenues unilateralis vel distichas laxè dispositis, bracteis carinatis arctè involutis anguste membranaceis acutis vel acutiusculis interiore quadruplo longiore, calycis tubo basi hirtulo, limbo tubo vix breviore lobis oblongis acutiusculis ♀. *S. cordata* Guss. Fl. Sic. Prodr. — Boiss. in D.C. Prodr. non L. — Rehb. Germ. tab. 94 optima. — Ic. Boec. Sic. tab. 34.

β. major. — Scapi crassiores elatiiores, spiculae sèpe bi et trifloræ, calycis limbus demum in lobos profunde partitus. — *S. dichotoma* Sm. Prodr. et Urv. En. non Cav.

Hab. in maritimis Græciæ prope Coron et Argos (herb. Fauché!), insulâ Melos (Urv!).

Fragilitate, spiculis tenuibus in spicas pectinatim dispositis, calycis lobis acutis a *S. virgata* formis semper distinguenda. *S. cordata* L. ex cl. Visiani in herb. Linn. inspectione est *S. pubescens* D.C.!

Ar. Geogr. Sicilia, Italia australis, Dalmatia.

15. **S. nuda** (Boiss. et Buhse Aufz. p. 184) glabra, rhizomate crasso lignoso pluricauli, foliis radicalibus cito exsiccatis congestis oblongo-spathulatis in petiolum eis longiore attenuatis, scapis tenuiter virgatis a basi dichotome ramosis ramis inferioribus sterilibus, spiculis binis vel ternis 3-4-floris in spicas terminales capituliformes confertis, bracteis fuscescentibus coriaceis pilosis late hyalino-membranaceis inferioribus binis minimis subrotundis interiore quadruplo majore ovatâ cymbiformi obtusa, paleis hyalinis obovatis flores æquantibus, calycis tubo ad costas longe piloso limbo purpurascente ad medium in lobos ovatos acutos eroso-dentatos fissò ♀. *S. Fischeri* Trautv. Stirp. Nov. v. p. 14!

Hab. in salsuginosis interioribus Armeniæ ad Alyndschæ (Radde!), Persiæ ad fontem Gendab prope Badkann jugi Elbrusensis et prope Kaschan (Buhse!). Persice *Schastarussion*.

Scapi 1-4 $\frac{1}{2}$ -pedales, capitula terminalia magn. avellanæ. Calyx 4 lineas longus. Ex affinitate *S. tenellæ* Turcz., folia majora, rami crassiores, etc. Ex sectione *Plathymenio* Boiss. in D.C. Prodr., a Sect. *Limonio* characteribus non sat firmis distinctâ.

16. **S. iatifolia** (Smith Trans. Linn. Soc. 1791) foliis indumento brevi stellato molliter velutinis amplis oblongo-ellipticis obtusis in petiolum longum attenuatis, scapis elatis subtereti-angulatis ad dichotomias squamis triangularibus acutis obsitis in paniculam amplissimam effusam abeuntibus, ramis sterilibus nullis, spiculis unifloris rarius bifloris remotiusculis in spicas laxas tenues secundas subre-curvas dispositis, bracteis binis exterioribus omnino hyalinis carinatis exteriore mucronulata, bracteæ interiore duplo majore imo tantum dorso herbaceâ hyalinâ obtusissimâ sèpe fissâ, calycis tubo obconico brevi glabro vel parce hirtulo, limbi tubo æquilongi lobis quinque triangularibus acutis ♀. Ic. Rehb. Crit. fig. 464. — *S. coriaria* Hoffm.

Hab. in campis aridis Tauriæ (M.B., Rehm!), Ciscaucasie (Stevl Hohen!).

Cl. Rehmann e campis Tauriæ sub *S. cespitosæ* nomine misit plantam characteribus inter *S. latifoliam* et *S. Caspiam* intermedium et earum prob. prolem hybridam.

Ar. Geogr. Bulgaria, Rossia australis, regio Uralensis.

17. S. Caspia (Willd. Enum. I, p. 336) glabra granulato-scabra, foliis parvis obovato vel lanceolato-spathulatis obtusis mutieis vel mucronulatis in petiolum attenuatis, scapis numerosis erectis squamis triangularibus margine membranaceis instructis fere a basi dichotome ramosissimis, ramis inferioribus sterilibus stricte capillaceo-multifidis articulatis, superioribus dense floriferis corymbosis, spiculis 2-3-floris in spicas confertas breves saepe subcapitatas distichas congestis, bracteis binis inferioribus hyalinis ovatis obtusis interiore duplo majore usque ad medium hyalinâ, calycis tubo brevi conico ad angulos ciliato, limbi nivei tubo longioris lobis 5 ovatis obtusis erosulis ♀. Ic. Rehb. Crit. fig. 335. — Ic. Germ. tab. 91. — *S. reticulata* M.B. I, p. 251. — Gmel. Sib. non L. — *S. bellidifolia* D.C. Fl. Fr. non alior.

Hab. ad littora maris, Attica (Heldr! Orph. Fl. Græc. N° 695!), Macedonia et Thracia (Griseb., Friv!), Tauria (Lebed., Rehm!).

Ar. Geogr. Gallia occidentalis, regio Mediterranea, Rossia australis, Son-garia chinensis.

18. S. Iconia (Boiss. et Heldr. in D.C. Prodr. XII, p. 661. — Diagn. Ser. I, 12, p. 92) glabra tota tuberculato-scabra, foliis minimis cito evanidis obovatis obtusis in petiolum brevem attenuatis, scapis humilibus a basi dichotome ramosissimis infracto-flexuosis squamis triangularibus late scariosis obsitis, ramis inferioribus sterilibus parce et breviter ramulosis superioribus divaricatim corymbosis, spiculis minutissimis bifloris in spicas breves secundas patulo-scropioideas dense imbricatis, bracteis externis hyalinis internâ vix innâ basi herbaceâ duplo longiore, calycis tubo brevissimo ad et inter costas piloso, limbi tubo longioris lobis ovatis obtusis ♀.

Hab. in salsuginosis planitiei ad Iconium Anatoliæ interioris (Heldr!).

Semipedalis, affinis *S. Caspiæ* et forsitan hujus varietas differt tamen bracteis fere omnino hyalinis, calycis fere tertia parte brevioris tubo toto hirsuto, ramis divaricatis.

19. S. pruinosa (L. Mant. p. 59 excl. synon.) tota tuberculis furfuraceo-puberulis pruinosa, foliis minimis cito evanidis ovato-orbiculatis basi attenuatis, scapis e rhizomate indurato erectis squamis fuscis triangularibus obsitis a basi ramosissimis infracto-flexuosis, ramis inferioribus sterilibus divaricatim ramulosis articulatis superioribus horizontaliter patentibus ramosis paniculatis, spiculis unifloris in spicas breves scorpioideas secundas imbricatis, bracteis rufescensibus albo-marginatis inferioribus ovatis acutiusculis interiore quadruplo majore convolutâ obtusissimâ, calycis tubo obconico glabro, limbo æquilongo niveo rotato obtusissime quinquelobo ♀. Ic. Viv.

Fl. Lib. Spec. tab I, fig. 27. — *S. aphylla* Forsk. Cat. pag. 60 non Poiret. — *S. tubiflora* Rœm. et Sch. p. 798 non Del.

Hab. in salsuginosis, Egyptus circa Alexandriam (Del., Ehr! Gaill! Martini!) et in deserto Arabico (Schweinf!), Arabia Petrea ad Ouadi Ghurundel (Boiss!), Palestina prope Askalon (Ky. 877!) et ad Mare Mortuum (Boiss!).

Ar. Geogr. Africa borealis.

20. **S. otolepis** (Schrenk Bull. Ac. Imp. Petrop. t. I, № 28) glabra glaucescens parce tuberculata, foliis parvis cito exsiccatis obovato-spathulatis obtusis in petiolum latum brevem attenuatis, scapis elatis teretibus jam infra medium ramosissimis, squamis ad dichotomias inferiores orbiculari-amplexicaulibus auriculatis cito abscissis et deciduis superioribus triangularibus, ramis inferioribus sterilibus tenuiter et stricte multifidis superioribus ample paniculato-corymbosis, spiculis unifloris minimis in spicas breves secundas subrecurvas imbricatis, bracteis omnino hyalinis inferioribus ovatis acutiusculis interiore triplo majore marginibus breviter connatis oblique cyathiformi calyceum includente, calycis tubo brevissimo ad nervos piloso limbi albi lobis 5 ovatis obtusis ♀.

Hab. in Turkestaniæ argillosis salsuginosis inter Buchara et Samarkand (Lehm!).

Bipedalis, folia pollicem vix excedentia. spicæ tenues 3-5 lineas longæ, calyx lineæ $\frac{1}{2}$, vix longus.

Ar. Geogr. Songaria.

21. **S. perfoliata** (C. A. Mey. et Karel in Bull. Mosq. 1839, p. 167) basi seabiuseula cæterum glabra glaucescens, foliis parvis cito evanidis oblongo-spathulatis obtusis basi attenuatis, scapis pluribus erectis elatis virgatis a basi patule ramosis et foliaceo-squamiosis in paniculam longam laxam abeuntibus, squamis caulinis et rameis foliaceis amplis amplexicaulibus orbiculatis auriculatis, summis parvis triangularibus acuminatis, ramis sterilibus paucis vel nullis, spiculis 2-3-floris in spicas breves secundas patulo-recurvatas imbricatis, bracteis binis inferioribus hyalinis ovatis carinatis acutis, interiore triplo majore a medio sursum hyalinâ obtusâ, calycis tubo ad 2 costas externas piloso, limbi tubo brevioris lobis ovatis acutiasculis ♀. *S. reniformis* Girard Ann. Sc. Nat. Ser. III, 2, p. 325.

Hab. in salsuginosis Persiae australis (Auch. 5246!), ad lacum Nemek Derja prope Schiraz (Ky. 464!), ad littus orientale maris Caspii (Kart!), inter Buchara et Samarkand (Lehm!), in Affghania (Griff!).

Bipedalis, insignis squamis caulinis diametro saepe 6-9 lineas latis, spicæ 5-6 lineas longæ.

22. **S. obovata** (Ledeb. Fl. Ross. III, p. 468) glabra, ramis crec-tis foliosis superne dichotome paniculato-corymbosis, foliis obovato-spathulatis obtusissimis mucronulatis basi cuneatis in petiolum revoluti basi subdilatatum non auriculatum al-euntibus, foliis caulinis

ad squamas parvas acutas reductis, paniculæ ramis in spicas breves secundas laxiusculas abeuntibus, floribus parvis, bracteis oblongo-lanceolatis acutis late membranaceis, calycis tubo hirtulo, limbo . . . 5.

Hab. in maritimis prov. Transeaucasie Talyisch prope Lenkoran (Haussk!). (Vid. in herb. Ledeb.)

Ex specimine juniori cuius flores non evoluti descripta, pedalis, folia cum petiolo 10-12 lineas longa 6-7 lata, panicula 4 poll. longa et lata multiflora. Species a sequentibus spiculis dissimilioris nec dense imbricatis vel capituliformibus distincta.

23. *S. suffruticosa* (L. Sp. I. p. 369) glabra glaucescens depresso-punctata suffruticosa ramosissima ramis ascendentibus brevibus foliosis, foliis carnosis oblongo-spathulatis obtusis in petiolam longe attenuatis, vaginâ petiolari ad utrumque petioli latus in auriculam oblongam hyalinam abeunte, scapis virgatis simplicibus vel dichotome ramulosis, spiculis 2-3-floris distiche et dense in spicas capituliformes secus inflorescentiæ ramulos flexuosos sessiles et remotas dispositis, bracteis ovatis carinatis obtusis rufescensibus albo-marginatis superiore duplo majore, calycis tubo glabro, limbi albi tubo brevioris profunde quinquelobi lobis oblongis obtusis 3. Ic. Gmel. Sib. 2. tab. 88. — *S. glauca* Less. in Linn. IX, p. 196.

Hab. in salsuginosis ad Caspium (C.A. Mey!), in Turcomaniâ ad Caspium (Kareli!), in Persiâ orientali et Affghanâ occid. inter Meschhed, Herat et Tebbes (Bge!).

Ar. Geogr. Rossia australis, Songaria, Turkestania borealis, Sibiria Uralensis et Altaica.

24. *S. leptophylla* (Schrenck in Bull. Phys. Ac. Pétersb. III, p. 211) glabra glaucescens depresso-punctata suffruticosa ramis brevibus erectis crebre foliosis, foliis carnosis linearis-semiteretibus acutis, vaginis apice rotundatis fere exauriculatis, scapis dichotome ramosis ad dichotomias inferiores foliosis, spiculis 2-3-floris minimis in spicas breves secundas subsessiles corymbosas imbricatis, bracteis rufescensibus anguste albo-marginatis inferioribus carinatis acutiusculatis interiorre duplo majore obovatâ obtusâ, calycis tubo glabro limbi tubo brevioris lobis ovatis 5.

Hab. in salsuginosis prope Kehris Demeh Affghaniae occid. inter Herat et Tebbes (Bge!).

S. leptophylla differt a *S. suffruticosa* et *S. carnosâ* quibus affinis foliis tenuissimis acutis linearis tertiam partem vix latis, inflorescentiâ magis corymbosâ. Specimina a cl. Bge collecta a tate nimis proiecta a Songarieis foliis abbreviatis 4-6 lineas nec pollicem longis tantum differre videntur.

Ar. Geogr. Songaria.

25. *S. carnosâ* (Boiss. in D.C. Prodr. XII, p. 663) glabra glaucescens depresso-punctata suffruticosa ramis erectis brevissimis crebre foliosis, foliis carnosis linearis-spathulatis subtus obtuse carinatis obtusiusculis longe attenuatis et linearis-triquetris, vaginâ petio-

lari anguste scariosa a petioli basi utrinque rotundata exauriculata, scapis erectis dichotome ramulosis, spiculis bifloris in spicas breves secundas secus ramos inflorescentiae sessiles et remotas imbricatis, bracteis rufescensibus anguste albo-marginatis obtusis inferioribus carinatis superiore duplo majore, calycino tubo glabro, limbi albi tubo brevioris lobis 5 rotundatis 5.

Hab. in prov. Aderb'djan Persiae prope Khoi (Auch. 25°51 Szovf).

Folia 6-9 lineas longa apice lineam lata. Affinis *S. suffruticosa* a qua præter folia angusta sat differre videtur auricularum ad petioli basin deficiencia, ramulis floriferis ab breviatis pollicem tantum longis.

26. *S. axillaris* (Forsk. Eg. Arab. p. 58), glabra glauca suffruticosa ramosa, ramis vetustioribus denudatis vaginarum reliquiis asperis, hornotinis brevibus erectis crassiuseulis crebre foliosis, foliis planis carnosis oblongo vel lanceolato-spathulatis obtusis vel acutiusculis longissime attenuatis, vaginâ petiolari fuscâ oblique truncatâ, scapis brevibus vel elongatis flexuosis parce ramosis anguste paniculatis articulatis fragillimis, spiculis bifloris spicas breviter pedunculatas vel sessiles scorpioideo-recurvatas densiusculas formantibus, floribus parvis, bracteis fuscis obtusis angustissime albo-marginatis inferioribus breviter ovatis superiore triplo majore involuta, calycis tubo recto oboconico imâ basi tantum pilosulo, limbo tubo breviore patulo obsolete angulato-lobato 5. Vahl Symb. I, p. 26, tab. 9. — *S. Bovei* Jaub. et Sp. Ill. Or. I, tab. 86.

Hab. ad littora Ægyptiaca maris Rubri in Eg. superiore (Acerbi!), ad Ouadi Lechuma (Schweinf!).

Ar. Geogr. Arab. felix (Forsk! Bové! Anders!).

27. *S. Arabica* (Jaub. et Sp. Ill. Or. I, t56, tab. 85) glabra glauca suffruticosa, ramis vetustioribus nudis hornotinis crebre foliosis, foliis planis carnosis obovato-spathulatis obtusissimis inferne longe attenuatis, vaginâ petiolari fuscâ oblique truncatâ, scapis axillaribus brevibus ramulosis in paniculam brevem abeuntibus, spiculis bifloris spicas densas breves secundas scorpioideas subsessiles formantibus, bracteis fuscis angustissime fusco-marginatis exterioribus ovalis acutis subcarinatis interiore quadruplo longiore oblongâ subcarinata involuta acutiusculâ, calycis tubo subincurvo glabriusculo, limbo tubo duplo breviore erecto lobis 5 obtusis crenulatis nervo rubro percursis 5.

Hab. in salsuginosis Arabiae Mascatensis (Auch. 5243!).

Valde affinis præcedenti, folia obtusiora interdum subretusa, panicula minus ramosa, spiculae subduplo maiores, bracteæ subcarinatae omnino fuscæ, calycis tubus subincurvus et limbis brevior non patulus.

28. *S. Stocksii* (Boiss. in D.C. Prodr. XII, p. 664) glauca suffruticosa, ramis vetustioribus denudatis, hornotinis foliosis, foliis carnosis planis glabris vel puberulis ovato-spathulatis obtusissimis

basi attenuatis, vaginâ petiolari oblique truncatâ, scapts brevissimis subangulatis fragilibus, panículâ brevi, spiculis billoris in spicas brevissimas fasciculiformes confertis, bracteis herbaceo-rubellis angustissime rubro-marginatis inferioribus ovatis subcarinatis obtusiusculis basi vaginantibus superiore triplo longiore oblongâ, calycis tubo ad costas dense et longe ciliato, iimbo patulo tubo breviore breviter et acutiusculo quinquelobo s. Hook. Ic. tab. 837.

Hab. in salsuginosis Belutschiae et prov. Scinde (Stocks).

Valde affinis *S. Arabica* a qua preter folia basi minus longe attenuata differt panicula ramis spicisque contractis, bracteis herbaceis, superiore fere non involutâ, calycis tubo longe ciliato.

S. cylindrifolia Forsk. Eg. Arab. p. 59. — Vahl Syrb. I, tab. 10 ex littoribus Arabiae Felicis, a precedentibus foliis longis teretibus linearis cylindricis distincta, eis floribus similis forsitan in littoribus Egypti superioris reperiatur.

SECT. III. SPHEROSTACHYS.

29. **S. globulifera** (Boiss. et Heldr. in p.c. Prodr. XII, p. 663) glabra glauca calcareo-punctata, foliis ad collum rosulatis coriaceis obovatis albomarginatis obtusis mucronulatis in petiolum brevem basi vaginantem attenuatis, scapis elatis subangulatis a medio dichotome paniculatis flexuosis, spiculis 2-3-floris in capitula sphaericâ basi bracteis inferioribus rufescensibus ovatis involucrata seenus paniculae ramos sessilia inferiora distantia superiore approximata densissime congestis, bracteâ superiore cæteris duplo majore ovato-orbiculari herbacea subcarinata late hyalinâ obtusa, paleis numerosis, calycis tubo brevi obconico toto dense albo-hirsuto, limbi tubo dimidio brevioris lobis brevibus subrotundis plicatis ♀. Ic. Tehih. As. Min. tab. 38.

Hab. in salsis hyeme inundatis planitiei Koniah Isauriae inter urbem et montem Karadagh (Heldr.). Fl. Jun.

Foliorum lamina $1\frac{1}{2}$ -pollicaris 8-9 lineas lata, scapi pedales, florum globuli magnitudine pisi in panicula subsecunda fere 15-20, flores minuti, calyx linea vix longior.

SECT. IV. SCHIZOPETALUM.

30. **S. macrorrhabdos** (Boiss. Diagn. Ser. II, 4, p. 67) perennis glabra glauca calcareo-punctata, foliis radicalibus rosulatis obovatis obtusis mucronulatis in petiolum brevem vaginantem attenuatis, scapo crasso valde elato squamis triangulari-ovatis acutis obsito a media parte stricte ramoso ramis elongatis virgatis paniculatis, spiculis 3-5-floris secus ramos paniculae valde distantibus, bracteis coriaceis rubellis ovatis late albo-marginatis exterioribus mucronulatis interiore sublongiore submuticâ, paleis truncatis præter nervum medium

rubrum albo-hyalinis, calycis basi obliqui et sub lente puberuli tubo tenui in limbum albo-membranaceum truncato-erosum dilatato φ .

Hab. in Aſſghania (Griff!), in Belutschia superiore (Stocks!).

Folia cum petiolo 2-3-pollicaria, scapus cum paniculæ ramis 8-12 pollicaribus 2-3-pedalis, calyx 5 lineas longus. Species foliis *Limoni* et inflorescentiæ *Acantholimonis* insignis, calyx ut in *Amheria* basi subcalcaratus, stigma oblongum *Statices*; a congeneribus petalis profunde bipartitis nec integris vel retusis distincta.

SECT. V. SCHIZYMENTUM.

31. **S. echioïdes** (L. Sp. 394) annua glabra, foliis paucis parvis obovatis vel oblongo-spathulatis obtusis in petiolum brevem attenuatis superne saepe tuberculatis, scapis tenuiter virgatis flexuosis dichotome paniculato-ramosis flexuosis, ramis elongatis patentibus fragilibus, spiculis unifloris rarius bifloris curvulis secus paniculæ ramos unilateralibus patentibus dissitis, bracteis inferioribus ovato-rotundis obtusis minimis, superiore quintuplo longiore coriacea involuta scabri angustissime membranacea, calycis tubo anguste angulato-cylindrico adpresse hirtulo, limbi tubo triplo brevioris truncati tandem lacero-lobati aristis 3 rubris truncato-recurvis ⊖. Ic. Rehb. Ic. Crit. 292 et Ic. Germ. 76. — Flora Græca, t. 299. — *S. aristata* Sm. Prodri. I, p. 213.

Hab. in maritimis et salsuginosis interioribus, Aegina (Sart!), Attica ad Phalerum et ad radices Hymetti (Heldr!), Eubea (Sprun!). Creta (Heldr!), Cyprus (Silb. Ky. 536!).

Variat interdum prolifera spicis e dichomiâ unaquaque inferiori pluribus

Ar. Geogr. Regio Mediterranea, Europæ et Africæ borealis.

32. **S. Owerini**, perennis glabriuscula glaucescens, foliis omnibus ad rhizomatis parce ramosi squamosi capita rosulatis minutissimis squamiformibus carnosulis sessilibus ovatis subtus carinatis margine denticulatis in cuspidem longisculam abeuntibus, scapis tenuibus erectis aphyllis inferne squamis paucis semiamplexicanilibus cuspidatis obsitis tere a basi pluries dichotomis, ramis filiformibus rigidulis divergentibus fragilibus, spiculis secus ramos valde dissitis subunilateralibus rectis unifloris vel inferioribus bifloris, bracteis inferioribus minimis trianguli-ovalis mucronatis superiore multo majore oblonga herbacea carinata obtusa mucronulata marginé membranacea, calycis tubo angustissimo striato glabriusculo, limbo tubo duplo breviore et vix latiore albo-membranaceo decemlobo lobis 3 trianguli-lanceolatis nervo excurrente aristulatis, 3 alternis enerviis obtusis quadruplo brevioribus, petalis roseis φ

Hab. in ditione Gumbet Caucasi Orientalis supra Tehirkal alt 4500' (Ower!).

Species ex unico specimine descripta sed characteribus insignis facie *S. echioïdem* referens. Folia sessquilinea vix longa, caules cum ramis 6-9-pollicares tenuissimi, spiculae intervallis carum longitudine longioribus dissitæ. Calyx tenuissimus tres lineas longus.

33. **S. Cabulica** (Boiss. in D.C. Prodr. XII, p. 666) annua? glabra glauca calcareo-pulverulenta, foliis crebris in rosulam densissime congestis cuneato-spathulatis obtusis mucronatis basi subsessili attenuatis 5-7-nerviis, scapis virgatis teretibus elatis dichotome et stricte ramosis ramis elongatis paniculatis, spiculis 1-2-floris secus ramos distantibus rachidiique adpressis laxissime spicatis, bracteâ inferiore coriaceâ triangulare mucronatâ, interiore triplo longiore involutâ anguste membranacea dorso excurrente aristulata, calycis tubo tenui cylindrico adpresso puberulo, limbo tubo breviore demum in 5 lacinias lineares patulo-recurvatas præter nervum validum membranaceas ad basin partito ⊖?

Hab. in regno Cabulico ad Karabagh et Mouklour (Griff!).

Folia eis *S. echooidis* numerosiora et majora pollicaria, scapi cum ramis $\frac{1}{2}$ -1-pedalibus 2-3-pedales, inflorescentia laxa *S. macrorrhizos*, calyx 5 lineas longus. Hæc species ex radice simplici verticali annua esse videtur.

SECT. VI. SIPHONANTHA.

34. **S. tubiflora** (Del. Fl. Eg. p. 251, tab. 25, fig. 2) glauca calcareo-tuberculata, rhizomate indurato breviter ramoso, foliis radiocalibus parvis ovato-rhombeis obtusis in petiolum subbreviore attenuatis, scapis humilibus flexuosis articulatis ramulosis fragillimis ramis inferioribus sterilibus simplicibus rigidis, corymbo terminali parvo, spiculis biploris elongatis in spicas breves secundas scorpioides fragillimas distiche congestis, bracteis subcoriaceis anguste membranaceis carinatis obtusis exteriore mucronata interiore quadruplo longiore longe oblongâ interdum subrecurvata, calycis tubo tenui glabro, limbo æquilongo in lacinias lineares bifidas inter cuspides nervo rufo excurrente aristatas ad basin partito ♀. *S. squamata* Poir. Suppl. 5, p. 237.

Hab. in arenosis Egypti prope Alexandriam (Del., Martins! Ehr! Samarit!).

Folia 4-5 lineas longa, scapi 3-4-pollicares, flores sex lineas longi, corolla ampla purpurea.

SECT. VII. PSYLLIOSTACHYS.

35. **S. spicata** (Willd. Sp. I, p. 1533) annua, foliis rosulatis glabris vel subtus hirtulis herbaceis oblongo-lanceolatis in petiolum brevem longe attenuatis obtusis mucronatis vel integris vel saepius in lacinias triangulare-oblongas saepe reversas versus folii basin diminutas basi saepe confluentes runcinato-pinnatipartitis, scapis folia subsuperantibus spicam terminalem vel saepius plures sessiles cylindricas densissimas infimas remotiusculas superiores ad terminalis basin congestas ferentibus, spiculis 2-4-floris, bracteis rufescenti-herbaceis dorso papilloso-hirtis inferiore oblongo-spathulata

incurvâ mucronatâ superiore latiore et subbreviore basi attenuatâ apice truncatâ anguste membranaceâ, calycis tubo a basi ad diuidiam partem vel ad $\frac{2}{3}$ pilis glandulosis hispido, limbi tubo æquilongi eroso-dentati nervis in aristas longas excurrentibus, corollæ roseæ vel albae lobis ovatis \odot . Gmel. Sib. tab. 91. — *S. lyrata* M. B. Casp. N° 38. — *S. sisymbriifolia* et *S. plantaginiflora* J. et Sp. Ill. tab. 87 et 88.

Hab. in salsis maritimis et interioribus Cis- et Transcaucasicæ ad Caspium (C.A. Meyt Hoht), Persiaë bor. (Steph!) et ad lacum Urmiah (Auch. 5244!), Persiaë australis ad Dalechi (Ky. 493!), Affghaniae (Griff!), Belutschiae (Stocksl.), Assyriaë inter Mossul et Bagdad (Oliv!) et prope Molsammera (Noë 932!), deserti Sindjar Mesopotamiae (Haussk!), Palestinaë ad mare Mortuum (Ky. 577! Schlumb!), Turcomanie (Kar!).

Ex speciminum copiâ note leviores quibus freti cl. Jaub. et Spach species binas distinxerunt evanuerunt.

Ar. Geogr. Sibiria Uralensis.

36. *S. leptostachya* (Boiss. Diagn. Ser. I, 7, p. 68) annua glabra glaucescens, foliis radicalibus breviter petiolatis linearibus in lacinias anguste lineares integras deflexas pectinatum et regulariter pinnatipartitis, scapis filiformibus foliis brevioribus in spicas simpli-cissimas longe et tenuiter cylindricas basi subinterruptas cæterum densifloras abeuntibus, spiculis 2-3-floris fasciculos inferius sub-distantes superius approximatos formantibus, bracteis glabris fuscescentibus inferiore minutissimâ triangulari acutâ, inferiore triplo maiore subcoriacea concavâ apice truncatâ incrassatâ et bigibbâ, calycis tubo adpresso hirtulo limbo in dentes lanceolatos acutos profunde parti-to \odot .

Hab. in salsis ad radices montis Sabst Buschom prope Schiras Persiaë australis (Ky. 367!).

Folia 1-1 $\frac{1}{2}$ -pollicaria 5-6 lineas lata, spicæ tenues 4-6-pollicares diametro 2 $\frac{1}{2}$ lineas latae, flores eis *S. spicate* dimidio minores a quâ spicis tenuissimis, bractearum glabrarum et limbi calycini structurâ eximie differt.

ARMERIA (Willd. Enum. p. 333).

Calyx infundibuliformis tubo inferne decemcostato super pedicel-lum apice subdilatum oblique inserto et basi calcarato, limbo se-rioso plicato quinque nervi quinquelobo. Petala quinque basi tantum in annulum brevissime coherentia. Filamenta sublibera. Styli quinque imâ basi concreti parte tertiatâ inferiore barbati. Stigmata filiformi-cylindrica tota superficie glandulosa. Utriculus membranaceus sub-indehisceens superne pentagonus et apice quinquegibbus basi irregu-lariter ruptilis. — Herbæ perennes interdum basi sufrutescentes, foliis linearibus, spiculis unibracteatis in capitulum hemisphericum terminale confertis.

1. A. Majellensis (Boiss. in D.C. XII, p. 685) cespitosa glabra, foliis biformibus externis brevibus patentibus late linearibus trinerviis saepe rigidulis, internis angustis elongatis, scapis saepius elongatis, capitulis majusculis, involueri pallide brunnei phyllis externis ovatis breviter cuspidiatis vel muticis interna obtusa subaequantibus, spiculis sessilibus, calycis tubo ad costas piloso calcare tubo quintuplo breviore foveolâ ovatâ vel ovato-oblongâ, limbi lobis breviter triangularibus in aristas longas attenuatis, corollâ purpureâ vel roseâ ♀.

Hab. in pascuis et rupestribus regionis subalpinæ et alpineæ, mons Olenos Peloponnesi (Heldr!), Parnassus (Guicci), Cytheron (Heldr. herb. Norm. 5061), Veluchi Etoliae (Samar!), Peristeri Macedoniae (Orph!).

β. brachyphylla. — Folia omnia abbreviata, interiora angusta pauca interdum nulla. — *A. Orphanidis* Boiss. Diagn. Ser. II, 4, p. 71.

Hab. cum tipo in regione alpine Græcia in eum saepe transiens, Chelmos et Malevo Peloponnesi (Orph!), Delphi Eubœa (Sarti Heldr!), Velnchi Etoliae (Samar!), in rupibus Athonis inter Lavra et Iviron (Janka!).

γ. leucantha. — Folia exteriora saepe sinnuato-undulata, interiora interdum angustissima, corolla ampla alba. — *Statice alliacea* Fl. Græc. tab. 294 non Cav. (foliis externis deficientibus). — *S. undulata* Bory et Chaub. Fl. Pelop. № 519, tab. 11 (optima). — *A. argyrocephala* Wallr.

Hab. in Taygeti regione alpine et subalpine (Heldr!), montibus Arcadiæ (Bory!), montibus Parnæs et Hymetto Atticae (Sibth., Sprn! Boiss! Heldr. Orph. Fl. exs. 696!) forma e locis calidioribus foliis interioribus saepe tenuioribus et capitulo minore), monte Onion Bœotiae (Heldr!), in Cariæ (Auch. 2495! mixta cum *A. Cariensis*), Libano in monte Sannin (Ehr!), supra Hadet (Bl!), in monte Hermone Antilibani supra Ainhata (Gail!).

Species ut plures e congeneribus valde polymorpha inter eas foliis primariis dissimilibus donatas *A. Nebrodensi* Guiss. magis affinis et foliis latioribus manifeste trinerviis nec uninerviis vel obsolete trinerviis discedens sed forsitan ad eam formis brachyphyllis (in Italiæ montibus quoque obviis) transiens. Varietas leucantha in Aprutio Neapolitano quoque observatur. — *A. canescens* Boiss. loc. cit. p. 686 (nomen specif. cl. Host improprium nam cl. Visiani observante planta glabra est et viridis) e Dalmatia et Montenegro, ab *A. Majellensi* non specifice differt.

Ar. Geogr. Italia australis, Dalmatia.

2. A. Cariensis (Boiss. in D.C. Prodr. XII, p. 677) cespitosa glabra, foliis biformibus uninerviis primariis linearibus seapo subfataribus, cæteris linearis-setaceis subcanaliculatis, scapis tenuibus, involueri pallide scariosi phyllis exterius triangulari-ovatis acutis interioribus sensim elongatis membranaceo-marginatis retusis dorso excurrente mucronulatis vel muticis, spiculis intra involuerum stipitatis, calycis tubo ad costas primarias pitoso cæterum glabro, calcare tubi dimidiâ vel tertiam partem æquante, foveolâ basilari linearis, limbi tubo longioris lobis oblongo-triangularibus in aristas longas attenuatis ♀.

Hab. in herbidis alpinis montis Cadmi Cariæ supra Gheyra (Boiss!), Timoli ad Bozdaghi (Bal. 231!), Phrygiae a Kaiageul Decè supra Ouchak (Bal!).

β. Rumelica. — Folia paulo latiora. — *A. Rumelica* Boiss. loc. cit. p. 677.

Hab. in regione superiori montis Korthiati Macedoniae (Heldr! Orph!), Thraciae montibus (Frix. exs. sub *A. Alpinâ!* et *A. scorzonericfoliâ!*), in monte Balkan supra Kalofer (Janka!).

γ. Thessala. — Folia linearia exteriora parum dissimilia, tubi calycini foveola paulo brevior oblongo-linearis. — *A. Thessala* Boiss. Diagn. Ser. II, 4, p. 70.

Hab. in pascuis alpinis Olympi Thessali (Heldr!).

Facies *A. Majellensis* sed folia uninervia et spiculae intra involucrum stipitatae; ab ea insuper calycis calcare elongato et foveolâ basilarî linearî remota et sectioni *Macrocentron* Boiss. potius adnumeranda.

LIMONIASTRUM (Mench Meth. 423.)

Calyx tubulosus membranaceus eostatus limbo minimo acute quinquelobo. Corolla infundibuliformis usque ad faucem gamopetala tubo longo tenui, limbi lobis obovatis. Filamenta usque ad faucem corollae adnata. Styli ad medium usque connati glabri. Stigmata filiformi-cylindrica. Utriculus membranaceus indehisens. Albumen parecum. — Suffrutices foliis carnosis, spiculae laxe spicatae rachidis latere complanato adpresso tribracteatae.

1. L. monopetalum (L Spec. I, p. 296 sub *Statice*) fruticosum glaucum calcareo-leprosum ramis ramulosissimis ramiulis crebre foliosis, foliis carnosis planis oblongo vel lanceolato-spathulatis obtusis in petiolam basi vaginante attenuatis, spieis strictis breviter particulatis, rachide flexuosa articulata fragillima, spiculis bifloris rachidi excavatae arete adpresso, bractea inferiore cyathiformi oblique truncata vaginante spiculam et rachidem amplectente, intermedia linea tri-triquetrâ apice subuliformi exerto intima longiore coriacea flores involvente, calycis tuba longo tenui membranaceo limbi minutis lobis acutis, corolle roseo tubo tenui exerto limbi magni subrotati partitionibus ovatis retusis. *Limoniastrum monopetalum* Boiss. in DC Prodr. XII, p. 689. — Ic. Bocc. Sic. tab. 47. -- Bot. Reg. 1842, tab. 59.

Hab. in maritimis Egypti ad Alexandriam et Port Said (Acerbi! Auch! Ehr!).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Sardinia, Sicilia, Italia meridionalis, Africa borealis.

TR. H. PLUMBAGEE Spach.

Calyx herbaceus vel glumaceus gamosepalus vel pentaphyllius. Styli ad apicem usque in unicum connati. Pericarpium a basi subalveum dehisens.

PLUMBAGO (L. Gen. 213).

Calyx gamosepalus herbaceus tubulosus stipitato-glandulosus. Corolla hypocraterimorpha tubo tenui limbo rotato quinquepartito. Filamenta libera basi in squamulas concaviusculas dilatata. Ovarium oblongum stylo filiforme. Stigmata quina filiformia introrsum glandulosa. Utriculus membranaceus basi circumscisse ruptus demum a basi ad medium valvatum fissus. — Herbae perennes vel suffrutices scandentes floribus subsessilibus spicatis, singulo tribracteato.

1. P. Europæa (L. Sp. I, p. 215) caulis elatis superne ramosis, ramis angulatis strictis gracilibus elongatis, foliis inferioribus obovatâ infimis in petiolum attenuatis, intermediis basi attenuatâ auriculato-amplexicanibus, superioribus et rameis diminitis anguste lanceolatis et linearibus minute et acutiuscule auriculatis, floralibus in spicas terminales prius subcapitatas dispositis, bracteis oblongis acutis glabris vel glandulosis dimidium calyceum non aquantibus inferiore sublongiore, calycis cylindrici glandulis stipitalis ad costas saepe biseriatis obsiti dectibus brevibus rectis, corollæ lilacine vel roseæ tubo calyce sesqui vel subduplo longiore limbi lobis obovatis obtusis ♀. Ie. Fl. Græc. tab. 191. — Rehb. Ie. Germ. tab. 87. — *P. lapathifolia* Willd. Enum. I, p. 198 et M.B. I, p. 444 (nullo modo diversa).

Hab. in siecis regionis inferioris et montanæ totius ditionis a Græciâ! Macedonia! Thraciâ! ad Anatoliam omnem! Syriam littoralem (Gall!) et interiorem (Ehr!), Palestinam! Kurdistanum! Armeniam! Transcaucasiom (Lede!, Hoht!). Persiam borealem (Szov! Buhse!), prov. Khorassan (Bge!).

Ar. Geogr. Europa australis tota a Lusitanâ ad Dalmatiâ, Africa borealis.

2. P. Zeylanica (L. Sp. I, p. 215) basi suffrutescens, caulis subscandentibus angulato-striatis flexuosis ramosissimis, foliis ovatis vel oblongis acutiusculis vel aenatis basi breviter et abrupte in petiolum brevem basi saepe minute auriculato-dilatatum abeuntibus, superioribus diminutis angustatis, floribus in spicas terminales densiusculas dispositis, rachide glandulosâ, bracteis oblongis acuminatis calyce triplo brevioribus inferiore aliis majore, calyce longe cylindrico totâ longitudine ad costas plurifariam glandulis longe stipitalis inaequalibus obsito apice uncinulatim et breviter quinquedentato post anthesin refracto, corollæ tubo calyce duplo longiore, limbi partitionibus emarginatis rebus mucronulatis 3.

β. glaucescens Boiss. in n.c. Prodr. p. 693. — Folia glaucescentia.

Hab. in Belutschia (Stocksl!), regno Cabulico (Lindl.).

Species subtropica vix ditionis nostræ, varietas x sie o a t (ο glaucescentia tautum differre videtur atque in Abyssinia (Schimp')). ditione teneatur quaque occurrit.

Ar. Geogr. India orientalis, Africa et Nova Hollandia tropicæ.

VOGELIA (Lam. Ill. tab. 149).

Calyx profunde quinquepartitus phyllis costâ latâ rufescente percursis, costarum marginibus inter se in tubum angustissimum cohærentibus, membranâ ad utrumque costarum latus in alas hyalinæ transverse undulatas verticales complicato-duplicatas et calycem gamosepalum pentapterum simulantes expansâ. Corolla infundibuliformis tubo longo tenui limbo quinquelobo. Filamenta libera. Ovarium fusiforme stylo filiformi. Stigmata filiformia introrsum glandulifera. Utriculus in valvas subfissilis. — Suffrutes ramosi facie *Plumbaginis*, flores breviter pedicellati spicati tribracteati.

V. Indica (Wight Ic. tab. 1075) fruticosa elata ramis teretibus virgatis elongatis, foliis sessilibus subcoriaceis sabintegris subtus prominule venosis magnis obovatis et ovato-orbiculatis auriculato-amplexicaulibus obtusis vel retusis mucronulatis, spicis gracillimis elongatis subsecundis, bracteis lanceolato-linearibus subinæqualibus pedicellum subæquantibus, calycis phyllis late lanceolatis transverse undulatim bullatis, corollæ lobis acutis ♂. *V. Arabica* Boiss. in D.C. Prodr. XII, p. 696.

Hab. in regno Mascate Arabiæ (Auch. 5285).

Frutex 6-10-pedalis, folia inferiora diam. 4-1½-pollicem lata, spicæ demum semipedales, calyx 5 lineas longus. Icon Wightiana ex ipso auctore valde imperfecta (quoad folia nimis lanceolata et acuta, minime reticulata et corollæ lobos truncato-retusos cum mucrone erroneè exhibitos.

Ar. Geogr. India Orientalis.

ORD. XCVI. PLANTAGINEÆ.

(Juss. Gen. p. 39.)

Flores hermaphroditi rarius dioici. Calyx quadripartitus persistens. Corolla monopetala hypogyna quadrifida regularis scariosa. Stamina 4 corollæ tubo inserta et ejus lacinia alterna, filamenta filiformia testivatione induplicatea, antheræ biloculares versatiles. Ovarium liberum 1-4-loculare loculis 1-8-ovulatis, stylus filiformis erectus longitrorsum bifariam stigmatosus, ovula peltata medio loculo vel in loculis uniovulatis basi affixa. Fructus capsularis rarius nucularis. Semina albuminosa, albumine corneo. Embryo rectus in centro albuminis.

PLANTAGO (L. Gen. 142).

Flores hermaphroditi spicati vel capitati, singuli sessiles bracteâ suffulti. Calycis laciniae conformes vel inæquales anticæ rarius in

unam coalitæ. Corolla tubulosa limbo quadripartito tandem refracto Stamina imo tubo inserta filamentis flaccidis. Capsula membranacea pyxidata versus basin circumcisso dehiscens placentam liberam 2-4-alatam includens eoque 2-4-loocularis. — Herbæ vel suffrutices foliis alternis vel oppositis.

Specierum Orientalium distributio.

SECT. I. EUPLANTAGO. — Acaules foliis omnibus radicalibus, rarius (conf. *P. amplexicaulem, cylindricam*) breviter caulescentes et tunc folia alterna!

§ 1. **Majores.** — Capsulæ bilocularis loculi 4-8-spermi. Semina angulata. Corolla glabra. — Perennis.

P. major.

§ 2. **Tenuifloræ.** — Capsulæ bilocularis loculi 4-8-spermi. Semina fusiformia. Corolla glabra. — Annuæ.

P. tenuiflora.

§ 3. **Mediae.** — Capsulæ bilocularis loculi dispermi sed saepius abortu monospermi. Semina facie exteriori subconvexa in'teriori planiuscula foveolâ notata. Corolla glabra. — Perennes.

P. media, gentianoides, Griffithii.

§ 4. **Cymbæformes.** — Capsula trilocularis loculis monospermis altero rarissime dispermo. Semina facie interiore sulcata vel cymbæformia.

* Corolla glabra.

+ Perennes.

P. montana, saxutilis, lanceolata, albicans.

++ Annuo-perennantes.

P. cylindrica, Stockssii.

+++ Annuæ.

P. amplexicaulis, Löflingii, minuta, Bellardi, Cretica, ovata, Haussknechtii, notata.

** Corollæ tubus glaber, lobi hirsuti. Annuæ.

P. lagopus, lagocephala, ciliata.

• • Corollæ tubus hirsutus lobi glabri. Annuæ.

P. lachnantha, evacina.

§ 5. **Biconvexæ.** — Capsula bilocularis loculis interdum bilocellatis. Semina plano-biconvexa. Corollæ tubus pilosus, lobi glabri.

* Capsula ob placentas 3-4-alatas 3-4-locellata 3-4-sperma.

P. Coronopus, crypsoides.

** Capsula placentis bialatis bilocularis disperma.

P. Serraria, maritima, carinata.

SECT. II. **PSYLLIUM.** — Caulis foliosus, folia opposita! Corolla glabra tubo saepius transverse rugoso. Semina cymbiformia.

* Perennes.

P. Arabica, Euphratica.

** Annuæ.

P. Psyllium, stricta, pumila, phæostoma, arenaria, squarrosa.

SECT. I. EUPLANTAGO.

§ 1. Majores.

1. **P. major** (L. Sp. 163) perennis acaulis, foliis petiolatis ovatis vel ovato-oblongis 3-7-nerviis integris vel irregulariter repandis aut glabris vel sparsim puberulis, scapis ascendentibus foliis æquilongis, spicæ linearι-cylindricæ elongatæ, bracteis ovatis concavis carinatis margine membranaceis calyci æquilongis, calycis laciniis aequalibus ovatis obtusis, corollæ lobis ovatis obtusiusculis, capsule ovata loculis 4-8-spermis, seminibus minimis angulatis interne planiusculis ȝ. Ic. Fl. Dan. tab. 461. — Rehb. Germ. t. 77.

Hab. in cultis humidis, ad vias totius ditionis a Græciâ! et Thraciâ! ad prov. Caucasicas! Persiam! Babyloniam! Arabiam petream (Schimp!).

P. minor. — Folia breviter et saepè vix petiolata minora saepius dentata, spicæ saepè abbreviata. — *P. intermedia* Gilib. Fl. Eur. I, p. 125. — *P. minima* D.C. Fl. Fr. 2297.

Hab. in Messeniâ (herb. Fauché!), in moute Delphi Enbeæ (Heldr!), Ponto Lazico ad Rhizé (Bal!), Affghaniâ (Griff.), Belutschia (Stocks!).

Ar. Geogr. Europa tota, Sibiria et inde in orbem sece totum diffusa.

§ 2. Tenuifloræ.

2. **P. tenuiflora** (W. K. Pl. rar. Hung. I, p. 37, tab. 39) annua acaulis pumila breviter hirtula vel glabrescens, foliis anguste linearibus integris carnosulis utrinque attenuatis acutis, scapis teretibus foliis subaequilongis, spicis erectis cylindricis elongatis laxiusculis, bracteis lanceolatis anguste membranaceis calyceem aequantibus vel (inferioribus) eo longioribus, calycis glabri laciniis ovatis dorso viridi carinatis cæterum late membranaceis, capsula elongato-conicâ calyce duplo longiore 8-16-spermâ, seminibus minimis fusiformibus ♂. Rehb. Germ. tab. 84.

Hab. in salsuginosis hyeme inundatis, Thracia inter Philippopolim et Stanimak (Janka!), Tauria (Stev.), Turkestanie deserta Karakum (Lehm!).

Ar. Geogr. Hungaria, Rossia australis, Lithuania, Sibiria altaica et insula Elaud Suecia si *P. minor* Fries huc recte adducitur.

§ 3. Mediæ.

3. **P. media** (L. Sp. 163) perennis acaulis, foliis pube septatâ breviter hirtis ovalis vel ellipticis expansis in petiolum brevem latum contractis nervis 7-9 convergentibus, scapis striatulis ascendentibus folia valde superantibus, spicâ oblongo-cylindricâ densâ, bracteis ovatis obtusis late scariosis, calycis laciniis ovatis obtusis bracteâ sublongioribus, corollæ argenteæ nitentis lobis lanceolatis acutiusculis, filamentis longe exsertis, capsulae ovatae loculis monospermis rarius dispermis, seminibus oblongis extus convexulis intus plano-concaviusculis ♀. Ic. Fl. Dan. tab. 551.

Hab. in pratis alpinis Macedoniae et Thraciae (Griseb.), circa Byzantium (ex Sibth.), Tauria (Stev.), Caucaso et regione subalpina et alpina Transcaucasie (Ledeb.), Persiae bor. prov. Aderbidjan (Szov. in herb. Petrop.).

Ar. Geogr. Europa borealis et media, in australioribus rarior vel nulla, Sibiria Altaica et Baikalensis.

4. **P. gentianoides** (Smith in Rees Cycl. V, p. 27) perennis acaulis glabra, foliis rosulatis 3-5-nerviis ovalis vel oblongo-spathulatis repandis vel dentatis breviter petiolatis, scapis teretibus ascendentibus folia triplo longioribus cum spicæ rachide glabris, spicis ovato-oblongis vel cylindraceis densiusculis, bracteis ovatis calyce subbrevioribus, calycis laciniis ovatis obtusis, corollæ fusce lobis breviter triangulari-lanceolatis, capsulae ovatae locuiis mono rarius dispermis, seminibus oblongis triconvexis illine planiusculis ♀. *P. uliginosa* Baumg. Fl. Transyl. I, p. 89.

Hab. in pratis humidis regionis abietinæ et superioris Olympi Bithyni (Sibth! Clem! Boiss! Pichl!).

β. Scardica. Griseb. Spie. 2, p. 303. — Minor. Spica abbreviata capituliformis.

Hab. in regione alpinâ Scardi Macedoniae 5200'-6500' (Griseb!).

Foliorum majorum lamina 12-15 lineas longa 6-9 lata, scapi 3-5-pollicares, spicæ 6-12 lineas longæ.

Ar. Geogr. Alpes Transylvaniae et Banatūs.

5. P1. Griffithii (Decaisne in D.C. Prodr. XIII, p. 700) perennis acaulis parce hirtula, foliis subcoriaceis ellipticis obtusis integris vel repandis in petiolum brevissimum latum contractis, scapis ascendentibus crassiusculis folia subsuperantibus cum spicæ rachide hirtis, spicis laxifloris elongatis tenuibus, bracteis breviter ovato-triangularris calyce brevioribus subcarinatis, calycis laciniis ovatis æqualibus late membranaceis, corollæ lobis ovatis acutis, capsulæ globosæ loculis monospermis, seminibus ovato-oblongis altero latere convexis altero plano-concaviusculis ♀.

Hab. ad Sir i Chuschma prope Cabul Affghaniae (Griff.).

Folia bipinnicaria, spicæ 2-4-pollicares, facies formarum minorum *P. majoris*. A *P. gentianoides* scapis rachideque hirtis nec glaberrimis, floribus laxis distincta. Specimina (juniora) ex montibus Kellal et Ssebekuh Persiae (Haussk!) ob rachidem hirtam hue spectare videntur.

§ 4. Cymbæformes.

* Corolla glabra. -

+ Perennes.

6. P. montana (Lam. Ill. 1670) perennis acaulis, foliis sessilibus breviter linear-lanceolatis integris vel remote denticulatis glabris vel villosulis 3-5-nerviis, scapis teretibus ascendentibus folio duplo longioribus, spicis rotundato-ovatis, bracteis late obovatis præter nervum virentem scariosis fuscescensibus obtusis obtuse et breviter mucronatis apice ciliatulis calyce longioribus, calycis laciniis membranaceis planis obtusis apice barbatis, corollæ lobis lanceolatis acutis, capsulæ oblongæ calycem superantis loculis monospermis, seminibus linear-oblongis hinc convexis illine concaviusculis ♀. Ic. Rchb. Germ. tab. 83. — *P. Alpina* Vill. non L.

Hab. in regione superiori montis Velugo Ætoliae (Nicolaidis!).

Ar. Geogr. Pyrenei, Alpes, Jurassus, Aprutium, montes regionis Danubialis.

7. P. saxatilis (M.B. Taur. Cauc. I, p. 109) perennis acaulis, foliis sessilibus lanceolatis acuminatis sæpius integris adpresso hirtis vel sericeis, scapis teretibus ascendentibus folia sæpius superantibus, spicis ovatis, bracteis fere omnino scariosis obovato-orbiculatis ciliatis calycem superantibus subcarinatis obtuse mucronatis, calycis scariosi laciniis ovatis acutis glabris vel apice barbatis, corollæ lobis ovatis acutis, capsulæ oblongæ calyce longiore, seminibus subrugosis ♀. *P. montana* Griseb. Spic. non Lam. — *P. montana* var. *Olympica* Boiss. exs. — *P. montana* β. *Caucasica* Dec. in D.C. Prodr.

Hab. in pratis regionis alpinæ, Macedonia in Seardo (Griseb!), Olympus Bithynus (Boiss!), mons Berydaghi Cataoniae (Haussk!), Armenia Turcica (Huet!) et Rossica (Buhse!), Caucasus omnis 2000'-8000' (M.B., C.A. Mey! etc.), Georgia Caucasicæ (Hoh!), montes Persiae bor. (Ky. 466!) et australis in alpibus Nur, Kellal, Ssebsseküh 12-13000' (Haussk!), montes Schahu et Avroman 9-10000' (Haussk!).

P. angustifolia. — Folia anguste linearia. — *P. Karakabanica* Boiss. et Huet MSS.

Hab. in montibus Armeniæ ad Erzerum (Auch. 5247! Huet!), in alpe Kuh Daëna Persiae austr. (Ky. 703!).

Species valde affinis *P. montanae* cuius est forsitan forma Orientalis; differt indumento magis sericeo, foliis plerumque longioribus et angustioribus, bracteis carinatis, calycis laciniis acutioribus.

8. ***P. lanceolata*** (L. Sp. 164) acaulis perennis interdum anno priori florens glabriuscula vel pilosa polymorpha, foliis oblongo-lanceolatis vel lanceolatis acutis basi in petiolum saepè brevem attenuatis 3-5-nerviis, scapis angulato-sulcatis ascendentibus et erectis folia superantibus, spicis ovato-oblongis vel cylindricis, bracteis ovatis acutis vel acuminatis glabris, calycis laciniis anticus glabris saepius in unicum binervem coalitis, posticis carinatæ saepius pilosulis, corollæ lobis ovatis acuminatis, capsulæ oblongæ loculis monospermis, semimibus oblongis laevibus hinc convexis illinc planiusculis ♀. Ic. Engl. hot. tab. 507.

Species valde polymorpha cuius formas vulgationes enumero.

α. genuina. — Scapi et folia elongata, ea glabra vel pilosa plus minus late lanceolata et longe petiolata.

Hab. in pratis et siccis totius ditionis a regione inferiori ad subalpinam!

β. attissima. — Procera saepius glabra, folia dilatata, spica elongata, *P. attissima* L. Sp. 164.

Hab. in humidis regionis praesertim inferioris, Græcia (Sprun!), prov. Caucasæ (Ledeb.), Affghania (Griffi) et alibi.

γ. eriophylla Decaisne. — Humilior, folia breviter petiolata vel subsessilia angusta plus minus hirsuta vel lanata, spicæ abbreviatæ ovatæ. — *P. lanuginosa* D.C. — *P. lanata* Portensch. — *P. intermedia* C. Koch Linn. XII, p. 711. — *P. Kurdica* Vatke Verh. Brand. 1874, p. 52.

Hab. in pascuis siccis montanis, Græcia (Orph!), Caucasus (Rehm!), Kurdistania (Haussk!), Persia borealis (Bge!) et australis (Haussk!).

δ. capitata Presl. — Facies formæ praecedentis sed spica globosa. — *P. capitata* Ten. — *P. ambigua* Guss. — *P. Byzantina* C. Koch Linn. XXI, p. 712.

Hab. in pascuis siccis praesertim montanis et alpinis, montes Græciae (Heldr! Orph!), Bithynia (Clem!), Libanus et Antilibanus (Boiss!), Persia (Buhse!).

Ar. Geogr. Europa tota, Africa borealis. In temperatis totius orbis nunc introducta.

9. **P. albicans** (L. Sp. 165) perennis molliter vel sericeo-lanata sœpe argentea, rhizomate ramoso cespitoso sublignoso, foliis in rhizomatis ramorum apice rosulatis linearibus vel lanceolato-spatiulatis acuminatis inferne longe attenuatis integris trinerviis planis vel sœpius undulatis, scapis teretibus numerosis ascendentibus folia superantibus, spicis cylindricis elongatis laxifloris basi sœpe interruptis, bracteis concavis oblongis eucullatis dorso herbaceis glabris vel breviter velutinis late membranaceis apice penicillato-pilosis calycem æquantibus, calycis laciniis ovatis apice pilosis dorso carinanti herbaceo excepto scariosis, corollæ lobis ovato-oblongis breviter acuminatis, capsulâ dispermâ, seminibus ovato-oblongis cymbiformibus ♀. Ic. Cav. tab. 124. — Fl. Græc. tab. 155. — *P. interrupta* Poir. Diet. — *P. argentea* Ten. Nap. Suppl. non D.C.

Hab. in arenosis et ad vias regionis calidæ, Græcia et ejus insulae!, Byzantium (Noë!), Rhodus (Bourg!), Cyprus (Ky!), Palestina Arabiae contermina (Boiss!), Egyptus inferior ad Alexandriam (Ehrenb! Gaill!), Persia australis ad Dalaki (Haussk!).

Ar. Geogr. Regio mediterranea Europæ et Africæ borealis.

++ Annuo-perennantes.

10. **P. cylindrica** (Forsk Eg. Arab. p. 31) annua et perennans acaulis vel brevissime caulescens collo interdum subramosa dense argenteo-lanata, foliis anguste aut latiusculæ linearibus vel lineari-oblongis elongatis basi longe attenuatis calloso-acutiusculis integris, pedunculis folio brevioribus interdum brevissimis tandem declinatis, spicis oblongis vel cylindricis, bracteis ovalis obtusis dorso herbaceis hirsutis margine late membranaceis ciliatis calycem subæquantibus, calycis laciniis oblongis obtusis dorso herbaceis margine et apice longe ciliatis, corolla glabra lobis ovato-lanceolatis acutis, capsulâ dispermâ, seminibus oblongis cymbiformibus dorso parce et irregulariter lineolato-suleatis ⊖ et ♀.

Hab. in arenosis Egypti inferioris ad Pyramides (Sieb! Ky!), circa Alexandria (Ky! sub *P. Victoriae*), ad Kahiram (Wiest 528!), in deserto inter Kahiram et Suez (Boiss!), Arabiæ Petreæ ad Ras Hamam (Schimp. 200!), planicie Ramleh (Boiss!). Arabiæ *Berchemi*.

Folia 3-4-pollicaria 1½-3 lin. lata, spicæ sex et denum 24-30 lineas longæ. Radice sœpius annuâ, pedunculis abbreviatis, spicis densifloris, bracteis hirsutis a *P. albicante* distincta.

11. **P. Stocksii** (Boiss. in D.C. Prodr. XIII, p. 720) annua vel perennans longe et patule sericeo-lanata, collo demum cespitoso et indurato, foliis lineari-vel oblongo-lanceolatis 3-5-nerviis inferne attenuatis apice longe attenuato-acuminatis, scapis tenuibus ascendentibus vel declinatis folia æquantibus vel subsuperantibus in spicas tenues longas remotifloras abeuntibus, bracteâ calyce subbreviore ovalâ obtusâ mucronulatâ membranaceo-marginatâ hirtâ vel glabratâ, calycis laciniis ovato-oblongis obtusis dorso viridibus hirtis vel gla-

bratis margine late membranaceis, corollæ glabrae lobis lanceolatis acutis, capsulâ oblongo-conicâ dispermâ, seminibus nitidis oblongo-linearibus cymbiformibus ♂ et ♀. *P. remotiflora* Stocks in Hook. Journ. 1852, p. 179.

Hab. in lapidosis Persiae prope Schahrud, inter Kerman et Ispahan (Bge!), prope Taft (Buhsel!), in Affghania ad Bolan (Griff!), in Belutschia (Stocks!).

Folia 1-2-pollicaria 2-3½ lineas lata, spicae 3-8-pollicares floribus 5-15 distantibus, semina 2 lineas longa. Species insignis ex *P. albicantha* affinitate

+++ Annuæ.

12. *P. amplexicaulis* (Cavan. Ic. II, p. 22, tab. 125) annua plus minus pilosa sicca nigricans acanlis vel caulescens collo sœpe ramosa, foliis lanceolatis acutis quinquenerviis in petiolum basi vaginantia-amplexicanlem angustatis, pedunculis axillaribus teretibus folio plerumque longioribus, spicis ovato-globosis, bracteis glabris ovato-orbiculares eucullatis obtusis præter dorsum virens membranaceis calyceem æquantibus, calycis glabri lacinias ovato-rotundis anticis dorso virenti carinatis posticis omnino membranaceis, corollæ lobis ovato-oblongis acutis, capsulâ dispermâ, seminibus magnis oblongis hinc convexis illine concavis margine anguste membranaceis ♂. *P. lagopoides* Desf. All. I, p. 135, tab. 39 — *P. Bauphala* Edgew. in Hook. Journ. Bot. 2, p. 285. — *P. salina* Decaisne in D.C. Prodr. XIII, p. 720.

Hab. in aridis plerumque salinis et gypsaceis, Græcia in monte Lycabæto Attice (Heeldr!) et ad Naupliam (Sprun!), insula Astypalea Cycladum (Urv!), Anatoliâ (ex D.C. Prodr.), Egypto inferiore (Ky! Schweinf!), Arabia petrea prope Nuckl (Boiss!), Persia australi prope Dalechi (Ky. 962! Haussk!), Affghania (Griff!), Belutschia (Stocks!).

Multum variat foliorum latitudine, caule nullo vel 4-6-pollicari, scapis interdum brevissimis.

Ar. Geogr. Insulæ Canarienses, Hispania australis, Africa borealis, India borealis, Arabia.

13. *P. Lœflingii* (L. Sp. 166) annua pumila acanlis patule pubescens, foliis linearibus vel anguste linear-lanceolatis subtrinerviis integris vel remote et tenuiter denticulatis attenuato-acuminatis, scapis tenuibus teretibus erectis vel decumbentibus folia æquantibus vel vix superantibus, spicis parvis ovatis vel oblongis, bracteis calyci æquilongis orbiculatis transverse latioribus præter dorsum lineare herbaceum in micronulum obtusum excurrens late membranaceis margine ciliatulis vel glabris, calycis lacinias omnino seariosis ovato-orbiculatis glabris, corollæ lobis breviter lanceolatis acutis, capsulâ dispermâ, seminibus oblongo-linearibus levibus facie internis sulcato-cymbiformibus ♂. Ic. Jacq. Vind. 2, tab. 126. — Trautv. Imag. tab. 18. — *P. minuta* G.A. Mey. Enum. non Pallas. — *P. Caspica* F. et M. Ind. Scin. 1837. — *P. nana* Bertol. Mise. I, p. 42!

* b in arenosis et ad cales aridos Syriae ad Meppo (Haussk!), ad Euphra-

tem (Chesney 35!), Mesopotamiæ (Haussk!), Georgiae et Iberiæ Caucasicæ (C.A. Mey! Hoh!), Persie australis prope Dalechi (Ky. 961!).

Ar. Geogr. Insulæ Canarienses, Hispania interior, Africa borealis.

14. *P. minuta* (Pall. It. Suppl. № 69, tab. E) annua pumila acaulis patule pubescens, foliis linearibus vel anguste linear-i-lanceolatis integris acuminatis uninerviis, scapis tenuibus teretibus ascendentibus folia æquantibus vel vix longioribus, spicis ovatis vel oblongis, bracteis calyce brevioribus calycisque laciniis glabris ovatis obtusis omnibus præter dorsum lineare herbaceum late albo-membranaceis, corollæ lobis triangulare-ovatis obtusiuseulis, capsula dispermâ, seminibus linear-i-oblongis cymbiformibus ⊖.

Hab. in Turkestaniâ ad mare Caspium (Kar!) et in desertis salsuginosis interioribus (Lehm!). In Iberiâ a Biébersteinio indicata sed prob. ex confusione cum *P. Löflingii*.

Valde affinis *P. Löflingii* a quâ tamen bracteâ calyce breviore et calycis glabri laciniis non omnino hyalinis sed dorso herbaceis distincta videtur.

Ar. Geogr. Rossia austro-orientalis, Songaria, Sibiria Uralensis.

15. *P. Bellardi* (All. Ped. I, p. 82, tab. 85) acaulis annua tota villosa, foliis lanceolatis vel lanceolato-linearibus acutis basi attenuatis trinerviis integris vel obsolele paucidentatis, pedunculis teretibus crassiuseulis erectis folia æquantibus vel superantibus, spicis densis ovato-oblongis vel cylindricis villosis, bracteis herbaceis villosis lanceolatis obtusiuseulis calycem æquantibus vel sublongioribus, calycis villosi laciniis oblongis exterioribus herbaceis interioribus præter nervum scariosis acuminatis, corollæ glabrae lobis oblongo-lanceolatis acuminatis. capsule ovata loculis monospermis, seminibus ovatis cymbiformibus ⊖. Ic. Fl. Græc. tab. 146. — Rehb. Germ. tab. 82. — *P. pilosa* Pourr. Tolos. 3. p. 324. — *P. holostea* Lam. Ill. № 1667.

Hab. in collinis siccis et arenosis, Graecia in Zacyntho (Marg!), Peloponneso (herb. Fauché! Orph!), Atticâ (Sprun!), Macedonia (Friv!), Byzantium (Pichl!), Anatolia orientalis ad Smyrnam (Bal!), Persia australis prope Gere (Ky. 62 exs! sub *P. Cretica*, Haussk!).

Ar. Geogr. Regio mediterranea totius Europæ et Africæ borealis.

16. *P. Cretica* (L. Sp. 165) acaulis annua tota villosa, foliis linear-i-lanceolatis vel linear-i-spathulatis acutiusculis basi sensim et longe attenuatis integris rarius subdentalis trinerviis, pedunculis teretibus crassis dense lanatis folio multo brevioribus junioribus erectis dein induratis recurvis, spicâ ovato-globosâ, bracteis herbaceis oblongo-lanceolatis dense lanatis calycem superantibus, calycis laciniis ovatis non carinalis superne lanatis exterioribus subherbaceis interribus membranaceis, corollæ glabrae lobis ovatis abrupte et breviter acuminatis, capsule ovata loculis monospermis, seminibus ovatis cymbiformibus ⊖. Ic. Fl. Græc. tab. 147.

Hab. in arenosis regionis inferioris, Creta (Sieb! Raul!), Byzantium (Noë!), Rhodus (Bourg!), Cyprus (Sibth!), Lydia ad Smyrnam (Auch. 2706! Bal!), Pamphylia (Heldr! Bourg!), Cilicia littoralis (Bal!), Syria littoralis (Bl!) et

interior ad Aleppo (Ky. 66! Haussk!) et Damascum (Gaill!), Palestina ad Hebron (Ky!) et Gaza (Boiss!).

Affinis *P. Bellardii* sed pedunculis brevibus demum recurvis, spicis abbreviatis, calycis laciñis subnudicis et corollæ lobis magis rotundatis distincta.

17. *P. ovata* (Forsk. Eg. Arab. p. 31) annua acaulis pumila molliter villosa lanata, foliis anguste linearibus vel lanceolatis integris vel obsolete calloso-denticulatis basi longe attenuatis, scapis gracilibus foliis æquiforis vel brevioribus sæpius declinatis, spicis subgloboso vel cylindrico-ovatis, bracteis glabris ovato-rotundis obtusis late scariosis fuscocentibus vel pallidis calycem æquantibus, calycis laciñis ovatis obtusis late scariosis glabris vel pubescentibus, corollæ lobis ovatis vel rotundis brevissime mucronulatis, capsulae lobulis monospermis, seminibus oblongis cymbiformibus margine anguste membranaceis ⊙. *P. decumbens* Forsk. Eg. Ar. p. 30 (non differt). — *P. argentea* Desf. Atl. II, p. 136. — *P. microcephala* Poir. Dict. IV, p. 378. — *P. villosa* Mœnch Méth. p. 459.

Hab. in arenosis et desertis Egypti inferioris et mediae (Forsk., Boiss Schweinf! etc.), Arabiae petreæ (Schimp. 203! Boiss!), Syriae interioris ad Palmyram (Bl!), Assyria ad Mossul (Noë 225!), Iberiae Caucasicæ et prov. Karabagh (C.A.M! Hoht), Persiae mediae orientalis (Bge!) et australis ad sinum Persicum (Ky. 61!), Turkestanijæ (Karl).

Ar. Geogr. Insulae Canarienses, Madera, Hispania austro-orientalis, Africa borealis, India boreali-occidentalis.

18. *P. Haussknechtii* (Vatke Verh. Bot. Brandenb. 1874, p. 53) annua acaulis pumila adpresso hirsuto-canescens, foliis integris linear-oblängis subtrinerviis basi longe attenuatis, scapis tenuibus substriatis foliis sublongioribus, spicis oblongis et oblongo-cylindricis, bracteis late ovato-orbiculatis obtusis dorso lanatis late membranaceo-marginatis calycem æquantibus, calycis laciñis superne hirsutis, corollæ lobis basi fuscocentibus ovatis acutis, capsulae loculis monospermis (altero interdum ex cl. Vatke dispermo), seminibus oblongis cymbiformibus ⊙. *P. phæopsis* Payne in Proceed. Palestine Explor. Soc. Appendix.

Hab. in Palestinâ ad orientem Jordani (Payne!), Mesopotamiâ (Oliv!) et prope Katt el Amara (Haussk!), in Persiâ australi ad fontes petroleum fundentes prope Dalechi (Ky!), prope Abuchir (Haussk!).

Scapi cum spicis 3-4-policlavæ, folia 1-3 lineas lata, facies *P. ovatae* a qua bracteis lanatis statim distinguitur, a *P. notatae* foliis integris, calyce superne nec basi hirsuto, corollæ lobis obtusioribus.

19. *P. notata* (Lagasc. Gen. et Sp. № 102) annua pumila acaulis plus minus hirsuta pallide virens, foliis rosulatis sessilibus lanceolatis vel linearibus remote lacinulatis trinerviis lacinulis anguste linear-lanceolatis basi pilorum fasciculo sæpins anctis, scapis teretibus declinatis folia vix æquantibus, spicis ovato-globosis rarius breviter cylindricis, bracteis ovato-orbiculatis dorso lanatis late membranaceis calycem æquantibus, calycis basi lanati laciñis glabriusculis

omnino membranaceis ovatis obtusis, corollæ lobis fuscis ovato-orbiculatis submucronulatis, capsulâ biloculari loculis monospermis, seminibus oblongis cymbiformibus ♂. *P. Syrtica* Viv. Libyc. p. 7, tab. 3. — *P. Olivieri* Decaisne in Barneoud Mon. Plant. p. 37. — *P. praecox* C A. Mey. Enum. p. 115.

Hab. in arenosis, Egyptus inferior prope Alexandriam (Oliv. Cadet! Samar!), Transcaucasia ad Baku (C.A. Mey!).

Planta 3-6-pollicaris facie *P. ovatae*, insignis foliis longe et anguste lacinulatis, calyce basi longe lanato omnino scarioso.

Ar. Geogr. Hispania austro-orientalis, Africa borealis.

** Corollæ tubus glaber, lobi hirsuti. Plante annua.

20. ***P. Lagopus*** (L. Sp. 165) acaulis annua, radice fibrosâ, collo hirsutissimo, foliis lanceolatis vel oblongo-spathulatis in petiolum attenuatis 3-5-nerviis glabrescentibus vel pubescenti-hirsutis integris vel obsolete denticulatis, scapis sulcatis erectis et ascendentibus pubescenti-sericeis folia 2-4-plo superantibus, spicis ovatis vel ovato-cylindricis densis, bracteis ovato-lanceolatis acuminatis praeter nervum viridem scariosis superne villosis, ealycis laciñiis apice villosis lateralibus carinatis, corollæ lobis ovatis acutis vel acuminatis extus secus nervum plus minus pilosulis, capsulae obovatae obtusæ loculis binis monospermis, seminibus oblongis cymbiformibus laevibus ♂. Ic. Fl. Gr. tab. 144. — Rehb. Ic. Germ. tab. 82. — *P. eriostachya* Ten. Prodr. p. 13. — *P. fornicata* C. Koch Linn. XXI, p. 713. — *P. glauca* C. A. Mey. Enum. p. 115!

Hab. in herbidis asperis siccis totius ditionis a Graeciâ (Orph. Fl. exs. 694!), Macedonia et Thraciâ (Griseb!), ad Taurium (Letz.), Transcaucasiam ad Caspium (C.A. Mey!), Persiam australē (Ky. 175!), Babyloniam (Noë!), Syriam et Palestinam! Egyptum inferiorem (Wiest! Schweinf!), et in Oasibus (Aschers!).

β. major. — Omnibus partibus major sepe subcaulescens. Folia latiora. — *P. Lusitanica* Willd. Sp. I, 644.

Hab. in Etolia (Nied!), Attica (Sprun!), Syria littorali (Gaill!), Mesopotamia (Noë!).

Ar. Geogr. Regio mediterranea totius Europæ et Africæ borealis, Insulæ Canarienses, Madera.

21. ***P. lagoccephala*** (Bunge pl. Lehm. p. 445) annua acaulis nana molliter pilosa, foliis anguste lineari-lanceolatis uninerviis basi et apice attenuatis acutis, scapis tenuibus arcuatis teretibus foliis subbrevioribus, spicis ovato-oblongis villosissimis, bracteis calycem æquantibus villosis ovatis longe acuminatis carinatis anguste membranaceis, ealycis laciñiis extus villosissimis margine membranaceis inæqualibus binis semiovatis inæquilateris binisque cum prioribus alternis linearı-oblongis acutiusculis, corollæ tubo glaberrimo laciñiis ovatis acutis extus undique hirsutissimis, capsulâ oblongâ dispermâ, seminibus navicularibus granulatis ♂.

Hab. in Turkestaniā ad fluvium Jan Darja (Lehm.).

Folia 1-2-pollices longa 1-1½ lin. lata, capitula 4-6 lineas longa. Facies *P. ovata*, ab ea et affinibus corollæ lobis extus hirsutissimis distincta.

22. *P. ciliata* (Desf. Atl. I, p. 137, tab. 39) annua vel perennans acaulis vel brevissime caulescens et collo ramosa humilius plus minus adpresso sericeo-argentea, foliis obovato vel lanceolato-spathulatis acutiusculis in petiolam saepe attenuatis, pedunculis crassiusculis folio æquilonigis vel brevioribus, spicis ovatis vel oblongo-cylindricis, bracteis ovatis obtusis calycem æquantibus dorso viridi hirtis late membranaceis et longe ciliatis, calycis laciniis membranaceis ovatis margine et apice longe villosis, corollæ tubo glabro lobis lanceolatis extus longe villosis, capsulæ dispermæ seminibus hinc convexis illine concavis hæribus ☽. *P. bellidifolia* Viv. Egypt. Dec. IV.

Hab. in arenosis deserti Egyptiaco-Arabici ad Ouadi Kuler inter Kenel et Kosseir (Schweinf!), Arabiæ petreæ in ditione Hanara (Schimp. 425!), ad Ouadi Useit (Boiss!).

β. lanata. — Indumentum minus adpressum! — *P. penicillata* Endl. et Fenzl Sert. Cabul. pag. et tab. I. — *P. eriantha* Decaisne in D.C. Prodr. XIII, p. 707.

Hab. in montosis Persiae orientalis ad Feisabad inter Chabbise et Kerman (Bge!), in Affghaniā (Honigb., Griff!), in Belutschia (Stocks!).

Ar. Geogr. Arabia borealis interior, Arabia tropica, India boreali-occidentalis.

*** Corollæ tubus hirsutus, lobi glabri. Annuae.

23. *P. lachnantha* (Bge Pl. Lehm. p. 446) annua acaulis nana tomentosa, foliis anguste linear-lanceolatis carnosulis acutiusculis uninerviis, scapis erectis firmis lanuginosis folio brevioribus, spicis globosis dense paucifloris, bracteis ovatis late membranaceis nervo viridi excurrente acutis glabris, sepalis elliptico-ovatis extus molliter lanatis anterioribus late interioribus angustius viridi-carinatis, corollæ tubo dense lanato lobis ovato-oblongis acutissimis margine obsolete ciliolatis, capsulæ dispermæ seminibus... ☽.

Hab. in Turkestaniā inter Ayatna et Karagata (Lehm.). Non vidi.

Junior et ex paucis speciminiibus tantum descripta sed corollæ tubo lanato insignis. Folia pollicem longa 1-1½ lin. lata, scapi semipollicares, spicae subquinquefloræ parvæ 2½ lineas fere in diametro.

24. *P. evaina*, annua acaulis nana tomentosa, foliis carnosulis anguste linearibus uninerviis, scapis erectis tandem recurvis crassis folio brevioribus spicæ vix brevioribus, spicis globosis densis pauci-floris, bracteis extus lanatis oblongo-cymbæformibus obtusiæculis calyce subbrevioribus, calycis laciniis extus lanatis ovatis obtusis concavis non carinatis dorso viridibus late membranaceis, corollæ tubo lanato lobis ovato-lanceolatis acutis glabris, capsulæ dispermæ, seminibus oblongis cymbæformibus ☽.

Hab. in Affghaniâ (Griff!), Persiâ orientali ad Schahrud, ad Kerman et prope Sserschah (Bunge!).

Facies *Evacis*, folia vix pollicem longa, capitula magnitudine Pisi. Ex descriptione affinis *P. lachnantha* quam non vidi et quæ capitulis ut videtur multo minoribus et bracteis glaberrimis differre videtur.

§ 5. Biconvexæ.

* Capsula ob placentas 3-4-alatas 3-4-locellata locellis monospermis.

25. *P. Coronopus* (L. Sp. 166) annua vel perennans acaulis plus minus et adpresso hirsuta, foliis oblongo-lanceolatis pinnatodentatis vel pinnatifidis vel bipinnatifidis laciniis acutis linearibus vel lanceolatis vel oblongis, scapis foliis longioribus vel brevioribus ascendentibus vel erectis, spicis cylindricis vel oblongis, bracteis breviter et adpresso hirtis anguste marginatis a basi ovalâ attenuatis acutis vel acuminatis, calycis segmentis anticus hirsutis ciliolatis positie dorso carinato-alatis ciliatis, corollæ tubo hirsuto lobis ovato-lanceolatis acuminatis, capsulae ovatæ loculis bilocellatis locellis monospermis, seminibus ovato-oblengis plano-biconvexis facie internâ umbonatis ♂ et ♀. Ic. Fl. Dan. tab. 272. — Rehb. Germ. t. 79. — *P. commutata* Guss Suppl. I, p. 46 (forma scapis abbreviatis spicæ subæquilongis).

Hab. in arenosis, aridis, ad vias, præsertim in maritimis totius ditionis a Græciâ! Macedoniâ et Thraciâ! ad Transcaucasiam! Persiam! Affghaniam (Griff!), Belutschiam (Stocks!), Egyptum!

P. simplex. — Folia anguste linearia integra vel paucidentata, spicæ st̄epe abbreviatæ. — *P. filiformis* C. Koch Linn. XXI, p. 709.

Hab. in Egypto (Ehr.), Græciâ (Auch. 2746!), Transcaucasiâ ad Baku (Seidl!), Babylonia ad Mohammera (Noë!), Persiæ australis salsis ad Dalechi (Ky. 197!).

Ar. Geogr. Europa media et australis a Daniâ et Gothiâ ad Germaniam et Dalmatiam, Africa borealis.

26. *P. crypsoides*, annua acaulis nana papillis adpresso scabrida, foliis linearibus vel lanceolatis acutis basi longe attenuatis subtrinerviis integris vel paucilacinulatis, scapis crassis spicæ subæquilongis foliis multo brevioribus tandem recurvis, spicis villosis ovatis paucifloris densis, bracteis laciniisque calycis coriaceis herbaceis hirsutis membranaceo-marginalis carinatis illâ subbreviore triangulari-ovatâ acutâ, hisce ovato-oblengis, corollæ tubo adpresso hirto 1 bis lanceolatis acutis, capsulae loculis bilocellatis locellis monospermis, seminibus ovalis plano-biconvexis marginatis minimis ♂. *P. Coronopus* var. *bombycinus* Decaisn. in p.c. Prodr. XIII, p. 732.

Hab. in arenosis Egypti circa Alexandriam (Ky., Schim.), Samar! Krâlik sub *P. Cretica*!).

Facies omnino *P. Cretica* a quâ seminibus, corollæ tubo hirto, etc., omnino

differt. Corollæ tubo hirto accedit ad *P. lachnantham* et *evacinam* quæ quoque seminibus cymbiformibus alienæ sunt. Folia 1-3-pollicaria, pedunculi 3-4 lineas longi, capitula piso vix majora. Differt a *P. Coronopi* formis spicis lanatis, scapis recurvis, seminibus triplo minoribus marginatis, etc.

• Capsula placentis bialatis bilocularis disperma.

27. *P. Serraria* (L. Sp. 166) peregrinis acaulis papillari-hirsuta, radice crassâ, foliis lanceolatis quinquenerviis acutis in petiolum planum brevem attenuatis in lacinias anguste linearis-lanceolatas incisoserratis, scapis teretibus foliis æquilongis vel longioribus, spicis cylindricis, bracteis ovatis apice attenuatis acutiusculis calyce brevioribus, calycis laciñis anticis ovatis, posticis dorso carinato-subulatis saepius ciliolatis, corollæ tubo hirto lobis ovatis acutis, capsulæ bilocularis loculis monospermis, seminibus plano-biconvexis ♀. Ic. Barrel. p. 749.

Hab. in pratis mari finitimis Zacynthi (Sibth., Heldr!), ad Liutzi Peloponnesi (Sart!).

Ar. Geogr. Lusitania, Hispania australis, regnum Neapolitanum, Sicilia, Dalmatia, Africa borealis.

28. *P. maritima* (L. Sp. 165) acaulis, rhizomate crasso carnoso saepè pluricipiti squamoso, foliis carnosis longe linearibus crassiusculis superne subcanaliculatis trinerviis nervis æquidistantibus integris vel remote denticulatis glabris rarius hirtis ad vaginas basilares saepius hirsutis vel lanatis, scapis teretibus adpresso hirtis foliis saepius longioribus, spicâ cylindrica gracili laxiusculâ floribus axi adpressis, bracteis ovatis concavis calyce brevioribus membranaceo-marginatis, calycis laciñis obtusis late scariosis, posticorum carinâ viridi in alam membranaceam angustam subexpansâ, corollæ tubo adpresso hirto lobis ovatis acutis, capsulae loculis binis monospermis, seminibus ovatis subcompressis ♀. Ic. Fl. Græc. tab. 148. — Rchb. Ic. Germ. tab. 80. — *P. salsa* Pall. — *P. teretifolia* Sieb. exs.

Hab. in salsis maritimis et interioribus, Attica ad Phalerum copiose (Sprun! Heldr! Orph. Fl. exs. 693!), Peloponnesus (Heldr!), Macedonia (Griseb.), Syria littoralis! Cyprus (Sibth., Kotschy!), Egyptus ad Alexandriam et Rosettam (Forsk., Ehrenb! Sieb! etc.), Tauria (Stev.), Persia borealis in prov. Aderbidjan prope Tabris (Buhse!).

γ. ciliata Koch Syn. — Folia asperula abbreviata, spicæ breviores. — *P. Wulfeni* M. et K. — *P. aspera* Gaud. — *P. Alpina* Sieb. exs.

Hab. in pascuis montanis, Armenia supra Baibut (Bourg!).

Ar. Geogr. Europa omnis, Sibiria.

29. *P. carinata* (Schrad. Cat. Gött. — Mert. et K. D. Flora I, p. 810) rhizomate cespitoso pluricipiti, foliis brevibus trigono-linearibus subtus carinatis basi glabriusculis vel sublanatis, scapis tenuibus teretibus folia multum superantibus adpresso hirtis, spicis breviter cylindricis interdum abbreviatis ovato-globosis, bracteis ovato-lanceo-

latis acutis calycem æquantibus, calycis laciniis late scariosis posticis carinatis. corollæ tubo adpresso hirto lobis ovato-lanceolatis acutis, capsulae loculis binis monospermis, semiibus ovatis subcompressis ȝ. Rchb. Germ. tab. 81. — *P. serpentina* Koch Syn. — *P. subulata* Wulf. Jacq. Collect. I. tab. 10, et Smith Prodr. — *P. capitellata* Ram. (forma spicis brevissimis). — *P. Alpina* var. *dioritica* (Ky. exs!).

Hab. in pascuis regionis subalpinæ et alpinæ, Græcia in monte Taygeto (Heldr!), Malevo (Orph. Fl. Gr. exs. 12!), Tymphresto Ætoliae (Sprui! Heldr. Herb. Norm. 671!), Macedonia supra Bitolia (Orph!), prope Perlepe (Griseb!), mons Rhodope Thraciæ (Friv!), Olympus Bithynus (Auch. 2705! Boiss!), Tmolus Lydia (Boiss! Bal. 288!), mons Berytdagh Cataoniae 6000' (Haussk!), Taurus Cilicus 8000' (Ky. 127! Bal!), Libanus (Bl!).

Ar. Geogr. Gallia, Hispania, Italia, Carnia, Dalmatia, Bosnia, Serbia.

SECT. II. PSYLLIUM.

* Perennes.

30. P. Arabica (Boiss. in Pinard exs. 1847. — Diagn. Ser. I, 12, p. 94) perennis dumulosa, caulinis suffrutescentibus ascendentibus ramosis inferne denudatis, ramis dense foliosis, foliis hispido-viscosissimis strictis linearis-triquetris superne sulcatis integerrimis obtusiusculis subincurvis, pedunculis ex axillis superioribus paucis corymbosis folia subsuperantibus, spicis ovato-sphaericis paucifloris, bracteis ovatis acuminatis vel acutis breviter hirto-viscidis calyce subbrevioribus, calycis laciniis anticis oblongis obtusiusculis hirtis, posticis carinato-plicatis paulo angustioribus, corollæ lobis ovato-lanceolatis acutis ȝ. *P. Psyllium* var. *Sinaica* Barn. Mon. p. 49. — *P. Psyllium* Decaisn. Fl. Sinaic. p. 7. — *P. Sinaica* Decaisn. in D.C. Prodr. XIII, p. 733.

Hab. in Arabiâ Sinaiticâ (Ehr! Bové! Auch. 1707! Schimp. 172!), in valle Arbain (Boiss!).

Dumi pedales, folia 1-1½-pollicaria ¼-1 lineam lata, capitula 5-7-flora piso submajora. A *P. Psyllio* caulinis fruticosis, etc., omnino aliena, affinis *P. Cynopi* a quo differt indumento viscido, bracteis non longe caudatis, etc. Cl. Barneoud nullam *P. Sinaican* (in Prodr. D.C. citatam) descriptam.

31. P. Euphratica (Decaisne in Barn. Mon. p. 51) perennis suffruticosa collo ramosa siccitate nigricans scabridula et pilosovillosula, ramis erectis, foliis linearibus elongatis longe attenuato-acuminatis planis integris, pedunculis ex axillis superioribus oppositis internodia et folia superantibus, spicis ovato-oblongis, bracteis ovatis acutiusculis membranaceo-marginatis dorso glabris vel hirtis, calycis laciniis oblongis posticis subcarinatis omnibus dorso hirtulis vel glabris, corollæ lobis ovato-lanceolatis acutis ȝ.

Hab. in Cappadociâ ad Euphratem (Auch. 2707!), in monte Begdaghi supra Malatia (Haussk!).

Pedalis, folia 2-3-pollicaria, pedunculi 2 pollices longi. A *P. Cynope* inter alias notas bracteis ut in *P. Arabicâ* ecaudatis distincta, ab ea foliis planis multo longioribus, pedunculis elongatis, indumento non viscido breviter scabridulo et pilis longis albidis intermixto aliena.

• • Annuæ.

32. P. Psyllium (L. Sp. 167) annua glandulifero-pubescent, caule herbaceo erecto simplici vel thyrsoideo-ramoso, foliis anguste linear-i-lanceolatis vel linearibus integris vel remote denticulatis, pedunculis ex axillis superioribus foliis sublongioribus corymboso-thyrsoideis, spicis ovato-sphaericis densifloris glanduloso-hirtis, bracteis e basi ovatâ lanceolatis acutis vel florum inferiorum acuminatis, calycis laciniis conformibus oblongis apice attenuatis, corollæ lobis lanceolatis acutis ♂. Ic. Fl. Græc. tab. 149. — Rchb. Germ. tab. 84.
— *P. Afra* L. Sp. 168.

Hab. in graminosis, cultis, vineis regionis inferioris et montanæ, Græcia et ejus insulæ! Macedonia et Thracia (Griseb.), Anatolia occidentalis, Cilicia (Bal!), Syria littoralis (Gaill!), Palestina (Noël), Egyptus in Oasi parvâ (Aschl!), Babylonia (Noël), Persia borealis et australis (Ky!).

Ar. Geogr. Regio mediterranea.

33. P. stricta (Schousb. Marocc. p. 35) annua superne minute pruinosa ad nodos papillari-hirtula, caule tenui herbaceo subsimplici, foliis anguste linearibus elongatis integris, pedunculis ex axillis superioribus tenuibus folia sæpe superantibus, spicis parvis ovatis densifloris sub lente minute scabrido-hirtis, bracteis lanceolatis florum inferiorum lanceolato-linearibus flore longioribus, calycis laciniis conformibus oblongis acutiusculis, corollæ lobis lanceolatis acutis ♂.
P. rugosa Hochst. in Schimp. Pl. Abyss.

Hab. in Egypto inferiore ad Belbeys (Ehrenbl!), Abuzabel (Schimp. 93!), ad Mataryeh (Kralik!), ad Ain el Beitha prope Aleppo (Haussk!), in Persiâ australi prope Dalechi (Ky!).

Verosimiliter varietas *P. Psyllii* a quo tantum differt indumento brevi non vel vix glanduloso-viscoso, pedunculis gracillimis, spicis minoribus glabriusculis, bracteis florum inferiorum elongatis. Exacte *P. Psyllio* et *P. pumilæ* intermedia.

Ar. Geogr. Croatia, Africa borealis, Abyssinia.

34. P. pumila (Willd. Enum. Hort. Ber. I. p. 162) annua glabriuscula, caulis tenuibus a basi decumbente erectis flexuosis ramosis, foliis subulato-capillaribus tenuissimis elongatis subrevolutis basi sæpe hirsutis, pedunculis ex axillis supremis filiformibus folia superantibus, capitulis globosis paucifloris minute puberulis, bracteis a basi sublatiore subulatis inferioribus spicæ æquilongis superioribus calyce longioribus, calycis laciniis oblongo-lanceolatis obtusinseulis, corollæ lobis ovato-lanceolatis acutis ♂. *P. Indica* Linn. ex descript.
— *P. exigua* Murr. Comm. Gött. 1778, p. 94, tab. 5 — *P. Rosetana* Poir. Dict. Suppl. IV, p. 433.

Hab. in Egypto superiore prope Girgeh (spontanea ex cl. Husson!), culta ex cl. Schweinfurth!), in regno Cabulico ad Schikarpore (Griff!).

Glabrescentia, caulis decumbentibus, foliis pedunculisque tenuissimis, spicis ob bracteas inferiores interdum omnes elongatas foliosis ab affini *P. stricta* dignoscenda.

35. *P. phæostoma* (Boiss. et Heldr. Diagn. Ser. II, 4, p. 71) annua tota glanduloso-puberula et papillari-hirta collo ramosa, caulis nanis lateratibus divaricatis saepe ramosis, foliis anguste linearibus integris, pedunculis foliis capitulisque æquilongis vel brevioribus, spicis oblongis densifloris breviter hirtis, bracteis oblongo-linearibus obtusis præter infimas calyce brevioribus, calycis lacinias oblongis obtusiusculis conformibus anguste membranaceis, corollæ lobis ovatis mucronatis fauce fuscâ ⊖.

Hab. in arenosis ad Alexandriam (Samar! Letourn!), ad Ouadi Kuser et Ouadi Askar deserti Ægyptiaco-Arabici (Schweinf!), ad Ouadi Mokatteb Arabiae petreæ (Boiss!).

Plantula 2-3-pollicaris saepius ramosissima, folia $\frac{1}{4}$ - $\frac{1}{2}$ -pollicaria, spicæ 6-8 lineas longæ 3 latæ. Affinis *P. strictæ* differt caulis nanis, pedunculis brevibus, bracteis abbreviatis.

36. *P. arenaria* (W. K. Pl. Rar. Hung. I, p. 51, tab. 54) annua pubescens, caule erecto vel diffuso stricto ramoso, foliis anguste linearibus vel filiformibus elongatis margine subrevolutis integris basi saepe villosulis, pedunculis axillaribus folio longioribus saepe subumbellato-thyrsoideis, spicis ovatis densis, bracteis dorso breviter papillari-hirtis infimis ovatis in cuspidem herbaceam flores superantem attenuatis superioribus spathulato-orbiculatis membranaceo-marginatis, calycis lacinias anticis oblique ovatis obtusissimis, posticis lanceolatis acutiusculis membranaceis, corollæ lobis ovato-lanceolatis acatis ⊖. Rchb. Germ. t. 85. — *P. Cynops* Sm. Prodr. I, p. 103 non L.

Hab. in campis arenosis saepe maritimis, Thessalia in maritimis (Heldr! Orph!), Thracia ad Philippopolin (Friv.), Bithynia (Sibth., Pichl!), insulæ Naxos et Thera (Sart!), Anatolia borealis (Wied!), Pontus (Bourgl!), Tauria (Stev.), Transcaucasia (Lebed., Becker!).

β. divaricata. — Divaricatum ramosa, folia abbreviata sublatiora. Forma maritima.

Hab. in arenosis maritimis ad Astros Pelopennesi (Orph!), in insula Andros (Sart!), in oasi Chargeh Egypti (Schweinf!).

Ar. Geogr. Hispania centralis, Gallia, Helvetia australis, Germania, Italia borealis, regio Danubialis, Rossia media et australis.

37. *P. squarrosa* (Murr. Comm. Gölt. 1781, p. 38, tab. 3) annua papillari-hirta, a collo saepius ramosa, caulis diffusis vel ascendentibus saepe ramulosis, foliis carnosulis linearibus vel lineario-oblongis abbreviatis acutis subcanaliculatis saepe recurvis, pedunculis ex axillis superioribus capitulo æquilongis vel brevioribus saepe subumbellatum 3-5 congestis, spicis pubescentibus oblongis vel oblongo-

cylindricis interdum laxiusculis, bracteis inferioribus binis spicam involucrantibus interdum elongatis oblongo-lanceolatis recurvis supra basin dilatatam substrangulatis, superioribus oblongis obtusiusculis calycrem æquantibus vel subsuperantibus, calycis laciniis anticis oblongo-spathulatis subobliquis, posticis oblongis carinato-subulatis, corollæ laciniis ovato-oblongis acutis ⊖. *P. Ægyptiaca* Jacq. Ic. Rar. tab. 28. — Collect. I, p. 45. — Fragm. tab. 81, fig. 2.

Hab. in arenosis maritimis, Syria ad Berythum (Bl! Gaill!), Palestina ad Joppem (Ky. 581).

β. brachystachys. — Spicæ globosæ sæpe glabrescentes, bracteæ involucrales crassiores et breviores. — *P. sarcophylla* Boiss. exs.

Hab. in Egypto ad Alexandriam (Ehrenb! Gaill!), ad Pyramides Gizeh (Hussont!), in oasi el Nuehl (Ky. 496!), Palestina prope Gaza (Auch. 2714! Boiss!), ad Joppem (Ky. 650!), ad Berythum (Bl!), in Pamphyliâ ad Side (Heldr!).

Variat prætereâ foliis ramis et pedicellis angustissimis hisce capitulo longioribus, foliisque lineam vel sesquilineam latis et pedicellis crassis brevisimis. Spicæ in formâ typicâ (prius ab auctoribus descriptâ sed rariore) interdum 9-10 lineas longæ, in varietate *β. pisum* æquantes vel majores.

Species incomplete descriptæ, certo certius præcedentium nonnullarum synonyma

P. polyatochya Viv. Eg № 31. — Scaposa, foliis linearibus elongatis uninerviis integris vel remote dentatis, spicis ovato-oblongis aliis subsessilibus aliis pedunculatis, bracteis ovato-acutis mucronulatis margine membranaceis, corollæ laciniis ovato-lanceolatis ⊖.

Affinis dicitur *P. Syrticæ* Viv. = *P. notatae* et lobis corollæ acutioribus tantum distincta. Verosimiliter eadem.

P. setosa Bertol. Misc. I, p. 30. — Scaposa, foliis scapo brevioribus obverse lanceolatis obtusis mucronulatis glabris ciliatis, spicâ crassâ cylindrâ obtusâ, bracteis ovatis acuminato-aristatis ciliatis margine membranaceis, laciniis corollinis breviter aristatis ⊖.

Hab. ad oras Euphratis (Chesney 136!).

An *P. cylindrica* vel *P. ciliata*?

CLASS. I. DICOTYLEDONEÆ.

Subclass. IV. MONOCHLAMYDEÆ D.C.

Perigonum calycinum vel corollinum sœpius simplex. (Exceptis nonnullis *Phytolaccaceis*, *Loranthaceis*, *Euphorbiaceis*, *Empetraceis*, etc., in quibus petala cum calyce adsunt.)—Subclassis nimis artificialis in posterum dilaceranda, ordinibus inter cohortes præcedentes naturalius distribuendis (Confer. d.c. Prodr. XIII, p. 1).

ORD. XCVII. PHYTOLACCACEÆ.

(Endl. Gen. p. 975.)

Flores in nostris hermaphroditi. Perigonum petaloideum 4-5-partitum lacinis aestivatione imbricatis. Stamina in fundo perigonii inserta ejus lacinis alterna et numero æqualia, vel plura libera, vel in cupulam basi coalita. Ovarium interdum unicappellatum, in nostris carpellis pluribus verticillatis connatis vel distinctis uniovulatis constans, ovulum erectum basifixum (in nostris) campylotropum. Styli tot quot ovarii loculi simplices. Fructus carpellis in baccam plurilocularem coalitis vel liberis constans. Semina erecta, embryo annularis radiculâ inferâ. Albumen farinaceum centrale rarius nullum. — Ordo ad *Ficoideas* accedens.

PHYTOLACCA (L. Gen. p. 233).

Flores hermaphroditi vel dioici, tribracteati. Perigonum 5-partitum lacinis petaloideis imbricatis æqualibus fructiferis reflexis. Stamina in fundo perigonii disco carnosulo inserta 5-12 subulata, antheræ biloculares. Carpella 5-12 in toro convexiusculo sessilia verticillata et inter se connata uniovulata, ovulo basifixo campylotropo. Styli 5-12 subulati breves in coronulam approximati sed liberi facie internâ stigmatosi. Fructus baccatus plurilocularis loculis inde-

hiscentibus monospermis. Semina verticalia breviter rostellata crustacea. Albumen centrale, embryo annularis periphericus. — Herbae rarius frutices.

1. P. decandra (L. Sp. 631) hermaphrodita glabra, caule crasso valde elato dichotome ramoso saepè purpurascente, foliis amplis tenuibus ovato-lanceolatis acutis basi cuneatis brevissime petiolatis, racemis oppositifoliis pedunculo petiolo longiore sussultis erectis multifloris, pedicellis patentibus flore sublongioribus, bracteâ inferiore lineari pedicello breviore superioribus minimis ovatis, perigonii laciinis ovatis concavis, staminibus subdenis, baccâ depresso-globosâ umbilicatâ sub-10-rostellata et 10 carpellari ♀. Ic. Lam. III. t. 393.

Hab. ex America boreali oriunda nunc variis locis ad fossas, vias subsponentanea, Græcia in insulis Zacynthio et Eubeâ (Sibth.), Baetia (Sprun!), Byzantii (Sibth.), Anatolia in Ponto (Tchih!), prope Trapezuntum et Rhize (Bourg! Bal!), Tauria et Caucaso (Ledeb.), in Assyriâ ad Diarbekir (Haussk!). Egypto (Del.).

Ar. Geogr. America borealis, Europa media.

2. P. pruinosa (Fenzl Advers. bot. Hort. Vindob. 1855, p. 6) dioica pruinosa glaucescens, caulis crassis erectis dichotome et striete ramosis, foliis carnosulis ellipticis et elliptico-lanceolatis acutiusculis basi cuneatis subsessilibus, racemis oppositifoliis erectis multifloris pedunculis folio multo brevioribus, plantæ feminine fructiferæ cum rachide valde incrassatis, pedicellis patentibus flore sublongioribus, bracteâ inferiore lineari-lanceolata pedicellum subæquante ceteris minimis, perigonii viridi-rubelli laciinis ovatis concavis, staminibus 11-16, baccâ depresso-globosâ umbellicatâ carpellis et sulcis 5-9 ♀.

Hab. in rupestribus regionis montanae et subalpinæ, Cilicia littoralis supra Sediching (Bal!), Tauri jugum Bulghardagh in regione Cedrorum 4-5000' (Ky. 271!), mons Masmenedagh Cappadociae (Bal!), in rupestribus porphyricis montis Amani Syriae (Ky. 123!), in sylvaticis Cypri circa Prodromo (Ky!).

Facies *P. dioicae*, indumentum pruinatum, folia angustiora acutiora subsessilia, racemi breviores.

Obs. *Pircinia dioica* Moq. arbor Americana in plateis regionum calidiorum culta frequenter occurrit.

GIESEKIA (L. Mant. 2. App. p. 362).

Flores hermaphroditi unibracteati. Perigonium quinquepartitum laciinis herbaceis margine membranaceis et saepè coloratis. Stamina 5-12 in liberia solitaria vel 2-3-na laciinis alterna, filamenta a basi latiore subulata. Carpella 5 vel abortu 3-4 in toro papillifero sessilia distincta verticillata uniovulata, ovulo basisixo campylotropo, stylus ex angulo centrali carpelli apice recurvus facie interna stigmatosus. Utriculi 4-5 in perigonio liberi monospermi indehiscentes. Semen verticale crustaceum rostellatum, albumen centrale, embryo annularis

periphericus. Herbæ (in nostris) annuæ prostratæ, foliis suboppositis, cymulis oppositifoliis. Genus nunc propter inflorescentiam non racemosam inter *Ficoideas* (ubi Fl. Orient. Vol. II, p. 764 omissum fuit) potius militat.

1. G. pharnaceoides (L. loc. cit.) annua glaucescens collo ramosa ramis procumbentibus dichotome ramosis, foliis breviter petiolatis ellipticis vel elliptico-lanceolatis obtusis, cymis ad dichotomias et terminalibus foliis supremis involucratis et superatis subsessilibus laxiuscule 10-20-floris, pedicellis flores minutos virides subæquantibus vel longioribus, perigonii laciniis ovatis concavis acutis subnerviis, staminibus quinis, carpellis ovato-sphæricis subcompresis verruculosis ♂. Ic. Roxb. pl. Corom. tab. 483. — *Pharnaceum occultum* Forsk. Eg. Arab. p. 58.

Hab. in arenosis Egypti (ex Schw. Beitr.), Affghaniæ ad Chuch (Griff!), Belutschiae (Stocks!).

Ar. Geogr. Senegalia, Nubia, Abyssinia, Arabia tropica, India, P. B. spei.

ORD. XCVIII. CYNOCRAMBEÆ.

(Necs Gen. Germ. fasc. 8. — Alph. d.c. Prodr. XVII, p. 32.)

Flores monoici. *Masc.* Perigonium bipartitum laciniâ anticâ et posticâ. Stamina 8-20, vel pauciora, imo calyci inserta. *Fem.* Perigonium tubuloso-infundibuliforme ovario adnatum limbo supero clavato-urceolato 4 vel 2-lobo mox caduco. Ovarium inferum uniloculare autice tumens et subglobosum unde perigonium laterale. Stylus lineariclavæformis lateralis ex ovario tumefacto fere gynobasicus a medio papillosus, ovulum unicum basilare campylotropum. Fructus recens subcarnosus tandem siccus nucularis indehiscens. Semen globosum testâ tenerâ, embryo hippocrepicus verticalis in axi albuminis farinacei. — Herba tenera stipulata. — Affinitas Ordinis multiplex, facies et characteres plures *Spinaclearum* inter *Salsolaceas*, sed ovarium omnino inferum, folia stipulata, etc.; ab *Urticaccis* differt ovario infero, stylo indiviso, a *Tetragoniaceis* ovulo erecto, etc. Ex cl. Alph. d.c. *Salsolacearum* tribus.

CYNOCRAMBE (Diosc. — Tournef. Cor. p. 52, tab. 485.
— *Theligonum* L. Gen. 864).

Character Ordinis. *Masc.* Perigonii laciniæ æstivatione valvatim admotæ dein revolutæ. Staminum filamenta tenuiter capillacea, antheræ longe et anguste lineares biloculares versatiles. *Fem.* Perigonii limbus hyalinus. Ovarium breviter stipitatum. Stylus post anthesin

cum perigonii limbo deciduu. Nuculæ pericarpium crassiusculum verruculis aciculas crystallinas occludentibus et lineis longitudinalibus obsitum.

1. **C. prostrata** (Görtn. Fr. I, p. 362, tab. 75) annua a collo sœpe ramosa ramis patulis flaccidis, foliis petiolatis ovato-rhombeis margine scabridis inferioribus oppositis superioribus ob alterum paris diminutum vel abortivum alternis, stipulis membranaceo-hyalinis integris vel dentatis foliorum oppositorum connatis alternorum amplexicaulibus, floribus minutis virentibus sessilibus, masculis geminis versus summos ramos foliis alternis oppositis nudis, feminis ad foliorum præsertim oppositorum axillas subternis bracteolatis ⊖. *Theligonum Cynocrambe* L. Sp. 1411. — Fl. Graeca, tab. 94t.

Hab. ad muros, in saxosis umbrosis regionis inferioris et montane, Græcia et ejus insulæ (Marg! Sprun! Heldr!), Thracia ad Byzantium (Griseb! Auch!), Anatolia littoralis occidentalis et australis (Fleisch! Bourg!), Syria littoralis (Ehrenb!) et ad Damascum (Gaill!) et Aleppo (Kyt!), Palestina (Boiss!), Cyprus (Ky!), Mesopotamia mons Sindjar (Haussk!).

Ar. Geogr. Regio mediterranea totius Europæ et Africæ borealis, insulæ Canarienses.

ORD. XCIX. SALSOLACEÆ.

(Moq. in D.c. Prodr. — *Chenopodeæ* Venten.)

Flores hermaphroditi polygami vel dioici. Perigonum herbaceum persistens 3-5-phylum interdum diphylum, phyllis plus minusve coalitis æstivatione imbricalis. Stamina quina vel pauciora hypogyna vel perigyna libera vel in cupulam coalita, staminodiis interdum interjectis. Discus hypogynus intra stamina vel nullus. Ovarium liberum vel inferne perigonio adnatum uniloculare uniovulatum ovulo basifixo, styli 2 rarius 3-4 discreti vel basi coaliti. Fructus indehiscens intra perigonum nunc immutatum nunc induratum nunc carnosum, nunc varie appendiculatum liber vel ejus tubo adhaerens. Pericarpium membranaceum vel crustaceum. Semen horizontale vel verticale (sœpe in eadem specie), testâ membranacâ vel crustacea. Embryo nunc circa albumen copiosum plus minusve perfecte anularis, nunc plano-spiralis albumine excentrico bipartito, nunc conico-spiralis albumine nullo; cotyledones angustæ, radicula hilo proxima. — Herbæ vel frutices, stipulis nullis, foliis oppositis vel alternis, floribus solitariis vel glomeratis axillaribus spicatis. Ordinem difficillimum parte longe maximâ ex notis, clavibus diagnosticis manuscriptis, summo labore præparatis et mihi benigne a cl. Prof. a Bunge communicatis, vel ex hujus *Anabasearum Revisione* (Pétersb. 1862) elaboravi. Viro æstimatissimo et amicissimo *Salsolacearum* peritissimo scrutatori gratias quam maximas agere mihi dulce est. In *Salicorniarum* descriptione egregiam monographiam cl. Ungern Sternberg pedetentim secutus sum.

SUBORDO I. CYCLOLOBEÆ Moq.

Embryo annularis vel hippocrepicus. Albumen copiosum centrale.

TR. I. CHENOPODIEÆ Moq.

Flores sæpissime hermaphroditæ homomorphi. Seminis integumentum crustaceum. Caulis continuus.

BETA (L. Gen. 310).

Flores hermaphroditæ tribracteolati. Perigonum quinquesfidum urceolatum basi semini subadbærrens tubo demum indurato. Stamina quinque annulo carnoso germen cingente inserta. Ovarium semiinferum, stigmata 2-3. Utriculus globosus perigonii tubo inclusus et basi adhærens. Semen horizontale. Embryo periphericus. — Herbæ annuae vel perennes, flores sessiles axillares vel terminales spicati solitarii vel 2-3 glomerulati basi connati et tunc fructum spurium exhibentes et conjuncti decidui.

1. **B. nana** (Boiss. et Heldr. Diagn. Ser. I, 7, p. 82) perennis glabra acaulis nana, radice verticali elongatâ, foliis omnibus radicalibus rosulatis oblongis acutiusculis integris in petiolum eis subæquilongum attenuatis, racemis e collo pedunculatis folia superantibus decumbentibus simplicibus laxis, floribus subsessilibus solitariis ad axillam bracteæ eis longioris oblongæ vel linearis sitis, perigonii herbacei laciniis oblongis demum carinatis sursum inflexis non membranaceo-marginatis, stigmatibus binis subulatis ♀.

Hab. in regione alpinâ Græciæ ad nivem deliquescentem, Taygetus loco Choupata dicto (Heldr!), Parnassus (Guicc! Heldr!). Fl. Aug., Sept.

Foliorum lamina 8-10 lineas longa 4-5 lata, racemi plures extrarosulares humi addressi 2-3-polices longi, perigonum fructiferum magn. seminis Cannabeos, tubo hemisphærico elevatim quinquecostato. Species insignis.

2. **B. vulgaris** (L. Sp. 322) annua vel perennans, caulis sub-simplicibus vel paniculatis, foliis radicalibus ovatis obtusis cuneatis interdum subcordatis, caulinis rhomboideo-ovatis, floribus 2-3-nis rarius solitariis glomerulatis, glomerulis in spicas longas tenues foliosas laxas vel aphyllas dispositis, perigonii herbacei laciniis oblongis apice cucullatis fructiferis carinatis et basi in tuberculum incrassatis, stigmatibus binis ⊖ et ⊙.

α. *typica*. Radix caulem unicum erectum edens, stigmata ovata. — *B. stricta* C. Koch Linn. XXII, p. 180.

Hab. in Egypto circa Kahiram (Schw!), in Babyloniam (Haussk!) et prob. alibi.

B. maritima. — Glabra rarius hirsuta. Caules e radice plures procumbentes, stigmata ovato-lanceolata. — *B. maritima* L. Sp. 322, Fl. Gr. tab. 254. — *B. Noëana* Bge in litt.

Hab. ad littora maritima Zacynthi (Sibth.), Atticae (Sprun! Heldr!), Ponti Euxini (Sibth.), maris Caspii (C.A. M!), Cyperi (Ky. 306!), Syriae littoralis (Auch. 2885! Bl! Gaill!), Egypti ad Alexandriam (Gaill!), in interioribus humidis Babyloniae ad Mohammera (Noë 78! et 910!).

Characteres quibus auctores *B. vulgare* (formam multo rariorem) et *B. maritiman* distinxerunt nempe radix annua vel perennans, caulis erectus unicus vel plures procumbentes fallaces et variabiles esse videntur. Am. Bge plantam ex Mohammera Babyloniae in litt. sub *B. Noëana* nomine distinxit ob lacinias perigonii fructu adpressas, stigmata magis ovata, etc. sed formas intermedias observo.

Ar. Geogr. Littora maritima Europæ borealis et australis, Africae borealis, Maderæ et insularum Canariensium.

3. ***B. trigyna*** (W. Kl. Pl. Rar. Hung. I, p. 34, tab. 35) perennis, radice longâ validâ fusiformi sâpe ramosâ, caulis erectis elatis angulato-sulcatis foliosis in paniculam pyramidatam lengam subaphyllam abeuntibus, foliis radicalibus longe petiolatis ovatis truncato-subcordatis subundulatis, superioribus breviter petiolatis deltaeis vel triangulari-lanceolatis acutis, spicis longis ante anthesin nutantibus aphyllis, floribus ternis glomerulatis lutescentibus, perigonii laciis oblongis obtusis concaviusculis late membranaceis, stigmatibus ternis linearibus ♂.

Hab. in cultis et pinguibus, Anatoliâ sept. (Wied! Auch. 2773!), Ponto Lazico ad Djimil (Bal!), Cappadocie monte Alidagh 5000' (Bal. exs. sub *B. longespicatâ*!), Armeniâ Turcicâ (Huet!), Tauriâ (Stev., Rehm!), prov. Caucasica (Ledebe).

Ar. Geogr. Hungaria.

4. ***B. lomatogona*** (F. et M. in Hohen. Talysch En. p. 124) perennis, caulis erectis elatis angulato-sulcatis in paniculas amplas aphyllas abeuntibus, foliis radicalibus longe petiolatis ovatis sâpe cordatis obtusis, caulinis superioribus subpetiolatis oblongo-lanceolatis et linearibus acutis, floribus solitariis sessilibus lutescentibus spicas longas laxas tenues formantibus, perigonii laciis oblongis obtusis concaviusculis albo-marginatis tandem conniventibus tubo fructifero quinquecostato, stigmatibus tribus linearibus ♂. *B. longespicata* Moq. Chenop. p. 13.

Hab. in campis regionis montanae et alpinæ, Olympus supra Broussam (Pichl!), Tmolus Lydie supra Philadelphiam et Caria in planicie Cybira (Boiss!), Lycia supra Elmalu (Bourg. 253!), Pisidia ad Buldur et Egirdir (Heldr!), Cappadocia (Ky. 192!), Taurus Cilicicus ad Gulek (Bal!), Aintab Syriae bor. (Haussk!), Armenia Turcica (Calv!), prov. Transcaucasica Talysch (Hoh!).

Ex am. a Bge sententiâ *B. lomatogonam* et *longespicatam* coniunxi, qui eas simillimas dicit; ego *B. lomatogone* specimina non vidi. Cl. Bge indicante

tubus perigonii fructiferi erronee sexcostatus ab auctoribus dicitur. Hæc species præter flores solitarios nec ternos a *B. trigynâ* facie simili mihi vix differre videtur.

5. **B. intermedia** (Bunge in litt.) perennis, caule erecto apice paniculato-ramoso, foliis radicalibus longe petiolatis ovatis sæpe subcordatis, superioribus oblongo-lanceolatis acutis, floribus spicatis solitariis vel 2-3-nis glomeratis, perigonii laciniis late hyalino-marginatis obtuse carinato-alatis in fructu erecto-conniventibus tubo quinquecostato, stigmatibus tribus lanceolato-linearibus divaricatis ♀.

Hab. in Anatoliâ prope Tokat, Amasia et Toptscha, inter Gheiwâ et Therakli (Wied!).

Planta elata facie *B. trigynæ* et *lomatogonæ* a quibus perigonii laciniis elevatim carinato-subalatis fere tantum differt. An hæc nota sat gravis?

6. **B. macrorrhiza** (Stev. in Mem. Soc. Mosq. III, p. 257) perennis, glabra, radice fusiformi validâ multicipiti, caulis prostratis, foliis coriaceo-carnosulis ovatis obtusissimis summis suborbiculatis, spicis simplicibus superne aphyllis, floribus ternis glomerulatis, perigonii laciniis oblongis obtusis patentibus demum carinatis, stigmatibus tribus linear-lanceolatis ♀.

Hab. in subalpinis ditionis Kubensis Caucasi prope Chinalug (Stev!), in districtu Khoi prov. Aderbidjan Persiae (Szov. exs. 342 ex Bge in litt.).

Vidi tantum specimen valde incompletum herbarii Steveniani hujus speciei de quâ judicium ferre non audeo.

CHENOPODIUM (L. Gen. 309).

Flores hermaphroditæ ebracteati. Perigonium quinquesidum vel partitum dorso inappendiculatum. Stamina quinque imo perigonio inserta. Stigmata bina. Utriculus depresso-globosus perigonio inclusus et ab eo liber. Pericarpium membranaceum tenuissimum. Semina horizontalia (interdum in nonnullis speciebus pauca verticalia). Embryo periphericus. — Herbæ floribus glomerulatis spicatis vel paniculatis.

SECT. I. CHENPODIASTRUM Moq. — Embryo perfecte annularis. Herbæ inodoræ vel foetidæ.

1. **C. polyspermum** (L. Sp. 321) annum inodorum viride vel rubellum interdum minute papillosum a collo sæpe ramosum caulis ramosis erectis vel diffusis, foliis alternis tenuiter petiolatis ovatis vel oblongis, rameis subsessilibus lanceolatis, floribus nunc glomeratis axillaribus ac in spicas terminales aphyllas dispositis, nunc cymulas divaricatas dichotomas axillares formantibus, perigonii laciniis membranaceis planis fructiferis patentibus, seminibus horizontalibus nitidis punctulatis ⊖. Ic. Fl. Dan. tab. 1153.

Hab. in cultis, glareosis prob. totius partis septentrionalis ditionis, Macedonia (Orph!), Byzantium (Sibth.), Pisidia ad lacum Beychelr (Heldr!), Pontus Lazicus (Bal!), Tauria (Stev.), Caucasus et Transcaucasia (Lede-b.).

Ar. Geogr. Europa borealis et media, Sibiria.

2. C. Vulvaria (L. Sp. 321) annum fætidissimum incano-cinerascens, caulis patent-ascendentibus ramosissimis, foliis alternis petiolatis rhombeo-ovatis integris, florum glomerulis racemiformibus axillaribus et terminalibus aphyllis, perigonii laciniis ecarinatis fructum tegentibus, semine horizontali punctulato margine acutiusculo ⊖. Ic. Fl. Dan. tab. 1152. — *C. olidum* Sm.

Hab. in ruderatis totius ditionis a Græciâ! ad Caucasum et prov. Transcaucasicas (Hohen!), Persiam medium (Haussk!). In australioribus montanum, Libanus (Ky. 341! Bl!), Egyptus.

Ar. Geogr. Europa omnis a Daniâ et Gothiâ ad Rossiam medium, Africa borealis.

3. C. ficifolium (Smith Brit. I, p. 276) annum virescenti-cinerascens, caule erecto striato ramoso, foliis alternis longe petiolatis strictis, inferioribus a basi cuneatâ oblongis vel oblongo-linearibus trilobo-hastatis dentatis lobo intermedio multo majore oblongo-lanceolato obtuso, superioribus linear-lanceolatis integris, florum glomerulis in racemos spiciformes axillares paniculam pyramidatam foliosam formantes congestis, perigonii laciniis acute carinatis fructum tegentibus, semine nitidulo nitido excavato-punctato margine obtuso ⊖. *C. viride* Curt. Lond. Fasc. 2, tab. 16 non L.

Hab. in cultis, Egyptus inferior circa Alexandriam (Gaill!), circa Byzantium (ex Sibth.).

Ar. Geogr. Anglia, Batavia, Gallia, Germania, Dalmatia, regio Danubialis, Rossia media et australis, Sibiria Altaica.

4. C. album (L. Sp. 219) annum albo-farinose vel viride, caulis erectis angulatis in paniculam strictam abeuntibus, foliis rhombeo-ovatis sinuatis vel dentatis rarius integris lanceolatis inferioribus petiolatis superioribus oblongo-lanceolatis acutis integris, florum glomerulis in racemos compactos vel interruptos simplices vel racemos basi exceptâ aphyllos dispositis, perigonii fructiferi clausi laciniis carinatis, seminis horizontalis rarius verticalis laevis nitidi margine acuto ⊖. *C. viride* L. Sp. 319 (forma viridis foliis sæpius integris, inflorescentiâ magis cymosâ). — *C. leiospermum* D.C. Fl. Pr. III, p. 392. — *C. concatenatum* Thuill.

Hab. in cultis et ruderatis regionis inferioris montanae et subalpinæ in totâ ditione a Græciâ! ad Egyptum! Persiam! et Afghaniam!

Specimina foliis sublobatis interdum cum *C. ficifolio* seminibus excavato-punctatis distincto confunduntur (v. gr. planta ex Assyriâ Ky. 450 b.).

Ar. Geogr. Totus orbis.

5. C. opulifolium (Schrad. in D.C. Fl. Pr. 5, p. 372) annum glauco-farinose, caule erecto vel ascendentem a basi sæpe ramoso,

foliis alternis petiolatis cuneatis subrotundo-rhombeis subtrilobis obtusis dentatis, superioribus conformibus diminutis interdum angustioribus, florum glomerulis in racemos vel cymas axillares paniculam strictam foliosam formantes dispositis, perigonii fructiferi laciniis carinatis fructum tegentibus, semine horizontali nitido subtiliter punctato margine obtuso ⊙. Ic. Vaill. Bot. tab. 7, fig. I.

Hab. in cultis et arenosis, Græcia in Atticâ et Cephallenia (Heldr!), regione media montis Kyllene (Heldr!), Syriâ littorali (Blî Gaill!) et circa Marasch (Haussk!), Transcaucasiâ (Hoh!), Persiâ boreali et occidentali (Buhse! Haussk!).

Ar. Geogr. Europa media et australis a Daniâ et Gothiâ ad regionem Danubialem, Africa borealis, Abyssinia.

6. C. murale (L. Sp. p. 318) annum viride, caulis ascendentibus angulato-sulcatis a basi ramosis, foliis omnibus petiolatis alternis rhomboe-ovatis vel oblongis acutis inæqualiter et acute dentatis, cymis axillaribus divaricatis laxiusculis, perigonii fructiferi clausi laciniis subcarinatis, seminibus opacis punctato-rugosis margine acutis ⊙. Ic. Fl. Dan. tab. 2048.

Hab. in cultis et ruderatis ubique a Græciâ et Egypto ad prov. Caucasicas, Persiam! Arabiam Petream! et Mascatensem! Affghaniam!

β. microphyllum. — Folia diminuta brevius dentata.

Hab. in littoribus marinis Eubeæ (Heldr!), Peninsulâ Athoâ (Friv!), ad Alexandriam Egypti (Gaill!).

Ar. Geogr. Totus orbis terrarum.

7. C. hybridum (L. Sp. 319) annum viride, caule erecto angulo parum ramoso, foliis majusculis petiolatis trianglari-ovatis truncatis vel subcordatis profunde sinuato-lobatis lobis triangularibus acutis, racemis patentim compositis erectis paniculam terminalem brevem subaphyllam formantibus, perigonii fructiferi laciniis carinatis patentibus, semine horizontali punctato-ruguloso margine acutiseculo ⊙. Ic. Fl. Dan. tab. 2049.

Hab. in ditione ut videtur rarius, Tauria (Stev.), Caucasus et Transcaucasia (M.B., C.A. Mey., Ledeb.).

Ar. Geogr. Europa borealis et media ad Galliam, Italiam, regionem Danubialem et Rossiam medium usque, Sibiria Uralensis, regio Himalaica, Africa borealis, America borealis.

8. C. urbicum (L. Sp. 318) annum, caule erecto angulato procero stricte et parce ramoso, foliis petiolatis erectis supra viridis nitidis subtus interdum pulverulentis carnosulis, inferioribus et mediis triangularibus basi in petiolum protractis acutis acute et inæqualiter dentatis, superioribus rhombeis et lanceolatis subintegris, florum glomerulis dense vel interrupte spicatis, spicis longis compositis elongatis cauli strictissimis thyrsum longum angustum inferne foliosum formantibus, perigonii fructiferi laciniis non cari-

natis fructu non adpressis, semine punctulato obtuso ⊖. Ic. Fl. Dan. tab. 1148. — *C. intermedium* Koch (forma foliorum dentibus triangulari-lanceolatis acuminatis).

Hab. in cultis ruderatisque, prope Athenas (Heldr!), in Macedonia et Thraciâ (Griseb.), Byzantii (Sibth.), Anatoliâ in Lycaoniâ ad Koniah (Heldr!), Cataoniâ (Haussk!). Tauriâ, Caucaso et Transcaucasia (Ledeb.), Kurdistanâ Persicâ 5800' (Haussk!), Persiâ (Belang.).

Ar. Geogr. Europa tota a Britannia et Scandinavia ad Rossiam, Sibiria, Dahuria.

9. **C. glaucum** (L. Sp. 320) annum, caulis a basi patentim ramosis erectis vel ascendentibus, foliis supra pallide virentibus subtus valde glauco-farinosis petiolatis oblongis sinuato-dentatis obtusis, florum glomerulis in racemos spiciformes axillares aphyllos foliis subbreviores dense congestis, perigonii fructiferi laciniis carinatis imperfecte elansis, seminibus levibus lucidis acute marginatis aliis verticalibus aliis numerosioribus horizontalibus ⊖. Ic. Fl. Dan. tab. 1151. — *Blitum glaucum* Koch D. Fl.

Hab. in ruderatis humidis, fimetis, Macedonia (Friv. ex Griseb.), Tauria, Caucasus et Transcaucasia (M.B., Hoh.), Turkestania ad Buchara (Lehm.), Afghanistan (Griff.).

Ar. Geogr. Europa fere omnis, Sibiria, Regio Himalaica.

SECT. II. BOTRYOIDES C.A. Mey. — Embryo ad $\frac{2}{3}$ annularis albumen imperfecte includens. — Herbæ aromaticæ.

10. **C. Botrys** (L. Sp. 320) annum glanduloso-pubescent, caule erecto rigidulo ramis erecto-patulis, foliis inferioribus petiolatis ovatis vel oblongis obtuse sinuato-subpinnatifidis superioribus diminutis spathulato-lanceolatis summis integris, cymis axillaribus brevibus erectis racemos longos elongatos angustos strictos subaphyllos formantibus, perigonii pubescenti-glandulosi laciniis ecarinatis, seminibus horizontalibus subglobosis levibus margine obtuso carinulato ⊖. Ic. Fl. Graec. tab. 253.

Hab. in arenosis et glareosis totius ditionis præsertim in regione montana a Greciâ, Macedonia et Thraciâ ad Libanum (Bl!), Egyptum infer. (Gaill!), Arabiam petream (Bové!), Persiam borealem (Auch!), Turkestaniam (Lehm.), regnum Cabulicum (Griff!).

Ar. Geogr. Europa australis a Belgio, Galliâ, Helvetiâ et Germaniâ australibus ad Rossiam medium, Sibiria Uralensis, Africa borealis et australis, America borealis.

11. **C. foetidum** (Schrad. Mag. Ges. Berl. 1808, p. 79) annum subglanduloso-pubescent, caule erecto sulcato-striato subsimplici, foliis petiolatis oblongis obtuse pinnatifido-sinuatibus superioribus conforniibus diminutis, cymis axillaribus iteratim divaricatiisque bisidis remotifloris foliis brevioribus paniculam strictam racemiformem

formantibus, perigonii hispiduli laciniis carinatis non clausis, seminibus horizontalibus globoso-lenticularibus margine obtuso ⊖.

Hab. in prov. Caucasicis prope Derbend (Eichw.), prope Kutais (Rupr!). An spontaneum?

Ar. Geogr. Abyssinia, Arabia tropica, America tropica.

12. C. ambrosioides L. Sp. 302) annum puberulum, caule sulcato erecto stricte ramoso, foliis breviter petiolatis oblongis et lanceolatis obtusis obsolete sinuato-dentatis subtus glandulosis superioribus integris, racemis spiciformibus elongatis saepius compositis paniculam longam angustam foliosam formantibus, perigonio fructifero clauso ecarinato, seminibus horizontalibus lœvibus lucidis obtusis ⊖. Ic. Barr. 1185.

Hab. ex Americâ oriundum nunc multis locis ad vias, in cultis subspontaneum, Peloponnesus (Pichl!), Attica et Cephalonia (Heldr!), Creta (Raul!), Syria littoralis ad Berythum (Bl!), Affghania (Griff.).

Ar. Geogr. Ex Americâ boreali et tropicâ in Europam australem, Africam borealem et australem, Indianam orientalem, Novam Hollandiam introductum.

BLITUM (L. Gen. 14 cum Emend.).

Flores hermaphroditi ebracteati. Perigonium 3-5-fidum vel partitum dorso inappendiculatum. Stamina 4-5 imo calyce incerta. Stigmata bina. Utriculus a perigonii liber depresso-globosus pericarpio membranaceo. Semina verticalia vel in sectione priori terminale uniuscujusque glomeruli horizontale. Embryo periphericus. — Herbæ facie *Chenopodiæ*.

SECT. I. ORTHOSPORUM C.A. Mey. — Perigonium fructiferum exsuccum. Flos terminalis glomeruli pentandrus semine horizontali, cæteri 2-3-andri semine verticali.

1. B. Bonus Henricus (L. Sp. 318 sub *Chenopodio*) perenne, caulis crassinsculis erectis sulcatis foliosis, foliis viridibus petiolatis trianguli-hastatis vel sagittatis integris undulatis, summis diminutis rhombis acuminatis, florum glomerulis in racemos densos aphyllos paniculam angustam strictam basi tantum foliatam formantes dispositis, perigonii laciniis non carinatis hand clausis, semine punctulato nitido margine obtuso ♀. *Bl. Bonus Henricus* C.A. Mey. Fl. Alt. I, p. 11. — *Agathophyllum Bonus Henricus* Moq.

Hab. ad vias, firmeta, in regione subalpinâ, mons Olenos Peloponnesi (Heldr!), Parnassus (Orph!), mons Delphi Eubœa (Heldr!), Macedonia (Friv!).

β. *nanum*. Caules 2-3-pollicares decumbentes.

Hab. in regione alpinâ Taygeti (Pichl!), Olympi Thessali (Anch. 2754!).

Ar. Geogr. Europa borealis et media in australioribus alpinum, Sibiria Uralensis.

2. B. rubrum (L. Sp. 318 sub *Chenopodio*) annum, caule a basi ramoso erecto vel procumbente, foliis petiolatis viridibus vel rubellis nitidis triangulari-rhombeis cuneatis sinuato-dentatis dentibus lanceolatis, spicis fere ex omnibus axillis compositis densifloris foliolosis strictis folio axillari brevioribus, perigonio fructifero imperfecte clauso non carinato, seminibus purpureis minimis nitidis punctulatis margine obtusis ⊖. Ic. Fl. Dan. tab. 1149. — *B. rubrum* Rehb. — *B. polymorphum* C.A. Meyer.

Hab. in ruderatis, ad pagos, circa Byzantium (Sibth.), in prov. Caucasicis (C.A. Mey!), Turkestaniā (Lehm.).

β. subintegrum. — Folia minora integra vel subintegra.

Hab. in Armeniā (Calv!), Babyloniā prope Bagdad (Noë!).

Ar. Geogr. Europa fere tota, Sibiria.

SECT. II. EUBLITUM. — Perigonium fructiferum baccatum. Stamen (in nostro) unicum. Semen verticale.

3. B. virgatum (L. Sp. 7) annum glabrum, caule ascendentē vel erecto ramoso ramis albidis rigidis, foliis viridibus petiolatis oblongo-triangularibus vel hastatis acuminatis profunde et acute dentatis, floralibus sensim diminutis lanceolatis glomerulos superantibus, glomerulis globosis ad omnes axillas sessilibus solitariis spicam longam laxam rigidam foliolosam formantibus, perigonio fructifero baccato rubro haud clauso ecarinato, semine lăviusculo margine obtuso ⊖. Ic. Lam. III. tab. 5.

Hab. in rupestribus regionis montanæ et alpinæ, Anatolia borealis et Pontus (Tchih!), Lycia supra Adalia (Bourg!), Taurus Lycaonicus (Heldr!) et Cilicieus (Bal!), Cataonia mons Berytdagh (Haussk!), Armenia Turcica (Huet!), Caucasus et Transcancasiania (C.A. M., Hoh!), Persia borealis in iugō Elbrus (Auch! Ky!), Persia australis et Kurdistania (Haussk!), Libanus ad Eden et Cedros (Boiss! Bl!), regnum Cabulicum (Griff!).

Ar. Geogr. Hispania media, Gallia, Dania et Suecia, Helvetia, Germania australis, Italia borealis, regio Danubialis, Rossia, Sibiria, India, Africa borealis.

TR. II. ATRIPLICEÆ C.A. Mey.

Flores diclini, monoici vel dioici, interdum polygamni. Seminis integumentum crustaceum vel coriaceum. Caulis continuus.

SPINACIA (L. Gen. 112).

Flores dioici ebracteati. *Masc.* Perigonium 4-5-partitum herbaceum laciniis non appendiculatis. Stamina 4-5 imo perigonio inserta filamentis capillaribus. *Fem.* Perigonium subglobosum apice 2-3-dentatum ovarium includens, fructiferum coriaceum nunc inerme inappendiculatum

nunc corniculis spinosis 2-3 e tubo ortis accrescentibus auctum. Stigmata 4 longa filiformia basi connata. Nucula perigonio adhæreens compressa. Semen verticale testâ tenui. Embryo periphericus albumen farinaceum cingens. — Herbæ annuæ, flores masculi in spicas terminales aphyllas dispositi, feminei ad foliorum axillas glomerati.

1. **S. tetrandra** (Stev. in Mém. Nat. Mosc. II, p. 182) annua glabra pumila, caulis a collo sæpius pluribus erectis vel ascendentibus, masculis in spicam interruptam simplicem vel ramoso-paniculatum abeuntibus, femineis simplicibus, foliis radicalibus longe petiolatis ambitu oblongis vel lanceolatis obtuse vel acute pinnatilobis partitis vel sinuato-repandis rarius integris, caulinis plantæ masculæ linear-lanceolatis dentatis vel integris subsessilibus fæmineæ sessilibus cordato-semiamplexicaulibus ovatis obtusissimis, glomerulis fæmineis axillaribus, perigonis fructiferis in glomerulo inter se con-natis spinoso-bicornutis, spinis late triquetris horizontalibus subulato-attenuatis pungentibus ♂. *S. minor* C. Koch Linn. XVII, p. 311, (forma foliis radiculibus integris).

Hab. in argillosis salsuginosis et pratis siccis Iberie Caucasicæ (Stev., Hoh!), Armeniae Rossicæ (Buhsel), ad littora Caspia prope Baku et Sallian (C.A. Mey!), Persiæ boreali (Szov!), Turkestania ad Buchara (Lehm.), Babyloniam ad Euphratem (Oliv! Hausskl!), Affghaniam (Griff.). Incolis olus *Schamum* dictum.

Planta 3-6-pollicaris, fæmineæ caules demum incrassati jam ex iuso collo interdum fructiferi. *S. oleracea* Mill. = *S. spinosa* Mœnch *inermis* Mœnch. = *S. glabra* Mill. sunt plantæ majores foliis 1-2 mm. in variis ditionis locis indicate sed non vere spontaneæ et quarum patria ignota. Prior differt a *S. tetrandra* glomeruli fructibus spinosis inter se liberis, posterior fructibus globosis inermibus liberis.

ATRIPLEX (L. Gen. 4153. — *Obione* Gærtn. — *Halimus* Wallr.).

Flores monoici vel subdioici, rarius hermaphroditi. Perigonium masc. vel hermaph. quinquepartitum, femineum phyllis binis liberis vel in thecam compressam plus minus coalitis constans. Stamina 3-5 in basi perigonii inserta. Ovarium ovatum compressum. Styli 2 setaceo-filiformes. Utriculus membranaceus phyllis perigonii inclusus. Semen verticale (in fl. herm. horizontale) testâ crustaceâ. Embryo periphericus radiculâ basin vel apicem ovarii spectante. — Herbæ, suffrutices vel frutices.

Specierum Orientalium distributio.

SECT. I. **DICHOSPERMUM** Dumort. — Flores polygami, feminei ad basin usque bipartiti semine verticali, hermaphroditi pauci quinquepartiti semine horizontali. Radicula infera ascendens.

A. hortense, nitens.

SECT. II. SCHIZOTHECA C.A. Mey. — Flores monoici raro subdioici, feminei bifidi vel bipartiti semine verticali, masculi quinquepartiti.

§ 1. **Annua.**

* Radicula infera ascendens.

+ Perigonii phylla ad basin usque discreta.

A. crassifolium, littorale, hastatum, patulum, dimorphostegium.

+ + Perigonii phylla ad medium vel altius connata

A. Tataricum, roseum.

** Radicula apicalis. Phylla connata.

A. Turcomanicum, thunbergiæfolium, pedunculatum, Moneta, Flabellum.

§ 2. **Perennia suffruticosa vel fruticosa.**

* Radicula apicalis. Perigonii phylla plus minus alte connata.

+ Folia opposita.

A. Chanykowii, verruciferum, pertulacoides.

+ + Folia alterna.

A. Belangeri, Palestinum, Alexandrinum, crystallinum, leucocladum, coriaceum.

** Radicula basilaris.

A. Halimus, Persicum, Griffithii, farinosum.

SECT. I. DICHOSPERMUM.

1. **A. hortense** (L. Sp. 1493) annum, caule erecto ramoso angulato, foliis alternis viridibus concoloribus cordato-triangularibus integris vel sinuato-dentatis petiolatis, superioribus ovato-lanceolatis, racemis ex axillis superioribus terminalibusque paniculatis elongatis laxiusculis, perigonii feminei fructiferi phyllis binis subliberis valde auctis integris membranaceis reticulato-venosis ovato-orbiculatis obtusis interdum mucronatis ⊖. Ic. Schk. Handb. t. 349.

Hab. in cultis, Byzantii (Silth.), in peninsula Athoā (Pichl!), Atticā a Phalerum (Heldr!), Anatoliā in pomeriis prope Koniah (Heldr!), prov. Caucasicae (Ledeb.).

A. microtheca Moq. est hujus varietas perigonio fructifero minore.

Ar. Geogr. Scandinavia, Rossia, Sibiria; cæterum sœpe cultum et area difficile delimitanda.

2. A. nitens (Rebent. Prodr. Neomarch. p. 126) annum, caule erecto angulato ramoso, foliis alternis petiolatis supra viridibus subtus argenteo-lepidotis cordato-triangularibus sinuato-dentatis rarius integris, superioribus anguste triangulari-lanceolatis longe acuminatis, racemis ex axillis superioribus terminalibusque paniculatis elongatis laxiusculis, perigoni feminei fructiferi phyllis binis subliberis auctis membranaceis reticulato-venosis integris ovatis obtusis vel acutiusculis ⊖. Ic. Schk. tab. 348, fig. 1. — *A. acuminatum* W. K. Pl. rar. tab. 103. — *A. Aucheri* Moq. Chen. En. p. 51.

Hab. in cultis, Cappadocia ad Karahissar (Ball!), Syria circâ Damascus et Palestina (Gaill!), Babylonia ad Bagdad (Noël!). Persia occidentalis (Auch. 3099!), Tauria, Caucasus et Transcaucasia (Ledeb.).

An ab *A. hortensi* sat distinctum?

Ar. Geogr. Germania, Regio Danubialis, Rossia, Sibiria.

SECT. II. SCHIZOTHECA.

§ 1. Annua.

* Radicula infera ascendens.

3. A. crassifolium (C.A. Mey. in Ledeb. Fl. Alt. 4, p. 309) annuum lepidoto-canescens, caule decumbente vel erecto flexuoso ramoso, foliis alternis carnosis petiolatis inferioribus ovato-oblongis obtusis sinuato-dentatis interdum subhastato-trilobis, supremis lanceolatis integris, florum glomerulis densissime spicatis spicis sœpe interruptis aphyllis stricte paniculatis, perigonii viridis herbacei fructiferi phyllis carnosis rhombeo-triangularibus hastato-trilobis margine denticulatis disco gibbis et lœvibus vel tuberculatis semine 1½-2-plo longioribus ⊖. Ic. Ledeb. tab. 42.

Hab. in salsuginosis deserti Karakum Turkestaniae (Lehm. ex Bge), Affghania (Griff. 1743 ex Bge).

A sequentibus foliis èt perigonis carnosis nec membranaceis distinctum.

Ar. Geogr. Sibiria Altaica.

4. A. littorale (Linn. Sp. p. 1494) annum viride, caule erecto flexuoso ramoso vel simplici, foliis membranaceis alternis petiolatis erectis anguste linear-i-lanceolatis vel linearibus basi non dilatatis integris vel denticulatis, florum glomerulis in spicas interruptas stricte paniculatas dispositis, perigonii fructiferi sessilis phyllis parvis subcoriaceis rhombeo-triangularibus integris vel rarius basi 1-2-dentatis disco muricatis rarius lœvibus ⊖. Ic. Fl. Dan. tab. 1287. — *A. marinum* L. Mant. II, p. 300.

Hab. in salsis maritimis et interioribus, Cappadocia prope Cæsaream (Ball!), Transcaucasia ad Caspium (C.A. M.) et prob. alibi.

Ar. Geogr. Scandinavia, Anglia, Gallia, Italia borealis, Dalmatia, regio Danubialis, Rossia, Sibiria, America borealis.

5. **A. hastatum** (L. Sp. 1494) annum viride vel lepidoto-canescens, caule erecto patule ramosissimo, foliis petiolatis alternis et oppositis inferioribus triangulari-hastatis basi truncatis integris vel sinuato-dentatis lobis basilaribus horizontalibus, superioribus hastato-lanceolatis, summis integris, florum glomerulis in spicas axillares et terminales interruptas inferne foliolosas superne imitas dispositis, perigonii fructiferi sessilis phyllis triangulari-deltoides inferne margine 1-2-dentatis facie levibus vel tuberculatis ♂. Ic. Schk. 348. — Rehb. Ic. I, tab. 16. — *A. patulum* Gris. Spic II, p. 308. — *A. triangulare* Willd. — *A. micranthum* C.A. Mey. in Led. Pl. Alt. IV, p. 308. — Ledeb. Ic. tab. 43.

Hab. in ruderatis et ad vias, Macedonia et Thracia (Griseb.), Tauria et Caucasia (Ledeb.) et prob. alibi.

A. salinum Wallr. — Sæpius procumbens lepidotum plus minus canescens vel canum, folia minora angustiora hastata sed integræ, perigonum diminutum. — *A. microcarpum* W. K. tab. 250. — *A. Græcum* Willd. Sp. IV, p. 958 et herb. (Specimen non fruticosum ut ait cl. auctor, sed inferne induratum!)

Hab. formâ typicâ latifoliâ et virente in ditione multo vulgatus, Græcia! et ejus insulæ! Anatolia! Syria (Bl! Gaill!), Egyptus (Samar! Guic!), regio Caucasia et Transcaucasia (Led., Hohen!), Persia (Haussk!), Afghanistan (Griff. 1737 ex Bge).

Ar. Geogr. Europa tota, Sibiria.

6. **A. patulum** (L. Sp. 1494) annum virenti-subcanescens, caule erecto clato patule ramosissimo, foliis breviter petiolatis alternis vel oppositis inferioribus a basi attenuatâ anguste oblongo-lanceolatis subintegratis, superioribus linearibus acutis, florum glomerulis in spicas interruptas rigidulas axillares et terminales basi solum foliolosas dispositis, perigonii phyllis hastato-rhomboideis basi cuneatis integris facie levibus rarius tuberculatis ♂. Ic. Schk. 347. — *A. Tataricum* Auct. non L. Sp. — *A. oblongifolium* W. K. pl. rar. tab. 211.

Hab. in ruderatis et ad vias prob. ubique, habeo ex Atticâ (Heldr!), Arianeâ (Haussk!), Kurdistaniâ!, prov. Caucasicis! Turkestaniâ (Lehm.).

Ar. Geogr. Europa tota, Sibiria, Africa borealis.

7. **A. dimorphostegium** (Kar. et Kir. Enum. Song. N° 714) annum, collo pluricaule caulibus herbaceis brevibus crassiusculis atbidis decumbentibus vel ascendentibus dichotome ramosissimis, foliis alternis membranaceis papillis crystallinis subtus presertim cinereis petiolatis a basi truncatâ vel breviter cuneatâ ovatis obtusiusculis integerrimis, floribus aliis ad axillas inferiores 2-3-nis perigonii fructiferi stipitati reflexi phyllis membranaceis orbiculari-cordatis dorso levibus margine repando-denticulatis, aliis in spicas

terminales breves foliis minoribus interstinetas congestis perigonii fructiferi multo minoris phyllis triangulari-ovatis dorso cristatis ⊖. *A. bracteosum* Trautv.

Hab. in arenosis salsis desertorum, Arabia petrea loco Hauara (Schimp. 212!), ad meridiem Gaza (Boiss!), Syria (Auch. 2758), Persia orientalis ad Schahrud (Bge!), deserta inter Djendack et Enarek (Buhse!), Turkestanica occidentalis regnum Cabulicum (Griff!), Belutschia ad Pushtkhana (Stocks!).

Caules semipedales, phylla florū inferiorum 3 lineas, superiorum vix 2 lin. diam. lata.

Ar. Geogr. Africæ borealis, Songariæ deserta.

8. A. Tataricum (L. Sp. p. 1053 ex Aschers. Ind. Sem. Berol. 1872, non Schkuhr nec Koch) annum deum saepe induratum sèpius lepidoto-canum, caulis erectis vel diffusis ramosissimis, foliis breviter petiolatis alternis inferioribus oppositis hastato-deltoideis dentatis incisis vel laciniatis rarius integris. superioribus lanceolatis et lanceolato-linearibus, florū glomerulis in spicas præter basin nudas elongatas inferne saepe interruptas congestis, perigonii fructiferi phyllis a basi ad medium cuneatis et albo-cartilagineis rhombis vel subtrilobis dentatis vel subintegris ⊖. *A. laciniatum* Koch Syn. —

ren. et Godr. Fl. Fr. — Fenzl Fl. Ross. et mult. Auct. non L. — *A. incisum* M.B. Taur. Cauc. III, p. 641. — *A. recurvum* Urv. Enum. N° 239. — *A. Grecum* Fl. Græc. X, tab. 963 non Willd. — *A. Veneratum* Willd. Sp. IV, p. 962.

Hab. in arenosis maritimis, desertisque salsis, Græcia et ejus insulæ, Macedonia (Orph!), Thracia (Friv!), Anatolia occid. (Pichler!), Tauria et Transcaucasia (M.B., Led., Hohen!), Egyptus inferior (Husson!), ditio Fayum (Kralik!).

β. virgatum. — Caules tenuius virgati albidi cum ramis erectis rigiduli, spicæ elongatæ graciles, folia oblongo-lanceolata vel lanceolato-linearia parce et breviter dentata sèpissime integra et basi cuneata nec hastata. — *A. Olivieri* Moq. Chenop. p. 52. — *A. pruinatum* Sieb. exs. — *A. lasianthum* Boiss. Diagn. Ser. I, 12, p. 95 (forma alpina pumila minus cana).

Hab. in ruderatis et salsis interioribus in regionem alpinam saepe ascendens, Armenia Turcica ad Baibut (Bourg!) et ad Van (Ky. Suppl. 563!), montes Lyciæ supra Elmalu (Bourg!), Syria ad Marasch (Haussk!) et in planicie ad occasum Antiochia (Boiss!), Libanus supra Eden (Boiss!), Palestina (Sieb!), mons Gara Kurdistaniae (Ky. 353!), montes Avroman et Schahu (Haussk!), cacumina montis Kuh Daëna Persiæ australis (Ky. 714!), Persia ad Teheran (Oliv! Auch. 3191!), Turkestanica deserta (Lehml!), Eamdem formam e Songariâ! India borealis! Tibetia! vidi.

Cl. Ascherson loc. cit. observante *A. laciniatum* L. Ed. I, p. 1053 littorum Europæ boreali-occidentalis incola = *A. arenarium* Woods = *A. farinosum* Dumortier est species inflorescentia foliosâ diversa, hanc notâ *A. roseo* valde accedens et quam ego pro hujus formâ boreali fructûs phyllis membranaceis minus cartilagineis facile haberem.

Ar. Geogr. Germania, regio Danubialis, Rossia media et australis, regio Mediterranea totius Europæ et Africæ borealis, Songaria, Sibiria.

9. **A. roseum** (L. Sp. 1493) annum demum induratum canescens vel canum, caulis erectis vel ascendentibus divaricatum ramosissimis, foliis breviter petiolatis alternis inferioribus ovato vel trianguli-rhombeis inaequaliter sinuatis vel dentatis, superioribus ovato-oblongis, florum glomerulis sessilibus axillaribus dissitis racemos foliosos formantibus, perigonii fructiferi phyllis ad medium usque connatis et cartilagineo-induratis trianguli-rhombeis acutiusculis angulatis vel dentatis disco levibus vel tuberculatis ♂. Ic. Schk. tab. 350 — Fl. Dan. tab. 1284. — *A. album* Scop.

Hab. in ruderatis et sabulosis maritimis regionis inferioris et montanae, Græcia et Cyclades (Heldr! Sart!), Macedonia et Thracia (Griseb.), Byzantii (Pichl!), Anatolia prope Koniali (Heldr!), Tauria et prov. Transcaucasicæ (C.A. Mey., Hohen.), Syria littoralis et interior (Bl! Gaill!), Egypti ditio Fayumi (Kralik! Aschers. 431).

Ar. Geogr. Europa maritima fere tota, Germania ad Rhenum, regio Danubialis, Sibiria Uralensis et Altaica.

** Radicula apicalis.

10. **A. Turcomanicum** (F. et M. in Karel. Enum. Bull. Mosq. 1839 nomen tantum) annum lepidoto-canescens vel canum, caulis tenuibus albidis ascendentibus et procumbentibus flexuosis divarication ramosissimis ramis elongatis tenuissimis intricatis, foliis parvis sparsis subsessilibus cuneatis subhastato-rhombeis utrinque obtuse 3-4-dentato-sinuatis, superioribus minimis suborbiculatis obtuse dentatis, glomerulis 2-3-floris axillaribus et terminalibus remotis spicas laxissimas formantibus, perigonii sessilis cuneato-obtriangularis papilloso coriacei phyllis ultra medium inter se connatis apice truncatis late et breviter 3-5-dentatis facie leviusculis vel tuberculatis ♂. *Obione Turcomanica* Bge pl. Lehm. p. 451. — *Atr. leptocladium* Boiss. Diagn. Ser. II, 4, p. 74. — *A. Lehmannianum* Bge pl. Lehm. p. 451.

Hab. in desertis et ruderatis, Babylonia ad Bagdad et Mohammera (Noë 312! et 933!), Persia ad Kasbin (Buhse!), prope Schurab et Ispahan (Haussk!), Persia orientalis (Bge!), Turkestania prope Buchara (Lehm!), Turcomania (Karel!).

Basi denum induratum, pedale vel procerius intricatum ramosissimum ramis flexuosis rigidulis tenuibus, glomeruli inferiores invicem remotissimi superiores magis approximati, perigonium fructiferum sessuilineam longum et apice latum.

11. **A. thunbergiæfolium** (Boiss. et Noë Diagn. Ser. II, 4, p. 74 sub *Obione*) annum, caulis erectis albidis tenuibus parce ramosis, foliis alternis membranaceis supra canescentibus subtus canis inferioribus petiolatis obtusissimis obtusissime auriculato-cordatis integris, superioribus subsessilibus hastato-triangularibus acutis subrepandis auriculis acutis, glomerulis 4-7-floris distantibus spicas laxas inferne tantum breviter foliosas stricte paniculatas formantibus,

perigonii sessilis cuneato-obtriangularis phyllis herbaceis ad medium usque connatis facie reticulatis apice truncatis denticulatis ⊖.

Hab. in pascuis Babyloniae prope Kutt (Noë 63!).

Planta 4-4½-pedalis, folia inferiora pollicem longa et lata, panicula parce et stricte ramosa semipedalis, perigonum sesquilineam longum et latum. Affine *A. Turcomanico* sed distinctum caule parum ramoso, foliis.

12. **A. pedunculatum** (L. Sp. p. 1675) annum lepidoto-canescens, caule pumilo erecto flexuoso a basi divaricatim ramoso, foliis alternis integerrimis carnosulis subpetiolatis oblongis vel lanceolato-spathulatis obtusis, glomerulis panicifloris in racemos breves ex axillis superioribus terminalesque dispositis, perigonii fructiferi longe pedicellati phyllis in thecam herbaceam obverse triangularem obtuse et divaricatim bilobam denticulo interjecto coalitis ⊖. Ic. Fl. Dan. tab. 304. — *Diotis atriplicoides* M.B. Taur. Cauc. II, p. 397. — *Obione pedunculata* Moq. — *Halimus pedunculatus* Wallr.

Hab. in maritimis salsis, Tauria (M.B.), Imeretia (Guld.).

Ar. Geogr. Anglia, Dania, Batavia, Gallia et Germania boreales, Bulgaria, Rossia media et australis, Turcomania, Songaria, Sibiria.

13. **A. Moneta** (Bge in litt.) annum albo vel sulfureo-farinosum, a basi ramosissimum diffusum humile, foliis crebris alternis cordato-suborbiculatis grosse et breviter sinuato-paucidentatis vel subintegris, spicis breviter paniculatis e foliis parum exsertis, perigonii fructiferi sessilis phyllis in unicum orbiculare integrum vel omnino coalitis vel marginibus membranaceis liberis, disco induratis crasse trinervibus inter nervos reticulatis ⊖.

Hab. in Persiae ditione Khorassan inter Riwed et Ssebsewar et ad Achmatabad non procul ab urbe Meschhed (Bge), in collibus Affghaniæ (Griff. 1750).

Specimina semipedalia, folia breviter petiolata e sicco carnosa pollicem et amplius longa et lata, superiora vix diminuta. Perigonium fructiferum 7-9 lineas latum et longum. Species fructu insignis.

14. **A. Flabellum** (Bge in litt.) annum virens vel albo-farinosum, caule erecto opposite ramoso, foliis plerisque oppositis cordatis argute dentatis vel 5-7-angulato-lobatis lobo terminali triangulari majore saepius in setam elongatam deciduum abente, inferioribus longe petiolatis superioribus sessilibus, supremis ad setam reductis, spicis paniculatis longis laxifloris, perigonii fructiferi stipitati phyllis omnino connatis in thecam transverse sublatiorem basi profunde cordatam lignosam radiatim septemnerviam æqualiter et obtuse septemlobam ad costas plicatam disco cristato-appendiculatam ⊖.

Hab. in montosis salsis ad orientem urbis Mesched inter Faz et Tabatküh prov. Khorassan (Bge), Turkestaniæ in valle Sarafschian prope Samarcand (Fedschenko).

Planta 1-1½-pedalis, folia inferiora 12-15 lin. longa et basi lata, spicæ graciles subaphyllæ. Perigonium fructiferum 4 lineas longum 5 latum. Species fructu insignis.

Ar. Geogr. Turkestaniæ orientalis.

§ 2. Perennia.

Radicula apicalis.

15. A. Chanikowii (Bge in litt.) fruticulosum patule et parce ramosum cano-farinosum, foliis oppositis subsessilibus ovatis acute sinuato-dentatis, perigonii fructiferi sessilis valde farinosi phyllis fere ad apicem usque connatis apice breviter herbaceo liberis patulis subtrilobis disco induratis et utrinque longitudinaliter verrucoso-bicristatis ♂.

Hab. ad littora lacūs Urmiensis Persiae occidentalis (Seidl.).

Specimen incompletum tantum vidi, gracile, pedale; facies *A. Turcomanica* sed basi fruticulosum, folia opposita 4-5 lineas longa, spicæ graciles nudæ, perigonium sesquilineam longum.

16. A. verruciferum (M.B. Taur. Cauc. II. p. 441) basi suffruticosum canum, caulis elongatis ascendentibus parce et stricte ramosis superne nudis, foliis petiolatis oppositis spatulato-ovatis vel oblongis obtusis integerrinis, superioribus linear-lanceolatis, glomerulis in spicas tenues laxiusculas paniculatas aphyllas dispositis, perigonio fructifero sessili subgloboso clauso undique verrucis obtusis dense obsesso ♂. Ic. Ledeb. Icones tab. 144. — *Obione verrucifera* Moq.

Hab. in salsis maritimis et interioribus, Cappadocia ad Cæsaream (Ball!), Tauria et Transcaucasia (M.B., Hohen!), Armenia Rossica ad Araxem (Szov!), Persia bor. (Buhse sub *Ob. portulacoides*!), prov. Aderbidjan (Szov!).

Ar. Geogr. Rossia australis. Songaria, Sibiria Uralensis ei Altaica.

17. A. portulacoides (L. Sp. 1493) basi suffruticosum lepidoto-argenteum, caulis elongatis basi procumbentibus dein ascendentibus ramis erectis, foliis oppositis breviter petiolatis basi attenuatis oblongis vel lanceolato-linearibus obtusis integerrimis, supremis parvis linearibus, glomerulis laxiuscule spicatis in paniculas terminales breves aphyllas dispositis, perigonii fructiferi canescens basi longe cuneati substipitati phyllis in thecam compressam obcuneatam levem apice trilobam coalitis dentibus brevibus triangularibus subæqualibus vel saepius intermedio minore mucroniformi ♂. Ic. Engl. bot. tab. 261. — *Obione portulacoides* Moq.

Hab. in maritimis salsis, Peloponnesus in Argolide (Sart!), Macedonia (Friv!), Byzantium (Cast! Janka!), Creta (Sieb!), Syria littoralis (Gail!), Egyptus ad Alexandriam (Samar!), Tauria (ex Ledeb. sed a Stevenio non visum).

Ar. Geogr. Europæ totius littora a Britanniâ et Daniâ ad Rossiam medium, Afria borealis.

18. A. Belangeri (Moq. in p.c. Prodr. XIII, p. 108) suffruticosum, caulis ascendentibus teretibus glabris ramosis ramis ascen-

dentibus, foliis alternis sessilibus parvis tenuibus supra glabris glancis subtus farinoso-subincanis rhomboideis obtusiusculis subsinuatis vel integris subtrinerviis nervis subtus puberulis, florum glomerulis axillaribus et terminalibus spicatis, perigonio fructifero in stipitem axi continuum abeundi persistenti. phyllis in thecam cuneato-spathulatam apice dilatatam acutiusculam et denticulatam lateraliter integrum disco coriaceam laevem nervosam coalitis 2.

Hab. in Persia (Bélanger N° 581), Affghania (Griff. 1747 ex Bge). Non vidi.

Caules pedem et amplius longi, folia 4-8 lineas longa 4-7 lata, thecae 2 lineas longae in duratae inferne cylindrico-conicae apice compressae foliaceae virescentes. Ex am. Bge qui fragmenta speciminis Belangeriani vidit perigonio fructifero persistente nec articulatum deciduo distinctissimum.

19. A. Palestinum (Boiss. Diagn. Ser. I, 12, p. 96) suffruticosum papilloso-canum ramosissimum dumulosum, caulis ascendentibus teretibus albidis crebre foliosis, foliis minutis sparsis a basi sessili attenuata oblongis et oblongo-linearibus obtusissimis integris vel obtuse sinuato-repandis. superioribus abbreviatis linearibus acutis, axillis saepe fasciculiferis, glomerulis sessilibus spicas terminales interruptas simplices aut paniculatas basi breviter foliolosas formantibus, perigonio fructifero sessili cano minuto oblongo-rhombeo basi cuneato. phyllis ad medium usque coalitis apice subherbaceis obtuse trianguli-ovatis facie coriaceis elevatim 3-4-tuberculatis 5.

Hab. in desertis Judeæ inter Hierosolymam et Jericho (Boiss!), ad basin collium Damasci (Boiss!), in fauce Ouadi Barrada (Gaill!), in deserto Palmyrae (Bl!), in deserto ad meridiem Gaza (Boiss!).

Caules semipedales crassiusculi rigiduli, folia majora 9-10 lineas longa 3-4 lata, fructus 2 lineas longus 1½ latus. Am. a Bunge in litt. fructus præcociiores majores magis herbaceos superioribus rariores indicat, sed eos in speciminibus meis frustra quæsivi; forsitan schedulae error.

20. A. Alexandrinum, suffruticosum adpresso canum ramosissimum dumulosum ramis diffusis, foliis subpetiolatis a basi cuneatâ ovatis oblongis obtusis obtuse et saepe obsolete sinuato-repandis margine subundulatis, superioribus abbreviatis, axillis fasciculiferis, glomerulis spicas terminales elongatas valde interruptas simplices vel parce ramosas præter partem inferiorem aphyllas formantibus, perigonio fructifero stipitato articulato minuto oblongo-rhombeo subtrilobo lobo terminali multo majore ovato. phyllis ad medium coalitis facie coriaceis acute cristato-tuberculatis 5.

Hab. in incultis et collibus arenosis ad Alexandriam Egypti (Ehrenb! Gaill! Samar! Aschers. 374 et 433! Letourne!).

Caules cum spicis tandem saepe semipedalibus 1-1½-pedales, folia 10-12 lineas longa tres lata, fructus 2 lineas longi stipe lineam longo suffulti. Haec planta quam pro *A. Palestini* forma prius habueram præter caules diffusos tenuiores ab eo fructibus stipitatis dorso echinatis differt. *A. parvifolium* Lowe et *A. Mauritanicum* Boiss. facie similia differunt fructu sessili formâ diverso.

Ar. Geogr. Regnum Tunetanum.

21. A. crystallinum (Ehrenb. MSS.) basi suffrutescens canum a collo dumulosum caulis brevibus diffusis vel ascendentibus, foliis minimis sessilibus orbiculatis integris, axillis fasciculigeris, florum glomerulis in spicis terminales tenues interruptas simplices ramulosas congestis, perigonio fructifero minimo ovato-rhombeo subtrilobo lobo terminali multo majore ovato, phyllis ad medium concretis disco parce cristato-tuberculato ♀. *A. glaucum* Moq. ex parte quoad pl. Ägyptiacam. — *A. rotundifolium* Boiss. in litt. non Moq.

Hab. in Egypto inferiore ad Alexandriam (Kralik! Samar!), ad Ramleh (Gaill!), inter Gabbari et Mariut (Duparq!).

Caules semipedales, folia 2-3 lineas diam. lata, fructus vix sessilineam longus. Facies omnino *A. glauci* Hispanici quod valde differt fructu duplo majore triangulari tuberculis crassis crebre obsito. *A. crystallinum* prius pro formâ brachyphyllâ *A. Alexandrinum* habui, sed præter foliorum formam fructus sessiles sunt nec stipitati, ambo tamen ulterius exploranda.

22. A. leucocladum (Boiss. Diagn. Ser. I, 12, p. 95) basi suffrutescens canum, caulis a collo numerosis ascendentibus albidis patule ramosis, foliis alternis a basi breviter cuneatâ vel truncatâ deltoideo-ovatis vel oblongis obtusis integris vel utrinque obtuse 1-2-dentato-sinuatis inferioribus breviter petiolatis cæteris sessilibus, supremis diminutis, glomerulis laxe spicatis spicis simplicibus sub-aphyllis vel patule paniculatis, perigonii fructiferi sessilis phyllis in thecam compressam basi cuneatam orbiculatam trilobam disco levem ad medium coalitis, lobis rotundatis obtusis intermedio submajore ♂.

Hab. in desertis Arabiae Petreæ ad Ouadi Hamme (Boyé! Schimp. exs. 434 sub *A. rosea*!), ad Ouadi Useit (Boiss!), in desertis Arabicis Egypti mediæ et in Oasi magnâ (Schweinf. 240! et 324!), in oasis Dachiel et Farafrah (Aschers. 2083! et 2084!). Arabice *Bargul* vel *Rocholle*.

Caules 1-1½-pedales, folia 6-9 lineas longa 5-6 lata, racemi 3-4-pollicares, perigonum fructiferum 2½ lineas longum et latum.

23. A. coriaceum (Forsk. Eg. Arab. p. 73 ex parte? — Del. Fl. Eg. p. 283, tab. 52!) fruticosum dense canum, caulis decumbentibus lignosis ramis ascendentibus crassiuseulis crebre foliosis, foliis coriaceis alternis subsessilibus a basi cuneatâ ovatis vel oblongis obtusis integerimis, glomerulis breviter spicatis spicis subinterruptis paniculam brevem densam terminalem formantibus, perigonii fructiferi parvi sessilis valvis in thecam breviter obtriangularem apice subæqua'iter tridentatam disco grosse tuberculatam ad medium usque coalitis 5. *A. ocytropifolium* Viv. Eg. p. 23.

Hab. in arenosis maritimis Egypti inferioris circa Alexandriam (Del! Ehrenb! Gaill! Samar!), in Egypto superiore (Ky. 977!).

Frutex *A. Halimo* ut videtur humilior, pedalis, magis prostratus. Forskahleus hanc speciem in suâ descriptione cum *A. Halimo* cui verba. « Calyces magni bivalves » convenient commiscuisse videtur.

• Radicula basilaris.

24. **A. Halimus** (L. Sp. 1492) fruticosum dumosum argenteo-canum, caulis lignosis elatis patentim ramulosis, foliis alternis breviter petiolatis a basi cuneatâ ovatis obtusis integris rarius basi obsolete repandis, florum glomerulis dense spicatis spicis subinterruptis in paniculam longam pyramidatam aphyllam dispositis, perigonii fructiferi sessilis basi cordati et breviter cuneati phyllis basi tantum connatis reniformibus transverse latioribus obtusissimis contiguis margine integris vel obsolete denticulatis facie laevibus reticulatis §. Ic. Fl. Græc. tab. 962. — *A. Kataf* Ehrenb. pl. Aeg. MSS. spec. florifera huc spectare videtur.

Hab. in maritimis Græciæ (Orph. Fl. exs. 274!), Atticæ et Achaiæ (Heldr!), Archipelagi (Urv!), Syriæ littoralis (Gaill!), Egypti ad Alexandriam (Duperq!), ad Ouadi Goreba (Ehrenb.)

β. *Schweinfurthii*. — Folia magis membranacea longius petiolata abrupte et brevissime cuneata sœpe subcordata.

Hab. in vallibus Ouadi Dugla et Ouadi Uaray deserti Aegyptiaco-Arabici (Schw!).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, Africa borealis.

25. **A. Persicum** (Boiss. Diagn. Ser. I, 12, p. 96) fruticosum adpresso canum, caule humili ramis tenuiter virgatis ascendentibus, foliis alternis breviter petiolatis a basi brevissime cuneatâ vel truncatâ deltaideo-oblongis obtusis integerrimis, racemis laxiuseulis aphyllis, perigonii fructiferi sessilis basi attenuati albo-velutini phyllis præter imam basin herbaceis basi exceptâ liberis ovatis integris vel utrinque obtuse unidentatis patentibus §.

Hab. in desertis Persiæ (Auch. 5268!), prope Murtchehar ad sept. Ispahan (Bge!).

Facies et folia *A. Halimi* sed humile, spicæ fructiferæ 1-1½-pollicares, fructus 3-4 lineas longus 2½-3 latus, basi attenuatus et longior quam latus dum in *A. Halimo* transverse latior et reniformi-cordatus est.

26. **A. Griffithii** (Moq. in D.C. Prodr. XIII, p. 102) suffruticosum canum a basi ramosum, caulis flexuosis patule ramulosis, foliis alternis breviter petiolatis rhombo-ovatis vel orbiculatis breviter cuneatis, floribus 2-3 in glomerulos laxe spicatos abbreviatos dispositis, involuci fructiferi phyllis amplis orbiculatis basi subcordatis et brevissime stipitatis liberis integris disco coriaceis et elevatis reticulatis §.

Hab. in regno Cabulico (Griff. 1751!), in Persiâ ad Ssertschah (Bge!).

β. *Stocksii*. — Folia oblongo-spathulata basi longius cuneata, fructus disco minus coriaceus et tenuiter reticulatus. — *A. Stocksii* Boiss. Diagn. Ser. II, 4, p. 73.

Hab. in Belutschia (Stocks!).

Caules sesquipedales flexuosi, folia 9-12 lineas longa et lata, fructus diam. 4-5 lineas lati.

27. A. farinosum (Forskahl Fl. Æg. Arab. p. CXXIII, № 302) fruticosum adpresso farinoso-canum raniosum ramis ascendentibus crassis crebre foliosis, foliis alternis majusculis brevissime petiolatis late ellipticis obtusis vel retusis integris basi sinu angusto cordatis obtuse biauriculatis, florum glomerulis secus paniculae terminalis basi exceptâ aphyllæ brevis compositæ patule ramosissimæ ramos breves tenues densiuscule et breviter spicatis, perigonii fructiferi cani sessilis oblongo-spathulati subcompressi integerrimi phyllis spongiosis a l medium usque connatis apice rotundatis disco inflato laevi 5. *A. Araticum* Ehrenb. MSS.

Hab. in littore Ægyptiaco maris Rubri loco Ouadi Lecnuma dicto inter Ras Benass et Kosseir (Schweinf!), ad Ssafage ad sept. Kosseir (Dr. Klun-singer!), in monte Gebel Nakus prope Tor Arabiae petreæ (Ehrenb!). In littore Ægyptiaco *Howæi*, in Arabia Osraëi andit.

Frutex bipedalis et forsas elatior, panicula terminalis 3-6 poll. longa ramis 1-2-pollicaribus, folia majora $1\frac{1}{2}$ -2-pollicaria 11-15 lin. lata et basi anguste cordata auriculis parvis obtusis approximatis, quâ notâ a congeneribus distinguitur. Fructus $2\frac{1}{2}$ -3 lin. longi $1\frac{1}{4}$ lati basi subconstricti et rotundati. Primus cl. Ascherson in hac plantâ speciem Forskahlei phrasi brevi sed characteristicâ descriptam agnovit.

Ar. Geogr. Arabia tropica.

Species non sat nota.

A. stylosum (Viv. Pl. Eg. p. 23, fig. 21) fruticosum furfuraceo-cinereum, foliis alternis integerrimis ovato-oblongis in petiolum attenuatis, perigonii fructiferi phyllis suborbiculari-ovatis integerimis dorso tuberculatis binis triangularibus auctis 5

Hab. in desertis Egypti.

Ex descriptione ex fructus analysi pro *A. Halimo* facile haberem sed fructus dorso tuberculatus dicitur.

EUROTIA (Adans. Fam. II, p. 260. — *Diotis* Schreb. non Desf.).

Flores monoici ebracteati. *Masc.* Perigonium quadripartitum lacinias hirtis. Stamina 4 in perigonii fundo inserta. *Fem.* Perigonium tubuloso-urceolatum apice bidentatum fructiferum auctum longe hispidum utriculum includens. Ovarium oblongum compressum, stigmata bina filiformia basi connata. Utriculi integumentum membranaceum hirtum. Semen verticale compressum testâ tenui. Embryo periphericus albumen cingens — Frutices.

1. E. ceratooides (L. Sp. 1389 sub *Axyride*) fruticosum totum indumentostellato adpresso cinereo vel fusco obsitum erectum ramosissimum ramis rigidulis tenuiter virgatis, foliis subcoriaceis linearioribus vel lanceolatis obtusiusculis integris apice subrevolutis basi breviter attenuatis subpetiolatis, floralibus angustatis abbreviatis,

spicis superne masculis inferne femineis axillaribus breviter pedunculatis paniculam longam angustam stricte spicæformem formantibus, perigonio femineo fructifero urceolato apice dentato pilis longis demum rufis hirsutissimo 5. *E. ceratoides* C.A. M. in Led. Fl. Alt. IV, p. 239.
 — *Diotis ceratoides* Willd. Sp. IV, p. 308. — Nees Gen. Germ. Ic.
 — *Achyranthes papposa* Forsk. Eg. Ar. p. 48 Jacq. Ic. Rar. tab. 189.
 — *Ceratospermum papposum* Pers.

Hab. in collinis sæpius gypsaceis, Cappadocia ad Cæsaream (Bal!), Armenia ad Erzerum (Tchih!), Caucasus et Transcaucasia (C.A. M! Hoh!), Persia borealis (Szov! Buhse!) et australis in montibus Kuh Delu et Sabst Buschom (Ky. 316!), Belutschia (Stocks!).

β. ferruginea. — Indumentum intensius ferrugineum, folia latiora basi rotundata vel truncata. — *Diotis ferruginea* Nees Gen. Germ. Ic. fig. 14-19. — *Eurotia ferruginea* Boiss. in Ky. exs.

Hab. in Persiâ australi prope Schiraz (Ky. 556!), Turkestaniâ prope Samarcand (Lehm!), regno Cabûlico (Griff!).

Species foliis anguste linearibus vel dilatatis valde varians, varietas ad typum intermediis manifeste transit.

Ar. Geogr. Hispania orientalis et australis, Austria et Moravia, Rossia media et australis, Songaria, Sibiria, Mongolia Chinensis, regio Himalaica.

CERATOCARPUS (Buxb. Comm. Petr. I, p. 244, tab. 9. — *Ceratoides* Tourn.)

Flores monoici ebracteati. *Masc.* Perigonium hyalinum cuneatum bilobum lobis obtusis carinatis. Stamen unicum imo perigonio insertum, filamentum filiforme. *Fem.* Perigonium tubulosum cuneatum demum valde accrescens fructiferum in thecam obcuneiformem plano-compressam apice truncatam bicarinatam angulis binis apice in spinam longam subulatam divergentem abeuntibus mutatum. Ovarium ovato-globosum, stigmata bina setacea basi coalita. Utriculus thecæ perigoniali inclusus, pericarpium membranaceum hirsutum. Semen clavatum verticale, testâ tenui cum pericarpio coalitâ. Embryo hippocrepicus periphericus. — Herba annua valde dichotoma floribus sessilibus femineis in axillis et dichotomiis solitariis, masculis in axillis superioribus 2-3 aggregatis.

1. C. arenarius (L. Sp. p. 1375) annuus hirtulus vel tomentoso-canescens fere a basi dichotome ramosissimus, foliis sessilibus subulatis rigidis coriaceis pungentibus margine revolutis inferioribus oppositis infimis cito evanidis latioribus oblongo-lanceolatis obtusis cæteris alternis, thecarum cornubus divergentibus thecæ æquilongis vel brevioribus ♂. Lam. Ill. tab. 741.

Hab. in arenosis et argillosis desertorum, Armenia Turcica ad Sbir (Tchih!), Tauria (Stev!), Transcaucasia tota (M.B., C.A. M! Hoh!), Persia borealis et media (Auch! Ky! Haussk!), Turkestania (Lehm!), regnum Cabûlicum (Griff!).

Variat multum foliorum longitudine, dichotomiarum intervallis brevissimis et longioribus, vel dense hemispherico-eriuaceus vel laxior. Flores feminei gemini ex axillis paris primarii foliorum obovati sunt, apice breviter biapiculati.

Ar. Geogr. Bulgaria, Rossia media et australis, Songaria, Sibiria.

TR. III. CAMPHOROSMEÆ Moq.

Flores hermaphroditi vel polygami homomorphi. Seminis integumentum simplex membranaceum. — Caulis continuus.

PANDERIA (Fisch. et Mey. Ind. Petrop. II, p. 45).

Flores polygami (vel hermaphroditi omnes) ebracteati. Perigonum urceolatum æqualiter quinquedentatum, fructiferi dentibus conniventibus faucom claudentibus fornicalis et dorso in tuberculum trigonum vel alam transversam minutam excrescentibus. Stamina 5 imo calyci inserta, antheris exsertis. Stylus brevis longe bicurvis. Utriculus perigonio clauso inclusus, pericarpium tenuissimum. Semen verticale. Embryo subannularis. — Herba annua facie Kochiae.

1. **P. pilosa** (F. et M. loc. cit.) annua pilosa collo simplex vel pluricaulis, caulis erectis virgatis fere tota longitudine breviter ramulosis, foliis alternis sessilibus oblongo-lanceolatis obtusis, floribus 2-3-nis sessilibus secus ramulos spicas breves densiusculas foliis breviter linearibus flores superantibus interstinctas formantibus ⊖. *P. dicaricata* C. Koch Linn. XXII, p. 186. — *Kirilowia pilosa* Bge Cat. Dorp. (ex ipso auctore). — *Kochia Noëana* Boiss. MSS.

Hab. in ruderatis, salsuginosis interioribus, Anatolia ad Koniah (Heldr!) et Cæsaream (Bal!), Armenia ad Erzerum (Huet! spec. floribus majoribus) et Van (Ky. Suppl. 526!), in sepulcreto Tarbet el Dahdah prope Damascum (Gaill!), in ruderatis Babyloniæ ad Bagdad (Noë 312! et 1124!), Georgia Caucasica et ditio Talysch circa pagos (Hoh!), Turkestania (Lehm!), Afghanistan (Griff!).

Facies *Kochia sedoides*, $\frac{1}{2}$ - $\frac{1}{2}$ -pedalis, perigonum fructiferum vel alis parvis oblongis aut quadratis vel præsertim in speciminiibus cultis tuberculis parvis obsitum.

Ar. Geogr. Songaria, Tibetia.

KIRILOWIA (Bge Del. Sem. Dorp. 1843, p. 7).

Flores polygami vel hermaphroditi ebracteati. Perigonum urceolatum æqualiter quinquedentatum demum immutatum. Stamina quinque imo calyci inserta antheris exsertis. Styli duo elongati filiformes. Utriculus membranaceus compressus liber perigonio inclusus.

Semen verticale. Embryo periphericus. — Herba annua facie *Axyridis* vel *Londesiæ*. — Genus a *Panderiâ* perigonio non appendiculato tantum diversum.

1. **K. eriantha** (Kar. et Kir. N° 423 sub *Londesiâ*) annua patule pilosa, caule tenui erecto simplici vel parce ramoso, foliis sparsis sessilibus oblongo-lanceolatis acutis, floribus hermaphroditis ad foliorum caulinorum axillam solitariis submajoribus, glomerulis trifloris ad ramulos axillares brevissimos dense foliolatos subcapitatis densissime villosis lateralibus minoribus femineis terminali hermaphrodito ♂. *K. eriantha* Bge loc. cit. — Pl. Lehn. p. 453.

Hab. in desertis argilloso-Turkestaniae (Lehm!), Affghaniæ (Griff. 1758).

Ar. Geogr. Songaria.

CAMPHOROSMA (L. Gen. 64).

Flores hermaphrodoti vel abortu polygami ebracteati. Perigonum breviter tubulosum compressum quadridentatum demum immutatum dentibus erectis duobus externis herbaceis carinatis duobus eis oppositis subminoribus membranaceis planis. Stamina 4, rarius unicum, basi perigonii inserta, antheræ exsertæ. Stylus filiformis stigmatibus filiformibus 2-3. Utriculus compressus perigonio inclusus, pericarpium membranaceum. Semen verticale testâ membranacea. Embryo annuliformis albumen parvum cingens. — Suffrutices vel herbæ floribus axillaribus sessilibus.

1. **C. Monspeliacum** (L. Sp. 178) perenne basi suffruticulosum et ramosissimum pubescens vel hirsuto-canescens, ramis cespitosis aliis sterilibus et fasciculatim foliosis, aliis elongatis a basi industrata prostrata ascendenti-erectis simplicibus vel paniculatis, foliis rigidis linearis-subulatis rectis vel recurvis, axillis fasciculigeris, floribus abortu polygamis in glomerulos densos steriles ovalis longe spicatos dispositis foliis breviter lanceolato-linearibus eis sublongioribus basi suffultis, perigonii basi glabri dentibus acutis inæqualibus 5. Ic. Nees Gen. Germ.

Hab. in arenosis siccis et salsis, Peloponnesus prope Sicyonem et Corinthum (Heldr.), Thracia (Friv!), Anatolia in salsis ad Iconium (Heldr!) et Cæsaream (Bal!), Cyprus (ex Sibth.).

Ar. Geogr. Gallia australis, Hispania, Italia, Dalmatia, Africa borealis.

2. **C. perenne** (Pall. Illustr. p. 66, tab. 57) perenne canescens basi fruticulosum, ramis sterilibus brevibus cespitosis floriferis ascendenter elongatis, foliis linearis-subulatis subcoriaceis, axillis fasciculigeris, floribus fere omnibus hermaphroditis in glomerulos densos sessiles ovalis longe et laxe spicatos dispositis foliis interfloralibus

lanceolato-subulatis multo brevioribus, perigonii omnino pilosi dentibus lateralibus longioribus ♂.

Hab. in argillosis subsalsis Tauriae (Stev., Rehm!), prov. Caucasicarum (Stev.), Persie bor. in prov. Ghilan (Auch. 5261!), in rupestribus ad Ispahan (Haussk!).

Facies omnino *C. Monspeliaci* a quo notis indicatis ex cl. Bunge specifice differt.

3. C. Ruthenicum (M.B. Taur. Cauc. III, p. 112) perenne basi suffruticosum pubescens a basi ramosissimum ramis aliis brevibus cespitosis densissime et fasciculatim foliosis, aliis floralibus elongatis ascendentibus subsimplicibus, foliis rigidis abbreviatis lineari-subulatis rectiusculis, axillis dense fasciculigeris, floribus in glomerulos densos sessiles ovatos dissitos longe spicatos dispositis fere omnibus hermaphroditis folia floralia ovata obtusa superantibus, perigonii pilosi dentibus subaequilongis apice breviter villosuiis ♂.

Hab. in Tauria circa Perecop (ex Stev.), ditione Daghestan Caucasi (Becker!).

Præter notas indicatas differt a *C. Monspeliaco* et *perenni* gracilitate majore, foliis abbreviatis, floribus et glomerulis minoribus. Ex am. a Bunge species distincta, ex claris Steven et Transvetter varietas *C. Monspeliaci*.

Ar. Geogr. Rossia media et australis, Sibiria Uralensis et Altaica.

4. C. polygamum (Bunge in litt.) perenne basi suffruticosum breviter et adpresso hirsutum ramis sterilibus brevibus cespitosis, floriferis elongatis ascendentibus, foliis rigidis lineari-subulatis brevissimis rectis, axillis dense fasciculigeris, floribus in glomerulos densos sessiles ovatos longe spicatos dispositis polygamis folia floralia acuta superantibus, perigonii hirsutissimi dentibus subaequilongis ♂.

Hab. in regni Cabulici alpinis ad Erak et Kullou (Griff. exs. 1763!), Belutschia prope Doubund (Stocks!).

Facie simillimum *C. Ruthenico* pro quo hoc prius habui. An characteres quibus species præcedentes hujus generis distinguntur sat firmi?

5. C. ovatum (W. K. Pl. rar. Hung. I, p. 64, tab. 63) annum hirtum vel pilosum a collo simplex vel ramosum ramis lateralibus decumbentibus, caule centrali vel unico erecto virgato, foliis carnosis angustissime lineari-subulatis, axillis non fasciculiferis, floribus tetrandris glabriusculis secus ramulos axillares numerosissimos terminalesque foliosos laxius spiciformes dispositis, foliis floralibus flores superantibus, perigonii dentibus obtusiusculis ♂. Ic. Nees Germ. fig. 15-16.

Hab. in paludosis salsis, Thracia borealis prope Karnabad (Jankal).

Amic. a Bunge in litteris *C. annum* Pall. Ill. p. 67, tab. 58 ex Rossia australi, Songariä oriundum floribus villosis imbricationis spicatis, perigonii dentibus acutissimis, lateralibus recurvis a *C. ovato* distinguit.

Ar. Geogr. Hungaria, Transylvania, Rossia australis?

6. **C. monandrum** (Bge in litt.) annum pusillum collo ramosum, ramis erecto-patulis filiformibus albidis pumilis remote foliosis, foliis brevibus linearibus longe rufesceni-villosis, floribus monandris in fasciculos axillares et terminales paucifloros longe rufo-vilosos foliis superatos dispositis, perigonii dentibus brevissimis subæqualibus ♂.

Hab. ad cacumen jugi Erak regni Cabulici (Griff. 1762!).

Planta triplicaris, folia remota 2-4 lineas longa inter se distantia, flores hirsutissimi. Facies *Panderiæ*.

CHENOLEA (Thunb. Nov. Gen. 1781, p. 9).

Flores hermaphroditi abortu interdum polygami ebracteati. Perigonium urceolatum breviter quinquesidum lobis carnosulis subcarinatis tomentosis demum nec induratis nec appendiculatis. Stamina 5 imo perigonio inserta antheris exsertis. Ovarium ovatum. Styli bini graciles filiformes inferne coaliti. Utriculus depresso perigonio immutato sed clauso inclusus, pericarpium membranaceum. Semen horizontale testâ membranaceâ. Embryo periphericus: albumen parcum vel nullum. — Suffrutescens incano-lanati.

1. **C. Arabica** (Boiss. Diagn. Ser. I, 12, p. 97) suffruticosa dumulosa, ramis floriferis abbreviatis ascendentibus breviter albo-lanatis simplicibus in spicas basi interruptas lanatas abeuntibus, foliis minimis sessilibus oblongo-linearibus obtusis canescensibus, floribus in glomerulos 5-15-floros bombycinos ovatos vel globosos sessiles dispositis, foliis floralibus ovato-oblongis glabrescentibus floribus subæquilongis, perigonii membranacei extus bombycini lobis obtusis hemisphericis ♂.

Hab. in declivibus aridis Judeæ circa Mar Saba et in desertis jugi Tih Arabiæ petreæ (Boiss!), in planicie elata Galala deserti Arabico-Ægyptiaci (Schweinf. exs. 242!). Fl. Mart.-Apr.

Durus semipedalis, folia 3-4 lineas longa lineam lata, spicæ 2-3-pollicares glomerulæ pisi magnitudine, superioribus confluentibus. Fructus 1 $\frac{1}{4}$ lin. longus. *C. lanata* Moq. species affinis Canariensis et Maroccana differt spicis tenuioribus, perigonii brevius tomentosi lobis ovatis acutis, semine exalbuminoso.

KOCHIA (Roth in Schrad. Journ. 1800, p. 307. — *Londesia* F. et M. — *Bassia* All. — *Echimspilon* Moq.).

Flores ebracteati hermaphroditi vel abortu feminei. Perigonium urceolatum quinquesidum lobis fructiferis in discum conniventibus dorso appendiculatis (racissime exappendiculatis, confer *K. latifoliae* var.). Stamina 5 imo calyce inserta. Styli duo setacei inferne coaliti.

Utriculus depresso perigonio inclusus. pericarpio membranaceo. Semen horizontale (raro in eadem specie paucissima verticalia) testa membranacea. Embryo subannularis albumen cingens. — Suffrutices vel herbae.

Specierum Orientalium dispositio.

§ 1. **Eukochiae.** — Loborum perigonii fructiferi appendices alaeformes.

• Perennes.

K. prostrata.

• Annuae.

K. arenaria, odontoptera, cana, Griffithii, scoparia.

§ 2. **Bassiæ.** — Loborum perigonii appendices tuberculiformes vel spiciformes.

K. monticola, hirsuta, sedoides, hyssopifolia, muricata, latifolia.

SECT. I. EUKOCHILE.

1. **K. prostrata** (L. Sp. 323 sub *Salsola*) suffruticosa adpresso et tenuiter pubescens basi ramosissima ramis virgatis ascendentibus erectis vel plus minusve prostratis, foliis linearibus vel subulatis planis acutis, floralibus abbreviatis florum glomerulos superantibus, floribus sessilibus solitariis vel saepius 3-5-glomerulatis in spicas longas interruptas vel densas simplices rarius ramosas dispositis, perigonii fructiferi lobis ovatis obtusis, alis dorsalibus semi-orbiculatis vel rhombeis obsolete crenatis calycis disco sublongioribus. *K. prostrata* Schrad. Journ. 1809, p. 85. — Jacq. Austr. tab. 294. — *Chenopodium Augustanum* All. Ped. II, p. 108. tab. 38.

Hab. in siccis arenosis, ad vias, interdum in subsalsis, Tauria, Caucasus et Transcaucasia (M.B., Hoh!), Turkestania (Lehm!).

β. canescens Moq. — Villosior canescens vel albo-villosa.

Hab. in ditione formâ typicâ copiosior, Anatolia in salsuginosis collum prope Iconium (Heldr!), circa Caesaream Cappadociae (Balt!), Armenia Turcica (Calv!). Transcaucasia! Persia borealis ad radices montis Demavend (Ky. 353!), Persia australis in alpe Kuh Daena (Ky. 707!).

Ar. Geogr. Hispania, Gallia australis, Italia, Germania australis et orientalis, regio Danubialis, Serbia, Rossia media et australis, Songaria, Sibiria, regio Himalaica. Africa borealis.

2. **K. arenaria** (W. K. Pl. rar. Hung. I, tab. 78. sub *Salsola*) annua pubescens vel hirsuta a collo simplex vel ramosa. caule erecto virgato simplici vel patule ramosissimo, foliis tenuiter subulato-tiliformibus subcarnosis subtus sulco exaratis, floralibus diminutis sed flores longe superantibus, floribus 1-3 ad axillas glomeratis sessilibus dense lanatis, perigonii fructiferi lobis conniventibus, alis disco calyeis longioribus oblongis rhombeis vel spatulato-ovatis ⊖. *K. arenaria* Roth in Schrad. Journ. — Nees Ic. Germ. — *K. dasyantha* Schrad. Journ. 1809. 3, p. 86.

Hab. in arenosis, Tauria (Leveillé), prov. Caucasicæ (M.B.), montes Persiae borealis (Buhse!).

Ar. Geogr. Gallia australis, Belgium. Germania, Italia, regio Danubialis, Rossia media et australis, Songaria.

3. **K. odontoptera** (Schrenk Bull. Pétersh. I, p. 361) annua patule canescens, caule erecto tenui a basi ramosissimo ramis erecto-patulis, foliis teneris cito marcescentibus planis linearibus acutiusculis, floralibus abbreviatis sed glomerulos superantibus rarius ea tantum æquantibus, floribus 2-3 glomerulatis axillaribus in spicas tenues laxas breves paniculatas dispositis, perigonii tomentelli globosi fructiferi alis cuneatis rhomboe-flabelatis inciso-dentatis disco perigonii longioribus ⊖. *K. stellaris* Moq. Chenop. Enum. p. 93.

Hab. in argillosis, Persie loco non notato (Belang!), deserti Karakum Turkestanie (Lehni.), regnum Cabulicum (Griff!).

Pedalis vel humilior, folia inferiora fere pollicaria linea $\frac{2}{3}$ lata. Valde distincta characteribus a *K. arenariæ* cuius eam cl. Moquin varietatem esse autem.

Ar. Geogr. Songaria, Tibetia (Hook et Thoms. exs. sub *Panderia pilosa* in herb. meo! sed ex eadem collectione sub eodem nomine amic. Bunge vidit *K. Griffithii*).

4. **K. cana** (Bge in litt.) annua adpressiuscula cano-hirsuta humilis collo ramosa. ramis lateralibus ascendentibus omnibus crebre foliosis, foliis breviter et latiuscula linearibus acutis superioribus vix abbreviatis glomerulos axillares subsuperantibus, perigonii tome:tos fructiferi alis ovatis integris hirsutis urceolatim conniventibus ⊖.

Hab. in Persia orientali inter Kerman et Ispahan (Bge!).

Specimen vidi non sat aetate proiectum canibus tripollicaribus, folia 3-4 linea longa vix lineam lata.

5. **K. Griffithii** (Bge in litt.) annua pilis crispis villosa ramosissima ramis tenuiter et longissime virgatis. foliis breviter lanceolatis acutiusculis floralibus ovatis obtusis glomerulos subsuperantibus et æquantibus. glomerulus axillaribus paucifloris villosissimus longissime et laxe spicatus, perigonii fructiferi lobis breviter ovatis obtusis conniventibus, alis carnosulis ovato-triangularibus acutiusculis ⊖.

Hab. in regno Cabulico (Griff. 1775). E ditione nondum vidi.

Huc refero specimen ex India boreali Herb. Wight N° 2479. Pedalis et sesquipedalis, rami laterales inferne sæpius ramulosi in spicas 6-10-pollices longas ahenentes. Species rufescens-villosissima perigonii alis non membranaceis sed crassiusculis subtriangularibus a congeneribus distineta.

Ar. Geogr. India borealis.

6. **K. scoparia** (L. Sp. 321 sub *Chenopodium*) annua pubescens vel villosa, caule erecto elato virgato firmo stricte ramosissimo, foliis viridibus basi in petiolum brevem attenuatis anguste lanceolatis vel linear-lanceolatis ciliatis, floralibus brevibus acuminatis flores superantibus, floribus 1-5 in glomerulos axillares sessiles paniculatim spicatos dispositis, perigonii lobis ovatis obtusis, carinâ in tuberculum triangulare acutum brevissimum interdum omnino abortivum rarius in alas breves incrassatâ ☽. *K. scoparia* Schrad. Hal. p. 18, tab. 1. — *Salsola scoparia* M.B.

Hab. in ruderatis et salsis humidis, Thracia littoralis (Griseb.), Anatolia prope Ieonium (Heldr!) et Caesaream (Ball!), Armenia Rossica (C. Koch), Transcaucasia (Hoh!), Persia prope Ispahan (Auch! Heldr!), Turkestanica (Lehm!). An vere spontanea? nam ubique colitur ad scopas utilissima.

Ar. Geogr. Hispania, Gallia, Italia et Germania austro-orientalis nbi sub-spicata, regio Danubialis, Rossia media et australis, Songaria, China, Japonia, India borealis.

SECT. II. BASSLE.

7. **K. monticola** (Boiss. in Ky. Pl. Pers. austr. 1845. — Moq. in D.C. Prodr. XIII, p. 133) annua pubescens vel hirsuta pusilla a cotto sæpius ramosa, ramis decurrentibus vel ascendentibus simpli-cibus filiformi-virgatis rigidulis, foliis inferioribus anguste linearibus basi attenuatis obtusiusculis, floralibus oblongis flores æquantibus vel superantibus, floribus 1-3-nis sessilibus axillaribus, spicis demum elongatis tenuibus laxissimis, perigonii ovato-oblongi submembranacei lobis ovatis conniventibus dorso in tuberculum breviter conicum viride ahenibus ☽. *Pentodon barbatus* Ehr. MSS.

Hab. locis nudis arenosis nive derelictis regionis alpinæ, jugum Makm-el 600'-800' (Ehr! Ky. 355!), supra Eden (Bl!), Antilhanus supra Bludan (Ky. 49!), Persia australis in montibus Kuh Daëna et Kuh Delu (Ky. 52! et 74!), mons Sawers ad nives 10-1200' (Haussk!), montes Persiae bor. (Bulse).

Plantula ad summum 4-5-pollicaris interdum vix pollicaris, folia 3-5 lineas longa. Perigonium immaturum tantum vidi, an ideo tuberculi conici in fructu magis evolvuntur? Ex amic. Bunge in litt. *Pandericæ* species et ab eo *Pandericæ pilosæ* dubitanter adnumeranda, sed habitus alienus, semen (junius) horizontale videtur, etc.

8. **K. hirsuta** (L. Sp. p. 323 sub *Salsola*) annua hirsuta, caule erecto ascendente vel procumbente a parte inferiore patule ramoso, foliis teretibus filiformibus carnosulis obtusiusculis albidis, glomerulis secus ramos tenues flexuosos axillaribus remotis hirsuto-lanatis, perigonii fructiferi appendicibus brevissime conicis pubescentibus

rectis vel subincurvis interdum inaequalibus ⊖. *K. hirsuta* Nolte. — *Suæda albida* Pall. Ill. tab. 45. — *Echin. hirsutus* Moq. — *Suæda crassifolia* Pall. Ill. p. 54, tab. 46. — *Ech. crassifolius* Moq. — *Schoberia Pallasiana* C A. Mey. Led. Fl. Alt. I, p. 402.

Hab. in maritimis districtus Siwasch Taurie (Rehm!), ad littora Persica maris Caspii Gmel. ex Pall., Turcomania.

Ex cl. a Bge auctoritate *Suædam crassifolium* Pall. huc retuli.

Ar. Geogr. Gallia, Belgium, Dania, Gothia, Italia borealis, Rossia australis, Songaria, Sibiria.

9. K. sedoides (Pall. Voy. p. 280, tab. 41 sub *Salsold*) annua, caule erecto virgato rigidulo canescenti-tomentoso, ramis numerosis tenuibus erecto-patulis, foliis carnosis semitereti-filiformibus pubescentibus pilis longioribus immixtis, floribus 2-3 glomeratis in spicas tenues laxas vel densifloras dispositis, foliis floralibus linearibus obtusis glomerulos superantibus, perigonii villosi fructiferi spinis conicis disco æquilonigis vel brevioribus inermibus sepe inaequalibus ⊖. *Suæda sedifolia* Pall. Ill. tab. 32-34. — *Salsola muricata* M.B. Casp. p. 148 non L. — *S. cinerea* W. K. pl. rar. tab. 106. — *Echin. sedoides* Moq.

Hab. in sterilibus salsis et salsuginosis, Tauria (Pall., M.B.), Iberia Cauca-sica (C. Koch).

Ar. Geogr. Regio Danubialis, Bulgaria, Rossia media et australis, Songaria, Sibiria.

10. K. hyssopifolia (Pall. Voy. p. 278, tab. 43 sub *Salsold*) annua hirsuta vel villosa collo simplex vel pluricaulis, caulis erectis albidis rigidulis patule ramosissimis, foliis oblongo-linearibus acutis planis tenuibus, floralibus abbreviatis oblongis flores superantibus basi lanatis, glomerulis 2-3-floris sessilibus interrupte et tenuiter spicatis spicis paniculatis, perigonii fructiferi spinis subulatis patentibus basi non dilatatis apice truncatis disco subæquilonigis vel longioribus ⊖. Ic. Nees Germ. fig. 18-24. — *Suæda hyssopifolia* Pall. Ill. tab. 36. — *Chenopodium villosum* Lam. Dict. I, p. 496. — *Echinosiphon lanatum* Moq. — *Ech. hyssopifolium* Moq. in p.c. Prodr. XIII, p. 135.

Hab. in arenosis et argillosis subsalsis, pagorum ruderatis, Cataonia ad Eski Malatia (Haussk!), Babylonia ad Bagdad (Noë 167! forma spinis elongatis), Armenia ad Erzerum et Baibut (Auch. 2795! Bourg!), Tauria (Brunn!), Transcaucasia (Hoh!), Persia borealis (Buhse!), Turkestania ad Samarkand (Lehn!).

Ar. Geogr. Italia, regio Danubialis, Rossia australis, Songaria.

11. K. muricata (L. Mant. p. 54 sub *Salsold*) annua villosa a collo saepius pluricaulis, caulis erectis vel ascendentibus a basi saepius ramosissimis ramisque erecto-patulis crassis, foliis linear-lanceolatis subearniosis planis, floribus 1-3 glomeratis sessilibus foliis

floralibus oblongis glomerulosis æquantibus vel superantibus suffulti spicas tenues laxas paniculatas formantibus, perigonii fructiferi spinis acicularibus flavidis explanatis rectis disco 2-3-plo longioribus ☽. *K. muricata* Schrad. Journ. 1809, p. 86. — *Bassia muricata* All. Misc. Taur. III, tab. 4. — *Sueda muricata* Pall. Ill. tab. 35. — *Cornulaca muricata* Del. Fl. Eg. Ill. p. 55. — *Echinopsis muricatus* Moq. in D.C. Prodr. XIII, p. 134. — *Salsola monobractea* Forsk. Eg. Arab. p. 55.

Hab. in arenosis et cultis ad deserti margines et in desertis, Egyptus inferior et media (Del! Ehrenb! Wiest! Boiss!), in Oasi magnâ (Schweinf!), Arabiae petreæ desertis (Bové! Schimp. 143! Boiss!).

Pedalis et sesquipedalis, folia 6-8 lineas longa præter floralia brevissima in plantâ fructiferâ evanida.

β. tenuifolia. — Caules et rami tenuiores, folia angustiora. — *K. ericifolia* Viv. Eg. p. 25.

Hab. cirea Alexandriam (Viv., Sieb!).

Ar. Geogr. Barbaria, Arabia tropica.

12. **K. latifolia** (Fresen. Beitr. p. 179) annua a collo saepius ramosa ramis erectis et ascendentibus pumilis fragilibus lanatis, foliis parvis planis virentibus parce et adpresse hirsutis inferioribus linearibus obtusis floralibus oblongo-linearibus vel ellipticis, floribus glo-merulatis dense spicatis lanâ copiosâ longâ demum valde accretâ involutis spicas densas formantibus, perigonii fructiferi spinis subulatis disco æquialongis apice incurvis ☽. *Echinopsis eriophorum* Moq. in D.C. Prodr. XIII, p. 135 sed ex cl. Bge in pl. Lehm. *Kochia eriophora* Schrad. Journ. 109, p. 86, tab. 3 est species diversa, (*K. hyssopifolia* vel forma proxime affinis). — *Salsola laniflora*, *eriantha* Hort.

Hab. in rupestribus et arenosis Arabiae petreæ ad Ouadi Hebran (Fresen., Schimp. 433!), in deserto tribùs Juda (Roth!), Syria in deserto Pahmyræ (Bl!), Mesopotamiâ et Babyloniam (Oliv!), Kutt ad Tigrem (Noë 10!). Persia australi ad Buschir (Haussk!). Belutschia (Stocks!).

β. inermis. — Flores nonnulli superiores hermaphroditi semine verticali, cæteri hermaphroditi et feminei steriles vel semine horizontali. Perigonium fructiferum exappendiculatum. — *Londesia eriantha* F. et M. Ind. II. Petr. p. 40.

Hab. in deserto Ægyptiaco-Arabico ad Ouadi Askar (Schweinf!), in Arabia petreæ (Auch. 272!) et Schimp. exs. 433 partim cum formâ typicâ ex Bge), deserto salso Persiae prope Djendack (Buhsei!), Turkestaniâ (Lehm!).

Varietas formæ typicæ præter perigonii spinas deficientes et semen alterum verticale nec horizontale ut ovum ovo simillima est et non fieri potest eas generice imo specifice differre. Omnibus notum est seminis situm in speciebus pluribus ordinis variare et perigonium appendiculatum vel exappendiculatum in *K. scopariâ* et aliis observatur.

AGRIOPHYLLUM (M.B. Taur. Cauc. III, 6).

Flores hermaphroditici cbraetecolati in axillis bractearum glomerulii solitarii. Perigonium 1-5-phylum interdum nullum, phyllis membranaceis. Stamina 1-5 nonnulla saepe sterilia filamentis subulatis. Ovarium obcompressum nunc in stylum bicrurem attenuatum nunc apice tridentatum dente intermedio biceruri. Utriculus compressus alatus vel apterus stylo persistente rostratus interdum præterea utrinque unidentatus, prope basin facie anticâ saepe quoque posticâ foramine orbiculari dehiscens. Semen verticale compressum testâ subcoriacea. Embryo annularis circa albumen centrale. — Herbae annuae coriaceæ glomerulis squarrosis.

1. A. lateriflorum (Lam. Encycl. Meth. IV, p. 756 sub *Eryngio*) annum glabriuscum, caule striato a basi divaricatim ramosissimo, foliis sessilibus anguste lanceolatis 5-7-nerviis pungentibus patentibus, glomerulis globosis axillaribus, bracteis a basi ovalâ triquetro-subulatis brevibus acerosis squarrosis, perigonii phyllo unico a basi ovata subulato, staminibus subquinis, fructu glabro membranaceo oblongo aptero utrinque foramine orbiculari dehidente in rostrum triaristatum abeunte aristis subulatis, mediâ sublongiore biceruri ♂. *A. lateriflorum* Moq. in n.c. Prodr. XIII, 2, p. 139. — *Gramen Orientale spicatum fruticosum spinosum spinis echinatis in capitulum congestis* Tourn. Cor. 39. — *A. Tournefortii* Fisch. MSS.

Hab. in Armenia (Tourn.), ad Araxem (Szov!).

Semipedale vel pedale, folia 1-1½-pollicaria 1-3 lineas lata, bracteæ tres lineas longæ. Fructus characteribus a congeneribus facile distinguendum.

2. A. arenarium (M.B. Taur. Cauc. III, p. 6) annum viride vel canescens pube stellata sparsâ plus minus adspersum, caule erecto angulato-striato crasso squarrose ramosissimo ramis flexuosis divaricatis rigidis, foliis sessilibus basi attenuatis lanceolatis longe acuminate-spinosis 5-15-nerviis patentibus-reflexis, florum glomerulis axillaribus, bracteis a basi ovata subulatis pungentibus squarrosis, perigonii phyllis 1-3 interdum nullis modo linearibus acutis modo obovatibus vel subquadrilateris, staminibus 3-2, fructu ovato anguste alato et margine hirsuto apice bicuspidato antice foramine orbiculari dehidente ♂. *A. squarrosum* Moq. in n.c. Prodr. XIII, p. 139. — *Corispermum squarrosum* Pall. Fl. Ross. II, p. 113, tab. 99 non Linn.

Hab. in arenosis prov. Caucasicarum ad Caspium (Stev., C.A. M.), ad littora Turcomanie (Karel.).

Ar. Geogr. Rossia australis, Songaria, Sibiria Altaica, China.

3 A. minus (F. et M. loc. cit. № 771. — Bunge pl. Lehm. p. 457) annum viride pube ramosâ sparsâ obsitum vel glabrum, caule striato

a basi divaricatum ramosissimo ramis tenuibus flexuosis, foliis sessilibus linearis subulatis 7-9-nerviis pungentibus reflexis, glomerulis axillaris subglobosis, bracteis a basi dilatata triquetro-subulatis acerosis squarrosis basi glabris, perigonii phyllis subternis linearibus et oblongo-linearibus, staminibus 3-5, fructu glabro utrinque operculatum dehiscente alato alis lanceolatis utrinque apice liberis et inciso-denticulatis, in rostrum tricuspidatum abeunte cuspidibus lateralibus divergentibus acicularibus medio bifido ⊖.

Hab. in arenosis mobilibus deserti Kisilkum Turkestaniae (Lehm!), Turkestaniae littora (Karelin).

4. **A. latifolium** (F. et M. in Karel. Enum. Bull. Mosq. 1839, p. 170) annum pube ramosa canescens, caule striato a basi diehotome et diffuse ramosissimo ramis tenuibus flexuosis, foliis petiolatis ovatis vel subcordatis longe pungenti-cuspidatis, glomerulis axillaris globosis, bracteis a basi ovata subulatis pungentibus brevibus squarrosis basi pilosis, perigonii phyllis subternis linearibus et linearis-oblongis membranaceo-hyalinis, staminibus 3-5, fructu superne canescente obovato ala denticulata superne in lobos breves rotundatos producta marginato, in rostrum lanceolatum semibifidum abeunte, utræque facie operculatum dehiscente ⊖.

Hab. in arenosis mobilibus desertorum circa Buchara et Agatme Turkestaniae (Lehm!), in Turcomaniâ (Karell), Belutschia (Stocks 1081¹).

CORISPERMUM (L. Gen. № 42).

Flores hermaphroditi ebracteati in bractearum axillis solitarii. Perigonium membranaceum 1-3-phyllum vel nullum. Stamina 1-5, plerumque terna, nonnulla saepe sterilia. Ovarium obcompressum ovatum, stylus bipartitus. Achenium indehiscens apice convexum postice planum circum circa alatum vel apterum styli basi bimucronatum, pericarpium induratum semini adhaerens. Semen verticale, embryo annularis albumen cingens. — Herbæ annuae.

1. **C. Orientale** (Lam. Dict. II, p. 111) annum plus minusve pubescens a basi divaricatum ramosissimum, foliis linearis-lanceolatis obtusiusculis subtrinerviis, spicis crassiusculis densiusculis, bracteis ovato-ellipticis albo-marginatis, perigonio plerumque monophyllo, staminibus 3-5, fructu ovato aptero apice integro mucronulato ⊖.
C. Pallasii Stev. — *C. spicatum* Willd. En. Suppl. p. 4.

Hab. in arenosis maritimis ditionis Transcaucasicae Talysh (Hoh.), in Persia (André).

Ar. Geogr. Rossia media, Songaria, Sibiria.

2. **C. Marschallii** (Stev. in Mém. Mosc. V, p. 336) annum adpresso pubescenti-flavescens, caule flexuoso a basi alternatim

ramoso, foliis angustissime linearibus elongatis acutis uninerviis, spicis terminalibus oblongis vel elongatis laxis vel densis, bracteis ovato-lanceolatis attenuato-acutis membranaceo-marginatis, perigonio saepius nullo, staminibus saepius ternis, fructu ovato-orbiculari alato ala dimidio semini subangustiore apice retusâ vel denticulatâ ⊖. *C. squarrosum* M.B. Taur. Cauc. I, p. 3.

Hab. forsitan in prov. Caucasicis e quibus autem nondum specimina vidi.

Ar. Geogr. Germania Rhenana, Italia superior, regio Danubialis, Rossia media et australis.

3. **C. Lehmannianum** (Bge Pl. Lehm. p. 458) annum humile pubescens demum glabratum a collo strictiuseule et fastigiatim ramosum, foliis a basi attenuata oblongo-spathulatis obtusis uninerviis, spicis elongatis gracilibus laxiusculis, bracteis oblongo-linearibus anguste marginatis cartilagineo-cuspidatis fructu duplo longioribus, perigonio monophyllo, stamine unico, fructu oblongo anguste marginato integro bicuspidato ⊖.

Hab. in collibus arenosis inter fluvios Jan et Kuwandarja Turkestaniae (Lehm!).

4. **C. nitidum** (Kit. in Schultz Oesterr. Fl. I, p. 7) annum pubescentem sparsim adspersum tandem glabrum, caule a basi divaricatum ramoso ramisque gracilibus, foliis patulis anguste linearibus acutiusculis obsolete uninerviis, spicis elongatis tenuissimis laxis, bracteis ovato-lanceolatis albo-marginatis, perigonio plerumque monophyllo, staminibus 3-5, fructu parvo ovato margine angustissime membranaceo apice integro bimucronulato ⊖. *C. purpurascens* Host.

Hab. in Tauria (Leveillé ex Moq.). Editione nondum vidi.

Ar. Geogr. Austria, regio Danubialis, Serbia, Rossia australis, Sibiria.

5. **C. hyssopifolium** (L. Sp. p. 6) annum hirtellum vel glabrum glaucescens a collo saepius pluricaule, caulis diffusis vel erectis flexuosis alternatim ramosis, foliis erectis linearibus obtusis mucronulatis uninerviis, spicis ad ramulos et caulem terminalibus oblongis vel cylindricis densis vel laxis, floribus floralibus (bracteis) linear-oblengis anguste albo-marginatis, perigonio monophyllo bilobo, staminibus 1-3, fructu suborbiculari ovato diaphane et angustissime marginato apice basi styli mucronato ⊖. Ic. Fl. Græc. tab. 1. — *Corispermum squarrosum* L. Sp. p. 6.

Hab. in arenosis et glareosis, Thracia maritima (Sibth.), Georgia Caucasicæ et prov. Talysh (Hoh!), Persia borealis maritima ad Enzeli (Buhse!), prope Kerman (Bge!).

Ar. Geogr. Gallia australis, Germania austro-orientalis, regio Danubialis, Rossia media et australis, Sibiria omnis, regio Himalaica, Tibetia, America borealis.

ANTHOCHLAMYS (Fenzl in Endl. Gen. p. 300).

Flores hermaphroditæ ebracteolati in bractearum axillis solitarii. Perigonium corollinum campanulato-urecolatum profunde quinquelobum lobis bilobis. Stamina quinque filamentis basi coalitis. Ovarium compressum substipitatum, styli bini apice coaliti. Achenium indehisrens orbiculare compressum alà scariosâ marginatum, pericarpium tenuë semini adhærens. Semen verticale. Embryo annularis albumen cingens. — Herba annua facie *Oligomeridis*.

1. **A. polygaloides** (F. et M. Index Petrop. 1835, p. 24 sub *Corispermo*) annua glaucescens præter apices subvillosos glabra, a collo sæpius ramosa, caule erecto flexuoso divaricatum ramosissimo, ramis virgatis longissimis, foliis infimis ovato vel oblongo-spathulatis subpetiolatis subuninerviis nervulosis, cæteris lanceolatis et linearibus, spicis elongatis primum confertis dein laxis, bracteis lanceolatis acutis fructu brevioribus, perigonii rosei vel albi laciniis cuneatis bilobis, fructus minutus alà plicatulâ diaphanâ dimidio semini subæquilateralè. *A. polygaloides* Moq. Chenop. N° 1. — Jaub. et Sp. Ill. Or. tab. 299.

Hab. in rupestribus montanis ditionis Transcaucasiae Talysh (Hoh!), in montibus Persiae circa Teheran (Auch. 5259! Haussk!).

Flores gratae odorati ex cl. Haussknecht.

TR. IV. SALICORNIEÆ Dumort.

Flores hermaphroditæ terni (rarius bini) amentaceo-spicati spatio concavo (*areæ floriferæ*) supra bractearum axillam immersi sessiles. Perigonium 3-4-phylum phyllis connatis apice tantum liberis rarius ad basin usque liberis, ob dentes conniventes et contiguos utrieuli-forme clausum demum spongiosum vertice scutiformi. Stamina 2 vel 1 anthesi ineunte inter dentes perigonii emergentia. Ovarium pyriforme, stylus basi simplex superius 2-3-cruris cruribus plumosis demum longe exsertis. Semen verticale, pericarpium membranaceum rarius lignosum. Albumen centrale interdum nullum. Embryo curvatus, semiannularis, annularis vel conduplicatus. — Herbæ, suffrutices vel arbusculæ, ramis et ramulis plerumque articulatis, articulis subclavatis. Folia vel opposita in vaginam corticis modo articulos vestientem laminâ destitutam et in cupulam brevissimam bilobam apice abeuntem connata, vel sparsa et tunc inter se libera parte vaginanti obliquâ laminâ vel rudimentariâ vel longiusculâ. In expositione tribus difficilimæ egregiam monographiam cl. Ungern Sternberg in Atti del Congresso di Firenze 1874, p. 259 pedentem secutus sum.

ARTHROCNEMUM (Moquin Chen. Enum. p. 444).

Caules et rami articulati aphylli. Folia omnia opposita per paria connata laminâ rudimentariâ. Bracteæ oppositæ connatæ persistentes. Flores spicati terni non vel vix connati areâ florigerâ (in nostrâ specie saltem) superficiali inserti. Perigonium apice trifidum obtuse pyramidatum. Semen albuminosum, embryo virguliformis, radicula infera. — Suffrutices.

1. A. glaucum (Del. Fl. Eg. Ill. p. 69 sub *Salicornia*) perenne fruticosum ramis erectis vel procumbentibus radicantibus, ramulis floriferis erectis, spicis crassis cylindricis obtusis, bracteis integerimis flores demum ex areâ semiexsertos non omnino obtegentibus, staminibus binis, pericarpio membranaceo. seminis testâ nigrâ nitidâ papillis brevibus conicis obsitâ 3. *A. glaucum* Ung. Sternb. — *S. Virginica* Forsk. Fl. Eg. Arab. — *S. macrostachya* Moric. Fl. Ven. I, p 2. Ic. Guss. Sic. tab. 4. — *Arthr. macrostachyum* Moris et Delponte Enum. Sem. Taur. 1854. — *Salic. fruticosa* et *Arthr. fruticosum* Moq. et alior. Auct. ex parte.

Hab. in maritimis salsis, Peloponnesus (Bory! Heldr!), Coreyra (Fried.), Cyclades (Bory!), Creta (Heldr! Raul!), Egyptus (Del., Ehrenb., Sieb. ex Ung. Sternb.), Arabia petrea ad Tor (Ehrenb.).

Caules primariae saepius expansi radicantes ramos erectos emittentes, rarius trunci erecti humanie attitudinis, spicæ crassæ (auct. cl. Ung. Sternb.) non ex apice caulinum primariorum erectorum sed ex ipsorum ramulis basilaribus oriundæ, interdum ramosæ et articulis inferioribus elongatis insignes. Sæpe confusum cum *Salic. fruticosa* qua preter characteres genericos differt spicis tenuioribus secus caules primarios axillibus et terminalibus, floribus in areâ florigerâ profunde immersis non exsertis etc. (Confer. Ung. Sternb. loc. cit. p. 275 et 287).

Ar. Geogr. Lusitania, Hispania Gaditana et Mediterranea, Gallia australis, Corsica, Sicilia, Sardinia, Italia australis, Dalmatia, Africa borealis.

SALICORNIA (L. Gen. 40 ex parte).

Caules et rami articulati aphylli. Folia omnia opposita per paria connata laminâ rudimentariâ. Bracteæ oppositæ connatæ persistentes. Flores terni spicati connati areâ florigerâ profunde excavata immersi et hujus parietibus adnati; areæ fructu delapso septulis in tres loculos divisæ cernuntur. Perigonium apice plano-convexum medio 3-4-denticulatum. Semen exalbuinosum, embryo conduplicatus cotyledonibus radiculæ inferæ fere contiguis. — Suffrutices vel herbæ annuæ.

1. S. fruticosa (L. Sp. p. 5) pereunis fruticosa, caulis et ramis primariis inferne saepè radicantibus decussatim ramosissima, spicis cylindraceis vel cylindraceo-clavatis obtusis, areæ florigeræ foveolis

tribus collateralibus, staminibus binis, perigonii floris medii vertice pentagono vel trapezoideo, seminiibus fulvis hirtis 5. Ic. Guss. Fl. Sie. tab. 3. — *S. radicans* Smith Engl. bot. tab. 1691. — *S. Alpini* et *S. anceps* Lag. — *S. Arabica* L. Sp. 3, p. 3. — *S. Europaea* Forsk. Eg. Arab. — *Arthr. fruticosum* Moq. ex parte.

Hab. in salsis maritimis, Græcia in Atticæ insula Salami (Orph!), Macedonia ad ostium Axii (Griseb.), insula Thera Archipelagi (Sart!), Anatolia ad Smyrnam (Forsk.), Syria littoralis (Rauwolf), Egyptus ad Alexandriam (Gaill.), et in oasi Ammonis (Ehr.), in Oasi Dachel (Aschers!).

Ar. Geogr. Littora Britanniae, Galliae, Lusitaniae, Hispaniae, Italiae, Dalmatiae, Arabiæ tropicae, P. B. Spei, Americæ totius.

2. **S. herbacea** (L. Sp. p. 5) annua erecta vel procumbens non radicans decussatim ramosissima, spicis breviter pedunculatis tenuiter cylindricis apice attenuatis, areæ florigeræ foveolis tribus in triangulum dispositis, staminibus saepius binis, perigonii floris medii vertice obovato subrhomboeo, seminiibus fulvis hirtis 5. Ic. Fl. Dan. tab. 303. — *S. annua* Engl. Bot. tab. 415. — *S. radicans* Koch Syn. non Smith. — *S. prostrata* et *S. acetaria* Pall. Ill. tab. 3 et 1.

Hab. in salsis maritimis et interioribus, Cyclades (Smith Prodr., Bory), Macedonia ad littora fluvii Axii (Griseb.), Cappadocia prope Cæsaream (Bal!), Tauria (Stev.), Transcaucasia (Hoh! Bunge), Persia in salsis prope Teheran (Haussk!), Turkestania (Lehm.), Egyptus ad lacum Mareotidem (Ehrenb!).

Ar. Geogr. Europa fere tota, Songaria, Africa borealis, P. B. Spei, America septentrionalis.

KALIDIUM (Moq. in D.C. Prodr. XIII, p. 446).

Ramuli oblique articulati. Folia sparsa inter se libera basi ramulo adnatâ cylindraceâ, laminâ manifestâ vel rudimentariâ. Braeteæ spirâliter dispositæ. Flores tres connati spicati areâ floriferâ immersi et ejus parietibus adnati; area fructu elapso septulis trilocularis evadit. Perigonium apice latiusculum planum vel convexum toto circuitu margine alæformi cinctum in medio 4-denticulatum. Albumen centrale copiosum, embryo hippocrepicus, radiculâ inferâ. — Suffrutices.

1. **K. Caspicum** (L. Sp. p. 4 sub *Salicornia* non Pall.) perenne frutescens ramosissimum dumosum, ramis ramulisque tenuibus exsiccatis fragilibus, foliorum alternorum parte adnatâ amplexicauli obovatâ apice rotundatâ laminâ plus minusve rudimentariâ, spicis alternis et terminalibus sessilibus oblongis vel cylindricis longioribus basi saepè remotifloris, areis florigeris trifloris hinc inde easu bifloris, perigonii vertice convexo subumbilicato floris intermedii obovato-subpentagono lateralium ovato-subtrigono 5. *K. Caspicum* Ung. Sternb. — *Kalidium Arabicum* Moq. — *Salicornia Arabica* Pall. non L. — *Salicornia Pallasii* C.A. Mey. in Talyseh Enum. p. 425. — Ic. Buxb. Cent. I, tab. 11, fig. 1.

Hab. in salsis, Armenia Rossica ad Araxem (Szovf!), prov. Karabagh et littora Caspia Georgiae (C.A. Mey! Hoh!), Turkestaniæ deserta Karakum et Kisilkum (Lehm!). In Arabiâ ex synonymis falsis erroneè indicatum.

Fruticulus vix pedalis, spicæ tenues 3-6 lineas longæ. Ramuli cum spicis sæpe insecti ietu in gallas strobiliformes mutati.

Ar. Geogr. Rossia austro-orientalis, Songaria, Sibiria Altaica, Mongolia Chinensis.

2. K. foliatum (Pall. Voy. App. p. 221, tab. 45 sub *Salicornia*) perenne frutescens ramosissimum erectum vel prostratum, ramis ramulisque tenuibus exsiccatis non fragilibus, foliorum alteriorum parte adnatâ semiamplexicauli laminâ semitereti breviter lanceolatâ vel linearî patente vel recurvâ, spicis sessilibus alternis et terminâlibus racemosis densifloris cylindricis obtusis, areis florigeris trifloris, perigonii vertice plano valde umbilicato floris intermedii pentagono-trapezoideo, florum lateralium subrhombico 5. *K. foliatum* Moq. — Pall. Ill. tab. 7. — *Sal. strobilacea* Gærtn. Fr. tab. 127, fig. 8 non Pall. — *Sal. Caspica* Pall. Ill. tab. 7.

Hab. in salsis Tauriæ (Pall.), Turcomaniæ (Karel.), Turkestaniæ (Lehm!).

Fruticulus humilis, foliorum lamina 2-3 lin. longa, spicæ 9-15 lin. longæ.

Ar. Geogr. Rossia austro-occidentalis, Songaria, Sibiria, Mongolia Chinensis.

HALOPEPLIS (Bunge Linn. XXVIII, p. 573).

Ramuli moniliformes vel nodulosi. Folia sparsa inter se libera, basi adnatâ foliorum superiorum amplexicauli. Bracteæ ordine spirali dispositæ. Flores tres plus minus connati spicati areâ florigerâ immersi et ejus parietibus adnati. Perigonii vertex tridenticulatus parvus convexiusculus circumcircâ non alatus. Albumen centrale copiosum, embryo uncinatus radiculâ ascendentæ. — Herbæ annuæ et suffrutices.

1. H. pygmæa (Pall. Ill. I, p. 8, tab. 2, fig. 2 sub *Salicornia*) annua, caule pumilo erecto a basi ramoso ramis erecto-patulis longiore, foliorum alteriorum parte vaginanti cordato-subglobosâ tractu inter-axillari subbreviore, laminâ rudimentariâ, spicis alternis et terminâlibus in racemum dispositis oblongis et oblongo-cylindricis, floribus 2-3 inferne coalitis et areæ florigeræ parietibus adnatis, staminibus binis, seminis orbicularis testâ papillis cylindraceis undique obsitâ ♂. *H. pygmæa* Bge ex Ung. Sternb. — *Halostachys Songarica* Schrenck Bull. Acad. Petersb. 1843, p. 361.

Hab. in Turcomaniâ (Cederholm), in Persiâ australi (Bunge ex Ung. Sternb.). Planta 4-5-pollicaris, spicæ 3-4 lineas longæ.

Ar. Geogr. Rossia austro-orientalis, Songaria.

2. H. amplexicaulis (Vahl Symb. II, p. 1 sub *Salicornia*) annua glauca, caule tandem inferne sæpe indurato ex ipsa basi

ramoso, ramis horizontalibus vel ascendentibus, ramulis adultis nodulosis, foliis alternis semiglobosis amplexicaulibus obtusis, laminâ rudimentariâ, spicis sessilibus alternis densilloris racemosis, floribus ternis ad verticem perigonii usque connatis et areæ florigeræ parietibus adnatis, stamine unico, seminis subovati testâ papillis cylindricis brevibus parte quâ embryoneum tegit obsoletâ ⊖. *H. amplexicaulis* Ung. Sternb. — *Salic. nodulosa* Del. Deser. Eg. vol. 19, p. 121. — *Salic. strobilacea* Del. Eg. tab. 3, fig. 2. — *Halostachys perfoliata* Moq. in D.C. Prodr. XIII, p. 148 ex parte.

Hab. in salsis maritimis, Cilicia ad ostium Iuv. Cydri (Baff!), Egyptus inferior ad mare Mediterraneum et lacum Mareotidem (Del., Sieb! Ehrenb! etc.), in deserto Libyco (Aschers!).

Ar. Geogr. Hispania australis, Italia australis et Sicilia, Africa borealis.

3. **H. perfoliata** (Forsk. Fl. Eg. Arab. p. 3 sub *Salicornia*) perennis fruticosa viridis ramosissima, ramis erectis vel diffusis flexuosis, foliis alternis excentrica perfoliatis ramulorum sterili subglobosis fertili hemisphericis laminâ nullâ, ramulis nodulosomoniliformibus, spicis alternis sessilibus oblongis racemosis, areæ florigeræ floribus ternis inferne coalitis, seminis orbicularis testâ juxta radiculam papillis brevissimis obsoletis vestitâ cæterum levâ. *Halopeplis perfoliata* Bge. — *Halostachys perfoliata* Moq. in D.C. Prodr. ex parte.

Hab. in littore Aegyptiaco maris Rubri ad sinum Berenices (Schweinf!).

Frutex pedalis.

Ar. Geogr. Nubia et Arabia tropicæ.

HALOSTACHYS (C.A. Mey. in Hohen. Talysh Enum. p. 425).

Caules aphylli articulati. Folia opposita connata laminâ rudimentariâ. Bractæ oppositæ inter se liberæ demum deidue. Flores terni non connati. Perigonum gamosepalum obpyramidalum vertice convexum ampli grossæ trilobum. Semen albuminosum, embryo incurvus, radicula ascendens. — Frutex.

1. **H. Caspica** (Pall. Voy. App. p. 480, tab. A, fig. 2 sub *Salicornia*) perennis frutescens vel arborescens trunco erecto ramosissimo, ramis in ramulos erecto-patulos articulates repetitive divisis articulis elongatis cylindricis vel clavatis, foliis oppositis apice in cupulam fere obsoletam abeuntibus, spicis cylindricis oppositis cerebellim racemosis inferioribus pedunculatis, bractearum laminis obtusis subcordatis, areis florigeris trifloris, stamine unico, semine rhombo-elliptico 5 vel 3. *H. Caspia* Meyer in Schrenck Bull. Petrop. 1841. — *Halocnemum Caspicum* M.B. Taur. Gaue. III, p. 3. — *Artrocnemum Caspicum* et *Halostachys Caspia* Moq. in D.C. Prodr.

Hab. in salsis Armeniae Rossicæ (Szov!), Georgiae Caucasicae ad Sallian et in valle fluv. Kur (G.A. Mey! Hoh!), Persia boreali (Buhse! Bge), Turkestania (Lehm!).

Frutex saepe humanam altitudinem superans, spicæ interdum pollicares densissimæ.

Ar. Geogr. Rossia austro-orientalis ad Caspium, Songaria.

HALOCNEMUM (M. B. Fl. Taur. Cauc. III, p. 3 ex parte).

Caulis articulati aphylli. Folia opposita connata laminâ rudimentariâ. Bracteæ oppositæ inter se liberæ demum deciduae. Flores terni non connati spicati. Perigonum triphyllum elavatum phyllis inter se liberis apice fornicate-inflexis. Semen albuminosum, embryo incurvus, radicula ascendens. — Suffrutex.

1. **H. strobilaceum** (Pall. It. App. p. 481, tab. B, fig. 1, 2 sub *Salicornid*) perenne frutescens ramosissimum distortum humile, ramis ranunculique assurgentibus brevissime articulatis gemmis sterilibus globosis onustis, articulis crassiusculis breviter cylindricis vel subelatatis apiee in eupulam foliorum obtusilobam abeuntibus, spicis oblongis et cylindricis sessilibus lateralibus et terminalibus, bractearum laminâ semiorbiculatâ, areis florigeris trifloris, stamine unico ♂. Pall. III, tab. 4. — *H. strobilaceum* Moq. in p.c. Prodr. XIII, p. 149. — *Salic. cruciata* Forsk. Fl. Eg. Arab. p. 59.

Hab. in salsis maritimis et interioribus, insula Coreyra (Frider.), Thracia ad littora Egra (Jankat!), Tauria (Rehm!), Egyptus inferior (Gaill! Schweinf!) et ad littora maris Rubri (Schweinf! Aschers!), Arabia petrea ad Tor (Bovél.), Georgia Caucasica (Hoh!). Persia borealis ad Caspium (Buhse!) et ad lacum Urmiah (Szov!), Tureomania ad Caspium (Kar. et Kir!), Turkestaniae deserta Kisylkum (Lehm!).

Sepius humile pedale, interdum autem orgyale; facies *Arthrocnemi glauci* a quo præter characteres genericos articulis brevioribus copiose gemmiferis statim dignoscitur.

Ar. Geogr. Sardinia, Sicilia, Rossia australis, Songaria, Africa borealis interior.

SUBORDO II. SPIROLOBE.E C.A. Mey.

Embryo spiralis. Albumen parcum bipartitum vel nullum.

Ta. V. SUEDEÆ Moquin.

Flores perfecti vel polygami minute bracteolati. Fructus utricularis. Semen horizontale vel verticale integumento saepius duplice crustaceo et membranaceo. Embryo plano-spiralis, albumen nullum vel massulus binas laterales embryone discretas formans.

SUÆDA (Forsk. Fl. Eg. Arab. p. 62. — *Schoberia* C.A. Mey. — *Chenopodina*, *Brezia*, *Belowia*, *Calvelia* Moq.).

Flores hermaphroditici vel polygami solitarii vel saepius glomerulati bracteolis minimis squamiformibus membranaceis suffulti. Perigonum quinquefidum vel partitum ab ovario liberum, laciniis fructiferis vel immutatis vel carnosis, vel spongioso-auctis, vel dorso carinatis aut corniculatis aut subulatis. Stamina 5 perigonio inserta. Stigmata 2-3. Utriculus perigonio inclusus ab eo liber. Semina (saepè in eadem stirpe) horizontalia vel verticalia, albumine nullo vel parco. Cl. Bunge observante in speciebus pluribus semina dimorpha adsunt, alia præcociora integumento crustaceo, alia integumento membranaceo. — Herbae vel frutices. — Amic. A de Bunge in litteris observante in plurimis speciebus annuis *Suæda* sero autumno lectis occurunt semina integumento membranaceo, albumine prorsus nullo, embryone viridi et ideo *Belowia paradoxa* Bge in Lehm. p. 162 est probab. *Suæda* enjusdam formam autumnalis.

Specierum Orientalium dispositio (ex cl. Bge).

* Perennes fruticosæ.

+ Florum glomeruli supra axillam petiolo adnati.

S. microphylla, *dendroides*, *Asphaltica*.

+ + Florum glomeruli axillares.

1. Perigonum fructiferum inflatum.

S. physophora.

2. Perigonum fructiferum non inflatum.

S. fruticosa, *vera*, *monoica*, *vermiculata*.

* * Annuæ, interdum inferne induratae.

+ Foliorum glomeruli petiola adnati

S. altissima, *heterocarpa*.

+ + Foliorum glomeruli axillares.

1. Perigonii fructiferi laciniæ apteræ vel longitudinaliter carinatae.

S. maritima, *salsa*, *setigera*, *arcuata*. *Transoxana*. *microsperma*.

2. Perigonii fructiferi laciniæ transverse alatae.

S. heterophylla.

* Perennes fruticosæ.

1. S. microphylla (Pall. Ill. p. 52, tab. 44) fruticosa crispato-hirta coesia, ramis erectis in ramulos filiformes divaricatissimos decomposite ramulosissimis, foliis minutissimis breviter linearis-oblongis vel clavatis basi attenuatis vel subpetiolatis obtusis carnosis patentissimis, floribus 1-3 glomerulatis laxe et tenuiter spicatis pedicello brevi petiolo breviter adnato, perigonio fructifero parvo sicco obtuse pentagono-sulcato laciniis apice inflexis conniventibus, semine horizontali nitido rostellato ȝ. *Chenopodium Biebersteinianum* R. et Sch. Syst. VI, p. 267. — *Chen. crassifolium* M.B. Taur. Cauc. III, p. 176. — *Chenopodina microphylla* et *Ch. Biebersteiniana* Moq. — *Schoberia microphylla* C.A. Mey.

Hab. in salsis sterilibus prov. Caucasicarum ad Kislar (C.A. M!), in Georgia ad Sallian et planitiem fluvii Kur (Hohen. sub *Schang. altissimâ!*), Armenia (Koch).

Folia 1-2 lineas longa, fructus semine *Cannabeos* triplo minor, planta divaricatissima ramulis floriferis tenuissimis.

Ar. Geogr. Sibiria Altaica.

2. S. dendroides (C.A. Mey Enum. p. 159 sub *Schoberia*) fruticosa glaberrima ramosissima ramis erecto-patulis, ramulis hornotinis virgatis simplicibus elongatis exsiccalis angulato-striatis albidis, foliis carnosis semitereti-filiformibus obtusis basi attenuatis patulis saepe incurvis, glomerulatis 1-3-floris sessilibus axillaribus sed petioli basi adnatis, perigonii fructiferi obovato-globosi laciniis ovatis concavis membranaceo-marginatis apice flexis conniventibus, seminibus horizontalibus nitidis tenuissime punctato-striatis ȝ. *Chenopodium microphyllum* M.B. Taur. Cauc. III, p. 175 excl. syn.

Hab. in campis et collibus argilloso-salsis Transcaucasiae ad Caspiuni a Derbend ad flumen Cyrum (C.A. Mey!), Armenia Rossica (ex C. Koch).

Facies *S. fruticosæ* sed folia basi magis angustata non solubilia 4-8 lineas longa 1/2-1 lata, florum glomeruli ut in *S. microphylla* petiolo quidem sed ad hujus basin nec superius adnati, sessiles nec pedicellati.

Ar. Geogr. In Sibiria Altaica indicata sed dubia.

3. S. Asphaltica (Boiss. Diagn. Ser. I, XII, p. 98 sub *Chenopodina*), fruticosa ramosissima ramis erectis tenuiter virgatis duris striatis cinerascentibus, ramulis hornotinis patentissimis foliosis simplicibus vel parce ramulosis, foliis squamis albidis obsitis carnosulis filiformibus obtusis basi attenuata subpetiolatis, floralibus abbreviatis, glomerulatis 1-3-floris sessilibus petiolo adnatis in spicas longas laxas foliosas dispositis, perigonii minimi fructiferi laciniis ovatis margine membranaceis obtusis immutatis, semine horizontali nitido laevi ȝ.

Hab. ad littora septentrionalia lacus Asphaltitis Judeæ (Boiss!).

Frutex ramis lignosis sed tenuibus, folia 5-7 lineas longa lineæ 2/3 lata, *S. dendroidi* ab amic. Bge in litt. adnumerata sed differt foliorum squamulosis, ramulis tenuioribus, floribus minoribus petiolo supra ejus basin adnatis; a *S. microphylla* foliis, etc., distineta.

4. S physophora (Pall. Ill. p. 51, tab. 53) glabriuscula viridis fruticosa ramis albidis erectis vel ascendentibus superne fastigiatim vel pyramidatim ramulosis tenuibus divaricatis, foliis semiteretibus filiformibus obtusis longis articulato-deciduis, floribus hermaphroditis sessilibus subsolitariis in cymulas racemiformes breves axillares laxifloras dispositis, stigmatibus 2 minutis, perigonio laciniis ovatis obtusis fructifero globoso spongioso-inflato rubello, semine horizontali punctulato nitido erostri 5. *Schoberia suffruticulosa* Lessing ex Bge. — *Chenop. physophora* Moq.

Hab. in Persia borealis salsis ad Gendab jugi Elbrus (Buhse!), Turcomaniâ australi (Kareli!), Turkestaniâ (Lehm!).

Frutex 2-4-pedalis, folia 1-1 $\frac{1}{2}$ -pollices longa linea $\frac{3}{4}$ lata, racemuli 6-9 lin. longi.

Ar. Geogr. Rossie australis desertum Caspicum, Sibiria Uralensis et Altaica.

5. S. fruticosa (L. Sp. p. 324 sub *Salsola*) fruticosa humilis viridis, ramis erectis vel ascendentibus crebre ramulosis ramulisque crebre foliatis, foliis carnosulis crassiusculis semiteretibus linearicylindricis obtusis latitudine 4-6-plo longioribus, floralibus abbreviatis flores superantibus, floribus hermaphroditis axillaribus 1-3-nis in spicas densiusculas foliatas dispositis, pergoni subglobosi laciniis ovatis concavis obtusis apice incurvis fructiferis siccis conniventibus, semine saepius verticali leví nitido rostellato 5. Ic. Engl. Bot. tab. 635. — *Salsola annularis* Poir. Diet.

Hab. in salsis Gracie (Guérin!), Atticæ (Heldr!), Cretæ (Sieb!), Cypri (Ky. 284!), Syrie littoralis (Bl!), Egypti inferioris (Ehrenb! Samar! Kral!).

β. brevifolia. — Folia abbreviata tres lineas sicciter longa. — Ic. Fl. Græc. tab. 253.

Hab. cum typo in Gracie Syriæ littorali et Egypto ad Alexandriam (Gaill!), regno Cabulico (Griff. 1770!).

Omnia specimenia *S. Indice* Willd. quæ hucusque vidi vel ad *S. fruticosam* vel ad *S. monican* spectabant. *S. nudiflora* W. affinis quoque, ex India, specie ob folia floralia floribus breviora subnudis differre videtur et in ditione non crescit, nam specimen ex Bagdad Auch. 2776 a Moquinio huc citatum est *S. vermiculata*!

Ar. Geogr. Anglia, Batavia, Belgium, regio Mediterranea Europæ et Africæ borealis, Russia australis, Arabia tropica, India borealis, Insulae Canarienses, Madera, Senegal, P. B. Spei, America.

6. S vera (Forsk. Eg. Arab. p. 69) fruticosa humilis farinâ detergibili aspersa demum virenti-casia, ramis elongatis crebre ramulosis, foliis minimis crebris saepe subimbricatis anguste carooso-vermiculatis latitudine 2-3-plo longioribus, floralibus conformibus flores sub-superantibus, tioribus hermaphroditis axillaribus solitariis vel 2-3-nis in spicas longas tenues densiusculas dispositis, perigonii fructiferi globosi laciniis ovatis obtusiusculis fructiferis siccis conniventibus, semine saepius horizontali leví nitido rostellato i. *Salsola farinosa* Poir. Diet. 7, p. 298. — *Suæda pruinosa* Lange Pugill. Hisp. p. 93.

Hab. in Egypto prope Alexandriam (Forsk., Del. in D.C. herb!).

Rami erecti pedales vel diffusi flexuosi, folia 1-1½, lineas longa (in specimibus ex Oran Bal. exs. 419 tamen majora tres lineas longa) eis *S. fruticosæ brevifolice* multo tenuiora et magis imbricata, perigonum fructiferum minus.

Ar. Geogr. Hispania austro-orientalis (Bourg. 14691 et 862 sub *S. fruticosâ!*, Lange N° 164!), Africa borealis ad Mostaganem (Bal. 381 et 419!).

7 ***S. monoica*** (Forsk. Eg. Arab. p. 70) fruticosa arborescens viridis, ramis erectis elatis crebre ramulosis ramulis erecto-patentibus crebre foliosis, foliis carnosulis late linearibus planiusculis obtusis, paniculâ terminali compositâ brevi, floribus polygamis 2-3-gloemerulatis in spicas terminales laxiusculas dispositis, foliis floralibus ellipticis basi valde attenuatis inferioribus flores superantibus, perigonii fructiferi obovato-oblongi laciniis ovatis siccis conniventibus, semine lævi verticali nitido subrostrato 5. *S. Indica* Thw. Ceyl. Enum. non Willd. — *S. fruticosa* Forsk. Fl. Eg. Arab. p. 70 huc forsan ex cl. Ascherson spectat.

Hab. in littoribus Aegyptiacis maris Rubri inter Kosseir et Ras Benass (Schweinf!), Oasi magna et Oasi Dachel (Schw! Aschers!), Arabia petrea in valle Unsert prope Noueba (Schimp. 721!). Ex Forskahleo prope Alexandriam sed ut videtur erroneè indicata. Arabice *Assal*.

Arbuscula 15-pedalis, folia fere *Taxi* 6-10 lineas longa 1½ lata.

Ar. Geogr. Nubia, Abyssinia, Arabia tropica, India.

8. ***S. vermiculata*** (Forsk. Fl. Eg. Arab. p. 70) fruticosa viridis a basi diffuse ramosissima ramis divaricatis intricatis, foliis succulentis parvis basi valde attenuatis obtusissimis inferioribus obovato-oblongis, superioribus subglobosis, floribus hermaphroditis axillaribus solitariis vel 2-3-nis folio superatis laxe et breviter spicatis, perigonio fructifero ovato laciniis siccis apice incurvis conniventibus, semine verticali nitido non rostellato 5. Forsk. Ic. tab. 18, fig. B. — *Salsola mollis* Desf. Atl. I, p. 218. — *Suæda mollis* Del. Fl. Eg. — *Salsola globulifolia* Poir. Dict. 7, p. 298.

Hab. in salsis maritimis et desertorum, circa Alexandriam (Del.), in desertis prope Suez (Schweinf!) et ad littora Aegyptiaca maris Rubri inter Kosseir et Ras Benass (Schw!), Arabia petrea ad Ouadi Ghurundel (Boiss!), Babylonitæ deserta (Auch. 2776! Noë 102!).

Frutex intricatus ¼-1-pedalis, folia inferiora 1½-3 lin. longa, omnia teretia valde succulenta, fere *Sedi*.

Ar. Geogr. Insulæ Canarienses, Africa borealis, Nubia, India orientalis (Bélang. ex Moq.).

• Annuae, interdum inferne induratæ.

9. ***S. altissima*** (L. Sp. p. 324 sub *Salsola*) annua glabra viridis vel glaucescens, caule erecto a basi strictiuscule et pyramidatum ramoso, foliis tenuiter filiformibus longis acutiusculis basi attenuatis subpetiolatis, floralibus abbreviatis tenuissimis flores superantibus,

floribus homomorphis, glomerulis pauei vel plurifloris sessilibus vel subpedicellatis petiolo adnatis, perigonii fructiferi parvi subglobosi laciiniis ovatis obtusis concavis immutatis fructum includentibus, semine horizontali vel verticali lævi nitido rostellato ⊖. *Suæda altissima* Pall. Ill. tab. 42. — *Chenopodium altissimum* L. et Auct. — *Suæda altissima* et *Chenopodium altissima* Moq. in D.C. Prodr. — *Schoberia altissima* C.A. M. — *Schob. leiosperma* C.A. M. Ledeb. Ic. tab. 45. — *Suæda maxima* C. Koch Linn. XXII.

Hab. in ruderatis, salsuginosis, Thessalia ad Volo (Orpl!), Thracia borealis (Jank!), Anatolia ad Ieonium (Heldr!) et ad Cesaream in salsis (Bal!), Armenia ad Van (Kyl!), ad Balut (Bourg!), Tauria ad Sudak (Rehni!), Transcaucasia (Hoh. exs. sub *Ch. salso*!), Persia borealis (Buhse!), Babylonica (Noë!), Egyptus.

Ar. Geogr. Hispania, regio Danubialis, Rossia australis, Sibiria Uralensis et Altaica.

10. *S. heterocarpa* (Fenzl in Ledeb. Fl. Ross. III, p. 781) annua glabra glaucescens, caule erecto patule ramoso ramis virgatis elongatis, foliis semiteretibus anguste linearibus obtusis basi attenuatis flaccidis exsiccatis flexuosis basi non articulatis, glomerulis rarius 1 saepius 3-6-floris pedicellatis folio axillari adnatis et eo superatis, floribus heteromorphis omnibus fertilibus aliorum perigonio fructifero inflato spongioso globoso apice quinquesulcato seriatim celluloso nitido, aliorum parvo immutato obtuse pentagono fructum arte ininclude, semine horizontali vel verticali globoso-depresso nitido sub lente subtilissime punctato ⊖.

Hab. in salsis Transcaucasiae prope Sallian (Hohen! sub *Schanginiâ altissimâ* partim), spec. floriferum ex deserto Persico Hawuz i Sultan (Haussk!) hue spectare videtur.

Facie affinis *S. altissima* quæcum facile confunditur, insignis perigonis fructiferis dimorphis, aliis spongiosis fuscescensibus 1-2 lineas diam. latis, aliis nigrescentibus lineam dimidiam tantum latis.

11. *S. maritima* (L. Sp. 321 sub *Chenopodio*) annua glabra glauco-virens erecta a collo saepe ramosa, caulis valde ramulosis ramisque crebre foliosis, foliis carnosis semitereti-filiformibus obtusis flaccidis, floralibus valde diminutis flores superantibus, glomerulis 2-3-floris axillaribus in spicas longas tenues foliatas laxas digestis, perigonii fructiferi globoso-depressi phyllis ovatis obtusis fructum includentibus dorso obscure carinatis, semine saepissime horizontali rostellato sub lente minutissime punctulato ⊖. *Suæda maritima* Dumort. — Ic. Fl. Dan. t. 489. — *Salsola salsa* Jacq. H. Vind. III, p. 44, tab. 83 non Pall. — *Schob. maritima* C.A. Mey.

Hab. in maritimis salsis, Græcia ad Naupliam (Sart!), ad Phalerum (Heldr!), Macedonia et Thracia (Friv!), Tauria, prov. Caucasicæ et Transcaucasia!, Persia borealis (Auch. 2286!).

Ar. Geogr. Europa tota maritima arcticis exceptis, Africa borealis, insulæ Canarienses, America borealis.

12. **S. salsa** (Pall. Ill. p. 46, tab. 39) annua glabra glaucescens, a basi saepe pluricaulis caule erecto stricte ramoso, foliis succulentis semitereti-filiformibus obtusiusculis inermibus, floralibus glomerulos superantibus; glomerulis axillaribus compactis 2-5-floris confertim vel laxe spicatis, perigonio fructifero globoso laciniis succulentis ovato-orbiculatis obtusis concavis dorso rotundatis vel subcarinatis apice subgibbis, semine horizontali nitido laevi ⊖. *Schoberia salsa* C.A. Mey.

Hab. in salsuginosis Tauriae (Stev.), Georgiae Caucasicæ et Somchetiae (C.A. M., Hoh.), ditionis Talysh (Hoh.). Specimen floriferum ex lacu Nemek-darja prope Schiras (Ky. 959!) huc spectare videtur.

Valde affinis *S. maritimæ* a qua hujus specimina exsiccata ægre distinguuntur: perigonii phylla carnosissima, semen non punctulatum note differentiales graviores esse videntur. Fx cl. Bunge flores (aestivales) ramorum inferiorum ut in congeneribus pluribus semina integumento crustaceo donata, superiores antem (autumnales) multo majores semina integumento membranaceo edunt. Specimina Hispanica, Italica pro *S. salsa* determinata ad *S. maritimam* referenda sunt.

Ar. Geogr. Rossia australis, Sibiria Uralensis et Altaica.

13. **S. setigera** (D.C. Catal. Hort. Monsp. p. 94 sub *Chenopodium*) annua lepidoto-pulverulenta ramis erectis vel diffusis ramosissimis elongatis saepius paniculatis, foliis carnosis teretiusculis cylindrico-subulatis exsiccatis semipellucidis acuminatis saepe setulâ pellucidâ facile evanescente terminatis, floralibus abbreviatis flores subsuperantibus, glomerulis axillaribus confertissime 3-5-floris in spicas longas inferne laxas digestis, perigonii fructiferi carnosò-inflati laciniis ovatis obtusis dorso convexis fructu adpressis, semine verticali vel horizontali nitido laevi ⊖. *Suæda setigera* Moq. — *Chenopodina setigera* Moq. — *Schoberia setigera* C.A. Mey. Fl. Alt. I, p. 402. — *Chenopodium Clemente* Spreng. Syst. — *Suæda Balansæ* Boiss. Diagn. Ser. II, 4, p. 75. — *Schoberia acuminata* C.A. Mey. loc. cit. p. 198 et Ledeb. Ic. tab. 44. — *Suæda gracilis* Moq. in D.C. Prodr. I. cit. p. 157. — *Salsola clavifolia* Hoh. exs. non Pall. — *Belovia haccifera* Moq. in D.C. Prodr. p. 168 (ex cl. Bge) forma netrocarpa. — *Suæda splendens* Gr. et Godr. Pl. Fr. III, p. 30. — *Suæda monantha* C. Koch Linn. XVII, n. 313. — *S. longifolia* G. Koch Linn. XXII, p. 188.

Hab. in maritimis Ætolie prope Missolonghos (Heldr. 1874!), ad Volo Thessaliae (Orphl.), in salsis Cappadociae prope Cæsaream (Bal!), Armeniae Rossicae (C. Koch!), in Egypto ad Damiatam (Sieb. sub *S. salsa* partim ex Fenzl.), Georgiae Caucasicæ ad Elisabethpol (Hohen!), Babyloniam ad Bagdad (Noe 311!).

Foliis acuminato-mucronatis vel setigeris exsiccatis subpellucidis a formis *S. maritimæ* primâ facie distinguenda. Sæpe ut in pluribus congeneribus adsunt semina dimorpha, alia præcociora crustacea, alia variora serotina integumento membranaceo.

Ar. Geogr. Hispania australis et orientalis, Gallia meridionalis, Sardinia, Turkestania borealis, Sibiria Altaica.

14. **S. arcuata** (Bge Pl. Lehm. p. 461) annua inferne indurata viridis glabra erecta a basi ramosissima, ramis ramulisque erecto-patulis tenuibus firmis, foliis linearis-teretibus basi attenuatis obtusiusculis arcuato-inflexis, glomerulis dense 3-6-floris sessilibus axillaribus folio florali superatis, floribus polygamis lateralibus minutissimis femineis quinquedentatis, cæteris hermaphroditis vel masculis quinquepartitis laciniis membranaceis trinerviis oblongis apice cucullatis, semine verticali lèvissimo nitido breviter rostellato ♂.

Hab. in Turkestaniâ ad Jan Darja (Lehm!).

Bipedalis vel elatior, folia 6-8 lineas longa.

15. **S. Transoxana** (Bge Pl. Lehm. p. 467) annua inferne indurata glaucescenti-cana ramosissima, ramis ramulisque erectis gracibus firmis, foliis planis linearis-oblongis basi attenuatis obtusis latitudine suâ 2-3-plo longioribus, glomerulis 4-5-floris axillaribus folio florali brevi superatis in racemos laxos dispositis, perigonii quinquepartiti laciniis sub anthesi herbaceis oblongis apice cucullatis dorso acute carinato-alatis, fructu . . . ♂.

Hab. in cultis ad Buchara Turkestaniæ (Lehm!).

Facies et racemi graciles *S. arcuatæ* sed folia plana, inferiora vix 3 lineas longa. Fructus ignotus sed jam florifera perigonii laciniis longitudinaliter carinato-cristatis insignis.

16. **S. microsperma** (C.A. Mey. in Eichw. Casp. p. 14, tab. 13 sub *Schoberiad*) annua a basi ramosissima erecta vel prostrata glaberrima glauca, ramia et ramulis erecto-patulis, foliis carnosis semiteret-linearibus superne latioribus obtusis, floralibus abbreviatis flores vix superantibus, floribus secus ramulos tenuissimos solitariis 2-3-nis glomerulatis minimis distinctis, perigonii depresso-globosi fructiferi exsucri laciniis ovatis eximie cucullato-carinatis obtusis, semine verticali vel horizontali lèvissimo nitidissimo ♂. *S. microsperma* Fenzl in Led. Fl. Ross.

Hab. ad oras lacūs Aralensis australioris ad ostium fluvii Oxus (Bge).

Ar. Geogr. Songaria.

17. **S. heterophylla** (Kar. et Kir. Enum. № 749 sub *Schoberiad*) annua glabra glauca, caule herbaceo erecto simplici vel stricte et simpliciter ramuloso. foliis semitereti-filiformibus acutis erectis vel incurvo-patulis, floribus 1-3 axillaribus sessilibus folio florali brevi superatis a ramulorum basi angustissime et laxiuscule spicatis, perigonii semiquinquefidi globosi demum aucti et subspongiosi laciniis ovatis concavis basi transversim gibbo compresso alæformi auctis, semine sæpius horizontali minutissime punctato nitido ♂. *Suæda heterophylla* Bge in litt. — *Brezia heterophylla* Moq. in D.C. Prodr. p. 167.

Hab. in valle Bamian regni Cabulici (Griff. 1771!).

Affinis *S. corniculatae* Bge = *Schoberiæ corniculatae* C.A. Mey. intra fines nostros non observatæ laciniis fructiferis apice fornicato-corniculatis distinctæ.

Ar. Geogr. Sibiria Altaica.

**SCHANGINIA (C.A. Mey. in Ledeb. Fl. Alt. I, p. 394.
— *Suædæ* Sp. Auct.)**

Flores polygami vel hermaphroditæ, solitarii vel sæpius glomerulati bracteolis minimis squamiformibus membranaceis suffulti. Perigonum urceolatum ovario adnatum laciniis quinque fructiferis tuberculo auctis aut inflatis vel immutatis. Stamina 5 tubo perigonii inserta. Stigmata 2-3. Utriculus membranaceus perigonio adnatus. Semen verticale crustaceum exalbuminosum basi rostratum. — Herbæ vel suffrutices facie *Suædæ*.

1. **Sch. linifolia** (Pall. Ill. p. 47, tab. 40 sub *Suædæ*) annua glaberrima glauca, caule erecto stricte folioso superne in paniculam elongatam abeunte, foliis alternis succulentis planis linearibus elongatis acutis floralibus brevissimis, floribus polygamis glomerulatis secus ramulos paniculæ laxe spicatis, perigonii laciniis ovatis concavis conniventibus demum infra apicem tuberculo auctis, perigonii fructiferi tubo minute pedicelliformi, semine pyriformi-globoso subtilater punctato-rugoso ♂. *Sch. linifolia* C.A. Mey. Fl. Alt. I, p. 395. — *Sch. tribracteatu* C. Koch Linn. XXII, p. 188! — *Chenopodium linifolium* Schult.

Hab. in salsis, districtus Chynys Armeniæ (C. Koch!), in salsis et cultis ad Buchara Turkestaniæ (Lehm!).

Folia 6-10 lineas longa vix lineam lata, perigonum fructiferum durum crustaceum semini arte adhærens semine *Cannabeos* triplo minus.

Ar. Geogr. Rossia australis, Sibiria Altaica.

2. **Sch. baccata** (Forsk. Eg. Arab. p. 69 sub *Suædæ*) annua glabra glauca plus minusve farinosa a collo ramosa ramis crebre foliosis lateralibus sæpe diffusis centrali erecto, foliis carnosis semi-teretibus linearci-cylindricis obtusis patentibus flexuosis, floralibus sensim abbreviatis flores longe superantibus, floribus hermaphroditis glomeratis spicas longas foliosas inferne laxas superne densifloras formantibus, perigonio fructifero turbinato inflato-baccato sicco spongioso laciniis ovatis valde concavis apice dorso incrassatis clausis, semine rostrato lævissimo ♂. *Sch. baccata* Moq. Chenop. p. 119. — *Salsola baccata* Poir. — *Enchylæna Ægyptiaca* Spreng. Syst. I, p. 923. — *Suæda setigera* β. *mutica* Schimp. exs. 438.

Hab. in cultis ad vias Egypti circa Alexandriam (Forsk., Del., Gaill!), ad Cahiram (Schweinf!), supra Theben (Boiss!), in oasi Dachel et prov. Fayoum (Asch!), in oasi Chargeh (Schw!), valle Hebron Arabiæ petreæ (Schimp. 438!).

Folia 6-9 lineas longa linea $\frac{1}{4}$ diam. lata nec ut *Forskahleus* ovata. Peri-

gonium fructiferum magn. seminis Cannabeos apice quinquegibbum inflatum spongiosum.

Ar. Geogr. Arabia tropica.

3. Sch. hortensis (Forsk. Fl. Eg. Arab. p. 71 sub *Suædd*) perennis suffruticosa glabra parce farinoso-glaucescens erecta patule et longe ramosa ramis albis induratis, foliis linearibus glabris acutiusculis elongatis, floribus hermaphroditis ad axillas glomerulatis spicas foliatas inferne laxas formantibus, perigonii demum basi attenuati laciniis concavis immutatis divergentibus, semine pyriformi rostrato nitido nigro obsolete punctato ♂. *Schanginia hortensis* Moq. Chenop. — *Sal-sola divergens* Poir. Dict. 7, p. 299. — *Chenopodium hortense* Rœm. et Sch. Syst. 6, p. 268.

Hab. in Egypto ad Birket el Hadj (Schweinf!), in deserto prope Bagdad (Noël), in hortis saepe culta.

Cubitalis, inferne fruticoso-indurata, differt a præcedentibus perigonio fructifero immutato non inflato nec tuberculato-gibboso.

Ar. Geogr. Arabia tropica.

BIENERTIA (Bge in litt. — *Schanginiæ* Sp. Fenzl.).

Flores abortu polygami, præcociores hermaphrodoti et nonnulli feminei supra basin imam folii axillaris glomerulati, cæteri omnes feminei in racemos breves glomerulum superantes dispositi. Perigonum ovario adnatum sub anthesi depresso-subglobosum apice lobis 5 planis conniventibus, fructiferum valde depresso circumcircum in alam horizontalem circularem spongiosam expansum lobis minutis spongiosis in medio disco verruciformibus. Stamina 5 inclusa. Stigma 2-3 brevissime clavata. Utriculus tenuiter spongiosus perigonio adnatus seminique arcte adhærens. Semina horizontalia crustacea tenuiter granulata, præcoccia turgida biconvexa, serotina inter cotyledones et radiculam canaliculata. Albumen in utroque embryonis latere parcum. — Herba annua facie *Suædæ* et *Schanginiæ*.

1. B. cycloptera (Bge in litt.) annua sed inferne lignescens a collo ramosa caulis basi induratis ascendentibus crassiusculis, foliis carnosis sessilibus linearis-subteretibus obtusis saepe recurvis, superioribus abbreviatis omnibus caducis, paniculæ angustæ racemis brevibus divaricatim et breviter ramulosis 6-10-floris carnosis exsiccatis ut et paniculæ rachis angulato-subalatis, fructus orbiculati ala circulari semine ipso subangustiore spongiosa cellulosa subradiata margine obtusa et obsolete erosula ⊖.

Hab. in Persia orientali frequens (Bge).

Semipedalis et pedalis, folia inferiora 6-10 lineas longa, panicula 5-6-pollicaris angusta racemis ramulosis rigidulis 1-1/2-pollicem longis. Perigonum fructiferum cum ala tres lineas diam. latum lenticulare, florum serotinorum

sæpe fatuum. Planta certe a *Suedâ bacciferâ* Pall. Ill. p. 48, tab. 41 ex descriptione, icono, loco natali omnino aliena, nec magis ut videtur ad *Schangi-niam baccifera* Fenzl in Ledeb. que nobis ignota est referenda.

TR. VI. SALSOLEÆ Moq.

Flores bibracteati, sæpissime hermaphroditi, rarissime polygami. Fructus utricularis rarius baccatus. Semen horizontale vel verticale integumento simplici membranaceo. Embryo conico-spiralis. Albumen nullum.

SUBTR. I. SODEÆ Moq.

Semen horizontale.

TRAGANUM (Del. Fl. Eg. III. № 342).

Flores hermaphroditi bibracteati axillares. Perigonium quinquefidum fructiferum inferne incrassatum et nucamentaceum antice et postice gibbo corniculiformi conico obtuso duro auctum, laciinis membranaceis apteris. Discus parvus. Staminodia nulla. Stamina 5 filamentis latis exsertis, antheræ sagittatæ acutæ. Stylus bifidus stigmatibus filiformibus. Utriculus subdepressus perigonio lignoso inclusus, pericarpio membranaceo. Semen horizontale. — Fruticulus ramosissimus non articulatus.

1. **T. nudatum** (Del. loc. cit. p. 230, tab. 22, fig. 1) suffruticosum patule et divaricatum ramosissimum ramulis alternis intricatis gracilibus albidis glabris vel minute seabridis, foliis alternis minutis distantibus carnosis oblongo-triquetris mucronatis subrecurvis supra plicatis subtus carinatis, axillis lanatis sæpe fasciculiferis, floribus 1-3 glomerulatis, bracteolis ovato-triquetris flore florali vix brevioribus, perigonii laciinis membranaceis rigidulis oblongo-lanceolatis obtusis erectis 5.

Hab. in arenosis desertorum Egypti (Del., Sieb!), ad Cahiram (Wiest 501!), ad Pyramides (Boiss!), in deserto Ægyptiaco-Arabico (Schw!), in Arabiæ petreæ deserto Tih (Boiss!), deserto Palestiniæ ad meridiem Gaza (Boiss!).

Fruticulus a basi intricatum ramosissimus tortuosus ramulis novellis elongatis, folia 2-2½ lineas longa. Nux (in icono Delileanâ pessime exhibita) ovata longitudinaliter striata basi rotundata corniculis binis rectis crassis obtusis inæquilibus laciinis subbrevioribus antice et postice instructa, 2½ lin. longa 1¼ diam. lata, matura facile decidua. An *T. Moquini* Webb e Canariis specificie differat nescio, hujus fructus ignotus.

Ar. Geogr. Africa borealis.

HORANINOWIA (Fisch. et Mey. in Schrenk Ber. p. 40).

Flores hermaphroditi bibracteati minuti axillares solitarii. Perigonum pentaphyllum phyllis fructiferis supra medium rectangule infractis angulo dorsali in gibberem horizontalem turgidum producto, infra gibbum suberoso-incerassatis superne scariosis conniventibus. Staminodia quinque orbiculata fimbriata in urceolum cum staminibus 5 connata, filamenta setacea inclusa, antherae muticæ vel appendiculatæ Styli 2 breves subeoaliti subelavati apice stigmatosi. Fructus utricularis perigonio inclusus. Semen horizontale. — Herbae annuae non articulatae, foliis subulatis pungentibus, floribus ramulorum florarium brevitate in glomerulos squarroso-foliolosos congestis.

1. H. ulicina (F. et M. loc. cit. p. 11) annua viridis scabrida a collo ramosa ramis tenuibus diffusis et erectis opposite ramulosis, internodiis inferioribus elongatis, foliis oppositis a basi dilatata subulato-triquetris patentibus vel recurvis, glomerulorum squarrosis longius spinosis, glomerulis ad ramulorum ortum remotis et secus ramulos magis approximatis sessilibas, bracteolis abbreviatis ovatis aristatis, perigonii phyllis ovato-lanceolatis acutis apice scarioso trianguli-ovato, autheris muticis ♂. *H. juniperina* Bge in pl. Lehm. 468 non C.A. Mey. (ex ipso).

Hab. in desertis arenosis Karakum et ad fluvium Kuwandarja Turkestaniae australis (Lehmi), Turcomania ad Caspium (Karel!).

Semipedalis vel pedalis, folia 3-6 lineas longa, glomeruli oppositi pisces minores, perigonium sub anthesi $1\frac{1}{4}$ lineam longum.

Ar. Geogr. Sibiria Uralensis et Altaica.

2. H. juniperina (C.A. Mey. in Schrenck Enum. p. 11) annua viridis scabrida a basi ramosa, ramis diffusis et erectis alternatim et tenuiter ramulosis, foliis alternis sed approximatis veluti verticillatis triquetro-subulatis pungentibus patentibus, floralibus brevioribus, bracteolis minutis ovalis mucronulatis vel muticis, floribus solitariis vel 2-3 minute glomerulatis secus ramulos breves subdistantibus, perigonii phyllis ovato-lanceolatis apice ovatis scariosis acutis, antheris muticis ♂. *Salsola anomala* C.A. Mey. in Eichw. Casp. p. 14, tab. 12. — *Hor. anomala* Moq.

Hab. in Turcomaniâ australi ad littus Caspii prope Krasnowodsk (Eichw.), in Persia orientali mediâ (Bge!).

Facies praecedentis, folia minora vix 3 lineas excedentia; ambo ab *H. minore* Schrenck in ditione non obvia antheris non micronatis differunt.

ARTHROPHYTON (Schrenck Bull. Acad. Petersb. II, p. 241).

Flores hermaphroditi bibracteati oppositi axillares sessiles. Perigonum pentaphyllum phyllis transverse gibbis. Staminodia quinque

cum staminibus 5 in urceolam connata, antheræ muticæ. Styli duo brevissimi. Fructus perigonio inclusus utricularis depressus. Semen horizontale. — Suffrutex nanus facie *Brachylepidis*, ramis ex trunko crasso brevibus articulatis. — Genus characteribus *Horaninowiae* affine sed habitu, ramis articulatis distinctum. Fructus immaturus tantum notus.

1. **A. subulifolium** (Schr. loc. cit.) minutissime cinerascens fruticulosum, ramis ex caudice lignoso crasso numerosis rectis tenuibus pumilis simplicibus ad articulos fragillimis, foliis oppositis tenuibus brevibus a basi sublatiore subulato-triquetris mucronatis, floribus parvis in axillis oppositis, bracteolis foliis floralibus brevioribus basi membranaceo-dilatatis, perigonii phyllis orbiculatis concavis late membranaceis ♂. *Anabusis affinis* Bge in Pl. Lehm. 480 ex ipsol non F. et M.

Hab. in deserto Kisylkum et monte Ssususkara Turkestaniæ australis (Lehm!).

Rami bipollicares, folia 3-4 lineas longa.

Ar. Geogr. Songaria.

HALOXYLON (Bge Pl. Lehm. p. 468. — *Salsolæ* et *Caroxyli* Auct. species articulatæ).

Flores hermaphroditæ bibracteolati. Perigonium quinquepartitum phyllis post anthesin auctis demum dorso in alas scariosas transversas omnes patulas excrescentibus. Staminodia quinque in urceolum cum filamentis connata apice libera. Antheræ muticæ. Stylus brevissimus vel nullus, stigmata 2-4. Utriculus turbinatus superne concavus carnosulus perigonio aperto inclusus. Semen horizontale. — Arbusculæ vel frutices ligno durissimo, ramulis articulatis, foliis sæpius oppositis connatis rudimentariis vel brevissimis, floribus solitariis axillaribus oppositis. — Genus a *Salsola* ramulis articulatis, staminodiis, filamentis monadelphis distinctum.

1. **H. Ammodendron** (C.A. Mey. Fl. Alt. I, p. 375 sub *Anabasi*) arborescens trunko crasso, ramulis annotinis pallide virentibus tenuibus, foliis ad articulorum brevium apicem squamiformibus breviter et obtuse triangularibus in cupulam intus breviter lanuginosam connatis, floribus remotis laxe et breviter spicatis, staminodiorum cupulâ ovarium superante, stigmatibus quaternis sessilibus, alis magnis suborbiculatis ♂. Ledeb. Ic. tab. 47. — Bge in Pl. Lehm. p. 469. — *H. Persicum* Bge in Buhse Aufz. p. 189. — *Pinus Orientalis* Falk Topogr. Beitr. 2, p. 260) determinatio ex ramulis annotinis insecti cuiusdam ictu deformatis strobiliformibus orta) Vulgo *Saxaul*.

Hab. in arenosis magni deserti salsi inter Djendack et Enarek (Buhse!), Turkestaniæ (Lehm!), Turcomaniæ littorali!

Arbuscula 12-20-pedalis, interdum tamen humilior.

Ar. Geogr. Sibiria Uralensis et Altaica.

2. **H. articulatum** (Cav. Ic. III, p. 43, tab. 284 non Forsk sub *Salsola*) glabrum viride fruticulosum humile, foliis ad articulorum apicem squamiformibus breviter triangularibus in cupulam intus breviter villosulam coalitis, staminodiis ovario brevioribus, stylo brevi, stigmatibus binis divergentibus, alis majusculis subæqualibus obovato-orbicularibus margine erosulis 5. *Curoxylon articulatum* Moq. in D.C. Prodr.

Hab. in deserto Tih Arabie Petreæ (Boiss! specimina nondum florifera), Egyptus (Redonté ex Moq.), prope Damascum (Gáilli! 2199 specimina juniora). Specimina mea floribus et fructu orbata subdubie ad *H. articulatum* refero, sed folia, habitus, omnino eadem.

Ar. Geogr. Arabia subtropica (Ehrenb!), Africa borealis ad Biskra, regnum Marocanum, Hispania austro-orientalis.

3. **H. multiflorum** (Moq. in D.C. Prodr. loc. cit. p. 212 sub *Anabasi*) fruticulosum glaucescens ramosissimum ramis erectis brevibus, foliis squamiformibus brevibus triangularibus crassiuseulis obtusis in cupulam coalitis, floribus dense et breviter spicatis, bracteolis amplis orbicularibus concavis cupulam superantibus, staminodiis brevibus subcartilagineis, stylo crasso stigmatibus lanceolatis, alis magnis semiorbiculatis 5. *H. multiflorum* Bge in litt.

Hab. in regno Cabulico (Griff. 1786!).

Fruticulus semipedalis, rami annotini bipollicares. Differt a precedentibus spicis densis pollicaribus, bracteolis magis evolutis.

4. **H. salicornicnm** (Moq. in D.C. Prodr. I. cit. p. 174) fruticosum glaucescens ramis erectis crassis crebre et stricte ramulosis paniculatis, foliis squamiformibus breviter triangularibus margine membranaceis in cupulam intus longe barbatam coalitis, floribus remotiusculis longe spicatis articulis subclavatis, bracteolis ovatis acutis basi lanatis, stylo brevi stigmatibus subclavatis, fructus alis majusculis obovato-orbicularis 5. *H. salicornicum* Bge in litt.

Hab. in regno Cabulico ad Baricab, Baissoul, Hugarnow (Griff. 1796!), in Belutschia (Stocks 1218!), in Persia Alghaniæ contermina ad Tebbes (Bge!).

Rami eis *H. multiflori* crassiores longiores paniculatum multispicati, spicæ tenues 2-3-pollicares. Ante a Bge prius plantam ex Tebbes ut speciem propriam sub *H. elegantis* nomine descripserat sed nunc ad *H. salicornicum* adducit.

5. **H. recurvum** (Wall. Cat. 6943 sub *Salsola*) glabrum pruinosum, ramis erectis crassis patentim et opposite ramuloso-paniculatis, foliis teretibus crassis brevibus mucronulatis vel obtusis patentibus vel recurvis, floribus dissitis longe spicatis, foliis floralibus ovatis bracteolas ovatas acutas superantibus, staminodiis cartilagineis a filamentorum cupula fere liberis, fructus alis mediocribus 5.

H. recurvum Bge in litt. — *Salsola Stocksii* Boiss. Diagn. Ser. II, 4, p. 75. — *Salsola Lana* Stocks exs. non Edgew. (quæ ex am. Bge est *Schoberia Indica*).

Hab. in Belutschia (Stocks!).

Paniculæ semipedales spicis fructiferis 1-1 $\frac{1}{2}$ -pollicaribus, alæ lineam longæ.

Ar. Geogr. Ditio Scinde, Indi atrópica.

6. ***H. Griffithii*** (Moq. in D.C. Prodr. I. c. p. 175 sub *Caroxylo*) albido-glaucescens fruticosum ramosissimum ramis erectis tenuibus rigidis albidis strictiuscule ramulosis, foliis breviter linearibus obtusiusculis erectis vel subincurvis superioribus ad squamas reductis omnibus oppositis, floribus remotiusculis laxe et tenuiter spicatis, staminodiis ovario brevioribus, stylo brevi, stigmate 2-4-lobo, fructus alis purpurascens obovato-reniformibus erosulis 5. *H. Griffithii* MSS.

Hab. in regni Cabulici iugno Kullou (Griff. 1792-93!), in Belutschia (Stocks 1491!).

Frutex ut videtur pedalis, folia 2 lineas longa superne concaviuscula subtus convexa. Species affinis sub nomine *H. Griffithii* ex herb. Ind. Comp. communicata e Tibetia est *H. Thomsonii* Bge in litt., foliis filiformibus 5-7 lineas longis, summis sparsis, stigmatibus binis elongatis ab *H. Griffithii* distincta.

SEIDLITZIA (Bunge in litt. — *Haloxylon* Sp. Fenzl).

Flores hermaphroditici bibracteati axillares oppositi solitarii vel 2-3-ni, laterales saepe tabescentes. Staminodia quina semiobtusiusculata glanduloso-ciliata in discum staminiferum coalita, filamenta 5 exserta, antheræ muticæ. Perigonii phylla quina basi connexa elliptica obtusa vel biloba, fructifera basi chartacea versus medium alis amplis membranaceis transversis absita, ala postica cochleariformis antice reflexa fructum tegens, cæteræ patentes. Stylus minutus apice bistigmatosus. Utriculus depresso. Semen horizontale. — Herba annua vel suffrutescens caule ramisque articulatis, foliis oppositis. — Genus *Haloxylo* affine ab eo potius habitu quam characteribus gravioribus distinctum.

1. ***S. florida*** (M.B. Taur. Cauc. I, p. 190 sub *Anabasi*) annua succulenta glaberrima glauca, caule erecto a basi brachiatim ramoso ramis simplicibus vel parce ramulosis patentibus vel decumbentibus, articulis inferioribus elongatis, foliis oppositis teretiusculis crassis brevibus obtusis patentissimis demum reflexis, braeoleolis ovatis obtusis folio florali superatis naviculari-concavis obtusis margine membranaceis, floribus parvis basi folii floralis immersis. Perigonii fructiferi alis roseis obovato-orbiculatis amplis basi angustatis, florum lateralium (cum adsunt) tabescientium alis minutis squamiformibus ☽. Ic. M.B. Cent. Ross. tab. 17. — *Seidlitzia florida* Bunge in litt. — *Salsola florida* Poiret Encyc. Suppl. V, p. 191. — *Anabasis cine-*

rea Moq. in D.C. Prodr. p. 214 ex Bge. — *Chenopodina pycnantha* C. Koch.

Hab. in aridis Armeniae Ibericæ ad fluv. Kur (M.B.), ad Erivan (Schuller!), in Somchetiâ (Eichw.), in Persia bor. prov. Aderbidjan (Szov!), circa Teheran (Oliv.), ad Murchehar prope Ispahan (Haussk!).

2. **S. Rosmarinus** (Ehrenb. MSS. sub *Suedii*) fruticosa opposite ramosissima ramis oppositis glabris albidis, foliis oppositis linearisemiteretibus obtusissimis flaccidulis, internodiis inferioribus elongatis floralibus abbreviatis, glomerulis axillaribus oppositis 3-5-floris folio florali abbreviato superatis, floribus villo brevi stipatis 5. *S. Rosmarinus* Bge in litt.

Hab. in insula Maksuk Arabiae (Ehr. № 1231).

Internodia inferiora pollicaria, folia inferiora 8-9 lin. longa lineam lata, floralia 2-3 lineas longa. Staminodia ex cl. Bge distincta, antheræ bipartite loculis apice tantum contiguis brevissime et obtuse apiculatis. Flores in speciminiis Ehrenbergianis juniores et fructus nulli, sed specimen fructiferæ facie et foliis simillima e Persia ad Hawuz i Sultan inter Ispahan et Teheran a cl. Haussknecht lecta ad speciem eamdem vel valde affinem spectare videntur. In illis aliae rufescentes obovato-orbiculatae subintegrae patentissimæ sesquilineam longæ et latæ tubo turbinato-hemisphaericō longiores.

SALSOLA (L. Gen. № 311 ex parte. — *Salsola* et *Caroxylon* Moq.).

Flores hermaphroditæ (rarissime polygamæ) axillares bibracteati saepius solitarii. Perigonium ultra medium vel fere ad basin quinquepartitum (rarissime 5-fidum), basi induratum vel membranaceum, laciinis demum auctis dorso transverse membranaceo-alatis (rarius carinatis). Discus hypogynus cupularis plus minus manifestus. Staminodia nulla. Stamina 5 (rarissime in flor. masc. 4) sub anthesi exserta, antheræ muticæ vel connectivo producto mucronatæ aut appendiculatae. Stylus bieruris. Fructus utricularis depresso-exsuetus rarius baccatus. Semen horizontale. — Herbae annuae vel frutices ramis non articulatis.

Specierum Orientalium dispositio.

SERIES I. — Annuæ.

* Oppositifoliae.

S. brachiata.

** Sparsifoliae. — Folia et rami alterna.

+ Perigonium fructiferum patens. Utriculus carnosus. Stigmata sessilia brevissima.

S. clavifolia.

+ + Perigonum fructiferum clausum. Utriculus siccus. Stigmata filiformia elongata.

1. Antheræ non appendiculatæ.

S. Soda, Sogdiana, Kali, spissa, tamariscina.

2. Antheræ appendiculatæ.

S. inermis, carinata, gossypina, crassa, lanata.

SERIES II. — Perennes.

* Oppositifoliæ. — Folia et rami opposita.

S. tetragona, longifolia.

** Sparsifoliæ. — Folia et rami alterna.

+ Perigonum basi induratum truncatum quinquefoveolatum.

S. Auricula, lancifolia, glauca, brachyphylla, subaphylla, hispidula.

+ + Perigonum basi membranaceum non foveolatum.

1. Antheræ punctiformi-apiculatæ vel muticæ.

S. Arbuscula, gemmascens, fletida, verrucosa, ericoides, vermiculata, larinica, rigida.

2. Antheræ distincte appendiculatæ.

S. canescens, aurantiaca, Persica.

SERIES I. — Annuæ.

* Oppositifoliæ.

1. **S. brachiata** (Pall. Ill. p. 30, tab. 22) annua pubescens serius glabrata prætereaque pilis patentibus plus minus hirsuta, caule erecto fere a basi opposite et brachiatim ramoso, ramis longe virgatis, foliis præter summa interdum alterna oppositis remotis semitereti-filiformibus carnosis acutis setâ fugaci terminatis recurvo-patentibus, floralibus abbreviatis recurvis bracteolas superantibus, floribus 2-3-nis fasciculos axillares oppositos formantibus, bracteolis ovato-lanceolatis, perigonii laciinis anguste lanceolatis longe acuminate, fructiferi alis late reniformi-obovatis marginibus imbricatis castaneis, antherarum appendiculâ oblongâ obtusâ cochleariformi loculis angustiore et breviore ⊖. *S. Pallasiana* Vest. — *S. pulla* C. Koch Linn. XXII, p. 190.

Hab. in salsis prov. Caucasicarum (C.A. Mey!), regni Cabulici (Griff!).

Ar. Geogr. Rossia australis, Sibiria Uralensis et Altaica.

• Sparsifoliae.

2. *S. clavifolia* (Pall. It. II, p. 486) annua glabra siccitate nigrescens, caule erecto crasso simplici vel saepius fere a basi ramosissimo ramis virgatis crebre foliatis erectis, foliis alternis carnosis breviter subclavato-cylindricis obtusissimis muticis vel obsolete micronatis patulis saepe incurvis, floralibus bracteolis et floribus longioribus, floribus 2-3-globeratis confertim spicatis, bracteolis ovato-ellipticis late membranaceis, antheris non appendiculatis, perigonii fructiferi patentis supra utriculum nou clausi tubo hemispherico alis infra apicem ortis reniformibus stramineis, phyllis supra alarum insertionem membranaceis semiorbiculatis, styli subnulli erubibus brevissimis, utriculo succoso ☽. *S. baccifera* Pall. Ill. tab. 23. — *S. foliosa* Schrad. in Schult. Syst. XI, p. 235. — *Anabasis foliosa* Pall. — *Anab. clavata* Gmel.

Hab. in Turcomaniâ littorali (Gmel.).

Ar. Geogr. Rossia austro-orientalis, Sibiria Uralensis et Altaica.

3. *S. Soda* (L. Sp. 323) annua glabra, caule a basi diffuse ramoso ramis oppositis vel alternis elongatis incurvo-ascendentibus, foliis infimis oppositis cæteris alternis a basi dilatata semiamplexicauli elongatis carnosis semiteretibus setulâ terminatis flaccide patentibus, floralibus bracteolas ovato-lanceolatas floribus longiores longe superantibus, floribus axillaribus solitariis remotis, perigonii ultra medium quinquepartiti lacinis lanceolatis membranaceis demum auctis, fructifero magno turgido membranaceo basi obsolete quinquefoveolato alis ad tuberculos vel cariunam transversam reductis, fructu maximo turgido ☽. Ic. Jacq. Vindob. tab. 68. — Pall. Ill. tab. 30.

Hab. in maritimis salsis, Graecia in Aëtolia (Reinh!), Cephalonia (Heldr!), Attica (Urv!), Transcaucasia! Persia borealis ad Caspium (Buhse!).

Ar. Geogr. Europa media et australis a Gallia occidentali et meridionali ad Rossiam australem, Sibiria Uralensis, Japonia.

4. *S. Sogdiana* (Bge Pl. Leh. p. 473) annua glaberrima, caule crassiunculo erecto a basi dense et pyramidatim ramoso ramis ascendentibus simplicibus, foliis alternis longis linearis-triquetris carnosis breviter cuspidato-subpungentibus, floralibus abbreviatis perigonio longioribus, floribus solitariis axillaribus remote spicatis, bracteolis ovatis acutis perigonio subbrevioribus, perigonis fructiferis cartilagineis basi hemisphericis infimis apteris cæteris alatis aliis valde inaequalibus tribus amplissimis late obovato-reniformibus sese invicem tegentibus cæteris minutis spathulatis vel dentiformibus, lacinis parte liberâ tenuissime membranaceis lanceolato-linearibus patentibus ☽. *S. heterocarpa* Bge MSS. olim.

Hab. in deserto Kisilkum Turkestaniæ non procul a Buchara (Lehm.), in salinis Uli Persiæ bor. (Seidl!), in sabulosis Persiæ austro-orientalis inter Kerman et Jezd et ad Bachtan (Bge!).

Planta characteribus *S. Kali* affinis sed habitu, foliis floralibus breviter cuspidatis non pungentibus *S. Sodæ* accedens. Semipedalis, folia inferiora sesquipollicaria, alæ majores latissimæ tres lineas longæ 5-6 latæ, alæ minores lineæ $\frac{3}{4}$ - $\frac{1}{4}$ longæ.

5. S. Kali (L. Sp. 322) annua hirta vel glabrescens, caule a basi divaricatum ramoso ramisque crassis hisce striatis præter centrale erectum procumbentibus vel ascendentibus, foliis alternis crassis a basi semiamplexicauli lanceolatis apice subulatis spinosis semiteretibus, floribus solitariis vel 2-3^o glomeratis axillaribus plus minus dense spicatis, bracteis a basi ovatâ pungentibus perigonio sublongioribus spinescentibus, perigonio fructifero cartilagineo basi rotundato, phyllis medio dorso alis obovatis vel reniformibus varie explicatis instrutis ♂. Ic Fl. Dan. 818. — Nees Gen. Germ. Ic. — *S. rosacea* Smith Prodr. non L.

Hab. in salsis maritimis et interioribus ditionis a Græciâ! et Macedoniâ! ad Egyptum! prov. Caucasicâ! Persiam! Turkestaniâ! Affghaniam!, in salsis interioribus Anatoliæ bor. (Wied!), Cappadociae (Bal!).

β. Tragus. — Sæpe calvescens, perigonii fructiferi alæ rudimentariae sæpe ad carinam transversam reductæ. — *S. Tragus* M.B. Casp. p. 142.

Hab. cum typo, Zacynthus (Orph!), Archipelagus (Sibth.), Pontus Lazicus (Bal!) et alibi.

γ. tenuifolia Moq. — Folia tenuiora filiformia elongata, perigonii alæ breves. — *S. Tragus* L. Sp. 322.

Hab. in Georgiâ Caucasicâ (Wilhm.), Turkestaniâ (Lehm!) et alibi.

Ar. Geogr. Europa omnis, Sibiria, Africa borealis et quoque Africa Asia tropicæ, America.

6. S. spissa (M.B. Mem. Mosc. I, p. 140) annua fructifera interdum basi indurata glanca villoso-pubescent tandem glabrescens, caule erecto tenui fere a basi alternatimi ramoso ramisque firmis, foliis alternis carnosis filiformibus obtusis muticis mox deciduis superioribus abbreviatis, floralibus late ovatis acutiusculis bracteolas subæquantiibus, floribus solitariis tenuiter dense vel laxiuscule spicatis, bracteolis dorso late carnosis orbiculatis subconcavis membranaceo-marginatis, perigonii glabri fructiferi laciniis ovato-lanceolatis acutiusculis denum discoideo-conniventibus, alis ex medio dorso ortis semiorbiculatis vel transverse latioribus rufescens ♂. *S. nitraria* Pall. Ill. tab. 15. — *S. incanescens* C A. Mey. in Eichw. Casp. p. 35 tab. 26 (forma partibus floralibus villosior incana). — *S. ericoides* var. *leucoxylon* Moq. in D.C. Prodr. p. 183 ex cl. Bge. — *S. suædoïdes* Moq. in D.C. Prodr. p. 162 ex desc? — *Halogeton Jaubertianus* Bal. pl. exs. — *Halogeton Olivieri* Moq. Chen. p. 159. — Jaub. et Sp. Ill. Or. tab. 134.

Hab. in salsis, Cappadocia ad occasum Cæsareæ (Balt), Armenia Rossica (Abich), Persia borealis ad Urmiah (Szovt!), in deserto inter Koum et Schurab (Haussk!), Persia australi ad Seytun et Bender Dilem (Haussk!), Afghanistan (Griff!).

Semipedalis vel paulo elatior ramosissima dumulosa, facies *S. scleranthæ* sed folia crassiora 3-4 lineas longa, bracteolæ orbiculatae ut et folia floralia membranaceo-marginatae, perigonium fructiferum cum alis tres lineas diametro latum.

Ar. Geogr. Rossia austro-orientalis, Sibiria Uralensis, Songaria.

7. *S. tamariscina* (Pall. It. III, p. 604) annua scabriuscula, caule erecto a basi vel paulo supra basin ramoso ramis virgatis alternis strictis, foliis alternis rigidis a basi semiamplexicauli subulatis spinulosis brevibus rectis vel recurvo-patulis, floralibus lanceolatis flores superantibus, floribus 1-3-globeratis secus ramos in spicas tenues longas plus minus densas dispositis, bracteolis triangulari-ovatis mucronatis perigonio subæquilongis, perigonii fructiferi glabri tubo breviter obconico alis erectiusculis brevibus seniorbiculatis flabellatim fuscescenti-nervosis ⊖. Ic. Pall. Ill. tab. 25 exel. fig. 2 (foliis nimis crassis imperfecta).

Hab. in salsis Tauriæ (ex Pallas), Transcaucasiæ ad Sallian (G.A. Mey.).

Pedalis, folia tenuissima 3-6 lineas longa, fructus alæ parvæ lineam circ. longæ.

Ar. Geogr. Rossia australis, Songaria, Sibiria Altaica.

8. *S. inermis* (Forsk. Fl. Eg. Arab. p. 57) annua demum induata valde pappuloso-farinosa, caulibus prostratis et ascendentibus alternatim ramosissimis ramis ramulisque tenuibus divergentibus, foliis minimis inferioribus cito deciduis semiamplexicaulibus breviter linearibus acutiusculis, superioribus et floralibus ovato-triquetris hisce imbricatis, floribus parvis secus ramulos solitariis laxe et tenuiter spicalis, bracteolis orbicularibus mucronulatis folio florali subæquilongis, perigonii fructiferi phyllis farinosis triangularibus in discum conniventibus, alis versus medium perigonium insertis explanatis obovatis margine sinuatis extus usque ad perigonii basin in nervum elevatum productis, antheræ appendice minutâ ovatâ obtusâ 5. Ic. Forsk. Icon. tab. 8, fig. B. — *S. villosa* Sieb. exs. non Del.

Hab. in collibus arenosis ad Alexandriam (Forsk., Sieb., Samar! Duparq!), in Syria (Oliv. ex Moq.). Arabice *Gummeli* vel *Nadcuva*.

Dumi 1-1 $\frac{1}{2}$ -pedales, folia infima pauca 2 lin. longa, floralia linea breviora, fructus cum alis diam. 2-2 $\frac{1}{2}$ lineas tantum latus.

9. *S. carinata* (G.A. Mey. in Eichw. Casp. p. 36, tab. 28) annua parte inferiori lanuginosa demum glabrata farinosa, caule humili tenui a basi ramosissimo demum cum ramis filiformibus tenuibus divaricatis intricatis durescente, foliis alternis et fasciculatis linear-setaceis obtusiusculis, caulinis abbreviatis, floralibus triangulari-ovatis bracteolis duplo brevioribus, floribus parvis solitariis lepidotis tenuiter et

remotiuscule spicatis, bracteolis late ovato-rhombeis acutis vel mucronato-acuminatis, perigonii lepidoti fructiferi laciniis lanceolatis acuminatis demum induratis in conum conniventibus, alis roseis subbasilaribus horizontalibus obovato-ellipticis acutiusculis basi unguiculato-angustatis totâ longitudine plus minus distincte carinalis inter se discretis, antheræ appendice ovato-oblongâ obtusâ loculis angustiore et breviore ⊖. *S. sclerantha* C.A. Mey. Casp. p. 35, tab. 27.

Hab. in littoribus Caspicis Turcomaniæ ad Krasnowodsk (Karel!), in deserto Kisylkum Turkestaniæ (Lehm!), in regno Cabulico ad Byani prope Turkistan (Griff. 1805!).

Dumulus intricatissimus $\frac{1}{2}$ -4-pedalis, folia tenuissima 3-4 lineas longa fructus cum alis $2\frac{1}{2}$ lineas diam. latus. Ex cl. a Bunge auctoritate species binas Meyerianas conjunxi.

Ar. Geogr. Songaria.

10. **S. gossypina** (Bge in litt.) annua undique gossypino-lanala divaricatum ramosa, foliis anguste linearibus acutis, florali bracteas ovatas obtusas subæquante, paniculâ corymbose ramosissimâ gracili flexuosa floribus solitariis spicatis, antheræ appendice ovato-lanceolatâ acutissimâ antheram ipsam subæquante, perigonio fructifero.... ⊖.

Hab. in montosis prope Achmetabad ad orientem urbis Mesched prov. Khorassan (Bge).

Junior tantum lecta, pilis bombycinis totam plantam præter caulem et ramos plus minus denudatos vestientibus insignis, folia in specimine jam fere destructa 5-6 lineas longa linea $\frac{2}{3}$ lata. Semipedolis, flores secus spicas subdissiti.

11. **S. crassa** (M.B. Mem. Soc. Nat. Mosc. I, p. 100) annua adpresso lanato-tomentosa vel cinereo-pubescentia, caule erecto a basi ascendentem vel diffuse ramosissimo ramis crassiusculis præter infimos alternis, foliis alternis a basi semiamplexicauli decurrente semiteretibus crasse filiformibus carnosis obtusissimis muticis patulis, floralibus abbreviatis, floribus solitariis densius vel laxius spicatis, bracteolis folio florali brevioribus late ovatis herbaceo-cuspidatis, antherarum appendice oblongâ loculis duplo breviore, perigonii fructiferi laciniis late triangularibus conniventibus alis latissime semiorbicularibus imbricatis purpureo-fuscis patentissimis perigonio ipso duplo longioribus, stigmatibus longe subulato-setaceis ⊖. *S. rosacea* M.B. Casp. App. p. 143 et Pall. Ill. tab. 18 (forma calvescens superne glabra) non Linn. — *S. cana* C. Koch Linn. XXII, p. 190.

Hab. in salsis maritimis et interioribus, Georgia Caucasia (C.A. M! Hoh!), prov. Karabagh (Hoh.), Armenia Rossica (C. Koch), Persiae borealis deserta salsa (Auch. 2744!) et ad lacum Ourmiah (Seidl!).

Pedalis et sesquipedalis, folia inferiora interdum 8-12 lin. longa, alæ 3 lin. longæ 4-5 latae.

Ar. Geogr. Rossia australis, Turcomania, Songaria, Sibiria Uralensis.

12. **S. lanata** (Pall. Voy. App. p. 282, tab. 51, fig. 2) annua albo-lanuginosa, caule erecto a basi patentim et alternatim ramoso rainis

inferioribus elongatis, foliis alternis semiamplexicaulibus semitereti-filiformibus carnosis obtusis, floralibus ovato-lanceolatis, floribus 1-3 glomeratis laxiuscule spicatis, bracteolis lanceolatis folio florali sub-brevioribus, perigonii fructiferi tubo hemispherico laciniis anguste linearie-lanceolatis, alis late reniformibus imbricatis laciniis sublongioribus fuscis horizontalibus, antherarum appendice loculis æquilongâ et latiore petaloideâ cochleariformi obovatâ, stigmatibus apice truncatis invicem applicatis ♀. Ic. Pall. Ill. tab. 21. — *S. laniflora* L. fil. Suppl. p. 172.

Hab. in salsis Persiae borealis ad lacum Urmiah (Seidli).

Planta $\frac{1}{2}$ -1-pedalis, folia inferiora 6-12 lineas longa, alæ eis *S. crassæ* cui affinis minores.

Ar. Geogr. Rossia australis, Songaria, Sibiria Altaica.

SERIES II. — Perennes.

* Oppositifoliae.

13. *S. tetragona* (Del. Fl. Eg. p. 228, tab. 21, fig. 4) fruticosa plus minus papilloso-pubescent et cana, ramis oppositis tortuosis inferioribus diffusis, foliis minimis tomentosis squamæformibus carnosis breviter triangulari-ovatis obtusis supra concaviusculis oppositis secus inflorescentiæ ramulos breves amentiformes tetragonos dense imbricatis, ramulis oppositis vel sparsis breviter cylindricis oppositis vel alternis longe spicatis, bracteolis ovatis obtusis concavis foliis similibus, floribus axillaribus solitariis, filamentis post anthesin persistentibus, antheris appendice punctiformi superatis, perigonii fructiferi phyllis parte liberâ lanceolatis elongatis in conum conniventibus extus sericeis, alijs infra medium insertis mediocribus patentibus margine subimbricatis ♂. *Caroxylon tetragonum* Moq. in D.C. Prodr. l. cit. p. 176. — *Halogenon tetragonum* Moq. Enum. p. 159.

β. tetrandra. — Sterilis, flores tetrameri staminibus quatuor. — *S. tetrandra* Forsk. Fl. Eg. Arab. p. 78. — *Halogenon tetrandrus* Moq. Enum. p. 160.

Hab. in Egypto ad Alexandriam (Forsk., Del., Ky. 5^o2!), ad Cahiram (Sieb sub *S. baccatâ*!), deserto Tih Arabiae Petreæ (Boiss!), Palestina ad mare Mortuum (Boiss!).

Species foliis lineam longis dense imbricatis insignis, spiculæ amentiformes 6-12 lineas longæ ut et flores vel oppositæ vel alternae. *S. tetrandra* Forsk. in quâ fructus nunquam observati est sexus masculus et sterilis speciei polygamæ

Ar. Geogr. Africa borealis.

14. *S. longifolia* (Forsk. Fl. Eg. Arab. p. 55) glabra parce farinosa fruticulosa, ramis ascendentibus vel erectis brevibus rigidis opposite et breviter ramulosis, foliis oppositis carnosis breviter linearibus tereli-trigonis obtusis, floralibus oblongis cucullatis obtusis

bracteolas orbiculatas concavas superantibus, floribus oppositis secus ramulos spicas longiusculas laxas vel breves confertas formantibus, perigonii fructiferi tubo subgloboso longitudinaliter et elevatim nervoso arolâ basilari subpentagonâ laciniis membranaceis brevissime et obtuse ovatis, aliis expansis basi attenuatis orbiculatis tubo vix duplo longioribus 5.

Hab. in collibus saxosis Egypti inferioris ad Alexandriam, Aboukir, Ram-leh (Forsk., Del., Auch. 2724! Ehr! Samar! etc.).

Suffrutex semipedalis, folia 4-5 lineas longa, fructus alatus diam. 5-6 lineas latus. Ab hac specificie diversissimam autumat am. Bunge *Salsola oppositifoliam* Desf. = *S. fruticosam* Cav. Ic. tab. 285, 3-4-pedalem, foliis tenuioribus sepe pollicaribus, ramis et spicis elongatis, fructu diam. cum alis amplissimis 9-11 lin. lato sed ejus praeterea characteres iidem. Si tantum varietas, areæ geographicæ *S. longifolia* Africa borealis, insule Canarienses, Hispania australis et Sicilia addenda sunt.

• • Sparsifoliae.

15. **S. Auricula** (Moq. Chen. Enum. p. 135) basi suffrutescens glabra glauca, caulis erectis elongatis virgatis foliosis superne paniculato-ramulosis, foliis alternis coriaceis semiamplexicaulinibus ovato vel cordato reniformibus interdum subrotundis obtusissimis vel breviter acutis, superioribus sensim diminutis interdum decurrentibus, floralibus bracteolisque concavis ovato-rotundis, floribus longe et laxiusculle spicatis, perigonii fructiferi magni laciniis membranaceis triangularibus acutis in conum conniventibus, alis supra medium insertis suborbiculatis integris amplis, antheris muticis 3. *Caroxylon Auricula* Moq. in D.C. Prodr. p. 172. — *Salsola Moquiniana* J. et Sp. III. tab. 137.

Hab. in Persiâ inter Ispahan et Teheran (Oliv.).

Æ. acutifolia Moq. — Folia ovato vel oblongo-lanceolata acuta fere omnia decurrentia, floralia bracteolæque acuta. — *Caroxylon acutifolium* Moq. in D.C. Prodr. — *Sals. decurrentis* J. et Sp. III. tab. 138.

Hab. in Persiâ boreali ad salinas Mamam (Seidl!), australi (Auch. 2956!), ad lacum Nemekejra prope Schiraz (Ky. 460), regno Cabulico (Griff!), Belutschijâ (Stocks!).

Caulis pedales et ses pripedales, folia in typo 5-9 lineas longa et lata, in varietate pollicem et amplius longa, fructus fere magnitudine Pisi. Varietas multo frequentior videtur et ad typum transire videtur.

16. **S. lancifolia** (Boiss. Diagn. Ser. I. 12, p. 98 sub *Caroxylo*) glabra glauca basi suffruticosa, caulis virgatis angulatis superne paniculatim ramulosis, foliis plus minus longe decurrentibus planis longe et anguste lanceolatis subulato-attenuatis, floralibus linearibus bracteolas triangulares acutas superantibus, floribus laxiusculle spicatis, perigonii fructiferi laciniis membranaceis triangularibus acutis in tubum conniventibus, alis medio dorso insertis orbiculatis crenulatis, antheris muticis 3.

Hab. ad littus septentrionale maris Mortui Palestinæ (Boiss!).

Pedalis et sesquipedalis, folia 1-1½-pollicaria basi 1-2 lineas lata. Aflinis *S. Auriculae acutifoliae* a quâ specifice differre videtur foliis angustissimis, floralibus anguste linearibus. Fructus subminor alis angustioribus.

17. *S. glauca* (M.B. Taur. Cauc. I, p. 185) glauca glabra suffruticosa, trunci ramis ascendentibus vel procumbentibus, ramis floriferis erectis elongatis teretibus superne paniculatis, foliis lineariteretiusculis basi non dilatatis obtusiusculis, floralibus linearibus brevibus bracteolas ovatas breviter caudatas carinatas superantibus, floribus solitariis longe et saepius laxè spicatis, perigonii fructiferi laciniis conniventibus triangulari-lanceolatis obtusiusculis, alis infra medium ortis stramineis amplis orbiculari-ovovatis imbricatis erosocrenulatis, antheris non appendiculatis 5. *S. spicata* Pall. Ill. p. 27, tab. 19 non Willd.

Hab. in salsis Armeniae Rossiae ad Kulp (Seidl!), Georgiae Caucasicæ in aridis collum Elisabethpol (Hohen!), Persiae bor. in prov. Aderbidjan (Szov!), ad salinas Uli et lacum (Seidl!), inter Ispahan et Teheran (Haussk!).

Truncus saepè crassitie digitii, rami floriferi pedales, folia 1-1½-pollicaria linearis vel sesquilineam crassa, perigonium fructiferum cum alis diam. 6 lineas latum.

18. *S. brachiphylla* (Boiss. et Haussk. non Spr.) glabra glauca basi suffruticosa, ramis floriferis erectis vel ascendentibus crebre foliatis in spicam simplicem vel ramulosam abeuntibus, foliis carnosis brevibus oblongo-linearibus inferne attenuatis obtusis mucronatis plerumque recurvis, floralibus bracteolas triangulares acutas superantibus, floribus laxiuscule spicatis, perigonii fructiferi laciniis membranaceis in conum conniventibus longe lanceolatis obtusis, alis stramineis infra medium ortis orbiculatis crenulatis, antheris muticis 5.

Hab. in Persiae deserto Hauz i Sultan inter Teheran et Ispahan (Haussk!), ad Hassara inter Kasbin et Teheran (Buhsel).

Aflinis *S. glaucae* specifice differre videtur foliis crassioribus recurvis parte inferiori attenuatis apice mucronatis 3-4 lineas tantum longis saepius recurvis, ramis floriferis semipedalibus, perigonii phyllis longe lanceolatis obtusis.

19. *S. subaphylla* (C.A. Mey. in Eichw. Casp., p. 34, tab. 24) fruticosa glabra a basi divaricatim ramosissima ramis patentim ramulosis paniculatis, foliis paucis remotis fragilibus cito exsiccatis et deciduis subcylindricis crassiusculis acutis mucronulatis, floralibus ac bracteolis æquilongis squamaeformibus late ovatis margine late scariosis obtusis, floribus longe et laxè spicatis, perigonii fructiferi laciniis triangularibus conniventibus, alis a medio dorso ortis stramineis semiorbiculari-ovovatis imbricatis margine erosulis, antheris muticis 5. *Caroxylon subaphyllum* Moq. I. cit. p. 173.

Hab. ad littus maris Caspii in Turcomaniâ meridionali ad Krasnowodsk (Kar!), in Turkestaniâ inter Buchara et Samarcand (Lehm!), Affghania (Griff!).

Frutex 1-1½-pedalis, folia rara cito evanida pollicem dimidium aut amplius

longa, floralia non ut in *S. glauca* bracteolas et flores excedentia. Species insuper ramis et ramulis divaricatis elongatis insignis.

20. **S. hispidula** (Bge Pl. Lehm. p. 469 sub *Caroxylo*) fruticosa glauca patule hispidula, ramis floriferis a trunko lignoso erectis elongatis stricte paniculatis, foliis alternis sessilibus semitereti-linearibus elongatis acutis, floralibus brevibus bracteolas ovato-triangulares acutiusculas superantibus, floribus solitariis longe et laxiuscule spicatis, perigonii fructiferi laciniis triangulari-lanceolatis obtusis in conum conniventibus, alis infra medium dorsum ortis late obovato-reniformibus stramineis ♂.

Hab. in deserto meridionali Kisilkum Turkestaniae (Lehm!), in regno Cabulico (Griff. 1809 ex Bge).

Rami 1-4½-pedales, folia 1-4½-pollicem longa 1½ lineas lata, perigonium fructiferum 5 lineas diametro latum. A *S. glauca* et *S. subaphylla* indumento distincta.

21. **S. arbuscula** (Pall. It. p. 412, tab. G, fig. 1) glabra, fruticosa trunko squarrose ramosissimo, ramis floriferis elongatis alternatim ramulosis, foliis alternis et fasciculatis linearibus obsolete trigonis obtusiusculis latitudine et longitudine variis, floralibus abbreviatis bracteolas orbiculari-ovatas margine membranaceas interdum breviter caudatas superantibus, floribus solitariis glabris laxe vel longiuscule spicatis, perigonii laciniis parte liberâ ovato-lanceolatis obtusis fructiferis depresso conniventibus, alis magnis semiobtusiusculatis infra medium ortis, antheris non appendiculatis ♂. Pall. Ill. tab. 27. — Eichw. t. 29 et 30. — *S. arborescens* L. fil. Suppl. p. 473.

Hab. in salsis, Persia bor. ad Tchechme Ali jugi Elbrus (Buhse!), in Turcomaniâ australi (Kareli).

β. angustifolia. — Folia angusta saepe longissima. — Eichw. tab. 31.

Hab. in Transcaucasiâ (Radde!), Turcomaniâ (herb. Petri!).

Folia variant 4 lineas-2½-pollices longa ½-2½ lin. lata.

Ar. Geogr. Rossia austro-orientalis, Songaria, Sibiria Altaica.

22. **S. gemmascens** (Pall. Ill. p. 24, tab. 16) fruticulosa pulvрrulento-cana, trunko humili distorto dumuloso, ramis erectis brevibus undique gemmis axillaribus globosis canis obsitis, foliis alternis cito deciduis a basi semiamplexicauli brevissime subulato-triquetris, floralibus squamæformibus ovatis concavis adpressis, floribus solitariis in spicas terminales rectas densiusculas dispositis, bracteolis concavis folia floralia duplo superantibus, perigonii fructiferi phyllis parte liberâ triangularibus acutis conniventibus, alis a basi ortis roseis late obovatis flabellato-nervosis imbricatis patentibus, antheræ appendice minutâ ♂. *S. verrucosa* C.A. M. Enum. p. 460 non M.B. (ex spec!). — *S. Passerina* Bge olim. — *Caroxylon nodulosum* Moq. d.c. Prodr. p. 177.

Hab. in salsis lutosis Armeniæ Rossicæ ad Erivan (Szov!), Georgiæ Cau-

casicæ ad Baku et Schemacha (C.A. Mey! Haussk!), ad Sallian et in valle fluvii Kur (Hoh. 2911!), Persiae borealis ad salinas Uli (Seidl!), inter Mendjil et Rudbar (Haussk!).

Diffr. a *S. verrucosâ* staturâ humili semipedali, ramis gemmis globosis crebre obsitis. Folia majora vix 2 lineas longa tenuia, perigonum cum alis 3-3 $\frac{1}{2}$, lin. diam. latum.

Ar. Geogr. Sibiria Uralensis, Mongolia Chinensis.

23. *S. fætida* (Del. Fl. Eg. Ill. № 310) fruticosa, furfuraceo-farinosa vel cinerea, ramis pubescentibus vel glabratris erectis elatis alternatim paniculato-ramosissimis erecto-patulis virgatis, foliis alternis minutissimis interdum glabratris orbiculatis carnosus margine membranaceis, floralibus imbricatis spicas brevissimas cylindricas formantibus, floribus solitariis confertis, bracteolis foliis floralibus similibus et subæqualibus, perigonii glabri fructiferi minimi phyllis parte liberâ ovato-triangularibus brevibus conniventibus alis albis versus medium insertis obovatis explanatis, stigmatibus exsertis elongatis divaricato-recurvis, antheræ appendice punctiformi 5. *Chenopodium baryosmon* Rœm. et Sch. Syst. VI, p. 269. — *Caroxylon fætidum* Moq. in d.c. Prodr. p. 178.

Hab. in desertis Egypti mediae et superioris, Ouadi Dachl deserti Arabici et in Oase magnâ (Selw!), ad Theben, Syenem (Del., Ehrenb! Boiss!), Belutschia (Stocks!). Arabice *Mulley* (ex Del.), *Cherit* (ex Schimp.).

Caules 1-2-pedales a basi paniculati spicis innumeris $\frac{1}{2}$ -1-pollicaribus diam. vix sesquilineam latis saepius densis, perigonum cum alis sesquilineam diam. latum.

Ar. Geogr. Nubia, Arabia tropica, India borealis, Senegalia.

24. *S. verrucosa* M.B. Mem. Soc. Nat. Mosc. I, p. 141) fruticosa interdum arborescens, ramis elongatis alternatim paniculato-ramosissimis ramulis tenuiter virgatis erecto-patulis, foliis alternis pubescentibus vel cano-tomentosis breviter semitereti-filiformibus tenuibus obtusis basi dilatata gibbis, floralibus brevissimis carnosus ovato-triquetris gibbis, floribus solitariis in spicas tenues dense congestis, bracteolis folio florali similibus longioribus cum eis albo-verrucosis, perigonii fructiferi phyllis parte liberâ triangulari-ovatis pubescentibus conniventibus, alis a medio dorso ortis semiobtusatis fuscis explanatis, stigmatibus brevibus conniventibus inclusis, antheræ appendice punctiformi 5. *S. dendroides* Pall. Ill. tab. 14 (folia nimis crassa). — *S. Georgica* Bunge in Ledeb. Fl. Ross. p. 814).

Hab. in salsis, Iberia et Georgia Caucasicæ (M.B., C.A. Mey! Hoh!), Armenia Rossica (Seidl!), Persia borealis (Seidl! Buhse! Haussk!).

Affinis *S. fætidae* a quâ præter alias notas foliis inferioribus breviter filiformibus ($\frac{1}{2}$ -2 lineas longis nec ovato-orbiculatis differt.

Ar. Geogr. Rossia austro-orientalis, Sibiria Uralensis.

25. *S. ericoides* (M.B. Mem. Soc. Natur. Moscou I, p. 141) suffruticosa glabriuscula cæsio-glauea, ramis annotinis brevibus, foliis

alternis carnosulis cylindraceis brevibus obtusis, floralibus abbreviatis bracteolas orbiculatas superantibus, floribus axillaribus solitariis remotiusculis breviter spicatis, perigonii fructiferi phyllis parte liberâ ovatis obtusis, alis versus medium ortis orbiculatis vel obovatis explanatis, antherarum appendice obsoletâ ȝ. *S. cæsia* F. et M. in Hohen. Talysh Enum. p. 121. — *S. glanca* C.A. Mey. Enum. p. 160 non M.B. — *S. ericoides* var. *foliolosa* Fenzl. in Led. III, p. 813.

Hab. in argillo-salsis Transcaucasiae ad Sallian et in deserto ad fluvium Kur (Hoh. exs. 2918).

Affinis *S. verrucosæ* differre videtur glabritie, staturâ humili et in primis foliis manifeste crassioribus (3 lin. longis 1/2 lin. crassis), spice pauciflorâ laxiores, flores majores, folia floralia bracteolas superantia.

26. *S. vermiculata* (L. Sp. p. 323) fruticosa pube lævi brevi pubescens vel canescens, ramis ascendentibus vel erectis paniculatum et crebre ramulosis, foliis a basi dilatatâ semiamplexicauli tereti-filiformibus brevibus, floralibus ovatis interdum breviter cuspidatis bracteolisque ovatis membranaceo-marginatis, floribus solitariis densius vel laxius spicatis, perigonii phyllis parte liberâ ovato-triangularibus obtusis hirtis conniventibus, alis infra medium insertis obovatis imbricatis patulis, antheræ appendice punctiformi ȝ. *S. microphylla* et *S. flarescens* Cav. Ic. tab. 287 et 288. — *S. brevifolia* Desf. Atl. I, p. 218. — *S. Agrigentina* Guss.

Hab. in rupibus maritimis insulæ Melos Archipelagi (Urv! Orph!).

Haec species ab auctoribus in Rossia australi, regione Caucasicâ, Sibirâ, Syria, Persia indicatur, sed ex commutatione cum *S. laricina* vel *S. rigidâ*.

Ar. Geogr. Lusitania, Hispania, Sardinia, Sicilia, Africa borealis.

27. *S. laricina* (Pall. Ill. p. 21, tab. 13) suffruticosa puberula glabrescens vel villosula superne glabra, caulinis e trunko indurato erectis vel ascendentibus virgatis rigidis longe paniculato-ramulosis, foliis persistentibus elongatis filiformi-setaceis muticis sæpe fasciculatis patulis, floralibus bracteolas æquantibus vel vix superantibus ovato-triquetris, floribus solitariis remotiusculis longe et tenuiter spicatis, bracteolis carnosis ovato-globosis dorso argute carinalis, perigonii fructiferi phyllis parte liberâ ovato-oblongis apice obtusis erosis in conum conniventibus, alis infra medium ortis reniformibus incurvo-patentibus fuscis, antheræ appendice punctiformi ȝ. *S. vermiculata* var. *luricina* Moq. in p.c. Prodr. p. 181.

Hab. in salsis Taurie prope Sudak (Stev.), prov. Caucasicarum ad Kislar (C.A. Mey.), Turcomaniæ ad Caspium (Karel.).

Pedalis et sesquipedalis, a *S. verrucosâ* et *gemmascente* foliis persistentibus 3-6 lineas longis, plantæ partis superioris glabritie, a *S. vermiculatâ* foliis tenuibus statim distinguenda.

Ar. Geogr. Rossia austro-orientalis, Sibiria Uralensis et Altaica.

28. *S. rigida* (Pall. Ill. p. 20, tab. 12) fruticulosa pilis flavescensibus sub lente horizontaliter denticulato-ramulosis hirsuta aut villosa

vel parte inferiori glabrescens, ramis vetustis prostratis vel ascenditibus, hornotinis virgatis rigidis anguste paniculato-ramulosis, foliis alternis filiformibus obtusis patulis vel recurvis, axillis fasciuliferis, floralibus abbreviatis flores subsuperantibus et aequalibus, floribus solitariis laxius vel densius anguste spicatis, bracteolis ovatis carnosò-carinatis sœpe herbaceo-cuspidatis, perigonii phyllis parte liberâ villosis triangulare-ovatis erectis, alis versus medium ortis late reniformibus imbricatis fuscis, antheræ appendice punctiformi ♂.

Hab. in salsis, Transcaucasia prope Tarki (Eichw.), Persia borealis (Buhse!), ad salinas Uli et Maman (Seidl!), Persia occid. ad Fîrrînd (Noë 1'33!), Turkestania in montosis Karatau prope Sainareand (Lehm!), Turcomania australis ad Caspium (Kar!), regnum Cabulicum (Griff.).

β. tenuifolia. — Folia tenuiora filiformi-setacea. — *S. vermiculata*
β. villosa Moq. in p.c. Prodr. p. 181. — *S. villosa* Del. Ill. N° 309 ?

Hab. in Syriâ (Auch.), circa Damascum (Gaill!) et Balbeck (Bl!), in Pales-tinâ (Boiss!), Egypto (Del. ex Moq.), Babylonia ad Bagdad (Noë!).

Species foliis crassiusculis vel tenuioribus valde variabilis, *S. vermiculata* valde affinis et ab eâ foliis longioribus, indumento pilis longioribus denticulatis constante an satis distincta.

Ar. Geogr. Sibiria Uralensis, Songaria.

29. *S. canescens* (Moq. in p.c. Prodr. p. 203 sub *Noëi*) pilis adpressiusculis canescens interdum glabrescens, suffruticosa trimeo lignoso tortuoso ramoso, ramis novellis aliis brevibus sterilibus crenatim foliosis aliis plus minus elongatis virgatis simplicibus in spicas longas laxas abeuntibus, foliis linearibus planis vel obsolete triquetris acutiusculis vel obtusis erectis, floralibus abbreviatis bracteolas subcoriaceas a basi ovatâ abrupte et breviter caudatas superantibus, floribus inajusculis dissitis, perigonii phyllis parte liberâ hirtis membranaceis rigidis oblongo-lanceolatis obtusis rectis bracteolas subequantibus, alis versus medium perigonium ortis transverse oblongis fuscis medio-eribus, antheræ appendice lanceolatâ acutissimâ dimidiâ antherâ longiore ♂. *Salsola tomentosa* Spach. in Ky. exs. 1843. — *Noëa tomen-tosa* ex parte et *N. Aucheri* Moq. in p.c. Prodr. 207.

Hab. in siccis regionis subalpinæ et alpinæ, Anatolia (Wiedt), Cataonia montes (Tchich! Haussk!), mons Maknel Libani 6500' (Ky!), montes Schahu, Pir Omar Gudrun, Dalehani, Kellal etc. Kurdistaniae et Persie orientalis (Haussk!), mons Gara Kurdistaniae (Ky. 346!), mons Kuli Daïna Persie australis (Ky. 646!), et mons Elbrus Persie hor. prope Passgala (Ky. 504!).

Rami floriferi 1/2-4-pedales et etiam longiores, folia 5-9 lineas longa lineam dimidiâ lata, spicæ terminales semipedales et longiores.

30. *S. aurantiaca* (Bunge in litt.) adpresso canescens vel tomentosa fruticulosa, ramis a collo pluribus erectis superne paniculatum et anguste ramulosissimis, foliis linearis-subtrigonis acutiusculis rectis vel incurvis, floralibus abbreviatis bracteolas subsuperantibus vel aequilongis, bracteolis a basi ovatâ breviter et obtuse caudatis, floribus secus paniculæ ramos breviter et dense spicatis, perigonii

phyllis parte liberâ membranaceis erectis oblongo-lanceolatis obtusis, alis latis aurantiacis transverse sublatiорibus, appendice triangulari-lanceolatâ acutâ antheram subæquante 5. *Halimocnemis tomentosa* et *Noëa tomentosa* Moq. ex parte.

Hab. in salsis Persiae, ad Teheran (Bél! Auch. 1205!), inter Teheran et Kasbin et ab Gendab jugi Elbrus (Buhse!), inter Kaschan et Kum atque prope Ispahan (Faussk!), ad salinas Uli (Seidl!), in regno Cabulico (Griff. 101! et 102!).

Diffr. a *S. canescens* ramis tenuioribus non simpliciter spicatis sed in paniculam multifloram abeuntibus, folia breviora sunt, flores minores conferti, alae multo latiores. Planta semipedalis vel paulo elatior.

31. S. Persica (Bge pl. exs.), fruticulosa dense et adpresso albotomentosa ramosissima ramis vetustis prostratis vel ascendentibus, annotinis erectis gracilibus brevibus simplicibus flexuosis, foliis carnis brevibus crassiusculis triquetris acutiusculis, axillis fasciculiferis, foliis floralibus bracteolisque breviter triangulari-lanceolatis crassiusculis hisce subbrevioribus, floribus solitariis laxa et breviter spicatis, perigonii fructiferi phyllis parte liberâ lanceolatis obtusis erectis rigidulis versus medium alis fuscouscentibus reniformibus obsitis, antheræ appendice obtusâ dimidiâ antherâ breviore 5.

Hab. ad salinas Persiae borealis (Seidl!).

Ex specimine humilis semipedalis, folia 3-4 lineas longe fere lineam lata, spica flexuosa laxa 3-4-pollicaris, flores majusculi. A *S. canescens* et *S. aurantiaca* foliis brevissimis crassis, antheræ appendice distincta.

Obs. *Salsola rosacea* L. et *S. collina* Pall. infra fines ditionis non crescere videntur. *Salsola Bottae* = *Halothamnus Bottae* Jaub. et Sp. ill. Or. tab. 136 in Arabia petreâ erroneâ indicata ex sched. Herb. Paris est species Arabiae tropicæ intra fines nostras non obvia.

SUBTR. II. ANABASEÆ Moq. Bunge Anabas. Revis. — Semen verticale.

Divis. I. Anabasideæ Bge. — Sepala exteriora tria, unum posticum, duo antica, duo lateralia interiora. Semen (in nostris) a dorso, nempe axi et bracteæ parallele compressum (In genere *Ofaisto* a ditione exsule semen axi et bracteæ contrarie compressum est).

NOËA (Moq. in D.C. Prodr. loc. cit. p. 207).

Flores hermaphroditæ solitarii bibracteati. Perigonii phylla quinque basi breviter connata, fructifera omnia alata. Staminodia nulla. Stamina quinque disco carnoso lobato inserta. Antheræ appendicula lanceolata. Stylus elongatus stigmatibus lanceolatis. Utriculus tenuiter membranaceus. Radiculae apex superus. — Suffrutices vel herbæ annuae, caule non articulato, foliis alternis. Genus ex am. Bge proximum *Salsola* et

ab eâ charactere nimis artificiali seminis verticalis nec horizontalis tantum distinctum.

* Perennes fruticulosæ.

1. N. spinosissima (Linn. fil. Suppl. p. 173) fruticosa glabra a basi squarrose ramosissima ramis erectis rigidis saepe flexuosis ramulis patentibus vel horizontalibus a basi ad medium florigeris apice spinosis, foliis alternis semiteretibus filiformibus mucronulatis, superioribus abbreviatis et tandem squamæformibus, floralibus ut et bracteolæ a basi ovalâ triquetro-subulatis, floribus in axillis solitariis ad ramuli basin saepe approximatis, perigonio glabro, antheræ appendiculâ ejus quartam partem æquante acutâ, perigoni fructiferi alis obovatis vel oblongis perigonio ipso sublongioribus 5. *N. spinosissima* Moq. — *Halogetum spinosissimum* C.A. Mey. En p. 159. — *Salsola Echinus* Labill. Syr. Dec. 2, p. 10, tab. 5. — Del. Eg. tab. 21. — *Salsola spinifex* Pall. Ill. tab. 24. — *Salsola mucronata* Forsk. Eg. Arab. p. 56.

Hab. in rupestribus siccis regionis inferioris montanae et subalpinae, Attica in prom. Sunio et Cyclades (Bory!), Anatolia bor. (Wied.), Lycaonia (Heldr!), Cappadocia (Bal!), Lydia in monte Sypilo (Boiss!), Libanus, Antilibanus et circa Damascum (Labill., Bl!), Palestina (Sieb! Roth!), Egyptus inferior ad Alexandriam (Ehr! Del! Samar! Gaill!), Armenia Turica (Haussk!), Transcaucasia (C.A. Mey! Hoh!), Persia bor. (Szov! Buhse!) et australis in alpe Kuh Daëna (Ky. 672!).

β. humilis. — *Erinacea depressa* 3-5-pollicaris.

Hab. in excelsioribus Libani (Ehr! Bl!), montis Gebel Scheick Antilibani (Bl! Gaill!).

Valde variat staturâ, ramis plus minus crassis rectis vel tortuosis, foliorum longitudine, etc.

Ar. Geogr. Turcomania, Affghania occidentalis, Barbaria.

2. N. Tournefortii (Spach in Ky. exs. Alep. Kurd. sub *Salsold*) suffruticosa seahro-puberula inermis basi ramosissima, ramis erectis elongatis simplicibus et parce ramulosis totâ longitudine florigeris, foliis alternis linearis-filiformibus elongatis mucronulatis, floralibus bracteolas æquantibus vel subsuperantibus ut eæ oblongo-lanceolatis carinatis, floribus solitariis laxe et longe spicatis, antheræ appendiculâ ejus quartam parte subæquante acutâ, perigonii fructiferi alis obovato-oblongis lacinias et bracteolas duplo superantibus 5. *Anabasis Tournefortii* Jaub. et Sp. Ill. tab. 132. — *Noëa Tournefortii* Moq. loc. cit. 208. — *Kali fruticosum Armenum flore magno purpureo*.

Hab. in rupestribus regionis subalpinæ et alpinæ, Armenia (Tourn., Tchih!) et in valle Gotchkar (Ky. 354!), circa Baibut (Bourg!) et Cappadocia in monte Masmeneudagh (Bal!), Syria borealis et Cataonia in montibus Ssossidagh, prope Adiaman et Malatia et in monte Berytdagh (Haussk!), mons Gar! Kurdistaniae (Ky. 345!), montes Persiæ occid. supra Kermanschah 5-8 000' (Haussk!) et prope Kirriad (Noë 1027!).

Rami floriferi graciles pedales et longiores.

3. **N? Daghestanica** (Turcz. in litt. ex Bge Anab. p. 26) fruticulosa tenuissime et patule puberula, ramis elongatis gracilibus ramulosis, foliis linearifiliformibus scabris acutis, floralibus abbreviatis sed bracteolas a basi dilatata naviculari subulatis acutissimas flore duplo longiores superantibus, floribus in ramulorum axillis fere omnibus solitariis laxe spicatis, perigonio glabro, antheræ loculis apice disjunctis connectivo in appendiculam ovato-lanceolatam acutam antheræ æquiloniam abeunti, fructu....§.

Hab. in Caucasi orientalis prov. Daghestan (Herb. Turcz. nunc Univ. Kar-kow). Non vidi.

Fructu ignoto genus cl. Bge subdubium quamvis proxima cum *N. Tournefortii* affinitas hauc *Noea* recte adnumerari suadeat.

4. **N. Griffithii** (Bunge Arab. p. 22) fruticulosa ramosa humilis hispidulo-velutina ramis patulis rigidis inermibus, ramulis floriferis brevibus, foliis carnosis triquetro-subulatis spinuloso-cuspidatis, superioribus abbreviatis, floralibus hamato-recurvis, floribus in axillis superioribus solitariis distantibus, bracteolis a basi concavâ ovata in cuspidem spinosam subito attenuatis folio florali subbrevioribus, perigonii phyllis extus puberulis ciliolatis, antheræ appendiculâ brevi obtusa æque latâ ac longâ antheræ sextam partem æquante, fructu....§.

Hab. in planiciebus arenosis Affghaniæ orientalis prope Dair Haj (Griff. 581). Non vidi.

Facies *Salsola Kali*, fructus ignotus et ideo subdubitanter *Noea* adnumerata.

** Herbaceæ annuae.

5. **N. minuta** (Boiss. et Bal. Diagn. Ser. II, 4, p. 76) annua glaucescens pubescens, caule nano erecto tenui subflexuoso parce et breviter ramuloso, foliis scabridis a basi late membranaceo-marginatâ subulatis albo-mucronatis spicas axillares æquantibus vel superantibus, floralibus abbreviatis ovato-lanceolatis adpressis, bracteolis ovato-lanceolatis naviculari-carinatis florem vix superantibus basi sublantatis, floribus solitariis arcte imbricatis spicas brevissimas densas axillares et terminalem formantibus, antheræ appendiculâ ejus quartam partem vix æquante obtusa, perigonii glabri fructiferi alis infra medium phyllorum ortis cuneatis truncato-retusis ♂.

Hab. in collibus aridis Cappadociæ inter Karahissar et Arapleu Kiöi 5000' (Ball).

Facies *Polychnemi*, bipinnatis, folia tenuissima majora 6-9 lin. longa.

6. **N. major** (Bunge Anab. p. 27) annua glaucescens glabra, caule depresso a basi in ramos flexuosos inferiores caulem æquantes diviso, foliis lanceolato-subulatis arcuato-recurvis subpungentibus superioribus abbreviatis, bracteolis subulatis hamato-recurvis subpungentibus flore longioribus folium florale subæquantibus, floribus solitariis laxe spicatis, antheræ appendiculâ acuminatâ antheram

dimidiati superante, perigonii glabri fructiferi alis oblongis brevibus ⊖.

Hab. in Affghania ad radices colli Kullou (Griff. 1113). Non vidi.

Caules 4-6-pollicares, folia inferiora semipollicaria, fructus a le 2 lineas longitudine non excedentes.

GIRGENSONIA (Bge in Lehm. p. 478).

Flores hermaphroditi solitarii bibracteati. Perigonii phylla quinque imâ basi connata fructifera exteriora 3 vel 2 rarius omnino alata. Staminodia quinque. Antheræ obtuse vel breviter acuminatæ. Ovarium apice glandulosum. Stylus abbreviatus, stigma capitato-bilobum. Utriculus membranaceus. Radiculæ apex superus. — Genus ab *Anabasis* radiculâ superâ tantum distinctum.

1. G. fruticulosa (Bunge Anab. p. 30) perennis glaberrima a caudice brevi lignoso pluricaulis, caulinis erectis a mediâ parte opposite ramosis ramis articulatis patulis interdum ramulosis, foliis oppositis basi connatis linearis-triquetris brevissimis mucronatis interdum curvulis, floralibus ut et bracteolæ oblongo-naviculäribus hisce subbrevioribus florem superantibus, floribus oppositis laxe spicatis, perigonii basi crispe lanati phyllis ovatis obtusis margine membranaceis apice inflexis, antheris exsertis obtusis non appendiculatis, stylo conico glanduloso, stigmate bilamellato, fructu (juniore) 5-alato 5.

Hab. in salsuginosis planitiae Karahissar Cappadociae (Bal!).

Caules 6-9-pollicares, folia caulina 2-2½ lin. longa, spicæ 1-2-pollicares. Fructus matus ignotus.

2. G. oppositiflora (Pall. It. II, App. № 103, tab. O et III, tab. 27 sub *Salsoli*) annua minute seabrido-pubescentia, caule erecto a basi opposite ramosissimo, foliis oppositis a basi connatâ dilatata subulato-triquetris longe mucronatis basi ciliatis, floralibus a basi ovata breviter cuspidatis, bracteolis oblongis navicularibus mucronatis folium florale et florem subsuperantibus, floribus oppositis interrupte spicatis, antheris inclusis abrupte et breviter apiculatis, stigmate capitato-bilobo, fructus phyllis acutis exterioribus tribus alatis alis orbiculatis vel triangulari-inaequilateris integris vel denticulatis ⊖. *G. oppositiflora* Fenzl in Led. Fl. Ross. III, p. 835. — *G. Pallasii*, *G. heteroptera* et *G. gypsophiloides* Bge Lehm. p. 303. — *Anabasis oppositiflora* M.B. — *A. heteroptera* J. et Sp. III, tab. 134. — *Chenopodium oppositiflorum* Willd. Spec. I, p. 1307.

Hab. in desertis Persie mediae (Oliv! Haussk!), orientalis (Bge!), monte Sindjar Mesopotamiae (Haussk!), Turkestaniae (Lehm!), Affghanie (Griffith!).

Valde variat humilis vel pedalis et altior, glauca vel virens, floribus omnibus vel inferioribus tantum remotis.

Ar. Geogr. Regio Aralo-Caspia, Songaria.

3. G. imbricata (Bunge Anab. p. 32) annua tenuiter scabriuscula, caule humili erecto a basi opposite ramosissimo, foliis oppositis a basi connatâ subulatis subrecurvo-patentibus mucronatis margine glaberrimis, floralibus late ovatis acutis, bracteolis folio subbrevioribus oblongo-navicularibus muticis, floribus oppositis secus ramos simplices a basi imbricatim spicatis, antheris inclusis longiuscule apiculatis, stigmate capitato, fructûs phyllis acutis exterioribus tribus alatis alis orbiculatis ♂.

Hab. in aridissimis Persiae mediæ orientalis in prov. Kuhistan inter Afris et Afsel Abad atque prope Birchand (Bge). Non vidi.

Planta cum ramis 2-4-pollicaris; affinis *G. oppositifloræ* sed præter alias notas flores in spicas hexastichas totam rami longitudinem occupantes dense imbricati.

4. G. diptera (Bge Pl. Lehm. p. 479) annua glabra, caule humili a basi stricte ramosissimo, foliis brevibus oppositis a basi latiore subulatis mucronatis, floralibus ovatis cuspidatis, bracteolis carinato-navicularibus folio florali brevioribus et angustioribus, floribus in spicas numerosas elongatas arte congestis imbricatis, antheris inclusis apiculatis, ovario apice dense glanduloso, stigmate capitato subbilobo, perigonii fructiferi phyllis binis anticis alatis cæteris apteris, aliis orbiculari-reniformibus grosse 2-3-dentatis ♂.

Hab. in limoso-salsis collium prope Samarcand Turkestaniæ (Lehm!).

Planta glauca 5-6-pollicaris, folia 2-3 lineas tantum longa. Spicae 1 $\frac{1}{2}$ -2-pollicares.

ANABASIS (L. Gen. 312).

Flores hermaphroditæ bibracteati solitarii rarius glomerati. Perigonii phylla ad basin usque libera. Antheræ exappendiculatæ rarius minute apiculatæ. Stylus ad basin usque in stigmata brevia obtusa divisus. Utriculus membranaceus interdum baccatus. Radiculæ apex inferus. — Fruticuli vel herbæ caulis et ramis articulatis, foliis oppositis stipe ad squamas reductis.

* Perennes caulis annuis herbaceis basi articulatâ deciduis.

1. A. macroptera (Moq. in D.C. Prodr. XIII, p. 213) glauca, radicis perennis crassæ lignosæ collo lato multiplici cicatricoso et lanato, caulis herbaceis cespitosis pumilis simplicissimis rectis vel ad articulos infracto-flexuosis, foliis præter cupulam subbilobam nullis, floribus in omnibus axillis solitariis oppositis distantibus, staminodiis sub anthesi dimidia phylla æquantibus obovatis glanduloso-simbriatis, perigonii basi breviter villosuli fructiferi aliis quinque purpureis magnis semiorbiculatis inæqualibus ♀.

Hab. ad rupes calcareas regni Cabulici prope Bamean (Griff. Journ. 1126). Non vidi.

Caules 3-4-pollicares. Valde officinalis *A. cretacea* Pall. Ill. tab. 9 a qua calyce fructifero pentaptero nec trialato differt.

2. A. brachiata (F. et M. in Karel. Enum. № 749 nomen. — Fenzl in Led. Fl. Ross. III, p. 825) flavicans, radicis perennis crassae collo crasso villoso cicatricoso pluricipiti, caulis herbaceis patule brachiato-ramosis, foliis reductis ad eupulam intus villosissimam bilobam lobis subulato-aristatis, floribus in axillis ramulorum solitariis oppositis distantibus terminalibus congestis, staminodiis suborbiculatis timbriatulis, perigonio basi lanato fructifero pentaptero alis purpureis obovatis integris inæqualibus ♀. *A. cretacea* G.A. Mey. Enum. non Pall.

Hab. in collibus argilloso-salsis, Transcaucasia prope Baku (G.A. Mey.), Persia borealis ad Gendab (Buhsei!), Turcomania ad Caspium (Karel!), Turkestania (Lehm!).

Caules 6-8-pollicares ramis inferioribus sœpe 1-2-pollicaribus superioribus abbreviatis.

Ar. Geogr. Deserta Aralo-Caspica.

•• Fruticulosæ.

3. A. aphylla (Linn. Amoen. Acad. II, p. 347) fruticulosa glabra, caulis a rhizomate crasso pluribus erectis parte inferiori lignosis persistentibus stricte fastigiato-ramosissimis ramis viridibus multiarticulatis demum deciduis, foliis ad eupulam brevem truncatam intus villosulam reductis, floribus solitariis oppositis in spicas terminales basi interruptas apice densiusculas dispositis, staminodiis ciliolatis, perigonio fructifero trialato aliis inæqualibus orbiculari-ovatis conniventibus ♂. *A. Tatarica* Pall. Ill. tab. 8.

Hab. in salsuginosis maritimis et interioribus, Transcaucasia ad flumen Kur (G.A. M! Hoh!), inter Baku et Schemacha (Haussk!), tota Persia media (Bge), Turkestania (Lehm!).

Caules 1-1½-pedales, facies *Ephedræ distachyeæ*.

Ar. Geogr. Rossia austro-orientalis, Songaria, Sibiria Uralensis et Altaica.

4. A. Haussknechti (Bge in litt.) fruticulosa glabra albido-glaucia, caulis hornotinis divaricatim ramosis, foliis junioribus acutiusculis in setam tandem deciduam productis, floribus solitariis oppositis in spicas laxas paniculatas dispositis, staminodiis ciliolatis, perigonio fructifero trialato aliis magnis inæqualibus patulis ♂.

Hab. abunde in collibus calcareis Persiae prope Ispahan (Haussk. exs. sub *A. aphyllæ*).

Suffrutex ramosissimus 6-9-pollicaris facie *A. aphyllam* referens sed præter folia setifera ab ea distinctus caulis non stricte erectis sed ascendentibus divaricatim rambulosis crassioribus albidis nec virentibus, siccis in articulis valde secundentibus, spicæ 6-8-floræ internodis fructu parum longioribus nec fructibus imbricatis, aliis tenuioribus multo majoribus (interdum 4 lineas latis) patulis nec conniventi-erectis.

5. A. phyllophora (Kar. et Kir. Enum. Altaica 1840, N° 735) fruticulosa glabra pallide virens multicaulis caulinis erectis superne pyramidatim ramosis, foliis triangulari-subulatis erectis, floribus in axillis oppositis solitariis dense spicatis, staminodiis ciliolatis, perigonio fructifero trialato alis inaequalibus conniventibus positâ majore 5. *Anabasis intermedia* Moq. in D.C. Prodr. p. 214. — *Anab. subulifolia* Schrenck. — *Brachylepis intermedia* Kar. et Kir. — *Brach. elatior* C.A. Mey.

Hab. in margaceis ad fluvium Kanawat inter El Raboué et Merré prope Damascum Syriæ (Gaill. 2201 specimenia tantum florifera et ideo subdubia).

Specimina florifera quæ ex amic. Bunge ad *A. phyllophoram* sed non absque dubitatione spectare videntur, fructifera investiganda. Planta Syriaca Songaricâ major 1-1 $\frac{1}{2}$ -pedalis. Patriæ diversissimæ.

Ar. Geogr. Songaria:

6. A. articulata (Forsk. Eg. Arab. p. 55, Icon. 8, fig. A sub *Salsola*) fruticosa vel arborescens ramosissima ramis erectis vel tortuosis, foliis ad cupulam brevissime bilobam intus villosam reductis, floribus oppositis solitariis superioribus in ramulorum apice approximatis, staminodiis orbiculari-ovovatis breviter ciliolatis, perigonio fructifero quinquealato alis striatis obovato'-orbiculatis sinuatis 5. *A. articulata* Moq. in D.C. Prodr. XIII, p. 212. — *A. crassa* Moq. Chenop.

Hab. in desertis arenosis, Egyptus (Forsk., Del.), ad Pyramides (Wiest!), ad Alexandriam (Samar!), circa Damiatam (Ehr!), et Suez (Bové!), in deserto Arabico (Husson!), Arabiæ petreæ (Bové!), prope Damascum inter El Raboué et Merré si specim. florifera (Gaill. exs. 2201 huc recte refero).

Variat ramis tortuosis internodiis abbreviatis vel ramis et ramulis elongatis rectiusculis.

Ar. Geogr. Africa borealis, Hispania?

7. A. setifera (Moq. in D.C. Prodr. XIII, 2, p. 214) fruticosa caulinis numerosis erectis vel ascendentibus brachiato-ramosis ramis elongatis, foliis oppositis carnosis crassis cylindricis subclavatis obtusissimis setâ subulatâ cito deciduâ terminatis, floribus in ramulorum axillis superioribus 3-7-nis glomeratis glomerulis inferioribus distantibus superioribus approximatis, staminodiis truncato-subrotundis fimbriatulis, perigonio fructifero pentaplero alis orbiculatis subintegratis subinæqualibus 5. *K. lutea* Moq. loc. cit. p. 215.

Hab. in desertis Egypti inter Kahiram et Suez (Ehr!), ad Ouadi Dugla et in planitie Galala deserti Arabici (Schweinf!), in salsis Persiae mediae (Oliv., Ege), ad Damgam (Buhsel), inter Hawuz i Sultan et Kenaragird (Haussk!).

Caules pedales; species foliis carnosis crassis horizontaliter patentibus 3-4 lineas longis 2-3 lati distinctissima.

Obs. *A. Ehrenbergii* Schweinf. MSS. = *Salsola arabica* Ehrenb. foliis parvis ovato-globosis obtusis ad ramulos steriles imbricatis, antheris appendiculatis insignis sed cuius fructus adhucdum ignotus in Arabiâ tropicâ et Nubiâ nec in nostrâ ditione crescit.

8. **A. annua** (Bge Anab. p. 46) annua, præter axillas parce villosulas glabra, caule a basi trichotome ramosissimio hemispherice dumoso, foliis carnosis linearis-oblongis apice subincurvatis et seta subulata incurvâ demum deciduâ terminatis, floribus 3-5 in omnibus axillis glomeratis numerosissimis, staminodiiis rotundatis ciliolatis, perigonio fructifero pentaptero alis reniformi-orbiculatis subinaequalibus ⊖.

Hab. in collibus sterilissimis Persiae mediæ orientalis circa Tun Birdschaud et Isfesar (Bge). Non vidi.

Planta 2-7-pollicaris foliis aequalibus *A. setiferæ*.

BRACHYLEPIS (C.A. Mey. Fl. Alt. I, p. 370).

Flores hermaphroditi bibracteati solitarii. Perigonii phylla ad basin usque libera in fructu immutata non alata. Staminodia quinque distincta fimbriolata. Antheræ muticæ. Stylus stigmataque abbreviata. Utriculus succosus vel baccatus. Radiculae apex inferus. — Fruticulus vel herbæ ramis articulatis: foliis oppositis. — Genus proximum *Anabasi* a quâ perigonio fructifero non alato tantum differt.

1. **B. salsa** (C.A. Mey. loc. cit. p. 372) caulis e rhizomate lignoso persistentibus procumbentibus, ramis erectis numerosissimis pumilis simplicibus vel fastigiatis, foliis nunc obsoletis nunc patentibus subcylindricis obtusis seta deciduâ apiculatis, floribus in spicas terminales dispositis inferioribus subremotis superioribus approximatis, perigonii phyllis membranaceis demum supra medium subinduratis chartaceis ovatis et oblongis obtusis 3. Ic. Ledeb. Icon. tab. 48. — *Br. humilis* Lessing Linn. 1839, p. 198. — *Microlepis salsa* Eichw. Casp. p. 2.

Hab. in desertis argillosis salsis Turkestaniae australis (Lehm!).

Fruticulus dumosus 3-4 rarius 5-6-pollicaris.

Ar. Geogr. Rossia austro-orientalis, Songaria, Sibiria Uriensis.

2. **B. eriopoda** (C.A. Mey. in Schrenck Bull. Acad. Petersb. 1843, p. 360) caulis ex caudice crasso brevi villosissimo pluribus basi articulatis herbaceis paulo supra basin brachiatim bis vel ter ramosissimis hemisphericis, articulis elongatis, foliis ovato-triangularibus recurvis in aristam deciduam acuminatis, floribus ad cupulas internodiorum oppositis valde distantibus, perigonii phyllis hyalinis rotundatis vel ovatis fructiferis fere immutatis, semine magno ♀.

Hab. in argilloso-salsis Persiae orientalis (Bge), Turkestaniae in deserto Melik inter Buchara et Termine (Lehm!).

Planta sicca fragillima 6-9-pollicaris.

Ar. Geogr. Deserta Aralo-Caspica et Songaria.

Div. 2. Halomocnemideæ. — Sepala exteriora duo, alterum anticum, alterum posticum, interiora tria lateralia.

Subdiv. 1. — Semen a dorso (axi et bracteæ parallele) compressum.

NANOPHYTUM (Less. Linn. 1834, p. 197.)

Flores hermaphroditi (summi serotini interdum abortu feminei) bibracteolati. Perigonii phylla libera fructifera accreta chartacea non appendiculata. Staminodia brevia. Antherarum connectivum brevissime apiculatum. Stylus elongatus, stigmata filiformia subulata. Utriculus membranaceus, radiculæ apex inferus. — Fruticulus dense pulvinaris, foliis minutis alternis confertis acerosis, ramis non articulatis.

1. N. erinaceum (Pall. Ill. tab. 48 sub *Polygnemo*) dense pulvinatum caudice ramisque vetustis lignosis tortuosis procumbentibus, ramulis hornotinis brevissimis confertissimis imbricatim foliatis in axillis gemmiferis, foliis semiamplexicaulibus a basi latiore triquetro-subulatis pungenti-mucronatis, floribus in ramulorum apice 1-7 approximatis, bracteolis foliis conformibus et minoribus florem sub anthesi superantibus, phyllis perigonii enerviis fructiferis auctis stramineis fructum eis multo minorem involventibus 5. *N. erinaceum* Bge Anab. p. 51. — *N. Caspicum* Less. loc. cit. — *N. juniperinum* C.A. Mey. — *N. macranthum* Fisch. et Mey. Bull. Petersb. 1844, 2, p. 134. — *Polygnemum juniperinum* M.B. Mem. Mosq. I, p. 154. — *Anabasis Sieversii* Willd. in R. et Sch. Syst. VI, p. 223.

Hab. in salsis desertorum Turkestaniæ prope Juss Chuduk (Lehm.).

Ar. Geogr. Deserta Aralo-Caspica, Turcomania borealis, Songaria.

PETROSIMONIA (Bge Anab. p. 52. — *Halimocnemidis* Sp. Auctorum).

Flores hermaphroditi solitarii bibracteolati axillares. Perigonii phylla libera, interiora tria vel omnia vel 1-2 interdum abortiva, fructifera immutata. Staminodia nulla. Stamina 5-1, antheræ apice sœpe coherentes breviter apiculatæ. Stylus longe bicurvis. Utriculus tenuiter membranaceus. Radiculæ apex inferus. — Herbæ annuae pube adpressâ bicuspitatâ vestitæ non articulatæ, foliis plerumque tantum infimis oppositis, interdum omnibus alternis.

* Oligandræ alternifoliæ (Bunge).

1. P. Volvox (Pall. Ill. tab. 50 sub *Polygnemo*) annua pube bicuspitatâ canescens erecta a basi divaricatim ramosa ramis infe-

rioribus oppositis cæteris sparsis, foliis omnibus alternis filiformi-teretibus longis acutis muticis vel obsolete mucronulatis, floribus alternis remote spicatis, bracteolis ovato-lanceolatis carinatis conniventibus florem et fructum involventibus et superantibus, perigonii phyllis staminibusque tribus, antheris fugacibus apice non cohaerentibus ⊖. *P. Volvox* Bge loc. cit. — *Halimocnemis Volvox* C.A. M. Fl. Alt. — *H. triandra* Moq. in D.C. Prodri.

Hab. in planitiebus subsalsis Tauriae ad Sebastopol!, prov. Caucasicarum (Stev.). In Transcaucasiam deesse videtur.

Ar. Geogr. Transylvania, tota Rossia orientalis, regio Caspica et Siliria Altaica

• Pentandrae alternifoliae (Bge).

2. **P. crassifolia** (Pall. Ill. p. 64, tab. 55 sub *Polyenemo*) annua pube bicuspidatâ canescens vel viridis glabrata a collo ramosa caulis diffusis vel ascendentibus dichotome vel opposite ramosissimis, foliis oppositis et sparsis carnosis crassis linear-teretiusculis obtusiusculis inferioribus elongatis, floribus alternis in spicas densiores vel laxiores flexuosa dispositis, bracteolis navicularibus apice acute carinatis fructiferis cartilagineis basi obtuse gibbis fructum amplexantibus et cum eo deciduis, perigonii phyllis fere tantum binis, staminibus quinis, antheris apice inter se coherentibus ⊖. *P. crassifolia* Bge loc. cit. p. 55. — *Halimocnemis crassifolia* C.A. Mey. — *H. alternifolia* Moq. in D.C. Psodr. XIII, p. 198. — *H. oppositifolia* Eichw. Casp. p. 154. — *Polyenemum oppositifolium* Pall. — *P. glaucum* Pall. Ill. tab. 53. — *Halimocnemis glauca* Bge in Lehmann. № 1188 et C.A. Mey. Fl. Alt.

Hab. in salsis Tauriae (Stev. sub *H. glauca*), Transeaucasie in Grusia (Herb. Petrop. ex cl. Bge), in deserto ad Murchehar prope Ispahan (Haussk. ex Bunge).

Multum variat indumento, foliis tenuioribus et brevioribus, rectis vel recurvis.

Ar. Geogr. Rossia australis, littora Caspica septentrionalia, Songaria, Sibiria, Turkestania borealis.

3. **P. glauca** (Pall. Ill. p. 63, tab. 54 sub *Polyenemo*) annua pube bicuspidatâ adpressâ cana, caule erecto a basi ramoso ramis inferioribus oppositis cæteris alternis omnibus ascendentibus cerebre et breviter ramulosis, foliis omnibus alternis semiteretibus filiformibus acutis mucronatis falcato-recurvatis, floralibus bracteolas superantibus, floribus breviter et dense spicatis, bracteolis ovatis acute carinatis basi gibbis perigonium fructiferum amplectentibus et æquidistantibus, perigonii phyllis semper quinis, staminibus quinis, antheris apice coherentibus ⊖. *P. glauca* Bge loc. cit. p. 58. — *Halimocnemis glauca* var. *pentandra* Fenzl in Led. Fl. Ross. III, p. 840 ex parte.

Hab. in limoso-salsis totius Persiae occidentalis, prov. Aderbidjan (Szovt Auch. 5256!), circa magnum desertum salsum (Buhse sub *H. squarrosum* Bge!).

Valde affinis formis quibusdum *P. crassifoliae* quâcum a multis confusa, floribus isomeris distincta.

Ar. Geogr. Littora Caspia septentrionalia (ex Pall.).

*** Pentandræ oppositifoliae (Bge).

4. **P. brachiata** (Pall. Ill. pag. 62, tab. 52 sub *Polyenemo*) annua pube adpressâ bicuspidatâ lèvi canescens, caule erecto sàpe elato a basi ramoso ramis omnibus oppositis elongatis patentibus ramulosis, foliis omnibus oppositis carnosis a basi dilatatâ semiteretifiliformibus subtus carinatis obtusiusculis, floralibus abbreviatis rectangule patentibus sàpe recurvis, floribus oppositis dense glomerato-spicatis, bracteolis naviculari-ovatis foliaceo-attenuatis sàpe recurvis basi demum coriaceis gibbis, perigonii phyllis staminibusque quinis, filamentis calycem vix excedentibus ☽. *P. brachiata* Bge loc. cit. p. 59. — *Halimocnemis brachiata* C.A. Mey. Fl. Alt. -- *H. Hohenackeri* Presl Bem. p. 109.

Hab. in salsis Graecie ad Athenas (Urv.), Macedoniae ad prom. Kara Burnu et Thracie ad Lagos (Jankal!), Tauriae (Stev!), Armenie Rossicae (Buhse!), Georgiae Caucasicæ (Hoh.), Persiae bor. ad Caspium (Buhse!).

Ar. Geogr. Rossia australis, Sibiria Uralensis et Altaica.

Subdiv. 2. — Semen a latere (axi et bracteæ contrarie) compressum.

A. *Physandreae* Bge. — Antheræ vesiculoso-appendiculatæ.
Staminodia nulla.

HALOCHARIS (Moq. in D.C. Prodr. XIII, 2, p. 201, ex parte.
— *Halimocnemidis* sp. Auct.).

Flores hermaphroditæ vibranteolati solitarii axillares spicati. Perigonii phylla 5 ad basin usque libera membranacea fructifera immutata. Staminodia nulla. Stamina 5 filamentis disco insertis, antherarum loculi ad apicem usque discreti et ibi connectivo brevi conjuncti, connectivo in vesiculam formâ variam producto. Stylus ad medium in crures subulatos divisus. Radicula infera ascendens. — Herbæ annuae non articulatae pilis rigidis basis fixis articulatis hispidæ pubesque breviori crispulâ pubescentes, foliis alternis apice piloso-barbatis.

1. **H. hispida** (C.A. Mey. in Schrenck Bull. Petrop. 1843, I, p. 30 sub *Halimocnemide*) annua humilis pilis basi tuberculatis hispidissima a collo saepius ramosa caulinibus tenuibus ramosis et breviter ramulosis, foliis brevibus semiæcylindricis obtusis apice præsertim et patule setoso-barbatis primariis cito emarginatis, floralibus ovatis flore

sæpe brevioribus, floribus in axillis inferioribus remotiusculis versus ramulorum apicem capitato-spicatis, bracteolis hirsutissimis cum perigonio deciduis obtuse carinatis, perigonii phyllis binis externis villosissimis, filamentis perigonum excedentibus, antherarum appendice luteâ sessili compressâ lanceolatâ acutissimâ loculis æquilatis dimidio breviore ⊖. *H. hispida* Bge loc. cit. p. 62, tab. I, fig. 1.

Hab. in salsis deserti Persiae mediae orientalis (Bge).

Planta 3-6-pollicaris, folia 2-4 lineas longa.

Ar. Geogr. Turkestania borealis, Songaria.

2. ***H. sulphurea*** (Moq. l. cit. p. 201 ex parte) annua pilis tuberculatis hispidissima, caulis erectis vel a basi in ramos diffusos divisis, ramulis floriferis abbreviatis, foliis carnosulis linearis-subtrigonis obtusis primariis cito emarcidis, floralibus oblongis bracteolas lanceolatas carinatas hirsutissimas vix æquantibus, floribus imbricatim et brevissime spicatis, perigonii phyllis binis externis villosissimis cæteris glabris, filamentis perigonio brevioribus, antherarum appendice sulphureâ sessili oblongâ obtusâ subcompressâ loculis latiore et æquilongâ ⊖. *H. sulphurea* Bge l. cit. p. 63 et tab. I, fig. 2.

Hab. in arenosis salsis Babyloniae ad Bagdad, Mohammera et Kutt (Noë 67! 1012!), in Persie desertis (Auch. 2786, Bunge), inter Kasbin et Teheran (Buhse!), prope Buchir (Haussk!).

Semipedalis et clator.

3. ***H. violacea*** (Bge loc. cit. p. 63) annua hispidissima, caule erecto vel a collo in ramos diffusos diviso, ramulis floriferis abbreviatis, foliis carnosulis linearis-subtrigonis obtusis primariis cito emarcidis, floralibus oblongis bracteolas lanceolatas carinatas hirsutissimas vix æquantibus, floribus imbricatim et dense subcapitato-spicatis, perigonio toto lanugine contortuplicatâ obsito, filamentis calyce brevioribus, antherarum appendice violacea sessili obovato-inflatâ loculis quadruplo latiore et eis sublongiore ⊖. Ic. Bge tab. I, fig. 3. — *H. sulphurea* Moq. (ex parte quoad pl. Affghanam).

Hab. in salsis Persiae austro-orientalis prope Kerman (Bge), regno Cabulico ad Pischbulak, Candahar, Ghazni (Griff!), Belutschia prope Ghurgina (Stocks!).

Valde affinis præcedenti a qua appendicis colore et formâ praesertim differt.

4. ***H. clavata*** (Bge loc. cit. p. 64) annua hispidissima a basi in ramos diffusos ex omnibus axillis ramuligeris divisa, foliis carnosulis linearis-subtrigonis obtusis primariis cito emarcidis, floralibus bracteolas dense hirsutas subsuperantibus flore subbrevioribus, floribus imbricatim et dense subcapitato-spicatis, phyllis perigonii externis basi longe barbatis intimis binis glabris, filamentis calyceem excedentibus, antherarum appendice purpurascente stipite tenui sensim clavato suffultâ clavato-oblongâ cum stipite loculis sublatiore et duplo

longiore ♂. Bge tab. I, fig. 4. — *Halocharis sulphurea* Moq. ex parte.

Hab. in arenosis Affghaniæ prope Karabagh et Jomy (Griff.).
Planta 2-5-pollicaris facie omnino præcedentium.

HALIMOCNEMIS (C.A. Mey. Fl. Alt. I, p. 381).

Flores hermaphroditæ bibracteolati in axillis solitarii. Perigonii phylla enervia quina rarius 3-4 tandem basi excrescente osseo-indurata coalescentia non appendiculata. Staminodia nulla. Stamina tot quot phylla disci margine inserta, antheræ vesicoso-appendiculatæ. Styli crura elongata obtusa vel lacero-truncata. Radiculae apex superus porrectus. — Herbae annuae carnosæ non articulatae, foliis præter infima alternis in cuspidem cartilagineam sæpius deciduam abeuntibus. Pubes duplex altera pilis elongatis articulatis, altera pilis brevibus constans, interdum evanida.

1. **H. Karelini** (Moq. l. cit. p. 196) annua glaberrima glauca. caule pumilo crassiusculo a collo ramosissimo ramis carnosis in hemispheram dispositis, foliis flaccidis basi semiamplexicaulibus semitereti-linearibus elongatis obtusis submuticis, floralibus late ovato-triangularibus acutis late membranaceis, floribus in omnibus axillis ob internodia brevissima approximatis, bracteolis folia floralia subsuperantibus late ovatis obtusis basi valde obliquis et late marginalis, perigonii phyllis quinis lanceolatis superne chartaceis longe acuminate in conum angustum conniventibus bracteolas longe superantibus, styli ramis crassiusculis obtusis, antheræ appendice subsessili globosâ loculis multo breviore ♂. Ic. Bge tab. 2, fig. 6.

Hab. in salsis deserti Turkestaniæ ad Kuwendarja (Lehm!), Turcomania ad Caspium (Karel!).

Plantula 2-5-pollicaris. Species valde affinis *H. sclerospermæ* Pall. Ill. tab. 56 in ditione non obviae et floribus 3-4-meris, bracteolis acutis, etc., distinctæ.

Ar. Geogr. Deserta Aralo-Caspica et Songaria.

2. **H. pilosa** (Moq. loc. cit. p. 195) annua pube dupli pilosa et canescens, caule a basi divaricatum ramosissimo ramis in hemispheram sæpius dispositis, foliis carnosulis semitereti-trigonis in cuspidem cartilagineam subpungentem deciduam abeuntibus, floralibus abbreviatis, floribus in axillis superioribus remotiusculis, bracteolis folio florali subbrevioribus late ovato-lanceolatis perigonio brevioribus ejus basi adnatis et cum eo deciduis in cuspidem acutissimam deciduam attenuatis, perigonii fructiferi basi rotundati glabri phyllis quaternis lanceolatis cuspidato-acuminatis apice in conum conniventibus, styli ramis obtusis, antheræ appendice roseâ stipitato-clavata obtusâ loculis longiore et latiore ♂. Ic. Bge t. I, fig. 7. — *H. villosa* Bge Pl. Lehm. p. 299 ex parte.

Hab. in desertis salsis Persiae borealis et mediæ, ad fontem Gendab jugi Elbrus (Buhsel!), circa Mesinan, Nichapur, Meschhed (Bge), ad salinas Nähend (Seidl!), in deserti Kisilkum Turkestaniæ collibus Ssusus Kara (Bge).

Tripollicaris interdum semipedalis, folia in planta vegetiore pollicem et amplius longa sæpius 7-8 lin. longa, perigonium fructiferum 5 lineas longum.

3. **H. villosa** (Kar. et Kir. Enum. 1841, N° 687) annua pube duplii alterà pilis longis articulatis alterà pilis brevibus simplicibus patule villosa, caule crassiusculo a basi ramosissimo ramis globum sæpius formantibus, foliis semiteretibus crassis pungentimucronatis rectis patentibus floralibus bracteolas superantibus, floribus in omnibus axillis solitariis racemos breves foliatos densiusculos formantibus, bracteolis carinatis basi gibbis flores superantibus ab eis liberis, perigonii phyllis 4 rarius 5 lanceolatis in conum demum conniventibus, fructifero usque ad basin globosam 4-5-loveolatam villosa, styli ramis dilatatis truncatis, antheræ appendice oblongâ sessili loculos subæquante ♂. Ic. Bge tab. I, fig. 8.

Hab. in desertis salsis ad fluvium Kuwandaria Turkestaniæ (Lehm!).

Dumulus diam. interdum pedalis vel multo minor, folia crassa exsiccatione rigida horizontalia pollicaria.

Ar. Geogr. Songaria.

4. **H. macranthera** (Bge Pl. Lehm. p. 475) annua glauca pube brevi canescens pilis elongatis subnullis, caule crasso carnoso a basi hemispherice ramosissimo, foliis semiteretibus cuspidato-pungentibus, floralibus flore longioribus, floribus in omnibus axillis solitariis racemos breves foliatos formantibus, bracteolis acute carinatis et basi gibbis cuspidatis a perigonio liberis et eo longioribus, perigonio fructifero basi valde indurato truncato margine calloso prominente phyllis dorso lanatis acutissimis in conum conniventibus, styli ramis apice dilatatis et flabellato-dentatis, antheræ appendice roseâ ovata loculis elongatis triplo breviore ♂. Ic. Bge t. I, fig. 9.

Hab. in monte Ssusus Kara deserti Kisilkum Turkestaniæ (Lehm!).

Facies et folia *H. villosæ*, robustior, perigonio fructifero et antheris ab eâ distinctissima.

5. **H. mollissima** (Bge Anab. p. 71) annua glauca breviter patule pubescens et pilis longissimis mollissimis strictis articulatis sericea, caule crasso a basi ramosissimo ramis globum amplum formantibus, foliis semitereti-trigonis cartilagineo-cuspidatis longis, floribus in omnibus axillis solitariis, bracteolis folio florali multo brevioribus florem pentamerum superantibus et cum perigonio deciduis, perigonii fructiferi basi osseâ globosâ margine calloso cinctâ phyllis sericeo-villosis acutis in conum conniventibus, styli ramis brevibus dilatatis lacero-dentatis, antheræ appendice aurantiacâ oblongâ sessili dimidios loculos æquante ♂. Ic. Bge tab. I, fig. 10.

Hab. in salsis montosis Tabat Kuh prope Meschhed Persiae boreali-orientalis (Bge). Non vidi.

Diametro pedalis, folia inferiora sesquipollucaria. Species indumento insignis, pilorum longiorum articuli constant cellulâ elongatâ cylindricâ apice in coronulam multifidam abeunte, more ramuli *Equiseti*.

HALOTIS (Bge Anab. p. 73).

Flores hermaphroditici bibracteolati in axillis solitarii. Perigonii phylla quina in fructu basi excrescentia coalescentia osseo-indurata, exteriora bina trinervia in fructu medio dorso auriculato-gibba. Staminodia nulla. Stamina quinque disci margini inserta. Antheræ vesicoso-appendiculatae. Styli rami breves dilatati laceri. Radiculae apex superus porrectus. — Herba annua. — Genus præter phylla externa demum dorso gibboso-auriculata *Halimocnemidi* simillimum et forsitan (ex ipso Bge) melius hujus sectio.

1. **H. occulta** (Bge loc. cit. p. 74) annua pubescenti simplici patula canescens pilis longioribus articulatis paucis evanidis intermixtis, caule humili a basi in ramos dichotomos angulatos diviso, foliis a basi semiamplexicauli tereti-trigonis in euspidem brevissimam deciduam abeuntibus patulis apice subrecurvis, floralibus conformibus, floribus axillaribus solitariis, bracteolis carnosulis late ovatis acuminatis carinatis valvatim conniventibus florem et fructum foventibus et occultantibus cum eo deciduis, gibbis phyllorum exteriorum perigonii fructiferi medio dorso sitis, phyllorum parte superiore membranaceâ in conum connivente, antheræ appendice ovato-oblongâ utrinque attenuatâ purpureâ loculis sublongiore ♂. Ic. Bge t. I, fig. 11.

Hab. in collibus salsis prov. Chorassan Persiae inter Tun et Birdchand prope pagum Afsel Abad (Bge). Non vidi.

Planta ad summum semipedalis, folia 8-6 lineas longa lineam crassa, perigonium sericeo-pubescentia sub anthesi 3-4 lin. longum.

PIPTOPTERA (Bunge in litt.).

Flores hermaphroditici bibracteolati in axillis solitarii. Perigonii phylla quina, anticum et posticum trinervia, interiora tria enervia; omnia in fructu basi indurata connata exterioribus latissime membranaceo-connatis alis articulatum et circumscisse deciduis. Fructus alarum basi superstite supra medium biauriculatus apicibus phyllorum connatis coronatus a latere subcompressus sexcostatus. Staminodia nulla. Antheræ membranaceo-appendiculatae. Styli erubes subulati conniventes. Radiculae apex superus. — Fruticulus ramosissimus pubescenti hispidulus habitu *Salsolæ* crassæ.

1. **P. Turkestanica** (Bge in litt.) fruticulosa patula et alternatim ramosissima breviter pubescentia, foliis alternis linearifiliformibus

crassis apiculatis, floralibus abbreviatis semiamplexicaulibus triangulare-lanceolatis fructu brevioribus, floribus sparsis solitariis in ramos spiciformes laxos inferne ramulosos dispositis, perigonio fructiferò alis flabellatis eo longioribus aucto ♂.

Hab. in argillosis salsis Turkestaniae prope Schurachan prope fortalitium Novo-Alexandrowsk (Korolkow).

Fragmenta tantum vidi, fructus piso paulo minor.

HALARCHON (Bge Anab. p. 75. — *Halocharidis* Moq. species).

Flores hermaphroditi bibracteolati in axillis solitarii. Perigonii phylla quina hyalina enervia libera, fructifera nec appendiculata nec basi indurata. Staminodia nulla. Stamina 5 disci minutis margini, inserta. Antheræ vesiculoso-appendiculatae. Stylus simplex versus medium verticillatim denticulis auctum, stigma breviter capitato-bilobum cinctum indusio membranaceo profunde radiato-lacero prius late infundibuliformi tandem peracta fœcundatione reverso pileiformi. Radiculæ apex superus. — Herba annua. — Genus ab *Halimocnemide* et *Halotide* calyce fructifero basi non indurato et styli stigmatisque structurâ distinctum.

1. **H. vesiculosus** (Moq. in D.C. Prodr. XIII, p. 202 sub *Halocharide*) annuus glauco-caesius pube simplici brevissimâ scabrido-canescens, caule a basi in ramos patentes vel decumbentes abeunte, foliis semitereti-subtrigonis præter ima opposita alternis pungenticepidatis, floralibus conformibus, floribus axillaribus solitariis inferioribus remotiusculis cæteris congestis, bracteolis cuspidatis foliis valde superatis, perigonii phyllis oblongo-lanceolatis acutissimis bracteolas superantibus, antheræ loculis linearibus angustissimis discretis, appendice roseâ obovatâ amplâ loculis sublongiore et multo latiore ⊖.
H. vesiculosus Bge l. cit. p. 75, tab. I, fig. 12, 25, 26.

Hab. in arenosis ad flavium Turmik Affghaniæ prope urbem Candahar (Griff. Journ. 718!).

Pedalis, folia 8-9 lineas longa, antheræ vesicula 2¹/₂-3 lineas longa. Species gregaria vesicularum colore pulcherrima.

GAMANTHUS (Bge Anab. p. 76. — *Halocharidis* Moq. spec.).

Flores hermaphroditi bibracteolati in axillis solitarii denique basibus foliorum floralium auctis connatis et induratis inclusi et cum eis decidui. Perigonii phylla 5 discreta membranacea vel chartacea fructifera non indurata. Staminodia nulla. Stamina 5 disco minutissimo inserta, antheræ vesiculoso-appendiculatae. Stylus elongatus bieruris erubibus subulatis. Radiculæ apex superus. — Herbae annuae.

1. G. gamocarpus (Moq. loc. cit. p. 201 sub *Halocharide*) annuus glabrescens vel pube densâ brevi canescens, caule a basi opposite vel dichotome ramosissimo ramis et ramulis divaricatis, foliis semiteretibus carnosis spinuloso-cuspidatis recurvis oppositis et sparsis, floralibus oppositis basi connato-vaginantibus et demum osseoinduratis, floribus in ramulorum apicibus geminatis oppositis axillis foliorum floralium omnino inclusis, bracteolis inclusis sed liberis, perigonii phyllis lanceolatis acutis cartilagineo-acuminatis basi breviter binerviis, antheræ appendice albâ oblongo-ellipticâ clavatâ stipite angustâ longiusculâ suffultâ loculis breviore et vix latiore ♂. *G. gamocarpus* Bge l. cit. p. 77, tab. 1, fig. 14.

Hab. in desertis salsis, in toto ambitu deserti magni Persiae mediae (Bél., Bge), in regno Cabulico ad Turnuk prope Kandahar (Griff. Journ. 719!).

Semipedalis et procerior, folia fere lineam diametro lata, inferiora pollicaria. Perigonium foliis floralibus inclusum et cum eis tandem deciduum sed cum eis et bracteolis non coalescens.

2. G. commixtus (Bge Anab. p. 76) annuus pube simplici villosulus, caule diffuso patentim a basi ramoso, foliis præter infima alternis patentissimis semitereti-linearibus spinuloso-cuspidatis, floralibus alternis congestis basibus incrassato-induratis confluentibus fructus foventibus, floribus 3-6 in apice ramulorum alternis congestis, bracteolis oblique lanceolatis carinatis apice spinulosis hamato-recursivis flore sublongioribus a folio florali liberis, perigonii phyllis chartaceis oblongo-lanceolatis acuminatis apice subpungentibus enerviis, antheræ appendice aurantiacâ obovato-oblongâ longe stipitatâ loculis longiore et duplo latiore ♂. Ic. Bge t. I, fig. 13. — *Halocharis vesiculosa* Moq. ex parte.

Hab. in salsis regni Cabulici prope Byam, Syghan, Chuschki et in montibus Gulguli (Griff. 1755!).

Semipedalis et longior, folia 6-8 lineas longa, calyx $2\frac{1}{2}$ lineas longus.

3. G. ovinus (Bge in litt.) annuus pube simplici adpressâ pubescentis tandem glabratus viridis, foliis præter infima alternis patentissimis semitereti-linearibus spinuloso-cuspidatis, floralibus in apice ramulorum congestis flores foventibus, floribus inferioribus alternantibus summis confertis saepe geminis suboppositis, perigonii phyllis acutis, antheræ appendice stipitatâ globosâ loculo breviore citrinâ ♂.

Hab. in deserto Kisilkum Turkestaniae (Bge). Ovium pabulum.

Fructus ignotus, affinis *G. commixto*, indumentum parcus et appendicis structura diversa. Antheræ $1\frac{1}{2}$ lin. longæ loculis basi discretis, connectivum productum in stipitem cavum lineæ $\frac{1}{2}$ longum, vesicula diam. lineam dimidiam lata.

4. G. pilosus (Pall. Ill. p. 28, tab. 20 sub *Salsold*). annuus carnosus glaucus pilis minutis adpresso aliisque articulatis patentibus sparsis obsitus, caule a basi ramosissimo ramis inferioribus oppositis

patentibus superioribus ramulisque alternis hisce fructiferis clavato-incrassatis, foliis plerisque alternis semiteretibus acutis vel breviter mucronatis rectis denique recurvis, floralibus alternis congestis abbreviatis demum basi incrassatis osseo-induratis flores alternos solitarios includentibus, bracteolis oblongo-lanceolatis subcarinatis basi gibbis florem æquantibus cum foliis floralibus basi coalescentibus, perigonii phyllis hyalinis linearis oblongis liberis apice retuso-bilobis subconvolutis exterioribus binis obsolete trinerviis interdum medio dorso tandem subgibbis, antheræ appendice sulphureâ globosâ sub sessili loculis quadruplo breviore \odot . *G. pilosus* Bge loc. cit. t. 1, p. 15. — *Polyneum malacophyllum* M.B. Mém. Mosc. I, p. 152. — *Halimocnemis malacophylla* C.A. Mey. Fl. Alt. I, p. 387. — *Halim. hirsuta* G. Koch Linn. XXII, p. 192.

Hab. in desertis salsis Iberiæ Caucasicæ et Grusiæ (C.A. M! Hoh! C. Koch). Planta demum pedalis, folia $\frac{1}{2}$ -1-pollicaria lineam crassa.

**HALANTHIUM (C. Koch Linn. XVII, p. 313.—*Halogetonis* sp. Fenzl.
— *Physogeton* Moq.).**

Flores hermaphroditæ bibracteolati in axillis solitarii. Perigonii phylla quinque ad basin usque libera, duo exteriora trinervia fructifera dorso alata, interiora immutata vel unum minute alatum. Discus minutissimus non accrescens. Staminodia nulla. Stamina 5 disco inserta. Antheræ vesiculoso-appendiculatae. Stylus et stigmata subulata elongata. Radiculae apex superus. — Herbæ annuae juniores pubescentes canescentes carnosæ.

1. **H. Mamamense** (Bge Anab. p. 82) annua glauca junior villosa, caule a basi in ramos divaricatos infimos oppositos cæteros alternos diviso, ramulis fructiferis tandem incrassatis, foliis præter infima alternis teretibus apice sæpe penicillato-pilosus obtusis breviter mucronatis brevibus recurvis cito evanidis, floralibus conformibus sensim abbreviatis, floribus in axillis superioribus ramulorum brevium solitariis paucis approximatis, bracteolis lanceolato-triquetris subcarinatis cuspidatis flore brevioribus, perigonii phyllis membranaceis oblongo-lanceolatis acuminatis duobus exterioribus dorso minute alatis tertio corniculato-gibboso, antheræ appendice flavescente denique albidâ subglobosâ loculos dimidios superante \odot . Bge t. I, fig. 18. — *Hal. molle* et *H. rigidum* Bge in litt. — *Halogeton* sp. Boiss. et Buhs. Enum. p. 190.

Hab. in salsis Persiæ occidentalis, ad Tabris (Buhse), ad salinas Manian (Seidl!).

Semipedalis vel humilior. Folia majora semipollicaria lineam vix crassa.

2. **H. Kulpianum** (C. Koch loc. cit. sub *Halimocnemide*) annum glaucum villosum demum glabratum, caule a basi in ramos

infimos sparsos divaricatim prostratos elongatos cæteros alternos soluto, foliis præter infima alternis semiteretibus obtusis muticis vel mucronulatis, floralibus abbreviatis bracteolas subæquantibus basi denique cartilagineo-gibbis incrassatis, foliis in omnibus axillis solitariis inferioribus remotis ad ramulorum apicem 3-4 approximatis, bracteolis fructiferis accretis basi cartilagineo-gibbis recurvis basi breviter inter se connatis, perigonii phyllis hyaliuis obtuse bilobis exterius in alam latam decolorem excrescentibus, antheræ appendice flavescenti-albidâ subglobosâ sessili loculos dimidios subæquante ⊖. *Halimocnemis pycnantha* C. Koch Linn. XXII, p. 193.

Hab. in salsis interioribus Armeniae Rossicie ad Araxem prope salinas Kulp (C. Koch! Seidl!), in ditione Pennek (C. Koch).

Folia $\frac{1}{2}$ -1-pollicaria diam. lineam dimidiata lata, bracteolæ 4-5 lineas, phylla perigonii 3 lineas longa.

3. ***H. rarifolium*** (C. Koch Linn. XVII, p. 314) annum canescens levatum vel glabratum a basi ramosissimum ramis infimis oppositis demum elongatis cæteris alternis, foliis semiteretibus obtusis muticis, floralibus late oblongis obtusissimis molliter cuspidatis flore multo brevioribus, floribus solitariis axillaribus inferioribus distantibus summis congestis demum sæpe elongato-spicatis, bracteolis ovato-oblongis concavis, perigonii phyllis distinctis linear-oblängis obtusissimis retuso-bilobis membranaceis exterioribus binis late alatis internis immutatis vel unico minute alato, antheræ appendice in stipitem longe attenuatâ transverse oblongâ ⊖. Ie. Bge tab. 1, fig. 19-22.

Species indumento, relativâ longitudine florum et fol. floralium, appendicis colore mire varians. Varietates sequentes proposuit am. Bge.

α. Aucherianum. — Appendices sulphureæ, folia floralia obtusa, indumentum densum. — *H. Aucheri* Moq. in D.C. Prodr. I. cit. p. 204. — *Halogenetum rarifolius* Fenzl. — *Halimocnemis Kowalewskyi* Szegl. Bull. Mosq. XXVI, tab. 5. — *Hal. robustum* Bge MSS.

Hab. in salsis Armeniae Rossicie prope Nachitschevan (Kowal! Szov!), prope Kulp (Szov! Seidl! Auch. 2771).

β. Abichii. — Appendices sulphureæ, folia floralia acuta, planta glabrescens vel glabra. — *Halocharis Abichii* C. Koch Linn. XXIV, p. 91.

Hab. cum precedente prope Nackitschevan et Kulp (Abich., Szov!).

γ. Belangeri. — Appendices albideæ, planta incano-lanata. — *Halimocnemis Belangeri* Moq. I. cit. p. 196. — *Hal. lanatum* Bge in litt. — *Halocharis pycnantha* C. Koch ex parte.

Hab. in Armenia Rossicie ad Erivan (Szov!), Persiæ (Belang.), prope Khoï prov. Aderbidjan (Szov!).

γ. lilacinum. — Appendices pallide lilaciæ, planta glabrata gracilis ramis mutantibus. — *Hal. lilacinum* Bge MSS. — *H. spicatum* Bge MSS.

Hab. ad lacum Urmiah Persiæ (Seidl!).

4. H. purpureum (Moq. Chenop. Enum. p. 158 sub *Halimocnemide*) annum breviter et molliter canescens pilis articulatis longioribus parce ciliatum, collo ramosum humile ramis erecto-patulis crassis flexuosis alternatim et breviter ramulosis, foliis praeter intima opposita alternis a basi cavâ teretibus crassis flavidо-cuspidatis patentibus vel recurvis, floralibus abbreviatis bracteolas et florem excedentibus, floribus solitariis secus ramulos breves subdistiche congestis, bracteolis lanceolatis, perigonii phyllis linearis-oblongis subconcavis obtusis retusis cum mucrone, antheræ appendice purpureâ obovatâ subsessili loculis apice connectivo brevi conjunctis tongiore ⊖. *Hal. p. purpureum* Bge tab. 1, fig. 23. — *Halogeton purpureus* Fenzl. — *Physogeton acanthophyllum* Jaub. et Sp. Ill. t. 135.

Hab. in Persia occid. prope Hamadan (Auch. 2777!), inter Seytun et Behshān (Haussk!).

Planta 3-6-pollicaris, folia crassa secca rigida et subpungentia, majora pollicaria, species antherarum vesiculâ purpureâ $2\frac{1}{3}$ linea longâ elegans.

B. Halogetonea. — Antheræ non vesiculoso-appendiculatae.
Staminodia distincta.

CORNULACA (Del. Eg. p. 234).

Flores hermaphroditi in axillis solitarii vel glomerati lanâ involuti, singuli bibracteolati vel in glomerulis externi 2-3-bracteolati interior ebracteolatus. Perigonii phylla quinque basi demum inerescente nuclei-formi indurata connata superne breviter libera hyalina denticulata, anticum fructiferum dorso in spinam eo multo longiore ex crescens. Staminodia quinque cum filamentis disco inserta eis connata et ovarium superantia. Stamina quinque antheris muticis vel apiculatis. Ovarium a latere compressum, stylus ad medium in erura subulata divisus. Radiculae apex superns. — Fruticuli rarius herbae annuae foliis subulatis vel spinosis, axillis setigeris.

1. C. Aucheri (Moq. in D.C. Prodr. I. cit. p. 218) annua, herbacea humilis a collo ramosissima minute seabrida, foliis crebris a basi subsemamplexicauli subulatis cuspidato-pungentibus patentibus, axillis fasciculigeris parce setiferis, floribus in axillis superioribus ternis glomeratis basi parce et breviter lanatis, perigonii floriferi phyllis spathulato-bovatis apice eroso-laceris, staminodiorum tubo phylla subaequante, perigonii fructiferi spinâ longe subulata acutissimâ ⊖.

Hab. In desertis Assyriæ (vel potius Persie) (Auch. 2801!), in Persia media occidentali circa desertum magnum salsum circa Rischim (Buhse!), inter Kaschan et Kum (Bge), in arenosis a l. B. Shir (Haussk!).

Herba 4-5-pollicaris, folia 3-4 linea tantum longa. Specimen floriferum

Aucherianum cum Buhseanis et Bungeanis ætate jam provectis conspecificum omnino videtur.

2. C. monacantha (Del. l. cit. p. 234, tab. 22) fruticosa glauca, ramis elongatis, foliis basi amplexicaulibus triquetro-subulatis pungentibus patulis, axillis fasciculigeris breviter lanatis, floribus in axillis 3-5 glomeratis lanâ densâ eos æquanti immersis, perigonii phyllis linearis-spathulatis obtusissimis apice eroso minute denticulatis, staminodiorum urceolo dimidium perigonum aequante, spinâ perigonii fructiferi subulatâ phyllis longiore ♂. *Salsola ferox* Lippi. — *Kochia monacantha* Steud.

Hab. in desertis Egypti ad Pyramides (Del., Sieb! Schweinf!), prope Suez (Bové!), in deserto ad Oasin Farafreh (Asch!), in deserto Ægyptiaco-Arabico et ad littus maris Rubri (Schweinf!), in Persia media orientali ad desertum salsum magnum et in Affghaniam occid. (Bge).

Suffrutex 3-5-pedalis, folia 2-3 lineas longa, flores 1½ lineas longi, spina perigonii fructiferi 2-3 lineas longa. Planta Persica quam non vidi ex am. Bunge paululum differt asperula nec laevis, foliis sublongioribus.

Ar. Geogr. Nubia.

3. C. amblyacantha (Bge Anab. p. 88) fruticosa glauca dense et congeste cespitosa ramis abbreviatis, foliis semiamplexicaulibus imbricatis ovato-triangularibus cuspidatis, axillis breviter et parce lanatis, floribus in axillis superioribus occultatis solitariis rarius ternis lanâ brevi parcâ cinctis, perigonii phyllis apice subtrilobolaceris, staminodiorum urceolo dimidium perigonum superante, perigonii fructiferi spinâ obtusâ phyllis breviore ♂.

Hab. in Persia australi loco non notato (Auch. 5263!).

Fruticulus nanus pulvinatus, folia eis præcedentis latiora breviora, huic characteribus affinis sed facie distinctissima. Species habitu et foliis affinis sed cuius flores adhucdum ignoti, *C. Ehrenbergii* Aschers. in Schweinf. Beitr. p. 189, in Arabiâ, Nubiâ et Abyssiniâ tropicis extra fines nostros crescit.

4. C. setifera (Moq. in D.C. Prodr. XIII, 2, p. 218) fruticosa pallide virens, ramis elongatis, foliis ramorum infimis squamæformibus cæteris subulato-spinosis sensim elongatis, axillis longe setiferis, floribus subsolitariis in axillâ occultatis lanâ brevi densâ stipatis, perigonii floriferi phyllis linearis-spathulatis lacero-dentatis, staminodiorum tubo sepalis breviore, perigonio fructifero.... ♂. *Astragalus setiferus* D.C. Prodr. II, p. 296.

Hab. in Mesopotamiâ inter Aleppo et Bagdad (Oliv!). V. in herb. D.C.

Species insignis facie omnino *Tragacanthæ* foliolis orbatae, a nemine post Olivierum collecta. Frutex pedalis, ramis annotini 3-4-pollicares, folia superiore ramulorum 12-15 lin. longa tenuia valde pungentia; fructus ignotus, tamen de genere (ex cl. Bge) nulla dubitatio.

HALOGETON (C.A. Mey. Fl. Alt. I, p. 378 ex parte.
— *Agathophora* Bge).

Flores abortu polygami in axillis glomerati lanâ densâ stipati, medius ebracteolatus exteriôres 2-3-bracteolati. Perigonii phylla quinque sublibera vel exteriora bina tantum vel tria aut omnia dorso alata vel gibba. Staminodia 4-5 cum filamentis connata. Stamina 5 rarius 3 vel 2, antheræ breviter vel non apiculatæ. Ovarium ovatum, stylus brevis, stigmata duo filiformia. Radicula supera. — Fruticui vel herbæ annuae foliis carnosis.

1. **H. alopecuroides** (Del. Descr. Eg. p. 222, tab. 21, sub *Salsola*) fruticosus humilis ramosissimus ramis elongatis virgatis patentibus vel curvatis albis glabris, foliis sparsis in ramulis vegetis distantibus semitereti-subclavatis supra planis subtus convexis crassis brevibus pungenti-mucronatis vel (ex el. Schweinf.) muticis, axillis lanatulis, floribus in axillis glomerulos globosos bracteolis orbiculari-ovalis intermixtos formantibus, perigonii phyllis sub anthesi hyalinis apice eroso-denticulatis, fructiferis antico et postico vel ternis vel omnibus suprà medium alâ ovato-flabellatâ auctis alis sœpius inæqualibus, staminibus et staminodiis quinis 3. *Anabasis alopecuroides* Moq. in D.C. — *Halogeton alopecuroides* Moq. Chenop. — *Agathophora alopecuroides* Fenzl, Bge. — *Traganum nudatum* Schimp. exs. non Del.

Hab. in desertis Egypti ad Pyramides (Del., Schimp!), ad Ouadi Dugla prope Kahiram et planicie Galala deserti Arabici (Schweinf!), Arabiâ petrea (Schimp! Boiss!). Arabice *El Hand*.

Fruetus pentapterus ut Schweinfurth observavit et ego in speciminiibus ab eo donatis observavi, generis *Agathophoræ* character essentialis igitur evanescit.

2. **H. glomeratus** (M.B. Mem. Mosq. I, p. 110 sub *Anabasi*) annuus glaber glaucus humilis a basi ramosissimus, ramis procumbentibus ascendentibus et erectis in hemisphæram densam excrescentibus, foliis brevibus carnosis setâ subulatâ mox evanidâ terminatis, glomerulis axillaribus approximatis spicatis, floribus glomerulii exterius feminis cæteris hermaphroditis diandris, bracteolis oblongis concavis, staminodiis 4, perigonii fructiferi phyllis sœpius omnibus alâ reniformi auctis, alis inæqualibus ⊖. *H. glomeratus* C.A. Mey. Fl. Alt. I, p. 378. — Ledeb. Ic. tab. 40.

Hab. in Turcomaniâ ad Caspium (Karell), Turkestaniâ prope Samarca d (Lehm!), regno Cabulico in valle Bamian (Griff. Journ. 1121!).

Planta ad summum semipedalis, folia majora 4 lineas longa.

Ar. Geogr. Sibiria Uralensis et Altaica.

SEVADA (Moq. in D.C. Prodr. XIII, p. 154).

Flores polygami 6-7 glomerati flore medio majore ebracteolato, laterilibus 2-3 bracteolatis minoribus binis flore medio proximis ebracteolatis cæteris abortu feminineis. Perigonii phylla sub anthesi ad medium usque concreta superne concava. Staminodia quinque obovata brevia carnosula glabra in floribus hermaphroditis inter se libera vix imâ basi cum filamentis cohærentia, in floribus feminineis nulla. Stamina quinque longe exserta, antheræ non appendiculatæ. Ovarium lageniforme, stylus subnnullus, stigmata breviter subulata dense papillosa. Fructus..... — Frutex facie *Suedæ vermiculatæ*. Genus fructu ignoto imperfecte descriptum ob ovarii formam semen verticale nec horizontale indicans et ex cl. Bunge inter *Anabaseas* juxta *Halogeton* militans.

1. **S. Schimperi** (Moq. loc. cit.) fruticulosa glabra humilis ramosissima, ramis primariis erassiusculis lignosis tortuosis cortice albida, annotinis rectis gracilibus rigidulis patentim et alternatim ramulosis, foliis parvis alternis vel suboppositis patulis obovato-oblongis basi attenuatis obtusissimis siccis nigricantibus, floralibus glomerulosis subsuperantibus vel æquantibus, axillis parce setiferis, glomerulis densilloris axillaribus in spicas breves basi interruptas approximatis, perigonii sub anthesi phyllis ovatis, perigonio fructifero..... 5. *Sueda platyphylla* Ehrenb. Ms. — *Sueda monoica* et *S. baccata* (Schimp. Un. Itin. № 867!).

Hab. in collibus granitosis et littore Maris Rubri prope Djedda Arabiæ tropicæ (Schimp!). Nondum intra fines ditionis nostræ observata.

ORD. C. AMARANTACEÆ.

(Juss. Genera 87.)

Flores hermaphroditi rarius polygami vel monoici. Perigonium scariosum 3-5-partitum aestivatione imbricatum. Stamina 3-5 phyllis opposita vel pauciora basi libera vel coalita, staminodiis interdum aucta. Ovarium liberum uniloculare uni vel pluriovulatum oyulis funiculis liberis e fundo ovarii erectis affixis ascendentibus vel pendulis amphitropis. Stylus simplex vel stigmata plura. Fructus mono vel polyspermus raro caryopsideus vel baccatus, saepissime utricularis membranaceus indehiscens vel circumscisse dehiscens. Semina saepius verticalia crustacea. Embryo areuatus vel annularis albumen farinaceum eingens. — Herbæ vel suffrutices foliis exstipulatis, floribus tribracteatis rarius bibracteatis.

TR. I. CELOSIEAE Endl. — Antherae biloculares. Ovarium multiovulatum. Folia alterna.

CELOSIA (L. Gen. p. 343).

Flores hermaphroditii tribraeatei. Perigonium phyllis quinque aequalibus erecto-patulis. Stamina quina inferne in cupulam connata, antherae biloculares. Staminodia nulla. Ovarium uniloculare multiovulatum, stigmata 2-3 minuta recurva. Utriculus ovatus transverse circumscissus polyspermus calyce involutus. Semina verticalia lenticulari-reniformia testa crustacea. Albumen centrale. Embryo annularis, radicula umbilico proxima. — Herbae floribus in spicas vel paniculas terminales et axillares dispositis nitidis, bracteis concavis coloratis persistentibus.

I. **C. argentea** (L. Sp. p. 296) annua glabra, caule erecto stricte ramoso, foliis subpetiolatis lineari-lanceolatis acutis, spieis longe pedunculatis oblongis vel cylindraceis demum obtusis nitidis albis, floribus sessilibus dignis, phyllis bracteas multo superantibus mucronulatis carinatis subtrinerviis, utricleis ovalo-pyriformibus ⊖.

Hab. in Affghanista prope Jalalabad (Griff!), in Egypto indicata sed an recte? Planta subtropica vix ditionis.

Ar. Geogr. Nubia, Abyssinia, India, China.

TR. II. ACHYRANTHEAE Endl. — Antherae biloculares. Ovarium uniovulatum. — Folia alterna vel opposita.

HABITZIA (Bieb. Cent. Ross. tab. 54).

Flores hermaphroditii vel abortu polygami tribraeatei. Perigonium quinquepartitum laeiniis aequalibus patentim rotatis. Filamenta quina filiformia libera, antherae biloculares ovatae. Staminodia nulla. Stigmata 2-3 subulata. Utriculus depresso-orbicularis calyce inferne stipatus transverse marginatus et circumscissus. Semen horizontale lenticulare testa crustacea, arillo nullo. Albumen centrale, embryo annularis. — Herba perennis volubilis facie *Tami*.

I. **H. tamnoides** (M.B. Taur. Cauc. III, p. 170) perennis glabrescens vel subrufescens, radice validâ oblongâ carnosa apice ramosâ, caulis elatis profunde sulcatis volubilibus simplicibus, foliis alternis longe petiolatis membranaceis ovatis acuminatis profunde cordatis integris, racemis ex omnibus axillis superioribus solitariis vel 2-3

folio brevioribus laxis ramulosis, floribus basi minute tribracteolatis subternis dichotomiali sessili cæteris pedicellatis, perigonii rotati laciniis linearis-oblongis obtusis concais trinerviis ♀. Rchb. Icon. bot. tab. 754. — Eichw. Casp. tab. 23.

Hab. in umbrosis faucium regionis inferioris Caucasi septentrionalis ad Pätilorsk, Wladikaukas (M.B., Stev! C.A. M!), ad Lars (Rehm!), in Georgia Caucasica (Wilms.), Colchide et Imeretiâ (Rupr!).

Facies omnino *Tamn communis*.

AMARANTUS (L. Gen. p. 490).

Flores polygamo-monoici tribracteati. Perigonum 5 vel triphyllum phyllis æqualibus erectis. Stamina totidem ac sepala, libera, filamentis subulatis, antheris bilocularibus. Staminodia nulla. Stylus nullus. Stigmata 2-3 subulato-filiformia. Utriculus ovatus apice attenuatus 2-3-rostris transverse circumscissus monospermus pericarpio membranaceo. Semen verticale lenticulari-reniforme crustaceum arillo nullo. Albumen centrale, embryo cyclicus radiculâ inferâ. — Herbæ annuæ, foliis alternis, floribus in spicas paniculatas vel glomerulos axillares dispositis.

* Pentandri, sepala quina.

1. **A. caudatus** (L. Sp. p. 1406) annuus glabrescens vel adpresso puberulus, caule erecto elato angulato, foliis longe petiolatis ovatis vel oblongo-lanceolatis utrinque attenuatis læte viridibus, paniculæ subsimplicis vel ramosæ spicis cylindricis pendulis terminali longissimâ, floribus arcte glomeratis coccineis vel rarius albidis, perigonio 5-sepalo bracteis subbreviore, utriculis calyce vix longioribus rugosiusculis apice 2-3-lobis ♂.

Hab. in ruderatis, Mesopotamia (Botta), Ægyptus inferior (Gaill!), in Affghanâ et Belutschia (ex Aith. Cat.). An vere spontaneus?

Ar. Geogr. Nubia, Abyssinia, India.

2. **A. hypochondriacus** (L. Sp. p. 1407) annuus glaber vel sebriuscus, caule erecto sulcato, foliis longe petiolatis oblongo-lanceolatis cuneatis acuminatis, paniculæ ramosæ spicis erectis cylindricis obtusis terminati longiore, floribus arcte glomeratis saturate purpureis, perigonio quinquesepalo bracteis longe aristatis breviore, utriculis calycem superantibus rugosiusculis apice 2-3-fidis ♂.

Hab. subspontaneus (ex Americâ introductus) in cultis et vineis, Armenia ad Karput et Syria borealis ad Marasch (Haussk!), Syria australis (Dr. Post!).

Ar. Geogr. America borealis.

3. **A. chlorostachys** (Willd. Amar. p. 34, tab. 10, fig. 19) annuus pallide virens minute et crispule pubescens, caule erecto

striato-sulcato, foliis petiolatis rhomboe-ovatis obtusiusculis, panicula strictæ ramosæ laxissimæ spicis cylindricis ascendentibus laxiusculis acutis terminali longissimo, bracteis lanceolatis longe subulato spinosis flore subdupo longioribus, floribus virentibus, perigonii sepalis quinis linear-i-oblongis retusis vel obtusis, utriculis calyce longioribus retusis apice 2-3-dentatis ⊖.

Hab. in ruderatis Pelopennesi prope Klakines (Orph!), Syrie prope Saida (Gaill!). Ex Americâ introductus.

Ar. Geogr. America borealis et tropica.

4. **A. patulus** (Bertol. Comum Neap. p. 171, tab. 12) annus intense viridis plus minus pubescens, caule erecto simplici vel ramoso, ramis inferioribus ascendentibus, foliis longiusculæ petiolatis deltoideo-ovatis obtusis, paniculae inferne foliatæ et laxæ superne compactæ spicis cylindricis ascendentibus acutis terminali multo longiore, bracteis lanceolato-spinulosis flore tertia parte longioribus, floribus viridibus, perigonii laciniis quinis ellipticis obtusis muticis vel mucronulatis, utriculis calyce longioribus retusis apice 2-3-dentatis ⊖. *A. Timeroyi* Jord.

Hab. in cultis, Egyptus circa Caniram (Gaill. exs. 23!) et prob. alibi.

Valde affinis *A. chlorostachydi* cuius forsitan varietas, differt colore intense viridi et bracteis brevius spinulosis flores parum superantibus.

Ar. Geogr. Ex Americâ ortus nunc in Hispaniâ! Italiâ, Gallia media et australi, Africâ boreali, in quilibus.

5. **A. retroflexus** (L. Sp. p. 1407) annus pallide virens minute pubescens, caule crasso erecto, foliis longe petiolatis ovato-rhombeis obtusiusculis, panicula terminalis compactæ spicis viridibus brevibus sessilibus inferioribus sœpe subremotis obtusis terminali vix longiore, bracteis lanceolatis longe subulato-spinulosis flore duplo longioribus, perigonii sepalis quinis linear-i-oblongis retusis mucronatis, utriculis calyce sublongioribus rugosis apice 2-3-dentatis ⊖. Ic. Willd. tab. 11.

Hab. (ex Americâ oriundus) nunc in cultis et ruderatis totius regionis a Graeciâ! et Macedoniâ! ad Caucasum! et Transcaucasicum! Syriam!

Ar. Geogr. America borealis et tropica, Europa media et australis, Africa borealis.

6. **A. paniculatus** (L. Sp. 1406) annus scabrinuscus, caule erecto sulcato, foliis longe petiolatis oblongo-ellipticis utrumque attenuatis, panicula ramosæ spicis erectis cylindricis acutiusculis terminali longiore, floribus glomeratis purpureis rarius pallidis, perigonio quinquesepalo bracteis aristatis breviore, utrantic calycem superantibus subrugosis apice 2-3-dentatis ⊖. Ic. Willd. Amar. tab. 2, fig. 4.

Hab. in cultis Palestine maritima (Gaill!), Babylonie ad Bagdad et Bassora (Noë!). Ut praecedens ex Americâ introductus.

Ar. Geogr. America borealis et tropica.

7. A. Gangeticus (L. Sp. p. 1403) annuus glabriusculus, caule erecto sulcato-striato, foliis longe petiolatis ovato-vel oblongo-rhombeis utrinque attenuatis obtusis, glomerulis axillaribus sessilibus subglobosis paniculam saepius ramosam laxam formantibus dissitis superioribus in spicam terminalem simplicem cylindricam approximatis, floribus densis virentibus, bracteis lanceolatis spinuloso-attenuatis perigonio subduplo longioribus, phyllis subquinis oblongo-lanceolatis spinuloso-attenuatis, utriculo rugoso apice longe 2-3-dentato calyce aequilongo ⊖. Ic. Willd. Amar. tab. 6.

Hab. prope Thebas Egypti (Auch. 2722!), ad Choubra prope Cahiram (Schw!).

In speciminibus a me observatis perigonium semper quinquesepalum nec trisepalum video.

Ar. Geogr. India Orientalis.

“ Triandri, sepala terna.

8. A. sylvestris (Desf. Cat. Hort. Par. 41) annuus glabriusculus, caule erecto profunde sulcato saepius ramoso ramis ascendentibus, foliis longe petiolatis rhombeo-ovatis oblongisque cuneatis obtusis, glomerulis paucifloris omnibus axillaribus sessilibus remotiusculis vel approximatis foliis longe superatis, bracteis lanceolatis acutis non pungentibus flori aequilongis, perigonii phyllis ternis linearibus carinatis mucronatis, utriculo rugoso apice 2-3 dentato calyce duplo longiore ⊖. *A. Blitum* Moq. in [p.c.] Prodr. — L. Sp. 1405 tantum ex parte. — *A. viridis* L. Sp. 1405 ex parte.

Hab. in ruderatis totius ditionis a Graeciâ! et Macedoniâ! ad Caucasum et Transcaucasiam! Syriam! Arabiam petræam! Egyptum! Persiam! Affghaniam (Griff!).

9. Græcizans. — Folia angustiora lanceolata. — *A. Græcizans* L. Sp. 1415. — *A. angustifolius* M.B.

Hab. in Cycladibus (Sart!), Egypto superiore (Boiss!), Babyloniâ (Noë!), Persia australi (Haussk!), Belutschîâ (Stocks!).

Ar. Geogr. Europa media et australis a Bataviâ, Belgio, Germaniâ media ad Rossiam medianam, Africa borealis, India.

9. A. albus (L. Sp. 1404) annuus glabriusculus, caule crassiusculo erecto demum scopario ramosissimo cum ramis tenuiter virgatis erecto-patulis rigido albido, foliis parvis longe petiolatis obovatis obtusis basi longe attenuatis superioribus et rameis lanceolato-oblongis aristulatis, glomerulis axillaribus sessilibus paucifloris saepè geminatis remotiusculis. bracteis subulato-attenuatis demum subpungentibus flores et utriculos superantibus, perigonii laciniis ternis oblongo-lanceolatis aristulatis, utriculo rugosissimo sepalis superato ⊖. Ic. Willd. Amar. tab. I.

Hab. (ex America oriundus) nunc in cultis et ruderatis introductus, Cephelonâ et Attica (Heldr!), Peloponnesus (herb. Fauché!).

Ar. Geogr. Ex Americâ boreali ad Lusitaniam, Hispaniam, Italianam, Africam borealem.

ALBERSIA (Kunth. Fl. Berol. II, p. 144. — *Euxolus* Rafin.).

Flores monoici tribracteati. Perigonum phyllis tribus æqualibus erectis. Stamina tria libera filamentis subulatis, antheris bilocularibus. Staminodia nulla, stylus nullus, stigmata tria filiformia. Utriculus membranaceus vel carnosus ovatus indehiscens monospermus. Semen verticale lenticulari-reniforme crustaceum, arillo nullo. Albumen centrale, embryo cyclicus, radicula infera. — Herbæ foliis alternis, floribus glomeratis vel spicatis, facie omnino *Amaranti*.

1. **A. polygama** (L. Annen 4, p. 294 sub *Amaranto*) annua glabriuscula, caule erecto angulato-sulcato ramuloso, foliis petiolatis ovato aut oblongo-rhombeis vel lanceolatis utrinque attenuatis sub-acuminatis, glomerulis omnibus axillaribus sessilibus geminatis vel solitariis remotiusculis vel approximatis foliis longe superatis, bracteis oblongis acutis flores vis æquantibus, phyllis lanceolatis acutis, utriculo ovato rugoso perigonum subæquante apice 2-3-dentato ⊖. Ic. Wight tab. 714. — *Euxolus polygamus* Moq. in D.C. Prodr. XIII, 272

Hab. in Egypto (Acerbi et Galopin ex Moq.).

Praeter characteres genericos facie *A. sylvestri* simillima; adest ut in hac specie forma foliis angustis.

Ar. Geogr. Nubia, Abyssinia, India.

2. **A. oleracea** (L. Sp. 1403 sub *Amaranto*) annua glabra, caule erecto vel ascendentem striato, foliis longe petiolatis ovatis obtusissimis vel retusis, glomerulis axillaribus sessilibus inferioribus remotis superioribus in spicam terminalem erectam brevem crassiusculam approximatis, floribus virentibus arcte approximatis, bracteis lanceolatis obtusis flores non æquantibus, phyllis albidis obtusiusculis, utriculo ovato laeviusculo acuto ⊖. *Euxolus oleraceus* Moq. loc. cit. p. 273.

Hab. in Egypto (ex Moq.). Editione non vidi.

Ar. Geogr. India Orientalis.

3. **A. Blitum** (Kunth Fl. Berol. II, p. 144) annua glabra, caulis diffusis vel ascendentibus, foliis viridibus longe petiolatis rhombo-ovatis vel oblongis obtusis vel retusis, glomerulis densifloris axillaribus inferioribus dissitis superioribus plerumque in spicam elongatam terminalem cylindricam subflexuosam approximatis viridibus, bracteis triangularibus acutis flores non æquantibus, phyllis anguste lanceolatis acutiusculis utriculo globoso acutiusculo calyce longiore ⊖. *Euxolus viridis* Moq. loc. cit. p. 273. — *Amarantus viridis* et *Am. Blitum* L. ex parte.

Hab. in ruderatis, prob. in totâ ditione, Macedonia (ex Griseb.), Transcaucasia (C.A. Mey.), Syria littoralis (Bl!), Egyptus ad Alexandriam (Gaill!), Persia Orientalis (Bge!).

Ar. Geogr. Europa omnis media et australis, Africa borealis, Abyssinia, India.

4. **A. caudata** (Jacq. Col!, II, p. 235. — Ic. rar. tab. 344 sub *Chenopodium*) annua glabriuscula, caulis erectis gracilibus, foliis longe petiolatis ovato vel oblongo-rhomboideis obtusis subtus albo-reticulatis, florum glomerulis præter infima pauca in paniculam terminalem spicis gracilibus subflexuosis constantem dispositis, spicâ terminali longiore, floribus viridibus laxiusculis, bracteis triangularris brevissimis, phyllis spathulato-linearibus acutiusculis membranaceis, utriculo subgloboso rrgosissimo calyce subdupo longiore ♂. *Euxolus caudatus* Moq. loc. cit. — *Amarantus gracilis* Desf. Tabl. p. 43.

Hab. in Belutschia (Stocks!).

Ar. Geogr. America tropica unde forsitan oriunda?, Insulae Canarienses, India.

5. **A. deflexa** (L. Mant. p. 295 sub *Amaranto*) perennis, caulis diffusis apice ascendentibus superne crispule puberulis, foliis petiolatis subtus ad nervos crispule puberulis oblongo-rhombeis vel lanceolatis acutiusculis, glomerulis inferioribus axillaribus cæteris in spicas erectiusculas cylindricas terminales sœpe subpaniculatas approximatis, bracteis ovato-triangularibus acutis flore brevioribus, phyllis lanceolatis mucronatis, utriculo ellipsoideo laevi acutiusculo calyce longiore ♀. *Amarantus prostratus* Balb. Misc. p. 44, tab. 10.

Hab. in ruderatis partis magis occidentalis ditionis, Græcia in insulâ Cephaliâ (Heldr!), insula Seiathos Cycladum (Orph!), Macedonia (Friv.).

Ar. Geogr. Europa australis a Gallia mediâ et Helvetia ad littorale Austria-cum et regionem Danubialem, Africa borealis.

ÆRVA (Forsk. Eg. Arab. p. 170).

Flores hermaphroditæ tribracteati. Perigonum subæqualiter pentaphyllum lanatum. Stamina quinque basi in cupulam coalita, filamenta subulata, antheræ biloculares. Staminodia interjecta dentiformia vel subulato-triangularia. Ovarium uniovulatum, stylus brevis, stigmata bina minuta. Utriculus subsphæricus indehiscent calyce inclusus. Semen verticale lenticulare testâ crustaceâ, arillo nullo, albumen centrale, embryo annularis radiculâ ascendente. — Herbæ vel suffrutescences albo-tomentosi foliis alternis.

1. **Æ. Javanica** (Juss. Ann. Mus. II, p. 131) perennis albo-tomentosa basi suffruticosa, caulis erectis alatis herbaceis teretiusculis stricte ramosis, foliis obovato vel oblongo-lanceolatis obtusis

mucronulatis basi longe attenuatis subpetiolatis, spicis cylindricis lanatis sessilibus in paniculam terminalem aphyllam saepe compositam dispositis terminali longiore, floribus nitidis, bracteis ovatis perigonio subbrevioribus mucronulatis exteriore pilosula, phyllis uninerviis, utriculo oblongo acutiusculo ♀. *Iresina Javanica* et *I. Persica* Burm. Ind. tab. 65. — *Celosia lanata* L. Sp. 298. — *Æ. tomentosa* Forsk. loc. cit. — *Achyranthes alopecuroides* Lam.

Hab. in lapidosis et desertis, Egyptus superior (Dell Boiss! Sieb!), in Oasi magna (Schw!) et Oasi Dachel (Asch!), Arabia petrea (Schiimp. 243!), Baby-
lonia (Auch. 3193! Noël), Persia australis (Gare.).

β. Borei Webb. Spic. Gorg. — Folia anguste lanceolata vel linearia
sæpe revoluta et interdum deflexa. — *Æ. Borei* Edgew.

Hab. in desertis Ægyptiaco-Arabicis prope Syenem (Kral!), et in littore
Ægyptiaco prope Kosseir (Schweinf!), in vallis Sinaiticis (Boiss!) et Omadi
Hebrae Petreæ (Bové! Schimp. 144!), Affghanâ (Griff!), Belutschia
(Stocks!).

Ar. Geogr. Africa borealis, Senegalalia, Insulae capitis viridis, Nubia, Abyssinia, Arabia, India.

2. *Æ. lanata* (L. Sp. p. 296 sub *Achyranthes*) annua vel perennans et inferne indurata tenuiter cinereo-tomentella, caulis herbaeis teretibus tenuibus patule ramosis, foliis obovatis obtusis in petiolum brevem attenuatis, spicis lanatis sessilibus omnibus dissitis ovatis vel breviter cylindricis inferioribus ad foliorum axillas superioribus secus ramos longe spicatis foliis minutis suffultis, bracteis triangulari-ovatis apiculatis extus pilosiusculis perigonio brevioribus, phyllis uninerviis, utriculo subgloboso ♀. *Æ. lanata* Juss. Ann. Mus. II, p. 131. — *Achyranthes villosa* Forsk. Eg. Ar. p. 48.

Hab. in Egypto (ex Forsk.), Persia ad sinum Persicum (Belanger).

Ar. Geogr. Nubia, Abyssinia, Arabia, India, Africa australis.

ACHYRANTHES (L. Gen. p. 443).

Flores hermaphroditici tribraeati. Perigonium 5 rarius 4-phylum glabrum phyllis subinæqualibus demum induratis. Stamina 5-4 basi in cupulam coalita. Staminodia quinque interjecta integra vel siue apice denticulata dorso interdum finibriato-appendiculata. Ovarium uniovulatum. Stylus subelongatus, stigma simplex capitatum. Utriculus ovatus vel subglobosus indehiscens calyce inclusus. Semen verticale suberustaceum, arillo nullo. Albumen parcum centrale, embryo annularis peripheriens, radicula ascendens — Herbae vel suffrutices, foliis et ramis oppositis bisecte articulatis, floribus spicatis.

1. *A. aspera* (L. Sp. p. 295. — Aschers. in Schw. Beitr. p. 172) fruticosa adpresso pubescens ramis elongatis ramosis ramulis virgatis erecto-patulis, foliis breviter petiolatis ovatis vel oblongis acutis vel

acuminatis superioribus parvis lanceolatis, spicis ex axillis superiores terminalibusque longis gracilibus demum laxis, floribus virentibus patulis demum deflexis, bractearum lateralium aristâ limbû suum ovatum æquante, phyllis bracteis subduplo longioribus anguste lanceolatis acutissimis obsolete trinerviis, utriculo oblongo ♂.
A. fruticosa Hort. Par.

Hab. in Egypto (Forsk., Sieb.), Belutschia (Stocks!).

β. argentea. — Flores subbreviores albidi vel purpurascentes, bractearum lateralium arista laminâ suâ subduplo longior. — *A. aspera* var. *Sicula* L. — *A. argentea* Lam. Dict. I, p. 545. — Ic. Fl. Græc. tab. 244.

Hab. in rupestribus, dumosis siccis, insula Rhodus (Ehr!), Syria littoralis inter Saida et Damur (Gaill!), Egyptus prope Cahiram (Ehr!), regnum Maccat Arabiae (Auch. 5254!).

Recte animadvertente cl. Ascherson transitus a typo ad varietatem manifesti sunt.

Ar. Geogr. Insulæ Canarienses, Azoricæ, Madera, Senegal, Africa borealis, Hispania australis, Sardinia, Sicilia, Calabria, Nubia et Abyssinia, Arabia, India, P. B. Spei.

DIGERA (Forsk. Fl. Eg. Arab. p. 65).

Flores hermaphroditi tribracteati ternati, intermedius fertilis laterales steriles in cristam ramulosam mutati. Perigonium glabrum phyllis 5 subinaequalibus erectis. Stamina 5 libera subulata, antheræ biloculares. Staminodia nulla. Ovarium uniovulatum. Stylus filiformis, stigmata bina recurva brevia. Caryopsis dura subglobosa calyce inclusa indehiscens. Semen verticale lenticulare testâ tenui membranaceâ. Albumen centrale, embryo annularis periphericus, radicula subascendens. Herbæ foliis alternis, floribus spicatis.

1. *D. alternifolia* (L. Mant. 50 sub *Achyranthes*) annua viridis glabra, caule erecto vel ascendentे flexuoso patule ramoso, foliis longe petiolatis ovatis vel oblongis acuminatis obtusis, spicis axillaribus gracilibus laxis pedunculatis folio longioribus, floribus viridibus, cristis (floribus lateralibus sterilibus) flabellatis 4-6-fidis, phyllis membranaceis oblongis acutiusculis ♂. *Digera alternifolia* Asch. in Schw. Beitr. p. 180. — *D. arvensis* β. *annua* Hochst. et St.

Hab. in Affghaniâ (Griff!).

D. arvensis α. *perennans* Moq. in D.C. Prodr. (Nubia Ky. 128) differre videatur foliis anguste lanceolatis, floribus albis.

Ar. Geogr. Arabia, Nubia, Abyssinia, India.

PUPALIA (Juss. Ann. Mus. 1803, p. 132. — *Desmochæta* D.C. Cat. Monsp. 1813, p. 101).

Flores hermaphroditi bibracteati ternati omnes fertiles, lateritium bracteæ in axillâ fasciculum aristarum uncinatarum vel acutatarum (flores abortivi?) ferentes. Perigonum phyllis quinis subæquilibus. Stamina quina basi in cupulam brevissimam connata, filamenta subulata, antheræ biloculares. Staminodia nulla. Ovarium uniovulatum, stylus filiformis, stigma capitatum. Utriculus oblongus calyce inclusus. Semen reniforme ellipticum crustaceum. Albumen centrale. Embryo annularis periphericus, radicula ascendens. — Suffrutices oppositifolii florum glomerulis interrupte spicatis.

1. **P. lappacea** (L. Sp. p. 204 sub *Achyranthes*) basi suffruticosa breviter tomentoso-flavescens, ramis teretibus virgatis elatis, foliis breviter petiolatis ovatis inferioribus obtusis cæteris acuminatis, spicis ex axillis superioribus oppositis terminalique pedunculatis longissimis, bracteis late scariosis triangulari-ovatis mucronatis villosis eis florum lateralium in axillâ fasciculum ramosum pilorum glochidiatorum demum auctorum divergentium edentibus. phyllis lanceolatis acuminatis bracteas superantibus, glochidibus florum lateralium paucis flavidis, utriculo acutiusculo 5. *P. lappacea* Moq. loc. cit. p. 331. — *Achyranthes echinata* Retz Obs. 2, p. 12. — *Desmochæta flavescens* D.C. — *D. xanthioides* A. Br. Flora 1841, p. 285, tab. 11.

Hab. in Belutschia culidiore (Stocks!).

Pili glochidiati demum elongati divaricati intricati calyces occultantes, glomeruli fructiferi decidui.

Ar. Geogr. Senegal, Nubia, Abyssinia, Arabia tropica, India.

POLYCNEMUM (L. Gen. 25).

Flores hermaphroditi bibracteati. Perigonum phyllis quinis æquilibus erectis glabris. Stamina tria (rarius 4-5) basi coalita, filamenta filiformia, antheræ biloculares. Ovarium uniovulatum. Stylus filiformis brevis, stigmata bina capillaria subpatula. Utriculus ovatus indehiscentes monospermus calyce inclusus. Semen verticale reniformi-lenticulare, embryo annularis periphæricus radiculâ ascendentem. — Herbae (nostræ) annuae ramosæ, foliis crebris alternis, floribus axillaribus sessilibus minutis.

1. **P. arvense** (L. Sp. p. 50) glabrum asperulum pallide virens collo multicaule caulis procumbentibus ramosis centrali sœpe erecto, foliis confertis strictis a basi subdilatata et membranaceâ anguste linearis-subulatis subtriquetris mucronatis subpungentibus,

floribus solitariis vel geminatis, bracteis scariosis a basi ovatâ subulatis perigonium vix æquantibus, phyllis ovato-acuminatis apice carinatis tenuissimis, utriculo calyce non longiore ⊖.

Hab. in cultis prov. Caucasicarum (M.B.). E ditione nondum vidi ubi forsitan per errorem ex confusione cum *P. majus* indicatum fuit.

Ar. Geogr. Europa media et australis a Belgio et Bataviâ ad Rossiam medium.

2. ***P. majus*** (Al. Braun in Koch Syn. II, p. 675) glabrum vel asperulum pallide virens a collo multicaule, caulis prostratis vel erectis elongatis stricte ramulosissimis, foliis confertis strictis a basi dilatata et membranacea anguste linearis-subulatis subtriquetris mucronatis subpungentibus, floribus solitariis vel binis, bracteis scariosis a basi lanceolata subulatis perigonio multo longioribus, phyllis ovato-lanceolatis acuminatis utriculo brevioribus ⊖. *P. arvense* Fl. Græc. Prodr. I, p. 23 non L.

Hab. in cultis, Tauria (herb. Fauché!), Byzantii (Sibth., Noë 2011) et prob. alibi.

Validius præcedente, omnibus partibus majus.

Ar. Geogr. Europa media in Gallia, Helvetiâ, Germaniâ, regione Danubiali.

3. ***P. verrucosum*** (Lang in Rehb. Fl. exs. p. 583) a collo multicaule, caulis crispule eanescentibus et verrucosis procumbentibus ramosis et centrali erecto, foliis confertis strictis a basi subdilatata et membranacea anguste linearis-subulatis pungentibus, floribus solitariis vel binis, bracteis scariosis a basi lanceolata setaceo-acuminatis perigonio brevioribus, phyllis ovato-lanceolatis acuminatis ⊖.

Hab. in regione alpinâ montis Masmeneudagh Cappadociæ (Ball sp. unicum), prob. alibi.

Ar. Geogr. Regio Danubialis.

TR. III. GOMPHRENEÆ Endl. — Antheræ uniloculares. Ovarium uniovulatum.

ALTERNANTHERA (Mart. Nov. Gen. II, p. 53).

Flores (in nostris) hermaphroditi tribracteati. Perigonum quinquesepalum. Stamina 5 filamentis filiformibus basi in cupulam connatis antheris unilocularibus. Staminodia interjecta denti-vel unguiformia minuta. Ovarium uniovulatum, stylus brevis, stigma capitatum vel bilobum. Utriculus ovatus indehiscens. Semen verticale sublenticulare crustaceum. Albumen centrale. Embryo annularis periphæricus, radicula ascendens. — Herbae oppositifoliae.

1. ***A. sessilis*** (L. Mant. p. 345 sub *Gomphrend*) annua? cauli-

bus inferne prostratis radicantibus dein erectis ramosis bifariam puberulis, foliis obovato-oblongis vel ovato-lanceolatis obtusis basi longe attenuatis subsessilibus, florum glomerulis ad axillas 3-4-nis sessilibus paucifloris subsphæricis, floribus albidis nitidis glabris, bracteis ovatis scariosis perigonio duplo brevioribus, phyllis scariosis glabris ovatis acuminatis, staminodiis subulatis filamenta subaequantibus, utriculo calyce longiore ⊖. *A. sessilis* R. Br. Pr. N. Holl. I, p. 417. — *A. denticulata* var. *major* Moq. in D.C. Prodr.

Hab. in uliginosis et madidis Egypti inferioris (Sieb! Ehrenb! Schweinf!), in oryzetis prov. Talysh ad Caspium prope Lenkorau (Hoht). in humidis prov. Ghilan Persie (Auch. 4521).

Sæpe confusa cum *A. nodiflora* R. Br. = *A. denticulata* R. Br. (Vide Asch. in Schweinf. Beitr. p. 175), foliis elongatis lanceolatis, floribus longioribus acutioribus, utriculo calyce breviore, etc., distincta.

Ar. Geogr. Hispania australis, Africa borealis et tropica, India, America tropica.

ORD. CL. POLYGONEÆ.

(Juss. Gen. p. 82.)

Flores hermaphroditi vel unisexuales. Perigonium inferum 3-5-6-partitum æstivatione imbricatum. Stamina definita numero varia basi perigonii inserta perigyna uniseriata vel biseriala, antheræ bilobulares. Ovarium liberum uniloculare uniovulatum ovulo erecto. Styli 2-3 liberi vel connati. Fructos indehiscentes nueamentaceus vel carnosus, nudus vel laciniis perigonii interioribus velatus. Embryo inversus (antitropus) in albumine farinoso aut carnosico, vel centralis rectiusculus, vel unilateralis curvatus. — Herbae vel frutices foliis alternis, stipulis vaginantibus.

TR. I. PTERYGOCARPÆ Meisn.

Achenii 2-4-goni anguli in alam expansi. Embryo axialis.

CALLIGONUM (L. Gen. 600. — *Pterococcus* Pall. — *Calliphysa* Fisch. et Mey.).

Flores hermaphroditi. Perigonium rotatum petaloideum non in crescens laciniis quinque plano-concavis patentibus. Stamina 10-16 basi connata. Ovarium tetragonum quadrisulcatum, styli quatuor liberi breves, stigmata capitata. Achenium vel quadrialatum, vel ad angulos præsertim setis 8-10-seriatis ramosis cancellato-intertextis echinatum.

vel undique setis intertextis in membranam achenium includentem
expansis dense obtectum. — Pericarpium sublignosum, semen quadri-
sulcatum. Embryo in axi albuminis carnosí rectus. — Frutices ramosissimi
ramis validis, ramulis hornotinis debilibus articulatis et siccis
facile secedentibus. Ochreæ membranaceæ brevissimæ bilobæ. Folia
sæpius minima et cito decidua. Flores pauci in ochrearum axillâ fasci-
culati laterales, pedicelli articulati. — Plantæ ex speciminibus siccis
nimis paucis et incompletis sæpe notæ et ideo nonnullæ species forsitan
in posterum reducendæ.

Specierum Orientalium distributio.

SECT. I. PTEROCOCCUS Bge in Pl. Lehm. — Achenium quadrialatum.

* Alæ in cristam unicam abeentes vel margine subintegræ.

C. Pallasia, Bungei, denticulatum, tetrapterum, stenopterum.

* Alæ margine bicristatae.

C. Persicum.

SECT. II. EUCALLIGONUM Bge. — Achenium setis apice liberis cancel-
lato-intertextis dense obsitum.

C. comosum, polygonoides, crinitum, eriopodium.

SECT. III. CALLIPHYSIA Bge. — Achenium setis intertextis apicibus in
membranam involventem expansis obsitum.

C. Calliphysa.

SECT. I. PTEROCOCCUS.

1. **C. Pallasia** (Lhér. Act. Soc. Linn. Lond. I, p. 180) fruticosum
glabrum, ramis flexuosis adultis fuscis, ramulis tenuissimis, foliis
linearibus basi cum ochreâ articulatis et cito deciduis, achenii eum
alis membranaceis orbiculati alis planis transverse multinerviis sim-
pliciter et minute denticulatis loculo æquilatis apice rotundatis stylo-
rum basi adnatis ȝ. *Pterococcus aphyllus* Pall. It. II, p. 332. — *Pal-
lassia Caspica* L. Suppl. p. 252. — *Calligonum polygonoides* Pall. It.
III, p. 530 non L.

Hab. in arenosis mobilibus ad mare Caspium (C.A. Mey.). Prob. ad septen-
trionem Caucasi et in ditione nostrâ nondum observatam.

Achenium cum alis orbiculatum 5-6 lineas diam. latum, denticuli lente tantum conspicui et interdum obsoleti.

Ar. Geogr. Rossia australis, Turkestania borealis, Songaria.

2. C. Bungei, fruticosum glabrum, ramis adultis valde flexuoso-infractis cinereis, novellis tenuibus glaucescentibus divaricatis, foliis . . . , achenii ad facies ovato-orbiculati alis membranaceis ternis vel quaternis membranaceis planiusculis transverse nervulosis margine obsolete repandis loculo sublato roribus apice rotundatis a styli basi liberis ȝ. *C. tripterum* Bge MSS.

Hab. in arenâ mobili Persiae orientalis inter Kerman et Yezd (Bge!).

Ramii tortuoso-infracti et facies *C. denticulati* a quo specifice differre videatur achenii alis subintegris et loculo vix latioribus; variant ternæ vel quaternæ.

3. C. denticulatum (Bge MSS.) fruticosum glabrum, ramis valde tortuosis cinereis, ramulis hornotinis valde flexuosis et divaricatis glaucescentibus, foliis . . . , achenii orbiculati alis membranaceis planiusculis vel undulatis transverse multinerviis loculo subduplo latioribus margine grosse denticulatis apice obtusis et a stylorum basi liberis ȝ.

Hab. in Persiâ orientali inter Kerman et Yezd (Bge!).

Achenia cum alis 7-8 lineas longa et lata, alæ fere tres lineas latæ. Differt a *C. Pallasiâ* et *C. Persico* alis multo latioribus.

4. C. tetrapherum (Jaub. et Sp. Ill. Or. V, tab. 471) fruticosum glabrum, ramis cinereis, ramulis hornotinis tenuibus teretibus glaucescentibus, foliis minimis brevissimis ochreæ adnatis, achenii ovati alis coriaceis loculo triplo angustioribus planis margine subdistiche spinulosis spinulis interdum 2-3 furcatis ȝ.

Hab. in Mesopotamiâ inter Bagdad et Aleppo (Oliv. in herb. Mus. Par.). Non vidi.

Ramuli novelli 4-10-pollicares, achenium cum alis 4-5 lineas longum 3-4 latum stylis persistentibus apiculatum, folia oculo nudo vix conspicua.

5. C. stenopterum (Bge MSS.) fruticosum glabrum glaucum, ramis flexuosis et tortuosis albidis, ramulis hornotinis tenuibus, foliis subulatis brevissimis, achenii oblongi alis membranaceis transverse nervulosis loculo angustioribus planis margine in spinulas subulatas subdistichas simplices abeuntibus ȝ.

Hab. in Persiâ orientali et Affghania occidentali, Kuh i Bochtan, prope Robat inter Herat et Tebbes (Bge!).

Incomplete notum ex specimine in quo achenia bina et nimis vetusta tantum vidi; alis angustis fructus accedit ad *C. tetrapherum* sed spinulis marginalibus tenuioribus simplicibus differre videtur.

6. C. Persicum (Boiss. et Buhse Aufz. p. 191 sub *Pterococco*) fruticosum glabrum, ramis adultis flexuosis albidis, ramulis horno-

tiniis tenuibus, foliis subulatis cum ochreâ articulatis cito deciduis, achenii cum alis membranaceis ovati alis planis loculo aequilatis transverse et elevatim multinerviis circumcirca acute denticulatis bicristatis eristâ exteriore breviore ȝ.

Hab. in lapidosis Persiae borealis ad fluvium Sefirud prope Mendschil et in desertis arenosis Enarek (Buhse!).

Differt a *C. Pallasii* ramis albidis, acheniorum alis bicristatis et profundius denticulatis.

SECT. II. EUALLIGONUM.

7. C. comosum (L'Her. in Trans. Soc. Linn. Lond. I, p. 180) fruticosum ramosissimum glabrum, ramis albidis tortuosis, ramulis tenuibus, foliis minutissimis breviter subulatis ochreæ adnatis, achenio ovato dense setoso setis 12-16-seriatis ramosis mollibus intertextis basi dilatata in quatuor alas breves plus minus conjunctis ȝ.

Hab. in desertis Ægyptiacis ad Kahiram et alibi (Sieb! Schimp! Boiss! etc.), in Oasi parvâ (Asch!), ad littora Ægyptico-Arabica (Schweinf!), desertis Arabie petree (Bové! Boiss!), Syriæ (Auch. 2561!), Persiae australis circa Isphahan, Kerman, Jezd (Bge!).

Frutex 10-15-pedalis, folia lineam vix excedentia.

Ar. Geogr. Africa borealis.

8. C. polygonoides (L. Sp. 478) fruticosum, glabrum, ramis albidis, ramulis tenuissimis, foliis tenuissime subulatis longiusculis ab ochreâ liberis et cito deciduis, achenio oblongo setoso setis rigidulis pluriseriatis ramosis basi dilatata in quatuor alas breves plus minus conjunctis ȝ. *Polygonoides Orientalis Ephedrae facie* Tourn. Coroll. 47. — It p. 356 Ic.

Hab. in arenosis subsalsis interioribus Armeniæ in monte Ararat (Tourn.), in Armeniâ (Auchi., Szov!), in prov. Aderbidjan Persiae inter pagos Seidkhodji et Tchorbulâ districtus Khoi (Szov!).

Ramuli eis speciei præcedentis magis elongati et flexuosi, folia longiora (3-5 lineas), fructûs setæ minus crebræ rigidiores. Ob specimenia sæpius imperfecta limites geographicæ inter hanc et *C. comosum* incomplete. Specimina nonnulla ab am. Bge in Persiâ orientali lecta ob ovaria juniora inter hanc et *C. Caputmedusæ* Schrenck incerta sunt.

9. C. crinitum (Boiss. Diagn. Ser. II, 4, p. 77) glaucum glabrum, ramis elongatis tenuibus subflexuosis striatis, ramulis tenuissimis elongatis, internodiis elongatis, foliis , acheniis angustè ellipticis apice attenuatis, setis mollibus longis octoseriatis rufis basi dilatatis et subconnatis supra medium bis vel ter dichotome ramosis ȝ.

Hab. in Belutschia (Stocks!).

Ex speciminiis imperfectis male notum, a præcedentibus diversum videatur ramis tenuioribus elongatis striatis, achenio angustiori absque setis 3-3½ lineas longo sesquialeam lato, sete tenuissimæ undique patentes punci-seriatæ 4-5 lineas longæ.

10. C. eriopodum (Bge pl. Lehm. p. 486) fruticosum ramis albidis gracilibus, ramulis hornotinis gracillimis, foliis linearis-subunitatis liberis cito deciduis, pedicellis subgeminis supra articulationem hispidulis, perigonii laciniis exterioribus extus hispidulis, ovario ovato setis densissimis octoseriatis ramosis basi liberis obsito, fructu....5.

Hab. in deserto Kisilkum Turkestaniae (Lehm.).

Species ex specimihiibus junioribus tantum nota, a præcedentibus glaberrimis parte superiore pedicelli et perigonii laciniis externis pilis brevibus crassiuseulis hirtis discrepans.

SECT. III. CALLIPHYSA.

11 C. Calliphysa (Bge Cat. Sem. Dorp. 1839, p. 7) fruticosum glabrum, ramis albidis tortuosis, ramulis tenuibus flexuosisimis, foliis basi ochreæ articulatis deciduis linearibus elongatis obtusis subuniternerviis, achenis ovatis 5. *Calliphysa juncea* Fisch. et Mey. Ind. Petrop. 1835, p. 24.

Hab. in Persia boreali inter Bedescht et Mianeh, orientali inter Kerman et Yezd (Bge!), in Turkestania ad Caspium (Karell!) et in desertis arenosis Karakum (Lehm!).

Ramuli saepe fasciculati, folia 10-12 lineas longa 1-1 1/4 lin. lata. Achenium 5-6 lineas longum membranæ fuscescente non reticulata inclusum.

PTEROHYRUM (Jaub. et Sp. III. II, p. 7).

Flores hermaphroditi. Perigonium rotatum petaloideum vix acre-scens quinquepartitum laciniis binis exterioribus demum reflexis ternis interioribus achenio adpressis. Stamina 8 perigonii fance inserta quorum tria sublongiora, filamenta basi barbulata, antheræ versatiles. Ovarium ovoideo-triquetrum, styli tres breves, stigmata capitata. Achenium coriaceum cordato-ovoideum rostratum alis tribus membranaceis transverse striatulis supra medium sinu profundo rotundato bipartitis. Semen loculum implens radiculâ longe rostratum, embryo in axi albaminis farinosi rectus, cotyledones semiteretes. Fructus alternatim ramosissimi, folia persistentia fasciculata, ochreæ membranaceæ brevissimæ truncatae vel bipartitæ. Pedicelli capillares paulo supra basin articulati singuli basi ochreolâ vaginati, fructiferi penduli.

SECT. I. STREPTOCARPA Jaub. et Sp.—Achenium ad quamvis faciem ad sinum alarum in plicam transversam hippocrepideam distortum et ideo alarum lobii superiores in eodem plano ac inferiores non siti. Semen infra rostrum transverse sinuosum.

1. P. Aucheri (Jaub. et Sp. Ill. II, tab. 107) fruticosum ramosissimum ramis ramulisque rigidis erecto-patulis cinereis, foliis fasciculatis glabris glaucescentibus tenuiter linearibus vel linearis-spathulatis margine revolutis teretiusculis, floribus in foliorum rosulis fasciculatis, alarum rosearum achenii lobis terminalibus ab inferioribus plus minus remotiusculis ♂.

Hab. in aridis et arenosis desertorum Persiae borealis, in valle Sefirud (Auch. 5269!), prope Bibersin et Enarek (Buhsel!), in ditione Khorassan inter Schahrud et Nichapur (Bge!), inter Meschhed et Herat (Bge!). Persice *Pia-rend*.

Folia 3-5 lineas longa lineam vix lata, pedicelli 2-3 lineas longi, achenium cum alis 3 lineas longum et latum.

2. P. ericoides, fruticosum ramosissimum ramis ramulisque elongatis cinerascentibus, foliis lente pruinoso-velutinis angustissime linearibus margine convolutis parte inferiore attenuatis, floribus in foliorum rosulis fasciculatis numerosis, achenii alarum lobis terminalibus ab inferioribus plus minusve remotiusculis ♀.

Hab. in Belutschia inferiore (Stocks № 419!), specimina sterilia ex Persiae australis monte Sawers et ex valle Chyrsan Elymaitidis huc quoque spectare videntur.

Facies et folia tenuissima *P. scoparii* sed velutina et achenium apice distortum. A *P. Aucheri* foliis velutino-pruinosis tenuioribus specifice distinctum videtur.

3. P. Olivieri (Jaub. et Sp. Ill. II, tab. 108) fruticosum ramosissimum, ramis ramentisque rigidis albidis erecto-patulis. foliis fasciculatis crassiusculis glaucis oblongo vel obovato-spathulatis obtusis margine subrevolutis glabris, floribus in foliorum rosulis fasciculatis, alarum rosearum achenii lobis terminalibus lobis inferioribus subcontiguis ♂. *P. Griffithii* Meisn. in D.C. Prodr. XIV, p. 31.

Hab. in Persia septentrionali circa Teheran (Oliv.), ad Schurab inter Teheran et Isfahan (Bge!), in Affghania prope Tarnuk et in jugo Bolan (Griff.).

β *gracile*. — Ramuli tenuiores elongati, folia minora elliptico-obovata, achenii aliae minores. — *P. gracile* Boiss. Diagn. Ser. I, 12, p. 102.

Hab. prope Sonak jugi Elbrusensis Persiae borealis (Ky. 664!), prope Schurab (Haussk!).

A *P. Aucheri* foliis latioribus praesertim distincta nam nota ex alarum lobis superioribus ab inferioribus remotis vel contiguis valde caduca videtur. *P. Griffithii* Meisn. achenium apice distortum nec rectum habet! ei huc spectat.

Ar. Geogr. Ditione Scinde Indiæ borealis.

4. P. Noëanum (Boiss. in Bot. Zeit 1853, p. 734 — Meisn. in D.C. Prodr. XIV, p. 31) fruticosum ramosissimum ramulis gracilibus patentibus fuscouscentibus, foliis minutis carnosulis viridibus breviter oblongo-lanceolatis obtusis margine subrevolutis, floribus creberrimis

in foliorum axillis fasciculatis, alarum rosearum angustarum lobis terminalibus et inferioribus subcontiguis ♂.

Hab. in Assyriâ prope Hanneky ad ripas fl. Dialæ et prope Mossul ad ripas Tigridis (Ky. 1002!).

Ex specimine unico mihi notum et ulterius explorandum. *Streptocaryæ* evidenter nec *Orthocaryæ* adsociandum. A precedentibus distinctum videtur foliis minoribus non glaucis, florum fasciculis secus ramulos numerosissimis, alis angustioribus.

SECT. II. ORTHOCARVA Jaub. et Sp. — Achenium non distortum alarum subpanduriformium lobus superior cum inferiore in eodem plano situs. Semen non sinuosum.

5. **P. scoparium** (Jaub. et Sp. Ill. II, tab. 109) fruticosum ramosissimum ramis ramulisque strictis albidis, foliis convoluto-filiformibus tereti-cylindricis glabris, floribus in foliorum rosulis fasciculatis, achenii alis rectis supra medium sinu profundo excisis, loculo continuo ♂.

Hab. in Arabiæ regno Mascate (Auch. 5270!).

RHEUM (L. Gen. 401).

Flores hermaphroditi. Perigonum herbaceum sexpartitum marcescens laciniis subæqualibus. Stamina novem imo calyci inserta. Ovarium liberum trigonum, styli tres breves patuli, stigmata integra discoidea. Achenium samaroideum trialato-triquetrum perigonum adpressum longe superans crustaceum. Semen erectum, embryo in albuminis farinosi axi rectus, radiculâ brevi superâ. — Herbæ rhizomatæ crasso, cœlibus annuis crassis sœpe subnullis, foliis amplis palminerviis.

1. **R. Tataricum** (L. fil. Suppl. p. 229) caule humili crasso aphylo fere a basi in paniculam ramosissimam divaricatam æquilongam ac latam abeunte, foliis glabris petiolis brevibus subcylindricis suffultis a basi angustissime cordatâ orbiculato-reniformibus transverse latioribus obtusissimis repandis minute spinuloso-denticulatis crasse trinerviis, pedunculis laxe racemosis geminis vel solitariis tenuibus fructu æquilongis supra medium articulatis, achenio ovato-elliptico basi cordato alis semine angustioribus ♀. *R. Caspicum* Pall. N. Act. Petrop. X, p. 382.

Hab. in deserto Karakum Turkestaniæ (Lehm!).

Ar. Geogr. Rossia australis, Songaria, Sibiria Altaica.

2. **R. Ribes** (Gronov. Fl. Orient. p. 130) caule elato crasso inferne folioso superne aphylo in paniculam amplam ramosissimam fastigia-

tam abeunte, foliis glaucescentibus coriaceis subcordato-orbiculatis vel reniformibus transverse latoribus margine minute denticulatis crasse subquinquencerviis nervis subtus saepius papilloso-scabris, pedicellis fasciculatis fructu subaequilongis filiformibus infra medium articulatis, achenio magno demum sanguineo subcarnoso-succulento cordato ovato ovato alis semine 2-3-plo angustioribus ♀. Ic. Dill. Eth. tab. 158. — Desf. Ann. Mus. II, tab. 94. — Pococke Orient. tab. 84. — Jaub. et Sp. Ill. tab. 470.

Hab. in regione subalpinâ et alpinâ, Palestina ad Kerek (Roth!), Libanus supra Ainet Colesyriæ (Boiss!) usque ad cacumina altiora (Bl!), Cataoniæ Berydagħ 7000' (Haussk!), Armenia prope Van (Deyrolle!) et ad Agerow-dagħ (Ky!), mons Pir Omar Gudrun Kurdistaniæ (Haussk!), Persia australis mons Kuh Daēna et ad Persepolin (Ky. 586! et 252!), prov. Aderbidjan Persia bor. (Auch. 5284!), Belutschia montes (Stocks!). Arabis et Persia Rivas audit. Caules et petioli novelli comeduntur.

OXYRIA (Hill. Veget. Syst. I, p. 24).

Flores hermaphroditici. Perigonum herbaceum quadripartitum marcescens laciniis duabus interioribus latoribus demum achenio adpresso, duabus exterioribus reflexis. Stamina sex, 4 geminata laciniis exterioribus et 2 solitaria interioribus opposita. Ovarium compressum, stigmata bina subsessilia penicillata. Achenium lenticulare membranaceum alâ latâ diaphanâ cinctum. Semen compressum erectum. Embryo in axi albuminis farinosi rectus radiculâ superâ. — Herbæ humiles teneræ.

1. **O. digyna** (L. Sp. 480 sub *Rumice*) perennis, cono pluricipiti ochreis et petiolis vetustis squamoso, caulis humilibus erectis subaphyllis in racenum subsimplicem vel paniculam stricte pauciramosam abeuntibus, foliis parvis longe petiolatis cordatis orbiculatis vel reniformibus interdum obsolete trilobis obtusissimis, racemis spiciformibus, pedicellis filiformibus versus medium articulatis fructu sublongioribus, achenio orbiculato basi cordato apice plus minus profunde inciso alis purpureis transverse venosis semini æquilateralis vel latoribus ♀. Ic. Fl. Dan. tab. 14. — *O. digyna* Campd. Mon. p. 155, tab. 3. — *O. reniformis* Hook. Pl. Scot. p. 111. Nees. Gen. Germ. Icon.

Hab. in frigidis regionis alpinæ, Olympus Bithynus (Sibth! Boiss!), mons Davrosdagħ Pisidiæ et Anemas Lycaoniæ (Heldr!), Taurus Cilicus (Ky! Bal!), Libanus (Post!), Cappadocia in monte Argæo (Tehih.), Armenia in montibus Techdagħ (Huett!) et Bingoldagh (Ky!), Cauc. occid. (C.A. M.) et orientalis (Rupr!), Persia borealis in jugo Elbrus (Ky. 4.5!), alpe Elamut (Auch. 5279!).

Specimen Persicum Aucheri a cl. Meisner erronee ad *O. elatiorem* R. Br. speciem Indicam relatum fuit.

Ar. Geogr. Montes Europæ borealis et mediae, Italiæ borealis, Corsiæ, regionis Danubialis, Sibiriae Dahuriæ, Americae arcticæ et borealis.

TR. II. APTEROCARPE Meisn.

Achenii anguli (in nostris) apteri. Embryo (in nostris) lateralis vel periphericus.

EMEX (Neck. El. II, p. 244).

Flores polygamo-monoici. *Masc.* Perigonum herbaceum 3-6-partitum laciiniis aequalibus patentibus. Stamina 4-6 sepalis alterna et per paria exteri tribus opposita. *Fem.* Perigonum herbaceum urceolato-trigonum sexdentatum dentibus tribus exterioribus demum spinescentibus subpatentibus, interioribus conniventibus. Ovarium trigonum ovulo basilari erecto, styli tres stigmatibus penicillatis. Achenium calycis tubo inclusum liberum. Semen erectum turbinato-triquetrum. Embryo circa albumen farinaceum periphericus, cotyledones foliaceæ albumen ambientes, radicula supera. — Herbae annuae dichotomae ochreis membranaceis truncatis laevis, floribus virentibus in pseudoverticillos axillares vel in racemos axillares breves aphyllos dispositis, inferioribus feminis subsessilibus, superioribus hermafroditis vel masculis pedicellis filiformibus non articulatis suffultis.

1. **E spinosus** (L. Sp. p. 481 sub *Rumice*) annuus glaber collo saepe pluricanlis caulis lateralibus saepe decombentibus, foliis petiolatis integris cordato vel truncato-ovatis aentis subundulatis, perigonii fructiferi dentibus interioribus ovato-triangularibus dorso tricostatis costis exterioribus marginantibus, exterioribus spinescentibus saepe arcuato-subrecurvis ♂. Ic. Fl. Græc. tab. 347. — *E. spinosa* Campd. Rum. p. 58, tab. 1, fig. 4. — *Rumex glaber* Forsk. Fl. Eg. Arab. p. 75.

Hab. in arenosis et campestribus regionis inferioris, Graecia et ejus insulae (Sibth., Sprn! Heldr! Urv!), Anatolia occidentalis et australis in Pamphyliâ (Bourg!), Ciliciâ (Balt!), Syria littoralis (Ehrenb!), Palestina circa Gaza (Boiss!), Egyptus inferior (Forsk., Auch! Boiss! etc.) et in Oasi Dachel (Aeh!).

Ar. Geogr. Lusitania, Hispania, Sardinia, Italia australis et Sardinia, Africa borealis.

RUMEX (L. Gen. 357).

Flores hermaphroditi vel abortu polygami vel dioici. Perigonum herbaceum sexpartitum inferum persistens partitionibus tribus interioribus post anthesin saepius auctis conniventibus achenium includentibus valvis dictis. exterioribus saepius minoribus patentibus vel reflexis rarius erectis. Stamina sex imo calyci inserta geminatim laci-

niis exterioribus opposita. Ovarium liberum ovulo basilari erecto, styli tres filiformes stigmatebus penicillatis. Achenium trigonum liberum valvis inclusum crustaceum. Semen erectum. Embryo lateralis albuminis farinosi angulo accumbens vel semiimmersus arcuatus, cotyledones foliaceæ, radicula supera. — Herbæ rarius frutices, folia alterna basi vaginantia vel ochreata, flores herbacei fasciculati vel pseudoverticillati in racemos simplices vel paniculas dispositi.

Specierum Orientalium distributio.

SECT. I. *LAPATHUM* Tourn. — Flores hermaphroditi rarius polygami. Perigonii laciniæ interiores, *valvæ*, demum auctæ coriaceo-membranaceæ nervo medio sæpe basi vel usque ad medium in *callum* incrassato. Styli liberi.

* Fruticosi.

R. Limoniastrum.

** Perennes.

+ Valvæ integræ vel subrepandæ (Conf. *R. Orientalem* β. *Græcum*).

1. Valvæ omnes ecallosæ.

R. Alpinus, aquaticus, Elbrusensis.

2. Valva anterior vel omnes calliferæ.

R. angustifolius, Patientia, Orientalis, crispus, conglomeratus, nemorosus.

+ + Valvæ serrulatae vel inciso-dentatae vel aculeatæ.

R. obtusifolius, Nepalensis, subtrilobus.

*** Monocarpici. Valvæ dentatae vel aculeatæ.

R. stenophyllum, pulcher, Raulini, Cassius, dentatus, palustris, comosus, bucephalophorus.

SECT. II. *ACETOSA* Tourn. — Flores dioici rarius hermaphrodito-polygami. Styli ovarii angulis superne adnati. Valvæ demum auctæ tenuiter membranaceæ nervo basi sæpe in squamulam deflexam incrassato, omnes integræ.

* Perennes.

+ Polygami.

R. scutatus, hastatus.

+ + Dioici.

R. Acetosa, Tmoleus, horizontalis, tuberosus.

Annui.

R. lacerus, vesicarius, roseus.

SECT. III. ACETOSELLA Meisn. — Flores dioici. Perigonii valvæ fructiferæ immutatae achenio adpresso non membranaceæ. Styli angulis ovarii adnati. Plantæ annuae.

R. Acetosella, acetoselloides.

SECT. I. LAPATHUM.

1. **R. Limoniastrum** (Jaub. et Sp. III. II, tab. 106) hermaphroditus fruticosus glaber, ramis erectis subflexuosis striatis superne stricte ramuloso-paniculatis, foliis glaucis carnosulis lanceolato-vel spathulato-oblongis obtusis subtrinerviis in petiolatum brevem attenuatis, ramulis in racemos simplices vel parce ramulosos laxiusculos basi exceptâ aphyllos abeuntibus, verticillastris 2-3-floris, floribus parvis, pedicellis tenuissimis ad medium articulatis perigonio subæquilongis, valvis membranaceis reniformibus transverse latioribus apice retuso-bilobis basi concavâ substipitatis non calliferis 5.

Hab. in monte Gebel Akadar regni Mascate Arabiæ (Auch. 5280!).

Folia 1-2-pollices longa 5-7 lineas lata, racemi 2-2½-pollicares, valvæ 2½ lineas latae. Affinis *R. nervoso* Vahl = *R. alismæfolio* Fresen. valvis orbiculatis integris basi squamulosis, etc. distincto.

2. **R. Alpinus** (L. Sp. 480) caule simplici in racemum compositum thyrsoideum subaphyllum abeunte, foliis amplis inferioribus petiolo longo superne canaliculato siccis profunde cordatis breviter ovatis obtusis, superioribus subsessilibus ovato-lanceolatis acutis, thyrsi ramis crebris strictis a basi floriferis fructiferis densis, pedicellis calyce longioribus quartâ parte inferiori articulatis, valvis cordato-ovatis integris vel subrepandis obtuse acuminatis tenuiter reticulatis omnibus ecallosis ♀. Ic. Moris. Sect. 5, tab. 27, fig. 2.

Hab. in pratis pinguisibus alpinis, Macedonia in montibus Nidge et Peristeri (Griseb.), Olympus Bithynus (Sibth., Boiss!), Armenia Turcica ad Mergasauk prov. Musch 8000! (Ky. Suppl. 582!), Pontus Lazicus supra Djimil (Bal!), Caucasus orientalis ad Didigwerdi Tuschetiae (Ruprl.).

β. subcalligerus. — Valvæ exterioris nervus medius basi obsolete callosus. — *R. Alpinus* M.B. Taur. Cauc. ex Ledeb. — *R. confertus* Willd. Enum. I, p. 397.

Hab. in Tauriâ (ex M.B.), in alpibus Tyfandagh et Kobi (Steven ex Ledeb.).

Ar. Geogr. Alpes et Pyrenæi, montes elatiores Hispaniæ borealis, Italiae Germaniae, regionis Danubialis. Varietas in Rossia media! Songariâ!

3. **R. aquaticus** (L. Sp. p. 479) caule elato simplici in paniculam strictam subaphyllam abeunte, foliis amplis petiolo canaliculato suffultis a basi latâ subcordatâ ovatis vel oblongis acutiusculis, superioribus oblongis vel lanceolatis, paniculæ ramis elongatis fructiferis densifloris, pedicellis calyce longioribus prope basin articulatis, valvis subcordato-ovatis obtusis integris vel subrepandis obtusis omnibus ecallosis ♀. Rehb. Ic. Crit. t. 369. — Fl. Dan. tab. 2388. — *R. Hippolapathum* Fries. — *R. Heleolapathum* Drejer.

Hab. ad aquas, Thracia ad fossas prope Enos (Griseb.), in palustribus agri Argolici et insulae Cypri (ex Sibth.). Editione nondum vidi.

Ar. Geogr. Scandinavia, Gallia borealis, Germania, regio Danubialis, Rossia, Siberia.

4. **R. Elbrusensis** (Boiss. Diagn. Ser. I, 12, p. 100) caule elato simplici in paniculam elongatam striete ramosam subaphyllam abeunti, foliis inferioribus petiolo planiusculo suffultis a basi inaequali concreata vel rotundata oblongo-lanceolatis acutiusculis tenuiter membranaceis, superioribus lanceolatis, paniculæ ramis elongatis laxifloris, pedicellis flore longioribus supra basin articulatis, valvis amplis a basi truncata orbiculatis tenuiter reticulatis circum circa obsolete erosio-denticalatis omnibus ecallosis ♀.

Hab. in jugo Elbrus Persiae borealis prope pagum Passgala (Ky. 294!).

Planta 2-3-pedalis, foliorum radicalium lamina 8-10-pollicaris, panicula pedalis, valvae fructiferæ 4 lineas diam. late. A *R. domesticus* (in Caucaso ut videtur erronee indicato) differt foliis tenuibus non undulatis, valvalis majoribus basi non cordatis.

5. **R. angustifolius** (Campd. Rum. p. 73, tab. 1, fig. 2) caule humili simplici in racemum simplicem vel basi ramulis 1-2 auctum elongatum laxum abeunte, foliis mediocribus inferioribus petiolatis lanceolatis acutis basi attenuatis vel rotundatis interdum oblongis, verticillastris discretis 5-8-floris inferioribus interdum folio linearilanceolato suffultis, pedicellis versus medium articulatis recurvis fructu brevioribus, valvis inaequalibus interioribus binis minoribus ovatis exteriore cordato-orbiculari omnibus integris erosulis elevatim reticulatis, nervo medio valvae majoris in callum plus minus manifeste incrassato ♀. *Lapathum angustifolium Cappadocicum magno fructu* Tourn. Cor. p. 38.

Hab. in pascuis alpinis Ponti Lazici supra Djimil 8400' (Bal!), supra Bai-but (Bourg!), in prov. Karabagh monte Arekligeduk (Szov!), in monte Zagros Kurdistanæ Persicæ (Haussk!).

β. macranthus. — Valvae minus inaequales, majores magis cordatae transverse latiores, exterior sæpe 7-9 lineas lata semper ecallosa. — *R. macranthus* Boiss. in Ky. exs. Diagn. Ser. II, 4, p. 80.

Hab. in pascuis alpinis Tauri Cilicici (Ky. 69! Bal!), in ditione Kassan Oghla Cilicie orientalis (Ky. 57!), in monte Berydaghi Cataoniae 7000' (Haussk!).

Species valvis fructiferis magnis inaequalibus inclita $\frac{1}{2}$ - $\frac{1}{2}$ -pedalis rarius sesquipedalis, folia lanceolata vel oblonga $1\frac{1}{2}$ - $2\frac{1}{2}$ -pollicaria. *R. macranthus* a cl. Meissner sectioni *Acetosæ* erronee adnumeratus formis super observatis ad *R. angustifolium* transit.

6. R. Patientia (L. Sp. I, p. 476) caule elato crasso in paniculam ramosam strictam elongatam basi excepta aphyllam abeunte, foliis amplis radicalibus petiolo canaliculato suffultis a basi attenuata vel rotundata oblongis vel lanceolatis acuminatis obtusis planis vel subundulatis, superioribus lanceolatis utrinque acutis, verticillastris approximatis, pedicellis perigonio aequilongis paulo supra basin articulatis, valvulis majusculis cordato-orbiculatis reticulatis margine repandis externa calliferæ φ , i.e. Schkuhr Handb. tab. 100. — *R. Olym-*
picus Boiss. Diagn. Ser. I, 5, p. 45.

Hab. in pratis subalpinis, mons Peristeri Macedoniae 4000' (Grisch.), Olym-

pus Bitinicus ad rivulos vallis Kirkbounar (Boiss!), Pontus Lazicus prope Djimil (Balt!).

β . Kurdiæ. — Folia anguste lanceolata pedalia basi longe atte-

nata, valvae callus multo major. Facies *R. Elbrusensis*.

Hab. in Kurdistanâ Persicâ ad Derbent i. Basian et in montibus Avromâ et Schâ'ru 5000' (Haussk!).

Ar. Geogr. Hispania, Sicilia, Austria, Dalmatia, regio Danubialis, Rossia sed an ubique sponte?

7. R. Orientalis (Bernh. in Schult. fil. Syst. VII, p. 1433) caule elato crasso in paniculam ramosam subaphyllum abeunte, foliis amplis radicalibus pedunculatis a basi plus minusve subcordata elongato-oblongis, superioribus oblongis et lanceolatis basi sessili cuneatis, racemis sessilibus densis, verticillastris subindistinetis, pedicellis perigonio subaequilongis infra medium articulatis, valvulis majusculis cordato-orbiculatis obtusis tenuiter reticulatis subintegris vel obsolete denticulatis unico crasse callifero φ .

Hab. in ruderatis Lydiee circa Ephesum et in pratis montis Tmol ad Boz-
dagh (Boiss!), Cariâ (Pinard!), valle Gorumse Cilicie Orientalis (Ky. Suppl.
435!), Cypro ad monasterium Troodina (Ky. 795!), Syria boreali ad Mæsænæ (Haussk!), Libano prope Eden (Bl!).

β . Græcæ. — Valvae profundius dentatae. — *R. Græcus* Boiss. et Heldr. Diagn. Ser. II, 4, p. 80.

Hab. in Græcia, Taygeti regio inferior prope Anavryti, Achæa prope Tri-
kala 3500' (Heldr.).

Hæc species fructu et facie *R. Patientia* simillima ab eo foliis basi plus
minus et anguste cordatis tantum differt et prob. hujus varietas.

8. R. crispus (L. Sp. I, p. 476) caule elato sulcato fere a basi
ramoso in paniculam longam angustam abeunte, foliis undulato-crispis
radicalibus longe petiolatis a basi attenuata lanceolatis acutiusculis
vel obtusis, superioribus et floralibus lanceolato-linearibus acutis,
verticillastris approximatis, pedicellis infra medium articulatis, valvulis

fructiferis ovato-orbiculatis subcordatis reticulatis integris vel basi subdentatis exteriore vel omnibus calliferis ♀. Ic. Fl. Dan. tab. 1334. — Engl. bot. tab. 1990.

Hab. in pratis pinguibus Græciae in Peloponneso (herb. Fauché!), Attica prope Caloreuma (Orph. 2568!), Macedonia (Frieder.), Anatolia boreali (Tchih!), monte Tmolo Lydiæ prope Bozdagh (Boiss!), Tauriæ et Transcaucasiæ (Ledeb.), Libano (Ehr., Ky. 957!).

β. elongatus. — Folia angustiora vix undulata valvæ angustiores ovatæ vel oblongæ. — *R. elongatus* Guss. pl. rar. p. 150 tab. 28. — *R. Turcicus* Boiss. Diagn. Ser. II, 4, p. 79.

Hab. in pratis alpinis Bozdagh Lydiæ cum typo (Bal!).

Ar. Geogr. Europa tota præsertim borealis et media, Sibiria, Africa borealis, Madera, America.

9. *R. conglomeratus* (Murr. Prodr. Gött. p. 52) caule sulcato elato paulo supra basi paniculato paniculæ foliosæ ramis gracilibus patentibus vel divaricatis, foliis integris vel repandis inferioribus petiolatis oblongis vel oblongo-lanceolatis basi rotundatis vel subcordatis, superioribus et rameis sessilibus anguste lanceolatis et linearibus acuminatis, verticillastris remotiusculis, pedicellis perigonio subæquilongis infra medium articulatis, valvis parvis linearis-oblongis obtusis integerrimis omnibus grosse calliferis ♀. *R. glomeratus* Schreb.— Rehb. Ic. Crit. t. 363.— *R. Nemolapathum* M.B. Taur. Cauc. non Wallr.

Hab. in pratis humidis præsertim regionis montanæ, Græcia in Laconia (Sart! Orph!), Zacynthus (Marg!), Macedonia et Thracia (Griseb., Orph!), Anatolia borealis!, Armenia (Tchih!), Cilicia (Péron!), Syria borealis in monte Amano (Ky!) et Cassio (Boiss!), Libanus et Antilibanus (Bl! Gailli!), Caucasus et Transcaucasia!

Ar. Geogr. Europa tota media et australis, Africa borealis, insulæ Canarienses, America borealis.

10. *R. nemorosus* (Schrad. ex Willd. En. I, p. 397) caule sulcato gracili elato superne in paniculam abeunte, paniculæ ramis gracilibus erecto-patulis aphyllis basi vel basi tantum foliosis, foliis integris inferioribus petiolatis cordato-oblongis superioribus lanceolato-linearibus, verticillastris distantibus, pedicellis prope basin articulatis perigonio brevioribus, valvis linearis-oblongis obtusis integris unicâ calliferâ ♀. *R. Nemolapathum* Wallr. Sched. Rehb. Ic. Crit. tab. 367. — *R. sanguineus* β. *viridis* Smith. — *R. Daghestanicus* C. Koch Linn. XXII, p. 209. — *R. condylodes* M.B. Taur. Cauc. I, p. 288.

Hab. in humidis Caucasi Orientalis circa Schamachi (M.B.), Georgiae Caucasiæ (Hoh.), prov. Talych (G.A. Mey.), Persiaæ borealis in valle Sefirud (Buhse!). Probab. etiam alibi.

Ar. Geogr. Europa præsertim borealis et media, America borealis et australis.

11. **R. obtusifolius** (L. Sp. I, p. 578) caule erecto striato cito in paniculam ramosam strictam parte inferiori exceptâ aphyllam abeunte, foliis obsolete repando-lobatis inferioribus petiolatis cordato-ovalis vel oblongis oblusis, superioribus utrinque attenuatis acuminatis lanceolatis, verticillastris remotiusculis, pedicellis elongatis prope basin articulatis, valvis ovato-triangularibus inferne dentatis in apicem oblongum obtusum integrum abeuntibus omnibus calliferis, dentibus acutis triangularibus vel subulatis ♀. Ic. Fl. Dan. tab. 1834. — Rehb. Ic. Crit. t. 366. — *R. Friesii* Gren. et Godr. Fl. Fr. III, p. 36. — *R. divaricatus* Fries Mant. — *R. Syriacus* Meisn. loc. cit. p. 53! — *R. dictyocarpus* Boiss. et Buhse Aufz. p. 192!

Hab. in ruderatis et pratis, Peloponnesus (Sibth.), Macedonia in peninsula Athoá (Griseb.), Byzantii (Sibth.), Tauria et prov. Caucasicæ et Transcaucasicæ (M.B., C.A. M., Buhse!), Libanus (Ehrenb!), Syria borealis ad Aintab (Haussk!), Persia borealis (Bge!), Affghania (Griff.), Belutschia (Stocks!).

Variat valvis minoribus quæ forma ex el. Grenier et Godron est forma typica Linnaei.

Ar. Geogr. Europa tota, Sibiria, Africa borealis, insulæ Canarienses et Madera.

Obs. *R. obtusifolio-crisp's.* — *R. pratensis* M. K. — *R. cristatus* Wallr. — *R. acutus* Auct. ex regione Caucasicæ indicatur.

12. **R. Nepalensis** (Spreng. Syst. II, p. 159) radicis fibris napiformi-incrassatis, caule elato superne ramoso ramis erecto-patulis, foliis oblongis inferioribus oblongis basi cordatis inferioribus oblongis basi cordatis petiolatis aculusculis, superioribus subsessilibus lanceolatis basi attenuatis, racemis elongatis strictis basi tantum foliiferis, verticillastris remotis, pedicellis infra medium articulatis tandem recurvis fructu subæquilongis, valvis ovato-orbiculatis valde reticulatis ad vel ultra medium in dentes crebros a basi sublatiore subulatos apice eximie hamatos fisis inæqualiter calligeris vel unicâ calliferâ ♀. *R. hamatus* Trevir. in Nov. Act. Leopold 13, p. 174 — *R. tuberosus* Roxb. — *R. Roxburghianus* Schult. — *R. peregrinus* Boiss. Diagn. Ser. I, 5, p. 46. — *R. uncinatus* Hort. — *R. hamulosus* Meisn. in p.c. Prodr. XIV, p. 55!

Hab. in pomariis regionis montanæ, Lydia in Tmolo ad Bozdagh (Boiss! Bal. 352!) et supra Philadelphiam (Boiss!), Taurus Cilicicus (Ky. semina sub nomine *R. Cedrorum*), Libanus circa Eden (Bl!), circa Gisr el Hajar (Ehrenb!).

Specimina Anatolica ab Indicis neutiquam distinguenda.

Ar. Geogr. Montes Indiæ, insulæ Java, P. B. Spei.

13. **R. subtrilobus** (Boiss Diagn. Ser. I, 5, p. 46) caule elato sulcato superne paniculato, foliis amplis undulato-subreundatis infimis petiolatis a basi rotundatâ vel subcordatâ oblongis acutiusculis, superioribus subsessilibus acutissimis omnibus subtus et ad nervos aspero-puberulis, paniculæ ramis elongatis laxiusculis basi exceptâ aphyllis, verticillastris subdistinctis, pedicellis infra medium articulatis fructu

æquilonoris tandem cernuis, valvis cordato-orbiculatis obsolete trilobis præter apicem triangularem integrum in dentes lanceolatos rectos numerosos ad $\frac{1}{3}$ vel $\frac{1}{4}$ usque fissis ④.

Hab. in pratis subalpinis Lydiæ ad Bozdagh (Boiss! Ball!).

Planta 4-5-pedalis, folia inferiora pedalia basi 4-5 pollices lata, paniculæ rami inferiores pedales, valvae $2\frac{1}{2}$ lineas diam. latæ. An inter *R. crispum* et *R. Nepalense* in consortio quorum crescit proles hybrida?

14. *R. stenophyllum* (Ledeb. Fl. Alt. II, p. 58. — Icon. tab. 399) caule erecto virgato in racemum basi interruptum densiusculum simplicem vel inferne parce et striete ramulosum abeunti, foliis lanceolatis utrinque attenuatis marginie subundulatis petiolatis, verticillastris inferioribus folio sessili linearis suffultis, pedicellis perigonio medioieri duplo longioribus prope medium articulatis, valvis trianguli-ovatis reticulatis subæqualiter calliferis margine plus minus dentieulatis vel serrulatis ②.

Hab. in humidis, Iberia Caucasicæ ad fluvium Kura prope Gori (Ower. in herb. Acad. Petrop!).

Hæc species ex radicis fibris elongatis attenuatis constantis indole certo certius monocarpica est

Ar. Geogr. Rossia austro-orientalis, Songaria, Sibiria Altaica.

15. *R. pulcher* (L. Sp. 477) biennis vel asperulus, caule erecto sulcato superne in paniculam flexuosam divaricatam rigidam abeunte, foliis parvis inferioribus petiolatis subcordatis ovato-oblongis vel panduriformibus obtusis, superioribus lanceolatis, racemis patentissimis elongatis, verticillastris remotis inferioribus folio linearis suffultis, pedicellis basi articulatis brevissimis tandem crassis cernuis, valvis ovato-oblongis acutiusculis utrinque apice excepto in 5-6 dentes rigidos rectos subspinescentes valvulae diametro breviores fissis omnibus inæqualiter calliferis ②. Ic. Engl. Bot. tab. 1576. — Rehb. Pl. Crit. tab. 486. — *R. divaricatus* L. Sp. I, p. 477 (forma foliis non strangulato-panduriformibus). — *R. tuberculatus* et *R. denticulatus* C. Koch Linn. XXII, p. 208!

Hab. in cultis et ad vias regionis inferioris et montanæ, Graecia (Sibth., Oph! Sart!), Thracia (Griseb.), Anatolia (Tchih.), Syria littoralis et Libanus (Bl! Gaill!), Palestina (Boiss!), Assyria (Haussk!), Egyptus in cultis Oasis parvae (Aschers!).

β. undulatus. — Folia subundulata. — *R. reticulatus* Besser Ind. Cremen, p. 1820. — *R. foreolatus* Hochst. in Lorent. Reise, p. 328 ex descriptione. Vix varietas.

Hab. in prov. Caucasicis (Hohen!).

Ar. Geogr. Europa tota australis et media a Britanniâ, Galliâ, Germaniâ australi ad regionem Danubialem et Rossiam australiem, Africa borealis, Maderâ et insulae Canarienses.

16. *R. Raulini* (Boiss. Diagn. Ser. I, 12, p. 100) biennis, caule erecto angulato-sulcato a basi pluries dichotomo, foliis parvis radica

libus longe petiolatis a basi breviter attenuata oblongo-obovatis obtusis margine subundulatis subtus ad nervos subpapillosum, verticillastris distantibus basi folio linear-lanceolato eis longiore suffultis, pedicellis fructiferis reflexis duris sub medio articulatis fructu subaequilongis, valvis triangulari-oblongis foveolatis utrinque in dentes 4-6 subulatos validos rectos spinosos ad $\frac{2}{3}$ latitudinis fissis omnibus graniferis, valvulae apice obtusiusculo denticulatae (2).

Hab. in cultis Cretæ ad Roumata et Khania (Raulin!).

Pedalis, affinis *R. pulchro* ab eo differt valvis paulo majoribus in spicas validiores et longiores multo profundius fissis, sed ex unico specimine descriptus et ulterius observandus.

17. R. Cassius, caulis ex rhizomate incrassato brevi erectis virgatis subsimplicibus, foliis parvis radicalibus petiolatis ovato-oblongis basi subcordatis vel rotundatis obtusis, superioribus brevius petiolatis oblongo-lanceolatis acutis, racemo longo parte inferiori excepta subaphyllo, verticillastris distantibus, pedicellis tandem crassis reflexis fructu brevioribus prope basin articulatis, valvis elevatim reticulatis coriaceis triangulari-deltoides acutis utrinque in 3-5 dentes subulatos apice subuncinatos ad medium fissis apice integris omnibus inæqualiter calliferis (2). *R. strictus* Meisn. in p.c. Prodr. XIV, p. 57 ex parte quoad pl. Syriacam non Link.

Hab. in Syriae monte Cassio (Boiss!).

Affinis *R. pulchro* a quo, caulis in racemum simplicem tenuem abeuntibus, valvis in dentes longiores uncinulatos et minus rigidos partitis differt. Erronee ad *R. strictum* Link relatus qui ex specim. herb. Berol. est varietas *R. dentati*. Ex aliis locis *R. stricti* in p.c. Prodr. indicatis Thracicus ad *R. pulchrum*, Arabicus ad *R. dentatum* spectant.

18. R. dentatus (L. Mant. p. 226) annus, caule erecto folioso simplici vel a basi ramoso, foliis inferioribus petiolatis basi rotundatis vel obsolete cordatis oblongis (interdum supra basin strangulatis panduriformibus) obtusis subundulatis, verticillastris subdiscretis omnibus folio lanceolato-lineari basi cuneato breviter petiolato suffultis longe racemosis, pedicellis fructu brevioribus reflexis, valvis parvis oblongo-lanceolatis acutis reticulatis omnibus callo saepius magno instructis utrinque 4-3-dentatis dentibus subulatis rectis patentibus valvae longitudinem aequantibus rarius obsoletis (3). *R. caulosissimus* Meisn. in p.c. Prodr. XIV, p. 57. — *R. Menzalensis* Ehr. exs!

Hab. in Egypti fossis et paludosis, (Del., Kraft! Schweinf! Boiss!), in magna Oase (Schweinf!), Oase Dachel (Aech!).

R. pleiodon. — Valvae 3-3-dentatae. — *R. strictus* Link Enum. Ber. p. 350 et herb. Berol! — *R. Ehrenbergii* Meisn. I. cit. p. 56! — *R. quadridentatus* (Ehrenb. exs! Spec. juniora callis minutis). — *R. Klotschianus* Meisn. loc. cit. p. 57 (saltem quoad plantam Atighanum!).

Hab. in Egypto ad lacum Menzaleh (Ehrenb!), Cataoniâ ad radices Beryt-dagh (Haussk!), Grusiâ (Wilhelms!), Assyriâ circa Mossul (Noë! Haussk!), Babyloniâ ad Bagdad (Oliv! Noë 1096!), Bassora (Haussk!), regno Mascate Arabie (Auch. 5341!), Affghanîa (Griff!), Belutschia (Stocks!).

Facile ab affini *R. palustri* foliis in petiolum non attenuatis distinctus. *R. Ucranicus* Besser in ditione non obvius est planta omnibus partibus multo tenuior, pedicellis tenuibus elongatis. Var. β . a cl. Meisner cum *R. pulchri* formis confusa fuit.

Ar. Geogr. India borealis.

19. *R. palustris* (Smith Fl. Brit. I, p. 394) biennis, caule erecto sulcato folioso paniculato, foliis radicalibus lanceolato-linearibus in petiolum brevem attenuatis acutis superioribus et floralibus verticillastris superantibus linear-lanceolatis acuminatis et linearibus, verticillastris racemosis remotiusculis, pedicellis reflexis fructu parvo sublongioribus infra medium articulatis, valvis rhombeis utrinque setaceo-bidentatis apice lanceolatis integris omnibus calliferis, dentibus valvae longitudine brevioribus ②. Ic. Fl. Dan. tab. 1872.

Hab. in limosis et paludosis, Macedonia (Friv!).

Ar. Geogr. Europa borealis et media a Scandinaviâ ad Rossiam, Galliâ, Germaniâ, Italiâ boreali, regione Danubiali, Songariâ, Sibiriâ.

20. *R. comosus* (Forsk. Fl. Eg. Arab. p. 76) annuus subacaulis a collo prolifere et breviter ramosus, foliis petiolatis basi cuneatis radicalibus oblongis obtusis, verticillastris in spicas densissimas cylindricas centralem in collo sessilem laterales ramos terminantes congestis omnibus folio lanceolato-lineari flores longe superanti suffultis, pedicellis tenuibus infra medium articulatis fructu minimo æquilongis, valvis triangularibus acuminatis reticulatis dentibus utrinque 2-3 setaceis valvâ multo longioribus, valvâ exteriore grosse callifera ⊖. *R. Ägyptiacus* L. Sp. I, p. 477.

Hab. ad limum exsiccatum Nili in Egypto media et inferiore (Ehrenb., Sieb! Boiss! etc.).

Obs. *R. Marschallianus* Rchb. species affinis sed verticillis remotis aliisque notis diversa in Tauriâ erronee indicata in ditione non crescit.

21. *R. bucephalophorus* (L. Sp. I, p. 479) annuus hermaproditus punitus a collo simplex vel sæpius pluricaulis, caulibus tenuibus erectis et ascendentibus in racemos longos tenues simplices aheuntibus, foliis petiolatis inferioribus ovato-spathulatis vel oblongo-lanceolatis summis linearibus, verticillastris remotiusculis 2-3-floris, pedicellis basi articulatis demum clavato-incrassatis complanatis et subtus sulcatis induratis arcuato-reflexis perigonio longioribus, valvis subecallosis a basi deltoideâ oblongo-lanceolatis apice integris inferne utrinque 2-4 dentatis dentibus patentibus subulatis spinosis vel rectis valvâ brevioribus vel longioribus uneinatis ⊖. Ic. Cav. tab. 41. — Fl. Graec. tab. 345.

Hab. in arenosis herbidis præsertim maritimis totius Græcie (Sibth., Sprun! Heldr!), Thracie ad Byzantium (Auehl Noël), Anatoliæ occidentalis

(Fleisch!) et australis (Bourg!), Cretæ (Sieb!), Rhodi (Bourg!), Syriae littoralis (Bl! Gailli!).

β. uncinatus. — Valvarum spinæ validiores uncinate. — *R. aculeatus* L. Sp. 481. Forumis intermediis ad typum transit.

Hab. in Cretā (Linn.), Palestinā (Boiss!).

Ar. Geogr. Hispania, Lusitania, Gallia australis, Italia, Africa borealis, insulæ Azoricæ et Canarienses.

SECT. II. ACETOSA.

22. **R. scutatus** (L. Sp. 480) perennis polygamus glaucescens glaber vel papillosus, radice repente, caulis numerosis ascendentibus flexuosis ramosis superne pluries dichotomis in racemos elongatos laxos abeuntibus, foliis carnosulis inferioribus longe petiolatis infimis cordatis ovatis obtusis, intermediis et superioribus breviter petiolatis hastato-panduriformibus acutis vel trilobis sinu profundo excisis lobo medio rhomboe-ovato transverse sublatiore lateralibus angustis, verticillastris 2-4-floris remotis, pedicellis filiformibus versus medium articulatis, valvis membranaceis basi cordatis ecallosis vix reticulatis achenio longioribus et latioribus ♀. Ic. Blackw. t. 506. — *R. hastifolius* M.B. Taur. Cauc. I, p. 290. — *R. pubescens* C. Koch Linn. XXII, p. 211.

Hab. in lapidosis saepius mobilibus, ad torrentes siccis regionis subalpinæ et alpine 5-6000', Græcia in Taygeto (Pichl!), Olympo Thessalo (Heldr! Orph!), Olympus Bithynus (Pichl!), Lydia in Tmolus ad Bozdagh (Bal!), montes Pisidiæ et Lycaoniæ (Heldr!), Cappadociæ (Bal!), Taurus Cilicus (Ky!), Pontus Lazicus et Armenia (Bal! Ky!), Caucasus et Transcaucasia (M.B., Rupr! Hohen!), Persia borealis in monte Elorus (Ky. 110!).

Forma hastifolia foliis profunde trilobis in Europâ occidentali æque ac in Oriente occurrit.

Ar. Geogr. Gallia centralis, Hispania borealis, Belgium, Germania australis, Helvetia, Italia, Dalmatia, regio Danubialis, Africa borealis.

23. **R. hastatus** (Don Prodr. Fl. Nepal. p. 74) polygamus fruticosus ramosissimus ramis novellis tenuibus striatis, foliis parvis glaucescentibus petiolatis aliis rhombis acutis aliis hastato-trilobis lobis linearibus lateralibus divergentibus, racemis paniculatis laxifloris. pedicellis infra medium articulatis, valvis mediocribus reniformi-orbiculatis obtusis vel retusis reticulatis basi sinu angusto cordatis et minute calliferis ♂.

Hab. in regni Cabulici alpinis (Griff!).

Caules inferne lignosi validi, rami annotini pedales et longiores, ioliorum lacinia 1-1½ lineas latæ intermedia 6-8 lineas longa. E^z affinitate *R. suffruticosi* Gay Hispanici.

Ar. Geogr. Jugum Himalaicum occidentale.

24. **R. Acetosa** (L. Sp. I, 481 exl. var.) perennis viridis glaber vel inferne puberulus, radicis fibris tenuibus, caulis erectis foliosis

in paniculam laxam elongatam strictam abeuntibus, foliis crassiusculis inferioribus longe petiolatis venosis ovato-oblongis obtusis basi sinu rotundato cordatis hastato sagittatis auriculis acutis, caulinis sessilibus lanceolatis acuminatis, stipulis laciniate-dentatis, verticillastris panicifloris aphyllis, pedicellis versus medium articulatis, valvis membranaceis reticulatis cordato-orbiculariatis basi squamulâ deflexâ obsoitis achenio longioribus, perigonii laciñiis externis triplo brevioribus reflexis ♀. Engl. bot. tab. 127. — *R. pseudacetosa* Bertol. Fl. It. 4. p. 252.

Hab. in pratis regionis montanae, Græcia in Peloponneso (Sibth.), in Olympo Thessaliam (Heldr!), Macedonia supra Bitolia (Orph.), Thracia (Griseb.), Byzantium (Sibth.), Tauria, Caucasus et Transcaucasia (Ledeb.), in ditione Talysh (C.A. Mey!).

R. alpinus. — Minor, collum incrassatum fibras crassiores pauciores elens, panicula brevior magis confertiflora, foliorum auriculæ subdivergentes. — *R. triangularis* Guss. Syn. I, p. 434.

Hab. in saxosis regionis alpinæ Græciæ, Kyllene 6000'-6500' (Heldr! Orph!), Parnassus 3000'-6300' (Orph! Heldr!).

Specimina Nebrodensibus et Calabriticis similia.

Ar. Geogr. Europa præsertim borealis et media a Scandinaviâ ad Rossiam, in australibus alpinus, Sibiria, America borealis.

25. **R. Tmoleus** (Boiss. Diagn. Ser. I, 12, p. 101) perennus glaucus glaber, radice, caulinis erectis sulcato-striatis in paniculam brevem dense multifloram thyrsoidem abeuntibus, foliis crassis oblongo-triangularibus acutis margine crisplis acutis basi hastatis auriculis divergentibus indivisis vel lobulo anetis inferioribus petiolatis superioribus sessilibus, stipulis laciniate-dentatis, pedicellis supra medium articulatis, valvis membranaceis reticulatis a basi rotundata ovatis basi squamulâ deflexâ obsoitis achenio longioribus, perigonii laciñiis externis triplo minoribus ♀.

Hab. in pratis subalpinis humidis ad Bozdagh montis Tmoli Lydiæ (Boiss!), ad pagum Gorumse ditionis Kassan Oglu Ciliciae Orientalis (Ky. 430!).

Validior *R. Acetosâ* a quo differt glauco-linea, panicula thyrsoidæ brevi densâ ramis strictissimis, valvis minoribus ovatis basi non cordatis.

26. **R. horizontalis** (C. Koch Linn. XXII, p. 211) perennus glaber glauco-serens, radice fibras cylindricas edente, caule crasso erecto striato-sulcato, foliis inferioribus petiolatis hastato-oblongis, superioribus sessilibus triangulari-lanceolatis acuminatis, stipulis lacero-dentatis, paniculae elongatae subaphyllæ ramis densifloris erecto-patulis, pedicellis versus medium articulatis, valvis membranaceis reticulatis basi cordatis callo deflexo basi obsoitis achenium superantibus, perigonii laciñiis externis deflexis parvis ♀.

Hab. in regione subalpina Ponti Lazici supra Djimil (C. Koch! Ball).

Caules crassiusculi et folia glauco-serens *R. Tmolei* sed panicula minus compacta ramis patentibus et valvæ valde cordatae. Hac ex loco natali

verosimiliter spectat *R. montanus* Ledeb. Fl. Ross. III, p. 510 non Desf. *R. montanus* Desf. = *R. arifolius* All. est species Europaea caule tenui, foliis viridibus membranaceis, stipulis integris, panicula laxissimam diversa.

27. *R. tuberosus* (L. Sp. I, p. 481) perennis viridis glaber, radice saepe incrassata fibras plus minus abrupte in tubera ovato-oblonga incrassatas edente, caulis erectis vel ascendentibus in paniculam brevem parum ramosam aphyllam abeuntibus, foliis membranaceis radicalibus longe petiolatis ovatis vel oblongis repandis obliuis basi rarius rotundatis saepius cordatis vel cordato-hastatis auriculis brevibus divergentibus obtusis vel acutis, foliis superioribus sessilibus hastatis linear-lanceolatis acutis, pedicellis supra medium articulatis, valvis membranaceis ovato-orbiculatis reticulatis basi cordatis rarius truncatis achenio longioribus et latioribus basi callo deflexo interdum deficiente auctis ♀. Ic. Fl. Graeca tab. 348.
— *R. Creticus* Boiss. Diagn. Ser. I, 12, p. 102.

Hab. in saxosis, pratis, regionis inferioris et montanae, Græcia ad Corinthum (Orph!), monte Parnes Atticæ et Onion Bœotiae (Heldr!), Cretâ (Raul!), Cycladibus (Sart!), Byzantii (Murman!), Lydia ad Smyrnam (Bal. 353!), Rhodo (Bourg!), Cypro (Sibth.), Syriâ ad Aleppo (Haussk!), Cappadociâ (Bal!), Georgiâ Caucasicâ (Hoh!), Persiâ boreali (Buhse!), Tauriâ (Stev.).

Ar. Geogr. Africa borealis.

28 *R. lacerus* (Balb. Miscell. p. 19) annuus, collo pluricaulis caulis herbaceis crassiulus decumbentibus et ascendentibus brevibus inferne dichotome ramosis, foliis parvis carnosulis petiolatis pinnatipartitis vel fidis inferioribus latioribus lobis lateralibus utrinque 1-2 ovato-triangularibus vel oblongo-linearibus parvis saepe deflexis interdum nullis terminali majore triangulari acuto, foliis summis lanceolato-linearibus integris, racemis axillaribus et terminalibus aphyllis subsimplicibus laxis, pedicellis filiformibus fructu longioribus prope basin articulatis, valvis tenuiter membranaceis reticulatis cordatis reniformi-orbiculatis callo oblongo instructis ♂. *R. pictus* Forsk. Eg. Arab. p. 77? — *Acetosa Aegyptia roseo seminis involucre, folio lacero* Lippi Schaw Specim. N° 5, tab. 1.

Hab. in arenosis desertorum et maritimis, Palestina ad meridiem Gaza (Boiss!), Egyptus ad Damiatam (Ehr! Sieb!), ad Alexandriam (Gaill! Samar!).

β. macrocarpus. — Valvulae majores ovatae.

Hab. in arenosis maritimis Syriae ad Ras Beirut (Labilll Bl!).

Species cum varietate lacerâ *R. Tingitani* L. non confundenda, in regno Tunetano et Algeriâ crescenti, radice perenni, foliis longe triangulari-lanceolatis acuminatis undulato-laceris distinctâ. *R. Tingitanus* L. erronee ut videtur in Smith Prodr. ex insulâ Cypro indicatur.

29. *R. vesicarius* (L. Sp. I, p. 479) annuus, a collo ramosus caulis herbaceis crassiulus striatis dichotomis, foliis carnosulis petiolatis a basi subcordata vel subhastata ovalis vel deltoideis integris, racemis oppositifoliis et terminalibus simplicibus et paniculatis

aphyllis, pedicellis supra medium articulatis tenuibus unifloris vel sæpius flores binos dorso connatos ferentibus, fasciculis paniculatis remotis, valvis tenuissime membranaceis reticulatis nervo non marginatis basi et apice sinu angusto cordatis ovalis subinæqualibus, majoribus margine sæpe replicatis ⊖. Ic. Campd. tab. 3.

Hab. in r. estribus maritimis Peloponnesi ad Naupliam (Sartorii), Egypti in desertis Kahirinis (Gailli!), littoris Egypciaco-Arabici inter Kosseir et Ras Benass (Schweinf!), ad Chargeh Oaseos magnæ (Schweinf!), in rupastribus apricis Persiae australis prope Gere (Ky. 33!).

Ar. Geogr. Insulæ Canarienses, Madera, Africa borealis.

30 R. roseus (L. Sp. I, p. 480) annuus, a collo ramosus caulinibus herbaceis crassiusculis pumilis striatis subsimplicibus ascendentibus vel erectis, foliis carnosulis petiolatis a basi truncatâ vel cuneatâ oblongo-deltoideis acutis summis lanceolatis, racemis oppositifoliis et terminalibus sæpius simplicibus aphyllis, fasciculis paniculatis remotis, pedicellis infra medium articulatis sæpius unifloris, valvis tenuiter membranaceis roseis reticulatis nervo tenui continuo interdum spinuloso-ciliato marginatis orbiculatis vel transverse latioribus basi et apice sinu profundo angusto incisis inæquilatis minoribus dorso sæpe papillosis ⊖. Ic. Flor. Graeca tab. 346.

Hab. in rupestribus Cypri (Sibth.) et ad Larnaca (Ky. 45!), Egypti ad Kahiram (Ky!), Arabiæ petreæ (Boiss!) et ad Ouadi Hebran (Schimp. 179!), Syria ad Gebel Khaisin prope Damascum (Gailli!), Kurdistaniæ ad Mt. Ieli (Noë 1158!), Persiae orientalis et Afganianæ (Bge!), Belutschia (S^t exs. sub *R. vesicario*!).

Valde et forsitan nimis affinis *R. vesicario*, humilior, minus ramosus, pedicelli inferius articulati et præsertim valvæ nervo marginatæ.

SECT. III. ACETOSELLA.

31. R. Acetosella (L. Sp. I, p. 481) dioicus, perennis sæpius multicaulis caulinibus tenuibus subsimplicibus in paniculam strictam subflexuosam aphyllam abeuntibus, foliis petiolatis hastato-tripartitis lanceolatis vel linearibus nonnullis vel omnibus interdum indivisis, lobis lateralibus triangularibus vel linearibus patentibus indivisis vel 3-5-fidis, verticillastris subdistantibus, floribus minimis, valvis internis cum achenio æquilongo arcte connatis ♀. Ic. Engl. Bot. tab. 1674. — Campd. Rum tab. 2. — *R. multifidus* L. Sp. I, p. 482 ex parte (forma foliis ob lacinias laterales fissas palmato-multifidis).

Hab. in regionibus magis borealibus ditionis sed ex confusione cum specie sequenti dubitanter et specimena authentica non vidi, prob. in Macedonia et Thraciæ borealibus, Tauriæ.

Ar. Geogr. Europa omnis, Africa borealis, Sibiria, insulæ Canarienses, P. B. Spei, America borealis et australis, Nova Hollandia.

32. R. acetoselloides (Bal. Bull. Soc. Bot. Fr. I, p. 282) diplocus, perennis multicaulis caulinibus erectis vel ascendentibus in panicul-

culam flexuosam abenntibus, foliis petiolatis hastato-tripartitis lacinia media oblongo-lanceolata majore lateralibus linearibus indivisis vel saepius multipartitis divaricatis, verticillastris paucifloris laxiusculis, floribus minimis, perigonii valvis ovatis obtusis achenio adpressis sed ab eo liberis phylla exteriora triplo superantibus ♀. *R. multifidus* L. Sp. I, p. 482 ex parte. — Sibth. Fl. Gr. tab. 349 et M.B. Taur. Cauc.

Hab. in collibus, agris derelictis, arenosis fere totius ditionis, Creta (Sieb!), Graeciae regio montana et subalpina in monte Malevo Laconiae (Orph!), et Velugo Aetolie (Heldr. Herb. Norm. 674!), Macedonia et Thracia (Friv! Pichler!), Byzantium (Noë!), Anatolia occid. ad Smyrnam et in monte Timolo (Bal. 352!), Anatolia borealis (Tchih!), Pontus Lazicus borealis et alpinus (Bal!), Armenia Turcica (Calv! Huet!), Cappadocia (Bal!), Cataonia mons Berydaghi (Haussk!), Caucasus et Transcaucasia (M.B., Hol!), Libanus (Bl!).

Simillimus *R. Acetosellae* et ab eo valvis cum fructu non concretis sed omnino liberis jantum distinguendus, haec nota distributione geographicæ suffulta gravis est, an satis firma sit ulterius observandum. Folia in utraque specie variant tripartita et multifida.

Ar. Geogr. Regnum Neapolitanum.

Species valde dubiae.

R. Armenus (C. Koch Linnaeana XXII, p. 210) caule elato striato crassiusculo in paniculam elongatam strictam abeunte, foliis radicalibus longe petiolatis elongato-oblongis acutis basi cordatis, caulinis sessilibus cordato-ovatis cauli adpressis, floralibus ad panicula ramos inferiores lanceolatis parvis, fructu.... ♀.

Hab. in ditione Daratchitchak Armeia Rossiae (C. Koch!).

Tripedalis, foliorum radicalium lamina fere pedalis tres pollices lata, panicula 12-15-polices longa. Ex specimina vix florenti descriptus et ideo affinitas inter *Lapathis* species perennes omnino dubia.

R. Lachanoides (Forsk. Eg. Arab. p. 209) foliis petiolatis radicalibus spathulatis obtusis, caulinis late lanceolatis, valvis integris granulatis.

Hab. in Anatolia (Forsk.).

An *R. conglomeratus*?

R. Chalepensis (Mill. Diet. № 11), caule ramoso, foliis cordato-oblongis acutis basi dentatis intimis amplis, floribus longe pedicellatis, valvis profunde spinoso-dentatis.

Hab. in Aleppo.

An *R. Nepalensis* vel *dentatus*?

'TRAPHAXIS (L. Gen. 356. — *Tragopyrum* M.B.).

Flores hermaphroditi. Perigonium petaloideum 4-5 partitum lacinias imbricatis internis demum accretis achenio adpressis. Stamina 6

vel 8 filamentis basi dilatata perigonii fundo insertis. Styli 2 vel 3. Ovarium lenticulare vel triquetrum pericarpio crustaceo. Semen loculum implens erectum lenticulare vel trigonum. Embryo in angulo seminis albumini farinaceo incumbens curvatus, radicula supera, cotyledones lineares. Frutices ramosissimi interdum spinescentes foliis alternis coriaceis vel crassiusculis cum pulvillo persistente articulatis. Ochreeæ scariosæ basi vaginantes. Pedicelli capillares articulati demum nutantes. Divisio in sectiones *Euatraphaxis* et *Tragopyrum* ob *A. Billardieri* in quæ flores tetrameri pentameris mixti observantur nimis artificialis est. eam tamen servavi. Artificialis quoque nota ex inflorescentiâ a claris Jaub. et Sp. deprompta, nam amic. Bunge observante eadem species vere racemulos ad ramos novellos et autumno laterales ad ramos jam lignescentes profert. Pedicelli demique ad basin in eadem specie ochreolâ propriâ cyathiformi donati vel orbati observantur.

Specierum Orientalium distributio.

SECT. I. EUATRAPHAXIS Jaub. et Sp. — Perigonium tetramerum. Stamina sex, stigmata bina. Achenium lenticulare.

A. spinosa, compacta, candida, Affghana.

SECT. II. TRAGOPYRUM. — *Tragopyrum* et *Tragatraphaxis* J. et Sp. Perigonium pentamerum. Stamina octo. Stigmata tria. Achenium triquetrum (confer *A. Billardieri*).

* Perigonii fructiferi sepala bina exteriora multo minora reflexa.

+ Folia subtus praesertim reticulata.

A. Billardieri, Tournefortii, laxifolia, pyrifolia.

+ + Folia anguste linearia uninervia avenia.

A. angustifolia, suedæfolia.

** Perigonii fructiferi sepala bina exteriora vix dissimilia non reflexa.

A. grandiflora, Aucheri.

SECT. I. EUATRAPHAXIS.

1. **A. spinosa** (L. Sp. I, p. 475) fruticosa, ramis gracilibus elongatis ramulisque patentibus vel stricti usculis spinescentibus vel inertibus cortice fuscescente vel albo-cinereâ, ochreis brevibus, foliis

parvis viridibus vel glaucis glabris ellipticis rhombis vel cordato-subrotundis reticulatis, pedicellis ad vel sub dimidiâ parte articulatis, floribus quadripartitis hexandris, perigonii laciniis ovato-orbiculatis binis externis reflexis interioribus binis auctis cordato vel reniformi-orbiculatis achenio plano ovato vel orbiculato accumbentibus coequatioribus et longioribus, stylo unico bistigmato δ . Species valde polymorpha eujus formas frequentiores numero altera in alteram manifeste transeuntes.

$\alpha. typica$. — Viridescens, folia elliptica vel oblonga. Haec est forma in hortis botanicis ubique culta. — *A. replicata* Lam. Dict. I, p. 329. — *A. Fischeri* Jaub. et Sp. III, II, p. 12.

Hab. in aridis lapidosis Transcaucasiae \ddagger ad Elisabethpol (Hohen!), Turcomaniae et Turkestaniae (Karel! Lehm!).

$\beta. rotundifolia$. — Folia ovato-rotunda vel orbiculata. — *A. Karelini* Jaub. et Sp. loc. citato. — *A. densiflora* C. Koch Linn. XXII, p. 212! — *A. Calverti* Boiss. Diagn. Ser. II, 4, p. 76.

Hab. in collibus Tauriae ad Sudak (Rehm! Stev.), Armeniae Turcicæ (C. Koch! Calvt!), Cataoniae inter Malatia et Karput (Haussk!), Persie austro-occidentalis in valle illuvii Chyrsan Elymaitidis (Haussk!), Turkestaniae (Lehm!).

$\gamma. Sinaica$. — Folia ovato-rotundata praecedentis sed minuta 2 lineas diam. lata, pedicelli abbreviati. — *A. Sinaica* Jaub. et Sp. I. cit. p. 12. — *Tragopyrum rotundifolium* Presl Benmerk. p. 109.

Hab. in cacumine montis Sinaitici Stae Catherineae (Bové! Schimp. 450!), in planicie elata Galala deserti Aegyptiaco-Arabici (Schweinf!), in Belutschia ad Kelat (Stocks!).

$\delta. glauca$. — Folia majuscula ovata vel oblonga saepè acuta et saepius glaucescentia, rami et ramuli canescentes vel cani.

Hab. in Armeniâ Rossicâ (Buhse!), prov. Karadagh et Aderbidjan (Buhse! Szov!), in valle Loura montis Elbrus (Ky. 541!), prov. Khorassan et Afghaniâ conterminâ (Bge!).

Ar. Geogr. Songaria.

2. A. compacta (Led. Fl. Alt. II, p. 55. Icon. tab. 411) fruticosa humilis ramis tortuosis abbreviatis, ramulis spinescentibus, ochreis laceris elongatis, foliis glaucescentibus obovatis vel ovatis obtusis reticulatis, pedicellis brevibus supra medium articulatis parte superiore laciniis externis vix æquilongâ, laciniis cordato-orbicularibus achenio orbiculato plano duplo longioribus et triplo latioribus δ .

Hab. in deserto Kisylkum Turkestaniae (Lehm!).

A formis rotundifoliis *A. spinosa* præter notas indicatas distincta ramis crassis brevibus tortuosis, ramulis confertim foliosis et floriferis.

Ar. Geogr. Songaria, Sibiria Uralensis.

3. A. candida (Boiss. et Haussk.) fruticosa subspinecens glabra, ramis elongatis ramulisque erecto-patulis elongatis gracilibus caudidis,

foliis parvis enerviis orbiculatis obtusissimis albido-cinereis, ochreis brevissimis, pediecellis infra medium articulatis, perigonii tetrameri laeiniis externis deflexis, internis fructiferis accretis cordato-reniformibus albis reticulatis achenio cordato-ovato nitido longioribus et la ioribus 5.

Hab. in glareosis ad radices montis Kuh Gerre ditionis Bachtiaris Persiae occidentalis (Haussk!).

Rami sesquipedales, folia 2½-3 lineas diam. lata, valvae interiores perigonii 3-4 lineas late. Species ramis foliisque candidis hisce enerviis ab *A. spinosæ* formis distincta.

4. A. Affghana (Meisn. in D.C. Prodr. XIV, p. 76) fruticosa glabra viridis ramosissima ramulis spinescentibus, ochreis brevibus fuscescenscentibus, foliis subsessilibus subnerviis ovatis acutis vel obtusis, pedicello valvis subbreviore infra medium articulato, perigonii tetrameri fructiferi valvis reniformi-orbiculatis tenuiter venosis achenio ovato plano dimidio longioribus 5.

Hab. in lapidosis regni Cabulici ad jugum Kullon (Griff.). Non vidi.

Folia 3-4 lineas longa 2-3 lata, valvae 4 lineas longæ et late. Distincta videtur ab *A. candida* formâ foliorum, ramorum colore, a formis *A. spinosæ* foliis obsolete nervosis. An tamen hujus forma?

SECT. II. TRAGOPYRUM.

5. A. Billardieri (Jaub. et Sp. Ill. Or. II, p. 44, tab. 44) fruticosa glabra pallide virens ramosissima ramulis abbreviatis saepius intricatis spinescentibus vel inermibus, ochreis brevibus bicuspidatis, foliis parvis reticulatis ovatis vel oblongis obtusis et acutis, pedicello ad vel infra medium articulato, floribus saepius omnibus pentameris octandris, perigonii laeiniis binis exterioribus ovatis deflexis interioribus ternis (rarius binis) accretis cordato-reniformibus achenio triquetro (rarius interdum plano) latioribus et longioribus, stylo 2 rarius trifido 5.

Hab. in rupestribus siccis regionis mediæ montis Hymetti Atticæ 2000' (Sart! Heldr. Herb. Norm. 1783!), Tymphresti Etoliae et montium Eubœa (Zucc.), Cretæ in latere meridionali montis Idæ supra Vorisa 4500' (Heldr!), Lycie supra Elmalu (Pest! Bourg. 252!), Phrygiæ ad Yachamichlar Keul supra Ouchak (Bal!), Ponti australis (Tchih!), Syriæ bor. in monte Ssoffdag (Haussk!), Antilibani supra Zebdani (Boiss!), in monte Garbi 4800' (Ky. 45!), ad Ouadi el Harir (Gaill!), Libani (Labill!).

β. heterantha. — Flores pauci (saepè in eodem racemulo) tetrameri achenio plano typicis pentameris achenio triquetro donatis commixti, nulla alia nota differentialis — *A. variabilis* Jaub. et Sp. loc. cit. p. 16, tab. 44.

Hab. in Syria (Auch. 2563!), in rupestribus montis Ssoffdag prope Balkis et Aintab (Haussk!), in lapidosis Cappadociae supra Karahissar 4000' (Bal!), in parte media montis Hermon Antil bani (Boiss!).

Species ab *A. spinosâ* floribus pentameris et achenio triquetro tantum distincta et etiam haec nota infirmatur varietate β . typo simillimâ in quâ flores tetrameris pentameris mixti occurunt.

6. A. Tournefortii (Jaub. et Sp. loc. cit. p. 14, tab. 112) fruticosa, ramulis subflexuosis inermibus novellis puberulis, ochreis in ligulam longam cuspidato-bifidam producens, foliis viridibus subpetiolatis a basi cuneatâ elliptico-oblongis cuspidulatis præter marginem crispulum et puberulum glabris, pedicellis infra medium articulatis articulo superiore laciniis externis vix longiore, perigonii pentameri valvis cordato-orbiculatis undulatis reticulatis achenio triquetro longioribus et multo latioribus δ . *Lapathum Orientale frutescens humilis* *flore pulchro* Tourn. Cor. p. 88.

Hab. in Armeniâ (Tourn. in herb. Mus. Par.). Non vidi.

Folia majora 9-10 lineas longa 2-4 lata, valvae demum 4 lineas longe $4\frac{1}{2}$ latæ. Species foliis cuspidatis et majoribus ab *A. Billardieri* formis dignoscenda.

7. A. buxifolia (M.B. Taur. Cauc. III, p. 284 sub *Tragopyro*) fruticosa hinnulîs lâte viridis, ramulis tenuibus flexuosis muticis junioribus pruinosis, ochreis fuscis brevibus breviter dentatis, foliis majusculis breviter petiolatis obovatis obtusis basi cuneatis margine crispulis tenuiter reticulatis, pedicelli juxta medium articulati articulo superiore laciniis exterioribus subduplo longiore, floribus pentameris, valvis cordato-orbiculatis roseis undulatis achenio trigono multo longioribus et latioribus δ . *A. buxifolia* Jaub. et Sp. loc. cit. p. 14. — *Polygonum buxifolium* M.B. Taur. Cauc. I, p. 300. — *Polygonum crispulum* Bot. Mag. t. 1065.

Hab. in Iberia Caucasicâ prope Tiflis (M.B., Hoh!), in Mingrelia (Eichw.). Folia interdum 9-10 lineas longa, valvae fructiferae demum $4\frac{1}{2}$ lin. latæ.

8. A. pyrifolia (Bge pl. Lelim. p. 483) fruticosa glabra, ramis elongatis albidis, ramulis patentibus rigidis brevibus valide spinosis, ochreis hyalinis subulato-bicuspidatis internodio longioribus, foliis majusculis longiuscule petiolatis a basi breviter cuneatâ obovato-orbiculatis obtusis mucronatis glaucescentibus subtus reticulatis, racemis lateralibus multi et densifloris, pedicelli articulo superiore inferiore et laciniis ellipticas reflexas duplo superante, perigonii pentameri valvis laciniis externas duplo superantibus orbiculatis, achenio trigono valvis dimidio breviore δ .

Hab. ad rupes vallis superioris Sarafchan prope Uramitau in Turkestania (Lehm!).

Frutex ut videtur majusculus ramis pedalibus, folia petiolo 3-4 lineas longo suffulta 8-12 lineas longa. Species racemis lateralibus nec terminalibus, valvis minoribus, ramulis spinosis ab *A. buxifoliâ* et *Tournefortii* distincta.

9. A. angustifolia (Jaub. et Sp. loc. cit. p. 15, tab. 115) suffruticosa ramosissima ramis strictis abbreviatis crebre foliatis, ochreis

bicuspidatis et laceris elongatis, foliis seabridis angustissime linearibus crasse uninerviis aveniis subtus margine revolutis floralibus brevissimis, floribus ex axillis superioribus 2-3-nis in racemos oblongos terminales densiusculos dispositis, pedicellis flore longioribus juxta medium articulatis, perigonii pentameri laciniis cordato-orbiculatis externis reflexis dimidio minoribus, achenio triquetro valvis duplo breviore et angustiore ȝ.

Hab. in Armeniâ (Auch. 2564!).

Suffrutex $\frac{1}{2}$ -1-pedalis, folia 1-1 $\frac{1}{2}$ -pollicaria $\frac{1}{2}$ -1 lineam lata, racemi 1-2-pollicares.

10. A. suædæfolia (Jaub. et Sp. loc. not. p. 15) fruticosa glabra, ramis sterilibus spinescentibus, foliis linearibus angustis acutis glaucis subtus revolutis aveniis nervo medio inconspicuo, ochreis brevibus bidentatis, pedicellis exsertis juxta medium articulatis, perigonii pentameri laciniis externis demum reflexis ovatis subbrevioribus, internis orbiculatis latioribus, achenio triquetro valvis multo angustiore et breviore ȝ.

Hab. in Persiæ prov. Aderbidjan prope Tabris (Auch. 5272 in herb. Mus Par.). Non vidi.

Ex descriptione brevi affinis videtur *A. angustifoliae* a quâ ramulis spinescentibus, foliis basi non angustatis obsolete uninerviis differt.

11. A. grandiflora (Willd. Sp. II, p. 440) fruticosum humile pallide virens, ramulis brevibus inermibus crebre foliatis, ochreis niveis internodio plerumque longioribus fimbriato-laceris, foliis subpetiolatis lanceolato-oblongis utrinque attenuatis acutis seabridis superne aveniis subtus subvenosis et margine revolutis, pedicellis ochreâ vix longioribus juxta apicem articulatis, perigonii pentameri laciniis cuneato-ovatis reticulatis exterioribus demum erecto patentibus triente minoribus, achenio triquetro valvis multo breviore et angustiore ȝ. *A. grandiflora* Jaub. et Sp. loc. cit. tab. 113. — *Tragopyrum grandiflorum* M.B. Taur. Cauc. III, p. 285.

Hab. in Armeniâ Turcicâ (Tourn., Auch. 2565!), in regione mediâ montis Aslandach Cappadociae (Ball).

Ramuli novelli e candice brevi lignoso tantum 2-4-pollicares, folia 6-10 lineas longa, flores breviter pedicellati in racemum ovatum densum foliosum dispositi.

12. A. Aucheri (Jaub. et Sp. loc. cit. p. 15, tab. 114) fruticosa, ramis elongatis ramulisque tenuissimis erectis, ochreis cuspidato-laceris internodio brevioribus, foliis anguste lanceolato-linearibus utrinque attenuatis acutis glabris subtus nervo remote nervuloso percursis et margine subrevolutis superioribus abbreviatis, floribus ex axillis superioribus 2-3-nis in racemum laxum brevem pauciflorum dispositis, pedicello flore breviore paulo infra apicem articulato, perigonii pentameri laciniis externis oblongo-ovatis apice demum subrecurvis, internis sublongioribus et latioribus ovatis omnibus reticulatis, achenio triquetro valvis subbreviore ȝ.

Hab. in aridis Persiae (Auch. 5271!), in montibus prope Pokideh (Buhset!).
Facies *Polygoni* e sect. *Avicularia*, rami 1-1½-pedales tenues, folia 1-2-pollicaria 1-2 lineas lata, racemi 1-3 pollices longi, interrupti.

POLYGONUM (L. Gen. 495).

Flores hermaphroditæ. Perigonium 4-5-sidum vel partitum semiherbaceum vel corollinum lobis subæqualibus persistentibus fructum includentibus. Stamina 8 biseriata vel rarius 5-4 imo perigonio inserta. Styli 2-3 liberi rarius connati. Ovarium liberum ovulo basilari erecto. Achenium trigonum vel lenticulare crustaceum, semen erectum. Embryo lateralis angulo albuminis accumbens vel semiimmersus arcuatus, cotyledones foliacæ, radicula supera. — Herbe annuae et perennes, vel suffrutices, petioli imâ basi in vaginam sæpius membranaceam (*ochream*) tubulosam vel ovato-oblongam expansi, bractæ interflorales *ochreolæ* dictæ.

Specierum Orientalium distributio.

SECT. I. BISTORTA Tourn. — Spica unica in apice caulis simplicis.

Styli tres liberi, stigmatibus parvis. Stamina 5-8. Achenium trigonum. Cotyledones accumbentes *Ochreæ* membranaceæ oblique truncatæ margine imberbes.

P. viviparum, *Bistorta*.

SECT. II. PERSICARIA Tourn. — Spicæ ramos caulis sæpius ramos terminantes. Styli 2-3 plus minus connati stigmatibus capitatis. Stamina 8-4 interjectis sæpe squamulis hypogynis. Achenium trigonum rarius lenticulare. Cotyledones accumbentes. *Ochreæ* cylindraceæ ore horizontaliter truncato sæpius setaceo-ciliatae.

* Perennia.

P. amphibium, *serrulatum*.

** Annua.

P. Hydropiper, *mite*, *flaccidum*, *minus*, *Persicaria*, *lapathifolium*, *lamerum*.

SECT. III. AMBLYGONUM Meisn. — Spicæ et babitus Sect. *Persicariae*. Styli 2 ad medium coaliti stigmatibus capitatis. Stamina 7 rarius 5-8 cum squamulis perigynis alterna. Achenium lenticulare. Cotyledones incumbentes lineares. *Ochreæ* cylindricæ.

P. Senegalense, *limbatum*.

SECT. IV. ACONOGONUM Meisn. — Flores in racemos paniculatos dispositi. Styli tres brevissimi stigmatibus capitatis. Stamina 8. Achenium triquetrum. Cotyledones accumbentes. Ochreæ oblique truncatæ margine imberbes.

P. Alpinum.

SECT. V. TINIARIA Meisn. — Flores in axillis fasciculati racemosi. Perigonii jacinæ externæ carinalæ vel alatae. Stamina 8. Squamæ nullæ. Stylus subnullus stigmatibus tribus capitatis. Achenium triquetrum. Cotyledones accumbentes. — Herbae volubiles.

P. Convolvulus, dumetorum.

SECT. VI. AVICULARIA Meisn. — Flores in axillis fasciculati vel solitarii spicas foliosas vel aphyllas formantes. Stamina 8 raro 5-6, filamenta trium interiorum basi ovata. Styli tres brevissimi stigmatibus capitatis minimis. Achenium triquetrum. Cotyledones incumbentes. Ochreæ basi amplexicaules breviter tubulosæ obliquæ bifidæ vel indivisæ, tandem apice laceræ. Ochreolæ florales cyathiformes.

§ 1. Annua.

* Ochreæ indivisæ tandem plus minusve dentato-laceræ.

P. tubulosum, polycnemoides, salsuginosum, corrigooides.

** Ochreæ bifidæ.

P. Cappadocicum, Bellardi, argyrocoleum, arenarium, pulchellum, aviculare.

§ 2. Perennia. — Rhizoma et caules parte inferiori plus minus indurata sed non suffruticosa.

* Ochreæ bifidæ.

P. equisetiforme, maritimum, Aleppicum.

** Ochreæ indivisæ apice tandem sœpe laceræ.

P. alpestre, hermiarioides.

§ 3. Fruticulosa. — Rhizoma lignosum multicaule.

* Ochreæ bifidæ.

P. setosum, luzuloides.

** Ochreæ indivisæ tandem laceræ.

P. radicosum, Kotschyanum, Libani, paronychioides, macrohymenium, Lemannianum, Affghanicum.

§ 4. **Spinescentia.**—Fruticulosa, ramuli post anthesin persistentes indurati rigidi. Pedicelli fructiferi nutantes, flores majusculi.

P. dumosum, aridum, salicornioides.

SECT. VII. PSEUDOMOLLIA. — Flores in glomerulos terminales dispositi. Perigonium non ut in sectionibus praeedentibus herbaceum vel corollinum sed hyalino-membranaceum profunde partitum laciniis binis externis navicularibus carinato-alatis. Stamina 7-5 basi dilatata. Styli bini. Acheneum compressum. Cotyledones incumbentes. Ochreæ stipuliformes. Ochreolæ bracteis ternis basi connatis constantes.

P. molliæforme, acaule

SECT. I. BISTORTA.

1. **P. viviparum** (L. Sp. 516) perenne, rhizomate tuberiformi, caule humili simplicissimo monostachyo, foliis subtus glaucescentibus margine subrevolutis inferioribus petiolo longo aptero suffultis ellipticis utrinque attenuatis, caulinis sessilibus lanceolatis, ochreis fuscis oblique truncatis, spicâ cylindricâ basi laxiusculâ sèpius bulbillis intermixtâ, ochreolis ovatis acuminatis, floribus breviter pedicellatis, staminibus exsertis, achenio ovato apice trigono ♀. Ic. Fl. Dan. tab. 13. — Engl. bot. tab. 669.

Hab. in pratis regionis alpinæ, montes Laconiae (ex Sibth.), Macedonia (Griseb.), Caucasus omissis 6-9000' (Ledeb.), Iberiæ et Guriæ montes (Ledeb., Owerin.).

Ar. Geogr. Alpes Europæ totius borealis et mediæ, montes Italiae borealis et mediæ, Dalmatiæ, Sibiriæ, Indiae, Americæ borealis.

2. **P. Bistorta** (L. Sp. 516) perenne, rhizomate crasso torto, caule elato simplici monostachyo, foliis subtus glaucescenti-puberulis margine subrevolutis inferioribus oblongo-ovatis basi truncatis vel cordatis petiolo longo superne alato suffultis. superioribus parvis lanceolatis acuminatis sessilibus amplexicaulibus, ochreis longissimis fuscis, spicâ densâ oblongâ obtusâ, ochreolis ovatis subulato-acuminatis, perigonio roseo, staminibus exsertis, achenio acute trigono ♀. Ic. Fl. Dan. tab. 421. — *P. carneum* C. Koch. Linn. XXII, p. 197.

Hab. in pratis alpinis Laconiae (Sibth.), Macedoniae, Thraciae (Griseb.), Ponti occid. (Tchih!), Ponti Lazici ad Djimil (Bal!), Armeniæ Turcicæ (Huett!), Caucasi totius et Iberiæ 4200'-9000' (Led. etc!).

P. angustifolium Meisn. — Folia radicalia oblongo-lanceolata vel linear-lanceolata imo linear-caricina in petiolum brevem sensim

angustata. Spicæ abbreviatæ ovatae. — *P. bistortoides* Boiss. Diagn. Ser. I, 5, p. 46. — *P. confusum* Meisn. — *P. Bistorta* var. *capitatum* C. Koch Linn. XIX.

Hab. in Olympo Bithyno et Tmolo Lydiae (Boiss!), monte Anemas Lycaoniæ (Heldr!), montibus Berydagħ et Aslandagh (Haussk! Ball!), Tauro Cilicico (Ky! Ball!), monte Elvend Persiae (Haussk!).

Ar. Geogr. Montes totius Europæ bor. et mediae ad Hispaniam borealem et Italiam medianam usque, Sibiria, regio Himalaica, America borealis.

SECT. II. PERSICARIA.

3. **P. amphibium** (L. Sp. 517) perenne, radice repenle, foliis oblongis vel lanceolatis, spicis terminalibus solitariis pedunculatis erectis oblongis densis, bracteis ovatis acutis pedicellos subæquanti-bus, floribus pentandris, perigonio eglantuloso roseo ♀. Variat.

α *natans*. — Caulis fluitans inferne radicans, ochreæ imberbes, folia natantia longe petiolata basi rotundato-subcordata lævia. — Ic. Fl. Dan. tab. 282.

β *terrestre*. — Caulis erectus, ochreæ breviter ciliatæ, folia sub-sessilia elongato-lanceolata angustiora et acutiora interdum adpresso scabrida.

Hab. α. in aquis, β in humidis probabiliter in totâ parte septentrionali ditionis, Byzantii et in Bithyniâ (Sibth.), Pisidiâ (Heldr!), Libanus (Bl!), Armenia (Calv!), Tauria, Caucasus et Transcaucasia (Ledeb!).

Ar. Geogr. Europa præsertim borealis et media a Scandinaviâ ad Rossiam, in australioribus rarius, Sibiria, China, Africa borealis, P. B. Spei, America borealis.

4. **P. serrulatum** (Lag. Gen. et Sp. 14), perenne, caulinibus inferne prostratis radicantibus dein erectis elatis ramosis, ochreis adpresso setosis margine ciliis sæpe eis æquilongis instructis, foliis subsessilibus a basi rotundatâ vel subcordatâ anguste linearis-sublan-ceolatis acuminatis margine adpresso ciliatis, spicis racemosis filiformibus laxiusculis erectis, floribus subhexandris, perigonio roseo eglantuloso, ochreolis crebre ciliatis ♀. — *P. salicifolium* Broussonet. Willd. Enum. p. 428.

Hab. in aquis fluentibus Atticæ ad Kephissum (Sprun! Heldr. Herb. Norm. 790!), Crete (Heldr!), Syriæ littoralis ad Sidonem (Bl! Gaill!), ad Tripoli (Boiss!), Ægypti ad Alexandriam (Samar!), Babylonia ad Korna (Noë!).

β. *salicifolium*. — Folia sublatiora et breviora, ochrearum setæ sæpe subbreviores. — *P. salicifolium* Del. Eg. (nomen solum). — *P. pseudohydropiper* Salzmann.

Hab. in Egypto inferiore ad Damiatam (Sieb!), ad Dimenete (Samar!).

Achenia ut in affinibus variant lenticularia et sæpius triquetra.

Ar. Geogr. Lusitania, Hispania australis et orientalis, Gallia australis, Italia, Sicilia, Africa borealis, Abyssinia, India borealis, insulæ Canarienses, Madera, P. B. Spei.

5. **P. Hydropiper** (L. Sp. 517) annum, acre, caulis erectis vel ascendentibus ramosis, ochreis brevibus breviter et parce ciliatis, foliis oblongo-lanceolatis et lanceolatis glabris vel ad nervum scabriusculis, spicis filiformibus interruptis arcuato-pendulis, floribus subhexandris, perigonio viridi-purpurascenti glanduloso - punctato, ochreolis ciliis destitutis ⊖. Ic. Fl. Dan. tab. 1376. — Rchb. Crit. tab. 394.

Hab. in humidis et ad fossas, Graecia (ex Sibth. Prodr.), Byzantii (Heldr!), Pontus Lazicus ad Rhizé (Ball), Syria prope Tripoli (Bl.), Persia in Susianá (Haussk!), Caucasus et Transcaucasia (Lebed.), Persia borealis in prov. Ghilan (Haussk!). Probab. fere ubique in ditione.

Ar. Geogr. Europa fere omnis præsertim borealis et media, Sibiria, Africa borealis, America borealis.

6. **P. mite** (Schrank Bair. Fl. I, p. 608) annum insipidum, caulis erectis vel ascendentibus ramosis, ochreis adpresso et parce hirtis apice ciliatis, foliis oblongo-lanceolatis lanceolatisque subtus scabridis, spicis filiformi-cylindricis interruptis plus minusve cernuis, floribus subhexandris, perigonio viridi-purpurascente eglanduloso, ochreolis ciliatis ⊖. *P. laxiflorum* Weihe Fl. 1826, p. 746.

Hab. ad fossas et aquas, prov. Transcaucasica Talysh prope Astara (Hoh.) et prob. alibi. An forma hybrida?

Ar. Geogr. Europa media ab Anglia et Gothia ad Rossiam medium, Italianam medium, Dalmatiam.

7. **P. flaccidum** (Roxb. Fl. Ind. -p. 291) annum? caulis erectis subflexuosis stricte ramosis, ochreis hirtis apice longe setosis, foliis pellucido-punctatis utrinque sed subtus copiosius adpresso hirtis breviter petiolatis basi attenuatis oblongis acuminatis, spicis stricte paniculatis filiformibus longissimis laxifloris nutantibus, perigonio glanduloso, ochreolis ciliatis ⊖. *P. Donii* Meisn. Polyg. p. 72.

Hab. in regno Cabulico (Griff.). E ditione non vidi.

Folia eis *P. hydropiper* multo longius acuminata, ochreæ longius ciliatæ, spicæ multo longiores, achenia dimidio minora.

Ar. Geogr. India.

8. **P. minus** (Huds. Engl. I, p. 148) annum, caulis diffusis vel procumbentibus tenuibus ramosis, ochreis adpresso hirtis longe ciliatis, foliis lanceolato-linearibus superne sensim attenuatis, spicis filiformibus laxis erectiuseculis, floribus subhexandris, perigonio purpureo eglanduloso, ochreolis ciliatis ⊖. Ic. Engl. bot. tab. 1043. — *P. pusillum* Lam. Fl. Fr. III, p. 235. — *P. strictum* All. Ped. 2, p. 207, tab. 68. — *P. serrulatum* Hoh. pl. exs. non Gussme.

Hab. in stagnis iestate exsiccatis, oryzetis, Transcaucasia ad Caspium prope Lenkoran et Astara (Hoh!) et prob. alibi.

Ar. Geogr. Europa media a Scandinavia ad Rossiam medium, Lusitania, Italia borealis, regio Danubialis, Sibiria, India, Japonia.

9. **P. Persicaria** (L. Sp. 518) annuum, caulis erectis vel ascenditibus patule vel stricte ramosis, ochreis adpresso hirsutis longe ciliatis, foliis oblongo-ellipticis vel lanceolatis, spicis pedunculatis vel sessilibus oblongo-cylindricis densis erectis scutulis, floribus subhexandris, perigorio virenti-roseo glanduloso, ochreolis ciliatis ♂. Rehb. Ic. Crit. fig. 684. — *P. biforme* Wahl. Succ. — *P. Niloticum* Meissn. in d.c. Prodr. XIV, p. 1191

Hab. ad aquas prob. totius ditionis. Vidi e Macedonia (Orph!), Bithynia (Thirke), Ponto Lazico ad Rhizé (Bal!), Syria circa Damascum (Gaill!), Egypto inferiore ad Damiatam (Ehr!). In Tauriā, Caucaso et Transcaucasia (Ledeb.)

Variat achenis lenticularibus et rarius trigonis

Ar. Geogr. Europa fere tota, Sibiria, India, America borealis.

10. **P. lapathifolium** (L. Sp. 517) annum, caulis erectis ramosis, ochreis truncatis parce et brevissimis ciliatis, foliis a basi attenuata ovato-ellipticis vel lanceolatis margine ciliatulis, spicis oblongo-cylindricis vel cylindricis densis erectis vel subnutantibus, floribus subhexandris, pedicellis perigonioque virenti albido vel rubello plus minus glanduloso-asperis, ochreolis non ciliatis ♂. Ic. Engl. bot. tab. 1382. — *P. nodosum* Pers. Ench. I, p. 44, et Rehb. Ic. Crit. tab. 492 (forma nodis caulis incrassatis). — *P. obtusatum* Steud. in Ky. Alepp. № 437. — *P. tenuiflorum* Presl (forma spicis tenuioribus).

Hab. ad aquas, fossas totius ditionis in regionem alpinam ascendens, ab Attica et Bœotia (Heldr. Herb. Norm. 789!), Macedonia in peninsula Athoā (Orph!), ad Ciliciam (Ky!), Syriam littoralem et Libanum (Bl!), A-syriam (Ky!), regionem Caucasicanum (Ledeb.).

P. incanum. — Folia subtus præsertim adpresso tomentosa. — *P. incanum* Schm. — *P. tomentosum* Schranck Fl. Bair.

Hab. in monte Malevo Laconie (Orph!), Pisidiā ad lacum Beychehr (Heldr!), Tauriā et Caucaso (M.B.).

Ar. Geogr. Europa omnis, Sibiria, Japonia, India, America.

11. **P. lanigerum** (B. Br. Prodr. 419) perenne, totum adpresso arachnoideo-lanatum canum, caulis parte immersa crebre et longe radicibus superne ramosis, ochreis longis adpresso lanatulis apice truncatis breviter ciliatis, foliis lanceolatis longe acuminatis in petiolam brevem attenuatis spicis tenuibus sessilibus vel breviter pedunculatis paniculatis erectis, ochreolis denticulato-ciliatis, pedicellis brevissimis, perigonio vix glanduloso, staminibus subsenis, achenio saepius lenticulari ♀. *P. lanatum* Roxb. — *P. arachnoideum* Klotsch.

Hab. in fossis oryzetorum Oasis parvæ Egypti mediae prope Anenah (Aschers. № 468! sterile).

Quamvis sterile tantum observatum omnino videtur *P. laniferum* jam a formis incanis *P. Persicariae* et *lapathifoli* radice perenni distinctum.

Ar. Geogr. Africa australis, India Orientalis, Nova Hollandia.

SECT. III. AMBLYGONUM.

12. P. Senegalense (Meisn. Mon. p. 54) perenne glabrum, caule crasso elato ramoso inferne fistuloso radicanti, ochreis longis latis subinflatis tenuiter membranaceis margine breviter ciliatis, foliis amplis petiolatis basi attenuatis late oblongo-lanceolatis longe acutatis punctatis margine sebredis, spicis racemoso-paniculatis linearibus elongatis densis erectis, ochreolis ciliatis, perigonio roseo glanduloso, achenio lenticulari ♀. *P. macrockatou* Fresen. Flora, 1838, p. 601. — *P. Persicaria* var. *macrophyllum* Ehrenb. herb!

Hab. in aquis Egypti interioris (Ehr.), ad canalem Mahmoudieh (Gail.!), ad Dimenete (Samar!), ad Ramleh (Duperq!).

Planta pluripedalis, folia saep 6-9-pollices longa interne 12-18 linea lati, spicæ 2-3-pollices longe. Haec species foliis magnis acuminatis et spicis tenuibus longis omnino refert *P. glabrum* Willd. ochreis non ciliatis, perigonio eglanduloso, embryonis cotyledonibus accumbentibus, etc., distinctissimum; hoc *P. glabrum* ex Affghania in p.c. Prodr. indicatur sed erroneè nam locus citatus (Loudianah) Indicus est.

Ar. Geogr. Abyssinia, Senegal, Borbonia.

13. P. limbatum (Meisn. in p.c. Prodr. XIV, p. 123), perenne adpresso et breviter pilosum, caulis ramosis ascendentibus, ochreis petiolo brevi longioribus arte cylindricis hispidis limbo herbaceo patente semiorbiculati superatis, foliis lanceolatis utrinque attenuatis, spicis solitariis vel corymbosis erectis cylindrico-linearibus densifloris strictis basi interruptis, ochreolis ciliolatis, floribus majusculis, perigonio roseo eglanduloso ♀.

Hab. ad ostia Nili (Ehrenb! in herb. Berol. Specimen sterile).

Ar. Geogr. Senegal, Africa tropica, India.

SECT. IV. ACONOGONUM.

14. P. Alpinum (All. Ped. I, p. 296, tab. 68) perenne glabrum vel puberulum, caulis erectis crassiusculis angulato-striatis flexuosis ramosis, ochreis hirtis brunnneis oblique truncatis laceris, foliis oblongo-lanceolatis vel lanceolatis basi cuneatis subsessilibus acuminatis, floralibus linearibus, paniculâ terminali folia superante racemis compositis multifloris. perigonio albido vel carneo basi saep glauco pedicello erecto sublongiore, achenio trigono levigato ♀. *P. polymorphum* Ledeb. Fl. Ross. ex parte. — *P. undulatum* Murr. Com. Götts. 5, p. 34 tab. 5.

Hab. in pratis pinguis regionis alpinae, Macedoniae mons Peristeri (Girib., Orph!), Rhodope Thraciae (Friv!), Olympus Bithynus (Auch! Boiss! Clem!), Tmolus Lydie (Boiss!), Armenia Turcica (Calv!), Pontus Lazicus (Bal!), mons Argaeus Cappadocie (Bal!), Caucasus (M.B., Rehm!), prov.

Transcaucasica Karabagh (Szov!), mons Karatau Turkestaniæ (Lehm!), Affghania (Griff.).

β latifolium. — Folia breviter petiolata late ovato oblonga majora 2 $\frac{1}{2}$ -4 poll. longa 1 $\frac{1}{2}$ -2 lata) nervis secundariis subtus magis prominentibus. — *P. latifolium* Ky. et Boiss. Sched.

Hab. in argillosis alpium prov. Musch Armeniæ 6500' (Ky. 462!).

Floriferum tantum visum sed a formâ typicâ non specifice diversum videtur.

Ar. Geogr. Hispania australis, Pyrenei, Alpes Galliæ et Helvetiæ Italiæque borealis, regionis Danubialis, Sibiræ, regionis Himalaicæ.

SECT. V. TINIARIA.

15. **P. Convolvulus** (L. Sp. p. 522) annum, caulis gracilibus angulatis striatis prostratis vel scandentibus, ochreis brevibus oblique truncatis, foliis cordato-sagittatis, floribus 3-6 pedicellis nutantibus suffultis ad axillas fasciculatis superioribus in spicas laxas approximatis, perigonii virenti-albidi laciniis ternis exterioribus subcarinatis, achenio trigono laciniis perigonii adpressis tecto ⊖.

Hab. in cultis ubique a Græciâ (Sibth., Heldr!), ad Caucasum et Transcaucasiam (Ledeb.), Libanum (Bl!), Persiam occidentalem (Haussk!), Affghaniam (Griff.).

Ar. Geogr. Europa tota, Sibiria, Asia borealis, Japonia, Africa borealis, America borealis.

16. **P. dumetorum** (L. Sp. 522) annum, caulis elongatis ramosis cylindricis volubilibus, ochreis brevibus oblique truncatis, foliis cordato-sagittatis, floribus 3-6 ad axillas fasciculatis pedicellis nutantibus suffultis fasciculis superioribus in spicas approximatis, perigonii virentis laciniis ternis exterioribus membranaceo-alatis, achenio trigono nitido ⊖. Ic. Fl. Dan. tab. 756.

Hab. in sepibus, dumetis, Græcia (ex Sibth.), Pontus Lazicus prope Trapezuntem (Ball), Tauria et Caucasus (M.B.), Transcaucasia ad Lenkoran (Hoh.), Affghania (Griff. ex Meisn.).

Ar. Geogr. Europa borealis et media, Asia borealis.

SECT. VI. AVICULARIA.

§ 1. Annua.

17. **P. tubulosum** (Boiss. in Ky. Sched. 1845) annuum flavovirens scabridulum vel glabrum, collo multicaule, caulis brevibus decumbentibus vel assurgentibus simplicibus vel parce ramosis angulo-triquetris tenuiter virgatis ochreis et foliis adpressis omnino vestitis, ochreis hyalinis subnerviis ovato-lanceolatis apice simbriatis, foliis carnosulis internodiis longioribus oblongo-linearibus uninerviis margine subrevolutis, floralibus conformibus flores et ochreas supe-

rantibus, floribus in axillis 1-3 brevissime pedicellatis longe et laxe spicatis, perigonii tubulosi tubo trigono lobis approximatis triplo longiore, achenio lucido leví ⊖. Diagn. Ser. I, 1, p. 83. — *P. rotibollioides* Jaub. et Sp. Ill. tab. 122 (1846).

Hab. in glareosis et arenosis montanis et alpinis Persiae, inter Teheran et Ispahan (Auch. 2590 bis!), circa Teheran (Ky. 481), in valle Lar jugi Elbrusensis 9000' (Buhsei!), in Persia australis montibus Sabst Buschon et Kuh Delu (Ky. 432! et 500!), in monte Sawers 8000' (Haussk!), in ditione Khorassan (Bge!).

Aline *P. polycnemoides* sed perigonii forma egregie distinctum, caules variant 1-7-pollicares.

18. ***P. polycnemoides*** (Jaub. et Sp. Ill. Or. tab. 120 cum deser.) annum glabrum a basi pluricaule caulibus prostratis gracilibus flaccidis angulatis, internodiis foliis brevioribus vel ea æquantibus, ochreis hyalinis subbenerviis apice laceris folio brevioribus, foliis parvis carnosulis subulato-linearibus uninerviis eveniis erecto-patulis, floribus in axillis 1-2 brevissime pedicellatis spicas longas tenues laxas formantibus, perigonii fructiferi ovati lobis tubo subbrevioribus, achenio nitido minute punctato ⊖. *P. Olivieri* Jaub. et Sp. loc. cit. tab. 121 forma vegetior foliis longioribus.

Hab. in glareosis et arenosis praesertim regionis subalpinæ et alpinæ, Cilicia Trachea in montibus Ermenek (Perl), Taurus Cilicicus ad Bulghar Maaden (Ky. 354! Bal!), Cappadocia prope Caesaream (Bal. 1120!), inter Malatia et Karput (Haussk!), Libanus ad Cedretum (Ky. 286! sub *P. rotibollioide*), supra Elen et Dinam (Bl!), alpes Kurdistanie Persicae et Persiae austro-occidentalis 7-9000' (Haussk!), Mesopotamia (Oliv. et Brug.), in Affghania (ex cl. Meisn. sed dubitanter).

Radicem annuam, caulinis flaccidis, foliis angustis, floribus subsolitariis, perigonii tubo longiore, achenis punctatis ab omnibus *P. herniarioides* formis distinguendum, rami in spicas tenues 3-5-pollicares abeunt.

19. ***P. salsugineum*** (M.B. Taur. Cauc. I, p. 304) annum, caule erecto tenui humili a basi strictiuscule ramoso flexuoso, internodiis elongatis, ochreis brevibus semihyalinis multinerviis simbriato-laceris, foliis in plantæ parte inferiori anguste linearibus internodia æquantibus cito deciduis, floribus solitariis axillaribus sessilibus spicas filiformes aphyllas laxissimas breves formantibus, perigonii minimi infundibuliformis tubo herbaceo lobis rubellis oblongo-linearibus longiore, achenio lanceolato-trigono opaco punctato perigonium superante ⊖.

Hab. in schistosis subsalsis bituminosis Caucasi Orientalis (M.B.). E ditione nondum vidi.

Plantula 4-6-pollicaris facies *Buffoniae*, spicæ remote 3-5-floræ 6-10 lineas longæ, perigonium sub anthesi lineam excedens, achenium non vidi.

Species perigonio tubuloso-infundibuliformi insignis.

Ar. Geogr. Rossia austro-orientalis.

20. ***P. corrigoioides*** (Jaub. et Sp. Ill. Or. tab. 124 cum deser.) annum glabrum, glaucescens vel subviolascens a collo ramosissimum

ramis elongatis flexuosis subangulatis vel ramulosis, internodiis inferioribus ochreâ folioque longioribus, ochreis breviter ovatis hyalinis enerviis dentatis vel laceris, foliis linear-i-spathulatis basi attenuatis carnosulis uninerviis eveniis, floribus axillaribus 4-6-nis pedicellis tenuibus infra medium articulatis eis triplo longioribus sussultis racemos tenues inferne interruptos superne densiusculos formantibus, perigonio albido-roseo fructifero ovato ultra medium lobato, achenio minimo levi nitido ⊖. *P. myrianthum* Boiss. MSS.

Hab. in Babyloniae exsiccatis circa Bagdad (nec ut videtur in Persia) (Oliv. et Brug., Noe 1007! Haussk!).

Icon citata plantam juniores in quâ folia florum proportione longiora sunt exhibet. Species facie *P. herniarioidem* quodammodo referens sed floribus minoribus albo-roseis longe pedicellatis egregie distincta.

21. *P. Cappadocicum* (Boiss. et Bal. Diagn. Ser. II, 4, p. 78) annum glaucum glabriuscum, caulis e collo pluribus erectis rigidulis, internodiis folio longioribus, ochreis argenteis enerviis elongatis fimbriatis elongatis sed folio et internodio brevioribus, foliis cito deciduis carnosulis uninerviis enerviis elliptico-linearibus basi attenuatis margine revolutis seabridulis, floribus solitariis vel binis sessilibus racemos breves laterales et terminales ochreis et foliis flores subsuperantibus intermixtos formantibus. perigonii albidi ovati tubo lobis breviore, achenio levissimo nitido ⊖.

Hab. inter segetes Cappadociae prope pagum Erek 4500' (Bal. exs. 4106!).

Affine *P. polycnemoidi* sed ab eo a cl. Balansa qui ambo viva observavit specie distinctum, caules erecti nec prostrati 4-5-pollicares, ochreæ magis evolute, internodia longiora, folia latiora et margine revoluta, ochreæ bifidæ, flores in racemos densos 4-5 linea longos laterales et terminalem nec in spicas longas laxas dispositi, achenia levia nec punctata.

22. *P. Bellardi* (All. Ped. II, p. 205, tab. 90, fig. 2) annum glabrum, caule erecto ramoso-paniculato ramisque gracilibus angulostriatis, internodiis elongatis, ochreis brevibus semihyalinis 6-7 nerviis laceris, foliis planis elliptico-lanceolatis uninerviis acutis subvenosis, floralibus anguste linearibus flore brevioribus, floribus 2-3-nis pedicellis calyci aequilongis vel longioribus sussultis spicas terminales longes tenues interruptas nudas formantibus, perigonio viridi rubro-marginato fructifero clauso achenium majusculum minute punctatum subnitidum arcte includente ⊖. *P. patulum* M.B. Taur. Cauc. I, p. 304. — *P. strictum* Ledeb. Fl. Alt. et Ic. tab. 444 (forma tenuior stricta). — *P. chlorocoleum* Steud. in Ky. Sched. — *P. reticulatum* C. Koch Linn. XXII, p. 200.

Hab. in cultis præsertim regionis montanæ, Graecia in montibus Kyllene 5000' (Heldr.) et Malevo 4000' (Orph. Fl. exs. 38!), Thessalia prope Volo (Orph!), Rhodus (Bourg!), Anatolia in Bithynia (Pichl!), Pisidiâ (Boiss! Heldr!), Cilicia in regione inferiore Tauri (Bal!), Armenia (Huet! Ky!), Syria ad Aleppo (Ky. 231!), in Libano usque ad Cedretum (Ky. 287! Bl!), Arabia petrea in jugo Sinai (Schimp. 388!), Egyptus inferior (Kral!), Egypti

Oases (Asch!), Tauria (Pall!), Transcaucasia (Hohen!), Persia borealis (Auct! Buhse!), Mesopotamia (Haussk!), Turkestania (Lehm!).

Ar. Geogr. Gallia, Hispania, Lusitania, Italia, Dalmatia, regio Danubialis, Rossia media et australis, Sibiria tota, India borealis, Africa borealis.

23. *P. argyrocoleum* (Steud. in Ky. Pl. Alepp. 440) annum viride, caulis erectis vel ascendentibus tenuiter virgatis ramosissimis, ramis in spicas filiformes nudas interruptas laxas abeuntibus, internodiis elongatis, ochreis brevibus basi fuscis 6-8-nerviis apice hyalinis laceris, foliis lanceolatis p'lanis utrinque attenuatis acutis uninerviis demum deciduis, floralibus linearibus floribus multo brevioribus, floribus 1-3-nis pedicellis perigonio sublongioribus suffultis, perigonio viridi albo-marginato fructifero clauso achenium parvum laevissimum nitidissimum arete includente ☽. *P. Noënum* et *P. deciduum* Boiss. in Noë pl. exs.

Hab. in cultis Assyria ad Mossul (Ky! Haussk!), prope Bagdad et Mohammera (Noë!), ad Bassora (Noë!), in monte Zagros Persiae occid. (Haussk!) et prope Kirrind (Noë!).

Facies *P. Bellardii*, folia angustiora, achenium dimidio minus nitidissimum laevissimum nec punctatum. Hæc notâ magis proximum *P. arenario* a quo præter ramos diffusos perigonio herbaceo albo-marginato nec fere toto corollino et profundius in lobos ampliores partito differt. Sub anthesi ochreolæ florales *P. Bellardi* non majores sunt sed per autumnum plantâ indurata et magis ramosâ sæpe augentur.

24. *P. arenarium* (W. K. Pl. rar. p. 69, tab. 67) annum, a basi ramosum ramis diffusis tenuibus flexuosis, internodiis elongatis, ochreis basi fuscis subsexnerviis superne hyalinis laceris, foliis linearilanceolatis uninerviis nervulosis acutis basi attenuatis, floralibus brevissimis, floribus 1-3-nis pedicellis æquilongis suffultis spicas terminales aphyllas breves laxiusculas formantibus, perigonii albo-rosei lobis magnis tubo longioribus, achenio laevissimo nitidissimo ☽. *P. elegans* Ten. Fl. Nap. II, p. 207, tab. 35. — *P. Venantianum* Clem. Sert. p. 83, tab. 3!

Hab. in arenosis Thraciae inter Kasanlik et Eskisagra (Janka!), Byzantii (Clem!), Caucaso et Transcaucasiâ (M.B., Hohen!).

Spicæ (sed non semper) eis *P. Bellardi* densiores, semper breviores, perigonium magis corollinum lobis majoribus.

Ar. Geogr. Italia australis, Regio Danubialis, Rossia media et australis, Sibiria.

25. *P. pulchellum* (Lois. Fl. Gall. I, p. 284, tab. 26) annum a basi ramosissimum caulis sæpius decumbentibus elongatis, internodiis elongatis, ochreis basi brunneis 4-6-nerviis superne hyalinis laceris, foliis oblongo-lanceolatis nervulosis acutis basi attenuatis, floralibus ad bracteas minimas reductis, floribus 1-2-nis pedicellis perigonio longioribus suffultis in spicas filiformes laxissimas terminales dispositis, perigonii albo-rosei lobis tubo longioribus, acheniis

sublucidis tenuiter rugulosis ♂. *P. arenarium* Auct. Fl. Gall. non W. K. — *P. Bellardi* var. *effusum* Meisn. in p.c. Prodr.

Hab. in cultis et ruderatis Atticæ (Heldr!), Macedoniâ (Friv.), prov. Musch Armenie inter Gumgum et Koweg 5200' (Ky. 288!).

Species pluribus confinis a *P. Bellardi* perigonio corollino, acheniis rugulosis dimidio minoribus distincta. *P. argyrocoleum* differt perigonio fere toto herbaceo, achenio laevissimo, *P. arenarium* inter omnes magis proximum achenii nitore et laevitie. Probabiliter confusum et alibi reperiendum.

Ar. Geogr. Gallia australis.

26. *P. aviculare* (Linn. Sp. I, p. 519) annum interdum subperennans, a collo saepius ramosum caulinibus procumbentibus rarius erectis ramulosis, ramulis tota longitudine foliosis, internodiis elongatis, ochreis internodio brevioribus basi fuscis 6-8-nerviis superne hyalinis demum laceris, foliis ellipticis vel lanceolatis venosis planis, floribus 3-5-floris axillaribus pedicellis brevissimis suffultis, perigonio herbaceo, acheniis ruguloso-striatulis parum nitidis ♂. Ic. Fl. Danica, tab. 803. — Engl. bot. 1252.

Hab. in cultis, ad vias totius ditionis a Græciâ! ad Egyptum! Affghaniam! Belutschiam! a regione inferiori ad subalpinam.

β. littorale. — Rami crassiores, folia magis carnosa. — *P. littorale* Link in Schrad. Journ. I, p. 54.

Hab. in littoribus maritimis Græciae et ejus insularum (Heldr!), Ponti (Tehih!), Syriae littoralis (Bl! Gaill!), Egypti inferioris (Schw! Samart!).

γ. alpinum. — Caules brevissimi subsimplices erecti vel decumbentes. — *P. aviculare* var. *nanum* Boiss. Voy. Esp.

Hab. in regionis alpinæ humidis vel exsiccatis, mons Delphi Eubeæ (Heldr!), Taurus Cilicieus 7-8000' (Ky. 354! Bal!).

Eudem varietas in regione arcticâ! Alpibus! Hispaniæ australis montibus occurrit.

Obs. *P. acetosum* M.B. species affinis et forsitan non sat distincta e Rossia australi et Sibiria erroneè in regione Caucasicâ et Affghanâ indicatur.

Ar. Geogr. Totus orbis terrarum.

§ 2. Perennia.

27. *P. equisetiforme* (Sibth. et Sm. Fl. Græc. I, p. 266, tab. 364) perenne, caulinibus e rhizomate indurato lignoso-incrassato longis decumbentibus duris inferne foliosis internodiis elongatis, ochreis fuscis multinerviis apice capillaceo-laceris, foliis subsessilibus oblongis et oblongo-lanceolatis acutis margine crispulis, floralibus anguste linearibus flores aequantibus vel vix superantibus, floribus in racemos terminales laxos elongatos vel confertos abbreviatis dispositis 2-3-nis pedicellis eis aequilongis suffultis, achenio nitido ♀. *P. equisetiforme* v. *Græcum* Meisn. Prodr. p. 86. — *P. controversum* Guss. Prodr. I, p. 472. — *P. Ehrenbergii* Meisn. p.c. Prodr. XIV, p. 86.

Hab. in siccis Graeciae (Link), Cephaloniæ (Heldr!), Cretæ a regione inferiori usque ad 1500' (Sieb! Raul! Heldr!), Rhodi (Bourg!), Ciliciae (Bal!), Syrie a regione littorali usque ad regionem subalpinam Libani (Boiss! Bl! Gaill!), Palestinæ ad Jordanum (Boiss!), Egypti ad Alexandriam, Kahiram, in districtu Fayum etc. (Ehrenb! Gaill! Samar! Kralik! etc.), Mesopotamia (Haussk!), Persie bor. (Buhse!), Afghanistan (Griff.).

Variat racemis densi-vel laxifloris, folia floralia in formâ densifoliâ breviora, forma laxiflora in iconâ Fl. Graecæ exhibetur; ibidem flores digni dicti prob. ex errore vel ex formâ monstrosa nam eos semper trigynos vidi. Planta ex Corsicâ et Sardinia est species diversa, *P. scoparium*.

Ar. Geogr. Lusitania, Hispania australis et orientalis, Sicilia, Africa borealis.

28. *P. maritimum* (L. Sp. I, p. 519) perenne, caulinibus ex rhizomate ramoso indurato procumbentibus totâ longitudine foliosis, internodiis brevibus, ochreis basi fuscis sub-12-nerviis superne longe hyalinis laceris, foliis glaucis subsessilibus ellipticis venosis margine revolutis, floralibus vix diminutis flores longe superantibus, floribus 1-3-nis axillaribus breviter pedicellatis in spicas foliosas longas laxas dispositis, achenio levè nitido ♀. Ic. Fl. Græc. tab. 363.

Hab. in littoribus maritimis Graecie (Sibth., Sprun! Heldr!), Cretæ, Cypri, Rhodi et fere omnium insularum (Sibth., Raul! Bourg!), Macedonia et Thracia (Friv! Cast!), Syrie littoralis (Bl!), Egypti infer. (Bové!), Tauræ (Rehm!), Ponti Lazici (Bal. exs. sub *P. aviculari!*!).

Ar. Geogr. Europæ regio mediterranea, Rossia australis, Africa borealis, insulae Azoricæ, Maderenses et Canarienses, America borealis et australis.

29. *P. Aleppicum* (Boiss. et Haussk.) glabrum perenne, caulinibus ex rhizomate tenui indurato decumbentibus vel ascendentibus tenuibus nanis simplicibus vel ramosis in racemos densiusculos abeuntibus, internodiis inferioribus foliis et ochreis longioribus, ochreis tubulosis a basi fuscescenti 6-8-nervi hyalinis laceris, foliis minimis elliptico-linearibus obtusis in petiolum brevem attenuatis subtus multinerviis, floralibus diminutis oblongis flores non æquantibus, floribus 3-4-nis pedicellis eis sublongioribus apice articulatis suffultis tubuloso-hypocrateriformibus tubo herbaceo lobis oblongis albidis longiore, achenis.... ♀.

Hab. ad vias prope Aleppo Syriæ (Haussk!).

Caules cum racemis 1-2-pollicaribus 3-4-pollicares, folia 2-3 lineas longa; floralibus longe pedicellatis affine *P. corrigioloidi* sed rhizoma perenne fere suffrutescens, flores majores, pedicelli apice nec infra medium articulati, perigonium non ovatum sed infundibuliforme tubo elongato sensim attenuato.

30. *P. alpestre* (G.A. Mey. Enum. p. 157) glabrum, rhizomate lignoso plus minus incrassato caudiculos duros squamosos caulesque numerosos breves crebre foliosos prostratos edente, ochreis hyalinis internodia brevia subæquantibus binerviis oblongo-lanceolatis apice demum bifidis vel dentatis, foliis internodio longioribus subearniosis ovato-oblongis in petiolum brevem attenatis acutis uninerviis eveniis margine scabriusculis, floralibus vix diminutis flores longe superan-

tibus, floribus ex fere omnibus axillis 5-8 breviter pedicellatis, perigonio sub anthesi infundibuliformi lobis patentibus tubo æquilongis, fructifero indurato trigono tubo lobis conniventibus longiore, achenio magno arete inclusu obtuse trigono laevi ♀. Ic. Jaub. et Sp. tab. 118. — *P. cognatum* ♂. *alpestre* Meisn. Polyg. Prodr. p. 91. — *P. monticolum* Steud. Nom. — *P. pluriflorum* G. Koch Linn. XXI, p. 202.

Hab. in siccis, rupestribus, ad vias regionis montanæ et alpinæ, Bithynia prope Prusiam (Pichl!), Phrygia ad Ouchak (Ball!), Lydia in monte Tmolo et Caria (Boiss!), Pisidia (Heldr!), Lycia supra Elmalu (Bourg. 254!), Taurus Cilicus (Ky. 422! Ball!), Libanus (Ehr! Bl!), Antilibanus et Hermon (Boiss! Ky!), ditio Golani Palestinae (Wetzst!), Aleppo ad vias et Mesopotamia (Haussk!), Armenia Turcica et Rossica (Ky! Bourg! Szov!), Caucasus centralis (C.A. Mey!) et orientalis (Beck!), Iberia Caucasicæ et ditio Talysh (C.A. M! Hoh!).

Nomen specificum Meisneri ad formam pedicellis longioribus, *P. rupestre* Kar. et Kir. spectat et ideo minus aptum.

P. ammannoides Meisn. in D.C. Prodr. — Internodia abbreviata, folia minora et angustiora. — *P. ammannoides* Jaub. et Spach Ill. Or. tab. 119.

Hab. in prov. Aderbidjan Persiae (Auch. 5273!).

Ar. Geogr. Songaria, regio Himalaica.

31. **P. herniarioides** (Del. Fl. Æg. Illustr. № 412) perenne a collo ramosissimum ramis prostratis seabriusculis totâ longitudine crebre foliatis elongatis ramulosis, interuodiis brevibus, ochreis brevibus laxis hyalinis laceris imâ basi fuseescens tibus paucinerviis, foliis oblongo vel linearis-spathulatis obtusis eveniis internodia æquantibus vel superantibus, floribus 3-6-nis axillaribus herbaceis breviter pedicellatis, perigonii fructiferi ovati basi angustati tubo lobis breviore, acheniis parvis basi apiceque attenuatis laevibus nitidissimis ♀. *P. illecebroides* Meisn. in D.C. Prodr XIV, p. 94? — *Pol. effusum* Meisn. ibid. p. 93.

Hab. in arenosis Egypti ad Nili ripas ab Alexandriâ et Kahirâ usque ad Thebas (Del., Boiss! Samar! Kral! Schweinf!), Syriae littoralis (ex Meisn.), Afghanistan (Griff.).

Variat foliis internodiisque plus minus abbreviatis, axillis pauci et plurifloris.

P. Roxburghii. — Folia mucronulata. Florum pedicelli perigonio æquilongi. — *P. Roxburghii* Meisn. loc. cit. p. 93. — *P. Dryandri* Spreng. — *P. elegans* Roxb. Fl. Ind. 2, pag. 291.

Hab. in regno Cabulico (Griff.).

Ar. Geogr. Italia australis, Abyssinia et Afr.ca tropica orientalis, Senegal, India.

§ 3. Fruticulosa.

32. **P. setosum** (Jacq. Obs. III, p. 8, tab. 57) glabrum, rhizomate suffruticoso multicauli, caulis erectis vel ascendentibus tenuiter

virgatis elongatis duris rigidulis striatis in racemos simplices vel paree ramoso-paniculatos abeuntibus, ochreis hyalinis longe tubulosis multinerviis apice fimbriato-laceris, foliis internodia superantibus vel æquantibus anguste lineari-lanceolatis elongatis margine subrevolutis subtus parallele multinerviis, floralibus abbreviatis flores superantibus, floribus 1-3-nis brevissime pedicellatis 1-3-nis racemos tenuissimos longos laxissimos formantibus, perigonii corollini albi tubo brevissimo lobis rotatis, achenio (ex cl. Meisn.) ovato nitido ȝ. Ic. Jaub. et Spach Ill. tab. 125. — *Pol. Orientale Caryophylli folio flore magno albo* Tourn. Cor. p. 39.

Hab. in regione alpinâ, Lydia in cacumine Tmoli supra Bozdagh (Balt!), Armenia (Tourn., Auch. 2588!), circa Baibut (Bourg!).

β. restionoides. — Folia angustissima, floralia floribus breviora et sic spicæ subundæ. — *P. restionoides* Boiss. et Haussk. MSS.

Hab. in aridis districtûs Warto prov. Musch Armeniæ 4800' (Ky. 309!), in valle Dehghan Tauri Cataonici (Haussk!).

Caulis cum racemis 10-15-pollicares, folia 10-15 lineas longa lineam dimidiata lata.

33. *P. luzuloides* (Jaub. et Sp. Ill. Or. tab. 126) glabrum, rhizomatis suffruticosi ramis interdum elongatis repentibus fibrilliferis, caulis numerosis erectis virgatis striatis rigidis stricte paniculatis, ochreis hyalino-fuscescentibus tubulosis elongatis multinerviis apice fimbriato-setosis, foliis longis late lanceolato-linearibus internodio sublongioribus inter nervum medium et marginales plus minus manifeste venosis, floralibus subulatis flore brevioribus, floribus 2-4-nis pedicellis perigonio æquidistantibus suffultis racemos longos filiformes laxos formantibus, perigonii corollini tubo brevissimo, lobis patentibus, achenio oblongo lævi ȝ. *P. dracunculifolium* Boiss. in pl. Noë. — *P. pachyrrhizum* Trautv. Stirp. Nov. V, p. 15!

Hab. in montibus Begdaghi et Baltandagh Cataonie (Haussk!), in Armenia prope Guungun 4000' (Ky. 565!), prope Karikiburun (Radde!), in Persia occid. montibus Elvend (Auch. 2589!), Dalechani (Haussk!), et in montibus supra Kirrind (Noë 1115!).

Valde affine *P. setoso* cuius forsitan varietas foliis latioribus (inf. 3-4 lineas latis). Differt tamen floribus longius pedicellatis et submajoriibus.

34 *P. radicosum* (Boiss. in Ky. Sched. 1845) glabrum, caulis e rhizomate crasso lignoso compacto plurimis herbaceis nanis procumbentibus vel ascendentibus simplicibus crebre foliosis, ochreis hyalinis ovatis subbinerviis apice dentatis, foliis parvis internodia æquantibus vel superantibus subcarnosis ovatis vel obovatis obtusis margine subrevolutis uninerviis in petiolum attenuatis, floralibus conformibus. floribus axillaribus 3-4-nis breviter pedicellatis, perigonio herbaceo infundibuliformi lobis margine rubellis demum conniventibus tubo subbrevioribus, achenio nitido lævi inclusu ȝ. *P. serpylaceum* Jaub. et Sp. Ill. tab. 117.

Hab. in Persiae borealis jugo Elbrusensi in alpe Elamut (Auch. 5274!), in alpe Hasartschal (4991!), in Persiae australis alpe Kuh Daëna (Ky. 779!) et ad nives alpis Kuh Nur 13000' (Haussk!).

Caules tenues 1-3-pollicares, folia *Serpilli* 2-4 lineas longa.

35. P. Kotschyanum (Boiss. in Ky. Sched. 1845) scabriuscum ramosissimum, caulis fruticulosus brevibus crassiusculis, ramis novellis rigidis simplicibus vix striatis in racemos simplices laxos abeuntibus, ochreis hyalinis subbinerviis oblongis fimbriato-laceris, foliis lanceolatis et lanceolato-oblongis margine revolutis uninerviis, floralibus diminutis sed flores superantibus, floribus 1-3-nis axillaris, perigonio herbaceo fructifero indurato ovato-angulato reticulato achenium leve nitidissimum includente 3. *P. thymifolium* Jaub. et Sp. Ill. tab. 116.

Hab. in Persia australi inter Fasa et Schiraz (Auch. 5276!), in regionibus superioribus alpis Kuh Daëna (Ky. 719!).

Rami novelli herbacei sed duri et rigiduli 3-4-pollicares, folia 4-5 lineas longa, affine *P. radicoso* differt scabritie, ramis rigidis etc.

36. P. Libani (Boiss. Diagn. Or. Ser. I, 12, p. 99) glabrum, rhizomate lignoso superne ramoso caules novellos herbaceos breves procumbentes vel ascendentes inferne denudatos superne foliis et ochreis imbricatis omnino vestitos edente, ochreis hyalinis ovatis ventricosis apice fimbriato-laceris internodia brevissima superantibus, foliis parvis elliptico-lanceolatis obtusis cum mucronulo cito evanido subtus elevatim multinerviis, floralibus flores superantibus, floribus in axillis subsolitariis subsessilibus spicas terminales breves densas foliosas formantibus, perigonii carnei ovali tubo virente lobis obovatis duplo breviore, acbenio... 3. *P. Hermonis* Ky. MSS.

Hab. in cacuminibus excelsis Libani 8000'-9000' in monte Makmel (Boiss!), Hermonis (Boiss! Ky. 188!).

β. Cedrorum. — Folia sublatiora, caules elongati interdum 3-4-pollicares, flores 2-3-ni longius pedicellati et ideo ex ochreolis longius exserti. forma vegetior. — *P. Cedrorum* Boiss. et Ky. Diagn. Ser. II, 4, p. 77.

Hab. in Libani regione subalpinâ ad Cedros 5800' et ad Hasrun (Bl! Ky. 344!), circa Balbek (Bl!). Huc quoque specimina substerilia ex alpinis montis Ida Cretae 4600' (Heldr. 3741!) spectare videntur.

Affine *P. paronychioidi* quod facie refert et *P. macrohymenio* sed characteribus distinctum.

37. P. paronychioides (C.A. Mey. Talysh. En. p. 120) fruticulosum glabrum, ramis novellis ex rhizomate crasso lignoso herbaceis brevissimis cespitosis ochreis et foliis imbricatis dense obtectis, ochreis hyalinis lanceolatis subnerviis apice laceris folia subæquantibus internodia brevissima longe superantibus, foliis anguste linearisubulatis subtus uninerviis margine revolutis longe mucronatis, floribus in axillis subsolitariis subsessilibus foliis et ochreolis fere

occultatis spicas densas breves subdistichas formantibus, perigonii lobis roseis tubo herbaceo brevioribus, achenio oblongo acute trigono laevissimo nitido 5. *P. Paronychia* C.A. Mey Enum. p 158 non Cham. et Schlecht. --- *P. Meyeri* Steud. Nom.

Hab. in lapidosis aridis montium ditionis Talysh 4000' (C.A. M1 Hoh!), in cæcumine montium Seidkhodji prope Khoi prov. Aderbidjan Persiae (Szov!), prov. Khorassan in regione alpinâ (Bge!).

Rami novelli 1-2-pollicares, spicae 6-8 lineas longæ, folia 3-4 lineas longa. Planta exsiccata fragillima ramis et spicis in articolos secedentibus.

38. ***P. macrohymenium***, suffruticosum glaucum, ramis e rhizomate indurato numerosis cespitosis herbaceis brevissimis ad angulos seabridis in spicas densas breves subdistichas foliis et bracteis imbricatis constantes cito abeuntibus, internodiis 2-3 inferioribus elongatis spicarum brevissimis, ochreis albis non nitidis latissime amplexieauli-ovatis acutis apice laceris floralibus subintegris, foliis linear-laneolatis obtusis mucronulatis margine revolutis, floribus in axillis solitariis et ochreis occultatis virenti-albidis sessilibus lobis tubo subæquilongis, acheniis . . . 5.

Hab. in Persia prope Ssof inter Teheran et Ispahan (Bge!).

Facies *P. paronychoides*, ramorum juniorum 2-3-pollicarium internodia 1-3 inferiora elongata, spicæ pollicares dense imbricatæ, ochreae multo latiores membranaceæ sed non nitentes, folia latiora floralia ochreis sublongiora. Planta exsiccata fragillima.

39. ***P. Lemannianum*** (Meisn. in D.C. Prodr. XIV, p. 91) glabrum suffruticosum, caulis ex rhizomate lignoso brevibus pluribus cespitosis filiformibus nanis ochreis undique occultatis, ochreis hyalinis enerviis internodia brevissima superantibus inferioribus oblongis acutis superioribus duplo latioribus ovatis obtusiusculis imbricatis diu integris, foliis ochreas æquantibus subulato-linearibus mucronulatis demum mutatis obtusis margine revolutis, floribus solitariis sessilibus minutis, perigonio corollino infundibuliformi semiquinquefido, acheniis . . . 5.

Hab. in alpinis regni Cabulici ad Koh-i-Baba et Yonutt 10-15000' (Griff. 1726!). Non vidi.

Canes 1-2½-pollicares basi lignescentes, ochreæ 3-5 lineas longæ, folia 3-4 lineas longa, perigonium magnit. *P. herniaroides*.

40. ***P. Affghanicum*** (Meisn. in D.C. Prodr. XIV, p. 90) suffruticosum caulis e rhizomate lignoso elongato numerosis erectis duriusculis subsimplicibus obsolete angulatis pruinoso-papillosis, ochreis hyalinis obsolete trinerviis oblongis setaceo-acuminatis demum laceris elongatis sed inferioribus internodio brevioribus, foliis glabris linearibus margine subrevolutis obsolete uninerviis eveniis setaceo-mucronatis, floribus solitariis axillaribus paucis, perigonio ad medium partito, acheniis . . . 5.

Hab. in regno Cabulico (Griff. 1724). Non vidi.

Caulis spithamei, ochreæ 4-6 lineas, folia 8-12 lineas longa $\frac{1}{2}$ -1 lata, flores magnit. *F. avicularis*.

§ 4. Spinescentia.

41. **P. dumosum** (Boiss. in Ky. Sched. 1845. — Diagn. Ser. I, 7, p. 83) tomentello-scabriuscum fruticulosum nanum ramosissime dumosum, ramis tenuibus rigidis subangulatis non striatis simplicibus rarius stricte et parce ramulosis in racemos tenues laxos abeuntibus mox aphyllis et persistentibus, ochreis hyalinis enerviis oblongis acuminatis internodia et folia subæquantibus, foliis linear-ellipticis acutis margine revolutis subtus uninerviis strictis internodio brevioribus, floralibus abbreviatis sed flore longioribus, floribus 1-2 axillaribus pedicellis eis æquilongis tandem nutantibus suffultis majusculis roseis, perigonio ad $\frac{3}{4}$ in lacinias oblongas partito, achenio... 5.

Hab. in Persiâ australi prope ruinas Persepolis (Ky. 242!).

Dumi hemispherici sesquipedales, folia 3-4 lineas longa lineam lata, flores sesquilineam longi.

42. **P. aridum** (Boiss. et Haussk.) lente scabriuscum fruticulosum ramosissime dumosum, ramis tenuibus elongatis dichotomis et ramulosis in racemos tenues longos laxissimos aphyllos abeuntibus demum persistentibus, ochreis breviter ovatis internodio multo brevioribus basi fuscescentibus 5-6-nerviis cæteris hyalinis, foliis in parte inferiori ramorum paucissimis linear-lanceolatis margine revolutis cito deciduis floralibus abortivis, floribus majusculis ex racemi ochreis 1-2-nis pedicellis brevibus tandem cernuis suffultis, perigonio roseo ad $\frac{3}{4}$ partito, achenio punctato subopaco 5.

Hab. in arenosis apricis montium Sawers et Nur Persiæ australis (Haussk!).

Dumus semipedalis vel pedalis, racemi 3-5-pollices longi tenues. Affine *P. dumosum* a quo differt ochreis abbreviatis basi plurinerviis nec enerviis, racemis elongatis aphyllis.

43. **P. salicornioides** (Jaub. et Sp. Ill. tab. 123 cum descr.) lente seabinuscum fruticulosum pumilum ramosissimum, ramis rigidis crassiusculis striatis patule ramulosis cito aphyllis ramulisque tandem induratis subspinosis, ochreis hyalinis brevibus truncatis laceris, foliis minimis carnosis internodia brevissima superantibus ovatis et oblongis uninerviis margine revolutis floralibus abortivis, floribus majusculis in axillis inferioribus 1-2-nis pedicellatis. perigonio roseo ad $\frac{3}{4}$ in lacinias ovatas partito. pedicello fructifero cernuo, acheniis crassis opacis punctatis 5. *P. oligophyllum* Boiss. in Ky. Sched. 488 nomen antiquius sed minus aptum.

Hab. in Persiâ inter Fasa et Schiraz (Auch. 5277), in alpe Kuh Delu (Ky. 468!), in apricis montium Sawers et Kuh Nur 10-13000' (Haussk!).

Ramis subspinescentibus, florū structurā, etc., affine *P. dumoso*, differt ramis crassioribus striatis patule ramulosis, ochreis brevissimis; ramuli in axillis tantum inferioribus flores paucos ferunt superne aphylli ochreis approximatis brevissimis obsiti. Achenium fere 2 lineas longum.

SECT. VII. PSEUDOMOLLIA.

44. **P. molliaeforme** (Boiss. Diagn. Ser. I, 7, p. 84) annum glabrum nanum simplex vel parce ramosum, caulis filiformibus brevissimis dichotomis vel subnullis florū glomerulos ochreis foliis quæ intermixtos inferne subdissitos superne agglomeratos edentibus, ochreis hyalinis nitidis enerviis a basi subventricosâ lanceolatis acuminatis setaceo-laceris, floralibus ovatis indivisis, foliis linearī setaceis mucronulatis 3-5-nerviis subrecurvis, floralibus ochreas et flores superantibus, floribus solitariis subsessilibus ochreis occultatis, perigonio hyalino fere ad basin quinquepartito lobis rotundatis, staminibus quinis 2-3 antheriferis, stylis divergentibus, achenio ovato compresso nitido ⊖.

Hab. in glareosis montis Kuh Daēna Persiae australis (Ky. 778!), Persiae bor. in monte Totschal prope Teheran (Ky. 573!), prope Schahrud (Bge!).

Plantula 1-2-pollicaris, caulis tenuissimis, internodiis inferioribus 3-4 lineas longis vel subnullis et tunc glomeruli onnes in unicum radicalem condensati, folia 4-5 lin. longa linea tertiam partem lata, perigonium vix lineam dimidiā longum.

45. **P. acaule**, annum glabrum, caule nullo, florū glomerulis in capitulum unicum radicale concretis, foliis linearī-setaceis rectis trinerviis mucronatis ochreas et flores subsuperantibus, ochreis hyalinis non nitentibus oblongis acutis apice dentato-laceris floralibus integris, floribus solitariis subsessilibus ochreis occultatis, perigonio membranaceo fere ad basin partito laciniis ovatis, staminibus 7-8, stylis rectis, achenio.... ⊖.

Hab. in Persia orientali inter Ischredabad et Tudisek inter Yezd et Ispan (Bge!).

Capitulum radicale avellanā minus glomerulis 3-5 sessilibus densifloris constans. Valde affine praecedenti differre videtur caule nullo, foliis abbreviatis, ochreis nec nitidis nec acuminatis, staminibus numerosioribus, etc.

ORD. CII. NYCTAGINACEÆ.

(Lindl. Syst.)

Flores (in nostris) hermaphroditi. Perigonium infundibuliforme tubo persistente fructum includente, limbo marcescente vel deciduo. Stamina numero varia disco cupulari circa ovarii basin inserta basi

sæpius connata. Ovarium liberum breviter stipitatum superum uniloculare, ovulum erectum anatropum, stylus simplex. Fructus basi indurata perigonii inclusus indeliscaens. Embryo incurvus vel conduplicatus vel longitudinaliter convolutus, albumen farinaceum includens. Radicula infera. — Ordo Valerianaceis affinis.

BOERHAAVIA (L. Gen. № 22).

Flores hermaphroditici exinvoluti. Perigonium infundibuliforme vel campanulatum parte inferiore demum accretâ et indurata fructum involvente et persistente, superiore quinquelobâ marcescenti sæpius decidua. Stamina 1-3 hypogyna basi in annulum coalita, antheræ subexsertæ biloculares. Ovarium acutum uniloculare, stylus filiformis, stigma obtusum. Fructus cylindrico-obconicus. Embryo conduplicatus. — Herbæ basi suffrutescentes perennes.

1. B. plumbaginea (Cav. Ic. II, p. 7, tab. 112. — Asch. et Schw. Beitr. p. 166) basi suffruticosa, caulis herbaceis sæpius elongatis parte inferiori dichotome ramosis, superiori subscandentibus, foliis a basi subcordata, truncata vel rotundato-cuneata ovatis margine integris inferioribus obtusis cæteris acutiusculis, pedunculis axillaribus folio longioribus umbellam terminalem et sepe verticillos 1-2-sessiles ferentibus, pedicellis minute glandulosis ovario æquilongis vel brevioribus, perigonii infundibuliformis hirti ovario triplo longioris tubo angusto in limbum sensim ampliato, fructu obconico-cylindrico basi attenuata adpresse hirtulo striato ♂. **B. scandens** Forsk. Deser. 12. — **B. dichotoma** Vahl En. I, p. 290. — **B. grandiflora** A. Rich.

2. glabrata. — Viridis, glabra vel minute puberula. — **B. verticillata** Boiss. in Ky. exs. non Poir.

Hab. in Palestinae planicie Esdraelon et prope Nazaret (Roth! Schlumberger!), ad Ascalon (Sieb!), circa Jericho (Boiss! Ky. 426!), in jugo Sinaitico (Auch. 2490!).

3. viscosa. — Tota viscosa-pubescent pallide virens.

Hab. ad Ouadi Esle peninsulae Sinaiticae (Ehr!), in Belutschia (Stocks).

Perigonium 3-4 lineas, fructus circ. tres lineas longi. Hujus speciei varietas foliis angulato-repandis **B. repanda** Willd. Sp. I, p. 22 ex India, Chinâ, etc. oriunda videtur.

Ar. Geogr. Hispania austro-orientalis, Nubia, Abyssinia, Arabia tropica, India, Nova Hollandia, P. B. Spei.

2. B. verticillata (Poir. Dict. V, p. 53) glabriuscula glaucescens basi suffruticosa, caulis albidis teretibus elatis dichotome opposite et divaricatim ramosissimis, foliis majusculis cordato-ovatis obtuse et late repando-sinuatibus, pedunculis filiformibus longis ex

axillis supremis et terminalibus 1-3 verticilos 3-5-floros subdistantes gerentibus, pedicellis capillaribus flore longioribus, perigonii infundibuliformis tubo brevissimo coarctato in limbum campanulatum abrupte ampliato, fructu adpressissime setuloso striato clavato tuberculis hemisphericis glabris sparsis et ad apicem truncatum in coronam approximatis obsito ♀. *B. repanda* Ky exs. non Willd.

Hab. in regno Mascate Arabiæ (Auch. 52501).

Folia sœpe 2-2 $\frac{1}{2}$ pollicem diam. lata, perigonum 1 $\frac{1}{2}$ lineam, fructus 2 lineas longi.

3. ***B. repens*** (L. Sp. 5) collo multicaulis, caulinis herbaceis diffusis dichotome ramosis alternatim ramulosis, foliis ovatis vel oblongis obtusis subsinuato-repandis vel lanceolato-linearibus, pedunculis axillaribus capillaribus umbellulas capituliformes 3-5-floras ferentibus paniculam foliosam formantibus, floribus minimis subsessilibus, perigonio brevi campanulato, fructu glanduloso oblongo-clavato quinquecostato et sulcato ♀. *B. sessiliflora* Steud.

Species valde variabilis cuius formæ frequentiores.

α. typica. — Adpresso papillari-puberula, folia ovata vel oblonga supra pallide virentia subtus albida. — Ic. Del. Eg. tab. 3, fig. 1. — *B. Maroccana* Ball.

Hab. in cultis Egypti superioris prope Syenem (Hussl) et in Oasi Dachel (Asch!), in Belutschia (Stocks!).

β. viscosa. — Folia ovato-oblonga, planta tota glanduloso-viscosa. — *B. ascendens* var. *pubescens* Schw. pl. exs.

Hab. in Egypto superiore Nubiâ finitima ad mare Rubrum (Schw!).

γ. diffusa. — Glabriuscula viridis, folia ovata majuscula concoloria. — *B. diffusa* L. Sp. 4. — *B. ascendens* Willd. Sp. I, p. 19.

Hab. in Egypto ad Belbeis (Ehr!), in Arabiâ Mascatensi (Auch. 52521).

δ. vulvariaefolia. — Adpresso et crispa hirta, folia parva anguste oblongo-elliptica subintegra. — *B. vulvariaefolia* Poir. Dict. V, p. 55.

Hab. in Egypto (Delile ex Poir.), in gypsaceis Persiæ australis ad Kotel Kumaredj (Haussk!).

Planta umbellulis capituliformibus, floribus subsessilibus, perigonio vix lineæ $\frac{1}{4}$ longo insignis; fructus sesquilineam circ. longus.

Ar. Geogr. Regnum Maroccanum, Nubia, Abyssinia, Arabia tropica, India.

4. ***B. elegans*** (Choisy in D.C. Prodr. XIII, p. 453) basi suffrutescens pluricaulis, caulinis glabris virgatis inferne foliosis superne in paniculam aphyllam dichotome et divaricatim ramosissimam nebulosam abeuntibus, foliis glancis sub lente minute puberulis ovato-oblongis, floribus ad dichotomias solitariis et terminalibus 1-3-nis pedicellis setaceis longissimis suffultis minimis, perigonio breviter campanulato, fructu oblongo-clavato quinquecostato inter costas sulcato et adpresso puberulo ♀. *B. rubicunda* Steud. in Schimp. exs.

B. stenophylla. — Folia anguste lineari-lanceolata.

Hab. in siccis Arabiae Mascatensis (Auch. 5250!), Belutschiae inferioris (Stocks!).

Folia 1-1½-pollicaria, 2-3 lineas in varietate tantum lata; species paniculæ aphyllæ divaricatum effusissimâ et pedicellis setaceis longissimis a *B. repentis* formis distincta.

Ar. Geogr. Nubia littoralis, Arabia tropica, regio Scinde Indiæ.

ORD. III. THYMELÆACEÆ.

(Meisn. d.c. Prod.)

Flores hermaphroditi vel rarius polygamo-dioici. Perigonium inferrum corollinum tubuloso-infundibuliforme, limbi 4 rarius 5-fidi lobis æstivatione imbricatis. Stamina perigonii tubo vel fauci inserta ejus lobis numero dupla. Squamulæ hypogynæ interdum obviæ sed in nostris nullæ. Ovarium liberum uniloculare uniovulatum ovulo pendulo anatropo, stylus unicus, stigma capitatum. Fructus siccus vel baccatus indehiscens semine unico pendulo, raphe linearis. Embryo rectus, radicula hilum spectans. Albumen nullum.

DAPHNE (L. Gen. p. 485).

Flores hermaphroditi. Perigonium quadrilobum infundibuliforme totum deciduum fauce esquamata. Antherae octo subsessiles dupli serie fauci inseritæ oblongæ subinclusæ. Stylus terminalis brevissimus, stigma capitatum. Baccæ carnosæ vel subsicca. Semen nucamentaceum testa crustacea. Albumen nullum. Embryo carnosus. — Frutices vel suffrutices.

* Folia herbacea decidua.

1. **D. Mezereum** (L. Sp. 509) fruticosa erecta stricte ramulosa, foliis hysteranthiis oblongo-lanceolatis obtusis in petiolum brevem attenuatis, fasciculis 3-5-floris lateralibus racemum interruptum infra gemmas foliorum formantibus, perigonii rosei lobis ovatis acutis tubum sericeum æquantibus, baccis ovatis rubris 5. Ic. Fl. Dan. tab. 268. — Rchb. Ic. Germ. tab. 556.

Hab. in regione subalpinâ et alpinâ, Macedonia in monte Peristeri (Orph!), Peloponnesus in Laconia (Sibth.), Pontus Lazicus ad Djimil 6000' (Bal!), Transcaucasia in Somchetia (Koch), Iberia Caucasica (Ower!).

Ar. Geogr. Europa borealis et media a Scandinaviâ ad Rossiam arcticam et ex Hispaniâ boreali et Italiâ ad regionem Danubialem et Dalmatiam, Sibiria.

2. D. Caucasica (Pall. Fl. Ross. I, p. 53) fruticosa erecta ramosa, ramis vetustis denudatis purpurascensibus novellis brevibus crenatis foliosis, foliis synanthiis oblongo-lanceolatis obtusis subapiculatis basi longe attenuatis sessilibus glabris, fasciculis terminalibus 5-10-floris inter folia suprema subsessilibus exinvolutatis, perigonii albi extus hirtelli lobis oblongis tubo duplo brevioribus, baccis nigris ȝ. *D. salicifolia* Lam. Dict. III, p. 438.

Hab. in Caucaso sept. (Guld.), Iberia (Stev! Hoh!), Cartiliniâ (Szov!), Imeretiâ et Mingrelia (Eichw!).

Frutex quadripedalis facie *Ligustri*, foliis saepe bipinnarib; flores magn. *Syringæ*.

Ar. Geogr. Ukrania.

•• Folia coriacea persistentia.

3. D. oleoides (Schreb. Dec. I, p. 13, tab. 7) fruticosa ramosissima, ramis vetustis denudatis 2-3-tome et crenatis dumulosis, ramulis rectis brevibus crenatis foliosis, foliis persistentibus coriaceis obovatis vel obovato-ellipticis obtusis saepe mucronulatis basi attenuatis subsessilibus utrinque adpresso et subtus copiosius pilosulis interdum glabratibus vel facie superiore glabris, floralibus terminalibus sessilibus fasciculatis albis ebracteatis, perigonii adpresso scripsi lobis patentibus triangulari-ovatis vel lanceolatis acutis tubi dimidiata longitudinem aequantibus superantibus vel brevioribus, ovario pubescente, baccâ exsuecâ ȝ. Ie. Jaub. et Sp. tab. 305. — *D. buxifolia* Vahl Symb. I, p. 29. — Fl. Græc. tab. 357. — Jaub. et Sp. tab. 306. — *D. glandulosa* Bert. Amen. p. 356. — Rehb. Ie. Germ. tab. 553. — *D. oleoides* ȝ. *brachyloba* et ȝ. *jasminea* Meisn. in p.c. Prodr. — *D. jasminca* Griseb. Spicil. II, p. 321 non Fl. Græc. — *D. alpina* Smith Prodr. non L.

Hab. in regione subalpinâ et alpinâ Græciae 3500'-6500' in Taygeto (Heldr!), Kyllene (Heldr!) Orph. Fl. Gr. exs. 35!), Delphi Eubeæ (Sprun! Heldr!), Velugo (Sprun!), Parnasso et Olympo Thessalo (Heldr!), Macedonia in Athone (Griseb!), Cretâ in montibus Sphacioticiis (Sieb! Raul!), Anatoliâ in Olympo Bithyno (Sibth., Boiss! Pichl!), Sipyllo (Auch!), Cadmo (Boiss!), Pisidâ et Ponio (Tchih!), montibus Galatiae (Noë!), Cappadocia in Argœo (Ky!), Cataoniae Berydaghi 8-9000' (Ball!), Armenia Turcicâ (Huet! Calvi Bourg!), Tauri Cilicico (Ky! Ball!), Libano (Boiss! Bl! Gaill!), Antilibano (Boiss!).

Species indumento plus minus raro, perigonii dentium cum tubo proportione maxime variabilis, varietates agere delimitandæ. Facie *D. alpinam* refert sed folia persistentia magis coriacea obsolete uninervia.

Ar. Geogr. Hispania australis, Corsica, Sardinia, Sicilia, Italia australis, Serbia, Africa borealis, regio Himalaica.

4. D. jasminea (Sibth. et Sm. Fl. Græc. IV, p. 50, tab. 358) fruticosa humilis glaucescenti-virens, ranis crassis depresso tortuosis ramosissimis, ramulis hornotinis brevibus intricatis demum subsquamoscentibus, foliis minimis oblongo-spathulatis carnosulis obtusis

mucronulatis basi attenuatâ sessilibus, floribus terminalibus 2-3-fasciculatis sessilibus ebracteatis pubescentibus extus purpurascensibus, limbi facie internâ albi lobis oblongo-lanceolatis tubo subtriplo brevioribus ♂. *D. microphylla* Meisn. in D.C. Prodr. XIV, p. 533.

Hab. in montibus Delphi Eubeæ (Sibth., Auch. 2518!), in monte Parnasso (Sibth.), in rupibus Kaki Scala Isthmi Corinthiaci prope Megaram (Boiss! Sart! Heldr!). Flor. videtur autumnum et iterum vere.

Suffrutex trunco brevi, facie *Salicis retusa*, folia 2-4 lineas tantum longa. Species ramis crassis tortuosis prostratis, foliis minimis, floribus extus roseis ab omnibus *D. oleoidis* formis distinctissima. Cl. Meisner certe iconem Flora Græca non vidit, nam si eam novisset pro *D. jasminæa* formas *D. oleoidis* non habuisset, veram *D. jasminæam* sub *D. microphyllæ* nomine describens.

5. ***D. sericea*** (Vahl Symb. Bot. I, p. 28) fruticosa, ramis elatis denudatis superne ramulosis, ramulis elongatis foliosis, foliis persistentibus elongatis ellipticis vel subspathulatis obtusis basi attenuatis subpetiolatis pallide viridibus supra glabris subtus adpresso pilosis margine subrevolutis, floribus sessilibus terminalibus capitatis et saepe insuper in axillis superioribus fasciculatis bracteatis, bracteis squamiformibus villosis perigonio multo brevioribus, perigonii extus dense sericeo-canis lobis ovatis obtusis roseis tubo subtriplo brevioribus, baccâ rubrâ ♂. Ic. Desf. Cor. tab. 9. — *D. collina* Smith Spicil. II, tab. 18. — Fl. Græc. tab. 359. — *D. australis* Cyr. pl. r.r. tab. 16. — *D. Neapolitana* Lodd. Bot. Cab. t. 719. — *D. oleæfolia* Lam. Dict. III, 440. — *D. buxifolia* Ledeb. Fl. Ross. III, p. 548! non Vahl.

Hab. in dumosis regionis montanæ, Cretæ 3-5000' (Sieb! Heldr! Raul!), Bithyniæ in regione inferiore ad Ghemlek, Brussa, Bolu, Sabandja (Griseb! Auch! Pest!), Lyciæ (Forbes), Pamphyliæ (Heldr!), Ciliciæ a regione inferiore ad 5-6000' (Ky! Bal!), ad Alexandrette (Ky. 2721) et Lataquié (Oliv.), Armenia Turciæ (C. Koch, Nordm!).

Ar. Geogr. Regnum Neapolitanum.

6. ***D. acuminata*** (Boiss. et Hoh. Diagn. Ser. I, 12, p. 103) fruticosa ramosa, ramulis hornotinis tenuibus rubellis minute velutinis crebre foliosis, foliis elongatis linear-lanceolatis glabris utrinque attenuatis sessilibus acutis mucronulatis præter nervum medium obsolete reticulato-venosis, floribus 5-7 brevissime pedunculatis corymbose capitellatis, bracteis squamiformibus minimis vel nullis, perigonii dense et breviter albo-velutini lobis ovatis albis tubo triplo brevioribus, baccâ aurantiacâ ♂. *D. mucronata* var. *Afghanica* Meisn. in D.C. Prodr. — *D. angustifolia* C. Koch Linn. XXII, p. 611.

Hab. in Armenia Rossicæ (C. Koch!), Kurdistaniæ Armeniæ ad Chana Putkie 7000' (Ky. Suppl. 571!), monte Gara Assyriæ (Ky. 333!), Kurdistaniæ Persicæ ad Mendeli (Noë!), in monte Fir Omar Gndrun (Haussk!), Persiæ australi in montibus Kuh Kilouyeh 6-10000' (Haussk!) et monte Kuh Delu (Ky. 551!), Persiæ boreali in prov. Aderbidjan (Szovl!) et in alpe Elamut jugi Elbrus (Auch. 5309!), Affghaniæ ad Chuggur (Griff!), Belutschia (Stocks!).

Folia 1-2½-pollicaria 2-3 linas lata, perigonum sub anthesi 3½ lineas longum.

Affinis *D. mucronatae* Royle Ill. p. 322, tab. 81 quæ floribus sessilibus roseis specificè distincta videtur.

7. *D. gnidiooides* (Jaub. et Sp. Ill. Or. IV, tab. 304) fruticosa ramosissima ramis vetustis denudatis erectis, hornotinis pruinosis strictiusculis crebre foliosis, foliis glaucescentibus pruinoso-puberulis demum minute pustulosis persistentibus rigidis basi sessili et apice attenuatis linear-lanceolatis pungenti-mucronatis, floralibus terminalibus et ad axillas supremas paucis fasciculatis breviter pedicellatis, bracteis 1-2 squamiformibus minimis hirsutis saepe oboletis, perigonii (ex auct. rosei) extus incano-sericei lobis patentibus ovatis obtusis tubo 2-3-plo brevioribus ȝ. *D. oleoides* Urv. Enum. p. 43. — Bot. Mag. tab. 1917 non L. Ieon mediocre sed quæ ob folia acuta huc potius quam ad *D. oleoidem* spectare videtur.

Hab. in insula Samos (Urv.), Cos (Auch. 2514), Lycia (herb. Zuccar.), Pamphyliâ (Oliv.), Cilicia Trachaeâ ad Anamour (Péron!).

Valde affinis *D. acuminatae* differt foliis longius mucronatis, floribus partim axillaribus roseis minoribus sub anthesi vix tres lineas longis.

8. *D. Gnidium* (L. Sp. 111) fruticosa erecta ramosa ramulis vetustis denudatis, hornotinis crebre foliosis pulverulento-velutinis, foliis sessilibus coriaceis persistentibus linear-lanceolatis cuspidato-acuminatis demum glabris, racemis ad axillas superiores terminalibusque simplicibus vel compositis paniculam brevem foliosam formantibus, pedunculis pedicellisque cano-tomentosis hisce patentibus perigonio brevioribus, perigonio adpresso sericeo cito deciduo albo lobis ovatis obtusis tubo subbrevioribus, baccâ ovalâ rubrâ dein nigrescente ȝ. Ic. Fl. Græc. tab. 356. — Rehb. Germ. t. 553. — *D. paniculata* Lam.

Hab. in fruticetis regionis inferioris Peloponnesi (Bory!), Atticae et Bœotiae (Sarti Orph!), Eubeæ (Heldr!), Anatoliae in Lycia et Cilicia (Tchih!). Fl. aest. et autumno.

Ar. Geogr. Regio mediterranea totius Europæ, Africa borealis, Insulæ Canarienses.

9. *D. glomerata* (Lam. Diet. III, p. 438) fruticosa humilis dumulosa a basi ramosa glaberrima, ramis crassiusculis inferne procumbentibus et radicantibus dein ascendentibus nudis cicatricosis simplicibus vel dichotomis apice confertim et rosulatim foliosis, foliis coriaceis synanthiis persistentibus oblongo-spathulatis obtusis basi attenuatis sessilibus, fasciculis 4-10-floris ex axillis supremis longis vel breviter pedunculatis in capitulum saepius arete congestis bracteis ovatis membranaceis stipatis, perigonii extus pallide rosei intus albidi glaberrimi lobis oblongis obtusiusculis tubo duplo brevioribus, ovario glabro ȝ. Ic. Jaub. et Sp. Ill. tab. 303. — *D. Imerica* C. Koch Linn. XXII, p. 613!

Hab. in regione alpinâ Pontus supra Trapezuntum (Tchih!), Pontus Lazicus supra Djamil 8'000' (Ball), in Armenia Turcicâ supra Koprubaciu 7-8'000'

(Huet!). in regione subalpinâ et alpinâ Caucasi 4500'-7000' (C.A. Mey.), Caucaesi orientalis 6-9000' (Rupr!), Iberia (M.B.), prov. Guriae et Karabagh (Szovl.).

Spithamea vel pedalis, folia 9-18 lineas longa florum capitulum involucrautia, perigonium sex lineas longum.

10. D. Laureola (L. Sp. 510) fruticosa, glaberrima, caule erecto dichotome ramoso ramis denudatis superne tantum foliosis, foliis magnis nitidis coriaceis persistentibus oblongo vel obovato-lanceolatis acutiusculis basi attenuatis sessilibus, floribus ad axillas superiores in racemos breves sessiles bracteatos subnutantes dispositis, bracteis membranaceis oblongis cucullatis caducis, pedicellis brevissimis, perigonii glabri viridis lobis ovatis acutis tubo triplo brevioribus, baccis nigris ȝ. Ic. Jacq. Austr. tab. 183. — Rehb. Germ. tab. 555.

Hab. in umbrosis regionis montanae Macedoniae (Friv!), Olympi Bithyni (ex Smithi Prodri.).

Ar. Geogr. Lusitania, Hispania, Anglia, Belgium, Gallia, Helvetia, Germania australis, Italia, Dalmatia, regio Danubialis, Africa borealis, insulae Azorice.

11. D. Pontica (L. Sp. 511) fruticosa glaberrima erecta ramis rubro-fuscis dichotomis superne foliosis, foliis majusculis coriaceis persistentibus obovato-lanceolatis acutiusculis basi cuneatis subsessilibus, floralibus valde diminutis sepe squamiformibus, pedunculis axillaribus plus minus elongatis subbifloris ebracteatis fructiferis mutantibus, pedicellis brevibus, perigonii glabri ochroleuco-virescentis lobis anguste lanceolatis attenuatis acutis patentibus vel deflexis tubo angusto vix brevioribus, ovario glabro, bacca nigra ȝ. Ic. Bot. Mag. tab. 1252.

Hab. in umbrosis et sylvaticis regionis inferioris, Thracia in sylva Belgrad prope Byzantium (Sibth., Barbey!). Bithynia in regione inferiore Olympi (Boiss!), ad Bolu (Pest!). Pontus (Tchih!), prope Trapezuntum (Bourg!), Pontus Lazicus ad Rhizé (Ball!), Transcaucasia in Imeretiâ, Mingrelia, Guria (Ledeb., Szovl.).

STELLERA (Gmel. in L. Acad. Amæn. 4, p. 299).

Flores hermaphroditi. Perigonum quadrilobum tubo cylindrico supra ovarium articulato parte inferiore persistente tumescente et fructum includente, superiore decidua, fauce esquamata, lobis patentibus. Stamina 8 duplice serie infra faucem inserta filamentis brevissimis. Ovarium apice barbulatum, stylus terminalis brevis, stigma ovoideum. Nucula calycis basi laxe tunicata, pericarpio crustaceo. Embryo albumine tenue inclusus. — Herbae perennes vel suffrutesc.

1. S Altaica (Thieb. in Pers. Ench. I, p. 436) basi suffruticosa, ramis numerosis erectis simplicibus virgatis crebre foliosis, foliis parce et crispule puberulis glabrisve subsessilibus elliptico-oblongis

utrinque attenuatis acutis, floribus in spicam terminalem ovato-oblongam dispositis, rachide glabrâ, floribus sessilibus erectis, perigonio infundibuliformi luteo glabro lobis quatuor oblongis tubo duplo brevioribus, ovario apice puberulo β . *Passerina racemosa* Wickstr. p. 320.
— Ledeb. Ic. Fl. Ross. tab. 374.

β . *minor*. — Rami abbreviati non pedales sed 4-5-pollicares, folia abbreviata 5-7 lineas tantum lata, spica depauperata, cæterum planta Sibiricæ similis.

Hab. in subalpinis Caneasi occidentalis ad rivulum Kassaut 5400'-6000' (G.A. Mey!), in Persiâ boreali orientali prope Schahrud et Tasch (Bge!). Specimina foliis angustioribus $1\frac{1}{2}$ lineas tantum latis.

Ar. Geogr. Sibiria Altaica.

2. **S. Lessertii** (Wikstr. Act. Holm. 1818, p. 346 sub *Passerini*) fruticosa ramis et trunco ramoso numerosis erectis tenuiter virgatis simplicibus vel ramulosis foliosis glabris superne pubescentibus, foliis subcoriaceis glabris vel plus minus hirtis ovato-oblongis subpetiolatis subtrinerviis et nervulosis acutiusculis vel obtusis, capitulo terminali mox in spicam oblongam vel cylindricam elongato, rachide tomentosâ, floribus sessilibus subhorizontalibus, perigonio infundibuliformi inferne albo-villosissimo superne adpresso pilosiusculo luteo lobis glabriusculis ovato-oblongis aentiusculis tubo triplo brevioribus, ovario apice plus minus hirto β . *Stellera Lessertii* C.A. Mey. I. eit. — Jaub. et Sp. Ill. tab. 301. — *Passerina Persica* Boiss. Diagn. Ser. I, 7, p. 85. — *Dais spicata* Endl. in Honigb. Sert. Cabul. p. 5, tab. 3. — *Stellera spicata* C.A. M. — *Stellera Griffithii* Meisn. in n.c. Prodr. XIV, p. 350 (forma foliis pilosiusculis).

Hab. in aridis totius Persie (Mich! Oliv! Auch. 2520 et 5287 a jugo Elbrusensi Persie bor. (Buhs! Ky. 210) ad prov. Khorassan (Bge!), monte Elvend 6000' et desertis Kermanschah, Ispahan, Tcheran (Haussk!), circa Schiraz (Ky. 321). Habeo tandem specimen ex Armeniâ Turcicâ prope Batum (Calv!).

β . *angustifolia*. — Folia anguste lanceolata. — *S. stachyoides* Schrenk Enum. 2, p. 16. — Ic. Jaub. et Sp. tab. 102. — *Passerina Ammodendron* Kar. et Kir.

Hab. in prov. Khorassan inter Schahrud et Nichapur (Bge!).

Nullum inter formas enumeratas discrimen specificum, forma typica cuius folia 3-6 lineas lata in varietatem β foliis 1-2 lineas tantum latis et saepè longioribus intermediis transit, flores omnium simillimi.

Ar. Geogr. Songaria, Sibiria Altaica.

LYGIA (Fasano Atti Ac. Neap. 1787, p. 235).

Flores hermaphroditi vel monoici. Perigonium urceolatum vel inflatibusiforme persistens tubo continuo, limbo quadrilido, fauce esquamata. Stamina octo summo tubo dupli serie inserta antheris

subsessilibus inclusis. Stylus brevis terminalis, stigma capitatum. Nucula calyce persistente inclusa. Albumen tenuissimum embryonem includens. — Herbæ annuae vel suffrutices floribus inconspicuis.

1. L. Passerina (L. Sp. 512 sub *Stellerd*) annua glabriuscula, caule erecto elongato simplici vel stricte ramoso ramisque tenuiter virgatis crebre foliosis, foliis linear-lanceolatis et linearibus sessilibus utrinque attenuatis, floribus hermaphroditis virentibus axillaribus sessilibus vel 2-3-nis spicas longas filiformes foliosas formantibus, bracteis fructu longioribus, perigonii adpresso pubescentis lobis ovatis obtusis erectis tubo duplo brevioribus, fructu pyriformi apice attenuato perigonio vix breviore ⊖. *Lygia Passerina* Fasan. loc. cit. — Ic. Rehb. Germ. 550. — *Passerina annua* Wickstr. — *Passerina arvensis* Lam.

Hab. in cultis et collibus regionis inferioris sed praesertim montane a Græcia ubi montana (Heldr! Orph!) et Macedonia (Friv!), ad Anatoliam! regiones Caucasicas! Syriam ad Aleppo (Ky!), Mesopotamiam (Ky. 353! Haussk!), Persiam hor. (Buhse! Bge!) et australiē ad Schiraz (Ky. 443!), Affghaniam (Griff!). Fl. Mai., Jun.

Ar. Geogr. Europa media et australis a Gallia et Germania media ad Ros-siam mediam, Sibiria Altaica, Africa borealis.

2. L. pubescens (Guss. Sic. Prodr. I, p. 466 sub *Passerina*) annua, pubescens, caule stricte ramoso ramisque elongatis tenuiter virgatis crebre foliosis, foliis linearibus vel lanceolatis sessilibus utrinque attenuatis, floribus axillaribus 1-3 hermaphroditis sessilibus virentibus tenuiter et longe spicatis, bracteis fructu non longioribus, perigonii adpresso pubescentis lobis ovatis obtusis erectis tubo duplo brevioribus, fructu pyriformi apice attenuato perigonio vix breviore ⊖. *Passerina arvensis* var. *pubescens* Ten. Syll. App. I, p. 365.

Hab. in argillosis et cultis, Syria prope Doumar (Gaill!), in monte Gebel Abiad deserti ad Palmyram (Bl!), Mesopotamia et Assyria (Oliv! Ky! Haussk!). Fl. Jul., Sept.

Forsan varietas præcedentis sed ex botanicis qui eam vivam observaverunt indumento et florescentia seriore distincta.

Ar. Geogr. Sicilia, Sardinia.

3. L. Aucheri (Meisn. in D.C. Prodr. XIV, p. 552 sub *Thymelæd*) perennis glabra, caulis e rhizomate suffrutescente numerosis erectis saepius simplicibus virgatis longissimis crebre et stricte foliosis, foliis linear-filiformibus acutis margine subrevolutis, floralibus sensim abbrevialis, floribus axillaribus 1-3 monoicis subsessilibus virentibus, bracteis ovatis minime basi villosis, perigonii adpresso pubescentis lobis in floribus masculis tubo tenui 2-3-plo brevioribus, in femineis tubum abbreviatum æquantibus, fructu pyriformi perigonio vix breviore ♂. *Stellera Hedgotis* Ehr. MSS. — *Lygia Amani* Ky. Sched.

Hab. in montanis siccis, Libanus (Ehr., Anch. 2532!), mons Torbol supra Tripoli (Boiss!), supra Sidonem (Bl!), vallis Barghoutiæ (Gaill!), prope

Becherre (Ky. 257!), in monte Amano (Ky. 407!), in collibus ad Antab Syriae hor. (Haussk!), Cappadocia ad Euphratem (Chesn.), Tauro Cilicie ad Gulek Boghas (Ky. 426! Bal!).

Caules cum spicis bi- vel tripedales, folia angustissima inferiora 6-10 lineas longa. Flores masculi eis *L. Passerinae* non longiores sed eorum tubus elongatus.

4. L. Cilicica (Meisn. in p.c. Prodr. XIV, p. 552 sub *Thymelaea*) glabra basi suffrutescens multicaulis, caulis ascendentibus tenuiter virgatis simplicibus brevibus foliatis, foliis breviter lanceolatis sessilibus utrinque attenuatis acutis, floralibus perigonio brevioribus, floribus in axillis superioribus 1-2 sessilibus moaocis breviter et laxiuscula spicatis, bracteis oblongis perigonio subbrevioribus, perigonii pubescentis lutei lobis rubellis in floribus masculis tubo duplo brevioribus in femineis tubo abbreviato paulo brevioribus, fructu elliptico perigonio breviore 5. *Lygia multicantis* Schott et Ky. in Ky. Taur. p. 351.

Hab. in regione subalpinâ Tauri Cilicie in rupestribus ad Karli Bogiaz 6500' (Ky! Bal!).

Cum spicâ hipollicari semipedalis, folia 3-4 lineas longa, flores eis *L. Passerinae* vix maiores, facies *Thymelaea thesioides*.

THYMELEA (Tourn. Inst. p. 394).

Flores hermaphroditi vel siepius polygamodioici. Perigonium infundibuliforme deciduum vel persistens tubo continuo limbo quadrifido, fave esquamata. Stamina octo summo tubo duplice serie inserta filamentis brevissimis. Stylus lateralis, stigma capitatum. Nucula calyce inclusa vel nuda. Albumen nullum. — Frutices.

* Calyx persistens. — *Chlamydanthus* C.A. Mey.

1. T. Tartonraira (L. 356 sub *Daphne*) fruticosa tota argenteo-sericea ramosissima ramis erectis inferne nudis cicatricosis superne cerebro foliosis, foliis obovatis vel spatulato-oblongis coriacibus crassis obtusiuseulis basi attenuatis persistentibus plurimerviis, floribus d' axillas aggregatis polygamodioicis rarius solitariis foliis saperatis, bracteis squamiformibus concavis perigonio multo brevioribus, perigonii extus sericei intus glabri lutei lobis tubo subbrevioribus 5. *Thymelaea Tartonraira* All. Ped. I, p. 133. — Ic. Fl. Graec. tab. 354. — Rehb. Germ. 532. — *Daphne tomentosa* Lam. Diet. III, p. 336, et *T. tomentosa* Endl. ex deser. — *Chlamydanthus Tartonraira* C.A. M.

Hab. in submontosis Atticae 8-1500' (Heldr. Herb. Norm. 435!, Orph. Fl. exs. 274!), Peloponnesi (Ky! Boiss!), Thraciae ad Pontum (Sibth.).

β. angustifolia Cuv. En. p. 42. — Folia anguste lanceolata et linearia, intermedia ad typum transit et vix varietas. — *D. argentea* Fl. Gr. tab. 355.

Hab. ad Corinthum et in Archipelago (Sibth., Urv!), Chios (Auch!), Rhodus (Bourg!), Creta (Sieb! Raul! Heldr!), Pamphylia (Bourg. 242!), insulæ Sinus Smyrnensis (Bal. 373!), Cyprus (Ky.).

Ar. Geogr. Hispania et Gallia australes, Italia merid., Sicilia, Sardinia, Corsica.

• • Calyx deciduus. — *Piptochlamys* C.A. Mey.

2. **T. hirsuta** (L. Sp. 559 sub *Passerina*) fruticosa fastigiata-ramosissima ramis tomentellis, ramulis nutantibus densissime et imbricatim foliosis, foliis minimis carnosulis ovatis vel oblongis obtusis enerviis supra convexis subtus concaviusculis junioribus tomentellis tandem glabris, floribus polygamо-dioicis 5-10 ad vel versus ramulorum apicem glomeratis ebracteatis, perigonii extus sericei intus glabri lutei campanulato-infundibuliformis lobis tubo subbrevioribus, fructu ovato glabro ♂. Ic. Fl. Græc. tab. 360. — Rchb. Germ. tab. 550. — *Th. hirsuta* Endl. — *Piptochlamys hirsuta* C.A. Mey.

Hab. in collibus regionis inferioris a maritimis ubi frequentior ad 1500', Græcia (Heldr! Orph. Fl. Gr. exs. 270!), Zacynthus (Marg!), Creta (Sieb! Heldr!), Hellespontus (Noë), Lycia (Forbes), Cilicia (Bal!), Palestina littoralis et Arabia petræa (Boiss!), Ægyptus inferior ad Alexandriam (Ehr., Gaill! Samar!).

Ar. Geogr. Europæ et Africæ borealis regio Mediterranea.

Species delendæ.

T. Orientalis Meisn. in D.C. Prodr. XIV, p. 557. — *Passerina Orientalis* Willd Sp. II, p. 43; species omnino fictitia ex synonymis duobus omnino obscuris, altero (Gronov. Orient. 126) ad plantam Orientalem et altero (Clus. Hist. I, p. 88) ad plantam Hispánicam spectantibus condita.

T. tomentosa Endl. Suppl. = *Daphne tomentosa* Lam. Dict. III, p. 436. Species obscura sed Hispánica! et nullo modo Orientalis.

DIARTHRON (Turez. Bull. Mosq. 1832, V, p. 204).

Flores hermaphroditi. Perigonium tubulosum supra ovarium constrictum articulatum parte inferiore persistente, limbi decidui quadridi lobis erectis conniventibus, fauce esquamata. Stamina 4 vel 8 et tunc biseriata, antheræ subsessiles. Stylus terminalis brevis, stigma parum incrassatum. Nucula calyce inclusa. Albumen parcum. — Herbæ annuae graciles.

1. **D. vesiculosum** (F. et M. Bull. Soc. Imp. Mosq. 1839, p. 170 sub *Passerina*) annum glabrum vel sub lente pilis paucis consper-

sum, caule erecto rigidulo dichotomie ramoso ramis filiformibus strictiusculis, foliis elliptico-lanceolatis vel linearibus acutiusculis, floribus minutis ebracteatis in racemos tenues terminales et oppositifolios dispositis, pedicellis brevissimis strictis clavatis, perigonii lutescentis glabri ad medium constricti tubo persistente membranaceo-vesiculoso carinato-octangulo, limbo subæquilongo deciduo lobis linearibus conniventibus obtusis, staminibus octo ♂. *D. vesiculosum* C.A. M. — *D. carinatum* Jaub. et Sp. Ill. Or. tab. 105.

Hab. in collibus argillosis Cappadociae ad Euphratem (Auch. 2534), Seid-khodji dist. Khoi prov. Aderbidjan Persiae (Szov!), prov. Khorassan prope Schahrud et Mesched (Bgel), Afghaniæ (Griff!), Belutschia (Stocks!).

Herba tenuis 1½-1-pedalis ramis erecto-patulis tenuissimis, folia 3-5 lineas longa, racemi 1-3-pollicares, perigonum sesquilineam longum.

Ar. Geogr. Songaria.

ORD. CIV. ELEAGNACEÆ.

(Lindl. Syst.)

Flores hermaphroditici interdum abortu dioici. Perigonum inferum 2-4-fiduīn aestivatione imbricatum. Discus perigonii tubum vestiens. Stamina perigonii fauci in se:ta ejus lacinis numero æqualia. Ovarium tubo perigonii inclusum liberum, ovulo unico erecto. Stylus simplex unistigmatosus. Fructus unilocularis monospermus tubo perigonii drupam mentiente induviatus, pericarpium crustaceum. Semen erectum anatropum, embryo erectus albumine angusto cinctus, radicula infera.

HIPPOPHAE (L. Gen. 4106).

Flores dioici spicati bracteati. Masc. in axillis bractearum ament lateralis sessiles. Perigonum diphylum. Stamina 4 filamentis brevisimis, antheræ biloculares. Fem. in axillis foliorum terminalium solitarii breviter pedicellati. Perigonum tubulosum limbo apice breviter bifido. Ovarium uniloculare uniovulatum cum stylo brevi perigonio inclusum, stigma elongatum longitudinaliter sulcatum. Fructus ex perigonio crescente succulento baccatus monopyremus. Pyrena dura altero latere leviter sulcata. Albumen nullum. Embryo rectus, radiculâ infera. — Frutices foliis alternis lepidotis.

1. H. rhamnoides (L. Sp. 1452) fruticosa elata patule ramosissima ramis demum apice spinescentibus lepidotis, foliis linearilanceolatis basi attenuatis subsessilibus obtusis supra viridibus subtus squamis lepidotis margine erosulis micantibus, amentis masculis

lateralibus saepe confertis, perigonii masculi phyllis subrotundis, baccâ pisiformi aurantiacâ ♂. Ic. Fl. Dan. tab. 265. — Rehb. Germ. t. 549.

Hab. in alveis arenosis rivorum. Hucusque observata Byzantii (Comi!), Ciliciâ (Tchih!), in Cappadociâ (Ball!), Ponto Lazico ad Rhizé (Ball!), Caucaso ad 6000' usque et Transcaucasiâ (Ledeb., Rupr!), Persiae borealis monte Elbrus (Ky!), Turkestania (Lehm.).

Ar. Geogr. Europa borealis et media ad Italiam medium Dalmatiam usque, Sibiria.

ELÆGNUS (L. Gen. 459).

Flores hermaphroditi vel masculi cum ovario abortivo. *Masc.* Perigonium campanulatum 4-6-fidum. Stamina tota quot perigonii dentes iisque alterna inclusa, filamentis fore totâ longitudine perigonio adnatis. *Antheræ biloculares* dorsifixæ, pistillum abortivum. *Herm.* Perigonium basi in tubum cylindricum attenuatum limbo 4-6-fido deciduo. Stamina ut in masculis. Discus glandulosus ad tubi faucem stylum cingens. Ovarium oblongum uniovulatum. Stylus elongatus, stigma clavatum. Fructus drupaceus, parte carnosâ et putamine sulkato ex perigonii tubo aucto et indurato ortis, pericarpium membranaceum monospermum styli basi apiculatum. Albumen parcissimum, embryo rectus, radicula infera. — Arbores vel frutices foliis alternis lepidotis vel stellatim pilosis.

1. E. hortensis (M.B. Taur. Cauc. II, p. 412) arborea inermis vel spinescens, foliis subtus vel utrinque argenteo-lepidotis in petiolum brevem attenuatis lanceolatis lanceolato-linearibus ellipticis vel ovatis, floribus axillaribus 1-3 pedicellis perigonii argentei tubo subæquilongis suffultis, limbi lobis triangularibus acutis intus flavis tubo brevioribus, stylo glabro, fructu minore secco insipido vel majore olivæformi sapido eduli ♂. *E. angustifolia* L. Sp. 476. — Ic. Fl. Graec. tab. 152. — Nees Ic. Germ. (Forma foliis angustioribus, ramis saepe spinescentibus, fructu non vel vix eduli). — *E. Orientalis* L. Mant. p. 41. Pall. Fl. Ross. tab. 5. (Forma saepius culta ramis inermibus, foliis latioribus, fructu majore eduli).

Hab. spontanea et culta in fere totâ ditione a Graeciâ! Macedoniâ! et Thraciâ! ad Caucasum et Transcaucasiam! Persiam! Turkestaniam! Mesopotamiam! Syriam! Egyptum! In australioribus v. gr. Persiâ australi usque ad 70°' secas torrentes (Hausk!), Affghania (Griff!), Belutschia (Stocks!). Turkestaniis *Dchida*, Persis *Sindschad* audit.

Omnis formae mire altera in alteram transeunt.

Ar. Geogr. Europeæ regio mediterranea (ubi forsitan introducta), Sibiria Uralensis et Altaica, China.

ORD. CV. LAURINEÆ.

(p.c. Fl. Pr. III, p. 361.)

Flores hermaphroditi vel unisexuales. Perigonium inferum 4-6-fidum vel partitum aestivatione imbricatum. Discus carnosus perigonii fundo adnatus. Stamina perigyna disci margini inserta laciniis perigonii opposita eis numero aequalia vel saepius 2-6-pla in series disposita. Antheræ filamento adnatae extrorsæ rarius externæ introrsæ biloculares vel loculis dissepimento transverso divisæ quadrilocellares locellis et loculis valvâ dehiscentibus. Ovarium liberum uniloculare, ovulo saepius unico ex apice loculi pendulo anatropo. Stylus simplex. Fructus monospermus baccatus vel drupaceus. Semen inversum, embryo orthotropus. Albumen nullum. — Arbores rarius suffrutices foliis alternis exstipulatis.

LAURUS (L. Gen. 503).

Flores dioici umbellati bracteis cœullatis decidui involuerati. Perigonium quadripartitum laciniis decidui. *Masc.* Stamina 8-12 bi vel triseriata filamentis versus medium biglandulosis. Antheræ introrsæ oblongæ biloculares operculo valvatim dehiscentes, connectivo lato. *Fem.* Stamina 4 sterilia filamentis apice utrinque glandulâ auctis hastæformibus. Ovarium uniovulatum ovulo pendulo. Stylus simplex, stigma obtusum. Fructus baccatus monospermus perigonii basi truncatâ suffultus. Semen e funiculo a basi fructûs ascendentem et superne testæ adnato pendulum. Albumen nullum. Cotyledones magnæ oleosæ, radicula inclusa supera. — Arbores sempervirentes foliis alternis.

1. *L. nobilis* (L. Sp. 529) arborea glabra, foliis breviter petiolatis supra nitidis subtus pallidioribus coriaceis oblongis vel lanceolatis acutis vel obtusiusculis margine subundulatis, umbellis 1-3-nis axillaribus breviter pedunculatis 4-6-floris, involueri squamis glabriusculis, perigonii laciniis obovatis obtusis, drupâ ellipticâ demum nigrâ §. Ic. Fl. Græc. tab. 365. — Nees Ic. Germ. — Rehb. Germ. 673.

Hab. in sylvaticis regionis sempervirentis 0-2-00' totius Graecie et ejus insularum (Heldr! Orph! Urv!), Macedonie et Thracie (Grisch!), totius Anatolie littoralis australis, occidentalis et septentrionalis ad Guriam!, et Imeretiam usque (Ruprl!), Syrie littoralis (Bl! Gaill!), Tauriae (Ledeb.).

Ar. Geogr. Regio mediterraea totius Europæ et Africæ borealis.

ORD. CVI. SANTALACEÆ.

(R. Br. Prodr. N. Holl. 350.)

Flores hermaphroditi vel polygamo-dioici. Perigonum superum 3-4-5-fidum æstivatione valvatum tubo a pedicello non distincto et ovario plus minus connato. Stamina 4-5 basi laciniarum perigonii inserta eisque opposita. Ovarium inferum uniloculare 2-4-ovulatum, ovula ad apicem placentæ centralis affixa pendula ad nucellum reducta. Stylus simplex, stigmata 1-4. Fructus nucamentaceus vel drupaceus monospermus, albumen carnosum, embryo intra apicem albuminis rectus brevis, radicula hilum spectans.

OSYRIS (L. Gen. 1401).

Flores dioici vel polygami. Perigonum 3-4-partitum lobis ovatis acutis caducis tubo in masc. gracili, in sem. obconico. Discus concavus lobis rotundatis partem superam et integrum perigonii vestiens. Stamina 3-4 ad margines disci inserta filamentis brevibus, antheræ biloculares. Ovarium carnosum inferum, stylus brevis, stigmata 3-4 lobis perigonii alterna. Placenta in cavitate minimâ ovarii brevissima apice 3-4 ovula gerens. Drupa sicca ovoideo-globosa vestigiis lobi-rum et disci coronata. Semen drupam implens, albumine copioso, embryone in parte superiori brevi, radiculâ superâ. — Frutices vel arbores ramis angulatis, floribus masculis in apice pedunculis con-gestis, femineis solitariis vel paucis.

1. O. alba (L. Sp. 1450) fruticosa sempervirens, foliis linearilanceolatis coriaceis acutis vel mucronatis, inflorescentiâ anguste et longe racemosâ, cymis in axi racemi lateralibus masculis multifloris floribus pedicellatis, femineis 1-paucifloris floribus subsessilibus, bracteis caducis, perigon trilobo, drupâ tandem rubrâ coriacea ȝ. Lam. Ic. tab. 802. — Rchb. Ic. Germ. tab. 518.

Hab. in fruticetis regionis inferioris, Græcia et ejus insulæ!, Macedonia et Thracia (Griseb!), Creta (Sieb!), Anatolia littoralis a Bithyniâ (Thirke!) ad Ciliciam (Balt), Syria littoralis (Lab! Gaill!), Palestina (Boiss!).

Ar. Geogr. Europa australis, Africa borealis.

Obs. Specimen junius et incompletum habeo *Osyridis* in montibus Hala Belutschiaæ alt. 5000' a cl. Stocks lectæ quæ prob. ad *O. arboream* Wallich referenda est.

COMANDRA (Nutt. Gen. I, p. 107).

Flores hermaphroditæ. Perigonium campanulatum ad medium quinquelobum. Discus epigynus perigonio adhaerens et eo dimidio brevior breviter et obtuse sinuato-quinquelobus lobis cum perigonii divisoriis alternis. Filamenta quinque perigonio inserta ejus lobis opposita basi araneoso-hirta. Antheræ ovatae loculis longitudinaliter dehiscentibus basi discretis. Ovarium inferum, stylus filiformis, stigma capitellatum, placenta filiformis erecta apice saepius triovulata. Nux subdrupacea ovoidea perigonii lobis coronata. Semen unicum inversum. Embryo in apice albuminis carnosí rectus. Radicula supera. — Suffruticuli foliis alternis penninerviis, cymâ corymbosâ terminali.

1. **C. elegans** (Rochel Banat. p. 36, tab. 4 sub *Thesio*) glabra, radice repente, caulinis numerosis inferne suffruticosis et stricte ramosis erectis totâ longitudine crebre foliosis, foliis oblongo-lanceolatis obtusiusculis basi sessili longe attenuatis, floribus 5-9 terminalibus corymbosis, perigonii laciniis ovato-triangularibus, nuce perigonii laciniis sessilibus coronatâ 5. Rchb. Hort. Bot. tab. 19. — *Hamiltonia elegans* Rchb. fil. Ic. Germ. fig. 1162.

Hab. in collibus arenosis Thraciae in monte Hæmo (Friv.), ad radices montis Rhodopes prope Stanimak (Janka!).

Ar. Geogr. Banatus, Serbia, Bulgaria, Moldavia.

THESIUM (L. Gen. 292).

Flores hermaphroditæ. Perigonium hypocraterimorphum vel infundibuliforme 4-5-fidum. Discus indistinctus. Filamenta basi cujusque lobi inserta basi barbata linearia. Antheræ oblongæ loculis longitudinaliter dehiscentibus. Ovarium inferum, stylus filiformis, stigma (in nostris) capitellatum, placenta filiformis erecta apice triovulata. Nux ellipsoidea caduca perigonii lobis saepius coronata. Semen unicum inversum. Embryo in apice albuminis carnosí rectus. Radicula supera. — Herbæ vel fruticuli floribus solitariis vel cymis axillaribus vel extraxillaribus. Plantæ subparasiticæ.

Specierum Orientalium dispositio.

SECT. I. CHRYSTHESIUM Jaub. et Sp. Ill. t. 104. — Perigonium hypocraterimorphum tubo elongato.

T. stelleroides, aureum.

SECT. II. EUTHESIUM A. D.C. — *Perigonum infundibuliforme tubo brevi vel campanulatum.* — Species omnes ditionis flores tribraeatos habent.

* Nux longitudinaliter nervosa nervulis inter nervos vel nullis vel obliquis nec transversis.

+ Pars supera perigonii fructiferi cum lobis erectis cylindrica nucem æquans.
T. Alpinum.

+ + Perigonii fructiferi lobî involuti in umbo nem nuce breviorem approximati. — Omnia perennia.

T. divaricatum, ramosum, Parnassi, brachyphyllum, Libanoticum, intermedium, montanum.

* * Nux inter nervos longitudinales nervulis obliquis et transversis reticulata,

+ Monocarpica.

T. compressum, heterophyllum, humile.

+ + Perennia.

T. Bergeri, procumbens, maritimum, impressum, Szovitsii, macranthum, Tauricolum.

SECT. I. CHRYSOTHESIUM.

1. **T. stelleroïdes** (Jaub. et Sp. Ill. Or. II, tab. 104) annum glabrum, caule erecto protero angulato stricte folioso superne stricte ramoso paniculato, foliis filiformibus subtrigonis obsolete uninerviis apiculatis, floribus subsessilibus bracteâ bracteolisque binis stipatis spicas terminales virgatas tandem elongatas formantibus, bracteâ bracteolisque flore multo brevioribus tructu sublongioribus subulatis, perigonii flavi tubulosò-hypocrateriformi tubo longo angusto parte ovario adhaerente lanato cæterum gibro lacinis angustè linearibus tubo 3-4-plto brevioribus, filamentis versns medias lacinias insertis, nuce ellipsoidea lanata perigonio maescente superata ⊖.

Hab. in Cappadociâ ad Euphratrem (Auch. 2554!).

Planta 4-4½-pedalis facie *Stellatae Passerinae*, folia 10-15 lineas longa, flor. 4-5 lineas longus.

2. **T. aureum** (Jaub. et Sp. Ill. Or. II, p. 104 et III, tab. 300) perenne glabrum sub lente parce asperulum e rhizomate suffrutescenti multicaule, caulis pumilis basi ramosis ramis erectis simpli-cibus crebre foliosis, foliis linearì-filiformibus uninerviis planis mar-

gine seabridulis mucronatis, floribus breviter pedicellatis terminalibus dense capitato-spicatis, bracteâ folio simili flore vix breviore bracteolis linearí-subulatis sublongiore, perigonii lutei glabri tubuloso-hypocraterimorphi lobis quinque oblongis erectis conniventibus concavis apice eneullato-incurvis et subrostellatis tubo quintuplo brevioribus, staminibus fauci insertis, fructu 5.

Hab. in Cappadociâ ad Euphratem (Auch. 2535!). Vid. in herb. Lessertiano.

Suffrutex cespitosus 3-5-pollicaris ramis novellis foliosis 2-3-pollicaribus, folia carnosula 4-7 lineas longa lineam dimidiâ lata, intima breviora, spicæ ovatae densinuscæ, perigonium 3½ lineas circiter longum, placenta in ovario filiformis tortuosa.

SECT. II. EUTHESIUM.

+ Nux longitudinaliter nervosa inter nervos non reticulata.

+ Pars supera perigonii fructiferi cylindrica cum lobis erectis nucem æquans.

3. **T. Alpinum** (L. Sp. 301) radice fusiformi, caulinis diffusis simplicibus vel parce ramosis in racemos elongatos unilaterales abeuntibus, foliis linearibus uninerviis, pedunculis unifloris abbreviatis erectis tandem subsecundis, bracteâ fructu multo longiore, bracteolis fructu sublongioribus, nuce breviter ellipsoideâ breviter stipitata, perigonio fructu tubuloso nucem æquante vel sublongiore in 4 lobos ovatos tandem conniventes et apice inflexos ad medium fisco ♀. Ic. Jacq. Austr. tab. 426. — Rehb. Germ. tab. 1131.

Hab. in montosis et alpinis siccis, in Ponti Lazieci valle Djimil (Bal!), in districtu Alagir Caucasi centrales ad moles glaciales Zei (Rupr!).

Ar. Geogr. Suecia meridionalis, Gallia, Germania, Hispania borealis, Italia, regio Danubialis, Bosnia, Serbia.

++ Perigonii fructiferi lobi involuti in umbonem nuce breviorem approximati.

4. **T. divaricatum** (Jan apud M. et K. Deutsch Fl. II, p. 286) radice descendente ramosâ multicauli, caulinis erectis vel ascendentibus elongatis pyramidatim paniculatis, paniculæ ramis subtrifloris patentibus vel divaricatis, foliis linearibus acutis uninerviis, bracteis margine sebris fructu brevioribus, nuce ellipsoideâ pedicello suo 2-3-plo longiore, perigonii lobis basi utrinque auriculâ dentatâ acutis, perigonio fructu involuto nuce triplo breviore ♀. Rehb. Ic. bot. tab. 436. — Rehb. fil. Ic. Germ. fig. 1155. — *T. hunifusum* Griseb. Spic. II, p. 323 non D.C. — *T. Byzantium* Alph. D.C. Prodr. XIV, p. 644!

Hab. in siccis regionis inferioris et montane, Macedonia et Thracia (Friv! Orph!), Byzantii (Oliv!), Anatolia borealis (Thirkel!), Lydia in monte Siplyo (Boiss! Bal. 379!), Tauras Cilicus (Ky!), Libanus ad Becherre (Ky. 934!), Tauria (Parr!).

Ar. Geogr. Gallia australis, Hispania, Italia, Carnia, Dalmatia, Hungaria, Serbia, Africa borealis.

5. **T. ramosum** (Hayne in Schrad. Journ. 1800, p. 30, tab. 7) radice fusiformi multicipiti, caulis erectis et ascendentibus a medio ramosis, ramis racemosis ramulis subunifloris erecto-patulis, foliis linearibus subtrinerviis, bracteis et bracteolis denticulato-scabris illâ fructu longiore hisce fructum æquantibus, nuce ellipsoideâ pedicello suo 3-4-plo longiore, perigonio fructus involuto nuce triplo breviore ♀. Rehb. Ic. Bot. fig. 646. — Rehb. fil. Ic. Germ. fig. 1154. *T. pratense* Ledeb. Fl. Ross. III, p. 542 non Ehrh. — *T. divaricatum* var. *expansum* Boiss. et Heldr. Diagn. Ser. II, 4, p. 81 forma ramulis elongatis magis expansis.

Hab. in regione alpinâ Ætoliae (Samar. et Guicc!), montibus Lyciae supra Elmali (Bourg!), Cappadociae monte Alidagh (Bal!), Ponto supra Trapezuntem et Baibut (Bourg!), Ponto Lazico supra Djimil 6300' (Bal!), monte Berydagh Cataoniae (Haussk!), Armeniaustrali in districtu Warto (Ky. Suppl. 577), Caucaso meridionali (Wilhm., Rehm!), Transcaucasiâ (Hoh.), Persia boreali prope Tabris et Siaret (Bge!).

β. *asperulum* Alph. d.c. Prodr. XIV, p. 664. — Totum ad foliorum margines, ramos, ramulos et bracteas asperulo-scabrum. — *T. asperulum* Boiss. et Buhse Aufz. p. 194.

Hab. in Armeniâ Rossica (Buhse!).

Species *T. divaricato* affinis differt ramulis floriferis strictioribus subunifloris, bractea fructu multo longiore. Perigonii lobi versus medium denticulo non semper obsiti.

Ar. Geogr. Austria, Dalmatia, Serbia, regio Danubialis, Rossia australis, Songaria.

6. **T. Parnassi** (Alph. d.c. Prodr. XIV, p. 643) radice tortuosâ stoloniferâ, caulis filiformibus pusillis ascendentibus simplicibus in racemos breves paucifloros abeuntibus, foliis anguste linearibus unnnerviis, ramulis floriferis brevibus erecto-patulis unifloris, bracteis margine lœvibus inferiore fructum superante, nuce ellipsoideâ pedicello suo atque perigonio quadruplo longiore ♀. *T. ramosum* Griseb. Spic. II, p. 223 non Hayne.

Hab. in regione superiore Parnassi supra Dakalia (Guicc. in Heldr. Fl. Gr. exs. 294!), Macedonia in monte Peristeri (Griseb.).

Caules cum racemo 2-3-pollicares. Inflorescentia et fructus *T. ramosi* a quo præter caules breves pusillos simplices et bracteas lœves nec scabridas radice stolones filiformes tenues edente specifice differre videtur. Magis affinis *T. brachyphyllum* differt caulis basi filiformibus et nudis, racemis laxis, bracteis lœvibus.

Ar. Geogr. Dalmatia, Herzegovina in monte Kom (Pancic), Aprutium Neapolitanum.

7. **T. brachyphyllum** (Boiss. Diagn. Ser. I, 5, p. 48) pallide virens sœpe scabridulum, rhizomate indurato ramoso stolonifero et radicante, caulis nanis prostratis et ascendentibus simplicibus in

racemos tenuis densifloros abeuntibus, ramulis floralibus brevissimis unifloris, foliis linearibus et oblongo-ellipticis basi attenuatis acutis obsolete uninerviis, bracteâ inferiore foliis simili nuce longiore cæteris brevibus, nuce ellipsoideâ pedicello demum incrassato-clavato longiore perigonio umboniformi brevissimo superatâ ♂.

Hab. in regione alpinâ montis Balkan Thracie (Pichler!). Tmolii Lydiae supra Bozdagh (Boiss!), montis Argæi Cappadocie (Ball!), Tauri Cilicie supra Bulghar Maaden (Ball!), Armeniae monte Bingöldagh 9800' (Ky. 388!) et in monte Techdagh (Huet! Calv!).

Caules 2-3-pollicares, folia 2-3 lineas longa. Planta collo indurata et suffrutescens.

8. T. Libanoticum (Ehrenb. MSS. — A. D.C. Prodr. XIV, p. 648) scabridum, rhizomate indurato pluricipiti, caulis brevissimis cespitibus simplicibus in spicas breves paucifloras abeuntibus, foliis turionum acicularibus cæteris linearibus acutis uninerviis, floribus in spicâ subsessilibus bracteâ fructu longiore bracteolisque brevioribus stipatis, nuce cum parte liberâ perigonii demum indurati ellipsoideâ lobis valde involutis breviter umbonantibus ♂.

Hab. in summo monte Makmel Libani (Ehrenb!), in cacuminibus inter Yamouny et Dimain (Bl!). Floriferum tantum vidi.

Cespites densi 1-2-pollicares, facie valde affinis *T. brachyphyllum*. An ab eo perigonii fructiferi parte liberâ induratâ (cl. Alph. D.C.) sat distinctum? Nucas non vidi.

9. T. intermedium (Schrad. Spic. Germ. p. 27) radice stoloniferâ, caulis erectis vel ascendentibus superne paniculatis, paniculae pyramidatae ramis divergentibus inferioribus trifloris, foliis linearilanceolatis acutis trinerviis, bracteis margine scabridis intermediâ fructu sublongiore, nuce oblongâ pedicello suo duplo longiore, perigonio fructu involuto nuce triplo breviore ♂. *T. linophyllum* Rehb. Ic. Germ. fig. 1160 et auct. plur. — Linn. ex parte.

Hab. in siccis Caucasi centralis (C.A. Mey.), Daghestaniae (Becker!), prov. Talysh (C.A. Mey.).

Ar. Geogr. Gallia, Germania, Hispania borealis, Italia, Dalmatia, regio Danubialis, Rossia media et australis.

10. T. montanum (Ehrh. herb. exs. № 2) radice descendente ramosâ apice multicauli, caulis erectis elatis a medio longe paniculatis, paniculae pyramidatae ramis tenuibus patentibus inferioribus dichotomis trifloris, foliis strictis lanceolatis longe acuminatis trinerviis, bracteis laeviusculis intermediâ fructum æquante vel sublongiore cæteris brevioribus, nuce ovoideâ pedicello 3-4-plo longiore, perigonio fructu involuto nuce triplo breviore ♂. Rehb. Ic. bot. tab. 452. — Rehb. fil. Ic. Germ. fig. 1161.

Hab. in regione sylvaticâ Olympi Thessali (Heldr!), in Olympo Bithyno (Auch. 2530!).

Ar. Geogr. Germania, Helvetia, Italia, Transylvania.

• Nux inter nervos longitudinales nervulis obliquis et transversis reticulata.
+ Monocarpica.

11. *T. compressum* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 104) monocarpicum, radice verticali, collo multicauli, caulis declinatis et intermedio erecto omnibus virgatis inferne parce ramosis ramorum decurrentiâ angulatis compressis in spicas longas tenuissimas laxas abeuntibus, foliis carnosulis enerviis linearibus acutis mucronatis, floribus minutis subsessilibus tubulosis præter infimos interdum 2-3-nos solitariis, bracteis squamiformibus lanceolatis acuminatis præter eas florum infimorum nuce multo brevioribus, nuce parvâ breviter pedicellatâ oblongâ albidâ inter nervos tenuiter reticulatâ perigonii lobis in umbonem angustum eâ quintuplo breviorem superatâ ♂.

Hab. in salsis hyeme inundatis Anatoliæ centralis, planities Iconii inter urbem et montem Karadagh (Heldr!), prata salsa ad occidentem Cæsareæ in Cappadociâ (Bal. 4039!).

Caules cum racemo 6-10-pollicari pedales, folia 6-10 lineas longa lineam lata, flores lineæ $\frac{3}{4}$ longi, nux vix lineam longa.

12. *T. heterophyllum* (Boiss. Ann. Sc. Nat. Ser. IV, vol. 2, p. 354) biennie, radice verticali, caulis e collo pluribus crassiusculis decumbentibus striato-sulcatis simplicibus vel parce et dichotome ramosis, ramulis brevissimis crassis confertim 3-5-floris in racemos laxisculos dispositis, foliis radicalibus rosulatis sessilibus ob longolanceolatis subenerviis, caulinis linearis-subfalcatis acutis margine denticulato-scabris, bracteâ bracteolisque lanceolatis scabridis illâ fructu multo longiore, nuculâ ellipticâ inter nervos reticulatâ stipite suo quadruplo longiore laciniis perigonii margine convolutis sed rectis conniventibus eâ dimidio brevioribus superatâ ♂. Ic. Tchih. Voy. 39.

Hab. in aridis calcareis regionis montanæ, Cappadocia ad Cæsaream (Bal!), Cataonia inter Albistan et Gurun (Tchih!) et prope Adiaman (Haussk!), Cilicia ad Gulek Boghaz (Bal!), Amanus supra Beilan 4000' (Ky. 63!).

β. Billardieri. — Folia radicalia diminuta ovata squamiformia. — *T. Billardieri* Boiss. Diagn. Ser. I, 5, p. 48. Vix varietas.

Hab. in Syriae collibus Elmalu (Bourg. 2441), in Phrygiæ monte Bulghas Dagh (Bal!), ad Tokat (Pestal!), in Antilibano (Labill!).

Caules 6-12-pollicares, folia radicalia sœpe pollicaria 3-4 lin. lata, flores ad ramulos florales brevissimos fasciculati, nux cum perigonio 2 lineas longa.

13. *T. humile* (Vahl Symb. 3, p. 43) monocarpicum, caulis e radice tenui verticali divisâ numerosis erectis et ascendentibus sœpius ramosis angulato-striatis cibbre foliosis in spicas elongatas tenues abeuntibus, foliis linearibus uninerviis serrulato-scabris inferiore longissimâ lateralibus fructum superantibus, floribus parvis, perigonii lobis breviter ovatis, nuculâ ellipsoideâ vix stipitatâ lobis perigonii

brevissime umbonatâ inter nervos longitudinales plus minus elevatim reticulatâ ♂. Guss. Pl. rar. tab. 200, fig. 2. — Rehb. Ic. bot. tab. 705. — *T. Græcum* Zucc. fasc. II, p. 14 (forma nuculis obsolete reticulatis).

Hab. in collibus et cultis regionis inferioris, Græcia et ejus insulæ (Sprun! Heldr! Urv.), Anatolia occidentalis (Bal!), Syria littoralis (Gailli!), Palestina (Boiss!), Egyptus inferior (Oliv., Samar!), Oases Egypti medie (Schweinf! Aschers!).

Forma nuculis obsolete et vix reticulatis occurrit cum typu mixta non in Græciâ tantum sed in Dalmatiâ, Italiaâ, etc. et specimenia omnino intermedia reperiuntur.

Ar. Geogr. Hispania australis, Sardinia, Italia australis, Sicilia, Africa borealis.

+ + Perennia.

14. **T. Bergeri** (Zuccar. pl. nov. fasc. 2, p. 16 et herb!) perenne glabrum, caulinis e radice verticali durâ numerosis gracilibus decumbentibus et erectis angulatis stricte ramosis in racemos longos tenues foliosos abeuntibus, foliis linearibus elongatis acutis uninerviis, racemi ramulis præter inferiores trifloros unifloris brevibus erecto-patalis, bracteâ foliis simili flore et fructu multo longiore, bracteolis multo brevioribus sed florem superantibus, floribus minimis, perigonii lobis breviter ovalis, nuce breviter stipitatâ ellipsoïdeâ inter nervos reticulatâ lobis in umbone brevissimum concretis superatâ ♀. — *T. Græcum* Boiss. et Sprun. Diagn. Ser. I, p. 47 non Zucc.

Hab. in collibus regionis inferioris et præsertim montanae, Bœotia (Berg.), Attica in montibus Parnes et Pateras (Boiss! Heldr!), Malevo Laconiae (Orph!), Creta in monte Idâ (Heldr!) et prov. Sphakia (Raul!), Rhodus (Bourg!), Lydia circa Smyrnam (Boiss!), Phrygia (Bal!), Lycia (Heldr!), Cilicia (Ky! Bal!), Cataonia in monte Sofdaghi (Haussk!), Syria bor. in Amano (Ky. 1454!) et Cassio (Boiss!), Syria littoralis (Bl! Gailli), Libanis ad Ed-en (Boiss!).

Caules 6-12-pollicares tenues, folia et bractea inferior elongata lineam dimidiâ lata, racemis sœpe semipedales, nucula vix sesquilinea longa. Species umbone perigoniali nuculæ minimo bracteisque longis insignis, a *T. humili* radice perenni distincta.

15. **T. procumbens** (C.A. Mey. Enum. p. 40) perenne, radice..., caulinis filiformibus brevibus procumbentibus ramosis, racemis brevibus laxis subsecundis, foliis linearibus elongatis uninerviis, pedunculis unifloris fructu sublongioribus, bracteis linearibus acutis laevibus inferiore nuce longiore lateralibus eam subæquantibus, nuce longuscule stipitatâ ovatâ inter nervos reticulatâ lobis perigonii ovatis in umbone eâ triplo breviorem approximatis superatâ ♀.

Hab. in herbidis montosis et subalpinis Caucaasi occidentalis ad rivulos Kisch Malka et Kassaut 2000'-6000' (C.A. Mey!), in Caucaasi orientalis Daghestania ad Schoralo versus fontes fluvii Samur 6200' (Rupr!).

Caules cum racemis 3-5-pollicares, folia 8-12 lineas longa vix lineam lata, nuces lineam longæ. A *T. Bergeri* cui nucularum reticulatione accedit caulinis pusillis, racemis brevibus, nuculâ longius umbonatâ distinctum. .

16. T. maritimum (C.A. Mey. Verz. p. 40) glaucum, caulis ex rhizomate ramoso indurato pluribus procumbentibus ramosis striatis, foliis carnosulis enerviis acutis margine scabridis, racemi elongati ramulis brevissimis crassis 3-1-floris, bracteis et bracteolis subæqualibus scabridis inferioribus fructu longioribus superioribus eo æquilongis, nuculâ globoso-ellipsoideâ inter nervos transverse reticulatâ lobis perigonii subinvolutis subconniventibus apice sub-infelix eæ æquilongis superatâ ȝ.

Hab. in littore arenoso maris Caspii prope Baku (C.A. Mey!).

Caules 4-8-pollicares, folia turionum abbreviata, caulina 6-9 lineas longa lineam lata, nux linea vix longior.

17. T. impressum (Steud. in Ky. pl. Alepp. Kurd. 1843) glabrum læviuscum basi suffrutescens multicaule, caulis erectis et ascendentibus crassiusculis angulato-striatis ramosis flexuosis in paniculam angustum abeuntibus, foliis lanceolato-linearibus et linearibus carnosulis 4-3-nerviis subtus parallele nervulosis, ramulis floriferis rigidis crassis inferioribus dichotomis superioribus abbreviatis 3-1-floris. bracteis lævibus inferiore fructum æquante vel superante cæteris brevioribus, nuce crassiusculâ oblongâ inter nervos transverse reticulatâ stipite brevissimo crasso suffultâ perigonio subæquilongo ultra medium in lacinias lineares margine convolutas rectiuseculas apice inflexas fisco superatâ ȝ.

Hab. in Armenia australi prope Karput (Noë 863!), in Cataonie monte Berytdagh 8000' (Bal!), in monte Gara Kurdistanie (Ky. 411!).

Caules 6-10-pollicares, folia 1-1½-pollicaria 1½-3 lineas lata, nux absque perigonio 2 lineas longa.

β. Kotschyanum. — Folia magis acuminata basi magis attenuata, bracteæ margine scabridulæ præsertim in floribus inferioribus racemi elongatæ, nux major 2½ lineas longa. — *T. Kotschyanum* Boiss. Diagn. Ser. I, 7, p. 86.

Hab. in Kurdistanie Persice montibus Avroman et Schahu 8000' (Haussk!), in Persiâ prope Ispahan (Auch. 5292! Bge!), ad Persepolin et in monte Sabst Buschom prope Schiraz (Ky. 416! et 818!), in montibus Bachtiaricis et jugis Sawers, Kuh Eschker 9-12000' (Haussk!).

Species affinis *T. maritimo* sed major magis suffruticosa, lieviscula, folia latiora subtus parallele multinervulosa, nux duplo et triplo major.

18. T. Szovitsii (Alph. D.C. Prodr. XIV, p. 649) perenne glabrum læve, radice descendente obliquâ fibrosâ, caulis erectis elatis firmis a medio paniculato-recurvatis, foliis anguste linearibus elongatis carnosulis enerviis, paniculæ ramis rigidulis erecto-patulis dichotome et divaricatim 3-5-floris cum flore in dichotomiâ, bracteâ florem æquante aut vix superante bracteolis minoribus, perigonii lobis linearie-longis parte integrâ brevioribus, nuce ellipsoideâ inter nervos reticulatâ breviter stipitatâ lobis crispato-involutis eâ dimidio brevioribus superatâ ȝ.

Hab. in Armeniâ Rossicâ (C.A. Mey! Buhse! sub *T. multicauli*), in prov. Aderbidjan (Szov. exs. 5761 sub *T. multicauli*).

Caules pedales et longiores, folia 1-2-pollicaria, facies *T. macranthi* cuius nux ignota sed quod perigonio multo profundius fisco et floribus 3 nec 2 lineas longis inflorescentiâque differt.

19. T. macranthum (Fenzl Flora 1843, p. 401) glabrum, caulis e radice lignosâ validâ erectis elatis firmis virgatis a medio ramosis, ramis tenuibus erecto-patulis in racemos laxifloros abeuntibus, ramulis floralibus brevissimis 1-3-floris, foliis elongatis anguste linearibus acutis enerviis, bracteâ bracteolisque lanceolatis illâ florem aequante, perigonii elongati laciñiis lineari-oblongis parte integrâ 2-3-plo longioribus margine inflexis apice incurvis, staminibus perigonium æquantibus, stylo perigonium subsuperante, nuce.... ♀.

Hab. in Mesopotamiâ inter Orfa et Suerek (Ky. 122!).

Caules plures 1-2-pedales sulcato-striati stricti. Folia 1-1½-pollicaria, flores tres lineas longi. Species floribus magnis insignis ob fructus ignotos quoad affinitatem subdubia.

20. T. Tauricum (Boiss. et Haussk. in Haussk. exs. 1863) glabrum pallide virens, suffruticosum caudice crasso lignoso multi-cauli, caulis tenuiter virgatis angulato-striatis simplicibus vel stricte ramosis in spicas longas laxiusculas gracillimas subunilaterales abeuntibus, foliis carnosulis subulatis elongatis mucronulatis, pedunculis unifloris strictis brevissimis vel subnullis, bracteis lanceolatis inferiore fructu breviore lateralibus multo brevioribus, perigonii floriferi laciñiis breviter ovatis, nuce ellipsoideâ breviter stipitata inter nerves reticulatâ perigonii lobis in umbonem eâ triplo breviorum et angustiorem involutis coronatâ ♂.

Hab. in montibus Berytdagh, Baltandagh et Begdaghi Cataoniae 7000 (Haussk!).

Caules cum racemis 4-6-pollicaribus fere pedales, folia pollicem et amplius longa vix dimidiata lineam lata, nux lineâ unâ parum longior.

ORD. CVII. LORANTHACEÆ.

(Lindl. Veg. Kingd.)

Flores unisexuales vel hermaphroditi. Calycis tubus ovario adnatus, limbus (in nostris) 4-3-fidus vel nullus. Corolla vel nulla vel petalis calycis tubo adnatis constans. Stamina tot quot petala et eis opposita. Ovarium inferum uniloculare uniovulatum, ovulo ad nucellum reducto erecto orthotropo; stylus simplex. Fructus baccatus. Semen pericarpio adhaerens. Embryo unicus (vel saepè plures) in albuminis carnosí superficiali cavitate nidulans inversus, radicula ab hilo remota. —

Frutices parasitici foliis oppositis vel nullis. — Ordo juxta *Santalaceas* ob ovuli structuram collocatus sed *Cornaceis* quoque affinis, inter *Monochlamydeas* corollæ præsentia frequentia anomalus.

VISCUM (L. Gen. 4403).

Flores (in nostris) dioici. Mase.: Calyx limbo 4-fido. Corolla nulla. Antheræ 4 sessiles mediis lobis calycis adnatae multicellulosæ multi-porosæ. Fem.: Calycis cum ovario coaliti limbus brevissimus obscure quadridentatus. Petala 4 carnosa squamiformia summo calyci inserta valvata. Fructus sessilis bacciformis monospermus. Stigma sessile obtusum. — Sufrutices parasitici ramis 2-3-tomis articulatis, floribus virentibus, foliis oppositis.

1. V. album (L. Sp. 1451) ab insertione ramosissimum, ramis teretibus, foliis coriacidis carnosis oblongis obtusis 5-6-nerviis basi attenuatis, floribus ad dichotomias superiores terminalibusque 5-6-nis sessilibus capitatis, baccis globosis albis 5.

Hab. ad arbores varias parasitica, Graecie regio montana in Taygeto ad Abetes (Boiss! Heldr!), mons Athos ad Abetes 4500'-5000' (Griseb.), circa Smyrnan ad Nymphio in Pomeceis (Bal!), Taurum Cilicie et monte Musmeneudagh Cappadociae ad *Abetum Cilicicum* (Ky! Bal!), Tauriæ et prov. Caucasicis (M.B.), Persia boreali in prov. Ghilan (Aucht!) et prope Asterabad ad Carpium (Bge!). Turcice *Bury Aghagü*.

Ar. Geogr. Europa omnis a Scandinavia meridionali ad Rossiam medianam, Siberia Uralensis et Amurensis, Japonia, Africa borealis.

2. V. cruciatum (Sieb. pl. Palest, exs. -- Boiss. Voy. Esp. p. 274) ab insertione ramosissimum ramis teretibus, foliis coriacidis carnosis oblongis obtusis obscure trinerviis basi attenuatis, floribus 2-4 ad dichotomias superiores terminalibusque in capitulum breviter pedunculatum dispositis, baccis globosis rubellis 5. *V. Orientale* D.C. Prodr. quoad pl. Palest. non Willd.

Hab. ad *Oleas* parasitica in Palestina circa Hierosolymam (Sieb! Boiss! Rot. J. Gail!), circa Naploune (Boiss!).

A *V. albo* et *V. laxe* foliis minoribus trinerviis, baccis pedunculatis coloris diversi, a *V. Orientale* Indico foliis non elongatis, baccis non purpureis, etc. distinctum.

Ar. Geogr. Hispania australis.

ARCEUTHOBIUM (M.B. Fl. Taur. Cauc. III, p. 629. — *Razumowskia* Hoffm.)

Flores dioici. Corolla nulla. Mase.: Fl. sessiles. Calyx involucello bipartito cinctus. 3-4-partitus lobis antheriferis, antheris patulis adnatis

unilocellatis rimâ dehiscentibus. Fem: Fl. pedicellati. Calyx ovario adnatus, ovatus compressus apice rimâ pertusus hians. Stigma sessile, ovulum inferum. Baccâ sicca elastice a pedicello sejuncta et semen projiciens. — Suffrutex parasiticus glaber dichotome et opposite ramosissimus aphyllus facie *Salicornia*.

1. **A. Oxycedri** (d.c. Fl. Fr. IV, p. 274 sub *Visco*) ramosissimum glabrum, ramulis oppositis dichotomisque strictis tereti-compressis breviter articulatis, foliis ad squamas triangulares in vaginulas connatas reductis, floribus masculis in ultimo articulo sessilibus, femineis 1-2 in axillis supremis terminalibusque brevissime pedicellatis, baccis ovoides ♀. Ic. Clus. Hist. 39. — *A. Oxycedri* M.B. Taur. Cauc. III, p. 629. — *Razumowskia Caucasica* Hofst.

Hab. in *Juniperis* (*Oxycedro, rufescens, drupaceâ*) parasitica in regione inferiori et montanâ, Gracia in Parnassi regione abietinâ (Guicci!), Macedonia australis (Friv!), Taurus Cilicus prope Gülek (Ky! Bal!), mons Akkerdaghs Tauri Cataonicus (Haussk!), Libanus supra Eden (Bl!), Tauria (Stev!), Iberia Caucasica (M.B.), Persia borealis in jugo Elbrus prope Radkhan in *Jun. Sabinâ* (Buhse!).

Ar. Geogr. Lusitania, Hispania, Gallia austro-orientalis, Istria, Dalmatia, Serbia, Africa borealis.

LORANTHUS (L. Gen. 443).

Flores hermaphroditi rarius dioici. Calycis tubus ovatus vel turbinate cum ovario connatus, limbus brevissimus truncatus vel dentatus. Petala 4-8 summo tubo calycis inserta libera vel basi concreta, valvata. Stamina petalis opposita, filamenta eis basi vel ad medium adnata. Antheræ biloculares. Stylus filiformis, stigma simplex. Baccâ ovata vel turbinata unilocularis monosperma. — Suffrutes parasitici oppositifolii facie *Visci*.

SECT. I. EULORANTHUS d.c. Prodr. IV, p. 286. — Petala 4-6 æqualia distincta. Antheræ ovato-oblongæ erectæ vel adnatae.

1. **L. Europæus** (L. Sp. 1672) glaber opposite ramosissimus ramis teretibus, foliis oppositis petiolatis oblongo-spathulatis obtusis parce venosis basi attenuatis, floribus parvis sæpius dioicis oppositis ochroleucis in spicas simplices laxas breves terminales dispositis, petalis sex linearibus recurvis calyce sesquilonioribus, baccâ ovatâ flavâ ♂. Ic. Jacq. Austr. tab. 30.

Hab. parasiticus ad *Castaneam vescam* in monte Malevo Laconiae 3000' (Heldr! Orph. Fl. Gr. exs. 290!), ad *Quercus* in Arcadiâ (Bory), montibus Eubeæ Delphi et Athos Macedoniae (Sibth.). Ad *Castaneam* in monte Timolo Lydiæ circa Bozdagh (Boiss! Bal. 387!), Kurdistanâ (Barré de Lancy!), Ciliciæ regio montana (Ky! Bal!), Taurus Cataonicus (Haussk!), montes

Avroman et Pendjavin Kurdistaniæ Persicæ 6000'-8000' (Haussk!). Græcis *Lazina*.

Abortu sæpius dioicus sed in speciminiibus fructiferis stamina saltem rudimentaria interdum reperi.

Ar. Geogr. Austria, Italia, Sicilia, regio Danubialis, Serbia, Bosnia, Rossia orientalis.

2. L. Grewinkii (Boiss. et Buhse Aufz. p. 106) glaber, opposite ramosissimus ramis brevioribus junioribus subquadrangulis, foliis oppositis anguste oblongis vel lineari-lanceolatis in petiolum brevem attenuatis vel inferioribus oblongis subtus obscure trinerviis, racemis brevissimis terminalibus 3-5-floris folio multo brevioribus, floribus minutis hermaphroditis oppositis ochroleucis, petalis sex oblongis carnosis calyce brevioribus, baccis ovatis e sicco rubellis ♂.

Hab. in Persiæ borealis jugo Elbrus prope pagum Tchehordeh ad *Pyrum communem* parasiticus (Buhse!), in Persiæ austro-occidentalis montibus Awiheng, Ataschke, Sawers, Nur, Eschker ad *Pyrum, Acerem Monspessulanum, Amygdalos* spinosos parasiticus 7000'-8000' (Haussk!).

Facies *Rhamni* vel *Pruni*, rami eis *L. Europæ* tenuiores et breviores, folia triplo minora et angustiora, flores hermaphroditæ minores et petala breviora.

SECT. II. SYMPHYANTHUS D.C. Prodr. I. cit. — Petala 4-6 æqualia in tubum fere ad medium usque connata. Antheræ ad apicem filamenti erectæ.

3. L. Acaciæ (Zuccar. Abh. Münch. Acad. III, p. 249, tab. 2, flg. 3) ramis teretibus elongatis, foliis suboppositis breviter petiolatis coriacieis glaucis glabris obovatis obtusissimis integris, floribus breviter pedicellatis 2-7 in fasciculos axillares dispositis supremis sæpe solitariis, calyce truncato, corollâ kermesinâ tubulosâ ultrâ medium in lacinias anguste lineari-spathulatas demum revolutas 4-5-fidâ, ad sinus quinquegibbâ, staminibus corollæ et stylo subæquilongis ♂.

Hab. ad *Rhamni* et *Zizyphi* species, *Acacia* species parasiticus, in Palestinâ ad Jericho (Roth, Boiss! Ky. 129!), circa Hebron (Roth).

Rami furcati nodulosi bipedales et longiores, folia 15-16 lineas longa 8-12 lata, flores fere biplicares, species elegans. *L. gibbosulus* A. Rich. ex Nubia et Abyssiniâ nostro affinis differre videtur foliis angustioribus elliptico-lanceolatis et calycis sub apice constricti limbo sublobato.

Ar. Geogr. Nubia.

ORD. CVIII. CYTINACEÆ.

(Lindl. Veg. Kingd. p. 91.)

Flores unisexuales rarius hermaphroditæ. Perigonum superum vel semisuperum phyllis vel lobis imbricatis rarius valvatis. Antheræ

numerose columnæ centrali affixæ 1-2-loculares. Ovarium inferum vel semiinferum uniloculare placentis parietalibus. Ovula numerosissima orthotropa vel anatropa. Fructus baccatus polyspermus. Albumen cellulosum embryonem minutum foveans. — Plantæ parasitæ.

CYTINUS (L. Gen. 1252).

Flores moneici vel dioici monochlamydei. Perigonium tubulosocampanulatum lobis 4-8 aestivatione imbricatis. *Masc.* Bracteæ binæ ad basin perigonii, rarius nullæ. Antheræ 8-10 ad apicem columnæ centralis sessiles uniseriatæ in capitulum connatæ biloculares extorsæ. *Fem.* Bracteæ binæ versus medium ovarium eo adnatae vel nullæ. Ovarium inferum uniloculare placentis parietalibus. Stylus cylindraceus, stigma globosum, ovula numerosissima orthotropa. Fructus baccatus pulposus unilocularis indehiscens polyspermus. Semina minuta albuminosa. — Herbæ radicibus parasiticæ, caulis brevibus carnosis imbricatim squamosis aphyllis.

1. C. Hypocistis (L. Syst. 826) monoicus, caulis brevibus cylindrico-clavatis solitariis vel aggregatis squamis ovatis vel ellipticis obtusis brevissime denticulato-ciliatis obsitis, floribus subsessilibus 5-10 in spicas fere capituliformes dispositis femineis lateralibus masculis centralibus, bracteis binis, perigonio pubescente quadri-lobo lobis ovato-ellipticis erectis, antheris 8, stigmate capitato 8-10-sulcato ♀. Ic. Fl. Græc. tab. 938. — Rchb. Germ. tab. 540.

Hab. ad radices *Cistorum* parasiticus in regione inferiore Græciae et ejus insularum (Sibth., Heldr!), Cretæ (Sibth.), Macedoniae, Thraciae et Bithyniae (Griseb!), Anatoliae meridionalis (Heldr!), Syriæ littoralis (Gaiill!), Cypri (Ky. 751!).

Planta squamis flavescentibus vel roseis vel kermesinis, floribus albidis vel flavidis.

Ar. Geogr. Regio Mediterranea Europæ et Africæ borealis, insulæ Canarienses.

PIOSTYLES (Guillem. Ann. Sc. Nat. Ser. II, vol. 2, p. 21.

Flores dioici. Bracteæ et perigonii phylla non distincta pluriseriata basi latâ persistentia imbricata. *Masc.* Antheræ numerosæ circa basin columnæ capituliformis stipitatæ vel sessilis 1-3-seriatim verticillatae uniloculares, pollen subglobosum viscosum. *Fem.* Ovarium inferum vel semiinferum uniloculare, stylus nullus vel unicus brevis, stigma sessile, ovula numerosissima placentis parietalibus undique inserta vel ad totam superficiem internam ovarii sparsa, orthotropa vel anatropa. *Bacca* carnosa polysperma, semina minuta testâ reticulatâ. — Plantæ minutæ globosæ ad rimas ramulorum parasiticæ.

1. P. Haussknechtii (Boiss. Archiv. Bibl. Univ. Genev. 1866, XXV, p. 259) bracteis et perianthii phyllis sub 14 exterioribus late elliptico-oblongis interioribus latoribus spathulatis erectis omnibus integris, columnâ antherifera capitata, antheris biseriatis, ovario semi-supero, stigmo umbonato, disco epigyno nullo, ovulis per totam ovarii cavitatem sparsis ⊖.

Hab. in ramulis variarum specierum spinosarum *Astragali* ad axillas petiolorum parasiticus. v. gr. in *Astr. rhodochroo*, *leioclado*, *rhodosemio*, *racemuloso*, *myriacanthon*, *Susiano*, etc. in regione alpinâ 60°-10000', mons Berytdagh Cataoniæ, Schahu Kurdistaniæ Persicæ, Eschker, Nur, Sawers, Parrow et montes Bachtiaris Persicæ austro-occidentalis (Haussk!).

Flores e cupulâ margine papyraceâ et lobato-lacerâ erumpentes globosi 2 lineas diametro lati, antherarum series superior ad basin capitulo sessilis, inferior annulum brevem crassum marginans, bacea globosa, semina elliptica subobliqua, testa cellulosa. Facie affinis speciebus congeneribus ex Americâ mediaâ et australi ad *Papilionaceas* varias quoque parasiticis.

ORD. CIX. BALANOPHORACEÆ.

(Rich. Ann. Mus. Par. 1822, p. 404.)

Flores polygami vel dioici in formam spadicis confertissimi. Perigonium imperfectum vel oligophyllum. Stamen (in nostro) unicum. Ovarium inferum vel semiinferum (in nostro) uniloculare, ovulo unico pendulo. Fructus indehiscens nucamentaceus vel subdrupaceus, semen solitarium fructus cavitatem implens, embryo minutus lateralis, albumen oleosum. — Herbæ parasitæ.

CYNOMORIUM (Micheli Gen. p. 47, tab. 42).

Flores polygami in eâdem spicâ permixti basi bracteolati. *Masc.* Perigonium phyllis 4-5 linearispathulatis constans. Stamen solitarius posticum filamento cylindrico, antherâ biloculari, loculis bilocellatis, ovarii rudimentum oblongo-clavatum canaliculatum in canaliculo filamentum antheræ excipiens. *Fem.* Perigonium phyllis 1-5 semisuperis vel omnino superis linearis-clavatis ovario adnatis. Ovarium sessile vel substipitatum stylo canaliculato et stigmate obtuso terminatum uniloculare uniovulatum, ovulum unicum funiculo breve ex apice loculi pendulum hemitropum. *Herm.* Rari plus minus imperfecti sed fertiles. Fructus ovario similis sed auctus nucamentaceus, pericarpio tenui subcoriaceo testae connato. Semen subglobosum, embryo intra albumen lateralis indivisus. — Planta carnosa phalloidea in radicibus parasitica.

1. C. coccineum (L. Sp. 1875), scapis e rhizomate ramoso

teretibus carnosis squamis sparsis unguiformibus tandem deciduis vestitis apice in spadicem clavatum crassum longum abeuntibus, cymis dichotomis multifloris in stratum spadicis confluentibus bracteis primariis peltatis in spicam dispositis prius imbricatis tandem remotis et denique deciduis basi suffultis, floribus masculis saepius in receptaculo communi sessilibus, feminis et hermaphroditis saepius secus cymas dispositis ♀. Ie. Rich. Mem. Mus. 8, tab. 21. — J. D. Hook. Trans Linn. Soc. 1856, tab 1. — *Fungus coccineus Melitensis typhoides* Bocc. Sic. p. 80, tab. 43.

Hab. in arenosis maritimis ad radices multarum plantarum vel perennium vel annuarum parasiticum, Creta! Palestina, Arabia petrea, Egyptus (Del.), planities Galala deserti Aegyptiaco-Arabici (Schweinf!), Persia orientalis (Bge!).

Ar. Geogr. Insulæ Canarienses, Hispania, Italia, Sardinia, Sicilia, Africa borealis, Songaria.

ORD. CX. ARISTOLOCHIACEÆ.

(Lindl. Veg. Kingd.)

Flores hermaphroditi monochlamydei. Perigonum superum indivisum irregulare vel regulare trifidum. Stamina definita libera et apice ovarii inserta vel cum stylo connata. Antheræ biloculares longitudinaliter dehiscentes. Ovarium inferum vel semiinferum sexloculare placentis centralibus multiovulatis; ovula anatropa raphe crassâ notata. Fructus capsularis plurilocularis. Embryo parvus ad basin albuminis carnosí vel carnei. — Herbæ (in nostris) foliis alternis simplicibus.

ASARUM (L. Gen. 589).

Perigonum campanulato-urceolatum inferne ovario adhaerens limbo trilobo aestivatione valvari. Stamina 12 summo ovario inserta, filamentis brevibus, antheris liberis apice concrecti appendiculatis. Ovarium semiinferum sexloculare, ovula plurima loculorum angulo centrali biseriatim affixa. Styli 6 in columnam connata, stigmata 6. Capsula coriacea calyce persistente coronata sexlocularis irregulariter dehiscens. Semina cymbiformia transverse rugulosa raphe strophiiformi carnosâ instructa. — Herbæ caulisibus repentibus radiculas edentibus.

1. **A. Europeum** (L. Sp. 633) pubescens, ramis novellis brevibus erectis, foliis hyeme persistentibus longe petiolatis geminis reniformibus obtusis vel retusis profunde cordatis basi squamis ternis

membranaceis caducis suffultis, petiolo unifloro inter folia bina terminali, perigonii utrinque villosi ad medium usque fissi lobis ovatis in cuspide brevem contractis ȝ. Fl. Dan. tab. 633. — Rchb. Ic. Germ. tab. 668.

Hab. in umbrosis partis occidentalis regionis, Laconia et circa Byzantium (Smith Prodr.).

β. Caucasicum Duchartre in D.C. Prodr. XV, p. 423. — Folia non obtusissima sed apice magis proiecta. — *A. Ibericum* Steven MSS.

Hab. in Ponto Lazico prope Rhizé (Bal!), Imeretia et Iberia (Nordm., Stev.).

Varicetas β (quam cæterum floriferam non vidi) foliis minus obtusis *A. Caucasicum* referunt, formam analogam ex Helvetiâ habeo.

Ar. Geogr. Europa media a Sueciâ et Rossia ad Italiam, Dalmatiam, regionem Danubialem, Sibiria Uralensis.

ARISTOLOCHIA (L. Gen. 1022).

Perigonium tubulosum ovario adnatum tubo supra ovarium ventricoso, apice saepius in ligulam dilatatum. Antheræ sex sessiles facie internâ columnae stylinæ adnatæ. Stylus brevis lobis (in nostris) sex stigmatosis patentibus. Ovarium inferum sexloculare ovulis plurimis angulo centrali loculorum biseriatim insertis. Capsula coriacea apice cicatrice limbi decidui umbilicata septicide basi vel apice sexvalvis. Semina uniseriata strophiolata. — Herbae interdum volubiles.

Specierum Orientalium distributio.

* Perigonium apice clausum poro sublaterali tantum apertum.

A. microstoma.

** Perigonium superne et postice in limbum (labium) unilaterale vel periphæricum sed basi non cordato-biauriculatum expansum.

+ Caules scandentes vel subscandentes.

A. sempervirens, altissima.

++ Caules non scandentes.

× Perigonii tubus rectus vel vix curvatus.

A. parvifolia, Tournefortii, Billardieri, Clematitis, rotunda, longa, pallida.

× × Perigonii tubus valde sigmoideo-curvatus.

A. incisa, hirta.

*** Perigonium superne et postice in labium unilaterale basi cordato-biauriculatum expansum.

× Perigonii tubus rectus.

A. auricularia.

× × Perigonii tubus valde sigmoideo-curvatus.

A. Olivieri, Maurorum, pæcylantha, Pontica, Iberica, Cretica.

* Perigonum apice clausum poro dehiscens.

1. **A. microstoma** (Boiss. Diagn. Ser. I, 5, p. 50) caulis ex caudice crasso cylindrico pluribus parte subterraneâ filiformibus parte exsertâ tenuibus erectis foliosis pumilis, foliis parvis sub lente parce hirtulis petiolo longioribus ovato-rotundis retusis basi profunde cordatis sinu subquadrato auriculis rotundatis, floribus parvis in axillis insimis caulinum solitariis pedunculo reflexo ovario æquilongo suffultis, perigonio obsolete puberulo fusco supra basin ventricosam ovatam subconstricto dein sensim ampliato subincurvo clavato obtusissimo oblique et obsolete subrostrato et infra apicem ostiolo circulari aperto, capsulâ sphæricâ ♀. *A. Clematitis* N° 5 Wheler It. Ic. mala. — *A. parafolia* Sprn. exs. non Sibth.

Hab. in lapidosis siccis et agris incultis regionis montanæ Græciæ, Hymettus et Corydalus (Sprn! Boiss! Heldr! Orph. Fl. exs. 128!), Citheron (Boiss!), Parnassus supra Rachova (Orph!).

Radix aromatica digiti crassitie, caules subramosi parte exsertâ 3-5-plicares, folia $\frac{1}{2}$ -1-pollicem longa et lata, perigonum cum ovario 9-10 lineas longum limbo labii loco in ampullam clavatum clausam oblique ostiolo minutissimo tantum dehiscentem dilatato. Species distinctissima.

** Perigonum in labium unilaterale vel periphericum basi non cordatum expansum.

+ Caules scandentes.

2. **A. sempervirens** (L. Sp. 961) radice... caulis tenuibus brevibus sexangularibus ramulosis debilibus prostratis vel subscandentibus, foliis persistentibus parvis subcoriaceis glabris lucidis breviter petiolatis oblongis basi sinu angusto cordatis auriculis rotundatis, floribus axillaribus solitariis pedunculis filiformibus petiolo triplo longioribus suffultis, perigonii lutescentis purpureo-vittati utriculo ovoideo, tubo incurvo superne ampliato in labium ovatum obtusiusculum eo non multo brevius abeunte, capsulâ.... ♀. Ic. Fl. Græc. tab. 934. — Bot. Mag. tab. 1116. — *Pistolochia altera sempervirens* Clus. pl. rar. 260 Ic. — Morison Sect. XII, tab. 17, fig. 16.

Hab. ad sepes et in muris Cretæ (Siebt!), ad Males, Malaxa, Askypous (Heldr!). Fl. Julio.

Planta *A. altissimâ* humilior, tenuior, caules pedales, folia minora 10-15 lineas longa, flores paulo minores.

3. **A. altissima** (Desf. Fl. Atl. II, p. 324, tab. 249) radice fibrosâ, caule inferne suffruticoso superne volubili alte scandente

sexangulari ramoso, foliis persistentibus lucidis et glabris breviter petiolatis ovato-oblongis acutis vel obtusiusculis basi sinu angusto cordatis auriculis rotundatis approximatis, floribus parvis axillaribus solitariis pedunculo petiolo triplo longiore suffultis, perigonii fusco-purpurei parce et minute hirtuli utriculo globoso, tubo curvato sensim ampliato in labium ovatum vel oblongum obtusiusculum tubo multo brevius, capsula oblongo-clavata pendulâ ♀.

Hab. in saxosis et ad fuentes umbras regionis inferioris et montanae, Grecia in monte Malevo Laconiae 3000' (Orph. Fl. exs. 129!) et in insula Cypro (Ky. 736! sub *A. sempervirenti*), Libani regio inferior (Bl! Gaill!), Antilibanus ad Rascheya (Ky. 736! sub *A. sempervirenti*).

Caulis adultus 12-18-pedalis digito sœpe crassior, folia bipinnata, perigonii pollicem fere longum. Affinis *A. sempervirenti* sed caules multo elatiiores et crassiores parte inferiore fruticosi, folia triplo majora, corolla non major sed fusca labio tubi proportione breviore.

Ar. Geogr. Sicilia, Africa borealis.

++ Caules non scandentes.

4. *A. parvifolia* (Sibth. et Sm. Fl. Gr. X, p. 27, tab. 936) caulis ex caudice fusiformi subterraneo numerosis filiformibus parte exsertâ procumbentibus vel ascendentibus foliosis pumilis, foliis parvis brevissime petiolatis sub lente subtus puberulis ovatis sinu subquadrato cordatis auriculis rotundatis, floribus ex axillis superiorebus brevissime pedunculatis solitariis, perigonii extus glabriuscui tubo suprâ partem ventricosam subconstricto vix incurvo in labium superne puberulum linear-lanceolatum tubo triplo longius abeunte, capsula sphærica ♀.

Hab. in Syriâ boreali ad Ain el Beitha (Haussk!), circa Berythum (Bl!), circa Hierosolymam (Roth exs. 4!), insula Cypro (Ky. 188!).

Caules semipedales, folia 5-7 lineas longa, perigonii tubus 7-8 lineas, labium 18 lineas longum. In Florâ Græcâ circa Athenas et in Graeciâ indicatur unde hanc speciem nunquam vidi, probabiliter ex commutatione cum *A. microstoma* cuius folia similia. Perigonii labium rectum video sed ex Florâ Græcâ iconie et descriptione tandem tortum fit.

5. *A. Tournefortii* (Jaub. et Sp. Ill. II, tab. 128) caulis ex caudice crasso oblongo vel fusiformi numerosis filiformibus parte subterraneâ nudis parte exsertâ prostratis vel ascendentibus foliosis pumilis, foliis parvis glabris brevissime petiolatis ovalis retusis basi cordatis sinu subquadrato auriculis rotundatis, floribus parvis axillaribus solitariis pedunculo ovario vix æquilongo suffultis, perigonii glabri olivacei tubo supra basin globoso-ventricosam angulo obtuso incurvo in labium linear-lanceolatum acutiusculum eo vix longius abeunte, capsula sphærica ♀. *A. longa folio minore subrotundo flore tenuissimo* Tourn. Cor. p. 8.

Hab. in insula Chios (Oliv.), in eadem insula prope Vrontados (Orph. Fl. exs. 708! sub *A. macroglossa*), in Troade (Oliv.), in herbidis insulae Kilsali sinu Smyrnensis (Bal. exs. 343 sub *A. parvifolia*), in Lyciâ prope Elmali (Bourg!).

Rami tenues 6-10-pollicares, folia majora 8-10 lineas longa, perigonium 10-12 lineas longum.

β. macroglossa. — Flores majores labio magis dilatato. — *A. macroglossa* Jaub. et Sp. Ill. tab. 127.

Hab. in insula Chios (Oliv. et Brug.).

Species foliis et habitu valde affinis *A. parvifoliae* Sibth. a qua foliis glabris et perigonii labio dimidio breviore superne non hirtulo differre videtur.

6. **A. Billardieri** (Jaub. et Sp. Ill. I, tab. 100) caulis erectis gracilibus simplicibus puberulis, foliis breviter petiolatis ovatis obtusis basi sinu ampio cordato-excisis sub lente utrinque papilloso-hispidulis, floribus axillaribus solitariis pedunculo hirsuto ovario hispido sublongiore suffultis, perigonio violaceo hirtello supra basin ventricosam incurvo superne vix ampliato in labium oblongo-linguaeforme obtusum æquilongum abeunte ♀.

Hab. in Syria loco non indicato (Labill. in herb. Webb et Delessert), in fissuris rupium ad Ain el Beitha inter Alexandrette et Aleppo (Haussk.).

Planta interdum pedalis, folia 1-1½-pollicaria, flores pollicem longi. Affinis *A. parvifoliae* et *Tournefortii*, procerior, folia duplo majora, ovarium et perigonium multo hirsutiora.

7. **A. Clematitis** (L. Sp. 1364) radice repente, caulis sulcatis simplicibus vel inferne parce ramosis erectis subflexuosis, foliis magnis longiuscula petiolatis deltoideo-ovatis obtusiusculis basi sinu profundo cordatis auriculis obtusis plus minus convergentibus, floribus ad axillas fasciculatis et breviter pedunculatis parvis, perigonii glabri ochroleuci utriculo globoso, tubo cylindrico rectiusculo labio ovato-lanceolato subbreviore, capsula magna ovala pendula ♀. Ic. Clus. Hist. II, p. 71. — Fl. Dan. tab. 1235. — Rchb. Ic. Germ. tab. 669.

Hab. in ruderatis, vineis et agrorum marginibus partis septentrionalis ditonis, Macedonia (Orph!), insula Tassos (Griseb.), Thracia et Byzantium (Sibth.), Anatolia tota borealis a Bithynia (Thirke!), ad Pontum Lazio (Bal!), Tauria et Caucasus (Ledeb., C.A. Mey.), Iberia (Eichw.).

Ar. Geogr. Europa media et australis ab Anglia et Belgio ad Rossiam medium.

8. **A. rotunda** (L. Sp. 1364) caudice globoso vel ovato, caulis erectis sulcatis simplicibus vel ramosis, foliis magnis glabris vel subtus parce puberulis subsessilibus ovatis obtusis basi sinu clauso profundo cordatis auriculis obtusis convergentibus, floribus in axillis solitariis pedunculo brevi sed petiolo longiore suffultis, perigonii lutei intus purpureo-striati extus breviter hirtuli utriculo basiliari ovato, tubo recto superne subampliato in labium oblongum obtusum vel retusum eo subæquilongum abeunte, capsula globosa reflexa ♀. Rchb. Ic. Germ. f. 1342.

Hab. in siccis et incultis regionis inferioris et montanæ, Græcia et ejus insulae (Sprun! Heldr! Orph!), Macedonia (Friv!), Bithynia (Tchili!).

Ar. Geogr. Regio mediterranea Europe et Africæ borealis.

9. **A. longa** (L. Sp. 1364) caudice verticali oblongo-fusiformi tandem longe cylindrico, caulis simplicibus vel saepius ramosis flexuosis saepius minute puberulis, foliis petiolo pedunculo sublongiore suffultis ovato-cordatis obtusis vel retusis basi sinu profundo lato cordatis auriculis rotundatis, floribus in axillis solitariis pedunculo ovarium subæquante suffultis, perigonii virenti-brunnei glabri utrículo subgloboso tubo longo recto superne subampliato in labium breviter lanceolatum eo duplo brevius abeunti, capsula ovoideâ ♀. Ic. Rehb. Germ. fig. 1344. — *A. pallida* β. *elongata* et *A. Attica*. Orph. Fl. Gr. exs!

Hab. ad muros, rupes in siccis regionis inferioris et montanæ Græciae, in Arcadiâ (Heldr!), monte Malevo (Orph!), Parnes Atticæ (Orph!), insulâ Andros (Sart!).

Ar. Geogr. Regio mediterranea Europæ et Africæ borealis, Madera, insulæ Canarienses.

10. **A. pallida** (Willd. Sp. IV, p. 162) caudice tuberoso globoso, caulis a basi subterraneâ filiformi nudâ ascendentibus subsimpli-cibus glabris, foliis petiolo pedunculo subæquilongo suffultis triangularibus vel orbicularibus obtusis vel retusis basi profunde cordatis auriculis rotundatis, floribus in axillis solitariis pedunculo ovario subbreviore suffultis, perigonii glabri virenti-flavidi purpureo-striati et intus maculati utriculo ovoideo tubo recto sensim dilatato in labium oblongo-lanceolatum obtusum eo subdimidio brevius abeunte, capsula oblongâ ♀. Ic. W. K. Pl. rar. Hung. tab. 240. — Fl. Græca tab. 936. — Rehb. Germ, fig. 1343. — *A. lutea* Desf. Choix, p. 12, tab. 8.

Hab. in regione montanâ, mons Delphi Eubœa (Sprun!), Byzantii (Counany!), Bithynia (Thirke), Asia minor (Auch. 5293!), mons Almadagh prope Ouchak Phrygiae (Bal!).

Ab *A. rotundâ* cui radice congruit et quâcum saepè confusa foliis petiolatis statim distinguenda, magis affinis *A. longæ* et specimenina exsiccata radice destituta saepè inter eas species incerta, caules in *A. pallidâ* tamen humiliores simplices, folia magis abbreviata, perigonii color diversus, labium magis obtusum.

Ar. Geogr. Helvetia meridionalis, Italia, Austria, Dalmatia, regio Danubialis.

11. **A. incisa** (Duchart. in p.c. Prodr. XV, p. 490) caulis gracilibus humilibus simplicibus vel ramosis, foliis breviter petiolatis parvis ovato-cordatis obtusis mucronulatis basi sinu obtuso profunde excisis auriculis obtusis incurvis utrinque verruculosis et subtus breviter hirtulis, floribus axillaribus solitariis pedunculo hirtulo deflexo petiolo longiore, perigonii atropurpurei glabri tubo supra utriculum subglobosum signoideo valde incurvo, labio ovato-oblongo obtusissimo subquadrato-bilobo tubo subdupo longiore ♀.

Hab. in Anatoliâ loco non indicato (Auch. Mus. Par. N° 2508 sub *A. Pistolochia*!).

Vix semipedalis, folia vix pollicem longa. A precedentibus perigonii tubo valde sigmoideo et labio lato apice bilobo distincta sed ex uno et incompleto specimine descripta et ulterius investiganda.

12. A. hirta (Linn. Sp. 961) undique hirta, radice crassâ elongatâ lignosâ, caulis erectis striatis flexuosis simplicibus, foliis majusculis petiolatis trianguli-vel subdeltoido-hastatis obtusis basi sinu profundo excisis auriculis latis obtusis intus incurvis, floribus pedunculo petiolo longiore apice recurvo suffultis axillaribus solitariis, perigonii atropurpurei extus et intus crebre hirti utriculo ovato, tubo stigmideo-recurvo superne sensim et parum ampliato, labio peripherico ovato concavo obtuso, capsula oblongâ hirtulâ ♀. *A. longa subhirsuta folio oblongo flore maximo* Tourn. Coroll. et Voy. 1, pag. 386. Ic. — Ic. Fl. Græc. tab. 937. — Jaub. et Sp. III. tab. 97.

Hab. in umbrosis, ad sepes, incultis derelictis, insula Chios (Orph. Fl. Gr. exs. 707!), Bithynia ad Broussam (Thirket Barbey!), ad Smyrnam (Boiss! Balt!), Phrygia (Balt!), Pamphylia et Pisidia (Heldr!), Cyprus (Sibth!).

Pedalis et procerior, affinis varietati latifoliae *A. Maurorum* a qua distinguitur indumento longiore et copioso, foliis obtusis, perigonio hirto et inprinisis labio peripherico basi obtuso integro vel retuso nec cordato-biauriculato.

*** Perigonium in labium unilaterale basi cordato-biauriculatum expansum.

13. A. auricularia (Boiss. Diagn. Ser. I, 5, p. 49) pilis crispa-
lis basi subincrassatis aspera, caudice...., caulis punilis parte
subterranea filiformibus nudis ad basin partis exsertae foliosae saepe
ramulosis, foliis intinis ad squamas sessiles ovatas reductis ceteris
breviter petiolatis ovato-cordatis acutiusculis secus nervum medium
longitudinaliter subplicatis margine erosulis basi sinu profundo
cordatis auriculis rotundatis, floribus majusculis in axillis supremis
solitariis 2-5 approximatis inferioribus longiuscule ceteris breviter
pedunculatis, perigonii sparsim hirti brunneo-nigri utriculo ovato,
tubo cylindrico recto vix ampliato utriculo 5-6-plo longiore, labio
late cordato-ovato obtusiusculo erosulo basi rotundatim biauriculato,
capsula ovato-sphaericâ deflexâ ♀.

Hab. in cultis Anatolie australis, Caria interior (Boiss! Pinard!), Lycia
prope Elmalu (Bourg!).

Caules sub anthesi 2-3-pollicares fructiferi vix semipedales, folia tandem
1245 lineas longa, flores sesquipollicares, perigonii labium 8-9 lineas longum
basi 7 lineas latum, capsula nuce paulo minor.

14. A. Olivieri (Collegno in Boiss. Diagn. Ser. I; 5, p. 50)
sub lente minute asperula, radice...., caule erecto crassiuseculo striato
flexuoso alternatim ramosissimo ramis erecto-patulis, foliis breviter
petiolatis trianguli-ovatis obtusiusculis mucronulatis basi sinu late
aperto cordatis auriculis obtusis intus curvatis, floribus axillaribus
solitariis pedunculo petiolo triplo longiore suffultis medioribus,
perigonii olivacei vel lividi extus glabriusculi utriculo ovato, tubo

valde incurvo superne subampliato, labio ovato-oblongo acuto basi obsolete auriculato, capsula.... ♀. *A. Bruguieri* Jaub. et Sp. Ill. II, tab. 129.

Hab. in Mesopotamiâ vel Assyriâ (Oliv. et Brug!).

Pedalis et elatior, folia sesquipollimicem longa et basi lata, flores fere pollicem longi. A formis *A. Maurorum* caule crasso flexuoso-sinuoso et valde ramoso, floribus minoribus, perigonio basi brevissime nec valde biauriculato distincta.

15. *A. Maurorum* (L. Sp. p. 1303) breviter asperulo-hirtella, caudice cylindrico tandem elongato, caulis a basi subterraneâ denu-datâ erectis simplicibus vel parce ramosis crebre foliosis foliis versus apicem sensim diminutis, foliis breviter petiolatis lanceolatis vel linearibus hastatis acutis basi profunde excisis sinu basilari amplio auriculis divergentibus basi angustatis oblongo-spathulatis obtusis, floribus solitariis axillaribus pedunculo longo apice nutanti suffultis, perigonii fulvi flavo-maculati sparsim hirtuli utriculo amplio ovato tubo valde infracto superne sensim ampliato, labio amplio oblongo apice acuto vel acuminato basi cordato obtuse biauriculato, capsula magnâ obovatâ ♀. *Aristolochia Rhazut* Ranw. Reis. 121, tab. 23. — *A. longa foliis angustis acuminatis* Moris. Hist. 12, tab. 17, fig. 11. — *A. Aucherii* Jaub. et Sp. Ill. 1, tab. 99. — *A. Anatolica* Boiss. MSS.

Hab. inter segetes regionis inferioris et præsertim montanæ, Phrygia ad Ouehak (Bal!), Pisidia ad Egirdir (Heldri), Caria (Pinard!), Cappadocia (Bal!), Cilicia Orientalis. (Ky. 51!), Syria circa Aleppo (Auch! Ky! Haussk!), Mesopotumia (Haussk!).

β. latifolia. — Folia latiora triangulari-hastata acuta vel obtusiuscula. — *A. Bottae* Jaub. et Sp. Ill. Or. tab. 98.

Hab. in agris derelictis, Cilicie ad pagum Alladagh (Bal. 766!), Assyriæ (Botta), Kurdistanæ in monte Avronian (Haussk!), Armeniæ Turcicæ ad Gümüşchikhanæ (Bourg!), Persiae borealis (Auch! Ky. 72!).

Variat multum foliis supra basin nunc vix 2 lineas latis nunc basi pollicem et amplius latis, acutis vel obtusiusculis, labio obtusiusculo vel acuminato. Capsula denum 2-2½-pollicaris.

16. *A. pæcilantha* (Boiss. Diagn. Ser. I, 12, p. 104) breviter hirtella, caudice oblongo vel cylindrico, caulis striatis simplicibus vel basi parce ramosis erectis, foliis longe petiolatis triangulari-hastatis obtusiusculis basi sinu lato cordatis auriculis obtusis intus recurvatis, floribus axillaribus solitariis pedunculo petiolo longiore apice nutanti suffultis, perigonii extus hirti intus glabri virescenti-flavidi maculis atropurpureis versus faucem pieti utriculo ovato, tubo valde curvato superne sensim ampliato, labio oblongo-triangulari obtuso basi cordato in auriculas binas oblongas concavas producendo, capsula oblonga magnâ pubescente ♀. *A. hirta* β. *pæcilantha* Duch. in D.C. Prodr. XV, p. 494.

Hab. in umbrosis Syriae et Palestine, Libanus (Bl! Gaill! Ehrenb!), ad Gisr el Hajar, Antilibanus ad Raschyea (Boiss!) et Zebdani (Ky. 234!), Palestina in tribu Ephraim (Boiss!) et ad Hierosolymam (Roth 5!).

β. scabridula. — Magis scabrida, folia breviora, pedunculi magis abbreviati, flores minores rubello-fuscescentes. — *A. scabridula* Boiss. Diagn. Ser. I, 2, p. 104.

Hab. in Antilibano prope Zebdani (Boiss!), in monte Garbi 5000' (Ky. exs. 30!).

Species ab *A. hirtâ* labii formâ omnino diversa sed *A. Maurorum* β *latifoliae* valde affinis, ab ea tamen foliis longe petiolatis, perigonii labio obtuso et intus glabro ut videtur distincta.

17. **A. Pontica** (Lam. Dict. I, p. 255) caudice tuberoso ramoso, caule elato flexuoso simplici, foliis amplis subtus minute hirtulis longe petiolatis ovatis obtusiusculis basi sinu lato cordatis auriculis amplis obtusis, floribus in axillis solitariis longe pedunculatis magnis, perigonii parce pilosi virenti-purpurei utriculo magno ovato, tubo late cylindrico valde incurvo sensim ampliato, labio ampio cordato-oblongo acuto basi late breviter et obtuse biauriculato φ . Ic. Buxb. Cent. I, p. 28, tab. 45 (pessima et forsan potius ad *A. hirtam* referenda). — *A. Orientalis humilis amplissimo folio* Tourn. Goroll. p. 9.

Hab. in sylvis Ponto Euxino conterminis (Tourn.), ad Rhizé Ponti Lazici (Ball).

Specimen descriptum bipedale foliis 5-7 quorum majora 6-pollices longa 5 lata. Flores 3-4-pollices longi 3-3 $\frac{1}{2}$ lati valde curvati labio cum auriculis 2 pollices longo.

18. **A. Iberica** (Fisch. et Mey. in Szov. exs.) caudice..., caule erecto flexuoso, foliis mediocribus subtus hirtis longiuscule petiolatis ovalis vel orbiculatis obtusis vel obtusiusculis subtus sinu lato cordatis auriculis obtusis, floribus mediocribus in axillis solitariis breviter pedunculatis, perigonii crebre pilosi utriculo ovato. tubo cylindrico valde incurvo vix ampliato, labio cordato-oblongo obtusiusculo basi late breviter et obtuse biauriculato φ . *A. Pontica* β . *parviflora* Duchartre in n.c. Prodr.

Hab. in fruticetis ad Malitzki Cartiliniae (Szov!), Imeretiâ (Stev!), Abkhasia (Wittm!), in littore arenoso ad St-Nicolas (Rupr!).

An cl. Duchartre suadente varietas *A. Pontica?* folia tamen multo minora tripollicaria, flores vix sesquipollicem longi labio obtusiore.

19. **A. Cretica** (Lam. Dict. I, p. 225) hirtula, caudice carnoso ramoso, caule erecto vel decumbente inferne interdum parce ramoso, foliis longiuscule petiolatis mediocribus reniformi-cordatis obtusissimis vel retusis basi sinu ampio cordatis auriculis rotundatis, floribus magnis in axillis solitariis pedunculo petiolo longiore suffultis, perigonii parce hirsuti pallide rubri utriculo ampio ovato, tubo late valde curvato a basi sensim et valde ampliato, labio ampio ovato-oblongo apice obtuso vel retuso basi in auriculas binas latas obtusas brevissimas fisco, capsulâ magnâ deflexâ oblongo-clavatâ pubescente φ . Ic. Desf. Choix Pl. p. 12, tab. 7.

Hab. in saxosis Cretæ (Tourn., Siebl), in fauce Hagio Janni prope Critza (Heldr!).

Pedalis et bipedalis, folia 1-2-pollices longa et lata, perigonum 4 pollices fere longum valde recurvum labio fere sesquipolicem longo. Affinis *A. Pontica* sed auriculæ ad labii basin multo breviores.

ORD. CXI. EUPHORBIACEÆ.

(Juss. Gen. 384.)

Flores unisexuales. Calyx liber inferus, quoad partium numerum et aestivationem varius, vel nullus. Petala calycis laciis alterna vel nulla. *Masc.* Stamina centro floris inserta filamentis liberis vel varie connatis. *Fem.* Ovarium liberum superum saepius triloculare, carpellis circa placentam centralem verticillatum dispositis 4-2-ovulatis, ovulis anatropis pendulis. Stylus divisus. Capsulae cocci a columnâ centrali secedentes elastice dissilientes. Semina testâ plus minus coriaceâ, albumine carnosâ, embryone recto axilli, radiculâ ad hilum versâ.

TR. I. EUPHORBIEÆ Bartl.

Flores apetalî, masculi plures monandri cum fæmineo unico involucro gamophyllo inclusi. Filamentum floris masculi cum pedicello articulatum. Ovarii loculi uniovulati. Cotyledones complanatæ.

EUPHORBIA (L. Gen. p. 243).

Involucrum subregulare campanulatum turbinatum vel hemisphericum lobis quinque (rarins 4-8) primariis membranaceis, aliisque totidem alternantibus secundariis glanduliformibus instructum. Flores masculi ecalyculati bracteolis ciliato-laceris interdum basi stipati. Flor. fæmineus centralis pedicellatus calyce 3-6-lobo saepè obsoleto instructus. Styli tres distincti vel plus minus coaliti indivisi vel bifidi. Semina pendula cum vel alsque carunculâ. — Frutices, suffrutices et herbæ succo acri lacteo turgidae. Involucri glandulæ membranâ marginali appendiculatæ vel eâ destitutæ.

Specierum Orientalium distributio.

SERIES I. — Stipulatæ.

SECT. I. ANISOPHYLLUM Ræper. — Folia omnia opposita basi obliqua. Stipulae lanceolatae vel lineares. Involucri glandulæ appendiculatæ.

E. Indica, Peplis, humifusa, granulata, Turcomanica, Chamæsyce, Aegyptiaca, sanguinea, thymifolia.

SECT. II. CHEIROLEPIDIUM Boiss. — Folia infima sparsa cætera opposita. Stipulae subulatæ. Styli indivisi. Glandulæ exappendiculatae palmatisidæ.

E. cheirolepis, Postii.

SECT. III. DIACANTHIUM Boiss. — Folia sparsa (interdum obsoleta). Stipulae aculeiformes. Glandulæ exappendiculatae. Frutices carnosæ pulvinis elevatis tuberculati vel costati.

E. nerifolia.

SERIES II. — Exstipulate

SECT. IV. TIRUCALLI Boiss. — Folia sparsa saepe obsoleta vel squamiformia, floralia opposita vel ternata. Glandulæ exappendiculatae. Frutices carnosæ.

E. Larica.

SECT. V. TITHYMALUS Boiss. — Folia opposita et saepius sparsa; floralia opposita. Glandulæ exappendiculatae integræ vel truncato-bicornes. Herbae, suffrutices vel frutices non carnosæ.

§ 1. Oppositifoliæ. — Folia omnia opposita. Glandulæ bicornes. Inflorescentia dichotoma.

E. pygmaea, *densa*, *sororia*, *Turczaninowii*.

§ 2. Crotonopsideæ. — Folia infimis sparsis exceptis oppositæ. Glandulæ quaternæ pectinatum 2-8-sidæ. Styli indivisi. Inflorescentia dichotoma.

E. lanata.

§ 3. Osyrideæ. — Folia sparsa. Glandulæ quinæ margine truncato-denticulatae. Inflorescentia lateralis. — Frutex ramis longis virgatis.

E. osyridaea.

§ 4. Pachycladæ. — Folia sparsa. Glandulæ integræ vel truncato-bicornutæ. Cymæ terminales umbellatae vel paniculæ. Frutices ramis crassis inferne denudatis cætrice superne foliosis.

E. dendroides.

§ 5. **Carunculares.** — Folia sparsa. Glandulæ integræ vel truncato bicornutæ. Inflorescentia umbellata. — Grex a sequentibus habitu proprio et præsertim carunculâ multo ampliore conicâ elatâ longitudinaliter sulcatâ distincta.

E. cornuta, megalantha, cæladenia, Stockiana, connata.

§ 6. **Galarrhæi.** — Folia sparsa, raro superiora verticillata. Glandulæ transverse ovatæ margine rotundatæ, nec bicornes. Bracteolæ inter flores masculos hirsutæ vel plumosæ. Inflorescentia umbellata ad involucra pauca vel solitarium interdum reducta.

* Semina lævia.

+ Capsula lævis vel sub lente minute et obscure elevato-punctata, non vero verrucosa.

1. Perennes

E. Bungei, Wittmanni, Cypria, pilosa, altissima.

2. Annuæ.

E. arguta, eriophora, Gaillardotii, subtuberculata, microsphæra.

+ + Capsula echinata vel verrucosa.

1. Annuæ.

E. Cybirensis, platyphylla, stricta, coniosperma.

2. Perennes.

E. palustris, Orientalis, aspera, macrocarpa, epithymoides, oblongata, condylocarpa, carlophylla, Apios, dulcis, Djmilensis, thamnooides, acanthothamnos, erinacea, Schottiana, capitulata.

** Semina minute tuberculata.

E. pubescens.

*** Semina areolatim reticulato-rugosa.

1. Perennes.

E. rhytidosperma.

2. Annuæ.

E. pterococca, helioscopia, Berythea, oxyodonta.

**** Semina transverse foveolato vel sulcato-exarata.

1. Perennes.

E. Cassia.

2. Annuæ.

E. phymatosperma.

§ 7. **Esulæ.** — Folia sparsa raro superiora verticillata. Glandulæ marginæ truncatæ, retusæ vel plus minus longe cornutæ. Bracteolæ inter flores masculos hirsutæ vel plumosæ. Inflorescentia umbellata raro dichotoma.

• Semina tuberculata. Omnes annuæ.

E. Aleppica, parvula, exigua, dracunculoides.

•• Semina transverse sulcata. Omnes annuæ.

E. falcata.

••• Semina longitudinaliter sulcata vel seriebus longitudinalibus foveolarum regulariter instructa. Omnes annuæ.

E. aulacosperma, Peplus, peploides, chamœpeplus.

•••• Semina irregulariter foveata, malleata vel reticulato-rugosa

1. Annuæ.

E. Ledebourii, Szovitsii, punctata, arvalis, Græca, Reuteriana, segetalis.

2. Perennes.

E. pinea, Pestalozzæ, erythrodon, obovata, Tcheranica, erythradenia, petrophila, Chesneyi, malleata, decipiens, cheiradenia, bothriosperma, striatella, plebeia, deflexa, Kurdica, bounophila, Aucheri, caudiculosa, herniariefoliaæ.

•••• Semina lœvia. — Species nostræ omnes perennes.

+ Folia floralia libera.

E. Terracina, Gerardiana, Buhsei, Cyparissias, Esula, leptocaula, virgata, hebecarpa, salicifolia, lucida, Iberica, agraria, Baselicis, Orphanidis, glareosa, tinctoria, Paralias.

++ Folia floralia connata.

E. amygdaloïdes, iteophylla, macroceras, rumicifolia, Kotschyana, macrostegia, Characias, Sibthorpii, Lycia, Heldreichii.

§ 8. **Myrsinitæ.** — Folia sparsa dense spiralia. Glandulæ marginæ pectinatæ vel bicornutæ. Bracteolæ inter flores obsoletæ. Inflorescentia umbellata.

* Semina vermiculato-rugosa.

E. Myrsinites, Anacampseros, denticulata.

** Semina tuberculata.

E. craspedia.

*** Semina lœvia.

E. Marschalliana, biglandulosa.

SECT. I. ANISOPHYLLUM.

1. **E. Indica** (Lam. Dict. II, p. 423) annua glaucescens sub lente adpresso et parce pubescens, caulinis elongatis ascendentibus vel decumbentibus ramulosis, foliis breviter petiolatis a basi rotundatâ obovatis retusis subtus pallidioribus remote et obsolete serrulatis, stipulis a basi dilatata setaceis ciliatis, cymis laxiusculæ capitatis axillaribus breviter pedunculatis basi sœpe 1-2 foliis minoribus suffultis, involucri turbinati extus et intus glabri lobis triangularibus fimbriatulis, glandulis ovatis appendice albâ ovatâ æquilitatâ vel angustiore marginatis, stylis brevibus bifidis, capsulæ adpresso hirtæ coccis subcarinatis, semine ovato-tetragono rugis 5-6 transversis fractis anastomosantibus ⊙. *E. granulata* β. *decumbens* Forsk. Arab. p. 112. — *E. decumbens* W. Suppl. — *E. cassioides* Presl Bemerk. p. 119.

Hab. in regno Mascate Arabiæ (Auch. 5301!).

Differt ab affini *E. hypericifolia* Americanâ indumento, capsulis majoribus, seminis sculpturâ.

Ar. Geogr. Africa tropica, Arabia tropica, India Orientalis.

2. **E. Peplis** (L. Amaen. Acad. III, p. 113) annua glabra carnosula glaucescens a collo ramosa, ramis crassiusculis prostratis dichotome ramulosis, foliis subsssilibus a basi valde inæquali obtuse uniauriculatâ oblongo-rhombeis præter basin interdum subdentatam integris obtusis, stipulis 2-3-fidis subulatis, involucris in axillis et dichotomiis solitariis pedicellatis campanulatis fauce hirtis lobis triangularibus ciliatulis, glandulis transverse oblongis concavis appendiceis angustiore integrâ vel lobulatâ, stylis breviter bilobis apice spathulatis, capsulæ magnæ coccis rotundatis, semine griseo ovato lœvi ⊙.

Hab. in arenosis maritimis totius Græciæ et ejus insularum (Sieb! Sprun., Sart!), Macedoniae (Friv!), Anatoliæ! Syriae (Bl! Gaill!), Ægypti ad Alexandriam (Auch! Gaill!), Tauriae et Transcaucasiæ ad Euxinum et mare Caspium (Lebed.), crescit quoque in interioribus, Armenia ad Gumuschkhane (Bourg!), Egyptus ad Benisouef (Ehr!).

Ar. Geogr. Insulae Canarienses et Azoricæ, Gallia occidentalis et Anglia, regio mediterranea Europæ totius et Africæ borealis.

3. **E. humifusa** (Willd. Enum. H. Berol. Suppl. p. 13) annua glaucescens glabra vel parce et patule hispidula, collo ramosa ramis prostratis brevibus dichotomis, foliis subsessilibus a basi valde obliquâ oblongo-ellipticis obtusis minute serrulatis, stipulis subulatis

lacinulatis, involueris in dichotomiis solitariis internodiorum superiorum brevitate saepe approximatis campanulato-turbinatis glabris lobis triangularibus saepe tridentatis, glandulis transverse oblongis appendice eis angustiore obtuse 2-3-lobulatis, stylis longis profunde bifidis, capsulae glabrae coccis obscure carinatis, semine oblongo-tetragono albido sub lente minute pustuloso ☉. *E. Chamaesyce* Fl. Alt. non L. — *E. pseudochamaesyce* F. et M. Ind. Petrop. X, p. 73. — *E. polygonisperma* Gren. et Godr. Fl. Fr. III, p. 75.

Hab. in Caucasi ditione Daghestan (Becker!).

Capsule eis *E. Peplis* multo minores, facies *E. Chamæsyces* a qua seminibus, etc. omnino differt. Forsan in ditione adventitia.

Ar. Geogr. Sibiria, Japonia, Mongolia, China. Variis aliis locis adventitia.

4. E. granulata (Forsk. Fl. Eg. Arab. p. 94) annua tota breviter et patule velutina cinerascens, collo ramosa ramis brevibus prostratis dichotome ramosissimis demum incrassatis ramulis fragilibus, foliis parvis integris a basi inaequali obovatis vel ellipticis obtusis, stipulis minimis lanceolatis ciliatulis, involueris minimis axillaribus subsessilibus racemulos brevissimos foliosos formantibus turbinatis velutinis fauce hirtulis lobis ovatis fimbriatulis, glandularum appendice eis æquilateralâ vel angustiore, capsulae adpresso hirtæ coccis rotundatis, semine acute oblongo-tetragono ruguloso-lacunoso ☉. *E. fragilis* Decaisne Flor. Sinaica. — *E. arillata* Edgew. Journ. Soc. Beng. XVI, p. 1218. — *E. Forskahlei* var. β et γ . J. Gay in Webb Phyt. Can. III, p. 242.

Hab. in Arabiâ petreâ (Bové!), Egypti deserto inter Kené et Kosseir (Schw!).

β . *glabrata*. — Boiss. in D.C. Prodr. — Praeter ramulos parce hirtulos glabra.

Hab. in regno Mascate Arabiæ (Auch. 5304!), Affghania (Griff!), Belutschia (Stocks!).

Species ab *E. Chamæsyce* foliis integerrimis, seminis angustioris sculptura crassiore distincta.

Ar. Geogr. Africa tropica, Algeria interior, Nubia, Arabia tropica, India borealis, insulae Capitis viridis.

5. E. Turcomanica (Boiss. Cent. Euph. p. 13. — D.C. Prodr. XV, p. 34) glabra vel breviter hirtula, collo ramosa ramis brevibus prostratis fragilibus dichotome ramosissimis, foliis parvis a basi valde obliquâ ellipticis subincurvis obtusissimis integris vel prope apicem obsolete denticulato-repandis, stipulis lanceolatis dentatis, involueris axillaribus patule hirtis intus glabris lobis triangularibus, glandulis anguste appendiculatis vel appendice destitutis, capsulae patule hirtæ coccis carinatis, semine oblongo acute tetragono irregulariter lacunoso paucirugoso ☉.

Hab. in Persiæ desertis (Auch. 3185!), Persiâ inter Ispahan et Teheran (Bgel), orientali prope Tebes et Sebsewar (Bgel), Turcomaniâ ad Caspium (Karell!).

β. Tigridis (Boiss. in D.C. Prodr.). — Semina ob rugulas magis anastomosantes scrobiculato-punctata. — *E. Tigridis* (Boiss. in sched.).

Hab. in Babyloniam prope Bagdad (Noë 931! et 1090!).

Inter *E. Chamæsyce* et *E. granulatum* media, a priore foliis subintegris formæ diversæ, ramis demum induratis, semine angustiore; a posteriore foliis basi magis obliquis uniauriculatis distincta.

6. E. Chamæsyce (L. Amæn. Acad. p. 115) annua plus minusve hirta vel dense canescens collo ramosa, ramis prostratis filiformibus raniulosis, foliis parvis a basi inæquali rotundatâ ovatis vel suborbiculatis obtusis obtuse crenulatis, stipulis setaceis basi denticulo sæpius auctis, involucris in dichotomiis et axillis supremis solitariis glabris vel hirtis fauce ciliatis lobis ovato-triangularibus fimbriatis, glandulis transversis cerinis subconeavis appendice æquilitatâ vel angustissimâ plus minusve trilobâ, capsulæ glabræ vel hirtæ coccis carinatis, seminis ovato-tetragoni albidi rugulis numerosis irregulariter anastomosantibus ☽. Ic. Fl. Græc. tab. 461. — Rehb. Germ. tab. 131. — *E. Massiliensis* D.C. Fl. Fr. V, p. 357. — *E. canescens* L. Sp. 652. — Cav. Ic. tab. 63.

Hab. in cultis et ad vias regionis inferioris et montane totius ditionis a Græciâ! et Thraciâ! ad Syriam! Babyloniam! Tauriam, Caucasum et Transcaucasiam, Persiam bor. (Szov!) et australem (Haussk!). Interdum alpina, Parnassi cacumen (Guic!).

Ar. Geogr. Regio mediterranea totius Europæ et Africæ borealis.

7. E. Ægyptiaca (Boiss. Cent. Euph. p. 43. — D.C. Prodr. XV, p. 35) annua crispule et breviter hirtula collo ramosa, ramis prostratis elongatis crassiusculis ramosissimis, foliis a basi obliquâ subcordatâ oblongo-ellipticis obtusis apice remote serrulato-crenatis crenis muticis vel mucronatis, stipulis lanceolatis fimbriatulis, involucris axillaribus solitariis internodiorum brevitate glomerulos vel racemulos breves foliosos formantibus turbinatis hirtis intus glabris lobis ovatis fimbriatis, glandulis rubris transverse ovatis coneavis appendice angustissimâ vel obsoletâ, capsulæ adpresso hirtæ coccis acutiuscule carinatis, semine albido oblongo-tetragono irregulariter lacunoso-rugoso ☽. *E. Forskahlei* var. α . J. Gay in Webb Phyt. III, p. 240.

Hab. in Egypto mediâ et superiore (Auch. 2033! Boiss! Kralik!), in Oasi magnâ (Schweinf!), in Oasibus Farafrah et Dachel (Asch!).

Rami saepe pedales eis *E. Chamæsyces* crassiores, folia majora saepe 6 lineas longa, involucra minora confertiora subsessilia.

Ar. Geogr. Senegal, Nubia, Abyssinia, Africa tropica, insulæ Capitis viridis, India borealis.

8. E. sanguinea (Hochst. et Steud. in Schimp. exs. — D.C. Prodr. XV, p. 35) annua glabra glaucescens collo ramosa, ramis elongatis prostratis raniulosis, foliis a basi valde inæquali elliptico-rhombeis subincurvis obtusissimis margine saepe calcareo-serrulatis,

slipulis elongatis a basi latiore saepe 2-3-fidâ subulatis, involucris in axillis supremis solitariis confertis racemulos foliosos formantibus turbinatis intus et extus glabris lobis linear-lanceolatis integris, glandulis appendice roseâ irregulariter et profunde 3-5-partitâ eis subaequilatâ marginatis, capsulae glabrae coccis carinatis, semine ovato-tetragono irregulariter paucirugoso ⊖.

Hab. in regno Cabulico (Griff).

Caules saepe semipedales et longiores, folia eis *E. Chamæsyces* majora, alata et affinis seminibus obscure rugulosis distincta.

Ar. Geogr. Nubia, Abyssinia, Arabia tropica, India, Africa australis.

9. **E. thymifolia** (Burm. Fl. Ind. p. 2) annua collo ramosa ramis prostratis filiformibus saepe elongatis ramosis crispulis hirtis, foliis breviter petiolatis a basi subinæquali oblongo-ellipticis obtusis obtuse crenulatis glabris vel subtus adpresso hirtis, stipulis elongatis lanceolatis fimbriato-dentatis, involucris axillaribus solitariis in racemulos breves foliosos axillares et terminales congestis turbinatis adpresso hirtis profunde fissis intus praeter lobos breves ciliatulos glabris, glandulis minimis stipitatis suborbiculatis appendice æquilatâ vel angustiore 2-3-lobâ, capsulis breviter stipitatis etiam junioribus erectis adpresso hirtis obtuse carinatis, semine rubello oblongo-tetragono transverse 4-5-sulcato ⊖. Ic. Thes Zeyl. tab. 103, fig. 2 — *E. Burmanniana* J. Gay in Webb Phyt. III, p. 239.

Hab. in Belu'schiâ (Frere ex Aitch. Catal.).

Ar. Geogr. Africa Orientalis tropica, India, America tropica.

SECT. II. CHEIROLEPIDIUM.

10. **E. cheirolepis** (F. et M. Bull. Soc. Mosq. 1839, p. 171) annua glabra vel hirtula, caule erecto alternatim ramuloso, foliis infimis alternis oblongis vel linear-lanceolatis superioribus oppositis ovatis vel obovatis omnibus remote et argute denticulatis, stipulis minutis dentiformibus subulatis, involucris axillaribus et terminalibus solitariis breviter pedicellatis campanulatis pubescensibus lobis ovatis fimbriato-dentatis, glandulis plus minus inæqualibus transverse ovatis 3-7-palmatis rarius subintegris, stylis ad medium usque coalitis indivisis circinnatis, capsulae membranaceæ tomentellæ coccis dorso rotundatis, semine oblongo minute verruculoso, carunculæ vesiculosæ erubibus binis descendilibus semini adpressis ⊖.

Hab. in sabulosis desertorum, prope Kerman Persiae australis (Bge!), in Turkestanie deserto Kisilkum (Lehm!), Turcomaniâ (Karelin!).

Demum pedalis vel sesquipedalis, folia inferiora pollicem et amplius longa 2-4 lineas lata, superiora breviora, capsula diam. 2 lineas lata. Species characteribus insignis.

11. **E. Postii**, annua, glauco-virens, caule erecto substriato a basi

alternatim ramoso ramis elongatis stricte plures dichotomis, foliis præter infima alterna oppositis breviter petiolatis ovatis et ovato-oblongis obtusis argute denticulatis, stipulis minimis subulatis utrinque binis, ramulis floriferis petiolisque junioribus hirtis, involucris breviter pedicellatis hemisphericis glabris lobis fimbriatis, glandulis longiuscule stipitatis transverse oblongis ultra medium 5-7 palmatisidis, stylis indivisis circinnatis, ovario juniore glabro, capsula..... ⊖.

Hab. inter Hamath et Maharadek deserti Syriaci (Dr. Post!).

Planta pedalis facie *E. lanatæ* sed glabrescens, folia 8-10 lineas longa. Ob stipularum naturam, involuci characteres juxta *E. cheirolepidem* collocanda caule multo tenuiori, ramis non dichotomis, involucro et ovario tomentellis, glandulis non stipitatis distinctam.

SECT. III. DIACANTHUM.

12. ***E. nerifolia*** (L. Hort. Cliff. p. 196) ex parte fruticosa vel arborea, ramis ob podaria obtusa basi vix confluentia spiraliter et obtuse pentagonis, aculeis stipularibus brevibus divergentibus nigris, foliis obovato-oblongis obtusis mucronulatis basi sensim in petiolum brevem attenuatis, cymis in axillis superioribus subsessilibus 1-3-cephalidis, foliolis sub involucro ovatis membranaceis, involuci hemispherici lobis magnis ovatis fimbriatis, glandulis transverse ovatis, capsula trigonâ ȝ.

Hab. in Belutschia (Stocks ex Aitch. Cat.). Vix ditionis nostræ civis.

Ar. Geogr. India Orientalis.

SECT. IV. TIRUCALLI.

13. ***E. Larica*** (Boiss. Cent. Euph. p. 24) fruticosa vix carnosa, ramis erectis teretibus dichotome et stricte ramosissimis junioribus pulvillis foliorum scutellæformibus remote cicatricosis, foliis in gemmis terminalibus citissime deciduis squamiformibus oblongo-linearibus acutiusculis, involucris terminalibus solitariis parvis subsessilibus foliolis binis oblongo-linearibus eis brevioribus suffultis campanulatis lobis ovatis fimbriatis, glandulis transverse oblongis immarginatis ȝ.

Hab. in prov. Laristan Persiae australis (Auch. 5299!).

Affinis *E. Schimperi* Presl ex Arabiâ tropicâ, rami tenuiores minus cicatricoso-asperi, regularius dichotomi, involucrum dimidio minus angustius. Species ex spec. incompletis tantum nota.

SECT. V. TITHYMALUS.

§ 1. Oppositifoliæ.

14. **E. pygmæa** (F. et M. in Bull. Mosq. 1839, p. 171) annua glabra, caule erecto tenui humili superne dichotome corymboso, foliis linearibus integris acutiusculis sessilibus oppositis, floralibus conformibus abbreviatis, involueri minuti sessilis campanulati lobis lanceolatis, glandulis transverse ovatis margine truncatis interdum brevissime bicornutis, capsulae coccis carinatis, semine hexagono ad facies sulco longitudinali percurso et transverse corrugato, carunculâ parvâ stipitatâ globosâ ⊖. Boiss. Euph. Ic. tab. 51. — *E. Inderiensis* Less. MSS.

Hab. in montosis prov. Khorassan Persiæ frequens (Bge!), Turcomania (Karel!), Turkestania (Lehm!).

Planta 3-6-pollicaris, folia 6-8 lin. longa vix lineam lata.

Ar. Geogr. Songaria.

15. **E. densa** (Schrenck Bull. Phys. Acad. Petrop. III, p. 308) annua glabra glaucescens pumila fere a collo dichotome ramosissima internodiis brevibus, foliis oppositis sessilibus basi breviter attenuatis suborbiculatis mucronulatis margine scabridis, floralibus conformibus, involucri subsessilis parvi turbinati lobis oblongis denticulatis, glandulis transverse oblongis 1-2 cornubus linearibus sœpe auctis, capsulae pedicello crasso suffultæ oblongæ coccis subcarinatis, semine albo quadrangulo reticulato-scrobiculato, carunculâ minimâ sessili rotundatâ ⊖.

Hab. saepius in salsuginosis, Persiæ orientalis prov. Khorassan et Kerman, Turkestania ad Kisylkum (Lehm!), Belutschia superior (Stocksl.).

Plantula 2-3-pollicaris 4-5-pollices lata facie *E. falcatae*.

Ar. Geogr. Songaria.

16. **E. sororia** (Schrenck Bull. Phys. Acad. Petrop. III, p. 308) annua glabra glaucescens a collo dichotome ramosissima pumila internodiis brevibus, foliis crebris oppositis anguste lanceolatis basi attenuatis, floralibus conformibus diminutis, involucris sessilibus turbinatis lobis ovatis denticulatis, glandulis transverse ovatis sœpe 1-2 cornubus linearibus auctis, capsulae coccis rotundatis, semine oblongo hexagono albo-leproso ad facies longitudinaliter sulcato transverse corrugato, carunculâ globosâ longe stipitatâ ⊖.

Hab. in desertis Persiæ mediae ad Ssof inter Ispahan et Teheran (Bge!).

Planta 4-8-pollicaris et totidem lata dense foliosa, folia infer. 8-10 lineas longa 2 lata.

Ar. Geogr. Songaria.

17. **E. Turczaninowii** (Kar. et Kir. Enum. Song. p. 178) annua glabra pumila a basi dichotoma internodiis brevissimis, foliis oppositis sessilibus a basi attenuatâ oblongo-lanceolatis acutis vel acuminatis inaequilateris plurinerviis margine seahridis, floralibus diminutis, involacri cylindrico-obconici lobis ovatis fimbriatulis, glandulis substipitatis transverse ovatis cornubus brevibus, capsulae coccis rotundatis, semine oblongo-pyramidalato albo-leproso facie exteriore et interiore subcarinato sulcis longitudinalibus ramulosis corrugato, carunculâ nullâ ⊖.

Hab. in collibus arenosis mobilibus Turkestaniæ ad fluvios Syr Darja et Kuwendarja (Lehm!).

Planta 2-5-pollicaris, folia majora pollicaria 4-5 lineas iata, ræcies *Anastaticæ*.

Ar. Geogr. Songaria.

§ 2. *Crotonopsidææ.*

18. **E. lanata** (Sieb. in Spreng. Syst. III, p. 792) annua adpresso-villosa canescens erecta a basi dichotoma, foliis petiolatis ovatis spinuloso-dentatis præter infima sparsa oppositis, floralibus sensim diminutis, involucris in dichotomiis et terminalibus breviter pedunculatis campanulato-hemisphericis extus et intus longe hirtis lobis triangulari-ovatis fimbriatis, glandulis concaviusculis transverse oblongis subtus hirtis margine pectinatum in lacinias setaceas fissis, capsulae magnæ ovatae lanatae coccis carinatis, semine nitido quadrangulo utrinque truncato grosse pustuloso, carunculâ stipitatâ horizontali oblongâ ⊖. Boiss. Ic. Euph. tab. 53. — *E. Syriaca* Spreng. Syst. III, p. 792. — *E. malacophylla* Clarke in Spreng. N. Entd. III, p. 161.

Hab. in cultis derelictis, vineis Syriae ad Aleppo (Auch!), ad Damascum et in Cælesyria (Boiss!), Mesopotamiæ et Assyriæ præsertim in regione montana (Ky. 433! Auch! Haussk!), Persiaustrali (Haussk!), media (Auch!), boreali (Ky. 605!) et in prov. Khorassan (Bge!).

Planta tandem sæpè sesquipedalis, folia 12-15 lin. longa. Capsula pisum fere æquans.

§ 3. *Osyrideæ.*

19. **E. osyridea** (Boiss. Diagn. Ser. I, 7, p. 87) fruticosa glauca breviter pruinosa-lomentella, ramis longis virgatis strictis striatis parce ramulosis inferne foliatis, foliis sparsis sessilibus lanceolato-linearibus acutiusculis integris, cymis breviter pedunculatis racemosis bifidis vel trifidis, involucro foliolis binis ovato-rhombeis suffulto tomentello turbinato lobis ovatis truncatis, glandulis transverse oblongis, capsula pedicellata tomentellâ globosâ trisulcâ, semine ovato-lævi, carunculâ minimâ depresso-ʒ. *E. lateriflora* Jaub. et Sp. Ill. Or. 2, tab. 130.

Hab. in Persiā australi (Auch. 5298!), in monte Buschom prov. Schiras (Ky. 111!), in Afḡhanistā occid. prope Herat et Anarderreh (Bge!) et orientali (Griff!), in Belutschia (Stocks!).

Rami crassitie penna anserina 2-3-pedales, folia 4-4½-pollicaria 1½-2 lineas lata, racemus pedalis vel longior angustus laxus, capsula piso parte tertia minor.

§ 4. Pachycladæ.

20. E. dendroides (L. Sp. 662) fruticosa glabra dichotome vel ternatim ramosa ramis rubellis nitidis vetustis, nudis annotinis crebris foliosis, foliis sessilibus a basi valde attenuatâ linearî-lanceolatis obtusis vel acutiusculis subtus pallidioribus, umbellis terminalibus 3-10-fidis radiis brevibus bifidis, foliis floralibus lutescentibus subcordato vel rhomboeo-semiorbiculatis involuerum superantibus, involueri campanulati lobis oblongis 2-3-fidis timbriatulis, glandulis peltatis transverse ovatis trunneatis vel semilunatis subtus hirtulis, capsula profunde trisulcæ coccis lateraliter subcompressis, semine ovato lœvi subcompresso, carunculâ sessili oblongâ cristæformi ♂. Ic. Fl. Græc. t. 470. — Rehb. Germ. fig. 4772

Hab. ad rupes præsertim maritimæ totius Peloponnesi (Sibth! Boiss! Orph. Fl. exs. 699!), Phocidæ ad Delphos (Heldr. Herb. Norm. 641!), Eubœa (Heldr!), Cycladum (Bory!), Cretæ (Sieb!). Fl. Mart., Apr.

Ar. Geogr. Hispania boreali-orientalis, Gallia australis, Italia, Africa borealis.

§ 5. Carunculares.

21. E. cornuta (Pers. Syn. 2, p. 47) annua vel perennans glabra glaucescens multicaulis, caulis ascendentibus demum induratis foliosis, foliis crassiusculis sessilibus linearibus vel obovato-oblongis acutis vel retusis plus minus argute serratis, umbellarum radiis 3 vel 2 pluries dichotomis, foliis floralibus a basi dilatatâ cordatâ pectinato-serrata caudato-acuminatis apice sepe recurvis, involueri campanulati lobis triangularibus timbriatis, glandulis concaviuseculis transverse ovatis margine breviter 2-4-cornutis rarius palmatim 6-10 fidis, capsulæ magnæ coccis rotundatis, semine albido lœvi, carunculâ magnâ conicâ profunde quadrisulcatâ ⊖. *E. retusa* Forsk. Fl. Eg. Arab. p. 93. — *E. serrata* var. β . Willd. Spec. II, p. 915.

Hab. in sterilibus et præsertim desertis Egypti (Forsk., Delile! Boiss! Schweinf! etc.), Arabiæ petreæ (Schimp. 337!).

Ar. Geogr. Africa borealis in regione Saharensi et agro Tunetano.

22. E. megalantha (Boiss. Diagn. Ser. I, 7, p. 95) perennis glauca, caulis ascendentibus striatis inferne ramos steriles

foliosos edentibus, foliis sessilibus crassiusculis lanceolatis acutis integris vel minute denticulatis, ramorum sessilium anguste linearibus, umbellæ radiis 5-6 semel vel pluries dichotomis, foliis floralibus a basi subdilatata cordata oblongo-lanceolatis acutis, involucri hemispherici magni glabri rarius velutini lobis longe linearibus glandulas superantibus, glandulis magnis stipitatis transverse ovatis integris vel truncatis vel breviter bicornutis, semine cæsio levii, carunculâ magnâ ovato-conicâ multisulcatâ latere raphen spectanti excavatâ ♀. Boiss. Euph. Ic. tab. 59. — *E. Ispahanica* Boiss. Diagn. Ser. I, 7, p. 91. — *E. coriacea* C. Koch Linn. XXI, p. 730.

Hab. in Armeniâ Rossicâ ad Erivan (C. Koch!), Persiâ boreali (Szov! Buhse! Ky. 56!), Persiâ media et australi (Bge! Auch. 5289! Ky. 270!), in monte Ataschke supra Sihna usque ad 9000' (Haussk!).

β . *gracilis*. — Caules tenuiores juncei, umbella 2-4-fida, folia et involucra diminuta. — *E. heteradena* Jaub. et Sp. Ill. Or. 2, tab. 131.

Hab. in Persiâ prope Ispahan (Auch. 5313!).

γ . *denticulata*. — Facies et inflorescentia depauperata varietatis præcedentis, folia parva oblongo-linearia versus apicem obtusiuscule denticulata, glandulæ cornua brevia subbiloba.

Hab. in Persiâ australi prope Khabbise et Kerman (Bge!).

Pedalis et procerior, foliorum formâ longitudine et latitudine valde variabilis sed lobi involucri longitudine et carunculae formâ semper recognoscenda. Capsula sæpe 3 lineas longa.

23. ***E. cæladenia*** (Boiss. Diagn.-Ser. II, 4, p. 81) annua glabra glaucescens erecta, foliis sessilibus linearibus obtusis argute et remotiuscule serratis, umbellæ radiis ternis elongatis pluries dichotomis, foliis umbellaribus et floralibus latioribus a basi subcordata oblongis et oblongo-triangularibus obtusis, involucri hemispherici intus hirti lobis truncatis bifidis, glandulis concaviusculis transverse ovatis integris, capsulae longe pedicellatae coccis rotundatis, semine albido, carunculâ sessili elongatâ angustâ conico-triquetrâ sulcatâ ♂.

Hab. in Belutschia ad Khozdar (Stocks!).

Pedalis, folia caulina 8-10 lineas longa 2 lata, radii 4-6-pollices longi, capsula 2 lineas circiter longa.

24. ***E. Stocksiana*** (Boiss. Cent. Euph. p. 34 d.c. Prodr. XV. p. 112) perennis glabra glaucescens, caulinis ascendentibus striatis inferne ramos steriles edentibus, foliis sessilibus a basi sublatiore linear-lanceolatis obtusis tenuiter et argute serrulatis, umbellæ radiis ternis brevibus bifidis, foliis floralibus abbreviatis triangulari-lanceolatis, involucri hemispherici intus hirtuli lobis elongatis acute bidentatis, glandulis stipitatis transverse ovatis integris, capsulâ.... ♀. *E. odontophylla* Boiss. Diagn. Ser. II, 4, p. 82 non Willd.

Hab. in Belutschia ad Kuam (Stocks 1069!), in Affghaniâ occidentali ad Ssertschah (forma foliis latioribus abbreviatis) (Bge!), in prov. Khorassan Persiæ (Bge!).

Subpedalis, facies *E. cornutæ* a quâ glandularum formâ et foliis floralibus non caudatis, etc., differt.

25. **E. connata** (Boiss. in p.c. Prodr. XV, p. 112) perennis glauca hispidula, caulis erectis crassiusculis striatis inferne parce ramulosis, foliis sessilibus a basi breviter cuneatâ vel subcordatâ ovato-orbiculatis obtusissimis profunde et acute subduplicato-dentatis, umbellæ trifidæ radiis bifidis, foliis umbellaribus et floralibus in unum orbiculare minute denticulatum ad medium usque connatis, involueri hemispherici hirti lobis lanatis lanceolatis apice indivisis vel laceris glandulas superantibus, glandulis stipitatis transverse oblongis, capsulæ ovatae profunde trisulcæ cocceis rotundatis, semine subgloboso, carunculâ conico-cylindricâ sulcatâ basi in lacinias dimidio semini æquilongas solutâ ♀. Boiss. Euph. Ic. tab. 61.

Hab. in Persiaustrali prope Kerman (Bge!).

Pedalis, umbellæ radii 2-3-pollicares, folia inferiora 6-9 lineas, superiora dilatata pollicem et amplius diametro longa, capsula pisum æquans.

§ 6. Galarrhæi.

* Semina laevia.

+ Capsula laevis vel minute et obscure sub lente tuberculata nec verrucosa.

1. Species perennes.

26. **E. Bungei** (Boiss. in p.c. Prodr. XV, p. p. 115) perennis glabra glauca, caulis a basi prostratâ caudiculosâ ascendentibus tenuibus simplicibus pumilis dense foliatis, foliis parvis crassis a basi sessili late cordatâ triangulari-ovatis acutiusculis margine crispato-undulatis et argute denticulatis, floralibus subconformibus, umbellæ parvæ radiis ternis brevibus, involueri membranacei breviter turbinati lobis profunde et acute bifidis glandulas superantibus, glandulis substipitatis transverse oblongis, capsulæ subsphæricæ coccis rotundatis, semine ovato, carunculâ unicâ depressâ ♀. Boiss. Ic. Euph. tab. 65.

Hab. in Persia boreali-orientali prope Schahrud et Ssebsewar (Bge!).

Caulis 4-5-pollicares, folia ab infimis squamæformibus sursum aucta, majora 8-9 lineas longa basi obtuse auriculata, radii tandem 1-2-pollicares.

27. **E. Wittmanni** (Boiss. Cent. Euph. p. 31 et p.c. Prodr. XV, p. 115) glabra e rhizomate suffrutescenti pluricaulis, caulis pumilis ascendentibus simplicibus foliatis, foliis membranaceis sessilibus margine sub lente cartilagineis et serrulatis inferioribus et ramorum sterilium oblongo-lanceolatis, floralibus oblongis, umbellæ radiis quinis tenuibus bifidis, involueri turbinato-campanulati lobis late ovatis ciliatulis, glandulis substipitatis transverse ovatis, stigmatibus

subsessilibus, ovario glabro, capsulâ sub lente minute granulato-asperulâ, semine parvo lævi, carunculâ depresso conicâ ♂.

Hab. in saxosis Imeretiae Caucasicæ ad Atskur et Usurgeti (Wittm!), ad Borschom 2500'-5200' (Radde exs. 631).

Caules tenues 5-6-pollicares, folia 6-9 lineas longa 1 $\frac{1}{2}$ -2 $\frac{1}{2}$ lata. Capsula magnitudine seminis *Cannabeos*. Facies *E. flavidomæ*.

28. E. Cypria (Boiss. in D.C. Prodr. XV, p. 115) glabra suffruticoso-erinacea a basi ramosissima ramis tenuibus rigidis dichotomis vel alternatim ramulosis, foliis paucis minimis linearibus coriaceis subintegris, floralibus ovato-oblongis in mucronulum recurvum abeuntibus, umbellis ad involucrum terminale solitarium inter folia floralia sessile turbinatum fauce velutinum reductis, lobis transverse ovatis truncatis vel retusis, glandulis transverse ovatis, stylis longis, ovario lævi, capsulâ . . . ♂.

Hab. in cultis derelictis insulæ Cypræ prope Catiga (Gaudry!). Vid. in herb. Mus. Par.

Suffrutex erinaceus sesquipedalis, ramuli annotini 1-2-pollicares nudiusculi, folia 2 lineas longa dimidiata sub lente obsolete crenulata, folia floralia opposita submajora 2 $\frac{1}{2}$ lin. longa. Facies *E. spinosæ* et *erinaceæ*, sed leioarpa videtur, capsulâ tamen non nota locus subincertus.

29. E. pilosa (L. Sp. I, p. 659) pilosa rarius glabrescens, caulis ex rhizomate crasso erectis sæpe proceris crassis foliosis ex axillis et sub umbellâ sæpe ramulosis, foliis sessilibus oblongis et oblongo-lanceolatis obtusis vel acutiusculis apice plus minus obsolete serrulatis, umbellæ radiis 5-6 abbreviatis trifidis dein bifidis, foliis floralibus lutescentibus ovatis obtusis mucronulatis, involucri campanulati extus hirti lobis ovatis ciliatis, glandulis ovatis, capsulâ subglobosâ glabrâ vel patule hirtâ lævi vel sparsim et minute tuberculatâ, carunculâ patellari depressâ ♀. Reich. Ic. Germ. fig. 4770. — *E. procera* M.B. Taur. Cauc. I, p. 378. — Rchb. Cent. II, tab. 270. — *E. villosa* W. K. Hung. I, p. 96, tab. 93.

Hab. in collibus Tauriæ (Stev.), radicibus Caucasi ad fl. Terek (M.B! C.A. M!).

Ar. Geogr. Hispania borealis, Gallia australis et occidentalis, Italia australis, Germania orientalis, regio Danubialis, Rossia media et australis, Sibiria Uralensis.

30. E. altissima (Boiss. Diagn. Ser. I, p. 52) hirsuta sæpe albo-lanata rarius glabrescens, caulis ex rhizomate crasso valde elatis crassis crebre foliosis in paniculam longam amplam abeuntibus, umbellis compositis 6-9-radiatis, foliis sessilibus longe lanceolatis acutissimis subtus nervo albo percursis a medio sursum minute serrulatis, floralibus ovatis mucronulatis, involucri campanulati pilosi lobis ovato-oblongis, glandulis ovatis, capsulæ globosæ trisulcæ longe pilosæ sub lente minute tuberculatæ coccis rotundatis, carunculâ minutâ depressâ ♀. Boiss. Euph. Ic. tab. 66.

Hab. ad aquas in Phrygiâ et Lydiâ (Boiss! Bal!), Ciliciâ ad Ermenek (Péron!), Syriâ bor. ad Marasch et Aintab (Haussk!), Antilibano (Ehr! Ky. 111!), Assyria (Ky. 367!). Specimina ex Marienfeld Kachetie Transcaucasicae sub *E. procerâ* ab Hohenacker distributa huc quoque spectare videntur.

Caules 5-10-pedales, affinis *E. pilosæ* differt foliis longis angustis acutis simis, paniculâ amplâ elongatâ nec corymboso-contractâ.

2. Species annue.

31. *E. arguta* (Soland. in Russ. Alepp. II, p. 252) annua patule et molliter hispidula glaucescens, caule erecto in umbellam simplicem vel sæpius in paniculam abeunte, foliis argute et profunde serratis oblongis et oblongo-lanceolatis infimis obtusis in petiolum brevem attenuatis cræteris sessilibus acutis, umbellæ radiis 3-5 semel vel his 3-5-fidis, foliis floralibus parvis rhomboe-ovatis, involueri turbinati extus glabri intus sub glandulis pilosuli lobis breviter ovatis fimbriato-dentatis, capsulæ sphæricæ depresso sub lente puncticulatae trisulcalæ coccis rotundatis, semine sphærico brunneo, carunculâ orbiculari depressâ ⊖. Ic. Fl. Græc. V, tab. 468. — *E. calendulaefolia* Del. Fl. Eg. tab. 30, fig. 1.

Hab. in cultis, Peloponnesus in Arcadiâ (Bory!), Cyprus (Sibth.), Cilicia (Bal!), Syria littoralis (Bl! Gaill!), Palestina (Boiss!), Egyptus inferior et superior (Del., Sieb! Boiss! etc.), in Oasi Dachel (Asch!).

Facies *E. helioscopiae* sed laxior et procerior, folia floralia minora.

32. *E. eriophora* (Boiss. Diagn. Ser. I, 5, p. 51) annua tota molliter hispida grisea, caule erecto humili ex axillis superioribus ramuloso breviter paniculato, foliis sessilibus basi attenuatis oblongo-lanceolatis acutis minute denticulatis, floralibus ovato-oblongis acuminate vel mucronatis, umbellarum radiis ternis bis vel ter dichotomis patulis, involueri parvuli turbinati glabri intus hirti lobis ovalis truncatis fimbriato-laceris, glandulis stipitatis transverse reniforme oblongis margine superiore arcuatis, capsulæ globosæ depresso patule hirtulæ coccis rotundatis, semine sphærico cæsio fusco-punctato, carunculâ nullâ ⊖. Boiss. Euph. Ic. tab. 68. — *E. lasiocarpa* C. Koch Linn. XXI, p. 721.

Hab. in cultis derelictis Cariæ interioris (Boiss!), Phrygiæ (Bal. 184!), Fisiæ (Heldr!), Armeniae Turcicæ (C. Koch! Bourg!), Syriae borealis ad Aintab, Nisib (Auch! Haussk!), Persiae Babyloniæ finitimæ ad Kirrind (Noël!).

Semipedalis, folia majora 1-1½-pollicaria 5-6 lin. lata, capsula magn. pist. Facies *E. lanatæ*.

33. *E. Gaillardotii* (Boiss. et Bl. Diagn. Ser. II, 4, p. 84) annua glabra glaucescens, caule erecto brevi sub umbellâ ramuloso ramis umbellæque radiis quinque demum valde elongatis dichotome et plures ramosissimis, foliis subsessilibus a basi attenuatâ elliptico-lanceolatis acutis serrulatis demum deflexis et cito deciduis, umbellaribus conformibus, floralibus suborbiculatis mucronatis ob internodia abbreviatis.

viata demum approximatis et subimbricatis, involucri turbinati intus ad faucem hirti lobis ovatis fimbriatis, glandulis transverse ovato-triangularibus, capsulae minutæ ovatae coccis rotundatis, semine ovato subquadrangulo nigrescente luteo-maculato, carunculâ nullâ ⊖. Boiss. Ic. tab. 69.

Hab. in cultis Cœlesyriæ et Antilibani (Gaill! Ky. 251!), Mesopotamiæ ad Beschhabur (Ky 398!), prope Orfa, Suerek, Kkarput (Haussk!).

Caulis 3-5-pollicaris sed umbella terminalis cum radiis demum pedalis, folia floralia ut in *E. falcatâ* sæpe imbricata, capsula vix lineam longa et lata. Glandularum formâ ad *Esulas* paululum accedit.

34. *E. subtuberculata* (C.A. Mey MSS. — Boiss. in D.C. Prodr. XV, p. 118), annua glabriuscula a collo sæpe pluricaulis caulis erectis, umbellæ radiis quinis 3-5-ffdis dein bifidis, foliis sessilibus serrulatis basi attenuatis infimis obovalis obtusis cæteris oblongolanceolatis acutis, floralibus ovato-orbiculatis mucronulatis ad nervum subtus hirtulis, involucri turbinati intus et extus hirti lobis ovatis truncatis fimbriatis, glandulis transverse oblongis, capsula pilosulâ demum glabrata globosa subtrisulcatâ sub lente minutissime tuberculato-rugulosa, semine ovato subcompresso rufescente, carunculâ minimâ transverse ovalâ ⊖.

Hab. juxta vias in Persiæ prov. Aderbidjan ad Khoï (Szov!). V. in herb. Petrop.

Facies *E. platiphyllæ* a qua capsula non verrucosa differt, aff. *E. microsphæræ* capsula exsulca levissimâ subindehiscenti diversæ.

35. *E. microsphæra* (Boiss. Diagn. Ser. I, 7, p. 87) annua glabriuscula, caule erecto ad collum interdum 1-2 ramulis aucto, umbellæ radiis quinis trifidis dein interdum bifidis, foliis sessilibus membranaceis apice serrulatis infimis basi attenuatis subpetiolatis oblongis obtusis, cæteris sessilibus acutis, floralibus ovato vel orbiculato-triangularibus obtusis, involucri turbinati extus et intus hirti lobis ovatis fimbriatis, glandulis flavis transverse ovatis, capsula longe stipitatâ juniore hirtâ maturâ glabrâ lævi sphæricâ non sulcatâ subindehiscenti, semine ovato subpresso, carunculâ minimâ patellari ⊖. *E. Mohammerensis* Boiss. MSS.

Hab. in cultis Babyloniæ ad Mohammera (Noë 39!), Persiæ australis prope Schiras (Ky. 448! Haussk!), Persiæ austro-orientali ad Chabbise (Bge!).

Similis formis macrocarpis *E. Cybirensis* a qua umbellæ radiis trifidis nec bifidis et præsertim capsula levissimâ nec echinata differt.

++ Capsula echinata vel verrucosa.

1. Species annuae.

36. *E. Cybirensis* (Boiss. Diagn. Ser. I, 7, p. 89 et 12, p. 109) annua glabra, caule erecto subsimplici, umbellæ radiis quinis demum elongatis et sæpe plures dichotomis, foliis sessilibus infimis obovato-

spathulatis obtusis cæteris ellipticis acutiusculis omnibus superne *serrulatis*, floralibus cordatis triangulari-deltoides, involuci sessilis *campanulati* glabri lobis ovatis dentatis, glandulis transverse oblongis, capsula magnâ stipite æquilongo suffulta sphæricâ vix sulcatâ subindehiscenti aculeis elongato-conicis apice setiferis plus minus copiose obsitâ, semine ovato compresso, carunculâ patellari sessili ☽. Boiss. Euph. Ic. tab. 7. — *E. verrucosa* L. Mant. II, p. 303 et herb! ex parte.

Hab. in cultis et vinetis, Rhodus ad Bastida (Bourg!), Caria (Boiss!), Syria littoralis (Bl! Gaill!) et interior ad Rascheya (Boiss!), Palestina ad Naplouse (Boiss!), Mesopotamia (Haussk!).

β. microcarpa. — Capsula minor.

Hab. in Ciliciâ ad Mersina (Bal. exs. 738!).

γ. dehiscens. — Folia acutiora, capsula dehiscens aculeis mollibus vix induratis. — *E. echinocarpa* Sieb. pl. Cret. exs. non Brot.

Hab. in insulâ Cretâ (Sieber!).

δ. longifolia. — Folia utrinque attenuata elongata 2-3-pollicaria, capsula (junior) valde echinata. — *E. valerianæfolia* Desf. Tourn. Coroll. p. 90, tab. 69.

Hab. in insulâ Chios (Oliv. in herb. Parl.).

Species foliorum formâ capsulaeque magnitudine (diam. 2-1 $\frac{1}{2}$ lineas latâ) variabilis, proxima *E. ahenocarpe* Guss. quæ capsula parce et breviter echinulata et in primis intra involucrum subsessili differt.

37. ***E. platyphylla*** (L. Sp. 660) annua plus minusve molliter hirta vel lanuginosa, caule erecto sub umbellâ 5-radiatâ sœpe ramulos floriferos edente, foliis sessilibus subcordatis a medio serrulatis infimis obovatis obtusis superioribus obverse lanceolatis acutis, radiis subtrifidis, foliis floralibus triangulari-ovatis subcordatis mucronatis vel cuspidatis, involuci campanulati intus hirti lobis ovato-oblongis fimbriatis, capsulae sphericæ obsolete trisulcæ verrucis hemisphericis, semine ovato, carunculâ minimâ patellari transverse ovatâ ☽. Jacq. Austr. tab. 376. — Rehb. Ic. Germ. fig. 4758.

Hab. in cultis et ruderatis Græciæ (Marg! Sprunt!), Anatoliæ bor. (Wied!). Tauriæ (Stev!).

β. literata. — Koch. Syn. — Folia subtus vel utrinque densius villosa sœpe fusco-maculata. Capsula hirtula. — *E. literata* Jacq. Coll. II. Ic. Rar. tab. 482.

Hab. in Græciâ formâ vulgari vulgatior (Sart! Heldr! Orph. Fl. exs. 704!). Macedonia (Friv!).

Ar. Geogr. Europa præsertim septentrionalis et media ad Rossiam usque Africa borealis, Madera.

38. ***E. stricta*** (L. Syst. II, p. 1040) glabra vel pilosula, caule erecto sub umbellâ 3-7-radiatâ sœpius ramuloso, radiis 3-5-fidis, foliis leuiter membranaceis a medio serrulatis oblongo-lanceolatis infimis basi attenuatis subpetiolatis obtusis superioribus subcordatis acutis

vel acuminatis, floralibus triangulari-ovatis, involueri turbinato-campanulati intus hirti lobis ovatis fimbriatis, capsulæ sphæricæ trisulcæ verrucis breviter cylindricis, semine oblongo, carunculâ minimâ patellari transverse ovatâ ♂. Ic. Rehb. Germ. fig. 4756-7. — Fl. Græc. tab. 469. — *E. micrantha* M.B. Taur. Cauc. p. 377. — *E. oblongata* C. Koch Linn. XIX, p. 17 non Griseb.

Hab. in umbrosis et ad sepes, Anatolia occidentalis in montosis (Sibth! Boiss! Balt!), Pontus (Balt!), prov. Transcaucasicae (M.B., G.A. Mey!), Persia bor. (herb. Petrop!).

Differt a præcedente foliis minoribus et magis teneris, capsulæ dimidio minoris verrucis cylindricis, semine oblongo triplo minore.

Ar. Geogr. Anglia, Gallia, Helvetia, Germania, Italia borealis, regio Danubialis.

39. **E. coniosperma** (Boiss. et Buhse Aufz. p. 196) annua glabra pumila a basi dichotome ramosissima, foliis a basi attenuatâ subtrinervi oblongis obtusis integris, floralibus ovatis acutiusculis margine obsolete erosulis, involucris dichotomialibus et terminalibus parvis turbinatis lobis triangulari-ovatis laceris, glandulis transverse ovato-oblongis subinæqualibus, capsulæ globosæ trisulcatæ verrucis conico-compressis acutis, semine ovato tenuiter velutino, carunculâ nullâ ♂.

Hab. in Persiâ boreali prope Gamarlu (Buhse).

Planta 3-5-pollicaris facie *E. punctatam* Del. referens, folia 5-7 lineas longa 4 lata, capsula magn. seminis Cannabeos. Ex unico specimine descripta.

2. Species perennes.

40. **E. palustris** (L. Sp. p. 662) glabra, caulis e rhizomate crasso pluribus crassis elatis ex axillis ramos steriles edentibus, umbelle multiradiatae radiis brevibus trifidis bifidis, foliis sessilibus integris vel reproto denticulatis lanceolatis obtusis ramorum linearilanceolatis, floralibus lutescentibus basi attenuatis ovato-ellipticis obtusis, involueri campanulati intus hirti lobis oblongis truncatis denticulatis, capsulâ crassiusculâ globoso-trisulcatâ depressâ verrucis breviter oblongo-cylindricis obsitâ, carunculâ hemispherico-conicâ ♀. Ic. Fl. Dan. tab. 866. — Rehb. Ic. Germ. fig. 4771.

Hab. in paludosis Græciae (Sibth.), Bithyniæ (Wied!).

Ar. Geogr. Europa borealis et media ad Galliam, Italiam borealem, regionem Danubialem et Rossiam medium usque, Sibitia Uralensis et Altaica.

41. **E. Orientalis** (Linn. Annaen. Ac. III, p. 123 et herb!) glabra, caulis crassis rigide virgatis erectis sub umbellâ ramulos floriferos edentibus, umbelle radiis quinis 4-5-fidis, foliis sessilibus rigidis strictis lanceolatis acuminatis integris vel versus apicem obsolete denticulatis, umbellaribus oblongo-lanceolatis, floralibus ovatis vel orbiculatis mucronulatis, involueri hemispherico-turbinati lobis trun-

catis ciliatis, capsula globoso-depressa trisulcata parce pilosula, coccis dorso iterum sulculatione ad utrumque sulculationem latus verrucis brevibus compressis conicis basi in rugulas confluentibus obsitis, semine plumbico ovato, carunculata patellari φ . Boiss. Euph. Ic. tab. 72. — *E. notadenia* Boiss. Diagn. Ser. I, 12, p. 111.

Hab. in humidis Armeniae Turcicie (Tourn!), ad Müküs (Ky!), Armenia Rossica ad Koschadara et in prov. Aderbidjan Persie (Szov!), in iugno Elbrus (Ky. 420!), in montibus calcareis Tchinar et Maregum Persiae austro-occidentalis (Haussk!).

Planta tripedalis, folia 3-3½ poll. longa 5-7 lineas lata formam et rigiditatem ea *Bupleuri Gibraltarici* referentia nervo albo subtus percursa. Capsula 2½ lin. lata.

42. *E. aspera* (M.B. Taur. Cauc. I, p. 377) pubescenti hirta, caulis e rhizomate nodoso-inerassato superne breviter ramulosis ramulis umbellaque radiis 5-8 tenuibus bis vel ter bifidis, foliis subtus intensius et adpresso hirtis brevissime petiolatis a basi rotundata vel subattenuata oblongis et ellipticis integris vel obsolete repandis inferioribus obtusis cæteris acutis, umbellaribus elliptico-rhombeis, floralibus subcordatis reniformi-deltoides glabris, involueri turbinati lobis ovatis dentatis, capsulae breviter ovatae trisulcae inter verrucas hirtæ verrucis cylindrico-filiformibus, carunculata ovato-trigonata φ . Boiss. Euph. Ic. tab. 74. — *E. squamosa* Willd. Sp. II, p. 918! (nomen antiquius sed non congruum). — *F. muricata* M.B. Taur. Cauc. I, p. 378.

Hab. in nemorosis Caucasi sept. in montibus Beschtan et Machuka (Hoh!), Transcaucasiae in Iberia, Cartalimia (M.B., Szov!), Persiae bor. (Buhse!), Capadociae (Tourn!).

β . *oligadenia* Boiss. in D.C. Prodr. — Folia glabrescentia tenuius membranacea, capsula tantum ad dorsum coccorum verrucis paucis brevioribus obsita. — *E. Talyschensis* Boiss. et Buhse Aufz. p. 196.

Hab. in ditione Talysch (Buhse!).

Planta 1-2-pedalis, folia majora 2½-pollicaria. Capsula piso dimidio minor cito dehiscens.

43. *E. macrocarpa* (Boiss. et Buhse Aufz. p. 197) patentim et molliter hirta, caulis erectis dense foliosis sub umbellata ramulosis, ramulis umbellaque radiis 5-7 simplicibus vel bifidis brevibus, foliis a basi sessili amplexicauli subcordata oblongo-lanceolatis obtusis vel acutiusculis subrepandis, floralibus reniformi-orbiculatis obtusis transverse latioribus, involueri breviter turbinati lobis amplis ovato-rotundis denticulatis ciliatis, capsulae magnae globosae subtrisulcae verrucis lutescentibus crebris cylindrico-filiformibus, semine globoso ecarunculato φ .

Hab. in Armenia (Auch. № 2254 in D.C. herb!), in Persia boreali prope Ssanam (Buhse!), in dumetis montium Avroman et Schahu Kurdistaniae Persiae (Haussk!).

Sesquipedalis, folia $1\frac{1}{2}$ -2-pollicaria, inflorescentia subcorymbosa, capsula 4 lineas longa et lata quo charactere et foliis basi cordatis ab affini *E. aspera* distinguitur.

44. *E. epiphyoides* (Jacq. Austr. IV, p. 23, tab. 344.—Linn. herb! et Sp. 656 ex parte) patule albido-villosula, caulis e rhizomate pluribus herbaceis erectis sub umbellâ rarius ramulosis, umbellæ radiis quinis brevibus breviter et compacte trifidis bifidis, foliis sessilibus subintegris oblongis obtusis superioribus subcordatis, umbellaribus conformibus radios æquantibus, floralibus ellipticis vel rotundatis obtusis flavidis, involuci campanulati membranacei lobis tubo æquilongis ovatis truncatis subdentatis, capsulâ ovato-depressâ trisulcâ verrucis filiformibus flavidis vel purpureis obsitâ, carunculâ semilunari ✘. Rehb. Ic. Germ. fig. 4764 (capsulâ male oblongâ). — Bot. Mag. 2258.

Hab. in Macedonia et monte Hæmo Thraciæ (Friv.), in monte Alemdagh Anatoliæ borealis (Noël!).

Differt ab *E. fragiferâ* quâcum a Linnæo sub *E. epiphyoides* confusa caulis herbaceis, involuci lobis elongatis, capsulâ dimidio minore.

Ar. Geogr. Germania orientalis, regio Danubialis, Bosnia et Serbia, Rossia austro-occidentalis.

45. *E. oblongata* (Griseb. Spicil. Rum. I, p. 136) patule villoso-sula, caulis erectis parte inferiore induratis simplicibus vel sub umbellâ ramulos paucos floriferos edentibus, umbellæ radiis quinis abbreviatæ compacte 3-5-fidis dein bifidis, foliis sessilibus oblongo-lanceolatis obtusiusculis versus apicem tenuiter serrulatis infimis basi attenuatis cæteris basi rotundatis, umbellaribus radios subæquantibus tandem deflexis, floralibus ovatis mucronatis, involuci turbinate-campanulati intus hirtuli lobis oblongis truncatis, capsulâ ovatâ verru'is hemisphærico-conicis obsitâ, semine ovato-complanato, carunculâ patellari latere interno retusâ ✘. Boiss. Euph. Ic. tab. 76. — *E. pilosa* Smith Prodr. I, p. 329 ex parte Bory Fl. Pelop. non L. — *E. erioclada* Boiss. et Heldr. Diagn. Ser. I, 7, p. 88.

Hab. in sylvaticis Peloponnesi in Messeniâ et Laconiâ (Bory! Boiss! Orph. exs. 701!), Olympi Thessali (Heldr!), montis Athos (Griseb!), Byzantii (Noël!), insula Naxos (Despr!), Cretæ (Raul!).

β. sessiliflora. — Boiss. in D.C. Prodr. — Humilior, folia abbreviata, umbella ob radios umbellatos abbreviatos capituliformis. — *E. sessiliflora* Boiss. et Sprun. Diagn. Ser. I, 5, p. 52.

Hab. in monte Delphi Eubeæ (Sprun!).

Facies *E. pubescens*, umbellæ parvæ compactæ, glandulæ involuci sæpe tantum 2-3. Capsula $1\frac{1}{2}$ lineas lata.

46. *E. condylocarpa* (M.B. Taur. I, p. 377) glabra, caule ex radice tuberosâ ovato-globosâ unico crebre folioso superne sub umbellâ ramuloso thyrsideo, ramulis umbellæque radiis 5-8 brevibus bifidis, foliis sessilibus a basi latiore cordato-amplexicauli lanceolatis

serrulatis, superioribus abbreviatis oblongo triangularibus, floralibus a basi truncatâ semiorbiculari-reniformibus transverse latioribus obtusissimis, involuci hemispherico-turbinati lobis latis brevibus transverse ovatis, capsulâ breviter stipitatâ subsphæricâ verrucis hemispherico-conicis obsitâ, carunculâ ovato-triangulari ♀. Boiss. Euph. Ic. tab. 77. — *E. amplexicaulis* Ledeb. Fl. Ross. 3, p. 567!

Hab. in coltibus aridis argillosis Caucasi et Transcaucasiae (M.B., C.A. Mey! Hohen!), Armeniæ (Gundelsh. in Berol!), Persie borealis (Buhse!).

Radix nuce major, planta 1-1 $\frac{1}{2}$ -pedalis, folia majora pollicem longa.

47. *E. cardiophylla* (Boiss. et Heldr. Diagn. Ser. I, 12, p. 107) glabra, caulis e rhizomate verticali apice pluricipiti tenuibus erectis sæpius ramulosis thyrsoideis, foliis sessilibus præter infima basi attenuata profunde cordato-auriculatis ovato-oblongis acutis mucronulatis obsolete serrulatis, floralibus orbiculatis obtusis transverse sublatioribus, umbellæ radiis 3-4 tenuibus bifidis, involuci hemispherici lobis transverse ovatis, capsulâ longiuscule stipitatâ ovatâ secus coccorum dorsum verrucis hemisphericis depresso obsitâ cæterum laevi, carunculâ subsphæricâ ♀.

Hab. in fruticetis regionis inferioris montis Solyma Lyciae infra Kartsibahir (Heldr!), in regione inferiori Olympi supra Brussam (Pichl!).

Planta 8-12-pollicaris, folia pollicaria 5-6 lineas lata; a præcedente distincta radice, capsulâ longiuscule stipitatâ, etc.

48. *E. Apios* (L. Syst. p. 375) sparsim hirtula vel glabrescens, radice napiformi caudiculum brevem subterraneum squamuulosum apice ramosum edente, ramis decumbentibus vel ascendentibus pumilis tenuibus foliosis sub umbellâ sæpe ramulosis, radiis 3-5 simplificibus vel bifidis, foliis parvis a basi sessili subattenuata vel rotundata elliptico-oblongis serrulatis obtusis, floralibus transverse ovatis vel rhombeis retusis, involuci turbinato-campanulati glabri lobis ovatis subdentatis, capsulæ ovatae trisulcæ verrucis breviter cylindricis purpurascensibus, semine ovato subcompresso rubello opaco, carunculâ orbiculari depressâ ♀. Ic. Clus. Hist. 2, p. 190. — Boiss. Euph. Ic. tab. 78. — *E. pubigera* Friv. exs.

Hab. in collibus et sylvaticis regionis montanæ Peloponnesi et Atticæ (Sprun! Boiss! Heldr!), Macedoniæ (Friv.), insulæ Chios (Oliv!), Cretæ (Sieb. forma depauperata).

β. lamprocarpa Boiss. in D.C. Prodr. — Folia subintegra, fructus verrucæ hemispherice magis sparsæ, semen fuscum nitidum.

Hab. in monte Akdaghi Cataoniæ (Auch. 2040!), in pinetis Cilicie (Bal!), Syria in monte Amano prope Beilan (Haussk!), Antilibano inter Rascheyâ et Damascum (Boiss!). Fl. vere.

Caules 4-6-pollicares, folia 4-6 lineas longa, capsula majuscula.

Ar. Geogr. Regnum Neapolitanum.

49. *E. dulcis* (Jacq. Austr. III, p. 8, tab. 213) glabra vel parce puberula, rhizomatis horizontalis crassiusculi articulis carnosis bre-

viter concatenatis, caulis 1-2 tenuibus teretibus subsimplicibus, umbellæ radiis quinis tenuibus bifidis, foliis insimis squamæformibus cæteris breviter petiolatis membranaceis integris vel versus apicem serrulatis inferioribus basi attenuatis obovatis cæteris a basi rotundata lanceolato-oblongis obtusis, floralibus triangulari-ovatis basi truncatis, involuci breviter turbinati lobis ovatis dentatis, glandulis purpureis vel virentibus, capsula depressa profunde trisulca verrucis purpureis inæqualibus breviter cylindricis sparsim obsitâ, carunculâ stipitatâ depressa ♀. Rchb. Ic. Crit. tab. 143. — *E. dulcis* L. Sp. 656 et herb. ex parte, — *E. viridiflora* W. K. tab. 280. — *E. alpigena* Kern.

Hab. in umbrosis fagetorum Chalcidices Macedoniæ (Griseb.). E ditione nondum vidi.

Ar. Geogr. Europa media in Galliâ, Hispaniâ boreali, Helvetiâ, Germaniâ, Italia boreali, regione Danubiali, Rossia mediâ et australi.

50. ***E. Djimilensis*** (Boiss. in Bul. exs. 1866) caulis e rhizomate pluricipiti erectis simplicibus minute et patule puberulis, umbellæ capituliformis radiis 3-5 brevissimis, foliis glabris parvis sessilibus a basi attenuatâ elliptico-spathulatis integris obtusis nervo albo percursis, floralibus ovatis flores æquantibus, involuci turbinati lobis dentatis, capsula longiuscule stipitatâ globosâ trisulca verrucis parvis in lineolas confluentibus obsitâ, semine albido, carunculâ minimâ patellari ♀.

Hab. in regione alpinâ Ponti Lazici prope Djimil 6600' (Bal!). Fl. Ju.

Planta pedalis, folia majora vix pollicaria 5-6 lineas lata, umbella diametro 6-12 lineas lata involucris 3-9, capsula 2 lineas circiter lata. Facies *E. verrucosa*, folia minora spathulata, caules patule pubescentes nec glabri, umbellæ minores magis coarctatae, capsula duplo major.

51. ***E. thamnoides*** (Boiss. Cent. Euph. p. 33. — D.C. Prodr. XV, p. 131) glabra suffruticosa a basi dumosa, plantæ junioris rhizomate napiformi, ramis virgatis rigidulis crebre foliatis simplicibus, umbellæ radiis quinis breviter 2-3-fidis, foliis subpetiolatis oblongis acutiusculis integris basi attenuatis, floralibus obovatis obtusissimis, involuci hemispherico-campanulati intus valde hirti lobis late ovatis denticulatis, stylis brevissimis bifidis, capsula sphæricâ trisulca verrucis conico-cylindricis crebre obsitâ, semine rufo nitido, carunculâ globoso-depressa ♂. Boiss. Euph. Ic. tab. 80. — *E. dumosa* Boiss. Diagn. Ser. I, 12, p. 110 non *E. Mey.* — *E. dulcis* Sibth. Fl. Græc. V, p. 50, tab. 464 non L.

Hab. in collibus regionis inferioris Anatoliæ circa Ephesum (Auch. 877!), Pamphyliæ ad Adalia (Pest! Bourg!), Cypri (Labill!), in totâ Syriâ littorali et regione inferiori Libani frequens (Boiss! Gaill! Bl! etc.).

♂. *Hierosolymitana* Boiss. in D.C. Prodr. — Folia obovata, obtusiora, capsula paulo major. — *E. Hierosolymitana* Boiss. Diagn. Ser. I, 12, p. 110.

Hab. ad Hierosolymam (Auch. 2035! Boiss!), in Antilibano (Gaill! Ky!).

Dumus 2-3-pedalis, napulus hornotinus obverse pyriformis basi in radicem longe attenuatus, folia 6-8 lineas sepius longa. Facies *E. spinosa* sed inermis, ab affini *E. Bivoneae* foliis brevibus, stylis valde abbreviatis distincta.

52. *E. acanthothamnos* (Heldr. et Sart. in Boiss. Diagn. Ser. II, 4, p. 86) glabra suffruticosa dumulosa ramosissima, ramis brevibus intricatis rigidis, umbellæ radiis 3-4 brevibus semel vel bis divaricatim bifidis anthesi peractâ in spinas dichotome plures furcatas persistentes mutatis, foliis minutis integris oblongo-lanceolatis obtusiusculis, floralibus obovatis, involueri breviter turbinali intus hirtali lobis truncatis subdentatis, capsulæ globosæ profundè trisulæ verrucis crebris hemispherico-cylindricis, semine ovato ȝ. *E. spinosa* Sibth. Fl. Græc., V, p. 49, tab. 463. Griseb. Spic. I, p. 137 non L.

Hab. in regione sempervirenti totius Græciae (Boiss! Heldr! Sart!), Macedoniae in Peninsulâ Athoâ (Griseb!), Archipelago (Urv!), Cretâ (Sieb!), Rhodi (Auch! Bourg!), Bithyniâ (Wied!).

Semipedalis et bipedalis, facies *Poterii spinosi*, folia 3-5 lineas longa. Ab *E. spinosa* egregie distincta spinis non ex extremitate indurata ramorum sed ex radiis oriundis, multo tenuioribus furcatis nec simplicibus, involueris longius pedunculatis.

53. *E. erinacea* (Boiss. et Ky. Diagn. Ser. II, 4. p. 87) glabra suffruticosa dumulosa ramosissima, ramis intricatis brevibus rigidis crassiusculis apice dichotomis, ramulis monocephalais post anthesin induratis et spinescentibus, foliis minutis oblongo-lanceolatis acutis argute serrulatis, floralibus diminutis, involueri hemispherico-turbinati intus hirsuti lobis ovatis ciliatis, stylis brevibus bilobis, capsulâ longe stipitatâ ovatâ trisulcâ verrucis depresso hemispherici sparsis obsitâ, semine ovato plumbeo, carunculâ subsphæricâ ȝ. Boiss. Euph. Ic. 81.

Hab. in regione alpinâ Antilibani, vallis Martsch supra Zebdani 5900' (Ky. 105!), mons Gebel Scheick 6-9000' (Gaill!).

Dumus humilis semipedalis, folia 4-6 lineas longa 2½ lata, involueri dichotomiarum pedunculus 2-4 lineas longus. Capsula major eâ *E. spinosa* a quâ ut et a præcedente foliis serrulatis et inflorescentiâ differt.

54. *E. Schottiana* (Boiss. Diagn. Or. Ser. II, 4. p. 85) glabra ex rhizomate lignoso cespitosa, ramis herbaceis cespitosis tenuibus brevibus ascendentibus basi squamosis caeterum foliosis monocephalais vel apice breviter bifidis dicephalais, foliis parvis petiolatis basi rotundatis ovatis obtusis minute serrulatis, floralibus conformatibus, involueri hemispherico-campanulati intus hirtuli lobis ovatis retusis ciliatis, stylis elongatis bifidis, capsulâ (juniore) globosâ longe stipitatâ verrucis hemispherici crebre obsitâ ȝ. *E. densa* Schott et Ky. in Sched. non Schrenk.

Hab. in alpinis jugi Metdesis Tauri Cilicici 10000' (Ky. 1471).

Rami creberrimi 2-4-pollicares, folia 3-4 lineas longa 2½-3 lata. Capsula majuscula. Facies *Thymi montani* vel *E. chamæbuxus*.

55. *E. capitulata* (Rchb. Fl. exc. 873) glabra, caudiculis longis fragilibus radicantibus subterraneis, ramis cespitosis herbaceis brevibus dense foliosis aliis sterilibus aliis apice monocephalitis, foliis minutis obovatis obtusis integris, involuci feliis summis suffulti hemispherici intus parce hirtuli lobis octo oblongis ciliatulis denticulatis, glandulis octo subtropitatis purpureis transverse oblongis, stylis brevissimis, capsula subsessili sphæricâ trisulcâ verrucis ovatis vel cylindricis inflato-vesiculosis obsitâ, semine nitido, carunculâ patellari latere interno retusâ ♀. Boiss. Euph. Ic. tab. 83. Rchb. Ic. fig. 4759. — *E. soliflora* Visian.

Hab. in rupestribus apricis regionis alpine et sylvaticæ superioris in monte Olympo Thessaliae 5000'-7000' (Heldr. Orph. Fl. exs. 703!).

Rami 2-4-pollicares, folia saepius imbricata 3-4 lineas longa, capsula fere magnit. Pisi. Species inflorescentia monocephala et involucro octolobo insignis.

Ar. Geogr. Alpes Dalmatiæ, Bosniæ, Herzegovinæ.

** Semina minute tuberculata.

56. *E. pubescens* (Vahl Symb. II, p. 55) perennis villosa, caulibus erectis vel ascendentibus inferne tandem induratis et denu-datis cæterum crebre foliatis, umbellæ radiis quinis trifidis dein bisidis, foliis sessilibus serrulatis oblongo-lanceolatis vel linearibus acutis vel acuminatis, floralibus a basi subcordata ovato-rhombeis, involueri campanulati extus hirsuti lobis ovatis truncatis ciliatis, glandulis extus et margine ciliatulis, capsula sphæricâ trisulcata inter verrucas crebras breviter cylindricas villigera, semine ovato lineolis brevibus tuberculiformibus tenuiter exasperato, carunculâ minimâ transverse ovatâ ♀. Ic. Jacq. Ecl. I, p. 98, tab. 76. — Rchb. Ic. Germ. fig. 4769. — *E. verrucosa* Linn. Sp. 658 et herb. ex parte. — *E. pilosa* Smitu Prodr. I, p. 315 non L.

Hab. in humidis, cultis regionis inferioris et montanae frequens, Græcia! Anatolia maritima borealis et australis! Syria! Palestina!

Hæc species primo anno saepè floret et inde a cl. Vahl erroneè annua. Seminis tuberculi interdum obsoleti.

Ar. Geogr. Regio Mediterranea Europæ et Africæ borealis, insulæ Canarienses.

*** Semina areolatim reticulato-rugosa.

1. Perennes.

57. *E. rhytidosperma* (Boiss. et Bal. Diagn. Ser. II, 4, p. 84) perennis, caulibus virgatis erectis proceris inferne ramosis denudatis et induratis fere fruticosis, umbellæ radiis quinis gracilibus trifidis dein bisidis, foliis laxe hispidulis minutissime serrulatis a basi attenuatâ subpetiolatâ oblongo-ellipticis obtusis, floralibus a basi truncato-cordata ovato-orbiculatis, involueri turbinati lobis oblongis truncatis ciliatis, capsula sphæricâ depressâ profunde trisulcata secus

cocorum dorsum bifarium verrucis conico-cylindricis, obsoletâ semine lineolis tenuibus in rugas reticuliformes dispositis obsoito, carunculâ patellari depressâ ♀. Boiss. Euph. Ic. tab. 85.

Hab. in fauibus Guzelderè Ciliciae littoralis (Balt.). Vid. sicc. sp. et viv. cult.

Bipe latis, rami tenues, folia 10-15 lineas longa 4-6 lata, radii 3-4-pollicares, facies *E. Bivoneæ*.

2. Annuae.

58. **E. pterococca** (Brot. Fl. Lusit. II, p. 312) annua glabra, caule tenui erecto, umbellæ radiis quinis tenuibus semel vel bis bifidis, foliis tenuiter membranaceis serrulatis oblongis vel obovatis obtusis inferioribus in petiolum brevem attenuatis, floralibus diminutis rhomboeo-ovatis, involuci campanulato-iurbinati lobis ovatis subtruncatis, glandulis transverse oblongis, capsule glabrae globoso-depressæ coecis dorso longitudinaliter bialato-cristatis, semine tenuissime areolato-reticulato ecarunculato ♂. Ic. Brot. Phyt. Lus. tab. 76. — Rehb. Pl. Crit. tab. 464. — *E. stellulata* Sa'zm. Flora 1821, p. 110 non Loisel. — *E. bialata* Link Enum.

Hab. in collibus herbidis regionis inferioris rara, in Zacyntho (Marg!) tantum lecta.

Ar. Geogr. Lusitania, Hispania australis, Corsica, Sardinia et Sicilia, Africa borealis, insule Canarienses.

59. **E. helioscopia** (L. Sp. 658) annua, cauibus pumilis crassiusculis erectis vel ascendentibus inferne tandem denudatis, umbellæ radiis quinis brevibus pubescentibus trifidis dein bifidis, foliis glabriusculis obovatis cuneatis vel spathulatis obtusis serrulatis in petiolum brevem attenuatis, umbellaribus magnis, floralibus ellipticis vel orbiculatis, involuci turbinati lobis oblongis fimbriatis, capsule glabrae laevis globosæ trisulæ coecis dorso rotundatis, semine ovoideo apice acuto ancipiti favoso-reticulato, carunculâ transverse ovata fere verticali ♂. Ic. Fl. Dan. tab. 725. — Rehb. Ic. Germ. fig. 4754.

Hab. in ruderatis et cultis totius ditionis a Græciâ! ad regiones Caucasicas! Persiam medium et australem (Bgel), Syriam littoralem et interiorem ad Damascum (Gaill!), Arabia petream (Schimpl!), Mesopotamiam (Haussk!), Beluchiam (Stocks!).

β. Haussknechti. — Folia umbellaria et superiora acutiora, capsula subminor, semen minus acutum carunculâ rudimentariâ. — *E. Haussknechti* Boiss. MSS.

Hab. in cultis circa Aleppo et inter Aintab et Nisib (Ky. 115! Haussk!).

Ar. Geogr. Europa omnis, Japonia, Africa borealis. In alias regiones v. gr. Americam borealem et tropicam, P.-B. Spei, etc. introducta.

60. **E. Berythea** (Boiss. et Bl. Diagn. Ser. II, 4, p. 82) annua, caulibus a collo 1-3 hispidulis pumilis, umbellæ radiis quinis brevibus bifidis hispidis, foliis subintegris secus nervum medium pilo-

sulis infimis obovatis obtusiusculis in petiolum brevem attenuatis, superioribus et umbellaribus subsessilibus oblongo-ellipticis acutis argute denticulatis, floralibus triangularibus diminutis, involuci turbinati intus hirsutissimi lobis latis truncatis ciliatis, capsula globosâ depressâ profunde trisulcâ pilis patulis hispidâ sub lente asperulâ, semine globoso tuberculis seriatis subconfluentibus alveolato-reticulato, carunculâ minimâ globoso-conicâ ♂. Boiss. Euph. Ic. tab. 86.

Hab. in rupestribus calcareis Syriæ littoralis ad Berythum et Sidonem (Blt Gail!).

Ab *E. helioscopiâ* umbellis minutis magis compactis, capsula pilosa, semine non acuto, reticulis seminis non continuis, carunculâ, distincta.

61. **E. oxyodonta** (Boiss. et Haussk. in Haussk. exs. 1867) annua pilis longis patulis sparsis hispidula, caulis e collo pluribus nanis erectis et decumbentibus simplicibus, foliis profunde et acute pectinato-dentatis infimis minoribus ovalo-spathulatis in petiolum brevem attenuatis caeteris sessilibus ellipticis, floralibus ovatis, involuci turbinati glabri lobis latis ciliatulis, capsula glabra globosâ depressâ profunde trisulcatâ, semine ovato obtuso elevatim favoso-reticulato, carunculâ hemisphericâ latere interiori retusa ♂.

Hab. in petrosis et fissuris calcareis ad Turmanin, Ain el Beitha, Gebel Muhsan prope Aleppo et in Mesopotamia prope Orfa ad Nimruddagh et Gebel Taktak (Haussk!). Fl. Mart.

Caules 3-4-pollicares, folia majora 6-8 lineas longa. Ab affini *E. Berytheâ* foliis acute pectinatis, capsula glabra, seminis cristis magis prominentibus distincta.

**** Semina transverse foveolato vel sulcato-exarata.

1. Perennes.

62. **E. Cassia** (Boiss. Diagn. Ser. I, 12, p. 108) glabra basi suffrutescens multicaulis, caulis tenuiter virgatis erectis rigidulis crebre foliosis subsimplicibus vel superne dichotome ramulosis, umbellis parvis breviter et inaequaliter 2-3-radialis interdum ad involucrum solitarium reductis, radiis brevissime bifidis interdum moncephalis, foliis parvis sessilibus strictis vernes apicem minute et remote serrulatis inferioribus oblongo-spathulatis obtusis superioribus lineari-lanceolatis acutis, umbellaribus radios subaequantibus, floralibus a basi valde attenuata ovato-rhombeis acuminatis, involuci turbinato-campanulati lobis late et breviter ovatis ciliatis, glandulis transverse oblongis, capsulae depressae laevis profunde trisulcate coccis dorso subcarinatis, semine flavido foveolis vel lineis transversis brevibus nigris superficialiter exarato, carunculâ minimâ vel nullâ ♀. Boiss. Euph. Ic. tab. 88.

Hab. in umbrosis regionis montanae Syriæ, mons Cassius inter Latakieh et Suadieh (Boiss!), Libanus ad Beckfaya (Blt!) et Beitkhaschbo (Gail!), supra Berythum (Ehr!), Cyprus (Ky. 923!).

Caules 1-1½-pedales, folia 5-9 lineas longa inferiora 1½-2 lata, inflorescentia macra radiis 1-2-pollicaribus saepe multo brevioribus, capsula lineam longa. Habitus *Esulæ* v. gr. *E. Gerardianæ*, sed glandulæ margine rotundatæ nec truncatae.

2. Annuæ.

63. E. phymatosperma (Boiss. et Gaill. Diagn. Ser. II, 4, p. 83) annua glabra glauca nana, caule erecto simplici, umbellæ radiis subternis brevissimis breviter bifidis, foliis parvis sessilibus obovatis obtusis integris, umbellaribus ellipticis submajoribus, floralibus diminutis ovato-rhombeis mucronulatis, involueri parvi campanulati lobis late ovatis obtusissimis, glandulis transverse oblongis, stylis longis breviter bifidis, capsulae ovatae coccis dorso rotundatis, semine ovato obtuse tetragono subcompresso ad unamquamque faciem soleis tribus transversis exarato prætereaque minute albo-leproso, carunculâ breviter stipitatâ depressâ ♂.

Hab. in monte calcareo Gebel Khaisoun prope Damascum (Gaill!), in lapidosis ad Aleppo (Auch! 2686!), in monte Nimiruddagh prepe Orfa et ad Terek Mesopotamie, in montibus Gebel Taktak et Pir Omar Guîrun (Haussk!).

Plantula 3-4-pollicaris, folia caulina tres lineas umbellaria 5 lineas longa, umbella contracta diametro 1-1½-pollicaris, semen eo *E. helioscopiae* duplo magius.

§ 7. Esulæ.

* Semina tuberculata. — Annuæ.

64. E. Aleppica (L. Amæn. III, p. 122) annua glauca glabra vel pruinosa-scabridula a collo simplex vel pluricaulis, caulinis erectis crassiœculis densissime foliosis inferne tandem denudatis et cicatricosis sub umbellâ saepius ramuligeris, umbellæ radiis quinis crassiœculis pluries bifidis divaricatis, foliis integris sessilibus mucronatis ab inferioribus setaceis ad superiora anguste linearia sensim dilatatis, umbellaribus lanceolatis, floralibus ovato-rhombeis aristatis, involueri turbinato-campanulati fance hirti lobis parvis oblongis fimbriatis, glandulis semilunatis transverse latioribus, capsulae parvæ depressæ profunde trisulcatæ coccis carinatis, semine atro ovato-quadrangulo minute et elevatim tuberculato ecarunculato ♂. Ic. Fl. Gr. tab. 462. — Rehb. Germ. fig. 4779. — *E. pinifolia* Willd. Suppl. — *E. condensata* Fisch. in M.B. III, p. 322.

Hab. in cultis et vinetis, Graecia in Argolide, Achaiâ, Bœotia (Sprunt! Heldr!), Cycladibus (Bory!), Cretâ (Heldr!), Rhodo (Bourg!), Macdoniâ (Friv!), Anatoliâ occidentali, meridionali et interiori (Boiss! Heldr! Wied!), Syriâ littorali et interiori! Palestinâ! Mesopotamia! Assyriâ! Transcaucasia (Hohen!), Persiâ occidentali (Noë!).

Ar. Geogr. Italia australis, Dalmatia.

65. E. parvula (Delile Egypt. p. 290, tab. 30, fig. 4) glabra nana, caule nano vel inferne ramulosso, umbellæ radiis binis vel ternis

brevibus plures bifidis, foliis integris obovato-spathulatis obtusis vel retusis mucronulatis, umbellaribus et rameis oblongo-spathulatis, floralibus breviter oblongo-lanceolatis acutis, involuci minimi turbinati lobis oblongis ciliatulis, glandulis minimis concaviusculis transverse ovatis brevissime bicornutis cornubus interdum obsoletis, capsulae minimaæ globoso-depressæ trisulcatæ coccis dorso rotundatis, semine ovato-globoso minute albo-tuberculato ecarunculato ♂.

Hab. in incultis Egypti inferioris prope Alexandriam (Dell! Ehrenb! Samar!).

Facies *E. exiguae* vel *Szovitsii* a quibus differt formâ foliorum caulinorum, cornubus brevissimis, semine non tetragono et carunculâ destituto.

66. ***E. exigua*** (L. Amæn. III, p. 418) glabra minuta a collo simplex vel ramosa, caulis crebre foliosis erectis, umbellæ radiis 3-5 plures dichotomis, foliis linearibus strictis acutis vel obtusiusculis, umbellaribus et floralibus a basi dilatata sâpe subcordatâ trianguli-lanceolatis acutis, involuci turbinati fauce hirtuli lobis ovatis dentatis, glandulis semilunatis transverse latioribus divergentibus latitudine glandulæ longioribus, capsula ovatâ trisulcâ, semine albido ovato-quadrangulo minute tuberculato, carunculâ depressâ ♂. Ic. Fl. Dan. tab. 592. — Rchb. Germ. fig. 477.

β . *retusa* Roth. — Folia apice truncata vel retusa. — *E. retusa* Rchb. Germ. fig. 4778.

γ . *tricuspidata* Koch Syn. — Folia apice dilatata tridentata. — *E. tricuspidata* Lapeyr. Abr. p. 271. — *E. rubra* D.C. non Cav. — *E. diffusa* Jacq. Misc. II, p. 311. Ic. Rar tab. 88.

Hab. typus cum varietatibus ubique in cultis et collibus siccis regionis inferioris et montanæ a Graciâ! et Macedonia! ad Syriam! et prov. Transcaucasias! Var. β et γ potius vernales.

Ar. Geogr. Europa tota media et australis ab Angliâ, Daniâ et Gothiâ ad Rossiam medium, Africa borealis, insulæ Canarienses.

67. ***E. dracunculoides*** (Lam. Dict. II, p. 428) annua, caulis sâpius pluribus erectis ex axillis sâpius ramulosis, umbellæ radiis 2-3 elongatis plures dichotomis, foliis sessilibus basi attenuatis anguste linear-lanceolatis acutiusculis, floralibus brevioribus . basi latiore lanceolatis, involuci turbinati intus hirti lobis ovatis denticulatis ciliatis, glandulis semilunatis transverse latioribus cornubus latitudine glandularum longioribus vel æquioribus, capsula depressâ profunde trisulcatâ, semine oblongo-nigrantic tuberculis albidis basi sâpe lineatim confluentibus leproso, carunculâ depressâ transverse oblongâ ♂. Boiss. Euph. Ic. tab. 91. — *E. lanceolata* Rottl. in Spr. Syst. III, p. 791. — *E. glauca* Ehrenb. MSS.

Hab. in cultis Arabiæ Mascatensis (Auch. 5305!), forsitan in Egypto superiore maritimâ. Species subtropica vix ditionis.

Planta 1-1½-pedalis, folia caulina 1-1½-pollicaria 1½-2 lin. lata.

Ar. Geogr. Nubia maritima, Arabia tropica, India.

** Semina transverse sulcata. — Annuæ

68. **E. falcata** (L. Sp. 654) annua glabra glaucescens simplex vel multicaulis pumila, caulis erectis vel ascendentibus, ramulis umbellæque radiis ternis rarius 4-5 iteratim bifidis, foliis integris margine scabriuseulis basi attenuatis infimis oblongo-spathulatis obtusis vel retusis cæteris lanceolatis acutis, umbellaribus oblongis, floralibus ovatis vel ellipticis mucronato-acuminatis obliquis, involueri turbinati lobis ovatis simbriatis, glandulis transverse latioribus cornubus longis, capsulae ovatae coccis carinatis, semine oblongo-quadrangulo lateraliiter compresso ad facies transverse et regulariter 4-6-sulcato, carunculâ minimâ vel saepius nullâ ☽. Ic. Jacq. Austr. II, tab. 121. — Rehb. Germ. fig. 4776. — *E. acuminata* Lam. Dict. II, p. 426. — *E. obscura* Loisel. Not. p. 76, tab. 5, fig. 2. — *E. mucronata* Lam.

Hab. in cultis totius ditionis a Græciâ! ad Tauriam! prov. Cauca-icas! Syriam! Persianam bor. et australiem! Affghaniam (Griff!) et Belutschiam (Stocks!).

β . *rubra*. — Minor, tota rubens. — *E. rubra* Cav. Ic. I, tab. 34.

Hab. in Græciâ circa Athenas (Sprun!).

γ . *Galilaea* Boiss. in D.C. Prodr. — Folia floralia parva dissita, capsula minor longius stipitata, seminis dimidio minoris sulci abbreviati ad foveolas reducti. — *E. Galilaea* Boiss. Diagn. Ser. I, 12, p. 116.

Hab. in cultis Anatolie ad Tokat (Wiedem!), Ciliciae (Bal!), Syriae ad Sidonem (Boiss!), Galileæ (Boiss!), prov. Caucasicarum (Szov!).

δ . *ecornuta*. — Glandulæ cornua brevissima vel obsoleta. — *E. caudata* Boiss. et Haussk. MSS.

Hab. in calcareis Gebel Muhsan prope Aleppo Syriae et in deserto fluvii Chabur Mesopotamiae (Haussk!).

Ar. Geogr. Europa media et australis a Gallia et Germania mediâ ad Dalmatiam et regionem Danubialem, Africa borealis.

*** Semina longitudinaliter sulcata vel seriebus longitudinalibus foveolarum regulariter instructa. — Annuæ.

69. **E. aulacosperma** (Boiss. Diagn. Ser. I, 12, p. 117) annua glabra pumila, caulis erectis saepius ramosis, umbelle radiis ternis plures dichotomis, foliis membranaceis obsolete denticulatis a basi attenuatâ obovatis obtusis vel retusis, floralibus triangulari-rhombeis obliquis saepè transverse latioribus, involueri turbinati intus hirti lobis ovatis truncatis ciliatis, glandularum cornubus ejus latitudine 2-3-plo longioribus, capsula ovata depressâ profunde trisulcâ, semine ovato truncato longitudinaliter nigro-sexsulcato sub lente tenuissime asperulo, carunculâ depressâ ☽. Boiss. Euph. Ic. tab. 92.

Hab. in cultis Ciliciae ad Mersina (Bal!), Syriae ad Berythrum et Tripoli (Bl!), Damascum (Gaill!), Judeæ ad Hierosolymam (Boiss!).

β . *fossulata* Boiss. in D.C. Prodr. — Minor, seminis abbreviati

crassioris sulci concolores. — *E. fossulata* Boiss. et Gaill. Diagn. Ser. II, 4, p. 87.

Hab. in lapidosis ad Damascum (Gaill !).

Planta facie *E. Pepli* similis ab *E. sulcatâ* Lens cui seminis sculpturâ proxima est foliis obovatis nec linearibus, cornubus elongatis, seminibus majoribus asperulis nec lèvibus distincta.

70. *E. Peplus* (L. Sp. 658) annua glabra ramosa erecta, umbellæ radiis ternis plures bifidis, foliis tenuiter membranaccis integris in petiolum attenuatis obovalis obtusis vel retusis, floralibus sessilibus ovatis subobliquis, involuci turbinati lobis ovatis fimbriatulis, glandulæ cornubus tenuibus ejus latitudine multo longioribus, stylis brevibus profunde bifidis, capsulæ breviter ovatae profunde trisulcæ coccis compressis dorso angustissime alato-bicarinatis, semine albido ovato-subhexagono ad facies binas interiores sulco exarato ad quatuor exteriore lineâ 3-4 foveolarum rotundarum exsculpto, carunculâ depresso conicâ ⊖. Ic. Fl. Dan. tab. 1100. — Rchb. Germ. fig. 4775.

Hab. in cultis Graecie! Macedoniæ! Anatoliæ bor. (Wiedem !), Armeniæ (Bourg !), Syriae littoralis (Bl! Gaill!), Arabiæ petreæ (Schimp. 249!), Egypti inferioris in Oasibus Farafrah et Dachel (Asch !), Persiæ orientalis et australis (Bge! Ky.!). Fl. aestate.

E. maritima Boiss. in p.c. Prodri. — Minor decumbens, folia crassiora magis rotundata. — *E. peploides* Griseb. Spicil. I, p. 138 non Gouan.

Hab. in maritimis Byzantii (Clementi !), Bithyniæ (Griseb !), ad Alexandriam (Samar !).

Ar. Geogr. Europa omnis ab Angliâ et Succiâ ad Rossian medium, Sibîria, Africa borealis, Insule Canarienses. In Americam bor. et tropicam P. B. Spei, etc., introducta.

71. *E. peploides* (Gou. Fl. Monsp. p. 174) glabra ramosa pumila, umbellæ radiis 3-5 abbreviatis plures bifidis, foliis integris suborbiculatis basi in petiolum attenuatis, floralibus sessilibus ovatis, involuci turbinati lobis ovatis ciliatis, glandulæ cornubus ejus latitudine sublongioribus, stigmatibus subsessilibus, capsulæ breviter ovatae trisulcæ coccis bicarinatis, semine albido breviter ovato-hexagono ad lineas facies interiores sulco exarato ad quatuor exteriore lineâ 2-3 foveolarum exsculpto, carunculâ depresso conicâ ⊖. Rchb. Ic. Germ. fig. 4775. — *E. rotundifolia* Lois. Not. p. 75, tab. 5, fig. 4. — *E. Peplus* β minor Willd. Sp. II, p. 903.

Hab. in umbrosis et cultis Atticæ (Sart!), Peloponnesi (Bory!), Cretæ (Raul!), Syriae littoralis (Bl!). Fl. primo vere.

Valde affinis precedenti cuius forsan forma vernalis, differt foliis rotundioribus, stylis subnulis, seminis dimidio minoris foveolis paucioribus; characteres culta servat.

Ar. Geogr. Gallia et Hispania australes, Italia maritima, Dalmatia, Africa borealis.

72. E. chamæpeplus (Boiss. et Gaill. Diagn. Ser. II, 4, p. 88) glabra, caulinis natis erectis, umbellæ minutæ radiis tribus brevibus bifidis, foliis minutis subsessilibus ovatis obtusis vel retusis, floralibus oblengis, involueri breviter turbinati lobis ovatis truncatis fimbriatulis glandularum cornubus tenuibus ejus latitudine sublongioribus, stylis ad medium bifidis, capsulae ovatae coccis dorso rotundatis, semine cylindrico-hexagono ad facies binas internas longitudinaliter sulcato ad 4 interiores lineâ 4-5 foveolarum rotundarum insculpto, carunculâ rotundatâ depressâ ⊖.

Hab. in Palestinâ circa Hierosolymam (Boiss!), in Syria circa Damascum (Gaill!).

β. Sinaica Boiss. in D. C. Prodr. — Seminis obscure hexagoni foveolæ subordinatum dispositæ. — *E. Sinaica* Hochst. in Sched. pl. Schimp.

Hab. in graniticis nontis Stæ Catherine Arabie petreæ (Schimp. 234!), in deserto Ægyptiaco prope Pelusium (Ky. 511!).

Species omnibus partibus *E. Peplo* et *peptoide* minor, ab eis capsulae longius ovatae coccis dorso non bicarinatis, stylis duplo longioribus, semine angustiore cylindrico discedens.

**** Semina irregulariter foveata, malleata vel reticulato-rugosa.

1. Annuæ.

73. E. Ledebourii (Boiss. Cent. Euph. p. 35. — D. C. Prodr. XV, p. 143) annua glabra, caule humili erecto rigidulo, umbellæ radiis ternis rarius 4-5-nis elongatis semel vel bis bifidis, foliis integris ab infimis linearispathulatis saepe retusis ad superiori linearia sensim elongatis, floralibus longis anguste linearibus acutiusculis, involueri minimi turbinati lobis ovatis, glandulae cornubus subconvergentibus ejus latitudine sublongioribus, capsulae ovatae sub lente minute granulosæ, semine ovato-quadrangulo inæqualiter et subconfluenter foveolato-facunoso, carunculâ minutâ conicâ ⊖. Boiss. Euph. Ic. tab. 95. — *E. pygmaea* Ledeb. Fl. Ross. III, p. 369 ex parte novi Fisch. Mey. nec Philippi. — *E. exigua* var. 3 Stev. Taur. Verz. p. 300.

Hab. in Tauriâ ad Sudak (Pallas! Stev!), prope Elisabethpol Transcaucasia (Kol!).

Planta 4-8-pollicaris, folia floralia saepe pollicaria 1-1 $\frac{1}{2}$ lin. lata, capsula eâ *E. exiguae* triplo major.

74. E. Szovitsii (F. et M. Index Petrop. I, p. 27) glabra glaucescens pumila, caule brevissimo in umbellam 3-5-radiatam dichotome et divaricatim ramosissimam abeundi, foliis minutis crassiœculis integris caulinis linearis-cuneatis, dichotomiarum linearis-oblongis obtusis saepe falcatim incurvis, involueri minimi turbinato campanulati lobis oblongis fimbriatis, glandulis purpureis transverse oblongis truncatis saepe brevissime biorantis, capsulae longiusculæ stipitatae

ovalæ trisulcæ coccis carinatis, seminis albidi oblongi quadranguli faciebus concaviusculis transverse et irregulariter paucirugosis, carunculâ globoso-depressâ sulculosâ ⊖. Boiss. Euph. Ic. tab. 96.

Hab. in lapidosis montanis et in cultis regionis præsertim montanae, Anatolia interior fere tota in Cariâ (Boiss!), Lyciâ (Bourg!), Cappadociâ (Bal!), Cataoniâ in monte Berytlagh 8000' (Haussk!), Galatiâ (Wied!), Ciliciâ (Ky. 3581), Armenia Turcicâ (Huet!) et Rossicâ (Szov.), Syria circa Aleppo (Ky! Haussk!), ad Balbeck (Post!), ditio Daghestan Caucasi (Becker!), ditio Talysh (Hoh!), Persia bor. (Szov! Ky!), orientalis (Bge!), australis ad Schiraz (Ky!).

Planta 2-3-pollicaris umbellâ corymbose ramosissimâ tota con-tans, folia caulina 3-4 lineas longa, floralia longiora, capsula vix lineam longa. A varis auctoribus *Galarrheo* adnumerata sed glandulæ margine semper plus minus truncatae.

75. **E. punctata** (Delile Fl. Eg. p. 289, tab. 30, fig. 3) glabra pallide virens nana a collo saepius in 2-3 ramos divisa, umbellæ radiis saepius ternis brevibus dichotomis, foliis minimis subsessilibus obovatis transverse latioribus vel retusis obtusis, floralibus submajoribus deltoideis obtusis mucronulatis, involuci minimi turbinati membranacei lobis fimbriatulis, glandulæ cerinæ cornubus albidis brevibus, capsulae ovalæ profunde trisulcæ coccis dorso secus lineam depresso obsolete tuberculato-bieristatis, semine ovato minimo latere interiore foveis binis longitudinalibus oblongis exteriori foveolis irregularibus notato, carunculâ nullâ ⊖.

Hab. in arvis siccis hordeo consitis Egypti ad Alexandriam (Del.), aī stationem Bulkeley prope Randeh (Letourneur!). Fl. Mart.

Planta vix pollicem excedens, folia caulina 4-1½ lineas longa. Facies *E. peploides* sed semina irregulariter nec seriatim foveolato-punctata.

75 bis. **E. arvalis** (Boiss. et Heldr. Diagn. Ser. I, XII, p. 416) glabra glaucescens, canle pumilo a basi ramuloso subcorymboso, ramis umbellæque radiis ternis dichotome ramosissimis, foliis sessilibus crassiusculis integris a basi attenuatâ obovato-oblongis obtusis, floralibus a basi subcordatâ vel breviter cuneatâ ovalis acutis, involuci minimi turbinati membranacei lobis oblongæ cornubus ejus latitudini æquialongis interdum obsoletis, capsule ovalæ depressæ profunde trisulcæ coccis dorso rotundatis, semine ovato albido obsolete quadrangulo irregulariter reticulato-rugoso vel ob rugas magis confluentes foveolato-punctato, carunculâ membranacea conicâ postice sulcatâ ⊖. *E. parvula* C. Koch Linn. XXI, p. 731 non Del. — *E. ruderalis* Scheele Linn. XVII, p. 343!

Hab. in cultis Anatoliae interioris in Pisidiâ (Heldr!), Phrygia (Bal!), Cappadociâ (Bal!), Armeniâ (Calv!), Ponto Lazico ad Djimil 600' (Bal!), ditione Talysh (Hoh!), Persiâ boreali (Buhse!).

Planta 3-5-pollicaris, involucelli phylla 4-6 lineas longa. Facies *E. Szovitsii* sed folia diversa, capsula major non carinata, semina aliter sculpta. A me prius cum *E. punctata* confusa plantâ multo minore et cujus semina multo minora carunculâ destituta foveolis irregularibus sed non rugulosis insculpta sunt.

76. E. Græca (Boiss. et Sprun. Diagn. Or Ser. I, 5, p. 53) anuua glabra glaucescens a collo simplex vel pluricaulis, caulinis pumilis erectis vel aseendentibus umbellæ radiis 3-5 bis vel ter bifidis, foliis inferioribus parvis breviter petiolatis cuneato-obcordatis cæteris oblongo-linearibus linearibusque obtusis vel retusis, floralibus a basi cordatâ triangulari-deltoides acutis, involueri campanulati glabrinseculi lobis ovatis fimbriatulis, glandulae cornubus setaceis divergentibus ejus latitudine duplo longioribus, stylis ultra medium bifidis, capsulâ breviter ovatâ latae trisulcâ minute granulatâ, semine albido ovato irregulariter et reticulatum foveolato, carunculâ conicâ postice sulcatâ ♂. *E. Dalmatica* Vis. Fl. Dalm. III, p. 228. — *E. Terracina* Griseb. Fl. Rüm. I, p. 139 non L.

Hab. in lapidosis regionis inferioris et montanae Graeciae in Atticâ, Argolidâ, Bœtiâ, Eubœâ (Sprt Boiss! Heldr. Fl. exs. 489!), Thraciâ (Griseb!), Anatoliâ in Lydiâ (Boiss! Bal. 347!) et Ciliciâ (Ky!), Tauriâ meridionali (Godet!).

Proxima *E. Taurinensi* differt staturâ tantum semipedali, stylis profunde nec breviter bifidis, glandulis longissime nec breviter et obsolete bicornutis.

Ar. Geogr. Dalmatia.

77. E. Reuteriana (Boiss. Diagn. Ser. I, 12, p. 445) glabra glaucescens, caule erecto subsimplici, umbellæ radiis 3-5 iteratim bifidis, foliis inferioribus minutis obocordatis cæteris linearibus obtusis vel retusis cum mucronulo, summis acutis, floralibus a basi breviter cuneatâ reniformibus transverse latioribus obtusissimis mucronulatis, involueri campanulati lobis ovatis fimbriatulis, glandulae cornubus ejus latitudine longioribus, stylis bifidis, capsule ovatae profunde trisulcâ leviusculâ coccis dorso retundatis, semine profundis foveolis excavato, carunculâ breviter conicâ ♂. Boiss. Euph. Ic. tab. 97.

Hab. in cultis Syriae littoralis, Libani et Antilibani (Bl Gaill!), Palestine (Boiss!).

Valde affinis *E. Græca* et forsitan ejus var. *macrocarpa*; planta procerior sæpe pedalis, folia floralia obtusa, capsula leviuscula 2 lineas longa et lata, semen duplo majus profundius foveolatum.

78. E. segetalis (L. Sp. p. 657) glabra glaucescens, caule erecto crebre folioso sub umbellâ ramulos floriferos saepius edente, umbelle radiis quinis iterato-bifidis tandem ramosissimis, foliis caulinis linearibus acutis vel acuminate laevi deflexis superioribus a basi dilatata lanceolatis, floralibus reniformi-triangularibus transverse latioribus mucronulatis, involueri campanulati intus hirti lobis ovatis fimbriatis, glandulae cornubus setaceis ejus latitudini aequilongis, capsule ovatae profunde trisulcâ coccis minute granulosis dorso subsulcatis, semine albido ovalo dorso subcarinato irregulariter et reticulatum foveolato, carunculâ conicâ postice sulcatâ ♂. Ic. Jacq. Austr. tab. 450. — Fl. Graec. tab. 467. — Rehb. Germ. fig. 4780. — *E. simplex* G. Koch Linn. XXI, p. 730!

Hab. in Cretâ (Sibth!).

Ar. Geogr. Lusitania, Hispania, Gallia australis, Italia, Germania australis, regio Danubialis, Africa borealis, Madera et insulae Canarienses.

2. Perennes.

79. **E. pinea** (L. Syst. p. 376) glabra glaucescens, caulis erectis parte inferiori induratis crebre foliosis sub umbellâ sœpe ramulosis, umbellæ radiis 5-7 semel vel bis bifidis, foliis crassiusculis linearibus obtusiusculis mucronulatis inferioribus fere imbricatis, umbellaribus ovatis floralibus reniformi-cordatis obtusis, involueri campanulati intus hirti lobis ovatis fimbriatis, glandulæ cornubus setaceis ejus latitudine longioribus, capsulâ ovatâ trisulcâ minute granulosâ, semine albido ovato dorso subcarinato irregulariter reticulato-foveolato ♀. *E. cespitosa* Ten. Fl. Nap tab. 143. — *E. Ragusana* Rehb. Ic. Germ. fig. 47-2 b. — *E. congesta* Willd. Enum. Suppl. p. 28! — *E. calcarea* Coss. et Dur.

Hab. in rupestribus sœpius maritimis, Bithynia (Wedem. in herb. Petrop!), probabiliter in Graciâ.

Affinis *E. Portlandiae* sed folia angustiora crebrius imbricata, caules elatiores; *E. segetali* sed perennis.

Ar. Geogr. Regio mediterranea Europæ a Lusitania ad Dalmatiam, Africa borealis, Madera.

80. **E Pestalozzæ** (Boiss. Diagn. Ser. I, 12, p. 114) glabra ex rhizomate indurato cespitoso-multicaulis, caulis nanis simplicibus crebre foliosis, umbellâ ad capitulum 3-5-cephalum reductâ, foliis minutis glaucescentibus crassiusculis integris ab inferioribus dense imbricatis elliptico-linearibus margine subrevolutis ad superiora laxiuscula obverse oblonga paululum auctis omnibus obtusis, floralibus semiobtusis, involueri majusculi campanulati lobis ovatis ciliolatis, floralibus semiobtusis, involueri majusculi campanulati lobis ovatis ciliolatis, glandulis aurantiacis semilunatis cornubus brevibus ovatis, ovario glabro, capsulâ . . . ♀.

Hab. in regione alpinâ montis Akdagl Lyciae (Pest! Bourg!).

Caules 3-4-pollicares, folia 2-4 lineas longa 1½-2 lata. Planta umbellâ ad capitulum nuce avellanâ minus reductâ notabilis. Semen ignotum et ideo locus inter species foveolatas et leiospermas subdubius.

81. **E. erythrodon** (Boiss. et Heldr. Diagn. Ser. I, 12, p. 114) tota sub lente minutissime pruinoso-hirtella glaucescens e rhizomate indurato cespitoso-multicaulis, caulis nanis simplicibus crebre foliosis, umbellâ parvâ 6-12-cephalâ dense capituliformi, foliis minutis imbricatis integris oblongo-ellipticis acutis, floralibus a basi cuneatâ ovatis mucronatis, involueri glabri lobis ovatis truncatis denticulatis, glandulis purpureis transverse oblongis truncatis obsolete cornutis, ovario velutino, capsulâ . . . ♀.

Hab. in rupestribus montis Davrosdagh Pisidiæ 5500' (Heldr!).

Caules 2-3-pollicares, habitus et folia praecedentis sed acuta, capitula majora; ab ea insuper indumento pruinoso et ovario velutino distincta. Locus quoque ob semina non nota subdubius.

82. **E. obovata** (Decaisn. Flor. Sinaic. in Ann. Sc. Nat. 1834, p. 241) glabra glauca, caulinibus ex rhizomate verticali lignoso punulis decumbentibus vel ascendentibus superne ramulosis ramulis umbellaeque radiis 3-5-nis iteratim bifidis, foliis parvis crassis brevissime petiolatis cuneato-obovatis obtusis, floralibus orbiculatis, involucri breviter campanulati lobis ovalis truncatis ciliatis, glandulæ transverse oblongæ cornubus divergentibus ejus latitudine longioribus, capsula ovato-oblonga subtrisulcata, semine oblongo tetragono sparsim malleato, carunculata ovata φ .

Hab. in arenosis deserti Sinaitici (Bové!), in valle sex leucas ad orientem montis Sinai sita (Schimp. 325!), ad Ouadi Esle (Ehrenb!).

Semipedalis, folia majora 3-4 lineas longa, capsula semine Cannabeos sub-major. Ab affinibus foliis petiolatis muticis distincta.

83. **E. Teheranica** (Boiss. in D.C. Prodr. XV, p. 153) glauca sub lente patule et breviter hirtula, caulinibus ex rhizomate lignoso ascendentibus vel decumbentibus tenuibus a basi stricte ramosis, ramis umbellaeque radiis 4-5 pluries dichotomis elongatis, foliis parvis sessilibus ovatis obtusis mucronulatis margine cartilagineo-denticulatis, floralibus orbiculatis obtusis mucronulatis, involucri hemispherico-turbinati lobis ovatis truncatis denticulatis, glandulis transverse oblongis truncatis brevissime bicornutis, capsula ovato-oblonga trisulcata, semine albido oblongo crebre foveolato, carunculata conicata sulcatâ φ . Boiss. Euph. Ic. tab. 103.

Hab. in lapidosis prope Teheran Persiae (Auch. 5312! Ky. 65! Bunge!).

Caules saepè pedales, radii stricti pollicem et amplius longi, folia 2-3 lineas tantum longa, capsula 2 lineas longa. Facies *E. obovatae* a qua foliis sessilibus denticulatis mucronulatis, etc., discedit et *E. erythradeniae* foliis integris, glandularum structurâ, etc.; alienæ.

84. **E. erythradenia** (Boiss. Diagn. Ser. I, 7, p. 92) glauca, caulinibus ascendentibus a basi scoparie ramosissimis paucifoliis ramisque tenuiter virgatis, umbellae radiis binis rarius ternis breviter bifidis, foliis parvis carnosulis integris sessilibus oblongo-ellipticis vel obovatis obtusis, floralibus minimis oblongis, involucri breviter campanulati intus hirti lobis late ovatis truncatis ciliatis, glandulis subquadratis pectinatum et breviter bi vel saepius quadricornutis, capsula ovato-oblonga trisulcata, semine albido ovato-tetragono crebre et minute foveolato, carunculata parva globosa φ . *E. heteradena* Boiss. et Buhse Verz. non Jaub. et Sp.

Hab. in Persie australis monte Kuh Ajub prope Persepolin (Ky. 390!), in montibus inter Jezd et Ispahan (Buhse Bge!).

Caules pedales stricte dichotomi, folia 3-4 lineas longa, capsula 2 lineas longa. Facie *E. megalanthæ* var. *gracili* similis sed characteribus diversissima.

85. *E. petrophila* (C.A. Mey. Beitr. p. 9) glabra glaucescens, caulis e rhizomate indurato pluribus pumilis erectis et ascendentibus supra basin denum denudatam cicatricosam crebre foliosis, foliis parvis crassiusculis sessilibus integris linearispathulatis vel oblongis obtusis, umbellæ radiis 3-5 brevibus simplicibus vel bifidis, foliis floralibus a basi subcordatâ suborbiculatis obtusis vel retusis mucronulatis, involuci campanulati intus hirti lobis ovatis 2-3-fidis, glandulis purpurascensibus truncatis cornubus brevibus, capsula ovato-oblongâ subtrisulcâ sub lente granulatâ, semine ovato-tetragono subcompresso albido lineolis brevibus vel punctis depresso obsito, carunculâ conicâ ♀. Boiss. Euph. Ic. tab. 101. — *E. sacatilis* M.B. Taur. Cauc. III, p. 919 non Jacq. — *E. Armeniaca* Boiss. MSS. — *E. Nicæensis* ♂ minor Ledeb. Fl. Ross III, p. 573.

Hab. in saxosis Tauriae circa Sudak, Sympheropol, Theodosiam (Stev!), in monte Tchatyrdagh (Rehm!), in monte Matschuka Ciscaucasiae (Hohen!), in Anatoliâ boreali ad Saframbol et Kastamouni (Wied!), Armeniae rupestibus ad Baibut (Bourg!).

Semipedalis, folia 6 lineas longa raro in ramis sterilibus fere pollicaria, floralia 4 lineas longa et lata, capsula sesquilinea longa.

86. *E. Chesneyi* (Kl. et Geke Tric. p. 99 sub *Tithymalo*) glabra glaucescens ex rhizomate indurato pluricaulis, caulis erectis vel ascendentibus pumilis simplicibus, umbellæ radiis 4-5 breviter bifidis, foliis parvis sessilibus elliptico-lanceolatis integris inferioribus obtusis superioribus acutis, floralibus ovato-orbiculatis mucronulatis, involuci campanulati lobis triangulari-ovatis ciliatis, glandulis transverse ovalis cornubus brevibus ovalis vel retusis, capsula oblongâ, semine albido ovato obsolete tetragono parce depresso-punctato. carunculâ conicâ ♀. *E. Chesneyi* Boiss. in D.C. Prodr. XV, p. 452.

Hab. in Mesopotamia ad Euphratem (Chesney 457!), ad Diarbekir (Ky 232!), in deserto fluvii Chabur (Haussk!), ad Snaa Farfel deserti ad Palmyra (Bl!) et in argillosis montis Schergi Antilibau (Ky. 532!).

Planta 6-9-pollicaris, facies *E. petrophila*, semina punctis depresso multo rarioribus obsita.

87. *E. malleata* (Boiss. in D.C. Prodr. XV, p. 452) glabra flavido glaucescens, caulis striatis elatis virgatis sub umbellâ stricte et breviter ramulosis, ramulis umbellâque radiis 7-10 semel vel bis bifidis, foliis coriaceis sessilibus basi attenuatis anguste lanceolatis acutis integris, superioribus brevibus ovato-oblongis, floralibus minutis cordatis ovatis mucronatis, involuci campanulati glabri fauce hirti lobis oblongis acutiusculis velutinis, glandulâ semilunatâ inter cornua longiuscula bifida margine crenulatâ, capsula ovata vix trisulcâ subindehiscente, semine ovato parce et parum profunde malato-depresso, carunculâ depressâ ♀.

Hab. in Persia australi ad Ischredabad inter Jezd et Ispahan (Bge!), prope Ssof et Kohrud inter Ispahan et Kaschan (Haussk!).

Facies *E. Gerardiana*, sesquipedalis, folia 1-1½-pollicaria 2-3 lineas lata,

umbellis multiradiatis et capsula medioeri subindehiscenti vix sulcata inter artines insignis. Umbellæ multiradiatae *E. decipientis* sed caules elati et semina diversa.

88. ***E. decipiens*** (Boiss. et Buhse Aufz. p. 197) glauca sub lente minute pruinosa e rhizomate suffrutescenti multicaulis, caulis ascendentibus pumilis crassiusculis duris simplicibus, umbellæ radiis 10-15 brevibus æqualibus bifidis, foliis crenatis subcordiaceis integris a basi sessili attenuatâ linearis-lanceolatis acutiusculis, umbellariis obovato-oblongis obtusis floralibus parvis obovatis, involuci breviter campanulati intus hirti lobis ovatis retusis, glandulis transverse ovalis truncalis brevissime bicornutis, capsula ovalo-oblonga, semine albido oblongo foveolis nigris in sulcos longitudinales irregulariter plus minus coalitis exarato, carunculâ conicâ ♀.

Hab. in Persia inter Kaschan et Ispahan (Bul. se!), in monte Sofi prope Ispahan (Hort. Petrop!), inter Kohrud et Ssof (Bge!).

E. major. — Folia majora pollicaria 3 lineas lata. — *E. polycaula* Boiss. Diagn. Ser. I, 12, p. 112.

Hab. in jugo Elbrus occidentali prope Asadbar (Ky. 453!).

Facies *E. saxatilis* vel *E. Characias* diminutæ, caules 4-6-pollicares, folia in formâ typicâ 6-8 lineas longa 2 lata, capsula 2 lineas longa. Varietas β ob fructus ignotos subtubilia specifica nunc non sat differre videtur.

89. ***E. cheiradenia*** (Boiss. et Holien. Diagn. Or. Ser. I, 12, p. 112) glabra glaucescens ex rhizomate suffrutescenti pluricaulis, caulis erectis simplicibus, umbellæ radiis 5-6 longis breviter bifidis, foliis crassiusculis sessilibus basi attenuatis integris acutiusculis ab infimis elliptico-linearibus vel obovatis ad superiora sensim auctis, floralibus a basi cordatâ transverse ovatis obtusis, involuci campanulati intus hirti lobis ovatis bifidis ciliatis, glandulis semilunatis inter cornua brevia lata obtusa saepius biloba corniculis accessoriis dentiformibus auctis, capsula ovato-oblonga, semine albido vel flavido leproso profunde et crebre foveolis saepius in sulcos transverse anastomosantes coalitis serobiculato, carunculâ conicâ ♀. *E. schizadenia* Boiss. MSS. olim.

Hab. in monte Elbrus Persiae borealis (Auch. 5307! Buhse! Ky. 482!), in aridis ad Sergendeh prope Teheran (Bge!), in monte Elvend (Haussk!), in montibus Sawers et Nur districtus Kuh Kilouyeh et in montibus Avroman et Schiru Kurdistaniae Persiae 10000'-12000' (Haussk. exs. sub *E. bothriosperma*), in monte Kuh Delu (Ky. mixta cum *E. striatella*!).

Facies *E. petrophila* sed omnibus partibus major, pedalis, folia floralia latiora, semen valde serobiculatum.

90. ***E. bothriosperma*** (Boiss. et Ky. Cent. Euph. p. 36 et d.c. Prodr. XV, p. 151) glabra glauca ex rhizomate suffrutescenti multicaulis, caulis induratis ascendentibus, umbellæ radiis quinis tandem elongatis bis vel ter bifidis, foliis coriaceis integris sessilibus basi attenuatis, ramorum steriliarum linearibus plicatis acutis subre-

curvis, caulinis lanceolato-spathulatis breviter mucronatis, floralibus latis a basi subcordatâ semiorbiculatis obtusis, involueri glabri lobis breviter et late ovatis argute denticulatis apice truncatis, glandulæ semilunatæ cornubus obtusis ejus latitudini æquilongis, capsulâ oblongâ apice attenuatâ, semine ovato-tetragono subcompresso crebre et profunde reticulato-foveolato, carunculâ conicâ ♀.

Hab. in Armeniae Turcicae prov. Musch ad Gumgun in campis fertilibus 4600' (Ky. 305!).

Facie et seminibus serobiculatis valde affinis *E. cheiradeniae*, folia longiora acutiora, glandulæ non pectinatæ. Observandum an formis intermediis in eam transit.

91. *E. striatella* (Boiss. in D.C. Prodr. XV, p. 152) glabra glaucescenti-lutescens ex rhizomate indurato multicaulis, caulinis erectis rigidulis tenuiter virgatis crebre striatis superne sub umbellâ breviter et stricte ramulosis, ramulis radiisque umbellæ subquinis brevibus semel vel bis breviter bifidis, foliis remotiusculis parvis sessilibus oblongo-ellipticis obtusis, floralibus acutiusculis ovato-orbiculatis, involueri hemispherico-turbinali intus velutini lobis ovato-triangularibus ciliatis truncatis vel retusis, glandulis triangulari-semiorbiculatis truncatis cornubus brevissimis saepè obsoletis, capsulâ oblongâ apice attenuatâ, semine oblongo parce et superficialiter punctato-depresso, carunculâ conico-pyramidalâ ♀. Boiss. Euph. Ic. tab. 102.

— *E. striata* Boiss. Diagn. Ser. I, 7, p. 91 non Thunb.

Hab. in regione alpinâ montium Kuh Delu et Kuh Barfi Persiae australis (Ky. 346! et 520!), in montibus Kellal et Sselbekh ditionis Bachtiaris (Haussk!), prope Ispahan et in montibus prope Schahrud et Rudbar Persiae borealis (Bge!).

B. microsciadìa Boiss. in D.C. Prodr. — Umbellæ radii pauciores abbreviati, folia floralia acutiora. — *E. microsciadìa* et *E. Persepolitana* Boiss. Diagn. Ser. I, 7, p. 89 et 92.

Hab. in montibus Kuh Delu et Kuh Daëna (Ky. 487! et 625!), in montibus Sawers, Nur et Dalechani supra Sungur 9000'-12000' (Haussk!), circa Persepolin (Ky. 244!).

Semipedalis vel humilior, folia 6-9 lineas longa 3-4 longa. Inter affines caulinis striatellis et glandulis obsolete cornutis insignis.

92. *E. plebeia* (Boiss. Diagn. Ser. I, 7, p. 93) glabra siccitate nigrescens, caulinis ex rhizomate indurato ramoso pluribus crassis tenuibus ascendentibus, umbellæ radiis ternis divaricatis bifidis, foliis minutis margine anguste cartilagineis a basi truncato-subcordatâ ovato-oblongis acutiusculis, floralibus conformibus acutioribus, involueri campanulati lobis ovatis obtusis ciliatis, glandulæ semilunatæ cornubus latitudine glandulæ longioribus, capsulâ majusculâ ovato-oblongâ apice attenuatâ, semine griseo oblongo-tetragono punctis vel lineolis brevibus transversis sparsim et superficialiter obsito, carunculâ conicâ longitudinaliter sulcatâ ♀.

Hab. in glareosis a'pis Kuh Daëna Persiae au'tralis (Ky. 4(9)!).

Caules 2-3-pollicares tenues sed firmi, folia majora 4-5 lin. longa, capsula $2\frac{1}{4}$ lineas longa. A formis *E. striatelle* cui affinis, foliis margine cartilagineis, glandulæ cornubus elongatis specificè differre videtur.

93. E. deflexa (Sibth. et Sm. Fl. Græc. V, p. 52, tab. 466) glabra glaucescens, rhizomate tenui multicauli, caulis decumbentibus tenuibus sub umbellâ saepius ramulosis, umbellæ radiis quinque tandem elongatis iteratim bifidis, foliis præsertim secus caules steriles abbreviatus erebris breviter petiolatis integris minutis obovatis vel elliptico-spathulatis obtusissimis tandem saepe deflexis, floralibus deltoides vel rhombicis transverse latioribus, involueri campanulati intus hirsuti lobis ovatis truncatis ciliatis, glandulae semilunatae cornubus tenuibus ejus latitudine longioribus, capsula ovato-depressa profunde trisulcata secus coecorum dorsum elevatin punctulata, semine albido ovato erebre foveolato, carunculâ minutâ conico-depressâ ȝ.

Hab. in lapidosis regionis montanae et alpine Graecie et Macedoniae, moas Athos (Friv! Gris! Pichler!), Olympus Thessalus (Heldr!), Delphi Euboë (Sprn! Heldr!), Parnassus (Sprn! Heldr! Orph!), mons Pateras Atticae (Heldr!).

Caules herbacei 6-10-pollicares, capsula Cannabeos semine paulo minor. Ab *E. petrophilâ* quâcum cl. Grisebach eam conjungit caulis tenuibus decumbentibus, foliis petiolatis, semine, aliena.

94. E. Kurdica (Boiss. et Haussk.) glabra pallide virens, rhizomate tenui indurato multicanti, caulis ex parte inferiori saepius subterraneâ caudiculosa tenuibus flaccidis procumbentibus vel ascendentibus simplicibus, umbellæ radiis quinque filiformibus longiusculis bis bifidis, foliis parvis sessilibus integris obtusis inferioribus minimis squamaeformibus oblongo-spathulatis, umbellaribus ellipticis floralibus parvis orbiculari-rhombeis mueronatis, involueri breviter campanulati lobis truncatis fimbriatis, glandulae semiorbiculatae cornubus tenuibus parallelis ejus latitudine longioribus, capsula majusculâ ovata obtusa profunde trisulcata longe stipitata, semine albido lineolis longitudinalibus sinuosis parce et irregulariter exarato, carunculâ conico-pyramidalata ȝ.

Hab. ad rupes calcareas montium Avroman, Pir Omar Gudrun et Selahu Kurdistanæ Persie in regione subalpinâ 5000' (Haussk!).

Caules cum caudiculis 6-10-pollicares, folia superiora 3-4 lineas longa, floralia minora, capsula 2 lineas longa. Caulibus elongatis, umbellæ radiis longis, foliis floralibus parvis, semine profundius exarato ab *E. bounophila* differt. Affinis *E. Kanaorica* Indicæ.

95. E. bounophila (Boiss. in D.C. Prodr. XV, p. 154) glabra glaucescens, caulis nanis fragilibus parte subterraneâ caudiculosis nudis, umbellæ brevissime vel oblitteratim triradiatae radiis monocephalis, foliis crassiuseulis parvis subsessilibus basi attenuatis ab infimis minimis obovatis ad superiora oblongo-spathulata auctis, floralibus orbiculato-rhombeis mueronulatis, glandulae semilunatae cornubus convergentibus ejus latitudine sublongioribus, capsula majusculâ

ovatâ obtusâ trisulcâ, somine albido oblongo-cylindrico obsolete et parce longitudinaliter malleato, carunculâ ovatâ ♀. *E. monticola* Boiss. Diagn. Ser. I. 7, p. 93 non Hochst.

Hab. in glareosis regionum altiorum montis Ku'i Daëna Persiae australis (Ky. 630!).

Rhizoma carnosulum crassitie peume anserinæ caudiculos et caules vix bipollulares edens s iperno sat crebre foliosos, folia superiora 5-6 lineas longa, capsula fere 2 lineas longa $1\frac{1}{2}$ lata.

96. ***E. Aucheri*** (Boiss. Diagn. Ser. I. 7, p. 94) sub lente pruinosa glaucescens, caulis nanis fragilibus parte subterraneâ caudiculosis, umbellæ radiis 4-5 brevibus monocephalis, foliis breviter petiolatis inferioribus minimis squameformibus oblongo-linearibus, floralibus majoribus deltoideo-rhombeis obtusis vel retusis margine cartilagineo rabello eroso-denticulato instruetis, glandulae transverse ovate cornubus tenuibus ejus latitudine longioribus, semine oblongo-cylindrico parce et superficialiter malleato, carunculâ breviter conicâ ♀. *E. bonnophila* var. *Aucheri* Boiss. in D. C. Prodr.

Hab. in alpe Elamout Persiae borealis (Auch. 5307!), in monte Totschal prope Teheran (Ky. 558!).

β *stenophylla*. — Folia caulina angustius linearis-spathulata, floralia eis formæ typicæ minora.

Hab. in rupestribus Persiae austro-occidentalis in montibus Sawers et Kuh Nur 4300' (Haussk!).

Species ex florim speciei diverse commixtione in Diagnosibus Or. erronee descripta, a finis *E. bonnophila* quæcum eam conjunxi, sed foliis subpetiolatis margine eroso-cartilagineis diversa videtur. Pollicaris vel bipollicularis, folia floralia in typo 3-4 lineas, in varietate 2 lineas tantum lata.

97. ***E. caudiculosa*** (Boiss. in D. C. Prodr. XV, p. 454) glabra pallide vires, caulis ex rhizomate verticali crassiunculâ fragilibus parte subterraneâ longe caudiculosis exsertâ brevissimis crebre foliosis, umbellâ capituliformi breviter triradiatâ tricephalâ vel monocephalâ, foliis minutis crassiunculis integris ab inferioribus subpetiolatis obovatis ad superiora oblongo-spathulata subauctis, floralibus ovato-rhombeis, involueri intus hirti lobis ovatis ciliatis, glandulae transverse ovatae cornubus ejus latitudini æquilongis, capsula majusculâ ovatâ profunde trisulcâ, semine griseo ovato dorso subcarinato præter 2-3 punctos ad faciem internam depresso levâ, carunculâ depressim postice sulcatâ ♀.

Hab. in summo monte Makmel Libani (Ehrenb! Boiss!), in cacumine Hermonis (Boiss! Ky. exs. 190 sub *E. pumilâ*!).

Rami parte ex-sertâ 1-1 $\frac{1}{2}$ -pollicares, folia majora 3 lineas longa, radii 4-6 lineas longi. Affinis *E. Aucheri* et *bonnophila*, a priori foliis integris, a posteriori semine non longitudinaliter malleato distincta. *E. herniarifolia* proxima quoque habet folia floralia minorâ ovato-elliptica, capsulae coccos dorso subbivalvata, seminis sculpturam diversam. Transitus ad species leiospermacæ mediante *E. Orphanidis* præbet.

98. **E herniariæfolia** (Willd. Sp. II, p. 902) pallide virens, rhizomate verticali tenui, caulis numerosissimis tenuibus parte inferiori saepius subterranea caudiculosis parte exserta prostratis vel ascendentibus brevibus dense foliosis, umbellæ brevissime 3-5-radiatae capituliformi saepe ad involuerum solitarium reductæ, foliis minimis glabris vel puberulis obovatis vel breviter ellipticis obtusis integris inferioribus subpetiolatis cæteris sessilibus floralibus conformibus vix acutis mucronulatis, involueri turbinati lobis ovatis truncatis denticulatis, glandulae transverse oblongæ cornubus tenuibus divergentibus, capsule glabra rarius velutinae ovato-oblongæ trisulcatae coecis dorso angustissime bialatis, semine albido ovato subtetragono facie interna longitudinaliter bisinuato, externa parec foveolato, carunculæ breviter conicæ ♀. *E. pumila* Fl. Græc. V, p. 47, tab. 460. — *E. pycnophylla* G. Koch Linn. XX, p. 17.

Hab. in lapidosis et glareosis regionis alpinæ, Græcia in monte Parnes Atticæ (Pichl!), Parnasso et Kyllene (Heldr!), Malevo Laconie et montibus iugulæ Chios (Orph. Fl. exs. 700!), Crète in montibus Idæ et Lassithi (Heldr! Raul!), Anatolia in Olympo Bithyno (Sibth! Boiss! etc.), Mesogi, Cadmo, Tmolos (Boiss!), montibus Lyciae (Bourg!), Pisidie (Heldr!). Cappadociæ in Argeo (Balt!), Tauro Cilicio (Ky! Balt!), Cataonia in monte Berytlagh (Haussk!) Armenia (Ky!), Ponto ad Djimil (Bal!).

Folia 1 $\frac{1}{2}$ -2 $\frac{1}{2}$ lineas tantum longa, species ab affinis coecis dorso elevatim bilineatis distincta. Non crescit in Libano ubi erroneè indicavi eam cum *E. caudiculosa* confundens.

***** Semina levia. — Species omnes perennies.

+ Folia floralia libera.

99. **E. Terracina** (L. Sp. p. 654) glabra basi ramosa, caulis inferne demum induratis erectis et ascendentibus superne et axillis saepe ramulosis, umbellæ radiis 3-5 tandem ramosissime iteratobifurcatis, foliis firmulis linearibus lanceolatis vel oblongo-linearibus obtusis mucronulatis a medio sursum serrulatis inferioribus et rameorum sterilius saepe truncato-retusis, umbellaribus ovalo-ellipticis, floralibus semiorbicularibus vel ovato-triangularibus, involueri campanulati lobis ovatis fimbriato-laceris, cornubus setaceis glandulae latitudine duplo longioribus, capsule depresso profunde et late trisulcatae coecis dorso subcarinatis, semine albido ovato dorso carinato, carunculæ postice elongatæ rostratæ ♀. *E. Provincialis* Willd. Sp. II, p. 914. — Rehb. Ic. Germ. 4790. — *E. Valentina* Orteg. — *E. seticornis* Poir. Diet. Suppl. II, p. 617 — *E. obtusifolia* Lam. Diet. II, p. 430. — *E. Neapolitana* Ten. Fl. Nap. tab. 42. — *E. heterophylla* Desf. Atl. I, p. 385, tab. 102. — *E. diversifolia* W. Enn.

Hab. ad vias, in cultis regionis inferioris, Græcia in Ætolia (Reinh!), Cycladicibus (Schmidt!), Crète (Baulin!), Syria littoralis (Bl! Gaill!), Palestina (Sieb!).

β . *prostrata* — Caules prostrati vel procumbentes, folia diminuta. — *E. leiosperma* Fl. Græc. V, p. 51, tab. 465. — *E. Portlandica* Sm.

Prodr. I, p. 327 non L. — *E. obliquata* Forsk. Eg. Arab. p. 93. — *E. Alexandrina* Del. Fl. Eg. p. 90, tab. 30. — *E. modesta* Boiss. Cent. Euph. p. 34.

Hab. in arenosis maritimis Græciæ (Sibth.), insulæ Thera (Sart!), ad Hierosolymam (Roth!), Arabiæ petreæ (Boiss!), Egypto ad Alexandriam (Del! Samar! Gaill!).

Species proetiformis, caulis erectis vel prostratis, folia formâ et magnitudine variabilis, floralia plus minus longe mucronata.

Ar. Geogr. Europæ regio mediterranea tota a Lusitaniâ ad Italianam, Africa borealis, insule Canarienses, Azoricæ, Madera.

100. ***E. Gerardiana*** (Jacq. Austr. V, p. 17, tab. 436) glabra glaucescens multicaulinis, rhizomate verticali sublignoso, caulis erectis virgatis duriusculis cærebre foliosis, umbellæ radiis numerosis tenuibus semel vel bis bifidis, foliis strictis linearibus vel lanceolato-linearibus mucronatis integris, umbellaribus brevibus acutis, floralibus a basi truncatâ vel cordatâ triangulari-ovatis mucronato-aristatis flavidis, involueri campanulati fance hirtuli lobis triangulari-ovatis ciliatis, glandulis transverse triangulari-ovatis truncatis ecornutis, capsula breviter ovatâ trisulcâ breviter et obsolete granulosâ, semine ovato, carunculâ depresso-conicâ sessili ♀. Rehb. Ic. Germ. fig. 4794. — *E. Cajogala* Ehrh. Beitr. II, p. 102. — *E. linariaefolia* Lam. Dict. II, p. 162. — *E. glaucescens* Willd. Enum. Suppl. p. 28.

Hab. in campis siccis praesertim montanis, Graecia in monte Malevo Laciniæ (Heldr!), Macedonia et Thracia (Griseb! Orph!), Byzantii (Commany!), Archipelagus (Urv!), Anatolia in Bithyniâ (Wied!) et Isauriâ (Heldr!), Transcaucasiâ (Hoh!), Persiae bor. montanis (Szov! Buhse! Bge!), prope Hamadan (Haussk!).

β. Hohenackeri. — Folia floralia elongata oblongo-lanceolata. — *E. Hohenackeri* Hochst. et St. MSS.

Hab. ad margines agrorum prope Helenendorf Georgiae Caucasicæ (Hohen!).

Species ob glandulas involueri prorsus eornutas et sœpe obscure truncatas *Galarrhæis* adnumerata sed ob faciem *E. Cyparissias* vel *Esulam* referentem potius *Esulis* adnumeranda.

Ar. Geogr. Gallia, Belgium, Helvetia et Germania australes, Hispania, Italia, regio Danubialis, Rossia media et australis, Sibiria Uralensis.

101. ***E. Buhsei*** (Boiss. in D.C. Prodr. XV, p. 167) glabra basi indurata dumulosa, caulis erectis ramosissimis striatis corymbosis, umbellæ radiis 5-7 semel vel bis bifidis, foliis parvis a basi sessili subattenuatâ elliptico-linearibus obtusiusculis integris, floribus rhomboe-orbiculatis obtusis, involueri campanulati intus breviter hirtuli lobis elongatis linear-i-quadratis truncato-retusis ciliolatis, glandulis transverse oblongis truncatis sœpe erosio-denticulatis, capsula ovalâ trisulcâ, semine ovato, carunculâ conicâ depresso-radiatim sulcatâ ♀. Boiss. Euph. Ic. tab. 410.

Hab. in Persiâ orientali ad Rischin (Buhse!) et prope Schahrud (Bge forma filamentis hirtulis!), austro-orientali inter Khabbise et Kerman (Bge!), australi (Ky. 878!).

Basi probabilitate suffruticosa, caules 6-10-pollicares penūt' anserinā tenuiores tota longitudine stricte ramulosi, folia 5-6 lineas longi lineam lata, floralia 3-4 lineas lata, capsula 2½ lineas longa. Ex affinitate *E. Gerardiana*.

102. **E. Cyparissias** (L. Sp. 661) glabra, rhizomate repente apice pluricauli, caulis erectis crebre foliosis inferne ramos breves steriles superne ramulos floriferos edentibus, radiis numerosis tenuibus bis bifidis, foliis late viridibus anguste linearibus integris, rameorum steriliū fere setaceis, floralibus lutescentibus a basi rotundatā vel subcordatā ovato-reniformibus obtusissimis mutatis vel mucronatis, involuci breviter campanulati lobis ovatis ciliatis saepe retusis, glandulis semilunatis brevissime attenuato-bicornutis, capsulā depressā profunde trisulcā punctulato-seabridā, carunculā orbiculari sessili ♀. Ic. Jaeq. Austr. V, t. 435. — Fl. Dan. tab. 2052. — Rehb. Ic. Germ. fig. 4793.

Hab. in pratis regionis montanæ et alpine Macedoniae occidentalis in monte Peristeri 2400'-4000' (Griseb., Orph!).

Ar. Geogr. Europa borealis et media a Suecia australi et Rossia media ad Hispaniam borealem, Italiam omnem in montanis, Dalmatiam.

103. **E. Esula** (L. Sp. 660) glaucescens, rhizomate repente graciili stolonifero, caulis erectis striatis ramulosis, ramulis umbellaque multifidae radiis bis bifidis, foliis a basi attenuatā linearibus lanceolatis vel cylindro-lanceolatis prope apicem saepe seabridis infimis saepe subpetiolatis, rameorum steriliū angustioribus, umbellariis breviter lanceolatis, floralibus rhombo-ovatis vel reniformibus transverse latioribus mucronatis, involuci campanulati lobis latis brevibus truncatis timbriatis, glandulis brevissime bicornutis, capsulā ovalā profunde trisulcā punctato-seabridā, carunculā orbiculari sessili ♀. Ic. Fl. Dan. tab. 1270. — Rehb Ic. Germ. fig. 4791.

Hab. in humidis regionis montanæ, Caucasus (C.A. Mey!), Persie prov. Khorassan inter Nichapur et Mesched (Bge!).

Ar. Geogr. Anglia, Scotia, Svecia meridionalis, Suecia, Gallia borealis et occidentalis, Hispania in montanis, Italia borealis et media, regio Danubialis, Rossia media et australis, Sibiria, China borealis.

104. **E. leptocaula** (Boiss. in D.C. Prodr. XV, p. 159) glabra glaucescens, rhizomate tenui, caulis tenuibus erectis ex axillis superioribus stricte ramulosis, ramulis radiisque 3-5-nis tenuissimis longe bifidis saepe inaequalibus, foliis rigidulis anguste et longe linearibus acutiusculis integris margine subrevolutis prominule uninerviis, rameorum steriliū subsetaceis, umbellaribus longis inaequalibus lanceolato-linearibus, floralibus parvis rhombo-reniformibus mucronatis, involuci breviter campanulati lobis breviter ovatis ciliatis, glandulis breviter bicornutis, capsulā majuscūlā profunde trisulcā elevato-punctatā, semine albido ovato, carunculā parvā subsphaericā ♀. Boiss. Ic. Euph. tab. 105. — *E. tenuifolia* M.B. Taur. Cauc. I, p. 372

non Lam. — *E. bupleuroides* Willd. herb! non Desf. — *E. longifolia* Guld. It. I. p. 192.

Hab. in campestribus Taurie (Stev!), in prov. Guriā Caucasī (Nordm!).

Planta ½-4-pedalis, folia 2-3-pollicaria 1-2 lineas lata, facies *Bupleuri cariciifoliae*; differt a formis angustifoliis *E. Esulae* radiis paucis tenuissimis elongatis, capsula majore, etc.

Ar. Geogr. Rossia australis occidentalis et regio Wolgensis.

105. ***E. virgata*** (W. K. Pl. rar. Hung. II, p. 176, tab. 162) glabra, radice verticali pluricipiti, caulis erectis virgatis induratis tenuiter striatis cibre foliosis superne ex axillis ramulosis, ramulis radiisque 5 tenuibus iterato-bifidis anguste thyrsoides, foliis strictis rigidulis linear-lanceolatis acutis integris basi vix attenuatis, umbellariibus brevibus ovatis, floralibus transverse latioribus mucronulatis, involueri campanulati lobis ovatis denticulatis, cornubus apice dilatato denticulatis glandulæ semilunatae latitudine longioribus, capsula breviter ovata profunde trisulcata elevato-punctata, semine ovato, carunculata orbiculari postice breviter rostrata ♀. Rehb. Ic. Germ. fig. 4792.

Hab. in pratis Macedoniæ (Friv. ex Griseb. Spic.), in prov. Daghestan Caucasī (Becker!).

g. Orientalis Boiss. in p.c. Prodr. I. cit. p. 160. — Ramosior, folia elongata sœpe 2-3½-pollicaria, capsula paulo major.

Hab. in Anatoliâ boreali (Wied!), Armeniâ Turcicâ (Huet!), Transcaucasî (Hoh!), Kurdistaniâ (Ky. 569! sub *E. Esula*), Persiâ boreali (Szovits! Bge!).

g. Uralensis. — Ramosissima, folia longa sœpe angustissima. — *E. Uralensis* Fisch. in Link Enum. II, p. 14.

Hab. in Transcaucasî ad Sallian (herb. Ledeb!), Armeniâ australi (Ky. 303!), Cappadociâ ad Euphratem (Auch. 2220).

Species radice, foliis integris strictis basi vix attenuatis, cornubus elongatis ab *E. Esulae* formis distincta.

Ar. Geogr. Moravia, Bohemia, Illyria, regio Danubialis, Rossia australis Songaria, Sibiria Altaica.

106 ***E. hebecarpa*** (Boiss. Diagn. Or. Per. I, 7, p. 90) canthibus erectis virgatis firmis striatis totâ longitudine stricte ramosis, radiis umbellæ centralis amplæ 7-8-nis lateralium 3-4-nis omnibus bifidis, foliis crassiusculis strictis glabris a basi breviter attenuatae linear-lanceolatis acutis integris, superioribus et umbellaribus dilatatis oblongo-lanceolatis, floralibus ovato-deltoides mucronulatis sœpius puberulis, involueri velutini lobis ovatis ciliatis, cornubus latitudine glandulæ brevioribus, capsula ovata profunde trisulcata velutinæ, semine ovato, carunculata orbiculari ♀.

Hab. in humidis ad radices montis Kuh Daëna Persiæ australis (Ky. 567!), in montibus Sawers, Nur, Dalechani (Haussk!).

Ab affini *E. virgata* var. *Orientali* foliorum floralium involueri et capsulae indumento, hujus coccis laevibus nec minute tuberculatis distincta.

107. *E. salicifolia* (Host Syn. p. 267) rhizomate longe repente, caulis erectis elatis minute puberulis sub umbellâ ramulosis, umbellæ multifidæ radiis brevibus iterato-bifidis, foliis majusculis pallide viridi-lutescentibus a basi breviter attenuatâ lanceolatis acuminatis obtusiusculis integris, umbellaribus ellipticis obtusis, floralibus luteis a basi subcordatâ reniformibus transverse latioribus obtusis, glandulis breviter bicornutis, capsulâ breviter ovalâ profunde trisulcâ elevato-punctatâ, semine ovato, carunculâ orbiculari sessili ♀. W. K. Hung tab. 55. — Rehb. Ic. Germ. fig. 4798. — *E. pallida* Willd. Sp. p. 923 ex parte.

Hab. in Rûmeliâ ad thermas prope Lia (Friv!).

Ar. Geogr. Hungaria et regio Danubialis, Bosnia, Rossia australis.

108. *E. lucida* (W. K. Pl. rar. Hung. p. 54. tab. 54) glabra, rhizomate longe repente, caulis crassis elatis striatis superne parce et breviter ramulosis, umbellæ multifidæ radiis brevibus iterato-bifidis, foliis firmis nitentibus a basi truncatâ vel cordatâ lanceolatis a medio apicem versus attenuato-acuminatis obtusis vel acutiusculis integris margine subrevolutis, floralibus ovato-rhombeis transverse latioribus mucronatis, involueri breviter campanulati intus hirtuli lobis magnis truncato-retusis, cornubus glandulæ latitudine æquialongis, capsulâ ovatâ punctato-scabrà, semine ovato-oblongo plumbeo, carunculâ oblongâ rostratâ ♀. *E. pallida* Willd. Sp. II, p. 923 ex parte.

Hab. in Anatoliâ bor. (Tehiht!).

Variat foliis angustioribus. — *E. lucida* β. *androsaemifolia* C. Koch Linn. XXI, p. 728. — Rehb. Ic. Germ. fig. 4797.

Hab. in Rûmeliâ boreali extra ditionem nostram (Noë!).

Ar. Geogr. Moravia, Silesia, Austria, regio Danubialis, Rossia australis.

109. *E. Iberica* (Boiss. Cent. Euph. p. 28. — D.C. Prodr. XV, p. 163) glabra, glaucescens, rhizomate longe repente, caulis erectis crassiœculis valde angulato-striatis superne rumulosis, umbellæ multifidæ radiis brevibus bifidis, foliis sessilibus firmulis supra lucidis subtus penninerviis margine scabridis et subrevolutis a basi rotundatâ oblongis acutiusculis, ramorum sterilium a basi truncatâ linearibus, umbellaribus breviter ovatis, floralibus a basi cordatâ reniformibus obtusis mucronatis, involueri campanulati lobis ovatis subtruncatis, glandulae cornubus brevibus divergentibus, capsula valde depressâ profunde trisulcâ elevato-punctatâ, semine ovato nigro, carunculâ transverse sublatiore ♀. Boiss. Euph. Ic. tab. 107. — *Tithym. Orientalis latissimo folio glanco et glabro* Tourn. Cor. — *E. salicifolia* β. *latifolia* Ledeb. Fl. Ross. III, p. 378. — *E. latifolia* Hohen. exs. non C.A. Mey.

Hab. in herbidis humidis Ponti Lazici supra Rhizé (Ball), Armeniae (Tourn. herb!), Iberie Caucasicæ (Hoh!), Persiae borealis (Buhse! Bge!).

Pedalis, folia caulina 4-4½-pollicaria 8-9 lin. lata, capsula 1¼ lin. longa 2 lata; differt ab *E. latifoliâ* Sibiricâ foliis firmis forme diversæ et magis accedit ad *E. lucidam* foliis longis longe attenuatis, capsulâ ovatâ, etc. distinctam.

110. *E. agraria* (M.B Taur. Cane. I, p. 375) glabra glaucescens, caulis e rhizomate repente erectis striatis crebre foliatis sub umbellâ breviter ramulosis thyrsoides, umbellæ multifidae radiis tenuibus bis bifidis, foliis rigidiusculis a basi sessili obtuse auriculato-cordatâ ovatis vel oblongis obtusis venulosis margine anguste cartilagineo-seabridis, caulinum sterulum ellipticis, floralibus reniformibus transverse latioribus obtusis mucronulatis, involuci turbinato campanulati intus hirti lobis ovatis ciliatis, glandulis truncatis brevissime bicornutis, capsulâ ovatâ profunde trisulcâ granulato-asperulâ, semine ovato, carunculâ conicâ ♀. Rehb. Ic. Crit. III, tab. 251. — *E. nitens* Trevir. — *E. thyrsiflora* Griseb. Spieil. I, p. 143 — *E. Transylvanica* Schur. — *E. subhastata* Vis. et Panc. Déc. I, p. 22, tab. 7.

Hab. in campis et al margines agrorum Thracie (Griseb!), Bithyniâ ad Seutari (Noe!), in montosis montis Mesogis Carie (Boiss!), Tauriâ (Stev!). Transcaucasiâ prope Elisabethgrad (Boschn!).

Planta 1½-3-pedalis, folia 4-4½-pollicaria, capsula Cannabeos semine vix major; foliis brevibus latis approximatis ad *Myrsinæas* accedit.

Ar. Geogr. Transylvania, Moldavia, Rossia australis.

111. *E. Baselicis* (Ten. Fl. Nap. IV, p. 265, tab. 143) glabra pallida, caulis ascendentibus parum elatis rigidulis, radiis 5-6 breviter bifidis, foliis sessilibus parvis basi attenuatis inferioribus ellipticis superioribus et umbellaribus ovatis basi truncato-cordatis omnibus mucronatis marginatis seabrido-serrulatis, floralibus semi-orbiculatis obtusissimis, involuci breviter campanulati intus hirtuli lobis ovalis ciliatis apice sœpe fisis, cornubus glandulæ latitudini æquilongis, capsulâ ovatâ minute granulatâ. semine ovato dorso carinato, carunculâ depressâ conicâ postice sulcatâ ♀. *E. rupestris* Friv. exs.

Hab. in rupestribus schistosis Macedoniæ et Thracie, in jnglo Babuna prope Perlepe (Griseb.), ad radices montis Rhodopes prope Banja (Friv!).

Specimina tantum florifera quæ ab *E. Baselicis* Neapolitanâ foliis evidenter serrulatis, superioribus minus dilatatis differre tantum videntur. Facies *E. saxatilis* foliis inferioribus majusculis basi longe attenuatis diversæ. Semper a formis pumilis *E. Nicavensis* foliis seabrido-serrulatis nec integris distinguuntur.

112. *E. Orphanidis* (Boiss. Diagn. Ser. II, 4, p. 89) glabra glaucescens, rhizomate ramoso caules numerosos parte inferiore subierraneâ longe caudiculosos nudos fragiles procumbentes parte exsertâ breves foliosos edente, ramulis floriferis axillaribus radiisque quinis brevibus bis bifidis, foliis parvis crassinervulatis integris a basi longe attenuatâ obovatis obtusis, umbellaribus breviter ovatis, floralibus a basi subcordatâ semiorbiculari-deltoides obtusis, involuci

campanulati intus dense velutini lobis late et breviter ovatis, cornibus subdivergentibus glandulae latitudine longioribus, capsula majuscule ovata obtuse trigonâ sub lente elevatim punctatâ, semine plumbeo breviter ovato, carunculâ radiatim lobatâ φ . Boiss. Euph. Ic. tab. 109. — *E. Hohenackeri* Orph. MSS. non Hochst.

Hab. in glareosis regionis alpinae Parnassi prope Lugari 5-6000' (Orph. Fl. exs. 407! Guicci!).

Caules parte exsertâ cum umbellâ 2-4-pollicares, folia majora 5-6 lineas longa, capsula pisum æquans; facies *E. deflexæ* sed omnibus partibus major.

113. *E. glareosa* (M.B. Taur. Cauc. I, p. 373) glabra pallide virens multicaulis, caulis erectis crassiusculis inferne tandem denuo induratis sub umbellâ ramulosis corymbosis, ramulis umbellæque multifidæ radiis semel vel bis bifidis, foliis crassiusculis a basi attenuatâ lanceolatis vel oblongo-lanceolatis prope apicem subtiliter crenulatis, umbellaribus ovatis, floralibus reniformi-deltoides mucronatis, involueri campanulati glabriusculi lobis apice truncato-retusis fimbriatis, glandulis ovato-triangularibus truncatis ecornutis, capsula ovata trisulcâ elevatim punctatâ, semine ovato, carunculâ conicâ postice suleatâ φ . *E. Nicæensis* β . *glareosa* Koch Syn. Ed. 2, p. 729.

Hab. in collibus siccis Anatoliæ borealis (Wied! Auch. 796!), Tauriæ (Stev!), Transcaucasiæ (Hohen!).

β . *lasiocarpa*. — Capsula velutina. — *E. Pannonica* Host Austr. II, p. 566. — Rchb. Germ. f. 4795.

Hab. in Rumelia (Noë!).

γ . *minor*. — Omnibus partibus minor.

Hab. in Armeniâ (Calvi!), Tauriâ (Stev!), Georgia Caucasicâ (Hoh. sub *E. saxatili*).

Species valde affinis *E. Nicæensi* a quâ formâ glandularum et capsula punctulatâ nec rugulosâ differt; specimina var. γ . sæpe cum *E. petrophilâ* commutantur.

Ar. Geogr. Austria, regio Danubialis, Rossia australis.

114. *E. tinctoria* (Boiss. et Huet Sched. 1853. — D.C. Prodr. XV, p. 106) lutescens sub lente pruinoso-hirtula multicaulis, rhizomate surculos repentes edente, caulis crassis albidis striato-sulcatis ramuloso-thyrsoideis, ramulis radiisque 7-9 bis bifidis, foliis coriacieis integris a basi attenuatâ lanceolatis acutis mucronatis nervis subtus prominulis, umbellaribus ovato-oblongis acutis, floralibus rhomboideo deltoideis longiuscule mucronatis, involueri intus velutini lobis ovatis truncatis, cornibus latitudine glandulae brevioribus, capsula ovata lœvi glabrâ vel tomentellâ obsolete trisulcâ φ . *E. Noëana* Boiss. in Noë exs. — *E. macroclada* Boiss. Diagn. Ser. I, p. 54 (forma foliis angustis). — *E. Syspirensis* C. Koch Linn. XXI, p. 725.

Hab. in siccis Anatoliæ borealis (Wied!), Cariæ (Boiss!), Lyciæ (Bourg!), Armeniæ (Huet!), Kurdistaniæ Persicæ (Haussk!), Persiæ bor. et occid. (Szov! Noë!).

β. schizoceras Boiss. in D. C. Prodr. — Glandularum cornua palmatim 2-3-loba et glandula ipsa saepe toto margine multifida. — *E. schizoceras* Boiss. Diagn. Ser. I, 5, p. 55. — *E. Damascena* Boiss. Diagn. Ser. I, 12, p. 112. — *E. tetraceras* Szov. exs. — *E. Lorentii* Hochst. in Lor. Wand. p. 344.

Hab. in Anatoliæ monte Aladagh (Wied!), Cappadociâ (Bal!), Syriâ bor. (Haussk!), Damasci (Boiss!), Libano et Antilibano (Bl! Gaill!).

Species ab *E. Nicœensi* affini surculis repentibus, caulinibus albidi striatis, foliis acutis subtus elevato-nervosis, capsulâ lèvissimâ distincta. Varietas β intermediis ad typum transit.

115. ***E. Paralias*** (L. Sp. 657) glabra glauca e radice ramosâ longâ multicaulis, caulinibus erectis duris dense et imbricatim foliatis, umbellæ confertæ radiis 3-5 crassis brevibus bis vel ter bifidis, foliis coriaceis integris inferioribus brevioribus linear-ellipticis obtusis superioribus oblongis acutiusculis, umbellaribus ovatis basi cordatis, floralibus reniformi-cordatis mucronulatis, involuci breviter campanulati intus hirti lobis oblongis ciliatis, glandularum cornubus brevibus divaricatis, capsulâ depresso profunde trisulcâ tenuiter granulata, semine albo ovato-globoso dorso subcarinato, carunculâ minimâ depressâ reniformi φ . Ic. Jacq. Vind. tab. 188. — Rchb. Germ. fig. 4789.

Hab. in arenosis maritimis Græcia! Archipelagi! Cretæ (Sieb!), Macedonia (Friv!), Anatoliæ bor. ad Samsan (Wied!) et ad Smyrnam (Ball), Syriæ littoralis (Gaill!), Egypti (Ehr! Schv!), in Guriâ littorali (Rupr!).

Ar. Geogr. Britannia, Gallia occidentalis et australis, Lusitania et Hispania, Italia, Dalmatia.

+ + Folia floralia connata. — Caules vel rami steriles hornotini hyeme foliati vere insequenti elongati et floriferi.

116. ***E. amygdaloïdes*** (L. Sp. p. 662) caulinibus floriferis teretibus erectis basi denudatâ induratis sub umbellâ ramulosis thyrsoides, radiis 5-19 bis bifidis, foliis puberulis vel glabratris inferioribus anni præcedentis majoribus dense congestis firmis obovato-oblongis in petiolum attenuatis, superioribus hornotinis remotis sessilibus oblongis, umbellaribus ovatis, floralibus semiobtusibulatis in patellam concaviusculam ultra medium coalitis, involuci campanulati intus hirtuli lobis ovatis dentatis, glandularum cornubus longis convergentibus acutis, capsulâ depresso profunde trisulcâ, semine nigro ovato, carunculâ minutâ subcarinatâ postice subrostratâ φ . Ic. Rchb. Germ. fig. 4799. — *E. sylvatica* Jacq. Austr. tab. 375. — *E. oblongifolia* C. Koch Linn. XXI, p. 726.

Hab. in sylvaticis Macedoniæ (Griseb!), Bithyniæ (Pest!), Anatoliæ bor. (Tchihi!), Phrygiæ (Bal!), Cappadociæ (Kyl!), Ponti Lazici (Bal!), Tauriæ et prov. Caucasicarum (Hoh!), Persiæ borealis in provinciis ad Caspium (Buhse! Begl.).

Ar. Geogr. Europa media et australis ab Angliâ, Belgio, Germaniâ, ad Ros-siam meridionalem, Turcomaniam.

117. **E. iteophylla** (Boiss. Cent. Euph. p. 39. — n.c. Prodr. XV, p. 171) glabra, caulis teretibus erectis crebre foliosis inferne tandem denudatis et cicatricosis, umbellæ radiis 5-6 brevibus bis bifidis, foliis sessilibus membranaceis a basi attenuatâ lanceolatis acutis integris ab inferioribus elongatis ad superiora oblonga sensim abbreviatis, floralibus in patellam parvam planiusculam orbicularem fere omnino connatis, glandularum cornubus longis convergentibus acutis, capsulâ depressâ elevatim punctulatâ, semine ovato ♀. Boiss. Euph. Ic. tab. 113. — *E. glaberrima* C. Koch Linn. XXI, p. 726 non Kotsch.

Hab. in Iberia Caucasicæ valle fluvii Aragwi (Hohen!), in Grusiâ (C. Koch!).

Affinis *E. amygdaloidi* differt caule etiam superne dense folioso, foliis formæ aliena, floralibus dimidio minoribus. Folia inferiora 3-4½-pollices longa 7-8 lineas lata.

118. **E. macroceras** (F. et M. Ind. IV, Petrop. p. 36) tota sparsim et patule hirtula, caulis herbaceis proceris sulcate-striatis sub umbellâ crebre ramulosis, ramulis radiisque quinis longis longe et iteratim bifidis, foliis amplis tenuiter membranaceis a basi petiolatâ cuneatâ ovato-oblongis acutis integris superioribus sensim diminutis, umbellaribus ovato-oblongis, floralibus in patellam orbicularem concaviusculam omnino connatis, involueri campanulati fauce velutini lobis dense ciliatis breviter ovatis truncato-retusis, cornubus convergenti-recurvis sensim attenuatis glandulæ latitudine longioribus, capsulâ depressâ elevatim punctatâ profunde trisulcâ, semine ovato-sphaerico, carunculâ sessili depressâ ♀. Boiss. Euph. Ic. t. 114.

Hab. in umbrosis subalpinis Iberia Caucasicæ (Hoh!), Cartaliniæ (Szov!), Guriæ (Nordm!), Caucasi orientalis (Rupr!).

Facies *E. asperæ* a quâ caracteribus longe distat, caules 2-3-pedales, radii 4-8-pollices longi, folia majora 3-4-pollicaria, capsula 2 lineas lata.

119. **E. rumicifolia** (Boiss. Cent. Euph. p. 39. — n.c. Prodr. XV, p. 171) pilosula vel glabrata, caulis erectis herbaceis striato-sulcatis crebre foliosis superne breviter ramulosis, radiis 7-9 brevibus tenuibus bis bifidis, foliis membranaceis sessilibus vel breviter petiolatis a basi obtuse cordato-biauriculatâ ovatis vel ovato-oblongis obtusis, superioribus et umbellaribus basi truncatis acutiusculis, floralibus in patellam orbicularem concaviusculam fere omnino coaliatis, involueri intus velutini lobis truncatis, glandulæ transverse latioris cornubus longis convergentibus, capsulâ elevato-punctatâ, semine ovalo, carunculâ planâ ♀. Boiss. Euph. Ic. tab. 115. — *E. cordifolia* C.A. Mey. MSS. in pl. Szov non alior.

Hab. in Guriâ (Szov!), Armeniâ Turcicâ (Huet!), Armeniâ australi ad Müküs (Ky!), Ponto Lazico in valle Djimil (Bal!), in sylvis supra Trapezuntem (Bourg!).

Differt ab *E. macrocerate* foliis basi auriculato-cordatis, radiis brevibus pollicaribus, foliis floralibus minoribus, glandulis transverse dilatatis.

120. E. Kotschyana (Fenzl Pugill. № 17. — Illustr. pl. Nov. p. 20) basi indurata, caulis crassis elatis glaucescentibus inferne tandem nudis suffruticosis sub umbellâ ramulosis thyrsoideis, ramulis radiisque 6-10 brevibus iterato-bifidis, foliis coriaceis integris glabris vel tomentellis supra lucidis subtus glaucescentibus breviter petiolatis basi attenuatis lanceolatis obtusiusculis mucronatis, superioribus et umbellaribus abbreviatis, floralibus in patellam orbicularē amplam planiuseculam ultra medium connatis, involueri extus glabri intus pilosi lobis late ovatis truncatis ciliatis cornubus latitudine glandulæ brevioribus apice interdum incrassatis, capsulâ glabrâ vel hirtulâ ovatâ basi et apice truncatâ profunde trisulcâ, semine cœsio ovato, carunculâ conicâ substipitatâ ♀. Boiss. Euph. Ic. tab. 116.

Hab. in alpinis Anatoliæ meridionalis in umbrosis, montes Pisidiæ (Heldr!), mons Akdag Lyciaæ (Bourg!), Taurus Cilicus (Ky! Bal!).

Facies *E. amygdaloïdis* a quâ differt caulis basi suffrutescentibus, foliis coriaceis lucidis, stylis longis nec brevibus, capsulâ duplo majore.

121. E. macrostegia (Boiss. in D.C. Prodr. XV, p. 171) basi suffrutescens, caulis elatis rubellis inferne tandem denudatis sub umbellâ ramulosis corymbosis, ramulis radiisque quinis elongatis iteratim bifidis, foliis coriaceis supra nitidis subtus pallidioribus glabris vel pilosulis subpetiolatis basi attenuatis ovato-oblongis obtusis, umbellaribus ovatis, floralibus in patellam amplam orbiculatam retusam connatis, involueri tomentelli lobis ovatis valde ciliatis, glandularum cornubus brevibus divergentibus, capsulâ ovatâ puberulâ vel glabrâ, semine plumbeo ovato, carunculâ conico-depressâ postice sulcatâ ♀. *E. erubescens* Boiss. Diagn. Ser. I, 7, p. 90 non *E. Mey.*

Hab. in umbrosis fauoris Guzelderé Ciliciæ supra Sedichig (Bal!), Libano circa Eden (Bl! Ky. 682!), Persiæ austro-occidentali in monte Kuh Delu (Ky. 533!), montibus Sawers, Eschker 7-9000' (Haussk!).

Valde affinis *E. Kotschyana* specifice differre videtur foliis abbreviatis, umbellæ radiis elongatis sœpe semipedalibus, foliis floralibus amplioribus, capsulâ seminibusque dimidio minoribus. An tamen hujus varietas?

122. E. Characias (L. Sp. 662) tota tomentella, caulis crassis inferne induratis et demum nudis cicatricosis superne crebre foliosis thyrsoideis, ramulis umbellæque multifidæ radiis brevibus bifidis, foliis firmis a basi longe attenuatâ oblongo-lanceolatis acutis superioribus sensim abbreviatis, umbellaribus ovato-oblongis mucronatis, floralibus virentibus ad medium usque in patellam oblongam concaviusculam connatis, involueri campanulati intus et extus hirsuti lobis oblongis bifidis, glandulis atropurpureis transverse latioribus truncatis sœpe crenulatis angulis in cornua brevissima interdum abeuntibus, capsulâ tomentosâ depressâ trisulcâ, semine ovato, carunculâ conico-depressâ postice sulcatâ ♀. Jacq. Ic. Rar. tab. 89. — Rchb. Germ. fig. 4800. — *E. eriocarpa* Bertol. Comm. p. 22, tab. 3. — *E. Cretica* Mill. Dict. I, № 28.

Hab. in regione inferiori insulæ Cretæ (Sieb!), in Eparchiâ Temenos ad radices Dycæi 1200'-1500' (Heldr!).

Species affinis forsitan in Graecia occidentali reperienda est *E. Wulfeni* Hoppe, umbellâ ampliore, patellâ profunde infundibuliformi, glandulis longe bicornutis distincta.

Ar. Geogr. Regio mediterranea Lusitaniæ, Hispaniæ, Galliæ, Italiæ.

123. *E. Sibthorpii* (Boiss. Cent. Euph. p. 39. — D.C. Prodr. XV, p. 172) tomentella, caulis crassis inferne induratis cicatricosis superne crebre foliosis thyrsoides, ramulis umbellisque multifidæ radiis brevibus bifidis, foliis firmis basi longe attenuatis anguste lanceolato-linearibus acutis utrinque sed subtus densius albo-velutinis, umbellariis brevissimis, floralibus virenti-flavidis in patellam infundibuliformem ultra medium connatis, involueri campanulati extus et intus hirti lobis oblongis bifidis, glandulis melleis transverse latioribus truncatis ad angulos acutos muticis vel brevissime cornuto-attenuatis, capsula tomentosâ depressâ trisulcâ, semine ovalo ♀. *E. Characias* Sm. Prodr. I, p. 332 non L.

Hab. in collibus saxosis, ad rivulos et in olivetis regionis inferioris Graeciæ ad Athenas (Sprun! Sart! Heldr. Herb. Norm. 530 sub *E. Wulfeni*!), in Peloponneso (Bory!), insulâ Chios (Oliv!). Fl. vere.

Similis *E. Wulfeni* Hoppe foliis latioribus, floralibus minoribus, involueri lobis indivisis, glandulis longe bicornutis distinctæ. *E. Characias* patella florali concaviusculâ nec infundibuliformi differt.

124. *E. Lycia* (Boiss. in D.C. Prodr. XV, p. 172) tota pubescenti-velutina, caulis crassiusculis totâ longitudine laxiusculæ foliatis, umbellis 2-4 superpositis 8-12-radiatis, foliis firmis basi attenuatis anguste lanceolato-linearibus obtusiusculis mucronulatis, umbellariis elliptico-oblongis, floralibus flavidis in patellam concavam oblongam ultra medium coalitis, involueri intus hispidissimi lobis bifidis, glandulis flavis semilunatis in cornua convergentia attenuatis, capsula tomentosâ profunde trisulcâ, semiine ovato, carunculâ conicâ intus retusâ ♀. Boiss. Euph. Ic. tab. 117.

Hab. in rupibus maritimis Pamphyliæ prope Adalia (Bourg!). Fl. vere.

Sesquipedalis, folia bipollucaria tres lineas lata, radii bipollulares verticillorum intervallo æquilongi vel longiores. Affinis *E. Sibthorpii* differt foliis obtusioribus non confertis, superioribus verticillatis, glandularum cornubus longioribus, etc.

125. *E. Heldreichii* (Orph. in Boiss. Diagn. Ser. II, 4, p. 90) caulis crassiusculis elatis inferne induratis et denudatis, umbellis 2-5 superpositis et distantibus, radiis 7-15 bis vel ter bifidis, foliis breviter hirtis caulinum sterilium sparsis obovato-oblongis in petiolam attenuatis fertilium verticillatis sessilibus oblongo-lanceolatis, floralibus parvis rhomboideis obtusis antice ad medium usque connatis postice liberis, involueri campanulati fauce hirti lobis ovatis longe ciliatis, cornubus convergentibus glandulæ latitudine longioribus,

capsulâ glabrâ lævi profunde trisulcâ, semine ovato nigro, carunculâ patellari postice subrostratâ ♀. Boiss. Euph. Ic. tab. 118. — *E. verticillata* Orph. olim.

Hab. in sylvaticis Olympi Thessali (Heldr!), montis Malevo Laconiæ 4000' (Orph. Fl. exs. 201!).

Caulæ floriferi 2-3-pedales, umbellæ 2-3-pollices remotæ. Capsula magn. *E. amygdaloidis*.

β. *Ræseri* Boiss. loc. cit. — Humilior, folia brevia oblonga. — *E. Ræseri* Orph. MSS.

Hab. in regione superiore montis Parnassi (Heldr!).

§ 8. Myrsiniteæ.

* Semina vermiculato-rugosa.

126. ***E. Myrsinites*** (L. Sp. 661) caulis ascendentibus et decumbentibus dense foliosis, umbellæ radiis 7-12 semel vel bis bifidis, foliis margine minute sebris obovatis vel obovato-oblongis abrupte et breviter mucronatis, floralibus reniformibus transverse latioribus obtusissimis mucronulatis, involuci campanulati hemisphaerici lobis magnis ovatis circumcirca ciliatulis, glandulis transverse oblongis, cornubus glandulæ latitudini æquilongis apice dilatatis et sublobatis, capsulæ ovatae lævis coccis carinatis, semine ovato griseo obsolete tetragono crebre et elevatim vermiculato-rugoso, carunculâ stipitatâ conico-depressâ apice concavâ ♀. Sibth. Fl. Gr. tab. 471. — Rchb. Germ. fig. 4796. — *E. curtifolia* Bor. et Ch. Exp. Mor. p. 30.

Hab. in lapidosis regionis inferioris et montanæ, Græcia in Atticâ et Bæotijâ (Boiss! Heldr!), Peloponneso (Bory!), Thraciâ (Friv!), Bithyniâ occid. (Wied!), Cypro in monte Prodromos (Ky. 399 sub *E. Marschalliana*!), Tauriâ (herb. Petrop!).

Ar. Geogr. Italia australis et Sicilia, Dalmatia, Bosnia, Serbia, Banatus.

127. ***E. Anacampseros*** (Boiss. Diagn. Or. Ser. I, 5, p. 55) caulis paucis erectis vel ascendentibus sub umbellâ interdum parce ramulosis, sterilibus valde abbreviatis dense foliosis, foliis margine anguste cartilagineis et denticulatis rhombeo-ovatis obtusis mucronatis, superioribus majoribus magis dissitis a basi breviter cuneatâ orbiculatis, floralibus a basi subcordatâ semiorbiculatis obtusis mucronatis, umbellæ radiis 3-5 brevibus bis bifidis, involuci hemisphaerici lobis late ovatis denticulato-erosis ciliatis, glandulis rubellis vel luteis transverse ellipticis corunbus brevissimis latis palmatim et obtuse dentatis, capsulæ lævis ovatae coccis obtuse undulatis, semine obtuse tetragono crebre et longitudinaliter vermiculato-rugoso, carunculâ stipitatâ depresse conicâ apice foveolatâ margine dilatatâ ♀.

Hab. in regione alpinâ Cariæ (Pinard!), Lydiæ in monte Tmolo (Boiss!), Paphlagoniæ (Wied!).

B. minor. — Folia minora, involucrum intense purpureum, cornua spathulata sed vix lobata.

Hab. in Cappadociæ valle Kamechly Tchaï (Bal!), monte Almadagh Phrygiae (Bal!).

Species affinibus minus multicaulis, umbellis pauciradiatis, foliis superioribus majoribus magis dissitis, seminis vermiculis magis longitudinalibus a præcedente diversa.

128. *E. denticulata* (Lehm. Dict. II, p. 431) caulis asecentibus interdum sub umbellâ parce ramulosus, umbellæ radiis 5-7 bis bifidis, foliis crebris margine anguste cartilagineo-scabridulis obtusis in mucronem abrupte attenuatis, floralibus ovato-orbiculatis obtusissimis subconcavis, involuci hemispherici lobis latis ovatis ciliatis, glandulis purpureis transverse latioribus margine truncatis breviter multidentato-pectinatis, capsulae magnæ trigonæ basi truncatæ coccis acute carinatis, semine oblongo obtuse tetragono rugis elevatis anastomosantibus vermiculato-serobiculato, carunculâ depressâ conicâ apice foveolatâ ♂. Boiss. Euph. Ic. t. 119. — Desf. Coroll. Tourn. tab. 68 (mala). — *E. Cilicica* Boiss. Diagn. Ser. II, 4, p. 88. — *E. cristata* Dietr. Syn. V, p. 324. — *E. rotundata* Hochst. in Lor. Reis. p. 344.

Hab. in collibus et lapidosis regionis montanæ et subalpinæ, Cappadocia ad Cæsaream (Tourn! Bal!), Taurus Cilicus in regione inferiori (Auch. 2083! Bal!), ditio Kassan Oghlu (Ky. 97!), Armenia prope Gumuschkhané (Bourg!), Syria borealis in monte Soffdagh et Mesopotamia ad Terek (Haussk!).

Affinis *E. Anacampseroti* quâcum sœpe confusa differt glandularum formâ, foliis floralibus concavis, etc.

• • Semina tuberculata.

129. *E. craspedia* (Boiss. Diagn. Ser. I, 7, p. 95) caulis brevibus asecentibus, umbellæ multifidæ radiis brevibus bifidis, foliis late cartilagineo-marginatis denticulatis obovato-rhombeis obtusis mucronulatis, umbellaribus ovatis, floralibus ovato-orbiculatis, involuci hemispherici lobis ovatis ciliatis, glandulis purpureis transverse ovatis margine pectinato-dentatis, capsulae majusculæ coccis acute carinatis, semine albido acute quadrangulo leproso-tuberculato, carunculâ stipitatâ conicâ apice foveolatâ longitudinaliter sulcatâ ♂.

Hab. ad rupes Assyriæ prope Mardin (Auch. 2708!), inter Mardin et Assuaner (Ky. 350!).

Affinis *E. denticulata* differt foliorum margine cartilagineo lineam fere lato, seminis sculpturâ; caulis semipedatis, umbellæ radii 1 $\frac{1}{2}$ -2-pollicares.

• • Semina laevia.

130. *E. Marschalliana* (Boiss. Diagn. Ser. I, 7, p. 94 excl. synon.) caulis asecentibus et decumbentibus crebre foliosis,

umbellæ multifidæ radiis brevibus bis bifidis, foliis margine cartilagineo-denticulatis elliptico-rhombeis basi et apice attenuatis acutis subpungentibus, umbellaribus ovatis floralibus a basi subcordatâ orbiculatis obtusis mucronulatis, involucri campanulati lobis coalitis denticulatis ciliatulis, glandulæ transverse ellipticæ cornubus latitudini glandulæ æquilongis crassis apice subdilatatis, capsulæ lœvis coccis obtuse angulatis, semine albido oblongo-tetragono lœviusculo, carunculâ conicâ acutâ sulcatâ ♀.

Hab. in montanis Armeniae (Auch. 5310!), prope Van (Ky. 1471), in Transcaucasiâ ad Atskur (Vittm!), prov. Talysh (Hoh. exs. sub *E. myrsinites*!), prov. Aderbidjan Persiae (Szovi!).

Folia eis *E. Myrsinites* minora acutiora et magis denticulata, seminibus ab eâ diversissima.

131. *E. biglandulosa* (Desf. Cor. Tourn. p. 88, tab. 67) caulinis crassis erectis striato-sulcatis, umbellæ multifidæ radiis brevibus semel vel bifidis, foliis imbricatis intagerrimis lanceolatis acutis subpungentibus, umbellaribus ovatis, floralibus a basi cordatâ reniformibus obtusis mucronulatis, involucri breviter et late campanulati lobis magnis ovatis ciliolato-denticulatis, glandulæ transverse oblongæ cornubus crassis apice capitatis lobulatis, capsulæ majusculæ trigonæ elevato-punctatæ coccis dorso angulatis, semine albido oblongo-tetragono lœvi, carunculâ depressâ lobulato-plicatâ ♀. *E. rigida* M.B. Taur. Cauc. I, p. 375. — Bot. Reg. t. 274 et vol. XXIV, tab. 43.

Hab. in collibus siccis regionis inferioris et montanæ, Græcia in monte Cœnos Cephaloniae (Heldr!), Arcadiâ, Laconia, prope Corinthum (Boiss! Sart! etc.), insula Cos (Urv!), Hellesponto (Herb. Mus. Par!), Anatoliâ in Cariâ (Auch. 2024!), Cappadociâ (Bal!), Ciliciâ (Ball!), Pamphylia (Bourg!), Syriâ boreali (Haussk! Gail!), Tauriâ (M.B. in herb. Petrop!).

Caules sœpe sesquipedales, capsula 3-3½ lin. longa. Nomen impropprium ad bina cornua glandulæ alludit.

Ar. Geogr. Sicilia.

Species obscuræ probab. præcedentium synonyma.

E. Belgradica (Forsk. Eg. Ar. p. 211) erecta, ramis alternis dichotomis, foliis oppositis sessilibus, caulinis oblongis serrulatis, superioribus semicircularibus, capsulis solitariis axillaribus tuberculatis.

Hab. in sylvâ Belgrad prope Byzantium (Forsk.).

Probab. forma æstivalis *E. pubescens* vel *platiphyllæ*.

E. densifolia (C. Koch Linn. XXI, p. 722) glabra, caule stricto dense folioso, umbellâ 3-5-radiatâ, foliis oblongis serrulatis subito in petiolum attenuatis, umbellaribus lanceolatis, floralibus ovato-acuminatis, glandulis transverse oblongis.

Hab. in Transcaucasiâ (Wilh.).

An *E. stricta*?

E. dicotoma (Forsk. Eg. Ar. p. 93) caule suberecto dichotomo spithameo; foliis oppositis ovato-lanceolatis sessilibus articulo brevioribus parvis integris, inferioribus majoribus margine denticulatis, anthodiis axillaribus solitariis sessilibus, capsula glabra.

Hab. in siccis Ras el Tin Egypti (Forsk.).

E. mucronata (Clarke in Spr. Neu Entd. II, p. 115) fruticosa glabra, foliis ovato-lanceolatis mucronatis integris, umbellaribus ovatis, floribus obovatis integrerrimis, petalis dentatis, capsulis verrucosis glabris.

Hab. in monte Gargaro Troadis (Clarkæ).

TR. II. PHYLLANTHEÆ Müll. Arg.

Florum involucrum nullum rarius unisexuale. Calyx 5-6-partitus laciiniis imbricatis. Petala calycis laciiniis alterna vel nulla. *Masc.* Stamina 5 vel plura antheris in alabastro erectis. *Fem.* Ovarium triloculare loculis biovulatis. Cotyledones latæ.

ANDRACHNE (L. Gen. 709).

Flores monoici. Calyx plus minus profunde quinquepartitus. Petala totidem membranacea calyce breviora, feminea minuta. Discus glandulis quinque (in nostris) inter se liberis petalis oppositis instructus. *Masc.* Stamina quinque filamentis basi plus minus connatis, antheræ breves biloculares, pistillum abortivum. *Fem.* Corolla diminuta vel abortiva. Ovarium glandulis disci basi suffultum triloculare loculis biovulatis, ovulis angulo centrali affixis. Styli tres bifidi vel bipartiti, stigmata subclavata. Capsula trilocularis tricocca coccis dispermis. Semina ecarunculata. Embryo in albumina carnosæ centralis, radicula hilum spectans. — Suffruticuli foliis minutis alternis stipulatis.

1. A. Colchica (Fisch. et Mey. MSS.) glabra glaucescens fruticosa ramosa ramis erectis flexuosis denudatis hornotinis crebre foliosis, foliis breviter petiolatis ovatis obtusis, stipulis minimis nigris triangularibus fimbriatis, floribus axillaribus longe et tenuiter pedicellatis masculis 2-3-nis feminineis solitariis, calycis laciiniis ad quintam partem usque coalitis obovatis obtusis, petalis filiformibus, disci glandulis obtuse bilobis calyce triplo brevioribus petala æquantibus, capsula depressâ glabra 5.

Hab. in saxosis cretaceis Imeretiæ et Mingreliaæ 800'-4000', in sylvaticis ad fluvium Zchenis Zkale prope Kutais (Szedinsky! Rupr!).

Frutex bipedalis et prob. elatior ramis inferioribus fere pennæ anserinæ

crassitie, folia tandem 9-10 lineas longa, capsula semine *Cannabeos* paulo major. Species a congeneribus ditionis caule fruticoso eximie distincta, a cl. Szedinsky ad *A. Chinensem* Bge relata ex descriptione foliis majoribus acutis, capsula multo majore sex millim. latâ aliquis notis alienam.

2. A. telephiooides (L. Sp. 1014) glabra glauca, caulis e rhizomate indurato vel suffrutescenti numerosis prostratis filiformibus simplicibus vel ramulosis crebre foliosis, foliis petiolatis obovatis vel ellipticis acutiusculis, stipulis triangularibus peltatis membranaceis laceris, floribus axillaribus breviter pedicellatis subsolitariis rarius 2-3-nis, calycis laciniis basi attenuatis rhombis acutiusculis anguste albo-marginatis, petalis lanceolatis calyce subbrevioribus, disci glandulis bilobis petalis triplo brevioribus, capsula depressâ glabrâ, seminibus angulatis punctatis dorso convexis 2. Ic. Fl. Græc. tab. 953. — Nees. Ic. Germ. — *A. rotundifolia* C.A. Mey. in Eichw. Casp. p. 18, tab. 20.

Hab. in incultis et collibus siccis regionis inferioris et montanæ totius ditionis a Græciâ et ejus insulis! Thraciâ! ad Tauriam et regionem Caucasum! Persiam borealem et australem, Syriam! Palestinam! Affghaniam (Griff!), Belutschiam (Stocks!).

A. rotundifolia forma est foliis paulo obtusioribus rotundatis, floribus interdum ut in formâ typicâ 2-3-nis.

Ar. Geogr. Hispania et Italia australes, Dalmatia, Africa borealis, insulæ Capitis Viridis.

3. A. fruticulosa (Boiss. Diagn. Ser. I, 7, p. 86) fruticulosa, ramis rigidis patule et intricatim ramosissimis paucifoliis inferne sub lente papillari-asperis, foliis oblongo-ellipticis subsessilibus margine sub lente asperis glaucis, stipulis membranaceis pulvilliformibus multifidis, floribus axillaribus subsolitariis, calycis laciniis oblongis obtusis albo-marginatis, petalis lanceolato-spathulatis subtruncatis calyce triente brevioribus, disci glandulis bifidis, capsula globosâ, seminibus angulatis oblongis 3.

Hab. in Persiâ australi prope Schiraz (348!), in deserto salso inter Teheran et Isfahan prope Robut Kerim (Haussk!), inter Kerman et Yezd (Bge!).

Ex speciminibus nimis incompletis nota sat ab *A. telephioide* differre videtur ramis crassioribus rigidulis duris aliquis notis in diagnosi indicatis. Capsula duplo major, semina magis elongata.

4. A. aspera (Spreng. Syst. III, p. 884) glauca papilloso-aspera vel hirtella, ramis a rhizomate lignoso numerosis tenuibus procumbentibus ramulosis saepe elongatis, foliis petiolatis reniformibus vel orbiculatis basi cordatis obtusissimis superioribus sensim diminutis sessilibus, pedunculis ad axillas superiores 3-4-floris inferioribus folio axillari longioribus paniculas angustas vel racemos simplices terminales laxos formantibus, calycis hirti laciniis lanceolatis obtusis, petalis calyce brevioribus, disci florum masculorum glandulis peltato-reniformibus, femineorum anguste obovatis truncatis, capsula hirtâ, seminibus angulatis oblongis 3.

Hab. in desertis Egypti superioris (Boiss! Schleinf!), Arabiae petreas (Schimp. 401! Auch! Boiss!), Mesopotamiae prope Orfa (Haussk!), Persiae austro-occidentalis (Ky. 953! Haussk!).

Variat foliis .inimis vel eis *A. telephoides* majoribus, caulis brevibus vel 1-1½-pedalibus.

Ar. Geogr. Nubia, Abyssinia, Arabia tropica.

PHYLLANTHUS (Sw. Fl. Ind. occid. II, p. 1101).

Flores monoici rarius dioici. Calyx 5-6-partitus. Petala nulla. Discus glandulifer interdum suppressis obsitus. Masc. Stam. 3-5 rarius plura filamentis in columnam connatis vel liberis. Fem. Ovarium triloculare loculis biovulatis. Styli tres saepius bifidi. Capsula trilocularis tricocca coccis dispermis. Semina ecarunculata. Embryo in albume carnosus centralis, radicula hilum spectans. — Frutices, suffrutes rarius herbæ, foliis alternis saepe distichis.

1. **P. rotundifolius** (Willd. Sp. IV, p. 584) annuus monoicus erectus glaber ramosissimus, runulis compresso-angulatis, floribus subgeminis breviter pedicellatis, calycis (fructiferi aceret) lacinis obovatis margine hyalinis, filamentis in columnam connatis, capsula levii, seminibus dorso longitudinaliter costatis ⊖.

Hab. in Egypto (Schimp. exs. 750).

Species tropica vix ditionis nostræ. *P. Niruri* L. species affinis ramulis teretibus distincta ex Belutschia in Aitches. Catal. indicatur.

Ar. Geogr. Africa tropica, India orientalis.

TR. III. CROTONEÆ Müll. Arg.

Flores non involucrati. Calycis laciniae subvalvares apice tantum quincunciales. Petala calycis lacinis alterna vel nulla. Masc. Stamina numero varia antheris in alabastro infracto-incurvis. Fem. Ovarium triloculare loculis uniovulatis. Cotyledones latæ.

CROZOPHORA (Neck. Elem. II, p. 337).

Flores monoici. Calyx quinquepartitus praefloratione valvatus. Discus glandulosus quinquelobus. Masc. Petala quinque cum calycis phyllis alterna. Stamina 5-15 1-3-verticillata filamentis basi in columnam centralem coalitis, antheræ biloculares connectivo excurrente mucronatae. Fem. Petala 5 angustiora interdum abortiva, staminodia 3 vel nulla. Ovarium globosum triloculare triovulatum. Styli tres bifidi vel bipartiti altero latere stigmatosi. Capsula ovato-globosa tricocca coccis monospermis bivalvibus. Semina apice attenuato tri-

gona ecarunculata. Embryo in albumine oleoso centralis, cotyledones amplæ, radicula brevis. — Herbæ pilis stellatis vel lepidotis instructæ foliis alternis stipulatis.

1. C. plicata (Vahl Symb. I, p. 75 sub *Crotone*) annua induimento stellato plus minus denso griseo tomentella vel tomentosa, ramis a collo prostratis elongatis dichotomis, foliis longe petiolatis a basi obliquâ subcordatâ ovatis obtusis repandis vel subtrilobis, stipulis linearibus, racemis axillaribus et terminalibus superne masculis inferne femineis, floribus masculis subsessilibus spicatis 15-10-andris, petalis lanceolato-ovatis, pedunculis femineis 1-3-floris demum deflexis, petalis femineis sublinearibus, capsulâ pilis stellatis tomentosâ, seminibus læviusculis ♂. *Crozophora plicata* A. Juss. Tent. Euph. p. 28. — Schweinf. pl. Nilot. tab. 4. — *Crozophora prostrata* Dalz. Bombay Fl. p. 233. — *Croton Rottleri* Geis. Cr. Mon. — *C. obliquifolium* Visian. pl. Egypt. t. 7.

Hab. in cultis Egypti inferioris et superioris (Auch. 2005! Wiest. 517! Boiss! etc.).

Ar. Geogr. Nubia, Abyssinia, Africa et Arabia tropicæ, regio Scinde et India.

2. C. gracilis (F. et M. in Kar. Enum. Turcom. 1839, p. 171) annua brevissime et stellatim tomentella grisea, ramis a collo diffusis ramosis, foliis longe petiolatis subcordato-ovatis acutiusculis integris basi obsolete 1-2-glandulosis, stipulis obsoletis, racemis axillaribus superne masculis inferne femineis, floribus masculis subsessilibus spicatis sub 5-andris petalis lanceolatis, pedunculis femineis 2-6-floris demum valde elongatis deflexis, petalis linearibus, capsulâ pilis stellatis tomentosâ, seminibus lævibus ♂. *Croz. sabulosa* Kar. et Kir. Bull. Mosq. 1842, p. 446.

Hab. in sabulosis Turkestaniae inter fluvios Kuwan et Jandarja (Lehm!).

Indumentum *C. tinctoria*, planta gracilior, folia integra, flores et capsulae minores.

Ar. Geogr. Songaria.

3. C. tinctoria (L. Sp. 1425 sub *Crotone*) annua, tenuiter et stellatim tomentosa vel hirta grisea, caule humili erecto subdivaricatim ramoso, foliis longe petiolatis a basi rotundatâ vel breviter cuneatâ integrâ rhombo-ovatis obtusis repandis subtus prope basin biglandulosis, stipulis anguste linearibus, racemis axillaribus et terminalibus superne masculis inferne fæmineis, floribus masculis subsessilibus spicatis sub 10-andris, calycis laciniis lineari-lanceolatis petalis linearibus, pedunculis femineis 1-4-floris ad racemi basin elongatis demum deflexis, capsulis lepidotis, -seminibus tuberculato-asperis ♂. *Crozophora tinctoria* Ad. Juss. — Ic. Fl. Gr. tab. 950. — Rehb. Ic. Germ. tab. 152. — Nees Ic. Germ.

Hab. in cultis regionis inferioris a Græcia! et ejus insulis! Macedonia (Friv!), ad Tauriam et Transcaucasiam (Ledeb.), Persiam borealem (Szov!) et austrialem (Haussk!), Syriam! Palestinam (Roth!), Egyptum (Ehr.).

β. subplicata Müll. Arg. in D.C. Prodr. — Folia acutiora sinuato-lobulata secus nervos interdum plicata. — An forma hybrida?

Hab. circa Kahiram (Bové! Husson!), prope Kosseir ad Mare Rubrum (Schweinf!).

Ar. Geogr. Regio mediterranea Europæ et Africæ borealis.

4. **C. verbascifolia** (Willd. Sp. IV, p. 539 sub *Crotone*) annua indumento stellato denso tomentoso-pannosa, caule humili erecto subdivaricatum ramoso, foliis longe petiolatis a basi rotundata vel subcordata ovatis vel oblongis saepe acutalis repandis subtus elevatim nervosis et prope basin biglandulosis, stipulis minutis linearibus, racemis axillaribus et terminalibus superne masculis floribus subsessilibus spicatis subpentandris inferne femineis pedunculis 1-4-floris tandem elongatis deflexis, calycis lacinias petalisque lineari-lanceolatis, capsulis lepidotis, seminibus tuberculato-asperis ♂. *Crozophora verbascifolia* Juss. — *Croton villosus* Fl. Græc. tab. 951. — *C. Hierosolymitana* Spreng. Syst. III, p. 850! — *C. Sieberi* Presl Bemerk. p. 109. — *C. integrifolia* Bge pl. Lehm. p. 490.

Hab. in cultis et sterilibus Græciae in Atticâ et Argolide (Sprun! Sart!), Cretæ (Oliv!), Anatoliæ occid. et australis (Boiss! Heldr!), Syriae littoralis et interioris (Auch! Bl!), Palestinæ (Siebl!), Mesopotamiae et Assyriæ (Ky. 441! Haussk!), Kurdistanæ Persicae ad 800' usque et Persie austro-occident. (Haussk!), Persie bor. (Buhsel!) et orientalis (Bgef!), Affghaniae (Griff!), Turkestaniae (Lehm!).

Ar. Geogr. Hispania austro-orientalis, Africa borealis.

5. **C. obliqua** (Vahl Symb. I, p. 78 sub *Crotone*) perennis stellatim tomentosa grisea fruticosa ramosa ramis erectis, foliis breviter petiolatis a basi rotundata vel breviter cuneatâ oblongo-lanceolatis acutiusculis vel acuminatis repando-subundulatis subtus prominenter nervosis et prope basin biglandulosis, stipulis linearibus mox deciduis, racemis axillaribus et terminalibus superne masculis inferne femineis, floribus masculis subsessilibus spicatis subpentandris petalis obovalo-lanceolatis, pedunculis femineis 1-2-floris demum deflexis, petalis linearibus, capsulis lepidotis, seminibus tuberculato-asperis ♀. *Crozophora obliqua* A. Juss. Tent. p. 28. — Schweinf. Pl. Nilot. tab. 3. — *Croton oblongifolium* Del. Fl. Eg. p. 379, tab. 51, fig. 1.

Hab. in desertis Egypti prope Suez (Del! Schweinf!), inter Kosseir et Ras Benass (Schw!), in Oasi Dachel (Aschers!), in Arabiâ petreæ ad Quadi Feiran (Auch. 2004! Schimp. 359! Boiss!), in Arabiâ Mascatensi (Auch. 5296!). Arabicæ *Sammua*.

Ar. Geogr. Nubia, Abyssinia, Africa et Arabia tropicæ, India borealis.

TR. IV. ACALYPHEÆ Müll. Arg.

Florum involucrum nullum vel unisexuale. Calycis masculi laciniaæ æstivatione valvares. Petala calycis lacinias alterna vel nulla.

Masc. Stamina numero varia antheris in alabastro erectis. *Fem.* Ovarium triloculare loculis uniovulatis. Cotyledones latæ.

MERCURIALIS (L. Gen. 1125).

Flores dioici rarius monoici. Calyx 3-4-partitus masculus valvaris, femineus imbricatus Petala nulla. *Masc.* Discus nullus. Stamina 8-12 in receptaculo elevato centralia. Antheræ loculi bini distincti connectivo globoso connexi. *Fem.* Discus evolutus. Ovarium biloculare loculis uniovulatis. Stylus brevis, stigmata bina elongata facie internâ stigmatosa. Stamina 2-3 ad filamentum reducta. Capsula bi rarius tricocca. Semen pendulum carunculatum, testa fragilis raphe longitudinali instructa. Embryo in albumine carnoso centralis, radicula supera. — Herbae oppositifoliae stipulatæ.

1. **M. perennis** (L. Sp. 1465) perennis, caulibus e rhizomate longe repente ad nodos radicanti erectis simplicibus basi nudis, foliis glabris vel hirtis breviter petiolatis elliptico-oblongis acutis serrulatis, floribus masculis in glomerulos secus pedunculos axillares laxos dispositis, femaleis 1-2 ad axillas pedunculatis, capsulâ breviter et sparsim pilosulâ, seminibus globoris subfoveolatis ♀. Ic. Nees Germ. — Rchb. Germ. t. 152.

Hab. in sylvis regionis montanæ, Parnassus (Fraas), mons Delphi Eubeæ (Heldr!), Thracia ad Byzantium (Sibth.), Pontus Lazicus (Bal!), Tauria et prov. Caucasicæ (M.B.), Persia boreali-orientalis ad Siaret (Bge!).

β. ovata Müll. in D.C. Prodr. — Folia ovata subsessilia. — *M. ovata* Sternb. et Hoppe.

Hab. in collibus Armeniæ prope Gumuschkhané (Bourg!).

Ar. Geogr. Europa omnis a Scandinaviâ ad Rossiam, Africa borealis. Varietas *ovata* in Alpibus Germanicis, montibus regionis Dauubialis, Rossia media.

2. **M. annua** (L. Sp. 1465) annua, caule erecto ramoso ad nodos incrassato, foliis glabris breviter petiolatis ovatis vel ovato-lanceolatis acutis margine crenatis vel obtuse serratis ciliatulis, floribus masculis in glomerulos approximatos vel dissitos secus pedunculos axillares folio longiores dispositis, floribus femaleis ad axillas solitariis vel binis subsessilibus, capsulâ tuberculis in papillas acutas albas abeuntibus obsitâ, seminibus ellipticis foveolato-tuberculatis ♂.

Hab. in cultis partis occidentalis ditionis a Græciâ! ad Syriam! Egyptum (Ehr!), Tauriam (Lebed.).

Varietatem monoicam (*Merc. ambigua* L.) e ditione nondum vidi.

Ar. Geogr. Europa media et australis, Africa borealis.

RICINUS (L. Gen. 1464).

Flores monoici. Calyx 3-5-partitus præfloratione valvatus. Petala nulla. Discus nullus. *Masc.* Stamina plurima e receptaculo plano-convexo in fasciculos polyandros superne multoties ramosos connata, antheræ biloculares didymo-globosæ. *Fem.* Ovarium ovato-globosum triloculare triovulatum. Stylus triseptus segmentis linearibus indivisis vel bipartitis aut bifidis hinc hirto-stigmatosis. Capsula ovato globosa tricocca coccis monospermis bivalvibus. Semina carunculâ bilobâ instructa. Albumen copiosum oleosum, cotyledones amplæ, radicula brevis. — Frutex.

1. **R. communis** (L. Sp. 1430) fruticosus ramis novellis et petiolis glauco-pruinosis, foliis alternis amplis petiolatis peltatis palmatim 7-11-lobatis dentatis, stipulis in unicum amplexicaulem connatis, floribus in racemos oppositifolios vel terminales iuferne masculos superne femineos dispositis, pedicellis ad medium articulatis, capsulâ plus minusve echinatâ ȝ. Ic. Fl. Græc. tab. 752.

Hab. ad sepes regionum calidiorum subsppontanea, Græcia, Anatolia occid. et australis! Syria! Egyptus! Persia australis! Belutschia (Stocks).

Ar. Geogr. Regio tropica et subtropica Asiæ, Africæ et Americæ.

ORD. CXII. BUXACEÆ.

(Klotzsch Tric. p. 12.)

Flores unisexuales. Calyx liber laciniis aestivatione imbricatis. Petala et discus nulla. *Masc.* Stamina in centro floris inserta. *Fem.* Ovarium liberum superum 2-3-loculare loculis 1-2-ovulatis ovoidis anatropis pendulis. Fructus baccatus vel capsularis et tunc loculicide dehiscens. Semina albuminosa embryone recto axili, radiculâ superâ. — Ordo Euphorbiaceis valde affinis a quibus capsulis loculicidis fere tantum differt.

BUXUS (L. Gen. 1053).

Flores monoici. Perigonum basi unibracteatum phyllis quatuor inæqualibus duobus, interioribus et duobus exterioribus invicem oppositis. *Masc.* Stamina 4 receptaculo inserta laciniis perigonii opposita, antheræ biloculares loculis connectivo incrassato insertis, pollinis grana non reticulata. *Fem.* Perigonum phyllis quatuor basi

trisquamatum. Ovarium triloculare loculis biovulatis, styli tres breves crassi, stigmata magna apice intus canaliculata et stigmatosa. Capsula coriacea stylis persistentibus trirostrata trilocularis loculicide trivalvis, valvae apice stylorum divisione lateraliter bimucronatae, loculi dispermi. Semina pendula albuminosa. — Frutices foliis coriaceis sempervirentibus oppositis, floribus ad axillas superiores fasciculatis.

1. *B. sempervirens* (L. Sp. 1394) fruticosa ramosissima, ramis erectis junioribus puberulis et foliosis, foliis ovatis vel ovato-oblongis, floribus sessilibus fasciculatis lateralibus masculis terminali fæmineo, perigonii lacinias ovatis, antheris oblongis, ovarii cornubus eo subduplo brevioribus ♂. Ic. Nees Gen. Germ.

Hab. in regione sylvaticâ in subalpinam et alpinam ascendens, Græcia borealis in Pindo (Sibth.), Olympo Thessalo (Heldr! Orph. Fl. Gr. exs. 697!), Macedonia (Griseb.), Byzantium (Cast!), Bithynia (Griseb.), Caria in montibus supra Denisleh (Ball), Caucasi latus meridionale, Transcaucasia, ditio Talyssch (Ledeb.), Persia boreali-orientalis supra Siaret (Bge!). Turcis *Gimschir*, græcis *Pyxari* audit.

Ar. Geogr. Europa media ab Angliâ et Belgio ad Germaniam medium et Dalmatiam, Hispaniam medium et Lusitaniam, Sibiria Uralensis, Japonia, Africa borealis.

2. *B. longifolia* (Boiss. Diagn. Ser. I, 12, p. 407) fruticosa ramosissima ramis erectis glabris foliosis, foliis majusculis elliptico-linearibus obtusis, floribus fasciculatis masculis lateralibus breviter pedicellatis, perigonii laciñiis oblongo-lanceolatis, antheris oblongo-sagittatis, ovarii cornubus eo vix brevioribus, capsula magnâ rostris apice recurvis longis ♂.

H.b. in regione montana Syriæ borealis circâ Antiochiam et versus montem Cassium (Boiss!).

Differt a *B. sempervirente* foliis subduplo majoribus 13-15 lineas longis 4-5 latis. Capsula fere duplo major longius rostrata cum rostris 6-7 lineas longa.

ORD. CXIII. EMPETRACEÆ.

(Lindl. Nat. Syst. 117.)

Flores dioici vel polygami Calyx liber 2-3-phyllos imbricatus herbarius. Corollæ petala hypogyna sepalis numero æqualia eis alterna receptaculo inserta. *Masc.* Stamina petalis numero æqualia et eis alterna cum pistilli rudimento, antheræ cordatae biloculares. *Fem.* Ovarium disco insidens carpidiis 2-9 marginibus introflexis columellæ centrali adnatis verticillatim connatis constans. Ovula in loculis solitaria ex basi anguli centralis ascendentia anatropa. Stylus brevis, stigma radiato-lobatum. Bacca sphærica pyrenis 2-9 axi cohærentibus

vel distinctis osseis monospermis. Semina erecta. Embryo rectus in albumine carnosō excentricus, radicula supera. Ordo *Euphorbiaceis* et *Celastrineis* affinis.

EMPETRUM (L. Gen. 1100).

Flores dioici. Calyx squamis imbricatis cinctus coriacens trisepalus. Petala tria sepalis longiora. Stamina tria exserta filamentis elongatis. Stylus unicus, stigma radiatum 6-9-partitum. Ovarium subglobosum 6-9-loculare. Drupa globosa bacciformis mesocarpio carnosō, pyrenis 6-9 cuneiformibus demum distinctis. — Fruticuli ericoidei floribus parvis sessilibus subsolitariis infra ramulorum apicem axillaribus basi squamis bractealibus 5-6 imbricatis suffultis.

1. E. nigrum (L. Sp. 1450) suffruticosum ramosissimum decumbens, ramis ascendentibus novellis crebre foliatis, foliis minimis linearī-oblongis obtusis rigidis obtusis margine revolutis et asperulis dorso lineato-depressis, baccis nigris 3. Rehb. Ic. Germ. t 158.

Hab. in saxosis frigilis regionis alpinæ, Pontus Lazicus supra Khabackar (Bal!), montes Guriæ (Nordm!), Caucasus centralis et occid. 7-9000' (C.A. Mey., Rehm!), orientalis 8-9000' (Becker! Rupr!). Probab. quoque in alpibus Macedonie.

Ar. Geogr. Europa arctica, sept. et media usque ad montes Hispaniæ borealis et Italiae mediæ, Sibiria, Japonia, America arctica et borealis.

ORD. CXIV. URTICACEÆ.

(Juss. Gen. p. 400.)

Flores monoici dioici vel polygamii. Perigonum sèpius herbaceum inferum varie partitum vel in floribus femineis integrum laciñiis aestivatione imbricatis vel valvatis. Stamina definita libera fundo perigonii inserta et ejus laciñiis opposita. Ovarium in nostris liberum uniloculare et uniovulatum, vel biloculare loculis uniovulatis. Styli 1-2. Fructus indehiscent, embryo rectus, curvatus vel spiratis, radicula supera. — Herbæ frutices et arbores foliis stipulatis, stipulis liberis sèpius caducis.

TRIB. I. URTICEÆ

Ovarium uniloculare ovulo erecto. Achenium perigonio tectum. Semen erectum, embryo in albuminis axi rectus — Herbæ.

URTICA (L. Gen. 1054).

Flores monoici vel dioici. *Masc.* Perigonium quadripartitum laciniis æqualibus concavis valvatis sub anthesi patentibus. Stamina 4 laciniis perigonii opposita earum basi inserta, antheræ oblongo-reniformes, interdum ovarii rudimentum cupuliforme. *Fem.* Perigonium tetraphyllum phyllis exterioribus (in nostris) minoribus, interioribus planis. Stigma sossile penicilliforme. Achenium oblongum compressum perigonio persistente et increscente tunicatum. — Herbæ oppositifoliæ, nostræ omnes pilis urentibus armatae.

1. U. urens (L. Sp. 1396) annua setosa monoica, caule erecte a basi ramoso, foliis ovalis argute inciso-dentatis petiolo subæquilongo suffultis, stipulis inter petiolos utrinque binis, racemis androgynis axillaribus simplicibus petiolo saepius brevioribus, floribus fæmineis numerosioribus ♂. Ic. Rehb. Germ. 652.

Hab. in ruderatis totus regionis a Græciâ! ad regionem Caucasicam! Syriam (Bl!), Egyptum (Schw! Aschers!).

Ar. Geogr. Europa omnis, Asia orientalis, Africa borealis, Abyssinia.

2. U. dioica (L. Sp. 1396) perennis setosa dioica, caulinis erectis subsimplicibus, foliis a basi cordatâ ovatis vel oblongo-lanceolatis acuminatis argute inciso-dentatis petiolo limbo breviore suffultis, stipulis inter petiolos utrinque binis, florum glomerulis in racemos axillares ramosos graciles a basi floriferos petiolo longiores dispositis ♀. Ic. Rehb. Germ. tab. 654. — *U. hispia* d.c. Fl. Fr.

Hab. in ruderatis ditionis magis borealis, Thracia (Silth.), Tauria (Stev.) et frequentior in saxosis umbrosis vel humidis regionis subalpinæ et alpinæ usque ad nives, Græcia in montibus Olenos (Heldr!), Parnassus (Sart!), Timolus (Bal!), Taurus Cilicus (Ky! Bal!), Libanus (Bl!), mons Hermon (Gaill!), Alpes Ponti Lazici (Bal!), Caucasus (Led.), mons Avroman Kurdistanie Persie (Haussk!).

β. *pubescens* Trautv. — Molliter pubescenti-canescens, stimulis fere omnino destituta. — *U. pubescens* Led. Fl. Ross. III, p. 638. — Eichw. Casp. tab. 39. — *U. subnitis* Boiss. in Bal. exs.

Hab. in Anatoliâ prope Smyrnam (Bal!), Caucaso ad Nartzana (Led.).

Ar. Geogr. Europa, Asia borealis, Africa borealis, America borealis.

3. U. Haussknechtii. perennis dioica, caulinis virgatis erectis opposite ramulosissimis adpresso puberulis et stimulis parce obsitis, foliis parvis rugulosis subtus ad nervos tres adpresso seabridis breviter petiolatis a basi rotundato-cuneatâ oblongo-lanceolatis acutis argute inciso-serratis, stipulis interpetiolaribus utrinque binis lanceolatis minimis, florum masculorum racemis axillaribus simplicibus 2-4-floris hispidis petiolo subæquilongis, perigonii flavigredi glabriuscili

laciiniis ovatis, floribus fæmineis.... ♀. *U. microphylla* Boiss. et Haussk. exs. non Sw.

Hab. ad mænia urbis Eski Malatia Cataoniae (Haussk!).

Caules 1-2-pedales ramis elongatis erecto-patulis, folia majora pollicaria inferne 5-6 lineas lata. Species subinermis viridis foliis subtus pallidioribus, racemis masculis foliisque angustis parvis insignis sed floribus fæmineis fructuque nondum observatis imperfecte nota.

4. *U. cannabina* (L. Hort. Ups. p. 282) perennis monoica, caulinibus erectis subsimplicibus subinermibus, foliis petiolis setosis suffultis sparsim setosis tripartitis segmentis pinnatipartitis, stipulis inter petiolos utrinque binis, racemis unisexualibus axillaribus petiolo longioribus inferioribus ramosis masculis superioribus simplicibus spiciformibus fæmineis ♀.

Hab. in Persiâ boreali prope fluvium Chahroot (Auch. 5314 ex Wedd.). Editione non vidi.

Ar. Geogr. Songaria, Sibiria, Dauria, China.

5. *U. pilulifera* (L. Sp. 1395) annua vel biennis monoica setosa cæterum pilosula vel glabra, caulinibus erectis sæpe ramosis, foliis petiolatis ovatis vel cordatis argute inciso-dentatis, stipulis inter petiolos utrinque binis, inflorescentiis masculis et fæmineis in eisdem axillis vel inferioribus omnibus masculis hisce pedunculatis racemosis ramosis petiolo longioribus, spicis fæmineis globosis pedunculatis in pedunculo solitariis rarius binis tandem nutantibus ♂ vel ♀. Ic. Rehb. Germ. tab. 653. — *U. Balearica* L. loc. cit. — *U. Chesneyana* Blume Mus. bot. Lugd. II, p. 152.

Hab. in ruderatis regionis inferioris totius ditionis a Graeciâ! Macedoniâ! ad Syriam (BH Gaill!), Palestinam (Boiss!), Mesopotamiam (Haussk!), Tauriam et Transcaucasiam (Ledeb.), Ægyptum.

Ar. Geogr. Europa media et australis ab Angliâ et Germaniâ med. à ad Transylvaniam et Dalmatiam, Africa borealis.

6. *U. membranacea* (Poir. Diet. IV, p. 638) aunua setosa monoica vel interdum dioica, caulinibus erectis vel ascendentibus ramosis, foliis petiolatis cordato-ovalis inciso-serratis, stipulis utrinque inter petiolos solitariis, floribus in spicas axillares geminalis unisexuales dispositis inferioribus fæmineis petiolo brevioribus rachide non dilatata, superioribus masculis petiolum superantibus rachide a medio membranaceo dilatata et facie superiore tautum floriferâ ♂. Ic. Rehb. Germ. tab. 652. — *U. canadensis* Vahl non Peir.

Hab. in ruderatis regionis inferioris Peloponnesi (Bory!), Atticae (Sart!), Zacynthi (Marg.), Crete (Sieb!), Panophyliae ad Adalia (Bourg!), Syria littoralis ad Sidonem (BH!).

Ar. Geogr. Regio mediterranea Europeæ et Africæ borealis.

MEMORIALIS (Hamilt. in Wall. Cat. № 1598).

Flores monoici bracteis scariosis stipati glomerulos axillares constituentes. *Masc.* Perigonium 4-5-parlitum laciñis ovalis apice abrupte infractis et plicâ transversâ saepè ciliatâ instruetis. Stamina 4-5 laciñis perigonii opposita. *Fem.* Perigonium tubulosum vel ovatum ore contracto 2-4-dentatum, fructiferum saepè alatum vel costatum. Ovarium inclusum liberum vel perigonio plus minusve cohærens. Stigma filiforme cum apice ovarii articulatum et cito deciduum altero latere villosum. Achenium ovario conforme pericarpio crustaceo nitido. Herbeæ perennes inermes, foliis oppositis, nervis basilaribus usque ad apicem limbi productis.

I. M. pentandra (Roxb. Ind. III, p. 583 sub *Urticā*) caulinibus erectis ramosis, foliis albo-tuberculatis trinerviis inferioribus oppositis a basi rotundato-subcordatâ linear-lanceolatis, superioribus et ramulorum alternis diminutis omnibus subtus ad nervos et margine adpresse hirtis, floribus masculis pentandris, perigonio fructifero late bialato ♀. *Mem. pentandra* Wedd. in D.C. Prodr. — *Pouzolzia pentandra* Benn. Pl. Javan. tab. 14. — *Urtica hippuroides* Griff. MSS.

Hab. ad fluvium Hussun Abdul Affghaniæ (Griff!). Planta subtropica vix ditionis.

Ar. Geogr. India Orientalis.

DEBREGEASIA (Gaudich. Bonit. tab. 90).

Flores monoici glomerulati vel capitulati bracteis scariosis stipati axillares. *Masc.* Perigonium quadripartitum segmentis ovatis breviter acuminatis. Stamina 4 laciñis opposita. *Fem.* Perigonium ventricos-tubulosum ore contracto quadridentato. Ovarium perigonio adnatum. Stigma sessile penicillatum. Fructus bacciformis e perigonio et pericarpi parte exteriore constans intus mucimentaceus. — Frutices inermes foliis alternis rugosis.

I. D. bicolor (Roxb. Ind. III, p. 589 sub *Urticā*) fruticosa ramulis elongatis junioribus canescensibus, foliis breviter petiolatis a basi cuneatâ lanceolatis acuminatis supra viridibus rugosis asperis subtus adpresse niveis, capitulis unisexualibus vel androgynis in axillis 1-4 sessilibus vel breviter pedunculatis 3. *D. bicolor* et *D. hypoleuca* Wedd. Urt. tab. 15. — *Boehmeria hypoleuca* Hochst.

Hab. in humidis ad Otipore Affghaniæ (Griff! Journ. № 47)

Ar. Geogr. Abyssinia, India.

PARIETARIA (L. Gen. 1452).

Flores polygami fasciculati vel cymosi bracteati. *Hermapl.* Perigonum laciniis 4 subæqualibus basi coalitis sub anthesi patentibus concavis, fructiferum plus minus aceretum. Stamina 4 libera basi perigonii inserta et ejus laciniis opposita, filamenta transverse plicata et inflexa sub anthesi elastice reflexa. Ovarium ellipticum. Stigma sessile penicillatum. *Fem.* Perigonum tubulosum apice quadridentatum persistens non accrescens, stylus exsertus, stigma penicillatum. Achenium ovatum subcompressum perigonio persistente tunicalatum. Herbæ perennes vel annuæ inermes, foliis alternis.

1. **P. officinalis** (L. Sp. 1492) perennis pubescens, caulis herbaceis erectis subsimplicibus, foliis longe petiolatis oblongo-lanceolatis acuminatis basi longe attenuatis triplinerviis integris, cymis axillaribus genuinatis glomeruliformibus densis multifloris, bracteis liberis non decurrentibus floribus subbrevioribus fructiferis immutatis, floribus dichotomiarum femineis terminalibus hermaphroditis perigonio campanulato demum vix elongato et stamna æquante ♀. *P. erecta* M. K. Deutsch. Fl. I. p. 825. — Ic. Rehb. Germ. fig. 1317.

Hab. in ruderatis, ad muros totius ditionis praesertim in parte magis septentrionali a Thraciâ (Friv! Auch!) ad regiones Caucasicas (Ledeb.), Cretam (Sieb!), Persiam bor. (Bge!).

Ar. Geogr. Europa media et australis ab Angliâ et Sueciâ meridionali ad Rossiam medium, Sibiria Uralensis.

2. **P. Judaica** (Linn. Sp. p. 1492) perennis pubescens, caulis e rhizomate indurato interdum suffrutescente diffusis vel ascendentibus simplicibus vel ramosis, foliis longe petiolatis a basi cuneatâ ovatis acutis triplinerviis integris, cymis axillaribus subsessilibus glomeruliformibus genuinatis, bracteis basi connatis secus axim decurrentibus flore subbrevioribus fructiferis immutatis, floribus dichotomiarum femineis cæteris hermaphroditis perigonio demum elongato stamna excedente ♀. Ic. Rehb. Germ. fig. 1318. — *P. officinalis* L. ex parte et Smith Brit. 189. — *P. diffusa* M. K. Deutsch. Fl. I. p. 827. — *P. punctata* Willd. Sp. IV. p. 953. — *P. elliptica*, *P. cryptarum* et *P. nitens* C. Koch Linn. XXII. p. 606.

Hab. ad muros et in fissuris rupium totius ditionis a Græciâ! Macedonia! et Thraciâ! ad Caucasum! et Transcaucasiam! Syriam littoralem et in Libano! Palestinam ad mœnia Hierosolymæ (Boiss!), Egyptum (Gaill!), montes Persiae australis (Ky. 2571 5101) et borealis (Buhse! Bge!), Turkestania (Lehm!), monte Chehel Tun Belutschiae (Stocks!).

β. brevipetiolata — Petioli breves. — *P. multicaulis* Boiss. et Heldr. Diagn. Ser. I, 12, p. 106. Huc speetat specimen *P. Judaica* herb. Linnæani!

Hab. in faucibus montium tophaceorum Pisidiæ prope Buldur (Heldr!), circa Ermenek Ciliciae Tracheæ (Per!), circa Orfa, Aintab, Marasch Syriæ borealis (Auch. 2794! Haussk!), in monte Gara Kurdistaniæ (Ky. 329!).

Species valde variabilis a *P. officinali* forsan non specifice distincta. In australioribus montanis et alpinis caules humiliores basi magis suffrutescentes fiunt, folia minora.

Ar. Geogr. Europa media et australis a Britanniâ, Belgio, Germaniâ media ad Dalmatiâ et Transylvaniâ, Africa borealis.

3. **P. Cretica** (L. Sp. p. 1492) perennis pubescens multicaulis, caulis diffusis simplicibus et ramulosis, foliis parvis longe petiolatis a basi rotundato-cuneatâ ovatis integris obscure triplineroviis, cymis axillaribus sessilibus trifloris flore intermedio femineo lateralibus hermaphroditis, floris feminei bracteâ linearî, floribus hermaphroditis vibranteatis, bracteis basi coalitis fructiferis valde auctis linearispathulatis inæqualibus involucrum rigidulum fuscum quinquepartitum formantibus ♀. Ic. Sibth. Fl. Græc. II, p. 44, tab. 154. — Wedd. Urt. tab. XVII, fig. 16 et 17.

Hab. in rupestribus regionis inferioris maritimæ, Creta ad Cydoniam (Sibth., Heldr!), Cimolus (Sibth.), Thera (Sartori!), Laconia prope Malta pagum (Heldr!), Attica ad Aeropolin Athenarum (Pichl!) et ejus insulæ (Sart! Heldr!), in insulis Pharmacus (Heldr!), insula Ponticonisi Eubeæ (Wild!).

4. **P. Lusitanica** (L. Sp. 1492) annua puberula, caulis tiliiformibus prostratis ramosis, foliis petiolatis ovatis utrinque attenuatis integris ex imâ basi trinerviis, cymis axillaribus subsessilibus glomeruliformibus, bracteis basi tantum coalitis oblongo-lanceolatis aequalibus post anthesin subimmutatis perigonia fructifera æquantibus, floribus hermaphroditis paucis sterilibus cæteris femineis, perigonio fructifero vix ancto rufescente indurato laciniis ovatis exacte conniventibus, achenii elliptici apiculo recto ⊖. Ic. Bocc. Sic. tab. 24, fig. B. — *P. parviflora* Frieder. et Blume Mus. Bot. Lugd. II, p. 249.

Hab. ad muros et rupes umbrosas Peloponnesi (Bory!), Atticæ (Heldr! Orph. Fl. exs. 303!), Zacynthi (Marg!), insulæ Tassos Macedoniæ (Griseb.), Chios (Orph!), Rhodus (Bourg!), Byzantii (Noël), circa Smyrnæ (Fleisch! Boiss!), Tauro Cilicio (Ky. 41 sub *P. micrantha*), Ciliciâ littorali (Bal!), Palestinâ (Boiss! Roth!), Mesopotamiâ in monte Sindjar (Haussk!), Tauriâ (Stev.), Caucaso et Transcaucasiâ (Ledeb.), Ossetiâ (Rupr!), Georgia Cauca-sicâ (Hohen!).

Ar. Geogr. Lusitania, Hispania, Gallia et Italia australes, Corsica, Sardinia, Serbia, Banatus, Rossia australis, Africa borealis.

5. **P. debillis** (Forst. Prodr. N° 387) annua puberula, caulis diffusis filiformibus flaccidis, foliis longe petiolatis ovatis vel cordatis integris ex imâ basi trinerviis, cymis axillaribus paucifloris breviter pedunculatis, bracteis linearibus vel lanceolatis post anthesin immutatis, floribus hermaphroditis paucis cæteris femine's, perigonii laciniis acuminatis fructiferis auctis patentibus, achenio oblique apicu-

Iato ⊙. Ic. Wedd. Urt. tab. XVII, fig. 24-25. — *P. micrantha* Ledeb. Fl. Alt. IV, p. 303. — Ic. Ross. t. 22.

Hab. in Belutschia (Stocks ex Wedd.). Editione nondum vidi.

Species *P. Lusitanicae* proxime affinis et ad quam præter synonymon indicatum Cl. Weddell plures alias conjungit.

Ar. Geogr. Sibiria, Abyssinia, regio Himalaica, Nova Hollandia et etiam (si *P. appendiculata* et aliae huic recte conjunguntur), insulae Canarienses, Madera, America tropica.

6. **P. alsinefolia** (Delile Fl. Eg. p. 137, tab. 50, fig. 2) annua puberula, caulibus filiformibus flaccidis diffusis, foliis parvis longe et tenuiter petiolatis a basi rotundata ovatis, cymis axillaribus trifloris inferiore hermaphrodito sterili bracteâ parvâ lanceolatâ sufluto cæteris fæmineis bracteis ternis demum valde auctis membranaceis cordato-ovatis suffultis, perigonio fructifero parum aucto ovato segmentis acutis conniventibus ⊙. *Freirea alsinefolia* Gaudich.

Hab. in umbrosis rupium Egypti inferioris et superioris in desertis (Del., Schweinf!), Arabiae petreæ et Palestinae ad Hebron, Hierosolymam etc. (Schimp! Boiss!), monte Gebel Khaisun ad Damascum (Gaill!), deserto ad Palmyram (Bl!), circa Aleppo (Ky. 1801), Persia australi (Haussk!) et orientali (Bgel), Belutschia (Stocks!).

FORSKAHLEA (L. Mant. p. 11. — *Caidbeja* Forsk. Eg. p. 82).

Flores monoici in involucris 4-6-phyllois intus dense lanatis numerosi masculi in circuitu, fæminei interiores; in involucris diphyllois rarioribus tantum fæminei. *Masc.* Perigonium monophyllum basi tubulosum superne dilatatum unilabiatum subtridentatum. Stamen unicum. *Fem.* Perigonium nullum. Ovarium ellipticum lanatum in stigma filiforme hispidum attenuatum. Achenium rectum lanatum compressum, albumen parcum. — Herbae vel suffrutices perennes foliis alternis, stipulis lateralibus liberis.

1. **F. tenacissima** (L. loc. cit.) perennis a collo saepè ligneo-scenti ramosa, ramis diffusis simplicibus vel ramulosis præter lanam adpressam stimulosis, foliis obovato-orbiculatis vel rhombatis breviter petiolatis supra basin cuneatam grosse crenatis vel serratis superne viridibus pilis rectis et aliis majoribus hamatis hispidis subtus præter nervos hispidos adpresso tomentoso-canis, involucris axillaribus globoso-turbinatis subsessilibus-4-nis et solitariis aliis 4-5-phyllois aliis rarioribus diphyllois foliolis lanceolatis acutis sericeo-hispidis ♀. Ic. Lam. III. tab. 388. — *Caidbeja adhærens* Forsk. loc. cit.

Hab. in rupestribus desertorum Egypti mediae et superioris! Arabiae petreæ (Schimp. 230! Boiss!), Persiae australis (Auch. 5315!), Affghaniæ (Griff!), Belutschiae (Stocks!).

Ar. Geogr. Hispania austro-orientalis, Africa borealis interior, Arabia tropica, India boreali-occidentalis.

TRIB. II. CANNABINEÆ.

Ovarium uniloculare ovulo pendulo. Achenium bivalve indehiscentia perigonio velatum. Embryo falcatus vel spiralis. Albumen nullum. — Herbæ.

HUMULUS (L. Gen. 446).

Flores dioici. *Masc.* Flores racemosi. Perigonum ad basin quinque-partitum. Stamina quinque basi laciniarum perigonii inserta erecta filamento brevissimo, antheris oblongis bilocularibus apiculatis. *Fem.* Flores amentacei ad basin bracteæ foliaceæ bini. Perigonum monophyllum apertum ovarium uniovulatum amplectens, stigmata bina subulata elongata pubescentia. Strobilus ovatus bracteis demum auctis foliaceis. Nucula subglobosa erecta pericarpio duro fragili. Embryo spiralis. — Herbæ perennes volubiles, foliis oppositis.

I. H. Lupulus (L. Sp. 1457) caulis elongatis sarmentosis volubilibus, foliis petiolatis superne retrorsum scabris cordatis palmatilobatis lobis 3-5 ovatis acuminatis serrulatis, foliis superioribus indivisis, racemis masculis oppositis axillaribus pedunculatis compotitis laxis, bracteis breviter lanceolatis, perigonii laciinis oblongis, stamentis femineis oppositis pedunculatis demum in strobilos ovatos abeuntibus, bracteis valde anctis scariosis extus glandulis luteis basi obsoitis ♀. Ic. Rehb. Germ. tab. 656. — Nees Gener. Germ.

Haec ad sepes et dumeta regionis inferioris et montanæ, Peloponnesus (Orph!), Thracia (Friv.), Bithynia (Sibth.), Pontus Lazicus ad Rhizé (Bal!), Tauria, Caucasus et Transcaucasia (Ledeb.), Syria circa Damascum (Orph!). Et prob. alibi.

Ar. Geogr. Europa fere omnis a Scandinaviâ ad Rossiam medium, Sibiria

CANNABIS (Tourn. Just. p. 555).

Flores dioici. *Masc.* Flores racemosi. Perigonum ad basin quinque-partitum. Stamina quinque basi perigonii laciniarum inserta, filamento nutante, antheris pendulis bilocularibus. *Fem.* Flores spicato-globosati. Perigonii loco bractea parva complicata persistens basi ventricosa ovarium globosum uniovulatum includens. Stigmata bina elongata filiformia pubescentia basi connata. Nucula ovata erecta pericarpio duro fragili subbivalvi sed subindehiscenti. Embryo falcato-plicatus. — Herbæ annuae inflorescentiæ glutinosâ.

I. C. sativa (L. Sp. 1457) annua, foliis oppositis petiolatis pinnatisectis segmentis 5-7 lanceolatis acuminatis serratis, racemis

masculis longe paniculatis, glomerulis femineis subsessilibus spicam terminalem formantibus ⊖. Ic. Nees Gen. Germ. — Rehb. t. 655. — *C. Orientalis*, *C. Indica*, *C. Chineensis* Auct.

Hab. ubique culta et in Persia austro-occid. aliisque locis subsppontanea (Haussk!), ex Sibiri vel regione Himalaiae verosimiliter oriunda.

TRIB. III. MOREÆ.

Ovarium 1-2-localare. Ovulum pendulum campylotropum. Semen pendulum. Albumen carnosum embryonem curvatum involvens. Fructus spurius carnosus e receptaculo vel e receptaculo et perigonii formatus. — Arbores.

MORUS (L. Gen. 4055).

Flores monoici raro dioici in spicas vel capitula unisexualia dispositi. *Masc.* Perigonium 4-partitum. Stamina 4 filamentis pefloratione inflexis. *Fem.* Perigonium 4-5-sepalum sepalis ovatis concavis mox succulentis ovarium amplexantibus. Ovarium biloculare. Stigmata bina. Fructus (syncarpium) ex utriculis capitato-spicatis abortu monospermis perigonio baccato inclusis inter se coalitis formatus. Seminis testa dura fragilis. Embryo uncinatus. — Arbores lactiflui foliis alternis.

1. **M. nigra** (L. Sp. 1398), foliis cordato-ovatis serrato-dentatis lobatis vel indivisis, spicis femineis subsessilibus pedunculo multo-ties longioribus, fructu eduli, perigonii laciiniis stigmatibusque margine villigeris ♂.

Hab. ubique culta, in sylvis districtus Transcaucasici Talysh ad Lenkoran (Hohen.) et Persiae borealis ad Caspium probab. spontanea.

Ar. Geogr. Ubique in Europæ mediâ et australi culta et hinc inde subsppontanea.

2. **M. alba** (L. Sp. 1398) foliis cordato-ovatis basi saepius inæquilibus integris serratis vel lobatis, pedunculis spicis femineas æquantes, perigonii laciiniis margine glabris, stigmatibus glabris breviter papillosis, fructu insapido ♂. *M. Indica* L. p. 1399.

Variat multum; fructus albi, vel nigri vel rubelli. Ex Chinâ oriunda dicitur.

Ar. Geogr. Ubique in regione mediterraneâ Europæ et Africæ borealis culta.

FICUS (L. Gen. p. 4486. — *Urostigma* et *Sycomorus* Gasparr.).

Flores monoici vel dioici parieti interno receptaculi carnosí cavi pyriformi-globosi basi bracteati apice umbilicati et ore angusto squa-

muloſo hiantis inserti. *Masc.* Perigonium 3-5-sepalum. Stamina 3-6 filamentis præfloratione rectis. *Fem.* Perigonium quinquefidum. Ovarium uniloculare, stylus lateralis, stigmata bina vel unicu[m]. Fructus receptaculo aucto carnoso utriculisque minimis crustaceis monospermis constans. Seminis testa membranacea. Embryo incurvus. — Arborcs lactifluæ foliis alternis.

1. F. Carica (L. Sp. 1513) arborea vel fruticosa, foliis deciduis supra scabris subtus pubescentibus vel velutinis ovatis basi cordatis 3-5-lobis, repandis, rarius indivisis, receptaculis ad ramos hornotinos solitariis axillaribus vel supra axillas ortis breviter pedunculatis pyriformibus vel globosis in stipitem eis breviorem attenuatis vel estipitatis 5. Species ex cl. Haussknecht observationibus in locis natalibus valde variabilis et ejus formis magis notabiles numero.

α. genuina. — Folia majuscula 3-5-loba rarius indivisa, receptaculi glabri pyriformes in stipitem attenuati.

Hab. in fissuris rupium regionis inferioris et montanæ, Græcia et Archipelagus! Macedonia et Thracia (Griseb!), Anatolia occidentalis et meridionalis (Tchih! Ball Kyl), Syria borealis et Mesopotamia ad Aintab, Orfa, etc. (Haussk!), Pontus Lazicus ad ostium fluvii Kalopotanios (Bal!), Tauria et Transcaucasia (Lebed!).

β. riparum Haussk. MSS. — Folia in lacinias 5-7 inferne angustatas spathulatas partita.

Hab. secus rivulos Syriae borealis prope Aintab, ad Giaurgöl prope Marasch (Haussk!).

γ. rupestris Haussk. MSS. — Folia indivisa ovata vel oblonga obtusa firmiora. Fructus pyriformes plus minus puberuli.

Hab. in fissuris rupium, faux Guzeldere Ciliciæ (Bal!), pagus Maaden Armeniæ Kurdiæ (Ky. 486!), Akkerdag prope Marasch et Aintab (Haussk!), Mesopotamia ad Orfa, Tcharmelik et in monte Sindjar (Haussk!), montes Kuh Kilouyeh Persie austro-occidentalis (Haussk!).

δ. globosa. — Folia triloba vel indivisa, receptacula globosa puberula stipite omnino destituta.

Hab. in fissuris rupium Cataoniæ in montibus Akdag et Begdag, circa Marasch, in deserto Mesopotamie inter Surug et Orfa (Haussk!).

ε. Johannis. — Folia parva supra aspera subtus tomentello-scabra profunde lobato-partita vel laciniata lobis et laciniis lobulatis undulato-crispis, fructus parvi velutini ovato-pyriformes vel globosi breviter stipitati. *F. Johannis* Boiss. Diagn. Ser. I, 7, p. 96.

Hab. ad rupes Persie australis, ad Gere et Schiraz (Ky. 91! et 316!) ad Teng Biresa prope Bebehan et Nackschi Timur prope Kaserun (Haussk!), Persie orientali inter Tebbes et Herat (Bge!), Belutschia ad Zahri (Stocksl!).

Ar. Geogr. Europa australis, Africa borealis, India septentrionalis.

2. F. Persica (Boiss. Diagn. Ser. I, 7, p. 97) fruticosa, ramis elongatis cinerascentibus, foliis deciduis ovato-orbiculatis basi rotundatis vel subcordatis ultra medium in lacinias spathulatas iterum

obtuse lobatas divisis superne pruinosis inferne præter nervos parce hirtulos glabris, receptaculis supra axillas solitariis subsessilibus glabris ovato-pyriformibus stipite eis longiore suffultis ♂.

Hab. in Persiâ austro-occidentali prope Dalechi et in rupestribus prope Gere (Ky. 91!).

Frutex 12-pedalis, folia 2-pollices circiter longa et lata, fructus maturus avellanâ major. An hæc species quoque formis *F. Caricæ* adnumeranda? sed formas intermedias nondum vidi. Ab eâ differt foliis glabriusculis et fructu stipite pollicem longo suffulto.

3. F. virgata (Roxb. Fl. Ind. III. p. 530) fruticosa, ramis virgatis, foliis petiolatis basi rotundatis vel late subcordatis ovatis obtusis vel acutiusculis serrato-crenatis subtus tomentosis, receptaculis supra axillas solitariis pedunculatis ovato-clavatis pubescentibus parvis ♂. Ie. Wight Icon. tab. 649.

Hab. ad rupes Belutschie (Stocks 609!), Affghaniæ (Griff.).

A formis integrifoliis *F. Caricæ* distincta videtur fructibus parvis ex cl. Roxburghi *Piso* vix majoribus. In Aitches. Punjab Catal. p. 138 est synonymon *F. Caricoidis* Roxb. quæ tamen ex Icone Wight tab. 634 diversissima videtur.

Ar. Geogr. India borealis.

4. F. pseudosycomorus (Decaisn. Flor. Sinaic. p. 4) arborea humilis, ramis junioribus pruinosis. foliis petiolatis ovatis vel ovato-orbiculatis profunde cordatis obtusis obtuse crenato-lobulatis aliis indivisis vel repandis aliis trilobis supra pallide virentibus scabris subtus pallidis tomentellis dein glabratis scabridis, involueris extra-axillaribus solitariis brevissime pedunculatis vel sessilibus turbinatis puberulis ♂. *F. palmata* Schweinf. exs. non Forsk. — *F. morifolia* Husson exs.

Hab. in Arabiâ Sinaiticâ (Bové 204!), in desertis Ægyptiaco-Arabicis ad Gebel Numreh (Husson!), ad Ouadi el Deyr (Figari!), ad Ouadi Dachel et Ouadi Natte (Schw!).

A formis *F. Caricæ* foliis multo profundius cordatis supra scaberrimis, fructibus subsessilibus distincta videtur. Fructus avellanâ majores. *F. palmata* Forsk. ex descriptione et Icone Vahl Symb. tab. 24 et specimine ex monte Maammera Arabiæ tropicæ (Botta 1837) longe differt foliis in lacinias elongatas acutas vel acuminatas partitis.

5. F. Sycomorus (L. Hort. Cliff. p. 471) arborea, foliis persistentibus petiolatis ovatis subcordatis obtusis integris præter nervos subtus elevatos et parce hirtulos glabris levibus, receptaculis in racemos tortuosos aphyllos e trunco vel ramis vetustis oriundos dispositis turbinatis breviter stipitatis tomentellis ♂. *F. Cypria* Rauw. It. 287, tab. 57. — *Sycomorus antiquorum* Gasparini.

Hab. in Egypti ambulacris (Del., Boiss! Ky. 405! Schweinf!).

Arbor ramis horizontalibus in comam latissimam extensis.

Ar. Geogr. Nubia, Abyssinia.

Obs. *Ficus elastica* L. et *F. religiosa* L. in Egypto et Belutschia frequenter coluntur. Cl. Haussknecht legit in insula Karak Sinus Persici *Ficum* 3-pedes

altam foliis coriaceis ovatis integerrimis, an spontaneam? an cultam? ex specimine valde incompleto ad *F. ramentaceam* Wight Ic. 657 vel ad *F. laccifera* W. Ic. 636 forsan referendam.

TR. IV. CELTIDEÆ.

Ovarium uniloculare uniovulum. Ovulum pendulum campylotropum. Fructus verus monopyrenus carnosus. Semen pendulum. Embryo curvatus cotylonibus conduplicatis. Album centrale parcum. — Arbores.

CELTIS (L. Gen. 1143).

Flores hermaphroditi et abortu polygami. Perigonii laciniæ 5-6 basi plus minus coalitæ ovatæ concavæ æst. imbricatæ. Stamina 5-6 laciniis opposita æstiv. incurva dein elastice erecta. Antheræ introrsæ. Stigmata bina elongata papillosa. Drupa ovata parum carnosa putamine osseo. — Arbores succo aqueo, foliis alternis, stipulis fugacissimis, floribus solitariis axillaribus pedunculatis.

1. **C. australis** (L. Sp. 1478) arborea, foliis a basi obliquâ rotundatâ vel cuneatâ ovato-lanceolatis longe acuminatis argute serratis subtus triplinerviis molliter pubescentibus, pedunculis fructiferis petiolo pluries longioribus, drupâ globosâ nigrâ nucleo reticulatourgoso ♂. Ic. Duham Arb. tab. 53.

Hab. in regione inferiori et præsertim montanâ, Grecia in Pentelico et Parnasso (Heldr!), Creta (Raul!), Macedonia (Orph!), Thracia (Griseb.), Anatolia borealis! Pontus Lazicus (Ball!), Imeretia ad Suckum Kalé (Ruprl!), Syria et Palestina (Boiss!), prov. Caucasica (ex Ledeb.), Persia borealis ad Asterabad (Karel.). Arbor saepè ob fructum sapidum culta et ideo forsan non ubique spontanea.

Ar. Geogr. Europa australis, Africa borealis.

2. **C. Caucasică** (Willd. Sp. 4, p. 994) arborea, foliis a basi obliquâ breviter cuneatâ ovato-rhombeis acuminatis simpliciter vel duplicitam serratis subtus triplinerviis junioribus subtus puberulis adultis glabratris, pedunculis fructiferis petiolo 2-3-plo longioribus, drupâ globosâ flavâ nucleo leviter reticulato basi non acuto ♂.

Hab. in Armeniâ Rossicâ et Iberiâ (Hohen! Szov!), Persia boreali (Auch. 5323! Ky. 126! et 633! Buhse!), montibus prov. Khorassan (Bge!), Persia austro-occid. (Ky. 512! Haussk!), regno Cabulico (Griff!), Belutschia (Stocks!).

Species *C. australi* et *C. Tournefortii* fere intermedia sed priori magis affinis et forsan hujus varietas Orientalis; differt foliis latioribus et brevioribus minus longe acuminatis et basi cuneatis subtus puberulis vel glabratris nec velutinis. A *C. Tournefortii* distinguitur foliis acuminatis majoribus, fructus nucleo non quadricostato nec basi acuto, trunco elatiore.

3. C. Tournefortii (Lam. Eneyel. 4, p. 138) arborea humiliis, foliis subcoriaceis a basi inaequali rotundata vel subcordata breviter rhomboeovatis acutis vel subacuminatis crenatis vel minute serratis glaucescentibus glabriusculis laevibus vel supra asperulis, pedunculis fructiferis petiolo subduplo longioribus, drupæ flavæ obovato-turbinata nucleo quadricarinato cæterum laeviusculo vel parce ruguloso basi acuto 3. Ic. Janb. et Sp. Ill. Or. tab. 400. — *Celtis Orientalis minor foliis minoribus et crassioribus, fructu flavo* Tourn. Voy. II, tab. 425. — *C. aspera* Steven Taur. p. 309.

Hab. in rupestribus Graecie in monte Kyllene (Orph. Fl. Gr. 204!), prope Trikala (Heldr!), Anatoliæ bor. in Galatiâ (Tourn.), Cappadociâ ad Cæsaream (Bal!), Cataoniâ (Haussk!), Mesopotamiâ ad Orfa (Haussk!), Assyriâ (Ky. 314!), Tauriâ (Stev.).

β. glabrata. — Folia glaberrima basi subcuneata nec cordata. — *C. glabrata* et *C. Kotschyana* Stev. Taur. p. 310 et 311.

Hab. in Lyciâ (Bourg. 250!), Ciliciâ ad Gülek (Bal!), Tauriâ (Stev.), Iberia Caucasicâ (Hoh. sub *C. Caucasicâ*), monte Gara Kurdistaniæ (Ky. 393!).

Frutex vel arbuscula *Pruno* non elatior, folia eis præcedentium crassiora magis coriacea vix vel breviter acutata.

Ar. Geogr. Siciliæ montes.

TR. V. ULMEÆ.

Ovarium dimerum sed abortu normali sœpe uniloculare. Ovulum pendulum anatropum. Fructus siccus unilocularis samaroideus vel utricularis. Semen pendulum. Embryo rectus. Albumen nullum. — Arbores et frutices.

ULMUS (L. Gen. 316).

Flores hermaphroditæ. Perigonum campanulatum 4-5-dentalum marcescens. Stamina 5-8 basi perigonii inserta filamentis præfloratione rectis. Ovarium biostre stigmatis binis elongatis intus villosis. Fructus samaroidens ntriculo circumcircâ alâ membranaceâ cincto. — Arbores foliis basi inaequalibus, stipulis caducis, floribus præco-cibus in fasciculos laterales dispositis.

1. U. campestris (L. Sp. 327) arborea, foliis ovato vel lanceo-lato-oblongis breviter acuminatis basi inaequalibus serratis, pubescen-tibus vel glabris, florum fasciculis subsessilibus pedicellis brevissimis, perigonii æqualiter 4-5-fidi laciñiis ciliatis, staminibus 4-5, samarâ obovatâ basi attenuatâ margine nudâ apice emarginatâ, semine sinu terminali contiguo 3. Ic. Nees. Gen. Germ. — *U. nuda* Ehrh. — *U. glabra* Mill. Engl. bot. tab. 2248. — *U. suberosa* Ehrh. (forma

minor microphylla ramorum cortice suberoso-alatâ). — *U. elliptica* C. Koch Linn. XXII, p. 599.

Hab. in campis ad rivos, sylvis regionis inferioris et præsertim montanæ, Græcia! Macedonia et Thracia (Griseb!), Bithynia (Schmidt!), in Anatoliâ boreali et Ponto frequenter (Tchih.), Armenia (Ky!), Syriâ ad Aleppo (Ky. 23!), Tauriâ, Caucaso et Transcaucasiâ (Ledeb.), Turkestanâ (Lehm!).

β. microphylla. — Folia multo minora (9-10 lin. longa) obtuse crenata subtus ad venus puberula. Samara minor (4 lineas longa). Facies *U. pumila* sed characteres *U. campestris*.

Hab. in Persiâ orientali (Bge!).

Ar. Geogr. Europa tota arcticâ exceptâ, Africa borealis, Sibiria, regio Himalaica.

2. *U. montana* (Sm. Engl. Fl. II, p. 22) arborea, foliis majusculis lanceolato-oblongis abrupte et longiusculæ acuminatis basi inæqualibus duplicatim serratis supra seberrimis, florum fasciculis subsessilibus pedicellis brevissimis, perigonii æqualiter 4-5-fidi lacinias ciliatis, staminibus 5-6, samarâ ovato-orbiculatâ margine nudâ apice emarginatâ, semine infra medium sito a sinn remoto 5. Engl. Bot. tab. 1887. — *U. major* Smith.

Hab. in sylvaticis montanis et alpinis, Peloponnesus (Bory), Anatolia borealis (Auch. 5321!), Pontus Lazicus ad Khabackar 5500' (Bal!), Cilicia in regione Abietinâ 4500'-5000' (Ky. 334! et 441!), Caucasus orientalis in Tuschetia 6000' (Rupr!).

Ar. Geogr. Europa tota arcticâ exceptâ, in australioribus montana, Africa borealis, Sibiria.

3. *U. pedunculata* (Foug. Mém. Ac. Sc. Paris 1787, tab. 2) arborea, foliis ovatis acuminatis basi inæqualibus duplicatim et argute serratis subtus pubescens, floribus longe pedicellatis laxe corymbo-fasciculatis pendulis, perigonii ore obliquo 5-8-lobo lobis anticis minoribus, staminibus sub 8, samaræ ovatae glabrae sed margine longe et molliter ciliatae emarginaturâ terminali semini subcontigua 5. *U. effusa* Willd. Prodr. Ber. 296. — *U. ciliata* Ehrh. Beitr. 6, 88. — *U. octandra* Schk. tab. 57.

Hab. in sylvis montanis Ponti Lazici (C. Koch), Tauriae (Stev.), Caucasi (M.B.). E ditione non vidi.

Ar. Geogr. Svecia meridionalis, Dania, Gallia, Germania, regio Danubialis, Rossia.

ZELKOUA (Spach Ann. Sc. Nat. 2 Ser. XV, p. 336. — *Abelicea* Endl. ex parte.).

Flores monoico-polygami, masculi in parte inferiore aphyllâ ramulorum annotinorum dense glomerati, hermaphroditi ad axillas foliorum juniorum solitarii. Perigonium campanulatum subseariosum

profunde 4-5-lobum. Discus in floribus fertilibus cupularis carnosus. Stamina 4-5 perigonio ad marginem disci inserta filamentis præfloratione rectis. Ovarium oblique ovatum stylis 2 dentiformibus intus papillosis coronatum. Nucula exsueca aptera gibboso-ovata crustacea rugoso-nervosa. — Arhores et arbusculæ foliis et stipulis caducis facie *Carpini*. Genus a *Planera* nuculâ non lamellato-crystatâ vix distinctum.

1. **Z. crenata** (Desf. Cat. Par. sub *Planera*) arborea. ramulis fertilibus elongatis, foliis oblongis basi subinæqualibus et brevissime petiolatis acutis acutinuscule crenato-serratis subtus concoloribus parce puberulis, floribus albidois inodoris, nuculis glabrescentibus 5. *Z. crenata* Spach loc. cit. — *Ulmus polygama* Rich. — *Rhamnus carpinifolia* Pall. Ross. II, p. 24, tab. 10. — *Planera Richardi* Michaux Fl. Bor. Am. II, p. 248.

Hab. sylvas in regione inferiore et montanâ Transcaucasiae formans, Colchis (Radde!), Imeretia (Parrot ex Led.), Iberia inter Gori et Uplis Ziche (Stev.), prov. Karabagh (Szovt.), prov. Talysh circa Lenkoran usque ad 5000' (C.A. Meyl Hoh!), totius Persiae septentrionalis ad Caspium (Auch, 5322! Buhse!), usque ad Asterabad (Karel!). Tatarice *Asadachatsch* et *Nilachatsch*.

2. **Z. Abelicea** (Sibth. et Sm. Prodr. I, p. 172 sub *Ulmo*) arbuscula, ramulis fertilibus abbreviatis, foliis parvis firmis subsessilibus ovalo-oblongis obtusis obtuse crenatis subtus albidois et pubescentibus, floribus albis suaveolentibus, nuculis pubescentibus 5. *Z. Cretica* Spach loc. cit. — *Quercus Abelicea* Lam. — *Planera Abelicea* R. et Sch. — *Pseudosantalum Creticum* Bauh. Pin. p. 393. — Poma Baldo 112 Ic. — Parkins. Th. Ic.

Hab. in declivibus saxosis montanis et subalpinis Cretæ 3500'-4500' (Belli, Tomnef, Sibth.), in montibus Sphacioticis circa Omalo (Heldr!). Fl. Martio. Incolis *Apchitza*.

Arbuscula 10-15-pedalis, folia eis præcedentis firmiora minora 7-12 lin. longa, nuculæ subminores.

ORD. CXV. JUGLANDACEÆ.

(Lindl. Nat. Syst. p. 427.)

Flores monoici masculi amentacei feminei aggregati spicati vel solitarii. *Masc.* Perigonum simplex bracteæ intus adnatum 6 vel abortu 2-3-lobum rarius nullum. Stamina numerosa in medio perigonii inserta filamentis brevissimis, antheris bilocularibus, connectivo siepe producto. *Fem.* Perigonum modo simplex cum ovario connatum, modo duplex externum involucriforme cum ovario plus minus connatum cupuliforme apice pluridentatum, interius cum ovario connatum apice quadridentatum. Ovarium primum uniloculare dein

dissepimentis 2-4 incomplete basi et apice 2-4-loculare. Stylus brevis. Stigmata sœpius bina. Placentarium centrale tandem cum dissepimentis connatum ovulum orthotropum erectum gerens. Drupa carnosa vel membranacea, pericarpio a perigonio formato, nuce ex ovario indurato ortâ basi et apice incomplete 2-4-loculari. Semen exalbuminosum. Embryo cotyledonibus carnosis bifidis gyroso-plicatis, radiculâ superâ, plumulâ intra cotyledones abditâ squamulis imbricatis sœpe instructâ. — Arbores foliis alternis compositis — Ordo juxta *Cupuliferas* et *Amentaceas* ex antiquâ normâ collocatum sed *Terebinthaceis* et *Myricaceis* magis affinis.

JUGLANS (L. Gen. 107).

Amenta mascula solitaria in parte inferiori ramulorum hornotinorum pendula. Perigonium 5-6-lobum. Stamina 2-multiseriata connectivo supra antheras producto. Flores feminei terminales solitarii vel 2-3 ni glomerati. Perigonium duplex cum ovario usque ad stylum connatum. Pericarpium carnosum indehiscens a nuce maturitate liberum irregulariter rumpens. Nux irregulariter rugosa in germinatione bivalvis — Arbores foliis imparipinnatis.

I. **J. regia** (L. Sp. 1415) arborea, foliis glabris 5-6-(raro 2)-jugis, foliolis ovatis vel oblongis integris apice attenuatis, nuce ovatâ vel globosâ reticulato-rugosâ dissepimentis qualuor, embryonis squamulis distictis 5. Nees Ic. Germ. Ic.

Hab. in sylvis regionis inferioris et montane Ponti et Armeniæ Turcicæ (Tchih.), Transcaucasiae in Imeretiâ (Eichw.), Georgia et prov. Karabagh (Hoh., Szov!), ditione Talysh (Hohen!), in Persia boreali (Buhsel!), Belutschia (Stocks!). Persice *Girdagan* vel *Wuhs*.

Ar. Geogr. India borealis.

PTEROCARYA (Kunth in Ann. Sc. Nat. 1824, p. 13).

Flores masculi dense amentacei. Perigonium 5-6-lobum. Stamina 9-10 biseriata, antherarum connectivum non productum. Flores feminei in spicas longas laxas dispositi. Perigonium duplex, exteriorus basi cupuliforme cum ovario connatum superne liberum lateraliter dipterum, interiorus basi cum ovario connatum tubiforme apice quadridentatum. Fructus lateraliter bialatus, pericarpium indehiscens scarioso-membranaceum, nux ossea laevis indehiscens. — Arbores foliis imparipinnatis.

I. **P. fraxinifolia** (Lam. Dict. IV, p. 502 sub *Juglante*) arborea, foliis sub-12-jugis, rachide glabrâ, foliolis sessilibus oblongis et oblongo-ellipticis acutis minute serratis subtus ad nervos pilosulis,

amentis masculis sessilibus ad ramorum extremitatem infra folia sitis, spicâ femineâ pedunculatâ terminali, nuculâ irregulariter turbinatâ subradiatim et elevatim nervulosâ alis obovatis multinerviis nuculæ diametro sublongioribus ȝ. *P. fraxinifolia* Spach Hist. Veg. II, p. 180. — Cas. n.c. in Ann. Sc. Nat. Ser. IV, vol. 18, fig. 7, 19, 20, 47-49. — *Juglans pterocarpa* Willd. Sp. IV, p. 455. — *Pt. Caucasica* C.A. Mey. Verz. p. 134. — *Rhus obscurum* M.B. Taur. Cauc. № 606.

Hab. in sylvis regionis inferioris ad mare Nigrum et Caspium, Pontus Lazicus ad Kemer Bournon inter Rhizé et Atina (Balt), Imaretia ad Poti (Rehm!), districtus Scheki (Stev.) et Talyseh prope Lenkoran et Astara (C.A. Mey! Hoht), Persiae bor. provinciis Ghilan et Mazanderan (Auch., Buhsei!), Asterabad (Bge!). Tataris *Motalagatch*, Persis *Kutschî*, *Lary*, vel *Chuol*.

Arbor 30-40-pedalis, folia 6-10-pollicaria foliolis 2-3-poll. longis, nuculæ absque alis *Piso* submajores.

ORD. CXVI. PLATANACEÆ.

(Lindl. Nat. Syst. p. 187.)

Flores monoici in capitula unisexualia globosa dispositi. Perigonium nisi squamulas minimas pilosulas saepe obsoletas pro eo sumas nullum. *Masc.* Stamina plurima squamis carnosis subelavatis (staminibus abortivis?), mixta filamentis brevissimis, antheris bilocularibus oblongis, connectivo truncato-peltato. *Fem.* Ovaria squamis clavatis (ovariis abortivis?) intermixta obconica basi pilosa unilocularia in stylum elongato-subulatum apice hinc stigmatosum abeuntia. Ovulum ex latere ventrali ovarii pendulum testâ tenui. Albumen nullum. Embryo rectus, radiculâ superâ. — Arbores.

PLATANUS (Tourn. Inst. p. 590, tab. 363).

Character ordinis. Arbores foliis alternis, petiolis basi inflatâ geminas foventibus. Pedunculi in ramulis junioribus quorum folia nondum evoluta terminales elongati penduli capitula 3-5 subsessilia moniliformiter ferentes, inferiores capitula mascula, superiores feminea.

1. **P. Orientalis** (L. Sp. pag. 417) arborea, foliis subtus tomentosis sed demum glabratis a basi cuneatâ vel rarius truncatâ vel subcordatâ plus minus profunde 3-5-fidis lobis lanceolatis iterum lobatis vel dentatis ȝ. Ic. Fl. Græc. tab. 945. — Pall. Fl. Græc. tab. 51.

Hab. in totâ fere ditione ad fontes regionis inferioris culta, sed spontanea in sylvis et ad torrentes regionis montane et subalpine, Græci e mortis seclus

rivos in consortio *Cercidis* (Boiss!), Macedonia, Thracia et Bithynia in Castanetis et sylvis montanis 0-2500' (Griseb!), Anatolia occidentalis et australis infra regionem Cedrorum usque ad 5000' (Tchih.), Creta et Rhodi montes (Heldr! Bourg!), Cyprus (Ky!), Libanus (Bl! Gaill!), Paphlagonia (Tchih!), Imeretia et Guria (Ledeb!), ditio Talysch (C.A. Mey.), Persia borealis usque ad Asterabad (Auch. 5330! Buhse!). Græcis *Platanos*, Persis *Tchanar*.

Cortex arboris Orientalis rugosus nec squamis latis caducis denudatis ut in plantâ in ambulaeris Europæ cultâ et cuius nomen et origo mihi incerta.

Ar. Geogr. Regnum Neapolitanum, Sicilia, regio Himalaica.

ORD. CXVI. CUPULIFERÆ.

(Rich. Anal. Fr.)

Flores monoici. *Masculi* in amenta cylindrica vel subrotunda bracteata vel ebraeata dispositi. Perigonium phyllis 5-9 inaequalibus vel nullum. Stamina 5-20 e fundo perigonii libera antheris bilocularibus rarius unilocularibus. *Feminei* solitarii vel aggregati involucro acrecenti foliaceo, cupulari vel coriaceo cincti. Perigonium ovario adnatum limbo denticulato sape evanescente. Ovarium 2-6-loculare loculis 1-2-ovulatis, ovulis pendulis. Styli 2-6 basi connati. Fructus involucro vario pericarpia includenti vel basi foventi fructum spurium formanti nucibusque abortu unilocularibus monospermis constans. Embryo rectus, radicula hilum spectans. Albumen nullum. — Frutices vel arbores foliis alternis stipulatis.

QUERCUS (Tourn. L. Gen. 1070).

Flores masculi in amenta filiformia pendula dispositi. Perigonium 5-9-partitum phyllis linearis-oblongis ciliatis. Stamina 5-20. *Flores feminei* globosi solitarii involucro bracteis imbricatis constanti tandem in cupulam acreto inclusi. perigonii tubus ovario adnatus, limbus parvus denticulatus superus. Ovarium superne uniloculare inferne placentis in centro vix coalitis triloculare loculis biovulatis, styli tres. Fructus grande abortu uniloculari monospermâ basi involucro cupulari inclusâ constans. Ovula abortiva nunc circa inferiorem nunc circa superiorem parlem seminis cum parietum vestigiis plus minusve persistentia. Cotyledones crassæ carnosæ.

Specierum Orientalium distributio.

SECT. LEPIDOBALANUS Endl. — Styli secus superficiem internam nec apice tantum stigmatosi. Capsula squamis sparsis imbricatis lecta. Ovula abortiva quina ad basin seminis.

§ 1. **Eulepidobalanus** Oerst. — Styli breves apice dilatati explanati. Fructus maturatio annua.

* Folia autumno aut vere decidua non sempervirentia.

+ Folia membranacea sinuata vel pinnatilobata lobis mucicis.

Q. pedunculata, *Armeniaca*, *sessiliflora*, *macranthera*, *conferta*.

++ Folia indivisa crenata vel dentata dentibus callosis vel mucronatis.

Q. Pontica, *Lusitanica*.

** Folia crassa sempervirentia plures annos persistentia.

Q. Ilex, *Balout*, *Suber*.

§ 2. **Cerris** Oerst. — Styli lineares apice subulati. Fructus maturatio biennis.

* Folia sempervirentia plures annos persistentia.

Q. alnifolia, *coccifera*, *Aucherii*

** Folia sero autumno aut tandem vere decidua.

Q. Cerris, *Ehrenbergii*, *Egilops*, *Macedonica*, *Look*, *Brantii*, *Persica*, *Libani*, *castaneofolia*.

§ 1. **Eulepidobalanus**.

* Folia non sempervirentia.

1. **Q. pedunculata** (Ehrh. Arbor. N° 77) arborea, foliis breviter petiolatis vel subsessilibus oblongo-ovovatis basi truncatis vel emarginatis glabris sinuato-lobatis vel pinnatifidis lobis rotundato-ovatis mucicis, pedunculis petiolo multoties longioribus, cupulae squamis tomentellis brevibus mucronatis adpressissimis 5. Ic. Rehb. Germ. t. 1313. — Kotsch. Eich. tab. 27. — *Q. ravenosa* Lam. Dict. I, p. 715 (excl. variet.). — *Q. pedunculiflora* C. Koch Linn. XXII, p. 324. — *Q. hypochrysa* Stev. Taur. p. 307.

Hab. in sylvaticis regionis montane et subalpinæ Græcie in Baetœ (Hedde!), Macedonia (Griseb.), Thracia (Webb.), Anatolia (Oliv!), Cappadocia (Ky!), Armenia Turcia in prov. Musch fre-piens (Ky!), Tauria (Stev!), prov. Caucasia et Transcaucasia (Ledeb., Ituprechtl.).

Æ. Haas A. D. C. Prodr. p. 5. — Folia subtus stellatum puberula nigra obtusissime lobata. Glandes magnæ cupulis latis. — *Q. Haas* Ky. Eich. t. 1, 2.

Hab. in Tauro Cilicie circa Gülek Boghas 3-5000' (Ky! Ba1!). Forma analoga aliis formis macrocarpis ejusdem speciei, *Q. Brutia* et *Q. Thomasii* Te.

γ . *pinnatipartita*. — Folia in laçinias numerosas oblongo-lineares profunde fissa vel partita. — *Q. Robur*. var. *laciniata* et *petiolaris* A. D.C. loc. cit.

Hab. in prov. Musch Armeniæ (Ky. 435! 436, 433! 551! 565!).

Species a nonnullis botanicis cum *Q. sessiliflora* conjuncta sed etiam saltuarii omnino contradicunt, differentias specificas in gemmis, ligni duritie, etc., agnoscunt. Area geographicæ multo magis versus septentrionem extenditur.

Ar. Geogr. Europa tota a Scandinaviâ et Rossia septentrionali.

2. Q. Armeniaca (Ky. Eich. tab. 25) arborea humilis glabra, foliis longiuscule petiolatis basi truncato-retusis obovatis vel obovato-oblongis margine grosse obtusissime et inæqualiter dentato-lobatis, pedunculis petiolo pluries longioribus, fructibus numerosis parvis, cupulæ squamis tomentellis brevibus latis in appendice triangularem adpressissimam abrupte abeuntibus §. *Q. Robur* var. *Armeniaca* Alph. D.C. loc. cit.

Hab. in sylvis montanis Ponti Lazici supra Trapezuntem (Bourg! Ky!), prope Rhizé (Bal!).

Arbor 24-3'-pedalis, folia eis *Q. pedunculatae* pro longitudine latiora obtuse dentato-lobata nec sinuato-pinnatifida, fructus numerosiores minores, cortex ramulorum nigricans. Characteribus species distincta videtur.

3. Q. sessiliflora (Sm. Brit. III, 1026) arborea elata, foliis longiuscule petiolatis a basi truncatâ vel cuneatâ obovalo-oblongis sæpius glabris vel glabrescentibus plus minus profunde sinuato-pinnatifobatis lobis obtusis mucicis, pedunculis vix petiolum æquantibus vel subnullis, cupulæ squamis tomentellis in appendicem triangularem brevem adpressissimam abeuntibus vel breviter lanceolatis obtusis § Ky. Eich. tab. 32. — *Q. Robur* L. ex parte. Rehb. Germ. tab. 1309. — *Q. Iberica* Stev. Mém. Mosq. IV, p. 70. — *Q. Æsculus* Griseb. Spic. II, 336 an Linn? — *Q. Hartwissiana*, *Q. erucæfolia* et *Q. longipes* Stev. Taur. p. 306.

Hab. in sylvis montanis Græciae 2-40° (Heldr!), Thraciæ (Jankal), Anatoliæ borealis, Armeniæ Turcicæ (Ky!), Tauriæ (Stev!), Caucasi et Transcaucasiæ (Ledebl! Ruprl!), prov. Karabagh (Szovl!), prov. Ghilan Persiæ bor. (Auch. 553!).

β *Cedrorum* A. D.C. loc. cit. p. 8. — Folia angustiora oblonga sæpe profundius pinnatifida. — *Q. Cedrorum* Ky Eich. tab. 37. — *Q. Abietum*, *Q. subalpina*, *Q. Iberica* Ky. pl. exs. — *Q. longifolia* C. Koch Linn. XXII, p. 327 (ex descript.).

Hab. in regione montana Tauri Cilicici (Ky. 373! 377! 375!), Libani supra Elen (Ky. 335!), Armeniæ Turcicæ (Ky 55!).

γ . *pinnatifida*. — Folia profundius, ultra medium pinnatifida vel partita. — *Q. Toza* Griseb. Spic. non Bosc. — *Q. pinnatifobata* C. Koch Linn. XXII, p. 326. — *Q. vulcanica* Boiss. et Heldr. MSS. — Ky. Eich. tab. 18.

Hab. in Macedoniâ supra Vodena (Orph!), Thraciâ (Griseb.), Arcadiâ ad Leondari (Heldr. exs. 3453!), monte Karadagh Lycaoniæ (Heldr!), Libano ad Danie 5000' (Ky. 336 sub *Q. Tozâ!*!).

δ. mannifera. — Folia intensius viridia supra lucida subtus præsertim ad nervos sæpe pubescentia parum profunde lobata, cupulæ squamæ magis lanceolatæ. — *Q. mannifera* Lindl. Bot. Reg. 1840, p. 40. — *Q. intermedia* Ky. exs. — *Q. Robur* var. *Buhseana* A. D.C. Prodr. p. 7. — *Q. lamprophyllos* C. Koch Linn. XII, p. 327. — *Q. Syspirensis* Alph. D.C. Prodr. l. cit. p. 12 non C. Koch.

Hab. in Ponto supra Trapezuntem et ad Gumuschkhané (Bourg! Ky. 36! 37! 559!), Imeretia, Colchis, Albania (Rupr!), Persia borealis (Buhse!).

ε. Tchorochensis A. D.C. loc. cit. — Folia oblonga glabra obtusissime sinuato-lobata et undulata. — *Q. Dchorochensis* Ky. Eich. tab. 39. — *Q. pseudodchorochensis* Ky. exs. — *Q. Robur* var. *bullata* A. D.C.

Hab. in Ponto et Arneniâ Turcicâ prope Trapezuntem et Gumuschkhané (Bourg! Ky. 38! 39! 40!).

ζ. pubescens. — Folia vernantia tomentosa adulta subtus pubescentia obovata basi sæpe emarginata obtuse lobata. — Ky. Eich. tab. 34. — *Q. pubescens* W. — *Q. lanuginosa* Thuill. — *Q. Syspirensis* C. Koch Linn. XII, p. 328 et herb! non Alph. D.C.

Hab. in collibus et sylvaticis, in Græciâ vulgaris, Arcadia (Guicci!), mons Kyllene (Orph!), montes Achaiæ circa Megaspileon (Heldr!), Parnes Atticæ (Heldr!), Parnassus (Orph. Fl. exs. 413! sub *Q. congestâ*), Olympus Thessalus (Heldr!), Creta (Heldr! Raul!), insula Chios (Heldr!), Armenia ad Ispir (Ky!), Tauria (Rehm!), Transcaucasia (ex Ledeb.).

η. brachyphylla Alph. D.C. l. cit. p. 11. — Folia ut in varietate præcedente subtus tomentosa sed abbreviata obtusissima et rotundatim pauciloba. — *Q. brachyphylla* Ky. Eich. tab. 9.

Hab. in Cretâ ad Retimo (Ky!), Apocorona (Raul!), Eubæâ (Ky. 50! Heldr!), Olympo Thessalo ad Litochorón (Orph!), ad Cordilio prope Smyrnam (Ky. 26! Bal. 289!). In Eubæâ Méradic, in Cretâ *Granitia* dicta.

Species valde ludibunda et cuius formæ altera in alteram confluunt, ejus varietates profundius pinnatisidæ cum *Q. Tozâ* Bosc. Europæ magis occidentalis incolâ foliis subtus tomentosis et cupulæ squamis elongatis distinctâ ron confundendæ sunt.

Ar. Geogr. Europa media et australis a Daniâ, Gothiâ ad Rossiam medium.

4. *Q. macranthera* (Fisch. et Mey. in Hohen. Talysh Enum. p. 29) arborea, foliis breviter petiolatis amplis obovatis basi rotundato-cuneatis vel subcordatis obtuse sinuato-multilobatis subtus pubescenti dense tomentosis, antheris filamento longioribus, spicis feminis brevibus subsessilibus hirsutissimis, cupulæ squamis pubescentibus lanceolatis adpressis 3.

Hab. in sylvis prov. Talysh prope Lenkoran, Astara, Swant (C.A. M!), in monte Sarial Georgie (Höht!), prov. Karabagh (C.A. Mey!), Armeniâ prope Nackitchewan (Szov!).

Affinis *Q. sessiliifloræ* præsertim ejus var. *manniferæ*, differt foliis ampliori-

bus brevius petiolatis subtus dense tomentosis et ex cl. auctore antheris duplo longioribus (quæ in spec. meis non sat evolutæ).

5. **Q. conferta** (Kitaib. in Rehb. Fl. Germ. exsicc. 1840) arborea elata. foliis sessilibus amplis a basi subcordatâ sensim dilatatis oblongo-ovatis spathulatis anguste multisinnatis crenatissimis lobis late linearibus integris vel obtuse lobulatis subtus plus minusve stellato-pubescentibus vel hirtellis, fructibus 2-5 in apice ramuli sessilibus, cupulæ squamis adpressis a medio sursum lanceolatis elongatis apice interdum subpatentibus 5. Ic. Rehb. Germ. tab. 1311. — Ky. Eich. tab. 14. — *Q. Farnetto* Ten. Cat. Nap. 1819, p. 65.

Hab. in sylvis montis Malevo Laconiae ad Hagios Petros (Orph!), Græciæ occidentalis (Ky. 62! et 63!), Byzantii prope Buyukderé (Janka!).

Specie a formis *Q. sessilifloræ* differre videtur foliis multo amplioribus subsessilibus a basi angustâ sensim dilatatis elongatis, glandibus omnino sessilibus confertis. Cupulæ squamæ longiores apice non adpressæ. Cum *Q. Tozæ* specie Europæ occidentalis non confundenda.

Ar. Geogr. Calabria, Banatus, Serbia.

6. **Q. Pontica** (C. Koch Linn. XXII, p. 319) fruticosa humilis, ramis glabris, foliis membranaceis ad ramorum hornotinorum brevium partem superiorem 4-5 approximatis magnis brevissime petiolatis a basi rotundato-subcuneatâ ellipticis obtusiusculis supra glabris subtus pallidioribus puberulis tandem glabratris parallele 15-18-nerviis circumcircâ denticulato-serratis denticulis acutis sursum uncinato incurvis, amentis masculis infra folia nascentibus longis pendulis, floribus feminis terminalibus solitariis sessilibus, squamis ad perigonii basin velutinis oblongis laxiusculis, perigonio sericeo, stylis apice dilatatis, fructu... 5.

Hab. in regione montanâ Ponti Lazici ad fontes fluvii Asferos (C. Koch), in mediâ viâ inter Andon et Djimil et in parte superiori vallis Of (Bal!).

Species insignis cuius tamen fructus adhucendum ignoti. Frutex tantum 9-10-pedalist., folia petiolo crasso 3-4 lineas longo suffulta 5-6-pollices longa 3-4 lata denticulis acutis sesquilineam longis, amenta numerosa 4-pollices longa. Ob fructus ex cl. Balansa maturatione annuos atque stigmatum formam minime inter *Cerrides* ut cl. C. Koch autumat sed inter *Eulepidobalanos* militat.

7. **Q. Lusitanica** (Lam. Diet. I, p. 719) arborea humilis vel fruticosa, foliis serius persistentibus tamen ante frondescientiam novam deciduis subcoriaceis petiolatis mediocribus a basi rotundatâ vel truncatâ ovalis vel ovalo-oblongis interdum sublanceolatis supra lucidis subtus stellato-pubescentibus vel glabratris obtusis vel acutis marginis dentitis vel crenatis dentibus et crenis acutis vel callosis, fructibus breviter pedunculatis sessilibusve solitariis vel paucis, cupulæ squamis tomentellis adpressissimis dorso longitudinaliter plus minus elevatim convexis infimis triangulari-ovalis cæteris breviter lanceolatis 5. *Q. faginea* Lam. Diet. 725.

Species polymorpha a *Q. sessilifloræ* formis foliis magis coriaceis diutius

persistenteribus nunquam obtuse sinuato-lobatis sed serratis vel crenatis dentibus callosis vel acutis semper distinguenda. Hujus formae intermediis altera in alteram transeunt, frequentiores enumero.

α. gennina. — Folia ovato-oblonga subtus pallidiora glabra vel rarius stellato-pubescentia margine dentata. — *Q. infectoria* Oliv. Voy. I, p. 252, tab. 14 et 15. — *Q. corymbifolia* Ehr. exs. — *Q. inermis* Ehr. ex parte exs. 295 et Ky. Cypr. pag. 216. — *Q. polycarpos* Ky.

Hab. in regione inferiori et montanâ Thracie ad Byzantium (Auch. 2884! Jankat!), Bithyniae (Griseb!), Lydiae (Heldr. 2185!), regione subalpinâ et alpinâ Armenie Turcicae (Ky. 563! 565! 567! 568! 570!), Cilicie circa Gulek (Ky. 370! Bal. exs. sub *Q. longitolia*!), Cypro (Ky!). Libano (Oliv., Ehrenb! Bl!), Kurdistanâ Persicâ in montibus Avroman et Schahu 6-7000' (Haussk!).

β. Boissieri Alph. n.c. loc. cit. — Folia subtus glabra vel glabrescentia oblonga et oblongo-lanceolata margine undulata regulariter et acute dentata. — *Q. Boissieri* Reut. in Boiss. Diagn. Ser. I, 12, p. 119. — *Q. rigida* C. Koch non Willd.

Hab. cum var. praeedente a quâ saepe non distinguenda, Thracia et Bithynia (Thirke! Heldr! Jankat!), Lydia (Boiss!), Armenia australis (Ky. 377! 369! 433!), Cataonia (Haussk!), Lycaonia (Heldr!), Cyprus (Ky. 374!), mons Amanus (Ky. 431!), mons Cassius (Boiss!), Libanus (Bl!), Kurdistanâ Persicâ (Haussk!).

Hujus varietatis et praeedentis specimina plura Hispanicis simillima sunt.

γ. petiolaris Alph. n.c. loc. cit. — Folia saepe longius petiolata paucicrenata vel integra oblonga vel ovata subtus glabrescentia. — *Q. petiolaris* Boiss. Diagn. I, 12, p. 120 non Blh. — *Q. leptocarpus* Ky. exs. 372! — *Q. Pfaffingeri* Ky. Eich. tab. 23. — *Q. tenuicola* Ky. Eich. tab. 10 (forma foliis acutius dentatis ad var. α transiens).

Hab. in Cariâ (Pinard!), Pamphyliâ (Heldr!), Ciliciâ circa Gulek (Ky. 322! 373! Bal!), monte Amano (Ky. 419! 209!), monte Cassio (Boiss!), Cypro (Ky. 761!), Kurdistanâ Persicâ inter Avroman et Masibin, prope Pendjavin 5-9000' (Haussk!).

δ. latifolia. — Folia late ovata margine saepe undulata obtuse crenata vel subintegra subtus saepius tomentella, glandes saepe grandes. — *Q. Cypria* Ky. ex parte non Spach. — *Q. inermis* Ehr. ex parte. — *Q. Syriaca* Ky. Eich. tab. 4. — *Q. Lusitanica* var. *Syriaca* Alph. n.c. — *Q. hypoleuca* Ky. exs. 99? ramus sterilis et ideo dubius.

Hab. in Cypro (Ky. 720! et 721!), Ciliciâ Gülek (Ky. 307!), Libano (Bl! Bal!), Antilibano (Ky. 364!).

Ar. Geogr. Lusitania, Hispania, Africa borealis in prov. Oran (Warion).

“ Folia sempervirentia.

8. **Q. Ilex** (L. Sp. 1412) arborea humilis, cortice non suberosâ, foliis persistentibus sempervirentibus coriaceis subtus pube stellatâ brevi incano-tomentosis superne glabris breviter petiolatis ovatis oblongis vel elliptico-lanceolatis acute dentatis vel integris acutis vel obtusis, antheris mucronulatis, racemis fructiferis brevibus breviter

pedunculatis, cupulæ squamis adpressissimis tomentosis triangulari-ovatis obtusis ȝ. Ic. Rehb. Germ. t. 1307. — Ky. Eich. tab. 38.

Hab. in regione inferiori et montana Græciæ (Sibth., Heldr! Ky!), Cephaloniae (Heldr!), Cretæ (Heldr! Ky!), Macedonia (Friv! Griseb.), Archipelagi (Auch.), Ponti ad Samsun (Ky. 71!), Syria littoralis (Ky!).

Ar. Geogr. Europæ regio Mediterranea, Africa borealis

9. Q. Balout (Griff. Itin. Not. p. 328) arborea mediocris a basi fere ramosa vix suberosa, foliis persistentibus sempervirentibus coriaceis supra glabris subtus glancis minutissime lepidotis breviter petiolatis basi rotundatis ovato-ellipticis obtusis integris, antheris muticis, cupulæ ore contractæ squamis minutis densissime imbricatis ȝ.

Hab. in montibus Bala Chugur Seraï 3000'-5500' (Griff. 4448!).

Folia 14-18 lin. longa 8-9 lata illis formarum integrifoliarum *Q. Illicis*: imilia sed indumentum faciei inferioris diversum squamulis minutissimis nec pilis stellatis constans. Fructus a me non visus. Ramulos steriles speciei affinis sed foliis longe aristato-dentatis subtus nudis ex Bharowul Afghaniæ (Griff. 23 sub *Q. Balout* var.) vidi.

10. Q. Suber (L. Sp. 1413) arborea, cortice suberoso, foliis persistentibus sempervirentibus coriaceis supra glabris subtus tomentoso-canis breviter petiolatis ovatis vel oblongis acutis acute paucidentatis rarius integris, antheris muticis, fructibus solitariis vel binis pedunculis brevibus crassis suffultis, cupulæ squamis tomentellis inferioribus brevibus adpressis superioribus elongatis acutis erectis laxiusculis ȝ. Ic. Rehb. Germ. tab. 1306.

Hab. in Græciâ raro (ex Fraas), in Albaniâ (Griseb.). E ditione nondum vidi.

Species ab auctoribus nonnullis cum *Q. Illice* confusa sed characteribus indicatis probe distincta. Ex cl. Oersted in Liebm. Chênes d'Amérique p. 45 et 18 *Q. Suber* stylis linearibus elongatis a *Q. Illice* stylis brevibus explanato differt et in aliâ sectione militat, ego tamen in speciminibus exsiccatis herbarii et in iconum analysibus in utrâque specie stylos breves apiceque recurvos et dilatatos vidi. *Q. Suber* Ky. Eich. t. 33 ex cl. Alph. D C. est *Q. occidentalis* J. Gay.

Ar. Geogr. Hab. in regione mediterraneâ totius Europæ et Africæ borealis.

§ 2. CERRIS.

* Folia sempervirentia.

11. Q. alnifolia (Poech Enum. Cypr. p. 12) fruticosa interdum arborescens, foliis breviter petiolatis coriaceis persistentibus sempervirentibus supra glabris subtus pubescentiæ adpressissime fulvo-tomentosis a basi rotundatâ vel subcordatâ ovatis vel orbiculatis obtusis vel acutiusculis integris vel a medio sursum parce denticulatis, nervis superne impressis subtus valde prominentibus, amentis masculis brevibus villosis, antheris pilosis, fructibus solitariis vel 2-3-nis

sessilibus, cupulae breviter cyathiformis squamis tomentellis inferioribus conicis adpressis, intermediis lanceolatis superioribus anguste linearibus longis uncinatim recurvis, glande cupulam 4-5-plo superante ♂. Ic. Ky. Eich. tab. 6. — *Q. Cypria* Jaub. et Sp. III. Or. tab. 56 non Kotschy (specim. juniora glande subabortivâ).

Hab. in montibus Cyperi 1600'-5000', supra Paphos (Auch. 2881!), in jugo Troodos (Ky. 4091 7561).

Frutex 4-1½-orgyalis, folia 9-15 lineas longa et lata ea specierum *Gaultheriae* referentia, glans 12-15 lineas longa. Fructus ex el. Gay maturatione annui sed ex el. Kotschy qui eos in loco natali observavit biennes quod melius quoque cum cupulae structurâ et stylis cylindricis congruit. Species distinctissima.

12. *Q. coccifera* (L. Sp. 1413. — J. D. Hook Trans. Linn. Soc. vol. XXIII, tab. 37) fruticosa ramosissima vel arborea, foliis semper-virentibus brevissime petiolatis coriaceis ovatis vel oblongis basi rotundatis vel cordatis saepius glaberrimis spinoso-dentatis planiusculis vel undulatis, fructibus solitariis saepius binis, cupulae hemisphericae squamis plus minus velutinis regulariter imbricatis parvis ab infinitis ovalis ad intermedias oblongo-lanceolatas et superiores angustiores lineares sensim auctis laxiuscule erectis vel squarrosis vel retroflexis ♂ et ♀.

Species suminopere variabilis cujus varietates magis notabiles quas numero formis intermediis conjuguntur.

α. genuina. — Suffruticosa vel fruticosa humilis ramis intricatis, folia parva ovata undulata valde spinosa, cupula breviter echinata. Ky. Eich. tab. 29.

Hab. in regione inferiore Graeciae (Heldr!), Zacynthi (Marg!), Cretæ (Ky. 309!), Macedoniae, Thraciae et Thraciae (Griseb., Noë!), Lydie ad Smyrnam (Fleisch!), Ponti prope Samisun (Tchih!).

β. integrifolia. — Fruticosa, folia parva vix denticulata vel integra, squamæ adpressæ vel breviter apice echinatae.

Hab. in Atticâ ad radices Hymetti (Heldr!), Cretâ in regione sylvaticâ montis Ida et in prov. Sphakia (Heldr!), Cataoniâ (Haussk!) et alii.

γ. calliprinos. — Fruticosa vel arborea, folia plana saepius majora, cupulae squamæ erectæ plus minus adpressæ. — *Q. calliprinos* Webb It. Hisp. p. 15. Jaub. et Sp. III. tab. 57. — *Q. pseudococcifera* Labill. Dec. V. p. 9, tab. 6 non Desf. — *Q. Fenzlii* Ky. Eich. tab. 24 (variatio foliis minoribus). — *Q. inops* Ky. vix forma fructu imperfecto.

Hab. in Tauro (Bal!), Libano (Labill! Bl!), monte Thabor Palestinæ (Gaill!).

δ. pseudococcifera. — Fruticosa elata vel arborea, folia planiuscula oblonga basi plerumque cordata majuscula var. γ, sed cupulae ob squamas plus minus patentes vel recurvas squarrosæ. — *Q. pseudococcifera* Desf. Fl. Atl. II, p. 349. Boiss. Voy. Esp. t. 165. — *Q. coccifera* Fl. Graec. tab. 944 et Rehb. Ic. Germ. 1308. — *Q. valida, dispar, echinata, brachybalanos, dipsacina, Sibthorpii* Ky. Sched.

Hab. præsertim in regione montanâ a Graeciâ (Heldr!), Bithyniâ (Griseb.),

Ponto australi (Tchih.), Cretā (Heldr!), Anatoliā australi et Ciliciā (Ky! Bal!) ad Libanum et Antilibanum (Ky! Gaill! Bl!), Palestinam in Carmelo (Gaill!).

Folia hujus formae 1-2-pollices longa dentata vel subintegra e Græciā! Libano! vidi, Arbores in Græciā 20-30-pedales vidi et in Bithyniā cl. Grisebach 80-pedales observavit.

ζ. Palestina. — Arborea, folia planiuscula varietatis præcedentis saepe elongata oblonga dentata vel subintegra. Cupulae squamae eximie retrofractæ — *Q. Palestina* Ky. Eich. tab. 19. — *Q. arcuata* Ky. Sched. — Ic. Trans. Linu. Soc. XXIII, tab. 36.

Hab. in regione montanā Ciliciæ (Ky! Bal!), Cretæ in montibus Sphaciotio's (Heldr!), Palestinā australi inter Gaza et mare Mortuum, ad Hebron, *Quercus Davidis* dictus, grandævus, coma diametro 90-pedali (Ky. 442! et 21!).

η. rigida Alph. p.c. l. cit. p. 56. — Fruticosa, folia parva ovato-oblonga valde spinosa rigida et undulata var. α sed subtus juniora tomentosa, adulta pruinoso-glauea, squamae echinato-recurvæ. — *Q. rigida* Willd. Sp. IV, p. 404. Ky. Eich. tab. 8.

Hab. in insulâ Chio ad Vrontado (Orph!), Lyciâ in collibus Elmalu et Pamphyliâ ad Adalia (Bourg!), regione montanâ Ciliciæ prope Gülek (Ky. 404!).

Cl. Kotschy in schedulis hujus speciei formas nullo modo distingendas aliis nominibus specificis plurimis designavit, quas consulto omitto.

Ar. Geogr. Regio mediterranea totius Europæ et Africæ borealis.

13. *Q. Aucheri* (Jaub. et Sp. Ill. Or. I, tab. 58) fruticosa ?, ramis junioribus fulvo-velutinis, foliis ovatis et ellipticis subsessilibus basi leviter cordatis rigidis aliis integris obtusis aliis spinoso-dentatis supra glabrescentibus subtus incano-tomentosis, fructibus sessilibus solitariis, cupulae turbinateæ cinereo-velutinæ squamis imbricatis inferioribus ovatis obtusis superioribus gradatim angustioribus extremitate recurvâ subpatentibus, glande vix exsertâ δ .

Hab. in insulâ Cos Archipelagi (Auch. 1879 herb. Par.). Non vidi.

Folia fere pollicaria petiolo 1-2 lineas longo suffulta, cupula 8-9 lineas longa 9-10 lata. An foliis subtus tomentosis et cupulâ turbinate sat a formis *Q. cocciferae* distincta inter quas var. *rigida* foliis subtus glaucis ad eam accedit sed cupulâ minore echinata differt?

** Folia sero autumno aut tandem vere decidua. — Grex cujus species limitibus certis ægre delimitantur. Plures ab auctoribus propositas reduxi et forsitan non sat audaci.

14. *Q. Cerris* (L. Sp. 1445) arborea, foliis deciduis membranaceis breviter petiolatis supra glabrinsculis subtus pubescentibus tandem concoloribus oblongis plus minus profunde pinnatifolatis lobis subintegratis triangularibus obtusis sœpius acutis semper apice callosis vel mueronatis, fructibus 1-3 subsessilibus, cupulae hemisphericae fuscescens echinatae squamis pubescentibus elongatis inferioribus lanceolatis patentibus superioribus linearis-subulatis erectis vel patenti-reflexis et contortis δ . Rchb. Ic. Gerin. t. 1316. —

Q. Austriaca Willd. Sp. IV, p. 454. — Ky. Eich. tab. 20. — *Q. crinita* Lam. Diet. I, p. 718. Oliv. Voy. tab. 12.

Hab. in sylvis regionis inferioris montanae et subalpinae. Macedonia (Griseb.), Lydia (Boiss!), Caria (Pinard!), Pamphylia in planicie et montibus usque ad 4500' (Heldr!), Cilicia (Ky. 385!), Syria borealis in monte Amano (Ky. 432!), Libano (Bl!) et circa Eden 550' (Ky. 328!), Anatolia borealis (Tehih!).

β *pseudocerris*. — Folia in lacinias lineares integras vel 3-5-lobas plus minus profunde pinnatipartita, cupulae squamme subvalidiores facie externa convexae. — *Q. pseudocerris* Boiss. Diagn. Ser. I, 12, p. 118. — *Q. Vallonea* Ky. Eich. tab. 7 non Kotschy exs.

Hab. in Lydiæ monte Timolo supra Philadelphium (Boiss!), montibus Cataoniae (Haussk!), vallibus Tauri Cilicie (Ky., Bal!), monte Cassio Syriæ (Boiss!). Cupulæ Graece *Velani*, Turcice *Balamut* dictæ ad nigrum colorem tincturæ inficiendum adhibentur.

Ar. Geogr. Gallia occidentalis et Jurassus!, Helvetia australis, Italia et Sicilia, Germania australis, Dalmatia, regio Danubialis.

15. Q. Ehrenbergii (Ky. Eich. tab. 45) arborea mediocris, foliis deciduis membranaceis petiolatis utrinque pilis stellatis tomentellis ambitu obtuse et breviter ovatis in lobos inaequales oblongos saepius acatos et apiculatos integros vel apice dentatos pinnatisectis vel profunde pinnatifidis, fructibus magnis solitariis subsessilibus, cupulae squamis laxiusculæ imbricatis tomentellis omnibus erectis et brevibus inferioribus et intermediis ovatis et oblongis acutis vel apiculatis, summis anguste linearibus vix longioribus, glande cupulâ sesquilociore 3.

Hab. in regione montanâ prope Mindos Gala prope Mersina Cilicie 2000' (Ky. 303!), in Libano (Oliv!), prope pagum Massa Antilibani (Ehrenb.), ad radices sept. montis Hermon (Ky.).

Differet a *Q. Cerri* foliis ea *Cratægi Orientalis* referentibus fere aequilongis et latis obtusissimis fere 2 pollices longis, cupulæ squamis laxissime imbricatis brevibus latis superioribus non recurvis.

16. Q. Ægilops (L. Sp. p. 1414) arborea elata trunco crasso cymâ dilatata rotundata, foliis subcoriaceis serius deciduis petiolatis supra glabratâ lucidis subtus tomentellis canescensibus a basi truncato-rotundata vel cordata ovatis vel oblongis grosse et inæqualiter dentato-serratis dentibus acuminatis vel mucronatis, fructibus solitariis rarius 2-3-nis crassis, cupulae globosæ squamis crassis tomentellis ovato-lanceolatis superioribus vel omnibus elongatis patentibus recurvis, glande cupulâ 2-3-plo longiore 3. Ie. Oliv. Voy. tab. 13. — Tehih. Voy. tab. 41. — J. D. Hook in Trans. Linn. Soc. XXIII, tab. 38. — *Q. Græca* Ky. Eich. tab. 30. Species summopere variabilis.

Hab. in Græciâ totâ a regione inferiore ad 3000' (Boiss! Heldr! Orph. Fl. exs. 301!), Cephallenâ (Heldr!), Cycladibus (Orph.).

β . *macrolepis*. — Vix variatio squamis omnibus elongatis et recurvis. — *Q. macrolepis* Ky. Eich. tab. 16.

Hab. in Cretâ circa Retimo (Sieb! Ky. 309!), in Græciâ (Ky. 65!).

γ. Ungerii. — Folia minora oblonga. Cupulæ squamæ intermediae interdum abbreviatæ et recurvæ, superiores elongatæ arcuatæ. — *Q. Ungerii* Ky. t. 43. — *Q. Vallonea* Alph. d.c. Prodr. l. cit. p. 46 non Ky. tab. 7. — *Q. Ægilops* var. *Oliveriana* Tchih. As. Min. tab. 42. — *Q. Trojana* Webb in Jaub. et Sp. Ill. tab. 57. — *Q. Goedelii* Ky. exs.

Hab. in Mysiâ et Phrygiâ (Webb, Jaub., Schmidt!), Anatoliâ meridionali (Heldr!), Lycânia et Ciliciâ (Ky. 387! 390! et № 394! 391! sub *Q. Vallonea*!).

δ. Ithaburensis. — Vix variatio. Folia late ovata formæ typicæ sed (adulta præsertim) brevius denticulata. Squamæ omnes recurvæ inferiores et intermediae abbreviatæ. — *Q. Ithaburensis* Decaisne Ann. Nat. Ser. 2, IV, p. 348. — Ky. Eich. tab. 12.

Hab. in montibus Thabor, Carmel et totâ Galilæâ (Bové! Boiss! Ky! Gaill!).

ε. Pyrami. — Folia ramulorum vernalium oblonga denticulata, vegetius evolutorum variantia latiora profundius lobata sæpe lyrata lobo terminali latiore rotundato basi cordato, squamæ inferiores et mediae ovatae arrectæ, superiores linearis-lanceolatæ retrofractæ. — *Q. Pyrami* Ky. Eich. tab. 3.

Hab. in planicie Cilicieâ ubi ad ostium Pyrami vastas sylvas format (Ky. 20 et 395!).

Species *Q. Cerri* proxima ab eo foliis etsi deciduis diutius persistentibus coriaceis plus minus dentatis sed nunquam pinnatifidis distincta. Cupulæ majores squamis crassioribus.

17. Q. Macedonica (Alph. d.c. Prodr. l. cit. p. 50) arborea, foliis subcoriaceis serius deciduis supra glabris lucidis subtus glabratris brevissime petiolatis a basi rotundata vel truncato-subcordata oblongis obtusiusculis denticulatis vel crenato-repandis dentibus ascendentibus mucronulatis, fructibus sessilibus 1-3-nis, cupulæ hemisphericae squamis pubescentibus infimis ovatis adpressis intermediis oblongis uncinato-recurvis apice rufo-sphacelatis, supremis longioribus inflexis, glande inclusâ aut breviter exsertâ ȝ. *Q. Æsculus* Griseb. Spic. II, p. 333 non L.

Hab. in sylvis Macedoniae supra Vodena *Q. Cerri* immixta (Griseb! Orph!), prope Prisidren et in Albaniâ (Griseb.).

Species ex paucis junioribusque speciminibus nota, an glandes maturæ cupulâ inclusæ? Valde affinis *Q. Ægilops* formis nonnullis præsertim var. *Trojanæ*, sed folia glabrata nec subtus tomentosa, squamæ cupulæ breviore.

18. Q. Look (Ky. Eich. tab. 31) arborea mediocris vel fruticosa, foliis breviter petiolatis rigidis subtus sub lente sparsim stellato-puberulis undulatis grosse crenato-dentatis dentibus mucronatis, fructibus brevissime pedunculatis solitariis vel binis, cupulæ subturbinalæ squamis breviter tomentellis majusculis imis ovatis subadpressis, intermediae ovato-lanceolatis apice infexo, superioribus elongatis linearibus uncinato-reflexis ȝ.

Hab. in latere orientali Hermonis Syriæ prope Rascheya 5000' (Ky. 172!), in districtu Hauran Palestinæ orientalis (Ky. 53!).

Planta ex speciminiis fructu immaturo imperfecte nota, affinis *Q. Aegilopi*.

19. Q. Brantii (Lindl. Bot. Reg. 1840, p. 41) arborea, comæ hemispherico-conicæ, ramulis et petiolis pulverulento-velutinis, foliis seruis deciduis coriaceis petiolatis supra glabratis subtus pilis stellatis pubescentibus late ovato-oblongis basi sœpius cordatis crenbre denticulatis dentibus brevibus acuminato-aristatis rarius crenatis crenis callosis, fructibus sessilibus solitariis maximis, cupulae obconico-hemisphericæ squamis inferioribus et intermediis late ovalis breviter acuminatis adpressis vel apice patentibus, superioribus longioribus linearibus erectis rarius reflexis 5. Ic. Ky. Eich. tab. 31.
— *Q. oophora* Ky. tab. 26! (forma foliis angustioribus fere *Q. Libani*).

Hab. in sylvis Armeniæ australis Kurdiæ ad meridiem urbis Bitlis et lacus Van 4000' (Brant, Ky. 221 533! 554! 560! 531!).

A *Q. Aegilope* cupulae squamis latis brevibus non recurvis, foliis aristato-dentatis, a *Q. Libani* varietatibus cui foliorum formæ accedit eis latioribus subtus tomentosis, squamis latis, etc., distincta.

20. Q. Persica (Jaub. et Sp. Ill. Or. I, tab. 55) arborea humilis vel fruticosa, foliis petiolatis griseis supra pubescentibus vel glabratis subtus tomentosis a basi subcordata vel truncata ovatis vel oblongis obtusis grosse vel repande crenato-serratis dentibus mucronatis, fructibus solitariis subsessilibus, cupulae turbinatae cinereo-velutinæ squamis inferioribus et intermediis late ovatis præter apicem triangularem mucronatum brevissimum patenti-recurvum adpressis adnatis, summis longioribus linearibus erectis vel sub-recurvis, glande cupulâ subdupo longiore 5. Ky. Eich. tab. 28.

Hab. in Mesopotamiâ circa Terek (Haussk!), Kurdistaniâ supra Mossul in monte Gara (Ky. 391!) et Amadiéh (Ky.), montibus Teriter, Schahu, Juanro Kurdistanie Persicae, monte Lolan supra Kermanschah et ubique in montibus ditionum Bachtiaris et Kuh Kilouyeh Persiae austro-occidentalibus 4-9000' (Haussk!), ad Perezend (Auch. 5331!), ad Gere (Ky. 415!), ad Mian Kotel prope Sc. iraz (Haussk!).

A formis *Q. Aegilopis*, et *Q. Brantii* specificè differre videtur cupulâ turbinata, squamis brevi-sime uncinatis.

21. Q. Libani (Oliv. Voy. tab. 32), arborea humilis vel frutex, ramis et petiolis glabris, foliis deciduis adultis utrinque glabratis superne lucidis petiolatis a basi rotundata vel cuneata lanceolatis regulariter et crenbre serratis dentibus ascendentibus triangularibus mucronato-aristatis, fructu solitario subsessili magno, cupulae squamis cinereis imbricatis adpressis breviter ovato-rhombeis supremis angustioribus sed non elongatis, glande breviter exsertâ 5. Ic. Ky. Eich. tab. 5. — *Q. Carduchorum* Ky. exs.

Hab. in sylvis regionis montanae Syrie (Labill.), montis Cassii (Boiss!), Ciliciæ (Ky. 380-383! Bal!), Cataoniae (Tchih!), montis Berydaghi (Haussk!),

Armeniæ australis (Ky. 20! 559! 562!), montis Gara Kurdistaniæ (Ky. 340!), montium Pendjavin et Teriter Kurdistaniæ Persicæ (Haussk!).

β. regia. — Folia ampliora dentibus subremotioribus et submajoris acutissimis aristatis, cupulae squamæ adpressissimæ superiores breviter mucronatae. — *Q. regia* Lindl. Bot. Reg. 1840, № 73. — Ky. Eich. t. 11.

Hab. in Kurdistaniæ Armeniæ (Brant), in prov. Musch Armeniæ (Ky. 432! Tchih!).

γ. vesca. — Formæ typicæ similis sed folia subtus canescentia tomentosa. — *Q. vesca* Ky. Eich. tab. 36.

Hab. in Armeniae Kurdiæ districtu Schirwan ad orientem urbis Bitlis 4500' (Ky. 555!), ad Nadschardagh Tauri Cataonici (Haussk!), in montibus Avroman et Schahu Karduchiae Persicæ 4-7000' (Haussk!).

δ. Tchihatchewii. — Similis var. *γ* sed folia angustiora, cupulae squamæ minores et obtusiores. — *Q. Tchihatchewii* Ky. MSS. — Alph. v.c. I. cit. p. 98. — Ie. Tchih. As. Min. tab. 40.

Hab. in Galatiâ inter Alizy et Mentiche (Tchih.).

Foliis angustis lanceolatis regulariter serratis dentibus aristatis et squamis brevibus adpressis *Q. Libani* a formis *Q. Egilopis* distinguitur.

22. Q castaneæfolia (C.A. Mey. Enum. p. 44 non Cosson) arborea mediocris, foliis deciduis petiolatis supra glabris subtus tomentellis a basi cuneatâ vel rotundatâ lanceolatis acutis grosse dentatis dentibus brevibus latis ascendentibus saepius mucronatis, fructibus 1-3 subsessilibus mediocribus, cupulae hemisphericæ tomentellæ squamis fructu maturo omnibus retroflexis inferioribus ovatis cuspidatis superioribus gradatim longioribus, glande cupulâ 2-3-plo longiore 3. Ky. Eich. tab. 40.

Hab. in sylvis prov. Transcaucasicae Talysch usque ad 5500' (C.A. Mey! Hoh!), in prov. Ghilan Persicæ (Auch. 5333! Buhsel).

A formis *Q. Libani* quibus foliis accedit cupulae squamis tenuioribus inferioribus non adpressis sed tandem retroflexis differre videtur. Icon. Jaub. et Spach. Ill. Orient. tab. 54 ab Aubrietio ex specimine imperfecto Tournefortii delineata non *Q. castaneæfoliam* sed prob. formam *Q. Egilopis* exhibet.

CASTANEA (Tournef. Instit. tab. 352).

Flores masculi in amentis glomerati, glomeruli bracteis cincti. Perigonum regulare sexpartitum. Stamina 40-20. *Flores feminei* intra involucrum quadrifidum plures saepius terni rarius solitarii. Perigonum ovario adnatum limbo supero 5-8-fido, ovarium lageniforme sexloculare loculis biovulatis, stigmata setiformia numero loculorum. Fructus involuero acerulo globoso aculeato denum 2-4-partito et nuculis 1-3 liberis constans. Pericarpium coriacenm. Semen abortu in nucula unicunum ovula abortiva prope verticem gerens. Cotyledones crassæ farinosæ undulato-rumivate.

I. C. vulgaris (Lam. Dict. I, p. 708) arborea elata, ramis patentibus junioribus puberulis cito glabratibus, foliis a basi obtusâ vel subcordatâ oblongo-lanceolatis breviter petiolatis magnis regulariter et acute serratis acuminateis adultis subtus glabralis vel plus minus tomentellis, nucibus magnis brunneis lucidis basi late areolatis albis. **C. vesca** Gaertn. Fr. tab. 3. — Rehb. Ic. Germ. t. 1305. — *Fagus Castanea* L. Sp. 1416.

Hab. sylvatica in regione montana et subalpinâ 2000'-4000' toton Graeciae! Macedoniae et Thraciae! Anatoliae septentr. et occidentalis (Tchih!), Imeretiae et Guriae (Ledebe.), prov. Karabagh (Szov!), ditionis Talyesch (G.A. Mey.), Persiae borealis (ex Ledebe.). Nec in Tauro (Bal!), nec in Libano spontanea videtur.

Ar. Geogr. Europa media et australis, Africa borealis, India borealis, Japonia, America borealis.

FAGUS (Tourn. Inst. p. 584, tab. 354).

Flores masculi in amenta globosa longe pedunculata globosa squamis minutis deciduis involuerata dispositi. Perigonium 5-6-fidum. Stamina 10-15. *Flores feminei* intra involuerum necolatum quadrilobum extus cum bracteis linearibus numerosis coalitum saepius bini. Perigonium ovario adnatum limbo parvo laciniato, ovarium trigonum triloculare loculis bioculatis, styli tres lineares secus rimam internam stigmatosi. Fructus involuero acreto ovoideo quadripartito lignoso nuculisque binis vel abortu solitariis trigonis perigonii limbo coranatis saepius monospermis constans. Pericarpium coriaceum. Cotyledones carnosae intus plieis irregularibus prominentibus instructae et arete coherentes. — Arbores vel frutices.

I. F. sylvatica (L. Sp. 1416) arborea elata, gemmis glabris lucidis, foliis deciduis in aestivatione secus nervos laterales plicatis a basi obtusâ vel subcordatâ ovatis vel oblongis acutis vel acuminateis plus minus manifeste sinuato-denticulatis junioribus margine et subtus ad nervos ciliato-sericeis, fructu fulvo subsericeo-tomentoso trigono laciniis linearibus erectis vel patentibus paucis basi reflexis. Ic. Fl. Dan. tab. 1284. — Rehb. Germ. t. 1304

Hab. sylvatica in regione montana, Graecia septentrionalis in montibus Pindo et Pelione (Fraas), Macedonia, Thracia et Bithynia 1500-4600' (Griseb! Orph!), Phrygia supra Gnediz (Bal!), Anatolia borealis (Tchih!), Pontus Lazicus (Bal!), Tauria, Caucasus et Transcaucasia (Ledebe! Rupr!), ditionis Talyesch (G.A. Mey.), Persia bor. in prov. Ghilan (Auch. 5325!) et Mazanderan (Bunge!).

Ar. Geogr. Europa media a Norvegia australi et Daniâ ad Rossiam mediam et Hispaniam boreali ad Italianam, Dalmatiam et regionem Danubialem, Japoniam

CORYLUS (Tourn. Inst. p. 581, tab. 347).

Flores masculi in amenta cylindrica compacta bracteata dispositi. Bracteæ imbricatæ cuneiformes, unaquaque bracteolas perigoniales binas interne supra basin fœvens cum eis basi coalita. Stamina 8 secus eristam brevem ad basin bracteolarum suturæ inserta, filamenta brevia, antheræ uniloculares sœpius apice barbatæ. *Flores feminei* in geminam paucifloram squamosam denum in ramulum exerescentem dispositi, squamæ imbricatæ inferiores steriles superiores bifloræ, flos unusquisque squamulis involucralibus minutis laceris villosis basi cohaerentibus cinctus. Perigonium ovario connatum limbo minimo denticalato. Ovarium biloculare loculis uniovulatis, stigmata bina elongata linearia. Nuentæ solitariæ vel 2-3 basi aggregatæ involucris valde aceretis foliaceis tubulosis apice laceris cinctæ uniloculares. Pericarpium lignosum. Cotyledones carnosæ obovatæ plano-convexæ. — Frutices.

1. **C. Avellana** (L. Sp. 1447) fruticosa, stipulis oblongis obtusis, foliis breviter petiolatis cordato-suborbiculatis duplicatim dentatis breviter acuminatis subtus pubescentibus, involucro uniseriato campanulato aperto apice patulo lacero-dentato nuce subbreviore, nuce ovoidicâ ȝ. Ic. Rehb. Germ. tab. 1300.

Hab. in regione montanâ et subalpinâ, Arcadia (Orph!), montes Thessaliæ, Macedonia, Thracia, Bithyniae (Sibth., Griseb., Orph. Fl. exs. 800 sub *C. Colurna*), Paphlagonia et Pontus (Tchih.), Pontus Lazicus in valle Khabakkar 5'000' (Bal!). Tauria, Caucasus et Transcaucasia (M.B., Ledeb.), ditio Talyssch (C.A. Mey.), Cilicia (Ky!), Syria bor. ad Aintab (Haussk!).

Ar. Geogr. Europa omnis a Scandinaviâ et Scotiâ ad Rossiam septentrionali, Africa borealis.

2. **C. Colurna** (L. Sp. p. 1417) fruticosa, stipulis lanceolatis acuminatis, foliis petiolatis cordatis ovatis vel obovatis acutis vel acuminatis duplicato-dentatis subtus ad nervos velutinis, antheris nudis, involuci biseriati phyllis externis ovato-lanceolatis laceris interioribus nucem superantibus multifidis lacinias longe acuminatis, nuce subreniformi ȝ. Ic. Rehb. Germ. t. 1303. — *C. Byzantina* Clus. Hist. I, p. 11. — *C. Pontica* C. Koch Linn. XXII, p. 229. — *C. Jacquemontii* Decaisne ex D.C.

Hab. in regione inferiore et montanâ, Macedonia et Thracia (Griseb.), Byzantii (Sibth.), mons Bulghasdaghi Phrygiae 4'000' (Bal!). Pontus (C. Koch).

Ar. Geogr. Banatus, India borealis.

3. **C. tubulosa** (Willd. Sp. IV, p. 470) fruticosa, stipulis oblongis oblongis, foliis ovatis cordatis acuminatis duplicatim serrato-dentatis subtus ad nervos plus minus puberulis, involucro uniseriato tubuloso-cylindraceo ad apicem nucis coarctato inciso-dentato ȝ. Ic. Rehb. Germ. t. 1301.

Hab. in fruticetis peninsulae Athoæ in Macedonia 1200'-2000' (Griseb.), Turacie (ex Bergg.), Ponti Lazici prope Rhizé (Bal!).

Ar. Geogr. Istria, Banatus.

CARPINUS (Tourn. Inst. I, p. 582, tab. 348).

Flores masculi in amenta cylindrica bracteata dispositi, bracteæ simplices ovatae. Stamina 6-12 in axillâ bracteæ inserta, antheræ disjunctim biloculares. *Flores feminei* amentacei squamis ternis bifloris exteriore deciduâ, interioribus binis involucralibus acercentibus indivisis vel trilobis unaquaque florem fovens. Ovarium perigonio ahaerens ejus limbo minute sexdentato coronatum bilocularis loculis uniovularis. Stigma bina filiformia. Nucula abortu unilocularis involucro foliaceo inclusa longitudinaliter costata. Pericarpium lignosum. Cotyledones carnosæ plano-convexæ. — Arbores vel frutices.

1. C. Betulus (L. Sp. 1416) arborea, foliis breviter petiolatis ovato-oblongis acutis vel acuminatis duplicato-serratis subtus ad nervos laterales crebros prominentes pubescentibus, amentis masculis sessilibus ad partem inferiorem ramorum, bracteis acutis ciliatis, strobilis feminis laxis terminalibus pedunculatis, involueri phyllis tripartitis laciis lanceolatis intermediâ aliis 2-3-plo longiore nuculam longe superante 5. Ic. Sekl. tab. 304. — Nees Germ. Ic. — Rehb. Germ. tab. 1296.

Hab. in sylvis regionis præsertim montanæ et in borealibus etiam inferioris, Peloponnesi montes (Bory), Graecia (Sibth.), Turacia et ad Byzantium (Friv!), Olympus Bithynus (Boiss!), Anatolia borealis (Tchih!), Pontus supra Trapezuntem (Bourgl.) et Rhizé (Bal!), Tauria, Caucasus et Transcaucasia (Ledeb., Rupr!), ditio Talyseh (C.A. Mey.), prov. Karadagh (Szov!), Persia borealis in Ghilan (Auch. 5329!).

Ar. Geogr. Europa sept. et media ab Angliâ, Gothiâ et Daniâ ad Rossiam medium, Galliam, Italiam australiæ, Dalmatiæ et regionem Danubialem.

2. C. Duinensis (Scop. Carn. II, p. 243, tab. 60) fruticosa, foliis breviter ovatis vel ellipticis acutis duplicato-serratis subtus ad nervos secundarios crebros pubescentibus, amentis masculis et feminis terminalibus illis sessilibus, strobilis feminis pedunculatis brevibus densisculis, involueri fructiferi phyllis dimidiato-ovatis indiuisis inæqualiter serratis nuculam parvam superantibus 5. *C. Orientalis* Lam. Dict. I, p. 707. — Rehb. Germ. Ic. 1298.

Hab. in regione montana et subalpinâ Arcadiae 3-4000' (Heldr! Orph. Fl. Gr. exs. 103!), Thessaliæ in Olympo (Heldr!), Macedonia australis et Turacie (Griseb., Friv!), Mysia, Bithynia et Anatolia bor. (Tchih!), Ponti ad Trapezuntem (Bourgl.), Pamphyliae montibus (Bourgr!), Cilicia in Tauri (Ky!), Cappadociâ (Ky!), Tauriæ, Caucaso, Transcaucasiâ, prov. Talyseh (Stev! Rehb! Hoh! Rupr!), Persia bor. in prov. Ghilan (Auch. 5328!) et prov. Astabad (Karell!).

Nuces eis speciei præcedentis triplo minores.

Ar. Geogr. Italia borealis et media, Sicilia, Istria, Dalmatia, regio Danubialis.

OSTRYA (Micheli Gen. p. 223, tab. 104).

Flores masculi in amenta cylindrica bracteata dispositi, bracteæ simplices ovatae. Stamina 6-12 in axillâ bracteæ inserta, antheræ loculis binis filamento fisco disjunctis. *Flores feminei* in amentum strobiliformem densum dispositi ad bracteæ caducæ axillam bini. Involucrum urceolatum membranaceum demum accrescens nuculam includens apice occlusum. Ovarium cum perigonio coalitum bilocularē biovulatum tubulo brevissimo ciliato perigonali coronatum. Stigmata bina filiformia. Nacula utriculo inclusa lœvis abortu monosperma testacea. Cotyledones carnosæ plano-convexæ. — Frutices elati.

1. **O. carpinifolia** (Scop. Carn. II, p. 243) fruticosa, foliis ovatis vel oblongis breviter petiolatis basi rotundatis vel subcordatis acuminatis argute serratis subtus ad nervos laterales utrinque 15-17 pubescentibus, strobilis ovatis pendulis, utriculis albidis ellipticis apice mucronatis nuculâ multoties majoribus 5. *Ostrya vulgaris* Willd. Sp. IV, p. 469. — Rchb. Ic. Germ. f. 1299. — *Carpinus Ostrya* L. Sp. 1417.

Hab. in regione inferiori sed sæpius montanâ, Græcia in montibus Malevo et Taygeto Laconiæ (Heldr! Orph!), monte Olenos et Kyllene (Orph. pl. exs. 286!), Eubœa et Cephaloniâ (Heldr!), Macedonia et Thraciâ (Friv!), Bithyniâ in Olympo (Sith.), Tauro Cilicio (Kyl Bal!). Syriæ borealis monte Annano (Kyl!) et Cassio (Boiss!), monte Akkerdagħ Cataoniae (Haussk!), Armeniâ Turcicâ (ex C. i. h), Transeaucasiâ occidentali (Radde!). Græcis *Ostrua* vel *Gavros*.

Ar. Geogr. Gallia, Cœlio-orientalis, Helvetia australis, Italia, Sicilia, Dalmatia, Carnia, Styria, Serbia.

ORD. CXVIII. BETULACEÆ.

(Endl. Gen. p. 272.)

Flores monoici amentacei. *Masc.* Amenta squamis pedicellatis appendiculatis sèpius trifloris constantia. Perigonum nunc squamiforme monophyllum nunc calyciforme 4-5-partitum vel -phyllum. Stamina 2-4 basi perigonii inserta. *Fem.* Amenta squamis inbricatis 2-3-floris constantia. Perigonium nullum. Ovarium bilocularē loculis uniovulatis, stigmata bina, ovulum paulo infra dissepimenti apicem suspensum. Fructus abortu unilocularis monospermus indehiscentis.

compressus membranaceus vel subcoriaceus sæpe alatus. Testa membranacea, albumen nullum, embryo rectus. — Arbores vel frutices.

ALNUS (Tourn. Inst. I, p. 387, tab. 359).

Flores monoici in amenta dispositi. *Masc.* Amentorum squamæ peltatae versus marginem inferiorem quadrilobulatae subtus sæpius trifloræ. Perigonum quadripartitum vel tetraphyllum. Stamina quatuor filamentis brevibus liberis, antheræ biloculares. *Fem.* Amentorum squamæ ovatae carnosæ ad axillam bifloræ fructiferæ auctæ lignosæ cuneato-ovovatae persistentes intus quadriappendiculatae. Ovarium minutum biloculare, stigmata bina filiformia. Nuculæ compressæ apteræ vel alâ marginatæ abortu uniloculares monospermæ. — Arbores vel frutices floratione vernali.

SECT. GYMNOTHYRSUS Spach. — Amenta foliis præcociora, feminea quoque ex geminis aphyllis prodeuntia. *Masc.* Squamæ trifloræ, perigonum quadripartitum. Nuculæ apteræ vel alâ angustâ coriacea rarius membranacea cinctæ.

1. **A. cordifolia** (Ten. Nap. Prodr. p. 54, tab. 99) arborea medioeris, foliis petiolatis a basi cordatâ ovato-orbiculatis vel ovato-oblongis acutis inæqualiter serrulatis subglutinosis supra glabris subtus ad axillas venarum pubescensibus vel glabratis, amentis feminis in gemmâ 1-3-nis ovato-oblongis pedunculis incurvis eis tenuioribus et 2-3-plo brevioribus suffultis, nuculis alâ angustâ submembranacea cinctis 3. Reg. Mon. tab. 16.

β. subcordata Regel Mon. p. 112. — Folia a basi subcordatâ ovato-oblonga sæpe subacuminata. — *A. subcordata* C.A. Mey. Ennun. p. 43.

Hab. in humidis ad rivulorum margines in regione inferiore et montana usque ad 3000'. Anatolia borealis (C. Koch), ditio Transcaucasica Talyssch (C.A. Mey! Hoh!), Persia borealis in prov. Ghilan (Anch. 532!). Reete a cl. Regel *A. cordifolia* haec varietas in Italiâ quoque obvia admuneratur.

Ar. Geogr. Corsica, Italia australis.

2. **A. Orientalis** (Decaisne Fl. Sinaie, in Ann. Sc. Nat. Ser. II, 4, p. 348) arborea humilis vel fruticosa, foliis longe petiolatis a basi subenneatâ rotundatâ rarius subcordatâ interdum subobliquâ ovato-oblongis vel ellipticis obtusis vel acuminatis irregulariter crenato-dentatis supra glabris subtus ad nervos prominentes interdum puberulis, amentis feminis e gemmâ 1-3-nis ovatis pedunculis æquolongis suffultis, nuculis apteris 3. Reg. Mon. tab. 17. — *A. oblongata* Ky. Sched.

Hab. ad ripas regionis inferioris Cilicie ad Mersina et Cydi ostre (Bal! Ky!), Cypri (Ky!). Syriae borealis ad radices montis Cassii (Boiss!), ad Bery-

thum et Sidonem (Bové! Bl! Gaill! Ky!), Persia boreali ad Astara (Buhsel!), prope Siaret (Bge!).

3. A. glutinosa (Willd. Sp. p. 334) arborea, foliis longiuscule petiolatis cuneato-ovoatis vel subrotundatis obtusissimis utrinque viridibus glabris subtus ad venarum axillas hirtis plus minus manifeste duplicatim crenatis vel denticulatis, amentis feminineis in quavis gemmâ racemosis ovatis pedunculis incurvis tenuibus eis longioribus suffultis, nuculis lenticularibus alâ coriaceâ angustissimâ cinctis 3. Ic. Rehb. Germ. tab. 631. — *Betula Alnus glutinosa* L. Sp. 1394. — *A. denticulata* C.A. Mey. Enum. p. 43 (forma foliis profundiis denticulatis). — *A. barbata* C.A. Mey. loc. cit. (forma foliis subtus ad nervos magis barbatis). — *A. nitens* C. Koch Linn. XXII, p. 334.

Hab. in paludosis et ad fluvios, Graecia omnis! Macedonia, Thracia et Bithynia, Anatolia borealis et Pontus! Lydia in Tmol (Boiss!), Armenia et Karduchia (Kyl!), Tauria, Caucasus et Transcaucasia (Ledeb., Ruprecht!).

Ar. Geogr. Europa fere omnis, Sibiria, Africa borealis.

4. A. incana (Willd. Sp. IV, p. 335) fruticosa, foliis petiolatis ovalis acutis vel subacuminatis argute duplicato-serratis superne glabris glutinosis inferne glancis pubescentibus vel tomentosis, amentis feminineis in quavis gemmâ racemosis ovatis in pedunculo communi crasso subsessilibus, nuculis lenticularibus alâ coriaceâ angustissimâ cinctis 5.

Hab. ad aquas, Caucasus usque ad 6000' (M.B., Hoh., C.A. Mey., ex Ledeb.), circa Kaischaur (Ower!), in Tuschetia et Daghestaniâ 4500'-5500' (Rupr!).

Ar. Geogr. Europa borealis et media a Scandinavia et Russia ad Galliam, Italiam borealem, regionem Danubialem usque, Sibiria, America borealis.

Obs. *Alnus glutinoso-incana* = *A. pubescens* Tausch forma hybrida in Caucaso a cl. Regel indicatur.

BETULA (Tourn. Inst. p. 588, tab. 360).

Flores monoici in amenta dispositi. *Masc.* Squamæ amentorum peltatae subtus utrinque squamulæ acutæ subtus trifloræ. Perigonium monophyllum squamiforme integrum. Stamina duo apice bifida antheris unilocularibus. *Fem.* Squamæ subintegrae saepius trilobæ imbricatae ad axillam trifloræ fructiferæ membranaceæ cum fructu deciduae. Ovarium biloculare, stigmata bina filiformia. Nuculæ minutæ samariformes utrinque alatae abortu uniloculares monospermæ. — Arbores et frutices floratione vernali.

1. B. alba (L. Sp. 1393) arborea, eidermide albâ argenteâ, ramis glabris junioribus gracillimis pendulis, foliis longe petiolatis rhombo-triangularibus acutis crenatis duplicato-serratis glabris subtus ramoso-

venosis, amentis fructiferis longe pedunculatis pendulis, squamarum trilobarum lobis lateralibus recurvatis, fructus alis nuce ellipticâ duplo vel triplo latioribus et latioribus apici stigmatum æquilonigis 5. Rehb. Germ. tab. 623. — *B. verrucosa* Ehrh. Beitr. VI, 96. — *B. alba* subsp. *verrucosa* Regel. Mon. — *B. ovala* C. Koch Linn. XXII, p. 333.

Hab. in alpinis et subalpinis, Thraciae mons Rhodope in regione subalpinâ supra Karlova (Friv.), Pontus Lazicus supra Djimil 6400' (Ball), Armenia Turcica ad radices Bingöldagh (Ky. 4201), Cappadocia in regione alpinâ Argæi (Ball), Caucasus et Transcaucasia (Led., Rupr!).

Ar. Geogr. Europa fere tota Graeciâ exceptâ, Sibiria omnis, Japonia.

2. **B. pubescens** (Ehrh. Beitr. VI, p. 98) arborea humilis vel fruticosa, ramis erectiusculis pubescentibus, foliis petiolatis subcordiacēis ovatis vel rhombēis acutis vel acuminatis duplicato-serratis subtus ramoso-venosis junioribus pubescentibus adultis glabratīs vel subtus ad axillas barbatis, amentis fructiferis longe pedunculatis pendulis, squamarum trilobarum lobis lateralibus patulis vel recurvatis, fructus alis nuci obovatæ subæquilatis apice stigmateatum brevioribus 5. *B. alba* subsp. *pubescens* Regel. Mon. — *B. glutinosa* Wallr. — *B. odorata* Bechst. Rehb. Germ. tab. 626.

Hab. in montibus Somchetiae Caucasicæ (Hoh. ex Ledeb.), in regione alpina montis Kasbek (Rehml.).

Ar. Geogr. Europa borealis et media ad Lusitaniam, Hispaniam, Italiam boreales, regionem Danubialem usque.

ORD. CXIX. SALICINEÆ.

(Rich. Elem. Ed. 4, p. 560).

Flores dioici amentacei, amenta squamis imbricatis constantia. Perigonii loco in squamarum axillâ glandula simplex aut duplex vel urecolus carnosus. *Masc.* Stamina bina vel numerosa; libera vel monadelpha. *Fem.* Ovarium liberum uniloculare multiovulatum placentis binis parietalibus brevibus, ovlis ascendentibus. Stylus unicus interdum nullus; stigma bina. Capsula bivalvis, semina plurima minuta exaluminosa, testa membranacea, comâ bombycinâ ex funculo brevi oriundâ tota involuta. Embryo rectus. — Arberes et fructes stipulatae, stipulis deciduis vel persistentibus.

SALIX (Tourn. Inst. p. 390, tab. 364).

Flores dioici in amenta dispositi basi squamis (bracteis) suffulti. Bracteæ integræ. Perigonii loco glandulae (nectarium) ad genitalium

basin, saepius unica antica interdum binæ altera postica. *Masc.* Stamina bina rarius 3-12, filamenta libera vel (cum gemina) interdum connata, antheræ biloculares. *Fem.* Ovarium sessile vel pedicellatum uniloculare vel phyllis introflexis incomplete bilocularis multiovulatum. Stylus nullus vel plus minus elongatus, stigmata bina integra saepius bifida. Capsula unilocularis bivalvis polysperma valvis post dehiscentiam retrorsum arcuatis. Semina numerosa valvis prope earum basin affixa erecta comosa. Albumen nullum, testa membranacea, radicula infusa. — Arbores vel frutices foliis alternis. — Genus ob formarum hybridarum (in ditione tamen riorum) copiam difficile elucidandum, nonnullas ab auctoribus in Oriente indicatas a me non visas consulto omisi.

Specierum Orientalium distributio.

§ 1. **Fragiles** Koch. — Amenta lateralia coëtanea. Squamæ concolores pallidæ ante capsulæ maturitatem deciduæ. Nectarii glandulæ binæ. Capsulæ glabrae.

* Stamina 5-10.

S. acmophylla, Persica, Sassafras, pentandra.

** Stamina bina rarius terrena.

S. Daviesii, fragilis, alba, Babylonica.

§ 2. **Amygdalinæ** Koch. — Omnia sectionis præcedentis sed bractæ persistentes. Stamina 2-3.

S. triandr.

§ 3. **Purp.** Koch. — Amenta lateralia præcoccia. Squamæ persistentes apice discolores. Glandula nectarifera unica. Filamenta in uniuersum plus minus longe connata. Antheræ purpuræ.

S. purpurea, angustifolia.

§ 4. **Caprææ** (*Caprææ, Prunozæ, Viminales* Koch). — Amenta lateralia præcoccia vel coëtanea. Squamæ apice discolores persistentes. Glandula unica. Stamina bina saepius libera rarius basi coalita. Antheræ flavæ.

* Filamenta inferne coalita.

S. incana, zygostemon.

** Filamenta libera.

+ Stylus subnillus vel brevis.

S. Caprea, cinerea, Silesiaca, pedicellata, aurita, repens.

++ Stylus elongatus.

S. nigricans, daphnoides, viminalis, hastata, arbuscula, apoda.

§ 5. **Glaciales** Koch. — Amenta terminalia. Squamæ apice discolores persistentes.

S. retusa.

§ 1. **Fragiles.**

1. **S. acmophylla** (Boiss. Diagn. Ser. I, 7, p. 98) arborea et fruticosa, ramis elongatis saepius rubellis, amentis rectis oblongo-cylindricis pedunculo brevi foliato insidentibus, squamis deciduis pallidis cucullato-ovatis crispatis pilosis, floribus masculis 4-5-andris, capsulis ovato-oblongis glabris pedicello glandulas 2-3-plo superante suffultis, stigmatibus sessilibus crassis, foliis glaberrimis longe linearilanceolatis acuminatis subtus glaucis, stipulis minimis 5. Ic. And. Mon. tab. 4, fig. 6.

Hab. in Persie australis alpe Kuh Daëna (Ky. 620!) et prope urbem Schiraz (Ky. 323!), inter Sihna et Kermanchah et ditione Bachtiaris (Haussk!), Persiae bor. jugo Elbrus prope Passgala (Ky. 421!) et prope Bibersin (Buhse!), Babyloniam inter Bagdad et Bassora ad Tigrin (Haussk!), Kurdistaniam Assyriacam prope Amadia (Ky. 400!).

Folia 4-6-pollices longa 4-6 linea lata. Foliis et habitu similis *S. Babylonica* a qua floribus polyandris, capsulis pedicellatis crassioribus differt. A *S. Safsaf* foliis longis longe acuminatis, capsula multo brevius pedicellata distincta.

2. **S. Persica** (Boiss. Diagn. Ser. I, 7, p. 99) arborea et fruticosa, ramis longis gracilibus, amentis brevissime foliato-pedunculatis oblongo-cylindricis, squamis pallidis deciduis oblongis acutiusculis basi lanatis apice nudis, floribus hexandris filamentis basi lanatis, capsulis oblongis acutis glabris pedicello nectarium bis superante suffultis, stylo brevi, foliis glaberrimis lanceolato-linearibus subtus subpallidioribus integris 5. Ic. And. Mon. tab. 4, fig. 7.

Hab. ad rivulos montanos Persie australis inter Abuchir et Schiraz (Ky. 401!), ad Schahpur (Haussk!), in Mesopotamiam ad ripas fluviorum Chabur et Sirgan atque prope Terek (Haussk!).

Affinis *S. acmophylle*, folia breviora minus acuminata, amenta breviora, squamae apice glabrae. Accedit quoque *S. Safsaf* a qua foliis integris, floribus hexandris, squamis apice glabris (an satis?) differt. Facies quoque *S. triandra*, folia breviora integra, stamina magis numerosa, capsulae brevius pedicellatae in amenta breviora densius congestæ.

3. **S. Safsaf** (Forsk. Cat. Eg. p. 76) arborea vel fruticosa, ramis rubello-fuscis, amentis rectis pedunculis foliolosis suffultis anguste cylindricis, squamis pallidis deciduis ovato-rotundatis vel oblongis

villosis. staminibus suboctonis filamentis basi villosis, capsulis ovato-globosis confertis glaberrimis pedicellis glandulâ urceolatâ demum quadruplo longioribus sussultis, stylo brevi stigmatibus brevissimis, foliis demum coriaceis elliptico-oblongis vel lanceolatis acuminatis serrulatis utrinquè glabris subtus subpallidioribus, stipulis in ramulis junioribus semicordatis ȝ. *S. subserrata* Willd. Sp. IV, p. 671. — *S. octandra* Sieb. — *S. Ägyptiaca* nonnull. non L. (quæ ex ejus herb. videtur *S. Capreae* forma).

Hab. in Egypto¹ inferiore media et superiore ad ripas Nili (Forsk., Del., Sieb! Boiss! Schweinf! etc.), in oasis Ägyptiacis (Aschers!).

Folia adulta 1¹/₂-2-pollicaria 5-10 lineas lata, amenta mascula pollicem et amplius longa.

Ar. Geogr. Nubia, Abyssinia, Senegalia.

4. ***S. pentandra*** (L. Sp. p. 1442) arborea elata rarius fruticosa, ramis junioribus glutinosis, amentis longe foliato-pedunculatis nutantipatulis densifloris obtusis, squamis pallide ovatis obovatis vel ovatis saltem apice glabris caducis, staminibus 5-10-nis, rachide glabrâ, filamentis basi tantum hirsutis, capsulis glaberrimis ex basi ovalâ attenuatis pedicello nectarium bis superante sussultis, stylo brevi, stigmatibus crass'usculis bilobis patulis, foliis junioribus glutinosis adultis glaberrimis lucidis ovatis et ovato-oblongis acuminatis glanduloso-serrulatis, stipulis ovato-oblongis, petiolo superne glanduligerō ȝ. Ic. Rehb. Germ. fig. 1268.

Hab. in humidis, prata alpina Ponti Lazici prope Djimil 5800' (Bal!), Transcaucasia occidentalis (Nordm. ex Ledeb.).

Ar. Geogr. Europa præsertim borealis et media ad Pyreneos et Italiam meridionalem usque, Sibiria omnis.

5. ***S. Daviesii*** (Boiss. Diagn. Ser. I, 7, p. 98) arborea, ramis glabris rubellis, amentis masculis rectis breviter cylindricis subsessilibus basi foliolatis, squamis pallidis deciduis late ovato-oblongis crenulatis dorso glaberrimis intus longe pilosis, staminibus binis rarius ternis filamentis imâ basi lanatis cæterum glabris, capsulis.... foliis glabris minute et obsolete serrulatis oblongis et oblongo-lanceolatis breviter acuminatis subtus albidis ȝ.

Hab. ad fontem montis Kuh Barfi prope Schiraz Persiae australis (Ky. 335!), ramus sterilis ex viciniis Gulbar in ditione Bachtiaris (Haussk!) huc quoque spectare videtur.

Ob flores fæmineos et fructus nondum observatos imperfecte nota, facies *S. albae* et *S. pentandrae*, a priori foliis glaberrimis, squamis multo latioribus, a posteriori staminum numero, foliis angustioribus vix denticulatis, etc., distincta.

6. ***S. fragilis*** (L. Sp. 1443) arborea, ramis subrectangule patentibus ad insertionem fragilibus, amentis pedunculo foliato insidentibus patulis vel recurvatis, squamis pallidis deciduis acutiusculis pilis longis basi præsertim hirsutis, staminibus saepius geminis fila-

mentis basi leviter pilosis, amentis femineis laxifloris, capsulis glaberrimis elongato-conicis attenuatis pedicello nectarium 2-3-plo superante suffultis, stylo mediocri, stigmatibus bipartitis divaricatis, foliis lanceolatis oblique acuminatis junioribus saepe subpilosis adultis glaberrimis ascendentibus serrulatis, stipulis semiordatis 3. Ic. Fl. Dan. tab. 2484. — Rehb. Germ. fig. 1264.

Hab. ad ripas et in bunnidis, Arcadia et Messenia (Sibth., Heldr!), Macedonia et Thracia (Griseb!), Byzantium (Sibth.), Bithynia (Schmidt!), Lycaen regio alpina supra Elmalu (Bourg!), Syria circa Aleppo (Ky. 81!), Libanum (Post!), Armenia australis (Tchih! Ky!), Tauria, Caucasus et Transcaeca in (Lebed.).

Ar. Geogr. Europa fere tota a Norvegia et Suecia ad Rossiam septentrionalem, Sibiria Uralensis et Altaica.

7. *S. alba* (L. Sp. 1449) arborea clata, ramis strictis parum fragilibus, amentis breviter foliato-pedunculatis flexuoso-patulis densifloris angustis, squamis flavidis apice glabriusculis deciduis, staminibus geminis, filamentis basi villosis, capsulis glabris ovato-conicis obtusis pedicello nectarium brevissimam subaequante suffultis, stylo subnullo, stigmatibus crassiusculis retusis, foliis lanceolatis recte acuminatis minute glanduloso-serrulatis utrinque plus minus adpresso sericeis, stipulis lanceolatis cito deciduis 3. Ic. Fl. Dan. 2552. — Rehb. Germ. fig. 1263. — *S. mirans* And. Mon. p. 49. — *S. Kassanogluensis* Ky. Sched. — *S. Libanotica* Boiss. MSS. (forma foliis latioribus et amentis masculis longissimi).

Hab. ad ripas in fere tota ditione a Graecia ad Caucasum! et Transcaucasiam (Szov!), Persiam borealem (Buhse!), Kurdistanam Persicam (Haussk!), Syriam littoralem, interiorem, et Libanum (Ehr! Bl! Gaill!), Turkestanum (Lehm!).

Ar. Geogr. Europa tota media et australis a Daniâ et Gothiâ ad Rossiam medium. Africa borealis.

8. *S. Babylonica* (L. Sp. 1473) arborea. ramis tenuibus longissimis arcuato-pendulis flexilibus, amentis breviter pedunculatis foliis pedunculi subsuperatis brevibus gracilibus curvatis, squamis lanceolatis stramineis caducis superne glabris, staminibus geminis, capsulis omnino sessilibus ovato-conicis glaberrimis pallide viridibus, stylo subnullo stigmatibus crassiusculis divaricatis integris, foliis anguste lanceolatis longissime et subobliquè acuminatis serrulatis utrinque glaberrimis, stipulis petiolo brevioribus subulatis 3. Ic. Gmel. tab. 34, fig. 2.

Hab. in provinciis Caucasicis ad Caspium (Led.), Persia boreali (Buhse!), Turkestanâ australi culta et subs spontanea sed ex Japonia verosimiliter oriunda. Specimina quedam foliifera sterilia ex Kurdistanâ et Mesopotamia a eis Kotschy lecta et a eis Anderson ad *S. Babylonica* relata potius ad *S. acmophyllum* referenda videntur.

Ar. Geogr. Japonia.

§ 2. Amygdalinæ

9. S. triandra (L. Sp. 1442) fruticosa rarius arborea, ramis tenuibus brunneis, amentis elongatis tenuibus patulis pedunculis brevibus foliolatis suffultis, squamis flavidis obovatis vel oblongis serius persistentibus pilosis vel apice glabrescentibus, capsulis pedicello glandulas 3 5-plo superante suffultis laxis ovato-conicis obtusis glabris, stylo brevissimo, stigmatibus brevibus recurvatis, foliis glaberrimis oblongis vel lanceolatis serrulatis, stipulis semicordatis ♂. *S. amygdalina* L. Sp. 1443. — Ic. Rehb. Germ. fig. 1256-7. — Fl. Dan. tab. 2558.

Hab. ad ripas et in humidis, Græcia (Sibth.), Bithynia (Griseb.), Taurus Cilicieus (Ky!), Caucasus et Transcaucasia (Ledeb!), Persia borealis (Szovt Buhsei!), montes Zagros Kurdistanæ Persicæ 5-9000' (Haussk!).

Specinina Orientalia plerumque ad formam foliis subtus glaucis spectant.

Ar. Geogr. Europa borealis et media ad Hispaniam borealem et Italiæ mediam usque, Sibiria tota, regio Amurensis.

§ 3. Purpureæ

10. S. purpurea (L. Sp. 1442) fruticosa, ramis vimineis glabris tenuibus, amentis sessilibus crebris interdum suboppositis basi bracteatis cylindricis gracilibus densifloris erectis masculis arcuatibus, filamentis in unicum biantheriferum connatis, antheris purpureo-nigris, squamis obovatis villosis atris, capsulis sessilibus ovato-conicis obtusis tomentosis albidis, stylo nullo, stigmatibus oblongis integris purpureis, foliis brevissime petiolatis lanceolatis parte inferiori attenuatis acuminatis argute serrulatis glabris planis glaucescentibus subtus pallidioribus, stipulis obsoletis ♂. Ic. Rehb. Germ. tab. 582 et 583. — *S. Helix* L. Sp. 1441. — *S. hippophaëfolia* Ledeb. Fl. Ross. III, pag. 601 forma foliis angustioribus acutissime cuspitatis.

Hab. secus ripas, Græcia in Arcadiâ (Orph! et Heldr!), Thracia ad Byzantium (Coun!), Anatolia borealis (Griseb!), Tauria, Caucasus et Transcaucasia (Ledeb., Wittm!), Armenia ad Erzerum (Calv!), Persia borealis (Auch! Ky! Buhse!), Turkestania (Lehm!).

β. purpureo-viminalis. — Filamenta non ad antheras usque coalita parte superiore libera, stylus elongatus, stigmata oblongo-linearia, folia elongata margine subrevoluta subtus pubescentia. — *S. rubra* Huds Engl. — Rehb. Germ. tab. 586. — *S. fissa* Hoffm.

Hab. in provinciis Caucasicis (Ledeb.).

γ. pallida. — Amenta breviter pedunculata, squamæ glabrescentes basi et margine pilosæ pallidæ apice tandem fusco-sphacelatæ, capsulae superne magis angustatae. — *S. pallida* Led. Fl. Alt. IV, p. 261. — Icon. tab. 454. — *S. Ledebouriana* Trautv. Mém. Petersb. III,

p. 631. — *S. tenuijulis* Ledeb. Fl. Alt. IV, p. 262. — Icon tab. 453. — *S. Elbursensis* Boiss. Diagn. Ser. I, 7, p. 117.

Hab. in Persiae borealis monte Elbrus prope Derbend secus rivos (Ky. 154!).

Hæc varietas, (an forma hybrida?) in Songariâ, Sibiriâ, regione Himalaica quoque occurrit. Amenta mascula nondum nota.

δ. amplexicaulis. — Sterilis tantum nota, folia sessilia vel subsessilia oblongo-lanceolata vel oblonga basi rotundato-truncata vel cordato-amplexicaulia. — *S. amplexicaulis* Bory et Chaub. Fl. Pelop. № 1586, tab. 36.

Hab. in Graecie Laconiâ secus fluvium Eurotas prope Helos (Bory), in Achaia inter Hagios Olasios et Calavryta et in pago Trikala 3500' (Heldr!).

Salix amplexicaulis Bory planta diu obscura ex cl. Buchinger in Steud. Nomi. ad *Apocynum Venetum* erroneè relata fuit; certissimè *Salicis* species! sed cuidam adnumeranda? Cl. Boutigny qui specimen Boryanum vidit eam *Salicis triandre* formam esse autem, ego autem praeante amico Reuter et omnibus perpensis eam ad *S. purpuream* adducere ratus existino.

Ar. Geogr. Europa media a Sueciâ mediâ ad Rossiam medium, Sibiria, Africa borealis.

11. **S. angustifolia** (Willd. Sp. IV, p. 699 non alior.) fruticosa, ramis tenuibus virgatis crebre et stricte foliosis junioribus sericeis, amentis subsessilibus erectiusculis anguste linearí-cylindricis densifloris, squamis pallide fuscescenti-flavidis lingulato-linearibus basi pilosis cæterum glabris, staminibus in singulom monadelphis antherâ fulvo-aureâ, capsulis sessilibus conicis adpresso sericeis squamas superantibus, stigmatibus sessilibus sanguineis divergentibus, foliis sessilibus rigidis angustissime linearibus acutis basi attenuatis integris vel obsolete calloso-denticulatis margine subitus revolutis utrinque sericeis rariis glabratis ȝ. *S. Withelmiana* M.B. Taur. Cane. III, p. 627. — *S. dracunculifolia* Boiss. Diagn. Ser. I, 7, p. 99.

Hab. ad rivos Armeniae Turcicæ ad Bainut (Bourg!), Tortum (Huet!), Armeniae Rossicæ ad Nackiteewan (Budiset!), Iberia et Georgia Caucasicis (Hoh!), Persiæ boreali in iugo Elbrus (Ky. 544!) australi circa Sebras et in monte Kuh Daïna (Ky. 621!), in ditione Kuh Kilouyeh (Haussk!).

Frutex 4-5-pedalis, folia 4½-3-pollicaria 2 lineas lata, amenta 6-9 lineas longa.

Ar. Geogr. Songaria, Tibetia.

§ 4. Capreæ.

12. **S. incana** (Sehrank Bai. Fl. I, 230) fruticosa, ramis elongatis rubro-fuseis glabris, amentis sessilibus basi bracteatis anguste cylindricis arcuatih, squamis flavido-fulvis oblongo-spathulatis obtusis vel truncatis glabriusculis, filamentis binis inferre connatis, antheris flavis, capsulis pedicello nectarium bis superante suffultis conicoleaneeolatis glaberrimis fuscescentibus, stylo brevi stigmatibus filiformibus bitidis divaricatis, foliis elongatis lanceolato-linearibus acumi-

natis margine revolto denticulatis supra viridibus subtus adpresso tomentoso-canis costâ mediâ fulvescente, stipulis nullis ♂. Ic. Rehb. Germ. tab. 566. — *S. Cyllenea* Boiss. et Orph. MSS.

Hab. in faucibus umbrosis regionis montanæ et subalpinæ, Arcadia in monte Kyllene supra Trikala c. 4000' (Heldr!), supra Flambouritza (Orph. Fl. Gr. exs 3!), Olympus Thessalus (Orph!), mons Rhodope Thracie (Friv!), Anatolia ad Aradach (Wied!).

Ar. Geogr. Hispania, Gallia, Helvetia, Germania, Italia, Dalmatia, regio Danubialis.

13. *S. zygostemon* (Boiss. Diagn. Ser. I, 12, p. 118) arborea vel fruticosa, ramis vitellinis junioribus puberulis, amentis grosse cylindricis brevissime pedunculatis masculis aureo-pilosus, squamis obovato-rotundatis apice nigricantibus villosis, filamentis binis ad medium fere coalitis antheris luteis, capsulis pedicello nectarium duplo superante suffultis cinereo-tomentosis ovato-conicis acutifusculis, stylo mediocri stigmatibus oblongis patentibus, foliis late obovato-lanceolatis acutis leviter crenulato-dentatis primum sericeo-villosis tandem glabratis, stipulis linearis-lanceolatis falcatis glanduloso-dentatis, gemmis glabris ♂.

Hab. ad cataractas Sir Abi Sehir jugi Elbursensis Persiae borealis prope Passgala (Ky. 185!).

Folia 1-2-pollicaria 6-8 lineas lata, amenta mascula villosissima 1½-pollicem, feminea 1½-2½-pollices longa. Ex grege *S. Caprea* et *cinerea*, filamentis ad medium usque connatis ab affinibus distincta. Ulterius observanda, an forsitan forma hybrida?

14. *S. Caprea* (L. Sp. 1448) arborea vel fruticosa, ramis diffusis, amentis sessilibus vel subsessilibus basi minute bracteolatis, squamis oblongis acutiusculis villosis apice nigris, capsulis pedicello nectarium 4-6-plo superante suffultis ex basi ovatâ elongato-lanceolatis tomentosis, stylo subnullo, stigmatibus ovatis bifidis conniventibus, foliis ovatis vel ellipticis vel lanceolato-obovatis abrupte breviter et saepius oblique acuminatis obsolete undulato-crenatis supra glabris subtus tomentosis, stipulis reniformibus, gemmis glabris ♂. Ic. Fl. Dan. 2601. — Rehb. Germ. fig. 2024.

Hab. in sylvaticis montanis, Graecia in Arcadiâ (Sibth.), Macedonia prope Bitolia et in monte Athos (Orph!), Anatolia bor. et Pontus (Tchih.), Pontus Lazicus (Bal!), Tauria, Caucasus et Transcaucasia (Ledeb., Rehm!), Armenia Rossica (Szovt!), Cilicia (Ky. 421!), Libanus (Bl! Post!), Persia bor. (Buhse!) et australis ad Persepolin (Ky. 802!).

Ar. Geogr. Europa tota, Sibiria, regio Amurensis.

15. *S. cinerea* (L. Sp. 1449) fruticosa, ramis rectiusculis tomentellis, amentis basi bracteatis masculis sessilibus brevibus femineis subsessilibus elongatis laxiusculis, squamis obtusiusculis fusco-nigris pilosis, capsulis pedicello nectarium 4-5-plo superante suffultis elongato-conicis acutiusculis cinereo-tomentosis, stylo brevissimo, stigmatibus ovatis bifidis, foliis elliptico vel lanceolato-obovatis acutis vel

breviter acuminatis undulato-serratis cinereo-viridibus supra pubescentibus vel glabratris subtus tomentoso-hirtis, interdum utrinque dense cinereo-canis, stipulis reniformibus, gemmis tomentello-canis ♂. Ic. Fl. Dan. 2601. — Rehb. Germ. fig. 2022. — *S. phlomoides* M.B. Taur. Cauc. II, p. 415.

Hab. in dumosis humidis, Græcia in Arcadiâ!, insulâ Cephallenâ! Parnasso (Heldr!), Macedonia (Griseb!), Byzantii (Coun!), Anatoliâ (Tchih! Wittm!), Armeniâ (Ky!), Tauro Cilicico (Ky! Ball), Caucaso et Transcaucasiâ (M.B., Rehm! Hoh.), Persiâ boreali (Buhse!).

β. Medemii. — Folia latiora ovato-oblonga saepe obtusa, amenta mascula interdum elongata cylindrica, sed haec forma hybrida videtur vel submonstrosa. — *S. Medemii* Boiss. Diagn. Ser. I, 7, p. 100.

Hab. in Persiâ australi ad Gere inter Abuchir et Schiraz (Ky. 413!), Persiâ occident. montibus Bachtiaris (Haussk!), Persiâ boreali orientali ad Siaret (Bge!).

Ar. Geogr. Europa sept. et media a Scandinaviâ et Rossia ad Hispaniam septentrionalem et Italiam medianam, Sibiria, Kamtchatka.

16. **S. Silesiaca** (Willd. Sp. IV, p. 660) fruticosa, amentis sessilibus basi vix bracteatis, squamis apice fuscis longe pilosis, capsulis pedicello nectarium 3-5-plo superante suffultis ex basi ovatâ conico-elongatis glabris vel sericeis, stylo mediocri, stigmatibus crassis bifidjs divaricatis, foliis obovatis vel obovato-ellipticis acuminatis serrulatis adultis utrinque glabris vel subtus subtomentosis nervis prominulis, stipulis reniformi-semicordatis, gemmis glabris ♂. Ic. Rehb. Germ. fig. 2019.

β. Caucasica And. Mon. p. 63. — Amentis breviter pedunculatis foliis parvis suffultis, foliis utrinque glaberrimis, stylo evidenti bifido.

Hab. in prov. Transcaucasiciis occidentalibus (Nordm. ex Ledeb.). Non vidi.

Species *S. Capreae* et *cinereae* affinis ab eis foliis minoribus adultis glabris discedens. Varietas β quam ex descriptione tantum novi a formâ typicâ amentis pedunculatis, capsulis brevius pedicellatis, stylo sublongiore differre videtur. An huic recte adnumerata?

Ar. Geogr. Silesia, Moravia, Carpathi, Bosnia, Transylvania.

17. **S. pedicellata** (Desf. Atl. II, p. 362) arborescens vel fruticosa, ramis junioribus pubescentibus, amentis breviter pedunculatis elongatis laxiusculis, squamis apice nigricantibus oblongis obtusis pilosis, capsulis glabris vel adpresso hirtis pedicello nectarium 4-6-plo superante suffultis ex basi ovatâ conicis elongatis, stylo brevi, stigmatibus bifidis, foliis breviter oblongo-lanceolatis vel obovatis sub-integris subtus reticulato-rugosis tomentosis vel tantum ad nervos hirtis, stipulis semiorbiculatis ♂.

Hab. in Libano ad Abeih (D' Post!) ad Djurd Berkacha (Bl!).

Ex cl. Moris Fl. Sardoa *S. pedicellata* est forma *S. cinereæ*. Huc referto specimen incompleta plantæ Algerensi et præsertim Sicula preter capsulas sericeo-hirtas nec glabras similia, sed capsulae indumentum in speciebus affinibus variare videtur. Planta tamen ulterius et ex speciminibus melioribus

observanda. Ad hanc speciem ex cl. Anders. probab. referenda *S. Ägyptiaca* L. Sp. 1444 et forsitan Sm. Prodr. II, 252.

Ar. Geogr. Hispania australis, Sardinia, [Sicilia, regnum Neapolitanum, Africa borealis.

18. *S. aurita* (L. Sp. 1446) fruticosa, ramis diffusis dense foliatis rufis adultis glabris, amentis sessilibus bracteis suffultis masculis ovatis feminis breviter cylindricis fructiferis subpedunculatis, squamis parce pilosis ferrugineis apice nigricantibus, capsulis pedicello nectarium 3-4-plo superante suffultis tomentosis ex basi ovata elongato-lanceolatis, stylo brevissimo, stigmatibus ovatis retusis, foliis mediocribus obovatis oblique recurvato-mucronatis undulato-crenatis rugosis superne obscure viridibus glabris subtus glaucis tomentosis, stipulis reniformibus, gemmis glabris ♂. Ic. Rehb. Germ. fig. 2020.

Hab. in umbrosis humidis, Macedonia in peninsulâ Chalcidice (Griseb.) Lydia prope Sardes (Bal!), Caucasus occidentalis in regione alpinâ (C.A. Mey.), Transcaucasia ad Lenkoran (Hoh.) et prob. alibi.

Ar. Geogr. Europa borealis et media, in australioribus rarior, Sibiria Altaica.

19. *S. repens* (L. Sp. 1447) fruticulosa humilis, ramis cinereis, amentis subsessilibus fructiferis ovali-cylindricis breviter pedunculatis foliis parvis suffultis, squamis spathulatis sericeo-pilosis fuscescentibus, capsulis pedicello nectarium 2-3-plo superante suffultis ex basi ovata lanceolatis saepius tomentosis, stylo brevi, stigmatibus ovatis bifidis, foliis parvis ovatis vel ellipticis interdum lanceolato-linearibus recurvo-apiculatis margine subrevolutis subintegris supra glabris, subtus interdum utrinque sericeis, stipulis lanceolatis saepе obsoletis ♂. Ic. Fl. Dan. 2489. — Rehb. Germ. fig. 2039.

Hab. in pascuis et paludosis spongiosis, ex ditione hucusque tantum in Anatoliâ boreali ad Samsun (Wiedem. ex Anders.) et in Turkestaniâ prope Samarkand (Lehm. ex Bge) indicata.

Ar. Geogr. Europa borealis et media ad Galliam, Italianam borealem, regionem Danubialem usque, Sibiria tota.

20. *S. nigricans* (Fries Nov. Mant. I, p. 52) arborea vel fruticosa, ramis divergentibus, ramulis incano-tomentosis, amentis sessilibus fructiferis breviter pedunculatis cylindricis patulis laxiusculis, squamis acutis pilosis apice nigricantibus, capsulis pedicello nectario 3-7-plo longiore suffultis ex basi ovata subulatis tomentosis vel glabris, stylo longo, stigmatibus bifidis, foliis ovatis vel obovatis rarior lanceolatis undulato-serratis junioribus pubescentibus adultis superne glabris inferne glancescenti-cinereis pilosulis vel glabratris. stipulis semicordatis ♂. Ic. Fl. Dan. 2553. — Rehb. Germ. fig. 2017. — *S. stylosa* D.C. — *S. stylaris* Ser.

Hab. ditionis dubia civis. specimen ex Syria boreali ad fluvium Nahr Belan montis Amani (Haussk!) hue magno cum dubio retero. Probabilius in Macedonia vel Thracia.

Ar. Geogr. Europa borealis et media ad Galliam, Italianam medium, regionem Danubialem usque, Sibiria, Kamtschatka.

21. *S. daphnoides* (Vill. Dauph. III, 765) arborea vel fruticosa, ramis junioribus cœsiis, amentis sessilibus ebracteatis vel minute bracteatis crassis densifloris albo vel lutescenti-lanatis masculis oblongo-cylindricis femineis cylindricis, squamis acutinensis longissime villosis apicem versus atratis, capsulis ovato-conicis glabris sessilibus, stylo elongato stigmatibus crassiusculis erecto-patulis integris, foliis oblongo-lanceolatis longe acuminatis glanduloso-seratis adultis glabris supra nitentibus subtus glandulis, stipulis semicordatis, gemmis magnis acutis 5. Ic. Rehb. tab. 602. — *S. præcor* Hoppe.

S. acutifolia And. — Folia anguste lanceolata et linearis-lanceolata acuminatissima, stipulae lanceolato-acuminatae. — *S. acutifolia* Willd. Sp. IV, p. 688. — *S. pruinosa* Wendl. — Rehb. Germ. tab. 603.

Hab. in montibus Karatau Turkestanie prope Samarkand (Lehm!). Species in ditione ut videtur rara, typus quoque in Turkestania ex ramis sterilibus a cl. Lehm. lectis crescere videtur.

Ar. Geogr. Scandinavia, Belgium, Gallia, Germania, regio Danubialis, Rossia media, Sibiria.

22. *S. viminalis* (L. Sp. 1448) fruticosa, ramis virgatis strictis elongatis valde toliatis, amentis sessilibus basi bracteatis cylindricis elongatis densifloris erecto-patulis, squamis obovato-oblongis fusco-fulvis apice obscurioribus dorso breviter villosis, stylo elongato stylis filiformibus, capsulis sessilibus tomentosis ex basi ovata lanceolatis, foliis longis anguste lanceolato-linearibus acuminatis supra viridisibus subtus argenteo-nigritibus margine subrevolutis integris, stipulis lanceolato-linearibus brevibus mox cadueis 5. Ic. Rehb. Germ. fig. 1248.

Hab. ad ripas et in dunosis humidis, Gracia et Byzantium (ex Sm. Prodri.), Olympus Bithynus (ex Tchih.), regio alpina Ponti Lazici prope Djimil 6300' (Bal!), Armenia ad radices Bingöldagh (Ky. Suppl. 574!) et prob. alibi.

Ar. Geogr. Europa tota borealis et media sed ubique culta et ideò area incerta, Songaria, Sibiria.

23. *S. hastata* (L. Sp. 1443) fruticosa humilis ramis patulis glabris, amentis pedunculo foliolato suffultis cylindraceis densifloris, squamis spathulatis obtusis fuscis longe albo-hirsutis, capsulis pedicello nectario subduplo longiore suffultis glabris ex basi ovata conico-attenuatis, stylo longo, stigmatibus bifidis, foliis ovatis vel elliptico-lanceolatis glabris crenulato-serrulatis subtus pallidioribus reticulato-venosis, stipulis semicordatis 5. Ic. Rehb. Germ. fig. 2013.

β? Pontica. — Capsulae brevius pedicellatae tomentello-sericea.

Hab. in regione alpinâ Ponti Lazici supra Djimil 6300' (Bal!).

Hujus plantæ specimen unicum femineum capsulis jam dehiscentibus videtur, squamae, folia, habitus cum formâ typicâ convenire videntur a qua differt indumento capsularum brevius pedicellatarum. Ulterius observanda.

Ar. Geogr. Montes Scandinavie, Pyrenæi, montes Hispanie Granatensis, Alpium tractus, Hungarie montes, Rossia arctica, Sibiria, regio Himalaica.

24. *S. arbuscula* (L. Sp. 1445) fruticulosa humilis, ramis divergentibus glabris, amentis sessilibus fructiferis breviter pedunculatis foliis parvis suffultis cylindricis densifloris erecto-patulis, squamis rotundato-ovatis obtusis rufescentibus hirsutis, capsulis subsessilibus vel breviter pedicellatis ovato-conicis tomentellis, stylo elongato, stigmatibus bifidis, foliis parvis glaberrimis breviter oblongo-lanceolatis acutis vel oblongo-ovoatis serrulatis supra viridibus nitidis subtus plus minus glaucis, stipulis obsoletis 5. Ic. Rehb. Germ. tab. 561. — *S. Waldsteiniana* Willd. Sp. IV, 679. — *S. prunifolia* Sm. Rehb. t. 562.

Hab. in alpinis totius Cauca-i 6000'-10000' (C.A. Mey., Rehm!), in Daghestania, Salataviâ, ditione Tindal (Rupr!).

Frutex in demissioribus 2-3-pedalis foliis majoribus 1-1 $\frac{1}{2}$ -pollicem longis, in regione superiori fruticulus depresso facie *S. retusa* foliis 5-8 lineas longis.

Ar. Geogr. Montes Lapponiae, Scandinaviae, Scotiae, Pyrenei, Alpes, Croatia, Transylvania, Rossia arctica, Sibiria Altaica, Dahuria.

25. *S. apoda* (Trautv. Ind. Sem. Hort. Petrop. 1865, p. 37) fruticosa ramis glabris, amentis coetaneis pedunculatis basi bracteatis, squamis atris lanatis, capsulis sessilibus lanceolatis glabris, stylo elongato stigmatibus brevibus integris, foliis ellipticis serrulatis glabris subtus glaucis, stipulis semicordatis 5.

Hab. in Caucaso prope Pari (Radde). Non vidi.

A cl. auctore *S. arbusculae* proxima dicitur, probabiliter hujus forma capsulis glabris.

§ 5. Glaciales.

26. *S. retusa* (L. Sp. 1445) suffruticosa, ramis tortuosis humi prostratis, amentis in ramulis lateralibus folii- et gemmiferis terminalibus erectis ovalis paucifloris, squamis obovato-spathulatis truncatis vel retusis flavidis apice nigricantibus demum glabris, capsulis pedicello nectarium duplo superante suffultis glabris ex basi ovalâ conicis obtusis, stylo brevissimo fulvo, stigmatibus crassis bipartitis, foliis parvis subsessilibus oblongo-ovatis obtusis vel retusis glabris nitidis siccis elevatim et longitudinaliter nervuloso-lineatis 5. Ic. Rehb. Germ. tab. 558.

Hab. in montibus Græciae (Sm. Prodr. II, p. 252). Locus proprius non indicatus sed verosimile est hanc speciem in Græcia borealis vel Macedoniae montibus occurrere.

Ar. Geogr. Scotia, Jurassus, montes Arverniae, Pyrenæi, Alpes, Apennini, montes Hungariae, Transylvaniæ, Croatiae, Bosniæ, Albaniæ.

POPULUS (Tourn. Inst. p. 592, tab. 365).

Flores dioici in amenta sessilia vel pedunculata, pendula dispositi. Squamæ bracteales crenatæ vel incisæ. Perigonum (discus) cupulare oblique truncatum. *Masc.* Stamina 8-12 libera perigonio inclusa,

antheræ biloculares. *Fem.* Ovarium sessile vel pedicellatum perigonio cinctum uniloculare multiovulatum, stylus brevissimus, stigmata bina bifurcata vel laciñiata. Capsula unilocularis bivalvis polysperma. Semina minuta dorso valvarum affixa longe comosa erecta. Albumen nullum. Embryo testâ tenui rugulosâ vestitus, radicula infera. Arbores sœpe proceræ, foliis alternis vernatione involutis, amentis præcocibus.

SECT. I. LEUCE Duby Bot. Gall. p. 427. — Gemmæ (saltem juniores) pubescentes. Ramuli juniores tomentosi vel hirti. Amenti squamæ ciliatae. Perigonium oblique truncatum integrum, stamina 4-8.

1. P. alba (L. Sp. 1463) arborea excelsa, ramis patentibus junioribus albidis, gemmis tomentosis, foliis petiolo longo insidentibus supra viridibus subtus niveo-tomentosis ramorum lateralium ovatis vel ovato-oblongis sinuato-angulosis, terminalium cordatis subrotundato-ovatis angulato-dentatis vel palmatim quinquelobis, amenti feminei squamis lanceolatis apice crenatis ciliatis 3. Engl. bot. tab. 1618. — Rehb. Germ. Ic. 1270. — *P. nivea* Willd.

Hab. in sylvis humidis et ad fossas totius ditionis a Græciâ et Macedonia ad prov. Caucasicas, Syriam littoralem et interiorem (Bl! Gaill!), Persiam bor. (Auch!) et australem (Haussk!).

Formæ inter *P. album* et *P. tremulam* hybridæ, *P. hybrida* M.B. Taur. Cauc. II, p. 422 et *P. canescens* Sm. Brit. passim, v. gr. in Syriâ boreali (Haussk!), Caucaso (M.B.), ditione Talyssch (C.A. Mey!), occurunt. — *P. Græca* Griseb. Spic. II, p. 345 ex Macedonia est quoque forma hybrida, sed *P. Græca* Ait. est *P. tremuloides* Mich. sp. Americana.

Ar. Geogr. Europa tota media et australis ab Angliâ et Daniâ ad Rossiam medianam, Sibiria Uralensis, Africa borealis.

2. P. tremula (L. Sp. 1464) arborea, ramis patulis glabris, gemmis viscidulis junioribus puberulis, foliis petiolo longo insidentibus ramorum juniorum et inferiorum ovatis acutis superiorum suborbiculatis dentatis junioribus interdum pubescentibus adultis utrinque glabris, amenti squamis digitato-incisis dense villoso-ciliatis 3. Rehb. Germ. f. 1274. — *P. Cilicica* Ky. Sched. — *P. australis* Ten.

Hab. in regione montanâ, Græcia in montibus Malevo et Olympo Thessalo (Orph!), Baotia (Sibth.), Macedonia et Thraciâ in monte Athone (Griseb.), Anatoliâ bor. (Tchih.), Lydia in monte Tmolo (Boiss!), Phrygiâ (Bal!), Armeniâ Kurdiçâ ad Boglan 4500' (Ky. 487!), Tauro Cilicio 4-5000' (Ky! Bal!), Tauriâ, Caucaso et Transcaucasiâ (Led., Rupr.).

Ar. Geogr. Europa borealis et media a Scandinavia et Rossiâ boreali ad Hispaniam borealem, Italiam, Dalmatiam, regionem Danubialem, Sibiria omnis, Davuria, Japonia, Africa borealis.

SECT. II. TURANGA Bge pl. Lehm. p. 498. — Gemmæ pubescentes. Ramuli juniores tenuissime puberulo-velutini. Amenti squamæ laceræ parce ciliatae. Perigonium urceolatum subæquale multidentatum, stamina 25-30.

3. **P. Euphratica** (Oliv. Voy. 3, p. 449, tab. 45-46) arborea, ramis patulis junioribus puberulis, foliis petiolis apice compressis insidentibus crassiulus glaucis glabris ramorum plantæ junioris et inferioribus sterilibus in eâdem arbore anguste linearibns elongatis, vel lanceolatis integris, ramorum superiorum fertilium a basi cuneatâ vel rotundatâ ovato-lanceolatis acutis vel triangularibus, vel rhomboideois vel deltoideois vel orbiculatis sæpe basi subcordatis vel transverse latioribus flabellatis integris aut parte superiori plus minus acute dentatis, capsulis magnis ovoideis ♂. Ic. folior. Wesmaël Monog. t. 10-13. — *P. diversifolia* Schrenk Enum. p. 15. — Trautv. Im. t. 16.

Hab. ad rivos et aquas regionis inferioris, Egyptus in Oasi parvâ (Aschers!), Palestina ad Jordanum non procul a Jericho (Bové! Boiss!), Syria borealis ad fluv. Aksu prope Marasch (Haussk!), Mesopotamia et Assyria (Ky. 401!), Babylonia ad Euphratem et Tigrin (Oliv! Noë!), Persia occidentalis (Haußk!), australis inter Abuchir et Schiraz (Ky. 233!), borealis in prov. Ghilan (Micheaux), prope Mendjil et Schahrud (Buhse!) et orientalis in prov. Khorassan (Bge!), Turkestania (Lehm!), regnum Cabulicum (Griff!), Belutschia (Stocks!). Persis Terängut vel Patta, Kliurguisis Turanga, Arabis Sâ'saf.

Species mire heterophylla, folia ramorum juniorum sæpe 3-4-pollices longa lineas 2-3 lata illis *S. hippophaefolia* similia, cetera sensim omnes formas a lanceolatâ ad transverse flabellatam præbentia. Capsula interdum trivalvis.

Ar. Geogr. amplissima, Africa borealis interior a prov. Oran, Sibiria Altaica, regio Scinde Indiæ, regio Himalaica.

SECT. III. AIGEIROS Duby loc. cit. — Gemmæ et ramuli glabri visidi. Amenti squamæ glabræ. Perigonum integerrimum. Stamina 12-30.

4. **P. nigra** (L. Sp. 1464) arborea elata, ramis patentibus, foliis petiolo longo apice compresso suffultis glabris nitidis concoloribus deltoideo vel rhomboideo-triangularibus longitudine latioribus minute serrulatis acuminatis ♂. Ic. Rchb. Germ. fig. 1275.

Hab. ad aquas regionis inferioris et montanæ fere in totâ ditione a Græciâ (Sibth.), ad Macedoniam et Thraciam (Orph., Griseb.), Tauriam, Caucasum et Transcaucasiam (Ledeb.), Persiam borealem (Szovits!), Syriam circa Damascum (Gailli).

Ar. Geogr. Europa media et australis ab Angliâ et Daniâ ad Rossiam mediam, Sibiria Uralensis et Altaica, Africa borealis.

5. **P. pyramidalis** (Rozier in Lam. Dict. 5, p. 235) arborea elata, ramis et ramulis erectis pyramidatis, foliis petiolo longo apice compresso suffultis glabris nitidis concoloribus rhombeis æquilongis et latis acuminatis basi truncatis denticulatis ♂. *P. fastigiata* Poir. Dict. — *P. dilatata* Ait. Kew (nomen ambiguum). — *P. Italica* Auct.

Hab. ad vias sæpe culta, ex Oriente ut dicitur, indigena, Macedonia (Griseb.), Anatolia omnis et Armenia (ex Tchih.), in vallibus Antilibani et Cœlesyriæ frequentissima (Boiss!) sed an spontanea ?, valles montis Karatau Turkestania (Lehm. ex Bge) ubi spontanea videtur. Persis *Ssofidar*.

Ar. Geogr. Ubique in Europâ mediâ et australi culta.

EMENDANDA ET ADDENDA

AD VOLUMEN QUARTUM

Pag. 56. Loco SOLENOSTOMA lege SOLENOSTEMMA.

Pag. 58. Post **Pentatropidem spiralem** adde

2. **P. microphylla** (Roxb. Fl. Ind. II, p. 35 sub *Asclepiade*) glabra volubilis, foliis lanceolatis mucronulatis subcarnosis breviter petiolatis, umbellis subsessilibus paucifloris, pedicellis filiformibus bracteolis setaceis basi stipatis, corollæ laciniis lanceolatis reflexis, gynostegio stipitato, coronae phyllis calcaratis horizontalibus apice acuminatis gynostegio adpresso 3. *P. microphylla* Wight et Arn. Contrib. p. 52. — Illustr. tab. 352.

Hab. ad fontes calidos Ouadi Zeska in littore orientali mari Rubri in Arabiâ petreâ (Kersten!).

Forma foliis brevibus ei; Iconis citatae angustioribus.

Ar. Geogr. India.

Pag. 63. Post **Leptadeniam pyrotechnicam** adde

2. **L. heterophylla** (Delil. in Caiet. Centur. p. 47, tab. 63 sub *Cynancho*) fruticosa foliosa, foliis petiolatis glabris infinitis ovato-deltoides subcordatis superioribus a basi cuneatâ oblongo-ellipticis acutis, umbellis 4-5-floris pedunculis petiolo brevioribus suffultis, pedicellis gracilibus flore triplo longioribus, corollæ laciniis introrsum tenuiter puberulis, fancis squamulis depresso brevibus, folliculis oblongis 3. *L. Delilei* Decaisne in n.c. Prodr. 8, p. 628.

Hab. in insulâ Philæ Egypti superioris (Ehrenb. ex Ascherson in litt.). Non vidi.

Ar. Geogr. Nubia et Abyssinia.

Ad pag. 64.

ORD. LXXIX bis LOGANIACEÆ.

(Endl. Gen. 574.)

Flores regulares hermaphroditi. Calyx inferus tubo campanulato saepius brevissimo, lobis 4-5 imbricatis. Corolla gamopetala tubulosa vel infundibuliformis limbi lobis 4-5 valvatis imbricatis vel contortis. Stamina tot quot corollæ lobis eis alterna fanci affixa. Ovarium superum bi rarissime 3-5-loculare. Stylus simplex vel bifidus. Ovula numerosa vel in loculis 1-2 anatropa vel amphitropa. Placentæ medio septo vel axi adnatae. Fructus capsularis vel baccatus. Semina albuminosa, embryo rectus saepius in axi albuminis. — Herbæ, frutices vel arbores, foliis oppositis lineâ vel membranâ transversâ connexis. — Ordo a *Rubiaceis* quibus valde affinis fere tantum ovario supero discedens, a *Serophulariaceis* quibus *Buddleia* olim associabatur floribus regularibus, staminibus corollæ lobis isomeris, petiolis membranâ connexis etc. distinctus.

BUDDLEIA (L. Gen. 57).

Calyx quadridentatus vel semi-quadrifidus. Corollæ tubus brevis vel elongatus, limbus quadrifidus. Stamina quatuor inclina vel ad fancem sessilia. Stylus integer apice clavatus capitatus vel bilobus. Capsula septicide bivalvis, valvis bifidis vel subintegris. Semina numerosa parva testâ laxiusculâ saepè in alam membranaceam expansâ. — Herbæ, frutices, vel arbores.

1. B. crispa (Benth. Scroph. Ind. p. 43) fruticosa tomento denso cana, foliis breviter petiolatis ovato-oblongis inferioribus subcordatis inciso-crenatis vel crispis, superioribus dinnutis subintegris, capitulis densis breviter pedunculatis in paniculas oblongas vel subracemosas dispositis, corollæ tubo calycem campanulatum subdupo superante 5.

Hab. in regno Cabulico prope Olipore et alibi (Griff!), in Belutschia ad Chehel Tun (Stocks!).

Affinis *B. paniculata* Wall. differt foliis inciso-crispis, capitulis paucifloris.

Ar. Geogr. Regio Himalaica, ditio Silhet Indiae borealis.

- Pag. 85. **Convolvulus** ad § 8, **Scandentes** adde « Ovarium glabrum. »
- Pag. 102. Ad **C. glomeratum** adde
Hab. in Palestinā ad Daunieh (Dr Kersten!).
- Pag. 103. Ad **C pilosellæfolium** adde
Hab. in magnā Oasi Egypti ad Djeumah (Schweinf!).
- Pag. 162. Ad **Anchusam Cesatianam**. Haec species ex autopsiā specimenis typici in herbario Vindobonensi est *Pulmonaria mollis* Wulf.!
- Pag. 178. Post **Podonosma Syriacum** adde
2. P. Galalense (Schweinf. in Sched.) totum glandulosum et setis basi tuberculatis sebroidum, ramis ex rhizomate indurato brevibus crebre foliosis, foliis sessilibus breviter oblongo-linearibus obtusis, bracteis foliosis demum laxis, bracteis lanceolatis pedicellisque calyce longioribus, calycis lacinias lanceolatis obtusis dimidiā corollā brevioribus, corollæ glabrae albidae lobis breviter lanceolatis demum reflexis, antheris ultra medium exsertis, nectario hirsuto ♀.
- Hab. in speluncis vallis Ouadi Natfe deserti Egyptiaco-Arabici (Schw!).
- Planta 4-5-pollicaris, differt a *P. Syriaco* foliis parvis 3 lineas longis lineam latis, calycis duplo brevioris (vix 2 lineas longi) lacinias latiusculis obtusis; nuculas non vidi.
- Pag. 203. Ad **Onosma Orientale** adde
Ex herbarii Linnean inspectione est *O. stellulati* W. K. forma, sed Linnei descriptio in Amoen. Acad. 4, p. 267 non omnino convenit.
- Pag. 242. Ad **Myosotidem involucratam** adde synonymon *Trigonocaryum prostratum* Trantv. Aliq. sp. novae N° 14.
- Pag. 258. Ad **Paracaryum leptophyllum** adde Syn. *P. laxiflorum* Trantv. Aliq. sp. novae 6, p. 8, montes Palantenken distr. Erzerum (Radde!).
- Pag. 402. Ad **Scrophulariam Michonianam** lege (Gosson et Kralik).
- Pag. 442. Ad **Veronicam multifidam** dele synonymon *V. purifolia* Vahl.
- Pag. 545. **Mentha Palegium** lege **Pulegium**.
- Pag. 867. Ad **Staticem obovatam** loco « Haussknecht » lege « Hansen. »
- Pag. 893. **P. polyatochya** lege **P. polystachya**.

- Pag. 923. Ad **Kochiam** loco « *spicæformes* » lege « *spinæformes*. »
- Pag. 1059. Ad **Comandram** eleganter adde « Mons Parnes Atticæ prope Dekeleiam. (Tatoi) (Holzmann!).
- Pag. 1135. Ad **Euphorbiæ denticulata** loco « Lehm.
Dict. » lege « Lamarck Dict. »
- Pag. 1167. Ad **Quercum Lusitanicum** pro « *Q. tenuicula* Ky. »
lege *Q. Tauricola* Ky.



ADDENDA AD VOLUMEN PRIMUM

Pag. 98.

ORD. I bis MENISPERMACEÆ.

(D.C.)

Flores dioici vel polygami parvi. Sepala sæpius sex biseriata, rarius 9-12 tri vel quadrilaterata, in unaquaque serie imbricata. Petala sæpius sex biseriatim imbricata. *Masc.* Stamina petalis numero æqualia et opposita, filamenta libera vel coalita, antheræ biloculares sæpius adnatæ extrorsæ. *Fem.* Carpella tria raris plura libera toro brevi imposita monostyla. Ovula in carpellis solitaria amphitropa suturæ ventrali affixa. Drupæ sessiles vel stipitatae monospermae oblique vel lunulatae. Albumen varium, embryo incurvus.

COCCULUS (D.C. Syst. I, 545).

Sepala sex biseriata, interiora majora. Petala sex sepalis breviora. Stamina sex libera antheris quadrilobis. Carpella tria stylis erectis vel recurvis indivisis. Drupæ a latere compressæ styli cicatrice a basi parum remotâ. Putamen dorso tuberculatum lateribus profunde excavatis, semen hippocrepiforme, albumen pareum, radicula brevis, cotyledones lineares planæ. — Frutices sarmentosi scandentes vel subscandentes.

1. C. Leœba (Del. Fl. Eg. III. 30, tab. 54 sub *Menispermo*) glaucus puberulus, ramis tenuibus elongatis subvolubilibus, foliis ovato-oblongis obtusis breviter petiolatis, cymis pauci-floris breviter pedunculatis axillaribus.

Hab. in desertis ad rupium parietes, Egyptus media et superior (Del., Ehr! Anch. 35! Ky. 602!), in jugo Sinaitico Arabiae Petreæ (Ehrenb.).

Ar. Geogr. Nubia, Abyssinia, Senegal, Arabia tropica.

Pag. 757.

ORD. XLVII bis. CERATOPHYLLEÆ.

(Gray Brit. II, p. 554.)

Flores monoici. Perigonium multipartitum. *Masc.* Involuci phylla linearia apice 2-3-dentata. Stamina 12-15 antheris subsessilibus oblongo-ovoatis apice emarginato bispinulosis bilocularibus loculis dissepimento incompleto semibifidis. *Fem.* Perigonii phylla linearia indivisa. Ovarium liberum ovatum uniloculare uniovulatum, ovulum pendulum, stylus rectus stigmate subulato. Nux monosperma indehiscentis coriacea stylo apiculata et sepe basi aculeata aculeis exercentibus. Embryo antitropus, radicula supera. cotyledonibus 4 verticillatis duobus latioribus, plumulâ polyphyllâ. Albumen nullum. — Ordo nulli proprius affinis, inter Dicotyledoneas ad *Podostemoneas* et *Callitrichineas*, inter Monocotyledoneas ad *Naiadeas* ascendens.

CERATOPHYLLUM (L. Gen. 1035).

Characteres ordinis. Plantæ aquatice immersæ ramosissimæ rigidulæ, caule ramisque nodoso-articulatis, foliis sessilibus verticillatis 2-3-tonne in laciniæ lineares integras vel incisas multifidis, floribus axillaribus solitariis sessilibus.

1. **C. demersum** (L. Sp. 1409) foliis dichotome in laciniæ filiformi-setaceas divisis, fructibus ovalibus apteris trispinosis spinis ad basin binis recurvatis, terminali fructum æquante vel superante ♀. Ic. Schleid. Linn. II, tab. 11. — Nees Gen. Germ.

Hab. in aquis stagnantibus, Thracia prope Philippopolin (Friv.), prov. Caucasæ (C.A. Mey.), Egyptus, Affghania (Griff.) et prob. alibi in ditione.

Ar. Geogr. Europa tota, Sibiria, Africa borealis.



INDEX

INDEX

ORDINUM, GENERUM, SPECIERUM ET SYNONYMORUM

	PAG.		PAG.
ACANTHACEÆ	518		
<i>Acanthodium spicatum</i> , Del.	520	<i>Acantholimon diapensioides</i> ,	
Acantholimon , Boiss.	825	<i>Boiss.</i>	830
<i>acerosum</i> , W.	857	<i>distachynum</i> , <i>Boiss.</i>	828
<i>acmostegium</i> , <i>B. et Buhse</i>	826	<i>Echinus</i> , <i>L.</i>	840
<i>androsaceum</i> , Boiss.	841	<i>Eschkerense</i> , <i>B. et Haussk.</i>	844
<i>Araxanum</i> , <i>Bge.</i>	854	<i>fasciculare</i> , <i>Boiss.</i>	851
<i>aristulatum</i> , <i>Bge.</i>	849	<i>Fausti</i> , <i>Trautv.</i>	842
<i>Armenum</i> , <i>B. et Huet.</i>	859	<i>festuaceum</i> , <i>J. et Sp.</i>	843
<i>Aspadanum</i> , <i>Bge.</i>	845	<i>flexuosum</i> , <i>B. et Haussk.</i>	845
<i>Assyriacum</i> , Boiss.	852	<i>gen stoides</i> , <i>J. et Sp.</i>	846
<i>Atropatanum</i> , <i>Bge.</i>	853	<i>glumaceum</i> , <i>J. et Sp.</i>	855
<i>Auganum</i> , <i>Bge.</i>	857	<i>Griffithianum</i> , <i>Boiss.</i>	829
<i>avenaceum</i> , <i>Bge.</i>	851	<i>Gulistanum</i> , <i>Bge.</i>	855
<i>Balansæ</i> , Boiss.	859	<i>Haussknechtii</i> , <i>Bge.</i>	859
<i>Baltanense</i> , <i>B. et Haussk.</i>	858	<i>Heratense</i> , <i>Bge.</i>	848
<i>Bodeanum</i> , <i>Bge.</i>	856	<i>Hohenackeri</i> , <i>J. et Sp.</i>	854
<i>brachiphyllum</i> , <i>Boiss.</i>	846	<i>horridum</i> , <i>Bge.</i>	851
<i>brachystachyum</i> , <i>B. et Haussk.</i>	844	<i>Huetii</i> , <i>Boiss.</i>	835
<i>bracteatum</i> , <i>Gir.</i>	827	<i>Ieopicum</i> , <i>B. et Heldr.</i>	859
<i>breviscapum</i> , <i>B. et Haussk.</i>	840	<i>incomptum</i> , <i>B. et Buhse</i>	850
<i>bromifolium</i> , <i>B. et Haussk.</i>	829	<i>Karelini</i> , <i>Scz.</i>	847
<i>Cabulicum</i> , <i>Boiss.</i>	828	<i>Kotschy</i> , <i>J. et Sp.</i>	840
<i>Cæsareum</i> , <i>B. et Bal.</i>	857	<i>Kotschy</i> var.	859
<i>Calverti</i> , <i>Boiss.</i>	851	<i>Kurdicum</i> , <i>Bge.</i>	845
<i>caryophyllaceum</i> , <i>Boiss.</i>	858	<i>latifolium</i> , <i>Boiss.</i>	828
<i>Cataonicum</i> , <i>Bge.</i>	856	<i>laxiflorum</i> , <i>Boiss.</i>	852
<i>cephalotes</i> , <i>Boiss.</i>	827	<i>laxiflorum</i> , <i>Boiss.</i> in	
<i>curviflorum</i> , <i>Bge.</i>	847	Bourg.	852
<i>cymosum</i> , <i>Bge.</i>	826	<i>lepturoides</i> , <i>J. et Sp.</i>	842
		<i>leucanthum</i> , <i>Boiss.</i>	850
		<i>Libanotiem</i> , <i>Boiss.</i>	840

	Pag.		Pag.
<i>Acantholimon Listoniiæ</i> , Boiss.	858	<i>Acanthus Syriacus</i> , Boiss.	522
<i>longiflorum</i> , Boiss.	849	Achyranthes , L.	993
<i>Lycaonicum</i> , <i>B. et Haussk.</i>	856	<i>alopécuroïdes</i> , Lam.	995
<i>melananthum</i> , Boiss.	845	<i>alternifolia</i> , L.	994
<i>oliganthum</i> , Boiss.	844	<i>argentea</i> , Lam.	994
<i>Peronini</i> , Boiss.	842	<i>aspera</i> , L.	993
<i>petraeum</i> , <i>B. et Haussk.</i>	852	<i>echinata</i> , Retz.	995
<i>Phrygium</i> , Boiss.	858	<i>fruticosa</i> , Hort.	994
<i>Pinardi</i> , Boiss.	858	<i>lanata</i> , L.	993
<i>polystachyum</i> , Boiss.	847	<i>lappacea</i> , L.	995
<i>pterostegium</i> , Bge.	826	<i>papposa</i> , Forsk.	918
<i>puberulum</i> , <i>B. et Bal.</i>	841	<i>villosa</i> , Forsk.	995
<i>quinquelobum</i> , Bge.	848	<i>Acinos acuminatus</i> , Friv.	582
<i>restiaceum</i> , Bge.	850	<i>Alpinus</i> , Mœnch	582
<i>roseum</i> , Boiss.	853	<i>ascendens</i> , Mœnch	582
<i>Sahendicum</i> , <i>B. et Buhse</i>	855	<i>erectus</i> , Friv.	585
<i>scabrellum</i> , <i>B. et Haussk.</i>	845	<i>icanus</i> , Griseb.	585
<i>Schahrudicum</i> , Bge.	855	<i>multiflorus</i> , C. Koch	579
<i>Schirazianum</i> , Boiss.	852	<i>rotundifolius</i> , Friv.	582
<i>scirpicum</i> , Bge.	829	<i>thymoides</i> , M.	585
<i>Scorpius</i> , Boiss.	850	Aerva , Forsk.	992
<i>Senganense</i> , Bge.	859	<i>Bovei</i> , Edg.	993
<i>splendidum</i> , Bge.	827	<i>Javanica</i> , Juss.	992
<i>Stocksii</i> , Boiss.	850	<i>lanata</i> , L.	993
<i>subulatum</i> , Boiss.	845	<i>tomentosa</i> , Forsk.	993
<i>Szovitsii</i> , <i>B. et Buhse</i>	847	<i>Agathophora alopecuroides</i> ,	985
<i>Talagonicum</i> , Boiss.	852	Fenzl	985
<i>Tataricum</i> , Boiss.	829	Agriophyllum , M. B.	928
<i>Tchihatchewii</i> , F. et M.	838	<i>arenarium</i> , M. B.	928
<i>tenuiflorum</i> , Boiss.	854	<i>lateriflorum</i> , Lam.	928
<i>tomentellum</i> , Boiss.	852	<i>latifolium</i> , F. et M.	929
<i>tragacanthinum</i> , <i>J. et Sp.</i>	849	<i>minus</i> , F. et M.	928
<i>truncatum</i> , Bge.	848	<i>squarrosum</i> , Moq.	928
<i>venustum</i> , Boiss.	831	<i>Tournefortii</i> , Fisch.	928
<i>viscidulum</i> , Boiss.	851	Ajuga , L.	798
<i>Wiedemannii</i> , Bge.	841	<i>bombycina</i> , Boiss.	805
Acanthus , L.	520	<i>bracteosa</i> , Wall.	799
<i>Dioscoridis</i> , L.	521	<i>Chamæpitys</i> , L.	802
<i>edulis</i> , Forsk.	520	<i>Chamæcistus</i> , Ging.	801
<i>hirsutus</i> , Boiss.	521	<i>Chia</i> , Poir.	802
<i>mollis</i> , L.	521	<i>euphrasioïdes</i> , Boiss.	801
<i>Raddei</i> , Trautv.	521	<i>Genevensis</i> , L.	799
<i>spinosisimus</i> , Desf.	525	<i>glabra</i> , Bth.	804
<i>spinosus</i> , Nees	522	<i>Iva</i> , L.	802
<i>spinosus</i> , L.	522	<i>intermedia</i> , B. et Orph.	802

	Pag.		Pag.
<i>Ajuga levigata</i> , Russ.	804	<i>Alkanna nonneiformis</i> , Gr.	224
<i>Laxmanni</i> , L.	804	<i>Orientalis</i> , L.	227
<i>Mesogitana</i> , Boiss.	805	<i>Pinardi</i> , Boiss.	229
<i>oblongata</i> , M. B.	801	<i>primuliflora</i> , Gr.	229
<i>Orientalis</i> , L.	800	<i>Pulmonaria</i> , Gr.	224
<i>Palestina</i> , Boiss.	805	<i>rostellata</i> , Boiss.	235
<i>pseudo-iva</i> , D.C.	802	<i>Sartoriana</i> , B. et Heldr.	229
<i>pyramidalis</i> , L.	799	<i>Scardica</i> , Griseb.	224
<i>reptans</i> , L.	798	<i>Sieberi</i> , A.D.C.	228
<i>salicifolia</i> , Stev.	801	<i>strigosa</i> , Boiss.	225
<i>salicifolia</i> , L.	800	<i>Syriaca</i> , Boiss.	226
<i>scoparia</i> , Boiss.	801	<i>tinctoria</i> , L.	227
<i>tridactylites</i> , Ging.	805	<i>tomentosa</i> , Griseb.	197
<i>vestita</i> , Boiss.	805	<i>tubulosa</i> , Boiss.	225
<i>Aipyanthus cchioides</i> , Stev.	211	<i>viscidula</i> , Boiss.	252
Albersia , Kth.	991	Alnus , Tourn.	1179
<i>Blitum</i> , Kth	991	<i>barbata</i> , C. A. M.	1180
<i>candata</i> , Jacq.	992	<i>cordifolia</i> , Ten.	1179
<i>deflexa</i> , L.	992	<i>denticulata</i> , C. A. M.	1180
<i>oleracea</i> , L.	991	<i>glutinosa</i> , W.	1180
<i>polygama</i> , L.	991	<i>incana</i> , W.	1180
<i>Alectorolophus grandiflorus</i> ,		<i>nitens</i> , C. Koch	1180
<i>Wallr.</i>	479	<i>oblongata</i> , Ky.	1177
<i>parviflorus</i> , Wallr.	479	<i>Orientalis</i> , Devaisn.	1179
Alkanna , Tausch	253	<i>subcordata</i> , C. A. M.	1177
<i>amplexicaulis</i> , D.C.	250	Alternanthera , Mart.	996
<i>areolata</i> , Boiss.	252	<i>denticulata</i> , Moq.	997
<i>Aucheriana</i> , D.C.	226 et 253	<i>sessilis</i> , L.	996
<i>azurea</i> , Boiss.	253	<i>Amaracus cordifolius</i> , Montb.	
<i>bicolor</i> , Boiss.	250	et Auch.	548
<i>brachysolen</i> , Boiss.	250	<i>Dictamnus</i> , Bth.	547
<i>bracteosa</i> , Boiss.	251	<i>Tournefortii</i> , Bth.	547
<i>Cappadocica</i> , B. et Bal.	226	AMARANTACEÆ.	986
<i>cordifolia</i> , C. Koch	227	Amarantus , L.	988
<i>filaginoides</i> , A.D.C.	250	<i>albus</i> , L.	990
<i>frigida</i> , Boiss.	251	<i>angustifolius</i> , M. B.	990
<i>Galilea</i> , Boiss.	251	<i>Blitum</i> , L.	991
<i>Græca</i> , Boiss. et Sp.	250	<i>Blitum</i> , Moq.	990
<i>hirsentissima</i> , A.D.C.	227	<i>candatus</i> , L.	988
<i>incana</i> , Boiss.	252	<i>chlorostachys</i> , W.	988
<i>Kotschyana</i> , D.C.	205	<i>deflexus</i> , L.	992
<i>macrophylla</i> , B. et Heldr.	235	<i>Gangeticus</i> , L.	990
<i>wacosiphon</i> , B. et Heldr.	225	<i>gracilis</i> , Desf.	992
<i>Mathioli</i> , Tausch	227	<i>Græcizans</i> , L.	990
<i>megacarpa</i> , A.D.C.	226	<i>hypochondriacus</i> , L.	988

	Pag.		Pag.
<i>Amarantus oleraceus</i> , L.	991	Anagallis , L.	6
<i>paniculatus</i> , L.	989	<i>arvensis</i> , L.	6
<i>patulus</i> , Bert.	989	<i>cærulea</i> , Lam.	6
<i>polygamus</i> , L.	991	<i>latifolia</i> , L.	6
<i>prostratus</i> , Balb.	992	<i>phœnicea</i> , Lam.	6
<i>retroflexus</i> , L.	989	<i>tenella</i> , L.	6
<i>sylvestris</i> , Desf.	990	Anarrhinum , Desf.	562
<i>Timeroyi</i> , Jord.	989	<i>ecalcaratum</i> , Fenzl	562
<i>viridis</i> , L.	991	<i>Orientale</i> , Bth.	563
<i>Amblyanthera chrysanthia</i> ,		<i>pubescens</i> , Fres.	565
Fisch.	475	<i>Anblatum Orientale</i> , Tourn.	518
Amphicome , Lindl.	80	Anchusa , L.	150
<i>Emodi</i> , Royle	80	<i>Egyptiaca</i> , L.	159
<i>Amsonia Orientalis</i> , Dec.	47	<i>aggregata</i> , Lehm.	157
Anabasis , L.	968	<i>alpestris</i> , Stev.	164
<i>affinis</i> , Bge.	948	<i>amplexicaulis</i> , Sm.	152
<i>alopecuroides</i> , Moq.	985	<i>angustifolia</i> , D C.	152
<i>Ammodendron</i> , C. A.		<i>angustifolia</i> , Fl. Gr.	154
Mey.	948	<i>arvalis</i> , Rehb.	152
<i>annua</i> , Bge.	971	<i>arvensis</i> , L.	160
<i>aphylla</i> , L.	969	<i>aspera</i> , Boiss.	154
<i>articulata</i> , Forsk.	970	<i>asperrima</i> , Del.	215
<i>brachiata</i> , F. et M.	969	<i>Aucheria</i> , A. D. C.	158
<i>cinerea</i> , Moq.	950	<i>azurea</i> , Rehb.	154
<i>clavata</i> , Gmel.	955	<i>Barrelieri</i> , All.	155
<i>crassa</i> , Moq.	970	<i>Cesatiana</i> , Fenzl	162 et Add.
<i>cretacea</i> , C. A. M.	969	<i>cespitosa</i> , Lam.	156
<i>Ehrenbergii</i> , Schw.	970	<i>clavata</i> , Viv.	153
<i>florida</i> , M. B.	950	<i>deflexa</i> , Lehm.	158
<i>foliosa</i> , Pall.	955	<i>echinata</i> , Lam.	155
<i>glomerata</i> , M. B.	985	<i>echioides</i> , M. B.	211
<i>Hausknechtii</i> , Bge.	969	<i>flava</i> , Forsk.	159
<i>heteroptera</i> , J. et Sp.	967	<i>Gmelini</i> , Led.	154
<i>intermedia</i> , Moq.	970	<i>hispida</i> , Forsk.	158
<i>lutea</i> , Moq.	970	<i>hyalotricha</i> , Boiss	158
<i>macroptera</i> , Moq.	968	<i>hybrida</i> , Ten.	152
<i>multiflora</i> , Moq.	949	<i>ineana</i> , Led.	154
<i>oppositiflora</i> , M.B.	967	<i>Italica</i> , Retz	154
<i>phyllophora</i> , Kar. et		<i>leiosperma</i> , Fl. Pcl.	164
Kir.	970	<i>leptophylla</i> , Led.	154
<i>setifera</i> , Moq.	970	<i>leucomelius</i> , Lehm.	250
<i>Sieversii</i> , W.	972	<i>limbata</i> , Boiss.	156
<i>subulifolia</i> , Sehr.	970	<i>linearifolia</i> , Urv.	154
<i>Tatarica</i> , Pall.	969	<i>Lorentii</i> , A.D.C.	154
<i>Tournefortii</i> , J. et Sp.	965	<i>Intea</i> , M. B.	165

Pag.		Pag.
	<i>Androsace bidentata</i> , C. Koch	15
	<i>cespitosa</i> , Lehm.	15
	<i>channejasme</i> , Host.	14
	<i>dasyphylla</i> , Bge.	14
	<i>elongata</i> , L.	17
	<i>filiformis</i> , Retz	17
	<i>hedrantha</i> , Griseb.	14
	<i>Jacquemontii</i> , Duby	15
	<i>incana</i> , Lam.	15
	<i>intermedia</i> , Ledeb.	16
	<i>Lehmanniana</i> , Spr.	14
	<i>longifolia</i> , C. Koch	16
	<i>longiscapa</i> , C. Koch	14
	<i>macrantha</i> , Boiss. et Huet.	16
	<i>maxima</i> , L.	18
	<i>multiscapa</i> , Duby	15
	<i>nanu</i> , Horn.	17
	<i>odoratissima</i> , Schreb.	14
	<i>Olympica</i> , Boiss.	14
	<i>penicillata</i> , Sch.	15
	<i>pleioscapa</i> , C. Koch	16
	<i>radiata</i> , Lehm.	17
	<i>septentrionalis</i> , L.	17
	<i>valerianoides</i> , Lehm.	16
	<i>villosa</i> , Jacq.	14
	<i>villosa</i> , L.	15
	<i>Wiedemannii</i> , Boiss.	16
	Anopanthus , Endl.	494
	<i>Biebersteinii</i> , Reut.	494
	<i>coccineus</i> , Marsch.	494
	<i>Tournefortii</i> , Reut.	494
	Anthochlamys , Fenzl	951
	<i>polygaloides</i> , F. et M.	951
	Anticharis , Endl.	422
	<i>Arabica</i> , St. et Hochst.	422
	<i>glandulosa</i> , Asch.	425
	<i>linearis</i> , Bth.	425
	Antirrhinum , L.	585
	<i>fugax</i> , Boiss.	584
	<i>glaucum</i> , Wight.	587
	<i>majus</i> , L.	585
	<i>Orontium</i> , L.	585
	<i>parriflorum</i> , Jacq.	375
	<i>pilosum</i> , Fl. Pol.	563
	<i>pterospermum</i> , Rich.	586

Anchusa macrocarpa , Boiss.	15
<i>et Hoh.</i>	154
<i>micrantha</i> , R. et Sch.	158
<i>Milleri</i> , Willd.	159
<i>myosotidiflora</i> , Lehm.	157
<i>neglecta</i> , A.D.C.	156
<i>nonneoides</i> , F. et M.	155
<i>obliqua</i> , Vis.	155
<i>ochroleuca</i> , M. B.	155
<i>officinalis</i> , L.	151
<i>Orientalis</i> , L.	161
<i>Orientalis</i> , L.	220
<i>ovata</i> , Lehm.	161
<i>paniculata</i> , Ait.	154
<i>Parnassica</i> , B. et Orph.	152
<i>parviflora</i> , Fl. Gr.	158
<i>parviflora</i> , W.	162
<i>perlata</i> , Lam.	161
<i>pieta</i> , M. B.	166
<i>Pinardi</i> , Boiss.	152
<i>rhizochroa</i> , Viv.	227
<i>rosea</i> , M. B.	164
<i>spinocarpas</i> , Forsk.	249
<i>Spruneri</i> , Boiss.	160
<i>strigosa</i> , Lab.	155
<i>stylosa</i> , M. B.	159
<i>Thessala</i> , Boiss. et Sp.	160
<i>tinctoria</i> , Desf.	227
<i>tomentosa</i> , Boiss.	154
<i>tuberculata</i> , Forsk.	227
<i>undulata</i> , L.	152
<i>variegata</i> , L.	161
<i>ventricosa</i> , Fl. Gr.	169
<i>ventricosa</i> , Viv.	165
<i>versicolor</i> , Stev.	164
Andrachne , L.	1157
<i>aspera</i> , Spr.	1158
<i>Colechica</i> , F. et M.	1157
<i>fruticulosa</i> , Boiss.	1158
<i>rotundifolia</i> , Mey.	1158
<i>telephioides</i> , L.	1158
Androsace , L.	15
<i>Albana</i> , Stev.	16
<i>arachnoidea</i> , Schott.	15
<i>Armeniaca</i> , Duby.	15

	Pag.		Pag.
Antirrhinum Rhodium , Boiss.	555	Armeria Cariensis , <i>Boiss.</i>	873
<i>rhytidospermum</i> , F. et M.	584	<i>Majellensis</i> , <i>Boiss.</i>	873
<i>spinescens</i> , Viv.	569	<i>Orphanidis</i> , <i>Boiss.</i>	873
<i>strictum</i> , Fl. Gr.	578	<i>Rumelica</i> , <i>Boiss.</i>	874
<i>supinum</i> , Fl. Gr.	578	<i>Thessala</i> , <i>Boiss.</i>	874
<i>tortuosum</i> , <i>Bosc.</i>	586		
APOCYNEÆ.	44	Arnebia , <i>Forsk.</i>	212
Apocynum , L.	48	<i>Bungei</i> , <i>Boiss.</i>	215
<i>Apocynum Syriacum</i> , Clus.	57	<i>calycina</i> , <i>Stev.</i>	214
<i>Venetum</i> , <i>L.</i>	48	<i>Cephalotes</i> , A.D.C.	212
AQUIFOLIACEÆ.	34	<i>cornuta</i> , <i>Led.</i>	213
Arceuthobium , M. B.	1068	<i>densiflora</i> , <i>Led.</i>	212
<i>Oxycedri</i> , <i>D.C.</i>	1069	<i>fimbriopetala</i> , <i>Stocks</i>	215
<i>Aretia longiflora</i> , <i>Fisch.</i>	19	<i>flavescens</i> , <i>Boiss.</i>	214
ARISTOLOCHIACEÆ.	1073	<i>Griffithii</i> , <i>Boiss.</i>	215
Aristolochia , L.	1074	<i>hispidissima</i> , <i>Spr.</i>	213
<i>altissima</i> , <i>Desf.</i>	1075	<i>linearifolia</i> , <i>D.C.</i>	214
<i>Anatolica</i> , <i>Boiss.</i>	1080	<i>longiflora</i> , <i>C. Koch</i>	211
<i>Aucherii</i> , J. et Sp.	1080	<i>ovovata</i> , <i>Bge.</i>	215
<i>auricularia</i> , <i>Boiss.</i>	1079	<i>tinctoria</i> , <i>Forsk.</i>	214
<i>Billardieri</i> , J. et Sp.	1077	<i>tingens</i> , A.D.C.	212
<i>Bruguieri</i> , J. et Sp.	1080	<i>Vivianii</i> , <i>Coss.</i>	214
<i>Clematitis</i> , <i>L.</i>	1077		
<i>Cretica</i> , <i>Lam.</i>	1081	Arthrocnemum , Moq.	952
<i>hirta</i> , <i>L.</i>	1079	<i>Caspicum</i> , <i>Moq.</i>	955
<i>hirta var.</i> , <i>Duch.</i>	1080	<i>fruticosum</i> , <i>Moq.</i>	932 et 955
<i>Iberica</i> , <i>F. et M.</i>	1081	<i>glaucum</i> , <i>Del.</i>	952
<i>incisa</i> , <i>Duch.</i>	1078	<i>macrostachyum</i> , <i>Mor.</i>	932
<i>longa</i> , <i>L.</i>	1078		
<i>lutea</i> , <i>Desf.</i>	1078	Arthropophytum , Schr.	947
<i>macroglossa</i> , J. et Sp.	1077	<i>subulifolium</i> , <i>Schr.</i>	948
<i>Maurorum</i> , <i>L.</i>	1080		
<i>microstoma</i> , <i>Boiss.</i>	1075	Asarum , L.	1073
<i>Olivieri</i> , <i>Coll.</i>	1079	<i>Europaeum</i> , <i>L.</i>	1073
<i>pallida</i> , W.	1078	<i>Ibericum</i> , <i>St.</i>	1074
<i>parvifolia</i> , <i>Sprun.</i>	1075		
<i>parvifolia</i> , <i>Fl. Gr.</i>	1076	ASCLEPIADEÆ.	49
<i>pœilantha</i> , <i>Boiss.</i>	1080	<i>Asclepias canescens</i> , W.	52
<i>Pontica</i> , <i>Lam.</i>	1081	<i>Cretica</i> , <i>Clus.</i>	52
<i>rotunda</i> , <i>L.</i>	1077	<i>fruticosa</i> , <i>L.</i>	61
<i>scabridula</i> , <i>Boiss.</i>	1081	<i>gigantea</i> , <i>Forsk.</i>	57
<i>semperflorens</i> , <i>L.</i>	1075	<i>laniflora</i> , <i>Del.</i>	58
<i>Tournefortii</i> , J. et Sp.	1076	<i>microphylla</i> , <i>Roxb.</i>	Add
Armeria , W.	872	<i>nigra</i> , <i>L.</i>	52
<i>argyrocephala</i> , <i>Wallr.</i>	875	<i>Orientalis</i> , <i>Buxb.</i>	52
		<i>procera</i> , W.	57
		<i>spiralis</i> , <i>Forsk.</i>	58
		<i>Vincetoxicum</i> , <i>L.</i>	54
		Asperugo , L.	275
		<i>Egyptiaca</i> , <i>L.</i>	159

	Pag.		Pag.
Asperugo procumbens , L.	275	Atriplex Halimus , L.	916
<i>verrucosa</i> , Lam.	159	<i>hastatum</i> , L.	909
Asterolinum , Link.	10	<i>hortense</i> , L.	907
<i>Linum stellatum</i> , L.	10	<i>incisum</i> , M. B.	910
Astragalus setiferus , D.C.	984	<i>Kataf</i> , Ehr.	916
Atraphaxis , L.	1019	<i>laeiniatum</i> , Koch	910
<i>Afghana</i> , Meisn.	1022	<i>lasianthum</i> , Boiss.	910
<i>angustifolia</i> , J. et Sp.	1023	<i>Lehmannianum</i> , Bge.	911
<i>Aucheri</i> , J. et Sp.	1024	<i>leptocladum</i> , Boiss.	911
<i>Billardieri</i> , J. et Sp.	1022	<i>leucocladum</i> , Boiss.	915
<i>buxifolia</i> , M.B.	1023	<i>littorale</i> , L.	908
<i>Calvertii</i> , Boiss.	1021	<i>marinum</i> , L.	908
<i>candida</i> , B. et Haussk.	1021	<i>micranthum</i> , C. A. M.	909
<i>compacta</i> , Led.	1021	<i>microcarpum</i> , W. K.	909
<i>densiflora</i> , C. Koch	1021	<i>Moneta</i> , Bge.	912
<i>Fischeri</i> , J. et Sp.	1021	<i>nitens</i> , Reb.	908
<i>grandiflora</i> , W.	1024	<i>oblongifolium</i> , W. K.	909
<i>Karelini</i> , J. et Sp.	1021	<i>ocymifolium</i> , Viv.	915
<i>pyrifolia</i> , Bge.	1023	<i>Olivieri</i> , Moq.	910
<i>replicata</i> , Lam.	1021	<i>Palestinum</i> , Boiss.	914
<i>Sinaica</i> , J. et Sp.	1021	<i>patulum</i> , Griseb.	909
<i>spinosa</i> , L.	1020	<i>patulum</i> , L.	909
<i>suædæfolia</i> , J. et Sp.	1024	<i>pedunculatum</i> , L.	912
<i>Tournefortii</i> , J. et Sp.	1023	<i>Persicum</i> , Boiss.	916
<i>variabilis</i> , J. et Sp.	1022	<i>portulacoides</i> , L.	915
Atriplex , L.	906	<i>pruinatum</i> , Sieb.	910
<i>acuminatum</i> , W. K.	908	<i>recurvum</i> , Urv.	910
<i>album</i> , Scop.	911	<i>roseum</i> , L.	911
<i>Alexandrinum</i> , Boiss.	914	<i>rotundifolium</i> , Boiss.	915
<i>Arabicum</i> , Ehr.	917	<i>Stocksii</i> , Boiss.	916
<i>Aucheri</i> , Moq.	908	<i>stylosum</i> , Viv.	917
<i>Belangeri</i> , Moq.	915	<i>Tataricum</i> , L.	910
<i>bracteosum</i> , Trautv.	910	<i>Tataricum</i> , Auct.	909
<i>Chanykowii</i> , Bge.	915	<i>thunbergiaefolium</i> , B.	
<i>coriaceum</i> , Forsk.	915	<i>et Noë</i>	914
<i>crassifolium</i> , C. A. Mey.	908	<i>triangulare</i> , W.	909
<i>erytallinum</i> , Ehr.	915	<i>Turcomanicum</i> , F. et M.	911
<i>dimorphostegium</i> , K. et		<i>Venatum</i> , W.	910
<i>Kir.</i>	909	<i>verruciferum</i> , M. B.	915
<i>farinosum</i> , Forsk.	917		
<i>Flabellum</i> , Bge.	912	Atropa , L.	290
<i>glaucum ex parte</i> , Moq.	915	<i>Belladonna</i> , L.	291
<i>Græcum</i> , W.	909	<i>Mandragora</i> , L.	291
<i>Graecum</i> , Fl. Gr.	910		
<i>Griffithii</i> , Moq.	916	Avicennia , L.	536
		<i>officinalis</i> , L.	536
		<i>tomentosa</i> , Wall.	537

	PAG.		PAG.
BALANOPHORACEÆ.			
Ballota , L.	1072	Beta <i>Noëana</i> , Bge.	899
<i>acetabulosa</i> , L.	771	<i>trigyna</i> , W. K.	899
<i>Arabica</i> , H. St.	772	<i>vulgaris</i> , L.	898
<i>Armena</i> , Boiss.	774	Betonica , L.	749
<i>Ancheri</i> , Boiss.	773	<i>Betonica Alopecuros</i> , Jacq.	750
<i>Damascena</i> , Boiss.	772	<i>Brantii</i> , Bth.	750
<i>fœtida</i> , Lam.	775	<i>glabrata</i> , C. Koch	752
<i>glandulifera</i> , Tr.	774	<i>Græca</i> , B. et Spr.	751
<i>integrifolia</i> , Bth.	775	<i>grandiflora</i> , W.	751
<i>Larendana</i> , Boiss.	774	<i>hirsuta</i> , C. A. M.	750
<i>macrodonta</i> , B. et Bal.	774	<i>Jacquinii</i> , Spr. et Godr.	749
<i>microphylla</i> , Bth.	776	<i>macrantha</i> , C. Koch	751
<i>nigra</i> , L.	775	<i>macrostachya</i> , Wend.	750
<i>obliqua</i> , Bth.	775	<i>nivea</i> , Stev.	750
<i>Orientalis</i> , Presl	776	<i>officinalis</i> , L.	752
<i>Persica</i> , Bth.	777	<i>Orientalis</i> , L.	750
<i>pseudodictamnus</i> , L.	772	<i>Orphanidea</i> , Heldr.	750
<i>rotundifolia</i> , C. Koch	774	<i>Scardica</i> , Griseb.	751
<i>rugosa</i> , Bth.	775	<i>serotina</i> , Host.	752
<i>Russelianæ</i> , Bth.	775	<i>stricta</i> , Anet.	752
<i>saxatilis</i> , Sieb.	775	BETULACEÆ.	1178
<i>Schimperi</i> , Bth.	776	Betula , Tourn.	1180
<i>undulata</i> , Fres.	773	<i>alba</i> , L.	1180
Barleria , L.	525	<i>glutinosa</i> , Wallr.	1181
<i>acanthoides</i> , Vahl	524	<i>odorata</i> , Bechst.	1181
<i>Aucheriana</i> , Nees	525	<i>ovata</i> , C. Koch	1181
<i>candida</i> , Nees	524	<i>pubescens</i> , Ehrh.	1181
<i>diandra</i> , Hochst.	525	<i>verrucosa</i> , Ehrh.	1181
<i>Hochstetteri</i> , Nees	525	Bienertia , Bge.	945
<i>Bartsia bicolor</i> , D.C.	474	<i>cycloptera</i> , Bge.	945
<i>irifida</i> , Spr.	472	BIGNONIACEÆ.	79
<i>Trixago</i> , L.	474	Bignonia <i>glauca</i> , Decaisn.	80
<i>versicolor</i> , Pers.	474	<i>undulata</i> , Roxb.	80
<i>Bassia muricata</i> , All.	927	Blepharis , Juss.	520
<i>Batatas littoralis</i> , Ch.	112	<i>edulis</i> , Forsk.	520
<i>Belowia baccifera</i> , Moq.	942	Blitum , L.	904
<i>sinuata</i> , Guss.	112	<i>Bonus Henricus</i> , L.	904
Beta , L.	898	<i>Bonus Henricus</i> , C.A. M.	904
<i>intermedia</i> , Bge.	900	<i>glaucum</i> , Koch	905
<i>lomatogona</i> , F. M.	899	<i>polymorphum</i> , C.A. M.	905
<i>longespicata</i> , Moq.	899	<i>rubrum</i> , L.	905
<i>macrorrhiza</i> , Stev.	900	<i>virgatum</i> , L.	905
<i>maritima</i> , L.	899	<i>Bæhmeria hypoleuca</i> , Hochst.	1148
<i>nana</i> , B. et Heldr.	898	Boerhaavia , L.	1044
		<i>ascendens</i> , W.	1045

	Pag.		Pag.
Boerhaavia dichotoma , Vahl	1044	Bungea , C. A. Mey.	471
<i>diffusa</i> , L.	1045	<i>trifida</i> , Vahl	471
<i>elegans</i> , Ch.	1045	Burgsdorffia rigida , Mænch	706
<i>grandiflora</i> , Rich.	1044	BUXACEÆ.	1143
Marroccana , Ball.	1045	Buxus , L.	1145
<i>plumbaginea</i> , Cav.	1044	<i>longifolia</i> , Boiss.	1144
<i>repanda</i> , Ky.	1045	<i>sempervirens</i> , L.	1144
<i>repens</i> , L.	1045	Caccinia , Sav.	276
<i>rubicunda</i> , St.	1045	<i>Actinobole</i> , Bge.	279
<i>scandens</i> , Forsk.	1044	<i>Ranwolffii</i> , C. Koch	277
<i>sessiliflora</i> , St.	1045	<i>Rauwolfii</i> , A.D.C.	279
<i>vulvariaefolia</i> , Poir.	1045	<i>Celsii</i> , Boiss.	277
BORRAGINÆ.	123	<i>glauca</i> , Sav.	277
Borrago , L.	150	<i>Kotschy</i> , Boiss.	278
<i>Africana</i> , L.	280	<i>Russeli</i> , Boiss.	278
<i>Arabica</i> , Ehr.	281	<i>strigosa</i> , Boiss.	278
<i>cordifolia</i> , Mænch	177	<i>Caidbeja adhærens</i> , Forsk.	1151
<i>crassifolia</i> , Vent.	277	Calamintha , Mænch	
<i>Cretica</i> , W.	177	<i>Acinos</i> , L.	582
<i>Indica</i> , L.	280	<i>Alpina</i> , L.	581
<i>macranthera</i> , Russ	279	<i>betulifolia</i> , B. et Bal.	576
<i>officinalis</i> , L.	150	<i>Byzantina</i> , C. Koch	578
<i>Orientalis</i> , L.	177	<i>canescens</i> , C. Koch	578
Boucerosia , Wight.	63	<i>Clinopodium</i> , Bth.	579
<i>Auchriana</i> , Dec.	63	<i>Cretica</i> , L.	578
<i>Sinaica</i> , Dec.	63	<i>depauperata</i> , Bge.	581
<i>Stocksiana</i> , Boiss.	63	<i>florida</i> , Boiss.	580
<i>Bovea Sinaica</i> , Dec.	425	<i>glabrescens</i> , Boiss.	580
Brachylepis , C. A. M.	971	<i>glomerulosa</i> , B. et Heldr.	579
<i>elatior</i> , C. A. Mey.	970	<i>grandiflora</i> , L.	576
<i>eriopoda</i> , C. A. M.	971	<i>graveolens</i> , M. B.	585
<i>humilis</i> , Less.	971	<i>Hausknechtii</i> , Bge.	580
<i>intermedia</i> , Kar.	970	<i>incana</i> , Fl. Gr.	578
<i>salsa</i> , C. A. M.	971	<i>maritima</i> , Bth.	585
<i>Brezia heterophylla</i> , Moq.	943	<i>Nepeta</i> , L.	577
Brunella , L.	691	<i>nepetoides</i> , Jord.	577
<i>alba</i> , Pall.	692	<i>nivea</i> , Boiss.	581
<i>grandiflora</i> , Mænch.	692	<i>officinalis</i> , Mænch	577
<i>laciniata</i> , L.	692	<i>organifolia</i> , Labill.	379
<i>vulgaris</i> , L.	691	<i>organifolia</i> , Vis.	575
<i>Brunnera myosotidiflora</i> , Stev.	157	<i>Pamphylica</i> , B. et Heldr.	577
<i>Buchnera Hermonthica</i> , Del.	469	<i>Patavina</i> , Jacq.	582
<i>Buglossum tinctorium</i> , Lam.	227	<i>Patavina</i> var. Gr.	582
Buddleia , L.	Add. 1198	<i>rotundifolia</i> , Rehb.	577
<i>crispa</i> , Bth.	Add. 1193		

	Pag.		Pag.
<i>Calamintha spicigera</i> , C. Koch	574	<i>Cardiotheca pubescens</i> , Ehr.	563
<i>Spruneri</i> , Boiss.	577	<i>virgata</i> , Ehr.	362
<i>staminea</i> , Boiss.	580	<i>Carissa</i> , L.	44
<i>suaveolens</i> , Sm.	582	<i>edulis</i> , Vahl	44
<i>umbrosa</i> , M. B.	578	<i>Caroxylon acutifolium</i> , Moq.	958
<i>Calligonum</i> , L.	997	<i>articulatum</i> , Moq.	949
<i>Bungei</i> , Boiss.	999	<i>Auricula</i> , Moq.	958
<i>Calliphysa</i> , Bge.	1001	<i>fætidum</i> , Moq.	961
<i>comosum</i> , L'hér.	1000	<i>Griffithii</i> , Moq.	950
<i>crinitum</i> , Boiss.	1000	<i>hispidulum</i> , Bge.	960
<i>denticulatum</i> , Bge.	999	<i>lanceifolium</i> , Boiss.	958
<i>eriopodum</i> , Bge.	1001	<i>nodulosum</i> , Moq.	960
<i>Pallasia</i> , L'hér.	998	<i>subaphyllum</i> , Moq.	959
<i>Persicum</i> , B. et Buhse	999	<i>tetragonum</i> , Moq.	957
<i>polygonoides</i> , Pall.	989	<i>Carpinus</i> , Tourn.	1177
<i>polygonoides</i> , L.	1000	<i>Betulus</i> , L.	1177
<i>stenopterum</i> , Bge.	999	<i>Duinensis</i> , Scop.	1177
<i>tetrapterum</i> , J. et Sp.	999	<i>Orientalis</i> , Lam.	1177
<i>tripterum</i> , Bge.	999	<i>Ostrya</i> , L.	1178
<i>Calliphysa juncea</i> , F. et M.	1001	<i>Castanea</i> , Tourn.	1174
<i>Calotropis</i> , R. Br.	57	<i>vulgaris</i> , Lam.	1175
<i>procera</i> , Willd.	57	<i>Celosia</i> , L.	987
<i>Calystegia</i> , R. Br.	111	<i>argentea</i> , L.	987
<i>sepium</i> , R. Br.	111	<i>lanata</i> , L.	993
<i>Soldanella</i> , R. Br.	112	<i>Celsia</i> , L.	349
<i>sylvatica</i> , W. K.	111	<i>acaulis</i> , B. et Ch.	554
<i>Campelepis viminea</i> , Falc.	51	<i>agrimoniæfolia</i> , C. Koch	559
<i>Camphorosma</i> , L.	920	<i>alpina</i> , B. et Ky.	551
<i>monandrum</i> , Bge.	922	<i>Arcturus</i> , L.	552
<i>Monspeliaenum</i> , L.	920	<i>Arcturus</i> , Jacq.	351
<i>ovatum</i> , W. K.	921	<i>Boissieri</i> , H. et Sart.	555
<i>perenne</i> , Pall.	920	<i>Aucheria</i> , Boiss.	560
<i>polygamum</i> , Bge.	921	<i>aurea</i> , C. Koch	361
<i>Ruthenicum</i> , M. B.	921	<i>brachysepala</i> , Fisch	360
<i>Camptocarpus Orientalis</i> ,		<i>bugulifolia</i> , Lam.	351
C. Koch	228	<i>Chrysurus</i> , Fenzl	359
<i>Campylanthus</i> , Roth	453	<i>Cilicia</i> , B. et Heldr.	358
<i>ramosissimus</i> , Wight	453	<i>collina</i> , Bth.	357
<i>salsoloides</i> , Stocks	453	<i>Coromandeliana</i> , Vahl	358
<i>Campylocaryon album</i> , D.C.	170	<i>coronopifolia</i> , B. et Bal.	356
<i>Syriacum</i> , Boiss.	226	<i>Cretica</i> , L. fil.	352
<i>Cannabis</i> , Tourn.	1152	<i>Cyllenea</i> , B. et Heldr.	355
<i>sativa</i> , L.	1152	<i>Dænzeri</i> , Bor. et Ch.	353
<i>Capraria Arabica</i> , H. et St.	422	<i>glandulosa</i> , Bosch.	350
<i>dissecta</i> , Del.	422	<i>heterophylla</i> , Desf.	359

	Pag.		Pag.
<i>Celsia incana</i> , Bth.	557	<i>Cerinthopsis Kurdica</i> , Ky.	271
<i>intricata</i> , Bth.	559	<i>Chæturus Marrubiastrum</i> ,	
<i>lanceolata</i> , Vent.	357	Ehrh.	754
<i>Lepturus</i> , Schult.	561	Chamæsphecos , Schr.	680
<i>Luciliæ</i> , Boiss.	556	<i>ilicifolius</i> , Schr.	680
<i>Lycia</i> , Boiss.	557	Chenolea , Thunb.	922
<i>Orientalis</i> , L.	360	<i>Arabica</i> , Boiss.	922
<i>parviflora</i> , Decaisn.	554	<i>Chenopodina altissima</i> , Moq.	941
<i>Persica</i> , C. A. Mey.	555	<i>Asphaltica</i> , Boiss.	938
<i>pinetorum</i> , Boiss.	358	<i>Biebersteiniana</i> , Moq.	938
<i>Pontica</i> , Boiss.	352	<i>microphylla</i> , Moq.	938
<i>pyroliformis</i> , B. et Heldr.	555	<i>physophora</i> , Moq.	959
<i>speciosa</i> , Fenzl	553	<i>pyrenantha</i> , C. Koch	951
<i>sublanata</i> , Jacq.	353	<i>setigera</i> , Moq.	942
<i>Suwarowiana</i> , C. Koch	557	Chenopodium , L.	900
<i>tomentosa</i> , Zucc.	556	<i>album</i> , L.	901
<i>viscosa</i> , Roth	559	<i>altissimum</i> , L.	941
Celtis , L.	1156	<i>ambrosioides</i> , L.	904
<i>aspera</i> , Stev.	1157	<i>Augustanum</i> , All.	925
<i>australis</i> , L.	1156	<i>baryosmum</i> , R. et Sch.	961
<i>Caucasica</i> , W.	1156	<i>Biebersteinianum</i> , R. et	
<i>glabrata</i> , Stev.	1157	Sch.	958
<i>Kotschyana</i> , St.	1157	<i>Bonus Henricus</i> , L.	904
<i>Tournefortii</i> , Lam.	1157	<i>Botrys</i> , L.	905
Ceratocarpus , Buxb.	918	<i>caulatum</i> , Jacq.	992
<i>arenarius</i> , L.	918	<i>Clemente</i> , Sp.	942
CERATOPHYLLEÆ .	1202	<i>concatenatum</i> , Th.	901
Ceratophyllum , L.	1202	<i>crassifolium</i> , M. B.	938
<i>demersum</i> , L.	1202	<i>ficifolium</i> , Sm.	901
<i>Ceratospermum papposum</i> ,		<i>fœtidum</i> , Schrad.	905
Pers.	918	<i>glaueum</i> , L.	903
Cerinthe , Tourn.	148	<i>hortense</i> , R. et Sch.	945
<i>Alpina</i> , Kit.	149	<i>hybridum</i> , L.	902
<i>aspera</i> , Roth	149	<i>intermedium</i> , Koch	903
<i>auriculata</i> , Ten.	148	<i>leiospermum</i> , D. C.	901
<i>Cilicica</i> , Ky.	148	<i>linifolium</i> , Schult.	944
<i>cleiostoma</i> , B. et Sp.	148	<i>maritimum</i> , L.	941
<i>glabra</i> , D. C.	149	<i>microphyllum</i> , M. B.	958
<i>macrophylla</i> , B. et Heldr.	148	<i>murale</i> , L.	902
<i>maculata</i> , M. B.	148	<i>olidum</i> , Sm.	901
<i>major</i> , Lam.	149	<i>oppositiflorum</i> , W.	967
<i>minor</i> , L.	148	<i>opulifolium</i> , Schrad.	901
<i>purpurea</i> , Viv.	150	<i>polyspermum</i> , L.	900
<i>retorta</i> , Fl. Gr.	149	<i>rubrum</i> , L.	905
<i>strigosa</i> , Rehb.	149	<i>setigerum</i> , D. C.	942

	Pag.		Pag.
<i>Chenopodium urbicum</i> , L.	902	<i>Convolvulus atriplicifolius</i> ,	
<i>villosum</i> , Lam.	926	Poir.	105
<i>viride</i> , Curt.	901	<i>Aucheria Chois.</i>	96
<i>Vulvaria</i> , L.	901	<i>batataefolius</i> , B. et	
<i>Chironia maritima</i> , W.	68	Haussk.	106
<i>Chlamydanthus Tartonraira</i> ,		<i>Besseri</i> , Spr.	97
Mey.	1053	<i>betonicifolius</i> , Mill.	105
Chlora , L.	66	<i>brachiphyllus</i> , Boiss.	90
<i>acuminata</i> , Koch	66	<i>brevispinus</i> , J. et Sp.	87
<i>perfoliata</i> , Willd.	66	<i>bryoniæfolius</i> , Rehb.	106
<i>serotina</i> , Koch	66	<i>Cairicus</i> , L.	113
Cicendia , Ad.	65	<i>Calverti</i> , Boiss.	94
<i>filiformis</i> , L.	66	<i>Cantabrica</i> , L.	95
Cionura , Griseb.	62	<i>capitatus</i> , Schimp.	105
<i>erecta</i> , L.	62	<i>cardiosepalus</i> , Boiss.	96
<i>Cissus arborea</i> , Forsk.	45	<i>Cataonicus</i> , B. et Haussk.	98
Clerodendron , L.	555	<i>cephalophorus</i> , Boiss.	102
<i>Acerbianum</i> , Vis.	556	<i>cephalopodus</i> , Boiss.	101
<i>Clinopodium Ægyptiacum</i> ,		<i>cespitosus</i> , Ehr.	97
Lam.	579	<i>chondrilloides</i> , Boiss.	92
<i>fruticosum</i> , Forsk.	777	<i>Cneorum</i> , Habl.	94
<i>plumosum</i> , Sieb.	579	<i>Cneorum</i> , L.	95
<i>vulgare</i> , L.	579	<i>Cneorum</i> , Forsk.	89
Cocculus , D.C.	Add. 1201	<i>cochlearis</i> , Gris.	98
<i>Leœba</i> , Del.	1201	<i>Cœlesyriacus</i> , Boiss.	109
<i>Colsmannia flava</i> , Lehm.	186	<i>commutatus</i> , Boiss.	94
Comandra , Nutt.	1059	<i>compactus</i> , Boiss.	98
<i>elegans</i> , Roch.	1059 et Add.	<i>Cyprius</i> , Boiss.	95
CONVOLVULACEÆ.	84	<i>Damascenus</i> , B. et Gaill.	107
Convolvulus , L.	84	<i>Dorycnium</i> , Boiss.	91
<i>acanthocladus</i> , Boiss.	86	<i>Eichwaldi</i> , Boiss.	87
<i>agrophilus</i> , C. Koch	104	<i>eremophilus</i> , B. et Buhse	90
<i>alceæfolius</i> , Fl. Pelop.	106	<i>erinaceus</i> , Led.	87
<i>althæoides</i> , L.	106	<i>erinaceus</i> , C. A. M.	90
<i>ambliocalyx</i> , Steud.	109	<i>evolvuloides</i> , Boiss.	91
<i>Ammocharis</i> , B. et Haussk.	104	<i>evolvuloides</i> , Desf.	111
<i>amaenus</i> , C. Koch	105	<i>farinosus</i> , L.	111
<i>arenarius</i> , Vahl	112	<i>Fatmensis</i> , Kze.	109
<i>argenteus</i> , Desr.	93	<i>Forskahlei</i> , Del.	89
<i>argyreus</i> , D.C.	107	<i>frutescens</i> , Pall.	87
<i>armatus</i> , Del.	88	<i>fruticosus</i> , Pall.	87
<i>Armenus</i> , Boiss. et Ky.	105	<i>Galaticus</i> , Rost.	104
<i>arvensis</i> , L.	108	<i>genistoides</i> , J. et Sp.	87
<i>ascendens</i> , St.	100	<i>Germaniciæ</i> , B. et Haussk.	104
<i>Assyriacus</i> , Gr.	99	<i>glomeratus</i> .	Add.

Pág.		Pág.	
Convolvulus gonoclados, Boiss.	102	Convolvulus reticulatus, Ch.	100
<i>Haussknechtii, Boiss.</i>	102	<i>Ruprechti, Boiss.</i>	96
<i>heterophyllus, Ehr.</i>	109	<i>sagittifolius, Mich.</i>	113
<i>hirsutus, Ten.</i>	106	<i>sagittifolius, Fl. Gr.</i>	105
<i>hirsutus, Stev.</i>	105	<i>salviaefolius, Sieb.</i>	100
<i>holosericeus, M. B.</i>	95	<i>sarothrocladus, Boiss.</i>	92
<i>humilis, Jacq.</i>	111	<i>saxatilis, M. B.</i>	94
<i>hystrix, Vahl</i>	88	<i>Scammonia, L.</i>	108
<i>Imperati, Vahl</i>	112	<i>Schimperi, Boiss.</i>	101
<i>intermedius, Lois.</i>	97	<i>Schirazianus, Boiss.</i>	94
<i>Italicus, R. et Sch.</i>	106	<i>Scindicus, Stocks</i>	90
<i>Kotschyianus, Boiss.</i>	101	<i>secundus, Desr.</i>	100
<i>lanatus, Vahl</i>	89	<i>sepinm, L.</i>	111
<i>lanuginosus, Vahl</i>	105	<i>sericeus, Burm.</i>	101
<i>lanuginosus, Led.</i>	94	<i>sericeus, Ch.</i>	89
<i>lasiophlæus, J. et Sp.</i>	86	<i>Sibthorpii, R. et Sch.</i>	105
<i>leiocalycinus, Boiss.</i>	86	<i>Siculus, L.</i>	109
<i>leptoclados, Boiss.</i>	91	<i>sinuatus, Pet.</i>	112
<i>Libanoticus, Boiss.</i>	97	<i>Sogdianus, Bge.</i>	103
<i>linearis, Bot. Mag.</i>	93	<i>Soldanella, L.</i>	112
<i>lineatus, Fl. Gr.</i>	93	<i>spicasfolius, Desr.</i>	97
<i>lineatus, L.</i>	98	<i>spinosis, Burm.</i>	87
<i>littoralis, L.</i>	112	<i>spinosis, Desr.</i>	87
<i>lycioides, Boiss.</i>	86	<i>stachydisfolius, Ch.</i>	107
<i>Mascatensis, Boiss.</i>	89	<i>stenophyllus, Boiss.</i>	106
<i>microphyllus, Sieb.</i>	103	<i>Stocksii, Boiss.</i>	110
<i>modestus, Boiss.</i>	95	<i>stoloniferus, Cyr.</i>	112
<i>nitens, C. Koch</i>	97	<i>strigulosus, Boiss.</i>	98
<i>oleæfolius, Desr.</i>	95	<i>suffruticosus, Desf.</i>	111
<i>oxyphyllus, Boiss.</i>	88	<i>svylvaticus, W. K.</i>	111
<i>oxysepalus, Boiss.</i>	90	<i>svylvestris, W.</i>	112
<i>Palestinus, Boiss.</i>	107	<i>tenellus, Stocks</i>	110
<i>Paleslinus, Boiss. var.</i>	106	<i>tenuissimus, Fl. Gr.</i>	107
<i>Parnassicus, Boiss.</i>	98	<i>tomentosus, Ch.</i>	105
<i>peduncularis, Boiss.</i>	105	<i>Tournefortii, Sieb.</i>	93
<i>pentapetaloides, L.</i>	110	<i>tricolor, L.</i>	111
<i>Persicus, L.</i>	99	<i>triqueter, Rehm.</i>	94
<i>ilosellæfolius, Desr.</i>		<i>ulicinus, Boiss.</i>	89
	103 et Add.	<i>undulatus, Cav.</i>	110
<i>pseudoscammonia, C.</i>		<i>verrucosus, Ehr.</i>	107
<i>Koch</i>	108	<i>virgatus, Boiss.</i>	88
<i>pubescens, Russ.</i>	105	<i>waltherioides, B. et</i>	
<i>pusillus, Pall.</i>	94	<i>Haussk.</i>	100
<i>pyrrhotrichus, Boiss.</i>	102	<i>Wheleri, Vahl</i>	115
<i>radicosus, Heldr. et Sart.</i>	99	<i>Cordia, L.</i>	123

	Pag.		Pag.
Cordia Myxa, L.	124	Crozophora Sieberi, Presl.	1141
Corispermum, L.	929	<i>tinctoria, L.</i>	1140
<i>hyssopifolium, L.</i>	930	<i>verbascifolia, W.</i>	1141
<i>Lehmannianum, Bge.</i>	930	Cunila capitata, L.	585
<i>Marschallii, Stev.</i>	929	CUPULIFERÆ.	1162
<i>nitidum, Kit.</i>	930	Cuscuta, L.	115
<i>Orientale, Lam.</i>	929	<i>acutissima, Buch.</i>	121
<i>Pallasii, Stev.</i>	929	<i>alba, Presl.</i>	116
<i>polygaloides, F. et M.</i>	931	<i>alpina, Hoh.</i>	118
<i>purpurascens, Hort.</i>	930	<i>approximata, Bab.</i>	116
<i>spicatum, W.</i>	929	<i>Arabica, Wight</i>	121
<i>squarrosum, M. B.</i>	930	<i>Arabica, Fres.</i>	120
<i>squarrosum, L.</i>	930	<i>astyla, Eng.</i>	112
<i>squarrosum, Pall.</i>	928	<i>aurantiaca, Req.</i>	121
Cornulaca, Del.	983	<i>Babylonica, Auch.</i>	118
<i>amblyacantha, Bge.</i>	984	<i>Balansæ, Boiss. et R.</i>	117
<i>Aucherii, Moq.</i>	983	<i>Boissieri, Stocks</i>	121
<i>monacantha, Del.</i>	984	<i>brachystyla, C. Koch</i>	118
<i>muricata, Del.</i>	927	<i>breviflora, Vis.</i>	121
<i>setifera, Moq.</i>	984	<i>brevistyla, A. Br.</i>	117
Corylus, Tourn.	1176	<i>Calliopes, Heldr.</i>	115
<i>Avellana, L.</i>	1176	<i>capillaris, Rehb.</i>	115
<i>Byzantina, Clus.</i>	1176	<i>Cesatiana, Bert.</i>	121
<i>Colurna, L.</i>	1176	<i>Chinensis, Lam.</i>	120
<i>Jacquemontii, Decaisn.</i>	1176	<i>chrysocoma, Welw.</i>	121
<i>Pontica, C. Koch</i>	1176	<i>ciliaris, Hoh.</i>	120
<i>tubulosa, W.</i>	1176	<i>Cretica, Tourn.</i>	116
Cressa, L.	114	<i>densiflora, Soy.</i>	118
<i>Cretica, L.</i>	114	<i>elegans, B. et Bal.</i>	119
Croton obliquifolium, Vis.	1140	<i>Epilinum, Weihe</i>	118
<i>obliquum, Vahl</i>	1141	<i>Epispartos, Boiss. et Orph.</i>	116
<i>oblongifolium, Del.</i>	1141	<i>Epithynum, L.</i>	115
<i>plicatum, Vahl</i>	1140	<i>Europaea, L.</i>	117
<i>Rottleri, Gris.</i>	1140	<i>gigantea, Griff.</i>	122
<i>tinctorum, L.</i>	1140	<i>globularis, Bert.</i>	117
<i>verbascifolium, W.</i>	1141	<i>globulosa, Boiss. et R.</i>	117
<i>villosum, Fl. Gr.</i>	1141	<i>hyalina, Boiss.</i>	118
Crozophora, Neck.	1139	<i>hyalina, Roth</i>	121
<i>gracilis, F. et M.</i>	1140	<i>hyalina, Wight</i>	120
<i>Hierosolymitana, Spr.</i>	1141	<i>Kotschyana, Boiss.</i>	119
<i>integrifolia, Bge.</i>	1141	<i>Kotschy, Desm.</i>	116
<i>obliqua, Vahl</i>	1141	<i>Kurdica, Eng.</i>	118
<i>plicata, Vahl</i>	1140	<i>Lehmaniana, Bge.</i>	112
<i>prostrata, Dalz.</i>	1140	<i>leucosphæra, B. et Heldr.</i>	116
<i>sabulosa, K. et Kir.</i>	1140	<i>lupuliformis, Krock.</i>	122

	Pag.		Pag.
Cyscanta macranthera , Heldr.	115	Cynanchum Argel , Del.	56
<i>major</i> , D. C.	118	<i>contiguum</i> , Koch	54
<i>micrantha</i> , Tin.	116	<i>erectum</i> , L.	62
<i>minor</i> , D. C.	115	<i>heterophyllum</i> , Del.	Add.
<i>monogyna</i> , Vahl	121	<i>laxum</i> , Bartl.	50
<i>monogyna</i> , Auct. Germ.	122	<i>longisolum</i> , Mart.	64
<i>obtusiflora</i> , H. B. K.	121	<i>medium</i> R. Br.	54
<i>Orientalis</i> , Tourn.	122	<i>minus</i> , C. Koch	56
<i>oxypetala</i> , Boiss.	121	<i>Monseliacum</i> , L.	60
<i>Palestina</i> , Boiss	116	<i>pyrotechnicum</i> , F.	63
<i>planiflora</i> , Ten.	116	CYNOCRABEÆ.	896
<i>Persira</i> , Decaisne	119	Cynocrambe , L.	896
<i>pulchella</i> , Eng.	120	<i>prostrata</i> , Gærtn.	897
<i>Rogowitschiana</i> , Trautv.	121	Cynoglossum , L.	263
<i>Schiraziana</i> , Boiss.	116	<i>angustifolium</i> , W.	263
<i>Schkuhriana</i> , Pf	118	<i>cristatum</i> , Lam.	258
<i>subuniflora</i> , C. Koch	123	<i>calycinum</i> , C. A. M.	276
<i>Tinei</i> , Jus.	121	<i>cheirifolium</i> , Scop.	265
<i>urceolata</i> , Kze.	116	<i>Columæ</i> , Biv.	265
Cyclamen , L.	10	<i>Creticum</i> , Vill.	265
<i>æstivum</i> , Park.	11	<i>Dioscoridis</i> , Led.	265
<i>Aleppicum</i> , Fisch.	12	<i>emarginatum</i> , Lam.	263
<i>Caucasicum</i> , Willd.	11	<i>Gileadense</i> , W.	269
<i>Cilicium</i> , Boiss. et Heldr.	11	<i>glastifolium</i> , W.	262
<i>Coum</i> , Mill.	11	<i>holosericeum</i> , Stev.	264
<i>Cyprium</i> , Ky.	13	<i>intermedium</i> , Fres.	255
<i>elegans</i> , Boiss. et Buhse	11	<i>lanatum</i> , Lam.	272
<i>Europæum</i> , L.	13	<i>longiflorum</i> , Lehm.	279
<i>ficariifolium</i> , Desm.	15	<i>macrostylum</i> , Bge.	266
<i>Græcum</i> , Link	13	<i>montanum</i> , Lam.	264
<i>hederæfolium</i> , Ait.	12	<i>myosotoides</i> , Lab.	257
<i>hederæfolium</i> , Sm.	12	<i>Nebrodense</i> , Guss.	265
<i>hederæfolium</i> , Willd.	15	<i>officinale</i> , L.	264
<i>Ibericum</i> , Stev.	11	<i>parvifolium</i> , C. Koch	265
<i>latifolium</i> , Fl. Gr.	12	<i>pellucidum</i> , Lup.	264
<i>Neapolitanum</i> , Ten.	12	<i>pictum</i> , Ait.	265
<i>Persicum</i> , Mill.	12	<i>pustulatum</i> , Boiss.	265
<i>Persicum</i> , Smith	13	<i>racemosum</i> , Schreb.	265
<i>repandum</i> , Fl. Gr.	12	<i>rugosum</i> , W.	221
<i>Romanum</i> , Griseb.	12	<i>Sphacioticum</i> , B. et Heldr.	263
<i>vernale</i> , C. Koch	11	<i>stamineum</i> , M. B.	269
<i>vernum</i> , Sweet	11	<i>stamineum</i> , Desf.	270
<i>vernum</i> , Lob.	12	<i>sylvaticum</i> , Heuke	264
Cynanchum , L.	60	Cynomorium , Mich.	1072
<i>acutum</i> , L.	60		

	Pag.		Pag.
Cynomorium coccineum , L.	1072	Debregeasia , Gaud.	1148
Cyphomattia , Boiss.	272	bicolor , Roxb.	1148
<i>lanata</i> , Lam.	272	<i>hypoleuca</i> , Wedd.	1146
CYRTANDRACEÆ.	81	Desmochæta <i>flavescens</i> , D.C.	995
CYTINACEÆ.	1070	<i>xanthiooides</i> , A. Br.	995
Cytinus , L.	1071	Diarthron , Turcz.	1054
<i>Hypocistis</i> , L.	1071	<i>carinatum</i> , J. et Sp.	1055
Dæmia , R. Br.	59	<i>vesiculosum</i> , F. et M.	1054
<i>cordata</i> , R. Br.	59	Diaptetera , Juss.	526
<i>incana</i> , Dec.	60	<i>buplevroides</i> , Nees	526
Dais spicata , Endl.	1054	<i>hirtula</i> , Nees	526
Daphne , L.	1046	<i>Roxburghiana</i> , Nees	526
<i>acuminata</i> , B. et Hoh.	1048	Digera , Forsk.	994
<i>Alpina</i> , Sm.	1047	<i>alternifolia</i> , L.	994
<i>angustifolia</i> , C. Koch	1040	<i>arvensis</i> , St. et H.	994
<i>argentea</i> , Fl. Gr.	1053	Digitalis , L.	429
<i>australis</i> , Cyr.	1048	<i>ambigua</i> , Murr.	432
<i>buxifolia</i> , Vahl	1047	<i>aurea</i> , Lindl.	429
<i>buxifolia</i> , Ledeb.	1048	<i>brachyantha</i> , Gr.	429
<i>Caucasica</i> , Pall.	1047	<i>Cariensis</i> , Boiss.	451
<i>collina</i> , Sm.	1048	<i>ciliata</i> , Trautv.	432
<i>glandulosa</i> , Bert.	1047	<i>ferruginea</i> , L.	429
<i>glomerata</i> , Lam.	1049	<i>grandiflora</i> , All.	452
<i>gnidium</i> , L.	1049	<i>Heldreichii</i> , J. et Sp.	431
<i>gnidioides</i> , J. et Sp.	1049	<i>lævigata</i> , W. K.	450
<i>jasminea</i> , Fl. Gr.	1047	<i>lævigata</i> , C. A. M.	450
<i>jasminea</i> , Griseb.	1047	<i>lanata</i> , Ehrh.	450
<i>Imerica</i> , C. Koch	1049	<i>leucophæa</i> , Fl. Gr.	451
<i>Laureola</i> , L.	1050	<i>lutea</i> , Sm.	432
<i>Mezereum</i> , L.	1046	<i>nervosa</i> , St. et Hochst.	430
<i>microphylla</i> , Meisn.	1048	<i>ochroleuca</i> , Jacq.	432
<i>mucronata</i> , Meisn.	1048	<i>orientalis</i> , Lam.	451
<i>Neapolitana</i> , Lodd.	1048	<i>viridiflora</i> , Lindl.	452
<i>oleæfolia</i> , Lam.	1048	Diocea hispidissima , Spr.	213
<i>oleoides</i> , Schreb.	1047	Dionysia , Fenzl	18
<i>oleoides</i> , Urv.	1049	<i>aretiooides</i> , Lehm.	19
<i>paniculata</i> , Lam.	1049	<i>Ancheri</i> , Duby	19
<i>Pontica</i> , L	1050	<i>bryoides</i> , Boiss.	21
<i>salicifolia</i> , Lam.	1047	<i>bryoides</i> , B. et Buhse	22
<i>sericea</i> , Vahl	1048	<i>cespitoso</i> , Duby	20
<i>Tartonraira</i> , L.	1053	<i>curviflora</i> , Bge.	22
<i>tomentosa</i> , Lam.	1053	<i>diapensiæfolia</i> , Boiss.	20
Datura , L.	292	<i>drabæfolia</i> , Bge.	20
<i>Stramonium</i> , L.	292	<i>Kotschy</i> , Bge.	21
		<i>Michauxii</i> , Duby	21

	PAG.	PAG.	
<i>Dionysia odora</i> , Fenzl	19	<i>Echinopsilon erassifolium</i> , Moq.	925
<i>revoluta</i> , Boiss.	18	<i>eriophorum</i> , Moq.	929
<i>rhaftodes</i> , Bge.	22	<i>hirsutum</i> , Moq.	926
<i>tapetodes</i> , Bge.	21	<i>hyssopifolium</i> , Moq.	926
Diospyros , L.	33	<i>lanatum</i> , Moq.	926
<i>Lotus</i> , L.	33	<i>muricatum</i> , Moq.	927
<i>Diotis atriplicoides</i> , M. B.	902	<i>sedoides</i> , Moq.	926
<i>ceratoides</i> , W.	918		
<i>ferruginea</i> , Nees	918		
<i>Diplophyllum veronicaeforme</i>		Echinospermum , Sw.	246
Lehm.	468	<i>barbatum</i> , M. B.	920
<i>Dipteracanthus longifolius</i> ,		<i>brachycentrum</i> , Led.	251
Stocks.	519	<i>Bungei</i> , Boiss.	252
<i>Disandra africana</i> , Camb.	429	<i>calathicarpum</i> , St.	254
<i>Distemon campanularis</i> , Ehr.	223	<i>Cariense</i> , Boiss.	250
Dodartia , L.	424	<i>Caspium</i> , F. et M.	252
<i>Orientalis</i> , L.	424	<i>filiforme</i> , God.	251
<i>Doratanthera lincaris</i> , Bth.	423	<i>heteracanthum</i> , Led.	249
Dorystoechas , Boiss.	590	<i>Ispahanicum</i> , Boiss.	249
<i>hastata</i> , B. et Heldr.	511	<i>Kotschyi</i> , Boiss.	251
Dracocephalum , L.	671	<i>laevigatum</i> , Kar. et Kir.	248
<i>aristatum</i> , Bert.	675	<i>Lappula</i> , L.	249
<i>Aucherii</i> , Boiss.	671	<i>maeranthum</i> , Led.	252
<i>Austriacum</i> , L.	673	<i>marginatum</i> , M. B.	252
<i>botryoides</i> , Stev.	671	<i>microcarpum</i> , Led.	251
<i>botryoides</i> , B. et Buhse	671	<i>oligacanthum</i> , Boiss.	250
<i>canescens</i> , L.	675	<i>putulum</i> , Lehm.	250
<i>Ibericum</i> , M. B.	674	<i>rigidum</i> , D.C.	251
<i>Inderiense</i> , Kar.	574	<i>semiglabrum</i> , Led.	251
<i>Kotschyi</i> , Boiss.	672	<i>sessilitorum</i> , Boiss.	253
<i>lamiifolium</i> , Desf.	757	<i>Sinaicum</i> , A.D.C.	251
<i>multicanthum</i> , M. et Auch.	672	<i>spinocarpus</i> , Forsk.	249
<i>ocymifolium</i> , Mill.	674	<i>squarrosum</i> , Rehb.	249
<i>odoratissimum</i> , P.	588	<i>Szovitsianum</i> , F. et M.	247
<i>peltatum</i> , L.	674	<i>Vahlianum</i> , Lehm.	249
<i>Royleanum</i> , Wall.	674		
<i>Ruyschiana</i> , L.	675	Echiocchilon , Desf.	210
<i>setigerum</i> , B. et Huet.	672	<i>fruticosum</i> , Desf.	211
<i>Tchihatchewii</i> , F.	598	<i>Echioides grandiflora</i> , Stev.	164
<i>teucrioides</i> , Bge.	642		
<i>thymiflorum</i> , L.	674	Echium , L.	204
		<i>acutifolium</i> , Lehm.	205
EBENACEÆ	35	<i>altissimum</i> , Jacq.	205
Ebolium , Kurz	525	<i>amœnum</i> , F. M.	209
<i>Linneanum</i> , Kurz	526	<i>arenarium</i> , Guss.	210
		<i>asperrimum</i> , M. B.	205
		<i>calycinum</i> , Viv.	210
		<i>claratum</i> , W.	205
		<i>Columnæ</i> , Ehr.	206

	Pag.		Pag.
Echium Creticum , Fl. Gr.	210	Elæagnus hortensis , M. B.	1056
<i>diffusum</i> , Fl. Gr.	207	<i>Orientalis</i> , L.	1056
<i>distachyrum</i> , Viv.	207	Elephas Columnæ , Guss.	478
<i>elegans</i> , Lehm.	207	<i>incurva</i> , Don.	478
<i>elongatum</i> , Poir.	206	<i>recta</i> , Don.	478
<i>expansum</i> , B. et Haussk.	207	Embelia Burmanni , Retz	45
<i>glomeratum</i> , Poir.	206	Emex , Neck.	1005
<i>glomeratum</i> , Led.	205	<i>spinuosus</i> , L.	1005
<i>hispidum</i> , Fl. Gr.	207	EMPETRACEÆ.	1144
<i>humile</i> , Desf.	158	Empetrum , L.	1145
<i>Italicum</i> , L.	205	<i>nigrum</i> , L.	1145
<i>linearifolium</i> , C. Koch	205	<i>Enchyliæna Ægyptiaca</i> , Spr.	944
<i>longiflorum</i> , Bert.	222	<i>Eranthemum Ecbolium</i> , And.	526
<i>longifolium</i> , Del.	208	Eremostachys , Bge.	793
<i>macranthum</i> , Viv.	209	<i>acanthocalyx</i> , Boiss.	795
<i>Orientale</i> , C. A. Mey.	209	<i>adenantha</i> , J. et Sp.	793
<i>Orientale</i> , L.	204	<i>Aralensis</i> , Bge.	796
<i>papillosum</i> , C. Koch	205	<i>glabra</i> , Boiss.	794
<i>parviflorum</i> , Mœnch	210	<i>hyoscyamoides</i> , B. et	
<i>pictum</i> , Jan	207	<i>Buhse</i>	796
<i>plantagineum</i> , L.	208	<i>laciiniata</i> , L.	793
<i>prostratum</i> , Del.	207	<i>lævigata</i> , Bye.	794
<i>prostratum</i> , Ten.	210	<i>Lehmanniana</i> , Bge.	795
<i>pustulatum</i> , Fl. Gr.	206	<i>loasæfolia</i> , Bth.	795
<i>pyramidalatum</i> , D.C.	205	<i>macrocheila</i> , J. et Sp.	795
<i>Pyrenaicum</i> , Desf.	205	<i>macrophylla</i> , M. et Auch.	797
<i>Rauwolfii</i> , Del.	208	<i>pulvinaris</i> , J. et Sp.	794
<i>rubrum</i> , Forsk.	207	<i>pyramidalis</i> , J. et Sp.	797
<i>rubrum</i> , Jacq.	204	<i>Stocksii</i> , Boiss.	795
<i>sericeum</i> , Vahl	207	<i>superba</i> , Bge.	796
<i>setosum</i> , Fl. Pelop.	207	<i>thyrsiflora</i> , Bth.	797
<i>setosum</i> , Vahl	209	<i>Tournefortii</i> , J. et Sp.	794
<i>Sibthorpii</i> , R. et Sch.	207	<i>Transoxana</i> , Bge.	796
<i>tinctorum</i> , Viv.	208	Eritrichium , Schrad.	242
<i>tuberculatum</i> , Link.	207	<i>nanum</i> , Vill.	242
<i>verecundum</i> , Viv.	210	<i>Persicum</i> , Boiss.	243
<i>violaceum</i> , Koch	209	Eryngium lateriflorum , Lam.	928
<i>vulgare</i> , L.	206	Erythrea , Ren.	67
<i>Edgeworthia buxifolia</i> , Falc.	32	<i>Anatolica</i> , C. Koch	67
Ehretia , L.	124	<i>Babylonica</i> , Griseb.	69
<i>floribunda</i> , Royl.	124	<i>Caspica</i> , Fisch.	67
<i>obtusifolia</i> , Hochst.	124	<i>Centaurium</i> , Pers.	68
ELEAGNACEÆ	1053	<i>latifolia</i> , Sm.	67
Elæagnus , L.	1053	<i>linearifolia</i> , Griseb.	68
<i>angustifolia</i> , L.	1056	<i>maritima</i> , Willd.	68

	Pag.		Pag.
<i>Erythrea Meyeri</i> , Bge.	67	<i>Euphorbia capitulata</i> , Rehb.	1106
<i>pulchella</i> , Fr.	67	<i>cardiophylla</i> , B. et Heldr.	1105
<i>ramosissima</i> , Pers.	68	<i>Cassia</i> , Boiss.	1108
<i>spicata</i> , L.	69	<i>cassioides</i> , Presl	1086
<i>tenuiflora</i> , Link	67	<i>caudata</i> , B. et Hanst.	1111
Eufragia , Gr.	473	<i>candidulosa</i> , Boiss.	1122
<i>latifolia</i> , L.	473	<i>cespitosa</i> , Ten.	1116
<i>viscosa</i> , L.	474	<i>charaepeplna</i> , B. et	
EUPHORBIACEÆ.	1082	<i>Gaill.</i>	1143
Euphorbia , L.	1082	<i>chamaesyce</i> , Fl. Alt.	1087
<i>Acanthothamnos</i> , H. et		<i>Chamæsyce</i> , L.	1088
<i>Sart.</i>	1105	<i>Characias</i> , Sm.	1133
<i>acuminata</i> , Lam.	1111	<i>Characias</i> , L.	1132
<i>Ægyptiaca</i> , L.	1088	<i>cheiradenia</i> , Boiss.	1119
<i>agraria</i> , M. B.	1109	<i>cheirolepis</i> , F. et M.	1089
<i>Aleppica</i> , L.	1109	<i>Cheaneyi</i> , Kl.	1118
<i>Alexandrina</i> , Del.	1124	<i>Cilicica</i> , Boiss.	1135
<i>alpigena</i> , Kern.	1104	<i>cœladenia</i> , Boiss.	1094
<i>altissima</i> , Boiss.	1096	<i>condensata</i> , F.	1109
<i>amplexicaulis</i> , Led.	1105	<i>condylocarpa</i> , M. B.	1102
<i>amygdaloïdes</i> , L.	1150	<i>congesta</i> , W.	1116
<i>Anacampseros</i> , Boiss.	1154	<i>eoniosperma</i> , B. et	
<i>Apios</i> , L.	1103	<i>Buhse</i>	1100
<i>arguta</i> , Sol.	1097	<i>counata</i> , Boiss.	1096
<i>arillata</i> , Edgew.	1087	<i>cordifolia</i> , C. A. M.	1151
<i>Armeniaca</i> , Boiss.	1118	<i>coriacea</i> , C. Koch	1094
<i>arvalis</i> , B. et Heldr.	1114	<i>cornuta</i> , Pers.	1095
<i>aspera</i> , M. B.	1101	<i>craspedia</i> , Boiss.	1133
<i>Aucherii</i> , Boiss.	1122	<i>Crelica</i> , Mill.	1152
<i>aulacosperma</i> , Boiss.	1111	<i>cristata</i> , Dietr.	1155
<i>Baselicis</i> , Ten.	1128	<i>curtifolia</i> , B. et Ch.	1134
<i>Belgradica</i> , Forsk.	1136	<i>Cybirensis</i> , Boiss.	1098
<i>Berythea</i> , B. et Bl.	1107	<i>Cyparissias</i> , L.	1125
<i>bialata</i> , Link	1107	<i>Cypria</i> , Boiss.	1096
<i>biglandulosa</i> , Desf.	1136	<i>Dalmatica</i> , Vis.	1115
<i>bothriosperma</i> , Boiss.	1119	<i>Damascena</i> , Boiss.	1130
<i>bounophila</i> , Boiss.	1121	<i>decipiens</i> , B. et Buhse	1119
<i>Bungei</i> , Boiss.	1095	<i>decumbens</i> , W.	1086
<i>buplevroides</i> , W.	1126	<i>deflexa</i> , Fl. Gr.	1121
<i>Burmanniana</i> , Gay.	1089	<i>dendroides</i> , L.	1093
<i>Buhsei</i> , Boiss.	1124	<i>densa</i> , Schr.	1091
<i>Cajogala</i> , Ehrh.	1124	<i>densa</i> , Sch. et Ky.	1105
<i>calcarea</i> , Coss.	1116	<i>densifolia</i> , C. Koch	1136
<i>calendulaefolia</i> , Del.	1096	<i>denticulata</i> , Lam.	
<i>canescens</i> , L.	1088	<i>1135 et Add.</i>	

Pag.		Pag
Euphorbia Djimilensis, Boiss. 1104		Euphorbia Hohenackeri, Hochst. 1124
<i>dichotoma</i> , Forsk. 1157		humifusa, W. 1086
<i>diffusa</i> , Jacq. 1110		Iberica, Boiss. 1127
<i>diversifolia</i> , W. 1125		Inderiensis, Less. 1091
<i>dracunculoides</i> , Lam. 1110		Indica, Lam. 1086
<i>dulcis</i> , Jacq. 1103		Ispahanica, Boiss. 1094
<i>dulcis</i> , Fl. Gr. 1104		iteophylla, Boiss. 1151
<i>dumosa</i> , Boiss. 1104		Kotschyana, Fenzl 1152
<i>echinocarpa</i> , Sieb. 1099		Kurdica, B. et Haussk. 1121
<i>epithymoides</i> , Jacq. 1102		lanata, Sieb. 1092
<i>erinacea</i> , B. et Ky. 1105		lanceolata, Rottb. 1110
<i>eriocarpa</i> , Ber. 1152		Larica, Boiss. 1090
<i>erioclada</i> , B. et Heldr. 1102		lasiocarpa, C. Koch 1096
<i>eriphora</i> , Boiss. 1097		lateriflora, J. et Sp. 1092
<i>erubescens</i> , Boiss. 1152		latifolia, Hoh. 1127
<i>erythradenia</i> , Boiss. 1117		Ledebourii, Boiss. 1115
<i>erythrodon</i> , B. et Heldr. 1116		leiosperma, Fl. Gr. 1123
<i>Esula</i> , L. 1125		leptocaula, Boiss. 1125
<i>exigua</i> , L. 1110		linariæfolia, Lam. 1124
<i>exigua</i> , var. Stev. 1115		literata, Jacq. 1099
<i>falcata</i> , L. 1111		longifolia, Guld. 1126
<i>Forskahlei</i> , J. Gay. 1087		Lorentii, Hochst. 1150
<i>Forskahlei</i> , var. Gay. 1088		lucida, W. K. 1127
<i>fossulata</i> , B. et Gaill. 1112		Lycia, Boiss. 1135
<i>fragilis</i> , Decaisn. 1087		maerocarpa, B. et Buhse 1101
<i>Gaillardotii</i> , B. et Bl. 1097		macroceras, F. et M. 1131
<i>Galilæa</i> , Boiss. 1111		macroclada, Boiss. 1129
<i>Gerardiana</i> , Jacq. 1124		macrostegia, Boiss. 1132
<i>glaberrima</i> , C. Koch 1151		malacophylla, Clarke 1092
<i>glareosa</i> , M. B. 1129		malleata, Boiss. 1118
<i>glaucia</i> , Ehr. 1110		Marschalliana, Boiss. 1155
<i>glaucescens</i> , W. 1124		Massiliensis, D. C. 1088
<i>Græca</i> , B. et Spr. 1115		megalautha, Boiss. 1095
<i>granulata</i> , Forsk 1087		microsciadia, Boiss. 1120
<i>granulata</i> , var. F. 1086		microsphæra, Boiss. 1098
<i>Haußknechtii</i> , Boiss. 1107		modesta, Boiss. 1124
<i>hebecarpa</i> , Boiss. 1126		Mohammerensis, Boiss. 1098
<i>Heldreichii</i> , Orph. 1153		monticola, Boiss. 1122
<i>helioscopia</i> , L. 1107		mucronata, Lam. 1111
<i>herniariæfolia</i> , W. 1125		mucronata, Cl. 1137
<i>heteradena</i> , B. et Buhse 1117		muricata, M. B. 1101
<i>heteradena</i> , J. et Sp. 1094		Myrsinites, L. 1134
<i>heterophylla</i> , Desf. 1125		Napolitana, Ten. 1125
<i>Hierosolymitana</i> , Boiss. 1104		neriifolia, L. 1090
<i>Hohenackeri</i> , Orph. 1129		

	PAG.
<i>Euphorbia Nicæensis v. gla-</i>	
<i>reosa</i> , K.	1129
<i>Nicæensis v. minor</i> ,	
<i>Led.</i>	1118
<i>nitens</i> , Trev.	1128
<i>Noëana</i> , Boiss.	1129
<i>notadenia</i> , Boiss.	1101
<i>obliquata</i> , Forsk.	1124
<i>oblongata</i> , Griseb.	1102
<i>oblongata</i> , C. Koch.	1100
<i>oblongifolia</i> , C. Koch.	1150
<i>ovovata</i> , Decaisn.	1117
<i>obscura</i> , Loïs.	1111
<i>obtusifolia</i> , Lam.	1125
<i>odontophylla</i> , Boiss.	1094
<i>Orientalis</i> , L.	1100
<i>Orphanidis</i> , Boiss.	1128
<i>osyridea</i> , Boiss.	1092
<i>oxyodonta</i> , B. et Haussk.	1108
<i>pallida</i> , W.	1127
<i>palustris</i> , L.	1100
<i>Pannonica</i> , Host	1129
<i>Paralias</i> , L.	1150
<i>parvula</i> , Del.	1109
<i>parvula</i> , C. Koch	1114
<i>Peplis</i> , L.	1086
<i>peploides</i> , Gou.	1112
<i>Peplus</i> , L.	1112
<i>Persepolitana</i> , Boiss.	1120
<i>Pestalozzae</i> , Boiss.	1116
<i>petrophila</i> , C. A. M.	1118
<i>phyllatosperrma</i> , B. et	
<i>Gaill.</i>	1109
<i>pilosa</i> , L.	1096
<i>pilosa</i> , Sm.	1106
<i>pilosa</i> , Sm.	1102
<i>pinea</i> , L.	1116
<i>pinifolia</i> , W.	1109
<i>platiphylla</i> , L.	1099
<i>plebeia</i> , Boiss.	1120
<i>polycaula</i> , Boiss.	1149
<i>polygonisperma</i> , Gr.	1087
<i>Portlandica</i> , Son.	1125
<i>Postii</i> , Boiss.	1089
<i>procera</i> , M. B.	1096

	PAG.
<i>Euphorbia Provincialis</i> , W.	1125
<i>pseudochamarsyce</i> , F.	
<i>et M.</i>	1087
<i>pterococca</i> , Brot.	1107
<i>pubescens</i> , Vahl	1106
<i>pubigera</i> , Friv.	1105
<i>pumila</i> , Fl. Gr.	1125
<i>punctata</i> , Del.	1114
<i>pycnophylla</i> , C. Koch	1125
<i>pygmaea</i> , Ledeb.	1115
<i>pygmæa</i> , F. et M.	1091
<i>Ragusana</i> , Rehb.	1116
<i>retusa</i> , Rehb.	1110
<i>retusa</i> , Forsk.	1093
<i>Reuteriana</i> , Boiss.	1145
<i>rigida</i> , M. B.	1156
<i>Röseri</i> , Orph.	1154
<i>rotundata</i> , Lor.	1155
<i>rotundifolia</i> , Loisel.	1112
<i>rubra</i> , Cav.	1111
<i>rubra</i> , D.C.	1110
<i>rud.</i> <i>atis</i> , Schlede	1114
<i>rumicitoria</i> , Boiss.	1151
<i>rupestris</i> , Friv.	1128
<i>rytidosperma</i> , B. et Bal.	1106
<i>salicifolia</i> , Host	1127
<i>salicifolia</i> var. <i>Led.</i>	1127
<i>sanguinea</i> , H. St.	1088
<i>saxatilis</i> , M. B.	1118
<i>schizoceras</i> , Boiss.	1150
<i>Sel ottiana</i> , Boiss.	1105
<i>segetalis</i> , L.	1115
<i>sessiliflora</i> , B. et Spr.	1102
<i>seticornis</i> , Poir.	1125
<i>Sibthorpii</i> , Boiss.	1155
<i>simplex</i> , C. Koch	1115
<i>Sinuica</i> , Hochst.	1115
<i>soliflora</i> , Vis.	1106
<i>sororia</i> , Schr.	1091
<i>spinosa</i> , Fl. Gr.	1105
<i>squamosa</i> , W.	1101
<i>stellulata</i> , Salzm.	1107
<i>Stockiana</i> , Boiss.	1094
<i>striata</i> , Boiss.	1120
<i>striatella</i> , Boiss.	1120

	Pag.		Pag.
<i>Euphorbia stricta</i> , L.	1099	<i>Euphrasia serotina</i> , Lam.	476
<i>subhastata</i> , V. et Panc.	1128	<i>Tatarica</i> , Fisch.	472
<i>subtuberculata</i> , C. A. M.	1098	Eurotia , Ad.	917
<i>sylvatica</i> , Jacq.	1150	<i>ceratoides</i> , L.	917
<i>Syriaca</i> , Spr.	1092	<i>ferruginea</i> , Boiss.	918
<i>Syssiprensis</i> , C. Koch	1129	<i>Euxolus caudatus</i> , Moq.	992
<i>Szovitsii</i> , F. et M.	1113	<i>oleraceus</i> , Moq.	991
<i>Talyschensis</i> , B. et Buhse	1101	<i>polygamus</i> , Moq.	991
<i>Teliranica</i> , Boiss.	1117	<i>viridis</i> , Moq.	991
<i>tenuifolia</i> , M. B.	1125	Evolvulus , L.	113
<i>Terracina</i> , Griseb.	1115	<i>alsinoides</i> , L.	113
<i>Terracina</i> , L.	1123	<i>fugacissimus</i> , Hochst.	113
<i>tetraceras</i> , Szov.	1150	<i>linifolius</i> , L.	113
<i>thamnoidea</i> , Boiss.	1104	<i>virgatus</i> , Chois.	92
<i>thymifolia</i> , Burm.	1089	<i>Exacum filiforme</i> , W. H.	66
<i>thyrsiflora</i> , Griseb.	1128		
<i>Tigridis</i> , Boiss.	1088	Fagus , Tourn.	1175
<i>tinctoria</i> , Boiss.	1129	<i>Castanea</i> , L.	1175
<i>Transylvanica</i> , Schur	1128	<i>sylvatica</i> , L.	1175
<i>tricuspidata</i> , Lap.	1110	<i>Faldermannia parviflora</i> , Tr.	587
<i>Tureomanica</i> , Boiss.	1087	<i>Taurica</i> , Tr.	588
<i>Turczaninowii</i> , K. et Kir.	1092	Ficus , L.	1153
<i>Uralensis</i> , Fisch.	1124	<i>Carica</i> , L.	1154
<i>Valentina</i> , Ort.	1126	<i>Cypria</i> , Rauw.	1155
<i>valerianæfolia</i> , Desf.	1096	<i>Johannis</i> , Boiss.	1154
<i>verrucosa</i> , L. part.	1105	<i>morisfolia</i> , Huss.	1155
<i>verrucosa</i> , L. Mant.	1099	<i>palmata</i> , Schw.	1155
<i>verticillata</i> , Orph.	1136	<i>Persica</i> , Boiss.	1154
<i>villosa</i> , W. K.	1099	<i>pseudosycomorus</i> , Dec.	1155
<i>virgata</i> , W. K.	1126	<i>Sycomorus</i> , L.	1155
<i>viridiflora</i> , W. K.	1104	<i>virgata</i> , Roxb.	1155
<i>Wittmanni</i> , Boiss.	1095		
Euphrasia , L.	472	Fontanesia , Labill.	38
<i>frutescens</i> , Sieb.	477	<i>phyllireoides</i> , Lab.	38
<i>glutinosa</i> , M. B.	475		
<i>gracilis</i> , Fr.	472	Forskahlea , L.	1151
<i>latifolia</i> , L.	475	<i>tenacissima</i> , L.	1151
<i>leianthera</i> , Fenzl	475		
<i>linifolia</i> , L.	475	Fraxinus , L.	39
<i>lutea</i> , L.	475	<i>Cappadocica</i> , Jan.	39
<i>minima</i> , Schl.	475	<i>coriariæfolia</i> , Scheele	40
<i>Odontites</i> , L.	476	<i>diversifolia</i> , Roch.	39
<i>officinalis</i> , L.	472	<i>excelsior</i> , L.	39
<i>pygmaea</i> , C. Koch	473	<i>lentiscifolia</i> , Desf.	40
<i>Salisburgensis</i> , Funk	473	<i>millelacuum</i> , C. Koch	39
		<i>Moercroftiana</i> , Wall.	41
		<i>Ornus</i> , L.	39
		<i>Ornus</i> , Pall.	40

	PAG.		PAG.
Fraxinus oxycarpa , Willd.	40	<i>Gentiana fimbriata</i> , W.	75
<i>oxyphylla</i> , M. B.	40	<i>gelida</i> , M. B.	75
<i>parvifolia</i> , Lam.	40	<i>Germanica</i> , var Gr.	70
<i>Persica</i> , Boiss.	41	<i>holosteoides</i> , Sch.	71
<i>petiolulata</i> , Boiss.	40	<i>humilis</i> , Stev.	72
<i>rostrata</i> , Gass.	40	<i>involucrata</i> , Fr.	71
<i>rotundifolia</i> , Lam.	59	<i>lutea</i> , L.	69
<i>Sogdiana</i> , Bge.	41	<i>minutissima</i> , Boiss.	71
<i>Syriaca</i> , Boiss.	40	<i>nivalis</i> , L.	74
<i>xanthoxyloides</i> , Wall.	41	<i>nutans</i> , Bge.	72
Freirea alsinefolia , Gaud.	1151	<i>obtusifolia</i> , Willd.	70
Galeobdolon luteum , Huds.	756	<i>Olivieri</i> , Gr.	76
Galeopsis , L.	752	<i>Pneumonanthe</i> , L.	74
<i>Galeobdolon</i> , L.	756	<i>prostrata</i> , Hänke	72
<i>glandulosa</i> , C. Koch	752	<i>punctata</i> , L.	69
<i>Ladanum</i> , L.	752	<i>Pyrenaica</i> , L.	71
<i>latifolia</i> , Hoffm.	752	<i>schistocalyx</i> , C. Koch	76
<i>pubescens</i> , Gr.	755	<i>septemfida</i> , Pall.	74
<i>Tetrahit</i> , L.	752	<i>Stelleriana</i> , Cham.	77
<i>versicolor</i> , Curt.	755	<i>tenella</i> , Clarke	71
Gamanthus , Bge.	979	<i>umbellata</i> , M. B.	71
<i>commixtus</i> , Bge.	980	<i>verna</i> , L.	75
<i>gamocarpus</i> , Moq.	980	Glechoma , L.	670
<i>ovinus</i> , Bge.	980	<i>erecta</i> , Roxb.	844
<i>pilosus</i> , Pall.	980	<i>hederacea</i> , L.	670
GENTIANEÆ	64	<i>hirsuta</i> , W. K.	670
Gentiana , L.	69	Gieseckia , L.	895
<i>aestiva</i> , R. et Sch.	75	<i>pharmaceoides</i> , L.	896
<i>Amarella</i> , Sm.	70	Girgensohnia , Bge.	967
<i>angulosa</i> , M. B.	75	<i>diptura</i> , Bge.	968
<i>aquatica</i> , Pall.	72	<i>fruticulosa</i> , Bge.	967
<i>aselepiadea</i> , G.	76	<i>gypsophyloides</i> , Bge.	967
<i>aurea</i> , L.	71	<i>heteroptera</i> , Bge.	967
<i>Biebersteinii</i> , Bge.	70	<i>imbricata</i> , Bge.	968
<i>Boissieri</i> , Sch.	75	<i>oppositillora</i> , Pall.	967
<i>calycina</i> , Boiss. et Haussk.	75	<i>Pallasii</i> , Bge.	967
<i>Caucasica</i> , M. B.	70	Glaux , L.	7
<i>ciliata</i> , L.	75	<i>maritima</i> , L.	7
<i>consobrina</i> , Sch.	71	GLORULARIEÆ	528
<i>cordifolia</i> , C. Koch	75	Globularia , L.	528
<i>eruciata</i> , L.	76	<i>Alypum</i> , Del.	551
<i>Djimilensis</i> , C.	72	<i>Alypnum</i> , L.	550
<i>filiformis</i> , L.	66	<i>Arabica</i> , J. et Sp.	550
<i>fimbriaplica</i> , C. Koch	75	<i>cordifolia</i> , L.	529
		<i>macrantha</i> , C. Koch	529

	Pag.		Pag.
Globularia Orientalis, L.	551	Halanthium	981
<i>pallida</i> , C. Koch	529	<i>rigidum</i> , Bge.	982
<i>stygia</i> , Orph.	550	<i>robustum</i> , Bge.	982
<i>trichocalyx</i> , St.	551	<i>spicatum</i> , Bge.	982
<i>trichosantha</i> , F. et M.	529	Haiarchon , Bge.	979
<i>vulgaris</i> , L.	528	<i>vesiculosus</i> , Moq.	979
<i>vulgaris v. Bithynica</i> , Gr.	529	Halimocnemis , C. A. M.	981
<i>Willkommii</i> , Nym.	529	<i>alternifolia</i> , Moq.	973
Glossonema , Decaisn.	60	<i>Belangeri</i> , Moq.	982
<i>Boveanum</i> , Del.	61	<i>brachiata</i> , C. A. M.	974
Gomphocarpus , R. Br.	61	<i>crassifolia</i> , C. A. Mey.	975
<i>fruticosus</i> , L.	61	<i>glaucia</i> , Bge.	975
<i>fruticosus</i> , Decaisn.	62	<i>hirsuta</i> , C. Koch	981
<i>Sinaicus</i> , Boiss.	61	<i>hispida</i> , C. A. M.	974
Gomphrena sessilis, L.	996	<i>Hohenackeri</i> , Presl	974
Goniolimon , Boiss.	855	<i>Karelini</i> , Moq.	976
<i>collinum</i> , Griseb.	855	<i>Kowalevskyi</i> , Scz.	982
<i>Dalmaticum</i> , Presl	855	<i>Kulpiana</i> , C. Koch	981
<i>graminifolium</i> , Ait.	854	<i>maeranthera</i> , Bge.	977
<i>Sartorii</i> , Boiss.	856	<i>malacophylla</i> , C. A. M.	981
<i>Tataricum</i> , L.	854	<i>mollissima</i> , Bge.	977
Gratiola , L.	426	<i>oppositiflora</i> , Eichw.	975
<i>officinalis</i> , L.	426	<i>pilosa</i> , Moq.	976
<i>Monnieria</i> , L.	426	<i>purpurea</i> , Moq.	983
<i>Gregoria aretioides</i> , Duby	19	<i>pycnantha</i> , C. Koch	982
<i>Aucherii</i> , Duby	19	<i>tomentosa</i> , Moq.	964
<i>Michauxii</i> , Duby	21	<i>triandra</i> , Moq.	975
Gymnandra , Pall.	527	<i>villosa</i> , Kar. et Kir.	976
<i>Armena</i> , Boiss.	528	<i>volvox</i> , C. A. M.	975
<i>stolonifera</i> , C. Koch	527	Halimus pedunculatus, Wallr.	912
Haberlea , Friv.	82	Halocharis , Moq.	974
<i>Heldreichii</i> , Boiss	85	<i>Abichii</i> , C. Koch	982
<i>Rhodopensis</i> , Fr.	82	<i>clavata</i> , Bge.	975
Hablitzia , M. B.	987	<i>gamocarpa</i> , Moq.	980
<i>tamnooides</i> , M. B.	987	<i>hispida</i> , C. A. M.	974
Halanthium , C. Koch	981	<i>pycnantha</i> , C. Koch	982
<i>Aucherii</i> , Moq.	981	<i>sulphurea</i> , Moq.	975
<i>Kulpianum</i> , C. Koch	981	<i>sulphurea</i> , Moq. ex parte	976
<i>lanatum</i> , Bge.	982	<i>vesiculososa</i> , Moq. part.	980
<i>lilacinum</i> , Bge.	982	<i>vesiculososa</i> , Moq.	979
<i>Manamense</i> , Bge.	981	<i>violacea</i> , Bge.	975
<i>molle</i> , Bge.	981	Halocnemum , M. B.	956
<i>purpureum</i> , Moq.	983	<i>Caspicum</i> , M. B.	935
<i>rarifolium</i> , C. Koch	982	<i>strobilaceum</i> , Pall.	956
		Halogeton , C. A. Mey.	985
		<i>alopecuroides</i> , Del.	985

INDEX.

1220

	Pag.		Pag.
Halogeton glomeratus , M. B.	985	Heliotropium Cabulicum , Bge.	143
<i>Jaubertianus</i> , Bal.	954	<i>calcareum</i> , Stocks	128
<i>Olivieri</i> , Moq.	954	<i>callosum</i> , Spr.	142
<i>purpureus</i> , Fenzl	985	<i>Carduchorum</i> , C. Koch	150
<i>rarifoliosus</i> , Fenzl	982	<i>Carmaticum</i> , Bge.	142
<i>spinosissimus</i> , C. A. M.	965	<i>Chorassanicum</i> , Bge.	129
<i>tetragonus</i> , Moq.	957	<i>circinatum</i> , Gr.	157
<i>tetrandrus</i> , Moq.	957	<i>commutatum</i> , C. Koch	150
Halopeplis , Bge.	954	<i>confertiflorum</i> , B. et Noë	154
<i>amplexicaulis</i> , Vahl	954	<i>Cordofanum</i> , H. et St.	144
<i>persfoliata</i> , Forsk.	955	<i>crassitolinum</i> , Boiss.	128
<i>pygmæa</i> , Pall.	954	<i>crispum</i> , Desf.	147
Halostachys , C. A. M.	955	<i>dasyarpum</i> , Led.	140
<i>Caspia</i> , Moq.	955	<i>dentieulatum</i> , B. et	
<i>Caspica</i> , Pall.	955	<i>Haussk.</i>	145
<i>persfoliata</i> , Moq.	955	<i>desertorum</i> , B. et Buhse	159
<i>Songarica</i> , Sch.	954	<i>dissimiliflorum</i> , Boiss.	152
<i>Halothamnus Bottæ</i> , J et Sp.	964	<i>ditrichum</i> , Stocks	143
Halotis , Bge.	978	<i>dolosum</i> , Not.	151
<i>oculata</i> , Bge.	978	<i>Eichwaldi</i> , St.	151
Haloxylon , Bge.	948	<i>ellipticum</i> , Led.	151
<i>Ammodendron</i> , C. A. M.	948	<i>eremobium</i> , Bge.	141
<i>artienatum</i> , Cav.	949	<i>eriocarpum</i> , Lehm.	142
<i>Griffithii</i> , Moq.	950	<i>eriocarpum</i> , Del.	145
<i>multiflorum</i> , Moq.	949	<i>Europeum</i> , L.	150
<i>Persicum</i> , Bge.	948	<i>gracillimum</i> , Bge.	155
<i>recurvum</i> , Wall.	949	<i>grandiflorum</i> , Auch.	155
<i>salicornicum</i> , Moq.	949	<i>Griffithii</i> , Boiss.	155
<i>Hamiltonia elegans</i> , Rehb.	1059	<i>Halame</i> , Boiss et Buhse	141
Heliocarya , Bge.	279	<i>Haussknechtii</i> , Bge.	157
<i>uonandra</i> , Bge.	279	<i>intermedium</i> , Andr.	155
<i>Heliophytum Kotschyi</i> , Bge.	147	<i>Kotschyanum</i> , Bge.	156
Heliotropium , L.	125	<i>Kowalevskyi</i> , Stegl.	152
<i>aenitiflorum</i> , Kar. et Kir.	159	<i>Kumentse</i> , Bge.	159
<i>Afghanum</i> , Boiss.	145	<i>Kunzei</i> , Lehm.	150
<i>Agdense</i> , Bge.	154	<i>lasiocarpum</i> , F. et M.	151
<i>Aleppicum</i> , Boiss.	129	<i>lineatum</i> , Del.	142
<i>Arbainense</i> , Fres.	146	<i>littorale</i> , Stev.	151
<i>Ancheri</i> , D C.	142	<i>Inteum</i> , Poir.	141
<i>biannulatum</i> , Bge.	155	<i>macrocarpum</i> , Guss.	151
<i>bicolor</i> , H. et St.	144	<i>Mammanense</i> , Bge.	156
<i>Bucconi</i> , Guss.	151	<i>Mesianum</i> , Bge.	157
<i>Bovei</i> , Boiss.	156	<i>mieranthos</i> , Pall.	158
<i>Brahuicum</i> , Stocks	140	<i>micranthum</i> , Bge.	158
<i>breviliimbé</i> , Boiss.	145	<i>minutiflorum</i> , Bge.	157

	Pag.		Pag.
<i>Heliotropium neglectum</i> , B. c. e.	137	<i>Heterocaryum lavigatum</i> ,	248
<i>Niloticum</i> , A.D.C.	130	A.D.C.	248
<i>Noëanum</i> , Boiss.	128	<i>pachypodium</i> , A.D.C.	248
<i>odorum</i> , Stev.	135	<i>rigidum</i> , A.D.C.	248
<i>ophioglossum</i> , Stocks	145	<i>Szovitsianum</i> , A.D.C.	248
<i>pallens</i> , Del.	132	<i>Hippophae</i> , L.	1055
<i>parviflorum</i> , Stev.	151	<i>rhamnoides</i> , L.	1055
<i>Persicum</i> , Burm.	147	<i>Horaninowia</i> , F. et M.	947
<i>Persicin</i> , Lam.	147	<i>anomala</i> , Moq.	947
<i>Prusianum</i> , Bge.	156	<i>juniperina</i> , C. A. M.	947
<i>radula</i> , F. et M.	144	<i>ulicina</i> , F. et M.	947
<i>ramosissimum</i> , Sieb.	147	<i>Hornemannia bicolor</i> , W.	424
<i>rariiflorum</i> , Stocks	144	<i>Hottonia</i> , L.	5
<i>rotundifelium</i> , Sieb.	144	<i>palustris</i> , L.	5
<i>samoliflorum</i> , Bge.	157	<i>Humulus</i> , L.	1152
<i>Schrenkianum</i> , Led.	140	<i>Lupulus</i> , L.	1152
<i>sericocarpum</i> , Bge.	142	<i>Hygrophila</i> , R. Br.	518
<i>Smyrneum</i> , Bge.	156	<i>polysperma</i> , Roxb.	519
<i>Sogdianum</i> , Bge.	146	<i>Hymenocrater</i> , F. M.	675
<i>Stevenianum</i> , Andr.	151	<i>Aucherii</i> , J. et Sp.	676
<i>strictum</i> , Led.	151	<i>bituminosus</i> , F. M.	676
<i>strigosum</i> , W.	145	<i>calycinus</i> , Boiss.	676
<i>subcanescens</i> , Stev.	150	<i>elegans</i> , Bge.	678
<i>suaveolens</i> , M. B.	155	<i>Haussknechtii</i> , Boiss.	677
<i>Sultanense</i> , Bge.	158	<i>ineanii</i> , Bge.	678
<i>supinum</i> , L.	127	<i>longifloris</i> , Bth.	677
<i>Syenites</i> , Spr.	150	<i>maerophyllus</i> , Bge.	677
<i>Szovitsii</i> , Stev.	152	<i>pallens</i> , Bge.	676
<i>Teheranicum</i> , Bge.	156	<i>paniciatius</i> , Bge.	677
<i>tenuiflorum</i> , Guss.	150	<i>secundiflorus</i> , J. et Sp.	676
<i>thymoides</i> , J. et Sp.	144	<i>sessilifolius</i> , Bth.	678
<i>Transoxanum</i> , Bge.	159	<i>Hyoscyamus</i> , L.	295
<i>trichostomum</i> , Bge.	140	<i>agrestis</i> , Kit.	295
<i>tuberculatum</i> , Boiss.	147	<i>albus</i> , L.	295
<i>Turcicum</i> , Friv.	155	<i>aurens</i> , L.	295
<i>undulatum</i> , Vahl	147	<i>aureus</i> , All.	296
<i>undulatum</i> , var. Boiss	147	<i>betæfolius</i> , Lam.	296
<i>villosum</i> , Willd.	155	<i>bipinnatisectus</i> , Boiss.	297
<i>Hemiadelphis polysperma</i> ,		<i>Camerarii</i> , F. et M	295
Nees	519	<i>Canariensis</i> , Ker.	296
<i>Herpestis</i> , Gærtn.	426	<i>Ceticus</i> , Tonrn.	296
<i>Monnieria</i> , L.	426	<i>Datora</i> , Forsk.	294
<i>Hesiodia bicolor</i> , Mœnch	707	<i>icanus</i> , Stocks	294
<i>Heterocaryum divaricatum</i> ,		<i>major</i> , Mill.	296
Stocks	25	<i>micranthus</i> , Led.	294

	PAG.		PAG.
Hyoscyamus minor , Mill.	296	Juglans fraxinifolia , Lam.	1160
<i>mutiens</i> , L.	295	<i>pterocarpa</i> , W.	1161
<i>niger</i> , L.	294	<i>regia</i> , L.	1160
<i>Orientalis</i> M. B.	295	Justicia , L.	525
<i>pallidus</i> , W. K.	295	<i>Chinensis</i> , Roxb.	526
<i>Persicus</i> , B. et Buhse	295	<i>Ecbolium</i> , L.	526
<i>pinuatifidus</i> , Schlecht.	295	<i>peploides</i> , Nees	525
<i>pungeus</i> , Gr.	294	<i>polysperma</i> , Roxb.	519
<i>purpureus</i> , Griseb.	295	<i>quinquangularis</i> , Roxb.	525
<i>pusillus</i> , L.	295	<i>Vahlii</i> , Roth	525
<i>reticulatus</i> , L.	295		
<i>Senecionis</i> , W.	297		
<i>Syssirensis</i> , C. Koch	295	Kalidium , Moq.	953
<i>varians</i> , Vis.	296	<i>Arabicum</i> , Moq.	953
Hypogomphia , Bge.	679	<i>Caspie nn</i> , L.	953
<i>nana</i> , Bth.	679	<i>foliatum</i> , Pall.	954
Hyssopus , L.	584	Kanahla , R. Br.	57
<i>angustifolius</i> , M. B.	584	<i>Dilili</i> , Decaisn	58
<i>Caucasicus</i> , Spr.	584	Kir ilowia , Bge.	919
<i>officinalis</i> , L.	584	<i>eriantha</i> , K. et Kir.	920
<i>Orientalis</i> , W.	584	<i>pilosa</i> , Bge.	919
Ilex , L.	54	Kochia , Roth.	922
<i>Aqnifolium</i> , L.	54	<i>arenaria</i> , W. K	914
Ipomæa , L.	112	<i>eana</i> , Bge	924
<i>littoralis</i> , L.	112	<i>dasyantha</i> , Schr.	924
<i>palmata</i> , Forsk.	115	<i>eriantha</i> , Bge.	920
<i>sagittata</i> , Desf.	115	<i>ericifolia</i> Viv.	927
<i>Senegalensis</i> , Lam.	115	<i>eriophora</i> , Schrad.	927
<i>Iresine Javanica</i> , Burm.	995	<i>Gritfithii</i> , Bge.	924
<i>Persica</i> , Burm.	995	<i>hirsuta</i> , L.	925
Jankæa , Boiss.	82	<i>hyssopifoli</i> , Pall.	926
<i>Heldreichii</i> , Boiss.	82	<i>latifolia</i> , Fres.	927
<i>Janthe bugulifolia</i> , Gr.	551	<i>monacantha</i> , St.	954
JASMINEÆ.	41	<i>monticola</i> , Boiss.	925
Jasminum , Tourn.	42	<i>muricata</i> , L	926
<i>fruticans</i> , L.	42	<i>Noëana</i> Boiss.	919
<i>humile</i> , L	42	<i>odontoptera</i> , Schr.	924
<i>officinale</i> , L	45	<i>prostrata</i> , L.	925
<i>revolutum</i> , Sims.	42	<i>scoparia</i> , L.	925
<i>Syriacum</i> , Boiss.	42	<i>sedoides</i> , Pall.	926
<i>Jiraseckia tenella</i> , Lam.	6	<i>stellaris</i> , Moq.	924
JUGLANDACEÆ	1159		
Juglans , L.	1160	LABIATÆ	557
		Lagochilus , Bge.	768
		<i>acutilobus</i> , Led.	771
		<i>alutaceus</i> , Bge.	770

	Pág.		Pág.
Lagochilus Aucheri, Boiss.	769	Lamium moschatum, Mill	765
<i>Cabulicus, Bth.</i>	769	<i>multifidum, L.</i>	767
<i>cuneatus, Bth.</i>	770	<i>nepetaefolium, Boiss.</i>	757
<i>glaberrimus, C. Koch</i>	771	<i>nivale, B. et Heldr</i>	757
<i>hispidus, Bél.</i>	769	<i>ochroleucum, Link</i>	765
<i>inebrians, Bge.</i>	770	<i>orthodon, Boiss.</i>	761
<i>insignis, Bél.</i>	770	<i>Persicum, B. et Buhse.</i>	765
<i>Kotschyannus, Boiss.</i>	769	<i>pictum, B. et Heldr.</i>	758
<i>macracanthus, F.</i>	770	<i>plebeium, Boiss.</i>	761
Lallemantia, F. M.	675	<i>plebeium, Boiss.</i>	758
<i>azurea, B. et Huet</i>	675	<i>Ponticum, B. et Bal.</i>	766
<i>canescens, L.</i>	675	<i>purpureum, L.</i>	766
<i>Iberica, M. B.</i>	674	<i>rectum, Sch.</i>	757
<i>peltata, L.</i>	674	<i>reniforme, M. et Auch.</i>	758
<i>Royleana, Wals.</i>	674	<i>Robertsoni, Boiss.</i>	764
<i>sulphurea, C. Koch</i>	675	<i>rugosum, Ait</i>	763
Lamium, L.	755	<i>stenosiphon, Boiss. et</i>	
<i>album, L.</i>	765	<i>Haussk.</i>	760
<i>Aleppicum, B. et Haussk.</i>	761	<i>striatum, Fl. Gr.</i>	757
<i>alpestre, Trautv.</i>	764	<i>tenuiflorum, F. et M.</i>	759
<i>amplexicaule, L.</i>	760	<i>tomentosum, W.</i>	764
<i>Armenum, Boiss.</i>	758	<i>truncatum, Boiss.</i>	762
<i>bifidum, Cyr.</i>	760	<i>veronicaefolium, Bth. Lab.</i>	758
<i>Bithynicum, Bth.</i>	736	<i>veronicaefolium, Bth.</i>	
<i>calycinum, Urv</i>	765	<i>Ann. Sc.</i>	758
<i>erinitum, M. et Auch.</i>	765	<i>vestitum, Bth.</i>	764
<i>Cylleneum, B. et Heldr</i>	757	Lantana, L.	532
<i>cymbalariaefolium, Boiss.</i>	759	<i>alba, Mill.</i>	532
<i>Ehrenbergii, B. et R.</i>	761	<i>collina, Dec.</i>	532
<i>eriocephalum, Bth.</i>	762	<i>dubia, Royle</i>	532
<i>erythrotrichum, Boiss.</i>	769	<i>Indica, Roxb.</i>	532
<i>filicaule, Boiss.</i>	765	Lathraea, L.	493
<i>galactophyllum, B. et</i>		<i>Phelipea, L.</i>	494
<i>Huet</i>	766	<i>Rhodopea, Dingl.</i>	493
<i>Galeobdolon, L.</i>	756	<i>squamaria, L.</i>	493
<i>Garganicum, var. Gr.</i>	757	LAURINEÆ	1057
<i>glechomoides, Sm.</i>	757	Laurus, L.	1057
<i>Gundelsheimeri, C. Koch</i>	763	<i>nobilis, L.</i>	1057
<i>Lycium, Boiss.</i>	772	Lavandula, Tourn.	540
<i>maculatum, L.</i>	765	<i>atriplicifolia, Bth.</i>	541
<i>macrodon, B. et Huet</i>	761	<i>Cariensis, Boiss.</i>	540
<i>Manganottianum, Clem.</i>	758	<i>coronopifolia, Poir.</i>	542
<i>mesogæon, Heldr.</i>	760	<i>multifida, L.</i>	541
<i>microphyllum, Boiss.</i>	759	<i>pedunculata v. Cariensis,</i>	
<i>molle, B. et Orph.</i>	756	<i>Bth.</i>	541

Pag.		Pag.	
Lavandula pubescens, <i>Decaisne</i>	541	Linaria <i>alsinefolia</i> , Bth.	566
<i>spectabilis</i> , C. Koch.	541	<i>Altaica</i> , Fisch.	575
Stœchas, <i>L.</i>	540	Armeniaea, <i>Chav.</i>	581
<i>striata</i> , Del.	542	<i>arvensis</i> , <i>L.</i>	575
<i>subnuda</i> , Bth.	542	Asealonica, <i>B. et Ky.</i>	582
LENTIBULARIÆ	1	Ancheri, <i>Boiss.</i>	572
Leonurus, <i>L.</i>	753	Balansæ, <i>B. et Reut.</i>	578
Cardiaca, <i>L.</i>	755	Biebersteinii, <i>Bess.</i>	573
<i>discolor</i> , C. Koch	755	<i>bombycinæ</i> , <i>Boiss.</i> et Bt.	567
marrubiastrum, <i>L.</i>	754	Cabulica, <i>Bth.</i>	568
<i>mollis</i> , Ehr.	721	<i>calycina</i> , <i>B. et Bal.</i>	576
Persicus, <i>Boiss.</i>	754	<i>Caucasica</i> , <i>Muss.</i>	567
Tataricus, <i>Tchih.</i>	755	Chalepensis, <i>L.</i>	581
<i>vilosus</i> , Desf.	755	<i>chloræfolia</i> , <i>Rehb.</i>	577
Lepidagathis, Willd.	524	<i>cirrhosa</i> , <i>L.</i>	568
<i>calycina</i> , <i>Hochst.</i>	525	<i>Cælesyriaca</i> , <i>Boiss.</i>	572
<i>strobilifera</i> , Stocks	525	<i>commutata</i> , <i>Bernh.</i>	568
Leptadenia, R. Br.	62	<i>concolor</i> , <i>Griseb.</i>	578
<i>Delilei</i> , Decaisne	1197	<i>confertiflora</i> , <i>Bth.</i>	571
heterophylla, <i>Del.</i>	1197	<i>coridifolia</i> , <i>Desf.</i>	579
<i>pyrotechnica</i> , <i>Forsk.</i>	65	<i>cryptarum</i> , <i>B. et Haussk.</i>	585
Leptorhabdos, Schr.	470	Dalmatia, <i>L</i>	576
<i>virgata</i> , <i>Bth.</i>	470	Damaseena, <i>B. et Gaill.</i>	575
Lesquerœuxia Syriaca, <i>B. et R.</i>	471	<i>Dchorocheensis</i> , C. Koch	579
Leucas, B. Br.	773	Elatine, <i>L</i>	567
<i>cephalotes</i> , <i>Spr.</i>	779	<i>eriocalyx</i> , <i>Boiss.</i>	572
<i>inflata</i> , <i>Bth.</i>	778	<i>fastigiata</i> , <i>Chav</i>	571
<i>urticæfolia</i> , <i>Vahl</i>	778	<i>fastigiata v. angustifolia</i> , Bth	572
Ligustrum, <i>L.</i>	57	<i>flavissima</i> , <i>Boiss.</i>	578
<i>vulgare</i> , <i>L.</i>	57	<i>floribunda</i> , <i>Boiss.</i>	565
Lilac Persica, Lam	58	<i>genis itoha</i> , <i>L.</i>	577
<i>vulgaris</i> , Lam.	58	<i>glauca</i> , Ehr.	582
Limnanthemum, Gou.	65	<i>Graeca</i> , <i>B. et Ch.</i>	567
<i>Indicum</i> , <i>L.</i>	65	<i>grandiflora</i> , <i>Desf.</i>	576
<i>nymphoides</i> , <i>L.</i>	65	Griffithii, <i>Bth.</i>	570
Limoniastrum, Mœnch	874	Hœlava, <i>Forsk.</i>	581
<i>monopetalum</i> , <i>L.</i>	874	<i>hians</i> , <i>Fenzl</i>	577
Limosella, <i>L.</i>	428	<i>junccea</i> , <i>Rehb.</i>	575
<i>aquatica</i> , <i>L.</i>	428	Iconia, <i>B. et Heldr.</i>	580
<i>calycina</i> , <i>Forsk.</i>	426	Kurdica, <i>B. et Hoh.</i>	571
Linaria, Juss.	565	<i>lanigera</i> , <i>Desf.</i>	566
<i>acerbiana</i> , <i>Boiss.</i>	566	<i>lineolata</i> , <i>Boiss.</i>	579
<i>Ægyptiaca</i> , <i>L.</i>	569	<i>linifolia</i> , <i>Auct.</i>	578
<i>albifrons</i> , <i>Fl. Gr.</i>	382		

	Pag.		Pag
<i>Linaria littoralis</i> v. <i>ptero-</i>		<i>Linearia simplex</i> , D. C.	575
<i>spora</i> , F. et M.	584	<i>Smithii</i> , Boiss. et Orph.	576
<i>longipes</i> , B. et Heldr.	365	<i>speciosa</i> , Ten.	572
<i>Læselii</i> , Schw.	373	<i>sphaerocarpa</i> , Bth.	387
<i>Macedonica</i> , Gr.	376	<i>spuria</i> , L.	566
<i>macilenta</i> , <i>Decaisne</i>	569	<i>striata</i> , Ledeb.	579
<i>macroura</i> , M. B.	575	<i>Syspirensis</i> , G. Koch	578
<i>Michauxii</i> , Chav.	574	<i>triphylla</i> , L.	580
<i>mierantha</i> , Cav.	375	<i>venosa</i> , Lindl.	573
<i>microcalyx</i> , Boiss	365	<i>vulgaris</i> , Mill.	572
<i>minor</i> , L.	585	Lindenbergia , Lehm.	424
<i>minutiflora</i> , C. A. M.	335	<i>fruticosa</i> , Bth.	425
<i>modesta</i> , Boiss.	582	<i>Sinaitica</i> , <i>Decaisne</i>	424
<i>monochroma</i> , B. et Heldr.	377	<i>urticæfolia</i> , Bth.	425
<i>Nurensis</i> , B. et Haussk.	374	Lindernia pyxidaria , All.	457
<i>odora</i> , M. B.	573	Lippia , L.	552
<i>orgyalis</i> , Reut.	377	<i>nodiflora</i> , L.	552
<i>ovata</i> , Boiss.	369	<i>repens</i> , Spreng.	553
<i>pallida</i> , Sprun.	378	Lithospermum , L	216
<i>Parnassica</i> , B. et Heldr.	378	<i>angustifolium</i> , Forsk.	219
<i>parviflora</i> , Desf.	376	<i>Apulum</i> , Friv.	229
<i>Pelisseriana</i> , L	375	<i>Apulum</i> , L.	218
<i>Peloponnesiaca</i> , B. et		<i>Arnebia</i> , Del.	215
Heldr.	378	<i>arvense</i> , L.	216
<i>Persica</i> , Chav.	584	<i>bicolor</i> , Bert.	216
<i>polyclada</i> , Fenzl	378	<i>callosum</i> , Vahl	219
<i>præalta</i> , B. et Bal.	377	<i>ciliatum</i> , Lehm.	219
<i>Prestandreae</i> , Tin.	367	<i>cornutum</i> , Led.	213
<i>pterospora</i> , Boiss.	384	<i>crassifolium</i> , Lehm.	220
<i>pulchella</i> , Hoh.	379	<i>cyanochroum</i> , Boiss.	212
<i>purpurea</i> , L.	580	<i>decumbens</i> , Vent.	214
<i>pyrenophylla</i> , B. et Bal.	372	<i>densiflorum</i> , Led.	212
<i>pyramidata</i> , Lam.	370	<i>digynum</i> , Forsk.	142
<i>ramosissima</i> , Wall.	368	<i>dispermum</i> , L.	242
<i>ramosissima</i> var. <i>ovata</i> ,		<i>divaricatum</i> , Sieb.	211
Bth.	369	<i>fruticosum</i> , Fl. Gr.	220
<i>reflexa</i> , L.	380	<i>fulvidulum</i> , Bert.	217
<i>rhytidosperma</i> , Boiss.	384	<i>Gasparrinii</i> , Heldr.	219
<i>rubrifolia</i> , Rob. et Cast.	385	<i>hirsutissimum</i> , Bert.	227
<i>rupestris</i> , C. A. M.	373	<i>hispidissimum</i> , Lehm.	213
<i>rupestris</i> , Guss.	385	<i>hispidulum</i> , Fl. Gr.	219
<i>segetalis</i> , C. Koch	381	<i>hispidum</i> , Forsk.	147
<i>Sibthorpiana</i> , B. et Heldr	378	<i>incrassatum</i> , Guss.	217
<i>Sieberi</i> , Rehb.	367	<i>Kotschy</i> , Boiss. et Hoh.	219
<i>sileneifolia</i> , Fisch.	377	<i>latifolium</i> , Forsk.	219

Pag.		Pag.	
Lithospermum Lehmanni, Tin.	227	<i>Lycium særumi</i> , Miers	289
<i>Leithneri</i> , Heldr.	216	<i>Tataricum</i> , Pall.	290
<i>Mesopotamicum</i> , D.C.	218	<i>Tureomanicum</i> , Turez	290
<i>micranthum</i> , Viv.	214	<i>Lycopsis Egyptiaca</i> , L.	159
<i>officinale</i> , L.	218	<i>alpestris</i> , Lehm.	163
<i>Orientale</i> , L.	227	<i>arvensis</i> , L.	160
<i>petreum</i> , A.D.C.	221	<i>bullata</i> , Cyr.	161
<i>purpureo-caeruleum</i> , L.	218	<i>Caspica</i> , Lehm.	166
<i>Pythicum</i> , Cl.	220	<i>ciliata</i> , C.A.M.	165
<i>retortum</i> , Pall.	242	<i>decurrans</i> , C.A.M.	168
<i>Rochelii</i> , Friv.	216	<i>Dioscoridis</i> , Rauw.	279
<i>rosmarinifolium</i> , Rehb.	221	<i>elongata</i> , Lehm.	153
<i>rosmarinifolium</i> , Ten.	220	<i>erecta</i> , Urv.	161
<i>rugosum</i> , D.C.	221	<i>flavescens</i> , C.A.M.	165
<i>selosum</i> , F. et M.	221	<i>glomerata</i> , Urv.	158
<i>Sibthorpiatum</i> , Gris.	216	<i>lutea</i> , Lam.	165
<i>strigosum</i> , M.B.	218	<i>micrantha</i> , Led.	161
<i>tenuiflorum</i> , L.	217	<i>Milleri</i> , Horn.	151
<i>tenuiflorum</i> , Fl. Gr.	217	<i>mollis</i> , Vis.	228
<i>tetrastrigma</i> , Lam.	215	<i>obtusifolia</i> , W.	164
<i>tinctorium</i> , L.	227	<i>Orientalis</i> , L.	161
<i>tinctorium</i> , Vahl	215	<i>pulla</i> , L.	166
<i>tubatum</i> , Bert.	214	<i>setosa</i> , Lehm.	165
<i>vestitum</i> , Wali.	215	<i>setosa</i> , Lehm.	211
LOGANIACEÆ.	Add.	<i>Sibthorpiana</i> , R. et S. h.	170
<i>Londesia eriantha</i> , F. et M.	927	<i>Taurica</i> , Stev.	161
LORANTHACEÆ.	1067	<i>variegata</i> , L.	161
Loranthus , L.	1069	Lycopus , L.	545
<i>Acaciæ</i> , Zucc.	1070	<i>decreseens</i> , C. Koch	545
<i>Europæns</i> , L.	1069	<i>Europæns</i> , L	545
<i>Grewinkii</i> , B. et Buhse	1070	<i>exaltatus</i> , L.	545
Lycium , L.	288	<i>pinnatifidus</i> , Pall.	545
<i>abeliaeflorum</i> , Rehb.	289	Lygia , Fas	1051
<i>Arabicum</i> , Schw.	289	<i>Amani</i> , Ky.	1052
<i>Barbarum</i> , L.	289	<i>Aucherii</i> , Meisn.	1052
<i>depressum</i> , Stocks	289	<i>Cilicia</i> , Meisn.	1055
<i>Europænum</i> , L.	288	<i>multicaulis</i> , Sch.	1055
<i>foliosum</i> , Stocks	289	<i>Passerina</i> , L	1052
<i>glaucum</i> , Miers	290	<i>pubescens</i> , Guss.	1052
<i>Indicum</i> , Wight	289	Lysimachia , L.	7
<i>Mediterraneum</i> , Dan.	289	<i>anagalloides</i> , Fl. Gr.	9
<i>Oriентale</i> , Miers	289	<i>atropurpurea</i> , L.	7
<i>Persicum</i> , Miers	289	<i>atropurpurea</i> , Murr.	8
<i>Ruthenicum</i> , Murr.	290	<i>Dahurica</i> , Ledeb.	8

PAG.		PAG.	
Lysimachia dubia, Ait.	8	Marrubium globosum, M. et Auch.	696
<i>Linum stellatum</i> , L.	10	<i>Hermonis</i> , Boiss.	696
<i>numinaria</i> , L.	9	<i>heterodon</i> , B. et Bal.	699
<i>punctata</i> , Jacq.	9	<i>Kotschy</i> , B. et Hoh.	695
<i>verticillata</i> , M. B.	9	<i>leonardoi</i> , Desr.	701
<i>vulgaris</i> , L.	9	<i>Libanoticum</i> , Boiss.	696
<i>Macrosiphonia cespitosa</i> , Duby	20	<i>Intescens</i> , Boiss.	693
Macrotomia , D.C.	211	<i>mieranthum</i> , B. et Heldr.	697
<i>cephalotes</i> , A.D.C.	212	<i>microphyllum</i> , Desr.	776
<i>cyanocroa</i> , Boiss.	212	<i>Noëanum</i> , Beiss.	696
<i>echioioides</i> , L.	211	<i>paniculatum</i> , Desr.	703
<i>Majorana crassifolia</i> , Bth.	555	<i>parviflorum</i> , F. M.	704
Mandragora , Juss.	291	<i>peregrinum</i> , C. A. Mey.	704
<i>autumnalis</i> , Spr.	291	<i>peregrinum</i> , L.	694
<i>officinarum</i> , L.	291	<i>Persicum</i> , C. A. M.	700
<i>officinarum</i> , Bert.	292	<i>Pestalozzæ</i> , Boiss.	703
<i>microcarpa</i> , Bert.	292	<i>plicatum</i> , Forsk.	700
<i>vernalis</i> , Bert.	291	<i>plumosum</i> , C. A. M.	701
Marrubium , L.	692	<i>polyodon</i> , Boiss.	704
<i>acetabulosum</i> , L.	772	<i>præcox</i> , Janka	702
<i>album</i> , B. et Bal.	703	<i>procерum</i> , Bge.	700
<i>Alysson</i> , L.	700	<i>propinquum</i> , F. et M.	701
<i>anisodon</i> , C. Koch	705	<i>pseudalyssum</i> , F. M.	697
<i>Apulum</i> , Ten.	705	<i>pseudodictamnus</i> , L.	772
<i>Astracanum</i> , Jacq	695	<i>pseudodictamnus</i> , Fl. Gr.	775
<i>Astracanicum</i> , M. B.	701	<i>parpureum</i> , Bge.	695
<i>ballotoides</i> , B. et Bal.	705	<i>radiatum</i> , Del.	704
<i>Bourgei</i> , Boiss.	698	<i>rotundifolium</i> , Boiss.	698
<i>candidissimum</i> , C. A. M.	700	<i>Thessalum</i> , B. et Heldr.	699
<i>candidissimum</i> , Friv.	702	<i>trachytienum</i> , Boiss.	698
<i>candidissimum</i> , C. Koch	705	<i>undulatum</i> , Fres.	775
<i>cataractarium</i> , Desr.	699	<i>velutinum</i> , Fl. Gr.	702
<i>cephalanthur</i> , B. et Noë	694	<i>velutinum</i> , var Bth.	699
<i>circinnatum</i> , Desr.	702	<i>vulgare</i> , L.	705
<i>cærulescens</i> , Bth.	701	Marsdenia erecta , R. Br.	82
<i>condensatum</i> , Boiss.	697	Mastostigma , Stocks	51
<i>crassidens</i> , Boiss.	705	<i>varians</i> , Stocks	51
<i>erisipum</i> , Sieb.	775	Mattia , Schult.	275
<i>Creticum</i> , Mill.	694	<i>Aucherii</i> , A.D.C.	260
<i>cuneatum</i> , Russ.	705	<i>brachyantha</i> , Boiss.	275
<i>Cylleneum</i> , B. et Heldr.	702	<i>bracteata</i> , Orph.	274
<i>depauperatum</i> , B. et Bal.	694	<i>Bungei</i> , Boiss.	275
<i>fascioides</i> , B. et Bal.	697	<i>canescens</i> , A.D.C.	272
<i>Friwaldskyanum</i> , Boiss.	702	<i>cespitosa</i> , A.D.C.	273

INDEX.

1237

Pag.

Mattia corymbiformis , D.C.	262	Melissa Pulegium , Gr.	575
<i>cristata</i> , Don.	258	<i>teucrioides</i> , Boiss.	642
<i>eriantha</i> , Led.	272	<i>umbrosa</i> , M. B.	578
<i>glaustifolia</i> , Don.	262	Melittis , L.	754
<i>Graea</i> , B. et Helv.	275	<i>grandiflora</i> , Son.	755
<i>incana</i> , Led.	262	<i>Melossophyllum</i> , L.	754
<i>lanata</i> , Sch.	272	Memorialis , Lam.	1148
<i>leptophylla</i> , A.D.C.	258	<i>pentandra</i> , Lorb.	1148
<i>punctata</i> , A.D.C.	272	MENISPERMACEAE	1201
<i>Schluembergeri</i> , Boiss.	274	<i>Menispermum</i> <i>Lerba</i> , Del.	1201
<i>Schmidti</i> , Heldr.	274	Mentha , L.	542
<i>staminea</i> , F. et Sch.	270	<i>ambigua</i> , Guss.	544
<i>umbellata</i> , C. Koch	272	<i>aquatica</i> , L.	544
Mazus , Lour.	425	<i>arvensis</i> , L.	544
<i>rugosus</i> , Lour.	424	<i>caespitosa</i> , Sieb.	545
Megacaryon , Boiss.	205	<i>eriantha</i> , C. Koch	544
<i>Orientale</i> , L.	204	<i>hirsuta</i> , Sm.	544
<i>Meissurnena tonacutosa</i> , R.		<i>incana</i> , W.	545
Br.	425	<i>lavandulacea</i> , W.	545
Melampyrum , L.	480	<i>macrostachya</i> , Ten.	545
<i>arvense</i> , L.	480	<i>microphylla</i> , C. Koch	545
<i>barbatum</i> , W. K.	480	<i>mollissima</i> , Borek.	545
<i>barbatum</i> , Led.	481	<i>neglecta</i> , Ten.	545
<i>Caneasicum</i> , Bye.	481	<i>nigrescens</i> , C. Koch	545
<i>ctinatum</i> , B. et Heldr.	481	<i>Noeana</i> , Boiss.	545
<i>eristatum</i> , L.	480	<i>Pulegium</i> , L.	545
<i>elatius</i> , Reut.	481	<i>rotundifolia</i> , L.	545
<i>Heracleoticum</i> , B. et		<i>Royleana</i> , Bth.	545
<i>Orph.</i>	482	<i>rugosa</i> , Lam.	545
<i>nemorosum</i> , L.	481	<i>saliva</i> , L.	544
<i>pratense</i> , L.	482	<i>Sieberi</i> , C. Koch	545
<i>stenophyllum</i> , Boiss.	482	<i>suavis</i> , Guss.	544
<i>sylvaticum</i> , L.	482	<i>sylvestris</i> , L.	545
<i>sylvaticum</i> , C. Koch	482	<i>tomentosa</i> , Urv.	545
Melissa , L.	584	<i>viridis</i> , L.	544
<i>altissima</i> , Fl. Gr.	584	Menyanthes , L.	64
<i>Calamintha</i> , L.	577	<i>nymphoides</i> , L.	65
<i>cordifolia</i> , Pers.	584	<i>trifoliata</i> , L.	64
<i>Cretica</i> , L.	578	Mercurialis , L.	1142
<i>grandiflora</i> , L.	576	<i>annua</i> , L.	1143
<i>graveolens</i> , Host.	584	<i>ovata</i> , Sternb.	1142
<i>marifolia</i> , Griseb.	574	<i>perennis</i> , L.	1142
<i>Nepeta</i> , L.	577	<i>Messerschmidia Arguzia</i> ,	
<i>officinalis</i> , L.	584	L. fil.	125
<i>perennis</i> , Forsk.	652	<i>Microcala filiformis</i> , Link	66

	Pag.		Pag.
<i>Microcarpaea cochlearifolia</i> , Sm.	428	Molucella , L.	768
<i>Microlepis salsa</i> , Eichw.	971	<i>acutiloba</i> , Led.	771
Micromeria , Bth.	568	<i>frutescens</i> , Fl. Gr.	776
<i>alternipilosa</i> , C. Koch	566	<i>laevis</i> , L.	768
<i>barbata</i> , B. et Ky.	574	<i>microphylla</i> , Del.	776
<i>Calverti</i> , Boiss.	571	<i>Sinaitica</i> , Ehr.	777
<i>congesta</i> , B. et Haussk.	575	<i>spinosa</i> , L.	768
<i>cremnophila</i> , B. et Heldr.	570	Monotheca , A.D.C.	52
<i>cristata</i> , Gris.	570	<i>Mascatensis</i> , A.D.C.	52
<i>cymuligera</i> , Boiss. et Haussk.	569	Morus , L.	1153
<i>Cypria</i> , Ky.	571	<i>alba</i> , L.	1153
<i>Dalmatica</i> , Bth.	575	<i>Indica</i> , L.	1153
<i>elliptica</i> , C. Koch	571	<i>nigra</i> , L.	1153
<i>Græca</i> , L.	571	<i>Munbya congregata</i> , Boiss.	212
<i>hispida</i> , B. et Heldr.	572	Myosotis , L.	254
<i>Juliana</i> , L.	569	<i>alpestris</i> , Schm.	238
<i>Libanotica</i> , Boiss.	573	<i>amœna</i> , Rup.	241
<i>marifolia</i> , Bth.	574	<i>amœna</i> , Schott	257
<i>microphylla</i> , Urv.	572	<i>Apula</i> , L.	218
<i>mollis</i> , Bth.	574	<i>arenaria</i> , Schr.	259
<i>myrtifolia</i> , Boiss.	670	<i>arvensis</i> , Roth	259
<i>nervosa</i> , Desf.	569	<i>barbata</i> , M. B.	250
<i>nummularifolia</i> , Boiss.	575	<i>Bœotica</i> , Reut.	236
<i>origanifolia</i> , Bth.	579	<i>Cadmea</i> , Boiss.	236
<i>origanifolia</i> , Vis.	575	<i>cespitosa</i> , Sch.	255
<i>Persica</i> , Boiss.	573	<i>collina</i> , Rehb.	239
<i>pubescens</i> , B. et Ky.	571	<i>Cretica</i> , B. et H.	237
<i>Reinholdi</i> , Heldr.	570	<i>eyanea</i> , Reut.	237
<i>serpyllifolia</i> , Bth.	574	<i>densiflora</i> , C. Koch	237
<i>Sinaica</i> , Bth.	572	<i>heteropoda</i> , Trantv.	239
<i>Sphaciotica</i> , B. et Heldr.	572	<i>hispida</i> , Sch.	239
<i>spicigera</i> , C. Koch	566	<i>Idæa</i> , Boiss. et Heldr.	256
<i>staminea</i> , B. et Hoh.	580	<i>incrassata</i> , Guss.	237
<i>xyloorrhiza</i> , B. et Heldr.	570	<i>intermedia</i> , Link	259
Moltzia , Lehm.	220	<i>involuta</i> , Stev.	242
<i>Anatolica</i> , Boiss.	221	<i>Kiesenwetteri</i> , Heldr.	256
<i>angustifolia</i> , D.C.	222	<i>Lappula</i> , L.	249
<i>aurea</i> , Boiss.	222	<i>latisiliqua</i> , B. et K.	238
<i>cærulea</i> , W.	221	<i>lingulata</i> , Lehm.	235
<i>Libanotica</i> , Zucc.	270	<i>lithospermifolia</i> , Guss.	237
<i>petrea</i> , Trutt.	221	<i>lithospermifolia</i> , Horn.	238
<i>punctata</i> , Lehm.	221	<i>littoralis</i> , Stev.	236
<i>Sendtneri</i> , Boiss.	225	<i>macrophylla</i> , Ad.	157
		<i>marginata</i> , M. B.	232
		<i>micrantha</i> , Guss.	235

	PAG.		PAG.
Myosotis micrantha , Pall.	259	Nepeta Andrica , B. et Heldr	662
<i>microsepala</i> , Schott.	258	<i>Argolica</i> , Bory	652
<i>minutiflora</i> , B. et R.	259	<i>aristata</i> , B. et Ky.	656
<i>modesta</i> , Schott	258	<i>betonicæfolia</i> , C. A. M.	658
<i>montana</i> , Bess.	257	<i>Bodeana</i> , Bge.	644
<i>nana</i> , Sm. Prod.	258	<i>bracteata</i> , Bth.	667
<i>nana</i> Vill.	242	<i>brevifolia</i> , C. A. M.	647
<i>obtusa</i> , W. K.	255	<i>Cadmea</i> , Boiss.	652
<i>Œnesia</i> , Heldr.	257	<i>Caesarea</i> , Boiss.	653
<i>Olympica</i> , Boiss.	258	<i>calaminthoides</i> , Bth.	644
<i>Orientalis</i> , Schenck	157	<i>Calverti</i> , Boiss.	646
<i>palustris</i> , Willd.	234	<i>calycina</i> , Fenzl	669
<i>phœnicea</i> , Ehr.	157	<i>camphorata</i> , B. et Heldr.	765
<i>platyphylla</i> , Boiss.	257	<i>Cataria</i> , L.	645
<i>Pontica</i> , C. Koch	257	<i>cephalotes</i> , Boiss.	641
<i>propinquia</i> , F. et M.	241	<i>chionophila</i> , B. et	
<i>pnsilla</i> , Lois.	256	<i>Haussk</i>	645
<i>pygmaea</i> , Bert.	259	<i>Cilicica</i> , Boiss.	661
<i>Pyrenaica</i> , Pourr.	259	<i>circinnata</i> , W.	751
<i>refracta</i> , Boiss.	240	<i>clinopodioides</i> , Bth.	644
<i>saxatilis</i> , Pall.	250	<i>colorata</i> , W.	659
<i>scorpioides</i> , L.	255 et	<i>concolor</i> , B. et Heldr.	658
<i>Sicula</i> , Guss.	255	<i>congesta</i> , F. M.	669
<i>sparsiflora</i> , Mik	241	<i>cordifolia</i> , C. Koch	658
<i> speluncicola</i> , Schott.	240	<i>crassifolia</i> , B. et Buhse	660
<i>squarrosa</i> , Retz	249	<i>erinita</i> , M. et Auch.	645
<i>stricta</i> , Link	259	<i>erispa</i> , W.	642
<i>suaveolens</i> , W. K.	258	<i>cryptantha</i> , B. et Haussk.	669
<i>sylvatica</i> , Hoffm.	257	<i>curvidens</i> , B. et Bal	669
<i>tenniflora</i> , Viv.	217	<i>curviflora</i> , Boiss.	662
<i>uliginosa</i> , Schrad.	25	<i>cyanæa</i> , Stev.	656
<i>uncata</i> , B. et Bal.	245	<i>Daëneusis</i> , Boiss.	667
MYRSINEACEÆ	51	<i>denudata</i> , Bth.	650
Myrsine , L.	51	<i>depanperata</i> , Bth.	650
<i>Africana</i> , L.	52	<i>Dirphyta</i> , Heldr.	652
<i>bifaria</i> , Wall.	52	<i>erodiifolia</i> , Boiss.	674
Nanophytum , Less.	972	<i>fissa</i> , C. A. M.	646
<i>Cuspicum</i> , Less.	972	<i>floccosa</i> , Bth.	641
<i>erinaceum</i> , Pall.	972	<i>glabra</i> , Pth.	646
<i>juniperinum</i> , Moq.	972	<i>glabrescens</i> , Boiss.	658
<i>macranthum</i> , F. et M.	972	<i>globithora</i> , Bge.	667
Nepeta , L.	657	<i>glomerata</i> , M. et Auch.	662
<i>acinifolia</i> , Spr.	656	<i>glomerulosa</i> , Boiss.	654
<i>alba</i> , Desf.	665	<i>gracilescens</i> , Boiss.	647
		<i>graciliflora</i> , B. et Haussk.	648

	Pag.		Pag.
<i>Nepeta grandiflora</i> , M. B.	659	<i>Nepeta petrea</i> , Bth.	665
<i>heliotropifolia</i> , Lam.	668	<i>Pisidica</i> , B. et Heldr.	662
<i>humilis</i> , Bth.	664	<i>podostachys</i> , Bth.	659
<i>hymenodonta</i> , Boiss.	665	<i>Pontica</i> , C. Koch	665
<i>junccea</i> , Bth.	651	<i>prostrata</i> , Bth.	641
<i>incana</i> , M. B.	656	<i>pubescens</i> , Bth.	640
<i>Isaurica</i> , B. et Heldr.	653	<i>pungens</i> , Bth.	666
<i>Ispananica</i> , Boiss.	666	<i>pusilla</i> , Bth.	666
<i>Italica</i> , L.	654	<i>pycnantha</i> , Bth.	662
<i>Italica</i> , Fl. Gr.	652	<i>racemosa</i> , Lam.	660
<i>Kotschy</i> , Boiss.	657	<i>raphanorrhiza</i> , Bth.	640
<i>lagopsis</i> , Bth.	640	<i>rariifolia</i> , C. Koch	646
<i>lamiaefolia</i> , Bth.	648	<i>Reichenbachiana</i> , F. M.	660
<i>lamiaefolia</i> , W.	645	<i>ruderalis</i> , Ham.	644
<i>lasiocephala</i> , Bth.	642	<i>rugosa</i> , Bth.	645
<i>taxa</i> , Bth.	646	<i>Ruprechtii</i> , Boiss.	661
<i>taxiflora</i> , Bth.	649	<i>saccharata</i> , Bge.	664
<i>leptantha</i> , B. et Haussk.	648	<i>satureoides</i> , Boiss.	667
<i>leptoelada</i> , Trautv.	648	<i>Schiraziana</i> , Boiss.	666
<i>leneostegia</i> , B. et Heldr.	652	<i>Scordotis</i> , L.	651
<i>longiflora</i> , C. A. M.	660	<i>Scorodonia</i> , Boiss.	653
<i>longituba</i> , Vent.	645	<i>septemcrenata</i> , Ehr.	656
<i>macrociphon</i> , Boiss.	647	<i>septemcrenata</i> v. <i>Persica</i>	657
<i>Maraeandia</i> , Bge.	640	<i>septemcrenata</i> , var.	650
<i>marifolia</i> , B. et Huet.	660	<i>serpyllifolia</i> , M. B.	574
<i>marrubioidea</i> , B. et Heldr.	665	<i>sessilifolia</i> , Bge.	649
<i>marrubioidea</i> , W.	654	<i>Sibirica</i> , M. B.	668
<i>melissæfolia</i> , Pers.	659	<i>Sibthorpii</i> , Bth.	652
<i>melissæfolia</i> , Lam.	659	<i>spathulifera</i> , Bth.	668
<i>menthoidea</i> , B. et Buhse	645	<i>speciosa</i> , B. et Noë	659
<i>Meyeri</i> , Bth.	664	<i>Spruneria</i> , Boiss.	654
<i>Meyeri</i> var.	665	<i>stenantha</i> , Ky. et B.	661
<i>mierantha</i> , Bge.	664	<i>subincisa</i> , Bth.	659
<i>Mussini</i> , Hæk.	660	<i>sulphurea</i> , C. Koch	665
<i>nuda</i> , L.	665	<i>supina</i> , Stev.	642
<i>obtusicrena</i> , Boiss.	647	<i>tenuicaulis</i> , Boiss.	661
<i>Orientalis</i> , Mill.	654	<i>tenerifolia</i> , W.	646
<i>Orphanidea</i> , Boiss.	654	<i>Tmolea</i> , Boiss.	655
<i>oxyodonta</i> , Boiss.	649	<i>Trautvetteri</i> , B. et Buhse	647
<i>pallida</i> , C. Koch	664	<i>trichocalyx</i> , B. et Haussk.	656
<i>paniculata</i> , Cr.	665	<i>trichodonta</i> , B. et Noë	669
<i>Pannonica</i> , Jacq.	665	<i>Tymphrestca</i> , H. et Sart.	654
<i>Parnassica</i> , H. et Sart.	654	<i>Ucranica</i> , L.	668
<i>parriflora</i> , M. B.	668	<i>Ucranica</i> , M. B.	665
<i>Persica</i> , Boiss.	657	<i>Vaillantii</i> , Bth.	670

	PAG.		PAG.
Nepeta violacea , Vill.	665	Nonnea pulmonarioides , B.	
<i>virgata</i> , C. Koch	646	<i>et Bal.</i>	168
<i>viscida</i> , Boiss.	655	<i>rosea</i> , M. R.	164
Nerium , L.	47	<i>setosa</i> , R. et Sch.	165
<i>Kotschy</i> , Boiss.	48	<i>Sibthorpiana</i> , Don.	170
<i>Mascatense</i> , A.D.C.	48	<i>sordida</i> , F. et M.	166
<i>odoratum</i> , Sol.	47	<i>stenosolen</i> , B. et Bal.	169
<i>Oleander</i> , L.	47	<i>Szovitsiana</i> , Stev.	166
Noëa , Moq.	964	<i>Tourica</i> , Led.	167
<i>Aucherii</i> , Moq.	965	<i>ventricosa</i> , Fl. Gr.	169
<i>Daghestanica</i> , Turez.	966	<i>versicolor</i> , Sweet.	164
<i>Griffithii</i> , Bge.	966	<i>Vivianii</i> , A.D.C.	165
<i>major</i> , Bge.	966	Nordmannia cordifolia , Led.	177
<i>minuta</i> , B. et Bal.	966	NYCTAGINACEÆ.	1045
<i>spinossissima</i> , L.	965		
<i>tomentosa</i> , Moq. ex parte	964	Obione pedunculata , Moq.	912
<i>tomentosa</i> , ex parte Moq.	965	<i>portulacoides</i> , Moq.	915
<i>Tournefortii</i> , Sp.	965	<i>Turcomanica</i> , Bge.	911
Nonnea , Med.	162	<i>verrucifera</i> , Moq.	915
<i>alba</i> , D.C.	170	Ocymum , L.	559
<i>alpestris</i> , Stev.	164	<i>Basileicum</i> , L.	559
<i>amplexicaulis</i> , Hochst.	170	<i>sanctum</i> , L.	559
<i>anchusoides</i> , B. et Buhse	168	Odontites , Hall.	474
<i>Armena</i> , B. et Hnet.	167	<i>Aucherii</i> , Boiss.	475
<i>Armena</i> , Stev.	165	<i>Bocconi</i> , Ky.	477
<i>atra</i> , Griseb.	167	<i>Caucasicæ</i> , C. Koch	472
<i>auriculata</i> , Boiss.	170	<i>Cratæca</i> , Boiss.	477
<i>Caspica</i> , Don.	166	<i>Cypris</i> , Boiss.	477
<i>deinrenns</i> , C. A. M.	168	<i>diregens</i> , Jord.	476
<i>diffusa</i> , B. et Buhse	166	<i>glutinosa</i> , M. B.	475
<i>Euphratica</i> , Boiss.	169	<i>ixodes</i> , B. et Heldr.	476
<i>flavescens</i> , F. et M.	165	<i>Linkii</i> , H. et Sart.	475
<i>intermedia</i> , Led.	167	<i>lutea</i> , L.	475
<i>lamprocarpa</i> , Gr.	164	<i>rubra</i> , Pers.	476
<i>lutea</i> , Lam.	165	<i>serotina</i> , Lam.	476
<i>lutea</i> , Fl. Pelop.	250	OLEACEÆ.	55
<i>macrosperrma</i> , Griseb.	170	Olea , L.	56
<i>macrosperrma</i> , B. et Heldr.	168	<i>cuspidata</i> , Wall.	56
<i>melanocarpa</i> , Boiss.	165	<i>Europæa</i> , L.	56
<i>obtusifolia</i> , W.	164	<i>Oleaster</i> , Hoff. et Link	56
<i>Persica</i> , Boiss.	167	Omphalodes , Tourn.	266
<i>Philistea</i> , Boiss.	166	<i>Cappadocica</i> , W.	266
<i>picta</i> , M. B.	166	<i>Cariensis</i> , Boiss.	257
<i>pulla</i> , L.	166	<i>cornifolia</i> , Lehm.	267
<i>pulla</i> var. <i>mollis</i> , A.D.C.	232	<i>glochidiata</i> , Bge.	255

	Pag.		Pag.
<i>Omphalodes hirsuta</i> , D.C.	257	<i>Onosma echioides</i> v. <i>parvi-</i>	
<i>intermedia</i> , D.C.	255	<i>florum</i> , Led.	182
<i>Luciliae</i> , Boiss.	267	<i>elegans</i> , C. Koch	186
<i>micrantha</i> , D.C.	255	<i>erectum</i> , Fl. Gr.	202
<i>myosotoides</i> , Fres.	256	<i>flavidum</i> , Boiss.	186
<i>myosotoides</i> , Schr.	257	<i>flavum</i> , Lehm.	186
<i>Persica</i> , Boiss.	256	<i>frutescens</i> , Lam.	190
<i>physodes</i> , Bge.	255	<i>fruticosum</i> , Lab.	185
<i>Pontica</i> , C. Koch	259	<i>gigantem</i> , Lam.	202
<i>rupestris</i> , Rupr.	267	<i>Gmelini</i> , Led.	181
<i>stricta</i> , C. Koch	256	<i>Gmelini</i> var. <i>microcar-</i>	
<i>Wittmanniana</i> , Stev.	267	<i>pum</i> , Led.	192
<i>Onosma</i> , L.	178	<i>gracile</i> , Trautv.	193
<i>albo-roseum</i> , F. et M.	198	<i>halophilum</i> , B. et Heldr.	188
<i>Aleppicum</i> , Boiss.	184	<i>hebebulbum</i> , D.C.	195
<i>angustifolium</i> , Lehm.	201	<i>Helveticum</i> , Boiss.	201
<i>arenarium</i> , W. K.	181	<i>heterophyllum</i> , Gr.	201
<i>Armenum</i> , D.C.	200	<i>incanescens</i> , D.C.	185
<i>asperifolium</i> , Lehm.	214	<i>Isanrium</i> , B. et Heldr.	196
<i>astroites</i> , Stev.	203	<i>Kilonyense</i> , B. et Haussk.	194
<i>Aucherianum</i> , D.C.	201	<i>Kotschy</i> , Boiss.	192
<i>aurantiacum</i> , Janka	201	<i>Laconicum</i> , Boiss.	201
<i>auriculatum</i> , Auch.	203	<i>lanceolatum</i> , B. et Haussk.	194
<i>Bodeanum</i> , Boiss.	187	<i>latifolium</i> , B. et Haussk.	198
<i>Bourgæi</i> , Boiss.	196	<i>liparioides</i> , D.C.	187
<i>brachysolen</i> , Boiss.	186	<i>microcarpum</i> , D.C.	192
<i>brevifolium</i> , D.C.	202	<i>microspermum</i> , Stev.	191
<i>bulbotrichum</i> , D.C.	184	<i>mite</i> , B. et Heldr.	200
<i>cærulescens</i> , Boiss.	198	<i>molle</i> , D.C.	197
<i>calycinum</i> , Stev.	181	<i>montanum</i> , Sm.	202
<i>Caspium</i> , Grun.	181	<i>mutable</i> , B. et Haussk.	199
<i>Cassinum</i> , Boiss.	192	<i>nanum</i> , D.C.	189
<i>cespitosum</i> , Ky.	190	<i>Olivieri</i> , Boiss.	195
<i>chlorotrichum</i> , B. et Neë	194	<i>oreodoxum</i> , Boiss.	195
<i>Chorassanicum</i> , Bge.	184	<i>Orientalis</i> , L. 203 et Add.	
<i>ciliatum</i> , D.C.	197	<i>Orientalis</i> , Lehm.	190
<i>concinnum</i> , D.C.	192	<i>Orpharideum</i> , Boiss.	201
<i>congestum</i> , D.C.	198	<i>ovalifolium</i> , Ky. et Boiss.	196
<i>dasytrichum</i> , Boiss.	197	<i>pachypodium</i> , Boiss.	187
<i>decipiens</i> , Sch. et Ky.	189	<i>pallidum</i> , Boiss.	201
<i>dichroanthum</i> , Boiss.	181	<i>paradoxum</i> , Janka	201
<i>diffusum</i> , Boiss.	191	<i>polyanthum</i> , D.C.	180
<i>echinatum</i> , D.C.	184	<i>polyphyllum</i> , Led.	188
<i>echioides</i> , Fl. Gr.	190	<i>procernum</i> , Boiss.	195
<i>echioides</i> , L.	181	<i>Rascheyanum</i> , Boiss.	199

	Pag.		Pag.
Onosma rigidum , Led.	201	Origanum Heracleoticum ,	
<i>rostellatum</i> , Lehm.	185	Rehb.	551
<i>Rousselii</i> , D.C.	200	<i>Heracleoticum</i> , Bth.	552
<i>rupestris</i> , M. B.	191	<i>levigatum</i> , Boiss.	550
<i>seaberrimum</i> , B. et Heldr.	201	<i>leptocephalum</i> , Boiss.	549
<i>serpicioides</i> , Huet.	201	<i>Libanotium</i> , Boiss.	548
<i>sericeum</i> , W.	185	<i>macrostachyum</i> , Link	551
<i>setigerum</i> , Spr.	185	<i>Marr.</i> , Fl. Gr.	553
<i>setosum</i> , Led.	181	<i>Marr.</i> , L.	553
<i>simplicissimum</i> , L.	188	<i>mieranthum</i> , Vogel	552
<i>Spruneri</i> , Boiss.	202	<i>microphyllum</i> , Sieb.	552
<i>stamineum</i> , Led.	184	<i>neglectum</i> , Vog.	552
<i>stellulatum</i> , W. K.	201	<i>nervosum</i> , Vog.	553
<i>strigosissimum</i> , Boiss	191	<i>normale</i> , Don.	551
<i>strigosum</i> , St.	181	<i>Onites</i> , L.	553
<i>Syriacum</i> , Lab.	178	<i>porviflorum</i> , Urv.	551
<i>Tauricum</i> , Pall.	201	<i>pruinosum</i> , C. Koch	551
<i>tenuiflorum</i> , W.	191	<i>pulegham</i> , Boiss.	547
<i>tinctorium</i> , Fl. Pel.	182	<i>pulegium</i> , B. et Heldr.	549
<i>Tournefortii</i> , Gr.	190	<i>rotundifolium</i> , Boiss	550
<i>trachyanthum</i> , Boiss.	192	<i>seabrum</i> , B. et Heldr.	549
<i>trachytrichum</i> , Boiss.	186	<i>Sipyleum</i> , L.	547
<i>Trapezuntinum</i> , Huet	201	<i>Smyrnæum</i> , Fl. Gr.	552
<i>tripartitum</i> , Hochst.	186	<i>Syriacum</i> , Sieb.	553
<i>Troodi</i> , Ky.	189	<i>Tournefortii</i> , Sibth.	547
<i>velutinum</i> , Boiss.	189	<i>vestitum</i> , Cl.	553
<i>versicolor</i> , Schott.	200	<i>virens</i> , C. A. Mey.	551
<i>Visanii</i> , Clem.	181	<i>vulgare</i> , L.	551
<i>xanthotrichum</i> , Boiss.	184		492
Origanum , L.	546	OROBANCHACEÆ.	502
<i>Ægyptiacum</i> , L.	553	Orobanche , L.	502
<i>albiflorum</i> , C. Koch	551	<i>Ægyptiaca</i> , Pers.	499
<i>angustifolium</i> , C. Koch	551	<i>alba</i> , Steph.	507
<i>calcaratum</i> , Juss.	547	<i>alpigena</i> , C. Koch	506
<i>ciliatum</i> , B. et Ky.	548	<i>amethystea</i> , Th.	512
<i>cordifolium</i> , Montb. et		<i>ammophila</i> , C. A. M.	515
Auch.	548	<i>Anatolica</i> , B. et R.	504
<i>Creticum</i> , L.	551	<i>angustisepala</i> , Sch.	517
<i>Dictamnus</i> , L.	547	<i>arenaria</i> , Borek.	493
<i>dubium</i> , Boiss.	555	<i>arenaria</i> , M. B.	515
<i>Ehrenbergii</i> , Boiss.	551	<i>arcuata</i> , Sch.	517
<i>gracile</i> , C. Koch	551	<i>Artemisia</i> , Vauch.	512
<i>Hausknechtii</i> , Boiss.	550	<i>atrorubens</i> , Schultz	508
<i>hirtum</i> , Auct.	551	<i>Attica</i> , Reut.	512
<i>hirtum</i> , Vogel	552	<i>borealis</i> , Turez.	496
		<i>brevisflora</i> , Sch.	517

	Pag.		Pag.
Orobanche <i>Buhsei</i> , Reut.	509	Orobanche <i>Indica</i> , Roxb.	499
<i>cæsia</i> , Rehb.	496	<i>Kotschyi</i> , Reut.	514
<i>cærultra</i> , Vill.	495	<i>Kurdica</i> , B. et Haussk.	505
<i>cærulescens</i> , Steph.	515	<i>latisquama</i> , Reut.	516
<i>cærulescens</i> , C. Koch	504	<i>lavandulacea</i> , Reh.	497
<i>camptolepis</i> , B. et R.	515	<i>Lazica</i> , Reut.	508
<i>caryophyllacea</i> , C. A. M.	512	<i>leucantha</i> , Griseb	507
<i>caryophyllacea</i> , Sm.	509	<i>leucopogon</i> , B. et Haussk.	504
<i>cernua</i> , Löfl.	514	<i>lilacina</i> , Sch.	517
<i>coccinea</i> , M. B.	494	<i>longiflora</i> , Pers.	495
<i>colorata</i> , C. Koch	504	<i>loricata</i> , Rehb.	512
<i>connata</i> , C. Koch	504	<i>Lycica</i> , Schultz	516
<i>condensata</i> , Gr.	512	<i>marginata</i> , Reut.	508
<i>crassistyla</i> , Schultz	506	<i>medicaginis</i> , Rehb.	515
<i>cruenta</i> , Bert.	505	<i>minor</i> , Sull.	512
<i>cruenta</i> var. <i>leiostemon</i> , Gr.	509	<i>Mutellä</i> , Sch.	499
<i>Cumana</i> , Wallr.	515	<i>nana</i> , Noë	499
<i>Cumana</i> , Mut.	515	<i>Palestina</i> , Reut.	506
<i>euprea</i> , B. et Bal.	510	<i>parviloba</i> , Sch.	516
<i>curviflora</i> , Viv.	515	<i>pedunculata</i> , Viv.	499
<i>Cypria</i> , Reut.	515	<i>polygonthera</i> , Reut.	515
<i>Delilei</i> , Decaisne	499	<i>pruinosa</i> , Lap.	505
<i>diaphana</i> , Sch.	517	<i>psilandra</i> , C. Koch	509
<i>elatior</i> , Rehb.	511	<i>pubescens</i> , Urv.	507
<i>Epithymum</i> , D.C.	509	<i>punctata</i> , Schultz	509
<i>Eryngii</i> , Dub.	512	<i>quadrisida</i> , C. Koch	505
<i>euglossa</i> , Rehb.	510	<i>ramosa</i> , L.	498
<i>ferruginea</i> , C. Koch	509	<i>rubens</i> , Wallr.	511
<i>fragrans</i> , Gr.	508	<i>segetum</i> , Sprun.	507
<i>fuliginosa</i> , Reut.	510	<i>speciosa</i> , D.C.	506
<i>Galii</i> , Dub.	508	<i>spectabilis</i> , Reut.	514
<i>Gallica</i> , Gr.	515	<i>Spruneri</i> , Schultz	506
<i>gamosepala</i> , Reut.	503	<i>squalida</i> , St.	517
<i>glabrata</i> , C.A. M.	511	<i>Stocksii</i> , Boiss.	505
<i>gracilis</i> , Sm.	505	<i>striata</i> , Reut.	512
<i>grandiflora</i> , Bor. et Ch.	507	<i>strobiligena</i> , Rehb.	509
<i>grandisepala</i> , Sch.	516	<i>subverticillata</i> , Sch.	516
<i>Grenieri</i> , Schultz	515	<i>Teuerii</i> , Schultz	508
<i>Grisebachii</i> , Reut.	512	<i>versicolor</i> , Schultz	508
<i>Hederæ</i> , Dub.	515	<i>vulgaris</i> , D.C.	509
<i>hians</i> , Reut.	508	<i>Wiedemannii</i> , Boiss.	510
<i>hians</i> , Stev.	517	<i>Ornus Europea</i> , Pers.	39
<i>Hispanica</i> , Boiss.	515	<i>Orthosiphon</i> , Bth.	539
<i>hyalina</i> , Sprun.	508	<i>pallidus</i> , Bth.	539
		<i>Ostrya</i> , Mich.	1178

INDEX.

4245

Pag.

Pag.

<i>Ostrya carpinifolia</i> , Scop.	1178	<i>Paracaryum incanum</i> , Ledeb.	262
<i>vulgaris</i> , W.	1178	<i>lamprocarpum</i> , Boiss.	258
Osyris , L.	1058	<i>leptophyllum</i> , A. D. C.	258
<i>alba</i> , L.	1058	<i>longipes</i> , Boiss.	244
Otostegia , Bth.	776	<i>microanthum</i> , Boiss.	255
<i>Aucherii</i> , Boiss.	778	<i>microcarpum</i> , Boiss.	255
<i>Kotschyii</i> , Boiss.	777	<i>modestum</i> , B. et Haussk.	260
<i>microphylla</i> , Boiss.	777	<i>myosotoides</i> , Lab.	257
<i>mollucoides</i> , Vahl	777	<i>Persicum</i> , Boiss.	256
<i>Persica</i> , Burm.	777	<i>Ponticum</i> , C. Koch	259
<i>scariosa</i> , Bth.	777	<i>Renterii</i> , B. et Haussk.	257
<i>Schimpieri</i> , Bth.	776	<i>rufiflorum</i> , Stocks	256
<i>Oxynepetia iurolucrata</i> , Bge.	670	<i>rugulosum</i> , D. G.	256
Oxyria , Hill.	1004	<i>salsum</i> , Boiss.	256
<i>digyna</i> , L.	1004	<i>stenolophum</i> , Boiss.	256
<i>reniformis</i> , Hook.	1004	<i>strictum</i> , C. Koch	256
<i>Oxyris ceratoides</i> , L.	917	<i>undulatum</i> , Boiss.	256
Oxystelma , R. Br.	59		
<i>Alpini</i> , Decaisne	59	Parietaria , L.	1149
Pæderota , L.	454	<i>alsinefolia</i> , Del.	1151
<i>Pontica</i> , Rupr.	454	<i>Cretica</i> , L.	1150
<i>Palladia atropurpurea</i> , J. et Sp.	8	<i>cryptarum</i> , C. Koch	1149
<i>Pallasia Caspica</i> , L.	998	<i>debelis</i> , Forst.	1151
Panderia , F. et M.	919	<i>diffusa</i> , M. K.	1149
<i>divaricata</i> , C. Koch	919	<i>elliptica</i> , C. Koch	1149
<i>pilosa</i> , F. et M.	919	<i>erecta</i> , M. K.	1149
Paracaryum , Boiss.	255	<i>Judaica</i> , L.	1149
<i>Aneyritanum</i> , Boiss.	266	<i>Lusitanica</i> , L.	1150
<i>angustifolium</i> , W.	265	<i>micrantha</i> , Led.	1151
<i>asperum</i> , Stocks	261	<i>multicaulis</i> , B. et	
<i>Aucherii</i> , A. D. C.	260	Heldr.	1149
<i>azureum</i> , B. et Heldr.	265	<i>nitens</i> , C. Koch.	1149
<i>calathicarpum</i> , Stocks	245	<i>officinalis</i> , L.	1149
<i>e lyceinum</i> , B. et Bal.	261	<i>parviflora</i> , Fried.	1150
<i>Cappadocicum</i> , B. et Bal.	259	<i>punctata</i> , W.	1149
<i>Cariense</i> , Boiss.	257		
<i>corymbiforme</i> , D. C.	262	<i>Passerina Ammodendrum</i> ,	
<i>corymbiforme</i> , Tchih.	265	K. et Kir.	1051
<i>cristatum</i> , Lam.	258	<i>annua</i> , Wickst.	1152
<i>denticulatum</i> , Boiss.	268	<i>arvensis</i> , Lam.	1052
<i>erigerifolium</i> , Schott.	265	<i>hirsuta</i> , L.	1054
<i>erysimifolium</i> , Boiss.	259	<i>Lessertia</i> , W.	1051
<i>glaucifolium</i> , W.	262	<i>Orientalis</i> , W.	1054
<i>hirsutum</i> , Boiss.	257	<i>Persica</i> , Boiss.	1051
		<i>pubescens</i> , Guss.	1052
		<i>racemosa</i> , Wickst.	1051
		<i>resiculosa</i> , F. M.	1055

	Pag.		Pag.
Pedicularis , L.	483	<i>Pergularia tomentosa</i> , L.	60
<i>achilleæfolia</i> , Hoh.	492	Periploca , L.	49
<i>acmodonta</i> , Boiss.	492	<i>angustifolia</i> , Lab.	50
<i>Araratica</i> , Bge.	488	<i>aphylla</i> , Decaisn.	50
<i>Armena</i> , B. et Huet	484	<i>gracilis</i> , Boiss.	50
<i>atropurpurea</i> , Nordm.	487	<i>Græca</i> , L.	49
<i>bicuspidata</i> , Gr.	492	<i>lævigata</i> , Ait.	50
<i>brachyodonta</i> , Schle.	491	<i>punicæfolia</i> , Cav.	50
<i>Cadmea</i> , Boiss.	485	<i>rigida</i> , Viv.	50
<i>campylotosiphon</i> , C. Koch	488	<i>Secamone</i> , Del.	59
<i>Caucasica</i> , M. B.	483	Perowskia , Kar.	588
<i>comosa</i> , L.	491	<i>abrotanoides</i> , Kar.	589
<i>condensata</i> , M. B.	487	<i>artemisioides</i> , Boiss.	589
<i>crassirostris</i> , Bge.	488	<i>atriplacifolia</i> , Bth.	589
<i>Elephas</i> , Boiss.	489	<i>serophulariæfolia</i> , Bth.	589
<i>flava</i> , Sm.	492	Petrosimonia , Bge.	972
<i>Friderici Augusti</i> , Tomm.	491	<i>brachiata</i> , Pall.	974
<i>Græca</i> , Bge.	490	<i>crassifolia</i> , Pall.	975
<i>Hacquetii</i> , C. Koch	488	<i>glauea</i> , Pall.	975
<i>jucunda</i> , Schott	492	<i>Volvox</i> , Pall.	972
<i>labellata</i> , Jacq.	489	<i>Pharnaceum occultum</i> , Forsk.	896
<i>leucodon</i> , Rehb. fil.	491	Phelipæa , Tourn.	494
<i>leucodon</i> , Gr.	490	<i>Egyptiaca</i> , Pers.	499
<i>Nordmanniana</i> , Bge.	489	<i>ambigua</i> , Bge.	501
<i>nudicaulis</i> , C. Koch	484	<i>arenaria</i> , Borck.	495
<i>occulta</i> , Janka	490	<i>Armena</i> , C. Koch	501
<i>ochroleuca</i> , Schle.	491	<i>cærulea</i> , Vill.	495
<i>Olympica</i> , Boiss.	486	<i>cæsia</i> , Gr.	499
<i>orthantha</i> , Gr.	486	<i>Cedreti</i> , Reut.	495
<i>Pontica</i> , Boiss.	485	<i>coelestis</i> , Reut.	496
<i>pyrenantha</i> , Boiss.	484	<i>Delilei</i> , Walp.	499
<i>rostrata</i> , C. A. Mey.	489	<i>emarginata</i> , Reut.	499
<i>rupestris</i> , B. et Orph.	490	<i>fissa</i> , C. A. Mey.	501
<i>Sibthorpii</i> , Boiss.	492	<i>flava</i> , C. A. Mey.	500
<i>subrostrata</i> , C. A. M.	488	<i>foliata</i> , Lamb.	494
<i>verticillata</i> , L.	486	<i>Fraasii</i> , Walp.	498
<i>verticillata</i> , Tchih.	486	<i>gigantea</i> , Kar.	504
<i>villobractea</i> , C. Koch	487	<i>Heldreichii</i> , Reut.	498
<i>Wilhelmsiana</i> , Fisch.	487	<i>hirtiflora</i> , Reut.	497
Pentodon <i>barbatus</i> , Ehr.	925	<i>Hohenackeri</i> , Reut.	499
Pentatropis , R. Br.	58	<i>incana</i> , Payne	504
<i>microphylla</i> , Roxb.	Add.	<i>Indica</i> , Reut.	499
<i>spiralis</i> , Forsk.	58	<i>lanuginosa</i> , C. A. M.	496
Peplidium , Del.	427	<i>lavandulacea</i> , Rehb.	497
<i>humilisum</i> , Del.	427	<i>longiflora</i> , Pers.	495

	Pag.		Pag.
<i>Phelipaea lutea</i> , Desf.	500	<i>Phlomis elliptica</i> , Bth	783
<i>macrantha</i> , C. Koch	499	<i>ferruginea</i> , Ten.	787
<i>Muteli</i> , Reut.	495	<i>flavescens</i> , Mill.	786
<i>obtusiloba</i> , Reut.	496	<i>floccosa</i> , Don.	786
<i>oxyloba</i> , Reut.	497	<i>fruticosa</i> , L.	784
<i>puleghella</i> , G. A. Mey.	496	<i>fruticosa</i> , Sieb.	786
<i>pyramidalis</i> , Reut.	498	<i>Ghilianica</i> , C. Koch	784
<i>ramosa</i> , L.	498	<i>glandulosa</i> , Schenk	788
<i>rufescens</i> , Gr.	499	<i>Haussknechtii</i> , Bge.	785
<i>salsa</i> , C. A. Mey	501	<i>Herbaventi</i> , L.	791
<i>Schullzii</i> , Mut.	490	<i>Herbarenti</i> , var. Russ.	788
<i>serotina</i> , Haussk.	499	<i>Iberica</i> , Vis.	795
<i>Syssirensis</i> , C. Koch	495	<i>imbricata</i> , Boiss.	785
<i>Tournefortii</i> , Desf.	494	<i>laciniflora</i> , L.	793
<i>tricholoba</i> , Reut.	498	<i>lanata</i> , W.	785
<i>tubulosa</i> , Schenk	500	<i>lanceolata</i> , B. et Hoh.	785
<i>Phleboanthe Larmanni</i> , Tausch		<i>linearis</i> , B. et Bal.	785
<i>Phillyrea</i> , Tourn.	56	<i>longifolia</i> , B. et Bal.	789
<i>latifolia</i> , Fl. Gr.	57	<i>lunariæfolia</i> , Fl. Gr.	785
<i>media</i> , L.	57	<i>lunariæfolia</i> , Sm. Prodr.	790
<i>pedunculata</i> , Bory	57	<i>lunariæfolia</i> , var.	788
<i>Vilmoriniana</i> , Boiss. et		<i>Lycia</i> , Don.	786
<i>Bal.</i>	57	<i>Mesopotamica</i> , Boiss.	791
<i>Phlomis</i> , L.	779	<i>mierophylla</i> , Sieb.	586
<i>angustifolia</i> , B. et Bal.	785	<i>moluccoides</i> , Vahl	777
<i>angustifolia</i> , Mill.	786	<i>Nissolia</i> , L.	781
<i>anisodonta</i> , Boiss.	789	<i>Olivieri</i> , Bth.	781
<i>Armeniaca</i> , W.	785	<i>oppositiflora</i> , B. et	
<i>Armeniaca</i> , Bth.	781	<i>Haussk.</i>	784
<i>Armeniaca</i> var. Boiss.	782	<i>Orientalis</i> , Mill.	751
<i>Aucheria</i> , Boiss.	780	<i>parvifolia</i> , Presl	786
<i>aurea</i> , Decaisn.	786	<i>Persica</i> , Boiss.	792
<i>bicolor</i> , W.	792	<i>pungens</i> , W.	790
<i>bicolor</i> , Bth.	786	<i>pungens</i> , var. Bth.	792
<i>Bourgatii</i> , Boiss.	787	<i>rigida</i> , Lab.	690
<i>brevilabris</i> , Ehr.	782	<i>Russeliania</i> , Lag.	788
<i>Bruguieri</i> , Desf.	780	<i>Samia</i> , L.	790
<i>cancellata</i> , Bge.	789	<i>Samia</i> , var. Viv.	786
<i>capitata</i> , Boiss.	782	<i>superba</i> , C. Koch	790
<i>Chorassanica</i> , Bge.	789	<i>Syriaca</i> , Boiss.	781
<i>chrysophylla</i> , Boiss.	787	<i>Tarsensis</i> , Boiss.	781
<i>clandestina</i> , B. et Ch.	711	<i>thapsoides</i> , Bge.	791
<i>cordata</i> , Boiss.	782	<i>tuberosa</i> , L.	792
<i>Cretica</i> , Presl.	786	<i>urticafolia</i> , Vahl	778
<i>cymifera</i> , Boiss.	792	<i>virens</i> , D. C.	788

	Pag.		Pag.
<i>Phlomis viscosa</i> , <i>Poir.</i>	788	<i>Plantago arenaria</i> , <i>W. K.</i>	892
Phyllanthus , <i>Sw.</i>	1159	<i>argentea</i> , <i>Desf.</i>	885
<i>rotundifolius</i> , <i>W.</i>	1159	<i>argentea</i> , <i>Ten.</i>	882
Physalis , <i>L.</i>	286	<i>aspera</i> , <i>Gand.</i>	889
<i>Alkekengi</i> , <i>L.</i>	287	<i>Bauphula</i> , <i>Edg.</i>	885
<i>arborescens</i> , <i>Th.</i>	287	<i>Bellardi</i> , <i>All.</i>	884
<i>flexuosa</i> , <i>L.</i>	287	<i>bellidifolia</i> , <i>Viv.</i>	887
<i>somnifera</i> , <i>L.</i>	287	<i>Byzantina</i> , <i>C. Koch</i>	881
Physoclaina , <i>Don.</i>	295	<i>capitata</i> , <i>Ten.</i>	881
<i>Orientalis</i> , <i>M. B.</i>	295	<i>capitellata</i> , <i>Ram.</i>	890
<i>Physogeton acanthophyllum</i> ,		<i>Caspica</i> , <i>F. et M.</i>	885
<i>J. et Sp.</i>	985	<i>carinata</i> , <i>Schrad.</i>	889
PHYTOLACCACEÆ	894	<i>ciliata</i> , <i>Desf.</i>	887
Phytolacca , <i>L.</i>	894	<i>commutata</i> , <i>Guss.</i>	888
<i>decandra</i> , <i>L.</i>	895	<i>Coronopus</i> , <i>L.</i>	888
<i>pruinosa</i> , <i>Fenzl.</i>	895	<i>Coronopus var.</i> <i>Decaisn.</i>	888
Pilosyles , <i>Guill.</i>	1071	<i>Cretica</i> , <i>L.</i>	884
<i>Hausknechtii</i> , <i>Boiss.</i>	1072	<i>erypoides</i> , <i>Boiss.</i>	888
Pinguicula , <i>L.</i>	1	<i>cylindrica</i> , <i>Forsk.</i>	882
<i>Albanica</i> , <i>Griseb.</i>	2	<i>Cynops</i> , <i>Sm.</i>	892
<i>cristallina</i> , <i>Fl. Gr.</i>	2	<i>decumbens</i> , <i>Forsk.</i>	885
<i>hirtiflora</i> , <i>Ten.</i>	2	<i>eriantha</i> , <i>Decaisn.</i>	887
<i>laeta</i> , <i>Pant.</i>	2	<i>eriostachya</i> , <i>Ten.</i>	886
<i>leptoceras</i> , <i>Rehb.</i>	2	<i>Euphratica</i> , <i>Decaisn.</i>	887
<i>Megaspilæa</i> , <i>Boiss. et</i>		<i>evacina</i> , <i>Boiss.</i>	890
<i>Heldr.</i>	2	<i>exigua</i> , <i>Murr.</i>	891
<i>vulgaris</i> , <i>L.</i>	2	<i>fornicata</i> , <i>C. Koch</i>	886
<i>Piptochlamys hirsuta</i> , <i>C. A. M.</i>	1054	<i>geatianoides</i> , <i>Sm.</i>	879
Piptoptera , <i>Bge.</i>	978	<i>glauca</i> , <i>C. A. Mey</i>	886
<i>Turkestana</i> , <i>Bge.</i>	978	<i>Griffithii</i> , <i>Decaisn.</i>	880
Planera <i>Abelicea</i> , <i>R. et Sch.</i>	1159	<i>Hausknechtii</i> , <i>Vatk</i>	886
<i>crenata</i> , <i>Desf.</i>	1159	<i>holostea</i> , <i>Lam.</i>	884
<i>Richardi</i> , <i>Mich.</i>	1159	<i>Indica</i> , <i>L.</i>	891
PLANTAGINEÆ	876	<i>intermedia</i> , <i>Gilib.</i>	875
Plantago , <i>L.</i>	876	<i>intermedia</i> , <i>C. Koch</i>	881
<i>Aegyptiaca</i> , <i>Jacq.</i>	895	<i>interrupta</i> , <i>Poir.</i>	882
<i>Afra</i> , <i>L.</i>	891	<i>Karakabanica</i> , <i>B. et Huet.</i>	881
<i>albicans</i> , <i>L.</i>	882	<i>Kurdica</i> , <i>Vatk.</i>	881
<i>Alpina</i> , <i>Sieb.</i>	889	<i>lachnantha</i> , <i>Bge.</i>	887
<i>Alpina</i> , <i>Vill.</i>	880	<i>lagoocephala</i> , <i>Bge.</i>	886
<i>Alpina v. dioritica</i> , <i>Ky</i>	890	<i>lagopoides</i> , <i>Desf.</i>	883
<i>altissima</i> , <i>L.</i>	881	<i>Lagopus</i> , <i>L.</i>	886
<i>ambigua</i> , <i>Guss.</i>	881	<i>lanata</i> , <i>Port.</i>	881
<i>amplexicaulis</i> , <i>Cav.</i>	885	<i>lanceolata</i> , <i>L.</i>	881
<i>Arabica</i> , <i>Boiss.</i>	890	<i>lanuginosa</i> , <i>d. c.</i>	881

	Pag.		Pag.
<i>Plantago Loeflingii</i> , L.	885	PLATANACEÆ	1161
<i>Lusitanica</i> , W.	886	Platanus , L.	1161
<i>major</i> , L.	878	<i>Orientalis</i> , L.	1161
<i>maritima</i> , L.	889	Pleurogyne , Esch.	77
<i>media</i> , L.	879	<i>Carinthiaca</i> , Wulf.	77
<i>microcephala</i> , Poir.	885	PLUMBAGINÆ	892
<i>minima</i> , D. C.	878	Flumbago , L.	875
<i>minuta</i> , C. A. M.	885	<i>Europæa</i> , L.	875
<i>minuta</i> , Pall.	884	<i>lapathifolia</i> , W.	875
<i>montana</i> , Lam.	880	<i>Zeylanica</i> , L.	875
<i>nana</i> , Bert.	885	Podonosma , Boiss.	178
<i>notata</i> , Lag.	885	<i>Galalense</i> , Schw.	1199
<i>Olivieri</i> , Decaisn.	886	<i>Syriaeum</i> , Lab.	178
<i>ovata</i> , Focke.	885	POLEMONIACEÆ	85
<i>penicillata</i> , Endl.	887	Polemonium , L.	85
<i>phœopsis</i> , Payne	885	<i>cæruleum</i> , L.	85
<i>phœostoma</i> , Boiss.	892	Polycnemum , L.	995
<i>pilosa</i> , Ponrr.	884	<i>arvense</i> , L.	995
<i>polystachya</i> , Viv.	895	<i>arvense</i> , Fl. Gr.	996
<i>præcox</i> , C. A. Mey.	886	<i>brachiatum</i> , Pall.	974
<i>Psyllium</i> , L.	891	<i>crassifolium</i> , Pall.	975
<i>Psyllium</i> var. <i>Sinica</i> ,		<i>erinaceum</i> , Pall.	972
Boiss.	890	<i>glaucum</i> , Pall.	975
<i>pumila</i> , W.	891	<i>juniperinum</i> , M. B.	922
<i>remotiflora</i> , Stocks	885	<i>majns</i> , A. Br	966
<i>Rosetana</i> , Poir.	891	<i>malacophyllum</i> , M. B	981
<i>rugosa</i> , Hochst.	891	<i>oppositifolium</i> , Pall.	975
<i>salina</i> , Decaisn.	885	<i>verneosinum</i> , Lang	996
<i>salsa</i> , Pall.	889	<i>Volvox</i> , Pall.	972
<i>sarcophylla</i> , Boiss.	895	POLYGONEÆ	997
<i>saxatilis</i> , M. B.	880	Polygonum , L.	1025
<i>serpentina</i> , C. Koch	890	<i>acaulis</i> , Boiss.	1045
<i>Serraria</i> , L.	889	<i>Afghanicum</i> , Meisn.	1041
<i>selosa</i> , Bert.	895	<i>Aleppicum</i> , B. et Haussk.	1057
<i>Sinica</i> , Decaisn.	890	<i>alpestre</i> , C. A. Mey.	1057
<i>squarrosa</i> , Murr.	892	<i>Alpinum</i> , All.	1051
<i>Stocksii</i> , Boiss.	882	<i>ampliibium</i> , L.	1028
<i>striata</i> , Schousb.	891	<i>arachnoideum</i> , Kl.	1050
<i>subulata</i> , Wulf.	890	<i>arenarium</i> , Fl. Gall.	1056
<i>Syrtica</i> , Viv.	886	<i>ar-narium</i> , W. K.	1055
<i>tenuiflora</i> , W. K.	879	<i>argyroeolum</i> , St.	1055
<i>teretifolia</i> , Sieb.	889	<i>aridum</i> , B. et Haussk.	1042
<i>uliginosa</i> , Baumg.	879	<i>avienlare</i> , L.	1056
<i>villosa</i> , Mœnch	883	<i>Bellardi</i> , All.	1024
<i>Wulfeni</i> , M. K.	889	<i>Bellardi</i> var. Meisn.	1056

	Pag.		Pag.
<i>Polygonum biforme</i> , Wahl.	1050	<i>Polygonum Meyeri</i> , St.	1041
<i>Bistorta</i> , L.	1027	<i>minus</i> , Huds.	1029
<i>bistortoides</i> , Boiss.	1028	<i>mite</i> , Schr.	1029
<i>buxifolium</i> , M. B.	1025	<i>mollieforme</i> , Boiss.	1045
<i>Cappadocicum</i> , Boiss.	1054	<i>monticolum</i> , St.	1058
<i>carneum</i> , C. Koch	1057	<i>myrianthum</i> , Boiss.	1054
<i>Cedrorum</i> , Boiss.	1040	<i>Niloticum</i> , Meisn.	1050
<i>chlorocoleum</i> , St.	1054	<i>nodosum</i> , Pers.	1050
<i>cognatum</i> , Meisn.	1058	<i>Noeanum</i> , Boiss.	1055
<i>confusum</i> , Meisn.	1028	<i>obtusatum</i> , St.	1050
<i>controversum</i> , Guss.	1056	<i>oligophyllum</i> , Boiss.	1042
<i>Convolvulus</i> , L.	1052	<i>Olivieri</i> , J. et Sp.	1055
<i>corrigioloides</i> , J. et Sp.	1053	<i>pachyrhizum</i> , Tr.	1059
<i>crispulum</i> , Bot. Mag.	1025	<i>Paronychia</i> , C. A. M.	1041
<i>deciduum</i> , Boiss.	1053	<i>paronychioides</i> , C.A.M.	1040
<i>Donii</i> , Meisn.	1029	<i>patulum</i> , M. B.	1054
<i>dracunculifolium</i> , Boiss.	1059	<i>Persicaria</i> , L.	1050
<i>Dryandri</i> , Spr.	1058	<i>pluriflorum</i> , C. Koch	1058
<i>dumetorum</i> , L.	1052	<i>polycenoides</i> , J. et Sp.	1055
<i>dumosum</i> , Boiss.	1042	<i>polymorphum</i> , Ledeb.	1051
<i>effusum</i> , Meisn.	1058	<i>pseudohydropiper</i> ,	
<i>Ehrenbergii</i> , Meisn.	1056	Salzm.	1028
<i>elegans</i> , Roxb.	1058	<i>pusillum</i> , Lois.	1055
<i>elegans</i> , Ten.	1055	<i>pusillum</i> , Lam.	1029
<i>equisetiforme</i> , Fl. Gr.	1056	<i>radioesnum</i> , Boiss.	1059
<i>flaccidum</i> , Roxb.	1029	<i>restionoides</i> , B. et	
<i>Hermonis</i> , Ky.	1040	Hanssk.	1050
<i>herniarioides</i> , Del.	1048	<i>reticulatum</i> , C. Koch	1054
<i>Hydropiper</i> , L.	1029	<i>rottbollioides</i> , J. et Sp.	1055
<i>illecebroides</i> , Meisn.	1058	<i>Roxburghii</i> , Meisn.	1058
<i>incanum</i> , Schleb.	1030	<i>salicifolium</i> , Br.	1028
<i>Kotschyianum</i> , Boiss.	1040	<i>salicornioides</i> , J. et Sp.	1042
<i>lanatum</i> , Roxb.	1050	<i>salsuginineum</i> , M. B.	1055
<i>lanigerum</i> , R. Br.	1050	<i>Senegalense</i> , Meisn.	1051
<i>lapathifolium</i> , L.	1050	<i>serpyllaceum</i> , J. et Sp.	1059
<i>latifolium</i> , Boiss.	1052	<i>serrulatum</i> , Lag.	1028
<i>laxiflorum</i> , Weih.	1029	<i>serrulatum</i> , Illohen.	1029
<i>Lemannianum</i> , Meisn.	1401	<i>setosum</i> , Jacq.	1058
<i>Libani</i> , Boiss.	1040	<i>strictum</i> , All.	1029
<i>limbatum</i> , Meisn.	1051	<i>strictum</i> , Led.	1054
<i>littorale</i> , Link	1056	<i>tenuiflorum</i> , Presl.	1050
<i>Iuzuloides</i> , J. et Sp.	1059	<i>thymifolium</i> , J. et Sp.	1040
<i>macrochiton</i> , F. et M.	1051	<i>tomentosum</i> , Sch.	1050
<i>maerohymeum</i> , Boiss.	1041	<i>tubulosum</i> , Boiss.	1052
<i>maritimum</i> , L.	1057	<i>undulatum</i> , Murr.	1051

	Pag.		Pag.
Polygonum Venantianum,		Primula glacialis, Ad.	28
Clem.	1055	<i>grandiflora,</i> Lam.	24
viviparum, L.	1027	<i>grandis,</i> Traub.	27
Populus, Tourn.	1192	<i>longifolia,</i> Bot. Mag.	28
<i>alba,</i> L.	1195	<i>luteola,</i> Rupr.	29
<i>australis,</i> Ten.	1195	<i>macrocalyx,</i> Bge.	25
<i>Cilicica,</i> Ky.	1195	<i>macrophylla,</i> C. Koch	28
<i>dilatata,</i> Ait.	1194	<i>megascaevolia,</i> Boiss. et	
<i>diversifolia,</i> Schr.	1194	<i>Bal.</i>	26
<i>Euphratica,</i> Oliv.	1194	<i>Meyeri,</i> Rupr.	26
<i>fastigiata,</i> Poir.	1195	<i>minima,</i> L.	51
<i>Graeca,</i> Ait.	1195	<i>nivalis,</i> Pall.	27
<i>Italica,</i> Auct.	1194	<i>officinalis,</i> Jacq.	24
<i>nigra,</i> L.	1194	<i>Pallasii,</i> Lehmann	26
<i>nivea,</i> W.	1195	<i>pycnorrhiza,</i> Led.	28
<i>pyramidalis,</i> Roy.	1194	<i>Sibthorpii,</i> Rehb.	24
<i>tremula,</i> L.	1195	<i>savoleolens,</i> Bertol.	23
Pouzolzia pentandra, Brum.	1140	<i>sylvestris,</i> Scop.	24
Prasium, L.	798	<i>Tournefortii,</i> Rupr.	28
<i>hirsutum,</i> Poir.	755	<i>Uralensis,</i> Fisch.	25
<i>majus,</i> L.	798	<i>veris,</i> L.	24
PRIMULACEAE.	4	<i>verticillata,</i> Decaisne	25
Primula, L.	22	Friva, Ad.	555
<i>acaulis,</i> Jacq.	24	<i>Abyssinica,</i> J. et Sp.	555
<i>algida,</i> Ad.	29	<i>dentata,</i> Juss.	555
<i>algida,</i> Janka	50	<i>leptostachya,</i> Aitch.	555
<i>amæna,</i> M. B.	26	Psilosstemon D.C.	177
<i>Aueherii,</i> Jaub. et Sp.	25	<i>Cretievii,</i> W.	177
<i>auriculata,</i> Pall.	28	<i>Orientalis,</i> L.	177
<i>Bayernii,</i> Rupr.	28	Pterocarya, Kth.	1160
<i>Boveana,</i> Decaisne	25	<i>Caucasicæ,</i> C. A. M.	1161
<i>Bungeana,</i> C. A. M.	29	<i>fraxinifolia,</i> Lam.	1160
<i>capitellata,</i> Boiss.	50	Pterococcus uphyllus, Pall.	998
<i>Caucasicæ,</i> C. Koch	29	<i>Persicus,</i> B. et Buhse	999
<i>Columnæ,</i> Ten.	25	Pteropyrum, J. et Sp.	1001
<i>cordifolia,</i> Rupr.	26	<i>Aueheri,</i> J. et Sp.	1002
<i>crassifolia,</i> Lehmann	28	<i>ericoides,</i> Boiss.	1002
<i>Darialica,</i> Rupr.	50	<i>gracile,</i> Boiss.	1002
<i>elatior,</i> Jacq.	25	<i>Griffithii,</i> Meisn.	1002
<i>farinifolia,</i> Rupr.	29	<i>Noëanum,</i> Boiss.	1002
<i>farinosa,</i> M. B.	29	<i>Olivieri,</i> J. et Sp.	1002
<i>farinosa</i> var. <i>Turcica,</i>		<i>scoparium,</i> J. et Sp.	1003
Friv.	50	Pulmonaria, L.	170
<i>floribunda,</i> Wall.	24	<i>augustifolia,</i> Wahl.	171
<i>frondosa,</i> Janka	50	<i>azurea,</i> Bess.	171

	Pag.		Pag.
<i>Pulmonaria mollis</i> , Wulf.		<i>Quercus Hartwissiana</i> , St.	1164
<i>170</i> et Add.		<i>hypochrysa</i> , Stev.	1165
<i>officinalis</i> , L.	170	<i>hypoleuca</i> , Ky.	1167
<i>Puneeria coagulans</i> , Stocks	288	<i>Iberica</i> , Stev.	1164
<i>Pupalia</i> , Juss.	995	<i>Ibicis</i> , Ky.	1164
<i>lappacea</i> , L.	995	<i>Ilex</i> , L.	1167
Quercus , Tourn.	1162	<i>incrassata</i> , Ebremb.	1167
<i>Abelicea</i> , Lam.	1159	<i>insectoria</i> , Oliv.	1167
<i>Abietum</i> , Ky.	1164	<i>inops</i> , Ky.	1169
<i>Aegilops</i> , L.	1171	<i>intermedia</i> , Ky.	1165
<i>Aegilops</i> , Gr.	1172	<i>Ithaburensis</i> , Dec.	1172
<i>Aesculus</i> , Griseb.	1164	<i>lamprophyllos</i> , Koch	1165
<i>alnifolia</i> , Poeh.	1168	<i>lanuginosa</i> , Th.	1165
<i>arcuata</i> , Ky.	1170	<i>leptocarpus</i> , Ky.	1167
<i>Armeniaca</i> , Ky.	1164	<i>Libani</i> , Oliv.	1173
<i>Aucherii</i> , J. et Sp.	1170	<i>longifolia</i> , C. Koch	1164
<i>Austriaca</i> , W.	1171	<i>longipes</i> , Stev.	1164
<i>Balout</i> , Griff.	1168	<i>Loox</i> , Ky.	1172
<i>Boissieri</i> , Reut.	1167	<i>Lusitanica</i> , Lam. 1166 et Add.	
<i>brachybalanos</i> , Ky.	1169	<i>Macedonica</i> , A.D.C.	1172
<i>brachyphylla</i> , Ky.	1165	<i>macranthera</i> , F. et M.	1165
<i>Brantii</i> , Lindl	1173	<i>macrolepis</i> , Ky.	1171
<i>calliprinos</i> , Webb.	1169	<i>mannisera</i> , Lindl.	1165
<i>Carduchorum</i> , Ky.	1175	<i>oophora</i> , Ky.	1175
<i>eastancæfolia</i> , C. A. M.	1174	<i>Palestina</i> , Ky.	1170
<i>Cedrorum</i> , Ky.	1164	<i>pedunculata</i> , Ehrh.	1165
<i>Cerris</i> , L.	1170	<i>pedunculiflora</i> , C. Koch	1163
<i>cocculera</i> , L.	1169	<i>Persica</i> , J. et Sp.	1175
<i>conferta</i> , Kit.	1166	<i>petiolaris</i> , Boiss.	1167
<i>erinita</i> , Lam.	1171	<i>Pfeffingeri</i> , Ky.	1167
<i>Cypria</i> , Ky.	1167	<i>pinnatilobata</i> , C. Koch	1164
<i>Cypria</i> , J. et Sp.	1169	<i>polycarpus</i> , Ky.	1167
<i>dipsacina</i> , Ky.	1169	<i>Pontica</i> , C. Koch	1166
<i>dispar</i> , Ky.	1169	<i>pseudocerris</i> , Boiss.	1171
<i>Dschorocensis</i> , Ky.	1165	<i>pseudocoecisera</i> , Lab.	1169
<i>echinata</i> , Ky.	1169	<i>pseudo-Dchorocensis</i> ,	
<i>Ehrenbergii</i> , Ky.	1171	Ky.	1165
<i>erucæfolia</i> , St.	1164	<i>pubescens</i> , W.	1165
<i>faginea</i> , Lam.	1166	<i>Pyrami</i> , Ky.	1172
<i>Farnetto</i> , Ten.	1166	<i>racemosa</i> , Lam.	1165
<i>Fenzlii</i> , Ky.	1169	<i>regia</i> , Lindl.	1174
<i>Gödelii</i> , Ky.	1172	<i>rigida</i> , C. Koch	1167
<i>Græca</i> , Ky.	1171	<i>rigida</i> , W.	1170
<i>Haus</i> , Ky.	1165	<i>Robur</i> , L.	1164
		<i>sessiliflora</i> , Sm.	1164

INDEX.

1253

Pág.

<i>Quercus Sibthorpii</i> , Ky.	1169
<i>subalpina</i> , Ky.	1164
<i>Suber</i> , L.	1168
<i>Syriaca</i> , Ky.	1167
<i>Syssirensis</i> , A.D.C.	1165
<i>Syssirensis</i> , C. Koch	1165
<i>Tauricola</i> , Ky.	1167
<i>Tchihatchewii</i> , Ky.	1174
<i>Toza</i> , Griseb.	1164
<i>valida</i> , Ky.	1169
<i>Vallonea</i> , A.D.C.	1172
<i>Vallonea</i> , Ky.	1171
<i>vesca</i> , Ky.	1174
<i>vulcanica</i> , Boiss.	1164
<i>Ramoudia Heldreichii</i> , Janka	85
<i>Razumowskia Caucasia</i> ,	
Hoffm.	1069
<i>Reptonia</i> , A.D.C.	52
<i>buxifolia</i> , Falc.	52
<i>buxifolia</i> , A.D.C.	55
<i>Rhamnus carpinifolia</i> , Pall.	1159
<i>Rhazya</i> , Decaisne	46
<i>Orientalis</i> , Decaisne	47
<i>Orientalis</i> , D.C.	47
<i>stricta</i> , Decaisne	46
<i>Rheum</i> , L.	1005
<i>Caspicum</i> , Pall.	1005
<i>Ribes</i> , Gron.	1005
<i>Tataricum</i> , L. fil.	1005
<i>Rhinanthus</i> , L.	479
<i>alpestris</i> , Wahl.	480
<i>Alpinus</i> , Baumg.	480
<i>Cristagalli</i> , L.	479
<i>Elephas</i> , L.	478
<i>major</i> , Ehrh.	479
<i>maximus</i> , W.	474
<i>minor</i> , Ehrh.	479
<i>Orientalis</i> , L.	478
<i>pubescens</i> , B. et Heldr.	479
<i>strictus</i> , C. Koch	478
<i>trifidus</i> , Vahl	471
<i>Rhus obscurum</i> , M. B.	1161
<i>Rhynchosoryx</i> , Gr.	477
<i>Elephas</i> , L.	473

<i>Rhynchosoryx Orientalis</i> , L.	478
<i>Ricinus</i> , L.	1145
<i>communis</i> , L.	1145
<i>Rindera</i> , Pall.	275
<i>cyclodonta</i> , Bge.	275
<i>Græca</i> , B. et Heldr.	275
<i>glutifolia</i> , R. et Sch.	262
<i>pubescens</i> , C. Koch	272
<i>Rivina pinniculata</i> , L.	45
<i>Rochelia</i> , Reh.	245
<i>ecellata</i> , B. et Bal.	245
<i>cardiosepala</i> , Bge.	246
<i>incana</i> , Kar. et Kir.	244
<i>leiocarpa</i> , Led.	245
<i>macrocalyx</i> , Bge.	246
<i>mollis</i> , Bge.	245
<i>peduncularis</i> , Boiss.	246
<i>Persica</i> , Bge.	244
<i>rectipes</i> , Stocks	246
<i>stellulata</i> , Rehb.	244
<i>stylaris</i> , Boiss.	245
<i>Rosmarinus</i> , L.	636
<i>offinalis</i> , L.	525
<i>Rostellaria peploides</i> , Nees	525
<i>Ruellia</i> , L.	519
<i>ciliaris</i> , L.	520
<i>longifolia</i> , Stocks	519
<i>Persica</i> , Burm.	520
<i>Rumex</i> , L.	1005
<i>Acetosa</i> , L.	1015
<i>Acetosella</i> , L.	1018
<i>acetoselloides</i> , Bal.	1018
<i>aculeatus</i> , L.	1015
<i>Egyptiacus</i> , L.	1014
<i>Alpinus</i> , L.	1007
<i>angustifolius</i> , Campd.	1008
<i>aquaticus</i> , L.	1008
<i>Armenus</i> , C. Koch	1019
<i>baecephalophorus</i> , L.	1014
<i>callosissimus</i> , Meisn.	1015
<i>Cassius</i> , Boiss.	1015
<i>C. tephensis</i> , Mill.	1019
<i>comosus</i> , Forsk.	1014
<i>condylodes</i> , M. B.	1010
<i>confertus</i> , W.	1007

	Pag.		Pag.
<i>Rumex conglomeratus, Murr.</i>	1010	<i>Rumex pulcher, L.</i>	1012
<i>Crelicus, Boiss.</i>	1017	<i>quadridentatus, Ehr.</i>	1015
<i>crispus, L.</i>	1009	<i>Raulini, Boiss.</i>	1012
<i>Daghestanicus, C. Koch</i>	1010	<i>reticulatus, Bess.</i>	1012
<i>dentatus, L.</i>	1015	<i>roseus, L.</i>	1018
<i>denticulatus, C. Koch</i>	1012	<i>Roxburghianus, Sch.</i>	1011
<i>dictyocarpus, B. et Buhse</i>	1101	<i>scutatus, L.</i>	1015
<i>digynus, L.</i>	1004	<i>spinosus, L.</i>	1005
<i>divaricatus, Fr.</i>	1011	<i>stenophyllus, Led.</i>	1012
<i>divaricatus, L.</i>	1012	<i>strictus, Link</i>	1015
<i>Ehrenbergii, Meisn.</i>	1013	<i>strictus, Meisn.</i>	1013
<i>Elbrusensis, Boiss.</i>	1008	<i>subtrilobus, Boiss.</i>	1011
<i>elongatus, Guss.</i>	1010	<i>Syriacus, Meisn.</i>	1011
<i>foveolatus, Hochst.</i>	1012	<i>Tmoleus, Boiss.</i>	1016
<i>Friesii, Gr. et Godr.</i>	1101	<i>triangularis, Guss.</i>	1016
<i>glaber, Forsk.</i>	1005	<i>tuberculatus, C. Koch</i>	1012
<i>glomeratus, Schr.</i>	1010	<i>tuberous, L.</i>	1017
<i>Græcus, B. et Heldr.</i>	1009	<i>tuberous, Roxb.</i>	1011
<i>hamatus, Trev.</i>	1011	<i>Turcicus, Boiss.</i>	1010
<i>hamulosus, Meisn.</i>	1011	<i>uncinatus, Hort.</i>	1011
<i>hastatus, Don.</i>	1015	<i>vesicarius, L.</i>	1017
<i>hastifolius, M. B.</i>	1015		
<i>Heleolapathus, Dreb.</i>	1008	Salicornia, L.	932
<i>Hippolapathum, Fr.</i>	1008	<i>acetaria, Pall.</i>	933
<i>horizontalis, C. Koch</i>	1016	<i>Alpini, Lag.</i>	935
<i>Klotschianus, Meisn.</i>	1013	<i>amplexicaulis, Vahl</i>	934
<i>lacerus, Balb.</i>	1017	<i>anceps, Lag.</i>	953
<i>Lachanus, Forsk.</i>	1019	<i>annua, Engl. Bot.</i>	955
<i>Limoniastrum, J. et Sp.</i>	1007	<i>Arabica, Pall.</i>	955
<i>macranthus, Boiss.</i>	1008	<i>Caspica, L.</i>	953
<i>Menzalensis, Ehr.</i>	1013	<i>Caspica, Pall. ex parte</i>	934 et 935
<i>multifidus, L.</i>	1018	<i>cruciata, Forsk.</i>	956
<i>multifidus, Fl. Gr.</i>	1019	<i>Europæa, Forsk.</i>	933
<i>Nemolapathum, M. B.</i>	1010	<i>foliata, Pall.</i>	934
<i>nemorosus, Schrad.</i>	1010	<i>fruticosa, L.</i>	952
<i>Nepalensis, Spr.</i>	1011	<i>glouca, Del.</i>	952
<i>obtusifolius, L.</i>	1011	<i>herbacea, L.</i>	952
<i>Olympicus, Boiss.</i>	1009	<i>macrostachya, Moric.</i>	932
<i>Orientalis, Bernh.</i>	1009	<i>nodulosa, Del.</i>	955
<i>palustris, Sm.</i>	1014	<i>Pallasii, C. A. M.</i>	955
<i>Patientia, L.</i>	1009	<i>persolliata, Forsk.</i>	955
<i>peregrinus, Boiss.</i>	1011	<i>prostrata, Pall.</i>	955
<i>pictus, Forsk.</i>	1017	<i>pygmæa, Pall.</i>	954
<i>pseudocetosa, Bert.</i>	1016	<i>radicans, Sm.</i>	955
<i>pubescens, C. Koch</i>	1015		

	PAG.		PAG.
Salicornia strobilacea , Gærtn.	934	Salix repens , L.	1190
<i>strobilacea</i> , Pall.	936	<i>retusa</i> , L.	1192
<i>strobilacea</i> , Del.	935	<i>rubra</i> , Huds.	1186
<i>Virginica</i> , Forsk.	932	<i>Safsaf</i> , Forsk.	1185
SALICINEÆ.	1181	<i>Silesiaea</i> , W.	1189
Salix , Tourn.	1181	<i>stylaris</i> , Ser.	1190
<i>aemophylla</i> , Boiss.	1185	<i>stylosa</i> , D.C.	1190
<i>acutifolia</i> , W.	1191	<i>sub serrata</i> , W.	1184
<i>Egyptiaca</i> , L.	1190	<i>tenuijulis</i> , Led.	1187
<i>Egyptiaca</i> , Auct.	1184	<i>triaurita</i> , L.	1186
<i>alba</i> , L.	1185	<i>viminalis</i> , L.	1191
<i>amygdalina</i> , L.	1186	<i>Waldsteiniana</i> , M. B.	1187
<i>angustifolia</i> , W.	1187	<i>Wilhelmsiana</i> , W.	1192
<i>apeda</i> , Trautv.	1192	<i>Zygostemon</i> , Boiss.	1188
<i>arbuseula</i> , L.	1192	SALSOLACEÆ	897
<i>aurita</i> , L.	1190	Salsola , L.	951
<i>Babylonica</i> , L.	1185	<i>Agrigentina</i> , Guss.	962
<i>Capræa</i> , L.	1188	<i>alopecuroides</i> , Del.	985
<i>cinerea</i> , L.	1188	<i>altissima</i> , L.	940
<i>Cyrene</i> , B. et Orph.	1188	<i>annularis</i> , Poir.	959
<i>daphnooides</i> , Vill.	1191	<i>anomala</i> , C. A. M.	947
<i>Daviesii</i> , Boiss.	1184	<i>Arabica</i> , Ehr.	970
<i>dracunculifolia</i> , Boiss.	1187	<i>arborescens</i> , L. fil.	960
<i>Elbursensis</i> , Boiss.	1187	<i>arbuscula</i> , Pall.	960
<i>fragilis</i> , L.	1184	<i>arenaria</i> , W. K.	924
<i>hastata</i> , L.	1191	<i>articulata</i> , Forsk.	970
<i>Helix</i> , L.	1186	<i>aurantiaca</i> , Bge.	965
<i>hippophaëfolia</i> , W.	1186	<i>Auriella</i> , Moq.	958
<i>incana</i> , Schr.	1187	<i>baccata</i> , Poir.	944
<i>Kasanogluensis</i> , Ky.	1185	<i>baccifera</i> , Pall.	955
<i>Ledebouriana</i> , Tr.	1186	<i>Baltw.</i> , J. et Sp.	964
<i>Libanotica</i> , Boiss.	1185	<i>brachiata</i> , Pall.	952
<i>Medemii</i> , Boiss.	1189	<i>brachiphylla</i> , B. et	
<i>micans</i> , And.	1185	<i>Hausk.</i>	959
<i>nigricans</i> , Fr.	1190	<i>brevifolia</i> , Desf.	962
<i>oetandra</i> , Sieb.	1184	<i>cassia</i> , F. et M.	962
<i>pallida</i> , Ledeb.	1186	<i>cana</i> , C. Koch	956
<i>pedicellata</i> , Desf.	1189	<i>canescens</i> , Moq.	965
<i>pentandra</i> , L.	1184	<i>carinata</i> , C. A. M.	951
<i>Persica</i> , Boiss.	1185	<i>cinerata</i> , W. K.	926
<i>phlomoides</i> , M. B.	1189	<i>clavifolia</i> , Pall.	953
<i>præcox</i> , Hoppe	1191	<i>clarifolia</i> , Homb.	542
<i>prunifolia</i> , Sm.	1192	<i>crassa</i> , M. B.	956
<i>pruinosa</i> , Wend.	1191	<i>decurvens</i> , J. et Sp.	958
<i>purpurea</i> , L.	1186	<i>dendroides</i> , Pall.	961

	Pag.		Pag.
<i>Salsola divergens</i> , Poir.	945	<i>Salsola pilosa</i> , Pall.	980
<i>echinus</i> , Lab.	963	<i>prostrata</i> , L.	925
<i>eriantha</i> , Host.	927	<i>pulla</i> , C. Koch	952
<i>ericoides</i> , M. B.	961	<i>recurra</i> , Wall.	949
<i>ericoides</i> Var. <i>leucoxylon</i> ,		<i>rigida</i> , Pall.	962
Moq.	954	<i>rosacea</i> , M. B.	956
<i>farenosa</i> , Poir.	959	<i>rosacea</i> , Sm.	954
<i>ferox</i> , Lip.	984	<i>salsa</i> , Jacq.	940
<i>flavescens</i> , Cav.	962	<i>scelerantha</i> , C. A. M.	986
<i>florida</i> , Poir.	950	<i>scoparia</i> , M. B.	925
<i>foetida</i> , Del.	961	<i>sedoides</i> , Pall.	926
<i>foliosa</i> , Schrad.	955	<i>Soda</i> , L.	915
<i>fruticosa</i> , L.	955	<i>Sogdiana</i> , Bge.	945
<i>gemmaeens</i> , Pall.	960	<i>spicata</i> , Pall.	949
<i>Georgica</i> , Bge.	961	<i>Spinifex</i> , Pall.	960
<i>glauca</i> , M. B.	959	<i>spissa</i> , M. B.	954
<i>glauca</i> , C. A. M.	962	<i>Stocksii</i> , Boiss.	950
<i>globulifolia</i> , Poir.	940	<i>suædoides</i> , Moq.	954
<i>gossypina</i> , Bge.	956	<i>subaphylla</i> , C. A. Mey.	959
<i>heterocarpa</i> , Bge.	955	<i>tamariscina</i> , Pall.	955
<i>hirsuta</i> , L.	925	<i>tetragona</i> , Del.	957
<i>hispidula</i> , Bge.	960	<i>tetrandra</i> , Forsk.	957
<i>hyssopifolia</i> , Pall.	926	<i>Tragus</i> , L.	954
<i>inecanescens</i> , C. A. M.	954	<i>vermiculata</i> , L.	962
<i>inermis</i> , Forsk.	955	<i>verrucosa</i> , C. A. Mey.	960
<i>Kali</i> , L	954	<i>verrucosa</i> , M. B.	961
<i>Lana</i> , Edgew.	950	<i>villosa</i> , Sieb.	955
<i>Lana</i> , Stocks	950	<i>villosa</i> , Del.	965
<i>lanata</i> , Pall.	956		45
<i>lanceolata</i> , Boiss.	958	SalVADORACEÆ	
<i>laniflora</i> , L. fil.	957	Salvadora , Gare.	45
<i>laniflora</i> , Host.	927	<i>oleoides</i> , D. C.	44
<i>laricina</i> , Pall.	962	<i>Persica</i> , Gare.	45
<i>longifolia</i> , Forsk.	957	Salvia , L.	590
<i>microphylla</i> , Cav.	962	<i>acetabulosa</i> , Vahl.	608
<i>mollis</i> , Desf.	940	<i>Ægyptiaca</i> , L.	651
<i>monobractea</i> , Forsk.	927	<i>Æthiopis</i> , L.	616
<i>Moquiniana</i> , Jaub. et Sp.	958	<i>albida</i> , Jacq.	622
<i>muclronata</i> , Forsk.	965	<i>Argaa</i> , B. et Bal.	601
<i>muricata</i> , L.	926	<i>argentea</i> , L.	618
<i>nitraria</i> , Pals.	954	<i>argentea</i> , var.	620
<i>Pallasiana</i> , Vest.	950	<i>aristata</i> , Auch.	617
<i>Passerina</i> , Bge.	960	<i>Armena</i> , C. Koch	622
<i>Persica</i> , Bge.	969	<i>atomaria</i> , Boiss.	619
		<i>Atropatana</i> , Bge.	619
		<i>Aucheria</i> , Bth.	395

Pag.		Pag.
596	<i>Salvia</i> Forskahlei, L.	609
626	<i>fragifera</i> , Etzl.	604
620	<i>frigida</i> , B. et Buhse	620
627	<i>frigida</i> , Boiss.	621
624	<i>fruticosa</i> , Mill.	596
600	<i>Gaillardottii</i> , Boiss.	628
610	<i>Gargantica</i> , Ten.	628
606	<i>glutinosa</i> , L.	609
625	<i>grandiflora</i> , Etzl.	595
605	<i>graveolens</i> , Vahl	615
605	<i>Habititziana</i> , W.	601
594	<i>Hanssknechtii</i> , Boiss.	605
604	<i>Heldreichiana</i> , Boiss.	690
626	<i>Hierosolymitana</i> , Boiss.	627
605	<i>horminoidea</i> , Pourr.	650
628	<i>Horninum</i> , L.	651
619	<i>Hydrangea</i> , D. C.	606
621	<i>hyemalis</i> , Brot.	650
612	<i>hypargyria</i> , F. et M.	612
602	<i>hypoachionwa</i> , Buhse.	620
596	<i>hypachionwa</i> , Boiss.	612
617	<i>hypoleuca</i> , Bth.	625
599	<i>Judaica</i> , Boiss.	685
618	<i>incarnata</i> , Etzl.	596
625	<i>incarnata</i> , Etzl.	601
622	<i>incarnata</i> , var.	602
629	<i>Indica</i> , L.	625
650	<i>Kochiana</i> , Kze.	616
616	<i>Kotschyii</i> , Boiss.	615
611	<i>Kurtaica</i> , B. et Hoh.	595
650	<i>laciniata</i> , W.	650
622	<i>lanigera</i> , Poir.	650
604	<i>leriaefolia</i> , Bth.	615
608	<i>leuconeura</i> , Boiss.	517
622	<i>Libanotica</i> , B. et Gaill.	594
596	<i>limbata</i> , C. A. M.	625
652	<i>Linkiana</i> , R. et Sch.	656
629	<i>longepetiolata</i> , C. Koch.	610
529	<i>Lorentii</i> , Hochst.	656
616	<i>macilenta</i> , Boiss.	655
606	<i>macrochlamys</i> , B. et Ky.	595
627	<i>macrosiphon</i> , Boiss.	615
652	<i>microstegia</i> , B. et Bal.	619
611	<i>modesta</i> , Boiss.	621
605	<i>Moluccar</i> , Bth.	608

	Pag.		Pag.
<i>Salvia Montbretii</i> , Bth.	611	<i>Salvia ringens</i> , Fl. Gr.	600
<i>multicaulis</i> , Vahl.	608	<i>rosæfolia</i> , Sm.	596
<i>multifida</i> , Fl. Gr.	630	<i>rotundifolia</i> , Vis.	594
<i>napifolia</i> , Jacq.	655	<i>rubifolia</i> , Boiss.	602
<i>nemorosa</i> , L.	629	<i>rugosissima</i> , Zucc.	630
<i>nivea</i> , Ehr.	619	<i>Ruprechtii</i> , Boiss.	624
<i>nudicaulis</i> , C. Koch	628	<i>Russeggeri</i> , Fenzl	599
<i>nutans</i> , L.	629	<i>Russelii</i> , Bth.	655
<i>oblonga</i> , C. Koch	628	<i>Sahendica</i> , B. et Buhse	624
<i>oblongata</i> , Vahl	629	<i>sautolinæfolia</i> , Boiss.	632
<i>odorata</i> , W.	622	<i>scabiosæfolia</i> , Lam.	600
<i>oligophylla</i> , Auch.	623	<i>Sclarea</i> , L.	616
<i>oreades</i> , Schott.	621	<i>Sheilei</i> , Boiss.	607
<i>orygialis</i> , Fenzl.	601	<i>Sibthorpii</i> , Bory	629
<i>Owerini</i> , Trantv.	617	<i>Sibthorpii</i> , Fl. Gr.	628
<i>pachystachys</i> , Tr.	398	<i>Sieberi</i> , Presl	615
<i>Palestina</i> , Bth.	614	<i>Sinaica</i> , Del.	615
<i>parviflora</i> , Vahl	610	<i>sinapifolia</i> , C. Koch	655
<i>pectinifolia</i> , F. et M.	598	<i>Sipylea</i> , Lam.	596
<i>Peloponnesiaca</i> , B. et		<i>Smyrnaea</i> , Boiss.	603
<i>Haussk.</i>	631	<i>Spielmanniana</i> , M. B.	629
<i>Persepolitana</i> , Boiss	614	<i>spinosa</i> , L.	615
<i>phlyctidea</i> , C. Koch	615	<i>spinulosa</i> , M. et Auch.	621
<i>ilantha</i> , Link	636	<i>spireæfolia</i> , B. et Hoh.	597
<i>pilifera</i> , M. et Auch.	602	<i>staminea</i> , M. et Auch.	626
<i>Pinardi</i> , Boiss.	609	<i>snfruticosa</i> , M. et Auch.	597
<i>pinifolia</i> , Pall.	601	<i>sylvestris</i> , L.	628
<i>pinnata</i> , L.	601	<i>Syriaca</i> , L.	610
<i>Pisidica</i> , B. et Heldr.	598	<i>Szovitsiana</i> , Bge.	607
<i>polyadenia</i> , Boiss.	624	<i>Tauricola</i> , Sch. et Ky.	608
<i>polyclonus</i> , Boiss.	611	<i>Tchiba'chewii</i> , Boiss.	598
<i>polymorpha</i> , Hoffm.	650	<i>Tebesana</i> , Bge.	635
<i>pomifer</i> , L.	604	<i>Tenorii</i> , Spr.	627
<i>potentillæfolia</i> , B. et		<i>Tmolea</i> , Boiss.	618
<i>Heldr.</i>	597	<i>trichocalycina</i> , Bth.	655
<i>præcox</i> , Vahl	614	<i>trichoclada</i> , Bth.	603
<i>præcox</i> , Savi	650	<i>trileba</i> , L. fil.	593
<i>pratensis</i> , L.	626	<i>truncata</i> , W.	631
<i>pseudæthiopis</i> , Boiss.	619	<i>Urmiensis</i> , Bge.	620
<i>pumila</i> , Bth.	652	<i>Vanensis</i> , B. et Noë	597
<i>Rascheyana</i> , Boiss.	607	<i>varia</i> , Vahl	610
<i>Rassami</i> , Boiss.	615	<i>verbascifolia</i> , M. B.	619
<i>recognita</i> , F. et M.	601	<i>Verbenaca</i> , L.	629
<i>Reuteriana</i> , Boiss.	614	<i>verticillata</i> , L.	654
<i>rhytidæ</i> , Bth.	610	<i>virgata</i> , Ait.	627

	Pag.		Pag.
Salvia viridis, L.	650	Sch. anginaria hortensis, Forsk.	945
viscosa, Jacq.	628	linifolia, Pall.	944
Wiedemannii, Boiss.	599	Schoberia acuminata, C. A. M.	942
hocheila, Boiss.	620	altissima, C. A. M.	941
Samolus, L.		dendroides, C. A. M.	958
Valerandi, L.	5	heterophylla, Kar. et Kir.	945
SANTALACEÆ.	1058	leiosperma, C. A. M.	941
Sarcostemma, R. Br.	58	maritima, C. A. M.	941
aphyllum, R. Br.	58	microphylla, C. A. M.	958
pyrotechnicum, R. Br.	63	microsperma, C. A. M.	945
Satureia, L.	562	Pallasiana, C. A. M.	926
alternipilosa, C. Koch	566	salsa, C. A. M.	942
approximata, Friv.	565	setigera, C. A. M.	942
Argæa, F. et M.	556	suffruticulosa, Less.	959
Atropatana, Bye.	567	Schweinfurthia, A. Br.	386
Bachtiarica, Bye.	567	papilionacea, Burm.	587
Boissieri, Haussk.	563	pterosperma, A. Br.	386
capitata, L.	560	Scoparia Boreana, Dun.	294
cuneifolia, Ten.	564	Daiora, Dun.	294
diffusa, Bth.	566	mutica, Dun.	294
filicaulis, Sch.	562	Scordium spinosum, Cav.	814
Græca, L.	571	SCROPHULARIACEÆ.	297
hortensis, L.	562	Scrophularia, L.	387
Juliana, L.	569	aestivalis, Cr.	397
intermedia, Bth.	563	Aintabensis, B. et Haussk.	415
intermedia, C. A. Mey.	564	alata, Gilib.	399
intermedia var. laxior,		amplexicaulis, Bth.	394
Bth.	566	Ani, C. Koch	418
laxiflora, C. Koch	562	aquatica, C. Koch	400
longitlora, B. et Haussk.	566	arguta, Sol.	595
montana, L.	563	anriiculata, Scop.	396
maerantha, C. A. Mey	566	Balbisii var. macro-	
mutica, F. et M.	565	phylla, Boiss.	400
nervosa, Desf.	569	Benthamiana, Boiss.	421
pachyphyllea, C. Koch	562	bicolor, Fl. Gr.	419
Parnassica, H. et Sart.	565	Boissieriana, J. et Sp	595
plumosa, Hampe	569	Byzantina, Bth.	591
spicigera, C. Koch	566	Cabulica, Bth.	420
spinosa, L.	563	cæsia, Fl. Gr.	408
stricta, Russ.	669	cæsia, Griseb.	409
subdentata, Boiss.	565	Calverti, Boiss.	595
Thymbra, L.	567	calycina, Boiss.	590
virgata, Vis.	564	canina, Fl. Gr.	405
Sch. anginaria, C. A. M.	944	canina, L.	419
baccata, Forsk.	944	capillaris, B. et Bal.	597

PAG.		PAG.
596	<i>Serophularia Castagueana</i> , Wydl.	408
407	<i>cutariæfolia</i> , Boiss. et <i>He'dr.</i>	402
418	<i>chamadryfolia</i> , B. et Hanssk.	415
599	<i>chlorantha</i> , Ky. et Boiss.	421
590	<i>clrysanthia</i> , J. et Sp.	
419	<i>chrysanthemifolia</i> , W.	417
408	<i>chrysanthemifolia</i> , Fl. Pel.	418
417	<i>cinerascens</i> , Boiss.	594
592	<i>clandestina</i> , Rupr.	418
591	<i>Claussii</i> , B. et Buhse	390
590	<i>congesta</i> , Stev.	417
420	<i>crassicaulis</i> , Boiss.	408
596	<i>crenophilis</i> , Boiss.	408
400	<i>Cretica</i> , B. et Heldr.	592
591	<i>cryptophila</i> , B. et Heldr.	404
400	<i>cymigera</i> , C. Koch	409
415	<i>decipiens</i> , Boiss. et Ky	409
596	<i>decumbens</i> , F. et M.	403
410	<i>depanp'ata</i> , Boiss.	421
414	<i>deserti</i> , Del.	418
597	<i>divaricata</i> , Led.	407
592	<i>ebulifolia</i> , M. B.	406
400	<i>Ehrharti</i> , Stev.	405
415	<i>expansa</i> , Rant.	405
405	<i>farnosia</i> , Boiss.	404
405	<i>fiticifolia</i> , Fl. Gr.	590
419	<i>flabibunda</i> , B. et Bal.	402
596	<i>fountainia</i> , Ky.	Add.
411	<i>frixiola</i> , Boiss.	
409	<i>frutescens</i> , Sm.	
597	<i>Georgica</i> , Bth.	
596	<i>glandulifera</i> , Cl.	
596	<i>glandulosa</i> , W. K.	
416	<i>glauca</i> , Decaisne	
405	<i>glauca</i> , Fl. Gr.	
596	<i>grandidentata</i> , Ten.	
410	<i>Griffithii</i> , Bth.	
415	<i>haematantha</i> , B. et Heldr.	
415	<i>Hellreichii</i> , Boiss.	
599	<i>Hemschinica</i> , C. Koch	
409	<i>heterophylla</i> , Fl. Gr.	
408	<i>Serophularia heterophylla</i> , W.	418
402	<i>Hierochuntina</i> , Boiss.	418
415	<i>hispidula</i> , B. et Bal.	415
421	<i>hypericifolia</i> , Wydl.	415
418	<i>hypsopifolia</i> , Boiss. et Haussk.	418
594	<i>Iwensis</i> , C. Koch	418
418	<i>incisa</i> , C. Koch	418
390	<i>Kotschyana</i> , Bth.	417
417	<i>lacinifolia</i> , C. Koch	417
408	<i>lacinifolia</i> , W. K.	408
592	<i>lateriflora</i> , Trautv.	592
404	<i>laxa</i> , B. et Heldr.	404
409	<i>Lazica</i> , Boiss.	409
403	<i>lepidota</i> , Boiss.	403
421	<i>leneocladia</i> , Bge.	421
418	<i>Libanotica</i> , Boiss.	418
405	<i>livida</i> , Fl. Gr.	405
406	<i>longiflora</i> , Bth.	406
405	<i>lucida</i> , L.	405
404	<i>lucida</i> , M. B.	404
590	<i>lunariaefolia</i> , B. et Bal.	590
598	<i>luridiflora</i> , F. et M.	598
599	<i>macrobotrys</i> , Led.	599
400	<i>macrophylla</i> , Boiss.	400
414	<i>mauginata</i> , Boiss.	414
596	<i>melissæfolia</i> , Gr.	596
407	<i>Mesogitana</i> , Boiss.	407
402	<i>Mesopotamica</i> , Boiss.	402
402	<i>Michoniana</i> , Cass.	402
408	<i>micrantha</i> , Urv.	Add.
590	<i>minima</i> , Bth.	408
595	<i>minima</i> , M. B.	590
409	<i>multifida</i> , W.	595
419	<i>multifida</i> , Lam.	409
418	<i>multifida</i> var. <i>densiflora</i> , Bth.	419
404	<i>myriophylla</i> , B. et Heldr.	418
400	<i>Neesii</i> , Wirtz.	404
407	<i>nepetaefolia</i> , B. et Heldr.	400
592	<i>nervosa</i> , Bth.	407
599	<i>nodosa</i> , L.	592
556	<i>nudicaulis</i> , Wydl.	599
596	<i>oligantha</i> , B. et Heldr.	556
409	<i>Oliveriana</i> , C. Koch	596

INDEX.

[33]

PAG.		PAG.
	<i>Serophularia Oliveriana</i> ,	
408	Wydl.	40
592	<i>Olivieri</i> , J. et Sp.	41
409	<i>Olympica</i> , Boiss.	41
592	<i>Orientalis</i> , L.	58
593	<i>Orientalis</i> , Boiss.	58
398	<i>oxysepaia</i> , Boiss.	58
593	<i>peregrina</i> , L.	58
411, 415 et 418	<i>Persica</i> , Bth.	681
417	<i>Pinardi</i> , Boiss.	681
400	<i>Pisidica</i> , B. et Heldr.	688
575	<i>polybotrys</i> , Boiss.	689
416	<i>prasiifolia</i> , B. et Haussk.	685
416	<i>pruinosa</i> , Boiss.	688
418	<i>puberula</i> , Boiss.	682
596	<i>puberula</i> , B. et Haussk.	682
401	<i>puilverlenta</i> , B. et Noë	685
407	<i>pyramidalis</i> , Wydl.	685
409	<i>pyrrholopha</i> , Boiss.	685
419	<i>ramosissima</i> , Urv.	685
412	<i>rostrata</i> , B. et Buhse	685
593	<i>rostrata</i> , Hœchst.	811
402	<i>rubricaulis</i> , Boiss.	689
418	<i>rupestris</i> , M. B.	687
418	<i>Ruprechtii</i> , Boiss.	688
404	<i>rutæfolia</i> , Boiss.	688
400	<i>Samaritanii</i> , B. et Heldr.	685
412	<i>seabiosæfolia</i> , Bth.	685
401	<i>seariosa</i> , Boiss.	685
593	<i>Scopolii</i> , Hoppe	685
596	<i>Scorodonia</i> , Host.	685
409	<i>Sibthorpiana</i> , Spreng.	685
414	<i>Sinica</i> , Bth.	685
596	<i>Smyrnæa</i> , Boiss.	685
405	<i>sphærocarpa</i> , B. et R.	685
415	<i>striata</i> , Boiss.	685
411	<i>subaphylla</i> , Boiss.	685
598	<i>sylvatica</i> , B. et Heldr.	685
420	<i>Syriaca</i> , Bth.	685
401	<i>tagetifolia</i> , B. et Haussk.	685
406	<i>Taygetea</i> , Boiss.	685
419	<i>thesioides</i> , B. et Buhse	685
596	<i>Tmolea</i> , Boiss.	685
412	<i>trichopoda</i> , B. et Bal.	685
418	<i>Urvilleana</i> , Decaisne	685
	<i>Serophularia Urvilleana</i> ,	
	Wydl.	40
	<i>variegata</i> , M. B.	41
	<i>vernalis</i> , L.	58
	<i>versicolor</i> , Boiss.	58
	<i>viscosa</i> , Boiss.	58
	<i>xanthoglossa</i> , Boiss.	58
	<i>xylorrhiza</i> , B. et Haussk.	409
	<i>Scutellaria</i> , L.	681
	<i>adenotricha</i> , B. et Haussk.	688
	<i>albida</i> , L.	689
	<i>Alpina</i> , L.	685
	<i>altissima</i> , L.	688
	<i>altissima</i> var. Mey.	687
	<i>bicolor</i> , Hœchst.	682
	<i>Caucasica</i> , Ham.	682
	<i>Chorassanica</i> , Bge.	685
	<i>Columnæ</i> , Host.	689
	<i>Columnæ</i> , All.	687
	<i>Columnæ</i> var. Bth.	688
	<i>cretacea</i> , B. et Haussk.	685
	<i>Cretica</i> , L.	811
	<i>Cretica</i> , Mill.	689
	<i>decumbens</i> , Sieb.	690
	<i>diffusa</i> , Bth.	686
	<i>fruticosa</i> , Desf.	684
	<i>galeruculata</i> , L.	690
	<i>glechomoides</i> , Boiss.	685
	<i>glutinosa</i> , Bth.	690
	<i>Gussonei</i> , Ten.	688
	<i>hastifolia</i> , L.	691
	<i>Hausknechti</i> , Boiss.	684
	<i>heterophylla</i> , M. et Anch.	687
	<i>hirta</i> , Sieb.	689
	<i>hirta</i> , Fl. Gr.	690
	<i>multicanalis</i> , Boiss.	685
	<i>nepetifolia</i> , Bth.	685
	<i>nigrescens</i> , Spr.	689
	<i>nummularia</i> , Fenzl	687
	<i>Orientalis</i> , L.	682
	<i>pallida</i> , Guss.	688
	<i>pallida</i> , M. B.	689
	<i>Parnassia</i> , H. et Sart.	688
	<i>pectinata</i> , M. et Anch.	685
	<i>peregrina</i> , L.	685

	PAG.		PAG.
Scutellaria peregrina , Fl. Gr.	688	Sideritis erythrantha , B.	
<i>peregrina</i> , W. K.	689	<i>et Heldr.</i>	709
<i>pinnatifida</i> , Ham.	683	<i>florida</i> , B. et Heldr.	710
<i>Pontica</i> , C. Koch	686	<i>Georgica</i> , C. Koch	710
<i>rubicunda</i> , Horn.	688	<i>hololeuca</i> , B. et Heldr.	709
<i>rupestris</i> , B. et Heldr.	690	<i>lanata</i> , L.	705
<i>salviæfolia</i> , Bth.	685	<i>Libanotica</i> , Lab.	712
<i>Sibthorpii</i> , B. et R.	688	<i>Lycia</i> , B. et Heldr.	709
<i>Sicberi</i> , Bth.	689	<i>microstegia</i> , B. et Haussk.	712
<i>Stocksii</i> , Boiss.	684	<i>montana</i> , L.	706
<i>Szovitsiana</i> , Bge.	685	<i>montana</i> , Fl. Gr.	707
<i>tomentosa</i> , Bert.	684	<i>mutica</i> , Boiss.	706
<i>Tournefortii</i> , Bth.	687	<i>nigricans</i> , Pers.	706
<i>utriculata</i> , Lab.	690	<i>Peloponnesiaca</i> , B. et	
<i>virens</i> , B. et Ky.	682	<i>Heldr.</i>	711
Seddera , Hochst.	114	<i>perfoliata</i> , L.	714
<i>latifolia</i> , Hochst.	114	<i>phlomoides</i> , B. et Bal.	711
Seidlitzia , Bge.	950	<i>Pisidica</i> , B. et Heldr.	712
<i>florida</i> , M. B.	950	<i>pullulans</i> , Vent.	713
<i>Rosmarinus</i> , Ehr.	951	<i>purpurea</i> , Talb.	706
SELAGINACEÆ.	527	<i>remota</i> , Urv.	707
SESAMEÆ.	81	<i>Rœseri</i> , B. et Heldr.	710
Sesamum , L.	81	<i>rhytidæa</i> , Spr.	718
<i>Indicum</i> , L.	81	<i>Romana</i> , L.	706
<i>Orientale</i> , Cham.	81	<i>rugosa</i> , Poir.	708
Sestinia calycina , Boiss.	676	<i>Scardica</i> , Griseb.	710
Sevada , Moq.	986	<i>Sipylea</i> , Boiss.	708
<i>Schimpéri</i> , Moq.	989	<i>stricta</i> , B. et Heldr.	713
Sibthorpia , L.	428	<i>Syriaca</i> , Bory	711
<i>Africana</i> , L.	428	<i>Syriaca</i> , L.	708
Sideritis , L.	705	<i>Syriaca</i> , Pall.	710
<i>Ægyptiaca</i> , Pers.	725	<i>Taurica</i> , M. B.	709
<i>Alopecuros</i> , Scop.	750	<i>Taurica</i> var. <i>procumbens</i>	715
<i>ambigua</i> , Fenzl	712	<i>thecezans</i> , B. et Heldr.	710
<i>arguta</i> , Boiss.	712	Siphonostegia , Bth.	470
<i>Balansæ</i> , Boiss.	707	<i>Syriaca</i> , B. et R.	471
<i>Bourgæi</i> , Boiss.	712	SOLANACEÆ.	283
<i>Cilicica</i> , B. et Bal.	711	Solanum , L.	285
<i>condensata</i> , Boiss. et		<i>assimile</i> , Friv.	285
<i>Heldr.</i>	713	<i>coagulans</i> , Forsk.	286
<i>Cretica</i> , Boiss.	708	<i>Dilleniæ</i> , Sch.	284
<i>Cretica</i> , Sm.	711	<i>Dulcamara</i> , L.	285
<i>distans</i> , W.	710	<i>flavum</i> , Kit.	284
<i>dura</i> , Bth.	710	<i>gracilipes</i> , Decaisne	286
<i>elegans</i> , Murr.	706		

	Pag.	Pag.
Solanum Hierochontinum,		
Dun.	286	<i>Stachys affinis</i> , Fres. 740
<i>hirsutum</i> , Dun.	284	<i>aleurites</i> , B. et Heldr. 753
<i>humile</i> , Bernh.	284	<i>Alpina</i> , L. 718
<i>Judaicum</i> , Bess.	284	<i>angustifolia</i> , M. B. 752
Kieseritzkyi, C. A. M.	285	<i>annua</i> , L. 745
<i>Memphiticum</i> , Dun.	284	<i>Arabica</i> , Horn. 747
<i>miniatum</i> , Bernh.	284	<i>Argolica</i> , Boiss. 755
<i>moschatum</i> , Presl.	284	<i>Armena</i> , Boiss. 694
<i>nigrum</i> , L.	284	<i>arvensis</i> , L. 747
<i>ochroleucum</i> , Bart.	284	<i>atherocalyx</i> , C. Koch. 750
<i>parviflorum</i> , Bad.	284	<i>Aucherii</i> , Bth. 758
<i>Persicum</i> , W.	285	<i>Balanse</i> , B. et Ky. 722
<i>puniceum</i> , Gmel.	284	<i>ballotaformis</i> , Vatk. 755
<i>sanctum</i> , L.	286	<i>Benthamiana</i> , Boiss. 754
<i>Sinaicum</i> , Boiss.	285	<i>Betonica</i> , Bth. 752
<i>suffruticosum</i> , Schomb.	284	<i>Eithynira</i> , Boiss. 720
<i>villosum</i> , Lam.	285	<i>Bodeana</i> , Bge. 741
<i>villosum</i> , Mill.	284	<i>bombycinia</i> , Boiss. 755
Solenanthus , Led.	268	<i>brachyodonta</i> , Coss. 746
<i>amplifolius</i> , Boiss.	270	<i>Brantii</i> , Bth. 750
<i>Biebersteinii</i> , D.C.	269	<i>Brantii</i> , Huet 722
<i>brachystemon</i> , F. et M.	268	<i>burgsdorffioides</i> , Boiss. 748
<i>cerinthoides</i> , Boiss.	271	<i>Byzantina</i> , C. Koch 718
<i>circinnatus</i> , Led.	270	<i>calycantha</i> , M. B. 745
<i>conglobatus</i> , A.D.C.	270	<i>candida</i> , B. et Ch. 742
<i>dubius</i> , F. et M.	269	<i>candida</i> v. <i>chrysanthia</i> , Bth. 742
<i>eriocalycinus</i> , B. et Buhse	268	<i>canescens</i> , Bory 728
<i>mollissimus</i> , Lehm.	269	<i>Cassia</i> , Boiss. 721
<i>petiolaris</i> , D.C.	270	<i>Caucasica</i> , C. Koch 731
<i>rumbicifolius</i> , Boiss.	270	<i>chamaesilieritis</i> , B. et Bal. 744
Tournefortii, D.C.	269	<i>chrysantha</i> , Boiss. 742
Solenostoma , Hayne		<i>ciliaris</i> , Boiss. 718
56 et Add.		<i>Cilicica</i> , B. et Bal. 746
Argel, Del.	56	<i>citrina</i> , B. et Heldr. 744
Spinacia , L.	905	<i>congesta</i> , Urv. 731
<i>minor</i> , C. Koch	906	<i>Cretica</i> , Sieb. 718
<i>tetrandra</i> , Stev.	906	<i>Cretica</i> , Fl. Gr. 719
Spiroconus glaucus , Stev.	281	<i>dasyanthes</i> , Heldr. 721
Stachys , L.	714	<i>decumbens</i> , Pers. 727
<i>abrotanifolia</i> , B. et Heldr.	752	<i>decumbens</i> , W. 745
<i>acerosa</i> , Boiss.	757	<i>distans</i> , Bth. 752
<i>acutifolia</i> , Link	724	<i>diversifolia</i> , Boiss. 755
<i>adenocalyx</i> , C. Koch	746	<i>Ehrenbergii</i> , Boiss. 721
<i>Egyptiaca</i> , Pers.	740	<i>elata</i> , C. Koch 725

	Pag.		Pag.
<i>Stachys floribunda</i> , M. et Auch.	744	<i>Stachys micrantha</i> , C. Koch	746
<i>fragilis</i> , Vis.	750	<i>mollissima</i> , Willd.	727
<i>fruticulosa</i> , M. B.	757	<i>Montbreti</i> , Bth.	717
<i>Garana</i> , Boiss.	719	<i>mueronata</i> , Sieb.	757
<i>Gernanica</i> , L.	720	<i>multicaulis</i> , Bth.	758
<i>Græca</i> , B. et Heldr.	723	<i>nepetæfolia</i> , Desf.	750
<i>grandiflora</i> , Bth.	751	<i>neurocalycina</i> , Boiss.	748
<i>Haussknechtii</i> , Vatk.	741	<i>nivea</i> , Labill.	759
<i>Heldreichii</i> , Boiss.	721	<i>obliqua</i> , W. K.	717
<i>Heraclea</i> , Bot. Græc.	725	<i>obscura</i> , B. et Bal.	747
<i>Heraclea v. lutea</i> , Bth.	717	<i>obtusata</i> , Boiss.	740
<i>Huetii</i> , Boiss.	722	<i>obtusierena</i> , Boiss.	759
<i>hydrophila</i> , Boiss.	726	<i>Olympica</i> , Poir.	749
<i>hypoleuca</i> , C. Koch	725	<i>Orientalis</i> , Forsk.	740
<i>Iberica</i> , M. B.	751	<i>Orientalis</i> , Vahl	716
<i>inflata</i> , Bth.	739	<i>Palæstina</i> , L.	740
<i>intermedia</i> , Ait.	720	<i>Palæstina</i> , Vahl	740
<i>Italica</i> , Bot. Græc.	719	<i>palustris</i> , L.	726
<i>Iva</i> , Griseb.	742	<i>paniculata</i> , Bth.	706
<i>ixodes</i> , B. et Haussk.	738	<i>Parolinii</i> , Vis.	750
<i>Kotschy</i> , Boiss.	741	<i>parviflora</i> , Bth.	740
<i>Kurdica</i> , B. et Hoh.	734	<i>patula</i> , Griseb.	750
<i>lætevirens</i> , Ky. et B.	734	<i>pauciflora</i> , Bth.	740
<i>lanata</i> , Mœnch	741	<i>pauciflora</i> , Vis.	717
<i>lanata</i> , Jacq.	718	<i>penicillata</i> , H. et Sart.	720
<i>lavandulæfolia</i> , Vahl	743	<i>Persepolitana</i> , Boiss.	749
<i>laxa</i> , B. et Buhse	742	<i>Persica</i> , Gmel.	724
<i>Lazica</i> , Boiss.	745	<i>pilifera</i> , Bth.	738
<i>leucoglossa</i> , Griseb.	750	<i>Pinardi</i> , Boiss.	728
<i>leucoglossa</i> var. Boiss.	751	<i>pinetorum</i> , B. et Bal.	722
<i>Libanotica</i> , Bth.	717	<i>Piridica</i> , B. et Heldr.	720
<i>linearifolia</i> , C. Koch	750	<i>plebeia</i> , Vatk.	755
<i>longespicata</i> , B. et Ky.	725	<i>plumosa</i> , Griseb.	746
<i>longiflora</i> , B. et Bal.	729	<i>polystachya</i> , Ten.	720
<i>longifolia</i> , Bth.	750	<i>procumbens</i> , Lam.	729
<i>Lusitanica</i> , B. et Ch.	723	<i>pubescens</i> , Ten.	745
<i>lycopsisformis</i> , C. Koch	725	<i>pumila</i> , Russ.	745
<i>Macedonica</i> , Don.	749	<i>ramosissima</i> , M. et Auch.	747
<i>macrocheilos</i> , Boiss.	737	<i>recta</i> , L.	729
<i>macrosperma</i> , Boiss.	747	<i>Reinerti</i> , Heldr.	720
<i>maritima</i> , Urv.	745	<i>rosea</i> , Desf.	725
<i>maritima</i> , L.	745	<i>rosea</i> , Hohen.	720
<i>menthoidea</i> , Ky. et B.	727	<i>rupestris</i> , M. et Auch.	744
<i>Mersinea</i> , Boiss.	719	<i>satureioides</i> , M. et Auch.	748
<i>Messeniaca</i> , Boiss.	728	<i>satureioides</i> var. Bth.	748

	Pag.		Pag.
Stachys Scyronica , Boiss.	755	Statice Balansæ , Boiss.	860
<i>sericea</i> , Ledeb.	724	<i>bellidifolia</i> , D.C.	865
<i>setifera</i> , C. A. M.	724	<i>bellidifolia</i> , Fl. Gr.	861
<i>Sibirica</i> , Link	720	<i>Besseriana</i> , R. Sch.	854
<i>sideritis</i> , Vill.	729	<i>Bovei</i> , J. et Sp.	868
<i>sideritoides</i> , C. Koch	730	<i>bracteata</i> , Gir.	827
<i>Sieberi</i> , C. Koch	718	<i>Cabulica</i> , Boiss.	871
<i>spectabilis</i> , Chois.	723	<i>cancellata</i> , Beruh.	865
<i>spinosa</i> , L.	736	<i>carnosa</i> , Schr.	867
<i>spinulosa</i> , Fl. Gr.	746	<i>caryophyllacea</i> , Boiss.	858
<i>Spruneri</i> , Boiss.	736	<i>Caspia</i> , Willd.	865
<i>stenostachya</i> , Boiss.	751	<i>collina</i> , Griseb.	853
<i>subnuda</i> , M. et Auch.	734	<i>cordata</i> , Desf.	865
<i>Swainsoni</i> , Bth.	735	<i>cordata</i> , Guss.	864
<i>sylvatica</i> , L.	726	<i>coriaria</i> , Hoffm.	864
<i>tenuifolia</i> , Pall.	732	<i>Corinthiaea</i> , B. et Heldr.	850
<i>tetragona</i> , B. et Heldr.	736	<i>Cosyrensis</i> , Guss.	865
<i>teucriifolia</i> , B. et Ky.	732	<i>Dalmatica</i> , Presl	853
<i>Thirkei</i> , C. Koch	719	<i>dianthifolia</i> , J. et Sp.	852
<i>Tmolea</i> , Boiss.	717	<i>dichotoma</i> , Sm.	864
<i>tomentosa</i> , Bth.	741	<i>echinus</i> , M.B.	854
<i>tomentosa</i> , Bge.	745	<i>echinus</i> , L. ex parte	840
<i>Tournefortii</i> , B. et Ch.	728	<i>echinus</i> , L. ex parte	854
<i>Tournefortii</i> , Poiret	718	<i>echoides</i> , L.	870
<i>Trapezuntea</i> , Boiss.	726	<i>echoides</i> , Sm.	862
<i>virgata</i> , B. et Ch.	731	<i>effusa</i> , Boiss.	859
<i>viridis</i> , B. et Heldr.	727	<i>erinacea</i> , J. et Sp.	853
<i>viscosa</i> , M. et Auch.	734	<i>ferox</i> , J. et Sp.	850
<i>viscosa</i> var. <i>Kotschy</i> , Bth.	735	<i>festucacea</i> , J. et Sp.	845
<i>viticina</i> , Boiss.	724	<i>Fischeri</i> , Trautv.	864
<i>Zuccarinii</i> , Bth.	731	<i>furfuracea</i> , Kehb.	865
Statice , L.	856	<i>genistoides</i> , J. et Sp.	846
<i>acerosa</i> , Hoh.	834	<i>glauca</i> , Less.	867
<i>acerosa</i> , M. B.	842	<i>globulariaefolia</i> , Desf.	860
<i>acerosa</i> , W.	857	<i>globulifera</i> , B. et Heldr.	869
<i>aciphylla</i> , J. et Sp.	835	<i>glumacea</i> , J. et Sp.	855
<i>Ægyptiaca</i> , Pers.	858	<i>Gmelini</i> , W.	859
<i>alata</i> , W.	858	<i>Græca</i> , Poir.	862
<i>alliacea</i> , Fl. Gr.	873	<i>graminifolia</i> , Ait.	859
<i>androsacea</i> , J. et Sp.	841	<i>hirsuta</i> , Presl	858
<i>aphylla</i> , Forsk.	866	<i>Hohenackeri</i> , J. et Sp.	854
<i>Arabica</i> , J. et Sp.	868	<i>Hohenackeri</i> , Ledeb.	855
<i>aristata</i> , Sm.	870	<i>horrida</i> , Gir.	855
<i>Aucheria</i> , Gir.	861	<i>hyssopifolia</i> , Gir.	862
<i>axillaris</i> , Forsk.	868	<i>hystrix</i> , J. et Sp.	840

	Pag.		Pag.
<i>Statice Jauberti</i> , Gir.	853	<i>Statice Smithii</i> , Ten.	863
<i>Iconica B. et Heldr.</i>	865	<i>Spachii</i> , Gir.	850
<i>incana</i> , M. B.	854	<i>spathulata</i> , Sieb.	861
<i>incana</i> , Vahl	855	<i>speciosa</i> , Forsk.	855
<i>incana</i> , Vis.	855	<i>spicata</i> , W.	871
<i>Karelini</i> , Scz.	847	<i>squamata</i> , Poir.	871
<i>Kotschyi</i> , J. et Sp.	840	<i>Stocksii</i> , Boiss.	868
<i>latifolia</i> , Sm.	864	<i>suffruticosa</i> , L.	867
<i>leptophylla</i> , Schr.	867	<i>Tatarica</i> , L.	854
<i>leptocephala</i> , Boiss.	872	<i>tenuifolia</i> , J. et Sp.	855
<i>lepturoides</i> , J. et Sp.	842	<i>Thouinii</i> , Viv.	858
<i>leucacantha</i> , J. et Sp.	850	<i>Tournefortii</i> , J. et Sp.	841
<i>lilacina</i> , Boiss.	859	<i>Tournefortii</i> , J. et Sp.	862
<i>Limonium</i> , L.	858	<i>tragacanthina</i> , J. et Sp.	849
<i>lyrata</i> , M. B.	872	<i>tripteris</i> , H. Par.	858
<i>macrorhabdos</i> , Boiss.	869	<i>tubiflora</i> , Del.	871
<i>Meyeri</i> , Boiss.	859	<i>tubiflora</i> , R. et Sch.	866
<i>monopetala</i> , L.	874	<i>ulicina</i> , W.	840
<i>nuda</i> , B. et Buhse	864	<i>undulata</i> , B. et Ch.	873
<i>obovata</i> , Ledeb.	867 et Add.	<i>venusta</i> , Fenzl, MSS.	852
<i>ocymifolia</i> , Poir.	861	<i>viminea</i> , Schrad.	863
<i>oleifolia</i> , Sm.	865	<i>virgata</i> , W.	863
<i>Oliveriana</i> , Andr.	862	Staurophragma , F.	
<i>Olivieri</i> , J. et Sp.	852	et M.	361
<i>otolepis</i> , Schr.	866	<i>Natolicum</i> , F. et M.	362
<i>Owerini</i> , Boiss.	870	Stellera , Gmel.	1050
<i>palmaris</i> , Fl. Gr.	862	<i>Altaica</i> , Th.	1050
<i>pauciflora</i> , J. et Sp.	841	<i>Griffithii</i> , Meisn.	1051
<i>perfoliata</i> , C. A. Mey.	866	<i>Hedyotis</i> , Ehr.	1052
<i>plantaginifolia</i> , J. et Sp.	872	<i>Lesserti</i> , Wick.	1051
<i>prolifera</i> , Urv.	862	<i>spicata</i> , C. A. M.	1051
<i>pruinosa</i> , L.	865	<i>stachyoides</i> , Schr.	1051
<i>pungens</i> , J. et Sp.	853	<i>Stemodia littoralis</i> , Vahl	425
<i>pyrenantha</i> , C. Koch	860	<i>muraria</i> , Roxb.	425
<i>Raddiana</i> , Boiss.	860	Striga , Lour.	469
<i>reniformis</i> , Cir.	866	<i>Hermonthica</i> , Del.	469
<i>reticulata</i> , M. B.	865	<i>Strophostoma sparsiflorum</i> ,	
<i>rorida</i> , Fl. Dalm.	865	T.	241
<i>rorida</i> , Fl. Gr.	862	STYRACACEÆ.	54
<i>scoparia</i> , M. B.	859	Styrax , Tourn.	35
<i>Scorpius</i> , J. et Sp.	850	<i>officinale</i> , L.	35
<i>serotina</i> , Rehb.	859	Suæda , Forsk.	957
<i>Sieberi</i> , Boiss.	861	<i>albida</i> , Pall.	626
<i>sinuata</i> , L.	857	<i>altissima</i> , L.	940
<i>sisymbriifolia</i> , J. et Sp.	872	<i>secuata</i> , Bge.	943

	Pag.		Pag.
Suaeda Asphaltica , Boiss.	958	<i>Swertia longifolia</i> , Boiss.	79
baccata , Schimp.	986	<i>obtusa</i> , C. Koch	79
<i>Balansæ</i> , Boiss.	942	<i>obtusa</i> , Led.	78
<i>crassifolia</i> , Pall.	926	<i>perennis</i> , M. B.	78
<i>dendroides</i> , C. A. M.	958	<i>Persica</i> , Griseb.	79
<i>fruticosa</i> , Forsk.	940	<i>punctata</i> , Baumg.	78
<i>fruticosa</i> , L.	959	<i>rotata</i> , Pall.	77
<i>gracilis</i> , Moq.	942	<i>stigmantha</i> , C. Koch	78
<i>heterocarpa</i> , Fenzl	941	<i>Sycomorus antiquorum</i> ,	
<i>heterophylla</i> , Bge.	945	Gusp.	1155
<i>heterophylla</i> , K. et Kir.	945	Symphytum , Tourn.	171
<i>Indica</i> , Thw.	940	<i>Anatolicum</i> , Boiss.	175
<i>Indica</i> , W.	959	<i>aspernum</i> , Bot. Mag.	175
<i>longifolia</i> , C. Koch	942	<i>asperrimum</i> , Urv.	175
<i>maritima</i> , L.	941	<i>asperum</i> , Lep.	175
<i>maxima</i> , C. Koch	941	<i>brachycalyx</i> , Boiss.	175
<i>mierophylla</i> , Pall.	938	<i>brociuum</i> , Fl. Pelop.	176
<i>microsperma</i> , C. A. M.	945	<i>bulbosum</i> , Sch.	176
<i>mollis</i> , Del.	940	<i>Caneasicum</i> , M. B.	172
<i>monantha</i> , C. Koch	942	<i>Clusii</i> , Gmel.	176
<i>monoica</i> , Forsk.	940	<i>cordatum</i> , M. B.	176
<i>monoica</i> , Schimp.	986	<i>echinatum</i> , Led.	175
<i>physophora</i> , Pall.	959	<i>filipendulum</i> , Bisch.	176
<i>platyphylla</i> , Ehr.	986	<i>grandiflorum</i> , D. C.	175
<i>pruinosa</i> , Lange	959	<i>Ibericum</i> , Stev.	176
<i>salsa</i> , Pall.	942	<i>Kurdicum</i> , Boiss. et	
<i>sedifolia</i> , Pall.	926	<i>Haussk.</i>	174
<i>setigera</i> , D. C.	942	<i>macrolepis</i> , Gay	176
<i>splendens</i> , G. et Godr.	942	<i>officinale</i> , L.	171
<i>Transoxana</i> , Bge.	943	<i>Orientale</i> , L.	171
<i>vera</i> , Forsk.	959	<i>Orientale</i> , M. B.	172
<i>vermiculata</i> , Forsk.	940	<i>Ottomanum</i> , Friv.	176
Suchtelenia , Kar.	276	<i>Palestinum</i> , Boiss.	175
<i>acanthocarpa</i> , Kar.	276	<i>peregrinum</i> , Led.	175
<i>calycina</i> , C. A. M.	276	<i>punctatum</i> , Gaud.	176
<i>uniseriatis</i> , Ledeb.	276	<i>racemosum</i> , Steph.	172
Sutera , Roth	422	<i>sepulcrale</i> , B. et Bal.	174
<i>dissecta</i> , Walp.	422	<i>sylvaticum</i> , Boiss.	172
<i>glandulosa</i> , Roth	422	<i>Tauricum</i> , Willd.	172
Vertia , L.	78	<i>tuberosum</i> , L.	175
<i>Aucherii</i> , Boiss.	78	<i>Zegheri</i> , Sch.	176
<i>Balansæ</i> , Boiss.	78	Syringa , L.	78
<i>Carinthiaca</i> , Wulf.	77	<i>Persica</i> , L.	78
<i>Iberica</i> , Fisch.	78	<i>vulgaris</i> , L.	58
<i>lactea</i> , Bge.	79		

	Pag.		Pag.
Tapeinanthus , Boiss.	679	Tenerium nivale , B.	808
Persiens, Boiss.	679	<i>Nuchense</i> , C. Koch	816
Brahniens, Boiss.	680	<i>Odontites</i> , B. et Bal.	815
Tecoma , Juss.	80	<i>Oliverianum</i> , Ging.	810
undulata, Roxb.	80	<i>Orientale</i> , L.	808
Teucrium , L.	805	<i>pæderotoides</i> , B. et Haussk.	814
alpestre, Fl. Gr.	819	<i>palustre</i> , Lam.	813
<i>Arduini</i> , Fl. Gr.	811	<i>Pannonicum</i> , Kern.	819
Aroanium, Orph.	807	<i>parviflorum</i> , Schreb.	810
brevifolium, Schreb.	807	<i>Persicum</i> , Boiss.	815
canum, F. et M.	816	<i>Pestalozzae</i> , Boiss.	807
canum var. Bth.	817	<i>pictum</i> , B. et Haussk.	809
capitatum, Fl. Gr.	821	<i>Polium</i> , L.	821
Chamaedrys, L.	816	<i>Polium</i> var.	822
Chamaepitys, L.	802	<i>procernum</i> , B. et Bl.	809
Charamoniense, Cav.	806	<i>pruinatum</i> , Boiss.	808
Chium, Poir.	802	<i>pseudohyssopus</i> , Sch.	821
coniortodes, B. et Bl.	818	<i>quadratulum</i> , Fl. Gr.	817
Creticum, Lam.	807	<i>regium</i> , Auct.	817
Creticum, L.	806	<i>rigidum</i> , Bth.	809
cuneifolium, Fl. Gr.	820	<i>rosmarinifolium</i> , Lam.	806
cuneifolium, Un. It.	820	<i>salicifolium</i> , L.	800
Cyprium, Boiss.	820	<i>scordioides</i> , Schreb.	815
densiflorum, Vis.	811	<i>Scordium</i> , L.	812
divaricatum, Sieb.	816	<i>Scorodonia</i> , Sm.	812
flavum, L.	816	<i>serratum</i> , Bth.	805
flavum var. Bth.	817	<i>Sibiricum</i> , L.	668
Hyreanicum, L.	811	<i>Sinaicum</i> , Boiss.	822
hyssopifolium, Sch.	806	<i>Smyrnium</i> , Boiss.	812
Kotschyanum, Pœch.	812	<i>Socinianum</i> , Boiss.	818
lævigatum, Russ.	804	<i>spinosum</i> , L.	814
lamifolium, Urv.	811	<i>Stocksianum</i> , Boiss.	821
lauviginosum, H. et Link	815	<i>supinum</i> , L.	819
Laxmanni, L.	804	<i>Syspirense</i> , C. Koch	810
leucocladum, Boiss.	820	<i>Taylori</i> , Boiss.	809
leucophyllum, M. et Auch.	817	<i>Theligonum Cynocrambe</i> , L.	897
lucidum, Fl. Gr.	817	Thesium , L	1059
maerum, B. et Haussk.	810	<i>Alpinum</i> , L	1061
Mascatense, Boiss.	819	<i>asperulum</i> , B. et Buhse	1062
Massiliense, L.	812	<i>aureum</i> , J. et Sp.	1060
malisoïdes, B. et Haussk.	815	<i>Bergeri</i> , Zucc.	1063
mierophyllum, Desf.	817	<i>Billardieri</i> , Boiss.	1064
montanum, L.	818	<i>brachiphyllum</i> , Boiss.	1062
Montbreti, Bth.	814	<i>Byzantinum</i> , A.D.C.	1061
multicaule, M. et Auch.	807		

Pág.		Pág.	
Thesium compressum, B. et Heldr.	1064	Thymus <i>Borei</i> , Bth.	556
divaricatum, Juss.	1061	<i>capitatus</i> , L.	560
<i>elegans</i> , Roch.	1059	<i>Cappadocius</i> , Boiss.	558
<i>Græcum</i> , B. et Sp.	1065	<i>Chamædrys</i> , Fr.	555
<i>Græcum</i> , Zucc.	1055	<i>Chaubardi</i> , B. et Heldr.	556
<i>heterophyllum</i> , Boiss.	1064	<i>cherlerioides</i> , Vis.	557
<i>humifusum</i> , Gr.	1061	<i>Ciliciensis</i> , B. et Bal.	559
<i>humile</i> , Vahl	1064	<i>collinus</i> , M. B.	555
<i>impressum</i> , St.	1066	<i>comptus</i> , Friv.	558
<i>intermedium</i> , Schr.	1095	<i>Creticus</i> , Brot.	560
<i>Kotschyanum</i> , Boiss.	1066	<i>decaenatus</i> , Bth.	559
<i>Libanoticum</i> , Ehr.	1065	<i>exiguus</i> , Fl. Gr.	585
<i>linophyllum</i> , Griseb.	1065	<i>fallax</i> , F. et M.	556
<i>maeranthum</i> , F	1067	<i>filiformis</i> , Sieb.	569
<i>maritimum</i> , C. A. Mey.	1066	<i>glabratus</i> , B. et Ch.	556
<i>montanum</i> , Ehr.	1065	<i>glaucus</i> , Friv.	558
Parnassi, A.D.C.	1062	<i>graveoleus</i> , Fl. Gr.	554
<i>pratense</i> , Led.	1062	<i>graveolens</i> M. B.	585
<i>procumbens</i> , C.A.M.	1065	<i>heterotrichus</i> , Gr.	556
<i>ramosum</i> , Griseb.	1062	<i>hirsutus</i> , M. B.	557
<i>ramosum</i> , Hayne	1062	<i>hirtus</i> , Viv.	652
<i>stelleroides</i> , J. et Sp.	1060	<i>icanus</i> , Fl. Gr.	578
Szovitsii, A.D.C.	1066	<i>inodorus</i> , Friv.	570
Tauricolum, B. et Haussk.	1067	<i>integer</i> , Griseb.	560
Thymbra , L.	561	<i>Kotschyanus</i> , B. et Hoh.	556
<i>ambigua</i> , Cl.	561	<i>lanceolatus</i> , Sm. Prodr.	556
<i>capitata</i> , Gr.	560	<i>lanceolatus</i> var. <i>angusti-</i>	
<i>hirsuta</i> , Pers.	568	<i>folius</i> , Bth.	559
<i>spicata</i> , L.	561	<i>Marschallianus</i> , W.	555
<i>verticillata</i> , L.	561	<i>Mastichina</i> , Sm.	569
THYMELÆACEÆ.	1046	<i>microphyllus</i> , Urv.	572
Thymelæa , Tourn.	1053	<i>montanus</i> , W. K.	555
<i>Cilicica</i> , Meissn.	1053	<i>nepetoides</i> , Don.	644
<i>hirsuta</i> , L.	1054	<i>nummularius</i> , M. B.	555
<i>Tartonraira</i> , L.	1053	<i>Ocheus</i> , H. et Sart.	555
<i>tomentosa</i> , Endl.	1054	<i>odoratissimus</i> , M. B.	558
Thymus , L.	554	<i>Pannonicus</i> , Bth.	555
<i>Acinos</i> , L.	582	<i>parvifolius</i> , C. Koch	556
<i>Alpinus</i> , L.	581	<i>pechinatus</i> , F. et M.	558
<i>augustifolius</i> , Pers.	556	<i>pruinosis</i> , Boiss.	558
<i>Argæus</i> , B. et Bal.	556	<i>pulchellus</i> , C. A. M.	555
<i>Balansæ</i> , Boiss.	556	<i>punctatus</i> , W.	585
<i>Billardieri</i> , Boiss.	560	<i>punctatus</i> , Vis.	556
		<i>quadranjulus</i> , Ehr.	559
		<i>rarifolios</i> , C. Koch	556

	Pag.		Pag.
<i>Thymus rigidus</i> , Sch. et Ky.	557	<i>Trixago rhinanthina</i> , Link	474
<i>Rægneri</i> , C. Koch	555	<i>viscosa</i> , Rehb.	474
<i>Serpillum</i> , L.	554		
<i>Sibthorpii</i> , Bth.	556	Ulmus , L.	1157
<i>Sipyleus</i> , Boiss.	556	<i>campestris</i> , L.	1157
<i>squarrosum</i> , F. et M.	557	<i>ciliata</i> , Ehrh.	1158
<i>striatus</i> , Vahl	557	<i>effusa</i> , W.	1158
<i>suaveolens</i> , Sm.	582	<i>elliptica</i> , C. Koch	1158
<i>Syriacus</i> , Boiss.	559	<i>glabra</i> , Mill.	1157
<i>Syrticus</i> , Spr.	652	<i>major</i> , Sm.	1158
<i>tenerioides</i> , B. et Sprun.	554	<i>montana</i> , Sm.	1158
<i>Tragoriganum</i> , L.	568	<i>nuda</i> , Ehrh.	1157
<i>vilosus</i> , Fl. Gr.	560	<i>octandra</i> , Schk.	1158
<i>vulgaris</i> , Sm. Pr.	557	<i>pedunculata</i> , Fenzl.	1158
<i>xylorrhizus</i> , Boiss.	556	<i>polygama</i> , Rich.	1159
<i>zygoides</i> , Gr.	557	<i>tuberosa</i> , Ehrh.	1157
<i>zygis</i> , Fl. Gr.	557		1145
<i>zygis</i> , Marsch.	558	URTICACEÆ.	
Tournefortia , L.	125	Urtica , L.	1146
<i>Arguzia</i> , L.	125	<i>Balearica</i> , L.	1147
<i>micranthos</i> , A. D. C.	158	<i>bicolor</i> , Roxb.	1146
Trachelanthus , Kze.	271	<i>cannabina</i> , L.	1147
<i>cerinthoides</i> , Boiss.	271	<i>caudata</i> , Vahl	1147
<i>Kurdieus</i> , Ky.	271	<i>dioica</i> , L.	1146
<i>Trachystemon Orientale</i> , Don.	177	<i>Chesneyana</i> , Bl.	1147
Traganum , Del.	946	<i>Haussknechti</i> , Boiss.	1146
<i>nudatum</i> , Schimp.	985	<i>hippuroides</i> , Griff.	1148
<i>nudatum</i> , Del.	946	<i>hispiata</i> , D. C.	1146
<i>Tragopyrum buxifolium</i> ,		<i>membranacea</i> , Poir.	1147
M. B.	1025	<i>microphylla</i> , B. et	
<i>grandiflorum</i> , M. B.	1024	Haussk.	1147
<i>rotundifolium</i> , Pr.	1021	<i>pentandra</i> , Roxb.	1148
Trichodesma , R. Br.	280	<i>pilulifera</i> , L.	1147
<i>Africanum</i> , L.	280	<i>pubescens</i> , Led.	1146
<i>Aucherii</i> , D. C.	282	<i>submitis</i> , Boiss.	1146
<i>Ehrenbergii</i> , Schw.	281	<i>urens</i> , L.	1146
<i>ineanum</i> , Bge.	282		
<i>Indicum</i> , L.	280	Utricularia , L.	
<i>molle</i> , D. C.	281	<i>inflexa</i> , Forsk.	5
<i>Stocksii</i> , Boiss.	282	<i>stellaris</i> , L. fil.	5
<i>Triguera baccata</i> , Dun.	551	<i>stellaris</i> , Willd.	5
Trixago , Stev.	474	<i>vulgaris</i> , L.	4
<i>Apula</i> , St.	474		
<i>carnea</i> , Griseb.	474	Vandellia , L.	427
<i>purpurea</i> , St.	475	<i>erecta</i> , Bth.	427
		<i>pyxidaria</i> , All.	427
		Verbascum , L.	298

	Pag.		Pag.
<i>Verbascum Adeliæ, Heldr.</i>	521	<i>Verbascum chrysorrhæcos,</i> <i>Boiss.</i>	540
<i>adenocaulon, B. et Bal.</i>	556	<i>Ciliciicum, Boiss.</i>	557
<i>adenophorum, Boiss.</i>	543	<i>Ceccarinianum, B. et</i> <i>Heldr.</i>	522
<i>alecoides, B. et Haussk.</i>	506	<i>compactum, M. B.</i>	506
<i>Aleppense, Bth.</i>	542	<i>crispum, Ehr.</i>	541
<i>alpestre, Boiss.</i>	517	<i>cuspidatum, Schrad.</i>	501
<i>alpigenum, C. Koch</i>	540	<i>cylindrocarpum, Gris.</i>	557
<i>alyssifolium Boiss.</i>	545	<i>Daenense, Boiss.</i>	518
<i>Amanum, Boiss.</i>	510	<i>Datmaceum, Boiss.</i>	528
<i>Antinori, B. et Heldr.</i>	516	<i>Delphicum, B. et Heldr.</i>	524
<i>Antiochium, Boiss.</i>	510	<i>denudatum, B. et Heldr.</i>	554
<i>Arcturus L.</i>	532	<i>detersile, B. et Heldr.</i>	516
<i>Armeniaeum, Huet</i>	561	<i>digitalifolium, B. et</i> <i>Haussk.</i>	542
<i>Armenum, B. et K.</i>	504	<i>diversifolium, Hochst.</i>	549
<i>asperulum, Boiss.</i>	525	<i>elegans, B. et Heldr.</i>	521
<i>astrophyllum, B. et Heldr.</i>	516	<i>epixanthinum, B. et</i> <i>Heldr.</i>	504
<i>auriculatum, Fl. Gr.</i>	515	<i>erianthum, Bth.</i>	519
<i>australe, Schr.</i>	502	<i>erioorrhaddon, Boiss.</i>	517
<i>Bannaticum, Gris.</i>	550	<i>Euphraticum, B. nth.</i>	545
<i>Barradense, Boiss.</i>	518	<i>fasciculatum, Ehr.</i>	518
<i>Basianicum, B. et Haussk.</i>	541	<i>ferrugineum, And.</i>	546
<i>Berytheum, Boiss.</i>	544	<i>floccosum, W. K.</i>	525
<i>betonicæfolium, Desf.</i>	506	<i>fœtidum, B. et Heldr.</i>	503
<i>Biebersteinii, Bess.</i>	524	<i>formosum, Fisch.</i>	506
<i>bipinnatifidum, Bot. Mag.</i>	512	<i>Gaillardotii, Boiss.</i>	522
<i>bisentellæfolium, Bth.</i>	546	<i>Gahlienii, Boiss.</i>	513
<i>Bithynicum, Boiss.</i>	545	<i>Garganicum var. Gr.</i>	505
<i>Blancheanum, Boiss.</i>	545	<i>geminiflorum, Hochst.</i>	549
<i>Blattaria, L.</i>	568	<i>Georgicum, Bth.</i>	506
<i>Boerrhavii, L.</i>	515	<i>Germanieæ, Haussk.</i>	554
<i>bombyciforme, Boiss.</i>	505	<i>glabratum, Friv.</i>	529
<i>brevidens, Bth.</i>	519	<i>globitorum, B. et Noë</i>	552
<i>bulbosum, Ehr.</i>	545	<i>glomeratum, Boiss.</i>	509
<i>Cæsareum, Boiss.</i>	514	<i>gnaphalodes, M. B.</i>	516
<i>Calaurium, Heldr.</i>	515	<i>gossypinum, M. B.</i>	548
<i>calvum, B. et Ky.</i>	558	<i>Græcum, Heldr. et Sart.</i>	556
<i>campestre, B. et Heldr.</i>	502	<i>Guicciardi, B. et Heldr.</i>	505
<i>candidissimum, D. C.</i>	515	<i>Heldreichii, Boiss.</i>	526
<i>Cedreti, Boiss.</i>	528	<i>heteromallum, Pare.</i>	505
<i>ceratophyllum, Schrad.</i>	512	<i>Hinkei, Friv.</i>	529
<i>cerinum, B. et Heldr.</i>	559	<i>Hohenackeri, F. et M.</i>	545
<i>cestroides, B. et Haussk.</i>	551		
<i>cheiranthifolium, Boiss.</i>	525		
<i>Chium, B. et Orph.</i>	509		
<i>chrysanthemum, Schrad.</i>	548		

	Pag.		Pag.
<i>Verbascum humile, Janka</i>	326	<i>Verbascum napifolium, Boiss.</i>	353
<i>hyoscyridifolium, Schrad.</i>	325	<i>nigrum, L.</i>	328
<i>hypoleucum, B. et Heldr.</i>	329	<i>obovatum, Hochst.</i>	349
<i>infidellum, B. et Haussk.</i>	342	<i>Olympicum, Boiss.</i>	323
<i>insigne, B. et Bal.</i>	508	<i>oreophilum, C. Koch</i>	349
<i>insulare, B. et Heldr.</i>	307	<i>orgyale, B. et Heldr.</i>	524
<i>Isauricum, B. et Heldr.</i>	555	<i>Orientale, M. B.</i>	550
<i>Khorassanicum, Boiss.</i>	319	<i>Osbeckii, L.</i>	351
<i>kochiaeforme, B. et</i>		<i>ovalifolium, Don.</i>	506
<i>Haussk.</i>	555	<i>oxycarpum, C. Koch</i>	317
<i>Köchelii, Fenzl</i>	544	<i>pannosum, Vis. et Pan.</i>	304
<i>Kotschyi, Boiss.</i>	511	<i>parviflorum, Lam.</i>	553
<i>lætum, B. et Haussk.</i>	358	<i>Pestalozzae, Boiss.</i>	348
<i>lagurus, F. et M.</i>	302	<i>petiolare, B. et Ky.</i>	527
<i>lasianthum, Boiss.</i>	319	<i>phœniceum, L.</i>	546
<i>leianthum, Benth.</i>	538	<i>phlomoides, L.</i>	501
<i>leptocladum, B. et Heldr.</i>	311	<i>phlomoides, Ft. Gr.</i>	514
<i>leuconeuron, B. et Heldr.</i>	344	<i>phlomoides, Urv.</i>	319
<i>leucophyllum, Griseb.</i>	520	<i>phyllostachyum, B. et</i>	
<i>longifolium, d.c.</i>	525	<i>Haussk.</i>	551
<i>longifolium, Ten.</i>	504	<i>pilematophoram, C. Koch</i>	518
<i>Lychnitis, L.</i>	324	<i>Pinardi, Boiss.</i>	559
<i>Lydium, Boiss.</i>	514	<i>pinnatifidum, Vahl</i>	512
<i>lyratifolium, Koch.</i>	544	<i>Pisidicum, B. et Heldr.</i>	525
<i>lyratum, Lam.</i>	552	<i>plieatum, Fl. Gr.</i>	512
<i>macrocarpum, Boiss.</i>	508	<i>protractum, Fenzl</i>	310
<i>macrophyllum, C. Koch</i>	529	<i>Prusianum, Boiss.</i>	505
<i>macrophyllum, Ky.</i>	515	<i>ptychophyllum, Boiss.</i>	515
<i>macrosepalum, B. et Ky.</i>	517	<i>pulverulentum, M. B.</i>	524
<i>macrurum, Ten.</i>	501	<i>pulverulentum, Vill.</i>	522
<i>maiale, d.c.</i>	514	<i>pumilum, B. et Heldr.</i>	520
<i>malacotrichum, B. et</i>		<i>Punaleuse, B. et Buhse</i>	507
<i>Heldr.</i>	505	<i>pycnostachyum, B. et</i>	
<i>mallophorum, B. et Heldr.</i>	523	<i>Heldr.</i>	517
<i>megaphlomos, Boiss.</i>	525	<i>pyramidalatum, M. B.</i>	540
<i>Mesopotamicum, Boiss.</i>	552	<i>racemiferum, B. et</i>	
<i>microcalycinum, Fenzl</i>	538	<i>Haussk.</i>	552
<i>microcarpum, Benth.</i>	509	<i>Rascheyanum, Boiss.</i>	328
<i>millelacum, Boiss.</i>	318	<i>repandum, W.</i>	508
<i>molle, C. Koch</i>	506	<i>rigidum, B. et Heldr.</i>	515
<i>montanum, Griseb.</i>	502	<i>ringens, Boiss.</i>	510
<i>monticolum, Boiss.</i>	525	<i>rubricaulis, B. et Heldr.</i>	555
<i>mucronatum, Lam.</i>	315	<i>rubiginosum, W. K.</i>	546
<i>myrianthum, Boiss.</i>	550	<i>saccatum, C. Koch</i>	506
<i>myriocarpum, B. et Heldr.</i>	527	<i>salviæfolium, Boiss.</i>	510

Pag.	Pag.
Verbascum Samaritanii, Heldr. 504	Verbena officinalis, L. 554
Sartorii, B. et Heldr. 502	<i>procumbens</i> , Forsk. 554
scaposum, Boiss. 547	<i>spina</i> , L. 554
Schimperianum, Boiss. 541	Veronica L. 454
Schneideri, Boiss. 517	<i>acinifolia</i> , L. 458
Schraderi, Mey. 501	<i>agrestis</i> , L. 466
Seytunense, Boiss. 525	<i>agrestis var. Byzantina</i> , Fl. Gr. 466
simplex, Lab. 511	Albanica C. Koch 462
Sinaiticum, Bth. 518	Aleppica , Boiss. 445
sinuatum, L. 522	amena , Stev. 462
Smyrneum, Boiss. 557	Anagallis , L. 457
speciosum, Schrad. 528	<i>anagaloides</i> , Guss. 457
spectabile, M. B. 307	<i>anisophylla</i> , C. Koch 445
sphenandroides, C. Koch 550	<i>aphylla</i> , L. 450
spinosum, L. 547	Armena , B. et Huet 441
splendidum, Boiss. 510	arvensis , L. 457
stachydifolium, B. et Heldr. 331	Aucheri, Boiss. 441
stachydiforme, B. et Buhse 519	Austriaca , L. 449
stenocarpum, B. et Heldr. 527	bartsiaefolia , Boiss. 464
stenophyllum , B. et Heldr. 526	Beccabunga , Boiss. 458
Sievenii, Boiss. et Buhse 529	bellidioides , L. 452
strictum, Cl. 549	Benthami , C. Koch 459
subnivale, B. et Haussk. 505	Billardieri , Vahl 445
Syriacum, Schrad. 511	biloba , L. 464
Szovitsianum, Boiss. 553	bombycinia , B. et Ky. 454
taraxacifolium, Boiss. 553	brevifolia , M. B. 455
Tauri, B. et Ky. 521	Bungacea , Janka 455
thapsiforme, Schrad. 501	Bungei , Boiss. 465
Thapsus, L. 501	Buxbaumiana , Pall. 451
Tiberiadis, Boiss. 341	Buxbaumii , Ten. 465
Triopolitanum, Boiss. 522	Calverti , Boiss. 450
triste, Sm. 546	campylopoda , Boiss. 464
undulatum, Lam. 512 et 515	Cariensis , Boiss. 450
vulcanicum, B. et Heldr. 514	Caucasica , M. B. 440
Wiedemannianum, F. <i>et M.</i> 507	ceratocarpa , C. A. M. 460
Wierzbickii, Heuff. 529	cespitosa , Boiss. 454
Wilhelmsianum, C. Koch 550	chamaedryoides , Fl. Pelop. 447
xanthophenicum, Gr. 547	chamaedrys , L. 446
VERBENACEÆ. 551	chamaepitys , Gr. 456
Verbena , L. 554	Chaubardi , B. et R. 461
<i>nodiflora</i> , L. 552	cinerrea , B. et Bat. 446
	conferta , Boiss. 459
	Cretica , Pall. 444

	Pag.		Pag.
<i>Veronica Crista-Galli, St.</i>	468	<i>Veronica maritima, M. B.</i>	455
<i>cuneifolia, Don.</i>	445	<i>maxima, Stev.</i>	447
<i>cymbalaria, Bod.</i>	467	<i>melissæfolia, Desf.</i>	447
<i>cymbalariafolia, Vahl</i>	467	<i>Michauxii, Lam.</i>	459
<i>cymbalarioides, Bl.</i>	468	<i>microcarpa, Boiss.</i>	441
<i>dentata, Schm.</i>	449	<i>microtheca, B. et Bal.</i>	464
<i>depauperata, W. K.</i>	450	<i>minuta, C. A. Mey.</i>	451
<i>dichrus, Schott.</i>	445	<i>monticola, Trautv.</i>	452
<i>didyma, Ten.</i>	466	<i>montoides, Boiss.</i>	457
<i>digitata, Lam.</i>	463	<i>multifida, L.</i>	442
<i>digitalis, Auct.</i>	456	<i>multifida, Jacq.</i> 449 et Add.	
<i>divaricata, B. et Bal.</i>	451	<i>nigricans, C. Koch</i>	459
<i>elatior, M. B.</i>	455	<i>Noëana, Boiss.</i>	445
<i>erinoïdes, Boiss.</i>	455	<i>nudiventris, Kar. et Kir.</i>	458
<i>euphrasiæfolia, Link</i>	450	<i>officinalis, L.</i>	451
<i>exilis, Schott.</i>	464	<i>Olympica, Pall.</i>	451
<i>filiformis, D.C.</i>	466	<i>opaca, Fries.</i>	467
<i>filiformis, Sm.</i>	466	<i>orbicularis, Fisch.</i>	450
<i>foliosa, W. K.</i>	455	<i>Orchidea, Cr.</i>	455
<i>fragilis, B. et Haussk.</i>	446	<i>Orientalis, Mill.</i>	445
<i>Galathaea, Boiss.</i>	448	<i>Ossetica, Stev.</i>	440
<i>gentianoides, Vahl</i>	451	<i>oxycarpa, Boiss.</i>	458
<i>glaberrima, B. et Bal.</i>	459	<i>Pæderotæ, Boiss.</i>	454
<i>glauca, Fl. Gr.</i>	461	<i>pallida, Horn.</i>	451
<i>Gorumsensis, B. et Ky.</i>	458	<i>paniculata, L.</i>	455
<i>Græca, Sprun.</i>	461	<i>Panormitana, Tin.</i>	467
<i>grandiflora, Don.</i>	462	<i>parviflora, Vahl</i>	445
<i>Griffithii, Bth.</i>	468	<i>pectinata, L.</i>	442
<i>Haussknechtii, Boiss.</i>	458	<i>peduncularis, M. B.</i>	459
<i>hederæfolia, L.</i>	468	<i>peduncularis, Fl. Pel.</i>	462
<i>hispidula, B. et Huet</i>	460	<i>pedunculata, Lab.</i>	460
<i>hospita, M. K.</i>	466	<i>Peloponnesiaca, B. et</i>	
<i>hybrida, M. B.</i>	455	<i>Orph.</i>	462
<i>Jacquini, R. et Sch.</i>	449	<i>perpusilla, Boiss.</i>	458
<i>ineana, L.</i>	456	<i>Persica, Poir.</i>	466
<i>ixodes, B. et Bal.</i>	459	<i>petrea, Stev.</i>	440
<i>Kotschyana, Bth.</i>	452	<i>phænicantha, C. Koch</i>	459
<i>Kurdica, Bth.</i>	445	<i>pinnatifida, Lam.</i>	457
<i>latifolia, L.</i>	449	<i>polifolia, Benth.</i>	444
<i>leiocarpa, Boiss.</i>	448	<i>politæ, Fries.</i>	466
<i>Liwanensis, C. Koch</i>	450	<i>præcox, All.</i>	465
<i>longifolia, L.</i>	455	<i>prostrata, Sm. Pr.</i>	449
<i>lysimachioïdes, Boiss.</i>	458	<i>pseudochomædrys, Jacq.</i>	419
<i>macropoda, Boiss.</i>	465	<i>pulchella, Bast.</i>	467
<i>macrostachya, Vahl</i>	444	<i>pumila, Cl.</i>	469

	Pag.		Pag.
Veronica punctata , Ham.	459	Vincetoxicum , Mœnch	56
<i>pusilla</i> , Bth.	457	<i>Alpinum</i> , Sch.	56
<i>recta</i> , Bth.	449	<i>amplifolium</i> , C. Koch	54
<i>reticulata</i> , C. Koch	461	<i>Anatolicum</i> , Boiss.	52
<i>Reuteriana</i> , Boiss.	459	<i>Arnottianum</i> , Wight	52
<i>Romana</i> , All.	458	<i>canescens</i> , Willd	52
<i>rubrifolia</i> , Boiss.	465	<i>funebre</i> , Boiss.	53
<i>Sartoriana</i> , B. et Heldr.	457	<i>fuseatum</i> , Rehb. fil.	53
<i>Scardica</i> , Gr.	459	<i>latifolium</i> , C. Koch.	53
<i>secundiflora</i> , C. Koch	459	<i>luridum</i> , Stocks	52
<i>serpyllifolia</i> , L.	455	<i>luteum</i> , Sieb.	52
<i>spicata</i> , L.	455	<i>medium</i> , Decaisne	53
<i>spuria</i> , L.	455	<i>nigrum</i> , L.	53
<i>spuria</i> , Sm.	455	<i>nivale</i> , Boiss. et Heldr.	53
<i>stenobotrys</i> , B. et Bl.	447	<i>officinale</i> , Mœnch	54
<i>succulenta</i> , All.	457	<i>parviflorum</i> , Decaisne	56
<i>sureculosa</i> , B. et Bal.	445	<i>Pisidicum</i> , B. et Heldr.	56
<i>Syriaca</i> , R. et Sch.	460	<i>pumilum</i> , Decaisne	54
<i>Syspirensis</i> , C. Koch	445	<i>Rehmanni</i> , Boiss.	53
<i>Taurica</i> , Stev.	445	<i>speciosum</i> , Boiss. et Spr.	53
<i>telephylloides</i> , Vahl	450	<i>stelliflorum</i> , Boiss.	56
<i>tenuifolia</i> , M. B.	442	<i>Thirkeanum</i> , C. Koch	53
<i>tenuis</i> , Led.	457	<i>Tmoleum</i> , Boiss.	52
<i>teucrioides</i> , Boiss. et Heldr.	444	<i>triste</i> , Griseb.	53
<i>Teucrium</i> , L.	448	<i>undulatum</i> , Heldr.	53
<i>Thessalica</i> , Benth.	455	Viscum , L.	1068
<i>thymifolia</i> , Fl. Gr.	444	<i>album</i> , L.	1068
<i>Tournefortii</i> , Guel.	466	<i>erectum</i> , Sieb.	1068
<i>trifolios</i> , L.	465	<i>Orientale</i> , D. C.	1068
<i>Tymphrestea</i> , Boiss.	444	<i>Oxycedri</i> , D. C.	1069
<i>umbrosa</i> , M. B.	440	Vitex , L.	553
<i>urticifolia</i> , L.	448	<i>Agnuscastus</i> , L.	553
<i>verna</i> , L.	456	<i>Negundo</i> , L.	555
<i>Vestiana</i> , R. et Sch.	447	Vogelia , L.	876
<i>viscosa</i> , Boiss.	457	<i>Arabica</i> , Boiss.	876
<i>Villarsia Indica</i> , Vent.	65	<i>Indica</i> , Wight	876
Vinca , L.	43	<i>Volkameria Acerbiana</i> , Vis.	556
<i>Bottae</i> , J. et Sp.	46	Wiedemannia , F. et M.	767
<i>herbacea</i> , W. K.	45	<i>erythrotricha</i> , B.	767
<i>Libanotica</i> , Zucc.	46	<i>multifida</i> , L.	767
<i>major</i> , L.	45	<i>Orientalis</i> , F. et M.	767
<i>minor</i> , L.	45	Withania , Panq.	287
<i>pubescens</i> , Urv.	45	<i>coagulans</i> , Stocks	288
<i>sessilifolia</i> , D. C.	46	<i>somnifera</i> , L.	287

INDEX.

	Pag.		Pag.
Wulfenia , Jacq.	433	Zizyphora <i>capitata</i> , <i>L.</i>	586
<i>Orientalis</i> , Boiss.	433	<i>cleonioides</i> , Boiss.	588
Yermoloffia hispida , Bel.	769	<i>elinopodiooides</i> , <i>M. B.</i>	585
<i>insignis</i> , Bel.	770	<i>compacta</i> , Friv.	586
Zappania nodiflora , Lam.	553	<i>dasyantha</i> , <i>M. B.</i>	586
Zataria , Boiss.	561	<i>fasciculata</i> , <i>C. Koch</i>	586
<i>braceleata</i> , Boiss.	562	<i>Gundelsheimeri</i> , <i>C. Koch</i>	586
<i>humilis</i> , Bth.	667	<i>Hispanica</i> , <i>L.</i>	586
<i>multiflora</i> , Boiss.	561	<i>lanceolata</i> , Boiss.	
Zelkoua , Sp.	1158	<i>nummularia</i> , <i>F. et M.</i>	386
<i>Abelicea</i> , <i>Fl. Gr.</i>	1159	<i>Persica</i> , <i>Bge.</i>	587
<i>crenata</i> , <i>Desf.</i>	1159	<i>pungens</i> , <i>Bge.</i>	666
Zietenia Orientalis , Gl.	743	<i>serpyllacea</i> , <i>M. B.</i>	585
Zizyphora , L.	585	<i>Taurica</i> , <i>M. B.</i>	587
<i>acutifolia</i> , <i>M. et Auch.</i>	587	<i>tenuior</i> , <i>L.</i>	587
<i>Brantii</i> , <i>C. Koch</i>	586	Zornia peltata , <i>Mœnch</i>	674
<i>canescens</i> , Bth.	585	Zwackhia , <i>Sendt.</i>	222
		<i>aurea</i> , <i>Sendtn.</i>	223



